

Contents

| | | |
|----------|--|-----------|
| 1 | Appendix A: Individually modifiable behaviours covered by the evidence review | 2 |
| 2 | Appendix B: Methods | 5 |
| 2.1 | Systematic searches | 5 |
| 2.1.1 | Stage 1 search: Broad search for systematic reviews | 5 |
| 2.1.2 | Stage 2 search: Targeted search for systematic reviews | 7 |
| 2.1.3 | Stage 3 search: Primary study searches | 8 |
| 2.1.4 | Sample search strategies | 8 |
| 2.2 | Selecting studies for inclusion | 13 |
| 2.2.1 | First pass appraisal | 17 |
| 2.2.2 | Second pass appraisal | 17 |
| 2.2.3 | Systematic review mapping | 19 |
| 2.2.4 | Full text appraisal | 20 |
| 2.2.5 | Systematic reviews generically assessing overweight and obesity prevention | 21 |
| 2.3 | Quality appraisal | 21 |
| 2.4 | Data extraction | 22 |
| 3 | Appendix C: Excluded study bibliography | 24 |
| 4 | Appendix D: Quality appraisal templates | 43 |
| 5 | Appendix E: Summary table of non-prioritised reviews | 49 |
| 6 | Appendix F: Evidence tables | 49 |

1 **Appendix A: Individually modifiable behaviours covered by the evidence review**

| Food and drinks | Energy and nutrients | Eating Patterns | Physical activity and exercise | Sedentary behaviour | Other factors |
|-----------------------------|--|---|---|---|---|
| 1. High energy dense foods | 1. Fat/protein/carbohydrate | 1. Eating speed /rate | 1. Active leisure/recreation | 1. Amount of sedentary time | 1. Sleep (amount and quality) |
| 2. (Low) energy dense foods | 2. Glycaemic index/ glycaemic load | 2. Eating/meal/snack frequency (eating occasions) | 2. Activities of daily living (e.g. housework, garden, DIY) | 2. Sitting | 2. Monitoring – e.g. weight, waist, clothes fit, pedometers, food diaries |
| 3. Whole grain | 3. Fibre | 3. Eating pattern - consistency, weekend vs. weekdays, energy intake split/timing through the day | 3. Incidental physical activity-(active habits) e.g. stairs | 3. Standing | 3. Over holiday/Christmas weight gain |
| 4. Refined grains | 4. Calorie control (watching what you eat) | 4. Setting or distractions (e.g. table vs. TV viewing) | 4. Walking (including steps – travel or leisure) | 4. TV and other screen - time, eating and viewing, displaced activity | 4. Stress minimising activities |
| 5. Fruit and vegetables | 5. Energy density | 5. Family meal (+eating with children) | 5. Sport | 5. Other sedentary activities – reading / commuting | 5. Support e.g. partner, social support, buddy |
| 6. Meat and fish | 6. Artificial sweeteners | 6. Portion size | 6. Active play (e.g. after school) | 6. Breaks in sedentary time (e.g. workplace breaks such as | 6. Avoiding screen advertising (e.g. advert free versus advert containing |

Bazian.

| Food and drinks | Energy and nutrients | Eating Patterns | Physical activity and exercise | Sedentary behaviour | Other factors |
|--|--|-------------------------------|--|---|---------------|
| | | | | meeting breaks, walking breaks) | viewing) |
| 7. Milk and other dairy | 7. Sugar, high fructose corn syrup, sucrose, glucose | 7. Snacking/snacks | 7. Active travel/commuting | 7. More active screen time (active versus passive gaming) | |
| 8. Nuts/legumes | 8. Caffeine | 8. Grazing/gorging | 8. Cycling (travel or leisure) | | |
| 9. Dietary pattern – specific combination of foods | 9. Catechins | 9. Eating out | 9. Strength/aerobic | | |
| 10. Vegan / vegetarian | | 10. Take away meals/fast food | 10. Intensity, time, frequency (total volume) 11. Intensity (same volume, high intensity vs. low intensity) | | |
| Drinks: 11. Sugar sweetened drinks 12. Fruit juice 13. Water 14. Alcohol – wine, beer, spirits 15. Tea and coffee 16. Artificially sweetened beverages | | 11. Meal planning | | | |

Bazian

| Food and drinks | Energy and nutrients | Eating Patterns | Physical activity and exercise | Sedentary behaviour | Other factors |
|--|----------------------|--|--------------------------------|---------------------|---------------|
| 17. Low-calorie drinks/Low-sugar drinks/Sugar-reduced-drinks | | | | | |
| | | 12. Meal skipping (including breakfast skipping) | | | |
| | | 13. Drinks with meals | | | |
| | | 14. Breakfast | | | |

2 Appendix B: Methods

2.1 Systematic searches

NICE and expert advisers provided potentially relevant references that were used to test the search strategy. These were also appraised for inclusion along with the search results. Systematic review searches covered the period 2005 to 2013, and primary study searches covered the period 1995 to 2013. Sample search strategies are reported in Section 2.1.4 below.

The first stage of systematic review searching took a broad approach. It combined three main facets to identify reviews relevant for Review 1:

- a facet targeting overweight and obesity prevention and healthy weight maintenance
- a facet targeting the broad areas such as diet, physical activity, and sedentary behaviour that the factors being assessed fall into
- a facet targeting systematic reviews (methodological filter).

The search also included a facet to identify reviews relevant for the complementary evidence review on message communication.

This strategy did not include specific terms focusing on each individual factor, but was intended to be broad enough to capture reviews addressing any factors such as these that were being assessed in the context of overweight and obesity prevention and healthy weight maintenance.

Scoping searches were carried out to identify which of the areas not covered by the first stage search looked most likely to provide additional reviews, to target second stage searching most efficiently. Potentially relevant reviews identified in these scoping searches were also assessed for inclusion.

2.1.1 Stage 1 search: Broad search for systematic reviews

Bibliographic databases (general and subject-specific) were searched to identify evidence from systematic reviews on the individually modifiable behaviours in the a priori list (Appendix A). An online search for systematic reviews in the grey literature (non-journal published papers) was also carried out on key websites.

Bibliographic database searches

The following bibliographic databases were searched:

- MEDLINE and MEDLINE In Process (OvidSP)

- Cochrane Database of Systematic Reviews
- HTA database (Centre for Reviews and Dissemination)
- Database of Abstracts of Reviews of Effects (DARE) (Centre for Reviews and Dissemination)
- Applied Social Sciences Index and Abstracts (ASSIA) (Proquest)
- Social Policy and Practice Database (Ovid)
- PsycINFO (Ovid)
- EPPI databases which include both journal published and grey literature: Bibliomap and DoPHER (Database of Promoting Health Effectiveness Reviews)
- healthevidence.org (registry of public health systematic reviews identified through searches of databases and websites including some covered in the list above, plus EMBASE, CINAHL, BIOSIS, SPORTDiscus, and Sociological Abstracts as well as reference list searches)

The SIGN systematic review filter was used for searching the MEDLINE, MEDLINE In Process and PsycINFO databases. The Medline search strategy was translated for the other databases, and adapted to take into account database size, coverage, available search facilities and available indexing terms.

A broad search on obesity prevention *only* without using the terms relating to the broad behavioural areas being targeted or the methodological filter was carried out in the following databases:

- Cochrane Database of Systematic Reviews
- HTA database (Centre for Reviews and Dissemination)
- Database of Abstracts of Reviews of Effects (DARE) (Centre for Reviews and Dissemination)
- DoPHER (Database of Promoting Health Effectiveness Reviews)
- healthevidence.org (registry of public health systematic reviews)

The same approach was used for some of the smaller databases such as Medline-In-Process database. The SIGN systematic review filter was used in the Medline-In-Process database. This search was adapted to take into account the more limited functionality and coverage of the Applied Social Sciences Index and Abstracts (ASSIA) and the Social Policy and Practice Database.

The search outputs were entered into Reference Manager.

Grey literature searches

Grey literature includes reports produced by government, academics, business and industry, theses or dissertations in electronic formats, but which are not controlled by commercial publishers/journals, i.e. where publishing is not the primary activity of the producing body.

We searched the following key websites for relevant grey literature reports:

- [Department of Health](#)
- [Public Health England](#)
- [National Obesity Observatory](#)
- [Harvard School of Public Health: The Obesity Prevention Source](#)
- [Centre for Diet and Activity Research](#)
- [WHO](#)
- [World Cancer Research Fund International \(WCRF\)](#)
- [Nutrition Evidence Library \(NEL\)](#) U.S. Department of Agriculture
- [Weight Management FAB approach \(Food, Activity, and Behavioural Support\)](#)
- Health Improvement and Innovation Resource Centre (New Zealand)
- [Food Standards Agency](#)
- [Sport England](#)
- [Obesity Learning Centre](#)
- [BASES 'British Association of Sport and Exercise Sciences'](#)
- [BASEM 'British Association of Sport and Exercise Medicine'](#)
- [European College of Sport Science](#)
- [CDC obesity http://www.cdc.gov/obesity/index.html](#)
- Foresight obesity <http://www.bis.gov.uk/foresight/our-work/projects/published-projects/tackling-obesities>
- [Institute of Medicine \(IOM\)](#)
- [National Weight Control Registry](#)
- [Scientific Advisory Committee on Nutrition \(SACN\)](#)
- [NICE Evidence](#)

2.1.2 Stage 2 search: Targeted search for systematic reviews

Based on gaps in the systematic review evidence identified in Stage 1 (see Section 2.2.3 for description of review mapping), the areas to be targeted in Stage 2 searching were discussed and agreed with NICE.

The search was focused on the following factors:

- Eating patterns (e.g. consistency, energy intake split through the day)
- Caffeine

- Coffee
- Holiday/Christmas weight gain
- Incidental physical activities (e.g. stair climbing)
- Breaks in sedentary time
- Sitting
- Stress-minimising activities
- Monitoring (e.g. self-weighing)
- Support

Stage 1 searches using only the broad obesity prevention facet should have identified all relevant reviews, and adding terms to this would not be anticipated to produce any additional hits. Therefore stage 2 searches were conducted only in the databases where Stage 1 searches were not limited to the broad obesity prevention facet.

2.1.3 Stage 3 search: Primary study searches

Based on gaps in the systematic review evidence identified in Stages 1 and 2 (see Section 2.2.3 for description of review mapping), the areas to be targeted in Stage 3 searching were discussed and agreed with NICE.

The search was focused on the following factors:

- Meal planning
- Holiday/Christmas weight gain
- Standing

The searches for Stage 3 were conducted in all databases excluding the secondary research databases:

- MEDLINE and MEDLINE In Process (OvidSP) (see below for Medline search strategy)
- Cochrane Central Register of Controlled Trials (CENTRAL)
- Applied Social Sciences Index and Abstracts (ASSIA) (Proquest, supplied by NICE)
- Social Policy and Practice Database (Ovid, supplied by NICE)
- PsycINFO (Ovid, supplied by NICE)
- EPPI databases which include both journal published and grey literature: Bibliomap and TRoPHI (Trials Register of Promoting Health Interventions)

2.1.4 Sample search strategies

Medline search strategy for Stage 1 broad systematic review search

Numbers in parentheses are # documents retrieved in MEDLINE (OvidSP).

Database: Ovid MEDLINE(R) <1946 to October Week 4 2013>

- 1 exp Obesity/ (141897)
- 2 Overweight/ (12114)
- 3 Weight Gain/ (23048)
- 4 Ideal Body Weight/ (115)
- 5 ((prevent* or reduc* or tackl* or address*) adj5 (obes* or "weight gain" or "excess weight" or overweight)).ti,ab. (18701)
- 6 ((maintain* or maintenance or prevent* or reduc* or control* or manag* or monitor* or healthy or normal or average) and (weight or bmi or body mass index or body fat or waist circumference or adiposity)).ti,ab. (392017)
- 7 (non obese or nonobese or non overweight or nonoverweight).ti,ab. (13360)
- 8 1 or 2 or 3 or 4 or 5 or 6 or 7 (503573)
- 9 Primary Prevention/ (14169)
- 10 Risk Factors/ (571499)
- 11 Health Promotion/ (53932)
- 12 Health Behavior/ (34680)
- 13 Health Education/ (52861)
- 14 Health Communication/ (497)
- 15 Information Dissemination/ (10367)
- 16 Marketing of Health Services/ (14042)
- 17 Health Knowledge, Attitudes, Practice/ (73857)
- 18 Risk Reduction Behavior/ (7539)
- 19 (promot* or advert* or marketing or program* or campaign* or scheme* or initiative* or strateg* or communicat* or message).ti,ab. (1831094)
- 20 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 (2444462)
- 21 Diet/ (109483)
- 22 exp *beverages/ or exp *food/ (683397)
- 23 exp Food Habits/ (21305)
- 24 Feeding Behavior/ (39759)
- 25 Energy Intake/ (31631)
- 26 (diet* or food* or eat*).ti. (211889)
- 27 exp Exercise/ (114912)
- 28 Motor Activity/ (79082)
- 29 Physical Fitness/ (22453)
- 30 (physical* or exercis* or fit* or aerobic*).ti. (184236)
- 31 Life Style/ (43434)
- 32 Sedentary Lifestyle/ (2568)
- 33 Size Perception/ (3877)
- 34 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 (1243671)
- 35 20 or 34 (3504940)
- 36 8 and 35 (191681)
- 37 Meta-Analysis as Topic/ (14079)
- 38 meta analy\$.tw. (57428)
- 39 metaanaly\$.tw. (1278)
- 40 Meta-Analysis/ (51298)
- 41 (systematic adj (review\$1 or overview\$1)).tw. (46645)
- 42 exp "Review Literature as Topic"/ (7635)
- 43 37 or 38 or 39 or 40 or 41 or 42 (114860)
- 44 cochrane.ab. (33111)
- 45 embase.ab. (29566)
- 46 (psychlit or psyclit).ab. (1189)
- 47 (psychinfo or psycinfo).ab. (9892)
- 48 (cinahl or cinhal).ab. (10882)
- 49 science citation index.ab. (2320)
- 50 bids.ab. (394)
- 51 cancerlit.ab. (739)
- 52 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 (50861)

- 53 reference list\$.ab. (11616)
- 54 bibliograph\$.ab. (11748)
- 55 hand-search\$.ab. (4216)
- 56 relevant journals.ab. (898)
- 57 manual search\$.ab. (2224)
- 58 53 or 54 or 55 or 56 or 57 (27351)
- 59 selection criteria.ab. (26048)
- 60 data extraction.ab. (10031)
- 61 59 or 60 (33618)
- 62 Review/ (1915234)
- 63 61 and 62 (25914)
- 64 Comment/ (534284)
- 65 Letter/ (804607)
- 66 Editorial/ (335541)
- 67 animal/ (5493002)
- 68 human/ (13649513)
- 69 67 not (67 and 68) (3962474)
- 70 64 or 65 or 66 or 69 (5154801)
- 71 43 or 52 or 58 or 63 (145675)
- 72 71 not 70 (136714)
- 73 36 and 72 (4330)
- 74 limit 73 to (english language and yr="2005 -Current") (3157)

Search strategies for stage 2 focused systematic review search

Database: Ovid MEDLINE(R) <1946 to November Week 3 2013>

- 1 exp Obesity/ (142945)
- 2 Overweight/ (12299)
- 3 Weight Gain/ (23205)
- 4 Ideal Body Weight/ (121)
- 5 ((prevent* or reduc* or tackl* or address*) adj5 (obes* or "weight gain" or "excess weight" or overweight)).ti,ab. (18862)
- 6 ((maintain* or maintenance or prevent* or reduc* or control* or monitor* or healthy or normal or average) and (weight or bmi or body mass index or body fat or waist circumference or adiposity)).ti,ab. (385493)
- 7 (non obese or nonobese or non overweight or nonoverweight).ti,ab. (13464)
- 8 1 or 2 or 3 or 4 or 5 or 6 or 7 (499364)
- 9 ((meal or eating or diet*) adj3 (pattern or habit or irregular)).ti,ab. (3528)
- 10 Coffee/ (4754)
- 11 coffee.ti,ab. (8263)
- 12 caffeine.ti,ab. (21680)
- 13 Caffeine/ (20280)
- 14 (holiday or Christmas).ti,ab. (3195)
- 15 Holidays/ (2159)
- 16 9 or 10 or 11 or 12 or 13 or 14 or 15 (42002)
- 17 stair*.ti,ab. (5622)
- 18 ((sedentary or sitting) adj5 (time or break*)).ti,ab. (1521)
- 19 (stress adj3 (reduc* or minimi*)).ti,ab. (15417)
- 20 (self-monitor* or self-weigh*).ti,ab. (4528)
- 21 Social Support/ (52277)
- 22 ((psychological or social) adj3 (support or network)).ti,ab. (28206)
- 23 17 or 18 or 19 or 20 or 21 or 22 (91604)
- 24 16 or 23 (133360)
- 25 8 and 24 (7725)
- 26 Meta-Analysis as Topic/ (14196)
- 27 meta analy\$.tw. (58408)
- 28 metaanaly\$.tw. (1286)
- 29 Meta-Analysis/ (52213)

- 30 (systematic adj (review\$1 or overview\$1)).tw. (47522)
- 31 exp "Review Literature as Topic"/ (7732)
- 32 26 or 27 or 28 or 29 or 30 or 31 (116583)
- 33 cochrane.ab. (33754)
- 34 embase.ab. (30194)
- 35 (psychlit or psyclit).ab. (1193)
- 36 (psychinfo or psycinfo).ab. (10152)
- 37 (cinahl or cinhal).ab. (11088)
- 38 science citation index.ab. (2372)
- 39 bids.ab. (395)
- 40 cancerlit.ab. (739)
- 41 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 (51910)
- 42 reference list\$.ab. (11809)
- 43 bibliograph\$.ab. (11863)
- 44 hand-search\$.ab. (4252)
- 45 relevant journals.ab. (914)
- 46 manual search\$.ab. (2265)
- 47 42 or 43 or 44 or 45 or 46 (27707)
- 48 selection criteria.ab. (26402)
- 49 data extraction.ab. (10177)
- 50 48 or 49 (34086)
- 51 Review/ (1924416)
- 52 50 and 51 (26304)
- 53 Comment/ (538304)
- 54 Letter/ (808271)
- 55 Editorial/ (337516)
- 56 animal/ (5513005)
- 57 human/ (13712248)
- 58 56 not (56 and 57) (3974347)
- 59 53 or 54 or 55 or 58 (5173095)
- 60 32 or 41 or 47 or 52 (147811)
- 61 60 not 59 (138759)
- 62 25 and 61 (205)
- 63 limit 62 to (english language and yr="2005 -Current") (151)

Sample search strategy for Stage 3: primary study search

Database: Ovid MEDLINE(R) In-Process & Other Non-Indexed Citations and OviD
 MEDLINE(R) <1946 to Present> Search Strategy:

- 1 exp Obesity/ (144151)
- 2 Overweight/ (12498)
- 3 Weight Gain/ (23356)
- 4 Ideal Body Weight/ (121)
- 5 ((prevent* or reduc* or tackl* or address*) adj5 (obes* or "weight gain" or "excess weight" or overweight)).ti,ab. (20763)
- 6 ((maintain* or maintenance or prevent* or reduc* or control* or manag* or monitor* or healthy or normal or average) and (weight or bmi or body mass index or body fat or waist circumference or adiposity)).ti,ab. (425240)
- 7 (non obese or nonobese or non overweight or nonoverweight).ti,ab. (14275)
- 8 1 or 2 or 3 or 4 or 5 or 6 or 7 (539354)
- 9 (holiday or Christmas).ti,ab. (3462)
- 10 Holidays/ (2171)
- 11 9 or 10 (4911)
- 12 8 and 11 (108)
- 13 ((meal* or menu) and plan*).ti,ab. (2643)
- 14 8 and 13 (493)
- 15 (stand or standing).ti. (10782)

- 16 8 and 15 (401)
- 17 Randomized Controlled Trials as Topic/ (103814)
- 18 Randomized Controlled Trial/ (395285)
- 19 Random Allocation/ (82260)
- 20 Double-Blind Method/ (132982)
- 21 Single Blind Method/ (19826)
- 22 Clinical trial/ (508008)
- 23 clinical trial, phase i.pt. (16502)
- 24 clinical trial, phase ii.pt. (27300)
- 25 clinical trial, phase iii.pt. (10407)
- 26 clinical trial, phase iv.pt. (1014)
- 27 controlled clinical trial.pt. (90572)
- 28 randomized controlled trial.pt. (395285)
- 29 multicenter study.pt. (185820)
- 30 clinical trial.pt. (508008)
- 31 exp Clinical Trials as Topic/ (299072)
- 32 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 (1088792)
- 33 (clinical adj trial\$.tw. (230108)
- 34 ((singl\$ or doubl\$ or treb\$ or tripl\$) adj (blind\$3 or mask\$3)).tw. (136258)
- 35 Placebos/ (33961)
- 36 placebo\$.tw. (170964)
- 37 randomly allocated.tw. (17431)
- 38 (allocated adj2 random\$.tw. (20044)
- 39 33 or 34 or 35 or 36 or 37 or 38 (447196)
- 40 32 or 39 (1241852)
- 41 case report.tw. (205280)
- 42 Letter/ (838323)
- 43 Historical Article/ (302092)
- 44 41 or 42 or 43 (1334094)
- 45 40 not 44 (1211323)
- 46 Epidemiologic studies/ (6316)
- 47 exp case control studies/ (674283)
- 48 exp cohort studies/ (1384167)
- 49 Case control.tw. (80852)
- 50 (cohort adj (study or studies)).tw. (90911)
- 51 Cohort analy\$.tw. (3850)
- 52 (Follow up adj (study or studies)).tw. (38340)
- 53 (observational adj (study or studies)).tw. (46443)
- 54 Longitudinal.tw. (147448)
- 55 Retrospective.tw. (278034)
- 56 Cross sectional.tw. (176313)
- 57 Cross-sectional studies/ (183409)
- 58 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 (1941703)
- 59 45 or 58 (2861180)
- 60 12 and 59 (42)
- 61 14 and 59 (194)
- 62 16 and 59 (136)
- 63 limit 60 to (english language and yr="1995 -Current") (36)
- 64 limit 61 to (english language and yr="1995 -Current") (182)
- 65 limit 62 to (english language and yr="1995 -Current") (129)

2.2 Selecting studies for inclusion

Studies were evaluated for inclusion against the criteria listed in the sifting protocol below. Broadly, systematic reviews of the association between the individually modifiable behaviours listed in Appendix A and healthy weight maintenance or overweight and obesity prevention were included. As many systematic reviews are unlikely to match the current review's inclusion criteria fully, potentially relevant reviews were included at early sifting stages unless it was clear that they were entirely non-relevant to the current scope.

| Parameters | Sifting criteria | Additional comments |
|---|--|--|
| Questions covered | <p>What individually modifiable behaviours may help children and young people to maintain a healthy weight or prevent excess weight gain?</p> <p>What individually modifiable behaviours may help adults to maintain a healthy weight or prevent excess weight gain?</p> | Reviews addressing other questions were tagged 'wrong question' |
| Exposures/ interventions covered | <p>Exposures: Individually modifiable behaviours listed in Appendix A that may help children, young people and adults to maintain a healthy weight or prevent excess weight gain</p> <p>Interventions: Interventions that target the individually modifiable behaviours listed in Appendix A and that are specifically aimed at maintaining a healthy weight or preventing excess weight gain</p> | Interventions should consist of an action that an individual can choose to perform themselves, rather than one requiring external intervention |
| Exposures/ interventions not covered | <p>Interventions to prevent obesity that are covered in other sections of NICE clinical guideline 43. That is, sections 1.1.2 to 1.1.7, and section 1.2, i.e. interventions that take place in/offered by:</p> <ul style="list-style-type: none"> - The NHS - Local authorities and partners in the community - Early years settings - Schools - Workplaces - Self-help, commercial and community programmes | <p>Records excluded on these criteria were tagged 'wrong intervention/exposure' at second (title and abstract) and third (full text) sift.</p> <p>NB Personal characteristics such as gender, SES and ethnicity of the populations included in studies where noted. If studies find that impact of the individual factors/ behaviours or interventions targeting them vary based on these</p> |

| Parameters | Sifting criteria | Additional comments |
|------------|---|---|
| | <p>Very low calorie diets</p> <p>Environmental factors beyond people's control (for example, the provision of cycle paths or content of school meals).</p> <p>Programmes, services or treatments for people who are overweight or obese (including lifestyle weight management services and pharmacological or surgical treatments).</p> <p>Management of medical conditions that may increase the risk of excess weight gain, being overweight or obese.</p> <p>Programmes, services or treatments for people who are underweight.</p> <p>Infant feeding (with breast milk or infant formula) and weaning.</p> <p>Complementary/non-mainstream therapies to prevent someone from becoming overweight or obese or to manage these conditions (for example, acupuncture, hypnotherapy, medicinal plants).</p> <p>Studies aiming to define 'overweight' and 'obese'.</p> <p>Related activities to combat obesity that are covered by other NICE guidance (such as breastfeeding).</p> <p>Parenting</p> <p>Health (or other) professional led interventions</p> <p>Work / school based interventions</p> <p>Working circumstances e.g. shift working</p> <p>Personal circumstances /characteristics that are not under solely under an individual's control (see Additional comment), e.g.:</p> <ul style="list-style-type: none"> - gender - marital status - parental weight (impact on their children) - menopause - puberty - birth weight | <p>characteristics this was reported.</p> |

| Parameters | Sifting criteria | Additional comments |
|--|--|--|
| | <ul style="list-style-type: none"> - socioeconomic status - ethnicity - self esteem Smoking | |
| Populations (groups) that will be covered | All adults and children who are not undergoing management or treatment for being overweight or obese. | The focus is on the general population (i.e. mixed populations) |
| Populations (groups) that will not be covered | Infants who have not been weaned Pregnant women Adults and children who are taking part in programmes or are receiving treatment for being overweight or obese (including lifestyle weight management programmes, pharmacological or surgical treatment). Adults and children who are taking part in programmes or receiving treatment for being underweight Populations that are selected solely on the basis of being overweight or obese, or having been overweight and obese and now reached a healthy weight (e.g. studies which follow groups who have taken part in weight loss interventions) Studies specifically in selected population subgroups (see Additional comments), e.g.: <ul style="list-style-type: none"> - post-pregnancy - learning difficulties - mental health conditions - disabilities - (NB people in these groups as well as those who are overweight or obese are part of the general population, but studies selecting participants on these characteristics exclusively will be excluded) | Records excluded on this criterion were tagged 'wrong population' at second (title and abstract) and third (full text) sift. (Weaning or 'complementary feeding' is the transition from an exclusively milk-based diet to a diet based on solid foods.) NB The presence of population subgroups such as those with learning difficulties, mental health conditions, or disabilities in included studies will be noted. If studies find that the impact of the individual factors/behaviours or interventions targeting them vary in these subgroups this will also be reported. |
| Comparators that will be covered | Studies of exposures: <ul style="list-style-type: none"> - No exposure - A different level of exposure (e.g. less versus more physical activity/calories/of a | |

| Parameters | Sifting criteria | Additional comments |
|---|--|---|
| | <p>food/nutrient)</p> <p>Intervention studies:</p> <ul style="list-style-type: none"> - No intervention - Usual care - Placebo/sham <p>A different level of the intervention (e.g. less versus more physical activity/calories/of a food/nutrient)</p> | |
| Comparators that will not be covered | <p>Intervention studies:</p> <p>Studies that compare interventions aimed at different behaviours e.g. physical activity intervention vs. diet based intervention</p> | <p>Records excluded on this criterion will be tagged 'wrong comparator' at second title and abstract and full text sift</p> |
| Outcomes that will be covered | <p>Weight and related outcomes</p> <p>Maintenance of weight or avoidance of weight gain in the short, medium and long term.</p> <p>Anthropometric measures such as BMI, waist circumference, percentage of healthy weight, or fat mass.</p> <p>Modifiable risk factor (behavioural) outcomes (to be extracted from intervention studies which also address weight and related outcomes)</p> <p>Diet</p> <p>Physical activity</p> <p>Frequency of weight monitoring</p> | <p>Only studies reporting weight or related outcomes will be included.</p> |
| Outcomes that will not be covered | <p>Determinant outcomes</p> <p>Psychological outcomes such as self-efficacy or motivation.</p> <p>Process measures such as acceptability of information (method or content) that aims to help people maintain a healthy weight.</p> | <p>Records excluded for on this criteria tagged 'wrong outcome at second title and abstract and full text sift.</p> |
| Study types/designs to be included | <p>Systematic Reviews (SRs)</p> <p>RCTs and cluster RCTs of any duration</p> <p>Prospective cohort studies lasting 12 months or longer</p> | <p>NB Due to the mixed nature of studies included in the reviews identified, reviews were not excluded if they included retrospective cohort studies, or cohort studies lasting less than 12 months</p> |

| Parameters | Sifting criteria | Additional comments |
|--|--|---|
| Studies types/designs that will not be included | <p>Studies that are not SRs, RCTs, or prospective cohorts.</p> <p>Studies published before 2005</p> <p>Economic analyses (references of these studies to be forwarded to NICE)</p> <p>Non-English language studies.</p> <p>Non-OECD countries studies.</p> <p>Citations without an abstract.</p> <p>Systematic reviews of systematic reviews</p> | <p>Records excluded for on this criteria tagged 'wrong study type' at second title and abstract and full text sift.</p> |

2.2.1 First pass appraisal

Evidence identified in the searches was first filtered at the title/abstract level by an Information Specialist to remove any clearly non-relevant material. Any queries regarding inclusion/exclusion were resolved by discussion with a second Information Specialist. Any outstanding queries were resolved by discussion with NICE. Studies were excluded at this stage if they were:

- Clearly non-relevant topics or populations or information (e.g. letters, animal studies, studies in specific settings or delivered by healthcare professionals)
- Non-relevant study design/type (i.e. not systematic review in Stage 1 or 2 searches, or not RCTs/prospective cohort studies in Stage 3 searches)

A random sample of 200 citations identified in the search for systematic reviews were double sifted by a second Information Specialist. A kappa of 0.60 or greater was considered to reflect good inter-rater reliability. The double sift of systematic reviews resulted in agreement above the agreed threshold (kappa=0.68). Ten per cent of the records identified in the search for primary studies were double sifted by a second Information Specialist. The double sift resulted in agreement above the agreed threshold (kappa=0.79).

This stage of screening acted as a “coarse filter” and erred on the side of inclusion, to avoid exclusion of studies that may be relevant. The filtered references were tagged in a Reference Manager database and passed on to a Research Analyst for second pass appraisal.

2.2.2 Second pass appraisal

A Health Research Analyst conducted a more detailed assessment of the title/abstract of records tagged during the first sift. Relevant studies were selected for full text appraisal. Studies were appraised using the sifting criteria described. As few systematic reviews were

anticipated to fully match the scope of the current review, those that had inclusion criteria that partially overlapped or had unclear overlap with the scope were included at this stage, e.g. the following:

- **Study designs:** reviews where included study designs were a mix of relevant and non-relevant study designs (RCTs and/or cohort studies plus other study designs) or where study designs were unclear
- **Settings:** reviews where included studies were in a mix of relevant and non-relevant settings (general community plus school/work settings) or where settings were unclear
- **Populations:** reviews including studies in a mix of general populations and selected obese/overweight populations and/or populations with a specific condition or where populations were unclear
- **Intervention/exposure:** reviews including studies of factor(s) of interest as well as other factors or where factors addressed were unclear
- **Comparator:** reviews where comparators were unclear or mixed
- **Outcome:** reviews had to mention measuring a weight related outcome for inclusion

Studies were not excluded based on duration of included studies, as this was difficult to judge at the title and abstract level. Reasons for exclusion of studies were recorded in the Reference Manager database at this stage (see Appendix C for excluded study bibliography). These reasons could include:

- Wrong question (i.e. studies not addressing the question about factors associated with weight maintenance, or interventions for healthy weight maintenance)
- Wrong population (e.g. studies with inclusion restricted to overweight or obese individuals)
- Wrong study design/type (i.e. not a systematic review in Stages 1 and 2; not an RCT or prospective cohort study in Stage 3)
- Wrong exposure/intervention (e.g. studies clearly not relating to an individually modifiable behaviour on the agreed list, or clearly in an entirely work/school-based setting)
- Wrong comparator (e.g. studies solely comparing the weight maintenance effects of interventions targeting different individually modifiable behaviours e.g. diet vs. physical activity)
- Wrong outcome (e.g. studies solely addressing knowledge outcomes or other non-weight-related outcomes)

Any queries regarding inclusion/exclusion were resolved by discussion with a second analyst. Any outstanding queries were resolved by discussion with NICE. If it was still unclear whether a study met inclusion/exclusion criteria, the full text was obtained. A 10% sample of citations were then double sifted by a second Health Research Analyst for reviews, which resulted in good inter-rater reliability ($\kappa=0.63$). For primary study searches the small number of

records selected for second pass appraisal (12 studies), all of the records were double screened by a second analyst and there was 100% agreement. Kappa was not calculated for this, as it may be under-estimated in sample sizes (Crewson 2005).

This stage of screening acted as a finer filter than the first pass appraisal, but again erred on the side of inclusion if details were not sufficient to allow decisions about the eligibility of the paper. Papers selected for full text appraisal were tagged in Reference Manager.

2.2.3 Systematic review mapping

In order to identify areas to be targeted for stage 2 and 3 searches, a rapid, rough mapping of the systematic reviews identified against the factors being assessed was carried out at the end of the second pass appraisal of Stage 1 and Stage 2 searches. As part of mapping at the end of Stage 1, for factors where no or few reviews were mapped, text word searches were carried out in the dataset to ensure that no potentially relevant reviews which had been missed.

Reviews were grouped according to which factors they appeared to cover based on title and abstract. The number of reviews that appeared to cover each factor was listed. An initial rapid quality assessment was designed to capture indications of review quality from the title and abstract. These were broadly based on criteria for definition of systematic reviews in the Database of Abstracts of Reviews of Effects (DARE). Reviews were given a score out of 4, by summing the total number of the following criteria it met:

- The review identifies itself in the title or abstract as a systematic review
- The review reports that it conforms to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) standards
- The abstract reports that two or more sources were searched
- The abstract reports the review's inclusion/exclusion criteria

Discussion and decision of which factors to target with searches was based around:

- Number of reviews covering the factor
- Quality of the reviews covering the factor
- Publication dates of reviews covering the factor (a proxy measure for being up-to-date)
- Likelihood of identifying additional reviews on the factor based on the number and type of hits identified in rapid scoping searches in PubMed (any potentially relevant reviews identified in these scoping searches were also added to the RefMan database)
- Potential for the factor to be covered by reviews on related factors (e.g. findings from reviews on energy density would also be relevant to the factors low and high energy dense foods)
- Match of the reviews identified to the factors being targeted

- Whether the factor was covered in by existing CG43 guidance (as adding nuance to existing guidance would be more likely for areas where there were no existing recommendations)
- Practical factors relating to search (e.g. specificity of terms involved)
- Whether the factor might be better suited to a primary study search (i.e. relatively newly researched topics might be less likely to be covered by systematic reviews)

The initial plan had been to assess match between the identified reviews' scope and that of the current review. In the end, this was not considered, as reviews that were completely outside of the scope were excluded, and of a sample of 63 reviews assessed, none were clearly a complete match to scope. Hence, this level of assessment would be unlikely to be a good way of differentiating the identified reviews. Expert advisers also provided input on any areas where they felt reviews might be missing.

In general,

- Factors where no reviews identified were considered first
- Factors where a review was identified were considered after this if:
 - There was only 1 SR on the factor, or
 - SRs identified did not appear to match well with the concept being searched for, or
 - All SRs identified had a lower quality score based on title and abstract (quality score 0 or 1).

2.2.4 Full text appraisal

The full text papers were appraised by a Research Analyst. Information on reason for exclusion was recorded (see Appendix C for excluded study bibliography). A 10% sample of full texts were planned to be double screened at this stage. This would have been a relatively low number of studies, and for low sample sizes the kappa statistic may be under-estimated (Crewson 2005). For this reason, alternative approaches were agreed with NICE.

For the reviews, to ensure that no reviews were excluded inappropriately, a more conservative approach was taken where all papers excluded at full text were assessed by a second analyst for inclusion/exclusion. Disagreements regarding inclusion/exclusion were resolved by discussion, with recourse to a third analyst as needed.

As anticipated at second pass appraisal, systematic reviews tended not to fully match the scope of the current review. Therefore, those that had inclusion criteria that partially overlapped with the scope were included at this stage. Reviews whose inclusion criteria or included studies fell completely outside of the scope of the current review in terms of study design, setting, population, intervention/exposure, comparator, or outcome were excluded.

Reviews were not excluded solely based on the duration of follow up of the included studies. Whether included cohort studies were prospective or retrospective was also not used as a reason for exclusion of reviews.

For the primary studies, all full texts were appraised by a second analyst.

2.2.5 Systematic reviews generically assessing overweight and obesity prevention

A number of reviews were identified which generically addressed overweight and obesity prevention, rather than specifically asking questions about the factors listed in Table 1 and describing results in a way that clearly separates results by factor.

The 'generic reviews' generally asked questions about effective overweight and obesity prevention interventions, or about the association between non-specific exposures rather than focusing on specific modifiable behaviours or factors (e.g. physical activity in general, with no further detail on the type, frequency, intensity or duration). They were sifted using the same criteria and process outlined in Sections 3.4.1 to 3.4.3. Due to the large volume of reviews identified which specifically addressed the factors listed in Table 1, these factor-specific reviews were prioritised for extraction, as they are more likely to provide clear answers regarding the relationship between the individually modifiable behaviours/factors on healthy weight maintenance.

The 'generic reviews' are listed in Appendix C.

2.3 Quality appraisal

Quality appraisal was carried out for all reviews selected at full text using a checklist based on the NICE systematic review quality checklist and the Critical Appraisal Skills Programme (CASP) systematic review quality checklist. Primary studies were assessed using the appropriate NICE quantitative study checklist. These checklists are provided in Appendix D. The ratings are broadly as follows:

[++] All or most of the checklist criteria have been fulfilled, indicating a high quality study.

[+] Some of the checklist criteria have been fulfilled, indicating a moderate quality study.

[-] Few or no checklist criteria have been fulfilled, indicating a low quality study.

A 10% sample of included reviews (n=14) was double quality appraised by a second research analyst. The kappa score obtained for this did not meet the required threshold for good inter-rater reliability (kappa=0.45; agreement 64%). This may in part have been due to the low number of studies double appraised. Areas where ratings did not agree were reviewed. In one

case, the lack of agreement was due to mis-entering of one of the question ratings into Excel, correcting this led to a kappa of 0.54 (agreement 71%). After discussion of the differences in ratings, other differences related to grey areas. Only one rating was changed due to presence of additional study tables in the supplementary information that had not been identified by the original reviewer.

A repeat sample of 14 papers was reviewed, and again agreement was 71%, but kappa was lower (kappa=0.20). This is likely to have been due to the low prevalence of the [-] and [++] categories in this sample (0 and 2 papers) - kappa can be under-estimated if any of the categories are uncommon as chance agreement is high (Sim and Wright, 2005). Therefore combining the samples still gave a low kappa (kappa=0.45; agreement 71%).

The 3 papers where ratings differed in the second sample all lay at the boundary between two rating categories, and differed by a single point difference on the total quality score (out of 8). To investigate whether underlying agreement was high, and to increase the sample size for the analysis, the kappa statistic was recalculated based on the answers to the eight individual questions for all 18 studies. This gave an agreement of 87%, and a kappa of 0.73, showing good inter-rater reliability. It was agreed with NICE that this indicated good underlying agreement and that further review was not required.

2.4 Data extraction

For factors covered by multiple reviews, the reviews were assessed and the highest quality, most up-to-date, and most relevant (i.e. best match for the scope) reviews covering children, young people and adults were selected for extraction. The aim was to have at least one review covering children and young people and at least one covering adults for each factor. More than one review of similar quality, search date, or relevance could be included if they covered pools of studies that did not overlap completely. For reviews not prioritised for extraction, reasons were recorded (see Appendix E), and review conclusions were extracted to give an idea of the direction of effect.

No reviews matched the scope of the current review exactly. The scope of the included reviews was assessed mainly in the following areas:

1. **Population** – reviews including some studies not matching the current review scope (e.g. overweight/obese people and/or people with specific conditions) were considered a partial match
2. **Study design** – reviews including some studies not matching the current review scope (e.g. cross sectional studies) were considered a partial match
3. **Setting** - reviews including some studies not matching the current review scope (e.g. school- or work-based studies) were considered a partial match

The intervention/exposures were not assessed for match, as these had to match the factors being assessed for inclusion. Time (i.e. follow up period) was also not assessed for match, as few reviews utilised duration of follow up as an inclusion criterion. Comparators were also not assessed for match as this did not provide an informative way of separating the reviews, few studies explicitly specified comparators in their inclusion criteria (particularly those including cohort studies), and the reviews including RCTs often included mixed comparators. No papers assessed at full text were excluded on this parameter. Although not specifically assessed for match with scope, any limitations relating to these areas were noted.

Where the duration of follow up was used as an inclusion criterion it was noted in the evidence tables, and where duration of follow up was a limitation this was also noted.

Many reviews did not provide in depth reporting of e.g. the level of exposures being compared in the included studies. This data was extracted where possible.

Where the review itself provided separate results and conclusions based on the different populations, settings, or outcomes it included, those most relevant to the current review (i.e. most closely matching the scope) were extracted. For example, if a review gave results for overweight and obese populations separately from those for general weight populations, the latter were reported. If results were not reported or summarised by the review separately, overall results and conclusions were reported. Due to limitations in the time available for the review, authors were not contacted for additional information, nor were underlying primary studies or related references obtained.

3 Appendix C: Excluded study bibliography

This appendix lists reasons for exclusion of studies at second (title and abstract) and third (full text) sift.

Wrong Population (15 studies)

1. Boutelle KN, Kirschenbaum DS, Baker RC et al. How can obese weight controllers minimize weight gain during the high risk holiday season? By self-monitoring very consistently. *Health Psychology*. 1999;18(4):364-8.
2. Boutelle KN. The efficacy of a self-monitoring intervention to promote weight management during the holidays. *Dissertation Abstracts International: Section B: The Sciences and Engineering*. 1111;.57(9-B).
3. Galani C, Schneider H. Prevention and treatment of obesity with lifestyle interventions: review and meta-analysis. *International Journal of Public Health*. 2007;52(6):348-59.
4. Hu T, Mills KT, Yao L et al. Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: a meta-analysis of randomized controlled clinical trials. *American Journal of Epidemiology*. 2012;176 Suppl 7:S44-S54.
5. Hursel R, Viechtbauer W, Westerterp-Plantenga MS. The effects of green tea on weight loss and weight maintenance: a meta-analysis. *International Journal of Obesity*. 2009 20090714 [Epub ahead of print];33(9):956-61.
6. Lepe M, Bacardi GM, Jimenez CA. Long-term efficacy of high-protein diets: a systematic review. *Nutricion Hospitalaria*. 2011;26(6):1256-9.
7. Musaiger AO. Overweight and obesity in eastern mediterranean region: prevalence and possible causes. *Journal of Obesity*. 2011 20110918 [Epub ahead of print];2011:407237.
8. Nordmann AJ, Nordmann A, Briel M et al. Effects of low-carbohydrate vs low-fat diets on weight loss and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Archives of Internal Medicine*. 2006;166(3):285-93.
9. Rego Costa AC, Rosado EL, Soares-Mota M. Influence of the dietary intake of medium chain triglycerides on body composition, energy expenditure and satiety: a systematic review. *Nutricion Hospitalaria*. 2012;27(1):103-8.
10. Schwingshackl L, Hoffmann G. Long-term effects of low glycemic index/load vs. high glycemic index/load diets on parameters of obesity and obesity-associated risks: a systematic review and meta-analysis. *Nutrition Metabolism & Cardiovascular Diseases*. 2013 20130617 [Epub ahead of print];23(8):699-706.
11. Steyn NP, Temple NJ. Evidence to support a food-based dietary guideline on sugar consumption in South Africa. *BMC Public Health*. 2012 20120704 [Epub ahead of print];12:502.
12. U.S. Department of Agriculture. Are high-protein (>35%) hypocaloric diets safe and effective for long term (more than six months) weight loss or maintenance? (DGAC 2010). Washington (DC): U.S. Department of Agriculture; 2010. Available from: http://www.nel.gov/conclusion.cfm?conclusion_statement_id=250328&highlight=hypocaloric&home=1.
13. U.S. Department of Agriculture. Are low-carbohydrate (less than 45%) hypocaloric diets safe and effective for long-term (more than six months) weight loss or maintenance?

Washington (DC): U.S. Department of Agriculture; 2010. Available from:
http://www.nel.gov/conclusion.cfm?conclusion_statement_id=250327.

14. U.S. Department of Agriculture. What is the relationship between diet self-monitoring and body weight? (DGAC 2010). Washington (DC): U.S. Department of Agriculture; 2010g. Available from:
http://www.nel.gov/conclusion.cfm?conclusion_statement_id=250321.
15. Weinheimer EM, Sands LP, Campbell WW. A systematic review of the separate and combined effects of energy restriction and exercise on fat-free mass in middle-aged and older adults: implications for sarcopenic obesity. *Nutrition Reviews*. 2010;68(7):375-88.

Wrong Exposure/Intervention (47 studies)

1. Adriaanse MA, Vinkers CD, de Ridder DT et al. Do implementation intentions help to eat a healthy diet? A systematic review and meta-analysis of the empirical evidence. *Appetite*. 2011 21056605 [Epub ahead of print];56(1):SP-183.
2. Barr-Anderson DJ, AuYoung M, Whitt-Glover MC et al. Integration of short bouts of physical activity into organizational routine a systematic review of the literature. [Review]. *American Journal of Preventive Medicine*. 2011;40(1):76-93.
3. Beets MW, Beighle A, Erwin HE et al. After school program impact on physical activity and fitness: A meta analysis. *American Journal of Preventive Medicine*. 2009 19362799 [Epub ahead of print];36(6):SP-527.
4. Bhopal RS, Rafnsson SB. Could mitochondrial efficiency explain the susceptibility to adiposity, metabolic syndrome, diabetes and cardiovascular diseases in South Asian populations?. *International Journal of Epidemiology*. 2009 20090507 [Epub ahead of print];38(4):1072-81.
5. Blaine B. Does depression cause obesity?: A meta-analysis of longitudinal studies of depression and weight control. *Journal of Health Psychology*. 2008;13(8):1190-7.
6. Chapman J, Qureshi N, Kai J. Effectiveness of physical activity and dietary interventions in South Asian populations: a systematic review. *British Journal of General Practice*. 2013;63(607):e104-e114.
7. Chillon P, Evenson KR, Vaughn A et al. A systematic review of interventions for promoting active transportation to school. *The International Journal of Behavioral Nutrition and Physical Activity*. 2011 21320322 [Epub ahead of print];8(1):SP-10.
8. Delavari M, Sonderlund AL, Swinburn B et al. Acculturation and obesity among migrant populations in high income countries--a systematic review. *BMC Public Health*. 2013 20130510 [Epub ahead of print];13:458.
9. DeMattia L, Lemont L, Meurer L. Do interventions to limit sedentary behaviours change behaviour and reduce childhood obesity: a critical review of the literature. *Obesity Reviews*. 2007;8(1):69-81.
10. Finlay SJ, Faulkner G. Physical activity promotion through the mass media: inception, production, transmission and consumption. *Prev Med*. 2005;40(2):121-30.
11. Fraser J, Skouteris H, McCabe M et al. Paternal influences on children's weight gain: A systematic review. *Fathering*. 2011;9(3):257-67.
12. Giskes K, Kamphuis CB, Van Lenthe FJ et al. A systematic review of associations between environmental factors, energy and fat intakes among adults: is there evidence

- for environments that encourage obesogenic dietary intakes?. *Public Health Nutrition*. 2007 20070222 [Epub ahead of print];10(10):1005-17.
13. Giskes K, van LF, Avendano-Pabon M et al. A systematic review of environmental factors and obesogenic dietary intakes among adults: are we getting closer to understanding obesogenic environments?. *Obesity Reviews*. 2011;12(5):e95-e106.
 14. Goodman C, Anise A. What is known about the effectiveness of economic instruments to reduce consumption of foods high in saturated fats and other energy-dense foods for preventing and treating obesity? 2006. Available from: http://www.euro.who.int/__data/assets/pdf_file/0010/74467/E88909.pdf.
 15. Grasser G, Van Dyck D., Titze S et al. Objectively measured walkability and active transport and weight-related outcomes in adults: a systematic review. *International Journal of Public Health*. 2013 20121206 [Epub ahead of print];58(4):615-25.
 16. Holsten JE. Obesity and the community food environment: a systematic review. *Public Health Nutrition*. 2009 20080514 [Epub ahead of print];12(3):397-405.
 17. Incedon E, Wake M, Hay M. Psychological predictors of adiposity: systematic review of longitudinal studies. *International Journal of Pediatric Obesity*. 2011 20110119 [Epub ahead of print];6(2-2):e1-11.
 18. Jensen JD, Hartmann H, de MA et al. Economic incentives and nutritional behavior of children in the school setting: A systematic review. *Nutrition Reviews*. 2011 22029832 [Epub ahead of print];69(11):SP-660.
 19. Jones A, Bentham G, Foster C et al. Tackling obesities: Future choices - Obesogenic environments - Evidence review. London: Government Office for Science; 2007. Available from: <https://www.bis.gov.uk/assets/foresight/docs/obesity/03.pdf>.
 20. Katz DL, O'Connell M, Yeh MC et al. Public health strategies for preventing and controlling overweight and obesity in school and worksite settings: a report on recommendations of the Task Force on Community Preventive Services. *Morbidity & Mortality Weekly Report*. 2005;Recommendations & Reports. 54(RR-10):1-12.
 21. Liber A, Szajewska H. Effects of inulin-type fructans on appetite, energy intake, and body weight in children and adults: systematic review of randomized controlled trials. *Annals of Nutrition & Metabolism*. 2013 20130723 [Epub ahead of print];63(1-2):42-54.
 22. Lubans DR, Morgan PJ, Cliff DP et al. Fundamental movement skills in children and adolescents: review of associated health benefits. *Sports Medicine*. 2010;40(12):1019-35.
 23. Ludy MJ, Moore GE, Mattes RD. The effects of capsaicin and capsiate on energy balance: critical review and meta-analyses of studies in humans. *Chemical Senses*. 2012 20111029 [Epub ahead of print];37(2):103-21.
 24. Luppino FS, de Wit LM, Bouvy PF et al. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry*. 2010;67(3):220-9.
 25. Michie S, Abraham C, Whittington C et al. Effective techniques in healthy eating and physical activity interventions: A meta-regression. *Health Psychology*. 2009 19916637 [Epub ahead of print];28(6):SP-690.
 26. Midei AJ, Matthews KA. Interpersonal violence in childhood as a risk factor for obesity: a systematic review of the literature and proposed pathways. *Obesity Reviews*. 2011 20110315 [Epub ahead of print];12(5):e159-e172.

27. Ohkawara K, Tanaka S, Miyachi M et al. A dose-response relation between aerobic exercise and visceral fat reduction: systematic review of clinical trials. *International Journal of Obesity*. 2007 20070717 [Epub ahead of print];31(12):1786-97.
28. Osei-Assibey G, Kyrou I, Adi Y et al. Dietary and lifestyle interventions for weight management in adults from minority ethnic/non-White groups: a systematic review. *Obesity Reviews*. 2010;11(11):769-76.
29. Peters J, Sinn N, Campbell K et al. Parental influences on the diets of 2-5-year-old children: Systematic review of interventions. *Early Child Development and Care*. 2012;182(7):SP-837.
30. Public Health England. Obesity and the environment briefing: increasing physical activity and active travel. 2013. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/256796/Briefing_Obesity_and_active_travel_final.pdf.
31. Public Health England. Obesity and the environment: regulating the growth of fast food outlets. 2013. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/256655/Briefing_Obesity_and_fast_food_final.pdf.
32. Sarri KO, Tzanakis NE, Linardakis MK et al. Effects of Greek Orthodox Christian Church fasting on serum lipids and obesity. *BMC Public Health*. 2003 20030516 [Epub ahead of print];3:16.
33. Skouteris H, McCabe M, Ricciardelli LA et al. Parent-child interactions and obesity prevention: A systematic review of the literature. *Early Child Development and Care*. 2012;182(2):153-74.
34. Soler RE, Leeks KD, Buchanan LR et al. Point-of-decision prompts to increase stair use. A systematic review update. *American Journal of Preventive Medicine*. 2010;38(2 Suppl):S292-S300.
35. Thamotharan S, Lange K, Zale EL et al. The role of impulsivity in pediatric obesity and weight status: a meta-analytic review. *Clinical Psychology Review*. 2013 20121220 [Epub ahead of print];33(2):253-62.
36. Trowman R, Dumville JC, Hahn S et al. A systematic review of the effects of calcium supplementation on body weight. *British Journal of Nutrition*. 2006;95(6):1033-8.
37. U.S.Department of Agriculture. What is the relationship between the environment, body weight and fruit/vegetable consumption? (DGAC 2010). Washington (DC): U.S. Department of Agriculture; 2010.
38. U.S.Department of Agriculture. What is the relationship between the intake of soy protein and body weight? (DGAC 2010). Washington (DC): U.S. Department of Agriculture; 2010.
39. Vainik U, Dagher A, Dube L et al. Neurobehavioural correlates of body mass index and eating behaviours in adults: a systematic review. *Neuroscience & Biobehavioral Reviews*. 2013 20121219 [Epub ahead of print];37(3):279-99.
40. Vamosi M, Heitmann BL, Kyvik KO. The relation between an adverse psychological and social environment in childhood and the development of adult obesity: a systematic literature review. *Obesity Reviews*. 2010 20090730 [Epub ahead of print];11(3):177-84.

41. van der Horst K, Oenema A, Ferreira I et al. A systematic review of environmental correlates of obesity-related dietary behaviors in youth. *Health Education Research*. 2007 20060721 [Epub ahead of print];22(2):203-26.
42. van Sluijs EM, Kriemler S, McMinn AM. The effect of community and family interventions on young people's physical activity levels: A review of reviews and updated systematic review. *British Journal of Sports Medicine*. 2011 21836175 [Epub ahead of print];45(11):SP-914.
43. Webber KJ, Loescher LJ. A systematic review of parent role modeling of healthy eating and physical activity for their young African American children. *Journal for Specialists in Pediatric Nursing: JSPN*. 2013 20130509 [Epub ahead of print];18(3):173-88.
44. Weng SF, Redsell SA, Swift JA et al. Systematic review and meta-analyses of risk factors for childhood overweight identifiable during infancy. *Archives of Disease in Childhood*. 2012 20121029 [Epub ahead of print];97(12):1019-26.
45. Westwood M, Fayter D, Hartley S et al. Childhood obesity: should primary school children be routinely screened? A systematic review and discussion of the evidence. *Archives of Disease in Childhood*. 2007;92(5):416-22.
46. Whiting S, Derbyshire E, Tiwari BK. Capsaicinoids and capsinoids. A potential role for weight management? A systematic review of the evidence. *Appetite*. 2012 20120522 [Epub ahead of print];59(2):341-8.
47. Winzenberg T, Shaw K, Fryer J et al. Calcium supplements in healthy children do not affect weight gain, height, or body composition. *Obesity*. 2007;15(7):1789-98.

Wrong Outcome (41 studies)

1. Atkin AJ, Gorely T, Biddle SJ et al. Interventions to promote physical activity in young people conducted in the hours immediately after school: A systematic review. *International Journal of Behavioral Medicine*. 2011 20658358 [Epub ahead of print];18(3):SP-176.
2. Ayala GX, Baquero B, Klinger S. A systematic review of the relationship between acculturation and diet among Latinos in the United States: implications for future research. *Journal of the American Dietetic Association*. 2008;108(8):1330-44.
3. Barnett A, Cerin E, Baranowski T. Active video games for youth: a systematic review. *Journal of Physical Activity & Health*. 2011;8(5):724-37.
4. Behringer M, Vom HA, Yue Z et al. Effects of resistance training in children and adolescents: A meta-analysis. *Pediatrics*. 2010 20974785 [Epub ahead of print];126(5):SP-e1199.
5. Biddiss E, Irwin J. Active video games to promote physical activity in children and youth: A systematic review. *Archives of Pediatrics & Adolescent Medicine*. 2010 20603468 [Epub ahead of print];164(7):SP-664.
6. Bird EL, Baker G, Mutrie N et al. Behavior change techniques used to promote walking and cycling: A systematic review. *Health Psychology*. 2013 23477577 [Epub ahead of print];32(8):SP-829.
7. Chapman CD, Benedict C, Brooks SJ et al. Lifestyle determinants of the drive to eat: a meta-analysis. *American Journal of Clinical Nutrition*. 2012 20120725 [Epub ahead of print];96(3):492-7.

8. Clark MJ, Slavin JL. The effect of fiber on satiety and food intake: a systematic review. *Journal of the American College of Nutrition*. 2013;32(3):200-11.
9. Conn VS, Hafdahl AR, Mehr DR. Interventions to increase physical activity among healthy adults: Meta-analysis of outcomes. *American Journal of Public Health*. 2011 21330590 [Epub ahead of print];101(4):SP-751.
10. Cruz-Ferreira A, Fernandes J, Laranjo L et al. A systematic review of the effects of Pilates method of exercise in healthy people. *Archives of Physical Medicine and Rehabilitation*. 2011 22030232 [Epub ahead of print];92(12):SP-2071.
11. De CM, De DE, De B, I et al. Correlates of energy balance-related behaviours in preschool children: a systematic review. *Obesity Reviews*. 2012;13 Suppl 1:13-28.
12. De MF, Van Lenthe FJ, Spittaels H et al. Interventions for promoting physical activity among European teenagers: A systematic review. *International Journal of Behavioral Nutritional and Physical Activity*. 2009 19961623 [Epub ahead of print];6:SP-82.
13. Fogelholm M. Physical activity, fitness and fatness: relations to mortality, morbidity and disease risk factors. A systematic review. *Obesity Reviews*. 2010 20090909 [Epub ahead of print];11(3):202-21.
14. Ford C, Ward D, White M. Television viewing associated with adverse dietary outcomes in children ages 2-6. *Obesity Reviews*. 2012 20120910 [Epub ahead of print];13(12):1139-47.
15. Hoehner CM, Ribeiro IC, Parra DC et al. Physical activity interventions in Latin America: Expanding and classifying the evidence. *American Journal of Preventive Medicine*. 2013 23415133 [Epub ahead of print];44(3 SUPPL. 3):SP-e31.
16. Kelley GA, Kelley KS. Aerobic exercise and lipids and lipoproteins in children and adolescents: a meta-analysis of randomized controlled trials. *Atherosclerosis*. 2007 20060630 [Epub ahead of print];191(2):447-53.
17. Kelley GA, Kelley KS. Impact of progressive resistance training on lipids and lipoproteins in adults: A meta-analysis of randomized controlled trials. *Preventive Medicine*. 2009 19013187 [Epub ahead of print];48(1):SP-9.
18. Kelley GA, Kelley KS, Roberts S et al. Efficacy of aerobic exercise and a prudent diet for improving selected lipids and lipoproteins in adults: A meta-analysis of randomized controlled trials. *BMC Medicine*. 2011 21676220 [Epub ahead of print];9:SP-74.
19. Kelley GA, Kelley KS, Roberts S et al. Comparison of aerobic exercise, diet or both on lipids and lipoproteins in adults: A meta-analysis of randomized controlled trials. *Clinical Nutrition*. 2012 22154987 [Epub ahead of print];31(2):SP-156.
20. Kelley GA, Kelley KS. Effects of diet, aerobic exercise, or both on non-HDL-C in adults: A meta-analysis of randomized controlled trials. *Cholesterol*. 2012 23198142 [Epub ahead of print];2012:840935.
21. Kelly N, Mazzeo S, Bean M. Systematic review of dietary interventions with college students: Directions for future research and practice. *Journal of Nutrition Education & Behavior*. 2013 23433967 [Epub ahead of print];45(4):SP-304.
22. Lachat C, Nago E, Verstraeten R et al. Eating out of home and its association with dietary intake: a systematic review of the evidence. *Obesity Reviews*. 2012 20111123 [Epub ahead of print];13(4):329-46.

23. Maitland C, Stratton G, Foster S et al. A place for play? The influence of the home physical environment on children's physical activity and sedentary behaviour. *International Journal of Behavioral Nutrition and Physical Activity*. 2013 23958282 [Epub ahead of print];10:SP-99.
24. Malik VS, Popkin BM, Bray GA et al. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care*. 2010 20100806 [Epub ahead of print];33(11):2477-83.
25. Maniccia DM, Davison KK, Marshall SJ et al. A meta-analysis of interventions that target children's screen time for reduction. *Pediatrics*. 2011 21708797 [Epub ahead of print];128(1):SP-e193.
26. Maniccia DM. Interventions to modify television viewing behavior of children: A review, meta-analysis, and key-informant interviews. *Dissertation Abstracts International: Section B: The Sciences and Engineering*. 2010;71:(5-B):3004.
27. Metcalf B, Henley W, Wilkin T. Effectiveness of intervention on physical activity of children: systematic review and meta-analysis of controlled trials with objectively measured outcomes (EarlyBird 54). *BMJ*. 2012 20120927 [Epub ahead of print];345:e5888.
28. Muller-Riemenschneider F, Reinhold T, Nocon M et al. Long-term effectiveness of interventions promoting physical activity: A systematic review. *Preventive Medicine*. 2008 18675845 [Epub ahead of print];47(4):SP-354.
29. Nettleton JA, McKeown NM, Kanoni S et al. Interactions of dietary whole-grain intake with fasting glucose- and insulin-related genetic loci in individuals of European descent: a meta-analysis of 14 cohort studies. *Diabetes Care*. 2010 20100806 [Epub ahead of print];33(12):2684-91.
30. Pearson N, Biddle SJ. Sedentary behavior and dietary intake in children, adolescents, and adults. A systematic review. *American Journal of Preventive Medicine*. 2011;41(2):178-88.
31. Peng W, Crouse JC, Lin JH. Using active video games for physical activity promotion: A systematic review of the current state of research. *Health Education & Behavior*. 2013 22773597 [Epub ahead of print];40(2):SP-171.
32. Schmidt ME, Haines J, O'Brien A et al. Systematic review of effective strategies for reducing screen time among young children. *Obesity*. 2012 20120105 [Epub ahead of print];20(7):1338-54.
33. Schubert MM, Desbrow B, Sabapathy S et al. Acute exercise and subsequent energy intake. A meta-analysis. *Appetite*. 2013 20121228 [Epub ahead of print];63:92-104.
34. Sepulveda MJ, Lu C, Sill S et al. An observational study of an employer intervention for children's healthy weight behaviors. *Pediatrics*. 2010 20101004 [Epub ahead of print];126(5):e1153-e1160.
35. Small L, Lane H, Vaughan L et al. A systematic review of the evidence: the effects of portion size manipulation with children and portion education/training interventions on dietary intake with adults. *Worldviews on Evidence-Based Nursing*. 2013 20120615 [Epub ahead of print];10(2):69-81.
36. Speck RM, Schmitz KH. Energy expenditure comparison: A pilot study of standing instead of sitting at work for obesity prevention. *Preventive Medicine: An International Journal Devoted to Practice and Theory*. 2011;.52(3-4).

37. Steeves JA, Thompson DL, Bassett DR et al. A review of different behavior modification strategies designed to reduce sedentary screen behaviors in children. *Journal of Obesity*. 2012 20110724 [Epub ahead of print];2012:379215.
38. Teixeira PJ, Carraca EV, Markland D et al. Exercise, physical activity, and self-determination theory: A systematic review. *International Journal of Behavioral Nutrition and Physical Activity*. 2012 22726453 [Epub ahead of print];9(78):SP.
39. U.S.Department of Agriculture. What are the health effects related to consumption of nuts? (DGAC 2010). Washington (DC): U.S. Department of Agriculture; 2010.
40. Verloigne M, Van Lippevelde W, Maes L et al. Family- and school-based correlates of energy balance-related behaviours in 10-12-year-old children: a systematic review within the ENERGY (European Energy balance Research to prevent excessive weight Gain among Youth) project. *Public Health Nutrition*. 2012 20120124 [Epub ahead of print];15(8):1380-95.
41. Wieringa NF, van der Windt HJ, Zuiker RR et al. Positioning functional foods in an ecological approach to the prevention of overweight and obesity. *Obesity Reviews*. 2008 20080226 [Epub ahead of print];9(5):464-73.

Wrong Study Type (51 studies)

1. Alio AP, Salihu HM, Berrings TJ et al. Obesity research and the forgotten African American child. *Ethnicity & Disease*. 2006;16(2):569-75.
2. Ben-Sefer E, Ben-Natan M, Ehrenfeld M. Childhood obesity: current literature, policy and implications for practice. *International Nursing Review*. 2009;56(2):166-73.
3. Bray GA, Popkin BM. Calorie-sweetened beverages and fructose: what have we learned 10 years later. *Pediatric Obesity*. 2013 20130429 [Epub ahead of print];8(4):242-8.
4. Butland B, Jebb S, Kopelman P et al. Tackling obesities: Future choices - Project report. London: Government Office for Science; 2007. Available from: <http://www.bis.gov.uk/assets/foresight/docs/obesity/17.pdf>.
5. Cappuccio FP, Taggart FM, Kandala NB et al. Meta-analysis of short sleep duration and obesity in children and adults. *Sleep*. 2008;31(5):619-26.
6. Centre for Diet and Activity Research (CEDAR). Evidence brief: Walking & cycling for transport. 2013. Available from: <http://www.cedar.iph.cam.ac.uk/wp-content/uploads/2012/05/Evidence-Brief-why-walk-and-cycle-v1.0.pdf>.
7. Cislak A, Safron M, Pratt M et al. Family-related predictors of body weight and weight-related behaviours among children and adolescents: a systematic umbrella review. *Child: Care, Health & Development*. 2012 20110713 [Epub ahead of print];38(3):321-31.
8. Conroy S, Ellis R, Murray C et al. An integrative review of Canadian childhood obesity prevention programmes. *Obesity Reviews*. 2007;8(1):61-7.
9. Department of Health. Healthy lives, healthy people: A call to action on obesity in England. 2011. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213720/dh_130487.pdf.
10. Department of Health. Physical activity guidelines in the UK: review and recommendations. London: Department of Health; 2010. Available from:

- https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213743/dh_128255.pdf.
11. Department of Health. Sedentary behaviour and obesity: review of the current scientific evidence. London: Department of Health; 2010. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213745/dh_128225.pdf.
 12. Department of Health. Start active, stay active: a report on physical activity from the four home countries' Chief Medical Officers. London: Department of Health; 2011. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216370/dh_128210.pdf.
 13. Elder SJ, Roberts SB. The effects of exercise on food intake and body fatness: a summary of published studies. *Nutrition Reviews*. 2007;65(1):1-19.
 14. Food Standards Agency. Saturated fat and energy intake programme. 2013. Available from: <http://www.food.gov.uk/multimedia/pdfs/satfatprog.pdf>.
 15. Forouhi NG, Sharp SJ, Du H et al. Dietary fat intake and subsequent weight change in adults: results from the European Prospective Investigation into Cancer and Nutrition cohorts. *American Journal of Clinical Nutrition*. 2009 20091014 [Epub ahead of print];90(6):1632-41.
 16. Gargallo FM, Quiles IJ, Basulto MJ et al. Evidence-based nutritional recommendations for the prevention and treatment of overweight and obesity in adults (FESNAD-SEEDO consensus document). The role of diet in obesity prevention (II/III). *Nutricion Hospitalaria*. 2012;27(3):800-32.
 17. Heikkila K, Fransson EI, Nyberg ST et al. Job strain and health-related lifestyle: findings from an individual-participant meta-analysis of 118000 working adults. *American Journal of Public Health*. 2013 20130516 [Epub ahead of print];103(11):2090-7.
 18. Horikawa C, Kodama S, Yachi Y et al. Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. *Preventive Medicine*. 2011 20110827 [Epub ahead of print];53(4-5):260-7.
 19. Institute of Medicine. Accelerating progress in obesity prevention: Solving the weight of the nation. 2012. Available from: <http://www.iom.edu/Reports/2012/Accelerating-Progress-in-Obesity-Prevention.aspx>.
 20. Institute of Medicine. Early childhood obesity prevention policies. 2011. Available from: <http://www.iom.edu/Reports/2011/Early-Childhood-Obesity-Prevention-Policies.aspx>.
 21. Institute of Medicine. Sleep disorders and sleep deprivation: An unmet public health problem. 2006. Available from: http://www.nap.edu/download.php?record_id=11617.
 22. Kaisari P, Yannakoulia M, Panagiotakos DB. Eating frequency and overweight and obesity in children and adolescents: a meta-analysis. *Pediatrics*. 2013 20130408 [Epub ahead of print];131(5):958-67.
 23. Kalra N, Newman M. A systematic map of the research on the relationship between obesity and sedentary behaviour in young people. 1706. 2009. Available from: <http://epi.ioe.ac.uk/cms/LinkClick.aspx?fileticket=fEdubxSgHd0%3d&tabid=2462&language=en-US>.

24. Maio GR, Manstead A, Verplanken B et al. Tackling obesity: Future choices - Lifestyle change - Evidence review. London: Government Office for Science; 2007. Available from: <http://www.bis.gov.uk/assets/foresight/docs/obesity/05.pdf>.
25. Maskarinec G, Takata Y, Pagano I et al. Trends and dietary determinants of overweight and obesity in a multiethnic population. *Obesity*. 2006;14(4):717-26.
26. McDonald SD. Management and prevention of obesity in adults and children. *Canadian Medical Association Journal*. 2007;176(8):1109-10.
27. Misra A, Sharma R, Gulati S et al. Consensus dietary guidelines for healthy living and prevention of obesity, the metabolic syndrome, diabetes, and related disorders in Asian Indians. *Diabetes Technology & Therapeutics*. 2011 20110413 [Epub ahead of print];13(6):683-94.
28. Monasta L, Batty GD, Cattaneo A et al. Early-life determinants of overweight and obesity: a review of systematic reviews. *Obesity Reviews*. 2010;11(10):695-708.
29. Moore CJ, Cunningham SA. Social position, psychological stress, and obesity: a systematic review. *Journal of the Academy of Nutrition & Dietetics*. 2012;112(4):518-26.
30. Muller MJ, Danielzik S, Pust S. School- and family-based interventions to prevent overweight in children. *Proc Nutr Soc*. 2005;64(2):249-54.
31. National Obesity Observatory. Preventing childhood obesity through lifestyle change interventions: A briefing paper for commissioners. 2009. Available from: http://www.noo.org.uk/securefiles/131122_1051//RF%20Child_Weight_Management_Services_prevention_paper_201109.pdf.
32. Piscopo S. The Mediterranean diet as a nutrition education, health promotion and disease prevention tool. *Public Health Nutrition*. 2009;12(9A):1648-55.
33. Pluim BM, Staal JB, Marks BL et al. Health benefits of tennis. *British Journal of Sports Medicine*. 2007 20070515 [Epub ahead of print];41(11):760-8.
34. Poobalan A, Taylor L, Clar C et al. Prevention of childhood obesity: A review of systematic reviews. Edinburgh: NHS Health Scotland; 1-1-2008. Available from: <http://www.healthscotland.com/documents/2819.aspx>.
35. Prentice-Dunn H, Prentice-Dunn S. Physical activity, sedentary behavior, and childhood obesity: a review of cross-sectional studies. *Psychology Health & Medicine*. 2012 20111014 [Epub ahead of print];17(3):255-73.
36. Sabate J, Wien M. Vegetarian diets and childhood obesity prevention. *American Journal of Clinical Nutrition*. 2010 20100317 [Epub ahead of print];91(5):1525S-9S.
37. Saneei P, Salehi-Abargouei A, Esmailzadeh A. Serum 25-hydroxy vitamin D levels in relation to body mass index: a systematic review and meta-analysis. *Obesity Reviews*. 2013 20130118 [Epub ahead of print];14(5):393-404.
38. Scientific Advisory Committee on Nutrition. Dietary reference values for energy. London: Scientific Advisory Committee on Nutrition; 2011. Available from: http://www.sacn.gov.uk/pdfs/sacn_dietary_reference_values_for_energy.pdf.
39. Scientific Advisory Committee on Nutrition. Statement on dietary fibre. London: Scientific Advisory Committee on Nutrition; 2008. Available from: http://www.sacn.gov.uk/pdfs/final_sacn_position_statement_for_website_dietary_fibre.pdf.

40. Shankar P, Ahuja S, Sriram K. Non-nutritive sweeteners: Review and update. *Nutrition*. 2013 20130708 [Epub ahead of print];29(11-12):1293-9.
41. Sherry B. Food behaviors and other strategies to prevent and treat pediatric overweight. *Int J Obes (Lond)*. 2005;29 Suppl 2:S116-S126.
42. Szajewska H, Ruszczynski M. Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Critical Reviews in Food Science & Nutrition*. 2010;50(2):113-9.
43. Tremblay MS, Leblanc AG, Carson V et al. Canadian Physical Activity Guidelines for the Early Years (aged 0-4 years). *Applied Physiology, Nutrition, & Metabolism = Physiologie Appliquee, Nutrition et Metabolisme*. 2012 20120327 [Epub ahead of print];37(2):345-69.
44. Tremblay MS, Leblanc AG, Carson V et al. Canadian Sedentary Behaviour Guidelines for the Early Years (aged 0-4 years). *Applied Physiology, Nutrition, & Metabolism = Physiologie Appliquee, Nutrition et Metabolisme*. 2012 20120327 [Epub ahead of print];37(2):370-91.
45. U.S.Department of Agriculture. Report of the Dietary Guidelines Advisory Committee on the Dietary Guidelines for Americans, 2010. Washington (DC): U.S. Department of Agriculture; 2010. Available from: <http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2010/DGAC/Report/2010DGACReport-camera-ready-Jan11-11.pdf>.
46. U.S.Department of Health and Human Services, U.S.Department of Agriculture. Dietary Guidelines for Americans, 2010 (7th Ed.). Washington (DC): U.S. Department of Agriculture; 2010. Available from: <http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2010/PolicyDoc/PolicyDoc.pdf>; <http://www.cnpp.usda.gov/DGAs2010-PolicyDocument.htm>.
47. Vimalaswaran KS, Berry DJ, Lu C et al. Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Medicine / Public Library of Science*. 2013 20130205 [Epub ahead of print];10(2):e1001383.
48. Wiseman M. The second World Cancer Research Fund/American Institute for Cancer Research expert report. Food, nutrition, physical activity, and the prevention of cancer: a global perspective. [Review] [7 refs]. *Proceedings of the Nutrition Society*. 2008 20080501 [Epub ahead of print];67(3):253-6.
49. World Cancer Research Fund, American Institute for Cancer Research. Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. 2007. Available from: http://www.dietandcancerreport.org/cancer_resource_center/downloads/Second_Expert_Report_full.pdf.
50. World Health Organization. Global strategy on diet, physical activity and health. 2004. Available from: http://www.who.int/dietphysicalactivity/strategy/eb11344/strategy_english_web.pdf.
51. Yost J, Krainovich-Miller B, Budin W et al. Assessing weight perception accuracy to promote weight loss among U.S. female adolescents: a secondary analysis. *BMC Public Health*. 2010 20100809 [Epub ahead of print];10:465.

Wrong Question (70 studies)

1. Allender S, Cowburn G, Foster C. Understanding participation in sport and physical activity among children and adults: a review of qualitative studies. *Health Education Research*. 2006;21(6):826-35.
2. Almiron-Roig E, Palla L, Guest K et al. Factors that determine energy compensation: a systematic review of preload studies. *Nutrition Reviews*. 2013 20130610 [Epub ahead of print];71(7):458-73.
3. Amireault S, Godin G, Vezina-Im LA. Determinants of physical activity maintenance: A systematic review and meta-analyses. *Health Psychology Review*. 2013;7:(1):55-91.
4. Boylan S, Louie JC, Gill TP. Consumer response to healthy eating, physical activity and weight-related recommendations: a systematic review. *Obesity Reviews*. 2012 20120308 [Epub ahead of print];13(7):606-17.
5. Broekhuizen K, Kroeze W, van Poppel MN et al. A systematic review of randomized controlled trials on the effectiveness of computer-tailored physical activity and dietary behavior promotion programs: an update. *Annals of Behavioral Medicine*. 2012;44(2):259-86.
6. Brown AW, Bohan Brown MM, Allison DB. Belief beyond the evidence: using the proposed effect of breakfast on obesity to show 2 practices that distort scientific evidence. *American Journal of Clinical Nutrition*. 2013 20130904 [Epub ahead of print];98(5):1298-308.
7. Brunton G, Thomas J, Harden A et al. Promoting Physical Activity amongst Children Outside of Physical Education Classes: A Systematic Review Integrating Intervention Studies and Qualitative Studies. *Health Education Journal*. 2005;64(4):323-38.
8. Burrows T, Golley RK, Khambalia A et al. The quality of dietary intake methodology and reporting in child and adolescent obesity intervention trials: a systematic review. *Obesity Reviews*. 2012 20120815 [Epub ahead of print];13(12):1125-38.
9. Centers for Disease Control and Prevention. Recommended community strategies and measurements to prevent obesity in the United States. Atlanta (GA): Centers for Disease Control and Prevention; 2009. Available from: <http://www.cdc.gov/mmwr/pdf/rr/rr5807.pdf>.
10. Cooper JA, Nguyen DD, Ruby BC et al. Maximal sustained levels of energy expenditure in humans during exercise. *Medicine & Science in Sports & Exercise*. 2011;43(12):2359-67.
11. Craigie AM, Lake AA, Kelly SA et al. Tracking of obesity-related behaviours from childhood to adulthood: A systematic review. *Maturitas*. 2011 20110915 [Epub ahead of print];70(3):266-84.
12. Cugelman B, Thelwall M, Dawes P. Online interventions for social marketing health behavior change campaigns: a meta-analysis of psychological architectures and adherence factors. *Journal of Medical Internet Research*. 2011 20110214 [Epub ahead of print];13(1):e17.
13. Doolen J, Alpert PT, Miller SK. Parental disconnect between perceived and actual weight status of children: a metasynthesis of the current research. *Journal of the American Academy of Nurse Practitioners*. 2009;21(3):160-6.

14. Dugas LR, Harders R, Merrill S et al. Energy expenditure in adults living in developing compared with industrialized countries: a meta-analysis of doubly labeled water studies. *American Journal of Clinical Nutrition*. 2011 20101215 [Epub ahead of print];93(2):427-41.
15. Enwald HP, Huotari ML. Preventing the obesity epidemic by second generation tailored health communication: an interdisciplinary review. *Journal of Medical Internet Research*. 2010 20100628 [Epub ahead of print];12(2):e24.
16. Fanning J, Mullen SP, McAuley E. Increasing physical activity with mobile devices: a meta-analysis. *Journal of Medical Internet Research*. 2012 20121121 [Epub ahead of print];14(6):e161.
17. Fisher E, Meidtner K, Angquist L et al. Influence of dietary protein intake and glycemic index on the association between TCF7L2 HapA and weight gain. *American Journal of Clinical Nutrition*. 2012 20120502 [Epub ahead of print];95(6):1468-76.
18. Flynn MA, McNeil DA, Maloff B et al. Reducing obesity and related chronic disease risk in children and youth: A synthesis of evidence with 'best practice' recommendations. *Obesity Reviews*. 2006 16371076 [Epub ahead of print];7(1 Suppl):SP-7.
19. Foley L, Maddison R, Olds T et al. Self-report use-of-time tools for the assessment of physical activity and sedentary behaviour in young people: systematic review. *Obesity Reviews*. 2012 20120320 [Epub ahead of print];13(8):711-22.
20. George ES, Kolt GS, Duncan MJ et al. A review of the effectiveness of physical activity interventions for adult males. *Sports Medicine*. 2012;42(4):281-300.
21. Giskes K, Avendano M, Brug J et al. A systematic review of studies on socioeconomic inequalities in dietary intakes associated with weight gain and overweight/obesity conducted among European adults. *Obesity Reviews*. 2010 20091103 [Epub ahead of print];11(6):413-29.
22. Gomersall SR, Rowlands AV, English C et al. The ActivityStat hypothesis: the concept, the evidence and the methodologies. *Sports Medicine*. 2013;43(2):135-49.
23. Hafekost K, Lawrence D, Mitrou F et al. Tackling overweight and obesity: does the public health message match the science?. *BMC Medicine*. 2013 20130218 [Epub ahead of print];11:41.
24. Hamel LM, Robbins LB. Computer- and web-based interventions to promote healthy eating among children and adolescents: a systematic review. *Journal of Advanced Nursing*. 2013 20120704 [Epub ahead of print];69(1):16-30.
25. Hardy R, Cooper R, Aihie SA et al. Body mass index, muscle strength and physical performance in older adults from eight cohort studies: the HALCYON programme. *PLoS ONE [Electronic Resource]*. 2013 20130220 [Epub ahead of print];8(2):e56483.
26. Health Council of the Netherlands, Gezondheidsraad. For fat and thin. Prevention of overweight and obesity and the risk of eating disorders. Health Council of the Netherlands; 2010. Available from: <http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?ID=32010001752>; <http://www.gezondheidsraad.nl/sites/default/files/201013E.pdf> .
27. Hendrie GA, Brindal E, Corsini N et al. Combined home and school obesity prevention interventions for children: what behavior change strategies and intervention characteristics are associated with effectiveness? *Health Education and Behavior*. 2012;39(2):159-71.

28. Hendrie GA, Brindal E, Baird D et al. Improving children's dairy food and calcium intake: can intervention work? A systematic review of the literature. *Public Health Nutrition*. 2013 22607694 [Epub ahead of print];16(2):SP-365.
29. Hinkley T, Crawford D, Salmon J et al. Review and special article: Preschool Children and Physical Activity A Review of Correlates. *American Journal of Preventive Medicine*. 2008;34(5):435-41.
30. Institute of Medicine. Bridging the evidence gap in obesity prevention: A framework to inform decision making. Institute of Medicine; 2010. Available from: <http://www.iom.edu/Reports/2010/Bridging-the-Evidence-Gap-in-Obesity-Prevention-A-Framework-to-Inform-Decision-Making.aspx>.
31. Institute of Medicine. Evaluating obesity prevention efforts: A plan for measuring progress. Institute of Medicine; 2013. Available from: <http://www.iom.edu/Reports/2013/Evaluating-Obesity-Prevention-Efforts-A-Plan-for-Measuring-Progress.aspx>.
32. Kang M, Marshall SJ, Barreira TV et al. Effect of pedometer-based physical activity interventions: A meta-analysis. *Research Quarterly for Exercise & Sport*. 2009 19791652 [Epub ahead of print];80(3):SP-648.
33. Karoliina Enwald HP, Aulikki Huotari ML. Preventing the obesity epidemic by second generation tailored health communication: An interdisciplinary review. *Journal of Medical Internet Research*. 2010;12:(2):205-23.
34. Katz DL, Murimi M, Pretlow RA et al. Exploring effectiveness of messaging in childhood obesity campaigns. *Childhood Obesity*. 2012;8(2):97-105.
35. Kitzman-Ulrich H, Wilson DK, St George SM et al. The integration of a family systems approach for understanding youth obesity, physical activity, and dietary programs. *Clinical Child and Family Psychology Review*. 2010;13(3):231-53.
36. Kremers S, Reubsaet A, Martens M et al. Systematic prevention of overweight and obesity in adults: a qualitative and quantitative literature analysis. *Obesity Reviews*. 2010;11(5):371-9.
37. Kremers SPJ. Theory and practice in the study of influences on energy balance-related behaviors. *Patient Education and Counseling*. 2010;79(3):291-8.
38. Krieger JW, Sitren HS, Daniels MJ et al. Effects of variation in protein and carbohydrate intake on body mass and composition during energy restriction: a meta-regression 1. *American Journal of Clinical Nutrition*. 2006;83(2):260-74.
39. Latimer AE, Brawley LR, Bassett RL. A systematic review of three approaches for constructing physical activity messages: What messages work and what improvements are needed? *The International Journal of Behavioral Nutrition and Physical Activity*. 2010;7(Art ID 36).
40. Lorenc T, Brunton G, Oliver S et al. Attitudes to walking and cycling among children, young people and parents: a systematic review. *Journal of Epidemiology & Community Health*. 2008;62(10):852-7.
41. Malik VS, Hu FB. Popular weight-loss diets: from evidence to practice. *Nature Clinical Practice Cardiovascular Medicine*. 2007;4(1):34-41.
42. Martin J, Chater A, Lorencatto F. Effective behaviour change techniques in the prevention and management of childhood obesity. *International Journal of Obesity*. 2013 20130612 [Epub ahead of print];37(10):1287-94.

43. Mernagh P, Paech D, Coleman K et al. Assessing the cost-effectiveness of public health interventions to prevent obesity: Overview report. Wellington: Victoria University of Wellington; 2010. Available from: <http://www.victoria.ac.nz/sog/researchcentres/health-services-research-centre/docs/downloads/CE-Obesity-Prevention-Overview-Report-publish.pdf>.
44. National Obesity Observatory. Physical activity in the Active People Survey: developing an indicator that includes walking and cycling for transport. National Obesity Observatory; 2009. Available from: http://www.noo.org.uk/uploads/doc/vid_4727_Active_People_Survey_Analysis_NOO_w_ebsite_FINAL.pdf.
45. National Obesity Observatory. Physical activity surveillance in England: what is measured and where are the gaps? National Obesity Observatory; 2009. Available from: http://www.noo.org.uk/uploads/doc/vid_3953_noo_PhysicalActivity1_210909.pdf.
46. National Obesity Observatory. Data sources: knowledge of and attitudes towards healthy eating and physical activity. National Obesity Observatory; 2010. Available from: <http://www.noo.org.uk/gsf.php5?f=6055&fv=6317>.
47. National Obesity Observatory. Data sources: environmental influences on physical activity and diet. National Obesity Observatory; 2011. Available from: http://www.noo.org.uk/uploads/doc/vid_10418_Environmental%20%20data%20sources%20FINAL_editedformatted_%20MG%20100311.pdf.
48. National Obesity Observatory. Knowledge and attitudes towards healthy eating and physical activity: what the data tell us. National Obesity Observatory; 2011. Available from: http://www.noo.org.uk/uploads/doc/vid_11171_Attitudes.pdf.
49. Nguyen B, Kornman KP, Baur LA. A review of electronic interventions for prevention and treatment of overweight and obesity in young people. *Obesity Reviews*. 2011 20110223 [Epub ahead of print];12(5):e298-e314.
50. Niemeier BS, Hektner JM, Enger KB. Parent participation in weight-related health interventions for children and adolescents: a systematic review and meta-analysis. *Preventive Medicine*. 2012 20120507 [Epub ahead of print];55(1):3-13.
51. Paech D, Coleman K, Mernagh P et al. Assessing the cost-effectiveness of public health interventions to prevent obesity: A systematic review of the effectiveness of 16 obesity prevention interventions. Wellington: Victoria University of Wellington; 2010. Available from: <http://www.victoria.ac.nz/sog/researchcentres/health-services-research-centre/docs/downloads/Obesity-prevention-systematic-review-publish.pdf>.
52. Parry LL, Netuveli G, Parry J et al. A systematic review of parental perception of overweight status in children. *Journal of Ambulatory Care Management*. 2008;31(3):253-68.
53. Pinard CA, Yaroch AL, Hart MH et al. Measures of the home environment related to childhood obesity: a systematic review. *Public Health Nutrition*. 2012 20110907 [Epub ahead of print];15(1):97-109.
54. Pocock M, Trivedi D, Wills W et al. Parental perceptions regarding healthy behaviours for preventing overweight and obesity in young children: a systematic review of qualitative studies. *Obesity Reviews*. 2010 20090923 [Epub ahead of print];11(5):338-53.
55. Public Health England. Obesity and disability – adults. Public Health England; 2013. Available from: http://www.noo.org.uk/uploads/doc/vid_18474_obesity_dis.pdf.

56. Rees R, Kavanagh J, Harden A et al. Young people and physical activity: A systematic review matching their views to effective interventions. *Health Education Research*. 2006 1704102 [Epub ahead of print];21(6):SP-806.
57. Rees R, Oliver K, Woodman J et al. The views of young children in the UK about obesity, body size, shape and weight: a systematic review. *BMC Public Health*. 2011 20110325 [Epub ahead of print];11:188.
58. Renzaho AM, Mellor D, Boulton K et al. Effectiveness of prevention programmes for obesity and chronic diseases among immigrants to developed countries - a systematic review. *Public Health Nutrition*. 2010 20090902 [Epub ahead of print];13(3):438-50.
59. Rice H, Fauth R, Reeves A. What works in combating childhood obesity: an anthology of the literature on effective whole-system approaches: report for the East Midlands Joint Regional Improvement and Efficiency Plan. London: Centre for Excellence and Outcomes in Children and Young People's Services (C4EO); 2013. Available from: <http://www.c4eo.org.uk/themes/files/childhood-obesity-anthology-may2011.pdf>.
60. Robertson L, Douglas F, Ludbrook A et al. What works with men? A systematic review of health promoting interventions targeting men. *BMC Health Services Research*. 2008 18598339 [Epub ahead of print];8(141):SP.
61. Seo DC, Sa J. A meta-analysis of psycho-behavioral obesity interventions among US multiethnic and minority adults. *Preventive Medicine*. 2008 20080116 [Epub ahead of print];47(6):573-82.
62. Seo DC, Sa J. A meta-analysis of obesity interventions among U.S. minority children. *Journal of Adolescent Health*. 2010 20100113 [Epub ahead of print];46(4):309-23.
63. Skouteris H, McCabe M, Swinburn B et al. Parental influence and obesity prevention in pre-schoolers: a systematic review of interventions. *Obesity Reviews*. 2011;12(5):315-28.
64. Sosa ET. Mexican American Mothers' Perceptions of Childhood Obesity: A Theory-Guided Systematic Literature Review. *Health Education & Behavior*. 2012;39(4):396-404.
65. Sung-Chan P, Sung YW, Zhao X et al. Family-based models for childhood-obesity intervention: a systematic review of randomized controlled trials. *Obesity Reviews*. 2013;14(4):265-78.
66. Teixeira FV, Pais-Ribeiro JL, Maia AR. Beliefs and practices of healthcare providers regarding obesity: a systematic review. *Revista Da Associacao Medica Brasileira*. 2012;58(2):254-62.
67. Tudor-Locke C, Hart TL, Washington TL. Expected values for pedometer-determined physical activity in older populations. *International Journal of Behavioral Nutrition & Physical Activity*. 2009 20090825 [Epub ahead of print];6:59.
68. van Gerwen M, Franc C, Rosman S et al. Primary care physicians' knowledge, attitudes, beliefs and practices regarding childhood obesity: a systematic review. *Obesity Reviews*. 2009 20081011 [Epub ahead of print];10(2):227-36.
69. Wofford LG. Systematic review of childhood obesity prevention. *Journal of Pediatric Nursing*. 2008;23(1):5-19.
70. Wycherley TP, Moran LJ, Clifton PM et al. Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized

controlled trials. *American Journal of Clinical Nutrition*. 2012 20121024 [Epub ahead of print];96(6):1281-98.

Excluded primary studies (4 studies)

1. Baker RC, Kirschenbaum DS. Weight control during the holidays: highly consistent self-monitoring as a potentially useful coping mechanism. *Health Psychology*. 1998;17(4):367-70.
2. Kobayashi M, Kobayashi M. The relationship between obesity and seasonal variation in body weight among elementary school children in Tokyo. *Economics & Human Biology*. 2006 20050908 [Epub ahead of print];4(2):253-61.
3. Phelan S, Wing RR, Raynor HA et al. Holiday weight management by successful weight losers and normal weight individuals. *Journal of Consulting and Clinical Psychology*. 2008;76(3):442-8.
4. Stevenson JL, Krishnan S, Stoner MA et al. Effects of exercise during the holiday season on changes in body weight, body composition and blood pressure. *European Journal of Clinical Nutrition*. 2013;67(9):944-9.

Generic Reviews (39 studies)

1. Barr-Anderson DJ, Adams-Wynn AW, DiSantis KI et al. Family-focused physical activity, diet and obesity interventions in African-American girls: a systematic review. *Obesity Reviews*. 2013 20121012 [Epub ahead of print];14(1):29-51.
2. Bleich SN, Ku R, Wang YC. Relative contribution of energy intake and energy expenditure to childhood obesity: a review of the literature and directions for future research. *International Journal of Obesity*. 2011 20101130 [Epub ahead of print];35(1):1-15.
3. Bluford DA, Sherry B, Scanlon KS. Interventions to prevent or treat obesity in preschool children: A review of evaluated programs. *Obesity*. 2007;15(6):SP-1356.
4. Bond M, Wyatt K, Lloyd J et al. Systematic review of the effectiveness of weight management schemes for the under fives. *Obesity Reviews*. 2011;12(4):242-53.
5. Branscum P, Sharma M. A systematic analysis of childhood obesity prevention interventions targeting Hispanic children: lessons learned from the previous decade. *Obesity Reviews*. 2011 20101026 [Epub ahead of print];12(5):e151-e158.
6. Branscum P, Sharma M. After-school based obesity prevention interventions: a comprehensive review of the literature. *International Journal of Environmental Research & Public Health* [Electronic Resource]. 2012;9(4):1438-57.
7. Brown T, Avenell A, Edmunds LD et al. Systematic review of long-term lifestyle interventions to prevent weight gain and morbidity in adults. *Obesity Reviews*. 2009 20090914 [Epub ahead of print];10(6):627-38.
8. Campbell KJ, Hesketh KD. Strategies which aim to positively impact on weight, physical activity, diet and sedentary behaviours in children from zero to five years. A systematic review of the literature. *Obesity Reviews*. 2007;8(4):327-38.
9. Ciampa PJ, Kumar D, Barkin SL et al. Interventions aimed at decreasing obesity in children younger than 2 years: a systematic review. *Archives of Pediatrics & Adolescent Medicine*. 2010;164(12):1098-104.

10. Conn VS, Phillips LJ, Ruppert TM et al. Physical activity interventions with healthy minority adults: Meta-analysis of behavior and health outcomes. *Journal of Health Care for the Poor and Underserved*. 2012 22643462 [Epub ahead of print];23(1):SP-59.
11. Connelly JB, Duaso MJ, Butler G. A systematic review of controlled trials of interventions to prevent childhood obesity and overweight: a realistic synthesis of the evidence. *Public Health*. 2007 20070430 [Epub ahead of print];121(7):510-7.
12. Friedrich RR, Schuch I, Wagner MB. Effect of interventions on the body mass index of school-age students. *Revista de Saude Publica*. 2012 20120424 [Epub ahead of print];46(3):551-60.
13. Hebden L, Chey T, Allman-Farinelli M. Lifestyle intervention for preventing weight gain in young adults: a systematic review and meta-analysis of RCTs. *Obesity Reviews*. 2012 20120314 [Epub ahead of print];13(8):692-710.
14. Hesketh KD, Campbell KJ. Interventions to prevent obesity in 0-5 year olds: an updated systematic review of the literature. *Obesity*. 2010;18 Suppl 1:S27-S35.
15. Hutchesson MJ, Hulst J, Collins CE. Weight management interventions targeting young women: a systematic review. *Journal of the Academy of Nutrition & Dietetics*. 2013 20130307 [Epub ahead of print];113(6):795-802.
16. Hutfless S, Maruthur NM, Wilson RF et al. Strategies to prevent weight gain among adults. *American Journal of Preventive Medicine*. 2013;45(6):e41-51.
17. Jimenez-Pavon D, Kelly J, Reilly JJ. Associations between objectively measured habitual physical activity and adiposity in children and adolescents: Systematic review. *International Journal of Pediatric Obesity*. 2010;5(1):3-18.
18. Kamath CC, Vickers KS, Ehrlich A et al. Clinical review: behavioral interventions to prevent childhood obesity: a systematic review and metaanalyses of randomized trials. *Journal of Clinical Endocrinology & Metabolism*. 2008 20080909 [Epub ahead of print];93(12):4606-15.
19. Kay SJ, Fiatarone Singh MA. The influence of physical activity on abdominal fat: a systematic review of the literature. *Obesity Reviews*. 2006;7(2):183-200.
20. Kesten JM, Griffiths PL, Cameron N. A systematic review to determine the effectiveness of interventions designed to prevent overweight and obesity in pre-adolescent girls. *Obesity Reviews*. 2011 20110816 [Epub ahead of print];12(12):997-1021.
21. Laska MN, Pelletier JE, Larson NI et al. Interventions for weight gain prevention during the transition to young adulthood: a review of the literature. *Journal of Adolescent Health*. 2012;50(4):324-33.
22. Lemmens VE, Oenema A, Klepp KI et al. A systematic review of the evidence regarding efficacy of obesity prevention interventions among adults. *Obesity Reviews*. 2008 20080219 [Epub ahead of print];9(5):446-55.
23. Lombard CB, Deeks AA, Teede HJ. A systematic review of interventions aimed at the prevention of weight gain in adults. *Public Health Nutrition*. 2009 20090804 [Epub ahead of print];12(11):2236-46.
24. Perez-Morales ME, Bacardi-Gascon M, Jimenez-Cruz A. Childhood overweight and obesity prevention interventions among Hispanic children in the United States: systematic review. *Nutricion Hospitalaria*. 2012;27(5):1415-21.

25. Rauner A, Mess F, Woll A. The relationship between physical activity, physical fitness and overweight in adolescents: a systematic review of studies published in or after 2000. *BMC Pediatrics*. 2013 20130201 [Epub ahead of print];13:19.
26. Reichert FF, Baptista Menezes AM, Wells JC et al. Physical activity as a predictor of adolescent body fatness: a systematic review. *Sports Medicine*. 2009;39(4):279-94.
27. Saunders KL. Preventing obesity in pre-school children: a literature review. *Journal of Public Health*. 2007;29(4):368-75.
28. Showell NN, Fawole O, Segal J et al. A systematic review of home-based childhood obesity prevention studies. *Pediatrics*. 2013 20130610 [Epub ahead of print];132(1):e193-e200.
29. Stehr MD, von LT. Preventing weight gain through exercise and physical activity in the elderly: a systematic review. *Maturitas*. 2012 20120229 [Epub ahead of print];72(1):13-22.
30. Stevens CJ. Obesity prevention interventions for middle school-age children of ethnic minority: a review of the literature. *Journal for Specialists in Pediatric Nursing: JSPN*. 2010;15(3):233-43.
31. Stice E, Shaw H, Marti CN. A meta-analytic review of obesity prevention programs for children and adolescents: the skinny on interventions that work. *Psychological Bulletin*. 2006;132(5):667-91.
32. Swedish Council on Technology Assessment in Health Care. Interventions to prevent obesity: a systematic review. Stockholm: Swedish Council on Technology Assessment in Health Care; 2005. Available from: http://www.sbu.se/upload/Publikationer/Content1/1/SBU_Obesity.pdf.
33. Taylor PJ, Kolt GS, Vandelanotte C et al. A review of the nature and effectiveness of nutrition interventions in adult males--a guide for intervention strategies. *International Journal of Behavioral Nutrition & Physical Activity*. 2013 20130129 [Epub ahead of print];10:13.
34. Timmons BW, Leblanc AG, Carson V et al. Systematic review of physical activity and health in the early years (aged 0-4 years). *Applied Physiology, Nutrition, & Metabolism = Physiologie Appliquee, Nutrition et Metabolisme*. 2012;37(4):773-92.
35. Wane S, van Uffelen JG, Brown W. Determinants of weight gain in young women: a review of the literature. *Journal of Women's Health*. 2010;19(7):1327-40.
36. Wang Y, Wu Y, Wilson RF et al. Childhood obesity prevention programs: comparative effectiveness review and meta-analysis. *Comparative Effectiveness Review*; 115. Rockville (MD): Agency for Healthcare Research and Quality; 2013. Available from: <http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?ID=12013032050>.
37. Wareham NJ, van Sluijs EM, Ekelund U. Physical activity and obesity prevention: A review of the current evidence. *The Proceedings of the Nutrition Society*. 2005 15960868 [Epub ahead of print];64(2):SP-229.
38. Waters E, de Silva-Sanigorski A, Hall BJ et al. Interventions for preventing obesity in children. *Cochrane Database of Systematic Reviews*. 2011 20111207 [Epub ahead of print];(12):CD001871.
39. Wilks DC, Besson H, Lindroos AK et al. Objectively measured physical activity and obesity prevention in children, adolescents and adults: a systematic review of prospective studies. *Obesity Reviews*. 2011;12(5):e119-e129.

4 **Appendix D: Quality appraisal templates**

Rapid title and abstract quality appraisal criteria for systematic reviews

Reviews were given a score out of 4, by summing the total number of the following criteria it met:

- The review identifies itself in the title or abstract as a systematic review
- The review reports that it conforms to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) standards
- The abstract reports that two or more sources were searched
- The abstract reports the review's inclusion/exclusion criteria

These questions were based on criteria for definition of systematic reviews in the Database of Abstracts of Reviews of Effects (DARE).

Full text systematic review quality appraisal template

The checklist was based on the NICE systematic review quality checklist ([‘Methods for the development of NICE public health guidance \(third edition\)’](#)) and the Critical Appraisal Skills Programme (CASP) systematic review quality checklist. Reviews were awarded one point for each criterion met, with those scoring 0-2 rated as low quality [-], those scoring 3-5 rated as moderate quality [+], and those scoring 6-8 rated as high quality [++].

| | | | | |
|---|---|---|----|---------|
| Study identification (author, year, REFID) | | Total score (score 1 for each criterion met): | | |
| Factor: | | | | |
| Checklist completed by: | | | | |
| | | In this review this criterion is met: | | |
| 1. | Does the review address an appropriate and clearly-focused question that is relevant to 1 or more of the guidance topic's key questions? <i>Answer Yes if the review aimed to look at one of the specific behaviours being assessed rather than general areas (e.g. physical activity/diet/sedentary/lifestyle interventions)</i> | Yes | No | Unclear |
| 2. | Does the review include the types of study/s relevant to the key research question/s? <i>Study types related to our review question are RCTs and cohort studies.</i> <ul style="list-style-type: none"> <i>Answer Yes if these are the only study types included (if only RCTs or only cohorts/longitudinal observational answer Yes)</i> <i>Answer No if other study types are included (e.g. RCTs and cross-sectional)</i> <i>(Not penalising cohorts that don't specify prospective design)</i> | Yes | No | Unclear |
| 3. | Is the literature search sufficiently rigorous to identify all the relevant studies? <i>Must meet following criteria for a Yes:</i> <ul style="list-style-type: none"> <i>At least 2 electronic sources should be searched</i> <i>Must include years and databases searched</i> <i>Key words must be stated</i> | Yes | No | Unclear |
| 4. | Is the study quality of included studies appropriately assessed and reported? <i>Must meet following criteria for a Yes:</i> <ul style="list-style-type: none"> <i>Methods of assessment provided</i> <i>Quality of included studies reported</i> <i>Quality of included studies considered in conclusions</i> | Yes | No | Unclear |
| 5. | Is an adequate description of the | Yes | No | Unclear |

| | | | | |
|----|--|-----|----|---------|
| | <p>analytical methodology used included, and are the methods used appropriate to the question?</p> <ul style="list-style-type: none"> • <i>e.g. if meta-analysis is used is it appropriate and is heterogeneity assessed and taken into consideration if it exists</i> • <i>if mixed study types are included, are these analysed separately in the results section?</i> | | | |
| 6. | <p>Were the characteristics of the included studies provided? <i>E.g. In an aggregated form such as a table, data should be provided on the participants, interventions/ exposures and outcomes.</i></p> | Yes | No | Unclear |
| 7. | <p>Were potential conflicts of interest reported? <i>Potential sources of support should be clearly acknowledged for the systematic review and considered for the included studies.</i></p> | Yes | No | Unclear |
| 8. | <p>Can the results be applied to the UK population?</p> <ul style="list-style-type: none"> • <i>Answer Yes if majority of studies in OECD countries</i> • <i>If country not specified, consider context of research question to UK</i> | Yes | No | Unclear |

Primary study quality assessment template

Primary studies were assessed using the NICE checklist for quantitative studies reporting correlations and associations ([‘Methods for the development of NICE public health guidance \(third edition\)’](#)). Individual questions in the checklist sections 1 to 4 are rated as follows (in section 5 NR and NA are not options):

| | |
|---------------------|--|
| ++ | Indicates that for that particular aspect of study design, the study has been designed or conducted in such a way as to minimise the risk of bias. |
| + | Indicates that either the answer to the checklist question is not clear from the way the study is reported, or that the study may not have addressed all potential sources of bias for that particular aspect of study design. |
| - | Should be reserved for those aspects of the study design in which significant sources of bias may persist. |
| Not reported (NR) | Should be reserved for those aspects in which the study under review fails to report how they have (or might have) been considered. |
| Not applicable (NA) | Should be reserved for those study design aspects that are not applicable given the study design under review (for example, allocation concealment would not be applicable for case-control studies). |

| | | |
|---|---------|-----------|
| Study identification: Include full citation details | | |
| Study design: | | |
| <ul style="list-style-type: none"> Refer to the ‘Methods for the development of NICE public health guidance (third edition)’ glossary of study designs and the algorithm for classifying experimental and observational study designs to best describe the paper's underpinning study design | | |
| Guidance topic: | | |
| Assessed by: | | |
| Section 1: Population | | |
| 1.1 Is the source population or source area well described? | Rating: | Comments: |
| <ul style="list-style-type: none"> Was the country (e.g. developed or non-developed, type of health care system), setting (primary schools, community centres etc), location (urban, rural), population demographics etc adequately described? | | |
| 1.2 Is the eligible population or area representative of the source population or area? | Rating: | Comments: |
| <ul style="list-style-type: none"> Was the recruitment of individuals, clusters or areas well defined (e.g. advertisement, birth register)? Was the eligible population representative of the source? Were important groups underrepresented? | | |
| 1.3 Do the selected participants or areas represent the eligible population or area? | Rating: | Comments: |
| <ul style="list-style-type: none"> Was the method of selection of participants from the eligible population well described? What % of selected individuals or clusters agreed to participate? | | |

| | | |
|---|---------|-----------|
| <p>Were there any sources of bias?</p> <ul style="list-style-type: none"> Were the inclusion or exclusion criteria explicit and appropriate? | | |
| Section 2: Method of selection of exposure (or comparison) group | | |
| <p>2.1 Selection of exposure (and comparison) group. How was selection bias minimised?</p> <ul style="list-style-type: none"> How was selection bias minimised? | Rating: | Comments: |
| <p>2.2 Was the selection of explanatory variables based on a sound theoretical basis?</p> <ul style="list-style-type: none"> How sound was the theoretical basis for selecting the explanatory variables? | Rating: | Comments: |
| <p>2.3 Was the contamination acceptably low?</p> <ul style="list-style-type: none"> Did any in the comparison group receive the exposure? If so, was it sufficient to cause important bias? | Rating: | Comments: |
| <p>2.4 How well were likely confounding factors identified and controlled?</p> <ul style="list-style-type: none"> Were there likely to be other confounding factors not considered or appropriately adjusted for? Was this sufficient to cause important bias? | Rating: | Comments: |
| <p>2.5 Is the setting applicable to the UK?</p> <ul style="list-style-type: none"> Did the setting differ significantly from the UK? | Rating: | Comments: |
| Section 3: Outcomes | | |
| <p>3.1 Were the outcome measures and procedures reliable?</p> <ul style="list-style-type: none"> Were outcome measures subjective or objective (e.g. biochemically validated nicotine levels ++ vs self-reported smoking -)? How reliable were outcome measures (e.g. inter- or intra-rater reliability scores)? Was there any indication that measures had been validated (e.g. validated against a gold standard measure or assessed for content validity)? | Rating: | Comments: |
| <p>3.2 Were the outcome measurements complete?</p> <ul style="list-style-type: none"> Were all or most of the study participants who met the defined study outcome definitions likely to have been identified? | Rating: | Comments: |
| <p>3.3 Were all the important outcomes assessed?</p> <ul style="list-style-type: none"> Were all the important benefits and harms assessed? Was it possible to determine the overall balance of benefits and harms of the intervention versus comparison? | Rating: | Comments: |
| <p>3.4 Was there a similar follow-up time in exposure and comparison groups?</p> <ul style="list-style-type: none"> If groups are followed for different lengths of time, then more events are likely to occur in the group followed-up for longer distorting the comparison. Analyses can be adjusted to allow for differences in length of follow-up (e.g. using person-years). | Rating: | Comments: |
| <p>3.5 Was follow-up time meaningful?</p> <ul style="list-style-type: none"> Was follow-up long enough to assess long-term benefits and harms? Was it too long, e.g. participants lost to follow-up? | Rating: | Comments: |
| Section 4: Analyses | | |
| <p>4.1 Was the study sufficiently powered to detect an intervention</p> | Rating: | Comments: |

| | | |
|---|---------|-----------|
| <p>effect (if one exists)?</p> <ul style="list-style-type: none"> • A power of 0.8 (i.e. it is likely to see an effect of a given size if one exists, 80% of the time) is the conventionally accepted standard. • Is a power calculation presented? If not, what is the expected effect size? Is the sample size adequate? | | |
| <p>4.2 Were multiple explanatory variables considered in the analyses?</p> <ul style="list-style-type: none"> • Were there sufficient explanatory variables considered in the analysis? | Rating: | Comments: |
| <p>4.3 Were the analytical methods appropriate?</p> <ul style="list-style-type: none"> • Were important differences in follow-up time and likely confounders adjusted for? | Rating: | Comments: |
| <p>4.6 Was the precision of association given or calculable? Is association meaningful?</p> <ul style="list-style-type: none"> • Were confidence intervals or p values for effect estimates given or possible to calculate? • Were CIs wide or were they sufficiently precise to aid decision-making? If precision is lacking, is this because the study is under-powered? | Rating: | Comments: |
| Section 5: Summary | | |
| <p>5.1 Are the study results internally valid (i.e. unbiased)?</p> <ul style="list-style-type: none"> • How well did the study minimise sources of bias (i.e. adjusting for potential confounders)? • Were there significant flaws in the study design? | Rating: | Comments: |
| <p>5.2 Are the findings generalisable to the source population (i.e. externally valid)?</p> <ul style="list-style-type: none"> • Are there sufficient details given about the study to determine if the findings are generalisable to the source population? • Consider: participants, interventions and comparisons, outcomes, resource and policy implications. | Rating: | Comments: |

5 **Appendix E: Summary table of non-prioritised reviews**

See attached document for Summary table of non-prioritised reviews.

6 **Appendix F: Evidence tables**

See attached document for evidence tables.