

Depression in adults: treatment and management

Appendix K: Clinical evidence flow charts

NICE Guideline <...>

Appendices

May 2018

Consultation draft

*Developed by the National Guideline
Alliance, hosted by the Royal College of
Obstetricians and Gynaecologists*

Disclaimer

Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and/or their guardian or carer.

Copyright

National Institute for Health and Care Excellence [2018]. All rights reserved. Subject to Notice of rights.


Contents

| | |
|---|----|
| Appendix K: Clinical evidence flow charts | 5 |
| K.1 Organisation and service delivery | 5 |
| K.1.1 Service delivery | 5 |
| K.1.2 Settings for care | 6 |
| K.2 RQ2.1/2.2 Acute treatment (NMA) | 7 |
| K.4 RQ2.1/2.2 Acute treatment (pairwise) | 8 |
| K.5 Further-line treatment | 9 |
| K.6 Chronic depressive symptoms | 10 |
| K.7 Complex depression | 11 |
| K.8 Psychotic depression | 12 |
| K.9 Relapse prevention | 13 |
| K.10 Access to services | 14 |

Appendix K: Clinical evidence flow charts


K.1 Organisation and service delivery

K.1.1 Service delivery




Update 2018

K.1.2 Settings for care




Update 2018

K.2 RQ2.1/2.2 Acute treatment (NMA)




2018

K.4 RQ2.1/2.2 Acute treatment (pairwise)




Update 2018

K.5 Further-line treatment




Update 2018

K.6 Chronic depressive symptoms




Update 2018


K.7 Complex depression



K.8 Psychotic depression




K.9 Relapse prevention



Update 2018

K.10 Access to services



Update 2018