

# Depression in adults: treatment and management

## NICE guideline: short version

### Draft for consultation, July 2017

**This guideline covers** identifying, treating and managing depression in people aged 18 and over. It recommends tailoring care and treatment based on the severity of a person's depression. It also includes advice on preventing relapse and managing complex and severe depression.

#### Who is it for?

- Healthcare professionals
- Other professionals who have direct contact with, or provide health and other public services for, people with depression
- Commissioners and providers of services for people with depression and their families and carers
- Adults with depression, their families and carers

This guideline will update and replace NICE guideline CG90 (published October 2009).

We have updated or added new recommendations on the treatment of new depressive episodes, further line treatment, treatment of chronic, psychotic and complex depression, preventing relapse and the organisation of and access to services.

You are invited to comment on the new and updated recommendations in this guideline. These are marked as:

- **[new 2017]** if the evidence has been reviewed and the recommendation has been added or updated **or**
- **[2017]** if the evidence has been reviewed but no change has been made to the recommended action.

You are also invited to comment on recommendations that NICE proposes to delete from the 2009 guideline.

We have not updated recommendations shaded in grey, and cannot accept comments on them. In some cases, we have made minor wording changes for clarification.

See [Update information](#) for a full explanation of what is being updated.

This version of the guideline contains the draft recommendations, context and recommendations for research. Information about how the guideline was developed is on the [guideline's page](#) on the NICE website. This includes the guideline committee's discussion and the evidence reviews (in the [full guideline](#)), the scope, and details of the committee and any declarations of interest.

1

2

# 1 Contents

2	Contents.....	3
3	Recommendations .....	4
4	1.1 Experience of care .....	4
5	1.2 Recognition, assessment and initial management .....	5
6	1.3 Access to services .....	9
7	1.4 General principles of care .....	10
8	1.5 First-line treatment for less severe depression.....	16
9	1.6 First-line treatment for more severe depression.....	19
10	1.7 Behavioural couples therapy for depression .....	20
11	1.8 Relapse prevention .....	21
12	1.9 Limited response and treatment-resistant depression.....	23
13	1.10 Treating chronic depression.....	26
14	1.11 Treating complex depression .....	27
15	1.12 Treating psychotic depression .....	28
16	1.13 Electroconvulsive therapy .....	28
17	1.14 Coordination and delivery of care .....	31
18	Terms used in this guideline .....	33
19	Putting this guideline into practice .....	35
20	Context.....	37
21	Reason for the update .....	38
22	More information.....	38
23	Recommendations for research .....	39
24	1 Effectiveness of peer support for different severities of depression.....	39
25	2 Mechanisms of action of psychological interventions .....	39
26	3 Rate of relapse .....	40
27	4 Group based psychological treatments for preventing relapse .....	40
28	5 Increased access to services.....	41
29	Update information .....	42
30	Recommendations that have been deleted or changed.....	43
31		
32		

## 1 Recommendations

People have the right to be involved in discussions and make informed decisions about their care, as described in [your care](#).

[Making decisions using NICE guidelines](#) explains how we use words to show the strength (or certainty) of our recommendations, and has information about prescribing medicines (including off-label use), professional guidelines, standards and laws (including on consent and mental capacity), and safeguarding.

### 2 **1.1 Experience of care**

#### 3 **Providing information and support**

4 **1.1.1** Make sure people with depression are aware of self-help groups,  
5 support groups and other local and national resources. [2004]

#### 6 **Advance decisions and statements**

7 **1.1.2** Consider developing advance decisions and advance statements  
8 collaboratively with people who have recurrent severe depression or  
9 depression with psychotic symptoms, and for those who have been  
10 treated under the Mental Health Act, in line with the **Mental Capacity**  
11 **Act**. Record the decisions and statements and include copies in the  
12 person's care plan in primary and secondary care, and give copies to  
13 the person and to their family or carer if the person agrees. [2009,  
14 amended 2017]

#### 15 **Supporting families and carers**

16 **1.1.3** When families or carers are involved in supporting a person with  
17 severe or chronic<sup>1</sup> depression, think about:

---

<sup>1</sup> Depression is described as 'chronic' if symptoms have been present more or less continuously for 2 years or more.

- 1 • providing written and verbal spoken information on depression
- 2 and its management, including how families or carers can support
- 3 the person
- 4 • offering a carer's assessment of their caring, physical and mental
- 5 health needs if needed
- 6 • providing information about local family or carer support groups
- 7 and voluntary organisations, and helping families or carers to
- 8 access them
- 9 • discussing with the person and their family or carer about
- 10 confidentiality and the sharing of information. [2009]

## 11 **Working with people from diverse ethnic and cultural backgrounds**

12 **1.1.4** Be respectful of, and sensitive to, diverse cultural, ethnic and

13 religious backgrounds when working with people with depression,

14 and be aware of the possible variations in the presentation of

15 depression these can cause. Ensure staff are competent in:

- 16 • culturally sensitive assessment
- 17 • using different explanatory models of depression
- 18 • addressing cultural and ethnic differences when developing and
- 19 implementing treatment plans
- 20 • working with families from diverse ethnic and cultural
- 21 backgrounds. [2009]

22 **1.1.5** Provide all interventions in the preferred language of the person with

23 depression if possible. [2004]

## 24 **1.2 Recognition, assessment and initial management**

25 **1.2.1** Be alert to possible depression (particularly in people with a past

26 history of depression or a chronic physical health problem with

27 associated functional impairment) and consider asking people who

28 may have depression if:

- 29 • during the last month, have they often been bothered by feeling
- 30 down, depressed or hopeless?

- 1 • during the last month, have they often been bothered by having  
2 little interest or pleasure in doing things? [2009]

3 1.2.2 If a person answers 'yes' to either of the depression identification  
4 questions (see recommendation 1.2.1) but the practitioner is not  
5 competent to perform a mental health assessment, refer the person  
6 to an appropriate professional who can. If this professional is not the  
7 person's GP, inform the person's GP about the referral. [2009]

8 1.2.3 If a person answers 'yes' to either of the depression identification  
9 questions (see recommendation 1.2.1) and the practitioner is  
10 competent to perform a mental health assessment, review the  
11 person's mental state and associated functional, interpersonal and  
12 social difficulties. [2009]

13 1.2.4 Consider using a validated measure (for example, for symptoms,  
14 functions and/or disability) when assessing a person with suspected  
15 depression to inform and evaluate treatment. [2009]

16 1.2.5 If a person has significant language or communication difficulties, (for  
17 example people with sensory or cognitive impairments), consider  
18 asking a family member or carer about the person's symptoms to  
19 identify possible depression. [2004, amended 2017]

20 (See also NICE's guideline on [mental health problems in people with](#)  
21 [learning disabilities.](#))

22 1.2.6 Conduct a comprehensive assessment that does not rely simply on a  
23 symptom count when assessing a person who may have depression.  
24 Take into account both the degree of functional impairment and/or  
25 disability associated with the possible depression and the length of  
26 the episode. [2009]

27 1.2.7 Think about how the factors below may have affected the  
28 development, course and severity of a person's depression in

1 addition to assessing symptoms and associated functional  
2 impairment:

- 3 • any history of depression and coexisting mental health or physical  
4 disorders
- 5 • any history of mood elevation (to determine if the depression may  
6 be part of bipolar disorder<sup>2</sup>)
- 7 • any past experience of, and response to, previous treatments
- 8 • the quality of interpersonal relationships
- 9 • living conditions, employment situation and social isolation. [2009,  
10 amended 2017]

## 11 Acquired cognitive impairments

12 1.2.8 When assessing a person with suspected depression:

- 13 • be aware of any acquired cognitive impairments
- 14 • if needed, consult with a relevant specialist when developing  
15 treatment plans and strategies. [2009, amended 2017]

16 1.2.9 When providing interventions for people with an acquired cognitive  
17 impairment who have a diagnosis of depression:

- 18 • if possible, provide the same interventions as for other people with  
19 depression
- 20 • if needed, adjust the method of delivery or length of the  
21 intervention to take account of the disability or impairment. [2009,  
22 amended 2017]

## 23 Depression with anxiety

24 1.2.10 When depression is accompanied by symptoms of anxiety, the first  
25 priority should usually be to treat the depression. When the person  
26 has an anxiety disorder and comorbid depression or depressive

---

<sup>2</sup> If needed, refer to NICE's guideline on [bipolar disorder: assessment and management](#).

1 symptoms, consult NICE guidance for the relevant anxiety disorder if  
2 available and consider treating the anxiety disorder first. [2004]

### 3 **Risk assessment and monitoring**

4 **1.2.11** Always ask people with depression directly about suicidal ideation  
5 and intent. If there is a risk of self-harm or suicide:

- 6 • assess whether the person has adequate social support and is  
7 aware of sources of help
- 8 • arrange help appropriate to the level of need
- 9 • advise the person to seek further help if the situation deteriorates.  
10 [2004]

11 **1.2.12** If a person with depression presents considerable immediate risk to  
12 themselves or others, refer them urgently to specialist mental health  
13 services. [2004]

14 **1.2.13** Advise people with depression of the potential for increased  
15 agitation, anxiety and suicidal ideation in the initial stages of  
16 treatment. Check if they have any of these symptoms and:

- 17 • ensure that the person knows how to seek help promptly
- 18 • review the person's treatment if they develop marked and/or  
19 prolonged agitation. [2004]

20 **1.2.14** Advise a person with depression and their family or carer to be  
21 vigilant for mood changes, negativity and hopelessness, and suicidal  
22 ideation, and to contact their practitioner if concerned. This is  
23 particularly important during high-risk periods, such as starting or  
24 changing treatment and at times of increased personal stress. [2004]

25 **1.2.15** If a person with depression is assessed to be at risk of suicide:

- 26 • take into account toxicity in overdose if an antidepressant is  
27 prescribed or the person is taking other medication; (if necessary,  
28 limit the amount of medicine available)



- 1 • consider increasing the level of support, such as more frequent
- 2 direct or telephone contacts
- 3 • consider referral to specialist mental health services. [2004]

#### 4 **Active monitoring**

5 **1.2.16** For people who do not want an intervention with less severe

6 depression, in particular those whose depressive symptoms are

7 improving, or people with subthreshold depressive symptoms:

- 8 • discuss the presenting problem(s) and any concerns that the
- 9 person may have
- 10 • provide information about the nature and course of depression
- 11 • arrange a further assessment, normally within 2 weeks
- 12 • make contact if the person does not attend follow-up
- 13 appointments. [2004]

### 14 **1.3 Access to services**

15 1.3.1 Commissioners and providers of mental health services should

16 consider using stepped care models for organising the delivery of

17 care and treatment of individuals with depression. Stepped care

18 pathways should:

- 19 • provide accessible information about the pathway, for example in
- 20 different languages and formats
- 21 • be accessible and acceptable to people using the services
- 22 • support the integrated delivery of services across primary and
- 23 secondary care
- 24 • have clear criteria for entry to the service
- 25 • have multiple entry points and ways to access the service,
- 26 including self-referral
- 27 • have agreed protocols for sharing information. [new 2017]

28 1.3.2 Commissioners and providers of mental health services should

29 ensure pathways are in place to support the coordination of care and

30 treatment of individuals with depression. Pathways should:

- 1 • promote easy access to, and uptake of, interventions in the
- 2 pathway
- 3 • allow for prompt assessment of adults with depression, including
- 4 assessment of severity and risk
- 5 • provide access to NICE-recommended interventions for
- 6 depression
- 7 • ensure coordination and continuity of care
- 8 • have routine collection of data on access to, uptake of, and
- 9 outcomes of the interventions in the pathway. [new 2017]

10 1.3.3 Commissioners and providers of mental health services should  
11 ensure pathways have the following in place for people with  
12 depression (in particular for men, older people, lesbian, gay, bisexual  
13 and transgender people and people from black, Asian and minority  
14 ethnic communities) to promote access and increased uptake of  
15 services:

- 16 • information about the pathway provided in a non-stigmatising way,  
17 using age and culturally appropriate language and formats
- 18 • services available outside normal working hours
- 19 • a range of different methods to engage with and deliver  
20 interventions, for example text messages, email, telephone and  
21 online
- 22 • services provided in community-based settings, for example in an  
23 individual's home, community centres, leisure centres, care  
24 homes, social centres and integrated clinics within primary care
- 25 • bilingual therapists or independent translators
- 26 • involvement of families/partners. [new 2017]

## 27 **1.4 General principles of care**

### 28 **All interventions**

29 1.4.1 Support people with depression to decide on their preferences for  
30 interventions by giving them:

- 1                   • information on what the interventions are, and the expected  
2                   outcomes
- 3                   • choice on the intervention type, how it will be delivered (face to  
4                   face or digitally), and where it will be delivered
- 5                   • the option, if possible, to choose the gender of the practitioner
- 6                   • information on what the next steps will be if the initial intervention  
7                   is not helpful.
- 8           1.4.2    Provide interventions for people with depression in a framework. This  
9                   should include:
- 10                   • an assessment of need
- 11                   • the development of a treatment plan
- 12                   • taking into account any physical health problems
- 13                   • regular liaison between healthcare professionals in specialist and  
14                   non-specialist settings
- 15                   • routine outcome monitoring and follow-up. [new 2017]
- 16           1.4.3    Use psychological and psychosocial treatment manuals<sup>3</sup> to guide the  
17                   form and length of interventions. [2017]
- 18           1.4.4    Consider using competence frameworks developed from treatment  
19                   manual(s) for psychological and psychosocial interventions to  
20                   support effective training delivery and supervision of interventions.  
21                   [2017]
- 22           1.4.5    For all interventions for people with depression:
- 23                   • use sessional outcome measures
- 24                   • review how well the treatment is working with the person
- 25                   • monitor and evaluate treatment adherence. [2017]

---

<sup>3</sup> Treatment manuals that have evidence for their efficacy from clinical trials are preferred.



- 1                   • how they may be affected when they stop taking antidepressant  
2                   medication, and how these effects can be minimised  
3                   • the fact that they cannot get addicted to antidepressant  
4                   medication. [2017]

5           1.4.9    Advise people taking antidepressant medication that although it is  
6                   not addictive, if they stop taking it, miss doses or don't take a full  
7                   dose, they may have discontinuation symptoms such as:

- 8                   • more mood changes  
9                   • restlessness  
10                  • problems sleeping  
11                  • unsteadiness  
12                  • sweating  
13                  • abdominal symptoms  
14                  • altered sensations.

15                   Explain that these discontinuation symptoms are usually mild and go  
16                   away after a week but can sometimes be severe, particularly if the  
17                   antidepressant medication is stopped suddenly. [2017]

18           1.4.10   When stopping an antidepressant medication, slowly reduce the  
19                   dose based on how long the person has been taking it. For example:

- 20                   • over several days if the person has been taking it for 2–8 weeks  
21                   • over several weeks if the person has been taking it for 2–12  
22                   months  
23                   • over several months if the person has been taking it for 12 months  
24                   or more. [new 2017]

25           1.4.11   If a person has discontinuation symptoms when they stop taking  
26                   antidepressant medication or lower their dose, reassure them that  
27                   they are not having a relapse of their depression. Explain that:

- 28                   • these symptoms are common

- 1                   • relapse does not usually happen as soon as you stop taking an  
2                   antidepressant or lower the dose
- 3                   • even if they start taking an antidepressant medication again or  
4                   increase their dose, the symptoms won't go away immediately.  
5                   [new 2017]
- 6           1.4.12    If a person has mild discontinuation symptoms when they stop taking  
7                   antidepressant medication:
- 8                   • monitor their symptoms  
9                   • keep reassuring them that such symptoms are common. [new  
10                   2017]
- 11          1.4.13    If a person has severe discontinuation symptoms, consider restarting  
12                   the original antidepressant medication at the dose that was  
13                   previously effective, or another antidepressant from the same class  
14                   with a longer half-life. Reduce the dose gradually while monitoring  
15                   symptoms. [new 2017]
- 16          1.4.14    When prescribing antidepressant medication for people with  
17                   depression who are under 30 years or are thought to be at increased  
18                   risk of suicide:
- 19                   • see them 1 week after starting the medication  
20                   • review them frequently until the risk of suicide is reduced. [2017]
- 21          1.4.15    Take into account toxicity in overdose when prescribing an  
22                   antidepressant medication for people at significant risk of suicide. Be  
23                   aware that:
- 24                   • tricyclic antidepressants (TCAs), except lofepramine, are  
25                   associated with the greatest risk in overdose
- 26                   • compared with other equally effective antidepressant medication  
27                   recommended for routine use in primary care, venlafaxine is  
28                   associated with a greater risk of death from overdose. [2017]

- 1            1.4.16      When prescribing antidepressant medication for older people (65  
2                                  years and over):
- 3                                  • consider prescribing them at a lower dose
  - 4                                  • take into account the person’s general physical health and  
5                                  possible interactions with any other medicines they may be taking
  - 6                                  • carefully monitor the person for side effects. [2017]
- 7            1.4.17      For people with depression taking lithium, monitor:
- 8                                  • renal and thyroid function and calcium levels before treatment and  
9                                  every 6 months during treatment, or more often if there is  
10                                  evidence of renal impairment
  - 11                                  • serum lithium levels 1 week after starting treatment and at each  
12                                  dose change until stable, and every 3 months after that. [2017]
- 13          1.4.18      Consider ECG monitoring in people taking lithium who have a high  
14                                  risk of cardiovascular disease. [2017]
- 15          1.4.19      For people with depression who are taking an antipsychotic<sup>4</sup>, monitor  
16                                  and review:
- 17                                  • weight, initially and then weekly for the first 6 weeks, then at  
18                                  12 weeks, at 1 year and then annually (plotted on a chart)
  - 19                                  • lipid and glucose levels at 12 weeks, at 1 year and then annually
  - 20                                  • adverse effects, for example extrapyramidal side effects and  
21                                  prolactin-related side effects with risperidone. [2017]
- 22          1.4.20      Advise people with winter depression that follows a seasonal pattern  
23                                  and who wish to try light therapy in preference to antidepressant or

---

<sup>4</sup> At the time of consultation (July 2017), antipsychotics did not have a UK marketing authorisation for this indication. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council’s Prescribing guidance: prescribing unlicensed medicines for further information.

See individual SPCs for full list of monitoring requirements.

1 psychological treatment that the evidence for the efficacy of light  
2 therapy is uncertain. [2009]

3 1.4.21 Although there is evidence that St John's wort may be of benefit in  
4 less severe depression, practitioners should:

- 5 • not prescribe or advise its use by people with depression because  
6 of uncertainty about appropriate doses, persistence of effect,  
7 variation in the nature of preparations and potential serious  
8 interactions with other drugs (including oral contraceptives,  
9 anticoagulants and anticonvulsants)
- 10 • advise people with depression of the different potencies of the  
11 preparations available and of the potential serious interactions of  
12 St John's wort with other drugs [2004].

13 1.4.22 Do not routinely provide medication management on its own as an  
14 intervention for people with depression. [new 2017]

## 15 **1.5 First-line treatment for less severe depression**

16 In this guideline the term less severe depression includes the traditional categories  
17 of subthreshold symptoms, mild depression, and the lower half of moderate  
18 depression.

### 19 **Lower intensity psychological interventions**

20 1.5.1 Offer group-based cognitive behavioural therapy (CBT) specific to  
21 depression as the initial treatment for people with less severe  
22 depression. [new 2017]

23 1.5.2 Deliver group-based CBT that is:

- 24 • based on a cognitive behavioural model
- 25 • delivered by 2 competent practitioners
- 26 • consists of up to 9 sessions of 90 minutes each, for up to 12  
27 participants
- 28 • takes place over 12–16 weeks, including follow-up. [new 2017]



- 1            1.5.3       Offer individual self-help with support for people with less severe  
2                            depression who do not want group CBT. [new 2017]
- 3            1.5.4       Follow the principles of CBT when providing self-help with support. It  
4                            should:
- 5                            • provide age-appropriate, written, audio or digital (computer or  
6    online) material
- 7                            • have support from a trained practitioner who facilitates the self-  
8    help intervention, encourages completion and reviews progress  
9    and outcome
- 10                           • consist of up to 6 sessions (face-to-face or by telephone or  
11    online), each up to 30 minutes
- 12                           • take place over 9–12 weeks, including follow-up. [2017]
- 13            1.5.5       Consider a physical activity programme specifically designed for  
14                            people with depression who do not want group CBT or self-help with  
15                            support. [new 2017]
- 16            1.5.6       Ensure physical activity programmes for people with less severe  
17                            depression:
- 18                            • are delivered in groups by a competent practitioner
- 19                            • consist of 45 minutes of aerobic exercise of moderate intensity  
20    and duration twice a week for 5 weeks, then once a week for a  
21    further 7 weeks
- 22                            • usually have 8 people per group. [new 2017]

23    **Pharmacological interventions**

- 24            1.5.7       Consider a selective serotonin reuptake inhibitor (SSRI) or  
25                            mirtazapine for people with less severe depression who choose not  
26                            to have psychological interventions, or based on previous treatment  
27                            history for confirmed depression had a positive response to SSRIs or  
28                            mirtazapine or had a poor response to psychological interventions.  
29                            [new 2017]

1 **Higher intensity psychological interventions**

2 1.5.8 Offer individual CBT or behavioural activation (BA) if a person with  
3 less severe depression:

- 4 • has a history of poor response when they tried group CBT, a  
5 physical activity programme, facilitated self-help or antidepressant  
6 medication before **or**
- 7 • has responded well to CBT or BA before **or**
- 8 • is at risk of developing more severe depression, for example they  
9 have a history of severe depression or the current assessment  
10 suggests a more severe depression is developing. [new 2017]

11 1.5.9 Consider interpersonal therapy (IPT) if a person with less severe  
12 depression would like help for interpersonal difficulties that focus on  
13 role transitions or disputes or grief and:

- 14 • has had group CBT, exercise or facilitated self-help,  
15 antidepressant medication, individual CBT or BA for a previous  
16 episode of depression, but this did not work well for them, **or**
- 17 • does not want group CBT, exercise or facilitated self-help,  
18 antidepressant medication, individual CBT or BA. [new 2017]

19 1.5.10 Provide individual CBT, BA or IPT to treat less severe depression  
20 over 16 sessions, each lasting 50–60 minutes, over 3–4 months.  
21 [new 2017]

22 1.5.11 When giving individual CBT, BA or IPT, also consider providing:

- 23 • 2 sessions per week for the first 2–3 weeks of treatment for  
24 people with less severe depression
- 25 • 3–4 follow-up and maintenance sessions over 3–6 months after  
26 finishing the course for all people who have had individual CBT,  
27 BA or IPT. [new 2017]

1 1.5.12 Consider counselling if a person with less severe depression would  
2 like help for significant psychosocial, relationship or employment  
3 problems and:

- 4 • has had group CBT, exercise or facilitated self-help,  
5 antidepressant medication, individual CBT or BA for a previous  
6 episode of depression, but this did not work well for them, **or**  
7 • does not want group CBT, exercise or facilitated self-help,  
8 antidepressant medication, individual CBT or BA. [new 2017]

9 1.5.13 Ensure counselling for people with less severe depression:

- 10 • is based on a model developed specifically for depression  
11 • consists of up to 16 individual sessions each lasting up to an hour  
12 • takes place over 12 to 16 weeks, including follow-up. [new 2017]

13 1.5.14 Consider short-term psychodynamic therapy (STPT) if a person with  
14 less severe depression would like help for emotional and  
15 developmental difficulties in relationships and:

- 16 • has had group CBT, exercise or facilitated self-help,  
17 antidepressant medication or individual CBT for a previous  
18 episode of depression, but this did not work well for them, **or**  
19 • does not want group CBT, exercise or facilitated self-help,  
20 antidepressant medication or individual CBT. [new 2017]

21 1.5.15 Ensure STPT for people with less severe depression:

- 22 • is based on a model developed specifically for depression  
23 • consists of up to 16 individual sessions each lasting up to an hour  
24 • takes place over 12 to 16 weeks, including follow-up. [new 2017]

## 25 **1.6 First-line treatment for more severe depression**

26 In this guideline the term more severe depression includes the traditional categories  
27 of the upper half of moderate depression and severe depression.

- 1 1.6.1 Offer individual CBT in combination with an SSRI or mirtazapine as  
2 the initial treatment for more severe depression. [new 2017]
- 3 1.6.2 If a person with more severe depression does not want to take  
4 medication, offer:
- 5 • group CBT, **or**
  - 6 • individual CBT or BA if the person does not want group therapy.  
7 [new 2017]
- 8 1.6.3 If a person with more severe depression does not want psychological  
9 therapy, offer an SSRI or mirtazapine. [new 2017]
- 10 1.6.4 Consider short-term psychodynamic psychotherapy, alone or in  
11 combination with an SSRI or mirtazapine, for a person with more  
12 severe depression who would like help for emotional and  
13 developmental difficulties in relationships and:
- 14 • has had individual CBT in combination with an SSRI, group CBT,  
15 or individual CBT or BA for a previous episode of depression, but  
16 this did not work well for them, **or**
  - 17 • does not want individual CBT in combination with an SSRI, group  
18 CBT, or individual CBT or BA. [new 2017]

19 **1.7 Behavioural couples therapy for depression**

- 20 1.7.1 Consider behavioural couples therapy for a person with depression  
21 who has problems in the relationship with their partner if:
- 22 • the relationship problem(s) could be contributing to their  
23 depression **or**
  - 24 • involving their partner may help in the treatment of their  
25 depression. [new 2017]
- 26 1.7.2 Ensure behavioural couples therapy for people with depression:
- 27 • follows the behavioural principles for couples therapy
  - 28 • provides 15–20 sessions over 5–6 months. [2017]

1        **1.8        *Relapse prevention***

2        1.8.1        Discuss the likelihood of having a relapse with people who have  
3                    recovered from depression. Explain:

- 4                    • that a history of previous relapse increases the chance of further  
5                    relapses
- 6                    • the potential benefits of relapse prevention. [new 2017]

7        1.8.2        Take into account that the following may increase the risk of relapse:

- 8                    • how often a person has had episodes of depression, and how  
9                    recently
- 10                    • any other chronic physical health or mental health problems
- 11                    • any residual symptoms and unhelpful coping styles. for example  
12                    avoidance and rumination)
- 13                    • how severe their symptoms were, risk to self and if they had  
14                    functional impairment in previous episodes of depression
- 15                    • the effectiveness of previous interventions for treatment and  
16                    relapse prevention
- 17                    • personal, social and environmental factors. [new 2017]

18        1.8.3        For people who have recovered from less severe depression when  
19                    treated with medication (alone or in combination with a psychological  
20                    therapy), but are assessed as having a higher risk of relapse,  
21                    consider:

- 22                    • psychological therapy (CBT) with an explicit focus on relapse  
23                    prevention, typically 3–4 sessions over 1–2 months
- 24                    • continuing their medication. [new 2017]

25        1.8.4        For people who have recovered from more severe depression when  
26                    treated with medication (alone or in combination with a psychological  
27                    therapy), but are assessed as having a higher risk of relapse, offer:

- 28                    • a psychological therapy [mindfulness-based cognitive therapy  
29                    (MBCT) or group CBT] in combination with medication, **or**

- 1                     • psychological therapy (MBCT or group CBT) with a focus on  
2                     relapse prevention if the person wants to stop taking medication.  
3                     [new 2017]
- 4             1.8.5     For people who have recovered from depression when treated with a  
5                     psychological therapy, but are assessed as having a higher risk of  
6                     relapse, offer further psychological therapy (see recommendation  
7                     1.8.93). [new 2017]
- 8             1.8.6     For people who are continuing with medication to prevent relapse,  
9                     maintain the same dose unless there is good reason to reduce it  
10                    (such as adverse effects). [new 2017]
- 11            1.8.7     For people continuing with medication to prevent relapse, hold  
12                    reviews at 3, 6 and 12 months after maintenance treatment has  
13                    started. At each review:
- 14                    • monitor mood state using a formal validated rating scale, for  
15                    example the [PHQ-9](#)  
16                    • review side effects  
17                    • review any personal, social and environmental factors that may  
18                    impact on the risk of relapse  
19                    • agree the timescale for further review (no more than 12 months).  
20                    [new 2017]
- 21            1.8.8     At all further reviews for people continuing with antidepressant  
22                    medication to prevent relapse:
- 23                    • assess the risk of relapse  
24                    • discuss the need to continue with medication. [new 2017]
- 25            1.8.9     Offer group CBT (or MBCT for those who have had 3 or more  
26                    previous episodes of depression) for preventing relapse to people  
27                    who are assessed as being at higher risk of relapse and who  
28                    recovered with medication but who want to stop taking it. [new 2017]

1 1.8.10 When choosing a psychological therapy for preventing relapse for  
2 people who recovered with initial psychological therapy, offer:

- 3 • 4 more sessions of the same treatment if it has an explicit relapse  
4 prevention component, **or**
- 5 • group CBT (or MBCT for those who have had 3 or more previous  
6 episodes of depression) if initial psychological therapy had no  
7 explicit relapse prevention component. [new 2017]

8 1.8.11 Re-assess a person's risk of relapse when they finish a  
9 psychological relapse prevention intervention. Discuss the need for  
10 continuing treatment with the person if necessary. [new 2017]

11 1.8.12 Deliver MBCT for people assessed as having a higher risk of relapse  
12 in groups of up to 15 participants. Meetings should last 2 hours once  
13 a week for 8 weeks, with 4 follow-up sessions in the 12 months after  
14 treatment ends. [new 2017]

15 1.8.13 Deliver group CBT for people assessed as having a higher risk of  
16 relapse in groups of up to 12 participants. Sessions should last 2  
17 hours once a week for 8 weeks. [new 2017]

## 18 **1.9 Limited response and treatment-resistant depression**

19 1.9.1 If a person with depression has had no response or a limited  
20 response to initial treatment (within 3–4 weeks for antidepressant  
21 medication or 4–6 weeks for psychological therapy or combined  
22 medication and psychological therapy), assess:

- 23 • whether there are any personal or social factors that might explain  
24 why the treatment isn't working
- 25 • whether the person has not been adhering to the treatment plan,  
26 including any adverse effects of medication.

27 Work with the person to try and address any problems raised. [new  
28 2017]







1 1.9.9 For people with depression whose symptoms have not adequately  
2 responded to a combination of medication and a psychological  
3 therapy after 12 weeks, consider:

- 4 • alternatives to combined treatment (see recommendation 1.10.2)
- 5 • switching to a different psychological therapy, such as cognitive  
6 behavioural analysis system of psychotherapy (CBASP), CBT or  
7 MBCT (see recommendation 1.10.1). [new 2017]

8 1.9.10 If a person finds that their antidepressant medication is helping them  
9 but they are having side effects, consider switching to another  
10 antidepressant with a different side effect profile. [new 2017]

## 11 **1.10 Treating chronic depression**

12 1.10.1 For people with symptoms of chronic depression, consider cognitive  
13 behavioural treatments (CBASP and CBT) in combination with  
14 antidepressant medication. The cognitive behavioural treatment  
15 should:

- 16 • have a focus on chronic depressive symptoms
- 17 • cover related maintaining processes, for example avoidance,  
18 rumination and interpersonal difficulties. [new 2017]

19 1.10.2 If a person with chronic depression chooses not to have combined  
20 treatment, offer:

- 21 • an SSRI alone, **or**
- 22 • cognitive behavioural treatments (CBASP and CBT) alone. [new  
23 2017]

24 1.10.3 For people with chronic depression who cannot tolerate, or have not  
25 responded to, SSRI treatment, consider alternative medication in  
26 specialist settings, or after consulting a specialist. Alternatives  
27 include:

- 28 • tricyclic antidepressants, **or**

- 1 • moclobemide, **or**
- 2 • amisulpride<sup>7</sup>. [new 2017]

3 1.10.4 For people with chronic depression who have been assessed as  
4 likely to benefit from extra social or vocational support, consider:

- 5 • befriending in combination with existing antidepressant  
6 medication or psychological therapy; this should be done by  
7 trained volunteers, typically with at least weekly contact for  
8 between 2–6 months
- 9 • a rehabilitation programme, if their depression has led to loss of  
10 work or their withdrawing from social activities over the longer  
11 term. [2017]

12 1.10.5 For people with chronic or treatment-resistant depression who have  
13 not responded to the interventions recommended in section 1.9 and  
14 1.10 consider referral to a specialist mental health services for advice  
15 and further treatment. [new 2017]

## 16 **1.11 *Treating complex depression***

17 1.11.1 For people with complex depression (depression comorbid with a  
18 personality disorder), consider referral to a specialist personality  
19 disorder treatment programme. See NICE guidance on borderline  
20 personality disorder for recommendations on treatment for  
21 personality disorder with coexisting depression. [new 2017]

22 1.11.2 For people with complex depression who have not been able to  
23 access, not been helped by or chosen not to be treated in a  
24 specialist personality disorder programme, consider a combination of  
25 antidepressant medication and CBT. [new 2017]

---

<sup>7</sup> At the time of consultation (July 2017), amisulpride did not have a UK marketing authorisation for this indication. The prescriber should follow relevant professional guidance, taking full responsibility for the decision. Informed consent should be obtained and documented. See the General Medical Council's Prescribing guidance: prescribing unlicensed medicines for further information.

1 1.11.3 When delivering antidepressant medication and CBT combination  
2 treatment for people with complex depression:

- 3 • give the person support and encourage them to carry on with the  
4 treatment
- 5 • provide the treatment in a structured, multidisciplinary setting
- 6 • extend the duration of treatment if needed, up to a year. [new  
7 2017]

## 8 **1.12 *Treating psychotic depression***

9 1.12.1 Refer people with depression with psychotic symptoms to specialist  
10 mental health services for a programme of coordinated multi-  
11 disciplinary care, which includes access to psychological  
12 interventions.[new 2017]

13 1.12.2 When treating people with depression with psychotic symptoms,  
14 consider adding antipsychotic medication to their current treatment  
15 plan. [new 2017]

## 16 **1.13 *Electroconvulsive therapy***

17 1.13.1 Consider electroconvulsive therapy (ECT) for acute treatment of  
18 more severe depression if:

- 19 • the more severe depression is life-threatening and a rapid  
20 response is needed, **or**
- 21 • multiple pharmacological and psychological treatments have  
22 failed. [2017]

23 1.13.2 For people whose depression has not responded well to ECT  
24 previously, only consider a repeat trial of ECT after:

- 25 • reviewing the adequacy of the previous treatment course
- 26 • considering all other options
- 27 • discussing the risks and benefits with the person or, if appropriate,  
28 their advocate or carer. [2017]

1           1.13.3     Make sure people with depression who are going to have ECT are  
2                     fully informed of the risks, and with the risks and benefits specific to  
3                     them. Take into account:

- 4                       • the risks associated with a general anaesthetic
- 5                       • any medical comorbidities
- 6                       • potential adverse events, in particular cognitive impairment
- 7                       • if the person is older, the possible increased risk associated with
- 8                             ECT treatment for this age group
- 9                       • the risks associated with not having ECT.

10                    Document the assessment. [2017]

11           1.13.4     Make the decision to use ECT together with the person with  
12                     depression if they have the capacity to give consent. Take into  
13                     account the requirements of the Mental Health Act 2007 (if  
14                     applicable), and make sure:

- 15                       • valid, informed consent is given without pressure or coercion from  
16                             the circumstances or clinical setting
- 17                       • the person is aware of their right to change their mind and  
18                             withdraw consent at any time
- 19                       • there is strict adherence to recognised guidelines on consent, and  
20                             advocates or carers are involved to help informed discussions.
- 21                       [2017]

22           1.13.5     If a person with depression cannot give informed consent, only give  
23                     ECT if it does not conflict with an advance treatment decision the  
24                     person made. [2017]

25           1.13.6     For a person with depression who is going to have ECT, assess their  
26                     cognitive function:

- 27                       • before the first treatment
- 28                       • at least every 3–4 treatments
- 29                       • at the end of the treatment course. [2017]



1         **1.14        Coordination and delivery of care**

2         **Collaborative care**

3           1.14.1     Consider collaborative care for all older people with depression, in  
4                     particular if they have significant physical health problems or social  
5                     problems. [new 2017]

6           1.14.2     Consider collaborative care as a method for the delivery of care for  
7                     people with more severe depression. [new 2017]

8           1.14.3     Ensure that collaborative care for people with more severe  
9                     depression covers:

- 10                   • patient-centred assessment and engagement
- 11                   • symptom measurement and monitoring
- 12                   • medication management
- 13                   • active follow-up by a designated case manager
- 14                   • delivery of psychological and psychosocial interventions within a  
15                    structured protocol, for example stepped care
- 16                   • taking any relevant physical health problems into account
- 17                   • regular liaison with primary and secondary care colleagues
- 18                   • supervision of practitioner(s) by an experienced mental health  
19                    professional. [new 2017]

20         **Specialist care planning**

21           1.14.4     Refer people to specialist mental health services for a programme of  
22                     coordinated multidisciplinary care if they have:

- 23                   • more severe depression with multiple complicating problems, for  
24                    example unemployment, poor housing or financial problems, **or**
- 25                   • significant coexisting conditions. [new 2017]

26           1.14.5     Ensure multidisciplinary care plans for people with more severe  
27                     depression with multiple complicating problems, or significant  
28                     coexisting conditions:

- 1 • are developed together with the person, their GP and other
- 2 relevant people involved in their care (with the person’s
- 3 agreement)
- 4 • set out the roles and responsibilities of all health and social care
- 5 professionals involved in delivering the care
- 6 • include information about 24-hour support services, and how to
- 7 contact them
- 8 • include a crisis plan that identifies potential crisis triggers, and
- 9 strategies to manage those triggers
- 10 • are updated if there are any significant changes in the person's
- 11 needs or condition
- 12 • are reviewed at agreed regular intervals
- 13 • include medication management (a plan for starting, reviewing
- 14 and discontinuing medication). [new 2017]

15 **Crisis care and home treatment and inpatient care**

16 1.14.6 Consider crisis and intensive home treatment for people with more

17 severe depression who are at significant risk of:

- 18 • suicide, in particular for those who live alone
- 19 • self-harm
- 20 • harm to others
- 21 • self-neglect
- 22 • complications in response to their treatment, for example older
- 23 people with medical comorbidities. [new 2017]

24 1.14.7 Ensure teams providing crisis resolution and home treatment (CRHT)

25 interventions to support people with depression:

- 26 • monitor and manage risk as a high-priority routine activity
- 27 • establish and implement a treatment programme
- 28 • ensure continuity of any treatment programme while the person is
- 29 in contact with the CRHT team, and on discharge or transfer to
- 30 other services when this is needed



- 1                   • have a crisis management plan in place before the person is  
2                   discharged from the team's care. [new 2017]
- 3           1.14.8    Consider inpatient treatment for people with more severe depression  
4                   who cannot be adequately supported by a CRHT team. [new 2017]
- 5           1.14.9    Make the full range of recommended psychological therapies (group  
6                   CBT, CBT or BA) available for people with depression in inpatient  
7                   settings. [new 2017]
- 8           1.14.10   When providing psychological therapies for people with depression in  
9                   inpatient settings:
- 10                   • increase the intensity and duration of the interventions  
11                   • ensure that they continue to be provided effectively and promptly  
12                   on discharge. [new 2017]
- 13           1.14.11   Consider using CRHT teams with people with depression who might  
14                   benefit from early discharge from hospital after a period of inpatient  
15                   care. [2017]

## 16 ***Terms used in this guideline***

### 17 **Depression severity**

18 In all recommendations in this guideline the terms *less severe depression* and *more*  
19 *severe depression* are used. Depression severity exists along a continuum and is  
20 essentially composed of three elements - symptoms (which may vary in frequency  
21 and intensity), duration of the disorder and the impact on personal and social  
22 functioning. Severity of depression is therefore a consequence of the contribution of  
23 all of these elements. Traditionally depression severity has been grouped under 4  
24 categories: *severe depression* which is characterised by a large number of  
25 symptoms with a major negative impact on personal and social functioning;  
26 *moderate depression* which has a smaller number of symptoms with a more limited  
27 negative impact on personal and social functioning; *mild depression* which has a  
28 small number of symptoms with a limited impact on personal and social functioning  
29 and *sub-threshold depressive symptoms* which do not meet criteria for a diagnosis of

1 depression and which typically have little impact on personal and social functioning.  
2 In the development of the recommendations for this guideline the GC was concerned  
3 to develop a way of representing the severity of depression in the recommendations  
4 which best represents the available evidence on the classification and facilitates the  
5 uptake of the recommendations in routine clinical practice. They therefore decided to  
6 use the terms *less severe depression* which includes the traditional categories of  
7 subthreshold symptoms, mild depression, and the lower half of moderate depression  
8 and *more severe depression* which includes the traditional categories of the upper  
9 half of moderate depression and severe depression.

### 10 **Chronic depression**

11 Chronic depression is when a person continually meets criteria for the diagnosis of a  
12 major depressive episode for at least two years.

### 13 **Collaborative care**

14 Collaborative care requires that the service user and healthcare professional jointly  
15 identify problems and agree goals for interventions, and normally comprises:

- 16 • case management which is supervised and supported by a senior mental health  
17 professional
- 18 • close collaboration between primary and secondary physical health services and  
19 specialist mental health services in the delivery of services
- 20 • the provision of a range of evidence-based interventions
- 21 • the long term coordination of care and follow-up.

### 22 **Medication management**

23 Medication management is giving a person advice on how to keep to a regime for  
24 the use of medication (for example, how to take it, when to take it and how often).  
25 The focus in such programmes is only on the management of medication and not on  
26 other aspects of depression.

### 27 **Routine outcome monitoring**

28 This is a system for the monitoring of the outcomes of treatments which involves  
29 regular (usually at each contact) assessment of symptoms and functioning using a  
30 valid scale. It can inform both service user and practitioner of progress in treatment.

1 It is often supported by computerised delivery and scoring of the measures which  
2 ensures better completion of the questionnaires and service level audit and  
3 evaluation. Alternative terms such as “sessional outcome monitoring” or sessional  
4 outcomes” may also be used which emphasise that outcomes should be recorded at  
5 each contact.

## 6 **Stepped care**

7 This is a system of delivering and monitoring treatments, so that the most effective,  
8 least intrusive and least resource intensive treatments are delivered first. Stepped  
9 care has a built in ‘self-correcting’ mechanism so that people who do not benefit from  
10 initial interventions can be ‘stepped up’ to more intensive interventions as needed.

## 11 **Putting this guideline into practice**

12 **[This section will be completed after consultation]**

13 NICE has produced [tools and resources](#) **[link to tools and resources tab]** to help you  
14 put this guideline into practice.

15 **[Optional paragraph if issues raised]** Some issues were highlighted that might need  
16 specific thought when implementing the recommendations. These were raised during  
17 the development of this guideline. They are:

- 18 • [add any issues specific to guideline here]
- 19 • [Use 'Bullet left 1 last' style for the final item in this list.]

20 Putting recommendations into practice can take time. How long may vary from  
21 guideline to guideline, and depends on how much change in practice or services is  
22 needed. Implementing change is most effective when aligned with local priorities.

23 Changes recommended for clinical practice that can be done quickly – like changes  
24 in prescribing practice – should be shared quickly. This is because healthcare  
25 professionals should use guidelines to guide their work – as is required by  
26 professional regulating bodies such as the General Medical and Nursing and  
27 Midwifery Councils.

1 Changes should be implemented as soon as possible, unless there is a good reason  
2 for not doing so (for example, if it would be better value for money if a package of  
3 recommendations were all implemented at once).

4 Different organisations may need different approaches to implementation, depending  
5 on their size and function. Sometimes individual practitioners may be able to respond  
6 to recommendations to improve their practice more quickly than large organisations.

7 Here are some pointers to help organisations put NICE guidelines into practice:

8 1. **Raise awareness** through routine communication channels, such as email or  
9 newsletters, regular meetings, internal staff briefings and other communications with  
10 all relevant partner organisations. Identify things staff can include in their own  
11 practice straight away.

12 2. **Identify a lead** with an interest in the topic to champion the guideline and motivate  
13 others to support its use and make service changes, and to find out any significant  
14 issues locally.

15 3. **Carry out a baseline assessment** against the recommendations to find out  
16 whether there are gaps in current service provision.

17 4. **Think about what data you need to measure improvement** and plan how you  
18 will collect it. You may want to work with other health and social care organisations  
19 and specialist groups to compare current practice with the recommendations. This  
20 may also help identify local issues that will slow or prevent implementation.

21 5. **Develop an action plan**, with the steps needed to put the guideline into practice,  
22 and make sure it is ready as soon as possible. Big, complex changes may take  
23 longer to implement, but some may be quick and easy to do. An action plan will help  
24 in both cases.

25 6. **For very big changes** include milestones and a business case, which will set out  
26 additional costs, savings and possible areas for disinvestment. A small project group  
27 could develop the action plan. The group might include the guideline champion, a  
28 senior organisational sponsor, staff involved in the associated services, finance and  
29 information professionals.

1 **7. Implement the action plan** with oversight from the lead and the project group.

2 Big projects may also need project management support.

3 **8. Review and monitor** how well the guideline is being implemented through the  
4 project group. Share progress with those involved in making improvements, as well  
5 as relevant boards and local partners.

6 NICE provides a comprehensive programme of support and resources to maximise  
7 uptake and use of evidence and guidance. See our [into practice](#) pages for more  
8 information.

9 Also see Leng G, Moore V, Abraham S, editors (2014) Achieving high quality care –  
10 practical experience from NICE. Chichester: Wiley.

## 11 **Context**

12 Each year 6% of adults in England will experience an episode of depression, and  
13 more than 15% of people will experience an episode of depression over the course  
14 of their lifetime. For many people the episode will not be severe, but for more than  
15 20% the depression will be more severe and have a significant impact on their daily  
16 lives. Recurrence rates are high: there is a 50% chance of recurrence after a first  
17 episode, rising to 70% and 90% after a second or third episode, respectively.

18 Women are between 1.5 and 2.5 times more likely to be diagnosed with depression  
19 than men. However, although men are less likely to be diagnosed with depression,  
20 they are more likely to die by suicide, have higher levels of substance misuse, and  
21 are less likely to seek help than women.

22 The symptoms of depression can be disabling and the effects of the illness  
23 pervasive. Depression can have a major detrimental effect on a person's personal,  
24 social and work life. This places a heavy burden on the person and their carers and  
25 dependents, as well as placing considerable demands on the healthcare system.  
26 Depression is expected to become the second most common cause (after ischaemic  
27 heart disease) of loss of disability-adjusted life years in the world by 2020.

28 Depression is the leading cause of suicide, accounting for two-thirds of all deaths by  
29 suicide.

1 Under-treatment of depression is widespread, because many people are unwilling to  
2 seek help for depression and detection of depression by professionals is variable.  
3 For example, of the 130 people with depression per 1,000 population, only 80 will  
4 consult their GP. Of these 80 people, 49 are not recognised as having depression.  
5 This is mainly because they have contacted their GP because of a somatic symptom  
6 and do not consider themselves as having a mental health problem (despite the  
7 presence of symptoms of depression).

### 8 ***Reason for the update***

9 This update of NICE clinical guideline CG90 was commissioned because a review  
10 identified new evidence that might potentially change the recommendations for:

- 11 • service delivery (collaborative care)
- 12 • lower intensity psychological interventions for depression
- 13 • higher intensity psychological interventions for depression
- 14 • pharmacological interventions for moderate to severe depression.

### 15 ***More information***

[The following sentence is for post-consultation versions only – editor to update hyperlink with guideline number] You can also see this guideline in the NICE pathway on [\[pathway title\]](#).

To find out what NICE has said on topics related to this guideline, see our web page on [\[developer to add and link topic page title or titles; editors can advise if needed\]](#).

[The following sentence is for post-consultation versions only – editor to update hyperlink with guideline number] See also the guideline committee's discussion and the evidence reviews (in the [full guideline](#)), and information about [how the guideline was developed](#), including details of the committee.

16

## 1 **Recommendations for research**

2 The guideline committee has made the following recommendations for research. The  
3 committee's full set of research recommendations is detailed in the [full guideline](#).

### 4 ***1 Effectiveness of peer support for different severities of*** 5 ***depression***

6 Is peer support an effective and cost effective intervention in improving outcomes,  
7 including symptoms, personal functioning and quality of life in adults as a stand-  
8 alone intervention in people with less severe depression and as an adjunct to other  
9 evidence based interventions in more severe depression?

#### 10 **Why this is important**

11 Not all people with depression respond well to first-line treatments and for some  
12 people the absence of good social support systems may account for the limited  
13 response to first line interventions. A number of models for the provision of peer  
14 support have been developed in mental health which aim to provide direct personal  
15 support and help with establishing and maintaining supportive social networks. Peer  
16 support is provided by people who themselves have personal experience of a mental  
17 health problem. However, to date few studies have established and tested peer  
18 support models for people with depression. Peer support models, including both  
19 individual and group interventions, should be tested in a series of randomised  
20 controlled trials which examine the effectiveness of peer support for different  
21 severities of depression alone or in combination with evidence-based interventions  
22 for the treatment of depression.

### 23 ***2 Mechanisms of action of psychological interventions***

24 What are the mechanisms of action of effective psychological interventions for acute  
25 episodes of depression in adults?

#### 26 **Why this is important**

27 Depression is a debilitating and highly prevalent condition in adults. Despite  
28 significant investment, the most effective and well-established treatments have only  
29 modest effects on depressive symptoms, and the majority of treatment is for

1 recurrent depressive episodes. Research is required to identify the mechanism of  
2 action of the effective individual psychological treatments for depression, which  
3 would allow for the isolation of the most effective components and the development  
4 of better treatments. The research will need to be able to fully characterise the  
5 nature and range of depressive symptoms experienced by people and relate these to  
6 any proposed underlying neuropsychological mechanisms. The studies will also  
7 need to take into account the impact of any moderators of treatment effect. This  
8 research is necessary to improve clinical outcomes and quality of life for patients, as  
9 well as to reduce the financial burden upon the NHS.

### 10 **3 Rate of relapse**

11 What is the rate of relapse in people with depression who present, and are treated,  
12 in primary and secondary care, and what factors are associated with increased risk  
13 of relapse?

#### 14 **Why this is important**

15 The current understanding of the rate of relapse in depression is that it is high and  
16 may be up to 50% after a first episode, rising to 80% in people who have had three  
17 or more episodes of depression. However, most studies have been undertaken in  
18 the secondary care setting and whether these figures represent the actual rate of  
19 relapse in primary care populations is uncertain. In addition, beyond the number of  
20 previous episodes and the presence of residual symptoms there is also considerable  
21 uncertainty about what other factors (biological, psychological or social) might be  
22 associated with an increased risk of relapse. This cohort study will enable clinicians  
23 to more accurately identify those at risk of relapse, and provide relapse prevention  
24 strategies for these individuals. Accordingly, this would improve clinical outcomes  
25 and quality of life in patients as well as facilitating more targeted use of NHS  
26 resources.

### 27 **4 Group based psychological treatments for preventing relapse**

28 What is the comparative effectiveness and cost effectiveness of group based  
29 psychological treatments in preventing relapse in people with depression (compared  
30 to each other and antidepressant medication) for people who have had a successful  
31 course of treatment with antidepressants or psychological therapies?



1 **Why this is important**

2 Depressive relapse is a frequent occurrence with implications for the wellbeing and  
3 quality of life for the individual and financial implications for the NHS.  
4 Antidepressants can be effective in preventing relapse but not all service users can  
5 tolerate them or wish to take them long-term. Two, group based psychological  
6 interventions (group CBT and mindfulness based cognitive therapy) have been  
7 developed and shown to be effective primarily in trials when compared to treatment  
8 as usual. However, they have not been compared with each other and only in a  
9 limited way against antidepressants. The randomised controlled trial should be  
10 designed to identify both moderators and mediators of treatment effect, have a  
11 minimum follow up period of two years, assess any adverse events and the relative  
12 cost-effectiveness of the interventions.

13 **5 Increased access to services**

14 What are the most effective and cost effective methods to promote increased access  
15 to, and uptake of, interventions for people with depression who are under-  
16 represented in current services?

17 **Why this is important**

18 There is general under-recognition of depression but the problem is more marked in  
19 certain populations. In addition, even where depression is recognised by the person  
20 with depression or by health professionals, access to treatment can still be difficult. A  
21 number of factors may relate to this limited access including a person's view of their  
22 problems, the information available on services and the location, design and systems  
23 for referral to services. A number of studies have addressed this issue and a number  
24 of strategies have been developed to address it but no consistent picture has  
25 emerged from the research which can inform the design and delivery of services to  
26 promote access. Little is also known about how these systems might be tailored to  
27 the needs of particular groups such as older people, people from black, Asian and  
28 minority ethnic communities, and people with disabilities who may have additional  
29 difficulties in accessing services.

## 1 **Update information**

2 This guideline is an update of NICE guideline CG90 (published October 2009) and  
3 will replace it.

4 New recommendations have been added on treatment of new depressive episodes,  
5 further line treatment, treatment of chronic, psychotic and complex depression,  
6 preventing relapse and the organisation of and access to services.

7 These are marked as:

- 8 • **[new 2017]** if the evidence has been reviewed and the recommendation has been  
9 added or updated
- 10 • **[2017]** if the evidence has been reviewed but no change has been made to the  
11 recommended action.

12 NICE proposes to delete some recommendations from the 2009 guideline, because  
13 either the evidence has been reviewed and the recommendations have been  
14 updated, or NICE has updated other relevant guidance and has replaced the original  
15 recommendations. [Recommendations that have been deleted or changed](#) sets out  
16 these recommendations and includes details of replacement recommendations.  
17 Where there is no replacement recommendation, an explanation for the proposed  
18 deletion is given.

19 Where recommendations are shaded in grey and **end [2009 or 2004]**, the evidence  
20 has not been reviewed since the original guideline.

21 Where recommendations are shaded in grey and end **[2009 or 2004, amended**  
22 **2017]**, the evidence has not been reviewed but changes have been made to the  
23 recommendation wording that change the meaning (for example, because of  
24 equalities duties or a change in the availability of medicines, or incorporated  
25 guidance has been updated). These changes are marked with yellow shading, and  
26 explanations of the reasons for the changes are given in 'Recommendations that  
27 have been deleted or changed' for information.

28 See also the [original NICE guideline and supporting documents](#).

1 **Recommendations that have been deleted or changed**

2 **Recommendations to be deleted**

Recommendation in 2009 guideline	Comment
<p>When working with people with depression and their families or carers: build a trusting relationship</p> <ul style="list-style-type: none"> <li>• work in an open, engaging and non-judgemental manner</li> <li>• explore treatment options in an atmosphere of hope and optimism</li> <li>• explain the different courses of depression, and that recovery is possible</li> <li>• be aware that stigma and discrimination can be associated with a diagnosis of depression</li> <li>• ensure that discussions take place in settings that respect confidentiality, privacy and dignity. (1.1.1.1)</li> </ul>	<p>The concepts in these recommendations are now covered by NICE guidance on <a href="#">Service user experience in adult mental health services</a></p>
<p>When working with people with depression and their families or carers:</p> <ul style="list-style-type: none"> <li>• provide information suited to their level of understanding about the nature of depression and the range of treatments available</li> <li>• avoid clinical language and if it has to be used make sure it is clearly explained</li> <li>• ensure that comprehensive written information is available in an appropriate language (and also in audio format if possible )</li> <li>• provide, and work with, independent interpreters (that is, someone who is not known to the person with depression) if needed. (1.1.1.2)</li> </ul>	
<p>Make every effort to ensure that a person with depression can give meaningful and informed consent before treatment starts. This is especially important when a person has severe depression or their treatment falls under the Mental Health Act or the Mental Capacity Act. (1.1.1.4)</p>	
<p>Ensure that consent to treatment is based on the provision of clear information (which should also be</p>	

<p>available in written form) about the intervention, covering:</p> <ul style="list-style-type: none"> <li>• what the intervention is</li> <li>• what is expected of the person while they are having it</li> <li>• likely outcomes (including any side effects). (1.1.1.5)</li> </ul>	
<p>Offer people with depression advice on sleep hygiene if needed, including:</p> <ul style="list-style-type: none"> <li>• establishing regular sleep and wake times</li> <li>• avoiding excess eating, smoking or drinking alcohol before sleep</li> <li>• creating a proper environment for sleep taking regular physical exercise. (1.4.1.2)</li> </ul>	<p>Replaced by:</p> <p><i>First line treatment for less severe depression</i></p> <p>Offer group-based cognitive behavioural therapy (CBT) specific to depression as the initial treatment for people with less severe depression. [new 2017] (1.5.1)</p> <p>Deliver group-based CBT that is:</p> <ul style="list-style-type: none"> <li>• based on a cognitive behavioural model</li> </ul>
<p>For people with persistent subthreshold depressive symptoms or mild to moderate depression, consider offering one or more of the following interventions, guided by the person's preference:</p> <ul style="list-style-type: none"> <li>• individual guided self-help based on the principles of cognitive behavioural therapy (CBT)</li> <li>• computerised cognitive behavioural therapy (CCBT)</li> <li>• a structured group physical activity programme. (1.4.2.1)</li> </ul>	<ul style="list-style-type: none"> <li>• delivered by 2 competent practitioners</li> <li>• consists of up to 9 sessions of 90 minutes each, for up to 12 participants</li> <li>• takes place over 12–16 weeks, including follow-up. [new 2017] (1.5.2)</li> </ul> <p>Offer individual self-help with support for people with less severe depression who do not want group CBT. [new 2017] (1.5.3)</p> <p>Follow the principles of CBT when providing self-help with support. It should:</p> <ul style="list-style-type: none"> <li>• provide age-appropriate, written, audio or digital (computer or online) material</li> </ul>
<p>CCBT for people with persistent subthreshold depressive symptoms or mild to moderate depression should:</p> <ul style="list-style-type: none"> <li>• be provided via a stand-alone computer-based or web-based programme</li> <li>• include an explanation of the CBT model, encourage tasks between sessions, and use thought-challenging and active monitoring of behaviour, thought patterns and outcomes</li> <li>• be supported by a trained practitioner, who typically provides limited facilitation of the programme and reviews progress and outcome</li> <li>• typically take place over 9 to 12 weeks, including follow-up. (1.4.2.3)</li> </ul>	<ul style="list-style-type: none"> <li>• have support from a trained practitioner who facilitates the self-help intervention, encourages completion and reviews progress and outcome</li> <li>• consist of up to 6 sessions (face-to-face or by telephone or online), each up to 30 minutes</li> <li>• take place over 9–12 weeks, including follow-up. [2017] (1.5.4)</li> </ul> <p>Consider a physical activity programme specifically designed for people with depression who do not want group CBT or self-help with support. [new 2017] (1.5.5)</p>

<p>Physical activity programmes for people with persistent subthreshold depressive symptoms or mild to moderate depression should:</p> <ul style="list-style-type: none"> <li>• be delivered in groups with support from a competent practitioner</li> <li>• consist typically of three sessions per week of moderate duration (45 minutes to 1 hour) over 10 to 14 weeks (average 12 weeks). (1.4.2.4)</li> </ul>	<p>Ensure physical activity programmes for people with less severe depression:</p> <ul style="list-style-type: none"> <li>• are delivered in groups by a competent practitioner</li> <li>• consist of 45 minutes of aerobic exercise of moderate intensity and duration twice a week for 5 weeks, then once a week for a further 7 weeks</li> <li>• usually have 8 people per group. [new 2017] (1.5.6)</li> </ul>
<p>Consider group-based CBT for people with persistent subthreshold depressive symptoms or mild to moderate depression who decline low-intensity psychosocial interventions (1.4.3.1)</p>	<p>Consider a selective serotonin reuptake inhibitor (SSRI) or mirtazapine for people with less severe depression who choose not to have psychological interventions, or based on previous treatment history for confirmed depression had a positive response to SSRIs or mirtazapine or had a poor response to psychological interventions. [new 2017] (1.5.7)</p>
<p>Group-based CBT for people with persistent subthreshold depressive symptoms or mild to moderate depression should:</p> <ul style="list-style-type: none"> <li>• be based on a structured model such as ‘Coping with Depression’</li> <li>• be delivered by two trained and competent practitioners</li> <li>• consist of ten to 12 meetings of eight to ten participants</li> <li>• normally take place over 12 to 16 weeks, including follow-up. (1.4.3.2)</li> </ul>	<p>Offer individual CBT or behavioural activation (BA) if a person with less severe depression:</p> <ul style="list-style-type: none"> <li>• has a history of poor response when they tried group CBT, a physical activity programme, facilitated self-help or antidepressant medication before or</li> </ul>
<p>Do not use antidepressants routinely to treat persistent subthreshold depressive symptoms or mild depression because the risk–benefit ratio is poor, but consider them for people with:</p> <ul style="list-style-type: none"> <li>• a past history of moderate or severe depression or</li> <li>• initial presentation of subthreshold depressive symptoms that have been present for a long period (typically at least 2 years) or</li> <li>• subthreshold depressive symptoms or mild depression that persist(s) after other interventions. (1.4.4.1)</li> </ul>	<ul style="list-style-type: none"> <li>• has responded well to CBT or BA before or</li> <li>• is at risk of developing more severe depression, for example they have a history of severe depression or the current assessment suggests a more severe depression is developing. [new 2017] (1.5.8)</li> </ul>
<p>For people with moderate or severe depression, provide a combination of antidepressant medication and a high-intensity psychological intervention (CBT or IPT). (1.5.1.2)</p>	<p>Consider interpersonal therapy (IPT) if a person with less severe depression would like help for interpersonal difficulties that focus on role transitions or disputes or grief and:</p> <ul style="list-style-type: none"> <li>• has had group CBT, exercise or facilitated self-help, antidepressant medication, individual CBT or BA for a previous episode of depression, but this did not work well for them, or</li> </ul>

<p>The choice of intervention should be influenced by the:</p> <ul style="list-style-type: none"> <li>• duration of the episode of depression and the trajectory of symptoms</li> <li>• previous course of depression and response to treatment</li> <li>• likelihood of adherence to treatment and any potential adverse effects</li> <li>• person’s treatment preference and priorities. (1.5.1.3)</li> </ul>	<ul style="list-style-type: none"> <li>• does not want group CBT, exercise or facilitated self-help, antidepressant medication, individual CBT or BA. [new 2017] (1.5.9)</li> </ul> <p>Provide individual CBT, BA or IPT to treat less severe depression over 16 sessions, each lasting 50–60 minutes, over 3–4 months. [new 2017] (1.5.10)</p> <p>When giving individual CBT, BA or IPT, also consider providing:</p>
<p>When prescribing drugs other than SSRIs, take the following into account:</p> <ul style="list-style-type: none"> <li>• The increased likelihood of the person stopping treatment because of side effects (and the consequent need to increase the dose gradually) with venlafaxine, duloxetine and TCAs.</li> <li>• The specific cautions, contraindications and monitoring requirements for some drugs. For example: <ul style="list-style-type: none"> <li>– the potential for higher doses of venlafaxine to exacerbate cardiac arrhythmias and the need to monitor the person’s blood pressure</li> <li>– the possible exacerbation of hypertension with venlafaxine and duloxetine</li> <li>– the potential for postural hypotension and arrhythmias with TCAs</li> <li>– the need for haematological monitoring with mianserin in elderly people.</li> </ul> </li> <li>• Non-reversible monoamine oxidase inhibitors (MAOIs), such as phenelzine, should normally be prescribed only by specialist mental health professionals.</li> <li>• Dosulepin should not be prescribed. (1.5.2.4)</li> </ul>	<ul style="list-style-type: none"> <li>• 2 sessions per week for the first 2–3 weeks of treatment for people with less severe depression</li> <li>• 3–4 follow-up and maintenance sessions over 3–6 months after finishing the course for all people who have had individual CBT, BA or IPT. [new 2017] (1.5.11)</li> </ul> <p>Consider counselling if a person with less severe depression would like help for significant psychosocial, relationship or employment problems and:</p> <ul style="list-style-type: none"> <li>• has had group CBT, exercise or facilitated self-help, antidepressant medication, individual CBT or BA for a previous episode of depression, but this did not work well for them, or</li> <li>• does not want group CBT, exercise or facilitated self-help, antidepressant medication, individual CBT or BA. [new 2017] (1.5.12)</li> </ul> <p>Ensure counselling for people with less severe depression:</p> <ul style="list-style-type: none"> <li>• is based on a model developed specifically for depression</li> <li>• consists of up to 16 individual sessions each lasting up to an hour</li> <li>• takes place over 12 to 16 weeks, including follow-up. [new 2017] (1.5.13)</li> </ul> <p>Consider short-term psychodynamic therapy (STPT) if a person with less severe depression would like help for emotional and developmental difficulties in relationships and:</p>
<p>For people started on antidepressants who are not considered to be at increased risk of suicide, normally see them after 2 weeks. See them regularly</p>	<p>Consider short-term psychodynamic therapy (STPT) if a person with less severe depression would like help for emotional and developmental difficulties in relationships and:</p>

<p>thereafter; for example, at intervals of 2 to 4 weeks in the first 3 months, and then at longer intervals if response is good (1.5.2.6)</p>	<ul style="list-style-type: none"> <li>• has had group CBT, exercise or facilitated self-help, antidepressant medication or individual CBT for a previous episode of depression, but this did not work well for them, or</li> </ul>
<p>If a person with depression develops side effects early in antidepressant treatment, provide appropriate information and consider one of the following strategies:</p> <ul style="list-style-type: none"> <li>• monitor symptoms closely where side effects are mild and acceptable to the person or</li> <li>• stop the antidepressant or change to a different antidepressant if the person prefers or</li> <li>• in discussion with the person, consider short-term concomitant treatment with a benzodiazepine if anxiety, agitation and/or insomnia are problematic (except in people with chronic symptoms of anxiety); this should usually be for no longer than 2 weeks in order to prevent the development of dependence. (1.5.2.8)</li> </ul>	<ul style="list-style-type: none"> <li>• does not want group CBT, exercise or facilitated self-help, antidepressant medication or individual CBT. [new 2017] (1.5.14)</li> </ul> <p>Ensure STPT for people with less severe depression:</p> <ul style="list-style-type: none"> <li>• is based on a model developed specifically for depression</li> <li>• consists of up to 16 individual sessions each lasting up to an hour</li> <li>• takes place over 12 to 16 weeks, including follow-up. [new 2017] (1.5.15)</li> </ul> <p><i>First line treatment for more severe depression</i></p> <p>Offer individual CBT in combination with an SSRI or mirtazapine as the initial treatment for more severe depression. [new 2017] (1.6.1)</p>
<p>People who start on low-dose TCAs and who have a clear clinical response can be maintained on that dose with careful monitoring. (1.5.2.9)</p>	<p>If a person with more severe depression does not want to take medication, offer:</p>
<p>If the person's depression shows some improvement by 4 weeks, continue treatment for another 2 to 4 weeks. Consider switching to another antidepressant as described in 1.8 if:</p> <ul style="list-style-type: none"> <li>• response is still not adequate or</li> <li>• there are side effects or</li> <li>• the person prefers to change treatment. (1.5.2.12)</li> </ul>	<ul style="list-style-type: none"> <li>• group CBT, or</li> <li>• individual CBT or BA if the person does not want group therapy. [new 2017] (1.6.2)</li> </ul> <p>If a person with more severe depression does not want psychological therapy, offer an SSRI or mirtazapine. [new 2017] (1.6.3)</p>
<p>For all high-intensity psychological interventions, the duration of treatment should normally be within the limits indicated in this guideline. As the aim of treatment is to obtain significant improvement or remission the duration of treatment may be:</p> <ul style="list-style-type: none"> <li>• reduced if remission has been achieved</li> <li>• increased if progress is being made, and there is agreement between the practitioner and the person with</li> </ul>	<p>Consider short-term psychodynamic psychotherapy, alone or in combination with an SSRI or mirtazapine, for a person with more severe depression who would like help for emotional and developmental difficulties in relationships and:</p> <ul style="list-style-type: none"> <li>• has had individual CBT in combination with an SSRI, group CBT, or individual CBT or BA for a</li> </ul>

<p>depression that further sessions would be beneficial (for example, if there is a comorbid personality disorder or significant psychosocial factors that impact on the person's ability to benefit from treatment). (1.5.3.1)</p>	<p>previous episode of depression, but this did not work well for them, or</p> <ul style="list-style-type: none"> <li>• does not want individual CBT in combination with an SSRI, group CBT, or individual CBT or BA. [new 2017] (1.6.4)</li> </ul>
<p>For all people with depression having individual CBT, the duration of treatment should typically be in the range of 16 to 20 sessions over 3 to 4 months. Also consider providing:</p> <ul style="list-style-type: none"> <li>• two sessions per week for the first 2 to 3 weeks of treatment for people with moderate or severe depression</li> <li>• follow-up sessions typically consisting of three to four sessions over the following 3 to 6 months for all people with depression. (1.5.3.2)</li> </ul>	<p><i>Behavioural couples therapy</i></p> <p>Consider behavioural couples therapy for a person with depression who has problems in the relationship with their partner if:</p> <ul style="list-style-type: none"> <li>• the relationship problem(s) could be contributing to their depression or</li> <li>• involving their partner may help in the treatment of their depression. [new 2017] (1.7.1)</li> </ul>
<p>For all people with depression having IPT, the duration of treatment should typically be in the range of 16 to 20 sessions over 3 to 4 months. For people with severe depression, consider providing two sessions per week for the first 2 to 3 weeks of treatment. (1.5.3.3)</p>	<p>Ensure behavioural couples therapy for people with depression:</p> <ul style="list-style-type: none"> <li>• follows the behavioural principles for couples therapy</li> <li>• provides 15–20 sessions over 5–6 months. [2017] (1.7.2)</li> </ul>
<p>For all people with depression having behavioural activation, the duration of treatment should typically be in the range of 16 to 20 sessions over 3 to 4 months. Also consider providing:</p> <ul style="list-style-type: none"> <li>• two sessions per week for the first 3 to 4 weeks of treatment for people with moderate or severe depression</li> <li>• follow-up sessions typically consisting of three to four sessions over the following 3 to 6 months for all people with depression. (1.5.3.4)</li> </ul>	
<p>For all people with persistent subthreshold depressive symptoms or mild to moderate depression having counselling, the duration of treatment should typically be in the range of six to ten sessions over 8 to 12 weeks. (1.5.3.6)</p>	
<p>For all people with mild to moderate depression having short-term psychodynamic psychotherapy, the</p>	



<p>duration of treatment should typically be in the range of 16 to 20 sessions over 4 to 6 months. (1.5.3.7)</p>	
<p>Do not routinely vary the treatment strategies for depression described in this guideline either by depression subtype (for example, atypical depression or seasonal depression) or by personal characteristics (for example, sex or ethnicity) as there is no convincing evidence to support such action. (1.6.1.1)</p>	
<p>For people with persistent subthreshold depressive symptoms or mild to moderate depression who have not benefited from a low-intensity psychosocial intervention, discuss the relative merits of different interventions with the person and provide:</p> <ul style="list-style-type: none"> <li>• an antidepressant (normally a selective serotonin reuptake inhibitor [SSRI]) or</li> <li>• a high-intensity psychological intervention, normally one of the following options:             <ul style="list-style-type: none"> <li>○ CBT</li> <li>○ interpersonal therapy (IPT)</li> <li>○ behavioural activation (but note that the evidence is less robust than for CBT or IPT)</li> <li>○ behavioural couples therapy for people who have a regular partner and where the relationship may contribute to the development or maintenance of depression, or where involving the partner is considered to be of potential therapeutic benefit. (1.5.1.1)</li> </ul> </li> </ul>	<p>Replaced by:</p> <p>If a person with depression has had no response or a limited response to initial treatment (within 3–4 weeks for antidepressant medication or 4–6 weeks for psychological therapy or combined medication and psychological therapy), assess:</p> <ul style="list-style-type: none"> <li>• whether there are any personal or social factors that might explain why the treatment isn't working</li> <li>• whether the person has not been adhering to the treatment plan, including any adverse effects of medication.</li> </ul> <p>Work with the person to try and address any problems raised. [new 2017] (1.9.1)</p> <p>If a person has had no response or a limited response to initial treatment after assessing the issues in recommendation 1.9.1, provide more support by increasing the number and length of appointments. Also consider:</p> <ul style="list-style-type: none"> <li>• changing to a combination of psychological therapy and medication if the person is on medication only, or</li> <li>• changing to psychological therapy alone, if the person is on medication only and does not want to continue with medication or</li> <li>• changing to a combination of 2 different classes of medication, in specialist settings or after consulting a specialist, if the person is on medication only or a combination of medication and psychological therapy</li> </ul>
<p>If the person's depression shows no improvement after 2 to 4 weeks with the first antidepressant, check that the drug has been taken regularly and in the prescribed dose. (1.5.2.10)</p>	
<p>If response is absent or minimal after 3 to 4 weeks of treatment with a therapeutic dose of an antidepressant, increase the level of support (for example, by weekly</p>	

<p>face-to-face or telephone contact) and consider:</p> <ul style="list-style-type: none"> <li>• increasing the dose in line with the Summary of Product Characteristics if there are no significant side effects or</li> <li>• switching to another antidepressant as described in Section 1.8 if there are side effects or if the person prefers. (1.5.2.11)</li> </ul>	<p>and does not want to continue with psychological therapy. [new 2017] (1.9.2)</p> <p>When changing treatment for a person with depression who has had no response or a limited response to initial medication, consider:</p> <ul style="list-style-type: none"> <li>• combining the medication with a psychological therapy (CBT, BA, or IPT), or</li> </ul>
<p>When reviewing drug treatment for a person with depression whose symptoms have not adequately responded to initial pharmacological interventions:</p> <ul style="list-style-type: none"> <li>• check adherence to, and side effects from, initial treatment</li> <li>• increase the frequency of appointments using outcome monitoring with a validated outcome measure</li> <li>• be aware that using a single antidepressant rather than combination medication or augmentation (see 1.8.1.5 to 1.8.1.9) is usually associated with a lower side-effect burden</li> <li>• consider reintroducing previous treatments that have been inadequately delivered or adhered to, including increasing the dose</li> <li>• consider switching to an alternative antidepressant. (1.8.1.1)</li> </ul>	<ul style="list-style-type: none"> <li>• switching to a psychological therapy alone (CBT, BA, or IPT) if the person wants to stop taking medication. [new 2017] (1.9.3)</li> </ul> <p>If a person has had no response or a limited response to initial medication and does not want to try a psychological therapy, and wants to try a combination of medications, inform them of the likely increase in their side-effect burden (including risk of serotonin syndrome). [new 2017] (1.9.4)</p> <p>If a person wants to try a combination of medications and is willing to accept an increased side-effect burden, consider:</p> <ul style="list-style-type: none"> <li>• adding an antidepressant of a different class to their initial medication, for example an SSRI with mirtazapine, in specialist settings or after consulting a specialist</li> <li>• combining an antidepressant with an antipsychotic or lithium in specialist settings or after consulting a specialist. [new 2017] (1.9.5)</li> </ul>
<p>When switching to another antidepressant, be aware that the evidence for the relative advantage of switching either within or between classes is weak. Consider switching to:</p> <ul style="list-style-type: none"> <li>• initially a different SSRI or a better tolerated newer-generation antidepressant</li> <li>• subsequently an antidepressant of a different pharmacological class that may be less well tolerated, for example venlafaxine, a TCA or an MAOI. (1.8.1.2)</li> </ul>	<p>When changing treatment for a person with depression who has had no response or a limited response to initial psychological therapy, consider:</p> <ul style="list-style-type: none"> <li>• combining the psychological therapy with an SSRI, for example sertraline or citalopram, or mirtazapine, or</li> <li>• switching to an SSRI, for example sertraline or citalopram, or mirtazapine if the person wants to stop the psychological therapy. [new 2017] (1.9.6)</li> </ul>
<p>Do not switch to, or start, dosulepin because evidence supporting its tolerability relative to other antidepressants is outweighed by the</p>	

<p>increased cardiac risk and toxicity in overdose. (1.8.1.3)</p>	<p>If a person has had no response or a limited response to initial medication and does not want a psychological therapy or a combination of medications, consider:</p>
<p>When switching to another antidepressant, which can normally be achieved within 1 week when switching from drugs with a short half-life, consider the potential for interactions in determining the choice of new drug and the nature and duration of the transition. Exercise particular caution when switching:</p> <ul style="list-style-type: none"> <li>• from fluoxetine to other antidepressants, because fluoxetine has a long half-life (approximately 1 week)</li> <li>• from fluoxetine or paroxetine to a TCA, because both of these drugs inhibit the metabolism of TCAs; a lower starting dose of the TCA will be required, particularly if switching from fluoxetine because of its long half-life</li> <li>• to a new serotonergic antidepressant or MAOI, because of the risk of serotonin syndrome</li> <li>• from a non-reversible MAOI: a 2-week washout period is required (other antidepressants should not be prescribed routinely during this period). (1.8.1.4)</li> </ul>	<ul style="list-style-type: none"> <li>• continuing with the current medication, with extra support, close monitoring and an increased dose if the medication is well tolerated, or</li> <li>• switching to a medicine of a different class, or</li> <li>• switching to medication of the same class if there are problems with tolerability. [new 2017] (1.9.7)</li> </ul>
<p>When using combinations of medications (which should only normally be started in primary care in consultation with a consultant psychiatrist):</p> <ul style="list-style-type: none"> <li>• select medications that are known to be safe when used together</li> <li>• be aware of the increased side-effect burden this usually causes</li> <li>• discuss the rationale for any combination with the person with depression, follow GMC guidance if off-label medication is prescribed, and monitor carefully for adverse effects</li> <li>• be familiar with primary evidence and consider obtaining a second opinion when using unusual combinations, the evidence for the efficacy of a chosen strategy is limited or the risk–benefit ratio is unclear</li> <li>• document the rationale for the chosen combination. (1.8.1.5)</li> </ul>	<p>If a person’s symptoms do not respond to a dose increase or switching to another antidepressant after 2–4 weeks, review the need for care and treatment and consider consulting with, or referring the person to, a specialist service. [new 2017] (1.9.8)</p> <p>For people with depression whose symptoms have not adequately responded to a combination of medication and a psychological therapy after 12 weeks, consider:</p> <ul style="list-style-type: none"> <li>• alternatives to combined treatment (see recommendation 1.10.2)</li> <li>• switching to a different psychological therapy, such as cognitive behavioural analysis system of psychotherapy (CBASP), CBT or MBCT (see recommendation 1.10.1). [new 2017] (1.9.9)</li> </ul> <p>If a person finds that their antidepressant medication is helping them but they are having side effects, consider switching to another antidepressant with a different side effect profile. [new 2017] (1.9.10)</p>

<p>If a person with depression is informed about, and prepared to tolerate, the increased side-effect burden, consider combining or augmenting an antidepressant with:</p> <ul style="list-style-type: none"> <li>• lithium or</li> <li>• an antipsychotic such as aripiprazole, olanzapine, quetiapine or risperidone or</li> <li>• another antidepressant such as mirtazapine or mianserin. (1.8.1.6)</li> </ul>	
<p>The following strategies should not be used routinely:</p> <ul style="list-style-type: none"> <li>• augmentation of an antidepressant with a benzodiazepine for more than 2 weeks as there is a risk of dependence</li> <li>• augmentation of an antidepressant with buspirone, carbamazepine, lamotrigine or valproate as there is insufficient evidence for their use</li> <li>• augmentation of an antidepressant with pindolol or thyroid hormones as there is inconsistent evidence of effectiveness. (1.8.1.9)</li> </ul>	
<p>For a person whose depression has not responded to either pharmacological or psychological interventions, consider combining antidepressant medication with CBT. (1.8.1.10)</p>	
<p>For a person whose depression has failed to respond to various strategies for augmentation and combination treatments, consider referral to a practitioner with a specialist interest in treating depression, or to a specialist service. (1.8.1.11)</p>	
<p>The assessment of a person with depression referred to specialist mental health services should include:</p> <ul style="list-style-type: none"> <li>• their symptom profile, suicide risk and, where appropriate, previous treatment history</li> <li>• associated psychosocial stressors, personality factors and significant relationship difficulties, particularly where the depression is chronic or recurrent</li> </ul>	

<ul style="list-style-type: none"> <li>• associated comorbidities including alcohol and substance misuse, and personality disorders. (1.10.1.1)</li> </ul>	
<p>In specialist mental health services, after thoroughly reviewing previous treatments for depression, consider reintroducing previous treatments that have been inadequately delivered or adhered to. (1.10.1.2)</p>	
<p>Medication in secondary care mental health services should be started under the supervision of a consultant psychiatrist. (1.10.1.4)</p>	
<p>Discuss antidepressant treatment options with the person with depression, covering:</p> <ul style="list-style-type: none"> <li>• the choice of antidepressant, including any anticipated adverse events, for example, side effects and discontinuation symptoms (see Section 11.8.7.2) and potential interactions with concomitant medication or physical health problems</li> <li>• their perception of the efficacy and tolerability of any antidepressants they have previously taken. (1.5.2.1)</li> </ul>	<p>Replaced by:</p> <p>When offering a person antidepressant medication:</p> <ul style="list-style-type: none"> <li>• explain the reasons for offering it</li> <li>• discuss the risks and benefits</li> <li>• discuss any concerns they have about taking the medication</li> <li>• ensure they have information to take away that is appropriate for their needs. [2017] (1.4.7)</li> </ul> <p>When prescribing antidepressant medication, give people information about:</p> <ul style="list-style-type: none"> <li>• how long it takes (typically 2–4 weeks) to begin to start to feel better</li> <li>• how important it is to follow the instructions on when to take antidepressant medication</li> <li>• how treatment might need to carry on even after remission</li> <li>• how they may be affected when they first start taking antidepressant medication, and what these effects might be</li> <li>• how they may be affected if they have to take antidepressant medication for a long time and what these effects might be, especially in people over 65</li> <li>• how taking antidepressant medication might affect their sense of resilience (how strong they feel and how well they can get over problems) and being able to cope</li> </ul>

	<ul style="list-style-type: none"> <li>• how taking antidepressant medication might affect any other medicines they are taking</li> <li>• how they may be affected when they stop taking antidepressant medication, and how these effects can be minimised</li> <li>• the fact that they cannot get addicted to antidepressant medication. [2017] (1.4.8)</li> </ul>
<p>Inform the person that they should seek advice from their practitioner if they experience significant discontinuation symptoms. (1.9.2.3)</p>	<p>Issue covered by the new recommendations in section 1.4</p>
<p>For people with severe depression and those with moderate depression and complex problems, consider:</p> <ul style="list-style-type: none"> <li>• referring to specialist mental health services for a programme of co-ordinated multiprofessional care</li> <li>• providing collaborative care if the depression is in the context of a chronic physical health problem with associated functional impairment. (1.7.1.2)</li> </ul>	<p>Replaced by:</p> <p>Refer people to specialist mental health services for a programme of coordinated multidisciplinary care if they have:</p> <ul style="list-style-type: none"> <li>• more severe depression with multiple complicating problems, for example unemployment, poor housing or financial problems, or</li> <li>• significant coexisting conditions. [new 2017] (1.14.4)</li> </ul> <p>Ensure multidisciplinary care plans for people with more severe depression with multiple complicating problems, or significant coexisting conditions:</p> <ul style="list-style-type: none"> <li>• are developed together with the person, their GP and other relevant people involved in their care (with the person's agreement)</li> <li>• set out the roles and responsibilities of all health and social care professionals involved in delivering the care</li> <li>• include information about 24-hour support services, and how to contact them</li> <li>• include a crisis plan that identifies potential crisis triggers, and strategies to manage those triggers</li> <li>• are updated if there are any significant changes in the person's needs or condition</li> <li>• are reviewed at agreed regular intervals</li> <li>• include medication management (a plan for starting, reviewing and</li> </ul>

	<p>discontinuing medication). [new 2017] (1.14.6)</p>
<p>Support and encourage a person who has benefited from taking an antidepressant to continue medication for at least 6 months after remission of an episode of depression. Discuss with the person that:</p> <ul style="list-style-type: none"> <li>• this greatly reduces the risk of relapse</li> <li>• antidepressants are not associated with addiction. (1.9.1.1)</li> </ul>	<p>Replaced by:</p> <p>Discuss the likelihood of having a relapse with people who have recovered from depression. Explain:</p> <ul style="list-style-type: none"> <li>• that a history of previous relapse increases the chance of further relapses</li> <li>• the potential benefits of relapse prevention. [new 2017] (1.8.1)</li> </ul>
<p>Review with the person with depression the need for continued antidepressant treatment beyond 6 months after remission, taking into account:</p> <ul style="list-style-type: none"> <li>• the number of previous episodes of depression</li> <li>• the presence of residual symptoms</li> <li>• concurrent physical health problems and psychosocial difficulties. (1.9.1.2)</li> </ul>	<p>Take into account that the following may increase the risk of relapse:</p> <ul style="list-style-type: none"> <li>• how often a person has had episodes of depression, and how recently</li> <li>• any other chronic physical health or mental health problems</li> <li>• any residual symptoms and unhelpful coping styles. for example avoidance and rumination)</li> </ul>
<p>For people with depression who are at significant risk of relapse or have a history of recurrent depression, discuss with the person treatments to reduce the risk of recurrence, including continuing medication, augmentation of medication or psychological treatment (CBT). Treatment choice should be influenced by:</p> <ul style="list-style-type: none"> <li>• previous treatment history, including the consequences of a relapse, residual symptoms, response to previous treatment and any discontinuation symptoms</li> <li>• the person's preference. (1.9.1.3)</li> </ul>	<ul style="list-style-type: none"> <li>• how severe their symptoms were, risk to self and if they had functional impairment in previous episodes of depression</li> <li>• the effectiveness of previous interventions for treatment and relapse prevention</li> <li>• personal, social and environmental factors. [new 2017] (1.8.2)</li> </ul>
<p>Advise people with depression to continue antidepressants for at least 2 years if they are at risk of relapse. Maintain the level of medication at which acute treatment was effective (unless there is good reason to reduce the dose, such as unacceptable adverse effects) if:</p> <ul style="list-style-type: none"> <li>• they have had two or more episodes of depression in the recent past, during which they experienced significant functional impairment</li> <li>• they have other risk factors for relapse such as residual symptoms, multiple previous episodes, or a</li> </ul>	<p>For people who have recovered from less severe depression when treated with medication (alone or in combination with a psychological therapy), but are assessed as having a higher risk of relapse, consider:</p> <ul style="list-style-type: none"> <li>• psychological therapy (CBT) with an explicit focus on relapse prevention, typically 3–4 sessions over 1–2 months</li> <li>• continuing their medication. [new 2017] (1.8.3)</li> </ul> <p>For people who have recovered from more severe depression when treated with medication (alone or in combination with a psychological therapy), but are</p>

<p>history of severe or prolonged episodes or of inadequate response</p> <ul style="list-style-type: none"> <li>the consequences of relapse are likely to be severe (for example, suicide attempts, loss of functioning, severe life disruption, and inability to work). (1.9.1.4)</li> </ul>	<p>assessed as having a higher risk of relapse, offer:</p> <ul style="list-style-type: none"> <li>a psychological therapy [mindfulness-based cognitive therapy (MBCT) or group CBT] in combination with medication, or</li> </ul>
<p>When deciding whether to continue maintenance treatment beyond 2 years, re-evaluate with the person with depression, taking into account age, comorbid conditions and other risk factors. (1.9.1.5)</p>	<ul style="list-style-type: none"> <li>psychological therapy (MBCT or group CBT) with a focus on relapse prevention if the person wants to stop taking medication. [new 2017] (1.8.4)</li> </ul> <p>For people who have recovered from depression when treated with a psychological therapy, but are assessed as having a higher risk of relapse, offer further psychological therapy (see recommendation 1.8.3). [new 2017] (1.8.5)</p>
<p>People with depression on long-term maintenance treatment should be regularly re-evaluated, with frequency of contact determined by:</p> <ul style="list-style-type: none"> <li>comorbid conditions</li> <li>risk factors for relapse</li> <li>severity and frequency of episodes of depression. (1.9.1.6)</li> </ul>	<p>For people who are continuing with medication to prevent relapse, maintain the same dose unless there is good reason to reduce it (such as adverse effects). [new 2017] (1.8.6)</p>
<p>People who have had multiple episodes of depression, and who have had a good response to treatment with an antidepressant and an augmenting agent, should remain on this combination after remission if they find the side effects tolerable and acceptable. If one medication is stopped, it should usually be the augmenting agent. Lithium should not be used as a sole agent to prevent recurrence. (1.9.1.7)</p>	<p>For people continuing with medication to prevent relapse, hold reviews at 3, 6 and 12 months after maintenance treatment has started. At each review:</p> <ul style="list-style-type: none"> <li>monitor mood state using a formal validated rating scale, for example the PHQ-9</li> <li>review side effects</li> </ul>
<p>People with depression who are considered to be at significant risk of relapse (including those who have relapsed despite antidepressant treatment or who are unable or choose not to continue antidepressant treatment) or who have residual symptoms, should be offered the following psychological interventions:</p> <ul style="list-style-type: none"> <li>individual CBT for people who have relapsed despite antidepressant medication and for people with a significant history of depression and residual symptoms despite treatment</li> <li>mindfulness-based cognitive therapy for people who are currently well but have experienced three or more previous episodes of depression. (1.9.1.8)</li> </ul>	<ul style="list-style-type: none"> <li>review any personal, social and environmental factors that may impact on the risk of relapse</li> <li>agree the timescale for further review (no more than 12 months). [new 2017] (1.8.7)</li> </ul> <p>At all further reviews for people continuing with antidepressant medication to prevent relapse:</p> <ul style="list-style-type: none"> <li>assess the risk of relapse</li> <li>discuss the need to continue with medication. [new 2017] (1.8.8)</li> </ul> <p>Offer group CBT (or MBCT for those who have had 3 or more previous episodes of depression) for preventing relapse to people who are assessed as being at higher risk of relapse and who recovered</p>



<p>For all people with depression who are having individual CBT for relapse prevention, the duration of treatment should typically be in the range of 16 to 20 sessions over 3 to 4 months. If the duration of treatment needs to be extended to achieve remission it should:</p> <ul style="list-style-type: none"> <li>• consist of two sessions per week for the first 2 to 3 weeks of treatment</li> <li>• include additional follow-up sessions, typically consisting of four to six sessions over the following 6 months. (1.9.1.9)</li> </ul> <p>Mindfulness-based cognitive therapy should normally be delivered in groups of eight to 15 participants and consist of weekly 2-hour meetings over 8 weeks and four follow-up sessions in the 12 months after the end of treatment. (1.9.1.10)</p>	<p>with medication but who want to stop taking it. [new 2017] (1.8.9)</p> <p>When choosing a psychological therapy for preventing relapse for people who recovered with initial psychological therapy, offer:</p> <ul style="list-style-type: none"> <li>• 4 more sessions of the same treatment if it has an explicit relapse prevention component, or</li> <li>• group CBT (or MBCT for those who have had 3 or more previous episodes of depression) if initial psychological therapy had no explicit relapse prevention component. [new 2017] (1.8.10)</li> </ul> <p>Re-assess a person's risk of relapse when they finish a psychological relapse prevention intervention. Discuss the need for continuing treatment with the person if necessary. [new 2017] (1.8.11)</p> <p>Deliver MBCT for people assessed as having a higher risk of relapse in groups of up to 15 participants. Meetings should last 2 hours once a week for 8 weeks, with 4 follow-up sessions in the 12 months after treatment ends. [new 2017] (1.8.12)</p> <p>Deliver group CBT for people assessed as having a higher risk of relapse in groups of up to 12 participants. Sessions should last 2 hours once a week for 8 weeks. [new 2017] (1.8.13)</p>
<p>When stopping an antidepressant, gradually reduce the dose, normally over a 4-week period, although some people may require longer periods, particularly with drugs with a shorter half-life (such as paroxetine and venlafaxine). This is not required with fluoxetine because of its long half-life. (1.9.2.2)</p>	<p>When stopping an antidepressant medication, slowly reduce the dose based on how long the person has been taking it. For example:</p> <ul style="list-style-type: none"> <li>• over several days if the person has been taking it for 2–8 weeks</li> <li>• over several weeks if the person has been taking it for 2–12 months</li> <li>• over several months if the person has been taking it for 12 months or more. [new 2017] (1.4.10)</li> </ul>
<p>Inform the person that they should seek advice from their practitioner if they experience significant discontinuation symptoms. If discontinuation symptoms occur:</p>	<p>Replaced by:</p> <p>If a person has discontinuation symptoms when they stop taking antidepressant medication or lower their dose, reassure them that they are not having a relapse of their depression. Explain that:</p>

<ul style="list-style-type: none"> <li>• monitor symptoms and reassure the person if symptoms are mild</li> <li>• consider reintroducing the original antidepressant at the dose that was effective (or another antidepressant with a longer half-life from the same class) if symptoms are severe, and reduce the dose gradually while monitoring symptoms. (1.9.2.3)</li> </ul>	<ul style="list-style-type: none"> <li>• these symptoms are common</li> <li>• relapse does not usually happen as soon as you stop taking an antidepressant or lower the dose</li> <li>• even if they start taking an antidepressant medication again or increase their dose, the symptoms won't go away immediately. [new 2017] (1.4.11)</li> </ul> <p>If a person has mild discontinuation symptoms when they stop taking antidepressant medication:</p> <ul style="list-style-type: none"> <li>• monitor their symptoms</li> <li>• keep reassuring them that such symptoms are common. [new 2017] (1.4.12)</li> </ul> <p>If a person has severe discontinuation symptoms, consider restarting the original antidepressant medication at the dose that was previously effective, or another antidepressant from the same class with a longer half-life. Reduce the dose gradually while monitoring symptoms. [new 2017] (1.4.13)</p>
<p>Use crisis resolution and home treatment teams to manage crises for people with severe depression who present significant risk, and to deliver high-quality acute care. The teams should monitor risk as a high-priority routine activity in a way that allows people to continue their lives without disruption (1.10.1.3)</p>	<p>Replaced by:</p> <p>Consider crisis and intensive home treatment for people with more severe depression who are at significant risk of:</p> <ul style="list-style-type: none"> <li>• suicide, in particular for those who live alone</li> <li>• self-harm</li> </ul>
<p>Consider inpatient treatment for people with depression who are at significant risk of suicide, self-harm or self-neglect. (1.10.2.1)</p>	<ul style="list-style-type: none"> <li>• harm to others</li> <li>• self-neglect</li> </ul>
<p>The full range of high-intensity psychological interventions should normally be offered in inpatient settings. However, consider increasing the intensity and duration of the interventions and ensure that they can be provided effectively and efficiently on discharge. (1.10.2.2)</p>	<ul style="list-style-type: none"> <li>• complications in response to their treatment, for example older people with medical comorbidities. [new 2017] (1.14.6)</li> </ul> <p>Ensure teams providing crisis resolution and home treatment (CRHT) interventions to support people with depression:</p>
<p>Consider crisis resolution and home treatment teams for people with depression who might benefit from early discharge from hospital after a period of inpatient care. (1.10.2.3)</p>	<ul style="list-style-type: none"> <li>• monitor and manage risk as a high-priority routine activity</li> <li>• establish and implement a treatment programme</li> <li>• ensure continuity of any treatment programme while the person is in</li> </ul>

	<p>contact with the CRHT team, and on discharge or transfer to other services when this is needed</p> <ul style="list-style-type: none"> <li>• have a crisis management plan in place before the person is discharged from the team's care. [new 2017] (1.14.7)</li> </ul> <p>Consider inpatient treatment for people with more severe depression who cannot be adequately supported by a CRHT team. [new 2017] (1.14.8)</p> <p>Make the full range of recommended psychological therapies (group CBT, CBT or BA) available for people with depression in inpatient settings. [new 2017] (1.14.9)</p> <p>When providing psychological therapies for people with depression in inpatient settings:</p> <ul style="list-style-type: none"> <li>• increase the intensity and duration of the interventions</li> <li>• ensure that they continue to be provided effectively and promptly on discharge. [new 2017] (1.14.10)</li> </ul> <p>Consider using CRHT teams with people with depression who might benefit from early discharge from hospital after a period of inpatient care. [2017] (1.14.11)</p>
<p>Teams working with people with complex and severe depression should develop comprehensive multidisciplinary care plans in collaboration with the person with depression (and their family or carer, if agreed with the person). The care plan should:</p> <ul style="list-style-type: none"> <li>• identify clearly the roles and responsibilities of all health and social care professionals involved</li> <li>• develop a crisis plan that identifies potential triggers that could lead to a crisis and strategies to manage such triggers</li> <li>• be shared with the GP and the person with depression and other relevant people involved in the person's care. (1.10.1.5)</li> </ul>	<p>Replaced by:</p> <p>Ensure multidisciplinary care plans for people with more severe depression with multiple complicating problems, or significant coexisting conditions:</p> <ul style="list-style-type: none"> <li>• are developed together with the person, their GP and other relevant people involved in their care (with the person's agreement)</li> <li>• set out the roles and responsibilities of all health and social care professionals involved in delivering the care</li> <li>• include information about 24-hour support services, and how to contact them</li> <li>• include a crisis plan that identifies potential crisis triggers, and strategies to manage those triggers</li> </ul>

	<ul style="list-style-type: none"> <li>• are updated if there are any significant changes in the person's needs or condition</li> <li>• are reviewed at agreed regular intervals</li> <li>• include medication management (a plan for starting, reviewing and discontinuing medication). [new 2017] (1.14.5)</li> </ul>
<p>For people who have depression with psychotic symptoms, consider augmenting the current treatment plan with antipsychotic medication (although the optimum dose and duration of treatment are unknown) (1.10.3.1)</p>	<p>Replaced by: Refer people with depression with psychotic symptoms to specialist mental health services for a programme of coordinated multi-disciplinary care, which includes access to psychological interventions.[new 2017] (1.12.1)</p> <p>When treating people with depression with psychotic symptoms, consider adding antipsychotic medication to their current treatment plan. [new 2017] (1.12.2)</p>
<p>Do not use ECT routinely for people with moderate depression but consider it if their depression has not responded to multiple drug treatments and psychological treatment. (1.10.4.2)</p>	<p>Replaced by: Consider electroconvulsive therapy (ECT) for acute treatment of more severe depression if:</p> <ul style="list-style-type: none"> <li>• the more severe depression is life-threatening and a rapid response is needed, or</li> <li>• multiple pharmacological and psychological treatments have failed. [2017] (1.13.1)</li> </ul>

1  
2

3 **Amended recommendation wording (change to meaning)**

Recommendation in 2009 guideline	Recommendation in current guideline	Reason for change
Make all efforts necessary to ensure that a person with depression can give meaningful and informed consent before treatment starts. This is especially important when a person has severe depression or is subject to the Mental Health Act. [2004] (1.1.1.4)	Make every effort to ensure that a person with depression can give meaningful and informed consent before treatment starts. This is especially important when a person has severe depression or their treatment falls under the Mental Health Act or the Mental Capacity Act. [2004, amended 2017] (1.1.4)	Amended to cite additional relevant legislation – the Mental Capacity Act.
For people with recurrent severe depression or depression with psychotic symptoms and for those who have been treated under the Mental Health Act, consider developing advance decisions and advance statements collaboratively with the person. Record the decisions and statements and include copies in the person's care plan in primary and secondary care. Give copies to the person and to their family or carer, if the person agrees. (1.1.2.1)	Consider developing advance decisions and advance statements collaboratively with people who have recurrent severe depression or depression with psychotic symptoms, and for those who have been treated under the Mental Health Act, in line with the Mental Capacity Act. Record the decisions and statements and include copies in the person's care plan in primary and secondary care, and give copies to the person and to their family or carer if the person agrees. [2009, amended 2017] (1.1.6)	Amended to cite additional relevant legislation – the Mental Capacity Act.
For people with significant language or communication difficulties, for example people with sensory impairments or a learning disability, consider using the Distress Thermometer <sup>8</sup> and/or asking a family member or carer about the person's symptoms to identify possible depression. If a significant level of distress is identified, investigate further. (1.3.1.5)	If a person has significant language or communication difficulties, (for example people with sensory or cognitive impairments), consider asking a family member or carer about the person's symptoms to identify possible depression. [2004, amended 2017] (See also NICE's guideline on mental health problems in people with learning disabilities.) (1.2.5)	Removed reference to use of the Distress Thermometer as this detail would be superseded by recommendations made in NICE's guideline on mental health problems in people with learning disabilities

<sup>8</sup> The Distress Thermometer is a single-item question screen that will identify distress coming from any source. The person places a mark on the scale answering: 'How distressed have you been during the past week on a scale of 0 to 10?' Scores of 4 or more indicate a significant level of distress that should be investigated further. (Roth AJ, Kornblith AB, Batel-Copel L, et al. (1998) Rapid screening for psychologic distress in men with prostate carcinoma: a pilot study. Cancer 82: 1904–8.) 1904–8.)

<p>In addition to assessing symptoms and associated functional impairment, consider how the following factors may have affected the development, course and severity of a person's depression:</p> <ul style="list-style-type: none"> <li>• any history of depression and comorbid mental health or physical disorders</li> <li>• any past history of mood elevation (to determine if the depression may be part of bipolar disorder)</li> <li>• any past experience of, and response to, treatments</li> <li>• the quality of interpersonal relationships</li> <li>• living conditions and social isolation.</li> </ul>	<p>Think about how the factors below may have affected the development, course and severity of a person's depression in addition to assessing symptoms and associated functional impairment:</p> <ul style="list-style-type: none"> <li>• any history of depression and coexisting mental health or physical disorders</li> <li>• any history of mood elevation (to determine if the depression may be part of bipolar disorder)</li> <li>• any past experience of, and response to, previous treatments</li> <li>• the quality of interpersonal relationships</li> <li>• living conditions, employment situation and social isolation. [2009, amended 2017] (1.2.7)</li> </ul>	<p>Added employment situation into the list of factors to consider as this would now be checked as standard</p>
<p>When assessing a person with suspected depression, be aware of any learning disabilities or acquired cognitive impairments, and if necessary consider consulting with a relevant specialist when developing treatment plans and strategies. (1.1.4.4)</p>	<p>When assessing a person with suspected depression:</p> <ul style="list-style-type: none"> <li>• be aware of any acquired cognitive impairments</li> <li>• if needed, consult with a relevant specialist when developing treatment plans and strategies. [2009, amended 2017] (1.2.8)</li> </ul>	<p>Removed reference to learning disabilities as there is now a separate NICE guideline on mental health problems in people with learning disabilities</p>

<p>When providing interventions for people with a learning disability or acquired cognitive impairment who have a diagnosis of depression:</p> <ul style="list-style-type: none"> <li>• where possible, provide the same interventions as for other people with depression</li> <li>• if necessary, adjust the method of delivery or duration of the intervention to take account of the disability or impairment. (1.1.4.5)</li> </ul>	<p>When providing interventions for people with an acquired cognitive impairment who have a diagnosis of depression:</p> <ul style="list-style-type: none"> <li>• if possible, provide the same interventions as for other people with depression</li> <li>• if needed, adjust the method of delivery or length of the intervention to take account of the disability or impairment. [2009, amended 2017] (1.2.9)</li> </ul>	<p>Removed reference to learning disabilities as there is now a separate NICE guideline on mental health problems in people with learning disabilities</p>
---	--	--

1  
2

3 **Changes to recommendation wording for clarification only (no change to**  
4 **meaning)**

Recommendation numbers in current guideline	Comment
All recommendations except those labelled [new 2017]	Recommendations have been edited into the direct style (in line with current NICE style for recommendations in guidelines) where possible. Yellow highlighting has not been applied to these changes.

5  
6

7 **ISBN:**