

NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE

Interventional procedures consultation document

Intravascular lithotripsy for calcified arteries in peripheral arterial disease

Peripheral arterial disease (PAD) is caused when a build-up of fatty substances (plaque) in the arteries restricts blood supply to parts of the body, usually the legs. The plaque can harden (calcify) and become like bone. If PAD is severe, the affected part of the body can die, needing amputation or causing the person to die. In this procedure, a long, thin, flexible tube called a catheter with a balloon at the end is inserted into the affected artery (intravascular). Ultrasound pulses are sent from the balloon to soften the calcified plaque (lithotripsy) and allow the artery to widen. The catheter and balloon are removed and sometimes a small metal tube, called a stent, is inserted to help keep the artery open. The aim of the procedure is to improve the blood flow and prevent amputations or death.

NICE is looking at intravascular lithotripsy for calcified arteries in peripheral arterial disease.

NICE's interventional procedures advisory committee met to consider the evidence and the opinions of professional experts with knowledge of the procedure.

This document contains the [draft guidance for consultation](#). Your views are welcome, particularly:

- comments on the draft recommendations
- information about factual inaccuracies
- additional relevant evidence, with references if possible.

NICE is committed to promoting equality of opportunity, eliminating unlawful discrimination and fostering good relations between people with particular protected characteristics and others.

This is not NICE's final guidance on this procedure. The draft guidance may change after this consultation.

After consultation ends, the committee will:

- meet again to consider the consultation comments, review the evidence and make appropriate changes to the draft guidance

- prepare a second draft, which will go through a [resolution process](#) before the final guidance is agreed.

Please note that we reserve the right to summarise and edit comments received during consultation or not to publish them at all if, in the reasonable opinion of NICE, there are a lot of comments or if publishing the comments would be unlawful or otherwise inappropriate.

Closing date for comments: 23 August 2023

Target date for publication of guidance: January 2024

1 Draft recommendations

- 1.1 Intravascular lithotripsy for calcified arteries in peripheral arterial disease should only be used with special arrangements for clinical governance, consent, and audit or research. Find out [what special arrangements mean on the NICE interventional procedures guidance page](#).
- 1.2 Clinicians wanting to do intravascular lithotripsy for calcified arteries in peripheral arterial disease should:
- Inform the clinical governance leads in their healthcare organisation.
 - Ensure that people (and their families and carers as appropriate) understand the procedure's safety and efficacy, and any uncertainties about these.
 - Take account of NICE's advice on [shared decision making](#), including [NICE's information for the public](#).
 - Audit and review clinical outcomes of everyone having the procedure. The main efficacy and safety outcomes identified in this guidance can be entered into [NICE's interventional procedure outcomes audit tool](#) (for use at local discretion).
 - Discuss the outcomes of the procedure during their annual appraisal to reflect, learn and improve.
- 1.3 Healthcare organisations should:
- Ensure systems are in place that support clinicians to collect and report data on outcomes and safety for everyone having this procedure.
 - Regularly review data on outcomes and safety for this procedure.
- 1.4 Patient selection should be done by a vascular multidisciplinary team including interventional radiologists and vascular surgeons.

1.5 NICE encourages further research into intravascular lithotripsy for calcified arteries in peripheral arterial disease. Research should include:

- details of patient selection, including the location and degree of stenosis
- longer-term outcomes, including mortality and amputation
- patient-reported outcomes including quality of life
- the need for revascularisation and amputation.

Why the committee made these recommendations

There is a moderate amount of evidence, including data from a randomised controlled trial, which suggests the procedure is safe. The evidence shows that blood vessel diameter is increased after the procedure, but because intravascular lithotripsy is sometimes done alongside other procedures it is difficult to know whether this is directly because of the intravascular lithotripsy procedure. The evidence suggests that the procedure is associated with a reduced need for a stent to keep the vessel open. But there is not enough long-term evidence, or evidence about how many amputations will be prevented by having this procedure. More evidence is also needed on patients' quality of life.

There may be groups of people who would particularly benefit from this procedure. These could include people with smaller vessels or with calcified arteries in a location unsuitable for a stent, but more evidence is needed.

2 The condition, current treatments and procedure

The condition

2.1 Peripheral arterial disease (PAD) is caused when a build-up of fatty substances (plaque) in the arteries restricts blood supply to the limbs, usually the legs. The plaque can calcify and become like

bone. This is known as intravascular calcification, and it is particularly common in people with diabetes mellitus or chronic kidney disease. The most common initial symptom of PAD is leg pain while walking, known as intermittent claudication. If blood flow is severely restricted, chronic limb-threatening ischaemia (CLTI, also known as critical limb ischaemia) can develop. Symptoms include severe pain at rest, ulceration or gangrene. CLTI is associated with a high risk of amputation and death, and the presence of arterial calcification increases these risks.

Current treatments

- 2.2 Management of PAD is described in [NICE's clinical guideline on peripheral arterial disease](#). For CLTI, revascularisation using percutaneous transluminal angioplasty (with or without stent placement) or a bypass graft is recommended. Atherectomy devices that vaporise, cut or grind away plaque within the artery are sometimes used alongside balloon angioplasty. If revascularisation is not an option, major amputation may be offered.

The procedure

- 2.3 Arterial access is established as for a standard angioplasty, usually through the femoral artery in the groin. An angioplasty balloon with ultrasound element is introduced and inflated next to the heavily calcified arterial plaques. The pressure exerted by the balloon is too low to expand the vessel but high enough to ensure good contact between the surface of the balloon and the arterial walls. Ultrasound pulses are then transmitted from the balloon, fracturing superficial and deep calcium within the arterial wall. A stent is sometimes inserted after the lithotripsy to keep the artery patent. The procedure is used as a preparatory treatment for balloon angioplasty or as an alternative to standard angioplasty.
- 2.4 The aim of intravascular lithotripsy is to improve the blood flow in the affected limb and prevent the need for amputation.

3 Committee considerations

The evidence

- 3.1 NICE did a rapid review of the published literature on the efficacy and safety of this procedure. This comprised a comprehensive literature search and detailed review of the evidence from 9 sources, which was discussed by the committee. The evidence included 1 systematic review and meta-analysis, 1 randomised controlled trial (reported in 2 papers), 1 prospective single-arm trial, 2 retrospective cohort studies, 1 prospective single centre registry and 1 case report. It is presented in the [summary of key evidence section in the interventional procedures overview](#). Other relevant literature is in table 5 of the overview.
- 3.2 The professional experts and the committee considered the key efficacy outcomes to be: quality of life, vessel patency, amputation rate, amputation-free survival, procedural success, revascularisation rate and need for a stent.
- 3.3 The professional experts and the committee considered the key safety outcomes to be: embolisation, perforation, dissection, worsening of pain and long-term procedure failure.
- 3.4 Patient commentary was sought but none was received.

Committee comments

- 3.5 Most of the evidence was on femoropopliteal lesions, but there was some evidence on iliac lesions. There are several ongoing studies, including 1 on arteries below the knee.
- 3.6 Clinical experts explained that the procedure is only used for chronic limb ischaemia and would not be used in an acute setting.
- 3.7 The clinical experts advised that there is no clear consensus for assessing the degree of calcification in a peripheral artery.

- 3.8 There is a company-funded registry of people having the procedure in the UK. But this is no longer recruiting.
- 3.9 There is an unmet need for a safe and effective endovascular option for treating heavily calcified arterial lesions when surgery is unsuitable.

James Tysome,
Vice chair, interventional procedures advisory committee
July 2023

ISBN: