

Subarachnoid haemorrhage

[T] Evidence review for investigating relatives of people with aneurysmal SAH

NICE guideline <number>

Evidence review underpinning

February 2021

Draft for consultation

*Developed by the National Guideline Centre,
hosted by the Royal College of Physicians*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and, where appropriate, their carer or guardian.

Local commissioners and providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to Notice of rights.

ISBN

[add for final publication version only, delete this text for consultation version]

Contents

1	Detecting intracranial arterial aneurysms in first-degree relatives	5
1.1	Review question: What is the clinical and cost effectiveness of investigations to detect intracranial arterial aneurysms in relatives of adults who have had a subarachnoid haemorrhage?	5
1.2	Introduction	5
1.3	PICO table	5
1.4	Clinical evidence	6
1.4.1	Included studies	6
1.4.2	Excluded studies	6
1.4.3	Summary of clinical studies included in the evidence review	7
1.4.4	Quality assessment of clinical studies included in the evidence review	7
1.5	Economic evidence	8
1.5.1	Included studies	8
1.5.2	Excluded studies	8
1.5.3	Summary of studies included in the economic evidence review	9
1.5.4	Unit costs	11
1.6	Evidence statements	11
1.6.1	Clinical evidence statements	11
1.6.2	Health economic evidence statements	11
1.7	The committee's discussion of the evidence	11
1.7.1	Interpreting the evidence	11
1.7.2	Cost effectiveness and resource use	12
1.7.3	Other factors the committee took into account	13
	Appendices	20
	Appendix A: Review protocols	20
	Appendix B: Literature search strategies	25
	B.1 Clinical search literature search strategy	26
	B.2 Health Economics literature search strategy	29
	Appendix C: Clinical evidence selection	33
	Appendix D: Clinical evidence tables	34
	Appendix E: Forest plots	35
	Appendix F: GRADE tables	36
	Appendix G: Health economic evidence selection	37
	Appendix H: Health economic evidence tables	39
	Appendix I: Excluded studies	47
	I.1 Excluded clinical studies	47
	I.2 Excluded health economic studies	50

1 **1 Detecting intracranial arterial aneurysms in** 2 **first-degree relatives**

3 Evidence review underpinning recommendation 1.4.14 and research recommendations in the
4 NICE guideline.

1.1 **5 Review question: What is the clinical and cost** 6 **effectiveness of investigations to detect intracranial arterial** 7 **aneurysms in relatives of adults who have had a** 8 **subarachnoid haemorrhage?**

1.2 **9 Introduction**

10 The reported prevalence of intracranial aneurysms varies between studies but is estimated to
11 range from 2% to 5% in the general population. The chance of detecting an aneurysm on
12 screening is reported to be around 10% in people with 2 first-degree relatives who have had
13 subarachnoid haemorrhage. People with a family history are also at higher risk of
14 subarachnoid haemorrhage than the general population.

15 Current practice varies, but some relatives of adults who have had a subarachnoid
16 haemorrhage are offered investigation and those found to have an aneurysm at risk of
17 rupture may be considered for intervention.

18 This review assessed evidence of clinical and cost-effectiveness of investigations to detect
19 intracranial arterial aneurysms in relatives of adults who have had a subarachnoid
20 haemorrhage.

1.3 **21 PICO table**

22 For full details see the review protocol in Appendix A:.

23 **Table 1: PICO characteristics of review question**

Population	First degree relatives of adults (16 and older) with a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm.	
Intervention(s)	Assessment of first degree relatives of people with aSAH <ul style="list-style-type: none">• Assessment with CTA or MRA	
Comparison(s)	To no routine assessment	
Outcomes	Critical outcomes: <ul style="list-style-type: none">• Mortality• Health and social-related quality of life (any validated measure)• Degree of disability or dependence in daily activities, (any validated measure e.g. Modified Rankin Scale and patient-reported outcome measures)• Subarachnoid haemorrhage	Important outcomes: <ul style="list-style-type: none">• Presence of cerebral aneurysm• Elective treatment
Study design	<ul style="list-style-type: none">• Randomised controlled trials (RCTs), systematic reviews of RCTs.• If insufficient RCT evidence is available, non-randomised studies will be considered if they adjust for key confounders (age), starting with prospective cohort studies.	

1.4 1 Clinical evidence

1.4.1 2 Included studies

- 3 No relevant clinical studies comparing assessment of first-degree relatives of people with
- 4 SAH to no routine assessment were identified.

- 5 See also the study selection flow chart in Appendix C:.

1.4.2 6 Excluded studies

- 7 See the excluded studies list in Appendix I:.

1.4.3 1 Summary of clinical studies included in the evidence review

2 No studies were included.

1.4.4 3 Quality assessment of clinical studies included in the evidence review

4 No studies were included.

5

1.5 1 Economic evidence

1.5.1 2 Included studies

- 3 Two health economic studies with the relevant comparison were included in this review^{24,3}.
- 4 These are summarised in the health economic evidence profile below (Table 2) and the
- 5 health economic evidence tables in Appendix H:.

1.5.2 6 Excluded studies

- 7 No relevant health economic studies were excluded due to assessment of limited
- 8 applicability and methodological limitations.
- 9 See also the health economic study selection flow chart in Appendix G:.

10

1.5.3 1 Summary of studies included in the economic evidence review

2 Table 2: Health economic evidence profile: Screening strategies compared to no screening

Study	Applicability	Limitations	Other comments	Incremental cost	Incremental effects	Cost effectiveness	Uncertainty
Hopmans 2016 ²⁴ [Netherlands]	Partially applicable ^(a)	Potentially serious limitations ^(b)	<ul style="list-style-type: none"> • Probabilistic Markov model based on estimated incidence of aneurysm and rupture risk throughout lifetime. • Cost-utility analysis (QALYs) • Population: People with one affected first degree relative with aneurysmal subarachnoid haemorrhage. • Comparators: <ul style="list-style-type: none"> ○ No screening ○ Multiple different screening strategies - every 5, 10 or 15 years between age bands 20-60, 20-70, 30-60, 30-70, 40-60,40-70 ○ One time screening strategies at age 30,35,40, 45, 50, 55 Time horizon: Lifetime	For full incremental analysis of all 25 strategies please see the full evidence table in Appendix H.		All screening strategies cost effective when compared to no screening. The most cost effective strategy is screening every 5 years aged 30 to 70 with an ICER of £16,409 per QALY gained.	Probability strategies cost effective (£20/£30K threshold): NR No further sensitivity analyses undertaken.
Bor 2010 ³ [Netherlands]	Partially applicable ^(c)	Potentially serious limitations ^(d)	<ul style="list-style-type: none"> • Probabilistic Markov model based on estimated incidence of aneurysm and rupture risk throughout lifetime. 	For full incremental analysis of the most efficient 10 strategies please see the full evidence table in Appendix H. ^(e)		All screening strategies cost effective when compared to no screening.	ANCOVA analysis suggested that utility of individuals who were not offered screening and did

Study	Applicability	Limitations	Other comments	Incremental cost	Incremental effects	Cost effectiveness	Uncertainty
			<ul style="list-style-type: none"> • Cost-utility analysis (QALYs) • Population: People with two or more first degree relatives with aneurysmal subarachnoid haemorrhage. • Comparators: <ul style="list-style-type: none"> ○ No screening ○ Multiple different screening strategies with different age and screening intervals • Time horizon: Lifetime 			The most cost effective strategy is screening every 2 years aged 20 to 80 with an ICER of £15,714 per QALY gained.	not experience SAH and utility of individuals having a negative screening most influenced health effectiveness.

- 1 Abbreviations: ANCOVA = Analysis of covariance; ICER= incremental cost-effectiveness ratio; NR= Not reported; QALY= quality-adjusted life years; RCT= randomised
2 controlled trial
3 (a) Dutch resource use data (mix 1997-2003, 1996-97) and unit costs (2012) may not reflect current NHS context. Costs and health effects were discounted at a non-
4 reference case rate (4%, 1.5% respectively). Non-NICE reference case utility measure used to estimate some utility values to calculate QALYs.
5 (b) Risk of aneurysm development and rupture were estimated from non-comparative cohort studies and assumptions made to estimate risk over time. Uncertainty not
6 explored through sensitivity analysis.
7 (c) Dutch resource use data (mix 1997-2003, 1996-97) and unit costs (2007) may not reflect current NHS context. Costs and health effects were discounted at a non-
8 reference case rate (1.5%). Unclear how estimates of utility values to calculate QALYs were determined.
9 (d) Treatment effects estimated primarily from non-comparative cohort studies and assumptions about aneurysm development and rupture risk over time.
10 (e) Due to the large volume of strategies in this study, the full incremental analysis in Appendix H does not include the interventions that were subject to dominance or
11 extended dominance.

12

1.5.4 1 Unit costs

2 Relevant unit costs are provided below to aid consideration of cost effectiveness.

3 Table 3: UK costs of screening

Description	Unit cost
Magnetic Resonance Imaging Scan of One Area, without Contrast, 19 years and over [NHS Reference cost code: RD01A]	£121
Neurosurgery consultant led outpatient follow up appointment [NHS Reference cost code: WF01B,150]	£228

4 Source: NHS Reference Cost 2018/19⁵⁰

1.6 5 Evidence statements

1.6.1 6 Clinical evidence statements

7 No relevant published evidence was identified.

1.6.2 8 Health economic evidence statements

- 9 • One cost–utility analysis found that different screening strategies (that varied by age
10 range and frequency of screening) were cost effective compared to no screening in
11 people with one affected first degree relative with aneurysmal subarachnoid haemorrhage.
12 When all the screening strategies compared in the model are considered, screening every
13 5 years between the age of 30 and 70 was the most cost-effective intervention (ICER:
14 £16,409 per QALY gained). This analysis was assessed as partially applicable with
15 potentially serious limitations.
- 16 • One cost–utility analysis found that different screening strategies (that varied by age
17 range and frequency of screening) were cost effective compared to no screening in
18 people with two or more first degree relatives with aneurysmal subarachnoid
19 haemorrhage. When all the screening strategies compared in the model are considered,
20 screening every 2 years between the ages of 20 and 80 was the most cost-effective
21 intervention (ICER: £15,714 per QALY gained). This analysis was assessed as partially
22 applicable with very serious limitations.

1.7 23 The committee’s discussion of the evidence

1.7.1 24 Interpreting the evidence

1.7.1.1 25 The outcomes that matter most

26 The critical outcomes were mortality; health and social-related quality of life; degree of
27 disability and subarachnoid haemorrhage. These outcomes were selected to understand the
28 impact of screening and potential treatment on a relative’s health and quality of life. The
29 important outcomes were presence of cerebral aneurysm and elective treatment.

1.7.1.2 30 The quality of the evidence

31 No relevant published clinical evidence was identified.

1.7.1.3 1 Benefits and harms

2 The committee acknowledged that the main objective of investigating relatives of adults who
3 have had a subarachnoid haemorrhage is detection of intracranial arterial aneurysms, some
4 of which may be considered to require treatment. Despite the potential risks, treatment to
5 secure an aneurysm could prevent death or other adverse consequences of subarachnoid
6 haemorrhage.

7 The committee discussed that any benefits of investigation of relatives must be balanced
8 against the risks and costs of investigation, including possible psychological burden of
9 people knowing that they have an aneurysm. Investigation may also reveal an aneurysm that
10 is not suitable for treatment, or the absence of aneurysms but the potential for development
11 of new aneurysms. In some patients such findings may lead to increased anxiety and a fear
12 of symptoms associated with subarachnoid haemorrhage, such as headache, potentially
13 increasing unnecessary primary care or emergency department attendances.

14 The committee noted that routine screening of relatives for intracranial arterial aneurysms is
15 currently not supported within NHS practice and any change in practice would have a
16 significant resource impact including additional investigations and interventions requiring
17 hospital consultations. The committee also agreed that the optimal timing and frequency of
18 investigations to detect and monitor intracranial aneurysms in relatives is unknown.
19 Nevertheless, the committee acknowledged that there is significant variation in national
20 practice and a small number of relatives of people with aSAH currently undergo MR
21 angiography to look for aneurysms. The committee noted that investigation would only be
22 recommended for people thought to have a significant risk of having a brain aneurysm that
23 could rupture at some point in the future, and the committee discussed this would apply to
24 people with two or more first-degree relatives who have had subarachnoid haemorrhage as
25 these individuals are at substantially increased risk of SAH. The committee acknowledged
26 this reflected the guidance on brain aneurysm provided by the NHS and in their experience
27 reflected practice at many UK neurosurgical centres. The committee also agreed that the risk
28 of SAH is recognised to be greater in individuals with 2 first degree relatives with aSAH than
29 those with 1 first degree relative with aSAH.

30 The committee noted the lack of evidence to support investigation, and decided to make a
31 consensus recommendation to explain to patients (and if appropriate their family members)
32 the uncertainties surrounding the risks and benefits of investigation for intracranial
33 aneurysms in relatives of people with aSAH. The committee also acknowledged that in their
34 experience some people do not want to be investigated for a potential aneurysm. The
35 committee noted information about the rationale for and timing of investigations for brain
36 aneurysms is available via the NHS website, and agreed this resource should be highlighted
37 within the recommendations. The committee added that the relatives' own risk of developing
38 an intracranial aneurysm should be taken into account when considering investigation, taking
39 into account risk factors such as smoking, high blood pressure, and autosomal dominant
40 polycystic kidney disease. The committee concluded that further research was needed.

1.7.241 Cost effectiveness and resource use

42 Two published economic evaluations were included in this review. Both studies assessed
43 multiple different screening strategies in first degree relatives of people with subarachnoid
44 haemorrhage from a Dutch perspective. One paper assessed screening strategies in people
45 with 1 first degree relative, and showed that all screening strategies were cost effective
46 compared to no screening; the most cost effective strategy was to screen every 5 years
47 between the ages of 30 and 70. This paper was assessed as partially applicable with
48 potentially serious limitations. One paper assessed screening strategies in people with 2 or
49 more first degree relatives with aSAH, which also suggested that all screening strategies
50 were cost effective compared to no screening; the most cost effective strategy was to screen

1 every 2 years between the ages of 20 and 80. This study was assessed as partially
2 applicable with very serious limitations.

3 The committee were concerned that this evidence is not reflective of NHS practice or the
4 NICE reference case. It was noted that any aneurysms identified on screening in England
5 are typically only treated if they measure at least 7mm in diameter, whereas in the models
6 the threshold for treatment was 5mm. Furthermore, the committee were concerned that there
7 are likely to be some additional costs of monitoring aneurysms that have been identified but
8 are too small to require treatment.

9 The committee considered the evidence used in the model to simulate the incidence of de
10 novo aneurysms, aneurysm growth, rupture risk, and outcomes of aneurysm treatment and
11 concluded the data used for the model were reasonable estimates. However, the committee
12 could not make recommendations with potentially large resource impact without the support
13 of comparative clinical evidence.

14 The committee noted that screening could sometimes be more detrimental (compared to no
15 screening), and lead to additional costs for the NHS due to increased patient concerns and
16 worries incurred as a result of screening. The committee acknowledged routine
17 investigations for aneurysms in first-degree relatives could lead to an increase in visits to
18 GPs and emergency departments when the NHS does not currently support routine
19 investigation.

20 Overall, this recommendation is not expected to have a substantial resource impact as it is
21 reflective of current practice in England.

1.7.32 Other factors the committee took into account

23 The committee discussed that requests for monitoring people with relatives who have had a
24 subarachnoid haemorrhage is increasing and this is an important concern for patients and
25 their families. Given the lack of evidence, the committee considered this to be an important
26 area for further research and so by consensus made a research recommendation for the
27 clinical and cost effectiveness of investigations to detect intracranial arterial aneurysms in
28 first-degree relatives of people who have had aSAH.

1 References

- 2 1. Bhat AR, Afzalwani M, Kirmani AR. Subarachnoid hemorrhage in Kashmir: causes,
3 risk factors, and outcome. *Asian Journal of Neurosurgery*. 2011; 6(2):57-71
- 4 2. Bhat AR, Wani MA, Kirmani AR, Ramzan AU, Alam S, Raina T et al. High incidence
5 of intracranial aneurysmal subarachnoid hemorrhage (SAH) in Kashmir, India.
6 *Biomedical Research*. 2012; 23(1):79-92
- 7 3. Bor AS, Koffijberg H, Wermer MJ, Rinkel GJ. Optimal screening strategy for familial
8 intracranial aneurysms: a cost-effectiveness analysis. *Neurology*. 2010; 74(21):1671-
9 1679
- 10 4. Bor AS, Rinkel GJ, Adami J, Koffijberg H, Ekbom A, Buskens E et al. Risk of
11 subarachnoid haemorrhage according to number of affected relatives: a population
12 based case-control study. *Brain*. 2008; 131(10):2662-2665
- 13 5. Bor AS, Rinkel GJ, van Norden J, Wermer MJ. Long-term, serial screening for
14 intracranial aneurysms in individuals with a family history of aneurysmal subarachnoid
15 haemorrhage: a cohort study. *Lancet Neurology*. 2014; 13(4):385-392
- 16 6. Bossuyt PM, Raaymakers TW, Bonsel GJ, Rinkel GJ. Screening families for
17 intracranial aneurysms: anxiety, perceived risk, and informed choice. *Preventive
18 Medicine*. 2005; 41(3-4):795-799
- 19 7. Broderick JP, Brown RD, Jr., Sauerbeck L, Hornung R, Huston J, 3rd, Woo D et al.
20 Greater rupture risk for familial as compared to sporadic unruptured intracranial
21 aneurysms. *Stroke*. 2009; 40(6):1952-1957
- 22 8. Broderick JP, Sauerbeck LR, Foroud T, Huston 3rd J, Pankratz N, Meissner I et al.
23 The Familial Intracranial Aneurysm (FIA) study protocol. *BMC Medical Genetics*.
24 2005; 6:17
- 25 9. Bromberg JE, Rinkel GJ, Algra A, Limburg M, van Gijn J. Outcome in familial
26 subarachnoid hemorrhage. *Stroke*. 1995; 26(6):961-963
- 27 10. Bromberg JEC, Rinkel GJE, Algra A, Van Duyn CM, Greebe P, Ramos LMP et al.
28 Familial subarachnoid hemorrhage: distinctive features and patterns of inheritance.
29 *Annals of Neurology*. 1995; 38(6):929-934
- 30 11. Brown BM, Soldevilla F. MR angiography and surgery for unruptured familial
31 intracranial aneurysms in persons with a family history of cerebral aneurysms.
32 *American Journal of Roentgenology*. 1999; 173(1):133-138
- 33 12. Brown RD, Jr., Broderick JP. Unruptured intracranial aneurysms: epidemiology,
34 natural history, management options, and familial screening. *Lancet Neurology*. 2014;
35 13(4):393-404
- 36 13. Chalouhi N, Chitale R, Jabbour P, Tjoumakaris S, Dumont AS, Rosenwasser R et al.
37 The case for family screening for intracranial aneurysms. *Neurosurgical Focus*. 2011;
38 31(6):E8
- 39 14. Chan DY, Abrigo JM, Cheung TC, Siu DY, Poon WS, Ahuja AT et al. Screening for
40 intracranial aneurysms? Prevalence of unruptured intracranial aneurysms in Hong
41 Kong Chinese. *Journal of Neurosurgery*. 2016; 124(5):1245-1249
- 42 15. Coppola AR. Familial intracranial aneurysm. *International Surgery*. 1973; 58(7):508
- 43 16. Crawley F, Clifton A, Brown MM. Should we screen for familial intracranial aneurysm?
44 *Stroke*. 1999; 30(2):312-316

- 1 17. Dippel DW, ter Berg JW, Habbema JD. Screening for unruptured familial intracranial
2 aneurysms. A decision analysis. *Acta Neurologica Scandinavica*. 1992; 86(4):381-
3 389
- 4 18. Edelson L, Caplan L, Rosenbaum AE. Familial aneurysms and infundibular
5 widening. *Neurology*. 1972; 22(10):1056-1060
- 6 19. Flahault A, Trystram D, Fouchard M, Knebelmann B, Nataf F, Joly D. Screening for
7 unruptured intracranial aneurysms in autosomal dominant polycystic kidney disease:
8 a survey of 420 nephrologists. *PloS One*. 2016; 11(4):e0153176
- 9 20. Flahault A, Trystram D, Nataf F, Fouchard M, Knebelmann B, Grunfeld JP et al.
10 Screening for intracranial aneurysms in autosomal dominant polycystic kidney
11 disease is cost-effective. *Kidney International*. 2018; 93(3):716-726
- 12 21. Fox JL, Ko JP. Familial intracranial aneurysms. Six cases among 13 siblings. *Journal*
13 *of Neurosurgery*. 1980; 52(4):501-503
- 14 22. Greebe P, Bromberg JEC, Rinkel GJE, Algra A, Van Gijn J. Family history of
15 subarachnoid haemorrhage: supplemental value of scrutinising all relatives. *Journal*
16 *of Neurology Neurosurgery and Psychiatry*. 1997; 62(3):273-275
- 17 23. Hashimoto I. Familial intracranial aneurysms and cerebral vascular anomalies.
18 *Journal of Neurosurgery*. 1977; 46(4):419-427
- 19 24. Hopmans EM, Ruigrok YM, Bor ASE, Rinkel GJE, Koffijberg H. A cost-effectiveness
20 analysis of screening for intracranial aneurysms in persons with one first-degree
21 relative with subarachnoid haemorrhage. *European Stroke Journal*. 2016; 1(4):320-
22 329
- 23 25. Huston IJ, Torres VE, Wiebers DO, Schievink WI. Follow-up of intracranial
24 aneurysms in autosomal dominant polycystic kidney disease by magnetic resonance
25 angiography. *Journal of the American Society of Nephrology*. 1996; 7(10):2135-2141
- 26 26. Jacoby G, Sams IR. Should people with a first-degree relative who died from
27 subarachnoid hemorrhage be screened for aneurysms? *Journal of Family Practice*.
28 2006; 55(1):59-60
- 29 27. Jain KK. Familial intracranial aneurysms; review of literature and presentation of six
30 new cases. *Acta Neurochirurgica*. 1974; 30(1-2):129-137
- 31 28. Kasuya H, Onda H, Takeshita M, Hori T, Takakura K. Clinical features of intracranial
32 aneurysms in siblings. *Neurosurgery*. 2000; 46(6):1301-1305; discussion 1305-1306
- 33 29. Kelly AG. Unruptured intracranial aneurysms: screening and management.
34 *CONTINUUM: Lifelong Learning in Neurology*. 2014; 20(2 Cerebrovascular
35 Disease):387-398
- 36 30. Kim DH, Van Ginhoven G, Milewicz DM. Familial aggregation of both aortic and
37 cerebral aneurysms: evidence for a common genetic basis in a subset of families.
38 *Neurosurgery*. 2005; 56(4):655-661; discussion 655-661
- 39 31. Kim DH, Van Ginhoven G, Milewicz DM. Incidence of familial intracranial aneurysms
40 in 200 patients: comparison among Caucasian, African-American, and Hispanic
41 populations. *Neurosurgery*. 2003; 53(2):302-308
- 42 32. Kojima M, Nagasawa S, Lee YE, Takeichi Y, Tsuda E, Mabuchi N. Asymptomatic
43 familial cerebral aneurysms. *Neurosurgery*. 1998; 43(4):776-781

- 1 33. Koshy L, Easwer HV, Premkumar S, Alapatt JP, Pillai AM, Nair S et al. Risk factors
2 for aneurysmal subarachnoid hemorrhage in an Indian population. *Cerebrovascular*
3 *Diseases*. 2010; 29(3):268-274
- 4 34. Kubota M, Yamaura A, Ono J. Prevalence of risk factors for aneurysmal
5 subarachnoid haemorrhage: results of a Japanese multicentre case control study for
6 stroke. *British Journal of Neurosurgery*. 2001; 15(6):474-478
- 7 35. Leblanc R. Familial cerebral aneurysms. *Canadian Journal of Neurological Sciences*.
8 1997; 24(3):191-199
- 9 36. Leblanc R, Melanson D, Tampieri D, Guttmann RD. Familial cerebral aneurysms: a
10 study of 13 families. *Neurosurgery*. 1995; 37(4):633-638; discussion 638-639
- 11 37. Lee JS, Park IS, Park KB, Kang DH, Lee CH, Hwang SH. Familial intracranial
12 aneurysms. *Journal of Korean Neurosurgical Society*. 2008; 44(3):136-140
- 13 38. Lin TJ, Hwang FC, Chen CJ, Chiu WT, Chang CK. Familial hypertensive intracerebral
14 hemorrhage and autosomal dominant polycystic kidney disease. *Journal of Clinical*
15 *Neuroscience*. 2005; 12(4):474-477
- 16 39. Lindgaard L, Eskesen V, Gjerris F, Olsen NV. Familial aggregation of intracranial
17 aneurysms in an Inuit patient population in Kalaallit Nunaat (Greenland).
18 *Neurosurgery*. 2003; 52(2):357-362; discussion 362-353
- 19 40. Lorenzo-Betancor O, Blackburn PR, Edwards E, Vazquez-Do-campo R, Klee EW,
20 Labbe C et al. PCNT point mutations and familial intracranial Aneurysms. *Neurology*.
21 2018; 91(23):E2170-E2181
- 22 41. Lozano AM, Leblanc R. Familial intracranial aneurysms. *Journal of Neurosurgery*.
23 1987; 66(4):522-528
- 24 42. Magnetic Resonance Angiography in Relatives of Patients with Subarachnoid
25 Hemorrhage Study Group. Risks and benefits of screening for intracranial aneurysms
26 in first-degree relatives of patients with sporadic subarachnoid hemorrhage. *New*
27 *England Journal of Medicine*. 1999; 341(18):1344-1350
- 28 43. Mensing L, Ruigrok Y, Rinkel G. Rupture risk for familial compared to sporadic
29 intracranial aneurysms. *European Stroke Journal*. 2017; 2(1 Suppl 1):77
- 30 44. Mensing L, Ruigrok Y, Rinkel GJ. Rupture risk for familial compared to sporadic
31 intracranial aneurysms. *European Journal of Neurology*. 2017; 24 (Suppl 1):34
- 32 45. Mensing LA, Greving JP, Verhoeff TA, Rinkel GJE, Ruigrok YM. Comparison of
33 rupture risk of intracranial aneurysms between familial and sporadic patients. *Stroke*.
34 2019; 50(6):1380-1383
- 35 46. Mensing LA, Rinkel GJ, Vlak MH, van der Schaaf IC, Ruigrok YM. Difference in
36 aneurysm characteristics between patients with familial and sporadic aneurysmal
37 subarachnoid haemorrhage. *PloS One*. 2016; 11(4):e0154281
- 38 47. Miller TD, White PM, Davenport RJ, Al-Shahi Salman R. Screening patients with a
39 family history of subarachnoid haemorrhage for intracranial aneurysms: screening
40 uptake, patient characteristics and outcome. *Journal of Neurology, Neurosurgery and*
41 *Psychiatry*. 2012; 83(1):86-88
- 42 48. Nagae K, Goto I, Ueda K, Morotomi Y. Familial occurrence of multiple intracranial
43 aneurysms. Case report. *Journal of Neurosurgery*. 1972; 37(3):364-367
- 44 49. National Institute for Health and Care Excellence. Developing NICE guidelines: the
45 manual. London. National Institute for Health and Care Excellence, 2014. Available

- 1 from:
2 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20overview>
- 3 50. NHS England and NHS Improvement. National cost collection for the NHS 2018-19.
4 2019. Available from: <https://improvement.nhs.uk/resources/national-cost-collection/>
5 Last accessed: 01/04/2020.
- 6 51. Okamoto K, Horisawa R, Kawamura T, Asai A, Ogino M, Takagi T et al. Family
7 history and risk of subarachnoid hemorrhage: a case-control study in Nagoya, Japan.
8 *Stroke*. 2003; 34(2):422-426
- 9 52. Organisation for Economic Co-operation and Development (OECD). **{CONFIRM**
10 **ACCESS DATE WITH HE}** Purchasing power parities (PPP). 2012. Available from:
11 <http://www.oecd.org/std/ppp> Last accessed:
- 12 53. Pei Shan W, Longstreth Jr WT, Koepsell TD. Subarachnoid hemorrhage and family
13 history: a population-based case- control study. *Archives of Neurology*. 1995;
14 52(2):202-204
- 15 54. Phillips RL. Familial cerebral aneurysms. Case reports. *Journal of Neurosurgery*.
16 1963; 20:701-703
- 17 55. Raaymakers TW. Aneurysms in relatives of patients with subarachnoid hemorrhage:
18 frequency and risk factors. MARS Study Group. Magnetic Resonance Angiography in
19 Relatives of patients with Subarachnoid hemorrhage. *Neurology*. 1999; 53(5):982-988
- 20 56. Raaymakers TW, Rinkel GJ, Ramos LM. Initial and follow-up screening for
21 aneurysms in families with familial subarachnoid hemorrhage. *Neurology*. 1998;
22 51(4):1125-1130
- 23 57. Raaymakers TWM. Functional outcome and quality of life after angiography and
24 operation for unruptured intracranial aneurysms. *Journal of Neurology Neurosurgery*
25 *and Psychiatry*. 2000; 68(5):571-576
- 26 58. Rinkel GJ. Intracranial aneurysm screening: indications and advice for practice.
27 *Lancet Neurology*. 2005; 4(2):122-128
- 28 59. Rinkel GJ, Djibuti M, Algra A, van Gijn J. Prevalence and risk of rupture of intracranial
29 aneurysms: a systematic review. *Stroke*. 1998; 29(1):251-256
- 30 60. Roberts G, Nanra J, Phillips J. Screening for familial intracranial aneurysm: resource
31 implications. *British Journal of Neurosurgery*. 1999; 13(4):395-398
- 32 61. Ronkainen A, Hernesniemi J, Puranen M, Niemitukia L, Vanninen R, Ryyananen M et
33 al. Familial intracranial aneurysms. *Lancet*. 1997; 349(9049):380-384
- 34 62. Ronkainen A, Hernesniemi J, Ryyananen M, Puranen M, Kuivaniemi H. A ten percent
35 prevalence of asymptomatic familial intracranial aneurysms: preliminary report on 110
36 magnetic resonance angiography studies in members of 21 Finnish familial
37 intracranial aneurysm families. *Neurosurgery*. 1994; 35(2):208-212; discussion 212-
38 203
- 39 63. Ronkainen A, Hernesniemi J, Tromp G, Weir BKA, Schievink WI, Piepgras DG.
40 Special features of familial intracranial aneurysms: report of 215 familial aneurysms.
41 *Neurosurgery*. 1995; 37(1):43-47
- 42 64. Ronkainen A, Miettinen H, Karkola K, Papinaho S, Vanninen R, Puranen M et al. Risk
43 of harboring an unruptured intracranial aneurysm. *Stroke*. 1998; 29(2):359-362
- 44 65. Ronkainen A, Niskanen M, Piironen R, Hernesniemi J. Familial subarachnoid
45 hemorrhage. Outcome study. *Stroke*. 1999; 30(5):1099-1102

- 1 66. Ronkainen A, Puranen MI, Hernesniemi JA, Vanninen RL, Kaarina Partanen PL,
2 Tapani Saari J et al. Intracranial aneurysms: MR angiographic screening in 400
3 asymptomatic individuals with increased familial risk. *Radiology*. 1995; 195(1):35-40
- 4 67. Ronkainen A, Vanninen R, Hernesniemi J. Familial aneurysms. *Headache Quarterly*.
5 1998; 9(1):34-38
- 6 68. Ruigrok YM, Rinkel GJ, Algra A, Raaymakers TW, Van Gijn J. Characteristics of
7 intracranial aneurysms in patients with familial subarachnoid hemorrhage. *Neurology*.
8 2004; 62(6):891-894
- 9 69. Sakai N, Sakata K, Yamada H, Yamamoto M, Aiba T, Takeda F. Familial occurrence
10 of intracranial aneurysms. *Surgical Neurology*. 1974; 2(1):25-29
- 11 70. Schievink WI, Schaid DJ, Rogers HM, Piepgras DG, Michels VV. On the inheritance
12 of intracranial aneurysms. *Stroke*. 1994; 25(10):2028-2037
- 13 71. Sundquist J, Li X, Sundquist K, Hemminki K. Risks of subarachnoid hemorrhage in
14 siblings: a nationwide epidemiological study from Sweden. *Neuroepidemiology*. 2007;
15 29(3-4):178-184
- 16 72. Takao H, Nojo T, Ohtomo K. Screening for familial intracranial aneurysms. Decision
17 and cost-effectiveness analysis. *Academic Radiology*. 2008; 15(4):462-471
- 18 73. Teasdale GM, Wardlaw JM, White PM, Murray G, Teasdale EM, Easton V. The
19 familial risk of subarachnoid haemorrhage. *Brain*. 2005; 128(7):1677-1685
- 20 74. ter Berg HW, Bijlsma JB, Veiga Pires JA, Ludwig JW, van der Heiden C, Tulleken CA
21 et al. Familial association of intracranial aneurysms and multiple congenital
22 anomalies. *Archives of Neurology*. 1986; 43(1):30-33
- 23 75. ter Berg HW, Dippel DW, Habbema JD, Bijlsma JB, van Gijn J, Tulleken CA et al.
24 Treatment of intact familial intracranial aneurysms: a decision-analytical approach.
25 *Neurosurgery*. 1988; 23(3):329-334
- 26 76. ter Berg HW, Dippel DW, Limburg M, Schievink WI, van Gijn J. Familial intracranial
27 aneurysms. A review. *Stroke*. 1992; 23(7):1024-1030
- 28 77. Toglia JU, Samii AR. Familial intracranial aneurysms. *Diseases of the Nervous*
29 *System*. 1972; 33(9):611-613
- 30 78. van der Voet M, Olson JM, Kuivaniemi H, Dudek DM, Skunca M, Ronkainen A et al.
31 Intracranial aneurysms in Finnish families: confirmation of linkage and refinement of
32 the interval to chromosome 19q13.3. *American Journal of Human Genetics*. 2004;
33 74(3):564-571
- 34 79. Verdure P, Gilard V, Guyant-Marechal L, Belien J, Cebula H, Hannequin D et al.
35 Familial intracranial aneurysm, the relationship of the aortic diameter. *Neuro-*
36 *Chirurgie*. 2015; 61(6):385-391
- 37 80. Vlak MH, Rinkel GJ, Greebe P, Algra A. Risk of rupture of an intracranial aneurysm
38 based on patient characteristics: a case-control study. *Stroke*. 2013; 44(5):1256-1259
- 39 81. Wermer MJ, Rinkel GJ, van Gijn J. Repeated screening for intracranial aneurysms in
40 familial subarachnoid hemorrhage. *Stroke*. 2003; 34(12):2788-2791
- 41 82. White PM. How to manage the patient with a family history of aneurysmal
42 subarachnoid haemorrhage. *Practical Neurology*. 2004; 4(2):88-103

- 1 83. Woo D, Khoury J, Haverbusch MM, Sekar P, Flaherty ML, Kleindorfer DO et al.
2 Smoking and family history and risk of aneurysmal subarachnoid hemorrhage.
3 Neurology. 2009; 72(1):69-72
- 4 84. Zuurbier C, Mensing L, Wermer M, Juvela S, Lindgren A, Jaaskelainen J et al.
5 Comparison of rupture risk between familial and sporadic intracranial aneurysms: an
6 individual patient data meta-analysis. European Journal of Neurology. 2019; 26(Suppl
7 1):282
- 8 85. Zuurbier CCM, Mensing LA, Wermer MJH, Juvela S, Lindgren AE, Jaaskelainen JE
9 et al. Comparison of rupture risk between familial and sporadic intracranial
10 aneurysms: an individual patient data meta-analysis. European Stroke Journal. 2019;
11 4(Suppl 1):75-76
- 12

1 Appendices

2 Appendix A: Review protocols

3 Table 4: Review protocol: Detecting aneurysms in relatives of people with SAH

Field	Content
PROSPERO registration number	CRD42019160102
Review title	What is the clinical and cost effectiveness of investigations to detect intracranial arterial aneurysms in relatives of adults who have had a subarachnoid haemorrhage?
Review question	What is the clinical and cost effectiveness of investigations to detect intracranial arterial aneurysms in relatives of adults who have had a subarachnoid haemorrhage?
Objective	To determine if investigations to detect intracranial arterial aneurysms in relatives of people with SAH is clinically and cost-effective.
Searches	<p>The following databases will be searched:</p> <ul style="list-style-type: none"> • Cochrane Central Register of Controlled Trials (CENTRAL) • Cochrane Database of Systematic Reviews (CDSR) • Embase • MEDLINE <p>Searches will be restricted by:</p> <ul style="list-style-type: none"> • English language only <p>The searches may be re-run 6 weeks before final committee meeting and further studies retrieved for inclusion if relevant.</p> <p>The full search strategies will be published in the final review.</p>
Condition or domain being studied	Aneurysmal subarachnoid haemorrhage
Population	<p>Inclusion: First degree relatives of adults (16 and older) with a confirmed subarachnoid haemorrhage caused by a ruptured aneurysm.</p> <p>Exclusion:</p> <ul style="list-style-type: none"> • Adults with subarachnoid haemorrhage caused by head injury, ischaemic stroke or an arteriovenous malformation. • Children and young people aged 15 years and younger.
Intervention/Exposure/Test	<ul style="list-style-type: none"> • Assessment of first degree relatives of people with aSAH • Assessment with CTA or MRA • Diagnostic accuracy of CTA and MRA has been shown to be high. DSA is invasive and so should not be used as a screening intervention on relatives who may not have aneurysms.
Comparator/Reference standard/Confounding factors	<p>Comparators:</p> <ul style="list-style-type: none"> • To no routine assessment
Types of study to be included	Randomised controlled trials (RCTs), systematic reviews of RCTs. If insufficient RCT evidence is available, non-randomised studies will be considered if they adjust for key confounders (age), starting with prospective cohort studies.
Other exclusion criteria	Exclusions:

Field	Content
	<ul style="list-style-type: none"> • Non-English language studies • Conference abstracts will be excluded as it is expected there will be sufficient full text published studies available.
Context	n/a
Primary outcomes (critical outcomes)	<ul style="list-style-type: none"> • Mortality • Health and social-related quality of life (any validated measure) • Degree of disability or dependence in daily activities, (any validated measure e.g. Modified Rankin Scale and patient-reported outcome measures) • Subarachnoid haemorrhage <p>Outcomes will be captured at or after the point of imaging follow-up. Outcomes will therefore be grouped at <1 year, 1-2 years and >2-5 years.</p>
Secondary outcomes (important outcomes)	<ul style="list-style-type: none"> • Presence of cerebral aneurysm • Elective treatment <p>Outcomes will be captured at or after the point of imaging follow-up. Outcomes will therefore be grouped at <1 year, 1-2 years and >2-5 years.</p>
Data extraction (selection and coding)	<ul style="list-style-type: none"> • EndNote will be used for reference management, sifting, citations and bibliographies. All references identified by the searches and from other sources will be screened for inclusion. 10% of the abstracts will be reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer. The full text of potentially eligible studies will be retrieved and will be assessed in line with the criteria outlined above. • EviBASE will be used for data extraction. • If not an intervention review, add: A standardised form will be used to extract data from studies (see Developing NICE guidelines: the manual section 6.4).
Risk of bias (quality) assessment	<p>Risk of bias will be assessed using the appropriate checklist as described in Developing NICE guidelines: the manual.</p> <p>Systematic reviews: Risk of Bias in Systematic Reviews (ROBIS)</p> <p>Randomised Controlled Trial: Cochrane RoB (2.0)</p> <p>Non randomised study, including cohort studies: Cochrane ROBINS-I</p> <p>10% of all evidence reviews are quality assured by a senior research fellow. This includes checking:</p> <ul style="list-style-type: none"> • papers were included /excluded appropriately • a sample of the data extractions • correct methods are used to synthesise data • a sample of the risk of bias assessments <p>Disagreements between the review authors over the risk of bias in particular studies will be resolved by discussion, with involvement of a third review author where necessary.</p>
Strategy for data synthesis	<p>Pairwise meta-analyses will be performed using Cochrane Review Manager (RevMan5).</p> <p>GRADEpro will be used to assess the quality of evidence for each outcome, taking into account individual study quality and the meta-</p>

Field	Content														
	<p>analysis results. The 4 main quality elements (risk of bias, indirectness, inconsistency and imprecision) will be appraised for each outcome. Publication bias is tested for when there are more than 5 studies for an outcome.</p> <p>The risk of bias across all available evidence was evaluated for each outcome using an adaptation of the 'Grading of Recommendations Assessment, Development and Evaluation (GRADE) toolbox' developed by the international GRADE working group http://www.gradeworkinggroup.org/</p> <p>Where meta-analysis is not possible, data will be presented and quality assessed individually per outcome.</p> <p>Subgroups will be investigated separately if meta-analysed results show heterogeneity.</p> <p>Heterogeneity between the studies in effect measures will be assessed using the I² statistic and visually inspected. An I² value greater than 50% will be considered indicative of substantial heterogeneity. Sensitivity analyses will be conducted based on pre-specified subgroups using stratified meta-analysis to explore the heterogeneity in effect estimates. If this does not explain the heterogeneity, the results will be presented pooled using random-effects.</p>														
Analysis of sub-groups	<p>Strata:</p> <ul style="list-style-type: none"> • n/a <p>Subgroups for heterogeneity:</p> <ul style="list-style-type: none"> • Extent of familial history: <ul style="list-style-type: none"> • single first degree relative • multiple first-degree relatives • Smoking status <ul style="list-style-type: none"> • Smoker • Non-smoker • Blood pressure <ul style="list-style-type: none"> • Hypertensive (>140/90) • Non-hypertensive (<140/90) • Polycystic kidney <ul style="list-style-type: none"> ○ Yes ○ No • Genetic causes present (identifiable connective tissue disorders e.g. Marfans, Ehlers-Danlos) <ul style="list-style-type: none"> ○ Yes ○ No 														
Type and method of review	<table border="1"> <tbody> <tr> <td><input checked="" type="checkbox"/></td> <td>Intervention</td> </tr> <tr> <td><input checked="" type="checkbox"/></td> <td>Diagnostic</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Prognostic</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Qualitative</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Epidemiologic</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Service Delivery</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Other (please specify)</td> </tr> </tbody> </table>	<input checked="" type="checkbox"/>	Intervention	<input checked="" type="checkbox"/>	Diagnostic	<input type="checkbox"/>	Prognostic	<input type="checkbox"/>	Qualitative	<input type="checkbox"/>	Epidemiologic	<input type="checkbox"/>	Service Delivery	<input type="checkbox"/>	Other (please specify)
<input checked="" type="checkbox"/>	Intervention														
<input checked="" type="checkbox"/>	Diagnostic														
<input type="checkbox"/>	Prognostic														
<input type="checkbox"/>	Qualitative														
<input type="checkbox"/>	Epidemiologic														
<input type="checkbox"/>	Service Delivery														
<input type="checkbox"/>	Other (please specify)														
Language	English														
Country	England														
Anticipated or actual start date															

Field	Content		
Anticipated completion date	3 February 2021		
Stage of review at time of this submission	Review stage	Started	Completed
	Preliminary searches	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Piloting of the study selection process	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Formal screening of search results against eligibility criteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Data extraction	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Risk of bias (quality) assessment	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Data analysis	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Named contact	5a. Named contact National Guideline Centre 5b Named contact e-mail SAH@nice.org.uk 5e Organisational affiliation of the review National Institute for Health and Care Excellence (NICE) and the National Guideline Centre		
Review team members	From the National Guideline Centre: <ul style="list-style-type: none"> • Ms Gill Ritchie • Mr Ben Mayer • Mr Audrius Stonkus • Mr Vimal Bedia • Ms Emma Cowles • Ms Jill Cobb • Ms Elizabeth Pearton • Ms Amelia Unsworth 		
Funding sources/sponsor	This systematic review is being completed by the National Guideline Centre which receives funding from NICE.		
Conflicts of interest	All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline.		
Collaborators	Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of Developing NICE guidelines: the manual. Members of the guideline committee are available on the NICE website.		
Other registration details			
Reference/URL for published protocol			

Field	Content										
Dissemination plans	NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as: notifying registered stakeholders of publication publicising the guideline through NICE's newsletter and alerts issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE.										
Keywords											
Details of existing review of same topic by same authors	None										
Current review status	<table border="1"> <tr> <td><input checked="" type="checkbox"/></td> <td>Ongoing</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Completed but not published</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Completed and published</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Completed, published and being updated</td> </tr> <tr> <td><input type="checkbox"/></td> <td>Discontinued</td> </tr> </table>	<input checked="" type="checkbox"/>	Ongoing	<input type="checkbox"/>	Completed but not published	<input type="checkbox"/>	Completed and published	<input type="checkbox"/>	Completed, published and being updated	<input type="checkbox"/>	Discontinued
<input checked="" type="checkbox"/>	Ongoing										
<input type="checkbox"/>	Completed but not published										
<input type="checkbox"/>	Completed and published										
<input type="checkbox"/>	Completed, published and being updated										
<input type="checkbox"/>	Discontinued										
Additional information											
Details of final publication	www.nice.org.uk										

1

2 **Table 5: Health economic review protocol**

Review question	All questions where health economic evidence applicable
Objectives	To identify health economic studies relevant to any of the review questions.
Search criteria	<ul style="list-style-type: none"> • Populations, interventions and comparators must be as specified in the clinical review protocol above. • Studies must be of a relevant health economic study design (cost–utility analysis, cost-effectiveness analysis, cost–benefit analysis, cost–consequences analysis, comparative cost analysis). • Studies must not be a letter, editorial or commentary, or a review of health economic evaluations. (Recent reviews will be ordered although not reviewed. The bibliographies will be checked for relevant studies, which will then be ordered.) • Unpublished reports will not be considered unless submitted as part of a call for evidence. • Studies must be in English.
Search strategy	A health economic study search will be undertaken using population-specific terms and a health economic study filter.
Review strategy	<p>Studies not meeting any of the search criteria above will be excluded. Studies published before 2003, abstract-only studies and studies from non-OECD countries or the USA will also be excluded.</p> <p>Each remaining study will be assessed for applicability and methodological limitations using the NICE economic evaluation checklist which can be found in appendix H of Developing NICE guidelines: the manual.⁴⁹</p> <p>Inclusion and exclusion criteria</p> <ul style="list-style-type: none"> • If a study is rated as both 'Directly applicable' and with 'Minor limitations' then it will be included in the guideline. A health economic evidence table will be completed and it will be included in the health economic evidence profile. • If a study is rated as either 'Not applicable' or with 'Very serious limitations' then it will usually be excluded from the guideline. If it is excluded then a health economic

evidence table will not be completed and it will not be included in the health economic evidence profile.

- If a study is rated as 'Partially applicable', with 'Potentially serious limitations' or both then there is discretion over whether it should be included.

Where there is discretion

The health economist will decide based on the relative applicability and quality of the available evidence for that question, in discussion with the guideline committee if required. The ultimate aim is to include health economic studies that are helpful for decision-making in the context of the guideline and the current NHS setting. If several studies are considered of sufficiently high applicability and methodological quality that they could all be included, then the health economist, in discussion with the committee if required, may decide to include only the most applicable studies and to selectively exclude the remaining studies. All studies excluded based on applicability or methodological limitations will be listed with explanation in the excluded health economic studies appendix below.

The health economist will be guided by the following hierarchies.

Setting:

- UK NHS (most applicable).
- OECD countries with predominantly public health insurance systems (for example, France, Germany, Sweden).
- OECD countries with predominantly private health insurance systems (for example, Switzerland).
- Studies set in non-OECD countries or in the USA will be excluded before being assessed for applicability and methodological limitations.

Health economic study type:

- Cost–utility analysis (most applicable).
- Other type of full economic evaluation (cost–benefit analysis, cost-effectiveness analysis, cost–consequences analysis).
- Comparative cost analysis.
- Non-comparative cost analyses including cost-of-illness studies will be excluded before being assessed for applicability and methodological limitations.

Year of analysis:

- The more recent the study, the more applicable it will be.
- Studies published in 2003 or later but that depend on unit costs and resource data entirely or predominantly from before 2003 will be rated as 'Not applicable'.
- Studies published before 2003 will be excluded before being assessed for applicability and methodological limitations.

Quality and relevance of effectiveness data used in the health economic analysis:

- The more closely the clinical effectiveness data used in the health economic analysis match with the outcomes of the studies included in the clinical review the more useful the analysis will be for decision-making in the guideline.

1

2 Appendix B: Literature search strategies

3 This literature search strategy was used for the following review;

- 4 • What is the clinical and cost effectiveness of investigations to detect intracranial
5 arterial aneurysms in relatives of adults who have had a subarachnoid haemorrhage?

6 The literature searches for this review are detailed below and complied with the methodology
7 outlined in Developing NICE guidelines: the manual⁴⁹

- 1 For more information, please see the Methods Report published as part of the accompanying
- 2 documents for this guideline.

B.1.3 Clinical search literature search strategy

- 4 Searches were constructed using a PICO framework where population (P) terms were
- 5 combined with Intervention (I) and in some cases Comparison (C) terms. Outcomes (O) are
- 6 rarely used in search strategies for interventions as these concepts may not be well
- 7 described in title, abstract or indexes and therefore difficult to retrieve. Search filters were
- 8 applied to the search where appropriate.

9 **Table 6: Database date parameters and filters used**

Database	Dates searched	Search filter used
Medline (OVID)	1946 – 26 June 2020	Exclusions
Embase (OVID)	1974 – 26 June 2020	Exclusions
The Cochrane Library (Wiley)	Cochrane Reviews to 2020 Issue 6 of 12 CENTRAL to 2020 Issue 6 of 12	None

10 **Medline (Ovid) search terms**

1.	exp Subarachnoid Hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	Intracranial Aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracerebral or intra-cerebral or intracranial or intra-cranial or brain) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.
15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16
18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23

25.	6 not 24
26.	(exp child/ or exp pediatrics/ or exp infant/) not (exp adolescent/ or exp adult/ or exp middle age/ or exp aged/)
27.	25 not 26
28.	limit 27 to English language
29.	Magnetic Resonance.ti,ab.
30.	(MRA or MRAs or MRI or MRIs or NMR or NMRs).ti,ab.
31.	Magnetic Resonance Angiography/ or Magnetic Resonance Imaging/
32.	(tomograph* or CTA or CTAs or CAT).ti,ab.
33.	tomography, x-ray computed/ or computed tomography angiography/
34.	diagnostic imaging/ or tomography/
35.	(angiogra* or arteriogra*).ti,ab.
36.	cerebral angiography/
37.	Mass screening/
38.	screen*.ti,ab.
39.	(scan* or image* or imaging).ti,ab.
40.	risk assess*.ti,ab.
41.	Risk assessment/
42.	or/29-41
43.	(family or families or familial or parent* or sibling*).ti,ab.
44.	(first degree adj2 (relative* or relation*)).ti,ab.
45.	(hereditary or heredity or heritability or inherit*).ti,ab.
46.	((genetic or gene or genes) adj2 (predispos* or factor* or influenc* or risk* or determin* or prognos* or test or tests or defect* or fault* or mutation or mutate* or mutant)).ti,ab.
47.	genetic predisposition to disease/
48.	family health/
49.	Heredity/
50.	Family/
51.	or/43-50
52.	28 and 42 and 51

1 Embase (Ovid) search terms

1.	*subarachnoid hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp intracranial aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	Case report/ or Case study/
11.	(letter or comment*).ti.
12.	or/7-11

13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	Nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental animal/
19.	Animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	(exp child/ or exp pediatrics/) not (exp adult/ or exp adolescent/)
25.	23 not 24
26.	limit 25 to English language
27.	Magnetic Resonance.ti,ab.
28.	(MRA or MRAs or MRI or MRIs or NMR or NMRs).ti,ab.
29.	nuclear magnetic resonance imaging/ or magnetic resonance angiography/
30.	(tomograph* or CTA or CTAs or CAT).ti,ab.
31.	computer assisted tomography/ or computed tomographic angiography/
32.	brain angiography/
33.	mass screening/
34.	screening test/
35.	screen*.ti,ab.
36.	(angiogra* or arteriogra*).ti,ab.
37.	(scan* or image* or imaging).ti,ab.
38.	diagnostic imaging/
39.	risk assess*.ti,ab.
40.	risk assessment/
41.	or/27-40
42.	(family or families or familial or parent* or sibling*).ti,ab.
43.	(first degree adj2 (relative* or relation*)).ti,ab.
44.	(hereditary or heritability or heredity or inherit*).ti,ab.
45.	((genetic or gene or genes) adj2 (predispos* or factor* or influenc* or risk* or determin* or prognos* or test or tests or defect* or fault* or mutation or mutate* or mutant)).ti,ab.
46.	genetic predisposition/
47.	family health/
48.	heredity/
49.	familial disease/ or "genetic and familial disorders"/
50.	family/
51.	family assessment/
52.	or/42-51
53.	26 and 41 and 52

1 Cochrane Library (Wiley) search terms

#1.	MeSH descriptor: [Subarachnoid Hemorrhage] explode all trees
-----	--

#2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) near/3 (hemorrhag* or haemorrhag* or bleed* or blood*)):ti,ab
#3.	(SAH or aSAH):ti,ab
#4.	MeSH descriptor: [Intracranial Aneurysm] this term only
#5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) near/3 (aneurysm* or aneurism* or hematoma* or haematoma*)):ti,ab
#6.	(or #1-#5)
#7.	Magnetic Resonance:ti,ab
#8.	(MRA or MRAs or MRI or MRIs or NMR or NMRs):ti,ab
#9.	MeSH descriptor: [Magnetic Resonance Angiography] this term only
#10.	MeSH descriptor: [Magnetic Resonance Imaging] this term only
#11.	(tomograph* or CTA or CTAs or CAT):ti,ab
#12.	MeSH descriptor: [Tomography, X-Ray Computed] this term only
#13.	MeSH descriptor: [Computed Tomography Angiography] this term only
#14.	MeSH descriptor: [Diagnostic Imaging] this term only
#15.	MeSH descriptor: [Tomography] this term only
#16.	(angiogra* or arteriogra*):ti,ab
#17.	MeSH descriptor: [Cerebral Angiography] this term only
#18.	MeSH descriptor: [Mass Screening] this term only
#19.	screen*:ti,ab
#20.	(scan* or image* or imaging):ti,ab
#21.	risk assess*:ti,ab
#22.	MeSH descriptor: [Risk Assessment] this term only
#23.	(or #7-#22)
#24.	(family or families or familial or parent* or sibling*):ti,ab
#25.	(first degree next/2 (relative* or relation*)):ti,ab
#26.	(hereditary or heredity or heritability or inherit*):ti,ab
#27.	((genetic or gene or genes) next/2 (predispos* or factor* or influenc* or risk* or determin* or prognos* or test or tests or defect* or fault* or mutation or mutate* or mutant)):ti,ab
#28.	MeSH descriptor: [Genetic Predisposition to Disease] this term only
#29.	MeSH descriptor: [Family Health] this term only
#30.	MeSH descriptor: [Heredity] this term only
#31.	MeSH descriptor: [Family] this term only
#32.	(or #24-#31)
#33.	#6 and #23 and #32

B.2.1 Health Economics literature search strategy

2 Health economic evidence was identified by conducting a broad search relating to
3 subarachnoid haemorrhage population in NHS Economic Evaluation Database (NHS EED –
4 this ceased to be updated after March 2015) and the Health Technology Assessment
5 database (HTA) with no date restrictions. NHS EED and HTA databases are hosted by the
6 Centre for Research and Dissemination (CRD). Additional searches were run on Medline and
7 Embase.

1 Table 7: Database date parameters and filters used

Database	Dates searched	Search filter used
Medline	2003 – 23 June 2020	Exclusions Health economics studies
Embase	2003 – 23 June 2020	Exclusions Health economics studies
Centre for Research and Dissemination (CRD)	HTA - Inception – 23 June 2020 NHSEED - Inception to March 2015	None

2 Medline (Ovid) search terms

1.	exp Subarachnoid Hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp Intracranial Aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter/
8.	editorial/
9.	news/
10.	exp historical article/
11.	Anecdotes as Topic/
12.	comment/
13.	case report/
14.	(letter or comment*).ti.
15.	or/7-14
16.	randomized controlled trial/ or random*.ti,ab.
17.	15 not 16
18.	animals/ not humans/
19.	exp Animals, Laboratory/
20.	exp Animal Experimentation/
21.	exp Models, Animal/
22.	exp Rodentia/
23.	(rat or rats or mouse or mice).ti.
24.	or/17-23
25.	6 not 24
26.	limit 25 to English language
27.	Economics/
28.	Value of life/
29.	exp "Costs and Cost Analysis"/
30.	exp Economics, Hospital/
31.	exp Economics, Medical/
32.	Economics, Nursing/

33.	Economics, Pharmaceutical/
34.	exp "Fees and Charges"/
35.	exp Budgets/
36.	budget*.ti,ab.
37.	cost*.ti.
38.	(economic* or pharmaco?economic*).ti.
39.	(price* or pricing*).ti,ab.
40.	(cost* adj2 (effective* or utilit* or benefit* or minimi* or unit* or estimat* or variable*)).ab.
41.	(financ* or fee or fees).ti,ab.
42.	(value adj2 (money or monetary)).ti,ab.
43.	or/27-42
44.	26 and 43

1 Embase (Ovid) search terms

1.	subarachnoid hemorrhage/
2.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)).ti,ab.
3.	(SAH or aSAH).ti,ab.
4.	exp intracranial aneurysm/
5.	((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial or brain or saccular or berry or wide-neck*) adj3 (aneurysm* or aneurism* or hematoma* or haematoma*)).ti,ab.
6.	or/1-5
7.	letter.pt. or letter/
8.	note.pt.
9.	editorial.pt.
10.	case report/ or case study/
11.	(letter or comment*).ti.
12.	or/7-11
13.	randomized controlled trial/ or random*.ti,ab.
14.	12 not 13
15.	animal/ not human/
16.	nonhuman/
17.	exp Animal Experiment/
18.	exp Experimental Animal/
19.	animal model/
20.	exp Rodent/
21.	(rat or rats or mouse or mice).ti.
22.	or/14-21
23.	6 not 22
24.	limit 23 to English language
25.	health economics/
26.	exp economic evaluation/
27.	exp health care cost/
28.	exp fee/

29.	budget/
30.	funding/
31.	budget*.ti,ab.
32.	cost*.ti.
33.	(economic* or pharmaco?economic*).ti.
34.	(price* or pricing*).ti,ab.
35.	(cost* adj2 (effective* or utilit* or benefit* or minimi* or unit* or estimat* or variable*)).ab.
36.	(financ* or fee or fees).ti,ab.
37.	(value adj2 (money or monetary)).ti,ab.
38.	or/25-37
39.	24 and 38


1 NHS EED and HTA (CRD) search terms

#1.	MeSH DESCRIPTOR Subarachnoid Hemorrhage EXPLODE ALL TREES
#2.	MeSH DESCRIPTOR Intracranial Hemorrhages EXPLODE ALL TREES
#3.	(((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (hemorrhag* or haemorrhag* or bleed* or blood*)))
#4.	((SAH or aSAH))
#5.	#1 OR #2 OR #3 OR #4
#6.	MeSH DESCRIPTOR Aneurysm EXPLODE ALL TREES
#7.	((aneurysm* or hematoma* or haematoma*))
#8.	#6 OR #7
#9.	MeSH DESCRIPTOR Intracranial Aneurysm EXPLODE ALL TREES
#10.	(((subarachnoid* or arachnoid* or cerebral or intracranial or intra-cranial) adj3 (aneurysm* or hematoma* or haematoma*)))
#11.	#9 OR #10
#12.	MeSH DESCRIPTOR Aneurysm, ruptured
#13.	(((ruptur* or weak* or brain or trauma*) adj3 (aneurysm* or hematoma* or haematoma*)))
#14.	#12 OR #13
#15.	(#5 or #8 or #11 or #14)

2

1 Appendix C: Clinical evidence selection

Figure 1: Flow chart of clinical study selection for the review of detecting aneurysms in relatives of people with SAH



2

3

1 **Appendix D: Clinical evidence tables**

2 No studies were included.

3

1 **Appendix E: Forest plots**

2 No studies were included.

3

4

1 **Appendix F: GRADE tables**


2 No studies were included.

3

4

1 **Appendix G: Health economic evidence**
2 **selection**
3

Figure 2: Flow chart of health economic study selection for the guideline



* Non-relevant population, intervention, comparison, design or setting; non-English language

¹ Appendix H: Health economic evidence tables

Study	Hopmans 2016																																																																																																										
Study details	Population & Interventions	Costs		Health outcomes	Cost effectiveness																																																																																																						
<p>Economic analysis: CUA (health outcome: QALYs)</p> <p>Study design: Probabilistic decision analytic model</p> <p>Approach to analysis: Markov model with health states: healthy no aneurysm, healthy with aneurysm, healthy with known small aneurysm, disabled, dead. 1 year cycles. Incidence of aneurysm and rupture risk constant throughout lifetime. Detected aneurysms >5mm treated with either clipping or coiling.</p> <p>Perspective: Dutch healthcare provider</p> <p>Time horizon: Lifetime</p> <p>Treatment effect duration: n/a</p> <p>Discounting: Costs: 4%; Outcomes: 1.5%</p>	<p>Population: People with one affected first degree relative with aneurysmal subarachnoid haemorrhage.</p> <p>Cohort settings: Start age: 15 Male: NR</p> <p>Interventions:</p> <ol style="list-style-type: none"> 1. No screening 2. Screening every 5 years, age 20 to 60 3. Screening every 10 years, age 20 to 60 4. Screening every 15 years, age 20 to 60 5. Screening every 5 years, age 20 to 70 6. Screening every 10 years, age 20 to 70 7. Screening every 15 years, age 20 to 70 8. Screening every 5 years, age 30 to 60 9. Screening every 10 years, age 30 to 60 10. Screening every 15 years, age 30 to 60 11. Screening every 5 years, age 30 to 70 	<p>Full incremental analysis^{(a)(b)}</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Intv</th> <th>Cost</th> <th>QALY</th> <th>Incr cost</th> <th>Incr QALY</th> <th>ICER</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>£0</td> <td>0</td> <td colspan="3" style="text-align: center;">Baseline</td> </tr> <tr> <td>25</td> <td>£76</td> <td>0.01</td> <td>£76</td> <td>0.01</td> <td>£7,600</td> </tr> <tr> <td>23</td> <td>£109</td> <td>0.011</td> <td colspan="3" style="text-align: center;">Dominated by 24</td> </tr> <tr> <td>24</td> <td>£90</td> <td>0.011</td> <td>£14</td> <td>0.001</td> <td>Extendedly dominated</td> </tr> <tr> <td>22</td> <td>£126</td> <td>0.013</td> <td>£50</td> <td>0.003</td> <td>Extendedly dominated</td> </tr> <tr> <td>20</td> <td>£184</td> <td>0.014</td> <td colspan="3" style="text-align: center;">Dominated by 21</td> </tr> <tr> <td>21</td> <td>£147</td> <td>0.016</td> <td>£71</td> <td>0.006</td> <td>Extendedly dominated</td> </tr> <tr> <td>16</td> <td>£215</td> <td>0.027</td> <td>£139</td> <td>0.017</td> <td>£8,176</td> </tr> <tr> <td>19</td> <td>£253</td> <td>0.028</td> <td>£38</td> <td>0.001</td> <td>Extendedly dominated</td> </tr> <tr> <td>15</td> <td>£279</td> <td>0.03</td> <td>£64</td> <td>0.003</td> <td>Extendedly dominated</td> </tr> <tr> <td>10</td> <td>£361</td> <td>0.035</td> <td colspan="3" style="text-align: center;">Dominated by 18</td> </tr> <tr> <td>18</td> <td>£318</td> <td>0.035</td> <td>£103</td> <td>0.008</td> <td>£12,875</td> </tr> <tr> <td>13</td> <td>£369</td> <td>0.036</td> <td>£51</td> <td>0.001</td> <td>Extendedly dominated</td> </tr> <tr> <td>14</td> <td>£430</td> <td>0.038</td> <td>£112</td> <td>0.003</td> <td>Extendedly dominated</td> </tr> <tr> <td>4</td> <td>£509</td> <td>0.043</td> <td colspan="3" style="text-align: center;">Dominated by 9</td> </tr> <tr> <td>9</td> <td>£460</td> <td>0.044</td> <td>£142</td> <td>0.009</td> <td>£15,778</td> </tr> </tbody> </table>				Intv	Cost	QALY	Incr cost	Incr QALY	ICER	1	£0	0	Baseline			25	£76	0.01	£76	0.01	£7,600	23	£109	0.011	Dominated by 24			24	£90	0.011	£14	0.001	Extendedly dominated	22	£126	0.013	£50	0.003	Extendedly dominated	20	£184	0.014	Dominated by 21			21	£147	0.016	£71	0.006	Extendedly dominated	16	£215	0.027	£139	0.017	£8,176	19	£253	0.028	£38	0.001	Extendedly dominated	15	£279	0.03	£64	0.003	Extendedly dominated	10	£361	0.035	Dominated by 18			18	£318	0.035	£103	0.008	£12,875	13	£369	0.036	£51	0.001	Extendedly dominated	14	£430	0.038	£112	0.003	Extendedly dominated	4	£509	0.043	Dominated by 9			9	£460	0.044	£142	0.009	£15,778
Intv	Cost	QALY	Incr cost	Incr QALY	ICER																																																																																																						
1	£0	0	Baseline																																																																																																								
25	£76	0.01	£76	0.01	£7,600																																																																																																						
23	£109	0.011	Dominated by 24																																																																																																								
24	£90	0.011	£14	0.001	Extendedly dominated																																																																																																						
22	£126	0.013	£50	0.003	Extendedly dominated																																																																																																						
20	£184	0.014	Dominated by 21																																																																																																								
21	£147	0.016	£71	0.006	Extendedly dominated																																																																																																						
16	£215	0.027	£139	0.017	£8,176																																																																																																						
19	£253	0.028	£38	0.001	Extendedly dominated																																																																																																						
15	£279	0.03	£64	0.003	Extendedly dominated																																																																																																						
10	£361	0.035	Dominated by 18																																																																																																								
18	£318	0.035	£103	0.008	£12,875																																																																																																						
13	£369	0.036	£51	0.001	Extendedly dominated																																																																																																						
14	£430	0.038	£112	0.003	Extendedly dominated																																																																																																						
4	£509	0.043	Dominated by 9																																																																																																								
9	£460	0.044	£142	0.009	£15,778																																																																																																						

12. Screening every 10 years, age 30 to 70	12	£498	0.044	Dominated by 9		
13. Screening every 15 years, age 30 to 70	17	£505	0.045	£45	0.001	Extendedly dominated
14. Screening every 5 years, age 40 to 60	7	£558	0.048	£98	0.004	Extendedly dominated
15. Screening every 10 years, age 40 to 60	3	£707	0.056	£247	0.012	Extendedly dominated
16. Screening every 15 years, age 40 to 60	6	£746	0.06	£286	0.016	Extendedly dominated
17. Screening every 5 years, age 40 to 70	8	£746	0.06	£286	0.016	Extendedly dominated
18. Screening every 10 years, age 40 to 70	11	£821	0.066	£361	0.022	£16,409
19. Screening every 15 years, age 40 to 70	2	£1,197	0.076	£376	0.01	Extendedly dominated
20. Screening once at age 30	5	£1,272	0.084	£451	0.018	£25,056
21. Screening once at age 35						
22. Screening once at age 40						
23. Screening once at age 45						
24. Screening once at age 50						
25. Screening once at age 55						

			<p>Currency & cost year: 2012 Euros (presented here as 2012 UK pounds^(c))</p> <p>Cost components incorporated:</p> <p>Costs of screening - MRA and outpatient consultation with a neurologist and specialised nurse.</p> <p>Costs of UIA treatment - hospitalisation, follow-up in outpatient clinic and imaging.</p> <p>Cost associated with being disabled (in nursing home).</p>	<p>Each individual screening strategy compared to no screening is cost effective at £20,000 per QALY gained.</p> <p>The most cost effective screening strategy when compared to each other is 11 (screening every 5 years, age 30 to 70).</p> <p>Analysis of uncertainty:</p> <p>No further sensitivity analyses undertaken.</p>
--	--	--	--	---

Data sources

Health outcomes: Calibrated the probability of unruptured intracranial aneurysm developing and rupture risk to ensure unruptured intracranial aneurysm prevalence and aSAH incidence as estimated in the model were similar to reported literature from Danish and Swedish cohort studies (Bor 2008 & Bromberg 1995) of 4.0% in first degree relatives and 20.7/100,000 per year in first degree relatives respectively. Mortality for other causes based on age-specific mortality rates in the general Dutch-population. **Quality-of-life weights:** Converted SF-36 scores to utilities using the method as described by Nichol 2001 or used available EQ5D scores. Assumed that the effect of screening on utility lasted one year and that utility changed each time screening was performed. **Cost sources:** Costs were derived from the Dutch manual for costing 2012, local data of Utrecht University Medical Centre and literature (Halkes 2006 & Roos 2002).

Comments

Source of funding: Grant of the Netherlands Organisation for Scientific Research (NWO). **Limitations:** Dutch resource use data (mix 1997-2003, 1996-97) and unit costs (2012) may not reflect current NHS context. Costs and health effects were discounted at a non-reference case rate (4%, 1.5% respectively). Non-NICE reference case utility measure used to estimate some utility values to calculate QALYs. Risk of aneurysm development and rupture were estimated from non-comparative cohort studies and assumptions made to estimate risk over time. Uncertainty not explored through sensitivity analysis. **Other:** None.

Overall applicability:^(d) Partially applicable **Overall quality:**^(e) Potentially serious limitations

- 1 *Abbreviations: CUA= cost-utility analysis; EQ-5D= Euroqol 5 dimensions (scale: 0.0 [death] to 1.0 [full health], negative values mean worse than death); ICER= incremental*
- 2 *cost-effectiveness ratio; n/a= not applicable; QALYs= quality-adjusted life year; SF-36=36-Item Short Form Health Survey*
- 3 *(a) Intervention number in order of least to most effective*
- 4 *(b) Full incremental analysis of available strategies: first strategies are ruled out that are dominated (another strategy is more effective and has lower costs) or subject to*
- 5 *extended dominance (the strategy is more effective and more costly but the incremental cost effectiveness ratio is higher than the next most effective option and so it*
- 6 *would never be the most cost effective option); incremental costs, incremental effects and incremental cost effectiveness ratios are calculated for the remaining strategies*
- 7 *by comparing each to the next most effective option.*
- 8 *(c) Converted using 2012 purchasing power parities⁵²*
- 9
- 10 *(d) Directly applicable / Partially applicable / Not applicable*
- 11 *(e) Minor limitations / Potentially serious limitations / Very serious limitations*
- 12

Study	Bor 2010																																																																							
Study details	Population & Interventions	Costs	Health outcomes	Cost effectiveness																																																																				
<p>Economic analysis: CUA (health outcome: QALYs)</p> <p>Study design: Decision analytic model</p> <p>Approach to analysis: Markov model with health states: healthy without aneurysm, healthy with aneurysm, healthy with known small aneurysm, disabled, and dead. 1 year cycles. Incidence of aneurysm and rupture risk constant throughout lifetime. If aneurysm suspected on MRA then DSA was undertaken to identify false positives. Detected aneurysms treated with either clipping or coiling unless too small (<3-5mm). 63%-67% considered too small for treatment.</p> <p>Perspective: Dutch healthcare provider</p> <p>Time horizon: Lifetime</p> <p>Treatment effect duration: n/a</p> <p>Discounting: Costs: 1.5%; Outcomes: 1.5%</p>	<p>Population: People with two or more first degree relatives with aneurysmal subarachnoid haemorrhage.</p> <p>Cohort settings: Start age: 15 Male: NR</p> <p>Interventions:</p> <ol style="list-style-type: none"> 1. No screening 2. Screening every 10 years, age 20 to 60 3. Screening every 15 years, age 20 to 60 4. Screening every 15 years, age 20 to 70 5. Screening every 2 years, age 20 to 80 6. Screening every 3 years, age 20 to 80 7. Screening every 5 years, age 20 to 80 8. Screening every 7 years, age 20 to 80 9. Screening every 10 years, age 20 to 80 10. Screening every 15 years, age 40 to 60 	<p>Full incremental analysis^{(a)(b)}</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Intv</th> <th>Cost</th> <th>QALY</th> <th>Incr cost</th> <th>Incr QALY</th> <th>ICER</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>£0</td> <td>0</td> <td colspan="3" style="text-align: center;">Baseline</td> </tr> <tr> <td>10</td> <td>£69</td> <td>0.060</td> <td>£69</td> <td>0.060</td> <td>£1,150</td> </tr> <tr> <td>3</td> <td>£143</td> <td>0.120</td> <td>£74</td> <td>0.060</td> <td>£1,233</td> </tr> <tr> <td>4</td> <td>£268</td> <td>0.150</td> <td>£125</td> <td>0.030</td> <td>£4,167</td> </tr> <tr> <td>2</td> <td>£370</td> <td>0.170</td> <td>£102</td> <td>0.020</td> <td>£5,100</td> </tr> <tr> <td>9</td> <td>£639</td> <td>0.210</td> <td>£269</td> <td>0.040</td> <td>£6,725</td> </tr> <tr> <td>8</td> <td>£955</td> <td>0.250</td> <td>£316</td> <td>0.040</td> <td>£7,900</td> </tr> <tr> <td>7</td> <td>£1,374</td> <td>0.280</td> <td>£419</td> <td>0.030</td> <td>£13,967</td> </tr> <tr> <td>6</td> <td>£2,257</td> <td>0.340</td> <td>£883</td> <td>0.060</td> <td>£14,717</td> </tr> <tr> <td>5</td> <td>£3,357</td> <td>0.410</td> <td>£1,100</td> <td>0.070</td> <td>£15,714</td> </tr> </tbody> </table> <p>Currency & cost year: 2007 Euros (presented here as 2007 UK pounds^(b))</p> <p>Cost components incorporated: Costs of screening - MRA and outpatient consultation with a neurologist and specialised nurse. Costs of UIA treatment - hospitalisation, follow-up in outpatient clinic and imaging. Cost associated with being disabled (in nursing home). Complications from recoiling not included.</p>				Intv	Cost	QALY	Incr cost	Incr QALY	ICER	1	£0	0	Baseline			10	£69	0.060	£69	0.060	£1,150	3	£143	0.120	£74	0.060	£1,233	4	£268	0.150	£125	0.030	£4,167	2	£370	0.170	£102	0.020	£5,100	9	£639	0.210	£269	0.040	£6,725	8	£955	0.250	£316	0.040	£7,900	7	£1,374	0.280	£419	0.030	£13,967	6	£2,257	0.340	£883	0.060	£14,717	5	£3,357	0.410	£1,100	0.070	£15,714	<p>All screening strategies cost effective when compared to no screening.</p> <p>Multiple other strategies (37 in total) were included in the analysis. Those reported are those that are most efficient (dominated or extendedly dominated strategies not reported here).</p> <p>The most cost effective screening strategy when compared to each other is 5 (screening every 2 years, age 20 to 80).</p>
Intv	Cost	QALY	Incr cost	Incr QALY	ICER																																																																			
1	£0	0	Baseline																																																																					
10	£69	0.060	£69	0.060	£1,150																																																																			
3	£143	0.120	£74	0.060	£1,233																																																																			
4	£268	0.150	£125	0.030	£4,167																																																																			
2	£370	0.170	£102	0.020	£5,100																																																																			
9	£639	0.210	£269	0.040	£6,725																																																																			
8	£955	0.250	£316	0.040	£7,900																																																																			
7	£1,374	0.280	£419	0.030	£13,967																																																																			
6	£2,257	0.340	£883	0.060	£14,717																																																																			
5	£3,357	0.410	£1,100	0.070	£15,714																																																																			

Analysis of uncertainty: To identify which input variables influence model outcomes the most an ANCOVA analysis was undertaken. Only suitable for models that are highly linear – in this model this only applied to the health effects. This analysis suggested that utility of individuals who were not offered screening and did not experience SAH and utility of individuals having a negative screening most influenced health effectiveness.

--	--	--	--

Data sources

Health outcomes: Due to limited evidence decided to use a probability of aneurysm development of 0.3%-0.7% per annum, and a probability of aneurysm rupture of 1.0 – 2.0% per annum. Small aneurysms were assigned a rupture rate of 0.5%-1.0% per annum, and an enlargement (>1mm) of 2.2%-4.9%. Small aneurysms were imaged every 2 years until they either enlarged or ruptured. Enlarged small aneurysms changed the next cycle to rupture rate of 1.0%-2.0%, and were always treated following the next imaging moment. With these probabilities approximately 27% of the study population would develop 1 or more aneurysms during their lifetime, and approximately 10% of the study population would develop SAH during their lifetime in the natural history (no screening) scenario. **Quality-of-life weights:** Not reported. **Cost sources:** College voor zorgverzekeringen kostenhandleiding 2004 (Dutch Health Care Insurance Board), Halkes 2006, Roos 2002.

Comments

Source of funding: Grant of the Netherlands Organisation for Scientific Research (NWO). **Limitations:** Dutch resource use data (mix 1997-2003, 1996-97) and unit costs (2007) may not reflect current NHS context. Costs and health effects were discounted at a non-reference case rate (1.5%). Unclear how estimates of utility values to calculate QALYs were determined. Risk of aneurysm development and rupture were estimated from non-comparative cohort studies and assumptions made to estimate risk over time. **Other:** None.

Overall applicability:^(c) Partially applicable **Overall quality:**^(d) Very serious limitations

1 Abbreviations: CUA= cost-utility analysis; EQ-5D= Euroqol 5 dimensions (scale: 0.0 [death] to 1.0 [full health], negative values mean worse than death); ICER= incremental
2 cost-effectiveness ratio; n/a= not applicable; QALYs= quality-adjusted life year; SF-36=36-Item Short Form Health Survey
3 (a) Converted using 2007 purchasing power parities⁵²
4 (b) Intervention number in order of least to most effective
5 (c) Full incremental analysis of available strategies: first strategies are ruled out that are dominated (another strategy is more effective and has lower costs) or subject to
6 extended dominance (the strategy is more effective and more costly but the incremental cost effectiveness ratio is higher than the next most effective option and so it
7 would never be the most cost effective option); incremental costs, incremental effects and incremental cost effectiveness ratios are calculated for the remaining strategies

- 1 *by comparing each to the next most effective option. Due to the large volume of strategies, interventions that were subject to dominance or extended dominance were not*
- 2 *reported here.*
- 3 *(d) Directly applicable / Partially applicable / Not applicable*
- 4 *(e) Minor limitations / Potentially serious limitations / Very serious limitations*

5

6

1 Appendix I: Excluded studies

I.1.2 Excluded clinical studies

3 Table 8: Studies excluded from the clinical review

Reference	Reason for exclusion
Bhat 2011 ¹	Incorrect intervention – no assessments to detect SAH in families
Bhat 2012 ²	Incorrect intervention – no assessments to detect SAH in families
Bor 2010 ³	Incorrect study design – economics Markov model
Bor 2008 ⁴	Incorrect intervention – no assessments to detect SAH in families
Bor 2014 ⁵	Incorrect study design – no comparison group for people with familial risk of SAH
Bossuyt 2005 ⁶	No relevant outcomes identified
Broderick 2009 ⁷	Incorrect study design – no comparison group for people with familial risk of SAH
Broderick 2005 ⁸	Incorrect study design – study protocol only
Bromberg 1995 ⁹	Incorrect intervention – no assessments to detect SAH in families
Bromberg 1995 ¹⁰	Incorrect intervention – no assessments to detect SAH in families
Brown 1999 ¹¹	Incorrect study design – no comparison group for people with familial risk of SAH
Brown 2014 ¹²	Incorrect study design – review / editorial (references checked)
Chalouhi 2011 ¹³	Incorrect study design – Systematic review (references checked)
Chan 2016 ¹⁴	Incorrect study design – no comparison group for people with familial risk of SAH
Coppola 1973 ¹⁵	No relevant outcomes identified
Crawley 1999 ¹⁶	Incorrect study design – modelling of potential investigation results
Dippel 1992 ¹⁷	Incorrect study design – economics paper
Edelsohn 1972 ¹⁸	No relevant outcomes identified
Flahault 2016 ¹⁹	Incorrect study design – opinions based on e-questionnaire
Flahault 2018 ²⁰	Incorrect study design – no comparison group for people with familial risk of SAH
Fox 1980 ²¹	No relevant outcomes identified
Greebe 1997 ²²	Incorrect intervention – no assessments to detect SAH in families
Hashimoto 1977 ²³	No relevant outcomes identified
Huston 1996 ²⁵	Incorrect intervention – investigation and follow up of intracranial haemorrhage for people with ADPKD
Jacoby 2006 ²⁶	Incorrect study design – review / editorial (references checked)
Jain 1974 ²⁷	No relevant outcomes identified

Reference	Reason for exclusion
Kasuya 2000 ²⁸	Incorrect study design – no comparison group for people with familial risk of SAH
Kelly 2014 ²⁹	Incorrect study design – Systematic review (references checked)
Kim 2003 ³¹	Incorrect study design – no comparison group for people with familial risk of SAH
Kim 2005 ³⁰	Incorrect study design – retrospective investigation of SAH in families (no screening)
Kojima 1998 ³²	Incorrect study design – no comparison group for people with familial risk of SAH
Koshy 2010 ³³	Incorrect study design – no comparison group for people with familial risk of SAH
Kubota 2001 ³⁴	Incorrect intervention – prevalence of risk factors for SAH
Leblanc 1997 ³⁵	Incorrect study design – Literature review (references checked)
Leblanc 1995 ³⁶	Incorrect study design – mixed methodology of screening (DSA/MRA); no comparison group for people with familial risk of SAH
Lee 2008 ³⁷	Incorrect study design – no comparison group for people with familial risk of SAH
Lin 2005 ³⁸	Incorrect study design – abstracts / case reports only
Lindgaard 2003 ³⁹	Incorrect study design – retrospective investigation of SAH in families (no screening)
Lorenzo-Betancor 2018 ⁴⁰	Incorrect intervention – gene mutation search study
Lozano 1987 ⁴¹	No relevant outcomes identified
Magnetic Resonance Angiography in Relatives of Patients with Subarachnoid Hemorrhage Study Group 1999 ⁴²	Incorrect study design – no comparison group for people with familial risk of SAH
Mensing 2017 ⁴³	Incorrect study design – abstract only
Mensing 2017 ⁴⁴	Incorrect study design – citation only
Mensing 2019 ⁴⁵	Incorrect study design – does not match protocol. Investigation of incidence of unruptured intracranial aneurysms compared to sporadic incidence of unruptured intracranial aneurysms (both groups have screening).
Mensing 2016 ⁴⁶	Incorrect intervention - comparison of shape between familial and sporadic
Miller 2012 ⁴⁷	Incorrect study design – incidence of SAH with specific risk factors; no comparison of familial screening compared to no screening
Nagae 1972 ⁴⁸	No relevant outcomes identified
Okamoto 2003 ⁵¹	Incorrect intervention – no assessments to detect SAH in families; data gained through questionnaire
Pei Shan 1995 ⁵³	Incorrect intervention – no assessments to detect SAH in families
Phillips 1963 ⁵⁴	No relevant outcomes identified
Raaymakers 1999 ⁵⁵	Incorrect study design – no comparison group for people with familial risk of SAH
Raaymakers 1998 ⁵⁶	Incorrect study design – no comparison group for people with familial risk of SAH

Reference	Reason for exclusion
Raaymakers 2000 ⁵⁷	Incorrect study design – no comparison group for people with familial risk of SAH
Rinkel 2005 ⁵⁸	Incorrect study design – review / editorial (references checked)
Rinkel 1998 ⁵⁹	Incorrect study design – Systematic review on incidence of ICA rupture
Roberts 1999 ⁶⁰	Incorrect intervention – no assessments to detect SAH in families; data gained through questionnaire
Ronkainen 1997 ⁶¹	Incorrect study design – no comparison group for people with familial risk of SAH
Ronkainen 1994 ⁶²	Incorrect study design – no comparison group for people with familial risk of SAH; unclear cohorts of familial SAH
Ronkainen 1995 ⁶³	Incorrect study design – special features of familial intracranial aneurysms
Ronkainen 1998 ⁶⁴	Incorrect intervention – incidence of familial SAH via autopsy
Ronkainen 1999 ⁶⁵	Incorrect study design – comparison between familial and sporadic SAH screening; no correct comparison group for people with familial risk of SAH
Ronkainen 1995 ⁶⁶	Incorrect study design – no comparison group for people with familial risk of SAH
Ronkainen 1998 ⁶⁷	Incorrect study design – Systematic review (references checked)
Ruigrok 2004 ⁶⁸	Incorrect study design – no comparison group for people with familial risk of SAH
Sakai 1974 ⁶⁹	No relevant outcomes identified
Schievink 1994 ⁷⁰	Incorrect study design – no comparison group for people with familial risk of SAH
Sundquist 2007 ⁷¹	Incorrect intervention – no assessments to detect SAH in families
Takao 2008 ⁷²	Incorrect study design – Economical Markov model
Teasdale 2005 ⁷³	Incorrect study design – no comparison group for people with familial risk of SAH
ter Berg 1986 ⁷⁴	Incorrect intervention – no assessments to detect SAH in families
ter Berg 1988 ⁷⁵	Incorrect intervention – screening for familial intracranial aneurysms via DSA (exclusion criteria)
ter Berg 1992 ⁷⁶	Incorrect study design – Literature review (references checked)
Toglia 1972 ⁷⁷	No relevant outcomes identified
van der Voet 2004 ⁷⁸	Incorrect study design – investigating genetic causes of familial SAH
Verdure 2015 ⁷⁹	Incorrect study design – investigating genetic causes of familial SAH
Vlak 2013 ⁸⁰	Incorrect study design – investigating risk factors for aneurysm rupture
Wermer 2003 ⁸¹	Incorrect study design – no comparison group for people with familial risk of SAH
White 2004 ⁸²	Incorrect study design – Literature review (references checked)
Woo 2009 ⁸³	Incorrect intervention – no assessments to detect SAH in families

Reference	Reason for exclusion
Zuurbier 2019 ⁸⁴	Incorrect study design – citation only
Zuurbier 2019 ⁸⁵	Incorrect study design – abstract only

I.2.1 Excluded health economic studies

2 Published health economic studies that met the inclusion criteria (relevant population,
3 comparators, economic study design, published 2003 or later and not from non-OECD
4 country or USA) but that were excluded following appraisal of applicability and
5 methodological quality are listed below. See the health economic protocol for more details.

6 **Table 9: Studies excluded from the health economic review**

Reference	Reason for exclusion
None.	

7