

# Osteoarthritis: assessment and management (update)

NICE guideline: methods

*NICE guideline <number>*

*Methods*

*April 2022*

*Draft for Consultation*



## **Disclaimer**

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

## **Copyright**

© NICE, 2022. All rights reserved. Subject to Notice of rights.

# Contents

<b>1</b>	<b>Development of the guideline</b> .....	<b>5</b>
1.1	Remit.....	5
1.2	What this guideline covers.....	<b>Error! Bookmark not defined.</b>
1.3	What this guideline does not cover .....	<b>Error! Bookmark not defined.</b>
<b>2</b>	<b>Methods</b> .....	<b>6</b>
2.1	Developing the review questions and outcomes .....	6
2.1.1	Stratification .....	9
2.2	Searching for evidence.....	9
2.2.1	Clinical and health economics literature searches .....	9
2.3	Reviewing evidence .....	10
2.3.1	Types of studies and inclusion and exclusion criteria .....	11
2.4	Methods of combining evidence .....	12
2.4.1	Data synthesis for intervention reviews .....	12
2.4.3	Data synthesis for prognostic risk factor reviews.....	13
2.4.4	Data synthesis for qualitative reviews .....	13
2.5	Appraising the quality of evidence by outcomes .....	13
2.5.1	Intervention reviews .....	13
2.5.3	Prognostic reviews.....	20
2.5.5	Qualitative reviews.....	21
2.6	Assessing clinical importance.....	25
2.7	Identifying and analysing evidence of cost effectiveness .....	25
2.7.1	Literature review .....	26
2.7.2	Undertaking new health economic analysis.....	28
2.7.3	Cost-effectiveness criteria .....	28
2.7.4	In the absence of health economic evidence.....	29
2.8	Developing recommendations .....	29
2.8.1	Research recommendations .....	30
2.8.2	Validation process.....	31
2.8.3	Updating the guideline .....	31
2.8.4	Disclaimer .....	31
2.8.5	Funding.....	31
2.9	General terms .....	31
	References.....	42

# 1 Development of the guideline

## 2 1.1 Remit

3 NICE received the remit for this guideline from NHS England. NICE commissioned  
4 the National Guideline Centre to produce the guideline.

5 The remit for this guideline is:

6 Osteoarthritis: care and management (update).

7 To see “What this guideline covers” and “What this guideline does not cover” please  
8 see the Osteoarthritis: assessment and management scope.

## 2 Methods

This guideline was developed using the methods described in the 2014 NICE guidelines manual, updated 2018.

Declarations of interest were recorded according to the NICE conflicts of interest policy.

Sections 2.1 to 2.3 describe the process used to identify and review evidence. Sections 2.1.1 and 2.7 describe the process used to identify and review the health economic evidence.

### 2.1 Developing the review questions and outcomes

The review questions developed for this guideline were based on the key areas and draft review questions identified in the guideline scope. They were drafted by the National Guideline Centre technical team and refined and validated by the committee and signed off by NICE. A total of 16 review questions were developed in this guideline and outlined in Table 1.

The review questions were based on the following frameworks:

- population, intervention, comparator and outcome (PICO) for reviews of interventions
- population, exposure and outcomes for prognostic reviews
- population, setting and context for qualitative reviews.

This use of a framework informed a more detailed protocol that guided the literature searching process, critical appraisal and synthesis of evidence, and facilitated the development of recommendations by the guideline committee. Full literature searches, critical appraisals and evidence reviews were completed for all the specified review questions.

**Table 1: Review questions**

Evidence report	Type of review	Review questions	Outcomes
1.1	Intervention	What is the additional benefit of imaging in the diagnosis of osteoarthritis in people with suspected osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Psychological distress</li> <li>• Healthcare utilisation (prescribing, investigations, hospitalisation or health professional visit)</li> <li>• Any alternative diagnosis</li> </ul>
2.1	Qualitative	What information on osteoarthritis, including the management of flare-ups, do people with osteoarthritis, their family and carers need during and after diagnosis?	N/A
3.1	Intervention	What is the clinical and cost-effectiveness of exercise therapy	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> </ul>

Evidence report	Type of review	Review questions	Outcomes
		for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Serious adverse events</li> </ul>
3.2	Prognostic	What is the benefit of weight loss for the management of osteoarthritis in overweight and obese people?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> </ul>
3.3	Intervention	What is the clinical and cost-effectiveness of manual therapy for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Minor adverse events</li> <li>Moderate/major adverse events</li> </ul>
3.4	Intervention	What is the clinical and cost-effectiveness of acupuncture for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Serious adverse events</li> </ul>
3.5	Intervention	What is the clinical and cost-effectiveness of electrotherapy for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Minor adverse events</li> <li>Moderate/major adverse events</li> </ul>
3.6	Intervention	What is the clinical and cost-effectiveness of devices (such as supports, splints and braces) for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Number of adverse events</li> </ul>
4.1	Intervention	What is the clinical and cost-effectiveness of oral, topical and transdermal medicines for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>Health-related quality of life</li> <li>Pain</li> <li>Physical function</li> <li>Psychological distress</li> <li>Osteoarthritis flares</li> <li>Serious adverse events 1A: Gastrointestinal (bleeding or perforation) adverse events</li> </ul>

Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> <li>• Serious adverse events 1B: Gastrointestinal (non-bleeding or perforation) adverse events</li> <li>• Serious adverse events 2: Cardiovascular adverse events</li> <li>• Serious adverse events 3: Hepatorenal adverse events</li> <li>• Serious adverse events 4: Central nervous system adverse events</li> </ul>
4.2	Intervention	What is the clinical and cost-effectiveness of intra-articular injections for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Psychological distress</li> <li>• Osteoarthritis flares</li> <li>• Serious adverse events</li> </ul>
5.1	Intervention	What is the clinical and cost-effectiveness of treatment packages (that include combinations of interventions) for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Psychological distress</li> <li>• Osteoarthritis flares</li> <li>• Discontinuation</li> </ul>
6.1	Intervention	Is regular follow-up and review needed for people with osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Psychological distress</li> <li>• Osteoarthritis flares</li> <li>• Residential service or hospital admission (including disability allowance)</li> <li>• Progression to joint replacement</li> </ul>
6.2	Intervention	What is the clinical and cost-effectiveness of using radiological investigations (for example: x-ray, ultrasound, MRI) to inform management choices for people with osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Changes to planned management</li> <li>• Psychological distress</li> <li>• Osteoarthritis flares</li> <li>• Number of adverse events</li> </ul>
7.1	Intervention	What is the clinical and cost-effectiveness of arthroscopic procedures for the management of osteoarthritis?	<ul style="list-style-type: none"> <li>• Health-related quality of life</li> <li>• Pain</li> <li>• Physical function</li> <li>• Progression to joint replacement</li> </ul>



Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> <li>• Psychological distress</li> <li>• Osteoarthritis flares</li> <li>• Serious adverse events</li> </ul>
8.1	Prognostic	What are the indicators for possible joint replacement surgery?	<ul style="list-style-type: none"> <li>• Progression to joint replacement</li> </ul>
8.2	Prognostic	Do people with osteoarthritis who are at less than or more than healthy weight have better outcomes after joint replacement surgery than people of healthy weight?	<ul style="list-style-type: none"> <li>• Mortality</li> <li>• Health-related quality of life</li> <li>• Post-operative patient-reported outcome measure</li> <li>• Total adverse events</li> <li>• Surgical site infection</li> <li>• Venous thromboembolic events</li> </ul>

1

## 2 2.1.1 Stratification

3 The following questions included specific stratifications:

- 4 • In questions 1.1 3.2, 3.6, 4.1 (topical [local] medicines only) and 4.2, analyses  
5 were stratified by joint site affected by osteoarthritis (hip, knee, ankle, foot, toe,  
6 shoulder, elbow, wrist, hand, thumb, finger, temporomandibular joint and  
7 multisite). In questions 8.1 and 8.2 this stratification was also used, but limited to  
8 the hip, knee and shoulder.
- 9 • In question 6.1, analyses were stratified by groups who may respond more to  
10 regular follow up (people starting a new pharmacological intervention, people  
11 starting a new non-pharmacological intervention and long-term condition  
12 management).

13 Where studies reported a mix of populations across strata, a threshold of 80% was  
14 agreed with the committee as a cut off for what would be acceptable to constitute  
15 a predominant group.

## 16 2.2 Searching for evidence

### 17 2.2.1 Clinical and health economics literature searches

18 Systematic literature searches were undertaken to identify all published clinical and  
19 health economic evidence relevant to the review questions. Searches were  
20 undertaken according to the parameters stipulated within the NICE guidelines  
21 manual 2014, updated 2020<sup>4</sup> (see <https://www.nice.org.uk/process/pmg20>).  
22 Databases were searched using relevant medical subject headings, free-text terms  
23 and study-type filters where appropriate. Searches were restricted to papers  
24 published in English. Studies published in languages other than English were not  
25 reviewed. All searches were updated on 17 November 2021. Papers published or  
26 added to databases after this date were not considered. Where new evidence was  
27 identified, for example in consultation comments received from stakeholders, the  
28 impact on the guideline was considered, and the action agreed between NGC and  
29 NICE staff with a quality assurance role.

1 Search strategies were quality assured by the following approaches. Medline search  
2 strategies were checked by a second information specialist following the PRESS  
3 checklist.<sup>3</sup> Searches were cross-checked with reference lists of relevant papers,  
4 searches in other systematic reviews were analysed, and committee members were  
5 requested to highlight key studies.

6 Searching for unpublished literature was not undertaken. The NGC and NICE do not  
7 have access to drug manufacturers' unpublished clinical trial results, so the clinical  
8 evidence considered by the committee for pharmaceutical interventions may be  
9 different from that considered by the MHRA and European Medicines Agency for the  
10 purposes of licensing and safety regulation.

11 The search strategies can be found as an appendix to each evidence review.

12 During the scoping stage, a search was conducted for systematic reviews in the  
13 following databases:

- 14 • Medline (Ovid)
- 15 • Cochrane Database of Systematic Reviews

## 16 **2.3 Reviewing evidence**

17 The evidence for each review question was reviewed using the following process:

- 18 • Potentially relevant studies were identified from the search results by reviewing  
19 titles and abstracts. The full papers were then obtained.
- 20 • Full papers were evaluated against the pre-specified inclusion and exclusion  
21 criteria set out in the protocol to identify studies that addressed the review  
22 question. The review protocols are included in an appendix to each of the  
23 evidence reports.
- 24 • Relevant studies were critically appraised using the preferred study design  
25 checklist as specified in the NICE guidelines manual.<sup>4</sup> The checklist used is  
26 included in the individual review protocols in each of the evidence reports.
- 27 • Key information was extracted about interventional study methods and results into  
28 'EviBase', NGC's purpose-built software. Summary evidence tables were  
29 produced from data entered into EviBase, including critical appraisal ratings. Key  
30 information about non-interventional study methods and results were manually  
31 extracted into standard Word evidence tables (evidence tables are included in an  
32 appendix to each of the evidence reports).
- 33 • Summaries of the evidence were generated by outcome. Outcome data were  
34 combined, analysed and reported according to study design:
  - 35 ○ Randomised data were meta-analysed where appropriate and reported in  
36 GRADE profile tables.
  - 37 ○ Prognostic data were meta-analysed where appropriate and reported in  
38 adapted GRADE profile tables.
  - 39 ○ Qualitative data were synthesised across studies using thematic analysis and  
40 presented as summary statements in GRADE CERQual tables.
- 41 • A minimum of 10% of the abstracts were reviewed by two reviewers, with any  
42 disagreements resolved by discussion or, if necessary, a third independent  
43 reviewer.
- 44 • All of the evidence reviews were quality assured by a senior systematic reviewer.  
45 This included checking:

- 1 ○ papers were included or excluded appropriately
  - 2 ○ a sample of the data extractions
  - 3 ○ a sample of the risk of bias assessments
  - 4 ○ correct methods were used to synthesise data.
- 5 Discrepancies will be identified and resolved through discussion (with a third  
6 reviewer where necessary).

### 7 **2.3.1 Types of studies and inclusion and exclusion criteria**

8 The inclusion and exclusion of studies was based on the criteria defined in the review  
9 protocols, which can be found in an appendix to each of the evidence reports.  
10 Excluded studies (with the reasons for their exclusion) are listed in an appendix to  
11 each of the evidence reports. The committee was consulted about any uncertainty  
12 regarding inclusion or exclusion.

13 Conference abstracts were not generally considered for inclusion. Literature reviews,  
14 posters, letters, editorials, comment articles, unpublished studies and studies not in  
15 published in English language were excluded.

#### 16 **2.3.1.1 Type of studies**

17 Randomised trials and other observational studies (including prognostic studies)  
18 were included in the evidence reviews as appropriate.

19 For intervention reviews, randomised controlled trials (RCTs) were included where  
20 identified as because they are considered the most robust type of study design that  
21 can produce an unbiased estimate of the intervention effects. Refer to the review  
22 protocols in each evidence report for full details on the study design of studies that  
23 were appropriate for each review question.

24 For prognostic review questions, prospective and retrospective cohort studies were  
25 included. Case–control studies were not included.

26 Systematic reviews and meta-analyses conducted to the same methodological  
27 standards as the NICE reviews were included within the evidence reviews in  
28 preference to primary studies, where they were available and applicable to the review  
29 questions and updated or added to where appropriate to the guideline review  
30 question. Individual patient data (IPD) meta-analyses were preferentially included if  
31 meeting the protocol and methodological criteria.

#### 32 **2.3.1.1.1 Qualitative studies**

33 In the qualitative reviews, studies using focus groups, or structured or semi-  
34 structured interviews were considered for inclusion. Survey data or other types of  
35 questionnaires were only included if they provided analysis from open-ended  
36 questions, but not if they reported descriptive quantitative data only.

#### 37 **Saturation of qualitative studies**

38 Data extraction in qualitative reviews is a thorough process. A common approach  
39 applied in systematic reviews of qualitative data is to stop extracting data once  
40 saturation has been reached. In an exploratory review, where themes are not  
41 predefined in the protocol, thematic or data extraction may be applied. For the

1 purposes of this review, extraction of information from relevant studies was stopped  
2 when data saturation was reached, i.e. no new information was emerging for a  
3 specific theme. This includes; studies that don't report any new themes additional to  
4 those already identified in the review as well as not contributing additional information  
5 to the existing themes, as well as studies which report a new theme but data from  
6 other themes in the study doesn't contribute to the existing review themes. In the  
7 latter scenario only the new theme data is extracted. These studies are not  
8 specifically excluded from the review as they nevertheless fit the criteria defined in  
9 the review protocol. Any studies for which data were not extracted due to data  
10 saturation having been reached, but that fit the inclusion criteria of the protocol, were  
11 listed in the table for studies 'identified but not extracted due to saturation'

## 12 **2.4 Methods of combining evidence**

### 13 **2.4.1 Data synthesis for intervention reviews**

14 Meta-analyses were conducted using Cochrane Review Manager (RevMan5)<sup>9</sup>  
15 software

#### 16 **2.4.1.1 Analysis of different types of data**

##### 17 ***Dichotomous outcomes***

18 Fixed-effects (Mantel–Haenszel) techniques were used to calculate risk ratios  
19 (relative risk, RR) for the binary outcomes. The absolute risk difference was also  
20 calculated using GRADEpro<sup>1</sup> software, using the median event rate in the control arm  
21 of the pooled results.

22 For binary variables where there were zero events in either arm or a less than 1%  
23 event rate, Peto odds ratios, rather than risk ratios, were calculated as they are more  
24 appropriate for data with a low number of events. Where there are zero events in  
25 both arms, the risk difference was calculated and reported instead.

##### 26 **Continuous outcomes**

27 Continuous outcomes were analysed using an inverse variance method for pooling  
28 weighted mean differences.

29 Where the studies within a single meta-analysis had different scales of measurement  
30 for the same outcomes, standardised mean differences were used (providing all  
31 studies reported either change from baseline or final values rather than a mixture of  
32 both); each different measure in each study was 'normalised' to the standard  
33 deviation value pooled between the intervention and comparator groups in that same  
34 study.

35 The means and standard deviations of continuous outcomes are required for meta-  
36 analysis. However, in cases where standard deviations were not reported, the  
37 standard error was calculated if the p values or 95% confidence intervals (95% CI)  
38 were reported, and meta-analysis was undertaken with the mean and standard error  
39 using the generic inverse variance method in Cochrane Review Manager  
40 (RevMan5)<sup>9</sup> software.

## 1 **Generic inverse variance**

2 If a study reported only the summary statistic and 95% CI the generic-inverse  
3 variance method was used to enter data into RevMan5.<sup>9</sup> If the control event rate was  
4 reported this was used to generate the absolute risk difference in GRADEpro.<sup>1</sup> If  
5 multivariate analysis was used to derive the summary statistic but no adjusted control  
6 event rate was reported no absolute risk difference was calculated.

## 7 **Complex analysis**

8 Where studies had used a crossover design, paired continuous data were extracted  
9 where possible, and forest plots were generated in RevMan5<sup>9</sup> with the generic  
10 inverse variance function. Forest plots were also generated in RevMan5<sup>9</sup> with the  
11 generic inverse variance function. If paired continuous or categorical data were not  
12 available from the crossover studies, the separate group data were analysed in the  
13 same way as data from parallel groups, on the basis that this approach would  
14 overestimate the confidence intervals and thus artificially reduce study weighting  
15 resulting in a conservative effect. Where a meta-analysis included a mixture of  
16 studies using both paired and parallel group approaches, all data were entered into  
17 RevMan5<sup>9</sup> using the generic inverse variance function.

## 18 **2.4.2 Data synthesis for prognostic risk factor reviews**

19 Adjusted odds ratios and mean differences, risk ratios, or hazard ratios, with their  
20 95% CIs, for the effect of the pre-specified prognostic factors were extracted from the  
21 studies. Studies were only included if the confounders pre-specified by the committee  
22 were either matched at baseline or were adjusted for in multivariate analysis.  
23 Prospective cohort studies reporting multivariable analyses that adjusted for key  
24 confounders identified by the committee at the protocol stage for that outcome were  
25 the preferred study design.

26 Data were not combined in meta-analyses for prognostic studies because they were  
27 not adjusted for the same confounders and were not similarly homogenous to pool.

## 28 **2.4.3 Data synthesis for qualitative reviews**

29 The main findings for each included paper were identified and thematic analysis  
30 methods were used to synthesise this information into broad overarching themes  
31 which were summarised into the main review findings. The evidence was presented  
32 in the form of a narrative summary detailing the evidence from the relevant papers  
33 and how this informed the overall review finding plus a statement on the level of  
34 confidence for that review finding. Considerable limitations and issues around  
35 relevance were listed. A summary evidence table with the succinct summary  
36 statements for each review finding was produced including the associated quality  
37 assessment.

## 38 **2.5 Appraising the quality of evidence by outcomes**

### 39 **2.5.1 Intervention reviews**

40 The evidence for outcomes from the included RCTs and, where appropriate, non-  
41 randomised intervention studies, were evaluated and presented using the 'Grading of  
42 Recommendations Assessment, Development and Evaluation (GRADE) toolbox'

1 developed by the international GRADE working group  
 2 (<http://www.gradeworkinggroup.org/>). The software (GRADEpro<sup>1</sup>) developed by the  
 3 GRADE working group was used to assess the quality of each outcome, taking into  
 4 account individual study quality and the meta-analysis results.

5 Each outcome was first examined for each of the quality elements listed and defined  
 6 in Table 2.

7 **Table 2: Description of quality elements in GRADE for intervention studies**

Quality element	Description
Risk of bias	Limitations in the study design and implementation may bias the estimates of the treatment effect. Major limitations in studies decrease the confidence in the estimate of the effect. Examples of such limitations are selection bias (often due to poor allocation concealment), performance and detection bias (often due to a lack of blinding of the patient, healthcare professional or assessor) and attrition bias (due to missing data causing systematic bias in the analysis).
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question.
Inconsistency	Inconsistency refers to an unexplained heterogeneity of effect estimates between studies in the same meta-analysis.
Imprecision	Results are imprecise when studies include relatively few patients and few events (or highly variable measures) and thus have wide confidence intervals around the estimate of the effect relative to clinically important thresholds. 95% confidence intervals denote the possible range of locations of the true population effect at a 95% probability, and so wide confidence intervals may denote a result that is consistent with conflicting interpretations (for example a result may be consistent with both clinical benefit AND clinical harm) and thus be imprecise.
Publication bias	Publication bias is a systematic underestimate or overestimate of the underlying beneficial or harmful effect due to the selective publication of studies. A closely related phenomenon is where some papers fail to report an outcome that is inconclusive, thus leading to an overestimate of the effectiveness of that outcome.
Other issues	Sometimes randomisation may not adequately lead to group equivalence of confounders, and if so this may lead to bias, which should be taken into account. Potential conflicts of interest, often caused by excessive pharmaceutical company involvement in the publication of a study, should also be noted.

8 Details of how the 4 main quality elements (risk of bias, indirectness, inconsistency  
 9 and imprecision) were appraised for each outcome are given below. Publication bias  
 10 was considered with the committee. There was no reason to suspect it was present  
 11 from the evidence available, so it was not explored with funnel plots.

### 12 2.5.1.1 Risk of bias

13 The main domains of bias for RCTs are listed in Table 3. Each outcome had its risk  
 14 of bias assessed within each study first using the appropriate checklist for the study  
 15 design (Cochrane RoB 2 for RCTs, or ROBINS-I for non-randomised studies or  
 16 ROBIS for systematic reviews). For each study, if there was no risk of bias in any  
 17 domain, the risk of bias was given a rating of 0; 'no serious risk of bias'. If there was  
 18 risk of bias in just 1 domain, the risk of bias was given a 'serious' rating of -1, but if  
 19 there was risk of bias in 2 or more domains the risk of bias was given a 'very serious'  
 20 rating of -2. An overall rating is calculated across all studies by taking into account  
 21 the weighting of studies according to study precision. For example, if the most

precise studies tended to each have a score of  $-1$  for that outcome, the overall score for that outcome would tend towards  $-1$ .

**Table 3: Principle domains of bias in randomised controlled trials**

Limitation	Explanation
Selection bias (sequence generation and allocation concealment)	If those enrolling participants are aware of the group to which the next enrolled patient will be allocated, either because of a non-random sequence that is predictable, or because a truly random sequence was not concealed from the researcher, this may translate into systematic selection bias. This may occur if the researcher chooses not to recruit a participant into that specific group because of: <ul style="list-style-type: none"> <li>• knowledge of that participant's likely prognostic characteristics, and</li> <li>• a desire for one group to do better than the other.</li> </ul>
Performance and detection bias (lack of blinding)	Patients, caregivers, those adjudicating or recording outcomes, and data analysts should not be aware of the arm to which the participants are allocated. Knowledge of the group can influence: <ul style="list-style-type: none"> <li>• the experience of the placebo effect</li> <li>• performance in outcome measures</li> <li>• the level of care and attention received, and</li> <li>• the methods of measurement or analysis</li> </ul> all of which can contribute to systematic bias.
Attrition bias	Attrition bias results from an unaccounted for loss of data beyond a certain level (a differential of at least 10% between groups). Loss of data can occur when participants are compulsorily withdrawn from a group by the researchers (for example, when a per-protocol approach is used) or when participants do not attend assessment sessions. If the missing data are likely to be different from the data of those remaining in the groups, and there is a differential rate of such missing data from groups, systematic attrition bias may result.
Selective outcome reporting	Reporting of some outcomes and not others on the basis of the results can also lead to bias, as this may distort the overall impression of efficacy.
Other limitations	For example: <ul style="list-style-type: none"> <li>• Stopping early for benefit observed in randomised trials, in particular in the absence of adequate stopping rules.</li> <li>• Use of unvalidated patient-reported outcome measures.</li> <li>• Lack of washout periods to avoid carry-over effects in crossover trials.</li> <li>• Recruitment bias in cluster-randomised trials.</li> </ul>

#### 4 2.5.1.2 Indirectness

Indirectness refers to the extent to which the populations, interventions, comparisons and outcome measures are dissimilar to those defined in the inclusion criteria for the reviews. Indirectness is important when these differences are expected to contribute to a difference in effect size, or may affect the balance of harms and benefits considered for an intervention. As for the risk of bias, each outcome had its indirectness assessed within each study first. For each study, if there were no sources of indirectness, indirectness was given a rating of 0. If there was indirectness in just 1 source (for example in terms of population), indirectness was given a 'serious' rating of  $-1$ , but if there was indirectness in 2 or more sources (for example, in terms of population and treatment) the indirectness was given a 'very serious' rating of  $-2$ . An overall rating is calculated across all studies by taking into account the weighting of studies according to study precision. For example, if the most

1 precise studies tended to have an indirectness score of -1 each for that outcome, the  
2 overall score for that outcome would tend towards -1.

### 3 **2.5.1.3 Inconsistency**

4 Inconsistency refers to an unexplained heterogeneity of results for an outcome  
5 across different studies. When estimates of the treatment effect across studies differ  
6 widely, this suggests true differences in the underlying treatment effect, which may  
7 be due to differences in populations, settings or doses. Statistical heterogeneity was  
8 assessed for each meta-analysis estimate by an I-squared ( $I^2$ ) inconsistency statistic.

9 Heterogeneity or inconsistency amongst studies was also visually inspected. Where  
10 statistical heterogeneity as defined above was present or there was clear visual  
11 heterogeneity not captured in the  $I^2$  value predefined subgrouping of studies was  
12 carried out according to the protocol. See the review protocols for the subgrouping  
13 strategy.

14 When heterogeneity existed within an outcome ( $I^2 > 50\%$ ), but no plausible  
15 explanation could be found, the quality of evidence for that outcome was  
16 downgraded. Inconsistency for that outcome was given a 'serious' score of -1 if the  $I^2$   
17 was 50–74%, and a 'very serious' score of -2 if the  $I^2$  was 75% or more. If a visual  
18 assessment indicated heterogeneity but the  $I^2$  was under 50% then the analysis was  
19 considered on a case by case basis. Where it appeared that imprecision was  
20 masking possible heterogeneity (leading to a reduced  $I^2$  value), then the outcome  
21 was downgraded for inconsistency.

22 If inconsistency could be explained based on pre-specified subgroup analysis (that is,  
23 each subgroup had an  $I^2 < 50\%$ ) then each of the derived subgroups were presented  
24 separately for that forest plot (providing at least 2 studies remained in each  
25 subgroup). The committee took this into account and considered whether to make  
26 separate recommendations based on the variation in effect across subgroups within  
27 the same outcome. In such a situation the quality of evidence was not downgraded.

28 If all predefined strategies of subgrouping were unable to explain statistical  
29 heterogeneity, then a random effects (DerSimonian and Laird) model was employed  
30 to the entire group of studies in the meta-analysis. A random-effects model assumes  
31 a distribution of populations, rather than a single population. This leads to a widening  
32 of the confidence interval around the overall estimate. If, however, the committee  
33 considered the heterogeneity was so large that meta-analysis was inappropriate,  
34 then the results were not pooled and were described narratively.

### 35 **2.5.1.4 Imprecision**

36 The criteria applied for imprecision were based on the 95% CIs for the pooled  
37 estimate of effect, and the minimal important differences (MID) for the outcome. The  
38 MIDs are the threshold for appreciable benefits and harms, separated by a zone  
39 either side of the line of no effect where there is assumed to be no clinically important  
40 effect. If either end of the 95% CI of the overall estimate of effect crossed 1 of the  
41 MID lines, imprecision was regarded as serious and a 'serious' score of -1 was  
42 given. This was because the overall result, as represented by the span of the  
43 confidence interval, was consistent with 2 interpretations as defined by the MID (for  
44 example, both no clinically important effect and clinical benefit were possible  
45 interpretations). If both MID lines were crossed by either or both ends of the 95% CI  
46 then imprecision was regarded as very serious and a 'very serious' score of -2 was



1 given. This was because the overall result was consistent with all 3 interpretations  
2 defined by the MID (no clinically important effect, clinical benefit and clinical harm).  
3 This is illustrated in Figure 1.

4 The value / position of the MID lines is ideally determined by values reported in the  
5 literature. 'Anchor-based' methods aim to establish clinically meaningful changes in a  
6 continuous outcome variable by relating or 'anchoring' them to patient-centred  
7 measures of clinical effectiveness that could be regarded as gold standards with a  
8 high level of face validity. For example, a MID for an outcome could be defined by the  
9 minimum amount of change in that outcome necessary to make patients feel their  
10 quality of life had 'significantly improved'. MIDs in the literature may also be based on  
11 expert clinician or consensus opinion concerning the minimum amount of change in a  
12 variable deemed to affect quality of life or health.

13 In the absence of values identified in the literature, the alternative approach to  
14 deciding on MID levels is to use the modified GRADE 'default' values, as follows:

- 15 • For dichotomous outcomes the MIDs were taken to be RRs of 0.8\* and 1.25. For  
16 'positive' outcomes such as 'patient satisfaction', the RR of 0.8 is taken as the line  
17 denoting the boundary between no clinically important effect and a clinically  
18 important harm, whilst the RR of 1.25 is taken as the line denoting the boundary  
19 between no clinically important effect and a clinically important benefit. For  
20 'negative' outcomes such as 'bleeding', the opposite occurs, so the RR of 0.8 is  
21 taken as the line denoting the boundary between no clinically important effect and  
22 a clinically important benefit, whilst the RR of 1.25 is taken as the line denoting the  
23 boundary between no clinically important effect and a clinically important harm.  
24 There aren't established default values for ORs and the same values (0.8 and  
25 1.25) are applied here but are acknowledged as arbitrary thresholds agreed by the  
26 committee.
  - 27 ○ In cases where there are zero events in one arm of a single study, or some or  
28 all of the studies in one arm of a meta-analysis, the same process is followed  
29 as for dichotomous outcomes. However, if there are no events in either arm in  
30 a meta-analysis (or in a single unpooled study) the sample size is used to  
31 determine imprecision using the following rule of thumb:
    - 32 – No imprecision: sample size  $\geq 350$
    - 33 – Serious imprecision: sample size  $\geq 70$  but  $< 350$
    - 34 – Very serious imprecision: sample size  $< 70$ .
  - 35 ○ When there was more than one study in an analysis and zero events occurred  
36 in both groups for some but not all of the studies across both arms, the  
37 optimum information size was used to determine imprecision using the  
38 following guide:
    - 39 – No imprecision:  $> 90\%$  power
    - 40 – Serious imprecision: 80-90% power
    - 41 – Very serious imprecision:  $< 80\%$  power.
- 42 • For continuous outcome variables the MID was taken as half the median baseline  
43 standard deviation of that variable, across all studies in the meta-analysis. Hence  
44 the MID denoting the minimum clinically important benefit was positive for a  
45 'positive' outcome (for example, a quality of life measure where a higher score  
46 denotes better health), and negative for a 'negative' outcome (for example, a  
47 visual analogue scale [VAS] pain score). Clinically important harms will be the  
48 converse of these. If baseline values are unavailable, then half the median  
49 comparator group standard deviation of that variable will be taken as the MID. As

1 these vary for each outcome per review, details of the values used are reported in  
2 the footnotes of the relevant GRADE summary table.

- 3 • If standardised mean differences have been used, where the GC are able to  
4 specify a priority measure, the results are back-converted to a mean difference on  
5 that scale for the assessment of imprecision and clinical importance. If it is not  
6 deemed appropriate to back-convert to a single scale, then the MID was set at the  
7 absolute value of +0.5. This follows because standardised mean differences are  
8 mean differences normalised to the pooled standard deviation of the 2 groups,  
9 and are thus effectively expressed in units of 'numbers of standard deviations'.  
10 The 0.5 MID value in this context therefore indicates half a standard deviation, the  
11 same definition of MID as used for non-standardised mean differences. Where  
12 there was only one study included 0.5 SD (SMD) was used.

13 \*NB GRADE report the default values as 0.75 and 1.25. These are consensus  
14 values. This guideline follows NICE process to use modified values of 0.8 and 1.25  
15 as they are symmetrical on a relative risk scale.


16 For this guideline, the following MIDs for continuous or dichotomous outcomes were  
17 found in the literature and adopted for use:

18 **Table 4: Published or pre-agreed MIDs**

Outcome measure	MID	Source
EQ-5D	0.03	Consensus pragmatic MID used in previous NGC NICE guidelines
SF36	Physical component summary: 2 Mental component summary: 3 Physical functioning: 3 Role-physical: 3 Bodily pain: 3 General health: 2 Vitality: 2 Social functioning: 3 Role-emotional: 4 Mental health: 3	User's manual for the SF-36v2 Health Survey, Third Edition <sup>2</sup>

19

**Figure 1:** Illustration of precise and imprecise outcomes based on the 95% CI of dichotomous outcomes in a forest plot (Note that all 3 results would be pooled estimates, and would not, in practice, be placed on the same forest plot)



### 1 2.5.1.5 Overall grading of the quality of clinical evidence

2 Once an outcome had been appraised for the main quality elements, as above, an  
 3 overall quality grade was calculated for that outcome. The scores (0, -1 or -2)  
 4 from each of the main quality elements were summed to give a score that could be  
 5 anything from 0 (the best possible) to -8 (the worst possible). However, scores were  
 6 capped at -3. This final score was then applied to the starting grade that had  
 7 originally been applied to the outcome by default, based on study design. RCTs start  
 8 at High, the overall quality became Moderate, Low or Very Low if the overall score  
 9 was -1, -2 or -3 points respectively. The significance of these overall ratings is  
 10 explained in Table 5. The reasons for downgrading in each case are specified in the  
 11 footnotes of the GRADE tables.

12 Non-randomised intervention studies started at Low, and so a score of -1 would be  
 13 enough to take the grade to the lowest level of Very Low. Non-randomised  
 14 intervention studies could, however, be upgraded if there was a large magnitude of  
 15 effect or a dose-response gradient.

16 **Table 5: Overall quality of outcome evidence in GRADE**

Level	Description
High	Further research is very unlikely to change our confidence in the estimate of effect
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate
Very low	Any estimate of effect is very uncertain

## 1 2.5.2 Prognostic reviews

2 An adapted GRADE profile was used for quality assessment per outcome. If data  
3 were meta-analysed, the quality for pooled studies was presented. If the data were  
4 not pooled, then a quality rating was presented for each study.

### 52.5.2.1.1 Risk of bias

6 The risk of bias for prognostic studies was evaluated according to the QUIPS  
7 checklist, the main criteria are given in **Table 6**.

8 **Table 6: Description of risk of bias criteria for prognostic studies**

Risk of bias	Aim of section
Study participation	To judge selection bias (likelihood that relationship between the prognostic factor and outcome is different for participants and eligible non-participants)
Study attrition	To judge the risk of attrition bias (likelihood that relationship between prognostic factor and outcome are different for completing and non-completing participants).
Prognostic factor measurement	To judge the risk of measurement bias related to how the prognostic factor was measured (differential measurement of prognostic factor related to the baseline level of outcome).
Outcome measurement	To judge the risk of bias related to the measurement of outcome (differential measurement of outcome related to the baseline level of prognostic factor).
Study confounding	To judge the risk of bias due to confounding (i.e. the effect of the prognostic factor is distorted by another factor that is related to the prognostic factor and outcome).
Statistical Analysis and Reporting	To judge the risk of bias related to the statistical analysis and presentation of results.

### 92.5.2.1.2 Inconsistency

10 Inconsistency was assessed as for intervention studies.

### 112.5.2.1.3 Imprecision

12 In meta-analysed outcomes, or for non-pooled outcomes, the position of the 95% CIs  
13 in relation to the null line determined the existence of imprecision. If the 95% CI did  
14 not cross the null line then no serious imprecision was recorded. If the 95% CI  
15 crossed the null line then serious imprecision was recorded.

### 162.5.2.1.4 Overall grading

17 Quality rating was assigned by study. However, if there was more than 1 outcome  
18 involved in a study, then the quality rating of the evidence statements for each  
19 outcome was adjusted accordingly. For example, if one outcome was based on an  
20 invalidated measurement method, but another outcome in the same study was not,  
21 the second outcome would be graded 1 grade higher than the first outcome.

22 Quality rating started at high for prospective and retrospective studies, and each  
23 major limitation brought the rating down by 1 increment to a minimum grade rating of  
24 very low, as explained for interventional reviews. For prognostic reviews prospective  
25 cohort studies with a multivariate analysis are regarded as the gold standard

1 because RCTs are usually an inappropriate design to answer the question for these  
 2 types of review. Furthermore, if the study is looking at more than 1 prognostic factor  
 3 of interest then randomisation would be inappropriate as it can only be applied to 1 of  
 4 the prognostic factors.

### 5 2.5.3 Qualitative reviews

6 Review findings from the included qualitative studies were evaluated and presented  
 7 using the 'Confidence in the Evidence from Reviews of Qualitative Research'  
 8 (CERQual) Approach developed by the GRADE-CERQual Project Group, a subgroup  
 9 of the GRADE Working Group.

10 The CERQual Approach assesses the extent to which a review finding is a  
 11 reasonable representation of the phenomenon of interest (the focus of the review  
 12 question). Each review finding was assessed for each of the 4 quality elements listed  
 13 and defined below in Table 7.

14 **Table 7: Description of quality elements in GRADE-CERQual for qualitative**  
 15 **studies**

Quality element	Description
Methodological limitations	The extent of problems in the design or conduct of the included studies that could decrease the confidence that the review finding is a reasonable representation of the phenomenon of interest. Assessed at the study level using the CASP checklist.
Coherence	The extent to how clear and cogent the fit is between the data from the primary studies and the review finding.
Relevance	The extent to which the body of evidence from the included studies is applicable to the context (study population, phenomenon of interest, setting) specified in the protocol.
Adequacy	The degree of the confidence that the review finding is being supported by sufficient data. This is an overall determination of the richness (depth of analysis) and quantity of the evidence supporting a review finding or theme.

16 Details of how the 4 quality elements (methodological limitations, coherence,  
 17 relevance and adequacy) were appraised for each review finding are given below.

#### 18 2.5.3.1 Methodological limitations

19 Each review finding had its methodological limitations assessed within each study  
 20 first using the CASP checklist. Based on the degree of methodological limitations,  
 21 studies were evaluated as having minor, moderate or severe limitations. A summary  
 22 of the domains and questions covered is given below.

23 **Table 8: Description of limitations assessed in the CASP checklist for**  
 24 **qualitative studies**

Domain	Aspects considered
Are the results valid?	<ul style="list-style-type: none"> <li>• Was there a clear statement of the aims of the research?</li> <li>• Is qualitative methodology appropriate?</li> <li>• Was the research design appropriate to address the aims of the research?</li> <li>• Was the recruitment strategy appropriate to the aims of the research?</li> <li>• Was the data collected in a way that addressed the research issue?</li> </ul>

Domain	Aspects considered
	<ul style="list-style-type: none"> <li>Has the relationship between researcher and participants been adequately considered?</li> </ul>
What are the results?	Have ethical issues been taken into consideration? Was the data analysis sufficiently rigorous? Is there a clear statement of findings?
Will the results help locally?	How valuable is the research?

1 The overall assessment of the methodological limitations of the evidence was based  
 2 on the limitations of the primary studies contributing to the review finding. The relative  
 3 contribution of each study to the overall review finding and of the type of  
 4 methodological limitation(s) were taken into account when giving an overall rating of  
 5 concerns for this component.

#### 6 **2.5.3.2 Relevance**

7 Relevance is the extent to which the body of evidence from the included studies is  
 8 applicable to the context (study population, phenomenon of interest, setting)  
 9 specified in the protocol. As such, relevance is dependent on the individual review  
 10 and discussed with the guideline committee.

#### 11 **2.5.3.3 Coherence**

12 Coherence is the extent to which the reviewer is able to identify a clear pattern  
 13 across the studies included in the review, and if there is variation present (contrasting  
 14 or disconfirming data) whether this variation is explained by the contributing study  
 15 authors. For example, if a review finding in 1 study does not support the main finding  
 16 and there is no plausible explanation for this variation, or if there is ambiguity in the  
 17 descriptions in the primary data, then the confidence that the main finding reasonably  
 18 reflects the phenomenon of interest is decreased.

#### 19 **2.5.3.4 Adequacy**

20 The judgement of adequacy is based on the confidence of the finding being  
 21 supported by sufficient data. This is an overall determination of the richness (and  
 22 quantity of the evidence supporting a review finding or theme. Rich data provide  
 23 sufficient detail to gain an understanding of the theme or review finding, whereas thin  
 24 data do not provide enough detail for an adequate understanding. Quantity of data is  
 25 the second pillar of the assessment of adequacy. For review findings that are only  
 26 supported by 1 study or data from only a small number of participants, the confidence  
 27 that the review finding reasonably represents the phenomenon of interest might be  
 28 decreased because there is less confidence that studies undertaken in other settings  
 29 or participants would have reported similar findings. As with richness of data, quantity  
 30 of data is review dependent. Based on the overall judgement of adequacy, a rating of  
 31 no concerns, minor concerns, or substantial concerns about adequacy was given.

#### 32 **2.5.3.5 Overall judgement of the level of confidence for a review finding**

33 GRADE-CERQual is used to assess the body of evidence as a whole through a  
 34 confidence rating representing the extent to which a review finding is a reasonable

1 representation of the phenomenon of interest. For each of the above components,  
2 level of concern is categorised as either;

- 3 • no or very minor concerns
- 4 • minor concerns
- 5 • moderate concerns, or
- 6 • serious concerns.

7 The concerns from the 4 components (methodological limitations, coherence,  
8 relevance and adequacy) are used in combination to form an overall judgement of  
9 confidence in the finding. GRADE-CERQual uses 4 levels of confidence: high,  
10 moderate, low and very low confidence. The significance of these overall ratings is  
11 explained in Table 9. Each review finding starts at a high level of confidence and is  
12 downgraded based on the concerns identified in any 1 or more of the 4 components.  
13 Quality assessment of qualitative reviews is a subjective judgement by the reviewer  
14 based on the concerns that have been noted. An explanation of how such a  
15 judgement had been made for each component is included in the footnotes of the  
16 summary of evidence tables.

17 **Table 9: Overall level of confidence for a review finding in GRADE-CERQual**

Level	Description
High confidence	It is highly likely that the review finding is a reasonable representation of the phenomenon of interest.
Moderate confidence	It is likely that the review finding is a reasonable representation of the phenomenon of interest.
Low confidence	It is possible that the review finding is a reasonable representation of the phenomenon of interest.
Very low confidence	It is not clear whether the review finding is a reasonable representation of the phenomenon of interest.

## 18 2.5.4 Enrichment design trials

### 19 **Selecting participants based on previous response to treatment.**

20 Clinical trial enrichment methods involve the selection of participants most likely to  
21 respond to treatment. Trials using this enrichment method could overestimate the  
22 true efficacy of treatment in the general population. An analysis of this overestimation  
23 could provide a clear precedent and justification for, where appropriate, excluding  
24 these trials from guideline decision-making. Inclusion criteria that, explicitly or  
25 otherwise, select a population that does not reflect the population for which the  
26 treatment is designed could result in flawed evidence for recommendations. The use  
27 of this evidence could influence the observed treatment effect and the subsequent  
28 cost-effectiveness analysis, which in turn could impact on the treatments  
29 recommended.

### 30 **Sensitivity analysis approach**

31 In the case of heterogeneity, sensitivity analysis should be conducted to exclude all  
32 trials utilising enrichment methods (described below). However, exploration of the  
33 effects of the **type** of enrichment study should also be explored given that they select  
34 participants based on previous response to varying extents. Sensitivity analysis  
35 approach should therefore be:

- 36 1. Sensitivity analysis 1: Exclude categories 1-4
- 37 2. Sensitivity analysis 2: Exclude category 1 only

3. Base recommendations on the sensitivity analysis that best explains heterogeneity (or neither if neither explain heterogeneity)
  - a. Be both consistent and conservative with approach throughout the evidence review – i.e. if one outcome shows that excluding all 3 categories of enrichment trials explains heterogeneity, then exclude all of these studies throughout the comparison regardless of whether heterogeneity is present

## Categories

### 1. Including only responders

1. Run-in treatment washout, excluding those that get better/stay the same when medication withdrawn
2. Run-in treatment washout, including only those whose pain increases on withdrawal
3. Run-in period with test treatment, inclusion criteria are only those who respond to the treatment
4. Inclusion criteria – known previous responders to the treatment being tested

### 2. Excluding non-responders (but not exclusively including only responders)

1. Run-in treatment washout, excluding those that get better when the treatment is withdrawn
2. Exclusion criteria – previous non-response/intolerance to the treatment under investigation

### 3. Selection of specific population

1. Inclusion criteria – using a specific therapy, exclusion criteria – responding to a treatment
2. Exclusion criteria – those who are currently well controlled with minimal adverse events
3. Those who have not responded to adequate trials of one or more other treatments
4. Run-in placebo period, excluding people that respond to placebo
5. Response criteria (either via run-in period, washout or inclusion criteria) related to a treatment **not** being tested (e.g. NSAID responders randomised to opioid versus placebo)

### 4. Unclear

1. Previous response to treatment is unclear, some implicit suggestion that selection based on response may have taken place

### 5. All treatment naïve

1. All participants are naïve to the treatment being tested

### 6. No response criteria

1. Population washed out of previous treatment, but no inclusion criteria based on withdrawal response

### 7. Mixed population

1. Inclusion criteria for people already receiving treatment fits into the above, but treatment naïve participants also included (and these accounted for >20% of the population)



1 *Note: Where a study states treatment naïve participants were included but the*  
2 *number is not stated, this is assumed to be <20% (and so study categorised into one*  
3 *of the above)*

## 4 **2.6 Assessing clinical importance**

5 The committee assessed the evidence by outcome in order to determine if there was,  
6 or potentially was, a clinically important benefit, a clinically important harm or no  
7 clinically important difference between interventions. To facilitate this, binary  
8 outcomes were converted into absolute risk differences (ARDs) using GRADEpro<sup>1</sup>  
9 software: the median control group risk across studies was used to calculate the  
10 ARD and its 95% CI from the pooled risk ratio.

11 The assessment of clinical benefit, harm, or no benefit or harm was based on the  
12 point estimate of absolute effect for intervention studies, which was standardised  
13 across the reviews. The committee considered for most of the dichotomous  
14 outcomes in the intervention reviews that if at least 100 more participants per 1000  
15 (10%) achieved the outcome of interest in the intervention group compared to the  
16 comparison group for a positive outcome then this intervention was considered  
17 beneficial. The same point estimate but in the opposite direction applied for a  
18 negative outcome. For mortality any reduction represented a clinical benefit. For  
19 adverse events 50 events or more per 1000 (5%) represented clinical harm.

20 For continuous outcomes if the mean difference was greater than the minimally  
21 important difference (MID) then this represented a clinical benefit or harm. For  
22 outcomes such as mortality any reduction or increase was considered to be clinically  
23 important.

24 Established MIDs found in the literature and were agreed to be used for EQ-5D and  
25 SF-36 (as measures of quality of life).

26 The published values used for imprecision and clinical importance are provided in  
27 **Table 4**. For continuous outcomes where the GRADE default MID has been used,  
28 the values for each outcome are provided in the comments column incorporated into  
29 the relevant GRADE tables.

## 30 **2.7 Identifying and analysing evidence of cost** 31 **effectiveness**

32 The committee is required to make decisions based on the best available evidence of  
33 both clinical effectiveness and cost effectiveness. Guideline recommendations should  
34 be based on the expected costs of the different options in relation to their expected  
35 health benefits (that is, their 'cost effectiveness') rather than the total implementation  
36 cost. However, the committee will also need to be increasingly confident in the cost  
37 effectiveness of a recommendation as the cost of implementation increases.  
38 Therefore, the committee may require more robust evidence on the effectiveness and  
39 cost effectiveness of any recommendations that are expected to have a substantial  
40 impact on resources; any uncertainties must be offset by a compelling argument in  
41 favour of the recommendation. The cost impact or savings potential of a  
42 recommendation should not be the sole reason for the committee's decision.<sup>4</sup>

43 Health economic evidence was sought relating to the key clinical issues being  
44 addressed in the guideline. Health economists:

- 1 • Undertook a systematic review of the published economic literature.
- 2 • Undertook new cost-effectiveness analysis in priority areas.

### 3 **2.7.1 Literature review**

4 The health economists:

- 5 • Identified potentially relevant studies for each review question from the health  
6 economic search results by reviewing titles and abstracts. Full papers were then  
7 obtained.
- 8 • Reviewed full papers against prespecified inclusion and exclusion criteria to  
9 identify relevant studies (see below for details).
- 10 • Critically appraised relevant studies using economic evaluations checklists as  
11 specified in the NICE guidelines manual.<sup>4</sup>
- 12 • Extracted key information about the studies' methods and results into health  
13 economic evidence tables (which can be found in appendices to the relevant  
14 evidence reports).
- 15 • Generated summaries of the evidence in NICE health economic evidence profile  
16 tables (included in the relevant evidence report for each review question) – see  
17 below for details.

#### 18 **2.7.1.1 Inclusion and exclusion criteria**

19 Full economic evaluations (studies comparing costs and health consequences of  
20 alternative courses of action: cost–utility, cost-effectiveness, cost–benefit and cost–  
21 consequences analyses) and comparative costing studies that addressed the review  
22 question in the relevant population were considered potentially includable as health  
23 economic evidence.

24 Studies that only reported cost per hospital (not per patient), or only reported average  
25 cost effectiveness without disaggregated costs and effects were excluded. Literature  
26 reviews, abstracts, posters, letters, editorials, comment articles, unpublished studies  
27 and studies not in English were excluded. Studies published before and studies from  
28 non-OECD countries or the USA were also excluded, on the basis that the  
29 applicability of such studies to the present UK NHS context is likely to be too low for  
30 them to be helpful for decision-making.

31 Remaining health economic studies were prioritised for inclusion based on their  
32 relative applicability to the development of this guideline and the study limitations.

33 about the assessment of applicability and methodological quality see **Table 10** below  
34 and the economic evaluation checklist (appendix H of the NICE guidelines manual<sup>4</sup>)  
35 and the health economics review protocol, which can be found in each of the  
36 evidence reports.

37 When no relevant health economic studies were found from the economic literature  
38 review, relevant UK NHS unit costs related to the compared interventions were  
39 presented to the committee to inform the possible economic implications of the  
40 recommendations.

### 1 2.7.1.2 NICE health economic evidence profiles

2 NICE health economic evidence profile tables were used to summarise cost and  
 3 cost-effectiveness estimates for the included health economic studies in each  
 4 evidence review report. The health economic evidence profile shows an assessment  
 5 of applicability and methodological quality for each economic study, with footnotes  
 6 indicating the reasons for the assessment. These assessments were made by the  
 7 health economist using the economic evaluation checklist from the NICE guidelines  
 8 manual.<sup>4</sup> It also shows the incremental costs, incremental effects (for example,  
 9 quality-adjusted life years [QALYs]) and incremental cost-effectiveness ratio (ICER)  
 10 for the base case analysis in the study, as well as information about the assessment  
 11 of uncertainty in the analysis. See **Table 10** for more details.

12 When a non-UK study was included in the profile, the results were converted into  
 13 pounds sterling using the appropriate purchasing power parity.<sup>8</sup>

14 **Table 10: Content of NICE health economic evidence profile**

Item	Description
Study	Surname of first author, date of study publication and country perspective with a reference to full information on the study.
Applicability	An assessment of applicability of the study to this guideline, the current NHS situation and NICE decision-making: <sup>(a)</sup> <ul style="list-style-type: none"> <li>• Directly applicable – the study meets all applicability criteria, or fails to meet 1 or more applicability criteria but this is unlikely to change the conclusions about cost effectiveness.</li> <li>• Partially applicable – the study fails to meet 1 or more applicability criteria, and this could change the conclusions about cost effectiveness.</li> <li>• Not applicable – the study fails to meet 1 or more of the applicability criteria, and this is likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.</li> </ul>
Limitations	An assessment of methodological quality of the study: <sup>(a)</sup> <ul style="list-style-type: none"> <li>• Minor limitations – the study meets all quality criteria, or fails to meet 1 or more quality criteria, but this is unlikely to change the conclusions about cost effectiveness.</li> <li>• Potentially serious limitations – the study fails to meet 1 or more quality criteria, and this could change the conclusions about cost effectiveness.</li> <li>• Very serious limitations – the study fails to meet 1 or more quality criteria, and this is highly likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.</li> </ul>
Other comments	Information about the design of the study and particular issues that should be considered when interpreting it.
Incremental cost	The mean cost associated with one strategy minus the mean cost of a comparator strategy.
Incremental effects	The mean QALYs (or other selected measure of health outcome) associated with one strategy minus the mean QALYs of a comparator strategy.
Cost effectiveness	Incremental cost-effectiveness ratio (ICER): the incremental cost divided by the incremental effects (usually in £ per QALY gained).
Uncertainty	A summary of the extent of uncertainty about the ICER reflecting the results of deterministic or probabilistic sensitivity analyses, or stochastic analyses of trial data, as appropriate.

15 (a) *Applicability and limitations were assessed using the economic evaluation checklist in appendix H of*  
 16 *the NICE guidelines manual*<sup>4</sup>

## 1 2.7.2 Undertaking new health economic analysis

2 As well as reviewing the published health economic literature for each review  
3 question, as described above, new health economic analysis was undertaken by the  
4 health economist in selected areas. Priority areas for new analysis were agreed by  
5 the committee after formation of the review questions and consideration of the  
6 existing health economic evidence.

7 The committee identified oral, topical and transdermal (OTT) medicines and  
8 electroacupuncture as the highest priority areas for original health economic  
9 modelling. Electroacupuncture was given a high priority since there were no studies  
10 identified during the economic review that compared it with usual care, and a positive  
11 recommendation will likely cause a significant resource impact for the NHS. OTT  
12 medicines were given a high priority for modelling since current practice for medical  
13 management of osteoarthritis is varied and the harms and benefits associated with  
14 different drug classes is difficult to assess without an economic model quantifying the  
15 costs and quality adjusted life years (QALYs) for each drug class.

16 The following general principles were adhered to in developing the cost-effectiveness  
17 analyses:

- 18 • Methods were consistent with the NICE reference case for interventions with  
19 health outcomes in NHS settings.<sup>4, 6</sup>
- 20 • The committee was involved in the design of the model, selection of inputs and  
21 interpretation of the results.
- 22 • Model inputs were based on the systematic review of the clinical literature  
23 supplemented with other published data sources where possible.
- 24 • When published data were not available committee expert opinion was used to  
25 populate the model.
- 26 • Model inputs and assumptions were reported fully and transparently.
- 27 • The results were subject to sensitivity analysis and limitations were discussed.
- 28 • The model was peer-reviewed by another health economist at the NGC.

29 Full methods and results of the cost-effectiveness analysis for the oral, topical and  
30 transdermal (OTT) medicines model and electroacupuncture model are described in  
31 a separate economic analysis reports.

## 32 2.7.3 Cost-effectiveness criteria

33 NICE sets out the principles that committees should consider when judging whether  
34 an intervention offers good value for money.<sup>4, 5, 7</sup> In general, an intervention was  
35 considered to be cost effective (given that the estimate was considered plausible) if  
36 either of the following criteria applied:

- 37 • the intervention dominated other relevant strategies (that is, it was both less costly  
38 in terms of resource use and more clinically effective compared with all the other  
39 relevant alternative strategies), or
- 40 • the intervention cost less than £20,000 per QALY gained compared with the next  
41 best strategy.

42 If the committee recommended an intervention that was estimated to cost more than  
43 £20,000 per QALY gained, or did not recommend one that was estimated to cost less  
44 than £20,000 per QALY gained, the reasons for this decision are discussed explicitly

1 in 'The committee's discussion of the evidence' section of the relevant evidence  
2 report, with reference to issues regarding the plausibility of the estimate or to factors  
3 set out in NICE methods manuals.<sup>4</sup>

4 If a study reported the cost per life year gained but not QALYs, the cost per QALY  
5 gained was estimated by multiplying by an appropriate utility estimate to aid  
6 interpretation. The estimated cost per QALY gained is reported in the health  
7 economic evidence profile with a footnote detailing the life-years gained and the  
8 utility value used. When QALYs or life years gained are not used in the analysis,  
9 results are difficult to interpret unless one strategy dominates the others with respect  
10 to every relevant health outcome and cost.

#### 11 **2.7.4 In the absence of health economic evidence**

12 When no relevant published health economic studies were found, and a new analysis  
13 was not prioritised, the committee made a qualitative judgement about cost  
14 effectiveness by considering expected differences in resource use between options  
15 and relevant UK NHS unit costs, alongside the results of the review of clinical  
16 effectiveness evidence.

17 The UK NHS costs reported in the guideline are those that were presented to the  
18 committee and were correct at the time recommendations were drafted. They may  
19 have changed subsequently before the time of publication. However, we have no  
20 reason to believe they have changed substantially.

## 21 **2.8 Developing recommendations**

22 Over the course of the guideline development process, the committee was presented  
23 with:

- 24 • Summaries of clinical and health economic evidence and quality (as presented in  
25 evidence reports [A–J]).
- 26 • Evidence tables of the clinical and health economic evidence reviewed from the  
27 literature. All evidence tables can be found in appendices to the relevant evidence  
28 reports.
- 29 • Forest plots (in appendices to the relevant evidence reports).
- 30 • A description of the methods and results of the cost-effectiveness analysis  
31 undertaken for the guideline (in a separate economic analysis report).

32 Decisions on whether a recommendation could be made, and if so in which direction,  
33 were made on the basis of the committee's interpretation of the available evidence,  
34 taking into account the balance of benefits, harms and costs between different  
35 courses of action. This was either done formally in an economic model, or informally.  
36 The net clinical benefit over harm (clinical effectiveness) was considered, focusing on  
37 the magnitude of the effect (or clinical importance), quality of evidence (including the  
38 uncertainty) and amount of evidence available. When this was done informally, the  
39 committee took into account the clinical benefits and harms when one intervention  
40 was compared with another. The assessment of net clinical benefit was moderated  
41 by the importance placed on the outcomes (the committee's values and preferences),  
42 and the confidence the committee had in the evidence (evidence quality). Secondly,  
43 the committee assessed whether the net clinical benefit justified any differences in  
44 costs between the alternative interventions. When the clinical harms were judged by  
45 the committee to outweigh any clinical benefits, they considered making a

1 recommendation not to offer an intervention. This was dependant on whether the  
2 intervention had any reasonable prospect of providing cost-effective benefits to  
3 people using services and whether stopping the intervention was likely to cause harm  
4 for people already receiving it.

5 When clinical and health economic evidence was of poor quality, conflicting or  
6 absent, the committee decided on whether a recommendation could be made based  
7 on its expert opinion. The considerations for making consensus-based  
8 recommendations include the balance between potential harms and benefits, the  
9 economic costs compared to the economic benefits, current practices,  
10 recommendations made in other relevant guidelines, patient preferences and equality  
11 issues. The consensus recommendations were agreed through discussions in the  
12 committee. The committee also considered whether the uncertainty was sufficient to  
13 justify delaying making a recommendation to await further research, taking into  
14 account the potential harm of failing to make a clear recommendation (see  
15 section 2.8.1 below).

16 The committee considered the appropriate 'strength' of each recommendation. This  
17 takes into account the quality of the evidence but is conceptually different. Some  
18 recommendations are 'strong' in that the committee believes that the vast majority of  
19 healthcare and other professionals and patients would choose a particular  
20 intervention if they considered the evidence in the same way that the committee has.  
21 This is generally the case if the benefits clearly outweigh the harms for most people  
22 and the intervention is likely to be cost effective. However, there is often a closer  
23 balance between benefits and harms, and some patients would not choose an  
24 intervention whereas others would. This may happen, for example, if some patients  
25 are particularly averse to some side effect and others are not. In these circumstances  
26 the recommendation is generally weaker, although it may be possible to make  
27 stronger recommendations about specific groups of patients.

28 The committee focused on the following factors in agreeing the wording of the  
29 recommendations:

- 30 • The actions health professionals need to take.
- 31 • The information readers need to know.
- 32 • The strength of the recommendation (for example the word 'offer' was used for  
33 strong recommendations and 'consider' for weaker recommendations).
- 34 • The involvement of patients (and their carers if needed) in decisions on treatment  
35 and care.
- 36 • Consistency with NICE's standard advice on recommendations about drugs,  
37 waiting times and ineffective interventions (see section 9.2 in the NICE guidelines  
38 manual<sup>4</sup>).

39 The main considerations specific to each recommendation are outlined in 'The  
40 committee's discussion of the evidence' section within each evidence report.

### 41 **2.8.1 Research recommendations**

42 When areas were identified for which good evidence was lacking, the committee  
43 considered making recommendations for future research. Decisions about the  
44 inclusion of a research recommendation were based on factors such as:

- 45 • the importance to patients or the population
- 46 • national priorities

- 1 • potential impact on the NHS and future NICE guidance
- 2 • ethical and technical feasibility.

### 3 **2.8.2 Validation process**

4 This guidance is subject to a 6-week public consultation and feedback as part of the  
5 quality assurance and peer review of the document. All comments received from  
6 registered stakeholders are responded to in turn and posted on the NICE website.

### 7 **2.8.3 Updating the guideline**

8 Following publication, and in accordance with the NICE guidelines manual, NICE will  
9 undertake a review of whether the evidence base has progressed significantly to alter  
10 the guideline recommendations and warrant an update.

### 11 **2.8.4 Disclaimer**

12 Healthcare providers need to use clinical judgement, knowledge and expertise when  
13 deciding whether it is appropriate to apply guidelines. The recommendations cited  
14 here are a guide and may not be appropriate for use in all situations. The decision to  
15 adopt any of the recommendations cited here must be made by practitioners in light  
16 of individual patient circumstances, the wishes of the patient, clinical expertise and  
17 resources.

18 The National Guideline Centre disclaims any responsibility for damages arising out of  
19 the use or non-use of this guideline and the literature used in support of this  
20 guideline.

### 21 **2.8.5 Funding**

22 The National Guideline Centre was commissioned by the National Institute for Health  
23 and Care Excellence to undertake the work on this guideline.

24

## 25 **2.9 General terms**

26

Term	Definition
Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.
Algorithm (in guidelines)	A flow chart of the clinical decision pathway described in the guideline, where decision points are represented with boxes, linked with arrows.
Allocation concealment	The process used to prevent advance knowledge of group assignment in an RCT. The allocation process should be impervious to any influence by the individual making the allocation, by being administered by someone who is not responsible for recruiting participants.
Applicability	How well the results of a study or NICE evidence review can answer a clinical question or be applied to the population being considered.
Arm (of a clinical study)	Subsection of individuals within a study who receive one particular intervention, for example placebo arm.

Term	Definition
Association	Statistical relationship between 2 or more events, characteristics or other variables. The relationship may or may not be causal.
Base case analysis	In an economic evaluation, this is the main analysis based on the most plausible estimate of each input. In contrast, see Sensitivity analysis.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable), with which subsequent results are compared.
Bayesian analysis	A method of statistics, where a statistic is estimated by combining established information or belief (the 'prior') with new evidence (the 'likelihood') to give a revised estimate (the 'posterior').
Before-and-after study	A study that investigates the effects of an intervention by measuring particular characteristics of a population both before and after taking the intervention, and assessing any change that occurs.
Bias	Influences on a study that can make the results look better or worse than they really are. (Bias can even make it look as if a treatment works when it does not.) Bias can occur by chance, deliberately or as a result of systematic errors in the design and execution of a study. It can also occur at different stages in the research process, for example, during the collection, analysis, interpretation, publication or review of research data. For examples see selection bias, performance bias, information bias, confounding factor, and publication bias.
Blinding	A way to prevent researchers, doctors and patients in a clinical trial from knowing which study group each patient is in so they cannot influence the results. The best way to do this is by sorting patients into study groups randomly. The purpose of 'blinding' or 'masking' is to protect against bias. A single-blinded study is one in which patients do not know which study group they are in (for example whether they are taking the experimental drug or a placebo). A double-blinded study is one in which neither patients nor the researchers and doctors know which study group the patients are in. A triple blind study is one in which neither the patients, clinicians or the people carrying out the statistical analysis know which treatment patients received.
Carer (caregiver)	Someone who looks after family, partners or friends in need of help because they are ill, frail or have a disability.
Case-control study	A study to find out the cause(s) of a disease or condition. This is done by comparing a group of patients who have the disease or condition (cases) with a group of people who do not have it (controls) but who are otherwise as similar as possible (in characteristics thought to be unrelated to the causes of the disease or condition). This means the researcher can look for aspects of their lives that differ to see if they may cause the condition. For example, a group of people with lung cancer might be compared with a group of people the same age that do not have lung cancer. The researcher could compare how long both groups had been exposed to tobacco smoke. Such studies are retrospective because they look back in time from the outcome to the possible causes of a disease or condition.
Case series	Report of a number of cases of a given disease, usually covering the course of the disease and the response to treatment. There is no comparison (control) group of patients.



Term	Definition
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinical effectiveness	How well a specific test or treatment works when used in the 'real world' (for example, when used by a doctor with a patient at home), rather than in a carefully controlled clinical trial. Trials that assess clinical effectiveness are sometimes called management trials. Clinical effectiveness is not the same as efficacy.
Clinician	A healthcare professional who provides patient care. For example, a doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence-based medicine databases including the Cochrane Database of Systematic Reviews (reviews of randomised controlled trials prepared by the Cochrane Collaboration).
Cohort study	A study with 2 or more groups of people – cohorts – with similar characteristics. One group receives a treatment, is exposed to a risk factor or has a particular symptom and the other group does not. The study follows their progress over time and records what happens. See also observational study.
Comorbidity	A disease or condition that someone has in addition to the health problem being studied or treated.
Comparability	Similarity of the groups in characteristics likely to affect the study results (such as health status or age).
Concordance	This is a recent term whose meaning has changed. It was initially applied to the consultation process in which doctor and patient agree therapeutic decisions that incorporate their respective views, but now includes patient support in medicine taking as well as prescribing communication. Concordance reflects social values but does not address medicine-taking and may not lead to improved adherence.
Confidence interval (CI)	A range of values for an unknown population parameter with a stated 'confidence' (conventionally 95%) that it contains the true value. The interval is calculated from sample data, and generally straddles the sample estimate. The 'confidence' value means that if the method used to calculate the interval is repeated many times, then that proportion of intervals will actually contain the true value.
Confounding factor	Something that influences a study and can result in misleading findings if it is not understood or appropriately dealt with. For example, a study of heart disease may look at a group of people that exercises regularly and a group that does not exercise. If the ages of the people in the 2 groups are different, then any difference in heart disease rates between the 2 groups could be because of age rather than exercise. Therefore, age is a confounding factor.
Consensus methods	Techniques used to reach agreement on a particular issue. Consensus methods may be used to develop NICE guidance if there is not enough good quality research evidence to give a clear answer to a question. Formal consensus methods include Delphi and nominal group techniques.
Control group	A group of people in a study who do not receive the treatment or test being studied. Instead, they may receive the standard treatment (sometimes called 'usual care') or a dummy treatment (placebo). The results for the control group are compared with those for a group

Term	Definition
	receiving the treatment being tested. The aim is to check for any differences. Ideally, the people in the control group should be as similar as possible to those in the treatment group, to make it as easy as possible to detect any effects due to the treatment.
Cost–benefit analysis (CBA)	Cost–benefit analysis is one of the tools used to carry out an economic evaluation. The costs and benefits are measured using the same monetary units (for example, pounds sterling) to see whether the benefits exceed the costs.
Cost–consequences analysis (CCA)	Cost–consequences analysis is one of the tools used to carry out an economic evaluation. This compares the costs (such as treatment and hospital care) and the consequences (such as health outcomes) of a test or treatment with a suitable alternative. Unlike cost–benefit analysis or cost-effectiveness analysis, it does not attempt to summarise outcomes in a single measure (like the quality-adjusted life year) or in financial terms. Instead, outcomes are shown in their natural units (some of which may be monetary) and it is left to decision-makers to determine whether, overall, the treatment is worth carrying out.
Cost-effectiveness analysis (CEA)	Cost-effectiveness analysis is one of the tools used to carry out an economic evaluation. The benefits are expressed in non-monetary terms related to health, such as symptom-free days, heart attacks avoided, deaths avoided or life years gained (that is, the number of years by which life is extended as a result of the intervention).
Cost-effectiveness model	An explicit mathematical framework, which is used to represent clinical decision problems and incorporate evidence from a variety of sources in order to estimate the costs and health outcomes.
Cost–utility analysis (CUA)	Cost–utility analysis is one of the tools used to carry out an economic evaluation. The benefits are assessed in terms of both quality and duration of life, and expressed as quality-adjusted life years (QALYs). See also utility.
Credible interval (CrI)	The Bayesian equivalent of a confidence interval.
Decision analysis	An explicit quantitative approach to decision-making under uncertainty, based on evidence from research. This evidence is translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Deterministic analysis	In economic evaluation, this is an analysis that uses a point estimate for each input. In contrast, see Probabilistic analysis
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Disutility	The loss of quality of life associated with having a disease or condition. See Utility
Dominance	A health economics term. When comparing tests or treatments, an option that is both less effective and costs more is said to be 'dominated' by the alternative.
Drop-out	A participant who withdraws from a trial before the end.

Term	Definition
Economic evaluation	<p>An economic evaluation is used to assess the cost effectiveness of healthcare interventions (that is, to compare the costs and benefits of a healthcare intervention to assess whether it is worth doing). The aim of an economic evaluation is to maximise the level of benefits – health effects – relative to the resources available. It should be used to inform and support the decision-making process; it is not supposed to replace the judgement of healthcare professionals.</p> <p>There are several types of economic evaluation: cost–benefit analysis, cost–consequences analysis, cost-effectiveness analysis, cost-minimisation analysis and cost–utility analysis. They use similar methods to define and evaluate costs, but differ in the way they estimate the benefits of a particular drug, programme or intervention.</p>
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	<p>A measure that shows the magnitude of the outcome in one group compared with that in a control group.</p> <p>For example, if the absolute risk reduction is shown to be 5% and it is the outcome of interest, the effect size is 5%.</p> <p>The effect size is usually tested, using statistics, to find out how likely it is that the effect is a result of the treatment and has not just happened by chance (that is, to see if it is statistically significant).</p>
Effectiveness	<p>How beneficial a test or treatment is under usual or everyday conditions, compared with doing nothing or opting for another type of care.</p>
Efficacy	<p>How beneficial a test, treatment or public health intervention is under ideal conditions (for example, in a laboratory), compared with doing nothing or opting for another type of care.</p>
Enrichment design trials	<p>Where a population for a trial are selected based on parameters which may change the effect of the intervention (for example: people who have previously responded positively to a class of medication being included in a trial of that class of medication).</p>
Epidemiological study	<p>The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (for example, infection, diet) and interventions.</p>
EQ-5D (EuroQol 5 dimensions)	<p>A standardised instrument used to measure health-related quality of life. It provides a single index value for health status.</p>
Evidence	<p>Information on which a decision or guidance is based. Evidence is obtained from a range of sources including randomised controlled trials, observational studies, expert opinion (of clinical professionals or patients).</p>
Exclusion criteria (literature review)	<p>Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.</p>
Exclusion criteria (clinical study)	<p>Criteria that define who is not eligible to participate in a clinical study.</p>
Extended dominance	<p>If Option A is both more clinically effective than Option B and has a lower cost per unit of effect, when both are compared with a do-nothing alternative then Option A is said to have extended dominance over Option B. Option A is therefore cost effective and should be preferred, other things remaining equal.</p>
Extrapolation	<p>An assumption that the results of studies of a specific population will also hold true for another population with similar characteristics.</p>
Follow-up	<p>Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been</p>

Term	Definition
	assessed in order to observe changes in health status or health-related variables.
Generalisability	The extent to which the results of a study hold true for groups that did not participate in the research. See also external validity.
Gold standard	A method, procedure or measurement that is widely accepted as being the best available to test for or treat a disease.
GRADE, GRADE profile	A system developed by the GRADE Working Group to address the shortcomings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE profile.
Harms	Adverse effects of an intervention.
Hazard Ratio	The hazard or chance of an event occurring in the treatment arm of a study as a ratio of the chance of an event occurring in the control arm over time.
Health economics	Study or analysis of the cost of using and distributing healthcare resources.
Health-related quality of life (HRQoL)	A measure of the effects of an illness to see how it affects someone's day-to-day life.
Heterogeneity or Lack of homogeneity	The term is used in meta-analyses and systematic reviews to describe when the results of a test or treatment (or estimates of its effect) differ significantly in different studies. Such differences may occur as a result of differences in the populations studied, the outcome measures used or because of different definitions of the variables involved. It is the opposite of homogeneity.
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.
Inclusion criteria (literature review)	Explicit criteria used to decide which studies should be considered as potential sources of evidence.
Incremental analysis	The analysis of additional costs and additional clinical outcomes with different interventions.
Incremental cost	The extra cost linked to using one test or treatment rather than another. Or the additional cost of doing a test or providing a treatment more frequently.
Incremental cost-effectiveness ratio (ICER)	The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another.
Incremental net benefit (INB)	The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the INB is calculated as: $(£20,000 \times \text{QALYs gained}) - \text{Incremental cost}$ .
Indirectness	The available evidence is different to the review question being addressed, in terms of PICO (population, intervention, comparison and outcome).
Intention-to-treat analysis (ITT)	An assessment of the people taking part in a clinical trial, based on the group they were initially (and randomly) allocated to. This is regardless of whether or not they dropped out, fully complied with the treatment or switched to an alternative treatment. Intention-to-treat analyses are often used to assess clinical effectiveness because they mirror actual

Term	Definition
	practice: that is, not everyone complies with treatment and the treatment people receive may be changed according to how they respond to it.
Intervention	In medical terms this could be a drug treatment, surgical procedure, diagnostic or psychological therapy. Examples of public health interventions could include action to help someone to be physically active or to eat a more healthy diet.
Intraoperative	The period of time during a surgical procedure.
Kappa statistic	A statistical measure of inter-rater agreement that takes into account the agreement occurring by chance.
Length of stay	The total number of days a participant stays in hospital.
Licence	See 'Product licence'.
Life years gained	Mean average years of life gained per person as a result of the intervention compared with an alternative intervention.
Long-term care	Residential care in a home that may include skilled nursing care and help with everyday activities. This includes nursing homes and residential homes.
Logistic regression or Logit model	In statistics, logistic regression is a type of analysis used for predicting the outcome of a binary dependent variable based on one or more predictor variables. It can be used to estimate the log of the odds (known as the 'logit').
Loss to follow-up	A patient, or the proportion of patients, actively participating in a clinical trial at the beginning, but whom the researchers were unable to trace or contact by the point of follow-up in the trial
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
Meta-analysis	A method often used in systematic reviews. Results from several studies of the same test or treatment are combined to estimate the overall effect of the treatment.
Multivariate model	A statistical model for analysis of the relationship between 2 or more predictor (independent) variables and the outcome (dependent) variable.
Net monetary benefit (NMB)	The value in monetary terms of an intervention net of its cost. The NMB can be calculated for a given cost-effectiveness threshold. If the threshold is £20,000 per QALY gained then the NMB for an intervention is calculated as: $(£20,000 \times \text{mean QALYs}) - \text{mean cost}$ . The most preferable option (that is, the most clinically effective option to have an ICER below the threshold selected) will be the treatment with the highest NMB.
Non-randomised intervention study	A quantitative study investigating the effectiveness of an intervention that does not use randomisation to allocate patients (or units) to treatment groups. Non-randomised studies include observational studies, where allocation to groups occurs through usual treatment decisions or people's preferences. Non-randomised studies can also be experimental, where the investigator has some degree of control over the allocation of treatments.  Non-randomised intervention studies can use a number of different study designs, and include cohort studies, case-control studies, controlled before-and-after studies, interrupted-time-series studies and quasi-randomised controlled trials.

Term	Definition
Number needed to treat (NNT)	<p>The average number of patients who need to be treated to get a positive outcome. For example, if the NNT is 4, then 4 patients would have to be treated to ensure 1 of them gets better. The closer the NNT is to 1, the better the treatment.</p> <p>For example, if you give a stroke prevention drug to 20 people before 1 stroke is prevented, the number needed to treat is 20. See also number needed to harm, absolute risk reduction.</p>
Observational study	<p>Individuals or groups are observed or certain factors are measured. No attempt is made to affect the outcome. For example, an observational study of a disease or treatment would allow 'nature' or usual medical care to take its course. Changes or differences in one characteristic (for example, whether or not people received a specific treatment or intervention) are studied without intervening.</p> <p>There is a greater risk of selection bias than in experimental studies.</p>
Odds ratio	<p>A measure of treatment effectiveness. The odds of an event happening in the treatment group, expressed as a proportion of the odds of it happening in the control group. The 'odds' is the ratio of events to non-events.</p>
Opportunity cost	<p>The loss of other healthcare programmes displaced by investment in or introduction of another intervention. This may be best measured by the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.</p>
Outcome	<p>The impact that a test, treatment, policy, programme or other intervention has on a person, group or population. Outcomes from interventions to improve the public's health could include changes in knowledge and behaviour related to health, societal changes (for example, a reduction in crime rates) and a change in people's health and wellbeing or health status. In clinical terms, outcomes could include the number of patients who fully recover from an illness or the number of hospital admissions, and an improvement or deterioration in someone's health, functional ability, symptoms or situation.</p> <p>Researchers should decide what outcomes to measure before a study begins.</p>
P value	<p>The p value is a statistical measure that indicates whether or not an effect is statistically significant.</p> <p>For example, if a study comparing 2 treatments found that one seems more effective than the other, the p value is the probability of obtaining these, or more extreme results by chance. By convention, if the p value is below 0.05 (that is, there is less than a 5% probability that the results occurred by chance) it is considered that there probably is a real difference between treatments. If the p value is 0.001 or less (less than a 1% probability that the results occurred by chance), the result is seen as highly significant.</p> <p>If the p value shows that there is likely to be a difference between treatments, the confidence interval describes how big the difference in effect might be.</p>
Perioperative	<p>The period from admission through surgery until discharge, encompassing the preoperative and postoperative periods.</p>
Placebo	<p>A fake (or dummy) treatment given to participants in the control group of a clinical trial. It is indistinguishable from the actual treatment (which is given to participants in the experimental group). The aim is to determine what effect the experimental treatment has had – over and</p>

Term	Definition
	above any placebo effect caused because someone has received (or thinks they have received) care or attention.
Polypharmacy	The use or prescription of multiple medications.
Posterior distribution	In Bayesian statistics this is the probability distribution for a statistic based after combining established information or belief (the prior) with new evidence (the likelihood).
Postoperative	Pertaining to the period after patients leave the operating theatre, following surgery.
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Preoperative	The period before surgery commences.
Prior distribution	In Bayesian statistics this is the probability distribution for a statistic based on previous evidence or belief.
Primary care	Healthcare delivered outside hospitals. It includes a range of services provided by GPs, nurses, health visitors, midwives and other healthcare professionals and allied health professionals such as dentists, pharmacists and opticians.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Probabilistic analysis	In economic evaluation, this is an analysis that uses a probability distribution for each input. In contrast, see Deterministic analysis.
Product licence	An authorisation from the MHRA to market a medicinal product.
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good prognosis is associated with low rate of undesirable outcomes; poor prognosis is associated with a high rate of undesirable outcomes.
Prospective study	A research study in which the health or other characteristic of participants is monitored (or 'followed up') for a period of time, with events recorded as they happen. This contrasts with retrospective studies.
Publication bias	Publication bias occurs when researchers publish the results of studies showing that a treatment works well and don't publish those showing it did not have any effect. If this happens, analysis of the published results will not give an accurate idea of how well the treatment works. This type of bias can be assessed by a funnel plot.
Quality of life	See 'Health-related quality of life'.
Quality-adjusted life year (QALY)	A measure of the state of health of a person or group in which the benefits, in terms of length of life, are adjusted to reflect the quality of life. One QALY is equal to 1 year of life in perfect health. QALYS are calculated by estimating the years of life remaining for a patient following a particular treatment or intervention and weighting each year with a quality of life score (on a scale of 0 to 1). It is often measured in terms of the person's ability to perform the activities of daily life, freedom from pain and mental disturbance.
Randomisation	Assigning participants in a research study to different groups without taking any similarities or differences between them into account. For example, it could involve using a random numbers table or a computer-generated random sequence. It means that each individual

Term	Definition
	(or each group in the case of cluster randomisation) has the same chance of receiving each intervention.
Randomised controlled trial (RCT)	A study in which a number of similar people are randomly assigned to 2 (or more) groups to test a specific drug or treatment. One group (the experimental group) receives the treatment being tested, the other (the comparison or control group) receives an alternative treatment, a dummy treatment (placebo) or no treatment at all. The groups are followed up to see how effective the experimental treatment was. Outcomes are measured at specific times and any difference in response between the groups is assessed statistically. This method is also used to reduce bias.
RCT	See 'Randomised controlled trial'.
Reporting bias	See 'Publication bias'.
Resource implication	The likely impact in terms of finance, workforce or other NHS resources.
Retrospective study	A research study that focuses on the past and present. The study examines past exposure to suspected risk factors for the disease or condition. Unlike prospective studies, it does not cover events that occur after the study group is selected.
Review question	In guideline development, this term refers to the questions about treatment and care that are formulated to guide the development of evidence-based recommendations.
Risk ratio (RR)	The ratio of the risk of disease or death among those exposed to certain conditions compared with the risk for those who are not exposed to the same conditions (for example, the risk of people who smoke getting lung cancer compared with the risk for people who do not smoke). If both groups face the same level of risk, the risk ratio is 1. If the first group had a risk ratio of 2, subjects in that group would be twice as likely to have the event happen. A risk ratio of less than 1 means the outcome is less likely in the first group. The risk ratio is sometimes referred to as relative risk.
Secondary outcome	An outcome used to evaluate additional effects of the intervention deemed a priori as being less important than the primary outcomes.
Selection bias	Selection bias occurs if: a) The characteristics of the people selected for a study differ from the wider population from which they have been drawn, or b) There are differences between groups of participants in a study in terms of how likely they are to get better.
Sensitivity analysis	A means of representing uncertainty in the results of economic evaluations. Uncertainty may arise from missing data, imprecise estimates or methodological controversy. Sensitivity analysis also allows for exploring the generalisability of results to other settings. The analysis is repeated using different assumptions to examine the effect on the results. One-way simple sensitivity analysis (univariate analysis): each parameter is varied individually in order to isolate the consequences of each parameter on the results of the study. Multi-way simple sensitivity analysis (scenario analysis): 2 or more parameters are varied at the same time and the overall effect on the results is evaluated.



Term	Definition
	<p>Threshold sensitivity analysis: the critical value of parameters above or below which the conclusions of the study will change are identified.</p> <p>Probabilistic sensitivity analysis: probability distributions are assigned to the uncertain parameters and are incorporated into evaluation models based on decision analytical techniques (for example, Monte Carlo simulation).</p>
Significance (statistical)	A result is deemed statistically significant if the probability of the result occurring by chance is less than 1 in 20 ( $p < 0.05$ ).
Stakeholder	<p>An organisation with an interest in a topic that NICE is developing a guideline or piece of public health guidance on. Organisations that register as stakeholders can comment on the draft scope and the draft guidance. Stakeholders may be:</p> <ul style="list-style-type: none"> <li>• manufacturers of drugs or equipment</li> <li>• national patient and carer organisations</li> <li>• NHS organisations</li> <li>• organisations representing healthcare professionals.</li> </ul>
State transition model	See Markov model
Stratification	When a different estimate effect is thought to underlie two or more groups based on the PICO characteristics. The groups are therefore kept separate from the outset and are not combined in a meta-analysis, for example; children and adults. Specified a priori in the protocol.
Sub-groups	Planned statistical investigations if heterogeneity is found in the meta-analysis. Specified a priori in the protocol.
Systematic review	A review in which evidence from scientific studies has been identified, appraised and synthesised in a methodical way according to predetermined criteria. It may include a meta-analysis.
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Transition probability	In a state transition model (Markov model), this is the probability of moving from one health state to another over a specific period of time.
Treatment allocation	Assigning a participant to a particular arm of a trial.
Univariate	Analysis which separately explores each variable in a data set.
Utility	In health economics, a 'utility' is the measure of the preference or value that an individual or society places upon a particular health state. It is generally a number between 0 (representing death) and 1 (perfect health). The most widely used measure of benefit in cost–utility analysis is the quality-adjusted life year, but other measures include disability-adjusted life years (DALYs) and healthy year equivalents (HYES).

## References

1. GRADE Working Group. The Grading of Recommendations Assessment, Development and Evaluation (GRADE) Working Group website. 2011. Available from: <http://www.gradeworkinggroup.org/> Last accessed:
2. Maruish M, Maruish M, Kosinski M, Bjorner JB, Gandek B, Turner-Bowker D et al. User's manual for the SF36v2 Health Survey. 2011.
3. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *Journal of Clinical Epidemiology*. 2016; 75:40-46
4. National Institute for Health and Care Excellence. Developing NICE guidelines: the manual [updated October 2020]. London. National Institute for Health and Care Excellence, 2014. Available from: <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20oerview>
5. National Institute for Health and Care Excellence. The NICE Charter. 2020. Available from: <https://www.nice.org.uk/about/who-we-are/our-charter> Last accessed: 10/02/2022.
6. National Institute for Health and Clinical Excellence. Guide to the methods of technology appraisal 2013. London. National Institute for Health and Clinical Excellence, 2013. Available from: <http://publications.nice.org.uk/pmg9>
7. National Institute for Health and Clinical Excellence. Social value judgements: principles for the development of NICE guidance. London. National Institute for Health and Clinical Excellence, 2008. Available from: <https://www.nice.org.uk/media/default/about/what-we-do/research-and-development/social-value-judgements-principles-for-the-development-of-nice-guidance.pdf>
8. Organisation for Economic Co-operation and Development (OECD). Purchasing power parities (PPP). 2021. Available from: <http://www.oecd.org/std/ppp> Last accessed: 10/02/2022.
9. Review Manager (RevMan) [Computer program]. Version 5. Copenhagen. The Nordic Cochrane Centre, The Cochrane Collaboration, 2015. Available from: <http://tech.cochrane.org/Revman>