

Stroke rehabilitation in adults (update)

NICE guideline: methods

NICE guideline GID-NG10175

Methods

April 2023

Draft for Consultation

*Evidence reviews were developed by the
Guideline Development Team at NICE*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE, 2023. All rights reserved. Subject to Notice of rights.

Contents

1	Development of the guideline	5
1.1	Remit.....	5
2	Methods	6
2.1	Developing the review questions and outcomes.....	6
2.1.1	Stratification	14
2.2	Searching for evidence.....	14
2.2.1	Clinical and health economics literature searches.....	14
2.3	Reviewing evidence	15
2.3.1	Types of studies and inclusion and exclusion criteria	15
2.4	Methods of combining evidence	17
2.4.1	Data synthesis for intervention reviews	17
2.4.2	Data synthesis for diagnostic reviews	18
2.4.3	Data synthesis for tool validity and reliability reviews	19
2.4.4	Data synthesis for qualitative reviews	19
2.4.5	Data synthesis for mixed methods reviews	19
2.5	Appraising the quality of evidence by outcomes.....	20
2.5.1	Intervention reviews	20
2.5.2	Diagnostic reviews.....	27
2.5.3	Tool validity and reliability reviews	28
2.5.4	Qualitative reviews.....	29
2.5.5	Mixed methods reviews.....	31
2.6	Assessing clinical importance.....	31
2.7	Identifying and analysing evidence of cost effectiveness.....	32
2.7.1	Literature review	32
2.7.2	Undertaking new health economic analysis.....	34
2.7.3	Cost-effectiveness criteria.....	35
2.7.4	In the absence of health economic evidence.....	35
2.8	Developing recommendations	36
2.8.1	Research recommendations	37
2.8.2	Validation process.....	37
2.8.3	Updating the guideline	37
2.9	General terms	37
	References.....	50

1 Development of the guideline

2 1.1 Remit

3 NICE received the remit for this guideline from NHS England.

4 The remit for this guideline is:

5 Stroke rehabilitation in adults

6 To see “What this guideline covers” and “What this guideline does not cover” please
7 see the Stroke rehabilitation in adults **scope**.

2 Methods

This guideline was developed using the methods described in the NICE guidelines manual⁴ as outlined in Table 1 below.

Table 1 Versions of the NICE guidelines manual followed during guideline development and guideline validation

Stage	2020 update	2022 update
Scoping	Y	
Development	Y	Y
Validation		Y

Declarations of interest were recorded according to the NICE conflicts of interest policy.

Sections 2.1 to 2.3 describe the process used to identify and review evidence. Sections 2.1.1 and 2.7 describe the process used to identify and review the health economic evidence.

2.1 Developing the review questions and outcomes

The review questions developed for this guideline were based on the key areas and draft review questions identified in the guideline scope. They were drafted by the technical team, refined and validated by the committee and signed off by NICE. A total of 17 review questions were developed in this guideline and outlined in Table 2.

The review questions were based on the following frameworks:

- population, intervention, comparator and outcome (PICO) for reviews of interventions (including test and treat and tool validity and reliability)
- population, index tests, reference standard and target condition for reviews of diagnostic test accuracy
- population, setting and context for qualitative reviews
- population, intervention, comparator and outcome (PICO) and population, setting and context for mixed methods reviews.

This use of a framework informed a more detailed protocol that guided the literature searching process, critical appraisal and synthesis of evidence, and facilitated the development of recommendations by the guideline committee. Full literature searches, critical appraisals and evidence reviews were completed for all the specified review questions.

Table 2: Review questions

Evidence report	Type of review	Review questions	Outcomes
A	Mixed methods	In people after stroke what is the clinical and cost effectiveness of early supported discharge compared with usual care? In people after stroke what factors are associated with effective	Quantitative outcomes (qualitative themes were extracted from the qualitative studies) <ul style="list-style-type: none"> • Mortality • Person/participant generic health-related quality of life

Evidence report	Type of review	Review questions	Outcomes
		delivery of early supported discharge care?	<ul style="list-style-type: none"> • Carer generic health-related quality of life • Physical dependency • Activities of daily living • Extended activities of daily living • Length of hospital stay • Caregiver strain index • Falls • Readmissions to hospital • Psychological distress/mood • Stroke-specific Patient-Reported Outcome Measures
B	Assess-and-treat and tool validity and reliability	In people after stroke, what is the optimal tool for assessment of fatigue?	<p>Assess-and-treat:</p> <ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Activities of daily living • Psychological distress (depression, anxiety, distress) • Stroke-specific Patient-Reported Outcome Measures • Participation in leisure activities/social groups scores • Withdrawal due to adverse events <p>Tool validity and reliability:</p> <ul style="list-style-type: none"> • Face/content/construct validity • Criterion/concurrent validity • Discriminant/convergent validity • Test-retest reliability • Internal consistency (intertest reliability, intratest reliability) • Inter-rater reliability • Responsiveness to change • Dimensions of fatigue considered
C	Intervention	What is the clinical and cost effectiveness of routine specialist orthoptist assessment for people after stroke?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life

Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Delayed diagnosis • Vision-related quality of life • Additional health care contacts • Hospitalisation • Activities of daily living • Stroke-specific Patient-Reported Outcome Measures
D	Diagnostic	In people after stroke, what is the optimal tool for assessment of hearing?	<p>Test-and-treat outcomes:</p> <ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Activities of daily living • Participation in leisure activities/social groups scores • Psychological distress (depression, anxiety, distress) • Stroke-related scales of cognition (non-spatial attention and working memory, spatial attention, memory, executive function) • Speech perception • Functional communication • Stroke-specific Patient-Reported Outcome Measures • Withdrawal due to adverse events <p>Test accuracy outcomes:</p> <ul style="list-style-type: none"> • Sensitivity • Specificity • Raw data to calculate 2x2 tables to calculate sensitivity and specificity • Area under the curve • Likelihood ratios • Positive predictive values • Negative predictive values • Intra-test and inter-test reliability
E	Mixed methods	In people after stroke, what is the clinical and cost effectiveness of more intensive rehabilitation	Quantitative outcomes (qualitative themes were extracted from the qualitative studies)

Evidence report	Type of review	Review questions	Outcomes
		<p>compared with standard rehabilitation?</p> <p>In people after stroke what factors are associated with effective delivery of more intensive rehabilitation?</p>	<ul style="list-style-type: none"> • Person/participant health-related quality of life • Carer generic health-related quality of life • Stroke outcome – modified Rankin scale • Activities of daily living • Physical function (upper limb, lower limb) • Communication – overall language ability • Communication – impairment specific measures (naming, auditory comprehension, reading, expressive language, dysarthria speech impairment) • Communication – functional communication • Psychological distress – depression • Stroke related scale of cognition (non-spatial attention and working memory, spatial attention, memory, executive function) • Swallow function and ability • Discontinuation from study
F	Intervention	In people after stroke, what is the clinical and cost effectiveness of self-care management and/or supported self-care management compared with usual rehabilitation?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Self efficacy • Activities of daily living • Participation restrictions • Psychological distress – depression • Stroke-specific Patient-Reported Outcome Measures • Health service usage (hospital readmissions, general practitioner attendance, emergency department visits) • Participant satisfaction • Adverse events (type and frequency)

Evidence report	Type of review	Review questions	Outcomes
G	Intervention	In people after stroke, what is the clinical and cost effectiveness of telerehabilitation compared with standard rehabilitation and as an adjunct to standard rehabilitation?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Activities of daily living • Mobility • Balance • Psychological distress – depression • Physical function – upper limb • Stroke-specific measures of cognition (non-spatial attention and working memory, spatial attention, memory, executive functions) • Swallow function and ability • Functional communication • Stroke-specific Patient-Reported Outcome Measures • Withdrawal due to adverse events
H	Intervention	In people after stroke, what is the clinical and cost effectiveness of community participation interventions compared with no intervention?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Return to work • Wellbeing score • Participation in leisure activities/social groups scores • Psychological distress (depression, anxiety, distress) • Activities of daily living • Discontinuation
I	Intervention	In people after stroke, what is the clinical and cost effectiveness of interventions for oral feeding compared with alternative interventions or usual care to reduce difficulties with oral feeding?	<ul style="list-style-type: none"> • Mortality • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Occurrence of chest infections • Occurrence of aspiration • Dysphagia present/return to normal diet

Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Discharge to residential service • Length of hospital stay • Re-admission • Swallowing ability • Nutrition • Hydration
J	Intervention	In people after stroke, what is the clinical and cost effectiveness of interventions to improve oral hygiene?	<ul style="list-style-type: none"> • Mortality • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Occurrence of pneumonia • Stroke outcome – modified Rankin scale • Requirement for enteral feeding support • Oral health outcome scales • Dysphagia severity • Presence of oral disease • Length of hospital stay • Re-admission • Stroke-specific Patient-Reported Outcome Measures
K	Intervention	In people with aphasia after stroke, what is the clinical and cost effectiveness of computer-based tools to augment speech and language therapy?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Communication – overall language ability • Communication – impairment specific measures (naming, auditory comprehension, reading, expressive language, dysarthria speech impairment and activity) • Communication – functional communication • Communication related quality of life • Psychological distress (depression, anxiety, distress) • Discontinuation
L	Intervention	In people after stroke, what is the clinical and cost effectiveness of group training to improve walking?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life

Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> • Carer generic health-related quality of life • 6-minute walk test • Walking speed • Functional mobility measures • Measures of standing balance • Measures of motor impairment • Activities of daily living • Stroke-specific Patient-Reported Outcome Measures • Length of hospital stay • Adverse event
M	Intervention	In people after stroke, what is the clinical and cost effectiveness of robot-assisted arm training to improve function and reduce disability?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Activities of daily living • Arm function • Arm muscle strength • Spasticity • Stroke-specific Patient-Reported Outcome Measures • Withdrawal for any reason • Adverse events (cardiovascular events, injuries and pain, other reported adverse events)
N	Intervention	In people after stroke, what is the clinical and cost effectiveness of music therapy to improve mood and activities of daily living?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Activities of daily living • Psychological distress (depression, anxiety, distress) • Wellbeing scores • Participation in leisure activities/social groups scores • Withdrawal due to adverse events
O	Intervention	In people with shoulder pain after stroke, what is the clinical and cost effectiveness of transcutaneous	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life

Evidence report	Type of review	Review questions	Outcomes
		electrical nerve stimulation, acupuncture, functional electrical stimulation and intra-articular steroid injection in reducing pain?	<ul style="list-style-type: none"> • Carer generic health-related quality of life • Pain • Physical function – upper limb • Activities of daily living • Stroke-specific Patient-Reported Outcome Measures • Withdrawal due to adverse events
P	Intervention	In people after stroke, what is the clinical and cost effectiveness of interventions (for example: oral baclofen, intrathecal baclofen, acupuncture and TENS [transcutaneous electrical nerve stimulation]), in reducing spasticity?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Spasticity outcome measures • Physical function (general, upper limb, lower limb) • Pain • Activities of daily living • Additional health care contacts • Hospitalisation • Stroke outcome – modified Rankin scale • Withdrawal due to adverse events
Q	Intervention	In people after stroke, what is the clinical and cost effectiveness of mirror therapy to improve motor function, visuospatial function and activities of daily living?	<ul style="list-style-type: none"> • Person/participant generic health-related quality of life • Carer generic health-related quality of life • Upper limb and hand motor function • Lower limb motor function • Global motor function • Fugl-Meyer Assessment Upper Extremity • Measures of motor impairment (upper limb and lower limb) • Activities of daily living) • Pain • Visuospatial neglect • Stroke-specific Patient-Reported Outcome Measures • Adverse events

Evidence report	Type of review	Review questions	Outcomes
			• Dropout rate

1

2 2.1.1 Stratification

3 The following questions included specific stratifications:

- 4 • In the question for report E, analyses were stratified by whether people had
- 5 communication difficulties or not.
- 6 • In the question for report P, analyses were stratified by the site of spasticity
- 7 (including focal, multifocal, segmental, generalised and mixed spasticity).

8 Where studies reported a mix of populations across strata, a threshold of 80% was
9 agreed with the committee as a cut off for what would be acceptable to constitute a
10 predominant group.

11 2.2 Searching for evidence

12 2.2.1 Clinical and health economics literature searches

13 The full strategy including population terms, intervention terms, study types applied,
14 the databases searched, and the years covered can be found in Appendix B of the
15 evidence review.

16 Systematic literature searches were undertaken to identify published clinical and
17 health economic evidence relevant to the review questions. These were run
18 according to the parameters as stipulated within the NICE guideline's manual,
19 <https://www.nice.org.uk/process/pmg20/chapter/identifying-the-evidence-literature-searching-and-evidence-submission>.

21 Databases were searched using relevant medical subject headings, free-text terms
22 and where appropriate study-type filters. Studies published in languages other than
23 English were not reviewed, and where possible, searches were restricted to English
24 language. Searches were updated on 08 January 2023 Papers published or added to
25 databases after this date were not considered. Where new evidence was identified,
26 for example in consultation comments received from stakeholders, the impact on the
27 guideline was considered, and the action agreed between the technical team and
28 NICE staff with a quality assurance role.

29 Searches were quality assured using different approaches prior to being run. Medline
30 search strategies were peer reviewed by a second information specialist using a QA
31 process based on the PRESS checklist³. Key (seed) papers if provided, were
32 checked if retrieved by the search.

33 Searching for unpublished literature was not undertaken.

34 Additional studies were added to the evidence base these consisted of references
35 included in relevant systematic reviews, and those highlighted by committee
36 members.

2.3 Reviewing evidence

The evidence for each review question was reviewed using the following process:

- Potentially relevant studies were identified from the search results by reviewing titles and abstracts. The full papers were then obtained.
 - Full papers were evaluated against the pre-specified inclusion and exclusion criteria set out in the protocol to identify studies that addressed the review question. The review protocols are included in an appendix to each of the evidence reports.
 - Relevant studies were critically appraised using the preferred study design checklist as specified in the NICE guidelines manual.⁴ The checklist used is included in the individual review protocols in each of the evidence reports.
 - Key information was extracted about intervention and qualitative study methods and results into EPPI reviewer version 5. Summary evidence tables were produced from data entered into EPPI Reviewer, including critical appraisal ratings. Key information about diagnostic and tool validity and reliability study methods and results were manually extracted into standard Word evidence tables (evidence tables are included in an appendix to each of the evidence reports).
 - Summaries of the evidence were generated by outcome. Outcome data were combined, analysed and reported according to study design:
 - Randomised data were meta-analysed where appropriate and reported in GRADE evidence profiles.
 - Diagnostic data were meta-analysed where appropriate or presented as a range of values in GRADE evidence profiles.
 - Tool validity and reliability data was combined where appropriate and presented in matrices.
 - Qualitative data were synthesised across studies using thematic analysis and presented as summary statements in GRADE CERQual tables.
 - Mixed methods data was treated separately as randomised and qualitative data (and presented as is appropriate for each) and then synthesised together where the results are presented in matrices and through a narrative summary.
 - A minimum of 10% of the abstracts were reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer.
 - All of the evidence reviews were quality assured by a senior systematic reviewer. This included checking:
 - papers were included or excluded appropriately
 - a sample of the data extractions
 - a sample of the risk of bias assessments
 - correct methods were used to synthesise data.
- Discrepancies will be identified and resolved through discussion (with a third reviewer where necessary).

2.3.1 Types of studies and inclusion and exclusion criteria

The inclusion and exclusion of studies was based on the criteria defined in the review protocols, which can be found in an appendix to each of the evidence reports. Excluded studies (with the reasons for their exclusion) are listed in an appendix to

1 each of the evidence reports. The committee was consulted about any uncertainty
2 regarding inclusion or exclusion.

3 Conference abstracts were not generally considered for inclusion. If abstracts were
4 included the authors were contacted for further information. Literature reviews,
5 posters, letters, editorials, comment articles, unpublished studies and studies not in
6 published in English language were excluded.

7 The exception for this were studies already included in Cochrane reviews that were
8 included in the reviews (Cochrane reviews were included for reviews A, F, G, I, L, M
9 and Q). If the authors of any of the included Cochrane reviews had included any of
10 the preceding material (for example: conference abstracts, translated studies,
11 unpublished studies) then this would still be included as further information was
12 obtained by the authors of the Cochrane reviews that allowed the study to be used in
13 the review.

14 **2.3.1.1 Type of studies**

15 Randomised controlled trials and observational studies (including diagnostic or tool
16 validity and reliability studies) were included in the evidence reviews as appropriate.

17 For intervention reviews, randomised controlled trials (RCTs) were included where
18 identified as because they are considered the most robust type of study design that
19 can produce an unbiased estimate of the intervention effects. Non-randomised
20 intervention studies were considered appropriate for inclusion if there was insufficient
21 randomised evidence for the committee to make a decision. In this case the
22 committee stated a priori in the protocol that either certain identified variables must
23 be equivalent at baseline or else the analysis had to adjust for any baseline
24 differences. If the study did not fulfil either criterion it was excluded. Refer to the
25 review protocols in each evidence report for full details on the study design of studies
26 that were appropriate for each review question.

27 For diagnostic review questions, diagnostic RCTs, cross-sectional studies and
28 retrospective studies were included. Case–control studies were not included. For tool
29 validity and reliability review questions, test and treat RCTs, cohort studies and
30 cross-sectional studies were included.

31 Systematic reviews and meta-analyses conducted to the same methodological
32 standards as the NICE reviews were included within the evidence reviews in
33 preference to primary studies, where they were available and applicable to the review
34 questions and updated or added to where appropriate to the guideline review
35 question. Individual patient data (IPD) meta-analyses were preferentially included if
36 meeting the protocol and methodological criteria.

37 **2.3.1.1.1 Qualitative studies**

38 In the qualitative reviews, studies using focus groups, or structured or semi-
39 structured interviews were considered for inclusion. Survey data or other types of
40 questionnaires were only included if they provided analysis from open-ended
41 questions, but not if they reported descriptive quantitative data only.

42 **Saturation of qualitative studies**

43 Data extraction in qualitative reviews is a thorough process. A common approach
44 applied in systematic reviews of qualitative data is to stop extracting data once

1 saturation has been reached. In an exploratory review, where themes are not
2 predefined in the protocol, thematic or data extraction may be applied. For the
3 purposes of this review, extraction of information from relevant studies was stopped
4 when data saturation was reached, i.e. no new information was emerging for a
5 specific theme. This includes; studies that don't report any new themes additional to
6 those already identified in the review as well as not contributing additional information
7 to the existing themes, as well as studies which report a new theme but data from
8 other themes in the study doesn't contribute to the existing review themes. In the
9 latter scenario only the new theme data is extracted. These studies are not
10 specifically excluded from the review as they nevertheless fit the criteria defined in
11 the review protocol. Any studies for which data were not extracted due to data
12 saturation having been reached, but that fit the inclusion criteria of the protocol, were
13 listed in the table for studies 'identified but not extracted due to saturation' in an
14 appendix to the qualitative evidence review.

15 **2.4 Methods of combining evidence**

16 **2.4.1 Data synthesis for intervention reviews**

17 Meta-analyses were conducted using Cochrane Review Manager (RevMan5)¹¹
18 software

19 **2.4.1.1 Analysis of different types of data**

20 ***Dichotomous outcomes***

21 Fixed-effects (Mantel–Haenszel) techniques were used to calculate risk ratios
22 (relative risk, RR) for the binary outcomes. The absolute risk difference was also
23 calculated using GRADEpro¹ software, using the median event rate in the control arm
24 of the pooled results.

25 For binary variables where there were zero events in either arm or a less than 1%
26 event rate, Peto odds ratios, rather than risk ratios, were calculated as they are more
27 appropriate for data with a low number of events. Where there are zero events in
28 both arms, the risk difference was calculated and reported instead.

29 **Continuous outcomes**

30 Continuous outcomes were analysed using an inverse variance method for pooling
31 weighted mean differences.

32 Where the studies within a single meta-analysis had different scales of measurement
33 for the same outcomes, standardised mean differences were used (providing all
34 studies reported either change from baseline or final values rather than a mixture of
35 both); each different measure in each study was 'normalised' to the standard
36 deviation value pooled between the intervention and comparator groups in that same
37 study.

38 The means and standard deviations of continuous outcomes are required for meta-
39 analysis. However, in cases where standard deviations were not reported, the
40 standard error was calculated if the p values or 95% confidence intervals (95% CI)
41 were reported, and meta-analysis was undertaken with the mean and standard error
42 using the generic inverse variance method in RevMan5¹¹.

1 **Generic inverse variance**

2 If a study reported only the summary statistic and 95% CI the generic-inverse
3 variance method was used to enter data into RevMan5.¹¹ If the control event rate was
4 reported this was used to generate the absolute risk difference in GRADEpro.¹ If
5 multivariate analysis was used to derive the summary statistic but no adjusted control
6 event rate was reported no absolute risk difference was calculated.

7 **Complex analysis**

8 Where studies had used a crossover design, paired continuous data were extracted
9 where possible, and forest plots were generated in RevMan5¹¹ with the generic
10 inverse variance function. When a crossover study had categorical data and the
11 number of subjects with an event in both interventions was known, the standard error
12 (of the log of the risk ratio) was calculated using the simplified Mantel–Haenszel
13 method for paired outcomes. Forest plots were also generated in RevMan5¹¹ with the
14 generic inverse variance function. If paired continuous or categorical data were not
15 available from the crossover studies, the separate group data were analysed in the
16 same way as data from parallel groups, on the basis that this approach would
17 overestimate the confidence intervals and thus artificially reduce study weighting
18 resulting in a conservative effect. Where a meta-analysis included a mixture of
19 studies using both paired and parallel group approaches, all data were entered into
20 RevMan5¹¹ using the generic inverse variance function.

21 **2.4.2 Data synthesis for diagnostic reviews**

22 Two separate sections in the review protocol were produced to reflect the 2 different
23 diagnostic study designs.

24 **2.4.2.1 Diagnostic RCTs**

25 Diagnostic RCTs (sometimes referred to as test and treat trials) are a randomised
26 comparison of 2 diagnostic tests, with study outcomes being clinically important
27 consequences of the diagnosis (patient-related outcome measures similar to those in
28 intervention trials, such as mortality). Patients are randomised to receive test A or
29 test B, followed by identical therapeutic interventions based on the results of the test
30 (so someone with a positive result would receive the same treatment regardless of
31 whether they were diagnosed by test A or test B). Downstream patient outcomes are
32 then compared between the 2 groups. As treatment is the same in both arms of the
33 trial, any differences in patient outcomes will reflect the accuracy of the tests in
34 correctly establishing who does and does not have the condition. Data were
35 synthesised using the same methods for intervention reviews (see section 2.4.1.1).

36 **2.4.2.2 Diagnostic accuracy studies**

37 For diagnostic test accuracy studies, a positive result on the index test was found if
38 the person had values of the measured quantity above or below a threshold value,
39 and different thresholds could be used. The thresholds were pre-specified by the
40 committee including whether or not data could be pooled across a range of
41 thresholds. The threshold of a diagnostic test is defined as the value at which the test
42 can best differentiate between those with and without the target condition. In practice
43 this usually varies across studies. If a test has a high sensitivity then very few people
44 with the condition will be missed (few false negatives). For example, a test with a
45 sensitivity of 97% will only miss 3% of people with the condition. Conversely, if a test

1 has a high specificity then few people without the condition would be incorrectly
2 diagnosed (few false positives).

3 Coupled forest plots of the agreed primary paired outcome measure for decision
4 making (sensitivity and specificity) with their 95% CIs across studies (at various
5 thresholds) were produced for each test, using RevMan5.¹¹ In order to do this, 2 by 2
6 tables (the number of true positives, false positives, true negatives and false
7 negatives) were directly taken from the study if given, or else were derived from raw
8 data or calculated from the set of test accuracy statistics.

9 **2.4.3 Data synthesis for tool validity and reliability reviews**

10 Data about validity and reliability was extracted from the studies identified in the sift.
11 This data was collated and presented in a matrix to compare the different tools to
12 each other, to allow for a qualitative and quantitative assessment of the results.

13 Face, content and construct validity were interpreted qualitatively from information
14 provided in the studies. Criterion, concurrent, convergent and discriminant validity
15 were interpreted by correlation against other scales. In review B, there were no gold
16 standard tests that measure the phenomenon of interest, so other tools that
17 supposedly measure the phenomenon of interest were considered. The stronger the
18 correlation (the closer the value to 1) the more associated the scales were, which
19 was considered positive for criterion, concurrent and convergent validity and negative
20 for discriminant validity (for which values closer to 0 were considered a positive
21 result).

22 This approach was also used for assessing internal consistency, test-retest reliability
23 and inter-rater reliability. Where Cronbach's alpha and intraclass correlation
24 coefficients were reported, values closer to 1 were considered to be better values,
25 while values closer to 0 were considered to be worse. Values above 0.75 were
26 considered to be good results, though the committee weighed these values against
27 the other components while assessing the quality of the tools and did not assign an
28 absolute threshold that a tool had to reach in order to be acceptable for use.

29 **2.4.4 Data synthesis for qualitative reviews**

30 The main findings for each included paper were identified and thematic analysis
31 methods were used to synthesise this information into broad overarching themes
32 which were summarised into the main review findings. The evidence was presented
33 in the form of a narrative summary detailing the evidence from the relevant papers
34 and how this informed the overall review finding plus a statement on the level of
35 confidence for that review finding. Considerable limitations and issues around
36 relevance were listed. A summary evidence table with the succinct summary
37 statements for each review finding was produced including the associated quality
38 assessment.

39 **2.4.5 Data synthesis for mixed methods reviews**

40 Analysis of individual studies according to methods relevant to whether they are
41 intervention or qualitative studies took place. After this, mixed methods synthesis
42 took place. Integration included:

- 43 • Narrative summary of the two evidence types,

- Matrices allowing for identification of elements of the qualitative themes within the quantitative studies,
- Sensitivity analyses of the quantitative studies to identify if including studies that only account for the qualitative themes mentioned accounts for any heterogeneity seen in the quantitative analyses.

These results were summarised and presented in the report documents and relevant discussion sections.

2.5 Appraising the quality of evidence by outcomes

2.5.1 Intervention reviews

The evidence for outcomes from the included RCTs and, where appropriate, non-randomised intervention studies, were evaluated and presented using the 'Grading of Recommendations Assessment, Development and Evaluation (GRADE) toolbox' developed by the international GRADE working group (<http://www.gradeworkinggroup.org/>). The software (GRADEpro¹) developed by the GRADE working group was used to assess the quality of each outcome, taking into account individual study quality and the meta-analysis results.

Each outcome was first examined for each of the quality elements listed and defined in Table 3.

Table 3: Description of quality elements in GRADE for intervention studies

Quality element	Description
Risk of bias	Limitations in the study design and implementation may bias the estimates of the treatment effect. Major limitations in studies decrease the confidence in the estimate of the effect. Examples of such limitations are selection bias (often due to poor allocation concealment), performance and detection bias (often due to a lack of blinding of the patient, healthcare professional or assessor) and attrition bias (due to missing data causing systematic bias in the analysis).
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question.
Inconsistency	Inconsistency refers to an unexplained heterogeneity of effect estimates between studies in the same meta-analysis.
Imprecision	Results are imprecise when studies include relatively few patients and few events (or highly variable measures) and thus have wide confidence intervals around the estimate of the effect relative to clinically important thresholds. 95% confidence intervals denote the possible range of locations of the true population effect at a 95% probability, and so wide confidence intervals may denote a result that is consistent with conflicting interpretations (for example a result may be consistent with both clinical benefit AND clinical harm) and thus be imprecise.
Publication bias	Publication bias is a systematic underestimate or overestimate of the underlying beneficial or harmful effect due to the selective publication of studies. A closely related phenomenon is where some papers fail to report an outcome that is inconclusive, thus leading to an overestimate of the effectiveness of that outcome.
Other issues	Sometimes randomisation may not adequately lead to group equivalence of confounders, and if so this may lead to bias, which should be taken into account.

Quality element	Description
	Potential conflicts of interest, often caused by excessive pharmaceutical company involvement in the publication of a study, should also be noted.

1 Details of how the 4 main quality elements (risk of bias, indirectness, inconsistency
2 and imprecision) were appraised for each outcome are given below. Publication bias
3 was considered with the committee. There was no reason to suspect it was present
4 from the evidence available, so it was not explored with funnel plots.

5 2.5.1.1 Risk of bias

6 The main domains of bias for RCTs are listed in Table 4. Each outcome had its risk
7 of bias assessed within each study first using the appropriate checklist for the study
8 design (Cochrane RoB 2 for RCTs, or ROBINS-I for non-randomised studies or
9 ROBIS for systematic reviews). For each study, if there was no risk of bias in any
10 domain, the risk of bias was given a rating of 0; 'no serious risk of bias'. If there was
11 risk of bias in just 1 domain, the risk of bias was given a 'serious' rating of -1, but if
12 there was risk of bias in 2 or more domains the risk of bias was given a 'very serious'
13 rating of -2. An overall rating is calculated across all studies by taking into account
14 the weighting of studies according to study precision. For example if the most precise
15 studies tended to each have a score of -1 for that outcome, the overall score for that
16 outcome would tend towards -1.

17 **Table 4: Principle domains of bias in randomised controlled trials**

Limitation	Explanation
Selection bias (sequence generation and allocation concealment)	If those enrolling participants are aware of the group to which the next enrolled patient will be allocated, either because of a non-random sequence that is predictable, or because a truly random sequence was not concealed from the researcher, this may translate into systematic selection bias. This may occur if the researcher chooses not to recruit a participant into that specific group because of: <ul style="list-style-type: none"> • knowledge of that participant's likely prognostic characteristics, and • a desire for one group to do better than the other.
Performance and detection bias (lack of blinding)	Patients, caregivers, those adjudicating or recording outcomes, and data analysts should not be aware of the arm to which the participants are allocated. Knowledge of the group can influence: <ul style="list-style-type: none"> • the experience of the placebo effect • performance in outcome measures • the level of care and attention received, and • the methods of measurement or analysis all of which can contribute to systematic bias.
Attrition bias	Attrition bias results from an unaccounted for loss of data beyond a certain level (a differential of at least 10% between groups). Loss of data can occur when participants are compulsorily withdrawn from a group by the researchers (for example, when a per-protocol approach is used) or when participants do not attend assessment sessions. If the missing data are likely to be different from the data of those remaining in the groups, and there is a differential rate of such missing data from groups, systematic attrition bias may result.
Selective outcome reporting	Reporting of some outcomes and not others on the basis of the results can also lead to bias, as this may distort the overall impression of efficacy.
Other limitations	For example:

Limitation	Explanation
	<ul style="list-style-type: none"> • Stopping early for benefit observed in randomised trials, in particular in the absence of adequate stopping rules. • Use of unvalidated patient-reported outcome measures. • Lack of washout periods to avoid carry-over effects in crossover trials. • Recruitment bias in cluster-randomised trials.

1 2.5.1.2 Indirectness

2 Indirectness refers to the extent to which the populations, interventions, comparisons
3 and outcome measures are dissimilar to those defined in the inclusion criteria for the
4 reviews. Indirectness is important when these differences are expected to contribute
5 to a difference in effect size, or may affect the balance of harms and benefits
6 considered for an intervention. As for the risk of bias, each outcome had its
7 indirectness assessed within each study first. For each study, if there were no
8 sources of indirectness, indirectness was given a rating of 0. If there was indirectness
9 in just 1 source (for example in terms of population), indirectness was given a
10 'serious' rating of -1, but if there was indirectness in 2 or more sources (for example,
11 in terms of population and treatment) the indirectness was given a 'very serious'
12 rating of -2. An overall rating is calculated across all studies by taking into account
13 the weighting of studies according to study precision. For example, if the most
14 precise studies tended to have an indirectness score of -1 each for that outcome, the
15 overall score for that outcome would tend towards -1.

16 2.5.1.3 Inconsistency

17
18 Inconsistency refers to an unexplained heterogeneity of results for an outcome
19 across different studies. When estimates of the treatment effect across studies differ
20 widely, this suggests true differences in the underlying treatment effect, which may
21 be due to differences in populations, settings or doses. Statistical heterogeneity was
22 assessed for each meta-analysis estimate by an I-squared (I^2) inconsistency statistic.

23 Heterogeneity or inconsistency amongst studies was also visually inspected. Where
24 statistical heterogeneity as defined above was present or there was clear visual
25 heterogeneity not captured in the I^2 value predefined subgrouping of studies was
26 carried out according to the protocol. See the review protocols for the subgrouping
27 strategy.

28 When heterogeneity existed within an outcome ($I^2 > 50\%$), but no plausible
29 explanation could be found, the quality of evidence for that outcome was
30 downgraded. Inconsistency for that outcome was given a 'serious' score of -1 if the I^2
31 was 50–74%, and a 'very serious' score of -2 if the I^2 was 75% or more. If a visual
32 assessment indicated heterogeneity but the I^2 was under 50% then the analysis was
33 considered on a case by case basis. Where it appeared that imprecision was
34 masking possible heterogeneity (leading to a reduced I^2 value), then the outcome
35 was downgraded for inconsistency.

36 If inconsistency could be explained based on pre-specified subgroup analysis (that is,
37 each subgroup had an $I^2 < 50\%$) then each of the derived subgroups were presented
38 separately for that forest plot (providing at least 2 studies remained in each
39 subgroup). The committee took this into account and considered whether to make

1 separate recommendations based on the variation in effect across subgroups within
2 the same outcome. In such a situation the quality of evidence was not downgraded.

3 If all predefined strategies of subgrouping were unable to explain statistical
4 heterogeneity, then a random effects (DerSimonian and Laird) model was employed
5 to the entire group of studies in the meta-analysis. A random-effects model assumes
6 a distribution of populations, rather than a single population. This leads to a widening
7 of the confidence interval around the overall estimate. If, however, the committee
8 considered the heterogeneity was so large that meta-analysis was inappropriate,
9 then the results were not pooled and were described narratively.

10 2.5.1.4 Imprecision

11 The criteria applied for imprecision were based on the 95% CIs for the pooled
12 estimate of effect, and the minimal important differences (MID) for the outcome. The
13 MIDs are the threshold for appreciable benefits and harms, separated by a zone
14 either side of the line of no effect where there is assumed to be no clinically important
15 effect. If either end of the 95% CI of the overall estimate of effect crossed 1 of the
16 MID lines, imprecision was regarded as serious and a 'serious' score of -1 was
17 given. This was because the overall result, as represented by the span of the
18 confidence interval, was consistent with 2 interpretations as defined by the MID (for
19 example, both no clinically important effect and clinical benefit were possible
20 interpretations). If both MID lines were crossed by either or both ends of the 95% CI
21 then imprecision was regarded as very serious and a 'very serious' score of -2 was
22 given. This was because the overall result was consistent with all 3 interpretations
23 defined by the MID (no clinically important effect, clinical benefit and clinical harm).
24 This is illustrated in Figure 1.

25 The value / position of the MID lines is ideally determined by values reported in the
26 literature. 'Anchor-based' methods aim to establish clinically meaningful changes in a
27 continuous outcome variable by relating or 'anchoring' them to patient-centred
28 measures of clinical effectiveness that could be regarded as gold standards with a
29 high level of face validity. For example, a MID for an outcome could be defined by the
30 minimum amount of change in that outcome necessary to make patients feel their
31 quality of life had 'significantly improved'. MIDs in the literature may also be based on
32 expert clinician or consensus opinion concerning the minimum amount of change in a
33 variable deemed to affect quality of life or health.

34 In the absence of values identified in the literature, the alternative approach to
35 deciding on MID levels is to use the modified GRADE 'default' values, as follows:

- 36 • For dichotomous outcomes the MIDs were taken to be RRs of 0.8* and 1.25. For
37 'positive' outcomes such as 'patient satisfaction', the RR of 0.8 is taken as the line
38 denoting the boundary between no clinically important effect and a clinically
39 important harm, whilst the RR of 1.25 is taken as the line denoting the boundary
40 between no clinically important effect and a clinically important benefit. For
41 'negative' outcomes such as 'bleeding', the opposite occurs, so the RR of 0.8 is
42 taken as the line denoting the boundary between no clinically important effect and
43 a clinically important benefit, whilst the RR of 1.25 is taken as the line denoting the
44 boundary between no clinically important effect and a clinically important harm.
45 There aren't established default values for ORs and the same values (0.8 and
46 1.25) are applied here but are acknowledged as arbitrary thresholds agreed by the
47 committee.

- 1 ○ In cases where there are zero events in one arm of a single study, or some or
2 all of the studies in one arm of a meta-analysis, the same process is followed
3 as for dichotomous outcomes. However, if there are no events in either arm in
4 a meta-analysis (or in a single unpooled study) the sample size is used to
5 determine imprecision using the following rule of thumb:
6 – No imprecision: sample size ≥ 350
7 – Serious imprecision: sample size ≥ 70 but < 350
8 – Very serious imprecision: sample size < 70 .
- 9 ○ When there was more than one study in an analysis and zero events occurred
10 in both groups for some but not all of the studies across both arms, the
11 optimum information size was used to determine imprecision using the
12 following guide:
13 – No imprecision: $> 90\%$ power
14 – Serious imprecision: $80-90\%$ power
15 – Very serious imprecision: $< 80\%$ power.
- 16 ● For mortality any change was considered to be clinically important and the
17 imprecision was assessed on the basis of whether the confidence intervals
18 crossed the line of no effect, that is whether the result was consistent with both
19 benefit and harm.
 - 20 ● For continuous outcome variables the MID was taken as half the median baseline
21 standard deviation of that variable, across all studies in the meta-analysis. Hence
22 the MID denoting the minimum clinically important benefit was positive for a
23 'positive' outcome (for example, a quality of life measure where a higher score
24 denotes better health), and negative for a 'negative' outcome (for example, a
25 visual analogue scale [VAS] pain score). Clinically important harms will be the
26 converse of these. If baseline values are unavailable, then half the median
27 comparator group standard deviation of that variable will be taken as the MID. As
28 these vary for each outcome per review, details of the values used are reported in
29 the footnotes of the relevant GRADE summary table.
 - 30 ● If standardised mean differences have been used, where the GC are able to
31 specify a priority measure, the results are back-converted to a mean difference on
32 that scale for the assessment of imprecision and clinical importance. If it is not
33 deemed appropriate to back-convert to a single scale, then the MID was set at the
34 absolute value of $+0.5$. This follows because standardised mean differences are
35 mean differences normalised to the pooled standard deviation of the 2 groups,
36 and are thus effectively expressed in units of 'numbers of standard deviations'.
37 The 0.5 MID value in this context therefore indicates half a standard deviation, the
38 same definition of MID as used for non-standardised mean differences.

39 *NB GRADE report the default values as 0.75 and 1.25 . These are consensus
40 values. This guideline follows NICE process to use modified values of 0.8 and 1.25
41 as they are symmetrical on a relative risk scale.


42 For this guideline, the following MIDs for continuous or dichotomous outcomes were
43 found in the literature and adopted for use:

44 **Table 5: Published or pre-agreed MIDs**

Outcome measure	MID	Source
6 minute walk distance	28 meters	Used in the 2013 version of the guideline

Outcome measure	MID	Source
Action Research Arm test	12 (dominant side), 17 (non-dominant side)	Used in the 2013 version of the guideline
Barthel index	1.85	Used in the 2013 version of the guideline
EQ-5D	0.03	Consensus pragmatic MID used in some previous NICE guidelines
Fugl-Meyer Assessment	10% of the total scale	Used in the 2013 version of the guideline
Functional Independence Measure	Motor subscale: 17 Cognition subscale: 3	Used in the 2013 version of the guideline
Motor Activity Log	1.0 (dominant side), 1.1 (non-dominant side)	Used in the 2013 version of the guideline
SF36	Physical component summary: 2 Mental component summary: 3 Physical functioning: 3 Role-physical: 3 Bodily pain: 3 General health: 2 Vitality: 2 Social functioning: 3 Role-emotional: 4 Mental health: 3	User's manual for the SF-36v2 Health Survey, Third Edition ²
Stairs test	15 seconds	Used in the 2013 version of the guideline
Timed up and go	10 seconds	Used in the 2013 version of the guideline
Walking speed	16 cm/second (acute stroke), 20 cm/second (chronic stroke)	Used in the 2013 version of the guideline
Wolf Motor Function Test	19 seconds (dominant side, 16% of the 120 second limit)	Used in the 2013 version of the guideline

Figure 1: Illustration of precise and imprecise outcomes based on the 95% CI of dichotomous outcomes in a forest plot (Note that all 3 results would be pooled estimates, and would not, in practice, be placed on the same forest plot)



1 2.5.1.5 Overall grading of the quality of clinical evidence

2 Once an outcome had been appraised for the main quality elements, as above, an
 3 overall quality grade was calculated for that outcome. The scores (0, -1 or -2) from
 4 each of the main quality elements were summed to give a score that could be
 5 anything from 0 (the best possible) to -8 (the worst possible). However, scores were
 6 capped at -3. This final score was then applied to the starting grade that had
 7 originally been applied to the outcome by default, based on study design. RCTs start
 8 at High, the overall quality became Moderate, Low or Very Low if the overall score
 9 was -1, -2 or -3 points respectively. The significance of these overall ratings is
 10 explained in **Table 6**. The reasons for downgrading in each case are specified in the
 11 footnotes of the GRADE tables.

12 **Table 6: Overall quality of outcome evidence in GRADE**

Level	Description
High	Further research is very unlikely to change our confidence in the estimate of effect
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate
Very low	Any estimate of effect is very uncertain

1 2.5.2 Diagnostic reviews

2 2.5.2.1 Diagnostic RCTs

3 Appraising the quality of evidence from diagnostic RCTs follows the same process as
4 section 2.5.1 for intervention reviews.

5 2.5.2.2 Diagnostic test accuracy

6 2.5.2.2.1 Risk of bias

7 Risk of bias and indirectness of evidence for diagnostic data were evaluated by study
8 using the Quality Assessment of Diagnostic Accuracy Studies version 2 (QUADAS-2)
9 checklists (see appendix H in the NICE guidelines manual 2014⁴). Risk of bias and
10 applicability in primary diagnostic accuracy studies in QUADAS-2 consists of 4
11 domains (see **Table 7**):

- 12 • patient selection
- 13 • index test
- 14 • reference standard
- 15 • flow and timing.

16 **Table 7 Summary of QUADAS-2 with list of signalling, risk of bias and**
17 **applicability questions.**

Domain	Patient selection	Index test	Reference standard	Flow and timing
Description	Describe methods of patient selection. Describe included patients (prior testing, presentation, intended use of index test and setting)	Describe the index test and how it was conducted and interpreted	Describe the reference standard and how it was conducted and interpreted	Describe any patients who did not receive the index test(s) and/or reference standard or who were excluded from the 2x2 table (refer to flow diagram). Describe the time interval and any interventions between index test(s) and reference standard
Signalling questions (yes/no/unclear)	Was a consecutive or random sample of patients enrolled?	Were the index test results interpreted without knowledge of the results of the reference standard?	Is the reference standard likely to correctly classify the target condition?	Was there an appropriate interval between index test(s) and reference standard?
	Was a case-control design avoided?	If a threshold was used, was it pre-specified?	Were the reference standard results interpreted without knowledge of the	Did all patients receive a reference standard?
	Did the study avoid			Did all patients receive the same reference standard?

Domain	Patient selection	Index test	Reference standard	Flow and timing
	inappropriate exclusions?		results of the index test?	Were all patients included in the analysis?
Risk of bias; (high/low/unclear)	Could the selection of patients have introduced bias?	Could the conduct or interpretation of the index test have introduced bias?	Could the reference standard, its conduct or its interpretation have introduced bias?	Could the patient flow have introduced bias?
Concerns regarding applicability (high/low/unclear)	Are there concerns that the included patients do not match the review question?	Are there concerns that the index test, its conduct, or interpretation differ from the review question?	Are there concerns that the target condition as defined by the reference standard does not match the review question?	

12.5.2.2.2 *Inconsistency*

2 Inconsistency refers to an unexplained heterogeneity of results for an outcome
3 across different studies. Inconsistency was assessed by visual inspection of the
4 primary outcome measures (sensitivity and specificity) using the point estimates and
5 95% CIs of the individual studies on the forest plots or the summary value if a
6 diagnostic meta-analysis had been conducted. The evidence was downgraded by 1
7 increment if there was no overlap of 95% confidence intervals or by 2 increments if
8 there was wide variability. Where only a single study reports an outcome,
9 inconsistency is rated as 'not detected'.

102.5.2.2.3 *Imprecision*

11 The judgement of precision was based on visual inspection of the confidence region
12 around the summary sensitivity and specificity point from the diagnostic meta-
13 analysis, if a diagnostic meta-analysis was conducted. Where a diagnostic meta-
14 analysis was not conducted, imprecision was assessed according to the range of
15 point estimates or, if only one study contributed to the evidence, the 95% CI around
16 the single study. The decision thresholds set by the committee were used to
17 determine whether imprecision is not serious, serious or very serious depending on
18 whether confidence intervals cross zero, one or two thresholds.

192.5.2.2.4 *Overall grading*

20 Quality rating started at high for prospective and retrospective cross-sectional
21 studies, and each major limitation (risk of bias, indirectness, inconsistency and
22 imprecision) brought the rating down by 1 increment to a minimum grade of very low,
23 as explained for intervention reviews. This was presented in a GRADE evidence
24 profile.

25 **2.5.3 Tool validity and reliability reviews**

26 In addition to the data synthesis of the tool validity and reliability, which assesses the
27 quality of the tools, each study was appraised for sources of bias. Each study was

appraised for bias arising from the population, the control population, recruitment/selection bias, sample size concerns and whether the metrics used are appropriate for the objective of the study. This was presented for each study separately and the committee took this into account qualitatively when assessing the results from each study.

These aspects of bias were taken into account in conjunction with the results of the study and assessed in their totality. As the nature of the tool validity and reliability review was to assess the quality of the tools, risk of bias was assessed as one component of this quality assessment alongside the other results from the studies and so the committee weighed these results against those of the others in order to determine the quality of the tools reported in the literature.

2.5.4 Qualitative reviews

Review findings from the included qualitative studies were evaluated and presented using the 'Confidence in the Evidence from Reviews of Qualitative Research' (CERQual) Approach developed by the GRADE-CERQual Project Group, a subgroup of the GRADE Working Group.

The CERQual Approach assesses the extent to which a review finding is a reasonable representation of the phenomenon of interest (the focus of the review question). Each review finding was assessed for each of the 4 quality elements listed and defined below in Table 8.

Table 8: Description of quality elements in GRADE-CERQual for qualitative studies

Quality element	Description
Methodological limitations	The extent of problems in the design or conduct of the included studies that could decrease the confidence that the review finding is a reasonable representation of the phenomenon of interest. Assessed at the study level using the CASP checklist.
Coherence	The extent to how clear and cogent the fit is between the data from the primary studies and the review finding.
Relevance	The extent to which the body of evidence from the included studies is applicable to the context (study population, phenomenon of interest, setting) specified in the protocol.
Adequacy	The degree of the confidence that the review finding is being supported by sufficient data. This is an overall determination of the richness (depth of analysis) and quantity of the evidence supporting a review finding or theme.

Details of how the 4 quality elements (methodological limitations, coherence, relevance and adequacy) were appraised for each review finding are given below.

2.5.4.1 Methodological limitations

Each review finding had its methodological limitations assessed within each study first using the CASP checklist. Based on the degree of methodological limitations, studies were evaluated as having minor, moderate or severe limitations. A summary of the domains and questions covered is given below.

Table 9: Description of limitations assessed in the CASP checklist for qualitative studies

Domain	Aspects considered
Are the results valid?	<ul style="list-style-type: none"> • Was there a clear statement of the aims of the research? • Is qualitative methodology appropriate? • Was the research design appropriate to address the aims of the research? • Was the recruitment strategy appropriate to the aims of the research? • Was the data collected in a way that addressed the research issue? • Has the relationship between researcher and participants been adequately considered?
What are the results?	<ul style="list-style-type: none"> • Have ethical issues been taken into consideration? • Was the data analysis sufficiently rigorous? • Is there a clear statement of findings?
Will the results help locally?	<ul style="list-style-type: none"> • How valuable is the research?

The overall assessment of the methodological limitations of the evidence was based on the limitations of the primary studies contributing to the review finding. The relative contribution of each study to the overall review finding and of the type of methodological limitation(s) were taken into account when giving an overall rating of concerns for this component.

2.5.4.2 Relevance

Relevance is the extent to which the body of evidence from the included studies is applicable to the context (study population, phenomenon of interest, setting) specified in the protocol. As such, relevance is dependent on the individual review and discussed with the guideline committee.

2.5.4.3 Coherence

Coherence is the extent to which the reviewer is able to identify a clear pattern across the studies included in the review, and if there is variation present (contrasting or disconfirming data) whether this variation is explained by the contributing study authors. For example, if a review finding in 1 study does not support the main finding and there is no plausible explanation for this variation, or if there is ambiguity in the descriptions in the primary data, then the confidence that the main finding reasonably reflects the phenomenon of interest is decreased.

2.5.4.4 Adequacy

The judgement of adequacy is based on the confidence of the finding being supported by sufficient data. This is an overall determination of the richness (and quantity of the evidence supporting a review finding or theme. Rich data provide sufficient detail to gain an understanding of the theme or review finding, whereas thin data do not provide enough detail for an adequate understanding. Quantity of data is the second pillar of the assessment of adequacy. For review findings that are only supported by 1 study or data from only a small number of participants, the confidence that the review finding reasonably represents the phenomenon of interest might be decreased because there is less confidence that studies undertaken in other settings or participants would have reported similar findings. As with richness of data, quantity

of data is review dependent. Based on the overall judgement of adequacy, a rating of no concerns, minor concerns, or substantial concerns about adequacy was given.

2.5.4.5 Overall judgement of the level of confidence for a review finding

GRADE-CERQual is used to assess the body of evidence as a whole through a confidence rating representing the extent to which a review finding is a reasonable representation of the phenomenon of interest. For each of the above components, level of concern is categorised as either;

- no or very minor concerns
- minor concerns
- moderate concerns, or
- serious concerns.

The concerns from the 4 components (methodological limitations, coherence, relevance and adequacy) are used in combination to form an overall judgement of confidence in the finding. GRADE-CERQual uses 4 levels of confidence: high, moderate, low and very low confidence. The significance of these overall ratings is explained in Table 10. Each review finding starts at a high level of confidence and is downgraded based on the concerns identified in any 1 or more of the 4 components. Quality assessment of qualitative reviews is a subjective judgement by the reviewer based on the concerns that have been noted. An explanation of how such a judgement had been made for each component is included in the footnotes of the summary of evidence tables.

Table 10: Overall level of confidence for a review finding in GRADE-CERQual

Level	Description
High confidence	It is highly likely that the review finding is a reasonable representation of the phenomenon of interest.
Moderate confidence	It is likely that the review finding is a reasonable representation of the phenomenon of interest.
Low confidence	It is possible that the review finding is a reasonable representation of the phenomenon of interest.
Very low confidence	It is not clear whether the review finding is a reasonable representation of the phenomenon of interest.

2.5.5 Mixed methods reviews

The quality of intervention studies and qualitative studies were assessed following the methods in the intervention qualitative review sections respectively. For more information, please refer to the Intervention reviews and Qualitative reviews sections.

2.6 Assessing clinical importance

The committee assessed the evidence by outcome in order to determine if there was, or potentially was, a clinically important benefit, a clinically important harm or no clinically important difference between interventions. To facilitate this, binary outcomes were converted into absolute risk differences (ARDs) using GRADEpro¹ software: the median control group risk across studies was used to calculate the ARD and its 95% CI from the pooled risk ratio.

1 The assessment of clinical benefit, harm, or no benefit or harm was based on the
2 point estimate of absolute effect for intervention studies, which was standardised
3 across the reviews. The committee considered for most of the dichotomous
4 outcomes in the intervention reviews that if at least 100 more participants per 1000
5 (10%) achieved the outcome of interest in the intervention group compared to the
6 comparison group for a positive outcome then this intervention was considered
7 beneficial. The same point estimate but in the opposite direction applied for a
8 negative outcome. For mortality any reduction represented a clinical benefit. For
9 adverse events 50 events or more per 1000 (5%) represented clinical harm.

10 For continuous outcomes if the mean difference was greater than the minimally
11 important difference (MID) then this represented a clinical benefit or harm. For
12 outcomes such as mortality any reduction or increase was considered to be clinically
13 important.

14 Established MIDs found in the literature and were agreed to be used for a range of
15 outcomes. The outcomes and the published values used for imprecision and clinical
16 importance are provided in **Table 5**. For continuous outcomes where the GRADE
17 default MID has been used, the values for each outcome are provided in the
18 footnotes of the relevant GRADE tables.

19 **2.7 Identifying and analysing evidence of cost** 20 **effectiveness**

21 The committee is required to make decisions based on the best available evidence of
22 both clinical effectiveness and cost effectiveness. Guideline recommendations should
23 be based on the expected costs of the different options in relation to their expected
24 health benefits (that is, their 'cost effectiveness') rather than the total implementation
25 cost. However, the committee will also need to be increasingly confident in the cost
26 effectiveness of a recommendation as the cost of implementation increases.
27 Therefore, the committee may require more robust evidence on the effectiveness and
28 cost effectiveness of any recommendations that are expected to have a substantial
29 impact on resources; any uncertainties must be offset by a compelling argument in
30 favour of the recommendation. The cost impact or savings potential of a
31 recommendation should not be the sole reason for the committee's decision.⁴

32 Health economic evidence was sought relating to the key clinical issues being
33 addressed in the guideline. Health economists:

- 34 • Undertook a systematic review of the published economic literature.
- 35 • Undertook new cost-effectiveness analysis in priority areas.

36 **2.7.1 Literature review**

37 The health economists:

- 38 • Identified potentially relevant studies for each review question from the health
39 economic search results by reviewing titles and abstracts. Full papers were then
40 obtained.
- 41 • Reviewed full papers against prespecified inclusion and exclusion criteria to
42 identify relevant studies (see below for details).
- 43 • Critically appraised relevant studies using economic evaluations checklists as
44 specified in the NICE guidelines manual.⁴

- 1 • Extracted key information about the studies' methods and results into health
2 economic evidence tables (which can be found in appendices to the relevant
3 evidence reports).
- 4 • Generated summaries of the evidence in NICE health economic evidence profile
5 tables (included in the relevant evidence report for each review question) – see
6 below for details.

7 **2.7.1.1 Inclusion and exclusion criteria**

8 Full economic evaluations (studies comparing costs and health consequences of
9 alternative courses of action: cost–utility, cost-effectiveness, cost–benefit and cost–
10 consequences analyses) and comparative costing studies that addressed the review
11 question in the relevant population were considered potentially includable as health
12 economic evidence.

13 Studies that only reported cost per hospital (not per patient), or only reported average
14 cost effectiveness without disaggregated costs and effects were excluded. Literature
15 reviews, abstracts, posters, letters, editorials, comment articles, unpublished studies
16 and studies not in English were excluded. Studies published before 2006 and studies
17 from non-OECD countries or the USA were also excluded, on the basis that the
18 applicability of such studies to the present UK NHS context is likely to be too low for
19 them to be helpful for decision-making.

20 Remaining health economic studies were prioritised for inclusion based on their
21 relative applicability to the development of this guideline and the study limitations.
22 However, in this guideline, no economic studies were excluded on the basis that
23 more applicable evidence was available.

24 For more details about the assessment of applicability and methodological quality
25 see **Table 11** below and the economic evaluation checklist (appendix H of the NICE
26 guidelines manual⁴) and the health economics review protocol, which can be found in
27 each of the evidence reports.

28 When no relevant health economic studies were found from the economic literature
29 review, relevant UK NHS unit costs related to the compared interventions were
30 presented to the committee to inform the possible economic implications of the
31 recommendations.

32 **2.7.1.2 NICE health economic evidence profiles**

33 NICE health economic evidence profile tables were used to summarise cost and
34 cost-effectiveness estimates for the included health economic studies in each
35 evidence review report. The health economic evidence profile shows an assessment
36 of applicability and methodological quality for each economic study, with footnotes
37 indicating the reasons for the assessment. These assessments were made by the
38 health economist using the economic evaluation checklist from the NICE guidelines
39 manual.⁴ It also shows the incremental costs, incremental effects (for example,
40 quality-adjusted life years [QALYs]) and incremental cost-effectiveness ratio (ICER)
41 for the base case analysis in the study, as well as information about the assessment
42 of uncertainty in the analysis. See **Table 11** for more details.

43 When a non-UK study was included in the profile, the results were converted into
44 pounds sterling using the appropriate purchasing power parity.¹⁰

1

Table 11: Content of NICE health economic evidence profile

Item	Description
Study	Surname of first author, date of study publication and country perspective with a reference to full information on the study.
Applicability	An assessment of applicability of the study to this guideline, the current NHS situation and NICE decision-making: ^(a) <ul style="list-style-type: none"> • Directly applicable – the study meets all applicability criteria, or fails to meet 1 or more applicability criteria but this is unlikely to change the conclusions about cost effectiveness. • Partially applicable – the study fails to meet 1 or more applicability criteria, and this could change the conclusions about cost effectiveness. • Not applicable – the study fails to meet 1 or more of the applicability criteria, and this is likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.
Limitations	An assessment of methodological quality of the study: ^(a) <ul style="list-style-type: none"> • Minor limitations – the study meets all quality criteria, or fails to meet 1 or more quality criteria, but this is unlikely to change the conclusions about cost effectiveness. • Potentially serious limitations – the study fails to meet 1 or more quality criteria, and this could change the conclusions about cost effectiveness. • Very serious limitations – the study fails to meet 1 or more quality criteria, and this is highly likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.
Other comments	Information about the design of the study and particular issues that should be considered when interpreting it.
Incremental cost	The mean cost associated with one strategy minus the mean cost of a comparator strategy.
Incremental effects	The mean QALYs (or other selected measure of health outcome) associated with one strategy minus the mean QALYs of a comparator strategy.
Cost effectiveness	Incremental cost-effectiveness ratio (ICER): the incremental cost divided by the incremental effects (usually in £ per QALY gained).
Uncertainty	A summary of the extent of uncertainty about the ICER reflecting the results of deterministic or probabilistic sensitivity analyses, or stochastic analyses of trial data, as appropriate.

2

(a) *Applicability and limitations were assessed using the economic evaluation checklist in appendix H of the NICE guidelines manual⁴*

3

4 2.7.2 Undertaking new health economic analysis

5

As well as reviewing the published health economic literature for each review question, as described above, new health economic analysis was undertaken by the health economist in selected areas. Priority areas for new analysis were agreed by the committee after formation of the review questions and consideration of the existing health economic evidence.

6

7

8

9

10

The committee identified two topics as the highest priority areas for original health economic modelling. These were:

11

12

- Intensity of physiotherapy (**Evidence review E**)
- Botulinum toxin for spasticity (**Evidence review P**)

13

14

These topics were modelled because, for each one, there was:

- a body of clinical effectiveness evidence,
- limited published cost effectiveness evidence,
- potential for a significant resource impact if recommended.

The following general principles were adhered to in developing the cost-effectiveness analyses:

- Methods were consistent with the NICE reference case for interventions with health outcomes in NHS settings.^{4,7}
- The committee was involved in the design of the model, selection of inputs and interpretation of the results.
- Model inputs were based on the systematic review of the clinical literature supplemented with other published data sources where possible.
- When published data were not available committee expert opinion was used to populate the model.
- Model inputs and assumptions were reported fully and transparently.
- The results were subject to sensitivity analysis and limitations were discussed.
- The model was peer-reviewed by another health economist.

Full methods and results of the two cost-utility analyses are described in a separate economic analysis reports.

2.7.3 Cost-effectiveness criteria

NICE sets out the principles that committees should consider when judging whether an intervention offers good value for money.^{4,6,8} In general, an intervention was considered to be cost effective (given that the estimate was considered plausible) if either of the following criteria applied:

- the intervention dominated other relevant strategies (that is, it was both less costly in terms of resource use and more clinically effective compared with all the other relevant alternative strategies), or
- the intervention cost less than £20,000 per QALY gained compared with the next best strategy.

If the committee recommended an intervention that was estimated to cost more than £20,000 per QALY gained, or did not recommend one that was estimated to cost less than £20,000 per QALY gained, the reasons for this decision are discussed explicitly in 'The committee's discussion of the evidence' section of the relevant evidence report, with reference to issues regarding the plausibility of the estimate or to factors set out in NICE methods manuals.⁴

When QALYs or life years gained are not used in the analysis, results are difficult to interpret unless one strategy dominates the others with respect to every relevant health outcome and cost.

2.7.4 In the absence of health economic evidence

When no relevant published health economic studies were found, and a new analysis was not prioritised, the committee made a qualitative judgement about cost effectiveness by considering expected differences in resource use between options and relevant UK NHS unit costs, alongside the results of the review of clinical effectiveness evidence.

1 The UK NHS costs reported in the guideline are those that were presented to the
2 committee and were correct at the time recommendations were drafted. They may
3 have changed subsequently before the time of publication. However, we have no
4 reason to believe they have changed substantially.

5 **2.8 Developing recommendations**

6 Over the course of the guideline development process, the committee was presented
7 with:

- 8 • Summaries of clinical and health economic evidence and quality (as presented in
9 evidence reports [A–Q]).
- 10 • Evidence tables of the clinical and health economic evidence reviewed from the
11 literature. All evidence tables can be found in appendices to the relevant evidence
12 reports.
- 13 • Forest plots (in appendices to the relevant evidence reports).
- 14 • A description of the methods and results of the cost-effectiveness analysis
15 undertaken for the guideline (in a separate economic analysis report).

16 Decisions on whether a recommendation could be made, and if so in which direction,
17 were made on the basis of the committee's interpretation of the available evidence,
18 taking into account the balance of benefits, harms and costs between different
19 courses of action. This was either done formally in an economic model, or informally.
20 The net clinical benefit over harm (clinical effectiveness) was considered, focusing on
21 the magnitude of the effect (or clinical importance), quality of evidence (including the
22 uncertainty) and amount of evidence available. When this was done informally, the
23 committee took into account the clinical benefits and harms when one intervention
24 was compared with another. The assessment of net clinical benefit was moderated
25 by the importance placed on the outcomes (the committee's values and preferences),
26 and the confidence the committee had in the evidence (evidence quality). Secondly,
27 the committee assessed whether the net clinical benefit justified any differences in
28 costs between the alternative interventions. When the clinical harms were judged by
29 the committee to outweigh any clinical benefits, they considered making a
30 recommendation not to offer an intervention. This was dependant on whether the
31 intervention had any reasonable prospect of providing cost-effective benefits to
32 people using services and whether stopping the intervention was likely to cause harm
33 for people already receiving it.

34 When clinical and health economic evidence was of poor quality, conflicting or
35 absent, the committee decided on whether a recommendation could be made based
36 on its expert opinion. The considerations for making consensus-based
37 recommendations include the balance between potential harms and benefits, the
38 economic costs compared to the economic benefits, current practices,
39 recommendations made in other relevant guidelines, patient preferences and equality
40 issues. The consensus recommendations were agreed through discussions in the
41 committee. The committee also considered whether the uncertainty was sufficient to
42 justify delaying making a recommendation to await further research, taking into
43 account the potential harm of failing to make a clear recommendation (see 2.8.1).

44 The committee considered the appropriate 'strength' of each recommendation. This
45 takes into account the quality of the evidence but is conceptually different. Some
46 recommendations are 'strong' in that the committee believes that the vast majority of
47 healthcare and other professionals and patients would choose a particular

1 intervention if they considered the evidence in the same way that the committee has.
 2 This is generally the case if the benefits clearly outweigh the harms for most people
 3 and the intervention is likely to be cost effective. However, there is often a closer
 4 balance between benefits and harms, and some patients would not choose an
 5 intervention whereas others would. This may happen, for example, if some patients
 6 are particularly averse to some side effect and others are not. In these circumstances
 7 the recommendation is generally weaker, although it may be possible to make
 8 stronger recommendations about specific groups of patients.

9 The committee focused on the following factors in agreeing the wording of the
 10 recommendations:

- 11 • The actions health professionals need to take.
- 12 • The information readers need to know.
- 13 • The strength of the recommendation (for example the word ‘offer’ was used for
 14 strong recommendations and ‘consider’ for weaker recommendations).
- 15 • The involvement of patients (and their carers if needed) in decisions on treatment
 16 and care.
- 17 • Consistency with NICE’s standard advice on recommendations about drugs,
 18 waiting times and ineffective interventions (see section 9.2 in the NICE guidelines
 19 manual⁵).

20 The main considerations specific to each recommendation are outlined in ‘The
 21 committee’s discussion of the evidence’ section within each evidence report.

22 2.8.1 Research recommendations

23 When areas were identified for which good evidence was lacking, the committee
 24 considered making recommendations for future research. Decisions about the
 25 inclusion of a research recommendation were based on factors such as:

- 26 • the importance to patients or the population
- 27 • national priorities
- 28 • potential impact on the NHS and future NICE guidance
- 29 • ethical and technical feasibility.

30 2.8.2 Validation process

31 This guidance is subject to a 6-week public consultation and feedback as part of the
 32 quality assurance and peer review of the document. All comments received from
 33 registered stakeholders are responded to in turn and posted on the NICE website.

34 2.8.3 Updating the guideline

35 Following publication, and in accordance with the NICE guidelines manual, NICE will
 36 undertake a review of whether the evidence base has progressed significantly to alter
 37 the guideline recommendations and warrant an update.

38 2.9 General terms

Term	Definition
Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.

Term	Definition
Algorithm (in guidelines)	A flow chart of the clinical decision pathway described in the guideline, where decision points are represented with boxes, linked with arrows.
Allocation concealment	The process used to prevent advance knowledge of group assignment in an RCT. The allocation process should be impervious to any influence by the individual making the allocation, by being administered by someone who is not responsible for recruiting participants.
Applicability	How well the results of a study or NICE evidence review can answer a clinical question or be applied to the population being considered.
Arm (of a clinical study)	Subsection of individuals within a study who receive one particular intervention, for example placebo arm.
Association	Statistical relationship between 2 or more events, characteristics or other variables. The relationship may or may not be causal.
Base case analysis	In an economic evaluation, this is the main analysis based on the most plausible estimate of each input. In contrast, see Sensitivity analysis.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable), with which subsequent results are compared.
Before-and-after study	A study that investigates the effects of an intervention by measuring particular characteristics of a population both before and after taking the intervention, and assessing any change that occurs.
Bias	Influences on a study that can make the results look better or worse than they really are. (Bias can even make it look as if a treatment works when it does not.) Bias can occur by chance, deliberately or as a result of systematic errors in the design and execution of a study. It can also occur at different stages in the research process, for example, during the collection, analysis, interpretation, publication or review of research data. For examples see selection bias, performance bias, information bias, confounding factor, and publication bias.
Blinding	<p>A way to prevent researchers, doctors and patients in a clinical trial from knowing which study group each patient is in so they cannot influence the results. The best way to do this is by sorting patients into study groups randomly. The purpose of 'blinding' or 'masking' is to protect against bias.</p> <p>A single-blinded study is one in which patients do not know which study group they are in (for example whether they are taking the experimental drug or a placebo). A double-blinded study is one in which neither patients nor the researchers and doctors know which study group the patients are in. A triple blind study is one in which neither the patients, clinicians or the people carrying out the statistical analysis know which treatment patients received.</p>
Carer (caregiver)	Someone who looks after family, partners or friends in need of help because they are ill, frail or have a disability.
Case-control study	A study to find out the cause(s) of a disease or condition. This is done by comparing a group of patients who have the disease or condition (cases) with a group of people who do not have it (controls) but who are otherwise as similar as possible (in characteristics thought to be unrelated to the causes of the disease or condition). This means the researcher can look for aspects of their lives that differ to see if they may cause the condition.

Term	Definition
	For example, a group of people with lung cancer might be compared with a group of people the same age that do not have lung cancer. The researcher could compare how long both groups had been exposed to tobacco smoke. Such studies are retrospective because they look back in time from the outcome to the possible causes of a disease or condition.
Case series	Report of a number of cases of a given disease, usually covering the course of the disease and the response to treatment. There is no comparison (control) group of patients.
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinical effectiveness	How well a specific test or treatment works when used in the 'real world' (for example, when used by a doctor with a patient at home), rather than in a carefully controlled clinical trial. Trials that assess clinical effectiveness are sometimes called management trials. Clinical effectiveness is not the same as efficacy.
Clinician	A healthcare professional who provides patient care. For example, a doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence-based medicine databases including the Cochrane Database of Systematic Reviews (reviews of randomised controlled trials prepared by the Cochrane Collaboration).
Cohort study	A study with 2 or more groups of people – cohorts – with similar characteristics. One group receives a treatment, is exposed to a risk factor or has a particular symptom and the other group does not. The study follows their progress over time and records what happens. See also observational study.
Comorbidity	A disease or condition that someone has in addition to the health problem being studied or treated.
Comparability	Similarity of the groups in characteristics likely to affect the study results (such as health status or age).
Concordance	This is a recent term whose meaning has changed. It was initially applied to the consultation process in which doctor and patient agree therapeutic decisions that incorporate their respective views, but now includes patient support in medicine taking as well as prescribing communication. Concordance reflects social values but does not address medicine-taking and may not lead to improved adherence.
Confidence interval (CI)	A range of values for an unknown population parameter with a stated 'confidence' (conventionally 95%) that it contains the true value. The interval is calculated from sample data, and generally straddles the sample estimate. The 'confidence' value means that if the method used to calculate the interval is repeated many times, then that proportion of intervals will actually contain the true value.
Confounding factor	Something that influences a study and can result in misleading findings if it is not understood or appropriately dealt with. For example, a study of heart disease may look at a group of people that exercises regularly and a group that does not exercise. If the ages of the people in the 2 groups are different, then any difference in heart disease rates between the 2 groups could be because of age rather than exercise. Therefore age is a confounding factor.
Content validity	The extent to which a test is subjectively viewed as measuring what it is meant to (for example: if a questionnaire is asking questions about

Term	Definition
	stroke fatigue, the questions ask about fatigue). Whether the test relates to underlying theoretical concepts. This includes face validity and construct validity.
Consensus methods	Techniques used to reach agreement on a particular issue. Consensus methods may be used to develop NICE guidance if there is not enough good quality research evidence to give a clear answer to a question. Formal consensus methods include Delphi and nominal group techniques.
Control group	A group of people in a study who do not receive the treatment or test being studied. Instead, they may receive the standard treatment (sometimes called 'usual care') or a dummy treatment (placebo). The results for the control group are compared with those for a group receiving the treatment being tested. The aim is to check for any differences. Ideally, the people in the control group should be as similar as possible to those in the treatment group, to make it as easy as possible to detect any effects due to the treatment.
Convergent and discriminant validity	Convergent validity is the extent to which a test shows that measures that should be related are related (for example: showing that a fatigue scale does measure fatigue). Discriminant validity the extent to which a test shows that measures that should not be related are not related (for example: showing that a fatigue scale does not measure pain).
Cost–benefit analysis (CBA)	Cost–benefit analysis is one of the tools used to carry out an economic evaluation. The costs and benefits are measured using the same monetary units (for example, pounds sterling) to see whether the benefits exceed the costs.
Cost–consequences analysis (CCA)	Cost–consequences analysis is one of the tools used to carry out an economic evaluation. This compares the costs (such as treatment and hospital care) and the consequences (such as health outcomes) of a test or treatment with a suitable alternative. Unlike cost–benefit analysis or cost-effectiveness analysis, it does not attempt to summarise outcomes in a single measure (like the quality-adjusted life year) or in financial terms. Instead, outcomes are shown in their natural units (some of which may be monetary) and it is left to decision-makers to determine whether, overall, the treatment is worth carrying out.
Cost-effectiveness analysis (CEA)	Cost-effectiveness analysis is one of the tools used to carry out an economic evaluation. The benefits are expressed in non-monetary terms related to health, such as symptom-free days, heart attacks avoided, deaths avoided or life years gained (that is, the number of years by which life is extended as a result of the intervention).
Cost-effectiveness model	An explicit mathematical framework, which is used to represent clinical decision problems and incorporate evidence from a variety of sources in order to estimate the costs and health outcomes.
Cost–utility analysis (CUA)	Cost–utility analysis is one of the tools used to carry out an economic evaluation. The benefits are assessed in terms of both quality and duration of life, and expressed as quality-adjusted life years (QALYs). See also utility.
Criterion validity	The degree to which a test corresponds to an external criteria that is known concurrently (for example: if a new test is shown to be comparable to a currently existing test).
Decision analysis	An explicit quantitative approach to decision-making under uncertainty, based on evidence from research. This evidence is

Term	Definition
	translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Deterministic analysis	In economic evaluation, this is an analysis that uses a point estimate for each input. In contrast, see Probabilistic analysis
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Disutility	The loss of quality of life associated with having a disease or condition. See Utility
Dominance	A health economics term. When comparing tests or treatments, an option that is both less effective and costs more is said to be 'dominated' by the alternative.
Drop-out	A participant who withdraws from a trial before the end.
Economic evaluation	An economic evaluation is used to assess the cost effectiveness of healthcare interventions (that is, to compare the costs and benefits of a healthcare intervention to assess whether it is worth doing). The aim of an economic evaluation is to maximise the level of benefits – health effects – relative to the resources available. It should be used to inform and support the decision-making process; it is not supposed to replace the judgement of healthcare professionals. There are several types of economic evaluation: cost–benefit analysis, cost–consequences analysis, cost-effectiveness analysis, cost-minimisation analysis and cost–utility analysis. They use similar methods to define and evaluate costs, but differ in the way they estimate the benefits of a particular drug, programme or intervention.
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	A measure that shows the magnitude of the outcome in one group compared with that in a control group. For example, if the absolute risk reduction is shown to be 5% and it is the outcome of interest, the effect size is 5%. The effect size is usually tested, using statistics, to find out how likely it is that the effect is a result of the treatment and has not just happened by chance (that is, to see if it is statistically significant).
Effectiveness	How beneficial a test or treatment is under usual or everyday conditions, compared with doing nothing or opting for another type of care.
Efficacy	How beneficial a test, treatment or public health intervention is under ideal conditions (for example, in a laboratory), compared with doing nothing or opting for another type of care.
Epidemiological study	The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (for example, infection, diet) and interventions.
EQ-5D (EuroQol 5 dimensions)	A standardised instrument used to measure health-related quality of life. It provides a single index value for health status.
Evidence	Information on which a decision or guidance is based. Evidence is obtained from a range of sources including randomised controlled trials, observational studies, expert opinion (of clinical professionals or patients).

Term	Definition
Exclusion criteria (literature review)	Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.
Exclusion criteria (clinical study)	Criteria that define who is not eligible to participate in a clinical study.
Extended dominance	If Option A is both more clinically effective than Option B and has a lower cost per unit of effect, when both are compared with a do-nothing alternative then Option A is said to have extended dominance over Option B. Option A is therefore cost effective and should be preferred, other things remaining equal.
Extrapolation	An assumption that the results of studies of a specific population will also hold true for another population with similar characteristics.
Follow-up	Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been assessed in order to observe changes in health status or health-related variables.
Generalisability	The extent to which the results of a study hold true for groups that did not participate in the research. See also external validity.
Gold standard	A method, procedure or measurement that is widely accepted as being the best available to test for or treat a disease.
GRADE, GRADE evidence profile	A system developed by the GRADE Working Group to address the shortcomings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE evidence profile.
Harms	Adverse effects of an intervention.
Health economics	Study or analysis of the cost of using and distributing healthcare resources.
Health-related quality of life (HRQoL)	A measure of the effects of an illness to see how it affects someone's day-to-day life.
Heterogeneity or Lack of homogeneity	The term is used in meta-analyses and systematic reviews to describe when the results of a test or treatment (or estimates of its effect) differ significantly in different studies. Such differences may occur as a result of differences in the populations studied, the outcome measures used or because of different definitions of the variables involved. It is the opposite of homogeneity.
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.
Inclusion criteria (literature review)	Explicit criteria used to decide which studies should be considered as potential sources of evidence.
Incremental analysis	The analysis of additional costs and additional clinical outcomes with different interventions.
Incremental cost	The extra cost linked to using one test or treatment rather than another. Or the additional cost of doing a test or providing a treatment more frequently.
Incremental cost-effectiveness ratio (ICER)	The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another.
Incremental net benefit (INB)	The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated

Term	Definition
	for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the INB is calculated as: $(£20,000 \times \text{QALYs gained}) - \text{Incremental cost}$.
Indirectness	The available evidence is different to the review question being addressed, in terms of PICO (population, intervention, comparison and outcome).
Intratest reliability	The degree of agreement among repeated administrations of a test performed by a single rater (whether consistent results are seen throughout the test when completed by a person). Relates to test-retest reliability.
Intention-to-treat analysis (ITT)	An assessment of the people taking part in a clinical trial, based on the group they were initially (and randomly) allocated to. This is regardless of whether or not they dropped out, fully complied with the treatment or switched to an alternative treatment. Intention-to-treat analyses are often used to assess clinical effectiveness because they mirror actual practice: that is, not everyone complies with treatment and the treatment people receive may be changed according to how they respond to it.
Internal consistency	The correlation between multiple items in a test that are intended to measure the same construct. Includes the intertest reliability and intratest reliability. This may be measured using: Cronbach's alpha and intraclass correlation coefficients.
Intertest reliability	The degree to which different people give consistent estimates of the same phenomena (test different people who experience fatigue and seeing if the test shows the same results for each of them)
Intervention	In medical terms this could be a drug treatment, surgical procedure, diagnostic or psychological therapy. Examples of public health interventions could include action to help someone to be physically active or to eat a more healthy diet.
Intraoperative	The period of time during a surgical procedure.
Kappa statistic	A statistical measure of inter-rater agreement that takes into account the agreement occurring by chance.
Length of stay	The total number of days a participant stays in hospital.
Licence	See 'Product licence'.
Life years gained	Mean average years of life gained per person as a result of the intervention compared with an alternative intervention.
Likelihood ratio	The likelihood ratio combines information about the sensitivity and specificity. It tells you how much a positive or negative result changes the likelihood that a patient would have the disease. The likelihood ratio of a positive test result (LR+) is sensitivity divided by (1 minus specificity).
Long-term care	Residential care in a home that may include skilled nursing care and help with everyday activities. This includes nursing homes and residential homes.
Logistic regression or Logit model	In statistics, logistic regression is a type of analysis used for predicting the outcome of a binary dependent variable based on one or more predictor variables. It can be used to estimate the log of the odds (known as the 'logit').
Loss to follow-up	A patient, or the proportion of patients, actively participating in a clinical trial at the beginning, but whom the researchers were unable to trace or contact by the point of follow-up in the trial

Term	Definition
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
Meta-analysis	A method often used in systematic reviews. Results from several studies of the same test or treatment are combined to estimate the overall effect of the treatment.
Multivariate model	A statistical model for analysis of the relationship between 2 or more predictor (independent) variables and the outcome (dependent) variable.
Negative predictive value (NPV)	In screening or diagnostic tests: A measure of the usefulness of a screening or diagnostic test. It is the proportion of those with a negative test result who do not have the disease, and can be interpreted as the probability that a negative test result is correct. It is calculated as follows: $TN/(TN+FN)$
Net monetary benefit (NMB)	The value in monetary terms of an intervention net of its cost. The NMB can be calculated for a given cost-effectiveness threshold. If the threshold is £20,000 per QALY gained then the NMB for an intervention is calculated as: $(£20,000 \times \text{mean QALYs}) - \text{mean cost}$. The most preferable option (that is, the most clinically effective option to have an ICER below the threshold selected) will be the treatment with the highest NMB.
Non-randomised intervention study	A quantitative study investigating the effectiveness of an intervention that does not use randomisation to allocate patients (or units) to treatment groups. Non-randomised studies include observational studies, where allocation to groups occurs through usual treatment decisions or people's preferences. Non-randomised studies can also be experimental, where the investigator has some degree of control over the allocation of treatments. Non-randomised intervention studies can use a number of different study designs, and include cohort studies, case-control studies, controlled before-and-after studies, interrupted-time-series studies and quasi-randomised controlled trials.
Number needed to treat (NNT)	The average number of patients who need to be treated to get a positive outcome. For example, if the NNT is 4, then 4 patients would have to be treated to ensure 1 of them gets better. The closer the NNT is to 1, the better the treatment. For example, if you give a stroke prevention drug to 20 people before 1 stroke is prevented, the number needed to treat is 20. See also number needed to harm, absolute risk reduction.
Observational study	Individuals or groups are observed or certain factors are measured. No attempt is made to affect the outcome. For example, an observational study of a disease or treatment would allow 'nature' or usual medical care to take its course. Changes or differences in one characteristic (for example, whether or not people received a specific treatment or intervention) are studied without intervening. There is a greater risk of selection bias than in experimental studies.
Odds ratio	A measure of treatment effectiveness. The odds of an event happening in the treatment group, expressed as a proportion of the odds of it happening in the control group. The 'odds' is the ratio of events to non-events.
Opportunity cost	The loss of other healthcare programmes displaced by investment in or introduction of another intervention. This may be best measured by

Term	Definition
	the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.
Outcome	The impact that a test, treatment, policy, programme or other intervention has on a person, group or population. Outcomes from interventions to improve the public's health could include changes in knowledge and behaviour related to health, societal changes (for example, a reduction in crime rates) and a change in people's health and wellbeing or health status. In clinical terms, outcomes could include the number of patients who fully recover from an illness or the number of hospital admissions, and an improvement or deterioration in someone's health, functional ability, symptoms or situation. Researchers should decide what outcomes to measure before a study begins.
P value	The p value is a statistical measure that indicates whether or not an effect is statistically significant. For example, if a study comparing 2 treatments found that one seems more effective than the other, the p value is the probability of obtaining these, or more extreme results by chance. By convention, if the p value is below 0.05 (that is, there is less than a 5% probability that the results occurred by chance) it is considered that there probably is a real difference between treatments. If the p value is 0.001 or less (less than a 1% probability that the results occurred by chance), the result is seen as highly significant. If the p value shows that there is likely to be a difference between treatments, the confidence interval describes how big the difference in effect might be.
Placebo	A fake (or dummy) treatment given to participants in the control group of a clinical trial. It is indistinguishable from the actual treatment (which is given to participants in the experimental group). The aim is to determine what effect the experimental treatment has had – over and above any placebo effect caused because someone has received (or thinks they have received) care or attention.
Polypharmacy	The use or prescription of multiple medications.
Positive predictive value (PPV)	In screening or diagnostic tests: A measure of the usefulness of a screening or diagnostic test. It is the proportion of those with a positive test result who have the disease, and can be interpreted as the probability that a positive test result is correct. It is calculated as follows: $TP/(TP+FP)$
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Prevalence	See Pre-test probability.
Primary care	Healthcare delivered outside hospitals. It includes a range of services provided by GPs, nurses, health visitors, midwives and other healthcare professionals and allied health professionals such as dentists, pharmacists and opticians.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Probabilistic analysis	In economic evaluation, this is an analysis that uses a probability distribution for each input. In contrast, see Deterministic analysis.
Product licence	An authorisation from the MHRA to market a medicinal product.

Term	Definition
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good prognosis is associated with low rate of undesirable outcomes; poor prognosis is associated with a high rate of undesirable outcomes.
Prospective study	A research study in which the health or other characteristic of participants is monitored (or 'followed up') for a period of time, with events recorded as they happen. This contrasts with retrospective studies.
Publication bias	Publication bias occurs when researchers publish the results of studies showing that a treatment works well and don't publish those showing it did not have any effect. If this happens, analysis of the published results will not give an accurate idea of how well the treatment works. This type of bias can be assessed by a funnel plot.
Quality of life	See 'Health-related quality of life'.
Quality-adjusted life year (QALY)	A measure of the state of health of a person or group in which the benefits, in terms of length of life, are adjusted to reflect the quality of life. One QALY is equal to 1 year of life in perfect health. QALYS are calculated by estimating the years of life remaining for a patient following a particular treatment or intervention and weighting each year with a quality of life score (on a scale of 0 to 1). It is often measured in terms of the person's ability to perform the activities of daily life, freedom from pain and mental disturbance.
Randomisation	Assigning participants in a research study to different groups without taking any similarities or differences between them into account. For example, it could involve using a random numbers table or a computer-generated random sequence. It means that each individual (or each group in the case of cluster randomisation) has the same chance of receiving each intervention.
Randomised controlled trial (RCT)	A study in which a number of similar people are randomly assigned to 2 (or more) groups to test a specific drug or treatment. One group (the experimental group) receives the treatment being tested, the other (the comparison or control group) receives an alternative treatment, a dummy treatment (placebo) or no treatment at all. The groups are followed up to see how effective the experimental treatment was. Outcomes are measured at specific times and any difference in response between the groups is assessed statistically. This method is also used to reduce bias.
RCT	See 'Randomised controlled trial'.
Receiver operated characteristic (ROC) curve	A graphical method of assessing the accuracy of a diagnostic test. Sensitivity is plotted against 1 minus specificity. A perfect test will have a positive, vertical linear slope starting at the origin. A good test will be somewhere close to this ideal.
Reference standard	The test that is considered to be the best available method to establish the presence or absence of the outcome – this may not be the one that is routinely used in practice.
Reliability	How dependably and consistently a test measures a characteristic.
Reporting bias	See 'Publication bias'.
Resource implication	The likely impact in terms of finance, workforce or other NHS resources.
Responsiveness to change	How much change in the test would lead to a clinically important difference in the experience of person.

Term	Definition
Retrospective study	A research study that focuses on the past and present. The study examines past exposure to suspected risk factors for the disease or condition. Unlike prospective studies, it does not cover events that occur after the study group is selected.
Review question	In guideline development, this term refers to the questions about treatment and care that are formulated to guide the development of evidence-based recommendations.
Risk ratio (RR)	<p>The ratio of the risk of disease or death among those exposed to certain conditions compared with the risk for those who are not exposed to the same conditions (for example, the risk of people who smoke getting lung cancer compared with the risk for people who do not smoke).</p> <p>If both groups face the same level of risk, the risk ratio is 1. If the first group had a risk ratio of 2, subjects in that group would be twice as likely to have the event happen. A risk ratio of less than 1 means the outcome is less likely in the first group. The risk ratio is sometimes referred to as relative risk.</p>
Secondary outcome	An outcome used to evaluate additional effects of the intervention deemed a priori as being less important than the primary outcomes.
Selection bias	<p>Selection bias occurs if:</p> <ul style="list-style-type: none"> a) The characteristics of the people selected for a study differ from the wider population from which they have been drawn, or b) There are differences between groups of participants in a study in terms of how likely they are to get better.
Sensitivity	<p>How well a test detects the thing it is testing for.</p> <p>If a diagnostic test for a disease has high sensitivity, it is likely to pick up all cases of the disease in people who have it (that is, give a 'true positive' result). But if a test is too sensitive it will sometimes also give a positive result in people who don't have the disease (that is, give a 'false positive').</p> <p>For example, if a test were developed to detect if a woman is 6 months pregnant, a very sensitive test would detect everyone who was 6 months pregnant, but would probably also include those who are 5 and 7 months pregnant.</p> <p>If the same test were more specific (sometimes referred to as having higher specificity), it would detect only those who are 6 months pregnant, and someone who was 5 months pregnant would get a negative result (a 'true negative'). But it would probably also miss some people who were 6 months pregnant (that is, give a 'false negative').</p> <p>Breast screening is a 'real-life' example. The number of women who are recalled for a second breast screening test is relatively high because the test is very sensitive. If it were made more specific, people who don't have the disease would be less likely to be called back for a second test but more women who have the disease would be missed.</p>
Sensitivity analysis	A means of representing uncertainty in the results of economic evaluations. Uncertainty may arise from missing data, imprecise estimates or methodological controversy. Sensitivity analysis also allows for exploring the generalisability of results to other settings. The analysis is repeated using different assumptions to examine the effect on the results.

Term	Definition
	<p>One-way simple sensitivity analysis (univariate analysis): each parameter is varied individually in order to isolate the consequences of each parameter on the results of the study.</p> <p>Multi-way simple sensitivity analysis (scenario analysis): 2 or more parameters are varied at the same time and the overall effect on the results is evaluated.</p> <p>Threshold sensitivity analysis: the critical value of parameters above or below which the conclusions of the study will change are identified.</p> <p>Probabilistic sensitivity analysis: probability distributions are assigned to the uncertain parameters and are incorporated into evaluation models based on decision analytical techniques (for example, Monte Carlo simulation).</p>
Significance (statistical)	A result is deemed statistically significant if the probability of the result occurring by chance is less than 1 in 20 ($p < 0.05$).
Specificity	<p>The proportion of true negatives that are correctly identified as such. For example in diagnostic testing the specificity is the proportion of non-cases correctly diagnosed as non-cases.</p> <p>See related term 'Sensitivity'.</p> <p>In terms of literature searching a highly specific search is generally narrow and aimed at picking up the key papers in a field and avoiding a wide range of papers.</p>
Stakeholder	<p>An organisation with an interest in a topic that NICE is developing a guideline or piece of public health guidance on. Organisations that register as stakeholders can comment on the draft scope and the draft guidance. Stakeholders may be:</p> <ul style="list-style-type: none"> • manufacturers of drugs or equipment • national patient and carer organisations • NHS organisations • organisations representing healthcare professionals.
State transition model	See Markov model
Stratification	When a different estimate effect is thought to underlie two or more groups based on the PICO characteristics. The groups are therefore kept separate from the outset and are not combined in a meta-analysis, for example; children and adults. Specified a priori in the protocol.
Sub-groups	Planned statistical investigations if heterogeneity is found in the meta-analysis. Specified a priori in the protocol.
Systematic review	A review in which evidence from scientific studies has been identified, appraised and synthesised in a methodical way according to predetermined criteria. It may include a meta-analysis.
Test-retest reliability	The reliability of the test measured over time (give the same person the same test at different times and see if the scores are the same).
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Transition probability	In a state transition model (Markov model), this is the probability of moving from one health state to another over a specific period of time.
Treatment allocation	Assigning a participant to a particular arm of a trial.
Univariate	Analysis which separately explores each variable in a data set.
Utility	In health economics, a 'utility' is the measure of the preference or value that an individual or society places upon a particular health

Term	Definition
	state. It is generally a number between 0 (representing death) and 1 (perfect health). The most widely used measure of benefit in cost–utility analysis is the quality-adjusted life year, but other measures include disability-adjusted life years (DALYs) and healthy year equivalents (HYEs).
Validity	The extent to which a test accurately measures what it is meant to measure.

1
2

References

1. GRADE Working Group. The Grading of Recommendations Assessment, Development and Evaluation (GRADE) Working Group website. 2011. Available from: <http://www.gradeworkinggroup.org/> Last accessed:
2. Maruish ME. User's Manual for the SF-36v2 Health Survey. 3rd ed. Quality Metric Incorporated. 2011. Available from: <https://books.google.co.uk/books?id=a0vYnQEACAAJ>
3. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *Journal of Clinical Epidemiology*. 2016; 75:40-46
4. National Institute for Health and Care Excellence. Developing NICE guidelines: the manual [updated January 2022]. London. National Institute for Health and Care Excellence, 2014. Available from: <https://www.nice.org.uk/process/pmg20>
5. National Institute for Health and Care Excellence. Developing NICE guidelines: the manual [updated October 2018]. London. National Institute for Health and Care Excellence, 2014. Available from: <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20oerview>
6. National Institute for Health and Care Excellence. The NICE Charter. 2020. Available from: <https://www.nice.org.uk/about/who-we-are/our-charter> Last accessed: 24/03/2022.
7. National Institute for Health and Clinical Excellence. Guide to the methods of technology appraisal 2013. London. National Institute for Health and Clinical Excellence, 2013. Available from: <http://publications.nice.org.uk/pmg9>
8. National Institute for Health and Clinical Excellence. Our principles. London. National Institute for Health and Clinical Excellence, 2020. Available from: <https://www.nice.org.uk/about/who-we-are/our-principles#introduction>
9. Novielli N, Cooper NJ, Abrams KR, Sutton AJ. How is evidence on test performance synthesized for economic decision models of diagnostic tests? A systematic appraisal of Health Technology Assessments in the UK since 1997. *Value in Health*. 2010; 13(8):952-957
10. Organisation for Economic Co-operation and Development (OECD). Purchasing power parities (PPP). 2012. Available from: <http://www.oecd.org/std/ppp> Last accessed: 20 January 2022.
11. Review Manager (RevMan) [Computer program]. Version 5. Copenhagen. The Nordic Cochrane Centre, The Cochrane Collaboration, 2015. Available from: <http://tech.cochrane.org/Revman>
12. WinBUGS [Computer programme] version 1.4. Cambridge. MRC Biostatistics Unit University of Cambridge, 2015. Available from: <http://www.mrc-bsu.cam.ac.uk/software/bugs/the-bugs-project-winbugs/>