

1 **Physical activity and the**
2 **environment update**

3 **Evidence Reviews – Appendix 4**

4 **GRADE Evidence Profiles**

5 **DRAFT**

6 *Jean Bennie, Olivia Crane, Adrienne Cullum, Karen*

7 *Peploe, Clare Wohlgemuth*

8 *July 2017*

9

10 *National Institute for Health and Care Excellence*

11

12 **Effectiveness and cost-effectiveness of public open space** 13 **and transport interventions: GRADE evidence profiles**

14 **1. Introduction**

15 As discussed at PHAC 1 (November 2016), the outcomes reported in the evidence reviews
16 are being appraised and presented using GRADE (Grading of Recommendations
17 Assessment, Development and Evaluation). This approach to assessing the quality of a
18 body of evidence and has been used in development of NICE clinical guidelines for a
19 number of years. The evidence is rated across studies for specific outcomes as opposed to
20 rating study by study. This approach assesses consistency of results across different
21 studies, provided the studies are measuring the same outcome.

22
23 GRADE looks at “the extent to which one can be confident that an estimate of effect of
24 association is close to the quantity of specific interest”¹. GRADE is concerned with how
25 certain we can be that the observed effect is close to the real effect. When using GRADE we
26 consider the risk of bias, consistency, directness, and precision of the studies reporting on a
27 particular outcome. The evidence regarding that outcome is then graded as either of **very**
28 **low, low, moderate, or high quality**.

29

30 **2. GRADE and Physical Activity Outcomes**

31 In order to apply GRADE consistently across outcomes, the PHAC considered the Minimal
32 Important Difference (MID), defined as *the smallest change in an outcome that is considered*
33 *important by patients or health care professionals*. At PHAC 2, it was agreed that for this
34 topic MID would be any change observed as a result of an intervention. It was discussed that
35 in certain population groups the smallest of changes in activity would benefit health and
36 wellbeing.

37

38 As highlighted in the protocol and [scope](#), the outcomes for Evidence Review 1 are:

39 *Primary outcomes*

- 40 • total physical activity (PA) (measured by, for example, time/ distance/ number of
41 steps/ levels of activity/ levels of recommended PA)
- 42 • total sedentary time (measured by time)
- 43 • domain-specific physical activity levels (active travel or physical activity in everyday
44 life, such as measures of walking, cycling or active play)
- 45 • public transport use (proxy measure of PA)

¹ Higgins JPT, Green S (editors). 12.2.1 The GRADE Approach. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.handbook.cochrane.org

Physical Activity and the Environment – Appendix 4: GRADE profiles

46 *Secondary outcomes*

- 47 • changes to road environment (such as introduction of traffic calming measures)
- 48 • changes to transport (such as changes in modal share)
- 49 • vehicle speeds
- 50 • car use

51 The committee agreed at PHAC 1 that all primary outcomes were to be regarded as critical
52 outcomes and all secondary outcomes were to be regarded as important outcomes.

53

54 The GRADE evidence profiles presented below show the appraised outcomes for Reviews
55 1, 2 and 3. All of the studies included in these reviews were non-randomised, therefore,
56 using GRADE, ratings start at “low” for evidence derived from observational studies..

57

58

59 Details of how the evidence for each outcome was appraised across each of the quality
60 domains is given below:

Quality domain	Description
Risk of bias	Limitations in study design and implementation may bias the estimates of the treatment effect. Major limitations in studies decrease the confidence in the estimate of the effect. Examples of such limitations are selection bias (often due to poor allocation concealment), performance and detection bias (often due to a lack of blinding of the participants, healthcare professional or assessor) and attrition bias (due to missing data causing systematic bias in the analysis). Where there are no study limitations, evidence is assessed as having ‘no serious’ risk of bias. Alternatively, evidence may be downgraded one level (‘serious’ risk of bias) or two levels (‘very serious’ risk of bias).
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question. Where the evidence is directly applicable to the PICO, it is assessed as having ‘no serious’ risk of indirectness. Alternatively, evidence may be downgraded one level (‘serious’ risk of indirectness) or two levels (‘very serious’ risk of indirectness).
Inconsistency	<p>Inconsistency refers to an unexplained heterogeneity of effect estimates between studies combined into the same GRADE profile due to presenting the same outcomes in the same way. If pooled in a meta-analysis, the I^2 statistic describes the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance).</p> <p>For the purposes of this review, as it was rarely possible to combine results into a meta-analysis, the committee agreed that the heterogeneity of the results be considered by the reviewers. Where heterogeneity could be explained by differences in study design, content of interventions and comparators, it was assessed as having ‘no serious’ inconsistency’. Where inconsistency could not be explained by these factors, results could be downgraded by one level for some heterogeneity, and by two levels for a large amount of unexplained heterogeneity.</p>

Physical Activity and the Environment – Appendix 4: GRADE profiles

Quality domain	Description
Imprecision	<p>95% confidence intervals denote the possible range of locations of the true population effect at a 95% probability, and so wide confidence intervals may denote a result that is consistent with conflicting interpretations (for example a result may be consistent with both public health benefit AND public health harm) and thus be imprecise. Imprecision can be caused by studies having relatively few individuals or few events.</p> <p>For the purpose of these reviews, the committee chose a Minimal Important Difference (MID) of any change, as any change at a population level was considered to be meaningful. Imprecision was judged using the following criteria;</p> <ul style="list-style-type: none"> • Where confidence intervals for an effect spanned the line of no effect, the result was downgraded for ‘serious’ imprecision because we are uncertain of where the true effect lies. • Where confidence intervals for an effect do not span the line of no effect, we are certain of where the true effect lies, and so there is no serious imprecision. <p>For the majority of studies reviewed for this guideline it was not possible to assess the degree of precision due to confidence intervals not being reported or able to be imputed from information provided in the paper. However, the committee considered it important to note that in public health, the approach often taken to assess whether an intervention is effective is one of hypothesis testing using P Values. P Values denote the probability of obtaining a given result assuming the null hypothesis (no effect) is true. For example, assuming the intervention has no effect, a p value of 0.05 means you would obtain the observed difference (or more) in 5% of studies due to random sampling error. The committee appreciated this is different from assessing the precision of an effect. In these instances, the P Value was used as follows;</p> <ul style="list-style-type: none"> • Where the P Value for an effect was >0.05, the confidence intervals would be likely to span the line of no effect. The result was downgraded for ‘serious’ imprecision. • Where the P Value for an effect was <0.05, the confidence intervals would be unlikely to span the line of no effect. The result was not downgraded as it had ‘no serious’ imprecision. <p>Where a study did not provide confidence intervals for an effect (or information to work this out), or P Values, confidence intervals of change over time in the intervention group and change over time in the control group were calculated and compared. Results were judged as follows:</p> <ul style="list-style-type: none"> • Where these confidence intervals overlapped each other, it was judged that the results were not significantly different between groups. The result was downgraded for ‘serious’ imprecision as we are not certain where the true effect is. • Where confidence intervals did not overlap each other, it was judged that the results were significantly different. The result was not downgraded as it had ‘no serious’ imprecision as we are certain where the true effect is. <p>Where a study did not provide confidence intervals, P-Values, or sufficient data to be able to calculate these, the review team could not be certain that the result was meaningful. Therefore the result was downgraded for serious imprecision.</p>

Physical Activity and the Environment – Appendix 4: GRADE profiles

Quality domain	Description
Other issues	<p>Sometimes randomisation may not adequately lead to group equivalence of confounders, and if so this may lead to bias, which should be taken into account. Potential conflicts of interest, often caused by excessive pharmaceutical company involvement in the publication of a study, should also be noted.</p> <p>The option to upgrade confidence in the evidence by one level for consistency was applied. Evidence is upgraded for consistency if a number of studies from different settings investigating the same intervention report the same outcome and show the same direction of effect.</p>

61

62

63

Physical Activity and the Environment – Appendix 4: GRADE profiles

64 **Review 1**

65 **Congestion charging**

66 [To note that all studies on congestion charging were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality								
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control										
Congestion charging																		
Total physical activity as measured by total time spent in physical activity (PA)																		
1 Bergman 2010	Non-randomised controlled study	No Serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	165	138	<p><u>Change in time spent in physical activity (self-reported using IPAQ questionnaire) (intervention vs control) (baseline to 5-month follow-up):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Control</th> <th>Intervention</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Overall PA</td> <td>No difference</td> <td>Increase</td> <td>0.015</td> </tr> </tbody> </table> <p>There was a statistically significant increase in physical activity compared to the control group</p>		Control	Intervention	P value	Overall PA	No difference	Increase	0.015	LOW
	Control	Intervention	P value															
Overall PA	No difference	Increase	0.015															
Total physical activity as measured by total time spent on moderate PA (4 METS)																		
1 Bergman 2010	Non-randomised controlled study	No Serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	165	138	<p><u>Change in time spent on moderate PA (self-reported using IPAQ questionnaire) (intervention vs control) (baseline to 5-month follow-up):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Control</th> <th>Intervention</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Moderate PA</td> <td>No difference</td> <td>Increase</td> <td>0.036</td> </tr> </tbody> </table> <p>There was a statistically significant increase in moderate physical activity compared to the control group</p>		Control	Intervention	P value	Moderate PA	No difference	Increase	0.036	LOW
	Control	Intervention	P value															
Moderate PA	No difference	Increase	0.036															
Total sedentary time as measured by the total time spent sitting																		

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Bergman 2010	Non-randomised controlled study	No Serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	165	138	<p><u>Change in total time spent sitting (self-reported using IPAQ questionnaire) (intervention vs control) (baseline to 5-month follow-up):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Control</th> <th>Intervention</th> <th>P value</th> <th>Effect size -r²</th> </tr> </thead> <tbody> <tr> <td>Sitting</td> <td>No difference</td> <td>Decreased</td> <td>0.009</td> <td>0.03</td> </tr> </tbody> </table> <p>Subjects in the intervention group reported statistically significant less time spent sitting compared to the comparator group</p>		Control	Intervention	P value	Effect size -r ²	Sitting	No difference	Decreased	0.009	0.03	LOW
	Control	Intervention	P value	Effect size -r ²																
Sitting	No difference	Decreased	0.009	0.03																
Changes to transport as measured by % of car drivers switching to public transport																				
2 Transport for London 2008 Karlstrom and Franklin 2009	Non-randomised uncontrolled studies	No serious risk of bias ³	No serious inconsistency	No serious indirectness	Serious imprecision ⁴	None	1550	n/a	<p><u>Percentage of car drivers switching to public transport (self-reported) (baseline to 18 months follow-up)</u></p> <p>About 25% of car drivers crossing the toll cordon (treated individuals) switch to public transport compared to 10% in the control group (car drivers not crossing the toll cordon). Initial car drivers crossing the toll cordon had a 15% higher rate of switching to public transport compared with those car drivers not crossing the cordon. There is a background rate of 8-11% of all travellers switching mode between baseline and follow-up, indicating a range of factors influence transport mode choice.</p> <p><u>Proportion of survey respondents switching transport mode post London congestion charge (follow-up period not clear) (self-reported) (intervention group only):</u></p> <p>Of residents living outside of charging zone, around half would not continue driving to the extension zone. Of these, 40% are estimated to have changed transport method as a result of the charge.</p>	VERY LOW										
Changes to transport as measured by traffic make-up on the road and bus patronage, various data collection methods, 12 months follow-up (important)																				
1 Transport for London	Non-randomised uncontrolled study	Serious risk of bias ⁵	Not assessable as single study	No serious indirectness	Serious imprecision ⁶	None	n/a		<p><u>Change in vehicles using the free passage route (percentage change between baseline and 12-month follow-up figures) (intervention only):</u></p> <table border="1"> <thead> <tr> <th>Chargeable</th> <th>2005-2006 & 2007 % change</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> </tr> </tbody> </table>	Chargeable	2005-2006 & 2007 % change			VERY LOW						
Chargeable	2005-2006 & 2007 % change																			

² An effect size of up to 0.1 is considered small and around 0.3 is considered moderate. Above 0.5 is considered a large effect, therefore effect size was small

³ Quality score allocated as (-) and (+) indicating some risk of bias – not downgraded.

⁴ Unable to tell whether intervention is effective as no measure of variance reported or p values – downgraded one level

⁵ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is in data collection methods.

⁶ Unable to tell whether intervention is effective as no measure of variance reported or p values – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

								<table border="1"> <tr> <td>Car and minicabs</td> <td>-3%</td> </tr> <tr> <td>Vans and lorries</td> <td>+7%</td> </tr> <tr> <td>Non chargeable</td> <td></td> </tr> <tr> <td>Licensed taxis</td> <td>+9%</td> </tr> <tr> <td>Two wheelers</td> <td>+12%</td> </tr> <tr> <td>Pedal cycles</td> <td>+18%</td> </tr> </table>	Car and minicabs	-3%	Vans and lorries	+7%	Non chargeable		Licensed taxis	+9%	Two wheelers	+12%	Pedal cycles	+18%																			
Car and minicabs	-3%																																						
Vans and lorries	+7%																																						
Non chargeable																																							
Licensed taxis	+9%																																						
Two wheelers	+12%																																						
Pedal cycles	+18%																																						
<p><u>Traffic make-up in vehicle-kilometres driven (% of total) within western extension zone during charging hours (direct observation):</u></p> <table border="1"> <thead> <tr> <th>Chargeable</th> <th>% change</th> <th>Baseline</th> <th>Follow-up</th> </tr> </thead> <tbody> <tr> <td>Car and minicabs</td> <td>-6%</td> <td>60%</td> <td>54%</td> </tr> <tr> <td>Vans, lorries and others</td> <td>+2.5%</td> <td>13%</td> <td>15%</td> </tr> <tr> <td>Non chargeable</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Licensed taxis</td> <td>+2%</td> <td>11%</td> <td>13%</td> </tr> <tr> <td>Buses and coaches</td> <td>+1%</td> <td>3%</td> <td>4%</td> </tr> <tr> <td>Two wheelers</td> <td>+1%</td> <td>5%</td> <td>6%</td> </tr> <tr> <td>Pedal cycles</td> <td>+1%</td> <td>5%</td> <td>6%</td> </tr> </tbody> </table> <p><u>Bus patronage changes between baseline and 1-year follow-up (record-kept data):</u> Bus passengers entering the charging zone increased by 6% (96,500/day to 102,000 /day) in charging hours, and 9% during morning peak period (34,100 to 37,200) (07:00-10:00). Increases for exiting the charging zone were 5% (90,100 to 94,200) and 2% (24,300 to 24,900) for charging hours and peak hours respectively. Percentages and absolute figures are slightly mismatched, likely to do with rounding of absolute figure</p>								Chargeable	% change	Baseline	Follow-up	Car and minicabs	-6%	60%	54%	Vans, lorries and others	+2.5%	13%	15%	Non chargeable				Licensed taxis	+2%	11%	13%	Buses and coaches	+1%	3%	4%	Two wheelers	+1%	5%	6%	Pedal cycles	+1%	5%	6%
Chargeable	% change	Baseline	Follow-up																																				
Car and minicabs	-6%	60%	54%																																				
Vans, lorries and others	+2.5%	13%	15%																																				
Non chargeable																																							
Licensed taxis	+2%	11%	13%																																				
Buses and coaches	+1%	3%	4%																																				
Two wheelers	+1%	5%	6%																																				
Pedal cycles	+1%	5%	6%																																				
<p>SUMMARY: see evidence statement 1.1</p>																																							

Physical Activity and the Environment – Appendix 4: GRADE profiles

68 **Guided busway⁷**

69 [To note that all studies on the guided busway were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality																							
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																									
Guided Busway																																	
Active travel as measured by the average time spent in active commuting																																	
1	Non-randomised controlled study	Serious risk of bias ⁸	Not assessable as single study	No serious indirectness	No serious imprecision	None	364	n/a	<p><u>Average time (mins in last 7 days) spent in active commuting and physical activity (self-reported) (intervention only) (baseline to 6-18 month follow up):</u></p> <table border="1"> <thead> <tr> <th rowspan="2">Activity</th> <th rowspan="2">% (N) Reporting Any Activity At Baseline</th> <th colspan="3">Time Spent In Activity (Min/M)</th> </tr> <tr> <th>Baseline</th> <th>Follow-Up</th> <th>P-Value</th> </tr> </thead> <tbody> <tr> <td>Active Weekly Commuting</td> <td>77.6 (364)</td> <td>120 (33-200)</td> <td>100 (33-200)</td> <td>0.001</td> </tr> <tr> <td>Walking</td> <td>27.8 (131)</td> <td>0 (0-20)</td> <td>0 (0-40)</td> <td>0.487</td> </tr> <tr> <td>Cycling</td> <td>56.6 (266)</td> <td>70 (0-160)</td> <td>40 (0-160)</td> <td>0.016</td> </tr> </tbody> </table> <p>Time spent on active commuting decreased significantly at follow-up, largely attributed to the decrease in median time spent on cycling</p> <p><u>Association between exposure to intervention (measured as proximity of participants' residence to the guided bus way) and PA:</u> Exposure to the busway was associated with a significantly greater likelihood of an increase in weekly cycle commuting</p>	Activity	% (N) Reporting Any Activity At Baseline	Time Spent In Activity (Min/M)			Baseline	Follow-Up	P-Value	Active Weekly Commuting	77.6 (364)	120 (33-200)	100 (33-200)	0.001	Walking	27.8 (131)	0 (0-20)	0 (0-40)	0.487	Cycling	56.6 (266)	70 (0-160)	40 (0-160)	0.016	VERY LOW
Activity	% (N) Reporting Any Activity At Baseline	Time Spent In Activity (Min/M)																															
		Baseline	Follow-Up	P-Value																													
Active Weekly Commuting	77.6 (364)	120 (33-200)	100 (33-200)	0.001																													
Walking	27.8 (131)	0 (0-20)	0 (0-40)	0.487																													
Cycling	56.6 (266)	70 (0-160)	40 (0-160)	0.016																													

⁷ The Guided Busway comprised a new bus network and an adjacent 22km traffic-free walking and cycling route

⁸ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

Physical Activity and the Environment – Appendix 4: GRADE profiles

									time (relative risk ratio [RRR] 1.34, 95% CI1.03, 1.76), but not for active commuting as a whole and walking.																										
Physical activity in everyday life as measured by the average time spent in recreational walking and cycling																																			
1	Panter et al 2016	Non-randomised controlled study	Serious risk of bias ⁹	Not assessable as 1 study	No serious risk of indirectness	Serious imprecision ¹⁰	None	364	n/a	<p><u>Change in average time (mins in past 7 days) spent in walking and cycling for recreation and physical activity (self-reported) (intervention only) (baseline to 6-18 month follow-up):</u></p> <table border="1"> <thead> <tr> <th rowspan="2">ACTIVITY</th> <th rowspan="2">% (N) REPORTING ANY ACTIVITY AT BASELINE</th> <th colspan="3">TIME SPENT IN ACTIVITY (MIN/WEEK) MEDIAN (IQR)</th> <th rowspan="2">P-Value</th> </tr> <tr> <th>Baseline</th> <th>Follow-Up</th> <th></th> </tr> </thead> <tbody> <tr> <td>RECREATION (TOTAL)</td> <td>83.3 (391)</td> <td>75 (28-150)</td> <td>79 (30,180)</td> <td>0.640</td> </tr> <tr> <td>Walking</td> <td>78.0 (366)</td> <td>57 (15-135)</td> <td>60 (0,150)</td> <td>0.551</td> </tr> <tr> <td>Cycling</td> <td>32.6 (153)</td> <td>0 (0-22.5)</td> <td>0 (0,19)</td> <td>0.416</td> </tr> </tbody> </table> <p>There was no significant difference between the time spent walking and cycling for recreation at follow up compared to baseline</p>	ACTIVITY	% (N) REPORTING ANY ACTIVITY AT BASELINE	TIME SPENT IN ACTIVITY (MIN/WEEK) MEDIAN (IQR)			P-Value	Baseline	Follow-Up		RECREATION (TOTAL)	83.3 (391)	75 (28-150)	79 (30,180)	0.640	Walking	78.0 (366)	57 (15-135)	60 (0,150)	0.551	Cycling	32.6 (153)	0 (0-22.5)	0 (0,19)	0.416	VERY LOW
ACTIVITY	% (N) REPORTING ANY ACTIVITY AT BASELINE	TIME SPENT IN ACTIVITY (MIN/WEEK) MEDIAN (IQR)			P-Value																														
		Baseline	Follow-Up																																
RECREATION (TOTAL)	83.3 (391)	75 (28-150)	79 (30,180)	0.640																															
Walking	78.0 (366)	57 (15-135)	60 (0,150)	0.551																															
Cycling	32.6 (153)	0 (0-22.5)	0 (0,19)	0.416																															
Changes to transport as measured by changes in proportion of journeys to work made by active travel (proximity)																																			
1	Heinen et al 2015	Non-randomised controlled study	Serious risk of bias ¹¹	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	470	n/a	<p><u>Change in proportion of all journeys to work in past 7 days made by active modes of travel as opposed to non-active modes such as car (self-reported) (intervention vs control) (baseline to 3-year follow-up) [Relative Risk Ratio and 95% confidence interval]:</u></p> <p>Proximity to the busway was a predictor of large and significant increase in active travel (1.80 [1.27 to 2.55] p<0.01) and reduced the likelihood of a small decrease in</p>	VERY LOW																								

⁹ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

¹⁰ No significant effect of intervention as P values greater than 0.05– downgraded one level

¹¹ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

Physical Activity and the Environment – Appendix 4: GRADE profiles

									active travel as a proportion of all journeys made (0.47 [0.28 to 0.81]) i.e. commuters living 4 km from the busway were almost twice as likely to report a substantial increase in active travel as a proportion of all journeys, and half as likely to report a small decrease, than those living 9 km away.	
Changes to transport as measured by changes in proportion of journeys to work made by active travel (urban / rural)										
1 Heinen et al 2015	Non-randomised controlled study	Serious risk of bias ¹²	Not assessable as measuring different outcomes	No serious indirectness	No serious imprecision	None	470	NA	<u>Urban / Rural (self-reported) (baseline to 3-year follow-up):</u> Living in villages or smaller settlements rather than urban areas predicted an increase in public transport trips as a proportion of all commuting trips (RRR 2.53 (1.06, 6.05), pp<0.05)	VERY LOW
Changes to transport as measured by changes in proportion of journeys to work made by active travel (baseline active commuting)										
1 Panter et al 2016	Non-randomised controlled study	Serious risk of bias ¹³	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	364	NA	<u>Active commuting changes by baseline level of active commuting:</u> The intervention had a significant effect on total active commuting only for those who reported the lowest levels of active commuting at baseline (RRR 1.76, 95% CI 1.16, 2.67).	VERY LOW
SUMMARY: see evidence statement 1.2										

70

71

72 **Upgrading of bus routes¹⁴**

73 [To note that all studies on upgrading bus routes were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment	No. of participants	Effect	Quality
--------------------	---------------------	--------	---------

¹² Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

¹³ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

¹⁴ This includes interventions to upgrade stops to show real-time passenger information, and increase bus frequency

Physical Activity and the Environment – Appendix 4: GRADE profiles

No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Upgrading of bus routes										
Public transport use (as a proxy of physical activity) as measured by bus use										
1 Loader and Stanley 2009	Non-randomised controlled study	Serious risk of bias ¹⁵	Not assessable as 1 study	No serious risk of indirectness	Serious risk of imprecision ¹⁶	None	Unclear ¹⁷	Unclear ¹⁸	<p><u>Change in bus use (patronage) between baseline and 10-year follow-up (counts) (intervention vs control):</u> Follow-up data shows intervention total bus patronage growth of 4.6% between baseline and 1-year follow-up. Control routes grew by 1.3% in the same period.</p> <p><u>Change in bus use (patronage) between baseline and 10-year follow-up (counts) by area (intervention vs control):</u> Greatest increases in use of intervention routes are seen in the Central Business District (CBD) and outer regions (13.8% and 10.8% respectively). A decrease is seen in usage of unchanged routes in the outer area (-0.9%).</p> <p><u>Bus patronage growth by time, Saturdays only (baseline to 1-year follow-up):</u> For intervention buses whose finishing times had previously been between 4pm and 5pm (n = 2), their afternoon validations “more than doubled” after extension of running hours. For buses whose previous finishing time was between 5pm and 6pm, afternoon demand increased by around 20%.</p>	VERY LOW
SUMMARY: see evidence statement 1.3										

74

75 **New light rail service**

76 [To note that all studies on the new light rail service were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as ‘low’].

Quality assessment	No. of participants	Effect	Quality

¹⁵ Quality score was (-) indicating high risk of bias - downgraded one level. Main bias was large loss to follow up.

¹⁶ No measures of variance provided – downgraded one lev

¹⁷ Participants numbers were not given

¹⁸ Participants numbers were not given

Physical Activity and the Environment – Appendix 4: GRADE profiles

No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control												
New light rail service																				
Public transport use (as a proxy measured of physical activity) as measured with train trips, walk trips and walk minutes																				
1 Boarnet 2013	Non randomised controlled study	No serious risk of bias	Not assessable as 1 study	No Serious indirectness	Serious imprecision ¹⁹	None	103	101	Change in train trips, walk trips, and walk minutes by household (intervention vs control) (baseline to 3-7 month follow-up) (self-reported data): Between group differences were not significantly different for train trips, walk trips, or walk minutes between baseline and follow-up in intervention group and control group. Both reportedly increased over time by a similar amount.	VERY LOW										
Public transport use (as a proxy measured of physical activity) as measured by 7-day accelerometer wear																				
1 Boarnet 2013	Non randomised controlled study	No serious risk of bias	Not assessable as single study	No serious indirectness	Serious imprecision ²⁰	None	103	101	<p><u>Moderate to Vigorous Physical Activity in average minutes over past 7 days (accelerometer data) (intervention vs control) (data from 3-7 month follow-up):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Intervention</th> <th>Control</th> <th>Mean difference</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>MVPA</td> <td>22.04</td> <td>18.73</td> <td>3.31</td> <td>0.674</td> </tr> </tbody> </table> <p>No difference between baseline and follow-up PA for either group</p>		Intervention	Control	Mean difference	P value	MVPA	22.04	18.73	3.31	0.674	VERY LOW
	Intervention	Control	Mean difference	P value																
MVPA	22.04	18.73	3.31	0.674																
SUMMARY: see evidence statement 1.4																				

77

¹⁹ Unable to tell whether the intervention had a significant effect as no difference between intervention and control – downgraded one level

²⁰ P values greater than 0.05, showing no significant effect of intervention – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

78 **New rail stop**

79 [To note that all studies on new rail stops were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality										
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control												
New Rail Stop																				
Public transport use (as a proxy measure of physical activity) as measured by rail ridership																				
1 ²¹ Brown and Werner (2007) Brown and Werner (2009)	Non-randomised uncontrolled studies	Serious risk of bias ²²	Not assessable is one study	No serious indirectness	No Serious imprecision	None	51	n/a	<u>Percentage of people travelling by rail in past 2 weeks (baseline to 7-11 month follow-up) (intervention only) (self-reported data):</u> <table border="1"> <thead> <tr> <th></th> <th>Before</th> <th>After</th> <th>% Change</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Participants Riding The Rail*</td> <td>50%</td> <td>68.75%</td> <td>37.5%</td> <td>0.001</td> </tr> </tbody> </table>		Before	After	% Change	P Value	Participants Riding The Rail*	50%	68.75%	37.5%	0.001	VERY LOW
	Before	After	% Change	P Value																
Participants Riding The Rail*	50%	68.75%	37.5%	0.001																
1 ²³ Brown and Werner (2007) Brown and Werner (2009)	Non-randomised uncontrolled studies	Serious risk of bias ²⁴	Not assessable is one study	No serious indirectness	Serious imprecision ²⁵	None	51	n/a	<u>Mean rail rides (baseline to 7-11 month follow-up) (intervention only) (self-reported data):</u> <table border="1"> <thead> <tr> <th></th> <th>Before</th> <th>After</th> <th>Mean difference (95% CI)*</th> </tr> </thead> <tbody> <tr> <td>Mean Rail Rides</td> <td>3.72 (Sd 6.46)</td> <td>5.02 (Sd 7.90)</td> <td>1.30 (-1.50, 4.10)</td> </tr> </tbody> </table>		Before	After	Mean difference (95% CI)*	Mean Rail Rides	3.72 (Sd 6.46)	5.02 (Sd 7.90)	1.30 (-1.50, 4.10)	VERY LOW		
	Before	After	Mean difference (95% CI)*																	
Mean Rail Rides	3.72 (Sd 6.46)	5.02 (Sd 7.90)	1.30 (-1.50, 4.10)																	
Evidence shows the addition of a rail stop significantly increased train use.																				
The mean difference over time was not significant *Calculated by reviewers.																				

²¹ One study but two publications

²² Quality score was (-) indicating high risk of bias – downgraded one level. Main bias was data collection methods and dropout.

²³ One study but two publications

²⁴ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias was data collection methods and dropout.

²⁵ Confidence intervals span the MID, therefore downgraded one level on imprecision

Physical Activity and the Environment – Appendix 4: GRADE profiles

Total physical activity as measured by an accelerometer per hour																		
1	As reported in Brown and Werner (2007) and Brown and Werner (2009)	Non-randomised uncontrolled studies	Serious risk of bias ²²	Not assessable as is one study	No serious indirectness	Serious imprecision ²⁶	None	51	<p>Change in number of moderate bouts²⁷ of activity per hour (accelerometer data) (baseline to 7-11 month follow-up) (intervention only):</p> <table border="1"> <thead> <tr> <th>(Participants)</th> <th>Before</th> <th>After</th> <th>Mean difference (95% CI)*</th> </tr> </thead> <tbody> <tr> <td>Moderate bouts*/hr (7-11 months follow up)</td> <td>0.06 (sd 0.09)</td> <td>0.06 (sd 0.08)</td> <td>0.00 (-0.03, 0.03)</td> </tr> </tbody> </table> <p>The mean difference over time was not significant.</p> <p>The moderate activity bouts* at baseline were related to bouts at follow-up, and follow-up rail rides ($r=0.46$, $\beta=0.39$, $p=0.01$) and larger households ($r=0.15$, $\beta=0.43$, $p=0.01$) accounted for the significant variance beyond the effects of baseline activity levels.</p> <p>Subgroup analysis shows that number of mean moderate activity bouts is significantly different between groups²⁸ ($p=0.03$). Non riders have the lowest (1.07 (SE 0.76)) followed by new riders (1.77 (SE 0.83)) and then continuing riders (3.68 (SE 0.60)).</p> <p>Proportion of bouts related to walking to a rail stop increased from 0.1 to 0.15 – no statistical significance reported.</p> <p>*Calculated by reviewers.</p>	(Participants)	Before	After	Mean difference (95% CI)*	Moderate bouts*/hr (7-11 months follow up)	0.06 (sd 0.09)	0.06 (sd 0.08)	0.00 (-0.03, 0.03)	VERY LOW
									(Participants)	Before	After	Mean difference (95% CI)*						
Moderate bouts*/hr (7-11 months follow up)	0.06 (sd 0.09)	0.06 (sd 0.08)	0.00 (-0.03, 0.03)															
<p>SUMMARY: see evidence statement 1.5</p>																		

80

81

²⁶ Confidence intervals span the MID, therefore downgraded one level on imprecision

²⁷ Moderate bouts defined as accumulations of 8 or more moderate minutes

²⁸ Subgroup analysis splits population into non-riders, new riders, and continuing riders. Baseline and follow-up results for these individuals are combined.

Physical Activity and the Environment – Appendix 4: GRADE profiles

82 **Complete Street interventions**²⁹

83 [To note that all studies on complete street interventions were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Complete Street interventions										
Total physical activity as measured by accelerometer (counts per minute)										

²⁹ New light rail, new bike lanes, improved pavements

Physical Activity and the Environment – Appendix 4: GRADE profiles

1	As reported in Brown et al 2015 Miller et al 2015	Non-randomised uncontrolled studies	Serious risk of bias ³⁰	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	537	n/a	<p><u>Activity, counts per minute (accelerometer data) (intervention only) (baseline to 7-11 month follow-up):</u></p> <table border="1"> <thead> <tr> <th>Riders (N)</th> <th>Baseline (SE)</th> <th>Follow-Up (SE)</th> <th>Beta³¹</th> <th>95% CI</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Never (393)</td> <td>308.36 (6.63)</td> <td>320.33 (7.11)</td> <td>-</td> <td>-</td> <td>-</td> </tr> <tr> <td>Former (41)</td> <td>391.05 (27.15)</td> <td>376.93 (23.18)</td> <td>-49.35</td> <td>-78.75, -19.94</td> <td>0.001</td> </tr> <tr> <td>Continued (51)</td> <td>361.08 (27.63)</td> <td>317.96 (25.73)</td> <td>-6.25</td> <td>-34.62, 22.12</td> <td>N.S.</td> </tr> <tr> <td>New (52)</td> <td>333.23 (20.75)</td> <td>381.04 (23.73)</td> <td>37.40</td> <td>10.41, 64.39</td> <td>0.007</td> </tr> </tbody> </table> <p>Former riders experienced a significant decline in PA compared to the never riders, new riders accrued statistically significant more PA compared to never riders</p> <p><u>Comparison of average minutes per day of physical activity for days using public transport versus days not using public transport: Point estimates and 95% confidence intervals</u></p> <table border="1"> <thead> <tr> <th>Group (n)</th> <th>Mean mins</th> <th>95% CI</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Overall PA</td> <td colspan="3">Public transport versus non-public transport days in 2012</td> </tr> <tr> <td>public transport days (207)</td> <td>19.65</td> <td>17.28, 22.02</td> <td>0.0001</td> </tr> <tr> <td>Non public transport days (285)</td> <td>9.59</td> <td>7.97, 11.21</td> <td></td> </tr> <tr> <td></td> <td colspan="3">Within-person differences for public transport users</td> </tr> <tr> <td>Public transport days (75)</td> <td>8.54</td> <td>5.00, 12.08</td> <td><0.0001</td> </tr> </tbody> </table> <p>Evidence suggests that subjects are more likely to have higher levels of physical activity on days when they use public transport compared to days when they do not. This difference is statistically significant for the group as well as within person</p>	Riders (N)	Baseline (SE)	Follow-Up (SE)	Beta ³¹	95% CI	P Value	Never (393)	308.36 (6.63)	320.33 (7.11)	-	-	-	Former (41)	391.05 (27.15)	376.93 (23.18)	-49.35	-78.75, -19.94	0.001	Continued (51)	361.08 (27.63)	317.96 (25.73)	-6.25	-34.62, 22.12	N.S.	New (52)	333.23 (20.75)	381.04 (23.73)	37.40	10.41, 64.39	0.007	Group (n)	Mean mins	95% CI	P value	Overall PA	Public transport versus non-public transport days in 2012			public transport days (207)	19.65	17.28, 22.02	0.0001	Non public transport days (285)	9.59	7.97, 11.21			Within-person differences for public transport users			Public transport days (75)	8.54	5.00, 12.08	<0.0001	VERY LOW
										Riders (N)	Baseline (SE)	Follow-Up (SE)	Beta ³¹	95% CI	P Value																																																		
Never (393)	308.36 (6.63)	320.33 (7.11)	-	-	-																																																												
Former (41)	391.05 (27.15)	376.93 (23.18)	-49.35	-78.75, -19.94	0.001																																																												
Continued (51)	361.08 (27.63)	317.96 (25.73)	-6.25	-34.62, 22.12	N.S.																																																												
New (52)	333.23 (20.75)	381.04 (23.73)	37.40	10.41, 64.39	0.007																																																												
Group (n)	Mean mins	95% CI	P value																																																														
Overall PA	Public transport versus non-public transport days in 2012																																																																
public transport days (207)	19.65	17.28, 22.02	0.0001																																																														
Non public transport days (285)	9.59	7.97, 11.21																																																															
	Within-person differences for public transport users																																																																
Public transport days (75)	8.54	5.00, 12.08	<0.0001																																																														

³⁰ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias was data collection methods and dropout.

³¹ Authors tested change scores of the 4 public transport ridership groups with 3 planned comparisons that compared never-riders with former, continuing, and new riders, respectively after adjustment for control variables

Physical Activity and the Environment – Appendix 4: GRADE profiles

Total physical activity as measured by moderate-vigorous physical activity (MVPA ³²)																																																		
1 As reported in Brown et al 2015 Miller et al 2015	Non-randomised uncontrolled studies	Serious risk of bias ³³	Not assessable as 1 study	No serious indirectness	Serious ³⁴ imprecision	None	537	n/a	<p><u>MVPA per 10 hours of accelerometer wear (baseline to 7-11 month follow-up) (intervention group only):</u></p> <table border="1"> <thead> <tr> <th>Riders (N)</th> <th>Beta (Se)</th> <th>95% Ci</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Never Riders (393)</td> <td>-</td> <td>-</td> <td>-</td> </tr> <tr> <td>Former (41)</td> <td>-6.37</td> <td>-10.32, -2.43</td> <td>N.S</td> </tr> <tr> <td>Continued (51)</td> <td>-0.81</td> <td>-4.62, 3.00</td> <td>N.S</td> </tr> <tr> <td>New (52)</td> <td>4.16</td> <td>0.54, 7.78</td> <td><0.05</td> </tr> </tbody> </table> <p>New riders gained 4.16 minutes per 10 hours compared to never riders. Evidence suggests that MVPA was significantly increased in new train users</p> <p><u>Within person differences average mins of PA per 10 hrs wear by year and public transport user group - Point estimates and 95% Cis</u></p> <table border="1"> <thead> <tr> <th>Riders (n)</th> <th>Point Estimate</th> <th>95% CI</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Never Riders (391)</td> <td>1.27</td> <td>-0.60, 3.14</td> <td>0.20</td> </tr> <tr> <td>Continued (51)</td> <td>-2.86</td> <td>-8.60, 2.88</td> <td>0.32</td> </tr> <tr> <td>Former (41)</td> <td>-5.54</td> <td>-11.88, 0.80</td> <td>0.085</td> </tr> <tr> <td>New (52)</td> <td>5.27</td> <td>-1.01, 11.55</td> <td>0.098</td> </tr> </tbody> </table> <p>Evidence suggests there was group to group changes but not within person changes in MVPA.</p>	Riders (N)	Beta (Se)	95% Ci	P Value	Never Riders (393)	-	-	-	Former (41)	-6.37	-10.32, -2.43	N.S	Continued (51)	-0.81	-4.62, 3.00	N.S	New (52)	4.16	0.54, 7.78	<0.05	Riders (n)	Point Estimate	95% CI	P Value	Never Riders (391)	1.27	-0.60, 3.14	0.20	Continued (51)	-2.86	-8.60, 2.88	0.32	Former (41)	-5.54	-11.88, 0.80	0.085	New (52)	5.27	-1.01, 11.55	0.098	VERY LOW
									Riders (N)	Beta (Se)	95% Ci	P Value																																						
Never Riders (393)	-	-	-																																															
Former (41)	-6.37	-10.32, -2.43	N.S																																															
Continued (51)	-0.81	-4.62, 3.00	N.S																																															
New (52)	4.16	0.54, 7.78	<0.05																																															
Riders (n)	Point Estimate	95% CI	P Value																																															
Never Riders (391)	1.27	-0.60, 3.14	0.20																																															
Continued (51)	-2.86	-8.60, 2.88	0.32																																															
Former (41)	-5.54	-11.88, 0.80	0.085																																															
New (52)	5.27	-1.01, 11.55	0.098																																															
Total physical activity as measured by light physical activity (LPA ³⁵) per 10 hours accelerometer wear																																																		

³² The cut off point for MVPA was at least 2020cpm

³³ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is selection bias.

³⁴ P values greater than 0.05, showing no significant effect of intervention– downgraded one level

³⁵ The cut off point for light PA was defined as less than 1000 counts per minute, ≤5 minutes

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Brown et al 2015	Non-randomised uncontrolled studies	Serious risk of bias ³⁶	Not assessable as 1 study	No serious indirectness	Serious Imprecision ³⁷	None	537	n/a	<u>LPA per 10 hours of accelerometer wear (baseline to 7-11 month follow-up) (intervention group only):</u>				VERY LOW
									Group (n)	Beta (SE)	95% CI	P Value	
									Never Riders (393)	-	-	-	
									Former Riders (41)	-9.99(±5.60)	-21, 1.01	0.075	
									Continuing Riders (51)	3.64(±5.41)	-6.98, 3.00		
New Riders (52)	8.67±5.14	-1.43, 18.77	0.092										
There were increases in light PA for new riders and continuing riders, however these were not statistically significant.													
Total sedentary time as measured by sedentary physical activity (SPA ³⁸)													
1 Brown et al 2015	Non-randomised uncontrolled studies	Serious risk of bias ³⁹	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	537	n/a	<u>SPA per 10 hours of accelerometer wear (baseline to 7-11 month follow-up) (intervention group only):</u>				VERY LOW
									Group (n)	Beta (SE)	95% CI	P Value	
									Never Riders (393)	-	-	-	
									Former Riders (41)	16.38 (+/-6.09)	4.41, 28.35	<0.01	
									Continuing Riders (51)	-2.84 (+/-5.88)	-14.39, 8.71		
New Riders (52)	-12.83 (+/-5.59)	-23.82, -1.85	<0.05										
There were significant increases in sedentary PA time for former riders and significant decreases for new riders													
Active transportation as measured by the number of bike trips undertaken													

³⁶ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias was selection bias.

³⁷ Intervention did not have a significant effect: Wide CI intervals that include the null hypothesis and p values greater than 0.05 – downgraded one level

³⁸ The cut off point for sedentary PA was not defined

³⁹ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias was selection bias.

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Brown et al 2016	Non-randomised uncontrolled studies	Serious risk of bias ⁴⁰	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁴¹	None	537	n/a	<p><u>Change in number of bike trips between baseline and 7-11 month follow-up (intervention group only) (self-reported data):</u></p> <p>For those living <800m away from the intervention, there was no significant difference in number bike trips between baseline and follow-up (baseline odds ratio when compared to follow-up 0.86 (95% CI 0.49 to 1.53), p≤0.62). There was also no significant difference in number of bike trips between near and far groups (odds ratio for far group: odds ratio 0.69 (95% 0.37 to 1.3), p≤0.25</p>	VERY LOW																											
Active travel as measured by the change in public transport related physical activity																																					
1 Brown et al 2016	Non-randomised uncontrolled studies	Serious risk of bias ⁴²	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	537	n/a	<p><u>Within person differences average mins of PA per 10 hrs accelerometer wear by year and public transport user group - Point estimates and 95% CIs:</u></p> <table border="1"> <thead> <tr> <th rowspan="2">Riders (n)</th> <th colspan="3">Change in public transport related PA 2012-2013</th> </tr> <tr> <th>Point Estimate</th> <th>95% CI</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Never Riders (391)</td> <td>N/A</td> <td>N/A</td> <td>N/A</td> </tr> <tr> <td>Continued (51)</td> <td>-1.15</td> <td>-3.03, -0.74</td> <td>0.23</td> </tr> <tr> <td>Former (41)</td> <td>-2.34</td> <td>-3.56, -1.08</td> <td>0.0005</td> </tr> <tr> <td>New (52)</td> <td>3.46</td> <td>2.20, 4.72</td> <td><0.0001</td> </tr> <tr> <td>Public transport days (75)</td> <td>8.54</td> <td>5.00, 12.08</td> <td><0.0001</td> </tr> </tbody> </table> <p>Evidence suggests that there were significant changes in public transport related physical activity in all types of riders except for continued riders</p>	Riders (n)	Change in public transport related PA 2012-2013			Point Estimate	95% CI	P Value	Never Riders (391)	N/A	N/A	N/A	Continued (51)	-1.15	-3.03, -0.74	0.23	Former (41)	-2.34	-3.56, -1.08	0.0005	New (52)	3.46	2.20, 4.72	<0.0001	Public transport days (75)	8.54	5.00, 12.08	<0.0001	VERY LOW
Riders (n)	Change in public transport related PA 2012-2013																																				
	Point Estimate	95% CI	P Value																																		
Never Riders (391)	N/A	N/A	N/A																																		
Continued (51)	-1.15	-3.03, -0.74	0.23																																		
Former (41)	-2.34	-3.56, -1.08	0.0005																																		
New (52)	3.46	2.20, 4.72	<0.0001																																		
Public transport days (75)	8.54	5.00, 12.08	<0.0001																																		
Physical activity in everyday life as measured by non-public transport related walking trips																																					

⁴⁰ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and lack of blinding.

⁴¹ Intervention had no significant effect: p value greater than 0.05 – downgraded one level

⁴² Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and lack of blinding.

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Miller et al 2015	Non-randomised uncontrolled studies	Serious risk of bias ⁴³	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁴⁴	None	537	n/a	<p><u>Within person differences average mins of PA per 10 hrs accelerometer wear by public transport user group (baseline to 7-11 month follow-up) (intervention only). Point estimates and 95% CIs:</u></p> <table border="1"> <thead> <tr> <th>Riders (n)</th> <th colspan="3">Change in non-public transport related PA</th> </tr> <tr> <th></th> <th>Point Estimate</th> <th>95% CI</th> <th>P Value</th> </tr> </thead> <tbody> <tr> <td>Never Riders (391)</td> <td>1.27</td> <td>-0.60, 3.14</td> <td>0.20</td> </tr> <tr> <td>Continued (51)</td> <td>-1.71</td> <td>-6.62, 3.20</td> <td>0.49</td> </tr> <tr> <td>Former (41)</td> <td>-3.20</td> <td>-9.36, 2.96</td> <td>0.30</td> </tr> <tr> <td>New (52)</td> <td>1.81</td> <td>-4.04, 7.66</td> <td>0.54</td> </tr> <tr> <td>Public transport days (75)</td> <td>8.54</td> <td>5.00, 12.08</td> <td><0.000</td> </tr> </tbody> </table> <p>Evidence suggests that there was no significant change in non-public transport related Physical Activity and significantly more physical activity was carried out on public transport days compared to non-public transport days</p>	Riders (n)	Change in non-public transport related PA				Point Estimate	95% CI	P Value	Never Riders (391)	1.27	-0.60, 3.14	0.20	Continued (51)	-1.71	-6.62, 3.20	0.49	Former (41)	-3.20	-9.36, 2.96	0.30	New (52)	1.81	-4.04, 7.66	0.54	Public transport days (75)	8.54	5.00, 12.08	<0.000	VERY LOW
Riders (n)	Change in non-public transport related PA																																					
	Point Estimate	95% CI	P Value																																			
Never Riders (391)	1.27	-0.60, 3.14	0.20																																			
Continued (51)	-1.71	-6.62, 3.20	0.49																																			
Former (41)	-3.20	-9.36, 2.96	0.30																																			
New (52)	1.81	-4.04, 7.66	0.54																																			
Public transport days (75)	8.54	5.00, 12.08	<0.000																																			
Public transport use (as a proxy measure of physical activity) as measured by public transport trips, including light rail, bus and/or commuter trip																																						
1 Brown et al 2016	Non-randomised uncontrolled studies	Serious risk of bias ⁴⁵	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	537	n/a	<p><u>Public transport trips (including light rail, bus, and/or commuter rail trip) (baseline to 12-month follow-up) (intervention only):</u></p> <p>For residents living <800m away from the intervention, public transport trips were significantly more likely at one-year follow-up compared to baseline (baseline odds ratio when compared to follow-up 0.61 (95% CI 0.4 to 0.93), p≤0.02). Residents living <800m away from complete streets intervention were more likely to take public transport trips than those living further away (odds ratio for far group 0.60 (95% 0.37 to 0.97), p≤0.04).</p>	VERY LOW																												
SUMMARY: see evidence statement 1.6																																						

⁴³ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and lack of blinding.

⁴⁴ The p value is greater than 0.05, showing no significant effect of intervention – downgraded one level

⁴⁵ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and lack of blinding.

Physical Activity and the Environment – Appendix 4: GRADE profiles

85 **Public transport fare integration**

86 [To note that all studies on public transport fare integration were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality								
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control										
Public transport fare integration and simplification of paying systems																		
Public transport use (as a proxy measure of physical activity) measured as change in daily passenger boarding																		
1	Non-randomised uncontrolled study	Serious risk of bias ⁴⁶	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁴⁷	None	253,200	NA	<p><u>Change in numbers of passengers per day (baseline 1 [6 years pre intervention]; baseline 2 [3 years before intervention] follow-up [11 months post-intervention]) (intervention group only) (count data):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline 1</th> <th>Baseline 2</th> <th>Follow-Up</th> </tr> </thead> <tbody> <tr> <td>Passengers per day</td> <td>236,100</td> <td>213,400</td> <td>253,200</td> </tr> </tbody> </table> <p>Daily passenger boarding increased by 7% between baseline 1 and follow-up, and 19% between baseline 2 and follow-up. Statistical significance not reported.</p>		Baseline 1	Baseline 2	Follow-Up	Passengers per day	236,100	213,400	253,200	VERY LOW
	Baseline 1	Baseline 2	Follow-Up															
Passengers per day	236,100	213,400	253,200															
Public transport use (as a proxy measure of physical activity) measured as daily passenger trips																		
1	Non-randomised uncontrolled study	Serious risk of bias ⁴⁸	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁴⁹	None	253,200	NA	<p><u>Change in daily passenger trips (baseline 1 [6 years pre intervention]; baseline 2 [3 years before intervention] follow-up [11 months post-intervention]) (intervention group only) (count data):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline 1</th> <th>Baseline 2</th> <th>Follow-Up</th> </tr> </thead> <tbody> <tr> <td>Daily passenger trips*</td> <td>182,700</td> <td>155,000</td> <td>167,000</td> </tr> </tbody> </table>		Baseline 1	Baseline 2	Follow-Up	Daily passenger trips*	182,700	155,000	167,000	VERY LOW
	Baseline 1	Baseline 2	Follow-Up															
Daily passenger trips*	182,700	155,000	167,000															

⁴⁶ Quality score was (-) indicating high risk of bias. Main bias is unrepresentative population, tools not validated or reliable – downgraded one level

⁴⁷ No measure of variance reported or p values – downgraded one level

⁴⁸ Quality score was (-) indicating high risk of bias. Main bias is unrepresentative population, tools not validated or reliable – downgraded one level

⁴⁹ No measure of variance reported or p values – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p><i>*Trips is lower than boarding as there may be multiple boardings per trip, if trip contains >1 leg.</i></p> <p>Daily passenger trips decreased by 9% between baseline 1 and follow-up, but increased by 9% between baseline 2 and follow-up.</p>
<p>SUMMARY: see evidence statement 1.7</p>									

87

Physical Activity and the Environment – Appendix 4: GRADE profiles

88 **Motorway extension**

89 [To note that all studies on motorway extensions were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality																														
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																																
Motorway Extension																																								
Changes to Transport measured as travel trips (all, bus, car, walking)																																								
1 Foley et al 2017	Non-randomised controlled study ⁵⁰	Serious risk of bias ⁵¹	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁵²	None	Cohort: 127 Cross-Sectional: 304	Cohort: 126 Cross-Sectional: 338	<p>Odds of Intervention individuals traveling, or using bus / car / walking at 2-year follow-up compared with control individuals at 2 year follow-up (self-reported data) (COHORT):</p> <table border="1"> <thead> <tr> <th></th> <th>All Travel</th> <th>Bus</th> <th>Car</th> <th>Walking</th> </tr> </thead> <tbody> <tr> <td colspan="5">Odds Ratio (95% Confidence Interval)</td> </tr> <tr> <td>Intervention</td> <td>2.1 (1.0, 4.2)*</td> <td>1.3 (0.6, 3.0)</td> <td>1.4 (0.7, 2.7)</td> <td>1.2 (0.6, 2.3)</td> </tr> <tr> <td>Control</td> <td colspan="4">Comparison for intervention</td> </tr> </tbody> </table> <p>* = P<0.05</p> <p>Results show that intervention participants were significantly more likely to undertake travel trip at follow-up than control participants. However, there were no significant differences between study areas for use of any mode of transport in particular.</p> <p>(REPEAT CROSS-SECTIONAL):</p> <table border="1"> <thead> <tr> <th></th> <th>All Travel</th> <th>Bus</th> <th>Car</th> <th>Walking</th> </tr> </thead> <tbody> <tr> <td colspan="5">Odds Ratio (95% Confidence Interval)</td> </tr> </tbody> </table>		All Travel	Bus	Car	Walking	Odds Ratio (95% Confidence Interval)					Intervention	2.1 (1.0, 4.2)*	1.3 (0.6, 3.0)	1.4 (0.7, 2.7)	1.2 (0.6, 2.3)	Control	Comparison for intervention					All Travel	Bus	Car	Walking	Odds Ratio (95% Confidence Interval)					VERY LOW
	All Travel	Bus	Car	Walking																																				
Odds Ratio (95% Confidence Interval)																																								
Intervention	2.1 (1.0, 4.2)*	1.3 (0.6, 3.0)	1.4 (0.7, 2.7)	1.2 (0.6, 2.3)																																				
Control	Comparison for intervention																																							
	All Travel	Bus	Car	Walking																																				
Odds Ratio (95% Confidence Interval)																																								

⁵⁰ This study also included a cohort analysis i.e. an analysis including only those who responded at both baseline *and* follow-up

⁵¹ Quality score was (-) indicating high risk of bias – downgraded one level [baseline variation between groups, tools not validated or reliable]

⁵² Although travel is marginally significant in cohort analysis, confidence intervals cross line of no effect in repeated cross-sectional analysis – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Intervention	1.0 (0.5, 1.9)	1.0 (0.6, 1.8)	1.1 (0.6, 2.0)	0.8 (0.5, 1.4)	
										Comparison for intervention				
										There were no statistically significant differences between groups for likelihood of undertaking any mode of travel, or travel as a whole, at follow-up.				
Changes to Transport measured as mins/day spent travelling the previous day (all, bus, car, walking)														
1	Non-randomised controlled study ⁵³	Serious risk of bias ⁵⁴	Not assessable as 1 study	No serious indirectness	Serious imprecision ⁵⁵	None	127	126	<u>Proportional change in minutes spent travelling the previous day for one unit change in exposure to intervention (incidence rate ratio, 95% CI). Intervention vs control. 2-year follow-up. COHORT:</u>					VERY LOW
Foley et al 2017							304	338						
										Incidence Rate Ratio (95% Confidence Interval)				
									Intervention	0.8 (0.5, 1.1)	1.0 (0.6, 1.7)	0.9 (0.6, 1.3)	0.9 (0.6, 1.4)	
										Comparison for intervention				
										There were no significant differences between intervention and control for time spent travelling in general, or time spent using any mode of transport in particular. Results are mirrored in the repeat cross-sectional analysis.				
SUMMARY: see evidence statement 1.8														

90
91
92

⁵³ This study also included a cohort analysis i.e. an analysis including only those who responded at both baseline *and* follow-up
⁵⁴ Quality score was (-) indicating high risk of bias – downgraded one level [baseline variation between groups, tools not validated or reliable]
⁵⁵ All Confidence Intervals overlap line of no effect – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

93 **Work travel plans** ⁵⁶

94 [To note that all studies on work travel plans were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality																		
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																				
New express bus route																												
Changes to transport as measured by changes in mode of travel to work																												
1 Collins and Agarwal 2015	Non-randomised controlled study	Serious risk of bias ⁵⁷	Not assessable as 1 study	No serious indirectness	Serious imprecision	None	656	n/a	<p><u>Change to proportion of all trips to work made passively / somewhat passively / by public transport / actively / varied by season. Intervention only. Self-reported data. Baseline to 1 year follow-up:</u></p> <table border="1"> <thead> <tr> <th>Commuter Group⁵⁸</th> <th>% At 13 Months Post-Intervention</th> <th>% Change</th> </tr> </thead> <tbody> <tr> <td>Exclusively Passive (N = ~267)</td> <td>40.7</td> <td>-0.6</td> </tr> <tr> <td>Somewhat Passive (N = ~56)</td> <td>8.5</td> <td>-0.7</td> </tr> <tr> <td>Public transport (N = ~56)</td> <td>8.5</td> <td>3.0 (P<0.01)</td> </tr> <tr> <td>Active (N = ~93)</td> <td>14.2</td> <td>-0.7</td> </tr> <tr> <td>Varies By Season (N = ~185)</td> <td>28.2</td> <td>-0.9</td> </tr> </tbody> </table> <p>The only noted change was in public transport users, whose increase was statistically significant.</p> <p><u>Characteristics of people shifting transport mode:</u></p>	Commuter Group ⁵⁸	% At 13 Months Post-Intervention	% Change	Exclusively Passive (N = ~267)	40.7	-0.6	Somewhat Passive (N = ~56)	8.5	-0.7	Public transport (N = ~56)	8.5	3.0 (P<0.01)	Active (N = ~93)	14.2	-0.7	Varies By Season (N = ~185)	28.2	-0.9	VERY LOW
Commuter Group ⁵⁸	% At 13 Months Post-Intervention	% Change																										
Exclusively Passive (N = ~267)	40.7	-0.6																										
Somewhat Passive (N = ~56)	8.5	-0.7																										
Public transport (N = ~56)	8.5	3.0 (P<0.01)																										
Active (N = ~93)	14.2	-0.7																										
Varies By Season (N = ~185)	28.2	-0.9																										

⁵⁶ Work travel plans included one study on new express bus Route to work (bus) with subsidised monthly pass and another study assessing the impact of University Transport Plan increasing parking charges and decreasing parking spaces, meanwhile improving facilities for active commuters

⁵⁷ Quality score was (-) indicating high risk of bias – downgraded one level. Main bias is large loss to follow up.

⁵⁸ Exclusively passive: drove own vehicle, or carpooled, or got dropped off, Somewhat passive: as above, but who parked off-campus and walked to University, Transit: public transport users, Active: walk or cycled to work, Varies by season: did not employ the same route all year round

Physical Activity and the Environment – Appendix 4: GRADE profiles

										There was significantly higher likelihood of 'shifting' modes if participants were female (p=0.036), lower household income (<0.001), did not have a driver's license (<0.001), had a transit pass (p<0.001), and did not have a permit to park at work (<0.001).					
Total physical activity as measured by self-reported survey															
1 Collins and Agarwal 2015	Non-randomised controlled study	Serious risk of bias ⁵⁹	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	656	n/a	<p><u>Physical activity by usual commute method, measured by self-reported data (at 1-year follow-up only) (post hoc group comparison):</u></p> <p>On the weekly commute, the commuter groups had significantly different levels of PA (F = 276.38, p<0.001), with active commuters showing the highest levels (140.3 mins ± 5.8 SE), public transport users showing lower (79.2 mins ± 6.4 SE) and exclusively passive commuters showing the lowest (no PA took place).</p> <p>When PA levels from the commute and recreational activities were combined, there was still a significant difference between groups (F = 52.56, p<0.001), with active commuters showing the highest levels (296.3 mins ± 10.9 SE), followed by somewhat passive commuters (237.4 mins ± 23.9 SE), public transport users (183.3 mins ± 15.5) and the lowest levels being amongst exclusively passive commuters (135.1 mins ± 7.8 SE)</p>			VERY LOW			
Bristol University Transport Plan⁶⁰															
Active Travel as measured by change in people's usual travel to work mode															
1 Brockman and Fox, 2011	Non-randomised uncontrolled study	Serious risk of bias ⁶¹	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	2,829	NA	<p><u>Active Travel as measured by change in people's (self-reported) usual travel to work mode (baseline to 9-year follow-up) (intervention only)</u></p> <table border="1"> <tr> <td>Usual form of transport at baseline (%)</td> <td>Usual form of transport to work at</td> <td>Significance of change (P Value)</td> </tr> </table>			Usual form of transport at baseline (%)	Usual form of transport to work at	Significance of change (P Value)	LOW
Usual form of transport at baseline (%)	Usual form of transport to work at	Significance of change (P Value)													

⁵⁹ Quality score was (-) indicating high risk of bias. Main bias is large loss to follow up – downgraded one level

⁶⁰ **Increasing parking charges and decreasing parking space**

⁶¹ Quality score was (-) indicating high risk of bias – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

97 **Review 2**

98 **Ciclovia/Street closures**

99 [To note that all studies on Ciclovia / street closures were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality															
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																	
Street Closures																									
Total physical activity as measured by mean daily minutes spent engaging in physical activity at the event																									
Torres et al 2016	Non randomised uncontrolled study	Serious risk of bias ⁶²	Not assessable as one study	No serious indirectness	Serious imprecision ⁶³	none	589	n/a	Participants meeting recommended PA (150 minutes) during street closure event (self-reported) 23.3% of survey respondents met the PA recommendation of doing 150 minutes or more of moderate to vigorous physical activity, during the ASA event. 20.0% met the recommendation in ASA2, and 16.4% in ASA5. The average over the three events was 19.4% . Follow-up period not applicable ⁶⁴ . Intervention group only.	VERY LOW															
Total physical activity as measured by mean daily minutes spent engaging in MVPA																									
1 D'Haese et al 2015	Non randomised controlled study	No Serious risk of bias	Not assessable as one study	No serious indirectness	No serious imprecision	none	51	71	<u>Moderate and Vigorous Physical Activity (MVPA⁶⁵) differences between intervention and control at follow-up (follow-up in same week as baseline data) (measured by accelerometer)</u> <table border="1"> <thead> <tr> <th>Mean minutes (SD)</th> <th>Intervention</th> <th>Control</th> <th>Time *condition B</th> <th>P</th> </tr> </thead> <tbody> <tr> <td>Baseline</td> <td>26.70(13.51)</td> <td>26.91(16.92)</td> <td>3.626</td> <td></td> </tr> <tr> <td>Follow up</td> <td>35.79(24.93)</td> <td>24.32(13.47)</td> <td></td> <td>0.057</td> </tr> </tbody> </table>	Mean minutes (SD)	Intervention	Control	Time *condition B	P	Baseline	26.70(13.51)	26.91(16.92)	3.626		Follow up	35.79(24.93)	24.32(13.47)		0.057	LOW
Mean minutes (SD)	Intervention	Control	Time *condition B	P																					
Baseline	26.70(13.51)	26.91(16.92)	3.626																						
Follow up	35.79(24.93)	24.32(13.47)		0.057																					

⁶² Quality score allocated as (-) indicating high risk of bias, main bias is variation in event location, time and duration – downgraded one level

⁶³ No measure of variance reported or p values, unable to determine whether or not more or less people achieved recommended MVPA pre intervention – downgraded one level

⁶⁴ Data was collected on the same day as each event – follow up not applicable

⁶⁵ MVPA not defined as how may METS or counts per minute etc.

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p>The intervention group showed a significantly greater increase in MVPA than the control group between baseline and follow-up ($p = 0.057$)⁶⁶. These changes remained significant when measured over the whole day (sedentary $p = 0.012$; MVPA $p = 0.010$) This was tested to ensure that intervention groups were not compensating for changes over the rest of the day (results are significant at ≤ 0.1. "Higher significance levels are used for interaction terms as they have less power").</p> <p>In intervention children, MVPA during intervention period contributed more to entire day Physical Activity (53.4%) than during normal period (48.6%). No significance stated.</p>													
Total sedentary time as measured by mean minutes of sedentary time per day																						
1	Non randomised controlled study	No Serious risk of bias	Not assessable as one study	No serious indirectness	Serious imprecision ⁶⁷	none	54	72	<p><u>Mean daily minutes of sedentary time / day: differences between intervention and control at follow-up (measured by accelerometer)</u></p> <table border="1"> <thead> <tr> <th>Mean minutes (SD)</th> <th>Intervention</th> <th>Control</th> <th>χ^2</th> </tr> </thead> <tbody> <tr> <td>Baseline</td> <td>146.30(38.36)</td> <td>156.49(41.69)</td> <td>3.896</td> </tr> <tr> <td>FOLLOW UP</td> <td>137.74(35.43)</td> <td>164.61(40.10)</td> <td></td> </tr> </tbody> </table> <p>Change between baseline and final follow-up in intervention (-8.56 [95% CI -22.49, 5.37]) and change between baseline and final follow-up in control (8.12 [95% CI -5.24, 21.48]) were not significantly different (-16.68 [95% CI -35.59, 2.23]) (calculated by reviewers).</p> <p>Intervention vs control. Baseline and follow-up measures collected in the same week.</p>	Mean minutes (SD)	Intervention	Control	χ^2	Baseline	146.30(38.36)	156.49(41.69)	3.896	FOLLOW UP	137.74(35.43)	164.61(40.10)		VERY LOW
Mean minutes (SD)	Intervention	Control	χ^2																			
Baseline	146.30(38.36)	156.49(41.69)	3.896																			
FOLLOW UP	137.74(35.43)	164.61(40.10)																				
Total sedentary time as measured by the proportion of people reporting that they would have been sedentary if they had not attended the event																						

0-1

⁶⁷Confidence intervals spanned the line of no effect (and therefore the MID) – downgraded one level.

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Torres et al 2016	Non randomised uncontrolled study	Serious risk of bias ⁶⁸	Not assessable as one study	No serious indirectness	Serious imprecision ⁶⁹	none	589	n/a	<u>Proportion of people reporting that they would have been sedentary if they had not attended the event</u>		VERY LOW
									<table border="1"> <tr> <td>Event 1</td> <td>34%</td> </tr> <tr> <td>Event 2</td> <td>49.0%</td> </tr> <tr> <td>Event 5</td> <td>54.4%</td> </tr> </table>	Event 1	
Event 1	34%										
Event 2	49.0%										
Event 5	54.4%										
Summary – see evidence statement 2.1											

100

101

⁶⁸ Quality score allocated as (-) indicating high risk of bias Main bias is variation in event location, time and duration – downgraded one level.

⁶⁹ No measure of variance reported or P values, unable to determine whether or not more or less people achieved recommended MVPA pre intervention – downgraded one level

⁷⁰ Data was collected on the same day as event – follow up not applicable

102 **Other Cycle Infrastructure**

103 [To note that all studies on “other cycle infrastructure” were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as ‘low’].

104 1.Improvement of cycle infrastructure for active commuting

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Improvement of cycle infrastructure for active commuting										
Active travel as measured by the number of active commuters observed										
1 Hendricks et al 2009	Non randomised uncontrolled study	Serious risk of bias ⁷¹	Not assessable as one study	No serious indirectness	Serious imprecision ⁷²	none	1853	n/a	<u>Total number of active commuters observed (counted) at one-year follow-up</u> At baseline, 1,028 active commuters were recorded. This increased to 1,853 at follow-up, an increase of 63%. At follow-up, 67% of active commuters were walking, 30% were biking, and 3% were using skateboard / rollerblades / another form of active transport.	VERY LOW
Summary- see evidence statement 2.3										

105

⁷¹ Quality score allocated as (-) indicating high risk of bias, main bias is unreliable count methods – downgraded one level.

⁷² No measure of variance reported or p values, only descriptive statistics provided – downgraded one level

106 2.Cycle demonstration towns⁷³

Physical activity in everyday life as measured by the proportion of individuals cycling (various methods)																		
2	Non randomised controlled studies	Serious risk of bias ⁷⁴	Not assessable as unit measures are too different for comparison	No serious indirectness	Serious imprecision ⁷⁵	none	>9000	Matched Comparison n = 969,605 Unfunded comparison n = 4,195,540 National Comparison Group n = 10,356,452	<p><u>Self-reported cycling for at least 30 minutes more than 12 times per month (n=6000) at up to 4-year follow-up</u> Proportion of adult cycle demonstration town (CDT) residents who cycled regularly (≥30 minutes ≥12 times per month)</p> <table border="1"> <thead> <tr> <th>2006</th> <th>2008</th> <th>%point increase</th> <th>% increase</th> </tr> </thead> <tbody> <tr> <td>2.6%</td> <td>3.5%</td> <td>0.9%</td> <td>34.6%</td> </tr> </tbody> </table> <p><u>Automatic cycle counters (n=3000) at up to 4-year follow-up</u> Data from automatic cycle counters shows that cycling levels (6 town average) increased by 27% between baseline and 1-3 year follow-up in the CDT towns, ranging from +6% to +29%</p> <p><u>Any cycling in a typical week (self-reported) (n =not provided)</u> The proportion of adult residents of the CDTs doing any cycling in a typical week in the previous year rose from 24.3% in 2006 to 27.7% in 2009, an increase of approximately 3.4%-points or 14%</p> <p><u>Cycling to work (self-reported) – percentage difference at 10 year follow-up compared to baseline (95% CI)</u> In intervention towns, cyclists as a proportion of commuters increased significantly more between baseline and follow up than all three comparison groups, as seen below (ratio of increase (with 95% CI):</p> <ul style="list-style-type: none"> • Intervention Compared with <i>Matched Comparison</i>: Relative intervention effect = 1.09 (1.07, 1.11) • Intervention Compared with <i>Unfunded Comparison</i>: Relative intervention effect = 1.18 (1.17, 1.20) • Intervention Compared with <i>National Comparison</i>: Relative intervention effect = 1.26 (1.25, 1.28) <p>Cycling increased in all quintiles of deprivation although smaller improvements were seen amongst most deprived areas.</p>	2006	2008	%point increase	% increase	2.6%	3.5%	0.9%	34.6%	VERY LOW
2006	2008	%point increase	% increase															
2.6%	3.5%	0.9%	34.6%															

⁷³ Cycle demonstration town interventions included school travel planning; cycle facilities at schools; pedestrian bridges

⁷⁴ Quality score allocated as (-) for both studies, indicating high risk of bias, main bias is lack of blinding and unclear baseline characteristics – downgraded one level.

⁷⁵ No measure of variance reported or p values, only descriptive statistics provided – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

Total sedentary time as measured by the proportion of respondents classed as inactive										
1 Sloman et al 2009	Non randomised controlled study	Serious risk of bias ⁷⁶	Not assessable as one study	No serious indirectness	Serious imprecision ⁷⁷	none	Not clearly defined	Not clearly defined	<p>The proportion of adult respondents classed as inactive A validated measure was used – EPIC, self-reported 4-level index. Proportions fell from 26.2% at baseline to 23.6% at 3 year follow-up, a fall of 2.6%-points or 10%.</p> <p>The proportion of people in medium urban areas who cycled 'less than once a year' or 'never' was stable at 68 or 67% in each year between baseline and follow-up.</p>	VERY LOW
Public transport use (as a proxy of physical activity) as measured by the change in proportion of respondents using public transport as their commute										
1 Goodman et al 2013a	Non randomised controlled study	No Serious risk of bias	Not assessable as one study	No serious indirectness	No Serious imprecision	none	1,266,337	Matched Comparison n = 969,605 Unfunded comparison n = 4,195,540 National Comparison Group n = 10,356,452	<p><u>Public Transport use and driving as commute (self-reported), intervention and control groups. Percentage difference at 10-year follow-up compared with baseline:</u></p> <p>In intervention towns public transport use increased by 0.32%-points (95% CI 0.24, 0.41), and driving decreased between baseline and follow up -3.01%-points (95% CI -3.13, -2.88). Absolute figures not reported, so % change cannot be calculated.</p> <p>The decrease in driving was significantly greater in the intervention towns than all comparison groups; changes in public transport were similar to comparison groups.</p>	LOW
Active travel as measured by the change in proportion of respondents walking as their commute										
1 Goodman et al 2013a	Non randomised study	No Serious risk of bias	Not assessable as one study	No serious indirectness	No Serious imprecision	none	1,266,337	Matched Comparison n = 969,605 Unfunded comparison n = 4,195,540	<p><u>Walking as commute (self-reported), intervention and control groups. Percentage difference at 10-year follow-up compared with baseline:</u></p> <p>In intervention towns, walking increased (1.71% (95% CI 1.62, 1.81))</p>	LOW

⁷⁶ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is lack of blinding and unclear baseline characteristics

⁷⁷ No measure of variance reported or p values, only descriptive statistics provided – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

								National Comparison Group n = 10,356,452	The increase in walking was significantly greater in the intervention towns than all comparison groups.	
Summary – see evidence summary 2.4										

107

108

109

3. Various on-street and off-street bicycle paths and bridge improvements

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Various on-street and off-street bicycle paths and bridge improvements										
Change in transport as measured by cycling as a proportion of all journeys										
1 Krizek et al 2009	Non randomised controlled study	Serious risk of bias ⁷⁸	Not assessable as only one study	No serious indirectness	Serious imprecision ⁷⁹	none	Unclear	Unclear	Grouped Interventions vs Control: Change in proportion of all journeys which are made by bicycle (between baseline and up to 10-year follow-up) (SD) Intervention area 1: The proportion of all journeys which were made by bicycle increased from 1.563% (baseline) to 1.775% (follow-up), a significant result (authors report that change is greater than 2 standard deviations (SDs) of the baseline proportion). This represents a 13.4% increase. Intervention area 2: The proportion of all journeys which were made by bicycle increased from 1.023% to 1.491% (statistically significant). This represents a 45.9% increase.	VERY LOW

⁷⁸ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias lack of blinding and non-similar baseline outcome measurements

⁷⁹ Though standard deviations are reported actual S.D. figures not provided in some instances- downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p>Control groups also increased from 0.510 to 0.627% (statistically significant).</p> <p><u>Bridges: Change in proportion of all journeys which are made by bicycle from baseline (1990) to follow-up (2000) (SD)</u></p> <p>Trips crossing the river by bicycle increased significantly (3.021% to 4.604% of all journeys crossing the river, 2SDs). This was in a context of generally increasing bicycle mode share: trips which both originated and terminated east of the river also increased (1.982% to 2.775%, 2SDs), as did those originating and terminating west of the river, although to a lesser extent (2.228% to 2.585%, 1 SD).</p> <p>Unclear whether data is self-reported: a Census Transportation Planning Package containing the data was used by the authors.</p>
<p>Summary – see evidence statements 2.6</p>									

110 4.A new greenway⁸⁰ for cyclists

Quality assessment							No. of participants		Effect	Quality						
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control								
A new greenway for cyclists																
Adverse outcome as measured by the number of accidents reported																
1 Poindexter et al 2007	Non randomised uncontrolled study	Serious risk of bias ⁸¹	Not assessable as one study	No serious indirectness	Serious imprecision ⁸²	none	Unclear	NA	<p>The number of accidents per year (crashes/year) reported at baseline and 1-2 year follow up (data from police-collated information) (no control):</p> <table border="1"> <thead> <tr> <th></th> <th>Baseline</th> <th>Follow up</th> </tr> </thead> <tbody> <tr> <td>No of crashes</td> <td>78.33(sd 8.33)</td> <td>50(nr)</td> </tr> </tbody> </table> <p>Authors report that this is a significant difference, but no p-value or SD given. When buffer area is stratified by distance from intervention greenway (0.5km categories), this decrease is only significant in 0.0km-0.5km and 0.5km-1.0km categories.</p>		Baseline	Follow up	No of crashes	78.33(sd 8.33)	50(nr)	VERY LOW
	Baseline	Follow up														
No of crashes	78.33(sd 8.33)	50(nr)														
Summary – see evidence statements 2.7																

111

⁸⁰ Green way defined as an off-street bicycle facility. Traffic free, with pedestrian lanes separated from cycling lanes.

⁸¹ Quality score allocated as (-) indicating high risk of bias Main bias is poor data collection methods – downgraded one level.

⁸² Though standard deviations are reported actual S.D. figures not provided in some instances – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

112 **Trails and Paths**

113 [To note that all studies on “Trails and Paths” were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as ‘low’].

114 **5.Extension of the existing Greenway⁸³**

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	control		
Extension of the existing Greenway										
Physical activity in everyday life as measured by the change in the mean number of days spent in at least 30 minutes of walking/week										
2 West and Shores 2011 West and Shores 2015	Non randomised controlled studies	Serious risk of bias ⁸⁴	No serious inconsistency	No serious indirectness	Serious imprecision ⁸⁵	none	225	141	<u>Self-reported change in mean number of past seven days participants did ≥30 mins of walking (follow-up times between 11 months and 1 year) (intervention and control* groups):</u> Both intervention and control groups showed increases in number of days (of the past 7) in which they achieved ≥30 mins of walking. The difference between intervention group increase and control group increase was not significant: (Mean difference in days [95% CI]: -0.19 [-0.68,0.29]). *Control groups lived further from intervention than intervention groups	VERY LOW
Total physical activity as measured by the mean number of days spent engaging in moderate or vigorous physical activity										

⁸³ Authors report that greenways are “open-space corridors reserved for recreational use or environmental preservation that connect urban centres”.

⁸⁴ Quality score allocated as (-) indicating high risk of bias. Main bias is contamination – downgraded one level

⁸⁵ Although mean number of days walking increased, there is no significant difference between intervention and control

Physical Activity and the Environment – Appendix 4: GRADE profiles

<p>2 West and Shores 2011 West and Shores 2015</p>	<p>Non randomised controlled studies</p>	<p>Serious risk of bias⁸⁶</p>	<p>No serious inconsistency</p>	<p>No serious indirectness</p>	<p>Serious imprecision⁸⁷</p>	<p>none</p>	<p>230</p>	<p>138</p>	<p><u>Self-reported change in mean number of past seven days participants engaged in moderate physical activity (follow-up times between 11 months and 1 year) (intervention and control* groups):</u> In one study, both intervention and control group showed small increases over time. In the other study, both intervention and control group showed small decreases over time. The difference between intervention group change and control group change was not significant: (Mean Difference in days is 0.07 [95% CI -0.35,0.50]). *Control groups lived further from intervention than intervention groups</p>	<p>VERY LOW</p>
<p>2 West and Shore 2011 West and Shore 2015</p>	<p>Non randomised study</p>	<p>Serious risk of bias⁸⁸</p>	<p>No Serious inconsistency</p>	<p>No serious indirectness</p>	<p>Serious imprecision⁸⁹</p>	<p>none</p>	<p>229</p>	<p>141</p>	<p><u>Self-reported change in mean number of past seven days participants engaged in vigorous physical activity (follow-up times between 11 months and 1 year) (intervention and control* groups):</u> In one study, both intervention and control group showed small increases over time. In the other study, both intervention and control group showed small decreases over time. The difference between intervention group change and control group change was not significant: Mean difference in days [95% CI]: 0.32 [-0.09, 0.73]). *Control groups lived further from intervention than intervention groups</p>	<p>VERY LOW</p>
<p>Summary – see evidence statement 2.8</p>										

115

116 6.Improvement to routes⁹⁰

⁸⁶ Quality score allocated as (-) indicating high risk of bias. Main bias is contamination– downgraded one level

⁸⁷ Results are not statistically significant – downgraded one level

⁸⁸ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is contamination.

⁸⁹ Results are not statistically significant – downgraded one level

⁹⁰ For example, infrastructural changes as well as interventions which are out of scope such as bulb planting.

Physical Activity and the Environment – Appendix 4: GRADE profiles

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Improvement to routes										
Physical activity in everyday life as measured by the total number of pedestrians using the route										
Adams and Cavill 2015	Non randomised uncontrolled study	Serious risk of bias ⁹¹	Not assessable as single study	No serious indirectness	Serious imprecision ⁹²	none	3541	NA	<p><u>Change in number of pedestrians using the intervention route over whole observation period (manual count data) (baseline to 3-19 month follow-up) (no control):</u></p> <p>Overall, there was a 14.9% increase of pedestrians using the routes.</p> <p>Increases were seen in all locations: London (856 to 964, 12.6%), Newcastle (129 to 205, 58.9%), Blackburn (621 to 732, 17.9%), Wolverhampton (280 to 378, 35.0%) and Rotherham (1197 to 1262, 5.4%).</p>	VERY LOW
See evidence statement 2.9										

117
118

7. Bicycle route⁹³ and off street bicycle facilities

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Bicycle boulevard and off street bicycle facility										

⁹¹ Quality score allocated as (-) indicating high risk of bias – downgraded one level

⁹² Unable to tell whether results are statistically significant as no measures of variance or P-values provided – downgraded one level

⁹³ These routes are stretches of street with traffic calming to increase cycle safety

Physical Activity and the Environment – Appendix 4: GRADE profiles

Physical activity in everyday life as measured by the total number of bike counts using the route																													
1 Rissel et al 2015	Non randomised controlled study	Serious risk of bias ⁹⁴	Not assessable as single study	No serious indirectness	Serious imprecision ⁹⁵	none	1396	NA	<p><u>Change in number of bicycles using the route at 4-month follow-up (Count data) (intervention and control):</u></p> <p>Intervention areas increased more than control areas, but no significance is reported. Bike counts increased by 23% (812 cyclists at baseline, and 1001 cyclists at 4-month follow-up) and 97% (201 cyclists at baseline, and 395 cyclists at 4-month follow-up) at the two bike count sites located on the intervention route. Rest of city (control): The change in rates of cycling between baseline and 4-month follow-up across the whole of the City of Sydney was a 3% increase. [to note: unable to calculate the percentage point change because the figures reported are numbers and not percentages, unable to determine the initial percent to calculate percentage point change)</p>	VERY LOW																			
Physical activity in everyday life as measured by the proportion of participants making bike trips																													
1 Dill et al 2014	Non randomised controlled study	Serious risk of bias ⁹⁶	Not assessable as one study	No serious indirectness	Serious Imprecision ⁹⁷	none	154	139	<p><u>Change in proportion of participants taking a bike trip at 2-12 month follow up compared with baseline (accelerometer data) (intervention and control):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline</th> <th>Follow-up</th> <th>Significance of change</th> </tr> </thead> <tbody> <tr> <td>Intervention</td> <td>61.1%</td> <td>58.2%</td> <td rowspan="2">>0.10</td> </tr> <tr> <td>Control</td> <td>55.4%</td> <td>52.9%</td> </tr> </tbody> </table> <p>No significance difference between groups at follow up</p> <p><u>Change in average number of bike trips made per person over study period at 2-12 month follow up compared with baseline (accelerometer data) (intervention and control):</u></p> <table border="1"> <thead> <tr> <th>BIKE TRIPS (SD)</th> <th>Baseline</th> <th>Follow-up</th> <th>Change (confidence interval)*</th> </tr> </thead> <tbody> <tr> <td>Intervention</td> <td>5.6 (4.9)</td> <td>4.4 (4.2)</td> <td>-1.2 (-2.22, -0.18)</td> </tr> </tbody> </table>		Baseline	Follow-up	Significance of change	Intervention	61.1%	58.2%	>0.10	Control	55.4%	52.9%	BIKE TRIPS (SD)	Baseline	Follow-up	Change (confidence interval)*	Intervention	5.6 (4.9)	4.4 (4.2)	-1.2 (-2.22, -0.18)	VERY LOW
	Baseline	Follow-up	Significance of change																										
Intervention	61.1%	58.2%	>0.10																										
Control	55.4%	52.9%																											
BIKE TRIPS (SD)	Baseline	Follow-up	Change (confidence interval)*																										
Intervention	5.6 (4.9)	4.4 (4.2)	-1.2 (-2.22, -0.18)																										

⁹⁴ Quality score allocated as (-) indicating high risk of bias Main bias is incomplete outcome data and low count validity – downgraded one level.

⁹⁵ Unable to tell whether results are statistically significant as no measures of variance or P values provided – downgraded one level

⁹⁶ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline outcome measures and incomplete outcome data

⁹⁷ Compared confidence intervals of difference between intervention over time and control over time. No significant differences between intervention and control group - downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Control	4.3 (3.8)	3.5 (3.3)	-0.8 (-1.64, 0.04)												
<p>Although the installation of a bicycle route was statistically significantly negatively correlated with number of bike trips ($p = 0.06$)⁹⁸, and the intervention group showed a significant decrease, the intervention and control change scores were not statistically significantly different. Therefore the result is not precise.</p> <p>*CALCULATED BY REVIEWERS</p>																								
Physical activity in everyday life as measured by the total number of bike trips taken for at least >10minutes																								
1	Non randomised controlled study	Serious risk of bias ⁹⁹	Not assessable as one study	No serious indirectness	Serious imprecision ¹⁰⁰	none	154	139	<p><u>Change in proportion of participants taking a bike trip lasting >10 minutes at 2-12 month follow up compared with baseline (accelerometer data) (intervention and control):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline</th> <th>Follow-up</th> <th>P</th> </tr> </thead> <tbody> <tr> <td>Intervention</td> <td>43.9%</td> <td>45.3%</td> <td></td> </tr> <tr> <td>Control</td> <td>39.7</td> <td>31.4%</td> <td>>0.10</td> </tr> </tbody> </table> <p>There was a slight non-significant increase in the intervention group from baseline to follow up. However the actual group mean minutes spent bicycling (of trips >10 minutes) decreased from 103.9 (SD 73.0) to 65.9 (SD 74.7) between baseline and 2-12 month follow-up</p> <p>>10 minutes spent biking was significantly negatively correlated with the installation of the bicycle route ($p = 0.00$).</p>				Baseline	Follow-up	P	Intervention	43.9%	45.3%		Control	39.7	31.4%	>0.10	VERY LOW
	Baseline	Follow-up	P																					
Intervention	43.9%	45.3%																						
Control	39.7	31.4%	>0.10																					
Physical activity in everyday life as measured by percentage of participants walking >20 minutes																								
1	Non randomised controlled study	Serious risk of bias ¹⁰¹	Not assessable as one study	No serious indirectness	Serious imprecision ¹⁰²	none	154	139	<p><u>Change in proportion of participants walking for >20 minutes/day at 2-12 month follow up compared with baseline (accelerometer data) (intervention and control):</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline</th> <th>Follow-up</th> <th>P</th> </tr> </thead> <tbody> <tr> <td>Intervention</td> <td>83.5%</td> <td>75.6%</td> <td></td> </tr> </tbody> </table>				Baseline	Follow-up	P	Intervention	83.5%	75.6%		VERY LOW				
	Baseline	Follow-up	P																					
Intervention	83.5%	75.6%																						

⁹⁸ Significant testing was considered at $p < 0.1$

⁹⁹ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline outcome measures and incomplete outcome data

¹⁰⁰ Unable to tell whether the intervention has had a significant effect – downgraded one level

¹⁰¹ Quality score allocated as (-) indicating high risk of bias Main bias is dissimilar baseline outcome measures and incomplete outcome data – downgraded one level.

¹⁰² Unable to tell whether the intervention has had a significant effect – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Control	79.3%	74.4%	>0.10	
Change between groups over time not statistically significant (p ≥0.10).													
Average minutes walked (of trips >20mins) per day also decreased in both groups (intervention 107.2 [SD 79.1] to 89.4 [SD 66.8], control 92.0 [SD 86.9] to 75.4 [SD 66.5]). Change between groups over time not statistically significant (p = 0.54).													
See evidence statement 2.10													

119

120 8.Trails with new way-finding signage¹⁰³

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
6 trails with new way-finding signage										
Physical Activity in everyday life as measured by the change in the mean number of trail users										
1 Clark et al 2014	Non randomised controlled study	No serious risk of bias	Not assessable as one study	No serious indirectness	Serious imprecision ¹⁰⁴	none	Unclear ¹⁰⁵	Unclear	Change in intervention trail usage between baseline and 1-9 month follow-up (intervention and control) (infra-red sensor provided data); Intervention trail usage increased by 35%, and control trails by 31%, both significant increases (p = <0.01).	VERY LOW

¹⁰³ This intervention also included a marketing campaign which is outside of the scope of this guideline but may have impacted outcomes

¹⁰⁴ Unable to tell whether the intervention had an effect (as the control group also showed increases which were not significantly different). Downgraded one level

¹⁰⁵ At baseline, approximately 6,454 individuals were counted. At follow-up, approximately 8,610 individuals were counted. Not separated by intervention and control.

Physical Activity and the Environment – Appendix 4: GRADE profiles

										There was no significant difference between the intervention and control groups (p = 0.3226)	
<p>Mid-intervention to 1-9 month follow-up change: Between mid-intervention and 1-9 month follow-up, control trail use did not change significantly (p = 0.69), but intervention trails did decrease significantly (141 mean users per day to 107) (p = <0.01).</p>											
<p>See evidence statement 2.11</p>											

121

122

9.Greenway/Path connecting residential and commercial areas

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
<p>Greenway connecting residential and commercial areas</p>										
<p>Total Physical activity as measured by the change in people counted walking and cycling</p>										
1 Fitzhugh et al 2010	Non randomised controlled study	No serious risk of bias	Not assessable as one study	No serious indirectness	No serious imprecision	none	Not reported	Not reported	<p>Change in people counted undertaking walking or cycling on the intervention route (intervention and control) (baseline to 14 month follow-up) (direct observations):</p> <p>Increase in physical activity counts were significantly higher in the intervention compared to control for total physical activity (from 4.5 to 13.0 counts of PA in intervention; 3.0 to 1.0 count of PA in control; p = 0.001). Intervention change and control change were significantly different for both pedestrian (p = 0.001) and cyclists (p = 0.038) counts.</p>	LOW
<p>Total physical activity as a measured by the change in the proportion of people engaging in moderate and vigorous physical activity</p>										

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Gustat et al 2012	Non randomised study	Serious risk of bias ¹⁰⁶	Not assessable as single study	No serious indirectness	No serious imprecision	none	336	356	<p><u>Change in people counted undertaking moderate or vigorous activity on the intervention route (intervention and control) (baseline to 10 month follow-up) (direct observations):</u></p> <p>[Note: Intervention neighbourhood split into 2 groups (I1 and I2) – I1 was area of path, I2 was area of playground. I2 is included in the analysis as the authors measure outcomes related to the path for this area as well, and both I1 and I2 are in the same neighbourhood]</p> <p>There were significant differences between the changes over time in the four groups (p = <0.001).</p> <p>Intervention area: A significant increase in the proportion of people engaged in moderate and vigorous activity was noted in I1 between baseline (36.7%) and follow-up (41.0%) (p = <0.001). No significant change in I2.</p> <p>Control areas: A significant decrease was seen in C1 (p = <0.001, no figures provided). No significant change in C2.</p>	VERY LOW
Total physical activity as a measured by the change in the proportion of people engaging in vigorous physical activity										
Gustat et al 2012	Non randomised study	Serious risk of bias ¹⁰⁷	Not assessable as only one study	No serious indirectness	No serious imprecision	none	336	356	<p><u>Change in people counted undertaking vigorous activity on the intervention route (intervention and control) (baseline to 10 month follow-up) (direct observations):</u></p> <p>[Note: Intervention neighbourhood split into 2 groups (I1 and I2) – I1 was area of path, I2 was area of playground. I2 is included in the analysis as the authors measure outcomes related to the path for this area as well, and both I1 and I2 are in the same neighbourhood]</p> <p>Intervention area: I1 underwent a significant increase in vigorous PA between baseline and 10-month follow-up (10.5% to 12.7%; p = <0.001). I2, C1 and C2 did not undergo significant changes: all decreased slightly but non-significantly</p>	VERY LOW
Physical activity in everyday life as measured by the number of individuals using the path/trail										

¹⁰⁶ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and selective outcome reporting.

¹⁰⁷ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar baseline characteristics and selective outcome reporting.

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Gustat et al 2012	Non randomised study	Serious risk of bias ¹⁰⁸	Not assessable as only one study	No serious indirectness	Serious imprecision ¹⁰⁹	none	336	356	<p><u>Change in percentage of people reporting trail use at baseline and 10-month follow-up (intervention and control) (self-reported survey):</u></p> <p>[Note: Intervention neighbourhood split into 2 groups (I1 and I2) – I1 was area of path, I2 was area of playground. I2 is included in the analysis as the authors measure outcomes related to the path for this area as well, and both I1 and I2 are in the same neighbourhood]</p> <p>Walking trail use increased slightly but non-significantly (from 21.9% to 29.6%). [To note, unclear from reported data whether this is I1 and I2 respondents combined].</p>	VERY LOW
Physical activity in everyday life as measured by the percentage of people reporting walking for recreation										
1 Gustat et al 2012	Non randomised study	Serious risk of bias ¹¹⁰	Not assessable as only one study	No serious indirectness	Serious imprecision ¹¹¹	none	336	356	<p><u>Percentage of people reporting walking (for leisure) at baseline and 10-month follow-up (intervention and control) (self-reported survey):</u></p> <p>[Note: Intervention neighbourhood split into 2 groups (I1 and I2) – I1 was area of path, I2 was area of playground. I2 is included in the analysis as the authors measure outcomes related to the path for this area as well, and both I1 and I2 are in the same neighbourhood]</p> <p>Increases were seen in I1 (60.0% to 65.3%), C1 (61.3% to 70.4%) and C2 (57.7% to 68.9%). I2 decreased (63.3% to 61.5%). There was no significant difference in the changes over time between groups (group by time effect; p value not reported).</p>	VERY LOW
Active travel as measured by the number of children engaging in active transport to school										

¹⁰⁸ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is self-reported outcome

¹⁰⁹ No measure of variance reported – downgraded one level

¹¹⁰ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is self-reported outcome

¹¹¹ No measure of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Fitzhugh et al 2010	Non randomised controlled study	No serious risk of bias	Not assessable as one study	No serious indirectness	Serious imprecision ¹¹²	none	Not reported			<u>Change in number of children engaging in active transport to school (intervention and control) (baseline to 14 month follow-up) (Direct observation):</u> There was no significance between intervention and control group change between baseline and follow up (p = 0.2061).	VERY LOW
Active travel as measured by the percentage of people reporting walking as transportation											
1 Gustat et al 2012	Non randomised controlled study	Serious risk of bias ¹¹³	Not assessable as one study	No serious indirectness	Serious imprecision ¹¹⁴	none	336	356		<u>Percentage of people reporting walking (transportation) at baseline and 10-month follow-up (self-reported survey):</u> Increases were seen in both intervention groups (29.3% to 34.8%; and 24.8% to 36.9%). Increases also seen in control groups (31.3% to 40.5%; and 19.8% to 31.1%). No between group comparison.	VERY LOW
Summary – see evidence statement 2.12											

123

124

10. Connect2 interventions including traffic free bridges and new riverside boardwalks

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Connect2 interventions (Cardiff, Kenilworth and Southampton) including traffic free bridges and new riverside boardwalks										
Physical activity in everyday as measured by walking along any of the intervention routes										

¹¹² Unable to tell whether the intervention had a significant effect as no difference between intervention and control – downgraded one level

¹¹³ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is self-reported outcome

¹¹⁴ Unable to tell whether the intervention had a significant effect as no measure of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Goodman et al 2013b	Non randomised uncontrolled study	Serious risk of bias ¹¹⁵	Not assessable as only one study	No serious indirectness	Serious imprecision ¹¹⁶	none	3516	NA	<p><u>Change in percentage of participants who had used the intervention route for walking (baseline to between 9 and 21 month follow-up) (intervention only) (self-reported):</u></p> <p>At follow-up, 29% of the total sample (92% of those who had actually used the intervention routes) had used the intervention routes for any kind of walking, rising to 35% at follow-up 2 (91%).</p> <p>The most common category of walking was walking for recreation, at 27% (84%) at follow-up 1, and 32% (85%) at follow-up 2. Walking for education, and walking for business were least popular: <1% at both follow-up 1 and 2 for both categories</p>	VERY LOW
Physical activity in everyday as measured by cycling along any of the intervention routes										
1 Goodman et al 2013b	Non randomised uncontrolled study	Serious risk of bias ¹¹⁷	Not assessable as only one study	No serious indirectness	Serious imprecision ¹¹⁸	none	3516	NA	<p><u>Change in percentage of participants who had used the intervention route for cycling (baseline to between 9 and 21 month follow-up) (intervention only) (self-reported):</u></p> <p>At follow-up 1, 13% (39% of those who had actually used the intervention routes) of respondents had used the intervention area for any form of cycling, rising to 16% (43%) at follow-up 2. Significance not stated</p> <p>The most popular form of cycling was recreational, with 12% (37%) using it for this purpose at follow-up 1, and 15% (39%) at follow-up 2. Education and business were again the least popular: <1% at both follow-up 1 and 2 for both categories.</p>	VERY LOW
Physical activity in everyday as measured by mean minutes per week spent walking and cycling										
1 Goodman et al 2014	Non randomised uncontrolled study	Serious risk of bias ¹¹⁹	Not assessable as only one study	No serious indirectness	Serious imprecision ¹²⁰	none	3516	NA	<p><u>Change in mean minutes per week of walking or cycling on route (baseline to between 9 and 21 month follow-up) (intervention only) (self-reported):</u></p> <p>Mean minutes per week increased by 4 minutes between baseline and follow-up 1, and 0 minutes</p>	VERY LOW

¹¹⁵ Quality score allocated as (-) indicating high risk of bias. Main bias is selection bias and drop-outs – downgraded one level.

¹¹⁶ No measure of variance reported – downgraded one level

¹¹⁷ Quality score allocated as (-) indicating high risk of bias. Main bias is selection bias and drop-outs – downgraded one level.

¹¹⁸ No measure of variance reported – downgraded one level

¹¹⁹ Quality score allocated as (-) indicating high risk of bias. Main bias is selection bias and drop-outs – downgraded one level.

¹²⁰ No measure of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p>between baseline and follow-up 2. No absolute numbers provided). Significance not stated.</p> <p><u>At one year follow up</u> There was no significant difference between proximity and time spend on walking or cycling 4.6 min/wk per km closer [CI -4.2, 13.4) Total physical activity: 0.9 min/wk per km closer [CI -6.8, 8.5, CI demonstrates no statistical significance)</p> <p><u>At 2 year follow up</u> Parameter estimates and 95% confidence interval (CI) for change in minutes/week, per kilometre closer to intervention (i.e. individual 1km away will have the following increases in activity compared with someone 2km away, and double the below compared with someone 3km away): Total walking and cycling*: +15.3 min/wk per km closer [CI 6.5, 24.2, $p = <0.001$]) Total walking and cycling**: +9.2 min/wk per km closer [CI 0.6, 17.9, CI demonstrates statistical significance]) *After adjusting for demographic, socioeconomic, and health characteristics, and walking and cycling time at baseline. ** Same as above, also excluding 65 outliers (those whose change score was ≥ 600 min/wk).</p>	
Total physical activity as measured by mean minutes per week spent on moderate or vigorous physical activity along any of the intervention routes										
1 Goodman et al 2014	Non randomised uncontrolled study	Serious risk of bias ¹²¹	Not assessable as only one study	No serious indirectness	Serious imprecision ¹²²	none	3516	NA	<p><u>Change in mean minutes per week spent in moderate or vigorous physical activity on routes (baseline to between 9 and 21 month follow-up) (intervention only) (self-reported):</u></p> <p>Moderate to vigorous intensity physical activity (MVPA) declined by 24 mins/week at 21 months follow up. Significance not reported</p> <p>There was no significant difference between proximity and time spend on total physical activity - 0.9 min/wk per km closer (CI -6.8, 8.5)</p> <p>Authors note that there were no significant changes at year 2 in forms of MVPA outside of walking and cycling (adjusted effect is 0.1min/wk, CI -6.2, 6.5), showing no</p>	VERY LOW

¹²¹ Quality score allocated as (-) indicating high risk of bias. Main bias is self-reported outcome– downgraded one level.

¹²² No measure of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									evidence that gains in walking and cycling are offset by reductions in other forms of activity.	
Physical activity in everyday life as measured by the proportion of respondents reporting use of the Connect2 routes										
1 Sahlqvist et al	Non randomised uncontrolled study	No Serious risk of bias	Not assessable as only one study	No serious indirectness	Serious imprecision ¹²³	none	3516	NA	<u>Use as measured by a face to face interview no follow (1 year follow up)</u> Cardiff: 2011 48%, 2012 52% Kenilworth: 2011 28%, 2012 37% Southampton: 2011 19%, 2012 22% The most common type of use (both within walking and within cycling) at all locations is recreation (higher than social/leisure, shopping, work and education combined).	VERY LOW
Awareness of the Connect2 towns intervention										
1 Sahlqvist et al	Non randomised uncontrolled study	No Serious risk of bias	Not assessable as only one study	No serious indirectness	Serious imprecision ¹²⁴	none	3516	NA	<u>Awareness as measured by a face to face interview no follow (1 year follow up)</u> Cardiff: 2011 86%, 2012 91% Kenilworth: 2011 57%, 2012 71% Southampton: 2011 47%, 2012 55%	Very LOW
Summary – See evidence statement 2.13										

125

126

¹²³ No measure of variance reported – downgraded one level

¹²⁴ No measure of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

127 **On-Street Cycle Lanes**

128 [To note that all studies on on-street cycle lanes were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality				
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control						
On-street Cycle Lanes														
Physical Activity in Everyday Life as measured by change in cyclists counted on streets														
4	4 non-randomised studies ¹²⁵	Serious risk of bias ¹²⁶	No serious inconsistency	No serious indirectness	No serious imprecision	Consistent and strong direction of effect – upgrade one level	Can't be calculated ¹²⁷	Can't be calculated ¹²⁸	Change in cyclists counted on the street (baseline to 3-11 month various follow-up) (Intervention only, or intervention vs control)		LOW			
Bjornskau et al 2012									INTERVENTION GROUP S	Mean number per day at baseline (SD)		Mean number per day at follow-up (SD)	Percent change (%)	Significance of change (P-Value)
Hunter et al 2009									Bjornskau	NR		NR	+50	NR
Parker et al 2011									Hunter	9.06		10.49	+17	<0.0001
Parker et al 2013									Parker 2011	90.9 (21.7)		142.5 (18.5)	+56.8	<0.0001
									Parker 2013	79.2 (30.5)	257.1 (50.9)	+224.6	<0.000	
									NR = not reported SD = standard deviation. Confidence intervals not calculated as participant numbers in Parker studies are unclear. Intervention streets saw significant increases in number of cyclists counted per day (three of the four studies – the fourth did not report significance). The two control groups (see evidence tables) both saw decreases – these were significant for the one study which reported significance (-33.1%, p = <0.000). No further data on the second study control group.					

¹²⁵ 2 controlled and 2 uncontrolled studies

¹²⁶ Observational studies, quality scores all allocated as (-) indicating high risk of bias – downgraded one level

¹²⁷ The number of participants cannot be calculated. Approximate trips (not participants) calculated by reviewers as >18,000 (Bjornskau did not report any figures so is not included)

¹²⁸ Participants cannot be calculated. Approximate trips (not participants) calculated by reviewers as >1,000 (for two studies with control groups: Parker 2011, Parker 2013).

Physical Activity and the Environment – Appendix 4: GRADE profiles

129

Adverse outcomes as measured by percentage change in cyclists riding with traffic (as opposed to in the wrong direction)															
2	Parker et al 2011 Parker et al 2013	2 non-randomised studies ¹²⁹	Serious risk of bias ¹³⁰	No serious inconsistency	No serious indirectness	No serious imprecision	Consistent and strong direction of effect – upgrade one level	5,209	1,088 ¹³¹	Change in percentage of cyclists riding with traffic (baseline to 3-6 month follow-up) (intervention only, or intervention vs control) ¹³²	LOW				
										With traffic (baseline) (%)		With traffic (follow-up) (%)	Change (%-point)	% change	Significance of change (P-Value)
										Parker 2011 Intervention	73.3	81.8	+8.5	+11.6	<0.001
										Parker 2013 Intervention	92.8	95.6	+2.8	+3.0 %	<0.003
										Parker 2013 control	96.6	93.5	-3.1	-3.2%	0.002
										The percentage of cyclists cycling with rather than against traffic increased significantly between baseline and follow-up, while the percentage in control streets decreased significantly.					
Adverse outcomes as measured by percentage change in cyclists riding on the pavements															

¹²⁹ 1 controlled study, 1 uncontrolled study

¹³⁰ Quality scores all allocated as (-) indicating high risk of bias – downgraded one level. Main bias is lack of blinding and dissimilar outcome measures at baseline

¹³¹ From one study only (second study uncontrolled)

¹³² This outcome measures counts rather than participants: one individual may have appeared multiple times

Physical Activity and the Environment – Appendix 4: GRADE profiles

								<u>Change in percentage of cyclists riding on the pavement (baseline to 3-6 month follow-up) (intervention only, or intervention vs control)</u>								
								Cyclists riding on the pavement (baseline) (%)	Cyclists riding on the pavement (follow-up) (%)	Change (%-point)	% change	Significance of change (P-Value)				
3	Bjornskau et al 2012 Parker et al 2011 Parker et al 2013	3 non-randomised studies ¹³³	Serious risk of bias ¹³⁴	No serious inconsistency	No serious indirectness	Serious imprecision ¹³⁵	None	5,209 ¹³⁶	1,088 ¹³⁷	Bjornskau intervention*	47, 22	23, 5	-24, -17	-51, -22.7	NR	VERY LOW
Parker 2011 Intervention										24.6	24.4	-0.2	-1.0	0.90		
Parker 2013 Intervention										93	93	0	0	0.81		
Parker 2013 control										0.5	2.2	+1.7	+340	<0.000		
*2 intervention sites which cannot be averaged are presented here NR = not reported																
Proportion of cyclists riding on the pavement did not significantly decrease in intervention groups between baseline and follow-up for two of the studies, and in the third no significance is given for the decrease. Proportion of cyclists riding on the pavement significantly increased in the control group between baseline and follow-up																

¹³³ 2 controlled studies, 1 uncontrolled study

¹³⁴ Quality scores all allocated as (-) indicating high risk of bias – downgraded one level. Main bias is lack of blinding and dissimilar outcome measures at baseline

¹³⁵ P-values are either not reported or are not significant for most studies – downgraded one level

¹³⁶ Bjornskau did not include the no of participants therefore total is from 2 / 3 studies

¹³⁷ Bjornskau did not include the no of participants therefore total is from 2 / 3 studies

Summary – See evidence statement 2.15

Physical Activity and the Environment – Appendix 4: GRADE profiles

131 **Safe Routes to School Interventions**¹³⁸

132 [To note that all studies on Safe Routes to School were non-randomised. In line with GRADE methodology all non-randomised studies were initially graded as 'low'].

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Safe Routes to Schools (SRTS) Programmes										
Active Travel as measured by change in active commute to school (ACS) between baseline and various follow-up periods										
2	Non-randomised studies ¹³⁹	Serious risk of bias ¹⁴⁰	No serious inconsistency	No serious indirectness	No serious imprecision	None	23 schools and 45 projects ¹⁴¹	34 schools ¹⁴²	<u>Change in rates of active commuting to school between baseline and various follow-up periods (intervention only, or intervention vs control). (self-reported)</u> Stewart reports that the intervention group (no control) had significantly increased rates of ACS between baseline and 1-4 month follow-up. Hoelscher reports that the increase in percentage of children actively commuting between baseline and 3-year follow-up was significantly higher in intervention than control for morning observations (p=0.024), but not for whole-day (p=0.078).	VERY LOW
Active Travel as measured by change in Walking to School										
2	Non-randomised uncontrolled studies	Serious risk of bias ¹⁴³	No serious inconsistency	No serious indirectness	Serious imprecision ¹⁴⁴	None	2 schools and 33 projects ¹⁴⁵	NA ¹⁴⁶	<u>Change in rates of walking to school between baseline and various follow-up periods (intervention only) (self-reported)</u> Stewart reports that the intervention group (no control) had significantly increased rates of walking to school between baseline and 1-4 month follow-up (9.0% to 11.7%; p = <0.0001).	VERY LOW

¹³⁸ Safe Routes to School interventions include improved pavements and crossings, speed reduction, traffic signals, car drop off zones etc.)

¹³⁹ 1 uncontrolled, 1 controlled

¹⁴⁰ All quality scores allocated as (-) indicating high risk of bias – downgraded one level

¹⁴¹ 45 projects may each contain one or more schools. Exact number of schools and therefore individuals is unknown.

¹⁴² Exact number of participants unknown.

¹⁴³ All quality scores allocated as (-) indicating high risk of bias – downgraded one level

¹⁴⁴ No measure of variance or significance reported for some results – downgraded one level

¹⁴⁵ Projects may each contain one or more schools. Exact number of schools and therefore individuals is unknown.

¹⁴⁶ Both studies uncontrolled

Physical Activity and the Environment – Appendix 4: GRADE profiles

Stewart et al 2014									Orenstein reports increases in incidence of walking to school of between 48.5% and 304.5% between baseline and follow-up (follow-up time not defined) in two intervention schools. Significance not reported													
Active Travel as measured by change in Cycling to School																						
2 Orenstein et al 2007 Stewart et al 2014	Non-randomised studies ¹⁴⁷	Serious risk of bias ¹⁴⁸	No serious inconsistency	No serious indirectness	Serious imprecision ¹⁴⁹	None	Unknown ¹⁵⁰	Unclear ¹⁵¹	<p><u>Change in rates of cycling to school between baseline and various follow-up periods (intervention only):(self-reported)</u></p> <p>Stewart reports that the intervention group (no control) had significantly increased rates of cycling to school between baseline and 1-4 month follow-up (1.6% to 2.4%; p = 0.011)</p> <p>Orenstein reports increases in incidence of cycling to school of between 0 and 160% between baseline and follow-up (follow-up time not defined) in two intervention schools. Significance not reported.</p>	VERY LOW												
Active Travel as measured by change in general or previous week cycling to school																						
1 Ostergaard et al 2015	Non-randomised controlled study	Serious risk of bias ¹⁵²	Not assessable as one study	No serious indirectness	Serious imprecision ¹⁵³	None	13 schools (1,296 children)	12 schools (1,105 children)	<p><u>Change in general or previous week cycling to school (baseline to 1-year follow-up) (intervention vs control): (self-reported)</u></p> <table border="1"> <thead> <tr> <th>COMMUTING TO SCHOOL BY CYCLING</th> <th>Beta-coefficient*</th> <th>95% Confidence Interval</th> <th>P-Value</th> </tr> </thead> <tbody> <tr> <td>Cycling as a general method</td> <td>-0.02</td> <td>-0.10, 0.05</td> <td>0.485</td> </tr> <tr> <td>Cycling as previous week method</td> <td>0.15</td> <td>-0.25, 0.54</td> <td>0.463</td> </tr> </tbody> </table> <p>* negative figures reflect a decrease, positive numbers reflect an increase</p> <p>There was no significant difference between changes in the control group and changes in the intervention group between baseline and 1-year follow-up for either measure</p>	COMMUTING TO SCHOOL BY CYCLING	Beta-coefficient*	95% Confidence Interval	P-Value	Cycling as a general method	-0.02	-0.10, 0.05	0.485	Cycling as previous week method	0.15	-0.25, 0.54	0.463	VERY LOW
COMMUTING TO SCHOOL BY CYCLING	Beta-coefficient*	95% Confidence Interval	P-Value																			
Cycling as a general method	-0.02	-0.10, 0.05	0.485																			
Cycling as previous week method	0.15	-0.25, 0.54	0.463																			

¹⁴⁷ 1 uncontrolled, 1 controlled

¹⁴⁸ All quality scores allocated as (-) indicating high risk of bias – downgraded one level

¹⁴⁹ No measure of variance or significance reported for some results – downgraded one level

¹⁵⁰ Number of participants unknown: Orenstein 2 schools; Stewart 33 projects.

¹⁵¹ One study uncontrolled, so no participants in control group. The second study does not give numbers of participants in control group.

¹⁵² Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is response bias and dissimilar baseline outcome measures

¹⁵³ Unable to tell whether intervention had an effect as P values greater than 0.05– downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

Adverse effects as measured by change in percentage of children aged 5-18 sustaining a traffic related injury between various follow-up periods (intervention vs control)																															
2	Non-randomised controlled studies	Serious risk of bias ¹⁶⁰	No serious inconsistency	No serious indirectness	Serious imprecision ¹⁶¹	None	13 schools (1,296 children) ¹⁶²	12 schools (1,105 children) ¹⁶³	<p><u>Change in incidence of children sustaining traffic injury (baseline to various follow-up periods) (intervention vs control) (self-reported data):</u></p> <table border="1"> <thead> <tr> <th>STUDY</th> <th>Baseline (I) (%)</th> <th>Baseline (C) (%)</th> <th>P-value</th> <th>Follow-up (I) (%)</th> <th>Follow-up (C) (%)</th> <th>P-value</th> </tr> </thead> <tbody> <tr> <td>Traffic injuries (all)</td> <td>23.8</td> <td>23.3</td> <td>0.878</td> <td>24.1</td> <td>23.6</td> <td>0.812</td> </tr> <tr> <td>Traffic injuries (severe)</td> <td>3.0</td> <td>3.5</td> <td>0.556</td> <td>4.2</td> <td>3.6</td> <td>0.521</td> </tr> </tbody> </table>	STUDY	Baseline (I) (%)	Baseline (C) (%)	P-value	Follow-up (I) (%)	Follow-up (C) (%)	P-value	Traffic injuries (all)	23.8	23.3	0.878	24.1	23.6	0.812	Traffic injuries (severe)	3.0	3.5	0.556	4.2	3.6	0.521	VERY LOW
									STUDY	Baseline (I) (%)	Baseline (C) (%)	P-value	Follow-up (I) (%)	Follow-up (C) (%)	P-value																
Traffic injuries (all)	23.8	23.3	0.878	24.1	23.6	0.812																									
Traffic injuries (severe)	3.0	3.5	0.556	4.2	3.6	0.521																									
<p>I = Intervention, C = Control</p> <p>Results show that there was no significant difference in incidence of any type of traffic injury between intervention and control groups at baseline, or between intervention and control groups at 1-year post-baseline follow-up.</p> <table border="1"> <thead> <tr> <th>STUDY 2</th> <th>Change between baseline and follow-up (%)</th> <th>95% Confidence Interval</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>-15</td> <td>NR</td> </tr> <tr> <td>Intervention</td> <td>-13</td> <td>-2, 23</td> </tr> </tbody> </table> <p>Results show that there was no significant decrease in injuries in the intervention group between baseline and 7-year follow-up</p>	STUDY 2	Change between baseline and follow-up (%)	95% Confidence Interval	Control	-15	NR	Intervention	-13	-2, 23																						
STUDY 2	Change between baseline and follow-up (%)	95% Confidence Interval																													
Control	-15	NR																													
Intervention	-13	-2, 23																													
Summary – See evidence statement 2.17																															

133

¹⁶⁰ Quality score for both allocated as (-) indicating high risk of bias – downgraded one level

¹⁶¹ P values greater than 0.05 or 95% CI overlaps 0, showing no significant effect of intervention – downgraded one level

¹⁶² Orenstein does not report figures, so this is from 1 / 2 studies only.

¹⁶³ Orenstein does not report figures, so this is from 1 / 2 studies only.

134 **Review 3**

135 **Parks**

136 To note, all studies reporting on parks interventions were observational, and therefore according to the GRADE process are initially classed as “Low”.

137 **11. Upgrading park facilities**

Quality assessment							No. of participants		Effect	Quality		
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control				
Upgrading park facilities(includes lighting, facilities, paths, greenery, gyms, landscaping, improved safety)												
Total physical activity as measured by moderate to vigorous physical activity												
5	Non-randomised studies ¹⁶⁴	Serious risk of bias ¹⁶⁵	Not assessable as measures too different to combine	No serious indirectness	Serious imprecision ¹⁶⁶	None	1892 ¹⁶⁷	NA	<u>Mean number of children engaged in MVPA per 2-hour observation period (SD) 4 month follow up Bohn-Goldbaum 2013</u> Children		VERY LOW	
Bohn-Goldbaum et al 2013									Intervention park a	Pre 1.17 (2.21)		Post 0.67 (1.18)
Veitch et al 2012									Control park b	2.86 (3.95)		1.98 (3.03)
Paton-Lopez et al 2014									Confidence interval cannot be calculated as number of participants not reported. The authors report there was significant differences between intervention parks and control parks at both time points. The mean number of children engaging in MVPA decreased at follow up in both groups			
Slater et al 2016									<u>Rate of activity among youth observed in park Paton-Lopez et al 2014</u> Children (3-11) n=370 <table border="1"> <thead> <tr> <th></th> <th>Pre intervention</th> <th>Post intervention</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> </tr> </tbody> </table>			
	Pre intervention	Post intervention										

¹⁶⁴ Controlled studies – Bohn-Goldbaum et al, Veitch et al, Slater et al and Tester and Baker et al; Uncontrolled Studies – Paton Lopez et al

¹⁶⁵ Quality score for all studies allocated as (-) indicating high risk of bias – downgraded one level. Main bias is dissimilar outcome measures at baseline and contamination

¹⁶⁶ The Mean number of children engaging in MVPA reduced at follow up for the intervention group but p values or measures of variance were not included, the control group also observed a decrease and the groups were significantly different at baseline unable to determine effect of the intervention (Bohn-Goldbaum study) and no measures of variance (Paton-Lopez study) – downgraded one level

¹⁶⁷ Only 2/4 studies reported the number of participants

Physical Activity and the Environment – Appendix 4: GRADE profiles

Tester and Baker 2009									<table border="1"> <tr> <td>Moderate physical activity</td> <td>53%</td> <td>54%</td> </tr> <tr> <td>Vigorous physical activity</td> <td>11%</td> <td>22%</td> </tr> </table> <p>Half of all activities observed among children (3-11years) were moderately active during both time periods. No p values or confidence intervals provided.</p> <table border="1"> <tr> <td>Adolescents* n=157</td> <td>Pre intervention</td> <td>Post intervention</td> </tr> <tr> <td>Moderate physical activity</td> <td>54%</td> <td>60%</td> </tr> <tr> <td>Vigorous physical activity</td> <td>11%</td> <td>21.9%</td> </tr> </table> <p>*adolescents cut-off ages were not provided</p> <p>The authors report that the results were not statistically significant – possibly due too few observations. No p values or confidence intervals provided.</p> <p><u>Park-Based moderate to vigorous physical activity MVPA 12 month follow up Slater et al 2016 (direct observations)</u></p> <p>Confidence interval cannot be calculated as number of participants not reported. Model 1 included control variables only and Model 2 also examined the effects of overall neighbourhood crime count, presence of park programs and park maintenance</p> <table border="1"> <thead> <tr> <th><u>Covariate</u></th> <th>Model 1 (Coefficient, SE)</th> <th>Model 2 (Coefficient, SE)</th> </tr> </thead> <tbody> <tr> <td>Group</td> <td>0.079 (0.121)</td> <td>-0.005 (0.126)</td> </tr> <tr> <td>Time</td> <td>0.262 (0.069) (P<0.05)</td> <td>0.306 (0.071)(p<0.05)</td> </tr> <tr> <td>Group + time</td> <td>0.174 (0.088)(p<0.05)</td> <td>0.199(0.089) (p<0.05)</td> </tr> </tbody> </table> <p>The results of Model 1 (0.17, P<0.05) and Model 2 (0.199, p<0.05) showed a significant increase in the number of people engaging in MVPA when comparing baseline with the 12 month follow-up.</p> <p><u>Mean number of males and females per observation for moderate and vigorous physical activity (Tester and Baker 2009)</u></p> <table border="1"> <thead> <tr> <th></th> <th>Baseline (2006)</th> <th>Follow-up (2007)</th> <th>P value (2-tailed) Males/females</th> </tr> </thead> <tbody> <tr> <td>Park A</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Moderate</td> <td>1.64</td> <td>7.8</td> <td><0.001/<0.001</td> </tr> <tr> <td>Vigorous</td> <td>1.04</td> <td>2.5</td> <td>0.04/0.05</td> </tr> <tr> <td>Park B</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Moderate</td> <td>3.22</td> <td>14.22</td> <td><0.001/<0.001</td> </tr> <tr> <td>Vigorous</td> <td>0.65</td> <td>4.18</td> <td><0.0001/0.03</td> </tr> <tr> <td>Park C(control)</td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	Moderate physical activity	53%	54%	Vigorous physical activity	11%	22%	Adolescents* n=157	Pre intervention	Post intervention	Moderate physical activity	54%	60%	Vigorous physical activity	11%	21.9%	<u>Covariate</u>	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)	Group	0.079 (0.121)	-0.005 (0.126)	Time	0.262 (0.069) (P<0.05)	0.306 (0.071)(p<0.05)	Group + time	0.174 (0.088)(p<0.05)	0.199(0.089) (p<0.05)		Baseline (2006)	Follow-up (2007)	P value (2-tailed) Males/females	Park A				Moderate	1.64	7.8	<0.001/<0.001	Vigorous	1.04	2.5	0.04/0.05	Park B				Moderate	3.22	14.22	<0.001/<0.001	Vigorous	0.65	4.18	<0.0001/0.03	Park C(control)			
	Moderate physical activity	53%	54%																																																																	
	Vigorous physical activity	11%	22%																																																																	
	Adolescents* n=157	Pre intervention	Post intervention																																																																	
	Moderate physical activity	54%	60%																																																																	
	Vigorous physical activity	11%	21.9%																																																																	
	<u>Covariate</u>	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)																																																																	
	Group	0.079 (0.121)	-0.005 (0.126)																																																																	
	Time	0.262 (0.069) (P<0.05)	0.306 (0.071)(p<0.05)																																																																	
	Group + time	0.174 (0.088)(p<0.05)	0.199(0.089) (p<0.05)																																																																	
	Baseline (2006)	Follow-up (2007)	P value (2-tailed) Males/females																																																																	
Park A																																																																				
Moderate	1.64	7.8	<0.001/<0.001																																																																	
Vigorous	1.04	2.5	0.04/0.05																																																																	
Park B																																																																				
Moderate	3.22	14.22	<0.001/<0.001																																																																	
Vigorous	0.65	4.18	<0.0001/0.03																																																																	
Park C(control)																																																																				

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p>Moderate 1.95 4.57 0.01/0.2</p> <p>Vigorous 1.45 1.48 0.83/0.53</p> <p>There were significant increase in the mean number of males and females observed engaging in either moderate or vigorous PA or a significant increase only for moderate PA in males in the control group. There was no between group comparison.</p> <p><u>Number of people observed walking (moderate PA) (intervention and control) Veitch et al 2012:</u> Intervention: baseline 155, 3-month follow-up 195, 8 month follow-up 369. Control: baseline 75, 3-month follow-up 92, 8 month follow-up 51. The results show there was a statistically significant increase in the number of people observed walking in the intervention park over time compared to the control park. There was a significant interaction between park and time for counts of people walking $F(2, 154) = 11.70, p = 0.0005$.</p> <p><u>Number of people observed being vigorously active (intervention and control) Veitch et al 2012:</u> Intervention: baseline 38, 3-month follow-up 137, 8 month follow-up 257. Control: baseline 5, 3-month follow-up 1, 8 month follow-up 0. The results show there was statistically significant increase in the number of people observed engaging in vigorous physical activity in the intervention park over time compared to the control park. There was a significant interaction between park and time for counts of people being vigorously active $F(2, 154) = 4.98, p = 0.008$.</p>													
Total physical activity as measured by % change in MET-hours expended in park																						
1 Cohen et al 2015	Non-randomised controlled study	Serious risk of bias ¹⁶⁸	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	Unclear ¹⁶⁹	Unclear	<p><u>MET hours expended in park at 3 years follow up (intervention and control) based on direct observations</u></p> <table border="1"> <thead> <tr> <th></th> <th colspan="2">Intervention parks</th> <th>Control parks</th> </tr> <tr> <th>*=$P < 0.001$ % change..</th> <th>Renovations complete Beta (SE)</th> <th>Under construction Beta (SE)</th> <th>No renovations Beta (SE)</th> </tr> </thead> <tbody> <tr> <td>met-hours expended in park</td> <td>254.8 (70.1)*</td> <td>28.2 (25.3)</td> <td>-53.1 (11.1)*</td> </tr> </tbody> </table> <p>Confidence interval cannot be calculated as number of participants not clear. The results show that there was a 250% increase in energy expended in the intervention parks compared to the baseline ($p < 0.001$).</p>		Intervention parks		Control parks	*= $P < 0.001$ % change..	Renovations complete Beta (SE)	Under construction Beta (SE)	No renovations Beta (SE)	met-hours expended in park	254.8 (70.1)*	28.2 (25.3)	-53.1 (11.1)*	VERY LOW
	Intervention parks		Control parks																			
*= $P < 0.001$ % change..	Renovations complete Beta (SE)	Under construction Beta (SE)	No renovations Beta (SE)																			
met-hours expended in park	254.8 (70.1)*	28.2 (25.3)	-53.1 (11.1)*																			
Total physical activity as measured by the proportion of individuals self-reporting meeting the recommended physical activity																						

¹⁶⁸ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is contamination, selective outcome reporting.

¹⁶⁹ Authors did not provide number of participants split by intervention/control groups just baseline and follow up

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Gidlow et al 2010	Non-randomised uncontrolled studies	Serious risk of bias ¹⁷⁰	Not assessable as 1 study	No serious indirectness	Serious imprecision ¹⁷¹	None	170	NA	<p><u>The proportion of individuals self-reporting physical activity (meeting the recommended 30 mins of moderate PA per day) 12 months follow up (self-reported)</u></p> <table border="1"> <thead> <tr> <th></th> <th>BASELINE (N=50)</th> <th>FOLLOW-UP (N=120)</th> <th></th> </tr> <tr> <th></th> <th>%</th> <th>%</th> <th></th> </tr> </thead> <tbody> <tr> <td>Meet PA recommendations</td> <td></td> <td></td> <td></td> </tr> <tr> <td>yes</td> <td>60.8%</td> <td>62.2%</td> <td>n</td> </tr> <tr> <td>no</td> <td>39.2%</td> <td>37.8%</td> <td>s</td> </tr> </tbody> </table> <p>The authors did not report on the actual p values for significant differences between the baseline and follow up. There was no significant differences between the number of days reported in engaging in at least 30 minutes of moderate physical activity and consequently there was no significant differences between baseline and followup in the proportion of respondents meeting the PA recommendations</p>		BASELINE (N=50)	FOLLOW-UP (N=120)			%	%		Meet PA recommendations				yes	60.8%	62.2%	n	no	39.2%	37.8%	s	VERY LOW
	BASELINE (N=50)	FOLLOW-UP (N=120)																												
	%	%																												
Meet PA recommendations																														
yes	60.8%	62.2%	n																											
no	39.2%	37.8%	s																											
1 Bohn-Goldbaum 2013	Non-randomised studies	Serious risk of bias ¹⁷²	Not assessable as 1 study	Serious indirectness ¹⁷³	Serious imprecision ¹⁷⁴	None	140	NA	<p><u>Total physical activity as measured by the proportion of children meeting a parental proxy questionnaire at 4 months follow up (self-reported)</u></p> <p>Sufficient physical activity was defined as attaining the recommended 60 minutes of at least moderate intensity physical activity daily. There were no significant differences between the reported physical activity in May and September.</p> <table border="1"> <thead> <tr> <th>Physical activity</th> <th>Total % (n=58)</th> <th>May % (n=34)</th> <th>September % (n=24)</th> <th>Chi-square value</th> </tr> </thead> <tbody> <tr> <td>Sufficient activity</td> <td>55.2 (32)</td> <td>58.8 (20)</td> <td>50.0 (12)</td> <td>0.44(0.51)</td> </tr> <tr> <td>Insufficient activity</td> <td>44.8 (26)</td> <td>41.2(14)</td> <td>50.0 (12)</td> <td></td> </tr> </tbody> </table>	Physical activity	Total % (n=58)	May % (n=34)	September % (n=24)	Chi-square value	Sufficient activity	55.2 (32)	58.8 (20)	50.0 (12)	0.44(0.51)	Insufficient activity	44.8 (26)	41.2(14)	50.0 (12)							
Physical activity	Total % (n=58)	May % (n=34)	September % (n=24)	Chi-square value																										
Sufficient activity	55.2 (32)	58.8 (20)	50.0 (12)	0.44(0.51)																										
Insufficient activity	44.8 (26)	41.2(14)	50.0 (12)																											
Physical activity in everyday life as measured by park use																														

¹⁷⁰ Quality score for all studies allocated as (-) indicating high risk of bias – downgraded one level. Main bias is results based on self-reported accounts.

¹⁷¹ The intervention did not has a significant effect – downgraded one level

¹⁷² Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is results based on self-reported accounts

¹⁷³ The authors used the reported measures of parents visiting the park to determine the sufficiency of physical activity carried out by the children – serious indirectness as proxy measure – downgraded one level

¹⁷⁴ No significant differences between the 2 time points –downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

<p>6 Bohn-Goldbaum 2013 Cohen et al 2009 Cohen et al 2015 Slater et al 2016 Tester and Baker 2009 Veitch et al 2012</p>	<p>Non-randomised controlled studies</p>	<p>serious risk of bias¹⁷⁵</p>	<p>Not assessable as unit measures are too different to combine</p>	<p>No serious indirectness</p>	<p>Serious imprecision</p>	<p>None</p>	<p>Unclear</p>	<p>unclear¹⁷⁶</p>	<p><u>Parental park use (%(n)) from survey (intervention park only) (Bohn-Goldbaum et al 2013)</u></p> <p>More than half of the parents visited the intervention park at least once per week. There was no significant difference in park visit frequencies between May (57.7%) and September (61.3%, p=0.47). Significantly lower proportion of survey respondents from September had visited the playground before the renovation (49.2%) than those from May (66.7%, P = 0.04)</p> <table border="1" data-bbox="1193 419 1944 746"> <thead> <tr> <th></th> <th>Total (n=140)</th> <th>May (n=75)</th> <th>September (n=65)</th> <th>Chi-square (P value)</th> </tr> </thead> <tbody> <tr> <td colspan="5">Playground visit frequency %(n)</td> </tr> <tr> <td>At least once per week</td> <td>59.4 (79)</td> <td>57.7 (41)</td> <td>61.3 (38)</td> <td>1.51 (0.47)</td> </tr> <tr> <td>1-2 per fortnight or less</td> <td>27.1 (36)</td> <td>31.0 (22)</td> <td>22.6 (14)</td> <td></td> </tr> <tr> <td>First time</td> <td>13.5 (18)</td> <td>11.3 (8)</td> <td>16.1 (10)</td> <td></td> </tr> <tr> <td colspan="5">Visited playground before renovation %(n)</td> </tr> <tr> <td>Yes</td> <td>58.6 (82)</td> <td>66.7 (50)</td> <td>49.2 (32)</td> <td>4.36 (0.04)</td> </tr> <tr> <td>No</td> <td>41.4 (58)</td> <td>33.3 (25)</td> <td>50.8 (33)</td> <td></td> </tr> </tbody> </table> <p><u>Overall park use at 3-5 years follow up (based on direct park observations) Cohen et al 2009</u></p> <p>The authors reported that overall park use (control and intervention) declined in all age groups bar 'teens' from baseline to follow up (14682 individuals used the 10 parks at follow up compared to 19579 at baseline). Impact on control and intervention parks separately not reported.</p> <p><u>Change in total park use at 3 years park use (Direct Observations) Cohen et al 2015</u></p> <table border="1" data-bbox="1193 1018 1944 1299"> <thead> <tr> <th></th> <th colspan="2">Intervention parks</th> <th>Control parks</th> </tr> <tr> <th>*=P<0.001 % change...</th> <th>Renovations complete Beta (SE)</th> <th>Under construction Beta (SE)</th> <th>No renovations Beta (SE)</th> </tr> </thead> <tbody> <tr> <td>Total park use</td> <td>233.1(55.9)*</td> <td>30.4(21.9)</td> <td>-48.6(10.3)*</td> </tr> <tr> <td>Children in park</td> <td>434.0 (112.7)*</td> <td>58.8 (33.5)*</td> <td>-7.4 (23.1)</td> </tr> <tr> <td>Teens in park</td> <td>-51.1 (10.4)*</td> <td>-7.3(19.7)</td> <td>0.3 (24.7)</td> </tr> </tbody> </table>		Total (n=140)	May (n=75)	September (n=65)	Chi-square (P value)	Playground visit frequency %(n)					At least once per week	59.4 (79)	57.7 (41)	61.3 (38)	1.51 (0.47)	1-2 per fortnight or less	27.1 (36)	31.0 (22)	22.6 (14)		First time	13.5 (18)	11.3 (8)	16.1 (10)		Visited playground before renovation %(n)					Yes	58.6 (82)	66.7 (50)	49.2 (32)	4.36 (0.04)	No	41.4 (58)	33.3 (25)	50.8 (33)			Intervention parks		Control parks	*=P<0.001 % change...	Renovations complete Beta (SE)	Under construction Beta (SE)	No renovations Beta (SE)	Total park use	233.1(55.9)*	30.4(21.9)	-48.6(10.3)*	Children in park	434.0 (112.7)*	58.8 (33.5)*	-7.4 (23.1)	Teens in park	-51.1 (10.4)*	-7.3(19.7)	0.3 (24.7)	<p>VERY LOW</p>
	Total (n=140)	May (n=75)	September (n=65)	Chi-square (P value)																																																																		
Playground visit frequency %(n)																																																																						
At least once per week	59.4 (79)	57.7 (41)	61.3 (38)	1.51 (0.47)																																																																		
1-2 per fortnight or less	27.1 (36)	31.0 (22)	22.6 (14)																																																																			
First time	13.5 (18)	11.3 (8)	16.1 (10)																																																																			
Visited playground before renovation %(n)																																																																						
Yes	58.6 (82)	66.7 (50)	49.2 (32)	4.36 (0.04)																																																																		
No	41.4 (58)	33.3 (25)	50.8 (33)																																																																			
	Intervention parks		Control parks																																																																			
*=P<0.001 % change...	Renovations complete Beta (SE)	Under construction Beta (SE)	No renovations Beta (SE)																																																																			
Total park use	233.1(55.9)*	30.4(21.9)	-48.6(10.3)*																																																																			
Children in park	434.0 (112.7)*	58.8 (33.5)*	-7.4 (23.1)																																																																			
Teens in park	-51.1 (10.4)*	-7.3(19.7)	0.3 (24.7)																																																																			

¹⁷⁵ All six studies with an allocated QA score of (-) indicating high risk of bias with some results based on self-reported accounts – downgraded one level

¹⁷⁶ Total of intervention and control groups unclear as a proportion of the 6 studies did not provide number of participants

Physical Activity and the Environment – Appendix 4: GRADE profiles

								<table border="1"> <tr> <td>Adults in park</td> <td>169.6 (39.9)</td> <td>29.8 (19.2)</td> <td>-53.7 (8.2)*</td> </tr> <tr> <td>Seniors in park</td> <td>25.4 (18.0)</td> <td>-8.8(13.1)</td> <td>-10.7(15.1)</td> </tr> </table> <p>Confidence intervals cannot be calculated as number of participants observed not reported. The results from the Cohen 2015 study show there was a 230% increase in park use in the intervention parks compared to the baseline (p<0.001). Significant increases were observed in children and adults at the renovated parks and a non-significant increase in the seniors age group (age cut offs not included). There was a significant 48% decrease in total park use in the control park at follow up compared to the baseline. The authors did not compare intervention park and control park</p> <p><u>Park usage at 1 year follow up (based on direct observations Slater et al 2016</u> Model 1 included control variables only and Model 2 also examined the effects of overall neighbourhood crime count, presence of park programs and park maintenance</p> <table border="1"> <thead> <tr> <th>Covariate</th> <th>Model 1 (Coefficient, SE)</th> <th>Model 2 (Coefficient, SE)</th> </tr> </thead> <tbody> <tr> <td>Group</td> <td>0.201(0.09) (p<0.05)</td> <td>0.056(0.096)</td> </tr> <tr> <td>Time</td> <td>0.031 (0.049)</td> <td>0.097 (0.052) (p<0.1)</td> </tr> <tr> <td>Group + time</td> <td>0.174 (0.062) (p<0.05)</td> <td>0.211 (0.063) (p<0.05)</td> </tr> <tr> <td>Park maintenance scale</td> <td></td> <td>-0.072(0.014)(p<0.05)</td> </tr> <tr> <td>Neighbourhood crime count (log)</td> <td></td> <td>0.359(0.104) (p<0.05)</td> </tr> <tr> <td>Park has programmes</td> <td></td> <td>0.159(0.199)</td> </tr> </tbody> </table> <p>Confidence intervals cannot be calculated as number of participants not reported. The results shows there was a statistically significant increase in park utilisation over time in intervention parks compared to control parks in both model 1 and 2. The results show that the only factor significantly associated with increased park use was low neighbourhood crime count (beta=0.359, p<0.05).</p> <p><u>Park use (1 year follow up) (direct observations) Taster and Baker 2009</u> There was a significant increase in the total number of visitors in observations on the intervention Park A (p=0.00) and B (p=0.00), but not for the control park (p=0.36)In the intervention parks the results show that there was significant increase in the number of children, adults and seniors visiting the parks at baseline and at follow-up. The teens' group was the only group whose visits decreased in the intervention parks, but increased in the control park. There were no significant changes in the control parks from baseline to follow-up apart from the male teens whose park visits significantly increased.</p> <p>The authors did not compare the intervention parks with the control park</p>	Adults in park	169.6 (39.9)	29.8 (19.2)	-53.7 (8.2)*	Seniors in park	25.4 (18.0)	-8.8(13.1)	-10.7(15.1)	Covariate	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)	Group	0.201(0.09) (p<0.05)	0.056(0.096)	Time	0.031 (0.049)	0.097 (0.052) (p<0.1)	Group + time	0.174 (0.062) (p<0.05)	0.211 (0.063) (p<0.05)	Park maintenance scale		-0.072(0.014)(p<0.05)	Neighbourhood crime count (log)		0.359(0.104) (p<0.05)	Park has programmes		0.159(0.199)
Adults in park	169.6 (39.9)	29.8 (19.2)	-53.7 (8.2)*																																		
Seniors in park	25.4 (18.0)	-8.8(13.1)	-10.7(15.1)																																		
Covariate	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)																																			
Group	0.201(0.09) (p<0.05)	0.056(0.096)																																			
Time	0.031 (0.049)	0.097 (0.052) (p<0.1)																																			
Group + time	0.174 (0.062) (p<0.05)	0.211 (0.063) (p<0.05)																																			
Park maintenance scale		-0.072(0.014)(p<0.05)																																			
Neighbourhood crime count (log)		0.359(0.104) (p<0.05)																																			
Park has programmes		0.159(0.199)																																			

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<table border="1"> <thead> <tr> <th>Intervention park A</th> <th>Baseline (2006)</th> <th>Follow-up (2007)</th> <th>P value (2-tailed) males/females</th> </tr> </thead> <tbody> <tr> <td>Children</td> <td>0.09</td> <td>3.55</td> <td>0.001/<0.001</td> </tr> <tr> <td>Teens</td> <td>0.64</td> <td>1.67</td> <td>0.813/0.008</td> </tr> <tr> <td>Adults</td> <td>4.07</td> <td>18.95</td> <td><0.001/<0.001</td> </tr> <tr> <td>Seniors</td> <td>0</td> <td>0.18</td> <td>0.003/0.16</td> </tr> <tr> <td colspan="4">Intervention park B</td> </tr> <tr> <td>Children</td> <td>0.42</td> <td>4.35</td> <td>0.006/0.003</td> </tr> <tr> <td>Teens</td> <td>1.37</td> <td>1.71</td> <td>0.931/0.116</td> </tr> <tr> <td>Adults</td> <td>2.69</td> <td>22.76</td> <td><0.001/<0.001</td> </tr> <tr> <td>Seniors</td> <td>0.4</td> <td>3.38</td> <td><0.001/<0.001</td> </tr> <tr> <td colspan="4">Control park</td> </tr> <tr> <td>Children</td> <td>0.27</td> <td>0.61</td> <td>0.257/0.042</td> </tr> <tr> <td>Teens</td> <td>1.32</td> <td>4.09</td> <td>0.00/0.27</td> </tr> <tr> <td>Adults</td> <td>6.97</td> <td>5.71</td> <td>0.37/0.478</td> </tr> <tr> <td>Seniors</td> <td>0.07</td> <td>0.04</td> <td>0.475/-</td> </tr> </tbody> </table> <p>Total number of users (Intervention and control) Veitch et al 2012: Intervention: baseline 235, 3-month follow-up 582, 8 month follow-up 985. Control: baseline 83, 3-month follow-up 114, 8 month follow-up 51. The results show that there was a statistically significant increase in park use for the refurbished park over time compared to the control park. There was a significant interaction between park and time for total counts of park users, $F(2, 154) = 14.99, p = 0.0005$</p>	Intervention park A	Baseline (2006)	Follow-up (2007)	P value (2-tailed) males/females	Children	0.09	3.55	0.001/<0.001	Teens	0.64	1.67	0.813/0.008	Adults	4.07	18.95	<0.001/<0.001	Seniors	0	0.18	0.003/0.16	Intervention park B				Children	0.42	4.35	0.006/0.003	Teens	1.37	1.71	0.931/0.116	Adults	2.69	22.76	<0.001/<0.001	Seniors	0.4	3.38	<0.001/<0.001	Control park				Children	0.27	0.61	0.257/0.042	Teens	1.32	4.09	0.00/0.27	Adults	6.97	5.71	0.37/0.478	Seniors	0.07	0.04	0.475/-
Intervention park A	Baseline (2006)	Follow-up (2007)	P value (2-tailed) males/females																																																																		
Children	0.09	3.55	0.001/<0.001																																																																		
Teens	0.64	1.67	0.813/0.008																																																																		
Adults	4.07	18.95	<0.001/<0.001																																																																		
Seniors	0	0.18	0.003/0.16																																																																		
Intervention park B																																																																					
Children	0.42	4.35	0.006/0.003																																																																		
Teens	1.37	1.71	0.931/0.116																																																																		
Adults	2.69	22.76	<0.001/<0.001																																																																		
Seniors	0.4	3.38	<0.001/<0.001																																																																		
Control park																																																																					
Children	0.27	0.61	0.257/0.042																																																																		
Teens	1.32	4.09	0.00/0.27																																																																		
Adults	6.97	5.71	0.37/0.478																																																																		
Seniors	0.07	0.04	0.475/-																																																																		
Sedentary behaviour																																																																					
2	Tester and Baker 2009	Non-randomised controlled studies	serious risk of bias ¹⁷⁷	Not assessable as measures too different to combine	No serious indirectness	Serious imprecision ¹⁷⁸	None	2614	597	<table border="1"> <thead> <tr> <th colspan="4">Mean number of males and females per observation (sedentary)</th> </tr> <tr> <th></th> <th>Baseline (2006)</th> <th>Follow-up (2007)</th> <th>P value (2-tailed) Males/females</th> </tr> </thead> <tbody> <tr> <td>Intervention park A</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Sedentary</td> <td>2.13</td> <td>14.01</td> <td><0.001/<0.001</td> </tr> <tr> <td>Intervention park B</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Sedentary</td> <td>0.84</td> <td>13.95</td> <td><0.001/<0.001</td> </tr> <tr> <td>Park (Control)</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Sedentary</td> <td>5.24</td> <td>4.39</td> <td>0.4/0.65</td> </tr> </tbody> </table> <p>There were significantly more people observed being sedentary in the intervention parks at follow up, however in the two intervention parks</p>	Mean number of males and females per observation (sedentary)					Baseline (2006)	Follow-up (2007)	P value (2-tailed) Males/females	Intervention park A				Sedentary	2.13	14.01	<0.001/<0.001	Intervention park B				Sedentary	0.84	13.95	<0.001/<0.001	Park (Control)				Sedentary	5.24	4.39	0.4/0.65																											
Mean number of males and females per observation (sedentary)																																																																					
	Baseline (2006)	Follow-up (2007)	P value (2-tailed) Males/females																																																																		
Intervention park A																																																																					
Sedentary	2.13	14.01	<0.001/<0.001																																																																		
Intervention park B																																																																					
Sedentary	0.84	13.95	<0.001/<0.001																																																																		
Park (Control)																																																																					
Sedentary	5.24	4.39	0.4/0.65																																																																		

¹⁷⁷ Quality score for all studies allocated as (-) indicating high risk of bias – downgraded one level. Main bias is some results based on self-reported accounts and contamination

¹⁷⁸ Some studies did not provide measures of variance included – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p>combined, there were 1681 physically active visitors in the follow-up week, compared to a total of 360 at baseline. There were statistically significant increases among males and females who were observed at each respective PA level in the intervention parks. The majority of visitors were however sedentary. Sedentary visitors increased 5+ fold</p> <p><u>Number of people observed lying/sitting (intervention and control) Veitch et al 2012:</u> Intervention: baseline 6, 3-month follow-up 119, 8 month follow-up 61. Control: baseline 0, 3-month follow-up 4, 8 month follow-up 0. Significance of interaction between park and time not reported</p> <p><u>Park-Based Sedentary behaviour</u></p> <table border="1"> <thead> <tr> <th>Covariate</th> <th>Model 1 (Coefficient, SE)</th> <th>Model 2 (Coefficient, SE)</th> </tr> </thead> <tbody> <tr> <td>Group</td> <td>0.409(0.119) (p<0.05)</td> <td>0.264(0.123) (p<0.05)</td> </tr> <tr> <td>Time</td> <td>-0.194(0.068) (p<0.05)</td> <td>-0.112(0.071)</td> </tr> <tr> <td>Group + time</td> <td>0.139(0.089)</td> <td>0.173(0.089) (p<0.054)</td> </tr> <tr> <td>Park maintenance scale</td> <td></td> <td>-0.090(0.019)(p<0.05)</td> </tr> <tr> <td>Neighbourhood crime count log</td> <td></td> <td>0.316(0.119) (p<0.05)</td> </tr> <tr> <td>Park has programmes</td> <td></td> <td>0.124(0.222)</td> </tr> </tbody> </table> <p>Confidence intervals cannot be calculated as number of participants not reported. The results show that intervention parks had significantly more people engaging in sedentary behaviour as well as a significant decrease in observed sedentary behaviour over time in the control group (beta = -0.19, p<0.05). The results also show that while provision of programmes did not influence sedentary behaviour, decreased park maintenance and increased neighbourhood crime were both associated with an increase in sedentary behaviour p<0.05.</p>	Covariate	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)	Group	0.409(0.119) (p<0.05)	0.264(0.123) (p<0.05)	Time	-0.194(0.068) (p<0.05)	-0.112(0.071)	Group + time	0.139(0.089)	0.173(0.089) (p<0.054)	Park maintenance scale		-0.090(0.019)(p<0.05)	Neighbourhood crime count log		0.316(0.119) (p<0.05)	Park has programmes		0.124(0.222)	
Covariate	Model 1 (Coefficient, SE)	Model 2 (Coefficient, SE)																													
Group	0.409(0.119) (p<0.05)	0.264(0.123) (p<0.05)																													
Time	-0.194(0.068) (p<0.05)	-0.112(0.071)																													
Group + time	0.139(0.089)	0.173(0.089) (p<0.054)																													
Park maintenance scale		-0.090(0.019)(p<0.05)																													
Neighbourhood crime count log		0.316(0.119) (p<0.05)																													
Park has programmes		0.124(0.222)																													
Perceived state of the park (safety, maintenance etc)																															
3 Cohen et al 2009 Cohen et al 2015	Non-randomised controlled studies	serious risk of bias ¹⁷⁹	Not assessable as units too different to combine	No serious indirectness	Serious imprecision ¹⁸⁰	None	Unclear	Unclear	<p><u>Perceptions of park safety (self-reported- survey responses) 3-5 years follow up, Cohen et al 2009 (intervention and control park)</u> Perceptions of park safety from baseline to follow-up improved among intervention park users and neighbourhood residents; while it decreased for the control parks. This was a significant change (p=0.007) ; however, it was not correlated with observed park use or self-reported exercise</p> <p><u>Survey Perceptions of safety (self-reported) 3 years follow up Cohen et al 2015 (intervention park only)</u></p>	VERY LOW																					

¹⁷⁹ Quality score for all studies allocated as (-) indicating high risk of bias – downgraded one level. Main bias is self-reported accounts, contamination, and selection bias.

¹⁸⁰ Some studies did not provide measures of variance included – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

Gidlow et al 2010									<p>Park renovations were associated with a significantly increased perception of park safety by park users (Beta estimate 1.43, p<0.01) and local residents (Beta estimate 0.42, p<0.01).</p> <p><u>Baseline and follow-up perceptions of the intervention park (self-reported) Gidlow et al 2010</u></p> <p>There were no significant changes in perception of the intervention park (design and appearance, ease of getting around and maintenance) between baseline and follow up.</p> <p>The authors did not seek any perceptions regarding the design and appearance, ease of getting around the park and maintenance of the control parks.</p>																																														
Total physical activity potential predictors determined by a multivariate model																																																							
1	Non-randomised studies	No serious risk of bias	Not assessable as only one study	No serious indirectness	No Serious imprecision	None	77	79	<p><u>Total physical activity potential predictors determined by a multivariate model (intervention and control parks)</u></p> <table border="1"> <thead> <tr> <th>Variables</th> <th>Intervention ratio of geometric means(95% ci)</th> <th>P value</th> <th>Control ratio of geometric means (95% ci)</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Exposure to playground (community of residence) intervention compared to control</td> <td>0.90 (0.69-1.16)</td> <td>0.417</td> <td>1.11(0.85,1.44)</td> <td>0.456</td> </tr> <tr> <td>BMI overall (per 1 z score unit increase)</td> <td>0.96 (0.87, 1.06)</td> <td>0.388</td> <td></td> <td></td> </tr> <tr> <td>BMI (control group)</td> <td></td> <td></td> <td>1.19(1.06,1.34)</td> <td>0.005</td> </tr> <tr> <td>BMI (intervention group)</td> <td></td> <td></td> <td>0.94(0.83,1.06)</td> <td>0.300</td> </tr> <tr> <td>Interaction: community by BMI z score</td> <td></td> <td></td> <td></td> <td>0.006</td> </tr> <tr> <td>Interaction sex & ethnicity</td> <td></td> <td></td> <td></td> <td>0.019</td> </tr> <tr> <td>Participant age (per 1 year increase)</td> <td>0.92(0.87, 0.97)</td> <td>0.004</td> <td>0.90(0.85,0.94)</td> <td><0.001</td> </tr> <tr> <td>Non-school day (ref school day)</td> <td>0.72(0.63, 0.81)</td> <td><0.001</td> <td>0.72(0.63,0.82)</td> <td><0.001</td> </tr> </tbody> </table>	Variables	Intervention ratio of geometric means(95% ci)	P value	Control ratio of geometric means (95% ci)	P value	Exposure to playground (community of residence) intervention compared to control	0.90 (0.69-1.16)	0.417	1.11(0.85,1.44)	0.456	BMI overall (per 1 z score unit increase)	0.96 (0.87, 1.06)	0.388			BMI (control group)			1.19(1.06,1.34)	0.005	BMI (intervention group)			0.94(0.83,1.06)	0.300	Interaction: community by BMI z score				0.006	Interaction sex & ethnicity				0.019	Participant age (per 1 year increase)	0.92(0.87, 0.97)	0.004	0.90(0.85,0.94)	<0.001	Non-school day (ref school day)	0.72(0.63, 0.81)	<0.001	0.72(0.63,0.82)	<0.001	LOW
Variables	Intervention ratio of geometric means(95% ci)	P value	Control ratio of geometric means (95% ci)	P value																																																			
Exposure to playground (community of residence) intervention compared to control	0.90 (0.69-1.16)	0.417	1.11(0.85,1.44)	0.456																																																			
BMI overall (per 1 z score unit increase)	0.96 (0.87, 1.06)	0.388																																																					
BMI (control group)			1.19(1.06,1.34)	0.005																																																			
BMI (intervention group)			0.94(0.83,1.06)	0.300																																																			
Interaction: community by BMI z score				0.006																																																			
Interaction sex & ethnicity				0.019																																																			
Participant age (per 1 year increase)	0.92(0.87, 0.97)	0.004	0.90(0.85,0.94)	<0.001																																																			
Non-school day (ref school day)	0.72(0.63, 0.81)	<0.001	0.72(0.63,0.82)	<0.001																																																			

Physical Activity and the Environment – Appendix 4: GRADE profiles

										Usually walking to school (ref: car or mixed)	1.18 (1.01, 1.39)	0.038	1.16(1.00,1.35)	0.046	
<p>There authors did not report on the mean total daily physical activity as measured by the accelerometer at baseline and follow-up but used in multivariate models to identify potential predictors of physical activity. The multivariate model found no evidence that participants in the intervention community had a statistically significant difference in their mean total daily physical activity (TDPA), compared to those living in the control community. The results show that living close to a playground (even after renovations) does not have a significant effect on total daily physical activity. There was evidence of statistically significant associations in the final model between follow-up physical activity and participant baseline age, school day, usual mode of travel to school, gender, and ethnicity.</p>															
<p>Summary - See evidence statement 3.1</p>															

138

139

12. New Parks

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
New parks										
Physical Activity in Everyday Life as measured by adults' frequency of park visits										
1 Cohen et al, 2014	Non-randomised controlled study	serious risk of bias ¹⁸¹	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	432	NA	Physical Activity in Everyday Life as measured by adults' self-reported frequency of park visits (intervention only) (baseline to 2 year post-baseline follow-up)	VERY LOW

¹⁸¹ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is results based on self-reported accounts and dissimilar characteristics and outcome measures at baseline

Physical Activity and the Environment – Appendix 4: GRADE profiles

									At follow up the percentage reporting visiting any park more than once per week tripled, (increased by 22.8 percentage points from 11.1% to 33.9%) a statistically significant change from baseline (p<0.0001).	
Physical Activity in Everyday Life as measured by average number of monthly visits										
1 King et al 2015	Non-randomised uncontrolled study	No serious risk of bias	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	4525	NA	<u>Physical Activity in Everyday Life as measured by average number of monthly visits (intervention only) (baseline to 2-year post-baseline follow-up)</u> The average monthly visits significantly increased by 362% at 2 year follow up.(from 180 to 651, p=0.02) When looking at the proportions of those who visited there was a significant increase in the proportion of teenagers visiting the park at follow up and a decrease in the proportion of children.	LOW
Physical Activity in Everyday Life as measured by proportion of adults exercising in the park										
1 Cohen et al, 2014	Non-randomised controlled study	Serious risk of bias ¹⁸²	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	432	NA	<u>Physical Activity in Everyday Life as measured by proportion of adults self-reporting exercising in the park (intervention only) (baseline to 2 year post-baseline follow-up)</u> The proportion of people exercising in the park significantly increased by 4.8 percentage points (from 9.6% at baseline to 14.4% at follow up, p<0.0395)	VERY LOW
Physical Activity in Everyday Life as measured by proportion of adults self-reporting engaging in leisure-time exercise										
1 Cohen et al, 2014	Non-randomised studies	serious risk of bias ¹⁸³	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	432	NA	<u>Physical Activity in Everyday Life as measured by proportion of adults self-reporting engaging in leisure-time exercise (intervention only) (baseline to 2 year post-baseline follow-up)</u> The proportion of people engaging in leisure time significantly increase by 9.9 percentage points (from 25.8% at baseline to 35.7% at follow up) (p<0.0025)	VERY LOW
Total Physical Activity as measured by energy expenditure levels										

¹⁸² Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is results based on self-reported accounts and dissimilar characteristics and outcome measures at baseline

¹⁸³ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is results based on self-reported accounts and dissimilar characteristics and outcome measures at baseline

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 King et al 2015	Non-randomised studies	No serious risk of bias	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	432	NA	<u>Total Physical Activity as measured by energy expenditure levels (intervention only) (baseline to 2-year post-baseline follow-up)</u> Pre- and post- comparisons between the non-park and park zones indicated a 38% decrease in energy expended in streets, alleys and parking lots and a 3-fold increase in energy expended within the park boundaries post-construction (P = 0.002).	LOW
Total Physical Activity as measured by change in proportion of people undertaking moderate and vigorous activity in the park										
1 King et al 2015	Non-randomised studies	No serious risk of bias	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	4525	NA	<u>Total Physical Activity as measured by change in proportion of people undertaking moderate activity or vigorous activity in the park (intervention only) (baseline to 2-year post-baseline follow-up) 2010 -2012</u> There was a significant increase in the proportion of individuals observed as engaging in vigorous physical activity from baseline (18.6%) to follow up (25.2%) an increase of 6.6 percentage points (p=0.04), however the proportion of individuals observed as engaging in moderate physical activity decreased from 43.4% to 40.8% at follow up, a decrease of 2.6 percentage points (p=0.007)	LOW
Sedentary behaviour										
1 King et al 2015	Non-randomised studies	No serious risk of bias	Not assessable as 1 study	No serious indirectness	Serious imprecision	None	4525	NA	The proportion of adults observed being sedentary decreased at follow up by 4 percentage points, no measures if variance provided.	Very low
Summary - See evidence statement 3.3										

140

141

13. Changing micro-environment

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		

Physical Activity and the Environment – Appendix 4: GRADE profiles

Movement of seats and picnic tables in a park													
Total physical activity as measured by METS expended by park visitors during intervention time													
									Condition				
									Adults		Children		
									N	Mets	N	Mets	
1	Non-randomised studies	serious risk of bias ¹⁸⁴	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	484	NA	summer 2012 (mean, ±standard error) Seating near (a ₁) 79 1.8±0.1 91 3.1±0.2 Seating far (b) 22 2.0±0.2 27 3.8±0.4 Seating near (a ₂) 55 1.4±0.1 57 3.1±0.3 A1 – tables nearer to the playground, B – tables further away from the playground, A2 – tables nearer to the playground again. Reviewers have calculated that: • For adults, METS expended is significantly higher in b when compared with A ₁ (mean difference 0.20, 95% CI 0.11, 0.29), and also when compared with A ₂ (mean difference 0.60, 95% CI 0.51, 0.69). For children, METS expended is significantly higher in b when compared with A ₁ (mean difference 0.70, 95% CI 0.54, 0.86), and also when compared with A ₂ (mean difference 0.70, 95% CI 0.53, 0.87). Children were more intensely active than adults (p=0.0001) METS Intensities were greater when seating was not accessible (B) than when seating was accessible (A1, A2) p<0.02).				VERY LOW
Total Physical Activity as measured by odds of adults engaging in MVPA													
1	Non-randomised uncontrolled studies	Serious risk of bias ¹⁸⁵	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	484	NA	Total Physical Activity as measured by odds of adults engaging in MVPA (intervention only) (unclear follow-up period) The odds of adults engaging in MVPA were at least 4.1 times significantly higher when the picnic tables s were moved away from the playground compared to when the tables were closer (p=0.03), the results remained consistent when the study was repeated at a later date.				VERY LOW
Total sedentary time as measured by odds of adults standing rather than sitting during intervention													

¹⁸⁴ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is presence of selection bias and confounders.

¹⁸⁵ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is presence of selection bias and confounders

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 Roemmich et al 2014	Non- randomised uncontrolled studies	Serious risk of bias ¹⁸⁶	Not assessable as 1 study	No serious indirectness	No Serious imprecision	None	484	NA	The odds of adults standing rather sitting were greater when the tables were further away from the playground compared to when the tables were nearer to the playground (OR – 9.4, 95% CI 2.5, 35.2, p value <0.0001) and the odds remained significantly great when the tables were moved back to the playground again (4.7, 95% CI 1.3,17.2; p value <0.02)	VERY LOW
Summary - See evidence statement 3.5										

142

¹⁸⁶ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is presence of selection bias and confounders

Physical Activity and the Environment – Appendix 4: GRADE profiles

143 **Effectiveness of neighbourhood interventions**

144 To note, all studies reporting on parks interventions were observational, and therefore according to the GRADE process are initially classed as
 145 “Low”.

Quality assessment							No. of participants		Effect	Quality																	
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																			
Moving to a new neighbourhood complying with Livable Neighbourhoods Guidelines (LNG) vs moving to conventional neighbourhood																											
Active Travel as measured by change in transportation-related walking (transport walking)																											
1 ¹⁸⁷ Christian et al 2013 Knuiman et al 2014	Non-randomised controlled studies ¹⁸⁸	No serious risk of bias	Not assessable as single study	No serious indirectness	Serious imprecision ¹⁸⁹	None	299 ¹⁹⁰	528	Mean minutes of transport walking per week (baseline is prior to move)(intervention only, or intervention vs control) (baseline to 1-, 3-, and 7-year post-baseline follow-up) <table border="1"> <tr> <td>CHRISTIAN ET AL 2013</td> <td>Baseline mean mins (SD)</td> <td>1-year follow-up mean mins (SD)</td> <td>3-year follow-up mean mins (SD)</td> </tr> <tr> <td>Intervention</td> <td>25.2 (55.33)</td> <td>15.2 (66.64)</td> <td>25.6 (70.90)</td> </tr> <tr> <td>Control</td> <td>28.1 (55.15)</td> <td>19.6 (50.55)</td> <td>25.7 (71.23)</td> </tr> </table> SD = standard deviation (calculated by reviewers) Change between baseline and final follow-up in intervention (-0.40 [95% CI -10.59, 9.79]) and change between baseline and final follow-up in control (2.4 [95% CI -5.28, 10.08]) were not significantly different (2.80 [95% CI -8.22, 13.93]) (calculated by reviewers). Prevalence of walking trips <table border="1"> <tr> <td>KNUIMAN ET AL 2014</td> <td>Baseline</td> <td>1-year follow-up</td> <td>3-year follow-up</td> <td>7-year follow-up</td> </tr> </table>	CHRISTIAN ET AL 2013	Baseline mean mins (SD)	1-year follow-up mean mins (SD)	3-year follow-up mean mins (SD)	Intervention	25.2 (55.33)	15.2 (66.64)	25.6 (70.90)	Control	28.1 (55.15)	19.6 (50.55)	25.7 (71.23)	KNUIMAN ET AL 2014	Baseline	1-year follow-up	3-year follow-up	7-year follow-up	VERY LOW
CHRISTIAN ET AL 2013	Baseline mean mins (SD)	1-year follow-up mean mins (SD)	3-year follow-up mean mins (SD)																								
Intervention	25.2 (55.33)	15.2 (66.64)	25.6 (70.90)																								
Control	28.1 (55.15)	19.6 (50.55)	25.7 (71.23)																								
KNUIMAN ET AL 2014	Baseline	1-year follow-up	3-year follow-up	7-year follow-up																							

¹⁸⁷ One study but two publications

¹⁸⁸ One publication splits data into intervention and control group, the other uses all data combined

¹⁸⁹ Confidence intervals spanned the line of no effect (and therefore the MID) – downgraded one level.

¹⁹⁰ One publication (Christian et al 2013) split participants into intervention (n=299) and control (n=528). The other publication (Knuiman et al 2014) treated all participants as intervention (n=1047)

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Percentage making a transport walking trip in a week (%)	37	28	29	36											
									Mean number of transport walking trips made per week	1.4	1.1	1.1	1.4											
Over time, proportion of participants making a walking trip and mean number of walking trips made increase to baseline levels.																								
Active Travel as measured by association between transport walking and public transport access, and number of possible destinations																								
1 Knuiman et al 2014	Non-randomised uncontrolled study	No serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	1,047	NA	<p><u>Active Travel as measured by association between transport walking and public transport access, and number of possible destinations (intervention only) (baseline to 1-, 3-, and 7-year post-baseline follow-up)</u></p> <p>Perceived (self-reported) and objective (GPS-identified) access to bus stops and railway stations are significantly associated with transport walking.</p> <p>Perceived number of types of destinations in the neighbourhood is more strongly associated with transport walking than objective measures of number of destinations present (both are significant when comparing presence of 8+ destinations with 0-3 destinations).</p> <p><u>Objective measures of public transport access and number of possible destinations in neighbourhood</u></p> <p>Association with objective walkability (OR, 95% CI)*</p> <table border="1"> <tr> <td>15-29 bus stops within 1600metres (compared with 0-14)</td> <td>1.63 (1.34, 1.98)</td> </tr> <tr> <td>≥30 bus stops within 1600metres (compared with 0-14)</td> <td>1.75 (1.39, 2.19)</td> </tr> <tr> <td>Railway station present within 1,600 metres of home</td> <td>1.34 (1.00, 1.81)</td> </tr> <tr> <td>4-7 types of destinations present (compared with 0-3)</td> <td>1.03 (0.87, 1.22)</td> </tr> <tr> <td>8-15 types of destinations present (compared with 0-3)</td> <td>1.29 (1.02, 1.64)</td> </tr> </table>					15-29 bus stops within 1600metres (compared with 0-14)	1.63 (1.34, 1.98)	≥30 bus stops within 1600metres (compared with 0-14)	1.75 (1.39, 2.19)	Railway station present within 1,600 metres of home	1.34 (1.00, 1.81)	4-7 types of destinations present (compared with 0-3)	1.03 (0.87, 1.22)	8-15 types of destinations present (compared with 0-3)	1.29 (1.02, 1.64)	LOW
15-29 bus stops within 1600metres (compared with 0-14)	1.63 (1.34, 1.98)																							
≥30 bus stops within 1600metres (compared with 0-14)	1.75 (1.39, 2.19)																							
Railway station present within 1,600 metres of home	1.34 (1.00, 1.81)																							
4-7 types of destinations present (compared with 0-3)	1.03 (0.87, 1.22)																							
8-15 types of destinations present (compared with 0-3)	1.29 (1.02, 1.64)																							

Physical Activity and the Environment – Appendix 4: GRADE profiles

									<p><u>Perceived measures of public transport access and number of possible destinations in neighbourhood</u></p> <p>Association with perceived walkability (OR, 95% CI)*</p> <p>Perceived access to bus stops (within 15-minute walk from home) 1.35 (1.10, 1.66)</p> <p>Perceived access to railway stations (within 15-minute walk from home) 1.44 (1.13, 1.85)</p> <p>3-6 types of destinations present (compared with 0-2) 2.07 (1.76, 2.43)</p> <p>7-11 types of destinations present (compared with 0-2) 2.32 (1.95, 2.77)</p>	
Active Travel as measured by association between transport walking and walkability										
1	Non-randomised uncontrolled study	No serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	1,047	NA	<p><u>Active Travel as measured by association between transport walking and walkability (intervention only) (baseline to 1-, 3-, and 7-year post-baseline follow-up)</u></p> <p>Walkability measure Association with perceived walkability (OR, [95% CI])* Association with objective walkability (OR, [95% CI])*</p> <p>Connectivity z-score 1.05 (0.99, 1.11) 1.09 (1.03, 1.15)</p> <p>Residential density z-score 1.04 (0.94, 1.15) 1.02 (0.92, 1.14)</p> <p>Land-use mix z-score 1.16 (1.08, 1.25) 1.21 (1.12, 1.30)</p> <p>*Odds Ratio, 95% confidence interval</p> <p>Objective (but not perceived) connectivity is significantly associated with transport walking. Neither perceived nor objective residential density mix is significantly associated with transport walking. Perceived and objective land-use mix is significantly associated with transport walking.</p>	LOW
Physical Activity in Everyday Life as measured by changes to time spent walking for recreation										
1	Non-randomised controlled study	No serious risk of bias	Not assessable as 1 study	No serious indirectness	Serious imprecision ¹⁹¹	None	299	528	<p><u>Mean minutes of recreational walking per week (baseline is prior to move) (intervention vs control) (baseline and 1- or 3-year follow-up)</u></p> <p>Baseline mean mins (SD) 1-year follow-up mean mins (SD) 3-year follow-up mean mins (SD)</p> <p>Intervention 65.9 (98.56) 85.4 (107.21) 95.1 (157.35)</p>	VERY LOW

¹⁹¹ Confidence intervals spanned the line of no effect (and therefore the MID) – downgraded one level.

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Control	77.3 (96.51)	91.4 (108.0)	86.2 (149.36)													
SD = standard deviation (calculated by reviewers) Change between baseline and final follow-up in intervention (29.20 [95% CI 8.15, 50.25]) and change between baseline and final follow-up in control (8.9 [95% CI -2.74, 20.54]) were not significantly different (20.30 [95% CI -3.67, 44.27]) (calculated by reviewers).																									
Physical Activity in Everyday Life as measured by travel mode to physical activity locations																									
1 Dunton et al 2012	Non- randomised controlled study	No serious risk of bias	Not assessable as 1 study	No serious indirectness	Serious imprecision ¹⁹²	None	46	48	<u>Physical Activity in Everyday Life as measured by travel mode to physical activity locations (children aged 9 – 13) (baseline to 6-12 month follow-up) (intervention vs control)</u> There was no difference between intervention and control changes over time (Group x time effect) for travel mode (walking or bicycling vs motorised transit) to place of physical activity (Adj. Wald F 0.46, p = 0.633).				VERY LOW												
Total Physical Activity as measured by change in time spent in MVPA per day																									
1 Dunton et al 2012	Non- randomised controlled study	No serious risk of bias	Not assessable as 1 study	No serious indirectness	Serious imprecision ¹⁹³	None	46	48	<u>Total Physical Activity as measured by change in time spent in MVPA per day (children aged 9 – 13) (baseline to 6-12 month follow-up) (intervention vs control)</u> <table border="1"> <thead> <tr> <th></th> <th>Baseline mean mins/day</th> <th>Follow-up mean mins/day</th> <th>Change in mins/day</th> </tr> </thead> <tbody> <tr> <td>Intervention</td> <td>32.75</td> <td>42.78</td> <td>10.03</td> </tr> <tr> <td>Control</td> <td>34.23</td> <td>38.40</td> <td>4.17</td> </tr> </tbody> </table> There was not a statistically significant difference between the change in intervention group and the change in control group (F Wald 0.44; p = 0.51)					Baseline mean mins/day	Follow-up mean mins/day	Change in mins/day	Intervention	32.75	42.78	10.03	Control	34.23	38.40	4.17	VERY LOW
	Baseline mean mins/day	Follow-up mean mins/day	Change in mins/day																						
Intervention	32.75	42.78	10.03																						
Control	34.23	38.40	4.17																						
Summary: See Evidence Statement 3.6																									

146
147
148

¹⁹² No difference in change in control and change in intervention – downgraded one level

¹⁹³ No significant difference in effect in control and effect in intervention – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
DIY-Streets (increasing safety and improving appearance of streets through planters, parking space provision and layout, and some traffic control methods)										
Physical Activity in Everyday Life as measured by self-reported activity levels (65+ years old)										
1 Ward Thompson et al 2014	Non-randomised controlled study	Serious risk of bias ¹⁹⁴	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	56 ¹⁹⁵	40 ¹⁹⁶	<u>Physical Activity in Everyday Life as measured by self-reported activity levels (65+ years old) (baseline to 2-year follow-up) (intervention vs control)</u> <u>Cross-sectional:</u> Self-reported frequency of summer outdoor activities significantly declined in the intervention group (p = 0.02) at 2 year follow-up. No significant differences for the comparison group (significance not reported). <u>Longitudinal:</u> Self-reported levels of outdoor activity in summer: did not increase significantly in either intervention or comparison groups (significance not reported). Participants in the intervention group perceived that they were more active at follow-up than baseline, significantly more so than those in the comparison group (p=0.04).	VERY LOW
Perceptions as measured by perceptions of safety and quality of facilities (65+ years old)										
1 Ward Thompson et al 2014	Non-randomised controlled study	Serious risk of bias ¹⁹⁷	Not assessable as 1 study	No serious indirectness	No serious imprecision	None	56 ¹⁹⁸	40 ¹⁹⁹	<u>Perceptions as measured by perceptions of safety and quality of facilities (65+ years old) (baseline to 2-year follow-up) (intervention vs control)</u> <u>Cross-sectional:</u>	VERY LOW

¹⁹⁴ Quality score was (-) indicating high risk of bias – downgraded one
¹⁹⁵ This is for cross-sectional data. 20 intervention participants for longitudinal analysis
¹⁹⁶ This is for cross-sectional data. 16 control participants for longitudinal analysis
¹⁹⁷ Quality score was (-) indicating high risk of bias – downgraded one
¹⁹⁸ This is for cross-sectional data. 20 intervention participants for longitudinal analysis
¹⁹⁹ This is for cross-sectional data. 16 control participants for longitudinal analysis

Physical Activity and the Environment – Appendix 4: GRADE profiles

								<p>In the intervention group, perceptions that “most of the streets and paths in my neighbourhood are safe to walk after dark” increased significantly ($p=0.04$). There was a significantly negative change in perceptions relating to “good outdoor facilities, including garden and parking, at home” ($p=0.02$). The comparison group saw no significant change over time.</p> <p><u>Longitudinal:</u> Responses to the statement ‘it is easy for me to walk on my street’ showed an increase in the intervention group, a change that was significant compared with the comparison group ($p=0.03$).</p>	
<p>SUMMARY: See Evidence Statement 3.7</p>									

149

Physical Activity and the Environment – Appendix 4: GRADE profiles

150 **Multicomponent**

151 To note, all studies reporting on parks interventions were observational, and therefore according to the GRADE process are initially classed as “Low”.

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Active living by design-Creation of city-level bike and pedestrian coordinator positions supporting implementation of environmental changes (crosswalks, park renovations etc), and extension of a walking path connecting intervention town with a city										
Total Physical Activity as measured by proportion of participants meeting either moderate or vigorous physical activity guidelines										
1 Chomitz et al 2012	Non-randomised controlled study	Serious risk of bias ²⁰⁰	Not assessable as single study	No serious indirectness	No serious imprecision	None	484	NA	<p><u>Total Physical Activity as measured by proportion of participants meeting either moderate physical activity (MPA) or vigorous physical activity (VPA) guidelines (baseline to 3-5 year post-baseline follow-up) (intervention only)</u></p> <p>The intervention is associated with significant increases in proportion of participants meeting MPA and/or VPA guidelines between baseline and 3-5 year follow-up in adults, from 40% at baseline to 62% at follow up, adjusted odds ratio 2.36 (95% CI 2.29,2.43) and high school students from 52% at baseline to 62% at follow up, adjusted odds ratio 1.61(1.34,1.92) but not in middle-school students who had a non-significant decrease</p>	VERY LOW
SUMMARY – See evidence statement 3.9										

152

153

²⁰⁰ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is selection bias and confounding

Physical Activity and the Environment – Appendix 4: GRADE profiles

154

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Implementation of "The District Approach": Improving green space through: redevelopment of existing parks; creation of public parks, natural playgrounds, community gardens, fishponds and public allotments.										
Physical Activity in Everyday Life as measured by proportion walking at least one a week										
1 Droomers et al 2016	Non-randomised controlled study	No serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	1018	1918 ²⁰¹ 3344 ²⁰² 46,885 ²⁰³ 229 ²⁰⁴	<p><u>Prevalence of respondents self-reporting leisure walking at least once/week:</u> Intervention and control groups, (baseline to 3.5 year post intervention implementation follow-up) Group x Time effect:</p> <p>When comparing intervention to each of the control groups, the difference in trend is only significant for a comparison with non-green neighbourhood control: -0.36 (95% CI -0.67, -0.05), indicating that the non-green neighbourhoods had significantly more positive change than intervention.</p> <p>Actual prevalence data not reported, only regression coefficients.</p>	LOW
Physical Activity in Everyday Life as measured by proportion making ≥1 leisure cycle trip/ week										
1 Droomers et al 2016	Non-randomised controlled study	No serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	1018	1918 ²⁰⁵ 3344 ²⁰⁶ 46,885 ²⁰⁷	<p><u>Prevalence of respondents self-reporting cycling for leisure at least once/week</u> Intervention and control groups, (baseline to 3.5 year post intervention implementation follow-up) Group x Time effect:</p>	LOW

²⁰¹ Narrow control: 1,918

²⁰² Broad control 3,344

²⁰³ Netherlands Control: 46,885

²⁰⁴ 12 non-green District Approach neighbourhoods Control: 229

²⁰⁵ Narrow control: 1,918

²⁰⁶ Broad control 3,344

²⁰⁷ Netherlands Control: 46,885

Physical Activity and the Environment – Appendix 4: GRADE profiles

								229 ²⁰⁸	There was no significant change in any of the groups when considering the prevalence of leisure cycling for at least once/week. The intervention group had a regression coefficient of -0.08 (95% CI -0.20, 0.04). The coefficient is very small suggesting the intervention had no effect on the prevalence of self-reported cycling. Actual prevalence data not reported, only regression coefficients.	
Physical Activity in Everyday Life as measured by proportion engaging in leisure sports at least once/week										
1 Droomers et al 2016	Non- randomised study	No serious risk of bias	Not assessable as single study	No serious indirectness	No serious imprecision	None	1018	1918 ²⁰⁹ 3344 ²¹⁰ 46,885 ²¹¹ 229 ²¹²	<p><u>Prevalence of respondents reporting engaging in sports for leisure at least once/</u></p> <p>Group x Time effect: There were no significant time, or group x time effects in any of the groups.</p> <p>The intervention group had a regression coefficient of -0.10 (95% CI -0.23,0.02), The small coefficient suggests the intervention had no effect on the prevalence of respondents reporting engaging in leisure sport for at least once/week.</p> <p>Actual prevalence data not reported, only regression coefficients</p>	LOW
SUMMARY – See evidence statement 3.10										

155

156

²⁰⁸ 12 non-green District Approach neighbourhoods Control: 229

²⁰⁹ Narrow control: 1,918

²¹⁰ Broad control 3,344

²¹¹ Netherlands Control: 46,885

²¹² 12 non-green District Approach neighbourhoods Control: 229

Physical Activity and the Environment – Appendix 4: GRADE profiles

157

Quality assessment							No. of participants		Effect	Quality
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control		
Smarter Choices, Smarter Places (SCSP) programme: introducing new bus services and shelters, ticketing improvements, promotional activity										
Total physical activity as measured by the proportion of participants meeting moderate physical activity (MPA) guidelines										
1 Norwood et al 2014	Non-randomised controlled study	serious risk of bias ²¹³	Not assessable as single study	No serious indirectness	No serious imprecision	None	7226	2316	<p><u>Proportion of participants meeting moderate physical activity (MPA) guidelines (intervention vs control; baseline vs 3-year follow-up):</u></p> <p>3-year follow-up: The proportion of participants meeting MPA guidelines was significantly greater in the intervention compared to control at 3 year follow up (p = <0.01; intervention = 30.8%; control = 24.9%).</p> <p>Change over time: Percentage of people meeting MPA guidelines was reduced in both groups between baseline and follow-up but this was greater in the control compared to the intervention (absolute reduction of 14.9% reduction in control vs. 3.4% reduction in intervention).</p> <p>Regression analysis, controlling for age, ownership of a car, employment status, health status, age, ethnicity, education level suggests that the likelihood of PA participation is significantly higher in the intervention areas relative to the control areas at 3 year follow up (p = <0.001, regression coefficient for area by year is 0.39.)</p>	Very LOW
Total physical activity as measured by the proportion of participants who were active at all										
1 Norwood et al 2014	Non-randomised controlled study	serious risk of bias ²¹⁴	Not assessable as single study	No serious indirectness	No serious imprecision	None	7226	2316	<p><u>Proportion of participants who were active at all (intervention vs control; baseline vs 3-year follow-up):</u></p> <p>3-year follow-up: The proportion of participants who were active at all was not significantly different between control and intervention areas (P value not reported; intervention = 69.9%; control = 70.1%).</p>	VERY LOW

²¹³ Quality score allocated as (-) indicating high risk of bias, main bias self reported outcome – downgraded one level.

²¹⁴Quality score allocated as (-) indicating high risk of bias, main bias was self reported outcome – downgraded one level.

Physical Activity and the Environment – Appendix 4: GRADE profiles

									Change over time: Proportions of participants who were active at all reduced in both groups between baseline and follow-up but this was greater in the control compared with the intervention (absolute reduction of 9.2% vs 0.7%). Those who are physically active are significantly more likely to meet physical activity guidelines in the intervention areas relative to the control areas (regression coefficient 0.13; p = <0.05).	
SUMMARY- See evidence statement 3.11										

158

159

Physical Activity and the Environment – Appendix 4: GRADE profiles

160

Quality assessment							No. of participants		Effect	Quality																								
No of studies	Design	Risk of bias	In-consistency	In-directness	Imprecision	Other considerations	Intervention	Control																										
Active England woodland projects (new play areas, visitor's centre, cycle tracks, walking trails, shower facilities, butterfly trail, climbing wall, promotional groups and events)																																		
Physical activity in everyday life as measured by the change in frequency of visits																																		
1 O'Brien and Morris 2009	Non-randomised uncontrolled study	No serious risk of bias	Not assessable as single study	No serious indirectness	Serious imprecision ²¹⁵	None	1467	NA	<p><u>Change in frequency of visits (as % of all visitors)</u> (follow up unclear – differed from each site) (1-3 years and 4-5 years)</p> <p>Total visitor numbers increased by between 47% and 2,143% between baseline and follow-up.</p> <table border="1"> <thead> <tr> <th></th> <th>Before</th> <th>After</th> </tr> </thead> <tbody> <tr> <td>Every day</td> <td>7.3</td> <td>2.2</td> </tr> <tr> <td>4-6 / week</td> <td>6.7</td> <td>3</td> </tr> <tr> <td>1-3 / week</td> <td>19</td> <td>19</td> </tr> <tr> <td>1-3 / month</td> <td>22.3</td> <td>27.6</td> </tr> <tr> <td>4-6 / year</td> <td>9</td> <td>19.2</td> </tr> <tr> <td>1-3 / year</td> <td>19.7</td> <td>18.8</td> </tr> <tr> <td>Less often</td> <td>15.9</td> <td>10.1</td> </tr> </tbody> </table> <p>Those visiting every day or 4-6 times per week declined as a proportion of all visitors. Those visiting 1-3 times per month and 4-6 times per year saw the greatest increase as a proportion of all visitors. Average visit time reportedly increased from 1.74 (standard error 0.04) to 2.33 (standard error 0.04).</p>		Before	After	Every day	7.3	2.2	4-6 / week	6.7	3	1-3 / week	19	19	1-3 / month	22.3	27.6	4-6 / year	9	19.2	1-3 / year	19.7	18.8	Less often	15.9	10.1	Very low
	Before	After																																
Every day	7.3	2.2																																
4-6 / week	6.7	3																																
1-3 / week	19	19																																
1-3 / month	22.3	27.6																																
4-6 / year	9	19.2																																
1-3 / year	19.7	18.8																																
Less often	15.9	10.1																																

²¹⁵No measures of variance reported – downgraded one level

Physical Activity and the Environment – Appendix 4: GRADE profiles

1 O'Brien and Morris 2009	Non-randomised uncontrolled study	No serious risk of bias	Not assessable as single study	No serious indirectness	Serious imprecision ²¹⁶	None	1467	NA	<p><u>Sub group analysis</u> In all three sites combined, there was no significant change in number of visitors with blue badges (actual numbers not given), however there was a decrease in proportion of visitors reporting having a long term illness (13.9% at baseline, 7.2% at follow-up; $p = <0.001$; actual numbers not reported). Black and Minority Ethnic (BME) individuals as a proportion of all visitors increased from 1.7% at baseline to 5.2% at follow up ($p = <0.001$).</p>	
Total Physical activity as measured by proportion of visitors taking ≥ 5 days exercise/week										
1 O'Brien and Morris 2009	Non-randomised uncontrolled study	serious risk of bias ²¹⁷	Not assessable as single study	No serious indirectness	Serious imprecision	None	1467	NA	<p><u>Total Physical activity as measured by proportion of visitors taking ≥ 5 days exercise/week</u> follow up unclear – differed from each site (1-3 years and 4-5 years) Proportion of visitors taking ≥ 5 days exercise/week declined from 55.9% to 36.1% between baseline and follow-up ($p = <0.001$).</p>	Very Low
Changes to perceived barriers to accessing forests for physical activity										
1 O'Brien and Morris 2009	Non-randomised uncontrolled study	serious risk of bias ²¹⁸	Not assessable as single study	No serious indirectness	Serious imprecision ²¹⁹	None	1467	NA	<p><u>Changes to perceived barriers to accessing forests for physical activity</u> follow up unclear – differed from each site (1-3 years and 4-5 years) [To note – Actual numbers and statistical significance not reported. NICE team derived this information from a bar chart with no number labels]. The largest changes in perceived barriers occurred in: lack of facilities, antisocial behaviour and lack of information (where there was a decrease in perceived barrier from baseline to follow-up). Compared with baseline, respondents were more likely to perceive weather as a barrier and have a preference for other countryside areas</p>	Very Low

²¹⁶No measures of variance reported – downgraded one level

²¹⁷ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is presence of selection bias and poor data collection methods

²¹⁸ Quality score allocated as (-) indicating high risk of bias – downgraded one level. Main bias is presence of selection bias and poor data collection methods

²¹⁹ No measures of variance reported – downgraded one level

SUMMARY – See evidence statement 3.12	
--	--

161