

Draft for consultation

# **Meningitis (bacterial) and meningococcal disease: recognition, diagnosis and management**

**[E2] Evidence review for antibiotics for bacterial  
meningitis caused by *Haemophilus influenzae***

*NICE guideline number tbc*

*Evidence review underpinning recommendations 1.6.4, 1.6.11  
and 1.6.16 in the NICE guideline*

*September 2023*

*Draft for consultation*

*This evidence review was developed by NICE*



### **Disclaimer**

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

### **Copyright**

© NICE 2023. All rights reserved. Subject to [Notice of rights](#).

ISBN:

## Contents

<b>Antibiotics for bacterial meningitis caused by Haemophilus influenzae .....</b>	<b>6</b>
Review question .....	6
Introduction .....	6
Summary of the protocol .....	6
Methods and process .....	7
Effectiveness evidence .....	7
Summary of included studies .....	8
Summary of the evidence .....	9
Economic evidence .....	9
Economic model .....	9
The committee's discussion and interpretation of the evidence .....	9
Recommendations supported by this evidence review .....	12
References – included studies .....	13
<b>Appendices .....</b>	<b>14</b>
<b>Appendix A    Review protocols .....</b>	<b>14</b>
Review protocol for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	14
<b>Appendix B    Literature search strategies .....</b>	<b>23</b>
Literature search strategies for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	23
<b>Appendix C    Effectiveness evidence study selection .....</b>	<b>30</b>
Study selection for: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	30
<b>Appendix D    Evidence tables .....</b>	<b>31</b>
Evidence tables for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	31
<b>Appendix E    Forest plots .....</b>	<b>39</b>
Forest plots for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	39
<b>Appendix F    GRADE tables .....</b>	<b>40</b>
GRADE tables for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	40
<b>Appendix G    Economic evidence study selection .....</b>	<b>43</b>
Study selection for: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	43
<b>Appendix H    Economic evidence tables .....</b>	<b>44</b>

	Economic evidence tables for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	44
<b>Appendix I</b>	<b>Economic model</b> .....	<b>45</b>
	Economic model for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	45
<b>Appendix J</b>	<b>Excluded studies</b> .....	<b>46</b>
	Excluded studies for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	46
<b>Appendix K</b>	<b>Research recommendations – full details</b> .....	<b>48</b>
	Research recommendations for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae? .....	48

# Antibiotics for bacterial meningitis caused by Haemophilus influenzae

## Review question

What antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae?

## Introduction

Bacterial meningitis is a rare but serious infection. The causative organism is usually confirmed by tests performed on cerebrospinal fluid or blood samples. Haemophilus influenzae is now a very rare cause of bacterial meningitis in the UK due to the success of the childhood vaccination programme.

The aim of this review is to determine what antibiotic treatment regimens are effective in treating bacterial meningitis caused by Haemophilus influenzae.

## Summary of the protocol

See Table 1 for a summary of the Population, Intervention, Comparison and Outcome (PICO) characteristics of this review.

**Table 1: Summary of the protocol (PICO table)**

<b>Population</b>	All adults, young people, children and babies (excluding neonates defined as aged 28 days old and younger) with confirmed bacterial meningitis caused by Haemophilus influenzae.
<b>Intervention</b>	Antibiotic agent of interest: <ul style="list-style-type: none"><li>• Cefotaxime</li><li>• Ceftriaxone</li><li>• Amoxicillin</li><li>• Ampicillin</li><li>• Meropenem</li><li>• Chloramphenicol</li><li>• Aztreonam</li></ul>
<b>Comparison</b>	<b>Stage 1 (all antibiotic agents of interest):</b> <ul style="list-style-type: none"><li>• Antibiotic agent A (single or combination) vs Antibiotic agent B (single or combination)</li></ul> <b>Stage 2 (antibiotic agents identified during stage 1 as most effective/for use where there are contraindications)</b> <ul style="list-style-type: none"><li>• Antibiotic agent A – Dose A vs Antibiotic agent A – Dose B</li><li>• Antibiotic agent A – Duration of administration A vs Antibiotic agent A – Duration of administration B</li><li>• Antibiotic agent A – Short infusion vs Antibiotic agent A – Extended infusion</li></ul>
<b>Outcome</b>	<b>Critical</b> Population: adults, children and infants <ul style="list-style-type: none"><li>• All-cause mortality (measured up to 1 year after discharge)</li><li>• Any long-term neurological impairment (defined as any motor deficits, sensory deficits [excluding hearing impairment], cognitive deficits*, or behavioural deficits*; measured from discharge up to 1 year after discharge)</li></ul> Population: adults <ul style="list-style-type: none"><li>• Functional impairment (measured by any validated scale at any time point)</li></ul>

Population: children and infants

- Severe developmental delay (defined as score of >2 SD below normal on validated assessment scales, or MDI or PDI <70 on Bayleys assessment scale, or inability to assign a score due to cerebral palsy or severity of cognitive delay; measured at the oldest age reported unless there is substantially more data available at a younger age)

#### Important

Population: adults, children and infants

- Hearing impairment (defined as any level of hearing impairment; measured from discharge up to 1 year after discharge)
- Serious intervention-related adverse effects leading to death, disability or prolonged hospitalisation or that are life threatening or otherwise considered medically significant
- CSF sterilisation (defined as treatment failure, time-to-sterilisation or delay)

Population: adults

- Intracranial collections as a complication (defined as abscess or empyema)

Population: children and infants

- Functional impairment (measured by any validated scale at any time point)

\*For infants and children below school-age, cognitive and behavioural deficits will be assessed at school-age.

1 CSF: cerebrospinal fluid; MDI: mental development index; PDI: psychomotor development index; SD: standard  
2 deviation

3 For further details see the review protocol in appendix A.

## 4 Methods and process

5 This evidence review was developed using the methods and process described in  
6 [Developing NICE guidelines: the manual](#). Methods specific to this review question are  
7 described in the review protocol in appendix A and the methods document (supplementary  
8 document 1).

9 Declarations of interest were recorded according to [NICE's conflicts of interest policy](#).

## 10 Effectiveness evidence

### 11 Included studies

12 Two studies were included for this review: 1 retrospective cohort study (Lapointe 1988), and  
13 1 randomised controlled trial (RCT: Molyneux 2011).

14 The included studies are summarised in Table 2.

15 One study compared cefotaxime to chloramphenicol (Lapointe 1988) and did not adjust for  
16 confounding factors. One study compared 5-day ceftriaxone therapy to 10-day ceftriaxone  
17 therapy (Molyneux 2011). All studies were conducted in babies and children (Lapointe 1988;  
18 Molyneux 2011).

19 See the literature search strategy in appendix B and study selection flow chart in appendix C.

### 20 Excluded studies

21 Studies not included in this review are listed, and reasons for their exclusion are provided in  
22 appendix J.

1 **Summary of included studies**

2 Summaries of the studies that were included in this review are presented in Table 2.

3 **Table 2: Summary of included studies**

Study	Population	Intervention	Comparison	Outcomes	Comments
Lapointe 1988  Retrospective cohort study  Canada	N=62  Babies and children with bacterial meningitis caused by $\beta$ -lactamase-producing and ampicillin-resistant Haemophilus influenzae  Age in months (mean): 16  Case-fatality: 0%	<u>Cefotaxime</u>  IV cefotaxime 200 mg/kg/day for 10 days*  *IV ampicillin 200 mg/kg/day also given in the first 48 h	<u>Chloramphenicol</u>  IV chloramphenicol 100 mg/kg/day for 10 days*  *IV ampicillin 200-400 mg/kg/day also given in the first 48 h	<ul style="list-style-type: none"> <li>All-cause mortality</li> <li>Any long-term neurological impairment</li> </ul>	Analyses unadjusted for confounding factors
Molyneux 2011  RCT  Bangladesh, Egypt, Malawi, Pakistan and Vietnam	H. influenzae N=266 (whole study N=1004)  Babies and children aged 2 months to 12 years with bacterial meningitis caused by H. influenzae*, S. pneumoniae, or N. meningitidis  Age in months (mean; SD): 38 (42)  Population treated with dexamethasone therapy: 44%  Case-fatality: 4%  *H. influenzae is causative organism of interest for this review	<u>5-day ceftriaxone therapy</u>  IV ceftriaxone 80-100 mg/kg once daily for 5 days	<u>10-day ceftriaxone therapy</u>  IV ceftriaxone 80-100 mg/kg once daily for 10 days	<ul style="list-style-type: none"> <li>All-cause mortality</li> <li>Any long-term neurological impairment</li> <li>Developmental delay</li> <li>Hearing impairment</li> <li>Serious intervention-related adverse effects</li> <li>CSF sterilisation</li> </ul>	Population is indirect for all outcomes except for all-cause mortality and CSF sterilisation due to 74% of population with meningitis caused by organisms other than H. influenzae

4 CSF: cerebrospinal fluid; IV: intravenous; RCT: randomised controlled trial; SD: standard deviation



1 See the full evidence tables in appendix D. No meta-analysis was conducted (and so there  
2 are no forest plots in appendix E).

### 3 **Summary of the evidence**

4 This section is a narrative summary of the findings of the review, as presented in the GRADE  
5 tables in appendix F. For details of the committee's confidence in the evidence and how this  
6 affected recommendations, see The committee's discussion and interpretation of the  
7 evidence.

8 The evidence was assessed as being very low quality due to risk of bias (arising from  
9 missing outcome data due to attrition, subjective measurement of the outcome, selective  
10 reporting, and failure to adjust for confounding factors), seriously imprecise findings and the  
11 inclusion of indirect populations and outcomes.

12 No important difference in mortality was shown between cefotaxime and chloramphenicol in  
13 the evidence reviewed, although cefotaxime was associated with a lower rate of neurological  
14 impairment.

15 The evidence showed no important differences between 5-day and 10-day ceftriaxone  
16 treatment for all-cause mortality, neurological impairment, developmental delay, hearing  
17 impairment, serious intervention-related adverse effects, or cerebrospinal fluid (CSF)  
18 sterilisation.

19 No eligible studies were identified that reported on functional impairment, or intracranial  
20 collections as a complication.

21 See appendix F for full GRADE tables.

### 22 **Economic evidence**

#### 23 **Included studies**

24 A single economic search was undertaken for all topics included in the scope of this  
25 guideline, but no economic studies were identified which were applicable to this review  
26 question.

#### 27 **Economic model**

28 No economic modelling was undertaken for this review because the committee agreed that  
29 other topics were higher priorities for economic evaluation.

### 30 **The committee's discussion and interpretation of the evidence**

#### 31 **The outcomes that matter most**

32 Bacterial meningitis is associated with high rates of mortality and morbidity, and antibiotics  
33 are the mainstay of treatment for bacterial meningitis. Therefore, all-cause mortality and  
34 long-term neurological impairment were prioritised as critical outcomes due to the severity of  
35 these outcomes. Severe developmental delay was prioritised over functional impairment in  
36 children and babies, as it is a more relevant and important outcome for this population.  
37 Functional impairment was prioritised as a critical outcome in adults due to the concern  
38 about the potential long-term limitations of bacterial meningitis on the ability to carry out  
39 certain activities of daily life.

40 In addition to functional impairment (in children and babies), hearing impairment, serious  
41 intervention-related adverse effects and CSF sterilisation were selected as important

1 outcomes in all age groups as these are relatively common after bacterial meningitis and  
2 may be related to antibiotic therapy. Intracranial collections as a complication was also  
3 included as an important outcome for adults as this is a rare but severe and life threatening  
4 complication of bacterial meningitis that may require prolonged antibiotic treatment.

#### 5 **The quality of the evidence**

6 The quality of the evidence was assessed using GRADE methodology. The evidence for all  
7 outcomes in this review was very low quality, and the main reasons for downgrading the  
8 evidence were risk of bias (arising from missing outcome data, subjective measurement of  
9 the outcome, selective reporting, and failure to adjust for confounding factors), imprecision  
10 (due to wide confidence intervals and small number of events), and indirectness (of either  
11 population, outcome, or both).

12 No evidence was found for functional impairment, or intracranial collections as a  
13 complication.

#### 14 **Benefits and harms**

15 The committee considered the evidence comparing cefotaxime and chloramphenicol for the  
16 treatment of meningitis caused by *Haemophilus influenzae* type b (Hib), that showed no  
17 important difference for mortality, but a lower rate of long-term neurological impairment  
18 associated with cefotaxime. However, the committee noted that this evidence came from a  
19 retrospective cohort study with a small sample size. No other evidence was identified  
20 comparing the effectiveness of different antibiotics for the treatment of Hib meningitis. Given  
21 the limitations of the evidence, the committee agreed to make recommendations based on  
22 their clinical knowledge and experience, and on current practice, and recommended  
23 ceftriaxone in line with the BNF (Joint Formulary Committee 2022) and BNFC (Paediatric  
24 Formulary Committee 2022), for the treatment of Hib meningitis.

25 The committee highlighted the potential practical and resource-use advantages associated  
26 with ceftriaxone because the long half-life means that it can be given only once a day. The  
27 committee acknowledged some concerns with once daily administration in that a second  
28 dose might need to be delayed if the first dose of ceftriaxone was administered outside of  
29 routine working hours; however, they were aware that a second dose can be given earlier, to  
30 shift the administration time, if there is a minimum of 12 hours between doses (Gbesemete  
31 2019).

32 The committee discussed some reasons why in clinical practice (particularly in intensive care  
33 units) cefotaxime might be given instead of ceftriaxone. For instance, to minimise the time  
34 that intravenous lines are being used for administering antibiotics, which might be needed for  
35 other medications, due to ceftriaxone typically being infused over 30 minutes intravenous  
36 and cefotaxime being given as a bolus. However, the committee agreed that this practice is  
37 not necessary, as ceftriaxone can be given as bolus. Sometimes there may be a reaction (for  
38 example, vomit reflex) if ceftriaxone is administered too quickly, but in the committee's  
39 experience this is relatively rare, which was supported by a recent study (Patel 2021). The  
40 committee agreed that ceftriaxone should be given as first-line treatment for Hib meningitis,  
41 unless contraindicated in which case cefotaxime can be considered. The committee  
42 highlighted contraindications around pre-term babies less than 41 weeks corrected  
43 gestational age and included a reference to these in the BNFC in the recommendation in  
44 order to raise awareness.

45 The committee agreed that advice from an infection specialist (a microbiologist or infectious  
46 diseases specialist) should be sought, given that Hib infection is now extremely rare in the  
47 UK due to vaccination.

48 The committee were aware that the previous NICE guideline on meningitis (NICE 2010)  
49 recommended 10-day antibiotic treatment for Hib meningitis. However, the committee noted

1 that the evidence reviewed showed no important difference between 5 and 10 days of  
2 ceftriaxone therapy, although this study was unlikely to have been adequately powered to be  
3 taken as definitive evidence of equivalence. The committee acknowledged that practice has  
4 changed since the previous NICE guideline, and that the previous recommendations were  
5 consensus rather than evidence based and pre-dated the widespread use of cephalosporins.  
6 The committee discussed that, in some instances, practice has moved to shorter (7-day)  
7 courses of antibiotics for the treatment of Hib meningitis without apparent impact on clinical  
8 outcomes, although they acknowledged that there is variation in practice. The committee  
9 were also aware of evidence from low- and middle-income countries, suggesting that shorter  
10 length of treatment may be effective. The committee recommended that people with  
11 meningitis caused by Hib should be treated for 7-10 days with ceftriaxone (or cefotaxime if  
12 ceftriaxone contraindicated). The committee agreed that recommending a range of 7 to 10  
13 days provided flexibility to stop at 7 days if the person had recovered or continue for 10 days  
14 if they had not. The committee agreed that further advice from an infection specialist should  
15 be sought if patients have not recovered after 10 days.

16 There was no evidence found on antibiotic use for Hib meningitis in people with a penicillin  
17 allergy, but the committee agreed it was important to make a recommendation for this  
18 population. Based on their knowledge and experience, the committee agreed that  
19 cephalosporin-induced anaphylaxis is rare, and the risk-benefit balance of a cephalosporin  
20 (relative to chloramphenicol as an alternative) is favourable in most patients with non-  
21 anaphylactic penicillin allergy. Therefore, the committee agreed that clinicians should seek  
22 information about the nature of the allergy and advice from an infection specialist before  
23 making a treatment decision. The committee acknowledged that it is important that treatment  
24 is not delayed; however, they agreed that information about the nature of allergy is often  
25 readily available from the patient's family. The committee agreed that a cephalosporin should  
26 still be considered if the nature of the allergic reaction they get is non-anaphylactic or non-  
27 severe, in accordance with the first line treatment recommended above. However, if the  
28 allergic reaction is anaphylactic or severe, alternatives to ceftriaxone or cefotaxime will be  
29 needed. The committee discussed that chloramphenicol is commonly used in the case of  
30 severe beta-lactam (penicillin, amoxicillin, or cephalosporin) allergy. Based on clinical  
31 knowledge and experience, the committee recommended chloramphenicol for Hib meningitis  
32 and anaphylactic or severe penicillin allergy.

33 The committee were aware that the previous NICE guideline on bacterial meningitis (NICE  
34 2010) recommended to treat people who have travelled outside the UK with vancomycin (in  
35 addition to the cephalosporin). However, they discussed that practice has changed since the  
36 previous NICE guideline. The committee were aware that current practice is to use rifampicin  
37 or linezolid in addition to a cephalosporin where the cephalosporin itself might be insufficient  
38 due to resistance. However, the committee highlighted that there is not enough evidence to  
39 support recommending them. Therefore, the committee recommended that, clinicians should  
40 seek advice from an infection specialist if cephalosporin resistance is suspected in people  
41 who have recently travelled abroad.

#### 42 **Cost effectiveness and resource use**

43 This review question was not prioritised for economic analysis and therefore the committee  
44 made a qualitative assessment of the likely cost-effectiveness of their recommendations. The  
45 committee considered that it would be cost-effective to facilitate the option of a shorter  
46 course of antibiotics than in previous NICE guidance (NICE 2010) for meningitis caused by  
47 Hib as some practice has moved in this direction without any apparent adverse impact on  
48 clinical outcomes. However, given the limitations of the evidence supporting shorter courses  
49 they did not want to mandate this, especially as they reasoned that the resource and  
50 antibiotic resistance implications of 3 days extra treatment would be minimal. The committee  
51 believed that by facilitating the option of a shorter course of antibiotics that their  
52 recommendation could lead to some small cost savings for the NHS.

1 **Recommendations supported by this evidence review**

2 This evidence review supports recommendations 1.6.4, 1.6.11, and 1.6.16. Other evidence  
3 supporting the recommendations 1.6.4 and 1.6.16 can be found in evidence reviews on  
4 antibiotic regimens for bacterial meningitis before or in the absence of identifying causative  
5 infecting organism (see evidence reviews D1 to D3) and for specific causative organisms  
6 (see evidence reviews E1 and E3 to E6).

7

## 1 **References – included studies**

### 2 **Effectiveness**

#### 3 **Lapointe 1988**

4 Lapointe, J. R. and Chicoine, L. (1988) Cefotaxime versus chloramphenicol for ampicillin-  
5 resistant Haemophilus influenzae meningitis. A retrospective study of 62 cases. *Drugs*  
6 35suppl2: 199-202

#### 7 **Molyneux 2011**

8 Molyneux, Elizabeth, Nizami, Shaikh Qamaruddin, Saha, Samir et al. (2011) 5 versus 10  
9 days of treatment with ceftriaxone for bacterial meningitis in children: a double-blind  
10 randomised equivalence study. *Lancet* (London, England) 377(9780): 1837-45

### 11 **Economic**

12 No studies were identified which were applicable to this review question.

### 13 **Other**

#### 14 **Gbesemete 2019**

15 Gbesemete, D., Faust, S. (2019). Prescribing in infection: antibacterials. In: Barker, C.,  
16 Turner, M., Sharland, M. (Eds.) *Prescribing Medicines for Children: From drug development*  
17 *to practical administration*, Pharmaceutical Press, London: UK

#### 18 **Joint Formulary Committee 2022**

19 Joint Formulary Committee. *British National Formulary* (online). London: BMJ Group and  
20 Pharmaceutical Press. Available at: <http://www.medicinescomplete.com> [Accessed  
21 04/04/2022]

#### 22 **NICE 2010**

23 National Institute for Health and Care Excellence (2010). *Meningitis (bacterial) and*  
24 *meningococcal septicaemia in under 16s: recognition, diagnosis and management*. Available  
25 at: <https://www.nice.org.uk/guidance/cg102> [Accessed 04/04/2022]

#### 26 **Paediatric Formulary Committee 2022**

27 Paediatric Formulary Committee. *BNF for Children* (online). London: BMJ Group,  
28 Pharmaceutical Press, and RCPCH Publications. Available at:  
29 <http://www.medicinescomplete.com> [Accessed 29/03/2022]

#### 30 **Patel 2021**

31 Patel, S., Green, H., Gray, J., Rutter, M., Bevan, A., Hand, K., Jones, C. E., Faust, S. N.  
32 (2021). Evaluating Ceftriaxone 80 mg/kg Administration by Rapid Intravenous Infusion—A  
33 Clinical Service Evaluation. *The Pediatric Infectious Disease Journal*, 40(2), 128-129

# 1 Appendices

## 2 Appendix A Review protocols

### 3 Review protocol for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis 4 caused by *Haemophilus influenzae*?

5 **Table 3: Review protocol**

Field	Content
PROSPERO registration number	CRD42021276561
Review title	Antibiotics for bacterial meningitis caused by <i>Haemophilus influenzae</i>
Review question	What antibiotic treatment regimens are effective in treating bacterial meningitis caused by <i>Haemophilus influenzae</i> ?
Objective	This review aims to find out what is the optimal antibiotic treatment regimen in improving outcomes for people with bacterial meningitis caused by <i>Haemophilus influenzae</i>
Searches	<p>The following databases will be searched:</p> <ul style="list-style-type: none"> <li>Cochrane Central Register of Controlled Trials (CENTRAL)</li> <li>Cochrane Database of Systematic Reviews (CDSR)</li> <li>Embase</li> <li>MEDLINE</li> </ul> <p>Searches will be restricted by:</p> <ul style="list-style-type: none"> <li>Date limitations: 1980</li> <li>English language</li> <li>Human studies</li> </ul> <p>The full search strategies for MEDLINE database will be published in the final review. For each search, the principal database search strategy is quality assured by a second information scientist using an adaptation of the PRESS 2015 Guideline Evidence-Based Checklist.</p>

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

---

Field	Content
Condition or domain being studied	Bacterial meningitis caused by Haemophilus influenzae
Population	<p>Inclusion: All adults, young people, children and babies (excluding neonates defined as aged 28 days old and younger) with confirmed bacterial meningitis caused by Haemophilus influenzae</p> <p>Exclusion:</p> <p>People:</p> <ul style="list-style-type: none"><li>• with known immunodeficiency.</li><li>• who have brain tumours, pre-existing hydrocephalus, intracranial shunts, previous neurosurgical procedures, or known cranial or spinal anomalies that increase the risk of bacterial meningitis.</li><li>• with confirmed viral meningitis or viral encephalitis.</li><li>• with confirmed tuberculous meningitis.</li><li>• with confirmed fungal meningitis.</li></ul>
Intervention/Exposure/Test	<ul style="list-style-type: none"><li>• Cefotaxime</li><li>• Ceftriaxone</li><li>• Amoxicillin</li><li>• Ampicillin</li><li>• Meropenem</li><li>• Chloramphenicol</li><li>• Aztreonam</li></ul>
Comparator/Reference standard/Confounding factors	<p>Stage 1 (all antibiotic agents of interest): Antibiotic agent A (single or combination) vs Antibiotic agent B (single or combination)</p> <p>Stage 2 (antibiotic agents identified during stage 1 as most effective/for use where there are contraindications)</p> <p>Comparisons:</p> <ol style="list-style-type: none"><li>1. Antibiotic agent A – Dose A vs Antibiotic agent A – Dose B</li><li>2. Antibiotic agent A – Duration of administration A vs Antibiotic agent A – Duration of</li></ol>

Field	Content
	administration B 3. Antibiotic agent A – Short infusion vs Antibiotic agent A – Extended infusion
Types of study to be included	<p>Include published full-text papers:</p> <ul style="list-style-type: none"> <li>• Systematic reviews of RCTs</li> <li>• RCTs</li> <li>• If insufficient RCTs: prospective cohort studies</li> <li>• If insufficient prospective cohort studies: retrospective cohort studies</li> </ul> <p>Non-randomised studies will be downgraded for risk of bias if they do not adequately adjust for the following covariates, but will not be excluded for this reason:</p> <ul style="list-style-type: none"> <li>• Co-morbidity</li> <li>• Severity of infection at presentation (including sepsis)</li> </ul> <p>Exclude:</p> <ul style="list-style-type: none"> <li>• Conference abstracts</li> </ul>
Other exclusion criteria	<p>Cohort studies from low income countries.</p> <p>Studies conducted prior to 1980 as currently used antibiotics were not in common usage prior to this date.</p> <p>Studies published not in English-language</p>
Context	This guidance will fully update the following: Meningitis (bacterial) and meningococcal septicaemia in under 16s: recognition, diagnosis and management (CG102)
Primary outcomes (critical outcomes)	<p>Adults</p> <ul style="list-style-type: none"> <li>• All-cause mortality (measured up to 1 year after discharge)</li> <li>• Any long-term neurological impairment (defined as any motor deficits, sensory deficits [excluding hearing impairment], cognitive deficits, or behavioural deficits; measured from discharge up to 1 year after discharge)</li> <li>• Functional impairment (measured by any validated scale at any time point)</li> </ul> <p>Children and infants</p> <ul style="list-style-type: none"> <li>• All-cause mortality (measured up to 1 year after discharge)</li> </ul>



Field	Content
	<ul style="list-style-type: none"> <li>• Any long-term neurological impairment (defined as any motor deficits, sensory deficits [excluding hearing impairment], cognitive deficits*, or behavioural deficits*; measured from discharge up to 1 year after discharge)</li> <li>• Severe developmental delay (defined as score of &gt;2 SD below normal on validated assessment scales, or MDI or PDI &lt;70 on Bayleys assessment scale, or inability to assign a score due to cerebral palsy or severity of cognitive delay; measured at the oldest age reported unless there is substantially more data available at a younger age)</li> </ul> <p>*For infants and children below school-age, cognitive and behavioural deficits will be assessed at school-age.</p>
Secondary outcomes (important outcomes)	<p>Adults</p> <ul style="list-style-type: none"> <li>• Hearing impairment (defined as any level of hearing impairment; measured from discharge up to 1 year after discharge)</li> <li>• Serious intervention-related adverse effects leading to death, disability or prolonged hospitalisation or that are life threatening or otherwise considered medically significant</li> <li>• CSF sterilisation (defined as treatment failure, time-to-sterilisation or delay).</li> <li>• Intracranial collections as a complication (defined as abscess or empyema)</li> </ul> <p>Children and infants</p> <ul style="list-style-type: none"> <li>• Hearing impairment (defined as any level of hearing impairment; measured from discharge up to 1 year after discharge)</li> <li>• Functional impairment (measured by any validated scale at any time point)</li> <li>• Serious intervention-related adverse effects leading to death, disability or prolonged hospitalisation or that are life threatening or otherwise considered medically significant</li> <li>• CSF sterilisation (defined as treatment failure, time to sterilisation or delay)</li> </ul>
Data extraction (selection and coding)	All references identified by the searches and from other sources will be uploaded into STAR and de-duplicated. Titles and abstracts of the retrieved citations will be screened to identify studies that potentially meet the inclusion criteria outlined in the review

Field	Content
	<p>protocol. Dual sifting will not be undertaken for this question. Full versions of the selected studies will be obtained for assessment. Studies that fail to meet the inclusion criteria once the full version has been checked will be excluded at this stage. Each study excluded after checking the full version will be listed, along with the reason for its exclusion. A standardised form will be used to extract data from studies. The following data will be extracted: study details (reference, country where study was carried out, type and dates), participant characteristics, inclusion and exclusion criteria, details of the interventions if relevant, setting and follow-up, relevant outcome data and source of funding. One reviewer will extract relevant data into a standardised form, and this will be quality assessed by a senior reviewer.</p>
Risk of bias (quality) assessment	<p>Quality assessment of individual studies will be performed using the following checklists:</p> <ul style="list-style-type: none"> <li>• ROBIS tool for systematic reviews</li> <li>• Cochrane RoB tool v.2 for RCTs and quasi-RCTs</li> <li>• Cochrane ROBINS-I tool for non-randomised (clinical) controlled trials and cohort studies</li> </ul> <p>The quality assessment will be performed by one reviewer and this will be quality assessed by a senior reviewer.</p>
Strategy for data synthesis	<p>Quantitative findings will be formally summarised in the review. Where multiple studies report on the same outcome for the same comparison, meta-analyses will be conducted using Cochrane Review Manager software. A fixed effect meta-analysis will be conducted and data will be presented as risk ratios if possible or odds ratios when required (for example if only available in this form in included studies) for dichotomous outcomes, and mean differences or standardised mean differences for continuous outcomes. Heterogeneity in the effect estimates of the individual studies will be assessed by visual inspection of the forest plots and consideration of the I<sup>2</sup> statistic. Heterogeneity will be explored as appropriate using sensitivity analyses and pre-specified subgroup analyses. If heterogeneity cannot be explained through sensitivity analysis then a random effects model will be used for meta-analysis, or the data will not be pooled if the random effects model does not adequately address heterogeneity. The confidence in the findings across all available evidence will be evaluated for each outcome using an adaptation of the 'Grading of Recommendations Assessment,</p>

Field	Content
	<p>Development and Evaluation (GRADE) toolbox' developed by the international GRADE working group: <a href="http://www.gradeworkinggroup.org/">http://www.gradeworkinggroup.org/</a></p> <p><b>Minimally important differences:</b></p> <ul style="list-style-type: none"> <li>• All-cause mortality: statistical significance</li> <li>• Serious intervention-related adverse effects: statistical significance</li> <li>• CSF sterilization: statistical significance</li> <li>• Intracranial collections: statistical significance</li> <li>• Validated scales: Published MIDs where available; if not GRADE default MIDs</li> <li>• All other outcomes: GRADE default MIDs</li> </ul>
Analysis of sub-groups	<p>Evidence will be stratified by:</p> <p><b>Stage 1</b> Age:</p> <ul style="list-style-type: none"> <li>• Younger Infants, older infants and children: &gt;28 days to &lt;18* years of age</li> <li>• Adults: ≥18* years of age</li> </ul> <p><b>Stage 2</b> Age:</p> <ul style="list-style-type: none"> <li>• Younger Infants: &gt;28 days to ≤3 months of age</li> <li>• Older infants and children: &gt;3 months to &lt;18* years of age</li> <li>• Adults: ≥18* years of age</li> </ul> <p>*There is variation in clinical practice regarding the treatment of 16 to 18 year olds. Therefore, we will be guided by cut-offs used in the evidence when determining if 16 to 18 year olds should be treated as adults or children.</p> <p>Evidence will be subgrouped by the following only in the event that there is significant heterogeneity in outcomes: Age:</p>

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

Field	Content		
	<ul style="list-style-type: none"> <li>• Young and middle aged adults</li> <li>• Older adults*</li> </ul> <p>*There is variation regarding the age at which adults should be considered older adults. Therefore, we will be guided by cut-offs used in the evidence when determining this threshold.</p> <p>Where evidence is stratified or subgrouped the committee will consider on a case by case basis if separate recommendations should be made for distinct groups. Separate recommendations may be made where there is evidence of a differential effect of interventions in distinct groups. If there is a lack of evidence in one group, the committee will consider, based on their experience, whether it is reasonable to extrapolate and assume the interventions will have similar effects in that group compared with others.</p>		
Type and method of review	<input checked="" type="checkbox"/>	Intervention	
	<input type="checkbox"/>	Diagnostic	
	<input type="checkbox"/>	Prognostic	
	<input type="checkbox"/>	Qualitative	
	<input type="checkbox"/>	Epidemiologic	
	<input type="checkbox"/>	Service Delivery	
	<input type="checkbox"/>	Other (please specify)	
Language	English		
Country	England		
Anticipated or actual start date	12/01/2021		
Anticipated completion date	07/12/2023		
Stage of review at time of this submission	Review stage	Started	Completed
	Preliminary searches	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
	Piloting of the study selection process	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

Field	Content												
	<table border="1"> <tr> <td>Formal screening of search results against eligibility criteria</td> <td><input checked="" type="checkbox"/></td> <td><input checked="" type="checkbox"/></td> </tr> <tr> <td>Data extraction</td> <td><input checked="" type="checkbox"/></td> <td><input checked="" type="checkbox"/></td> </tr> <tr> <td>Risk of bias (quality) assessment</td> <td><input checked="" type="checkbox"/></td> <td><input checked="" type="checkbox"/></td> </tr> <tr> <td>Data analysis</td> <td><input checked="" type="checkbox"/></td> <td><input checked="" type="checkbox"/></td> </tr> </table>	Formal screening of search results against eligibility criteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Data extraction	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Risk of bias (quality) assessment	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	Data analysis	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
Formal screening of search results against eligibility criteria	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>											
Data extraction	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>											
Risk of bias (quality) assessment	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>											
Data analysis	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>											
Named contact	<p>Named contact: National Guideline Alliance</p> <p>Named contact e-mail: meningitis&amp;meningococcal@nice.org.uk</p> <p>Organisational affiliation of the review: National Institute for Health and Care Excellence (NICE) and National Guideline Alliance</p>												
Review team members	National Guideline Alliance												
Funding sources/sponsor	This systematic review is being completed by the National Guideline Alliance which receives funding from NICE.												
Conflicts of interest	All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline.												
Collaborators	Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of <a href="#">Developing NICE guidelines: the manual</a> . Members of the guideline committee are available on the NICE website: <a href="https://www.nice.org.uk/guidance/indevelopment/gid-ng10149">https://www.nice.org.uk/guidance/indevelopment/gid-ng10149</a> .												
Other registration details	None												

Field	Content
Reference/URL for published protocol	<a href="https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021276561">https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021276561</a>
Dissemination plans	NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as: <ul style="list-style-type: none"> <li>• notifying registered stakeholders of publication</li> <li>• publicising the guideline through NICE's newsletter and alerts</li> <li>• issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE.</li> </ul>
Keywords	Bacterial meningitis, antibiotic, anti-bacterial, mortality, impairments
Details of existing review of same topic by same authors	None
Current review status	<input type="checkbox"/> Ongoing <input checked="" type="checkbox"/> Completed but not published <input type="checkbox"/> Completed and published <input type="checkbox"/> Completed, published and being updated <input type="checkbox"/> Discontinued
Additional information	None
Details of final publication	<a href="http://www.nice.org.uk">www.nice.org.uk</a>

1 CDSR: Cochrane Database of Systematic Reviews; CENTRAL: Cochrane Central Register of Controlled Trials; CSF: cerebrospinal fluid; GRADE: Grading of  
2 Recommendations Assessment, Development and Evaluation; MDI: mental development index; MEDLINE: Medical Literature Analysis and Retrieval System Online; MID:  
3 minimally important difference; NICE: National Institute for Health and Care Excellence; PDI: psychomotor development index; PRESS: Peer Review of Electronic Search  
4 Strategies; RCT: randomised controlled trial; RoB: risk of bias; ROBINS-I: risk of bias in non-randomised studies – of interventions; ROBIS: risk of bias in systematic reviews;  
5 SD: standard deviation

## 1 Appendix B Literature search strategies

### 2 Literature search strategies for review question: What antibiotic treatment 3 regimens are effective in treating bacterial meningitis caused by Haemophilus 4 influenzae?

#### 5 6 Clinical Search

7 This was a combined search to cover both this review (E2) and D1, D2, D3, E1, E3, E4, E5,  
8 E6 and F1 on antibiotic regimens for bacterial meningitis (before or in the absence of  
9 identifying causative infecting organism and for specific causative organisms) and  
10 meningococcal disease.

#### 11 12 Database(s): Medline & Embase (Multifile) – OVID interface

13 **Embase Classic+Embase** 1947 to 2022 November 09, **Ovid MEDLINE(R) and Epub**  
14 **Ahead of Print, In-Process & Other Non-Indexed Citations and Daily** 1946 to November  
15 09, 2022

16 Date of last search: 10 November 2022

17 *Multifile database codes: emczd = Embase Classic+Embase; ppez = MEDLINE(R) and Epub Ahead of*  
18 *Print, In-Process & Other Non-Indexed Citations and Daily*

#	Searches
1	Meningitis/ or Meningitis, Bacterial/ or Meningitis, Escherichia Coli/ or Meningitis, Haemophilus/ or Meningitis, Listeria/ or Meningitis, Meningococcal/ or Meningitis, Pneumococcal/ or Meningoencephalitis/
2	1 use ppez
3	meningitis/ or bacterial meningitis/ or haemophilus meningitis/ or hemophilus influenzae meningitis/ or listeria meningitis/ or meningococcal meningitis/ or pneumococcal meningitis/ or meningoencephalitis/
4	3 use emczd
5	((bacter* or infect*) adj3 (meningit* or meninges* or leptomeninges* or subarachnoid space?)).ti,ab.
6	(meningit* adj3 (e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococc* or pneumococc* or gram-negativ* bacill* or gram negativ* bacill* or streptococc* or group B streptococc* or GBS or streptococcus pneumon* or s pneumon* or septic* or sepsis* or bacter?emi?)).ti,ab.
7	((e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococc* or pneumococc* or gram-negativ* bacill* or gram negativ* bacill* or streptococc* or group B streptococc* or GBS or streptococcus pneumon* or s pneumon*) adj3 (septic* or sepsis* or bacter?emi?)).ti,ab.
8	(meningit* or mening?encephalitis?).ti,ab.
9	exp Neisseria meningitidis/ use ppez
10	neisseria meningitidis/ use emczd
11	(Neisseria* mening* or n mening*).ti,ab.
12	or/2,4-11
13	Meningococcal Infections/ use ppez
14	meningococcosis/ or meningococcemia/
15	14 use emczd
16	(meningococc* adj3 (sepsis* or septic* or toxic* or endotoxic* or disease? or infection?)).ti,ab.
17	(meningococcus* or meningococci* or meningococc?emi?).ti,ab.
18	or/13,15-17
19	exp Anti-Bacterial Agents/ or exp Penicillins/ or exp Cephalosporins/ or exp Cefotaxime/ or exp Amoxicillin/ or exp Ampicillin/
20	19 use ppez
21	exp antibiotic agent/ or antibiotic therapy/ or exp penicillin derivative/ or exp cephalosporin derivative/
22	21 use emczd
23	(anti?biotic* or anti?bacterial* or anti?biotherap*).ti,ab.
24	(empiric* adj2 (therap* or treatment?)).ti,ab.
25	(abbocillin or adimicin or alcomicin or alpen or amblosin or amcill or amfipen or aminobenzylpenicillin* or aminoglycosid* or amox?cillin* or amoxil* or ampicillin* or ancef or anticepim or apogen or axepim* or ayercillin or azithrom?cin* or benzo?penicillin* or benzy?penicillin* or bicillin or binotal or biomox or bmy 28142 or bmy?28142 or bristagen or bristamox or carbapenem* or cedax or ceftazidim* or ceftriaxon* or cefepim* or cefixim* or cefizox or cefobid* or cefotan or cefotaxim* or ceftaroline* or ceftin or ceftolozane* or ceftriaxon* or ceftriazon* or cefuroxim* or cezil or cepazin* or cephalosporin* or cephotaxim* or cepuroxim* or cepim?x or chloramphenicol* or ciprofloxacin* or claforan or clamoxyl or clarithromycin* or clindamycin* or collistin* or compocillin or cosmopen or cotrimoxazol* or co trimoxazol* or cristicillin or delaflaxacin* or deripen or dexamethasone or diatracin or doktacillin or duricef or elobact or erythromycin* or flucloxacillin* or fluoroquinolon* or fosfomycin* or gelacillin or gentam?cin* or gent?mycin* or gentamyl* or gentamytrex or gentaplus or gentarad or gentaso* or gentasporin or gentatrim or gent?cin* or gent?cyn* or geocillin* or geomycin* or glycopeptid* or guicitrin* or hexam?cin* or hiconcil or histocillin or ibiamox or imacillin or jenamicin or kefurox or kefzol or klaforan or lendacin or levofloxacin* or linezolid* or longacef or longaceph or lyphocin or macrolide* or mandol or maxcef or maxipim* or mefoxin or megacillin or meropen* or miram?cin* or monocid or moxacin or moxifloxacin* or ofloxacin* or oftagen* or omnipen or optigen* or pefloxacin* or penbritin* or penbrock or penicillin? or penicline or pentids or pentrex or pentrexl or pentrexyl or permapen or pfizerpen or polycillin or polymox or

Meningitis (bacterial) and meningococcal disease: recognition, diagnosis and management: evidence reviews for antibiotics for bacterial meningitis caused by Haemophilus influenzae  
DRAFT (September 2023)

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

#	Searches
	polymyxin* or primafen or principen or quinolon* or refobacin* or ribom?cin* or rifampicin or rifampin* or rocefalin or rocefin or rocephin* or roscillin or rufloxacin* or sagestam* or spectrobid or sulm?cin* or supen or tazobactam* or terram?cin* or tetracycline* or tobramycin* or totacillin or totapen or trimox or u?gencin* or ukapen or ultrabion or vamyisin or vancam* or vancocostacin or vancin or vancom* or vancomycin* or vankom* or velosef or vetramox* or viccillin or voncon* or wycillin or zimox or zinacef or zin?at).mp.
26	or/20,22-25
27	(12 or 18) and 26
28	(controlled clinical trial or pragmatic clinical trial or randomized controlled trial).pt. or drug therapy.fs. or (groups or placebo or randomized or randomly or trial).ab.
29	crossover procedure/ or double blind procedure/ or randomized controlled trial/ or single blind procedure/ or (assign* or allocat* or crossover* or cross over* or ((doubl* or singl*) adj blind*) or factorial* or placebo* or random* or volunteer*).ti.ab.
30	meta-analysis/
31	meta-analysis as topic/
32	systematic review/
33	meta-analysis/
34	(meta analy* or metanaly* or metaanaly*).ti.ab.
35	((systematic or evidence) adj2 (review* or overview*)).ti.ab.
36	((systematic* or evidence*) adj2 (review* or overview*)).ti.ab.
37	(reference list* or bibliograph* or hand search* or manual search* or relevant journals).ab.
38	(search strategy or search criteria or systematic search or study selection or data extraction).ab.
39	(search* adj4 literature).ab.
40	(medline or pubmed or cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or science citation index or bids or cancerlit).ab.
41	cochrane.jw.
42	((pool* or combined) adj2 (data or trials or studies or results)).ab.
43	letter/
44	editorial/
45	news/
46	exp historical article/
47	Anecdotes as Topic/
48	comment/
49	case report/
50	(letter or comment*).ti.
51	43 or 44 or 45 or 46 or 47 or 48 or 49 or 50
52	randomized controlled trial/ or random*.ti.ab.
53	51 not 52
54	animals/ not humans/
55	exp Animals, Laboratory/
56	exp Animal Experimentation/
57	exp Models, Animal/
58	exp Rodentia/
59	(rat or rats or mouse or mice).ti.
60	53 or 54 or 55 or 56 or 57 or 58 or 59
61	letter.pt. or letter/
62	note.pt.
63	editorial.pt.
64	case report/ or case study/
65	(letter or comment*).ti.
66	61 or 62 or 63 or 64 or 65
67	randomized controlled trial/ or random*.ti.ab.
68	66 not 67
69	animal/ not human/
70	nonhuman/
71	exp Animal Experiment/
72	exp Experimental Animal/
73	animal model/
74	exp Rodent/
75	(rat or rats or mouse or mice).ti.
76	68 or 69 or 70 or 71 or 72 or 73 or 74 or 75
77	60 use ppez
78	76 use emczd
79	77 or 78
80	28 use ppez
81	29 use emczd
82	80 or 81
83	(or/30-31,34,36-41) use ppez
84	(or/32-35,37-42) use emczd
85	83 or 84
86	27 not 79
87	limit 86 to English language



DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

#	Searches
88	limit 87 to yr="1980 -Current"
89	limit 88 to (conference abstract or conference paper or conference review or conference proceeding) [Limit not valid in Ovid MEDLINE(R),Ovid MEDLINE(R) Daily Update,Ovid MEDLINE(R) In-Process,Ovid MEDLINE(R) Publisher; records were retained]
90	89 use emczd
91	88 not 90
92	82 or 85
93	91 and 92 [SR/RCT data]
94	91 not 93 [Non-RCT data]

1  
2  
3  
4  
5

**Database(s): Cochrane Library – Wiley interface**  
**Cochrane Database of Systematic Reviews**, Issue 11 of 12, November 2022, **Cochrane Central Register of Controlled Trials**, Issue 11 of 12, November 2022  
 Date of last search: 10 November 2022

#	Searches
#1	MeSH descriptor: [Meningitis] this term only
#2	MeSH descriptor: [Meningitis, Bacterial] this term only
#3	MeSH descriptor: [Meningitis, Escherichia coli] this term only
#4	MeSH descriptor: [Meningitis, Haemophilus] this term only
#5	MeSH descriptor: [Meningitis, Listeria] this term only
#6	MeSH descriptor: [Meningitis, Meningococcal] this term only
#7	MeSH descriptor: [Meningitis, Pneumococcal] this term only
#8	MeSH descriptor: [Meningoencephalitis] this term only
#9	MeSH descriptor: [Neisseria meningitidis] explode all trees
#10	((bacter* or infect*) near/3 (mening* or leptomening* or subarachnoid space*)):ti,ab,kw
#11	((("e coli" or "escherichia coli" or haemophilus or hemophilus or hib or (h next influenz*) or listeria* or pneumococ* or (gram next negativ* next bacill*) or streptococ* or GBS or (s next pneumon*)) near/3 (septic* or sepsis* or bacteraemi* or bacteremi* or infect*)):ti,ab,kw
#12	(meningit* or mening?encephalitis* or (mening* next encephalitis*)):ti,ab,kw
#13	((neisseria* next mening*) or (n next mening*)):ti,ab,kw
#14	MeSH descriptor: [Meningococcal Infections] this term only
#15	meningococ*:ti,ab,kw
#16	{or #1-#15}
#17	MeSH descriptor: [Anti-Bacterial Agents] explode all trees
#18	((antibiotic* or antibacterial* or antibiotherap* or "anti biotic*" or "anti bacterial*" or "anti biotherap*"):ti,ab,kw
#19	((empiric* near/2 (therap* or treatment*)):ti,ab,kw
#20	((abocillin or adimicin or alcomicin or alpen or amblosin or amcill or amfipen or aminobenzylpenicillin* or aminoglycosid* or amox?cillin* or amoxil* or ampicillin* or ancef or anticepim or apogen or axepim* or ayercillin or azithrom?cin* or benzo?penicillin* or benzy?penicillin* or bicillin or binotal or biomox or bmy 28142 or bmy?28142 or bristagen or bristamox or carbapenem* or cedax or ceftazidim* or cefatriaxon* or cefepim* or cefixim* or cefizox or cefobid* or cefotan or cefotaxim* or ceftaroline* or ceftin or ceftolozane* or ceftriaxon* or ceftriaxon* or cefuroxim* or cefzil or cepazin* or cephalosporin* or cephotaxim* or cephiroxim* or cepim?x or chloramphenicol* or ciprofloxacin* or claforan or clamoxyl or clarithromycin* or clindamycin* or colistin* or compocillin or cosmopen or cotrimoxazol* or co trimoxazol* or crysticillin or delafloxacin* or deripen or dexamethasone or diatracin or doktacillin or duricef or elobact or erythromycin* or flucloxacillin* or fluoroquinolon* or fosfomicin* or gelaclillin or gentam?cin* or gent?mycin* or gentamyl* or gentamytrex or gentaplus or gentarad or gentaso* or gentasporin or gentatrim or gent?cin* or gent?cyn* or geocillin* or geomycin* or glycopeptid* or guicitrin* or hexam?cin* or hiconcil or histocillin or ibiamox or imacillin or jenamicin or kefurox or kefzol or klaforan or lendacin or levofloxacin* or linezolid* or longacef or longaceph or lyphocin or macrolide* or mandol or maxcef or maxipim* or mefoxin or megacillin or meropen* or miram?cin* or monocid or moxacin or moxifloxacin* or ofloxacin* or oftagen* or omnipen or optigen* or pefloxacin* or penbritin* or penbrock or penicillin? or peniciline or pentids or pentrex or pentrexl or pentrexyl or permapen or pfizerpen or polyicillin or polymox or polymyxin* or primafen or principen or quinolon* or refobacin* or ribom?cin* or rifampicin or rifampin* or rocefalin or rocefin or rocephin* or roscillin or rufloxacin* or sagestam* or spectrobid or sulm?cin* or supen or tazobactam* or terram?cin* or tetracycline* or tobramycin* or totacillin or totapen or trimox or u?gencin* or ukapen or ultrabion or vamsyn or vancam* or vancocostacin or vancin or vancom* or vancomycin* or vankom* or velosef or vetramox* or viccillin or voncon* or wycillin or zimox or zinacef or zin?at):ti,ab,kw
#21	{or #17-#20}
#22	#16 and #21
#23	"conference":pt or (clinicaltrials or trialsearch):so
#24	#22 not #23

6  
7  
8  
9

**Database(s): Database of Abstracts of Reviews of Effects (DARE); HTA Database – CRD interface**  
 Date of last search: 12 February 2021

#	Searches
1	MeSH DESCRIPTOR meningitis IN DARE,HTA
2	MeSH DESCRIPTOR meningitis, bacterial IN DARE,HTA
3	MeSH DESCRIPTOR Meningitis, Escherichia coli IN DARE,HTA
4	MeSH DESCRIPTOR Meningitis, Haemophilus IN DARE,HTA

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

#	Searches
5	MeSH DESCRIPTOR Meningitis, Listeria IN DARE,HTA
6	MeSH DESCRIPTOR Meningitis, Meningococcal IN DARE,HTA
7	MeSH DESCRIPTOR Meningitis, Pneumococcal IN DARE,HTA
8	MeSH DESCRIPTOR Meningoencephalitis IN DARE,HTA
9	MeSH DESCRIPTOR Meningococcal infections IN DARE,HTA
10	((((bacter* or infect*) NEAR3 (meningit* or meninges* or leptomeninges* or "subarachnoid space*")))) IN DARE, HTA
11	(meningit*) IN DARE, HTA
12	(((meningoencephalitis* or meningoencephalitis*))) IN DARE, HTA
13	(((meningococc* NEAR3 (sepsis* or septic* or toxic* or endotoxic* or disease or diseases or infection or infections)))) IN DARE, HTA
14	(((meningococcus* or meningococci* or meningococcaemia* or meningococemia*))) IN DARE, HTA
15	((Neisseria* NEAR1 mening*)) IN DARE, HTA
16	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15
17	MeSH DESCRIPTOR Anti-Bacterial Agents EXPLODE ALL TREES IN DARE,HTA
18	MeSH DESCRIPTOR Penicillins EXPLODE ALL TREES IN DARE,HTA
19	MeSH DESCRIPTOR Cephalosporins EXPLODE ALL TREES IN DARE,HTA
20	MeSH DESCRIPTOR Cefotaxime EXPLODE ALL TREES IN DARE,HTA
21	MeSH DESCRIPTOR Amoxicillin EXPLODE ALL TREES IN DARE,HTA
22	MeSH DESCRIPTOR Ampicillin EXPLODE ALL TREES IN DARE,HTA
23	(((antibiotic* or antibacterial* or antibiotherap* or anti-biotic* or anti-bacterial* or anti-biotherap* or "anti biotic*" or "anti bacterial*" or "anti biotherap*")) IN DARE, HTA
24	(((empiric* NEAR2 (therap* or treatment*))) IN DARE, HTA
25	(((abbocillin or adimicin or alcomicin or alpen or amblosin or amcill or amfipen or aminobenzylpenicillin or amox?cillin or amoxil* or ampicillin or ancef or anticepim or apogen or axepim* or ayercillin or benzo?penicillin* or benzy?penicillin* or bicillin or binotal or biomox or bmy 28142 or bmy-28142 or bmy28142 or bristagen or bristamox or cedax or ceftriaxon* or cefepim* or cefixim* or cefizox or cefobid* or cefotan or cefotaxim* or ceftin or ceftriaxon* or ceftriaxon* or cefuroxim* or cefzil or cepazin* or cephotaxim* or cepuroxim* or cepim*x or chloramphenicol or claforan or clamoxyl or compocillin or cosmopen or cotrimoxazol* or co trimoxazol* or co-trimoxazol or crysticillin or deripen or dexamethasone or diatracin or doktacillin or duricef or elobact or gelacillin or gentam?cin* or gent?mycin* or gentamyl* or gentamytrex or gentaplus or gentarad or gentaso* or gentasporin or gentatrim or gent?cin* or gent?cyn* or geocillin* or geomycin* or guicitrin* or hexam?cin* or hiconcil or histocillin or ibiamox or imacillin or jenamicin or kefurox or kefzol or klaforan or lendacin or longacef or longaceph or lyphocin or mandol or maxcef or maxipim* or mefoxin or megacillin or meropen* or miram?cin* or monocid or moxacin or oftagen* or omnipen or optigen* or penbritin* or penbrock or penicillin? or penicline or pentids or pentrex or pentrexl or pentrexyl or permepen or pfizerpen or polycillin or polymox or primafen or principen or refobacin* or ribom?cin* or rifampicin or rocefallin or rocefin or rocephin* or roscillin or sagestam* or spectrobid or sulm?cin* or supen or terram?cin* or totacillin or totapen or trimox or u?gencin* or ukapen or ultrabion or vamysin or vancam* or vancocostacin or vancin or vancom* or vancomycin or vankom* or velosef or vetramox* or vicillin or voncon* or wycillin or zimox or zinacef or zin?at)) IN DARE, HTA
26	#17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25
27	#16 AND #26

1  
2

1

2 **Economic Search**

3 One global search was conducted for economic evidence across the guideline.

4

5 **Database(s): NHS Economic Evaluation Database (NHS EED), HTA Database – CRD**  
6 **interface**

7 Date of last search: 11 March 2021

#	Searches
1	MeSH DESCRIPTOR meningitis IN NHSEED,HTA
2	MeSH DESCRIPTOR Meningitis, Bacterial IN NHSEED,HTA
3	MeSH DESCRIPTOR Meningitis, Escherichia coli IN NHSEED,HTA
4	MeSH DESCRIPTOR Meningitis, Haemophilus EXPLODE ALL TREES IN NHSEED,HTA
5	MeSH DESCRIPTOR Meningitis, Listeria IN NHSEED,HTA
6	MeSH DESCRIPTOR Meningitis, Meningococcal IN NHSEED,HTA
7	MeSH DESCRIPTOR Meningitis, Pneumococcal IN NHSEED,HTA
8	MeSH DESCRIPTOR Meningoencephalitis IN NHSEED,HTA
9	((bacter* or infect*) NEAR3 (meningit* or meninges* or leptomeninges* or subarachnoid space*)) IN NHSEED, HTA
10	((meningit* NEAR3 (e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococ* or pneumococ* or gram-negativ* bacill* or gram negativ* bacill* or streptococ* or group B streptococ* or GBS or streptococcus pneumon* or s pneumon* or septic* or sepsis* or bacter?emi?)) IN NHSEED, HTA
11	((e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococ* or pneumococ* or gram-negativ* bacill* or gram negativ* bacill* or streptococ* or group B streptococ* or GBS or streptococcus pneumon* or s pneumon*) NEAR3 (septic* or sepsis* or bacter?emi?)) IN NHSEED, HTA
12	((meningencephalitis* or meningoencephalitis* or meningit*) IN NHSEED, HTA
13	MeSH DESCRIPTOR Meningococcal Infections IN NHSEED,HTA
14	MeSH DESCRIPTOR Neisseria meningitidis EXPLODE ALL TREES IN NHSEED,HTA
15	((meningococ* NEAR3 (sepsis* or septic* or toxic* or endotoxic* or disease* or infection*)) IN NHSEED, HTA
16	((meningococcus* or meningococci* or meningococcaemia* or meningococcemia*) IN NHSEED, HTA
17	((Neisseria* NEXT mening*) IN NHSEED, HTA
18	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17

8

9 **Database(s): Medline & Embase (Multifile) – OVID interface**10 **Embase Classic+Embase 1947 to 2022 November 09, Ovid MEDLINE(R) and Epub**  
11 **Ahead of Print, In-Process & Other Non-Indexed Citations and Daily 1946 to November**  
12 **09, 2022**

13 Date of last search: 10 November 2022

14 *Multifile database codes: emczd = Embase Classic+Embase; ppez= MEDLINE(R) and Epub Ahead of*  
15 *Print, In-Process & Other Non-Indexed Citations and Daily*

#	Searches
1	Meningitis/ or Meningitis, Bacterial/ or Meningitis, Escherichia Coli/ or Meningitis, Haemophilus/ or Meningitis, Listeria/ or Meningitis, Meningococcal/ or Meningitis, Pneumococcal/ or Meningoencephalitis/
2	1 use ppez
3	meningitis/ or bacterial meningitis/ or haemophilus meningitis/ or listeria meningitis/ or pneumococcal meningitis/ or meningoencephalitis/
4	3 use emczd
5	((bacter* or infect*) adj3 (meningit* or meninges* or leptomeninges* or subarachnoid space?)).ti.ab.
6	(meningit* adj3 (e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococ* or pneumococ* or gram-negativ* bacill* or gram negativ* bacill* or streptococ* or group B streptococ* or GBS or streptococcus pneumon* or s pneumon* or septic* or sepsis* or bacter?emi?)).ti.ab.
7	((e coli or escherichia coli or h?emophilus or hib or h?emophilus influenz* or h influenz* or listeria* or meningococ* or pneumococ* or gram-negativ* bacill* or gram negativ* bacill* or streptococ* or group B streptococ* or GBS or streptococcus pneumon* or s pneumon*) adj3 (septic* or sepsis* or bacter?emi?)).ti.ab.
8	(mening?encephalitis* or meningit*).ti.ab.
9	or/2,4-8
10	Meningococcal Infections/ or exp Neisseria meningitidis/
11	10 use ppez
12	Meningococcosis/ or Meningococcemia/ or Neisseria Meningitidis/
13	12 use emczd
14	(meningococ* adj3 (sepsis* or septic* or toxic* or endotoxic* or disease? or infection?)).ti.ab.
15	(meningococcus* or meningococci* or meningococ?emi?).ti.ab.
16	(Neisseria* mening* or n mening*).ti.ab.
17	or/11,13-16
18	Economics/ use ppez
19	Value of life/ use ppez

Meningitis (bacterial) and meningococcal disease: recognition, diagnosis and management: evidence reviews for antibiotics for bacterial meningitis caused by Haemophilus influenzae  
DRAFT (September 2023)

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

#	Searches
20	exp "Costs and Cost Analysis"/ use ppez
21	exp Economics, Hospital/ use ppez
22	exp Economics, Medical/ use ppez
23	Economics, Nursing/ use ppez
24	Economics, Pharmaceutical/ use ppez
25	exp "Fees and Charges"/ use ppez
26	exp Budgets/ use ppez
27	health economics/ use emczd
28	exp economic evaluation/ use emczd
29	exp health care cost/ use emczd
30	exp fee/ use emczd
31	budget/ use emczd
32	funding/ use emczd
33	budget*.ti,ab.
34	cost*.ti.
35	(economic* or pharmaco?economic*).ti.
36	(price* or pricing*).ti,ab.
37	(cost* adj2 (effective* or utilit* or benefit* or minimi* or unit* or estimat* or variable*).ab.
38	(financ* or fee or fees).ti,ab.
39	(value adj2 (money or monetary)).ti,ab.
40	or/18-39
41	Quality-Adjusted Life Years/ use ppez
42	Sickness Impact Profile/
43	quality adjusted life year/ use emczd
44	"quality of life index"/ use emczd
45	(quality adjusted or quality adjusted life year*).tw.
46	(qaly* or qal or qald* or qale* or qtime* or qwb* or daly).tw.
47	(illness state* or health state*).tw.
48	(hui or hui2 or hui3).tw.
49	(multiattribute* or multi attribute*).tw.
50	(utilit* adj3 (score*1 or valu* or health* or cost* or measur* or disease* or mean or gain or gains or index)).tw.
51	utilities.tw.
52	(eq-5d* or eq5d* or eq-5* or eq5* or euroqual* or euro qual* or euroqual 5d* or euro qual 5d* or euro qol* or euroqol* or euro quol* or euroquol* or euro quol5d* or euroquol5d* or eur qol* or eurqol* or eur qol5d* or eurqol5d* or eur?qul* or eur?qul5d* or euro* quality of life or european qol).tw.
53	(euro* adj3 (5 d* or 5d* or 5 dimension* or 5dimension* or 5 domain* or 5domain)).tw.
54	(sf36 or sf 36 or sf thirty six or sf thirtysix).tw.
55	(time trade off*1 or time tradeoff*1 or tto or timetradeoff*1).tw.
56	Quality of Life/ and ((quality of life or qol) adj (score*1 or measure*1)).tw.
57	Quality of Life/ and ec.fs.
58	Quality of Life/ and (health adj3 status).tw.
59	(quality of life or qol).tw. and Cost-Benefit Analysis/ use ppez
60	(quality of life or qol).tw. and cost benefit analysis/ use emczd
61	((qol or hrqol or quality of life).tw. or *quality of life/) and ((qol or hrqol* or quality of life) adj2 (increas* or decreas* or improv* or declin* or reduc* or high* or low* or effect or effects or worse or score or scores or change*1 or impact*1 or impacted or deteriorat*).ab.
62	Cost-Benefit Analysis/ use ppez and cost-effectiveness ratio*.tw. and (cost-effectiveness ratio* and (perspective* or life expectanc*).tw.
63	cost benefit analysis/ use emczd and cost-effectiveness ratio*.tw. and (cost-effectiveness ratio* and (perspective* or life expectanc*).tw.
64	*quality of life/ and (quality of life or qol).ti.
65	quality of life/ and ((quality of life or qol) adj3 (improv* or chang*).tw.
66	quality of life/ and health-related quality of life.tw.
67	Models, Economic/ use ppez
68	economic model/ use emczd
69	care-related quality of life.tw,kw.
70	((capability\$ or capability-based\$) adj (measure\$ or index or instrument\$)).tw,kw.
71	social care outcome\$.tw,kw.
72	(social care and (utility or utilities)).tw,kw.
73	or/41-72
74	(9 or 17) and 40
75	(9 or 17) and 73
76	letter/
77	editorial/
78	news/
79	exp historical article/
80	Anecdotes as Topic/
81	comment/
82	case report/
83	(letter or comment*).ti.
84	76 or 77 or 78 or 79 or 80 or 81 or 82 or 83

## DRAFT FOR CONSULTATION


### Antibiotics for bacterial meningitis caused by Haemophilus influenzae

---

#	Searches
85	randomized controlled trial/ or random*.ti,ab.
86	84 not 85
87	animals/ not humans/
88	exp Animals, Laboratory/
89	exp Animal Experimentation/
90	exp Models, Animal/
91	exp Rodentia/
92	(rat or rats or mouse or mice).ti.
93	86 or 87 or 88 or 89 or 90 or 91 or 92
94	letter.pt. or letter/
95	note.pt.
96	editorial.pt.
97	case report/ or case study/
98	(letter or comment*).ti.
99	94 or 95 or 96 or 97 or 98
100	randomized controlled trial/ or random*.ti,ab.
101	99 not 100
102	animal/ not human/
103	nonhuman/
104	exp Animal Experiment/
105	exp Experimental Animal/
106	animal model/
107	exp Rodent/
108	(rat or rats or mouse or mice).ti.
109	101 or 102 or 103 or 104 or 105 or 106 or 107 or 108
110	93 use ppez
111	109 use emczd
112	110 or 111
113	74 not 112
114	limit 113 to English language
115	75 not 112
116	limit 115 to English language
117	114 or 116

1  
2

- 1 **Appendix C Effectiveness evidence study selection**
- 2 **Study selection for: What antibiotic treatment regimens are effective in treating**
- 3 **bacterial meningitis caused by Haemophilus influenzae?**
- 4 **Figure 1: Study selection flow chart**



- 5
- 6
- 7
- 8

## 1 Appendix D Evidence tables

### 2 Evidence tables for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused 3 by Haemophilus influenzae?

#### 4 Table 4: Evidence tables – effectiveness evidence

##### 5 Lapointe, 1988

**Bibliographic Reference** Lapointe, J. R.; Chicoine, L.; Cefotaxime versus chloramphenicol for ampicillin-resistant Haemophilus influenzae meningitis. A retrospective study of 62 cases; Drugs; 1988; vol. 35suppl2; 199-202

6

##### 7 Study details

<b>Country/ies where study was carried out</b>	Canada
<b>Study type</b>	Retrospective cohort study
<b>Study dates</b>	Not reported
<b>Inclusion criteria</b>	Babies and children with bacterial meningitis caused by $\beta$ -lactamase-producing and ampicillin-resistant Haemophilus influenzae
<b>Exclusion criteria</b>	Not reported
<b>Patient characteristics</b>	N=62 Age (months in mean): 16 Etiology: Haemophilus influenzae: 62 (100%)
<b>Intervention(s)/control</b>	Cefotaxime: Ampicillin 200 mg/kg/day IV plus cefotaxime 200 mg/kg/day IV in the first 48 hours followed by cefotaxime 200 mg/kg/day IV alone. Total duration of treatment was 10 days.

	Chloramphenicol: Ampicillin 200-400 mg/kg/day IV plus chloramphenicol 100 mg/kg/day IV in the first 48 hours followed by chloramphenicol 100 mg/kg/day IV alone. Total duration of treatment was 10 days
<b>Duration of follow-up</b>	During hospitalisation (both groups), 3.4 months (cefotaxime group), and 12.2 months (chloramphenicol group)
<b>Sources of funding</b>	Industry funded
<b>Sample size</b>	N=62

1 IV: intravenous

### 3 Outcomes

#### 4 Cefotaxime versus chloramphenicol: All-cause mortality and any long-term neurological impairment

Outcome	Cefotaxime, N = 36	Chloramphenicol, N = 26
<b>All-cause mortality (during hospitalisation)</b>	n = 0	n = 0
No of events		
<b>Any long-term neurological impairment (neurological sequelae including motor deficit, hearing loss and minor anomaly of the auditory potentials; up to 12.2 months after discharge)</b>	n = 4	n = 11
No of events		

### 6 Critical appraisal - ROBINS-I

Section	Question	Answer
1. Bias due to confounding	Risk of bias judgement for confounding	Critical <i>(Data not adjusted for confounding factors. The demographic, laboratory and clinical data of the study groups were comparable except for severity of infection at presentation (89% of</i>



Section	Question	Answer
		<i>cefotaxime group were in poor condition compared with 42% of chloramphenicol group) and cerebrospinal fluid leucocytes count, which was significantly lower in chloramphenicol group. Rate of comorbidity that could have caused confounding was not reported.)</i>
2. Bias in selection of participants into the study	Risk of bias judgement for selection of participants into the study	Low <i>(All eligible participants were included and followed up in the trial)</i>
3. Bias in classification of interventions	Risk of bias judgement for classification of interventions	Moderate <i>(Interventions clearly defined but assignment of intervention status was determined retrospectively)</i>
4. Bias due to deviations from intended interventions	Risk of bias judgement for deviations from intended interventions	Low <i>(No deviations from intended interventions)</i>
5. Bias due to missing data	Risk of bias judgement for missing data	Low <i>(No missing data)</i>
6. Bias in measurement of outcomes	Risk of bias judgement for measurement of outcomes	Serious <i>(Low risk for all-cause mortality as outcome measurement would not be influenced by knowledge of the intervention received, and serious risk for any long-term neurological impairment as it was not clearly described how this outcome was measured and this outcome could be somewhat subjective and therefore influenced by knowledge of assigned intervention.)</i>
7. Bias in selection of the reported result	Risk of bias judgement for selection of the reported result	Moderate <i>(No indication of selection of the reported analysis from among multiple analyses)</i>
Overall bias	Risk of bias judgement	Critical
Overall bias	Risk of bias variation	None

Section	Question	Answer
	across outcomes	
Overall bias	Directness	Directly applicable (All-cause mortality is directly applicable. However, any long-term neurological impairment is indirect outcome as it is a composite outcome including hearing loss and minor anomaly of the auditory potentials.)

1 ROBINS-I: risk of bias in non-randomised studies – of interventions

2

### 3 Molyneux, 2011

**Bibliographic Reference** Molyneux, Elizabeth; Nizami, Shaikh Qamaruddin; Saha, Samir; Huu, Khanh Truong; Azam, Matloob; Bhutta, Zulfiqar Ahmad; Zaki, Ramadan; Weber, Martin Willi; Qazi, Shamim Ahmad; Group, C. S. F. Study; 5 versus 10 days of treatment with ceftriaxone for bacterial meningitis in children: a double-blind randomised equivalence study; Lancet (London, England); 2011; vol. 377 (no. 9780); 1837-45

4

### 5 Study details

<b>Country/ies where study was carried out</b>	Bangladesh, Egypt, Malawi, Pakistan and Vietnam
<b>Study type</b>	Randomised controlled trial (RCT)
<b>Study dates</b>	September 2001 – December 2006
<b>Inclusion criteria</b>	Babies and children aged 2 months to 12 years with bacterial meningitis caused by Haemophilus influenzae, Streptococcus pneumoniae, or Neisseria meningitidis who were alive on day 5 after the start of treatment and clinically stable or improving. Meningitis caused by the study organisms was defined as positive CSF culture or latex agglutination, or positive blood culture plus >10 white blood cells per mL of CSF, or >100 white blood cells per mL of CSF with >50% granulocytes plus CSF glucose <1.66 mmol/L or <50% of blood glucose, or >100 white blood cells per mL of CSF with 75% polymorphonuclear leukocytes.
<b>Exclusion criteria</b>	Exclusion criteria at enrolment: Age ≤2 months, body weight ≤3 kg, pre-existing neurosurgical conditions, cerebral palsy,

	<p>seizure disorders, degenerative neurological conditions, skull fractures, active viral infections, known immunodeficiency, symptomatic AIDS, known hypersensitivity reaction to cephalosporins, cyanotic congenital heart disease, inaccessibility for follow-up, children treated with any parenteral antibiotics for 24 h before admission, and children randomly assigned on more than one occasion in this study</p> <p>Exclusion criteria for random assignment at day 5: Criteria listed above, ceftriaxone-resistant bacteria, serious adverse reactions to the drug given, presence of or growth of bacteria from cerebrospinal fluid taken 48 to 72 h after admission, pyogenic brain abscess, intracranial suppurative thrombophlebitis, subdural empyema, presence of another infection during admission that needed another injectable antibiotic, and meningitis caused by any bacteria other than <i>Haemophilus influenzae</i>, <i>Streptococcus pneumoniae</i>, or <i>Neisseria meningitidis</i></p>
<b>Patient characteristics</b>	<p><i>Haemophilus influenzae</i> N=266 (study also included bacterial meningitis with other causes (N=738) but these were not of interest for the current review); 5-day ceftriaxone therapy: 134; 10-day ceftriaxone therapy: 132</p> <p>Age<sup>1</sup> (months in mean; SD in parentheses): 38 (42)</p> <p>Sex<sup>1</sup>: male: 565 (56%); female: 439 (44%)</p> <p>Etiology: <i>Haemophilus influenzae</i>: 266 (27%); <i>Streptococcus pneumoniae</i>: 335 (33%); <i>Neisseria meningitidis</i>: 73 (7%); unknown: 330 (33%)</p> <p>Children infected with HIV<sup>2</sup>: 117 (12%)</p> <p><sup>1</sup>Reported for whole study, not based on causative organism</p> <p><sup>2</sup>Although HIV is listed as an exclusion criterion in the protocol, the evidence was not considered indirect as those with HIV accounted for &lt;25% of the population</p>
<b>Intervention(s)/control</b>	<p>5-day ceftriaxone therapy: Intravenous ceftriaxone 80-100 mg/kg once daily for 5 days followed by placebo for 5 days</p> <p>10-day ceftriaxone therapy: Intravenous ceftriaxone 80-100 mg/kg once daily for 10 days</p>
<b>Duration of follow-up</b>	Daily during hospitalisation, at discharge on day 10, and on day 40 and day 190 after enrolment
<b>Sources of funding</b>	Industry funded
<b>Sample size</b>	N=1004 <sup>1</sup>

<sup>1</sup>Reported for whole study, not based on causative organism

**Other information**

437 children (5-day ceftriaxone therapy: 209; 10-day ceftriaxone therapy: 228) received dexamethasone therapy<sup>1</sup>

<sup>1</sup>Reported for whole study, not based on causative organism

<sup>1</sup> *AIDS: acquired immunodeficiency syndrome; CSF: cerebrospinal fluid; HIV: human immunodeficiency virus; SD: RCT: randomised controlled trial; standard deviation*

1  
2

**3 Outcomes**

**4 5-day ceftriaxone therapy versus 10-day ceftriaxone therapy: All-cause mortality, any long-term neurological impairment, developmental delay, hearing impairment, serious intervention-related adverse effects and CSF sterilisation**

5

Outcome	5-day ceftriaxone therapy, N = 496	10-day ceftriaxone therapy, N = 508
<b>All-cause mortality (up to 6 months after discharge)</b>	4/134	5/132
Custom value		
<b>Any long-term neurological impairment (neurological sequelae including motor deficit, cranial nerve palsy and afebrile seizures; up to 6 months after discharge)</b>	21/496	30/508
Custom value		
<b>Any long-term neurological impairment (visual loss; up to 6 months after discharge)</b>	4/496	10/508
Custom value		
<b>Developmental delay (assessed using the age and stages questionnaire; up to 6 months after discharge)</b>	25/496	33/508
Custom value		
<b>Hearing impairment (up to 6 months after discharge)</b>	105/496	106/508
Custom value		

Outcome	5-day ceftriaxone therapy, N = 496	10-day ceftriaxone therapy, N = 508
<b>Serious intervention-related adverse effects (adverse events to the study drug; up to 6 months after discharge)</b>	0/496	0/508
Custom value		
<b>CSF sterilisation (positive cerebrospinal fluid or blood cultures on days 6-40)</b>	6/134	5/132
Custom value		

1 CSF: cerebrospinal fluid

2

### 3 Critical appraisal - Cochrane RoB2

Section	Question	Answer
Domain 1: Bias arising from the randomisation process	Risk of bias judgement for the randomisation process	Low <i>(Computer-generated randomisation, process of allocation controlled by an external unit, sealed opaque envelopes used for allocation concealment. No substantial differences between groups at baseline.)</i>
Domain 2a: Risk of bias due to deviations from the intended interventions (effect of assignment to intervention)	Risk of bias for deviations from the intended interventions (effect of assignment to intervention)	Low <i>(Participants and intervention staff were not aware of intervention. Appropriate analysis was used.)</i>
Domain 3. Bias due to missing outcome data	Risk-of-bias judgement for missing outcome data	Some concerns <i>(Loss to follow up greater in 10-day ceftriaxone therapy compared with other arm for all outcomes (5.3% vs 4% on day 40; 11% vs 6.5% at 6-month follow-up) and could be related to participants' health status or death.)</i>
Domain 4. Bias in measurement of the outcome	Risk-of-bias judgement for measurement of the outcome	Low/High <i>(Methods of measuring the outcomes were appropriate, and no difference in measurement of the outcomes between intervention groups. No information if</i>

Section	Question	Answer
		<i>outcome assessors were blinded to intervention status. Low risk for all-cause mortality, hearing impairment, CSF sterilisation and serious intervention-related adverse effects outcomes as outcome measurement would not be influenced by knowledge of assigned intervention, and high risk for neurological impairment, including visual loss, and developmental delay outcomes as they are somewhat subjective and may be influenced by knowledge of assigned intervention.)</i>
Domain 5. Bias in selection of the reported result	Risk-of-bias judgement for selection of the reported result	Low <i>(There is clear evidence that all eligible reported results for the outcome correspond to all intended outcome measurements and analyses in the protocol.)</i>
Overall bias and Directness	Risk of bias judgement	Some concerns for all-cause mortality, hearing impairment, CSF sterilisation and serious intervention-related adverse effects.  High risk for any long-term neurological impairment and developmental delay.
Overall bias and Directness	Overall Directness	Indirectly applicable <i>(All outcomes other than all-cause mortality and CSF sterilisation, data is presented for the whole study rather than the subgroup of interest. Therefore, population is very seriously indirect (&lt;50% had bacterial meningitis caused by organism of interest). Also, developmental delay is an indirect outcome (because no indication of severity).)</i>
Overall bias and Directness	Risk of bias variation across outcomes	Yes, see "Risk of bias judgement" cell above.

1 CSF: cerebrospinal fluid; RoB: risk of bias

2

1 **Appendix E Forest plots**

2 **Forest plots for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused by**  
3 ***Haemophilus influenzae*?**

4 No meta-analysis was conducted for this review question and so there are no forest plots.

5

1 **Appendix F GRADE tables**2 **GRADE tables for review question: What antibiotic treatment regimens are effective in treating bacterial meningitis caused**  
3 **by Haemophilus influenzae?**4 **Table 5: Evidence profile for comparison: cefotaxime versus chloramphenicol**

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Cefotaxime	Chloramphenicol	Relative (95% CI)	Absolute		
<b>All-cause mortality (unadjusted analyses): babies and children</b>												
1 (Lapointe 1988)	observational studies	very serious <sup>1</sup>	no serious inconsistency	no serious indirectness	very serious <sup>2</sup>	none	0/36 (0%)	0/26 (0%)	RD 0 (-0.06 to 0.06)	0 fewer per 1000 (from 60 fewer to 60 more) <sup>3</sup>	VERY LOW	CRITICAL
<b>Any long-term neurological impairment (neurological sequelae, including motor deficit, hearing loss and minor anomaly of the auditory potentials; unadjusted analyses): babies and children (follow-up 0-12.2 months)</b>												
1 (Lapointe 1988)	observational studies	very serious <sup>1</sup>	no serious inconsistency	very serious <sup>4</sup>	no serious imprecision	none	4/36 (11.1%)	11/26 (42.3%)	RR 0.26 (0.09 to 0.73)	313 fewer per 1000 (from 114 fewer to 385 fewer)	VERY LOW	CRITICAL

5 *CI: confidence interval; RD: risk difference; RR: risk ratio*6 <sup>1</sup> *Very serious risk of bias in the evidence contributing to the outcomes as per ROBINS-I*7 <sup>2</sup> *Sample size <200*8 <sup>3</sup> *Absolute effect calculated based on risk difference*9 <sup>4</sup> *Outcome is indirect as it is a composite outcome including hearing loss and minor anomaly of the auditory potentials and measured up to 12.2 months*

10

11 **Table 6: Evidence profile for comparison: 5-day ceftriaxone therapy versus 10-day ceftriaxone therapy**

Quality assessment							No of patients		Effect		Quality	Importance
No of	Design	Risk of	Inconsistency	Indirectness	Imprecision	Other	5-day	10-day	Relative	Absolute		



DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

studies		bias				considerations	ceftriaxone therapy	ceftriaxone therapy	(95% CI)			
<b>All-cause mortality: babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	serious <sup>1</sup>	no serious inconsistency	no serious indirectness	very serious <sup>2</sup>	none	4/134 (3%)	5/132 (3.8%)	RR 0.79 (0.22 to 2.87)	8 fewer per 1000 (from 30 fewer to 71 more)	VERY LOW	CRITICAL
<b>Any long-term neurological impairment (neurological sequelae including motor deficit, cranial nerve palsy and afebrile seizures): babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	very serious <sup>3</sup>	no serious inconsistency	very serious <sup>4</sup>	serious <sup>5</sup>	none	21/496 (4.2%)	30/508 (5.9%)	RR 0.72 (0.42 to 1.23)	17 fewer per 1000 (from 34 fewer to 14 more)	VERY LOW	CRITICAL
<b>Any long-term neurological impairment (visual loss): babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	very serious <sup>3</sup>	no serious inconsistency	very serious <sup>6</sup>	very serious <sup>7</sup>	none	4/496 (0.81%)	10/508 (2%)	RR 0.41 (0.13 to 1.3)	12 fewer per 1000 (from 17 fewer to 6 more)	VERY LOW	CRITICAL
<b>Developmental delay (assessed using the age and stages questionnaire): babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	very serious <sup>3</sup>	no serious inconsistency	very serious <sup>8</sup>	very serious <sup>7</sup>	none	25/496 (5%)	33/508 (6.5%)	RR 0.78 (0.47 to 1.29)	14 fewer per 1000 (from 34 fewer to 19 more)	VERY LOW	CRITICAL
<b>Hearing impairment: babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	serious <sup>1</sup>	no serious inconsistency	very serious <sup>6</sup>	very serious <sup>7</sup>	none	105/496 (21.2%)	106/508 (20.9%)	RR 1.01 (0.8 to 1.29)	2 more per 1000 (from 42 fewer to 61 more)	VERY LOW	IMPORTANT
<b>Serious intervention-related adverse effects (adverse events to the study drug): babies and children (follow-up 0-6 months)</b>												
1 (Molyneux 2011)	randomised trials	serious <sup>1</sup>	no serious inconsistency	very serious <sup>6</sup>	no serious imprecision	none	0/496 (0%)	0/508 (0%)	RD 0.00 (-0.0039 to 0.0039)	0 fewer per 1000 (from 3.9 fewer to 3.9 more) <sup>9</sup>	VERY LOW	IMPORTANT
<b>CSF sterilisation (positive cerebrospinal fluid or blood cultures on days 6-40): babies and children</b>												
1 (Molyneux 2011)	randomised trials	serious <sup>1</sup>	no serious inconsistency	serious <sup>10</sup>	very serious <sup>2</sup>	none	6/134 (4.5%)	5/132 (3.8%)	RR 1.18 (0.37 to 3.78)	7 more per 1000 (from 24 fewer to 105 more)	VERY LOW	IMPORTANT

1 CI: confidence interval; CSF: cerebrospinal fluid; RD: risk difference; RR: risk ratio

2 <sup>1</sup> Serious risk of bias in the evidence contributing to the outcomes as per RoB 2

3 <sup>2</sup> <150 events

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

---

- 1 <sup>3</sup> Very serious risk of bias in the evidence contributing to the outcomes as per RoB 2
- 2 <sup>4</sup> Outcome is indirect as it is a composite outcome including afebrile seizures, and population is indirect due to 74% of population with meningitis caused by organisms other than
- 3 *H. influenzae*
- 4 <sup>5</sup> 95% CI crosses 1 MID
- 5 <sup>6</sup> Population is indirect due to 74% of population with meningitis caused by organisms other than *H. influenzae*
- 6 <sup>7</sup> 95% CI crosses 2 MIDs
- 7 <sup>8</sup> Outcome is indirect as it is a composite outcome that could include mild or moderate developmental delay, and population is indirect due to 74% of population with meningitis
- 8 caused by organisms other than *H. influenzae*
- 9 <sup>9</sup> Absolute effect calculated based on risk difference
- 10 <sup>10</sup> Outcome is indirect as it is a composite outcome including positive blood culture

11

12


## 1 Appendix G Economic evidence study selection

### 2 Study selection for: What antibiotic treatment regimens are effective in treating 3 bacterial meningitis caused by Haemophilus influenzae?

4 A global economic search was undertaken for the whole guideline, but no economic  
5 evidence was identified which was applicable to this review question (see Figure 2).

6 **Figure 2: Study selection flow chart**

7



8

9

10

11

12

13

14

1 **Appendix H Economic evidence tables**

2 **Economic evidence tables for review question: What antibiotic treatment**  
3 **regimens are effective in treating bacterial meningitis caused by Haemophilus**  
4 **influenzae?**

5 No evidence was identified which was applicable to this review question

6

1 **Appendix I Economic model**

2 **Economic model for review question: What antibiotic treatment regimens are**  
3 **effective in treating bacterial meningitis caused by Haemophilus influenzae?**

4 No economic analysis was conducted for this review question.

5

## 1 Appendix J Excluded studies

### 2 Excluded studies for review question: What antibiotic treatment regimens are 3 effective in treating bacterial meningitis caused by Haemophilus influenzae?

#### 4 Excluded effectiveness studies

5 The excluded studies table only lists the studies that were considered and then excluded at  
6 the full-text stage for this review (N=9) and not studies (N=181) that were considered and  
7 then excluded from the search at the full-text stage as per the PRISMA diagram in Appendix  
8 C for the other review questions in the same search.

9 **Table 7: Excluded studies and reasons for their exclusion**

Study	Code [Reason]
Baker, C. J. and Long, S. S. (2019) 50 Years Ago in the Journal of Pediatrics: Ampicillin in the Treatment of Meningitis due to Haemophilus influenzae: An Appraisal after 6 Years of Experience. Journal of Pediatrics 208: 37	- Study design does not meet inclusion criteria
Boisivon, A., Berardi-Grassias, L., Guiomar, C. et al. (1987) Bacteriostatic and bactericidal effect of ceftriaxone on Hemophilus, Neisseria meningitidis, and Proteus. Chemioterapia : international journal of the Mediterranean Society of Chemotherapy 6(2suppl): 83-84	- No comparison of interest
Congeni, B. L. (1984) Comparison of ceftriaxone and traditional therapy of bacterial meningitis. Antimicrobial agents and chemotherapy 25(1): 40-44	- Population does not meet inclusion criteria
Goldwater, Paul N. (2005) Cefotaxime and ceftriaxone cerebrospinal fluid levels during treatment of bacterial meningitis in children. International journal of antimicrobial agents 26(5): 408-11	- No outcomes of interest
Lebel, M. H.; Hoy, M. J.; McCracken Jr, G. H. (1989) Comparative efficacy of ceftriaxone and cefuroxime for treatment of bacterial meningitis. Journal of Pediatrics 114(6): 1049-1054	- No comparison of interest
Peltola, H.; Anttila, M.; Renkonen, O. V. (1989) Randomised comparison of chloramphenicol, ampicillin, cefotaxime, and ceftriaxone for childhood bacterial meningitis. Finnish Study Group. Lancet (london, england) 1(8650): 1281-1287	- Population does not meet inclusion criteria
Roine, I., Ledermann, W., Foncea, L. M. et al. (2000) Randomized trial of four vs. seven days of ceftriaxone treatment for bacterial meningitis in children with rapid initial recovery. Pediatric infectious disease journal 19(3): 219-222	- Population does not meet inclusion criteria
Singhi, P., Kaushal, M., Singhi, S. et al. (2002)	- Population does not meet inclusion criteria

DRAFT FOR CONSULTATION

Antibiotics for bacterial meningitis caused by Haemophilus influenzae

Study	Code [Reason]
Seven days vs. 10 days ceftriaxone therapy in bacterial meningitis. Journal of tropical pediatrics 48(5): 273-279	
Toltzis, P. (2019) 50 Years Ago in the Journal of Pediatrics: Relapse of Hemophilus influenzae Type b Meningitis during Intravenous Therapy with Ampicillin. Journal of Pediatrics 208: 182	- Study design does not meet inclusion criteria

1

2 **Excluded economic studies**

3 No studies were identified which were applicable to this review question.

4

- 1 **Appendix K Research recommendations – full details**
- 2 **Research recommendations for review question: What antibiotic treatment**
- 3 **regimens are effective in treating bacterial meningitis caused by Haemophilus**
- 4 **influenzae?**
- 5 No research recommendation was made for this review.