

1

2

3

4

5

6

7 **The management of hip fracture in adults**

8

9

10

11

METHODS, EVIDENCE & GUIDANCE

14

15

16

17 Produced by the National Clinical Guideline Centre

1

2
3
4

5

6

7

8

9

10

11

12
13

14

15

16

17
18
19
20
21
22

23
24
25

26
27

28
29
30

31

Published by the National Clinical Guideline Centre at The Royal College of Physicians, 11 St Andrews Place, Regent's Park, London, NW11 4LE

First published 2011

© National Clinical Guideline Centre 2011

Apart from any fair dealing for the purposes of research or private study, criticism or review, as permitted under the Copyright, Designs and Patents Act, 1988, no part of this publication may be reproduced, stored or transmitted in any form or by any means, without the prior written permission of the publisher or, in the case of reprographic reproduction, in accordance with the terms of licences issued by the Copyright Licensing Agency in the UK. Enquiries concerning reproduction outside the terms stated here should be sent to the publisher at the UK address printed on this page.

The use of registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant laws and regulations and therefore for general use.

The rights of the National Clinical Guideline Centre to be identified as Author of this work have been asserted by them in accordance with the Copyright, Designs and Patents Act, 1988.

Citation: National Clinical Guideline Centre, ([year of publication on NICE website]). [The Management of Hip Fracture in Adults]. London: National Clinical Guideline Centre. Available from: www.ncgc.ac.uk

1 Contents

2	CONTENTS	1
3	GUIDELINE DEVELOPMENT GROUP MEMBERS	1
4	NCGC STAFF MEMBERS OF THE GUIDELINE DEVELOPMENT GROUP	2
5	EXPERT ADVISORS	2
6	ACKNOWLEDGEMENTS	3
7	ACRONYMS AND ABBREVIATIONS	4
8	1 INTRODUCTION	7
9	2 DEVELOPMENT OF THE GUIDELINE	10
10	2.1 WHAT IS A NICE CLINICAL GUIDELINE?	10
11	2.2 REMIT	11
12	2.3 WHO DEVELOPED THIS GUIDELINE?	11
13	2.4 WHAT THIS GUIDELINE COVERS.....	12
14	2.5 WHAT THIS GUIDELINE DOES NOT COVER.....	13
15	2.6 RELATIONSHIPS BETWEEN THE GUIDELINE AND OTHER NICE GUIDANCE	13
16	3 METHODS	15
17	3.1 DEVELOPING THE REVIEW QUESTIONS AND OUTCOMES	15
18	3.2 SEARCHING FOR EVIDENCE.....	20
19	3.3 EVIDENCE OF EFFECTIVENESS.....	21
20	3.4 EVIDENCE OF COST-EFFECTIVENESS.....	27
21	3.5 DEVELOPING RECOMMENDATIONS	30
22	3.6 VALIDATION PROCESS	31
23	3.7 UPDATING THE GUIDELINE	31
24	3.8 DISCLAIMER.....	31
25	3.9 FUNDING	31
26	4 GUIDELINE SUMMARY	32
27	4.1 MAP OF RECOMMENDATIONS	32
28	AN ALGORITHM WILL BE ADDED BEFORE PUBLICATION. KEY PRIORITIES FOR IMPLEMENTATION	32
29	KEY PRIORITIES FOR IMPLEMENTATION	33
30	4.2 FULL LIST OF RECOMMENDATIONS.....	35
31	4.3 RESEARCH RECOMMENDATIONS	39
32	5 IMAGING OPTIONS IN OCCULT HIP FRACTURE	46
33	5.1 INTRODUCTION	46
34	5.2 REVIEW QUESTION.....	47
35	5.3 RADIOGRAPHS	47
36	5.4 RADIONUCLIDE BONE SCAN (RNS)	48
37	5.5 ULTRASOUND (US).....	49

1	5.6	COMPUTED TOMOGRAPHY (CT)	50
2	5.7	RESEARCH RECOMMENDATION ON IMAGING OPTIONS IN OCCULT HIP FRACTURE	54
3	6	TIMING OF SURGERY	55
4	6.1	INTRODUCTION	55
5	6.2	RESEARCH RECOMMENDATIONS ON TIMING OF SURGERY	69
6	7	ANALGESIA	70
7	7.1	INTRODUCTION	70
8	7.2	SYSTEMIC ANALGESIA	71
9	7.3	NERVE BLOCKS COMPARED TO SYSTEMIC ANALGESIA	71
10	7.4	RESEARCH RECOMMENDATIONS ON ANALGESIA	82
11	8	REGIONAL (SPINAL OR EPIDURAL) VERSUS GENERAL ANAESTHESIA.....	83
12	8.1	INTRODUCTION	83
13	8.2	REGIONAL VERSUS GENERAL ANAESTHESIA	83
14	8.3	RESEARCH RECOMMENDATION ON ANAESTHESIA	88
15	9	SURGEON SENIORITY	90
16	9.1	INTRODUCTION	90
17	9.2	SURGEON SENIORITY	90
18	9.3	RECOMMENDATIONS AND LINK TO EVIDENCE	92
19	10	SURGICAL PROCEDURES	96
20	10.1	INTRODUCTION	96
21	10.2	SURGERY WITH REGARD TO EARLY MOBILISATION	96
22	10.3	DISPLACED INTRACAPSULAR FRACTURES	97
23	10.4	USE OF CEMENT IN ARTHROPLASTY	114
24	10.5	SURGICAL APPROACH TO HEMIARTHROPLASTY	122
25	10.6	EXTRACAPSULAR FRACTURE FIXATION	125
26	11	MOBILISATION STRATEGIES.....	135
27	11.1	INTRODUCTION	135
28	11.2	EARLY VS. DELAYED MOBILISATION	136
29	11.3	INTENSITY OF PHYSIOTHERAPY	139
30	12	MULTIDISCIPLINARY MANAGEMENT.....	146
31	12.1	INTRODUCTION	146
32	12.2	HOSPITAL-BASED MULTIDISCIPLINARY REHABILITATION VERSUS USUAL CARE	148
33	12.3	RESEARCH RECOMMENDATIONS ON HOSPITAL MULTIDISCIPLINARY REHABILITATION	165
34	12.4	COMMUNITY-BASED MULTIDISCIPLINARY REHABILITATION VERSUS USUAL CARE	167
35	12.5	RESEARCH RECOMMENDATIONS ON COMMUNITY MULTIDISCIPLINARY REHABILITATION.....	177
36	13	PATIENT AND CARER VIEWS AND INFORMATION.....	179
37	13.1	INTRODUCTION	179
38	13.2	PATIENT AND CARER VIEWS.....	179
39	13.3	INFORMATION FOR PATIENTS	186
40	13.4	CARER INVOLVMENT.....	188
41	13.5	RESEARCH RECOMMENDATIONS	188
42		GLOSSARY	190
43		APPENDICES.....	203
44		APPENDIX A: SCOPE	210
45	13.6	GUIDELINE TITLE	210
46	13.7	THE REMIT	210
47	13.8	CLINICAL NEED FOR THE GUIDELINE.....	210

1	13.9	THE GUIDELINE	213
2	13.10	RELATED NICE GUIDANCE	216
3	13.11	FURTHER INFORMATION.....	217
4	14	APPENDIX B: DECLARATIONS OF INTEREST	219
5	14.1	INTRODUCTION	219
6	14.2	DECLARATIONS OF INTERESTS OF THE GDG MEMBERS.....	220
7	14.3	DECLARATIONS OF INTERESTS OF THE NCGC MEMBERS.....	233
8	14.4	DECLARATIONS OF INTERESTS OF THE EXPERT ADVISORS.....	234
9	15	APPENDIX C: REVIEW PROTOCOLS.....	235
10	15.1	REVIEW PROTOCOL – IMAGING IN OCCULT HIP FRACTURE.....	235
11	15.2	REVIEW PROTOCOL – TIMING OF SURGERY.....	237
12	15.3	REVIEW PROTOCOL – ANALGESIA- SYSTEMIC MEDICATIONS.....	239
13	15.4	REVIEW PROTOCOL – ANALGESIA- NERVE BLOCKS COMAPARED TO SYSTEMIC ANALGESICS.....	241
14	15.5	REVIEW PROTOCOL – ANAESTHESIA	243
15	15.6	REVIEW PROTOCOL – SURGEON SENIORITY.....	245
16	15.7	REVIEW PROTOCOL – CEMENT.....	246
17	15.8	REVIEW PROTOCOL – INTRACAPSULAR FRACTURES.....	248
18	15.9	REVIEW PROTOCOL – SURGICAL APPROACH	250
19	15.10	REVIEW PROTOCOL – HEMIARTHROPLASTY STEM DESIGN.....	252
20	15.11	REVIEW PROTOCOL – EXTRACAPSULAR FRACTURES	254
21	15.12	REVIEW PROTOCOL – MOBILISATION STRATEGIES.....	256
22	15.13	REVIEW PROTOCOL – MULTIDISCIPLINARY REHABILITATION.....	258
23	15.14	REVIEW PROTOCOL – CARER INVOLVEMENT	264
24	15.15	REVIEW PROTOCOL – HEALTH ECONOMICS.....	266
25	16	APPENDIX D: LITERATURE SEARCH STRATEGIES.....	268
26	16.1	SEARCH STRATEGIES.....	268
27	16.2	SEARCH TERMS.....	270
28	17	APPENDIX E: EVIDENCE TABLES - CLINICAL STUDIES.....	291
29	17.1	EVIDENCE TABLE 1: IMAGING OPTIONS IN OCCULT HIP FRACTURE.....	293
30	17.2	EVIDENCE TABLE 2: TIMING OF SURGERY	296
31	17.3	EVIDENCE TABLE 3: OPTIMAL ANALGESIA	314
32	17.4	EVIDENCE TABLE 4: ANAESTHESIA	317
33	17.5	EVIDENCE TABLE 5: SURGEON SENIORITY.....	320
34	17.6	EVIDENCE TABLE 6: DISPLACED INTRACAPSULAR FRACTURES.....	324
35	17.7	EVIDENCE TABLE 7: SURGERY – CEMENT VERSUS NO CEMENT.....	333
36	17.8	EVIDENCE TABLE 8: EXTRACAPSULAR FIXATION	335
37	17.9	EVIDENCE TABLE 9: SURGICAL APPROACH TO HEMIARTHROPLASTY.....	383
38	17.10	EVIDENCE TABLE 10: MOBILISATION STRATEGIES	385
39	17.11	EVIDENCE TABLE 11: MULTIDISCIPLINARY REHABILITATION	394
40	17.12	EVIDENCE TABLE 12: PATIENT VIEWS.....	418
41	18	APPENDIX F: EVIDENCE TABLES - ECONOMIC STUDIES	434
42	18.1	EVIDENCE TABLE 13: GENERAL VERSUS REGIONAL ANAESTHESIA	435
43	18.2	EVIDENCE TABLE 14: DISPLACED INTRACAPSULAR FRACTURES.....	437
44	18.3	EVIDENCE TABLE 15: CEMENTED ARTHROPLASTIES	441
45	18.4	EVIDENCE TABLE 16: MULTIDISCIPLINARY REHABILITATION	443
46	19	APPENDIX G: FOREST PLOTS	456
47	19.1	RADIOLOGY	459
48	19.2	TIMING OF SURGERY.....	460
49	19.3	ANALGESIA.....	466
50	19.4	ANAESTHESIA.....	473
51	19.5	SURGICAL INTERVENTIONS	478

1	19.6	MOBILISATION STRATEGIES	517
2	19.7	INTENSIVE EXERCISE OR PHYSIOTHERAPY VS. USUAL CARE	519
3	19.8	MULTIDISCIPLINARY REHABILITATION	524
4	19.9	HOME-BASED MDR VERSUS USUAL INPATIENT REHABILITATION	529
5	20	APPENDIX H: HEALTH ECONOMIC ANALYSIS.....	532
6	20.1	COST ANALYSIS OF NERVE BLOCKS, NON-OPIOID ANALGESICS AND OPIOID ANALGESICS.....	532
7	20.2	HOURLY WAGE COSTS FOR A PLANNED TRAUMA LIST	535
8	20.3	PRICES FOR SLIDING HIP SCREWS AND SHORT AND LONG INTRAMEDULLARY NAILS	538
9	20.4	COST ANALYSIS OF THE INTERVENTIONS FOR INTENSIVE MOBILISATION STRATEGIES.	539
10	20.5	COST-EFFECTIVENESS ANALYSIS OF HOSPITAL INVESTMENT VERSUS NO HOSPITAL INVESTMENT FOR EARLY SURGERY	
11		542	
12	20.6	COST-EFFECTIVENESS ANALYSIS OF HOSPITAL MDR VS USUAL CARE	559
13	20.7	COST-EFFECTIVENESS ANALYSIS OF COMMUNITY MDR VS USUAL CARE	593
14	20.8	COST ANALYSIS OF CEMENTED VS. UNCEMENTED IMPLANTS (NEWER DESIGNS OF ARTHROPLASTY)	604
15	21	APPENDIX I: HIGH PRIORITY RESEARCH RECOMMENDATIONS.....	610
16	21.1	IMAGING OPTIONS IN OCCULT HIP FRACTURE	610
17	21.2	ANAESTHESIA.....	613
18	21.3	DISPLACED INTRACAPSULAR HIP FRACTURES	616
19	21.4	INTENSIVE REHABILITATION THERAPIES AFTER HIP FRACTURE	620
20	21.5	EARLY SUPPORTED DISCHARGE IN CARE HOME PATIENTS	623
21	22	APPENDIX J: EXCLUDED STUDIES	627
22		BIBLIOGRAPHY.....	636
23			

1 Guideline development group members

2

Professor Cameron Swift (Chair)	Emeritus Professor of Health Care of the Elderly
Mr Tim Chesser	Consultant Trauma and Orthopaedic Surgeon
Mr Anthony Field	Patient Member
Dr Richard Griffiths	Consultant Anaesthetist
Mr Robert Handley	Consultant Trauma and Orthopaedic Surgeon
Mrs Karen Hertz	Advanced Nurse Practitioner, Locomotor Directorate
Dr Sally Hope	General Practitioner
Dr Antony Johansen	Consultant Orthogeriatrician
Professor Sarah (Sallie) Lamb	Professor of Rehabilitation, Director of Warwick Clinical Trials Unit, Professor of Trauma Rehabilitation
Professor Opinder Sahota	Consultant Physician
Mrs Tessa Somerville	Patient Member
Mrs Heather Towndrow	Clinical Manager , Day Rehabilitation and Falls Prevention
Mr Martin Wiese	Consultant in Emergency Medicine

1 NCGC staff members of the guideline

2 development group

Dr Saoussen Ftouh	Senior Research Fellow / Project Manager
Ms Joanna Ashe	Information Scientist
Miss Elisabetta Fenu	Senior Health Economist
Dr Jennifer Hill	Operations Director
Dr Antonia Morga	Health Economist
Dr Sarah Riley	Research Fellow
Mr Carlos Sharpin	Senior Information Scientist / Research Fellow

3 Expert advisors

Professor Judy Adams	Consultant and Honorary Professor of Diagnostic Radiology
Mrs Pamela Holmes	Practice Development Manager (SCIE)
Mr Martyn Parker	Consultant Orthopaedic Surgeon
Dr Luigi Siciliani	Senior Lecturer in Health Economics

1 Acknowledgements

2 The development of this guideline was greatly assisted by the following people

3 NCGC

4 Kate Lovibond, David Wonderling, Abigail Jones, Maggie Westby, Laura Sawyer,
5 Clare Jones, Dr Lee-Yee Chong, Tamara Diaz

6 External

7 Dr Stuart White, Professor Maria Crotty, Dr Michelle Miller, The members of the
8 “Hip Fracture Anaesthesia” NHS Network, especially Dr Michael McBrien, Mr.
9 David Chittenden, Mr. John Ellington, Mr. David Drust, Ms Liz Hedinger and Ms
10 Sally Lambert

11

12

13

1 Acronyms and abbreviations

ADL	Activities of Daily Living
ANOVA	Analysis of variance
AO	Arbeitsgemeinschaft für Osteosynthesefragen
BNF	British National Formulary
CCA	Cost-consequences analysis
CEA	Cost-effectiveness analysis
c.f.	Confer (refer to)
CI / 95% CI	Confidence interval / 95% confidence interval
CT	Computed tomography
CUA	Cost-utility analysis
DH	Department of Health
DSA	Deterministic Sensitivity Analysis
ED	Emergency Department
ESD	Early Supported Discharge
EQ-5D	EuroQoL-5D
GA	General anaesthesia
GORU	Geriatric Orthopaedic Rehabilitation Unit
GDG	Guideline Development Group
GP	General Practitioner
GRADE	Grading of Recommendations Assessment, Development and Evaluation
HES	Hospital Episode Statistics
HFP	Hip fracture programme
HR	Hazard Ratio
HRQoL	Health-related quality of life
HTA	Health technology assessment
IC	Intermediate care
ICD-10	International Classification of Diseases, 10 th edition
ICER	Incremental cost-effectiveness ratio

IQR	Interquartile range
INMB	Incremental Net Monetary Benefit
IRR	Inter-rater reliability
ITT	Intention to treat
LOS	Length of Stay
LR⁺	Positive likelihood ratio
LR⁻	Negative likelihood ratio
LY	Life-Year
MD	Mean Difference
MDR	Multi-Disciplinary Rehabilitation
MARU	Mixed Assessment and Rehabilitation Unit
MRI	Magnetic resonance imaging
NCGC	National Clinical Guideline Centre
N/A	Not applicable
NHS	National Health Service
NHSEED	The NHS Economic Evaluation Database
NICE	National Institute for Health and Clinical Excellence
NNT	Number needed to treat
NPV	Negative predictive value
NSAID	Non steroidal anti-inflammatory drugs
OR	Odds ratio
PICO	Framework incorporating patients, interventions, comparison and outcome
POMA	Tinetti's performance oriented mobility assessment
PFF	Proximal femoral fracture
PPP	Purchasing Power Parity
PPV	Positive predictive value
p.r.n	Pro re nata
PSA	Probabilistic sensitivity analysis
QALY	Quality-adjusted life year
QUADAS	Quality assessment tool for diagnostic accuracy studies
RA	Regional anaesthesia
RCT	Randomised controlled trial
RNS	Radionuclide scan
ROC	Receiver operating characteristic
RR	Relative risk

SCIE	Social Care Institute for Excellence
SD	Standard deviation
SE	Standard error
SPC	Summary of product characteristics
SR	Systematic review
US	Ultrasound
WTP	Willingness to pay

1 Introduction

2 Hip fracture is the plain English term for a proximal femoral fracture or PFF. It refers to a
3 fracture occurring in the area between the edge of the femoral head and 5 centimetres
4 below the lesser trochanter (Figure 1). These fractures are generally divided into two main
5 groups depending on their relationship to the capsule of the hip joint. Those above the
6 insertion of the capsule are termed intracapsular, subcapital or femoral neck fractures.
7 Those below the insertion are extracapsular. The extracapsular group is split further into
8 trochanteric (inter- or pertrochanteric and reverse oblique) and subtrochanteric as shown.
9 The division into intra and extracapsular fractures relates to both the blood supply of the
10 femoral head and the mechanics of fixation.

11 Hip fracture is a major public health issue due to an ever increasing ageing population.
12 About 70,000 to 75,000 hip fractures (proximal femoral fractures) occur annually in the
13 UK³⁹, with a cost (including medical and social care) amounting to about £2 billion a year.
14 Demographic projections indicate that the UK annual incidence will rise to 91,500 by 2015
15 and 101,000 in 2020³⁹, with an associated increase in annual expenditure. The majority of
16 this expenditure will be accounted for by hospital bed days and a further substantial
17 contribution will come from health and social aftercare. At present about a quarter of
18 patients with hip fracture are admitted from institutional care, and about 10–20% of those
19 admitted from home ultimately move to institutional care.

20 Hip fracture is the commonest reason for admission to an orthopaedic trauma ward and is
21 usually a ‘fragility’ fracture¹ caused by a fall affecting an older person with osteoporosis or
22 osteopaenia (a condition in which bones lose calcium and become thinner, but not as much
23 as in osteoporosis). The National Hip Fracture Database reports the average age of a person
24 with hip fracture as 84 years for men and 83 for women, 76% of fracture occur in women.
25 Mortality is high – about 10% of people with a hip fracture die within 1 month and about
26 one third within 12 months. Most of the deaths are due to associated co morbidities
27 (including bronchopneumonia) and not just to the fracture itself reflecting the high
28 prevalence of comorbidity in people with hip fracture. It is often the occurrence of a fall
29 and fracture that signals underlying ill health. Thus, hip fracture is by no means an
30 exclusively surgical concern. Its effective management requires the co-ordinated
31 application of medical, surgical, anaesthetic and multidisciplinary rehabilitation skills and a
32 comprehensive approach covering the full time course of the condition from presentation
33 to subsequent follow-up, including the transition from hospital to community.

¹ The strict definition of a fragility fracture is one caused by a fall from standing height or less. For the purposes of this guidance, the definition is slightly more flexible to encompass all hip fractures judged to have an osteoporotic or osteopaenic basis

1 Although hip fracture is predominantly a phenomenon of later life, it may occur at any age
2 in people with osteoporosis or osteopenia, and this guidance is applicable to adults across
3 the age spectrum. Skills in its management have, however been accrued, researched and
4 reported especially by collaborative teams specialising in the care of older people (using the
5 general designation 'orthogeriatrics'). These skills are applicable in hip fracture irrespective
6 of age, and the guidance includes recommendations that cover the needs of younger
7 patients by drawing on such skills in an organised manner.

8 This guidance covers the management of hip fracture from the point of admission to
9 secondary care through to final return to the community and discharge from specific
10 follow-up. It assumes that anyone clinically suspected of having a hip fracture will be
11 referred for immediate hospital assessment other than in exceptional circumstances. It
12 excludes (other than by cross-reference) aspects covered by parallel NICE guidance, most
13 notably primary and secondary prevention of fragility fractures, but recognises the
14 importance of effective linkage to these closely related elements of comprehensive care.

15 The diagnosis of hip fracture is easily missed and in a small minority of patients the fracture
16 may not be apparent on a plain X-ray. In view of the serious nature of hip fracture the
17 guidance has sought to identify the most cost-effective imaging strategies to ensure this
18 does not happen.

19 Although not a structured service delivery evaluation, the Guideline Group was required to
20 extend its remit to cover essential implications for service organisation within the NHS
21 where these are fundamental to hip fracture management, and this has been done. In
22 general it is the case that suboptimal care and/or fragmentation of care result in longer
23 periods of dependency and/or hospitalisation leading to greater cost as well as inferior
24 outcome. There is substantial variation and lack of clarity in the UK in the extent, timing,
25 manner and organisation of the necessary collaborative and multidisciplinary elements of
26 effective management, including the timely achievement of rehabilitation after surgery
27 according to individual need. A further concern is the occurrence of delay before necessary
28 surgery is carried out. Prompt surgery has been generally recognised to be important, but
29 surgery is sometimes delayed for administrative or clinical reasons. Emerging evidence from
30 the National Hip Fracture database indicates substantial variation across centres in England
31 and Wales in this and other indicators of clinical and service quality. Such variation has
32 potentially profound economic implications, and priority has been given where appropriate
33 to underpinning recommendations with any available evidence of cost-effectiveness in the
34 NHS. Since work began on the guideline the Department of Health in England has launched
35 a high priority Best Practice Tariff initiative targeting a range of performance variables for
36 hip fracture, and the GDG have been aware of this contextual change as well as of
37 humanitarian issues in evaluating the evidence and formulating recommendations.


38 At all stages of hip fracture management, the importance of optimal communication with,
39 and support for, patients themselves and those who provide or will provide care –
40 including unpaid care family members or others – has been a fundamental tenet of
41 guidance development.

42 The view of the GDG is that an exceptional contemporary window of opportunity exists in
43 the NHS to achieve major improvements in the delivery of hip fracture care, to the benefit
44 not only of patients but of the system as a whole in terms of efficiency and cost. It is hoped
45 that implementation of this guidance will be instrumental to that end.

1

2

Figure 1: Types of hip fracture (Parker M & Johansen A, 2006)^{259,270}



Classification of hip fractures. Fractures in the blue area are intracapsular and those in the red and orange areas are extracapsular

3

4

5

Reproduced from BMJ, Parker, M., Johansen, A., 333(7557), 27-30, 2006 with permission from BMJ Publishing Group Ltd

2 Development of the guideline

2.1 What is a NICE clinical guideline?

NICE clinical guidelines are recommendations for the care of individuals in specific clinical conditions or circumstances within the NHS – from prevention and self-care through primary and secondary care to more specialised services. We base our clinical guidelines on the best available research evidence, with the aim of improving the quality of health care. We use predetermined and systematic methods to identify and evaluate the evidence relating to specific review questions.

NICE clinical guidelines can:

- provide recommendations for the treatment and care of people by health professionals
- be used to develop standards to assess the clinical practice of individual health professionals
- be used in the education and training of health professionals
- help patients to make informed decisions
- improve communication between patient and health professional

While guidelines assist the practice of healthcare professionals, they do not replace their knowledge and skills.

We produce our guidelines using the following steps:

- Guideline topic is referred to NICE from the Department of Health
- Stakeholders register an interest in the guideline and are consulted throughout the development process.
- The scope is prepared by the National Clinical Guideline Centre (NCGC)
- The NCGC establishes a guideline development group
- A draft guideline is produced after the group assesses the available evidence and makes recommendations

- 1 • There is a consultation on the draft guideline.
- 2 • The final guideline is produced.
- 3
- 4 The NCGC and NICE produce a number of versions of this guideline:
- 5 • the **full guideline** contains all the recommendations, plus details of the methods
- 6 used and the underpinning evidence
- 7 • the **NICE guideline** lists the recommendations
- 8 • the **quick reference guide** (QRG) presents recommendations in a suitable format
- 9 for health professionals
- 10 • information for the public (**'understanding NICE guidance' or UNG**) is written using
- 11 suitable language for people without specialist medical knowledge.

12 This version is the full version. The other versions can be downloaded from NICE

13 www.NICE.org.uk and the NCGC website www.ncgc.ac.uk.

14 **2.2 Remit**

15 NICE received the remit for this guideline from the Department of Health. They

16 commissioned the NCGC to produce the guideline.

17 The remit for this guideline is:

18 To prepare a clinical guideline on the management of fractured neck of femur.

19 **2.3 Who developed this guideline?**

20 A multidisciplinary Guideline Development Group (GDG) comprising professional group

21 members and consumer representatives of the main stakeholders developed this guideline

22 (see section on Guideline Development Group Membership and acknowledgements).

23 The National Institute for Health and Clinical Excellence funds the National Clinical

24 Guideline Centre (NCGC) and thus supported the development of this guideline. The GDG

25 was convened by the NCGC and chaired by Professor Cameron Swift in accordance with

26 guidance from the National Institute for Health and Clinical Excellence (NICE).

27 The group met every 6-8 weeks during the development of the guideline. At the start of the

28 guideline development process all GDG members declared interests including

29 consultancies, fee-paid work, share-holdings, fellowships and support from the healthcare

30 industry. At all subsequent GDG meetings, members declared arising conflicts of interest,

31 which were also recorded.

32 Members were either required to withdraw completely or for part of the discussion if their

33 declared interest made it appropriate. The details of declared interests and the actions

34 taken are shown in Appendix B.

1 Staff from the NCGC provided methodological support and guidance for the development
2 process. The team working on the guideline included a project manager, systematic
3 reviewers, health economists and information scientists. They undertook systematic
4 searches of the literature, appraised the evidence, conducted meta analysis and cost
5 effectiveness analysis where appropriate and drafted the guideline in collaboration with the
6 GDG.

8 **2.4 What this guideline covers**

9 The population of this guideline covers:

- 10 a) Adults aged 18 years and older presenting to the health service with a clinical
11 diagnosis (firm or provisional) of fragility fracture of the hip.
- 12 b) People with the following types of hip fracture:
- 13 • intracapsular (undisplaced and displaced)
 - 14 • extracapsular (trochanteric and subtrochanteric).
- 15 c) Those with comorbidity strongly predictive of outcome, and those without such
16 comorbidity. The influence (if any) of advanced age or gender on clinical decision-
17 making, management and outcome will be specifically evaluated.

18 For further details please refer to the scope in Appendix A and review protocols in
19 Appendix C.

20 Key clinical areas in this guideline are:

- 21 a) Using alternative radiological imaging to confirm or exclude a suspected hip
22 fracture in patients with a normal X-ray.
- 23 b) Involving a physician or orthogeriatrician in the care of patients presenting with hip
24 fracture.
- 25 c) Early surgery (within 48 hours).
- 26 d) Optimal preoperative and postoperative analgesia (pain relief), including the use of
27 nerve blockade.
- 28 e) Regional (spinal – also known as ‘epidural’) versus general anaesthesia in patients
29 undergoing surgery for hip fracture.
- 30 f) Surgeon experience and seniority
- 31 g) For displaced intracapsular fracture:
- 32 • Internal fixation versus arthroplasty (hip replacement surgery)
 - 33 • Total hip replacement versus hemiarthroplasty (replacing the head of the
34 femur only).

- 1 h) Choice of surgical implants - Sliding hip screw versus intramedullary nail for
2 trochanteric extracapsular fracture.
- 3 i) Choice of surgical implants - Sliding hip screw versus intramedullary nail for
4 subtrochanteric extracapsular fracture.
- 5 j) Cemented versus non-cemented arthroplasty implants.
- 6 k) Hospital-based multidisciplinary rehabilitation for patients who have undergone hip
7 fracture surgery.
- 8 l) Early transfer to community-based multidisciplinary rehabilitation for patients who
9 have undergone hip fracture surgery.

10 **2.5 What this guideline does not cover**

11 The population of this guideline does not cover:

- 12 a) People younger than 18 years.
- 13 b) People with fractures caused by specific pathologies other than osteoporosis or
14 osteopaenia (because these would require more condition-specific guidance).

15 Clinical areas not included in this guideline are:

- 16 a) Primary and secondary prevention of fragility fracture.
- 17 b) Prevention and management of pressure sores.
- 18 c) Prophylaxis for venous thromboembolism.
- 19 d) Prevention and management of infection at the surgical site.
- 20 e) Nutritional support.
- 21 f) Selection of prostheses for hip replacement.
- 22 g) Complementary and alternative therapies.

23 **2.6 Relationships between the guideline and other NICE guidance**

24 **Related NICE Health Technology Appraisals:**

25 Alendronate, etidronate, risedronate, raloxifene, strontium ranelate and teriparatide for
26 the secondary prevention of osteoporotic fragility fractures in postmenopausal women
27 (amended). NICE technology appraisal guidance TA161 (2011). Available from
28 www.nice.org.uk/TA161

29 Alendronate, etidronate, risedronate, raloxifene and strontium ranelate for the primary
30 prevention of osteoporotic fragility fractures in postmenopausal women (amended). NICE
31 technology appraisal guidance TA160 (2011). Available from www.nice.org.uk/TA160

- 1 The selection of prostheses for primary total hip replacement. NICE technology appraisal
2 guidance TA2 (2000). Available from www.nice.org.uk/TA2
- 3 **Related NICE Interventional Procedures Guidance:**
- 4 Minimally invasive hip replacement. NICE interventional procedure guidance (2010).
5 Available from www.nice.org.uk/guidance/IPG363
- 6 **Related NICE Clinical Guidelines:**
- 7 Surgical site infection. NICE clinical guideline CG74 (2008). Available from
8 www.nice.org.uk/CG74
- 9 Nutrition support in adults. NICE clinical guideline CG32 (2006). Available from
10 www.nice.org.uk/CG32
- 11 The management of pressure ulcers in primary and secondary care. NICE clinical guideline
12 CG29 (2005). Available from www.nice.org.uk/CG29
- 13 Falls. NICE clinical guideline CG21 (2004). Available from www.nice.org.uk/CG21
- 14 Preoperative tests. NICE clinical guideline CG3 (2003). Available from www.nice.org.uk/CG3
- 15 Venous thromboembolism – reducing the risk. NICE clinical guideline CG92 (2010). Available
16 from <http://guidance.nice.org.uk/CG92>
- 17 Delirium: diagnosis, prevention and management of delirium. NICE clinical guideline CG103
18 (2010). Available from <http://guidance.nice.org.uk/CG103>
- 19 Dementia: supporting people with dementia and their carers in health and social care. NICE
20 clinical guideline CG42 (2006). Available from www.nice.org.uk/CG42
- 21 NICE Related Guidance currently in development:
- 22 Osteoporosis. NICE clinical guideline. Publication date to be confirmed.

1 **3 Methods**

2 This guidance was developed in accordance with the methods outlined in the NICE
3 Guidelines Manual 2009²³³

4 **3.1 Developing the review questions and outcomes**

5 Review questions were developed in a PICO framework (patient, intervention, comparison
6 and outcome) for intervention reviews, and with a framework of population, index tests,
7 reference standard and target condition for reviews of diagnostic test accuracy. This was to
8 guide the literature searching process and to facilitate the development of
9 recommendations by the guideline development group (GDG). They were drafted by the
10 NCGC technical team and refined and validated by the GDG. The questions were based on
11 the key clinical areas identified in the scope (Appendix A). Further information on the
12 outcome measures examined follows this section.

Chapter	Review question	Outcomes
Radiology	In patients with a continuing clinical suspicion of hip fracture, despite negative radiographic findings, what is the clinical and cost-effectiveness of additional imaging (radiography after at least 48 hours), Radionuclide scanning (RNS), ultrasound (US) and computed tomography (CT), compared to magnetic resonance imaging (MRI), in confirming, or excluding, a hip fracture?	<ul style="list-style-type: none"> ▪ Sensitivity ▪ Specificity ▪ Positive and negative predictive values ▪ Positive and negative likelihood ratios
Timing of surgery	In patients with hip fractures what is the clinical and cost effectiveness of early surgery (within 24, 36 or 48 hours) on the incidence of complications such as mortality, pneumonia, pressure sores, cognitive dysfunction and increased length of hospital stay?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge ▪ Place of residence (compared with baseline) 12 months after fracture ▪ Functional status (30 days, 3 months, 1 year) ▪ Quality of life (30 days, 3 months, 1 year) ▪ Complications (including pressure ulcers)
Analgesia	In patients who have or are suspected of having a hip fracture, what is the clinical and cost effectiveness of nerve blocks compared to systemic analgesia in providing adequate pain relief and reducing side effects and mortality?	<ul style="list-style-type: none"> ▪ Pain Need for 'breakthrough' analgesia ▪ Mortality ▪ Adverse effects
Anaesthesia	In patients undergoing surgical repair for hip fractures, what is the clinical and cost effectiveness of regional (spinal/epidural) anaesthesia compared to general anaesthesia in reducing complications such as mortality, cognitive dysfunction thromboembolic events, postoperative respiratory morbidity, renal failure and length of stay in hospital?	<ul style="list-style-type: none"> ▪ Patient preference ▪ Early mortality up to 1 month ▪ Functional status up to 1 year ▪ Pain Adverse effects

Surgeon seniority	Does surgeon seniority (consultant or equivalent) reduce the incidence of mortality, operative revision and poor functional outcome?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Reoperation rate ▪ Dislocations ▪ Wound infection
Cement	In hip fracture patients undergoing total hip replacement what is the clinical and cost effectiveness of cemented total hip replacement versus uncemented total hip replacement on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?	<ul style="list-style-type: none"> ▪ Perioperative mortality ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture ▪ Wound healing complications
Intracapsular fractures	In patients undergoing repair for intracapsular hip fractures what is the clinical and cost effectiveness of internal fixation compared to hemiarthroplasty compared to total hip replacement on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture
Surgical approach	In patients having surgical treatment for intracapsular hip fracture with hemiarthroplasty what is the clinical and cost effectiveness of anterolateral compared to posterior surgical approach on mortality, number of reoperations, dislocation, functional status, length of hospital stay, quality of life and pain?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of hospital stay ▪ Reoperation rate ▪ Dislocations ▪ Functional status ▪ Quality of life ▪ Pain

Hemiarthroplasty stem design	In patients undergoing surgery for hip fracture what is the clinical and cost effectiveness of 'OEDP 10A rating' designs of stems in preference to Austin Moore or Thompson stems when inserting a hemiarthroplasty on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture
Extracapsular fractures	In patients undergoing repair for trochanteric extracapsular hip fractures what is the clinical and cost effectiveness of extramedullary sliding hip screws compared to intramedullary nails on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation (operative or postoperative fracture of the femur, cut-out and non-union) ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Wound healing complications
Extracapsular fractures	In patients undergoing repair for subtrochanteric extracapsular hip fractures, what is the effectiveness of extramedullary sliding hip screws compared to intramedullary nails on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation (operative or postoperative fracture of the femur, cut-out and non-union) ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Wound healing complications

Mobilisation strategies	In patients who have undergone surgery for hip fracture, what is the clinical and cost effectiveness of early mobilisation (<48 hours after surgery) compared to late mobilisation on functional status, mortality, place of residence/discharge, pain and quality of life?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Discharge destination
Mobilisation strategies	In patients who have undergone surgery for hip fracture, what is the clinical and cost effectiveness of intensive physiotherapy compared to non intensive physiotherapy on functional status, mortality, place of residence/discharge, pain and quality of life?	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Discharge destination ▪ Mobility
Multidisciplinary rehabilitation	In patients with hip fracture what is the clinical and cost effectiveness of 'orthogeriatrician' involvement in the whole pathway of assessment, peri-operative care and rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge ▪ Place of residence (compared with baseline) 12 months after fracture ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Multidisciplinary rehabilitation	In patients with hip fracture what is the clinical and cost effectiveness of hospital-based multidisciplinary rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge ▪ Place of residence (compared with baseline) 12 months after fracture ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)

Multidisciplinary rehabilitation	In patients with hip fracture what is the clinical and cost effectiveness of community-based multidisciplinary rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge ▪ Place of residence (compared with baseline) 12 months after fracture ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Carer involvement	In patients who have been discharged after hip fracture repair, what is the clinical and cost effectiveness of having a non paid carer (e.g. spouse, relative, friends) on mortality, length of stay, place of residence/discharge, functional status, hospital readmission and quality of life?	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge ▪ Place of residence (compared with baseline) 12 months after fracture ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)

1

2

3

3.2 Searching for evidence

4

3.2.1 Clinical literature search

5

Systematic literature searches were undertaken to identify evidence within published literature in order to answer the review questions as per The Guidelines Manual²³³.

6

Clinical databases were searched using relevant medical subject headings, free-text terms and study type filters where appropriate. Studies published in languages other than English were not reviewed. Where possible, searches were restricted to articles published in English language. All searches were conducted on core databases, MEDLINE, Embase and *The Cochrane Library*. Additional subject specific databases were used for some questions: PsycInfo for patient views and patient education questions; Cinahl for every question except those on anaesthesia, analgesia and the surgical procedures. All searches were updated on the 31st August 2010. No papers after this date were considered.

7

8

9

10

11

12

13

14

15

1 Search strategies were checked by looking at reference lists of relevant key papers,
2 checking search strategies in other systematic reviews and asking the GDG for known
3 studies. The questions, the study types applied, the databases searched and the years
4 covered can be found in Appendix D.

5 During the scoping stage, a search was conducted for guidelines and reports on the
6 websites listed below and on organisations relevant to the topic. Searching for grey
7 literature or unpublished literature was not undertaken. All references sent by
8 stakeholders were considered.

- 9 • Guidelines International Network database (www.g-i-n.net)
- 10 • National Guideline Clearing House (www.guideline.gov/)
- 11 • National Institute for Health and Clinical Excellence (NICE) (www.nice.org.uk)
- 12 • National Institutes of Health Consensus Development Program (consensus.nih.gov/)
- 13 • NHS Evidence (www.evidence.nhs.uk/)

14

15 **3.2.2 Health economic literature search**

16 Systematic literature searches were also undertaken to identify health economic evidence
17 within published literature relevant to the review questions. The evidence was identified by
18 conducting a broad search relating to the guideline population in the NHS economic
19 evaluation database (NHS EED) and health technology assessment (HTA) database with no
20 date restrictions. Additionally, the search was run on MEDLINE and Embase, with a specific
21 economic filter, to ensure recent publications that had not yet been indexed by these
22 databases were identified. This was supplemented by additional searches that looked for
23 economic papers specifically relating to the radiological imaging question on MEDLINE,
24 Embase, NHS EED and HTA databases, and the Health Economic Evaluations Database
25 (HEED) as it became apparent that some papers in this area were not being identified
26 through the first search. Studies published in languages other than English were not
27 reviewed. Where possible, searches were restricted to articles published in English
28 language.

29 The search strategies for health economics are included in Appendix D. All searches were
30 updated on the 31st August 2010. No papers published after this date were considered.

31

32 **3.3 Evidence of effectiveness**

33 The Research Fellow

- 34 • Identified potentially relevant studies for each review question from the relevant search
35 results by reviewing titles and abstracts – full papers were then obtained.
- 36 • Reviewed full papers against pre-specified inclusion / exclusion criteria to identify studies
37 that addressed the review question in the appropriate population and reported on
38 outcomes of interest (review protocols are included in Appendix C).

- 1 • Critically appraised relevant studies using the appropriate checklist as specified in The
2 Guidelines Manual²³³.
- 3 • Extracted key information about the study's methods and results into evidence tables
4 (evidence tables are included in Appendix E).
- 5 • Generated summaries of the evidence by outcome (included in the relevant chapter
6 write-ups):
- 7 ○ Randomised studies: meta analysed, where appropriate and reported in GRADE
8 profiles (for clinical studies) – see below for details
 - 9 ○ Observational studies: data presented as a range of values in GRADE profiles
 - 10 ○ Diagnostic studies: data presented as a range of values in adapted GRADE profiles
 - 11 ○ Qualitative studies: each study summarised in a table where possible, otherwise
12 presented in a narrative.

13

14 3.3.1 Inclusion/exclusion

15 See the review protocols in Appendix C for full details.

16

17 3.3.2 Methods of combining clinical studies

18 *Data synthesis for intervention reviews*

19 Where possible, meta-analyses were conducted to combine the results of studies for each
20 review question using Cochrane Review Manager (RevMan5) software. Fixed-effects
21 (Mantel-Haenszel) techniques were selected to calculate risk ratios (relative risk) for the
22 binary outcomes. The continuous outcomes were analysed using an inverse variance
23 method for pooling weighted mean differences and where the studies had different scales,
24 standardised mean differences were used.

25 Statistical heterogeneity was assessed by considering the chi-squared test for significance
26 at $p < 0.05$ or an I-squared inconsistency statistic of $> 50\%$ to indicate significant
27 heterogeneity. Where significant heterogeneity was present, we carried out predefined
28 subgroup analyses as defined in the protocol for each question (Appendix C). Sensitivity
29 analysis based on the quality of studies was also carried out if there were differences, with
30 particular attention paid to allocation concealment, blinding and loss to follow-up (missing
31 data).

32 Assessments of potential differences in effect between subgroups were based on the chi-
33 squared tests for heterogeneity statistics between subgroups. If no sensitivity analysis was
34 found to completely resolve statistical heterogeneity then a random effects (DerSimonian
35 and Laird) model was employed to provide a more conservative estimate of the effect.

36 For binary outcomes, absolute event rates were also calculated using the GRADEpro
37 software using event rate in the control arm of the pooled results.

1 *Data synthesis for diagnostic test accuracy review*

2 For diagnostic test accuracy studies, the following outcomes were reported: sensitivity,
3 specificity, positive predictive value, negative predictive value and positive and negative
4 likelihood ratios. In cases where the outcomes were not reported, 2 by 2 tables were
5 constructed from raw data to allow calculation of these accuracy measures. Summary
6 receiver operative characteristic (ROC) curves were not generated as we did not explore the
7 effect of different cut-off thresholds on sensitivity and specificity for the imaging questions.

8

9 **3.3.3 Appraising the quality of evidence by outcomes**

10 The evidence for outcomes from the included RCT and observational studies were
11 evaluated and presented using an adaptation of the 'Grading of Recommendations
12 Assessment, Development and Evaluation (GRADE) toolbox' developed by the international
13 GRADE working group (<http://www.gradeworkinggroup.org/>). The software (GRADEpro)
14 developed by the GRADE working group was used to assess the quality of each outcome,
15 taking into account individual study quality and the meta-analysis results. The summary of
16 findings was presented as two separate tables in this guideline. The "Clinical/Economic
17 Study Characteristics" table includes details of the quality assessment while the "Clinical
18 /Economic Summary of Findings" table includes pooled outcome data, where appropriate,
19 an absolute measure of intervention effect and the summary of quality of evidence for that
20 outcome. In this table, the columns for intervention and control indicate the sum of the
21 sample size for continuous outcomes. For binary outcomes such as number of patients with
22 an adverse event, the event rates (n/N: number of patients with events divided by sum of
23 number of patients) are shown with percentages. Reporting or publication bias was only
24 taken into consideration in the quality assessment and included in the Clinical Study
25 Characteristics table if it was apparent. Each outcome was examined separately for the
26 quality elements listed and defined in Table 3-1 and each graded using the quality levels
27 listed in

28 Table 3-2. The main criteria considered in the rating of these elements are discussed below
29 (see section 3.3.4 Grading of Evidence). Footnotes were used to describe reasons for
30 grading a quality element as having serious or very serious problems. The ratings for each
31 component were summed to obtain an overall assessment for each outcome.

32 Table 3-3. The GRADE toolbox is currently designed only for randomised trials and
33 observational studies but we adapted the quality assessment elements and outcome
34 presentation for diagnostic accuracy studies.

1 **Table 3-1: Descriptions of quality elements in GRADE for intervention studies**

Quality element	Description
Limitations	Limitations in the study design and implementation may bias the estimates of the treatment effect. Major limitations in studies decrease the confidence in the estimate of the effect
Inconsistency	Inconsistency refers to an unexplained heterogeneity of results
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question, or recommendation made
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of the effect relative to the clinically important threshold
Publication bias	Publication bias is a systematic underestimate or an overestimate of the underlying beneficial or harmful effect due to the selective publication of studies

2

3 **Table 3-2: Levels for quality elements in GRADE**

Level	Description
None	There are no serious issues with the evidence
Serious	The issues are serious enough to downgrade the outcome evidence by one level
Very serious	The issues are serious enough to downgrade the outcome evidence by two levels

4

5 **Table 3-3: Overall quality of outcome evidence in GRADE**

Level	Description
High	Further research is <i>very unlikely</i> to change our confidence in the <i>estimate of effect</i>
Moderate	Further research is <i>likely</i> to have an important impact on our confidence in the <i>estimate of effect</i> and may change the estimate
Low	Further research is <i>very likely</i> to have an important impact on our confidence in the estimate of effect and is likely to change the estimate
Very low	<i>Any estimate of effect is very uncertain</i>

6

7 **3.3.4 Grading the quality of clinical evidence**

8 After results were pooled, the overall quality of evidence for each outcome was considered.

9 The following procedure was adopted when using GRADE:

- 10 1. A quality rating was assigned, based on the study design. RCTs start HIGH and
11 observational studies as LOW, uncontrolled case series as LOW or VERY LOW
- 12 2. The rating was then downgraded for the specified criteria: Study limitations,
13 inconsistency, indirectness, imprecision and reporting bias. These criteria are detailed
14 below. Observational studies were upgraded if there was: a large magnitude of effect,
15 dose-response gradient, and if all plausible confounding would reduce a
16 demonstrated effect or suggest a spurious effect when results showed no effect. Each

1 quality element considered to have “serious” or “very serious” risk of bias were rated
 2 down -1 or -2 points respectively.
 3
 4 3. The downgraded/upgraded marks were then summed and the overall quality rating
 5 was revised. For example, all RCTs started as HIGH and the overall quality became
 6 MODERATE, LOW or VERY LOW if 1, 2 or 3 points were deducted respectively.
 7 4. The reasons or criteria used for downgrading were specified in the footnotes.
 8 The details of criteria used for each of the main quality element are discussed further in the
 9 following sections 4.3.5 to 4.3.8.

10 3.3.5 Study limitations

11 The main limitations for randomised controlled trials are listed in Table 3-4.

12 The GDG accepted that investigator blinding in surgical intervention studies was impossible
 13 and participant blinding was also impossible to achieve in most situations. Therefore, open-
 14 label studies for surgery were not downgraded in the quality rating across the guideline.
 15 Studies were downgraded for unclear or inadequate allocation concealment. .

16 Table 3-4 lists the limitations considered for randomised controlled trials.

17 **Table 3-4: Study limitations of randomised controlled trials**

<i>Limitation</i>	<i>Explanation</i>
Allocation concealment	Those enrolling patients are aware of the group to which the next enrolled patient will be allocated (major problem in “pseudo” or “quasi” randomised trials with allocation by day of week, birth date, chart number etc.)
Lack of blinding	Patient, caregivers, those recording outcomes, those adjudicating outcomes, or data analysts are aware of the arm to which patients are allocated
Incomplete accounting of patients and outcome events	Loss to follow-up not accounted and failure to adhere to the intention to treat principle when indicated
Selective outcome reporting	Reporting of some outcomes and not others on the basis of the results
Other limitations	For example: <ul style="list-style-type: none"> • stopping early for benefit observed in randomised trials, in particular in the absence of adequate stopping rules • use of unvalidated patient-reported outcomes • carry-over effects in cross-over trials • recruitment bias in cluster-randomised trials

18

19 3.3.6 Inconsistency

20 Inconsistency refers to an unexplained heterogeneity of results. When estimates of the
 21 treatment effect across studies differ widely (i.e. heterogeneity or variability in results), this
 22 suggests true differences in underlying treatment effect. When heterogeneity was

measured at either Chi square $p < 0.05$ or I-squared inconsistency statistic of $> 50\%$, but no plausible explanation can be found, the quality of evidence was downgraded by one or two levels, depending on the extent of uncertainty to the results contributed by the inconsistency in the results. In addition to the I-square and Chi square values, the decision for downgrading was also dependent on factors such as whether the intervention is associated with benefit in all other outcomes or whether the uncertainty about the magnitude of benefit (or harm) of the outcome showing heterogeneity would influence the overall judgment about net benefit or harm (across all outcomes).

If inconsistency could be explained based on prespecified subgroup analysis, the GDG took this into account and considered whether to make separate recommendations based on the identified explanatory factors, i.e. population and intervention. Where subgroup analysis gives a plausible explanation of heterogeneity, the quality of evidence would not be downgraded.


3.3.7 Indirectness

Directness refers to the extent to which the populations, intervention, comparisons and outcome measures are similar to those defined in the inclusion criteria for the reviews. Indirectness is important when these differences are expected to contribute to a difference in effect size, or may affect the balance of harms and benefits considered for an intervention.

3.3.8 Imprecision

The sample size, event rates and the resulting width of confidence intervals were the main criteria considered. Where the minimal important difference (MID) of an outcome is known, the optimal information size (OIS), i.e. the sample size required to detect the difference with 80% power and $p \leq 0.05$ was calculated and used as the criteria. The criteria applied for imprecision are based on the confidence intervals for pooled or the best estimate of effect as illustrated in Figure 3-1 and outlined in **Error! Reference source not found.**

Figure 3-1: Illustration of precise and imprecise outcomes based on the confidence interval of outcomes in a forest plot



1

2 *MID = minimal important difference determined for each outcome. The MIDs are the threshold for*
3 *appreciable benefits and harms. The confidence intervals of the top three points of the diagram were*
4 *considered precise because the upper and lower limits did not cross the MID. Conversely, the bottom three*
5 *points of the diagram were considered imprecise because all of them crossed the MID and reduced our*
6 *certainty of the results. Figure adapted from GRADEPro software.*

7

8 The following are the MID for the outcomes and the methods used to calculate the OIS in
9 this guideline:

- 10
- Any statistically significant difference in mortality
- 11
- The default confidence intervals in GRADE for relative risk of 0.75 and 1.25 for all
- 12 other outcomes.

13

14 **3.4 Evidence of cost-effectiveness**

15 Evidence on cost-effectiveness related to the key clinical issues being addressed in the guideline
16 was sought. The health economist:

- 17
- Undertook a systematic review of the economic literature
- 18
- Undertook new cost-effectiveness analysis in priority areas

19

20 **3.4.1 Literature review**

21 The Health Economist:

- 22
- Identified potentially relevant studies for each review question from the economic search
- 23 results by reviewing titles and abstracts – full papers were then obtained.
- 24
- Reviewed full papers against pre-specified inclusion / exclusion criteria to identify
- 25 relevant studies (see below for details).
- 26
- Critically appraised relevant studies using the economic evaluations checklist as specified
- 27 in The Guidelines Manual²³³.
- 28
- Extracted key information about the study's methods and results into evidence tables
- 29 (evidence tables are included in Appendix F).
- 30
- Generated summaries of the evidence in NICE economic evidence profiles – see below for
- 31 details.

32

33 **3.4.1.1 Inclusion/exclusion**

- 1 Full economic evaluations (cost-effectiveness, cost-utility, cost-benefit and cost-
2 consequence analyses) and comparative costing studies that addressed the review
3 question in the relevant population were considered potentially applicable as economic
4 evidence.
- 5 Studies that only reported cost per hospital (not per patient), or only reported average
6 cost effectiveness without disaggregated costs and effects, were excluded. However,
7 studies reporting the cost per hospital were included when it was possible to ascertain the
8 cost per patient of each intervention. Abstracts, posters, reviews, letters/editorials,
9 foreign language publications and unpublished studies were excluded. Studies judged to
10 have had an applicability rating of 'not applicable' were excluded (this included studies
11 that took the perspective of a non-OECD country).
- 12 Remaining studies were prioritised for inclusion based on their relative applicability to the
13 development of this guideline and the study limitations. For example, if a high quality,
14 directly applicable UK analysis was available other less relevant studies may not have
15 been included. Where exclusions occurred on this basis, this is noted in the relevant
16 section.
- 17 For more details about the assessment of applicability and methodological quality see the
18 economic evaluation checklist (The Guidelines Manual, Appendix H²³³ and the health
19 economics research protocol in Appendix C.
- 20 When no relevant economic analysis was found from the economic literature review,
21 relevant UK NHS unit costs related to the compared interventions were presented to the
22 GDG to inform the possible economic implication of the recommendation to make.
- 23
- 24 **3.4.2 NICE economic evidence profiles**
- 25 The NICE economic profile has been used to summarise cost and cost-effectiveness
26 estimates. The economic evidence profile shows, for each economic study, an assessment
27 of applicability and methodological quality, with footnotes indicating the reasons for the
28 assessment. These assessments were made by the health economist using the economic
29 evaluation checklist from The Guidelines Manual, Appendix H²³³. It also shows
30 incremental costs, incremental outcomes (e.g. QALYs) and the incremental cost-
31 effectiveness ratio from the primary analysis, as well as information about the assessment
32 of uncertainty in the analysis. See Table 3-5 for more details.
- 33 If a non-UK study was included in the profile, the results were converted into pounds
34 sterling using the appropriate purchasing power parity²⁴⁶.

35 **Table 3-5: Content of NICE economic profile**

<i>Item</i>	<i>Description</i>
Study	First author name, reference, date of study publication and country perspective.
Limitations	An assessment of methodological quality of the study*: <ul style="list-style-type: none"> • Minor limitations – the study meets all quality criteria, or the study fails to meet one or more quality criteria, but this is unlikely to change the conclusions about cost effectiveness. • Potentially serious limitations – the study fails to meet one or more quality criteria, and this could change the conclusion about cost

	effectiveness <ul style="list-style-type: none"> • Very serious limitations – the study fails to meet one or more quality criteria and this is very likely to change the conclusions about cost effectiveness. Studies with very serious limitations would usually be excluded from the economic profile table.
Applicability	An assessment of applicability of the study to the clinical guideline, the current NHS situation and NICE decision-making*: <ul style="list-style-type: none"> • Directly applicable – the applicability criteria are met, or one or more criteria are not met but this is not likely to change the conclusions about cost effectiveness. • Partially applicable – one or more of the applicability criteria are not met, and this might possibly change the conclusions about cost effectiveness. • Not applicable – one or more of the applicability criteria are not met, and this is likely to change the conclusions about cost effectiveness.
Other comments	Particular issues that should be considered when interpreting the study.
Incremental cost	The mean cost associated with one strategy minus the mean cost of a comparator strategy.
Incremental effects	The mean QALYs (or other selected measure of health outcome) associated with one strategy minus the mean QALYs of a comparator strategy.
ICER	Incremental cost-effectiveness ratio: the incremental cost divided by the respective QALYs gained
Uncertainty	A summary of the extent of uncertainty about the ICER reflecting the results of deterministic or probabilistic sensitivity analyses, or stochastic analyses of trial data, as appropriate.

1 *Limitations and applicability were assessed using the economic evaluation checklist from The Guidelines
2 Manual, Appendix H²³³

3 When no cost-effectiveness evidence was available, the cost of the interventions being
4 evaluated has in some cases been determined by conducting original cost analyses there
5 were reported in Appendix H. Alternatively, the GDG was presented with the cost figures
6 from relevant sources, such as the NHS reference cost for England and Wales.

7 3.4.3 Undertaking new health economic analysis

8 As well as reviewing the published economic literature for each review question, as
9 described above, new economic analyses were undertaken by the Health Economist in
10 priority areas. Priority areas for new health economic analysis were agreed by the GDG
11 after formation of the review questions and consideration of the available health
12 economic evidence.

13 Additional data for the analysis was identified as required through additional literature
14 searches undertaken by the Health Economist, and discussion with the GDG. Model
15 structure, inputs and assumptions were explained to and agreed by the GDG members
16 during meetings, and they commented on subsequent revisions.

17 See Appendix H for details of the health economic analyses undertaken for the guideline.

18

1 3.4.4 Cost-effectiveness criteria

2 NICE's report 'Social value judgements: principles for the development of NICE
3 guidance'²³² sets out the principles that GDGs should consider when judging whether an
4 intervention offers good value for money. In general, an intervention was considered to
5 be cost effective if either of the following criteria applied (given that the estimate was
6 considered plausible):

7 a) The intervention dominated other relevant strategies (that is, it was both less costly
8 in terms of resource use and more clinically effective compared with all the other
9 relevant alternative strategies), **or**

10 b) The intervention cost less than £20,000 per quality-adjusted life-year (QALY) gained
11 compared with the next best strategy.

12 If the GDG recommended an intervention that was estimated to cost more than £20,000
13 per QALY gained, or did not recommend one that was estimated to cost less than £20,000
14 per QALY gained, the reasons for this decision are discussed explicitly in the 'from
15 evidence to recommendations' section of the relevant chapter. This is written with
16 reference to the issues regarding the plausibility of the estimate or to the factors set out
17 in the Social value judgements report ²³².

18

19

20 3.5 Developing recommendations

21 Over the course of the guideline development process, the GDG was presented with:

- 22 • Evidence tables of the clinical evidence (Appendix E) and economic evidence
23 (Appendix F) reviewed from the literature.
- 24 • Summary of clinical and economic evidence and quality (as presented in chapters 5
25 to 13).
- 26 • Forest plots (Appendix G)
- 27 • A description of the methods and results of the cost-effectiveness analysis
28 undertaken for the guideline (Appendix H)

29 Recommendations were drafted on the basis of the GDG interpretation of the available
30 evidence, taking into account the balance of benefits, harms and costs. When clinical and
31 economic evidence was of poor quality, conflicting or absent, the GDG drafted
32 recommendations based on their expert opinion. The considerations for making consensus
33 based recommendations include the balance between potential harms and benefits,
34 economic or implications compared to the benefits, current practices, recommendations
35 made in other relevant guidelines, patient preferences and equality issues. The consensus
36 recommendations were done through discussions in the GDG.

1 **3.5.1 Research recommendations**

2 When areas were identified for which good evidence was lacking, the guideline development
3 group considered making recommendations for future research. Decisions about inclusion
4 were based on factors such as:

- 5 • the importance to patients or the population
- 6 • national priorities
- 7 • potential impact on the NHS and future NICE guidance
- 8 • ethical and technical feasibility

9

10 **3.6 Validation process**

11 The guidance is subject to an eight week public consultation and feedback as part of the
12 quality assurance and peer review of the document. All comments received from registered
13 stakeholders are responded to in turn and posted on the NICE website when the pre-
14 publication check of the full guideline occurs.

15

16 **3.7 Updating the guideline**

17 Following publication, and in accordance with the NICE guidelines manual, NICE will ask a
18 National Collaborating Centre or the National Clinical Guideline Centre to advise NICE's
19 Guidance executive whether the evidence base has progressed significantly to alter the
20 guideline recommendations and warrant an update.

21

22 **3.8 Disclaimer**

23 Health care providers need to use clinical judgement, knowledge and expertise when deciding whether
24 it is appropriate to apply guidelines. The recommendations cited here are a guide and may not be
25 appropriate for use in all situations. The decision to adopt any of the recommendations cited here
26 must be made by the practitioners in light of individual patient circumstances, the wishes of the
27 patient, clinical expertise and resources.

28 The National Clinical Guideline Centre disclaim any responsibility for damages arising out of the use or
29 non-use of these guidelines and the literature used in support of these guidelines.

30

31 **3.9 Funding**

32 The National Clinical Guideline Centre was commissioned by the National Institute for Health
33 and Clinical Excellence to undertake the work on this guideline.

34

1 **4 Guideline summary**

2 **4.1 Map of recommendations**

3 An algorithm will be added before publication.

1 **Key priorities for implementation**

2 The GDG identified ten key priorities for implementation. The decision was made after
3 discussion and voting by the GDG. They selected recommendations that would:

- 4 • Have a high impact on outcomes that are important to patients (**A**)
- 5 • Have a high impact on reducing variation in care and outcomes (**B**)
- 6 • Lead to a more efficient use of NHS resources (**C**)
- 7 • Promote patient choice (**D**)
- 8 • Promote equalities (**E**)
- 9 • Mean patients reach critical points in the care pathway more quickly (**F**).

10 In doing this the GDG also considered which recommendations were particularly likely to
11 benefit from implementation support. They considered whether a recommendation:

- 12 • Requires changes in service delivery (**W**)
- 13 • Requires retraining of professionals or the development of new skills and
14 competencies (**X**)
- 15 • Affects and needs to be implemented across various agencies or settings (complex
16 interactions) (**Y**)
- 17 • May be viewed as potentially contentious, or difficult to implement for other
18 reasons (**Z**)

19 For each key recommendation listed below, the selection criteria and implementation
20 support points are indicated by the use of the letters shown in brackets above and are
21 shown in the linking evidence to recommendations sections in the relevant chapters.

- 22 ➤ Perform surgery on the day of, or the day after, admission. (A, B, C, F, W, Y and Z).
- 23 ➤ Identify and treat correctable comorbidities immediately so that surgery is not
24 delayed by:
 - 25 • anaemia
 - 26 • anticoagulation
 - 27 • volume depletion
 - 28 • electrolyte imbalance
 - 29 • uncontrolled diabetes
 - 30 • uncontrolled heart failure
 - 31 • correctable cardiac arrhythmia or ischaemia

- 1
 - acute chest infection
- 2
 - exacerbation of chronic chest conditions (A, B, C, F, Y and Z).
- 3
 - Schedule hip fracture surgery on a planned trauma list (A, B, C, F, W, and Z).
- 4
 - Perform replacement arthroplasty (hemiarthroplasty or total hip replacement) in
- 5
 - patients with a displaced intracapsular fracture (A, B, C, F and Z).
- 6
 - Offer total hip replacements to patients with a displaced intracapsular fracture who:
- 7
 - were able to walk independently out of doors with no more than the use of a
- 8
 - stick and
- 9
 - are not cognitively impaired and
- 10
 - are medically fit for anaesthesia and the procedure (A, B, C, X, and Z).
- 11
 - Use extramedullary implants such as a sliding hip screw in preference to an
- 12
 - intramedullary nail in patients with trochanteric fractures above and including the
- 13
 - lesser trochanter (AO classification types A1 and A2) (A, B, C, and Z).
- 14
 - Offer patients a physiotherapy assessment and, unless medically or surgically
- 15
 - contraindicated, mobilisation on the day after surgery (A, B, C, D, E, F, W, X, Y and Z).
- 16
 - Offer patients mobilisation at least once a day and ensure regular physiotherapy
- 17
 - review (A, B, F, and W).
- 18
 - From admission, offer patients a formal, acute, orthogeriatric or orthopaedic ward-
- 19
 - based Hip Fracture Programme that includes all of the following:
- 20
 - orthogeriatric assessment
- 21
 - rapid optimisation of fitness for surgery
- 22
 - early identification of individual goals for multidisciplinary rehabilitation to
- 23
 - recover mobility and independence, and to facilitate return to prefracture
- 24
 - residence and long-term well-being.
- 25
 - continued co-ordinated orthogeriatric and multidisciplinary review
- 26
 - liaison or integration with related services, particularly mental health, falls
- 27
 - prevention, bone health, primary care and social services.
- 28
 - clinical and service governance responsibility for all stages of the pathway of
- 29
 - care and rehabilitation, including those delivered in the community.
- 30
 - (A,B,C,D,E,F,W,X,Y and Z).
- 31
 - Consider early supported discharge as part of the Hip Fracture Programme, provided
- 32
 - the Hip Fracture Programme multidisciplinary team remains involved, and the
- 33
 - patient:
- 34
 - is medically stable and

- 1 • has the mental ability to participate in continued rehabilitation and
- 2 • is able to transfer and mobilise short distances and
- 3 • has not yet achieved their full rehabilitation potential, as discussed with the
- 4 patient, carer and family (A,B,C, E,F,W, and Z).

5

6 **4.2 Full list of recommendations**

7 Some aspects of hip fracture management are already covered by NICE guidance
 8 and are therefore outside the scope of this guideline. In order to ensure
 9 comprehensive management and continuity, the following NICE guidance should be
 10 referred to when developing a complete programme of care for each patient::
 11 osteoporotic fragility fracture prevention (TA 160, 161 & 204)²³⁴⁻²³⁶, falls (CG21)²²⁷, pressure
 12 ulcers (CG29)²²⁸, nutrition support (CG32)²²⁹, dementia (CG42)²³⁰, surgical site infection
 13 (CG74)²³¹, venous thromboembolism (CG92)²³⁷ and delirium (CG103)²³⁰.

14 **4.2.1 Imaging options in occult hip fracture**

- 15 ➤ Offer magnetic resonance imaging (MRI) if hip fracture is suspected despite negative
- 16 anteroposterior pelvis and lateral hip X-rays. If MRI is not available within 24 hours or
- 17 is contraindicated, consider computed tomography (CT).

18 **4.2.2 Timing of surgery**

- 19 ➤ Perform surgery on the day of, or the day after, admission.
- 20 ➤ Identify and treat correctable comorbidities immediately so that surgery is not
- 21 delayed by:
 - 22 • anaemia
 - 23 • anticoagulation
 - 24 • volume depletion
 - 25 • electrolyte imbalance
 - 26 • uncontrolled diabetes
 - 27 • uncontrolled heart failure
 - 28 • correctable cardiac arrhythmia or ischaemia
 - 29 • acute chest infection
 - 30 • exacerbation of chronic chest conditions.

31 **4.2.3 Analgesia**

- 32 ➤ Assess the patient's pain:

- 1 • immediately upon presentation at hospital and
- 2 • within 30 minutes of administering initial analgesia and
- 3 • hourly until settled on the ward and
- 4 • regularly as part of routine nursing observations throughout admission.
- 5 ➤ Offer immediate analgesia to patients presenting at hospital with suspected hip
6 fracture, including people with cognitive impairment.
- 7 ➤ Ensure analgesia is sufficient to allow movements necessary for investigations (as
8 indicated by the ability to tolerate passive external rotation of the leg), and for
9 nursing care and rehabilitation.
- 10 ➤ Offer paracetamol every 6 hours preoperatively unless contraindicated.
- 11 ➤ Offer additional opioids if paracetamol alone does not provide sufficient preoperative
12 pain relief.
- 13 ➤ Consider adding nerve blocks if paracetamol and opioids do not provide sufficient
14 preoperative pain relief, or to limit opioid dosage. Nerve blocks should be
15 administered by trained personnel. Do not use nerve blocks as a substitute for early
16 surgery.
- 17 ➤ Offer paracetamol every 6 hours postoperatively unless contraindicated.
- 18 ➤ Offer additional opioids if paracetamol alone does not provide sufficient
19 postoperative pain relief.
- 20 ➤ Non steroidal anti-inflammatory drugs (NSAIDs) are not recommended.

21 **4.2.4 Anaesthesia**

- 22 ➤ Offer patients a choice of spinal or general anaesthesia after discussing the risks and
23 benefits.
- 24 ➤ Consider intraoperative nerve blocks for all patients undergoing surgery.

25 **4.2.5 Planning the theatre team**

- 26 ➤ Schedule hip fracture surgery on a planned trauma list.
- 27 ➤ Consultants or senior staff should supervise trainee and junior members of the
28 anaesthesia, surgical and theatre teams when they carry out hip fracture procedures.

29 **4.2.6 Surgical procedures**

- 30 ➤ Operate on patients with the aim to allow them to fully weight bear (without
31 restriction) in the immediate postoperative period.
- 32 ➤ Perform replacement arthroplasty (hemiarthroplasty or total hip replacement) in
33 patients with a displaced intracapsular fracture.

- 1 ➤ Offer total hip replacement to patients with a displaced intracapsular fracture who:
- 2 • were able to walk independently out of doors with no more than the use of a
- 3 stick and
- 4 • are not cognitively impaired and
- 5 • are medically fit for anaesthesia and the procedure
- 6 ➤ Use a proven femoral stem design rather than Austin Moore or Thompson stems for
- 7 arthroplasties. Suitable designs include those with an Orthopaedic Data Evaluation
- 8 Panel rating of 10A, 10B, 10C, 7A, 7B, 5A, 5B, 3A or 3B.
- 9 ➤ Use cemented implants in patients undergoing surgery with arthroplasty.
- 10 ➤ Consider an anterolateral approach in favour of a posterior approach when inserting a
- 11 hemiarthroplasty.
- 12 ➤ Use extramedullary implants such as a sliding hip screw in preference to an
- 13 intramedullary nail in patients with trochanteric fractures above and including the
- 14 lesser trochanter (AO classification types A1 and A2).
- 15 ➤ Use an intramedullary nail to treat patients with a subtrochanteric fracture.

16 **4.2.7 Mobilisation strategies**

- 17 ➤ Offer patients a physiotherapy assessment and, unless medically or surgically
- 18 contraindicated, mobilisation on the day after surgery.
- 19 ➤ Offer patients mobilisation at least once a day and ensure regular physiotherapy
- 20 review.

21 **4.2.8 Multidisciplinary management**

- 22 ➤ From admission, offer patients a formal, acute orthogeriatric or orthopaedic ward-
- 23 based Hip Fracture Programme that includes all of the following:
- 24 • orthogeriatric assessment
- 25 • rapid optimisation of fitness for surgery
- 26 • early identification of individual goals for multidisciplinary rehabilitation to
- 27 recover mobility and independence, and to facilitate return to prefracture
- 28 residence and long-term wellbeing.
- 29 • continued, coordinated, orthogeriatric and multidisciplinary review
- 30 • liaison or integration with related services, particularly mental health, falls
- 31 prevention, bone health, primary care and social services.
- 32 • clinical and service governance responsibility for all stages of the pathway of
- 33 care and rehabilitation, including those delivered in the community.

- 1 ➤ If a hip fracture complicates or precipitates a terminal illness, the multidisciplinary
2 team should still consider the role of surgery, as part of a palliative care approach
3 that:
- 4 • minimises pain and other symptoms and
- 5 • establishes patients' own priorities for rehabilitation and
- 6 • considers patients' wishes about their end-of-life care.
- 7 ➤ Healthcare professionals should deliver care that minimises the patient's risk of
8 delirium and maximises their independence, by:
- 9 • actively looking for cognitive impairment when patients first present with hip
10 fracture
- 11 • reassessing patients to identify delirium that may arise during their admission
- 12 • offering individualised care in line with 'Delirium' (NICE clinical guideline 103).
- 13 ➤ Consider early supported discharge as part of the Hip Fracture Programme, provided
14 the Hip Fracture Programme multidisciplinary team remains involved, and the
15 patient:
- 16 • is medically stable and
- 17 • has the mental ability to participate in continued rehabilitation and
- 18 • is able to transfer and mobilise short distances and
- 19 • has not yet achieved their full rehabilitation potential, as discussed with the
20 patient, carer and family.
- 21 ➤ Only consider intermediate care (continued rehabilitation in a community hospital or
22 residential care unit) if all of the following criteria are met:
- 23 • intermediate care is included in the Hip Fracture Programme and
- 24 • the Hip Fracture Programme team retains the clinical lead, including patient
25 selection, agreement of length of stay and ongoing objectives for
26 intermediate care and
- 27 • the Hip Fracture Programme team retains the managerial lead, ensuring that
28 intermediate care is not resourced as a substitute for an acute hospital
29 Programme.
- 30 ➤ Patients admitted from care or nursing homes should not be excluded from
31 rehabilitation programmes in the community or hospital, or as part of an early
32 supported discharge programme.

1 **4.2.9 Patient and carer information**

2 ➤ Offer patients (or, as appropriate, their carer and/or family) verbal and printed
3 information about treatment and care including:

- 4 • diagnosis
- 5 • choice of anaesthesia
- 6 • choice of analgesia and other medications
- 7 • surgical procedures
- 8 • possible complications
- 9 • postoperative care
- 10 • rehabilitation programme
- 11 • long-term outcomes
- 12 • healthcare professionals involved.

13

14 **4.3 Research recommendations**

15 The GDG identified the following priority areas for research:

- 16 • Imaging options in occult hip fracture
- 17 • Anaesthesia
- 18 • Displaced intracapsular hip fracture
- 19 • Early supported discharge
- 20 • Physiotherapy

21 **4.3.1 Research recommendation on imaging options in occult hip fracture**

22 ➤ In patients with a continuing suspicion of a hip fracture but whose radiographs are
23 normal, what is the clinical and cost effectiveness of computed tomography (CT)
24 compared to magnetic resonance imaging (MRI), in confirming or excluding the
25 fracture?

26 **Why this is important**

27 The GDG's consensus decision to recommend CT over a radionuclide bone scan as an
28 alternative to MRI to detect occult hip fractures reflects current NHS practice but assumes
29 that advances in technology have made the reliability of CT comparable with that of MRI. If
30 modern CT can be shown to have similar reliability and accuracy to MRI, then this has
31 considerable implications because of its widespread availability out of hours and lower cost.

1 It is therefore a high priority to confirm or refute this assumption by direct randomised
2 comparison. The study design would need to retain MRI as the 'gold standard' for cases of
3 uncertainty and to standardise the criteria, expertise and procedures for radiological
4 assessment. Numbers required would depend on the degree of sensitivity and specificity
5 (the key outcome criteria) set as target requirement for comparability, but need not
6 necessarily be very large.

7 **4.3.2 Research recommendation on anaesthesia**

- 8 ➤ What is the clinical and cost effectiveness of regional versus general anaesthesia on
9 postoperative morbidity in patients with hip fracture?

10 **Why this is important**

11 No recent randomised controlled trials were identified that fully address this question. The
12 evidence is old and does not reflect current practice. In addition, in most of the studies the
13 patients are sedated before regional anaesthesia is administered, and this is not taken into
14 account when analysing the results. The study design for the proposed research would be
15 best addressed by a randomised controlled trial. This would ideally be a multi-centre trial
16 including 3000 participants in each arm. This is achievable given that there are about
17 70,000 to 75,000 hip fractures a year in the UK³⁹. The study should have three arms that
18 look at spinal anaesthesia versus spinal anaesthesia plus sedation versus general
19 anaesthesia; this would separate those with regional anaesthesia from those with regional
20 anaesthesia plus sedation. The study would also need to control for surgery, especially type
21 of fracture, prosthesis and grade of surgeon.

22 A qualitative research component would also be helpful to study patient preference for
23 type of anaesthesia.

24 **4.3.3 Research recommendation on displaced intracapsular hip fracture**

- 25 ➤ What is the clinical and cost effectiveness of large-head total hip replacement versus
26 hemiarthroplasty on functional status, reoperations and quality of life in patients with
27 displaced intracapsular hip fracture?

28 **Why this is important**

29 Large-head total hip replacement is a development of traditional total hip replacement,
30 where a larger head makes the joint more stable and hence reduces the risks of dislocation.
31 Three small trials have shown traditional small-head total hip replacement to have better
32 outcomes and function, albeit with an increased dislocation rate in selected groups of
33 patients. The drawback with large-head arthroplasty is the additional implant cost and
34 theatre time. This cost can account for up to 20% of current NHS tariff (up to £2000) and
35 the study aims to address whether this translates to improved patient outcome. The study
36 design for the proposed research would be best addressed by a randomised controlled trial.
37 This would have two arms to compare current standard care (using hemiarthroplasty) with
38 using large-head total hip replacement for patients sustaining displaced intracapsular hip
39 fractures. The primary outcome would be patient mobility at 1 year and secondary
40 outcomes would include functional outcomes, quality of life and cost effectiveness of the
41 intervention.

42 It would be expected that a sample size of approximately 500 patients would be required to
43 show a significant difference in the mobility, hip function and quality of life (assuming 80%

1 power, $p < 0.05$). By recruiting through a trauma research network it is estimated that 10
2 centres would be able to recruit 20 patients per month (from 45 eligible patients) giving a
3 recruitment period of 25 months.

4 **4.3.4 Research recommendation on early supported discharge**

5 ➤ What is the clinical and cost effectiveness of early supported discharge on mortality,
6 quality of life and functional status in patients with hip fracture who are admitted
7 from a care home?

8 **Why this is important**

9 Residents of care and nursing homes account for about 30% of all patients with hip fracture
10 admitted to hospital. Two-thirds of these come from care homes and the remainder from
11 nursing homes. These patients are frailer, more functionally dependent and have a higher
12 prevalence of cognitive impairment than patients admitted from their own homes. One-
13 third of those admitted from a care home are discharged to a nursing home and one-fifth
14 are readmitted to hospital within 3 months. There are no clinical trials to define the optimal
15 rehabilitation pathway following hip fracture for these patients and therefore represent a
16 discrete cohort where the existing meta-analyses do not apply. As a consequence, many
17 patients are denied structured rehabilitation and are discharged back to their care home or
18 nursing home with very little or no rehabilitation input.

19 Given the patient frailty and comorbidities, rehabilitation may have a limited effect on
20 clinical outcomes for this group. The fact that they already live in a home where they are
21 supported by trained care staff, however, clearly provides an opportunity for a systematic
22 approach to rehabilitation. Early multidisciplinary rehabilitation based in care homes
23 or nursing homes would take advantage of the day-to-day care arrangements already in
24 place and provide additional NHS support to deliver naturalistic rehabilitation, where
25 problems are tackled in the patient's residential setting.

26 Early supported multidisciplinary rehabilitation could reduce hospital stay, improve early
27 return to function, and affect both readmission rates and the level of NHS-funded nursing
28 care required.

29 The research would follow a two-stage design: (1) an initial feasibility study to refine the
30 selection criteria and process for reliable identification and characterisation of those
31 considered most likely to benefit, together with the intervention package and measures for
32 collaboration between the Hip Fracture Programme team, care-home staff and other
33 community-based professionals, and (2) a cluster randomized controlled comparison (with
34 two or more intervention units and matched control units) set against agreed outcome
35 criteria. The latter should include those specified above, together with measures of the
36 impact on care-home staff activity and cost, as well as qualitative data from patients on
37 relevant quality-of-life variables.

38 **4.3.5 Research recommendation on physiotherapy**

39 ➤ What is the clinical and cost effectiveness of additional intensive physiotherapy
40 and/or occupational therapy (for example progressive, resistance training) after hip
41 fracture?

42 **Why this is important**

1 The rapid restoration of physical and self care functions is a critical to recovery from hip
2 fracture, particularly where the goal is to return to the patient to preoperative levels of
3 function and residence. Approaches that are worthy of future development and
4 investigation include progressive resistance training, progressive balance and gait training,
5 supported treadmill gait re-training, dual task training, and activities of daily living training.
6 The optimal time point at which these interventions should be started requires clarification.

7 The ideal study design is a randomised controlled trial. Initial studies may have to focus on
8 proof of concept and be mindful of costs. A phase III randomised controlled trial is required
9 to determine clinical effectiveness and cost-effectiveness. The ideal sample size will be
10 around 400 to 500 patients, and the primary outcome should be physical function and
11 health related quality of life. Outcomes should also include falls. A formal sample size
12 calculation will need to be undertaken. Outcomes should be followed over a minimum of 1
13 year, and compare if possible, either the recovery curve for restoration of function or time
14 to attainment of functional goals.

15 **4.3.6 Additional research recommendations**

16 The following research questions were selected by the GDG but were not prioritised in the
17 top five recommendations for research.

18 **4.3.6.1 Analgesia**

19 The GDG recommended the following research question:

- 20 ➤ What is the clinical and cost effectiveness of preoperative and postoperative nerve
21 blocks in reducing pain and achieving mobilisation and physiotherapy goals sooner in
22 patients with hip fracture?

23 **Why this is important**

24 Nerve blocks may potentially find an important role in the management of hip fracture
25 pain, both pre- and postoperatively, because of their potential to reduce the requirement
26 for opioids and their associated unwanted effects. Economically there are considerations
27 for staff training, but also for the potential benefits in terms of duration of stay and early
28 mobilisation. It is not possible from the existing literature to determine this with any
29 confidence and there is a pressing need for a definitive trial comparing these outcomes
30 with nerve blocks against a defined protocol of systemic opioid use.

31 **4.3.6.2 Timing of surgery**

32 The GDG recommended the following research question:

- 33 ➤ What is the clinical and cost effectiveness of surgery within 36 hours of admission
34 compared to surgery later than 36 hours from admission in mortality, morbidity
35 and quality of life in patients with hip fracture?

36 **Why this is important**

37 Early and appropriate surgery for hip fractures is the most effective form of pain relief,
38 potentially quickening the rehabilitation and reducing complications. Within the current
39 literature no specific time interval threshold has been identified (up to 24hr) below which a
40 reduction in delay has shown no benefit. In addition to the evidence of the cost

1 effectiveness below 48hr, pragmatic, organisational and humanitarian considerations have
2 been utilised to arrive at the recommendation to operate not later than the day after
3 admission. A formal study within the NHS based on an arbitrary but realistic 36hr threshold
4 would provide additional important data to that already available, in order to inform more
5 precisely the forward clinical and cost-effectiveness of the strategy. For ethical reasons, the
6 research design would be an observational cohort study, correcting for confounding
7 variables, possibly set in the context of the National Hip Fracture Database and examining
8 the effect of the time to surgery and its cost on key outcomes, including mortality,
9 complications, length of stay, time taken to rehabilitate and qualitative aspects of the
10 experiences of patients.

11 **4.3.6.3 Reverse oblique trochanteric fractures**

12 The GDG recommended the following research question:

- 13 ➤ What is the clinical and cost effectiveness of intramedullary versus extramedullary
14 total hip replacement on mortality, functional status and quality of life in patients
15 with reverse oblique trochanteric hip fracture?

16 **Why this is important**

17 Reverse oblique trochanteric fractures account for approximately 5 % of all trochanteric hip
18 fractures. This means it affects approximately over 1000 patients per year in the UK.
19 Presently there is little evidence as to which is the preferable implant (which can be either
20 extramedullary – outside the bone, or intramedullary - inside the bone). The potential
21 biomechanical advantage of intramedullary advantage may be offset by increased cost
22 (which can be over £1000 more expensive). A randomised trial comparing the two implants
23 using patient mobility, function and re-operation would allow a more informed choice of
24 treatment for this injury.

25 **4.3.6.4 Designated hip fracture units**

26 The GDG recommended the following research question:

- 27 ➤ What is the clinical and cost effectiveness of a designated hip fracture unit within
28 the trauma ward compared to units integrated into acute trusts on mortality,
29 quality of life and functional status in patients with hip fracture?

30 **Why this is important**

31 The increasingly structured approach to hip fracture care has led to a number of UK units
32 considering or establishing a specific 'hip fracture ward' as a specialist part of their acute
33 orthopaedic service.

34 Designated hip fracture wards may prove an effective means of delivering the whole
35 programme of coordinated perioperative care and multidisciplinary rehabilitation which
36 this NICE Guidance has proposed, but at present there is no high quality evidence of their
37 clinical effectiveness when compared to such care within general orthopaedic or trauma
38 beds.

39 It may not be practical to run an RCT within a trauma unit, but there is certainly potential
40 for cohort studies to explore the effect of such units on individual patients' mobility,

1 discharge residence, mortality and length of stay. Units considering the establishment of
2 hip fracture wards should be encouraged to consider performing such trials.

3 **4.3.6.5 Care/nursing home residents**

4 The GDG recommended the following research question:

- 5 ➤ Do patients admitted to hospital with a fractured hip who live permanently in a
6 care/nursing home have equal access to multidisciplinary rehabilitation as patients
7 admitted from home?

8 **Why this is important**

9 The existing literature on the effectiveness of multidisciplinary rehabilitation typically
10 excludes patients who live in care/nursing homes. From an equality perspective it
11 hypothesised that this group of people do not have access to the same multidisciplinary
12 rehabilitation as patients who are returning home as it is assumed patients returning to
13 care/nursing homes will have their care needs met by the home. The research design would
14 be a prospective observational cohort study to determine the extent and quality of
15 rehabilitation services available to this group in comparison to patients returning to their
16 own homes.

17 **4.3.6.6 Patient and carer quality of life**

18 The GDG recommended the following research question:

- 19 ➤ What quality of life value do individual patients and their carers place on different
20 mobility, independence and residence states following rehabilitation?

21 **Why this is important**

22 It is important in evaluating future priorities for intervention to determine whether the
23 perceived clinical and health economic benefits of rehabilitation outcomes in the research
24 literature are matched over the same time-frame by the quality of life judgements,
25 aspirations and expectations of patients themselves and their carers. There is currently no
26 evidence.

27 **4.3.6.7 Patient experience**

28 The GDG recommended the following research question:

- 29 ➤ What is the patient's experience of being admitted to hospital with a hip fracture in
30 relation to surgery, pain management, timeliness of information given, and
31 rehabilitation?

32 **Why this is important**

33 No studies from NHS populations were identified where patients commented specifically on
34 their surgery, their pain management and rehabilitation programme. There were comments
35 in the patient views studies about not being kept informed about the management of their
36 condition, however there was no information identified about the appropriate time to be
37 told. It may be that different patients want the information at different times. The studies
38 suggest that patients suffer from fear, pain and delirium until after surgery and it is

- 1 important to learn what (if anything) can be done to alleviate this which for many will be
- 2 considered the worst stage in their treatment.

5 Imaging options in occult hip fracture

5.1 Introduction

The occult, or 'hidden', hip fracture is one in which the clinical findings are suggestive of a fracture but this is not confirmed by radiographs.

Most hip fractures can be readily diagnosed using radiographs, consisting of an antero-posterior (AP) and a lateral projection of the hip, whenever the clinical suspicion of a fracture first arises. Importantly, no clinical decision rule has yet become available that would allow clinicians to exclude a hip fracture without imaging. To avoid misdiagnosis with hip pain being attributed erroneously to soft tissue injury and the patient being discharged, a high index of clinical suspicion of hip fracture is required. This applies in all patients presenting with a typical history - usually hip pain following trauma, e.g. a fall – as certain typical features, such as the inability to bear weight or a shortened, abducted and externally rotated leg, may be absent.

Achieving an accurate diagnosis as soon as possible is advantageous for a variety of reasons. The primary reason is that without an accurate diagnosis it is not possible to formulate a proper management plan. A fracture which is not obviously evident on radiographs is likely to be undisplaced. Once the hip fracture is demonstrated early diagnosis may allow for a simple procedure to fix the fracture in situ. Should it be confirmed that no hip fracture is present then other diagnoses may be sought, there is less chance of the patient being kept unnecessarily immobile and the patient may not need to stay in hospital.

Hip radiographs have an estimated sensitivity of between 90% and 98%, and the initial films will therefore miss only a small proportion of hip fractures. It is, however, essential to ensure that the radiographs are of satisfactory quality. In particular, if the initial AP film of the entire pelvis together with the lateral hip projection (taken in the position of comfort) show no fracture, a third film should be taken centred on the hip with the hip in 10 degrees of internal rotation to position the femoral neck at 90 degrees to the x-ray beam and ensure an optimum view of this area. All subsequent discussion and recommendations assume radiographs of this standard to have been obtained before characterising a suspected but undetected fracture as occult.

The prevalence of occult hip fractures is estimated to be around 3 – 4%; up to 9% in some series (though a proportion of this may reflect radiographs of inadequate standard as discussed above). Bone resorption around the fracture site, or cortical displacement, will render most occult hip fractures visible if radiographs are repeated after a few days. This is due to bone resorption occurring along the fracture line making it radiographically more

1 obvious, but displacement or impaction may occur during this interval due to the patient
2 having walked with the fracture. Delays in surgery due to late diagnosis are associated with
3 prolonged suffering and poorer health outcomes for patients, and expose clinicians to the
4 risk of litigation.

5 Optimal strategy for patient selection and timing of secondary imaging strategies to ensure
6 early diagnosis of occult hip fractures, while avoiding over investigation of patients with
7 soft tissue injury only, is yet to be determined. However, the inability to weight bear on the
8 day following the injury, in spite of adequate analgesia, should prompt clinicians to re-
9 evaluate the patient and have a high index of suspicion of hip fracture.

10 Imaging modalities used to assist in the early detection of occult hip fractures include
11 computed tomography (CT), radionuclide scan (RNS), magnetic resonance imaging (MRI)
12 and, rarely, ultrasound scanning (US). The type of secondary imaging modalities used locally
13 is often determined by considerations of access, particularly outside normal working hours,
14 and radiological expertise available. MRI is usually considered to be the reference standard,
15 as numerous studies have found MRI to have the highest accuracy (100% sensitivity and
16 between 93% and 100% specificity, depending on experience and skill of radiologist
17 interpreting the images).

18 In this chapter we consider the clinical and cost-effectiveness of a number of alternative
19 imaging modalities that can be used to detect an occult hip fracture when MRI is
20 unavailable or precluded for safety or technical reasons.

21

22 **5.2 Review question**

23 In patients with a continuing clinical suspicion of hip fracture, despite negative radiographic
24 findings, what is the clinical and cost-effectiveness of additional imaging (radiographs after
25 at least 48 hours, RNS, US and CT, compared to MRI, in confirming, or excluding, a hip
26 fracture?

27 **5.3 Radiographs**

28 **5.3.1 What is the diagnostic accuracy of additional radiographs (X-Rays) after 48 hours** 29 **compared to MRI in the diagnosis of occult hip fractures**

30 Radiographs are the most widely available imaging technique (in- and out-of hours) utilised
31 for diagnosis of hip fracture. They can be acquired quickly (5 minutes) and experience in
32 image interpretation is widespread.

33 A hip fracture not visible on the original radiographs may become evident on films taken a
34 few days later because of bone resorption (reduced bone density) along the fracture line,
35 impaction (fracture line becomes more dense) or displacement.

36 **5.3.1.1 Clinical evidence**

37 No studies were identified.

38 **5.3.1.2 Economic evidence.**

1 No studies were identified.

2 **5.3.1.3 Recommendations and link to evidence**

3 See Section 5.6.2

4 **5.4 Radionuclide bone scan (RNS)**

5 For a RNS of the skeleton a short-life radio-isotope (technetium 99m) is linked to methylene
6 diphosphonate (MDP) which is taken up in areas of bone formation (osteoblastic activity)
7 resulting in 'hot spots'. The isotope is injected intravenously and then there has to be a
8 delay of three hours before scanning, using a gamma camera and which takes 30 minutes,
9 will detect increased uptake in the skeleton. Other causes of high bone turnover such as
10 arthritis, synovitis and tumor may lead to false positive results and these are more frequent
11 in patients over the age of 70. It is common practice to defer RNS until 72 hours after injury
12 to avoid false negative scans but some authors suggest that the modern three-phase
13 technique may give accurate results after only 24 hours.

14 **5.4.1 What is the diagnostic accuracy of RNS compared to MRI in the diagnosis of occult 15 hip fractures**

16 Two RCTs with a total of 99 participants were identified. See Evidence Table 1, Appendix E.

17

18 **5.4.1.1 Clinical evidence**

19 **Table 5-6: Bone scanning – Quality assessment**

<i>Outcome</i>	<i>Number of studies</i>	<i>Design</i>	<i>Limitations</i>	<i>Inconsistency</i>	<i>Indirectness</i>	<i>Other considerations</i>
Diagnostic accuracy ^{88,286}	2	Cross sectional study	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Bone scanning was carried out up to 72 after admission

20 (a) Evans 1994⁸⁸ study did not clearly report patient demographics

21 (b) Not clear who interpreted the results and whether they were blind to the results of the reference
22 standard test

23

24

25 **Table 5-7: - Clinical summary of findings**

<i>Outcome</i>	<i>Sensitivity (%)</i>	<i>Specificity (%)</i>	<i>NPV (%)</i>	<i>PPV (%)</i>	<i>Likelihood Ratio (+ve)</i>	<i>Likelihood Ratio (-ve)</i>	<i>Quality</i>
Diagnostic accuracy	75-98	100	93-96	100	0	0.02-0.25	Low

26

27 **5.4.2 Economic evidence**

28 No studies were identified. The cost of the procedures in England and Wales were
29 presented to the GDG: a category 3 RNS costs £205, and an MRI (one area, no contrast)

1 costs £206 (source: National schedule of reference costs 2008-09; NHS trusts and PCTs
2 combined).

3

4 **5.4.2.1 Evidence statement(s)**

Clinical The sensitivity of bone RNS compared to MRI ranged from 75% to 98% and specificity was 100%. This means that between 2% and 25% of those who have a fracture, the fracture will have been missed. However, all patients who tested positively do actually have a fracture. (LOW QUALITY)

Economic No studies were identified on the cost-effectiveness of the diagnostic accuracy of RNS compared to MRI in the diagnosis of occult hip fractures.

5

6 **5.4.3 Recommendations and link to evidence**

7 *See section 5.6.2*

8

9 **5.5 Ultrasound (US)**

10 In ultrasound (US) imaging a probe emits ultrasound waves which are reflected off surfaces
11 and recored to form the image. Good contact is required between skin and probe
12 (coupling), generally achieved with gel, but may be problematic if there is pain or soft tissue
13 swelling in the site being scanned, which may be the case in hip fracture. US is widely
14 available, both in- and out-of-hours, does not use ionising radiation and is relatively
15 inexpensive. However, it takes considerable skill and expertise to acquire optimum images
16 and for interpretation of the appearances. Currently this kind of US scanning is performed
17 by a minority of specialised musculo-skeletal radiologists in the UK.

18 Ultrasound scanning of the hip may detect bone surface changes, effusions or haemorrhage
19 in patients with fractures but the results are non-specific and usually require confirmation
20 by MRI or CT. The technique is highly operator-dependent.

21 **5.5.1 Diagnostic accuracy of ultrasound (US) compared to MRI in the diagnosis of occult hip 22 fractures**

23 One study with 30 participants was identified. See Evidence Table 1, Appendix E and forest plot
24 G1 in Appendix G

25 **5.5.1.1 Clinical evidence**

26 **Table 5-8: Ultrasound (US) – Quality assessment**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations
Diagnostic accuracy ²⁹⁷	1	Cross sectional	No serious limitations	No serious inconsistency	No serious indirectness	Sonographic examinations were performed by highly experienced musculoskeletal radiologists

1

2 **Table 5-9: Ultrasound (US) - Clinical summary of findings**

Outcome	Sensitivity (%)	Specificity (%)	NPV (%)	PPV (%)	Likelihood Ratio (+ve)	Likelihood Ratio (-ve)	Quality
Diagnostic Accuracy	100	65	100	59	2.85	0	Moderate

3

4 **5.5.1.2 Economic evidence**

5 No studies were identified. The costs of the procedures in England and Wales were
6 presented to the GDG: ultrasound (US) costs £48 for a procedure lasting less than 20
7 minutes, and £62 for a procedure lasting more than 20 minutes. The cost of an MRI (one
8 area, no contrast) is £206 (source: National schedule of reference costs 2008-09; NHS trusts
9 and PCTs combined)

10 **5.5.1.3 Evidence statement(s)**

Clinical The sensitivity of ultrasound (US) compared to MRI was 100% and specificity was 65%. This means that none of the patients who had a fracture have been missed. However, of those who tested positive 35% do not actually have a fracture – i.e. there is a high percentage of false positives (sonographic abnormalities indistinguishable from those attributable to conditions other than fracture) (LOW QUALITY)

Economic No studies were identified on the cost-effectiveness of the diagnostic accuracy of ultrasound (US) compared to MRI in the diagnosis of occult hip fractures.

11 **5.5.2 Recommendations and link to evidence**12 *See section 5.6.2*

13

14 **5.6 Computed tomography (CT)**

15 CT uses rings of sensitive detectors and an X-ray tube which rotates around the patient to
16 acquire transverse axial images through the body. CT is a readily available imaging
17 modality but its value for the detection of occult hip fractures has not been extensively
18 evaluated. There is evidence that undisplaced fractures running parallel to the axial plane
19 can be missed and limited resolution of osteoporotic trabecular bone may make the
20 technique less reliable for the detection of fractures of the hip than of other areas of the

1 body. However, technical developments in CT (spiral, multi-detector referred to as MDCT)
 2 have enabled thin 2 dimensional (2D) sections to be acquired very rapidly and from which
 3 3D volumetric reconstructions can be acquired and displayed at bone, or a variety of soft
 4 tissue, settings. This has greatly enhanced the potential application of CT to imaging occult
 5 hip fractures. The scan is rapid (2minutes)(slice thickness 1.25mm; MAs between 100 to
 6 355 depending on patient size/weight; field of view 36cm) and from which coronal, sagittal
 7 and other planar/3D reformations can be generated. CT is particularly good for imaging
 8 bone, but does not show the marrow changes (oedema) which occur in hip fracture
 9 adjacent to the fracture line.

10 **5.6.1.1 Clinical evidence**

11 No studies that meet our inclusion criteria were identified.

12 **5.6.1.2 Economic evidence**

13 No studies were identified. The costs of the procedures in England and Wales were
 14 presented to the GDG: the cost for a CT scan (one area, no contrast) is £101. The cost of an
 15 MRI (one area, no contrast) is £206 (source: National schedule of reference costs 2008-09;
 16 NHS trusts and PCTs combined)

17 **5.6.1.3 Evidence statement(s)**

Clinical No studies were identified directly comparing the diagnostic accuracy of CT
 with MRI and that meet our inclusion criteria.

Economic No studies were identified on the cost-effectiveness of the diagnostic
 accuracy of CT compared to MRI in the diagnosis of occult hip fracture.

18

19 **5.6.2 Recommendations and link to evidence**

Recommendation

Offer magnetic resonance imaging (MRI) if hip fracture is suspected despite negative anteroposterior pelvis and lateral hip X-rays. If MRI is not available within 24 hours or is contraindicated, consider computed tomography (CT).

Relative values of different outcomes

Reliability (in terms of diagnostic accuracy) was considered the primary outcome of interest. A false positive diagnosis carries the risks either of unnecessary surgery or of delay and increased cost caused by the need for additional radiographic investigation; a false negative result carries the risks associated with subsequent fracture displacement and its consequences as well as avoidable prolonged immobility and pain. It is therefore important for the selected method to minimise both false positives and false negatives.

Trade off between clinical benefits and harms

MRI cannot be used in patients with certain types of metallic implants but does not otherwise have known harmful effects other than the potential to cause claustrophobia due to the need for patients to remain in a confined space for a considerable length of

time. MRI was considered to be the first choice option in view of its superior diagnostic accuracy (up to 100% specificity and sensitivity).

If limitations in the local availability of MRI lead to unacceptably prolonged delay to diagnosis offering an RNS or CT may have a net benefit to the patient even though both carry the risks of exposure to ionising radiation. A delay of several days may, however, be required for RNS to achieve the required sensitivity, it is also generally unavailable out-of-hours (a further cause of delay), and may provide less precise information for surgical planning.

Repeat radiographs after 48 hours have limited sensitivity and carry the risks of displacement during the intervening period, as well as those of delay to surgery.

Ultrasound (US) has no known harms but its low specificity means that further imaging confirmation (with resulting delay) is required to determine whether a positive US represents a fracture, thus limiting its use. Conversely, a negative US reliably excludes fracture and could in theory enable immediate discharge of this small subset of patients from Emergency departments.

The advent of MRI has enabled the accurate early identification of occult hip fractures that would previously have been missed. The precise natural history of such occult fractures (and therefore the precise place of surgical intervention) has therefore only begun to be fully clarified. It is at least theoretically possible that a proportion of occult fractures might not require surgery. At the same time techniques of fracture fixation have also become less traumatic and invasive. Unless and until these issues of benefit/harm are fully resolved, precise and reliable early diagnosis as a basis for surgical decision making remains a clinical priority.

Economic considerations

In England and Wales, the cost of a radionuclide scan (RNS) and of an MRI is very similar: a category 3 RNS costs £205, and an MRI (one area, no contrast) costs £206. However, an MRI is cost saving compared to an RNS, as the latter may result in a longer length of hospital stay (and the possible consequences of delay to surgery) before the fracture is diagnosed.

The GDG also considered MRI to be cost-effective compared to US, since in the case of a positive US, its low specificity would still necessitate additional imaging (notably MRI or CT) to confirm the diagnosis. The possible consequences of delay to surgery would need to be added to those of additional imaging.

Quality of evidence

Two cross sectional studies comparing RNS to MRI were identified. These studies had serious methodological limitations due to the limited reporting of patient demographics and lack of clarity as to whether the assessors were blinded to the results of the index test when interpreting the results of the reference standard and vice versa.

One cross sectional study comparing ultrasound (US) to MRI was identified. This study was of moderate quality. The GDG considered

that the reproducibility was a potential limitation as the sonographic readings were performed by highly experienced musculoskeletal radiologists. There were no serious inconsistencies or indirectness in any of the identified studies.

The assumption that MRI is the gold standard for detecting occult hip fracture and the recommendation advising use of CT as an alternative to MRI were based on unanimous GDG consensus.

Other considerations

The diagnosis and management of occult hip fracture is still very much an evolving area of practice. In the absence of an evidence-based clinical decision rule clinicians must exert clinical judgement to decide when suspicion of hip fracture after normal plain radiographs is great enough to warrant additional imaging.

Before radiographs are regarded as excluding a hip fracture one should ensure that radiographic quality is optimized. When AP pelvic or hip radiographs are performed the leg should be a little internally rotated with the great toes of the feet overlapping so as to bring the anteverted femoral neck parallel to the X-ray table. In this position little of the lesser trochanter should be visible medial to the femoral cortex (the more externally rotated is the leg the more obvious is the lesser trochanter). Optimising the positioning enables the greater trochanter to be better visualized and not obscured behind the femur. When a hip fracture is present it may prove impossible to position the leg in this optimum position because of pain, but this may be compensated for by appropriate X-ray tube angulation. It should also be ensured that the X-ray exposure factors are optimum to demonstrate both the entire pelvis, to check that fractures are not present in sites additional to the hip, and also for the hip suspected of fracture. To attain this separate exposures and radiographs may be required.

Whilst the GDG considered that MRI was the best test to use to detect occult hip fracture and that this should be the first choice, they noted that there may be occasions where MRI is not available and thought it was important to give guidance as to which test to use in these circumstances. The GDG's consensus decision to recommend CT over RNS is based on greater availability, especially outside the working week, and shorter delay to diagnosis. It also reflects current NHS practice.

In addition, the technical aspects of RNS of bone (a 3 hour delay after radionuclide is given until gamma emission can be recorded; also increased uptake of radionuclide depends on increased osteoblastic activity which may take several days to occur following fracture; lack of availability out of hours) makes this the least appropriate now for imaging occult hip fractures and is now not often used in this scenario, since the advent of CT and MRI.

The GDG were also aware that rapid advances in CT technology, such as 64-slice scanners and sophisticated 3 dimensional reconstruction algorithms, may well overcome the limitations of CT reported in the published literature about its value for detection of occult hip fractures.

1

2 **5.7 Research recommendation on imaging options in occult hip** 3 **fracture**

4 The GDG recommended the following research question:

5 In patients with a continuing suspicion of a hip fracture but whose radiographs are normal,
6 what is the clinical and cost effectiveness of computed tomography compared to magnetic
7 resonance imaging, in confirming or excluding the fracture?

8 **Why this is important**

9 The GDG's consensus decision to recommend CT over a radionuclide bone scan as an
10 alternative to MRI to detect occult hip fractures reflects current NHS practice but assumes
11 that advances in technology have made the reliability of CT comparable to that of MRI. If
12 modern CT indeed can be shown to have similar reliability and accuracy to MRI, then this
13 has considerable implications because of its widespread availability out of hours and lower
14 cost. It is a high priority, therefore, to confirm or refute this assumption by direct
15 randomised comparison. The study design would need to retain MRI as "gold standard" for
16 cases of uncertainty and would clearly need to standardise the criteria, expertise and
17 procedures for radiological assessment. Numbers required would depend on the degree of
18 sensitivity/specificity (the key outcome criteria) set as target requirement for comparability,
19 but need not necessarily be very large.

1 6 Timing of surgery

2 6.1 Introduction

3 The timing of treatment for patients sustaining fractures of the proximal femur remains one
4 of the biggest challenges to a health care system. It involves multidisciplinary co-ordination
5 between accident and emergency departments, acute orthopaedic trauma services,
6 orthogeriatricians, anaesthetists, as well as the availability of appropriate theatre space
7 with trained staff and relevant equipment. In the past these patients were given low
8 priority in the hospital system, which led to many delays and repeated periods of
9 starvation. It is recognised that it is not only the time a patient takes to get to surgery that
10 is important, but that the patient has to be medically optimised, with the anaesthetic,
11 surgical and theatre team being appropriately experienced. When planning any emergency
12 care it is not always possible to predict the number of cases which can present, so any
13 system which is set up must have the flexibility to adapt to the peaks and troughs of
14 admissions. This can lead to potential free theatre capacity in quieter periods.

15 As it would be unethical to enforce an unnecessary delay for patients sustaining fractures of
16 the proximal femur, all studies reported are retrospective cohort studies. As such the level
17 and quality of the evidence is poor.

18 The timing of surgery is an early marker of a patient's progress following a hip fracture. The
19 surgery does not stand alone. The pathway to safe, timely surgery includes proper
20 organisation and expertise in diagnosis, medical optimisation and anaesthesia. In the last
21 decade many orthopaedic trauma emergencies are now treated on dedicated planned
22 trauma lists. A planned trauma list is one with a rostered senior anaesthetist, senior
23 surgeon and dedicated theatre time. These by their nature usually concentrate the
24 expertise required.

25 There are sometimes legitimate reasons for delay and it is important to look at the
26 excluded patients in these studies. In a few patients delay to surgery is unavoidable.
27 However, it should be anticipated that many patients with hip fractures will be frail and
28 have comorbidities. The following would be common findings in patients presenting with
29 hip fractures:

- 30 • Anaemia
- 31 • Anticoagulation
- 32 • Volume depletion
- 33 • Electrolyte imbalance

- 1 • Uncontrolled diabetes
- 2 • Uncontrolled heart failure
- 3 • Correctable cardiac arrhythmia or ischaemia
- 4 • Acute chest infection
- 5 • Exacerbation of chronic chest conditions
- 6

7 Provided these problems are sought and measures initiated to correct them are taken
8 promptly the majority can be optimised within 24 hours.

9 When looking at the timings measured it is generally accepted the time of diagnosis should
10 be the initial time recorded and the time to the start of the anaesthetic procedure be the
11 index time measured. Objective outcomes used to compare timing of surgery include early
12 and late mortality, length of hospital stay, return to mobility, complications including chest
13 infections and pressure sores, change of residence and other surgical complications. What
14 has not been measured in the past is the pain and suffering experienced with prolonged
15 delay and what is the ethical time period the elderly, who are often very frail, should wait
16 for treatment.

17 **6.1.1 Review question**

18 In patients with hip fractures what is the clinical and cost effectiveness of early surgery
19 (within 24, 36 or 48 hours) on the incidence of complications such as mortality, pneumonia,
20 pressure sores, cognitive dysfunction and increased length of hospital stay?

21 10 studies met the inclusion criteria for this question, with a total of 193,793 participants.
22 Data are given for studies where outcomes have been adjusted for confounding factors
23 such as comorbidity and age using logistic regression (7 studies). A separate subgroup is
24 given which excludes patients who are unfit for surgery i.e. reason for delay is due to
25 unavailability of staff, theatres or equipment (3 studies). Delay to surgery in the identified
26 studies was from time to admission. All studies report surgical delay versus early surgery to
27 investigate the harm of delaying surgery.

28 The cut-off for delay to surgery in this analysis is 24, 36 and 48 hours.

29 See evidence table 2, Appendix E and forest plots G2 to G22 in Appendix G.

30

1 **6.1.1.1 Clinical evidence**2 **Table 6-10: Late (>24h) versus early surgery for hip fracture – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality – In hospital ^{19,351}	2	Observational	No serious limitations	No serious inconsistency	No serious indirectness (b, d)	Serious imprecision (e)
Mortality – 30 days ³⁰	2	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness (a, b, d)	Serious imprecision (e)
Mortality – 3 months ³⁵¹	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness (b)	Serious imprecision (e)
Mortality – 4 months ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision (e)
Mortality – 1 year ³⁵¹	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness (b)	Serious imprecision (e)
Return to independent living ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision (e)
Pressure ulcers ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	No serious imprecision
Major complications ^(c) ₁₉	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness (d)	No serious imprecision

- 3 (a) In Bottle and Aylin, 2006³⁰ baseline data, such as age is given for the entire cohort and also
4 stratified by type of surgery e.g. fixation, replacement, other procedure. No baseline data stratified
5 by delay to surgery. Patients were all admitted from their own home.
6 (b) In Weller et al., 2005³⁵¹ baseline data, such as age is stratified per hospital. No baseline data
7 stratified by delay to surgery.
8 (c) Severe complications were defined as cerebrovascular accident, cardiorespiratory complications,
9 digestive complications except unspecific paralytic ileus, and dialysis.
10 (d) The comparison is 24-48h vs. 0-24 h time to surgery for Bergeron 2006¹⁹
11 (e) The wide confidence intervals around the estimate make it difficult to determine and effect size for
12 this outcome.
13
14
15

1 **Table 6-11: Late (>24 hours) versus early surgery for hip fracture - Clinical summary of findings**

Outcome	Late surgery ^(a)	Early surgery ^(a)	Adjusted Odds Ratio	Absolute effect	Quality
Mortality – in hospital	325	523	0.88 (0.55 - 1.41)	N/A	Very low
Mortality – in hospital	25320	20303	1.17 (1.08 - 1.26)	N/A	Low
Mortality – 30 days	45862	69080	1.25 (1.19 - 1.31)	N/A	Very low
Mortality – 3 months	25320	20303	1.11 (1.05 - 1.17)	N/A	Very low
Mortality – 4 months	225	209	1.07 (0.67 - 1.70)	N/A	Very low
Mortality – 1 year	25320	20303	1.13 (1.05 - 1.22)	N/A	Very low
Return to independent living	225	209	0.86 (0.45 - 1.65)	N/A	Very low
Pressure ulcers	225	209	2.19 (1.21 - 3.96)	N/A	Low
Major complications	325	523	0.87 (0.58 - 1.29)	N/A	Low

(a) Numbers of patients in each study arm. No event data is given as the data provided is odds ratios adjusted using logistic regression for confounding factors.

2
3
4
5

Table 6-12: Late (>36h) versus early surgery for hip fracture – Clinical study characteristics

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality – in hospital ¹⁸⁹	1	Observational	No serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)
Minor complications ¹⁸⁹	1	Observational	No serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)
Major complications ¹⁸⁹	1	Observational	No serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)
Pressure ulcers ¹⁸⁹	1	Observational	No serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)
Mortality – 4 months ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)
Pressure ulcers ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	No serious imprecision
Return to independent living ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(a)

(a) Baseline data given for entire cohort not by time to surgery.

(b) Late surgery is between 24-48h with early surgery defined as <24h.

(a) The wide confidence intervals around the estimate make it difficult to determine and effect size for this outcome.

6
7
8
9
10

1 **Table 6-13: Late (>36 hours) versus early surgery for hip fracture - Clinical summary of findings**

Outcome	Late surgery ^(a)	Early surgery ^(a)	Adjusted Odds Ratio	Absolute effect	Quality
Mortality – in hospital	264	245	0.82 (0.42 - 1.62)	N/A	Very low
Minor complications	264	245	1.53 (1.05 - 2.22)	N/A	Very low
Major complications	264	245	0.96 (0.52 - 1.75)	N/A	Very low
Pressure ulcers	264	245	1.23 (0.71 - 2.12)	N/A	Very low
Mortality – 4 months	194	550	1.5 (0.63 – 1.74)	N/A	Very low
Pressure ulcers	194	550	3.42 (1.94 – 6.03)	N/A	Low
Return to independent living	194	550	0.44 (0.21 – 0.91)	N/A	Very low

2 (a) Numbers of patients in each study arm. No event data is given as the data provided is odds ratios
 3 adjusted using logistic regression for confounding factors.
 4
 5

5 **Table 6-14: Late (>48h) versus early surgery for hip fracture – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality – In hospital ^{19,189,351}	3	Observational	No serious limitations	No serious inconsistency	No serious indirectness ^(b,d)	Serious imprecision ^(e)
Mortality – 30 days ^{30,125}	2	Observational	Serious limitations ^(a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(e)
Mortality – 3 months ³⁵¹	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness ^(b)	No serious imprecision
Mortality – 4 months ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(e)
Mortality – 1 year ³⁵¹	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness ^(b)	No serious imprecision
Return to independent living ⁴	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(e)
Pressure ulcers ^{4,125,189}	3	Observational	No serious limitations	No serious inconsistency	No serious indirectness	No serious imprecision
Major complications ^{(c)19,189}	2	Observational	No serious limitations	No serious inconsistency	No serious indirectness ^(d)	Serious imprecision ^(e)
Minor complications ¹⁸⁹	1	Observational	No serious limitations	No serious inconsistency	No serious indirectness ^(d)	No serious imprecision

- (a) In Bottle and Aylin, 2006³⁰ baseline data, such as age is given for the entire cohort and also stratified by type of surgery e.g. fixation, replacement, other procedure. No baseline data stratified by delay to surgery. Patients were all admitted from their own home.
- (b) In Weller et al., 2005³⁵¹ baseline data, such as age is stratified per hospital. No baseline data stratified by delay to surgery.
- (c) In Bergeron 2006¹⁹, severe complications were defined as cerebrovascular accident, cardiorespiratory complications, digestive complications except unspecific paralytic ileus, and dialysis.
- (d) The comparison is >48h vs. 0-24 h time to surgery
- (e) The wide confidence intervals around the estimate make it difficult to determine the effect size for this outcome.

Table 6-15: Late (>48 hours) versus early surgery for hip fracture - Clinical summary of findings

Outcome	Late surgery ^(a)	Early surgery ^(a)	Adjusted Odds Ratio	Absolute effect	Quality
Mortality – In hospital ¹⁹	129	848	1.16 (0.64 - 2.13)	N/A	Very low
Mortality – in hospital ¹⁸⁹	98	509	0.93 (0.38 - 2.33)	N/A	Very low
Mortality – In hospital ³⁵¹	7314	20303	1.60 (1.42 - 1.80)	N/A	Low
Mortality – 30 days ³⁰	24391	90551	1.36 (1.29 - 1.43)	N/A	Very low
Mortality – 30 days ¹²⁵	3805	4578	0.71 (0.45 - 1.10)	N/A	Very low
Mortality – 3 months	7314	20303	1.40 (1.28 - 1.54)	N/A	Low
Mortality – 4 months	98	646	0.86 (0.44 - 1.69)	N/A	Very low
Mortality – 1 year	7314	20303	1.58 (1.26 - 1.99)	N/A	Low
Return to independent living	98	646	0.33 (0.14 - 0.78)	N/A	Very low
Pressure ulcers ⁴	98	646	4.34 (2.34 - 8.04)	N/A	Low
Pressure ulcers ¹²⁵	3805	4578	1.20 (0.9 - 1.6)	N/A	Very low
Pressure ulcers ¹⁸⁹	98	509	2.29 (1.19 - 4.40)	N/A	Low
Major complications ¹⁹	129	848	1.32 (0.79 - 2.20)	N/A	Very low
Major complications ¹⁸⁹	98	509	2.21 (1.01 - 4.34)	N/A	Very low
Minor complications	98	509	2.27 (1.38 - 3.72)	N/A	Low

(a) Numbers of patients in each study arm. No event data is given as the data provided is odds ratios adjusted using logistic regression for confounding factors.

Table 6-16: Late (>48h) versus early surgery for hip fracture (length of hospital stay outcomes)– Clinical study characteristics

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Postoperative length of hospital stay ¹⁹	1	Observational	No serious limitations ^(a)	No serious inconsistency	No serious indirectness	No serious imprecision
Postoperative length of hospital stay; without comorbidity ¹⁹	1	Observational	No serious limitations ^(a)	No serious inconsistency	No serious indirectness	No serious imprecision
Postoperative length of hospital stay (including	1	Observational	No serious limitations ^(a)	No serious inconsistency	No serious indirectness	No serious imprecision

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
rehab) ³⁰⁸						

(a) Mean and standard deviations are not provided, only median or mean and 95% confidence interval.

Table 6-17: Late (>48h) versus early surgery for hip fracture - Clinical summary of findings; length of hospital stay

Outcome	Late surgery ^(c)	Early surgery ^(c)	Median (days) Late surgery	Median (days) Early surgery	Quality
Postoperative length of hospital stay ^(a)	129	848	28	18	Low
Postoperative length of hospital stay; without comorbidity	30	248	20	16	Low
Postoperative length of hospital stay (including rehab)	174	3454	36.5 ^(b)	21.6 ^(b)	Low

(a) Data is unadjusted for co-morbidity, which is more frequent in the delayed surgery study arm.

(b) Mean number of days given, 95% confidence interval = 5.7 to 16.0, $p < 0.0001$.

(c) Numbers of patients in each study arm. No event data is given as the data provided is odds ratios adjusted using logistic regression for confounding factors.

1 **Table 6-18: Late (>24h) versus early surgery for hip fracture (exclusion of patients unfit for**
 2 **surgery) – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality 30 days ²¹⁵	1	Observational	Serious limitations (a, b)	No serious inconsistency	No serious indirectness	Serious imprecision (c)
Mortality and needing total assistance in locomotion at 6 months ²⁵⁰	1	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision (c)
Major postoperative complications ²⁵⁰	1	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision (c)

3 (a) Baseline data not reported separately for the restricted cohort.

4 (b) No protocol for determining which patients were unfit for surgery and anaesthesia, therefore
 5 variation between clinicians.

6 (c) The wide confidence intervals around the estimate make it difficult to determine and effect size for
 7 this outcome.

8

9

10

Table 6-19: Late (>24 hours) versus early surgery for hip fracture (exclusion of patients unfit for
surgery) - Clinical summary of findings

Outcome	Late surgery	Early surgery	Risk Ratio	Absolute effect	Quality
Mortality 30 days	85/1166	85/982	0.84 (0.63 - 1.12)	N/A	Very low
Mortality and needing total assistance in locomotion at 6 months	509		0.62 (0.35 - 1.08) (a)	N/A	Very low
Major postoperative complications	273		0.26 (0.07 - 0.95) (a)	N/A	Very low

11 (a) Adjusted odds ratio

12

13

1 **Table 6-20: Late (>48h) versus early surgery for hip fracture (exclusion of patients unfit for**
 2 **surgery) – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality 30 days ²¹⁵	1	Observational	Serious limitations (a, b)	No serious inconsistency	No serious indirectness	Serious imprecision (c)
Mortality at 1 year ³⁰⁸	1	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness	No serious imprecision
Change in residence (more dependent) ³⁰⁸	1	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness	Serious imprecision (c)
Return to original residence ³⁰⁸	1	Observational	Serious limitations (a)	No serious inconsistency	No serious indirectness	No serious imprecision

3 (a) Baseline data not reported separately for the restricted cohort.

4 (b) No protocol for determining which patients were unfit for surgery and anaesthesia, therefore
 5 variation between clinician decisions.

6 (c) The wide confidence intervals around the estimate make it difficult to determine and effect size for
 7 this outcome.

8

9

10 **Table 6-21: Late (>48 hours) versus early surgery for hip fracture (exclusion of patients unfit for**
 11 **surgery) - Clinical summary of findings**

Outcome	Late surgery	Early surgery	Risk Ratio	Absolute effect	Quality
Mortality 30 days	36/497	134/1651	0.89 (0.63 – 1.27)	N/A	Very low
Mortality at 1 year	24/174	238/3454	0.5 (0.34 – 0.74)	N/A	Very low
Change in residence (more dependent)	22/174	240/3454	0.55 (0.37 – 0.83)	N/A	Very low
Return to original residence	128/174	2974/3454	1.17 (1.07 – 1.28)	N/A	Very low

12

13 **6.1.1.2 Economic evidence**

14 One study^{304,304} was found which calculated the mean hospital costs for hip fracture
 15 patients who had received surgery at different points in time from admission. This study
 16 was excluded because of serious methodological limitations, as no reason was given as to
 17 why patients had faced delays before receiving surgery (whether it was because of medical
 18 or administrative reasons)

19 An original decision analytical model was developed to compare the cost-effectiveness of a
 20 strategy consisting in adding extra half-day operating lists to increase the proportion of
 21 patients operated within 48 hours from admission against a non-investment strategy.
 22 Please see Appendix H, section 8.5 for further details.

23 **Table 6-22: Early versus late (>48h) surgery for hip fracture - Economic study characteristics**

Study	Limitations	Applicability	Other Comments
NCGC decision model	Minor limitations (a)	Partial applicability (b)	

24 (a) Cost-effectiveness analysis based on a Markov model.

(b) The findings of the model may not be generalized to the whole UK NHS because its treatment effects and cost data are based on evidence from two specific hospital settings. The addition of extra operating lists may not be feasible for those providers where no spare theatre capacity is available.

Table 6-23: Early versus late (>48h) surgery for hip fracture - Economic summary of findings

Study	Incremental cost (£)	Incremental effects (QALYs)	ICER	Uncertainty
NCGC decision model	1) £1,000 for the first year of implementation of extra operating lists ^(a)	1) 0.0425 for the first year of implementation of extra operating lists	1) £22,542/QALY for the first year of implementation of extra operating lists	95% CI: cost saving – dominated (both in the first and in the second year of implementation of extra operating lists ^(d))
	2) £ 800 for the second year of implementation of extra operating lists ^(b)	2) 0.094 for the second year of implementation of extra operating lists ^(c)	2) £8,933/QALY for the second year of implementation of extra operating lists	

(a) In the first year of implementation of extra operating lists, the mean costs for investment in extra operating lists early surgery were £47.4, and for the non-investment strategy £46.4.

(b) For the second year, the mean costs associated with the strategy of investment for early surgery were £47.3, and for the non-investment strategy £46.4.

(c) In the first year of implementation of extra operating lists, the mean effectiveness for the strategy of investment for early surgery was 2.3637, and for the non-investment strategy 2.3212. In the second year, they corresponded to 2.415 and 2.321 respectively.

(d) 95% CI of ICERs calculated from the 10,000 Monte Carlo simulations. The high uncertainty of the model is due to all the types of variables, including the effectiveness of interventions. We have tested the uncertainty of all categories of inputs in the model (costs, utilities, relative risks), by making probabilistic one category at a time while keeping the others deterministic, and under all scenarios the findings showed great uncertainty, with a 95% CI cost saving – dominated”.

6.1.1.3 Evidence statement (s)

Clinical All patients

Early surgery (<24h) shows a statistically significant and clinically significant reduction in mortality (in 4 out of 7 studies) (VERY LOW QUALITY) and reduction in pressure ulcers (LOW QUALITY) with early surgery compared to late surgery. No statistically significant difference shown for return to independent living or major complications (LOW QUALITY).

Early surgery (<36h) – statistically significant and clinically significant reduction in pressure ulcers with early surgery compared to late surgery (LOW QUALITY). Statistically significant, but not clinically significant increased return to independent living (VERY LOW QUALITY). No statistically significant difference in mortality at 4 months (VERY LOW QUALITY).

Early surgery (<48h) shows a statistically significant and clinically significant reduction in mortality (in 4 out of 8 studies) (VERY LOW QUALITY), increased return to independent living (VERY LOW QUALITY), reduced pressure ulcers (LOW QUALITY), reduced major and minor complications with early surgery compared to late surgery (VERY LOW QUALITY).

Exclusion of patients unfit for surgery

Early surgery (<24h) – Statistically significant, but not clinically significant reduction in major postoperative complications with early surgery compared to late surgery. No statistically significant difference in mortality, with early surgery compared to late surgery. (VERY LOW QUALITY)

Early surgery (<48h) – Statistically significant, and clinically significant reduction in mortality at 1 year and patients changing residence (more dependent) and increased return to original residence (VERY LOW QUALITY). No statistically significant difference in mortality at 30 days with early surgery compared to late surgery. (VERY LOW QUALITY).

Economic Investing in adding extra operating lists as a way to increase the proportion of patients operated within 48 hours from admission is only marginally above the £20k/QALYs threshold in the first year of implementation, but becomes clearly cost-effective from the second year onwards.

This evidence has minor limitations and partial applicability.

1

2 6.1.2 Recommendations and link to evidence

Recommendation	Perform surgery on the day of, or the day after, admission.
Relative values of different outcomes	<p>The GDG recognised that hip fracture surgery was often disproportionately delayed in comparison with other operations, and that this in part reflected a lack of sufficient priority afforded to this group of patients.</p> <p>On humanitarian criteria alone, initiatives to avoid delay were considered to be of high priority in developing the guidance. It was considered that surgery was the best form of pain relief, and that to spend more than one night in hospital without operation was generally unacceptable.</p> <p>Postponement of surgery carries increased risk of complications, as well as prolongation of pain, and the need for repeated preoperative fasting.</p> <p>Of the outcomes derived from the literature, mortality, return to independent living, occurrence of specific complications (notably pressure ulcers) and duration of hospital stay were all considered of parallel and inter-related importance as indicators of care standard and efficacy.</p>
Trade off between clinical benefits and harms	<p>There was no instance in the literature of any advantage in delaying surgery, nor of disadvantage in reducing delay.</p> <p>Although the range of studies utilised a range of arbitrary or pragmatic time thresholds (governed to some degree by service context and organisation), there was no definitive cut-off point (up to and including 24 hours) beyond which further reduction of delay ceased to confer measurable benefit in one or more outcomes.</p> <p>Therefore the GDG considered it could not be prescriptive about</p>

the precise time threshold from the literature alone.

The trade off between early surgery and harms relate to the difficulties and infrastructure required to treat this population who present as emergencies. It is recognized surgery is the best form of analgesia and as over 30% present with cognitive impairment, it can be otherwise difficult to assess patients suffering. It is also considered humane not to leave this frail patient group waiting treatment (often being repeatedly starved). The potential harm of earlier surgery include the risks of not medically resuscitating and optimizing the patients health prior to a further surgical insult and ensuring the surgical team is experienced and available. A delay up to 36 hours allows for appropriate assessment and planning. It allows patients to be operated on in planned trauma lists and should allow most hospitals to cope with peaks in emergency admissions.

Only one study⁴ looked at complications, return to independent living and pressure sores. Whilst this study did report a small benefit in protecting against pressure sores it did not demonstrate any additional benefits. Regarding mortality one study³⁵¹ showed a small difference in mortality at one year, though again the difference and numbers were small.

Alani et al., 2008⁴ is the only study which looked at the 36 hour time frame. It failed to show improvement in mortality at four months yet showed a slight benefit in return to independent living and avoidance of pressure ulcers.

When comparing surgery at 48 hours, again the data is limited. The overall number of patients included is small and there is a reported decrease in mortality in two out of the five studies included^{30,351}. Apart from the benefits already reported in Alani's study, other outcomes were either not reported or did not show any difference.

Economic considerations

To be able to offer surgery for hip fracture patients by an experienced surgical team, within the recommended time period, it is recognized there may have to be an investment in infrastructure, specifically planned trauma operating lists with experienced surgical, anaesthetic and theatre teams. Generally these should occur in the normal working day. As admission numbers, including peaks and troughs, cannot be always predicted then this capacity may not always be utilised.

The potential costs of reducing delay to surgery were recognised- such as additional theatre time, out-of-hours staffing (including senior staff), out-of-hours lists and planned trauma lists.

These costs will be at least partially offset by potential savings from reduced length of stay, reduced complications and enhanced return to independent living.

There was no definitive health economic study for any time threshold in the literature. The guideline group therefore considered that an original decision model was crucial to inform the broad economic feasibility of any recommendation on reducing

surgical delay. As discussed in Appendix H, the GDG agreed that, out of the evidence included in the clinical review, the outcome data to undertake this analysis were adequate only to provide a model based on a 48hr threshold, and as a consequence this specific cut-off point was selected for the economic analysis.

The economic model demonstrates that investing to add extra operating lists in order to undertake surgery within 48 hours from admission is only marginally above the £20k/QALYs threshold in the first year of implementation, but becomes clearly cost-effective in the following years.

Furthermore, the implementation of extra operating lists will also achieve a more equitable distribution of health care resources in favour of patients that had previously been made to wait for surgery as other cases were given higher priority.

However, the model does not capture the possibility that the extra operating lists could potentially be used to treat cases in addition to hip fracture patients (thus resulting in an increase of activity for the hospital trust and subsequent QALYs gains for the patients treated).

In addition, our cost-effectiveness estimates are also conservative in that we do not look at the impact that early surgery has on the pain relief of our population.

Quality of evidence

The available clinical evidence covering this issue is of low quality, but in aggregate supports the avoidance of surgical delay.

For this reason there is an element of consensus in the wording of the recommendation which, in addition to the evidence of clinical benefit and NHS economic feasibility, also reflects a strong humanitarian case. The consensus was unanimous within the GDG.

The health economic analysis reported in Appendix H showed that surgery performed with 48 hours was cost effective.

Although the evidence base for this question is predominantly retrospective, cohort studies of low quality (all low or very low) it is not considered ethical to conduct an RCT to answer this question.

The main studies included were cohort studies that adjusted for confounding factors by logistic regression, which although were low quality were considered higher quality than cohort studies without any adjustment. The subgroup studies did not adjust for confounding factors, but were considered as similar quality to those studies using logistic regression as the population excluded those unfit for surgery.

Other considerations

The context of implementation has changed during guideline development in such a way as to highlight the relevance and feasibility of the recommendation, in that the Department of Health has introduced a Best Practice Tariff initiative to achieve hip fracture surgery within 36 hours of admission.

<p>Recommendation</p>	<p>Identify and treat correctable comorbidities immediately so that surgery is not delayed by:</p> <ul style="list-style-type: none"> • anaemia • anticoagulation • volume depletion • electrolyte imbalance • uncontrolled diabetes • uncontrolled heart failure • correctable cardiac arrhythmia or ischaemia • acute chest infection • exacerbation of chronic chest conditions.
<p>Relative values of different outcomes</p>	<p>The most important outcomes considered here were mortality, length of stay in hospital and postoperative complications.</p>
<p>Trade off between clinical benefits and harms</p>	<p>Patients should not be delayed for routine tests which will not affect the surgical or anaesthetic procedure. It has been shown in the majority of patients that longer delay leads to an increase in complications and length of stay in those medically fit.</p> <p>A number of medical conditions that might pose a concern to the surgeon or the anaesthetist are so commonly encountered among patients presenting with hip fracture that their occurrence should be anticipated, and admission assessment and management protocols designed that will expedite their management and so prevent their delaying surgery. The process of pro-actively seeking to identify such conditions will also help in identifying other less common potential concerns that might need more individual assessment - by experienced physicians (often orthogeriatricians) or anaesthetists - when a medical delay may be required.</p>
<p>Economic considerations</p>	<p>The early identification and treatment of patients' comorbidities may require additional resources in terms of personnel's rounds and ad-hoc tests. These costs would be at least partially off-set by savings linked with a lower length of hospital stay associated with the possibility of performing surgery at an earlier stage.</p>
<p>Quality of evidence</p>	<p>The evidence included in this chapter did not cover treatment of comorbidities. The main studies adjusted for these factors and the subgroup excluded patients unfit for surgery.</p>
<p>Other considerations</p>	<p>There should be the availability of experienced orthogeriatricians / physicians and anaesthetists to assess patients who may require further optimization. Regular review and communication with the surgical team is essential.</p>

1 **6.2 Research recommendations on timing of surgery**

2 **6.2.1 Surgery within 36 hours**

3 The GDG recommended the following research question:

- 4 ➤ What is the clinical and cost effectiveness of surgery within 36 hours of admission
5 compared to surgery later than 36 hours from admission in mortality, morbidity
6 and quality of life in patients with hip fracture?

7 **Why this is important**

8 Early and appropriate surgery for hip fractures is the most effective form of pain relief,
9 potentially quickening the rehabilitation and reducing complications. Within the current
10 literature no specific time interval threshold has been identified (up to 24hr) below which a
11 reduction in delay has shown no benefit. In addition to the evidence of the cost
12 effectiveness below 48hr, pragmatic, organisational and humanitarian considerations have
13 been utilised to arrive at the recommendation to operate not later than the day after
14 admission. A formal study within the NHS based on an arbitrary but realistic 36hr threshold
15 would provide additional important data to that already available, in order to inform more
16 precisely the forward clinical and cost-effectiveness of the strategy. For ethical reasons, the
17 research design would be an observational cohort study, correcting for confounding
18 variables, possibly set in the context of the National Hip Fracture Database and examining
19 the effect of the time to surgery and its cost on key outcomes, including mortality,
20 complications, length of stay, time taken to rehabilitate and qualitative aspects of the
21 experiences of patients.

7 Analgesia

7.1 Introduction

Pain is a major component of the patient experience following a hip fracture. Fracture and postoperative pain, along with fracture and surgical site blood loss, constitute the major physiological stresses facing these patients. Fear of pain is a major concern to them and their relatives. The best form of analgesia is surgical repair, but there will usually be a period when assessment is taking place when some analgesia is needed. Prompt and adequate relief of pain has long been identified as a major priority in the management of hip fracture, and one that has not always historically been achieved.

Pain relief is obviously important for simple humanitarian reasons and for acute nursing care, but also improves patients' wellbeing, reduces the risk of delirium, and facilitates the return to mobility and independence.

It is often difficult to assess the need for analgesia when the patients are lying still. They may require more pain relief when moved passively for investigations, such as radiological procedures and subsequently for the active mobilisation essential to their successful recovery. Many patients with hip fracture may be unable to express their pain, either because of cognitive impairment, acute delirium or an underlying expressive dysphasia.

Systemic analgesics act through the bloodstream on the whole body rather than on a localised area or region. They are still the most widely used drugs for providing pain relief in acute painful situations. Systemic analgesics used for pain relief in hip fracture include simple analgesics such as paracetamol, and a wide range of opioids. Non-steroidal anti-inflammatory drugs are usually avoided or used with caution because of their side effects. These include upper gastrointestinal bleeding, nephrotoxicity and fluid retention – to all of which the older population and are well known to exhibit increased susceptibility.

The nerves supplying the proximal femur may also be blocked by injecting local anaesthetic around the femoral nerve. These injections are referred to as nerve blocks and are sometimes administered to patients to reduce pain if simple analgesics and opioids have not proven to be sufficient. They are also thought to improve pain scores and mobility and to help avoid excessive opioid usage.

The aim of this chapter is to identify optimal preoperative and postoperative analgesia including the use of nerve blocks as adjuncts or alternatives to simple analgesics such as paracetamol and opioids.

1 The use of nerve blocks as with anaesthesia is covered in Chapter 8 on regional compared
2 to general anaesthesia.

3 **7.2 Systemic analgesia**

4 **7.2.1 Review question**

5 In patients who have or are suspected of having a hip fracture, what is the comparative
6 effectiveness and cost effectiveness of systemic analgesics in providing adequate pain relief
7 and reducing side effects and mortality?

8 **7.2.1.1 Clinical evidence**

9 No studies on the effectiveness of these drugs in hip fracture patients were identified.

10 **7.2.1.2 Economic evidence**

11 No relevant studies were identified. We conducted a cost analysis of a nerve block, non-
12 opioids and other analgesics. We found that a nerve block would cost approximately
13 £54.66. The average cost for opioids controlled drugs is £11.84 (where £1.34 is the average
14 cost per dose of the drugs and £10.50 the personnel cost of two trained nurses required for
15 the administration of the drugs). The price of opioids non-controlled drugs is estimated at
16 £1.96 per doses. The cost of non-opioids analgesics is less than £0.1p per dose. Please see
17 Appendix H section 8.1 for further details.

18 **7.2.2 Recommendations and link to evidence**

19 In order to present the recommendations in a logical manner and retain their sequential
20 order, the recommendations for this section are presented below in section 7.3.2

21

22 **7.3 Nerve blocks compared to systemic analgesia**

23 **7.3.1 Review question**

24 In patients who have or are suspected of having a hip fracture, what is the clinical and cost
25 effectiveness of nerve blocks compared to systemic analgesia in providing adequate pain
26 relief and reducing side effects and mortality?

27 One systematic review²⁶² was identified including 17 RCTs with a total of 888 participants.
28 See evidence table 3, Appendix E and forest plots G23 to G37 in Appendix G.

29

30 **7.3.1.1 Clinical evidence**

31 The review considered any nerve block that affects the nerves supplying the proximal
32 femur. These include the subcostal nerve, the lateral cutaneous nerve of the thigh, the
33 femoral nerve, psoas (lumbar plexus), fascia iliaca compartment block (FICB) and triple
34 (femoral, obturator and sciatic) nerve.

1 The literature search retrieved one Cochrane review (Parker et al 2002)²⁶². A further update
 2 search was then conducted to look for any papers that may have been published since the
 3 publication of this review. No additional studies were retrieved and therefore the clinical
 4 evidence presented in this chapter is based on the Parker et al results with the addition of
 5 the GRADE analysis.

6

7 **Table 7-24: Nerve blocks versus systemic analgesia – Clinical study characteristics**

Outcome	Number of studies	Design^f p)	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Pain ^{116,182,220}	3	RCT	Serious limitations ^(a)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Unsatisfactory pain control preoperatively or need for 'breakthrough' analgesia ^{51,98,116,182,220}	5	RCT	Serious limitations ^(b)	No serious inconsistency	No serious indirectness	No serious imprecision
Unsatisfactory pain control postoperatively ^{51,62}	2	RCT	Serious limitations ^(c)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Nausea and/or vomiting ^{62,98,116,220,318,331}	6	RCT	Serious limitations ^(d)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Need for anti-emetics ³³¹	1	RCT	Serious limitations ^(e)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Wound infection ⁹⁹	1	RCT	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Pneumonia ^{95,99,129,207,352}	5	RCT	No serious limitations	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Any cardiac complication ^{99,207}	2	RCT	Serious limitations ^(f)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Myocardial infarction ²⁰⁷	1	RCT	Serious limitations ^(g)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Puritis ³³¹	1	RCT	Serious limitations ^(h)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Pulmonary embolism ^{99,129}	2	RCT	No serious limitations	No serious inconsistency ^(m)	No serious indirectness	Serious imprecision ^(o)
Deep vein thrombosis ^{62,95,99,129,352}	5	RCT	Serious limitations ⁽ⁱ⁾	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Mortality ^{62,95,99,129,153,165,207,352}	8	RCT	Serious limitations ^(j)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)
Pressure sores ^{62,129,182}	3	RCT	Serious limitations ^(k)	No serious inconsistency ⁽ⁿ⁾	No serious indirectness	Serious imprecision ^(o)
Confusional state ^{62,182,352}	3	RCT	Serious limitations ^(l)	No serious inconsistency	No serious indirectness	Serious imprecision ^(o)

8 (a) One study (Gille 2006)¹¹⁶ did not state the method of randomisation. All 3 studies were not adequately
 9 blinded.

10 (b) High risk of bias due to lack of allocation concealment. 2 (Gille 2006 and Chudinov 1999)^{51,116} out of the
 11 5 studies did not specify their method of randomisation.

- 1 (c) One study (Chudinov 1999)⁵¹ did not clearly report its randomisation method and did not report any
2 allocation concealment.
- 3 (d) Low risk of bias. 2 out of the 6 studies did not clearly report randomisation method and allocation
4 concealment.
- 5 (e) High risk of bias due to unclear reporting of the method of randomisation
- 6 (f) One of the 2 studies (Matot 2003)²⁰⁷ has a high risk of selection bias due to unclear methods of
7 concealment and randomisation.
- 8 (g) This study has a high risk of selection bias due to unclear methods of concealment and randomisation
- 9 (h) This study has a high risk of selection bias due to unclear methods of concealment and randomisation. It
10 also had a very short follow up (24 hours).
- 11 (i) One of the 5 studies (White 1980)³⁵² has a high risk of selection bias due to unclear methods of
12 concealment and randomisation.
- 13 (j) Two of the studies (white and Hood)^{153,352} had a high risk of selection bias due to unclear methods of
14 concealment and randomisation. One study also had a high number of drop outs in one the trial arms.
- 15 (k) studies has a high risk of selection bias due to unclear methods of concealment and randomisation
- 16 (l) One of the studies (white 1980)³⁵² had a high risk of selection bias due to unclear methods of
17 concealment and randomisation. One study also had a high number of drop outs in one the trial arms.
- 18 (m) There was some non statistically significant heterogeneity $I^2 = 31\%$ $p=0.23$.
- 19 (n) There was some non statistically significant heterogeneity $I^2 = 30\%$ $p=0.23$.
- 20 (o) The wide confidence intervals around the estimate make the result imprecise. Consequently, it is
21 difficult to determine the true effect size for this outcome.
- 22 (p) The following studies included nerve blocks in conjunction with general anaesthesia: Foss et al (2005)⁹⁹,
23 Tuncer et al (2003)³³¹, Spansberg et al (1996)³¹⁸, Hood et al (1991)¹⁵³, Jones et al (1985)¹⁶⁵, White et al
24 (1980)³⁵².

25
26
27
28 **Table 7-25: Nerve blocks versus systemic analgesia - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk (95% confidence interval)	Absolute effect	Quality
Pain	106	104	N/A	SMD -0.52 (-0.8 to -0.25)	Low
Unsatisfactory pain control preoperatively or need for 'breakthrough' analgesia	18/150 (12%)	47/148 (31.8%)	RR 0.37 (0.23-0.61)	200 fewer per 1000 (from 124 fewer to 245 fewer)	Low
Unsatisfactory pain control postoperatively	1/20 (5%)	10/20 (50%)	RR 0.1 (0.01-0.71)	549 fewer per 1000 (from 177 fewer to 604 fewer)	Low
	15/21 (71.5%)	15/21 (71.5%)	RR 1 (0.68-1.47)		
Nausea and/or vomiting	18/141 (12.8%)	25/159 (15.7%)	RR 1.05 (0.63-1.75)	8 more per 1000 (from 58 fewer to 118 more)	Moderate
Need for anti-emetics	0/20 (0%)	5/20 (25%)	RR 0.09 (0.01-1.54)	227 fewer per 1000 (from 248 fewer to 135 more)	Low
Wound infection	0/28 (0%)	2/27 (7.4%)	RR 0.019 (0.01-3.85)	60 fewer per 1000 (from 73 fewer to 164 more)	Moderate
Pneumonia	12/129 (9.3%)	25/130 (19.2%)	RR 0.49 (0.26-0.94)	98 fewer per 1000 (12 fewer to 142 fewer)	Moderate

Any cardiac complication	3/62 (4.8%)	12/62 (19.4%)	RR 0.25 (0.07-0.84)	145 fewer per 1000 (from 31 fewer to 180 fewer)	Low
Myocardial infarction	1/34 (3%)	4/34 (12%)	RR 0.25 (0.03-2.12)	88 fewer per 1000 (from 114 fewer to 132 more)	Low
Pruritis	0/20 (0%)	5/20 (25%)	RR 0.09 (0.01-1.54)	227 fewer per 1000 (from 248 fewer to 135 more)	Low
Pulmonary embolism	1/53 (1.9%)	2/52 (3.8%)	RR 0.66 (0.11-3.86)	13 fewer per 1000 (31 fewer to 110 more)	Low
Deep vein thrombosis	7/116 (6%)	7/137 (5.1%)	RR 1.12 (0.43-2.93)	6 more per 1000 (29 fewer to 99 more)	Low
Mortality	9/189 (4.8%)	19/205 (9.3%)	RR 0.59 (0.29-1.21)	38 fewer per 1000 (66 fewer to 99 more)	Low
Pressure sores	3/86 (3.5%)	9/106 (8.5%)	RR 0.51 (0.11-2.39)	42 fewer per 1000 (76 fewer to 118 more)	Low
Confusional state	15/77 (19.5%)	34/101 (33.7%)	RR 0.63 (0.37-1.06)	125 fewer per 1000 (212 fewer to 20 more)	Low

1

2 7.3.1.2 Economic evidence

3 No relevant studies were identified. We conducted a cost analysis of a nerve block, non-
4 opioids and other analgesics. We found that a nerve block would cost approximately
5 £54.66. The average cost for opioids controlled drugs is £11.84 (where £1.34 is the average
6 cost per dose of the drugs and £10.50 the personnel cost of two trained nurses required for
7 the administration of the drugs). The price of opioids non-controlled drugs is estimated at
8 £1.96 per doses. The cost of non-opioids analgesics is less than £0.1p per dose. Please see
9 Appendix H section 8.1 for further details.

10 7.3.1.3 Evidence statement (s)

Clinical There is a statistically significant but not clinically significant reduction in pain when using nerve blocks compared to systemic analgesia. (LOW QUALITY). There is a statistically significant but not clinically significant reduction in pneumonia when using nerve blocks compared to systemic analgesia (MODERATE QUALITY).

There is no statistically significant difference between nerve blocks and systemic analgesia in all other outcomes (LOW QUALITY).

Economic No studies on the cost-effectiveness of nerve blocks for hip fracture patients

were identified.

1 7.3.2 Recommendations and link to evidence

Recommendation	<p>Assess the patient's pain:</p> <ul style="list-style-type: none"> • immediately upon presentation at hospital and • within 30 minutes of administering initial analgesia and • hourly until settled on the ward and • regularly as part of routine nursing observations throughout admission.
Relative values of different outcomes	This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.
Trade off between clinical benefits and harms	Regular assessments mean that the patients benefit from analgesia that is tailored to their needs and ensure that the analgesic agents have taken effect. There are no identifiable harms associated with this.
Economic considerations	The GDG agrees that the additional costs linked with the staff time required for regular pain assessment are likely to be offset by the beneficial outcomes of ensuring adequate analgesia.
Quality of evidence	There have been no studies of this approach to achieving adequate analgesia. The recommendation is based on GDG consensus.
Other considerations	<p>Satisfactory and timely pain relief can only be ensured by regular re-assessment.</p> <p>To maintain an adequate level of pain relief, analgesia should be administered routinely and not 'on demand'. It is good practice to re-assess a patient in severe pain after 30 minutes, as analgesia will have taken effect in this time and the need (or not) for additional analgesia can be determined. The 30-minute interval also reflects the pharmacokinetic/pharmacodynamic profiles of morphine and its active metabolite morphine-6-glucuronide. Adequate analgesic response is usual by 15 minutes after administration and should invariably be achieved by 30 minutes. Upward dose titration is otherwise required. The duration of effect varies, ranging from 2 to 24 hours reflecting inter-individual variability in morphine-6-glucuronide clearance and response. If further analgesia is required, the need for subsequent hourly reassessment is justified not only by the need to ensure a satisfactory response, but also to assess any unwanted effects. This hourly interval is also partly pragmatic, consistent with safe, common good clinical practice, and in line with CEM recommendations. For these reasons, the</p>

GDG felt that the recommended 30-minute check to ascertain and achieve initial response, and hourly observation thereafter to determine its duration, together with any adverse effects, are appropriate. The same intervals apply to dosage switches.

Some patients may be unable to express their need for pain relief to health care professionals. Regular assessment of pain and tailoring of medication accordingly will reduce the risk of these patients suffering because of inadequate pain control.

The GDG also considered evidence on patient views. Two studies in which patients mentioned pain management were identified (Section 13.2). In one, pain management did not seem to be a problem³¹⁴. However, in the other the patient had to keep asking for pain relief after surgery²⁷⁴. This highlights the importance of regular assessment.

Additional broad guidance on the assessment of pain in general in older people is given in a joint British Pain Society and British Geriatrics Society document to be found at:
[http://www.bgs.org.uk/Publications/Publication%20Downloads/Se
 p2007PainAssessment.pdf](http://www.bgs.org.uk/Publications/Publication%20Downloads/Se%20p2007PainAssessment.pdf)

<i>Recommendation</i>	Offer immediate analgesia to patients presenting at hospital with suspected hip fracture, including people with cognitive impairment.
Relative values of different outcomes	This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.
Trade off between clinical benefits and harms	Immediate pain control not only improves patients' wellbeing but may reduce the risk of delirium, and facilitate rehabilitation and a return to mobility and independence. The risks of pain relief are the side effects of the individual agents used to achieve it (see below).
Economic considerations	The GDG agrees that the costs of providing immediate and adequate analgesia are likely to be offset by the improvement in patients' wellbeing.
Quality of evidence	There have been no studies on the timing of analgesia on patient outcome. The evidence for efficacy is that of each agent. The recommendation is based on GDG consensus.
Other considerations	It is a humanitarian necessity that these patients receive adequate analgesia, even if cognitively impaired, or limited in their ability to express pain. Particular skill and sensitivity may be required in the management of pain in those who also show signs of delirium (see NICE delirium Guideline ²²⁴)

It must be remembered that patients may require more analgesia for investigations such as X Rays.

1

2

Recommendation	Ensure analgesia is sufficient to allow movements necessary for investigations (as indicated by the ability to tolerate passive external rotation of the leg), and for nursing care and rehabilitation.
Relative values of different outcomes	This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.
Trade off between clinical benefits and harms	Providing adequate levels of analgesia is essential in improving the patients' wellbeing and minimising their discomfort whilst clinical investigations are being carried out. Gentle rotation of the leg may be associated with some degree of pain but would not otherwise cause any additional harm to the patient. There are no other identifiable harms from carrying out this assessment.
Economic considerations	The beneficial outcomes of ensuring that adequate analgesia is provided to allow patients' movements are likely to offset the staff time required).
Quality of evidence	There have been no studies of this approach to achieving adequate analgesia. The recommendation is based on GDG consensus.
Other considerations	In both the pre and postoperative periods if the patient can tolerate passive rotation of the leg then this gives an indication they will be comfortable for preoperative radiographs as well as initial postoperative mobilisation. This procedure should adequately predict the adequacy of analgesia when patients subsequently have to be moved (e.g. on and off examination surfaces) for investigational procedures, such as X-rays.

3

Recommendation	Offer paracetamol every 6 hours preoperatively unless contraindicated.
Relative values of different outcomes	This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.
Trade off between clinical benefits and harms	Simple regular prescribed analgesia such as paracetamol is not associated with any significant harm or side effects. However, it

should be avoided or used with caution in patients with known hypersensitivity to paracetamol and in liver and renal disease.

Economic considerations

The cost of paracetamol is minimal (Appendix H, section 8.1). The administration of paracetamol would be part of routine drug rounds, and therefore it will not involve additional staff or administrative costs.

Quality of evidence

There are no placebo-controlled trials of the efficacy of preoperative administration of paracetamol in hip fracture patients as these are unethical. In a randomised controlled trial, Cuvillion et al 2007⁶² have shown that intravenous paracetamol can be as effective as nerve blocks or morphine in the postoperative phase. However, the dose of paracetamol used in the Cuvillion study was 2mg which exceeds the maximum recommended in the BNF. Therefore, the recommendation could not be solely based on this evidence. There were no studies comparing paracetamol, orally or via the rectal route. Therefore the recommendation was based on consensus.

Other considerations

Complications are especially more likely to develop when stronger analgesia is administered in the elderly. Regular paracetamol is first-line unless contra-indicated.

This and subsequent recommendations follow a logical hierarchy for the use of analgesic agents as indicated in the World Health Organisation pain relief ladder.

1

Recommendation

Offer additional opioids if paracetamol alone does not provide sufficient preoperative pain relief.

Relative values of different outcomes

This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated or by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.

Trade off between clinical benefits and harms

Repeated use of opioids may cause dependence and tolerance. While this should be borne in mind, it should not deter the achievement of effective pain relief in the acute situation of hip fracture. In those for whom the fracture is an incident within the pathway of a terminal illness, the palliative context of that illness should also be an important consideration. In particular, if there is a history of previous opioid use, the existence of acquired tolerance may necessitate the use of higher doses to relieve hip fracture pain. Many older patients may have impaired respiratory function and opioids should be used with caution in these patients. Smaller doses may be required in older patients.

Harm may come from excessive opioid administration:

- Some patients may develop nausea and constipation from stronger opioids and codeine. Regular laxatives may need to be administered.
- Severe constipation may exacerbate other chronic conditions like diverticulitis.
- The significant sedation from even mild opioids in this vulnerable group may slow down their postoperative mobilisation, and upset their balance.

There is a trade off between using stronger analgesia with more side effects and the benefit of better pain relief. Elderly patients are more susceptible to the harmful effects of opioid analgesics.

Opioids and NSAIDs can both cause harm in elderly patients with comorbidities. Most elderly hip fracture patients do have multiple chronic conditions such as decreased renal function, hiatus hernia or previous gastric or duodenal erosions, vertigo, diverticulitis, or mild cognitive problems that may be exacerbated by these forms of analgesia.

Economic considerations	The administration of some opioids requires two trained nurses for approximately 15 minutes. Please see Appendix H section 8.1 for further details. The GDG agrees that the additional costs are likely to be offset by the beneficial outcomes of ensuring adequate analgesia (see Recommendation 1).
Quality of evidence	No studies on the effectiveness of opioids compared to placebo or to other drugs in hip fracture patients were identified.
Other considerations	None

1

Recommendation	Consider adding nerve blocks if paracetamol and opioids do not provide sufficient preoperative pain relief, or to limit opioid dosage. Nerve blocks should be administered by trained personnel. Do not use nerve blocks as a substitute for early surgery.
Relative values of different outcomes	This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important. Adequate pain relief is beneficial. Reduction in the required administration of opioids and the associated side effects may also be an important outcome.
Trade off between clinical benefits and harms	Local nerve blocks are effective and may serve as a means of reducing the need for, and side effects of, opioids and other analgesia. However, as there they are associated with a very rare

incidence of nerve damage, administering them in a busy casualty department may require a rolling programme of training junior doctors or nurses to be competent with this technique.

Economic considerations

The additional cost of nerve blocks versus the cost of opioid drugs may be offset by savings in the resources that would be required to treat the side effects of opioids. The GDG agrees that the additional costs are likely to be offset by the beneficial outcomes of ensuring adequate analgesia.

Quality of evidence

There are a limited number of clinical trials that have examined the effectiveness of nerve blocks in conjunction with general anaesthesia. Some studies have looked at the impact of inserting nerve blocks before the surgical procedure, to see if this may reduce analgesic requirements and improve pain management. These studies show that nerve blocks reduce the degree of pain compared to systemic analgesia alone and that they may have fewer side effects compared to systemic analgesia.

Other considerations

Although studies have shown that nerve blocks are better than systemic analgesia at relieving pain, the GDG considered that this should not be the first line treatment. The GDG wished to ensure that the administration of analgesics is done in a step wise approach as some patients may benefit from simple analgesics such as paracetamol and therefore avoid the more serious side effects of stronger analgesics.

1

2

Recommendation

Offer paracetamol every 6 hours postoperatively unless contraindicated.

Relative values of different outcomes

This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. It is also of central importance in achieving early mobilisation postoperatively. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.

Trade off between clinical benefits and harms

Paracetamol administered first-line and regularly in standard dosage at this frequency is commonly effective and lacks the unwanted effects of second-line systemic agents (see below). It should be avoided or used with caution in patients with known hypersensitivity to paracetamol and in liver and renal disease.

Economic considerations

The cost of paracetamol is minimal. The administration of paracetamol would be part of routine drug rounds, and therefore it will not involve additional staff or administrative costs. (Appendix H, section 8.1.

Quality of evidence

Cuvillion et al have shown that intravenous paracetamol is as

<p>Other considerations</p>	<p>effective as nerve blocks or morphine in the postoperative phase.</p> <p>Paracetamol should be the first option as opioids often sedate patients when they need to be alert to understand and remember important instructions from the physiotherapist on early effective mobilisation. Also opioids may make patients feel dizzy and unconfident about their balance.</p> <p>Postoperatively active mobilisation may require additional pain relief. Pain may be a critical barrier to be overcome for effective early mobilisation.</p>
------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

1

<p>Recommendation</p>	<p>Offer additional opioids if paracetamol alone does not provide sufficient postoperative pain relief.</p>
<p>Relative values of different outcomes</p>	<p>This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. It is also of central importance in achieving early mobilisation postoperatively. Therefore, the GDG considered pain relief (for example as indicated by Visual Analogue Scales or by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.</p>
<p>Trade off between clinical benefits and harms</p>	<p>Opioids do have significant side effects of sedation, nausea, dizziness and constipation. However, pain is also a significant barrier to early mobilisation. Getting the analgesia right at each step of the hip fracture pathway is a skilled judgement for each individual patient until they are discharged.</p> <p>Often opioids sedate patients when they need to be alert to understand and remember important instructions from the physiotherapist on early effective mobilisation. Also opioids may make patients feel dizzy and unconfident about their balance.</p>
<p>Economic considerations</p>	<p>The GDG believe that the side-effects of opioids and additional costs are likely to be offset by the benefits of pain relief.</p>
<p>Quality of evidence</p>	<p>No studies on the effectiveness of opioids compared to placebo or to other drugs in hip fracture patients were identified. This recommendation is based on GDG consensus.</p>
<p>Other considerations</p>	<p>None.</p>

2

<p>Recommendation</p>	<p>Non steroidal anti-inflammatory drugs (NSAIDs) are not recommended.</p>
<p>Relative values of different outcomes</p>	<p>This group of patients is most commonly elderly and frail and pain is one of the main physiological and psychological stresses they face. It is also of central importance in achieving early mobilisation</p>

postoperatively. Therefore, the GDG considered pain relief (for example as indicated by the need for 'breakthrough analgesia') to be the most important outcome. The GDG also considered adverse events outcomes to be important.

Trade off between clinical benefits and harms

The benefits of pain relief are outweighed by the potential side effects of these drugs particularly (but not exclusively) in the elderly population. There is a known age-related increase in susceptibility to the harmful effects of NSAIDs including upper gastrointestinal bleeding, nephrotoxicity and fluid retention.

Economic considerations

The use of NSAIDs is expected to result in a QALY loss, mainly associated with the side effects and adverse events of NSAIDs in our population. The incremental cost savings would have to be considerably high to outweigh these negative benefits, and given the recommended interventions this is highly unlikely.

Quality of evidence

No RCTs on the effectiveness of NSAIDs compared to placebo or to other drugs in hip fracture patients were identified. This recommendation is based on GDG consensus.

Other considerations

The side effects of these drugs are too great in the elderly. Therefore, the GDG decided that they should be avoided as there are other safer alternatives available such as paracetamol and opioids.

As discussed, many of these patients have comorbidities of hiatus hernia, gastric or duodenal erosions, or chronic renal impairment, which can all be made worse by regular use of NSAIDs.

1 **7.4 Research recommendations on analgesia**

2 The GDG recommended the following research question:

- 3 ➤ What is the clinical and cost effectiveness of preoperative and postoperative nerve
4 blocks in reducing pain and achieving mobilisation and physiotherapy goals sooner
5 in patients with hip fracture?

6 **Why this is important**

7 Nerve blocks may potentially find an important role in the management of hip fracture
8 pain, both pre- and postoperatively, because of their potential to reduce the requirement
9 for opioids and their associated unwanted effects. Economically there are considerations
10 for staff training, but also for the potential benefits in terms of duration of stay and early
11 mobilisation. It is not possible from the existing literature to determine this with any
12 confidence and there is a pressing need for a definitive trial comparing these outcomes
13 with nerve blocks against a defined protocol of systemic opioid use.

1 **8 Regional (spinal or epidural) versus general** 2 **anaesthesia**

3 **8.1 Introduction**

4 Patients who have a proximal femoral fracture are usually offered surgery to treat the
5 injury. The vast majority of these operations will require some type of anaesthesia.
6 Anaesthesia may be general anaesthesia or regional anaesthesia.

7 General anaesthesia involves complete loss of consciousness. This may be achieved by
8 either inhalational agents or intravenous anaesthetic agents. Regional anaesthesia is
9 conducted by numbing the nerves that supply sensation to the lower limbs, with the
10 injection of local anaesthetic solution into the fluid surrounding the spinal cord. There are
11 two types of regional anaesthesia, spinal and epidural. During a spinal, local anaesthetic
12 drugs, sometimes in combination with opioid painkillers are injected directly into the
13 cerebro-spinal fluid of the spinal cord. The majority regional anaesthesia administered to
14 hip fracture patients is spinal anaesthesia rather than epidural.

15 Hip fracture patients are generally elderly and have significant comorbidities. This increases
16 the risks from all types of anaesthesia. At present both regional and general anaesthesia are
17 administered but the eventual choice is the preference and experience of the anaesthetist
18 in discussion with the patient and their carers.

19 The aim of this review is to identify whether regional anaesthesia confers any benefit
20 compared to general anaesthesia with regards to reducing complications and improving
21 patient outcomes after surgery.

22 **8.2 Regional versus general anaesthesia**

23 **8.2.1 Review question**

24 In patients undergoing surgical repair or replacement for hip fractures, what is the clinical
25 and cost-effectiveness of regional (spinal/epidural) anaesthesia compared to general
26 anaesthesia in reducing complications such as mortality, cognitive dysfunction,
27 thromboembolic events, postoperative respiratory morbidity, renal failure and length of
28 stay in hospital?

29 **8.2.1.1 Clinical evidence**

1 The literature search retrieved one Cochrane review (Parker et al 2004)²⁶⁶ including 22 RCTs
 2 with a total of 2567 participants. A further update search was then conducted to search for
 3 any papers that may have been published since the publication of this review. No additional
 4 studies were retrieved and therefore the clinical evidence presented in this chapter is based
 5 on the Parker et al results with the addition of the GRADE analysis.

6 In addition, we conducted a systematic review on patient views to look for evidence on
 7 patient preference as this was one of the main outcomes.

8 See evidence table4, Appendix E, forest plots G38 to G49.

9 **Table 8-26: General vs. regional anaesthesia – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality (early up to 1 month) ^{1,20,23,65,66,167,210,211,277,334,339}	11	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious Imprecision (c)
Mortality at 1 month ^{20,65,66,167,210,211,277,339}	8	RCT	Serious limitations (a), (b)	No serious inconsistency (d)	No serious indirectness	Serious Imprecision (c)
Length of stay in hospital ^{210,277}	2	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious imprecision
Vomiting ^{23,211}	2	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious Imprecision (c)
Acute confusional state ^{20,23,46,169,277}	5	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious imprecision (c)
Pneumonia ^{1,20,23,65,66,167,210,211,277}	9	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious Imprecision (c)
Myocardial infarction ^{65,66,167,210,211,277}	6	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious Imprecision (c)
Pulmonary embolism ^{1,20,23,36,65,66,210,211,277}	9	RCT	Serious limitations (a), (b)	No serious inconsistency (e)	No serious indirectness	Serious Imprecision (c)
Deep vein thrombosis ^{36,65,210,211}	4	RCT	Serious limitations (a), (b)	No serious inconsistency	No serious indirectness	Serious Imprecision (c)

- 10 (a) Some of the studies did not report definite allocation concealment
 11 (b) None of the trials clearly stated whether it was an intention to treat analysis
 12 (c) The relatively few events and few patients give wide confidence intervals around the estimate of effect.
 13 This makes it difficult to know the true effect size for this outcome.
 14 (d) Pooling of the results showed some but not statistically significant heterogeneity: $I^2 = 31%$ ($p = 0.18$)
 15 (e) The results of pooling all pulmonary embolism events showed statistical heterogeneity $I^2 = 47%$ ($p = 0.06$).
 16 The authors suggest this is mainly due to the significantly different in trials presenting results for
 17 fatal and non fatal pulmonary embolism. These were subsequently analysed in separate meta-analyses.
 18
 19
 20

1 **Table 8-27: General vs. regional anaesthesia - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk (95% CI)	Absolute effect	Quality
Mortality (early up to 1 month)	64/912 (7%)	93/966 (9.6%)	RR 0.73 (0.54-0.99)	26 fewer per 1000 (from 1 fewer to 44 fewer)	Low
Mortality at 1 month	56/811 (6.9%)	86/857 (10%)	RR 0.69 (0.50, 0.95)	31 fewer per 1000 (from 5 fewer to 50 fewer)	Low
Length of stay in hospital	108	110	N/A	Mean Difference 0.21 (-5.21-4.78)	Low
Vomiting	2/46 (4.3%)	3/49 (6.1%)	RR 0.7 (0.12-3.94)	18 fewer per 1000 (from 54 fewer to 179 more)	Low
Acute confusional state	11/117 (9.4%)	23/120 (19.2%)	RR 0.5 (0.26-0.95)	96 fewer per 1000 (from 10 fewer to 142 fewer)	Low
Pneumonia	21/574 (3.7%)	29/612 (4.7%)	RR 0.76 (0.44-1.3)	11 fewer per 1000 (from 26 fewer to 14 more)	Low
Myocardial infarction	5/502 (1%)	11/531 (2.1%)	RR 0.55 (0.22-1.37)	9 fewer per 1000 (from 16 fewer to 8 more)	Low
Pulmonary embolism	9/605 (1.5%)	13/640 (2%)	RR 0.88 (0.32-2.39)	2 fewer per 1000 (from 14 fewer to 28 more)	Low
Deep vein thrombosis	39/129 (30.2%)	61/130 (36.9%)	RR 0.64 (0.48-0.86)	169 fewer per 1000 (from 66 fewer to 244 fewer)	Low

2

3 **8.2.1.2 Economic evidence**

4 One study was identified. Chakladar 2010⁴⁸ is a cost study of general vs. spinal anaesthesia based
5 on a survey. Please see Economic Evidence table 6.1 in Appendix F for further details.

6 **Table 8-28: General anaesthesia vs regional anaesthesia- Economic study characteristics**

Study	Limitations	Applicability	Other Comments
Chakladar 2010⁴⁸	Potentially serious limitations (a)	Partially applicable ^(b)	Cost analysis of general anaesthesia vs. spinal anaesthesia.

7 (a) *Not a full economic evaluation – costs but not health effects. Cost analysis based on responses to a*
8 *questionnaire, not on a direct audit of equipment usage. Overhead costs and cost of treating side*
9 *effects were not included. No sensitivity analysis.*

10 (b) *UK study but does not estimate QALYs.*

11

1 **Table 8-29: General anaesthesia vs regional anaesthesia - Economic summary of findings**

Study	Incremental cost (£)	Incremental effects	ICER	Uncertainty
Chakladar 2010 ⁴⁸	76.77 ^(a)	NA	NA	NR

2 (a) General anaesthesia more costly than regional anaesthesia (SD):£270.58 (44.68) vs 193.81
3 (44.68); $p < 0.0001$

4 **8.2.1.3 Evidence statement (s)**

Clinical There is a statistically and clinically significant reduction in early mortality (up to 1 month) in patients having regional anaesthesia compared to those having general anaesthesia (LOW QUALITY).

There is a statistically significant but not clinically significant improvement in postoperative confusion and reduction in incidence of deep vein thrombosis in patients receiving regional compared to general anaesthesia (LOW QUALITY).

There were no statistically significant differences in length of stay in hospital, vomiting, pneumonia, myocardial infarction and pulmonary embolism (LOW QUALITY).

Economic One study found general anaesthesia to be more costly than spinal anaesthesia. This evidence has very serious limitations since it did not evaluate effectiveness and may not have included all important cost differences.

5 **8.2.2 Recommendations and link to evidence****Recommendation**

Offer patients a choice of spinal or general anaesthesia after discussing the risks and benefits.

Relative values of different outcomes

The GDG considered early mortality (up to 1 month) and patient preference to be the most important outcomes.

Trade off between clinical benefits and harms

Most clinical benefit was seen in patients undergoing regional anaesthesia. However, there is a small chance of nerve damage following regional anaesthesia.

Potential benefits with regional also include, reduction in venous thromboembolic (VTE) complications but studies were conducted in patients not receiving VTE prophylaxis which may lead to some false positive results. However, this finding is supported by a more comprehensive review of DVT and PE across all surgical patients in the NICE guideline on venous thromboembolism prophylaxis²²⁵.

A potential benefit of general anaesthesia includes lack of awareness throughout the surgical procedure. Indeed some patients perceive unconsciousness during general anaesthesia as a benefit. However, others fear the loss of control. A potential disadvantage of general anaesthesia is that recovery on the first postoperative day may be slower.

Economic considerations	<p>The GDG felt that because of the potentially serious limitations of the study included as economic evidence there were insufficient data to claim that the overall costs of the general and regional anaesthesia are substantially different.</p> <p>However, there was agreement in acknowledging that spinal anaesthesia usually involves lower costs for drugs, anaesthesia equipment and airway equipment than general anaesthesia.</p> <p>Nevertheless, these lower costs of regional anaesthesia could be offset by its longer administration time. The GDG debated at length whether regional anaesthesia required more time to be administered compared to general anaesthesia, but no agreement was reached.</p>
Quality of evidence	<p>The studies comparing the two types of anaesthesia were mainly of low methodological quality. They included small numbers of participants and only reported a few outcome measures. These varied between studies making pooling of the data difficult. The studies lacked methodological rigour in particular regarding allocation concealment, assessor blinding and intention to treat analysis. The studies are now considered to be out of date and no longer relevant to current anaesthesia and perioperative care. In addition, they do not account for the advances in safety in the field of anaesthesia. For example in some of the studies patients allocated to general anaesthesia did not receive thromboprophylaxis as part of routine care.</p> <p>The economic evidence has very serious limitations, as it is based on responses to a questionnaire on a hypothetical anaesthetic technique, and not a direct audit of actual equipment usage. Moreover, the analysis did not look at whether there are any potential savings linked to a reduction in the cases of confusion when regional anaesthesia is used.</p>
Other considerations	<p>The GDG also considered the evidence for other outcomes such as length of stay in hospital and adverse events including vomiting, acute confusional state and respiratory and cardiac complications. In the absence of any strong evidence favouring one method over the other, the GDG decided that the choice of anaesthesia should be based on the patient preference after being given sufficient information about the options available and the expertise of the anaesthetist.</p>

1

Recommendation	Consider intraoperative nerve blocks for all patients undergoing surgery.
Relative values of different outcomes	The GDG considered pain relief, postoperative mobility and reduction in opioid usage to be the main outcomes.

Trade off between clinical benefits and harms	As discussed in chapter 7 on using nerve blocks for hip fracture analgesia, local nerve blocks may serve as a means of reducing the need for, and side effects of, opioids and other analgesia. However, they are associated with a very rare incidence of nerve damage and must be administered by trained health care professionals.
Economic considerations	The GDG agreed this likely to be cost-effective because the administration of nerve blocks avoids the complications and side effects of opioids, and therefore might result in a reduced length of hospital stay. Please see the analgesia chapter for evidence on the cost-effectiveness of nerve blocks in general.
Quality of evidence	The evidence that nerve blocks reduce the degree of pain and the requirement for opioid analgesics compared to other forms of analgesia alone, and that they may have fewer side effects compared to systemic analgesia, is presented under Analgesia (Chapter 7). This includes several studies that have investigated the effectiveness of nerve blocks in conjunction with general anaesthesia to determine if this reduces the requirements for opioid analgesics and improve pain management. These studies show that nerve blocks reduce the degree of pain compared to systemic analgesia alone and that they may have fewer side effects compared to systemic analgesia. However, these studies could not be subgrouped in a meaningful way as they looked at different outcomes and differed in the way they reported them. Therefore, this recommendation was partly based on consensus.
Other considerations	Nerve blocks are often administered before a spinal anaesthetic, in order to position the patient. They are usually administered before a general anaesthetic and many are now conducted using ultrasound guidance. This reduces the chance of complications, such as an intraneural injection and also enables the dose of local anaesthetic administered to be lower. The use of nerve blocks in surgery has increased in recent years and has almost become routine practice. Therefore, studies to show any benefit may now be difficult to conduct, as withholding analgesia from such patients may be unethical. Administration of nerve blocks should not delay surgery.

1

2

3

8.3 Research recommendation on anaesthesia

4

The GDG recommended the following research question:

5

- What is the clinical and cost effectiveness of regional versus general anaesthesia on postoperative morbidity in patients with hip fracture?

6

7

Why this is important

1 No recent randomised controlled trials were identified that fully address this question. The
2 evidence is old and does not reflect current practice. In addition, in most of the studies the
3 patients are sedated before regional anaesthesia is administered and this is not taken into
4 account when analysing the results. The study design for the proposed research would be
5 best addressed by a randomised controlled trial. This would ideally be a multi-centred trial
6 including 3,000 participants in each arm. This is achievable if one considers that there are
7 70, 000 hip fractures a year in the UK³⁹. The study should have three arms which look at
8 spinal anaesthesia versus spinal anaesthesia plus sedation versus general anaesthesia, this
9 would separate those with regional anaesthesia from those with regional anaesthesia plus
10 sedation. The study would also need to control for surgery, especially type of fracture,
11 prosthesis and grade of surgeon.

12 A qualitative research component would also be helpful to study patient preference for
13 type of anaesthesia.

14

15

1 **9 Surgeon seniority**

2 **9.1 Introduction**

3 As a general observation of life one would conclude that to have a job completed
4 thoroughly, effectively and efficiently it would be appropriate to give the task to somebody
5 with adequate training and experience. Whether this can be extrapolated to the
6 relationship of the management of hip fractures to the seniority of the surgeon involved is
7 the purpose of this chapter.

8 The historical background of this question has to be considered in relation to the
9 environment in which hip fracture patients were treated. In the United Kingdom hip
10 fractures were commonly regarded as the surgical material for trainee surgeons to gain
11 their experience. In the past much of this work would have been unsupervised, and in the
12 main the trainees would have enjoyed the challenge and responsibility this gave them.

13 The operations were often performed outside of scheduled list times as extra or emergency
14 cases. Under these circumstances it was more likely that the anaesthetist involved in the
15 procedure would be more junior and the nursing scrub team not specifically from a trauma
16 theatre.

17 Any variations in outcome which may be simply labelled as related to surgeon seniority may
18 in fact have multiple underlying causes. A more senior surgeon is more likely to be
19 operating on a scheduled list, with more senior anaesthetists and a regular nursing scrub
20 team.

21 **9.2 Surgeon seniority**

22 **9.2.1 Review question**

23 What is the clinical and cost effectiveness of surgeon seniority (consultant or equivalent) in
24 reducing the incidence of mortality, the number of patients requiring reoperation, and poor
25 outcome in terms of mobility, length of stay, wound infection and dislocation? (See
26 evidence table 5, Appendix E and forest plots G50 and G51 in Appendix G).

27 **9.2.2 Clinical evidence**

28 No randomised evidence was identified. Three prospective cohorts including 2018
29 participants that adjusted for some confounding factors were identified.

1

2 **Table 9-30: Junior/less senior surgeon vs. senior surgeon – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Reoperations (follow up 6 months) ²⁵⁶	1	Cohort	serious limitations (a,b)	no serious inconsistency	serious indirectness ^(c,d,e)	serious imprecision ^(h)
Dislocation in hemiarthroplasty (follow up 0 to 10 years) ⁸⁵	1	Cohort	serious limitations (b)	no serious inconsistency	serious indirectness (f,g)	serious imprecision ^(h)
Dislocation in total hip replacement (follow up 0 to 11 years) ⁸⁵	1	Cohort	serious limitations (c)	no serious inconsistency	serious indirectness (f,g)	serious imprecision ^(h)

3

(a) Senior surgeons operated on significantly more patients with a poor pre-fracture mobility score and performed significantly more arthroplasties and significantly fewer osteosyntheses.

4

5

(b) Only a limited number of confounders were included in the analysis. No adjustment or mention of the anaesthetists experience or grade.

6

7

(c) Surgeon seniority measured by years experience rather than the grade of surgeon. Experienced surgeons with more than 3 years orthopaedic surgical experience either performing surgery or supervising junior registrars were compared unsupervised orthopaedic junior registrars with less than 3 years orthopaedic surgical experience.

8

9

10

(d) Only the technically demanding fractures were included in the analysis, not all surgery for hip fractures.

11

12

(e) Reoperation rate only measured at 6 months, not longer.

13

14

(f) The focus of the study is on surgical approach therefore baseline data by surgeon seniority is not reported.

15

16

(g) Dislocation is not a primary outcome.

17

(h) The wide confidence intervals make the estimate of effect imprecise.

18

19

Table 9-31: Junior/less senior surgeon vs senior surgeon – Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Reoperations (follow up 6 months)	16/56 (28.6%)	47/309 (15.2%)	multivariate odds ratio 2.01 (1.01 to 4.02)	289 more per 1000 (from 3 more to 864 more)	Very low
Dislocation in hemiarthroplasty (median follow up 4.3 (0 to 10) years)	37/404 (9.2%)	8/135 (5.9%)	multivariate odds ratio 1.3 (0.6 to 3)	18 more per 1000 (from 24 fewer to 118 more)	Very low
Dislocation in total hip replacement (median follow up 2.3 (0 to 11) years)	37/636 (5.8%)	8/77 (10.4%)	multivariate odds ratio 0.9 (0.3 to 2.8)	10 fewer per 1000 (from 73 fewer to 187 more)	Very low

20

21 **9.2.2.1 Economic evidence**

22

No studies were identified on the cost-effectiveness of junior/less senior surgeon vs. senior surgeon. However, we conducted a cost-analysis around the personnel cost of

23

1 a planned trauma list compared to the personnel cost of a general emergency
 2 theatre. We found that a planned trauma list involves an additional cost per hour of
 3 £94, See Appendix H section 20.2 for further details.

4

5 **9.2.2.2 Evidence statement (s)**

Clinical There is a statistically significant, but not clinically significant increased reoperation rate at 6 months with unsupervised junior orthopaedic registrars with less than 3 years experience than with experienced surgeons with more than 3 years experience. (VERY LOW QUALITY).

There is no statistically significant difference between Swedish post registrars and registrars in dislocation rate at a median follow up of 2.3 years after hemiarthroplasty in patients with hip fracture. (VERY LOW QUALITY).

There is no statistically significant difference between Swedish post registrars and registrars in dislocation rate at a median follow up of 2.3 years after total hip replacement in patients with hip fracture. (VERY LOW QUALITY).

There was no evidence identified for mortality, mobility, length of stay or wound infection.

Economic No studies were identified on the cost-effectiveness of junior/less senior surgeon vs. senior surgeon. However, we conducted a cost-analysis around the personnel cost of a planned trauma list compared to the personnel cost of a general emergency theatre. We found that a planned trauma list involves an additional cost per hour of £94, See Appendix H section 20.2 for further details.

6

7 **9.3 Recommendations and link to evidence**

<i>Recommendation</i>	Schedule hip fracture surgery on a planned trauma list
Relative values of different outcomes	Mortality, reoperation rate, dislocations, length of stay in secondary care and wound infection were considered the main outcomes. Complications, pain and functional status were also considered.
Trade off between clinical benefits and harms	No RCTs were identified evaluating a planned trauma list. Evidence is extrapolated from the surgeon seniority data. This shows a significantly higher reoperation rate with unsupervised/junior orthopaedic surgeons with less than 3 years experience than senior more experienced surgeons. There was no statistically significant difference in dislocation rates. No other outcomes were reported.
Economic considerations	A planned trauma list consists of a period of time allocated to the surgical management of patients with unplanned admissions following musculoskeletal injury. For this period there will be an

adequate operating theatre, with supporting equipment including an image intensifier. The responsible senior surgical, anaesthetic and theatre staff will have work plan allocating time to the list to carry out procedures or supervise their junior staff. Thus, a planned trauma list implies allocation and involvement of senior staff, who will either carry out the necessary procedures in the operating theatre or will adequately supervise the junior staff.

The GDG suggested that a possible comparator for a planned trauma list could be a **general emergency theatre**, shared by many different specialities, often occurring outside of normal working hours and staffed by trainees.

If we consider the case of a planned trauma list where operations are performed by a consultant surgeon and a consultant anaesthetist and if we take as comparator a general emergency theatre where both surgeon and anaesthetist are registrars, and we assume no other difference in the professional grade of the remaining staff involved in the operation, then the planned trauma list would result in an additional personnel cost per hour of £94 over the general emergency theatre. In particular, the personnel cost per hour for a planned trauma list with a consultant surgeon and consultant anaesthetist correspond to £337, and for a general emergency list with a registrar surgeon and a registrar anaesthetist (and with a consultant surgeon and consultant anaesthetist on call), to £243 (please see Appendix H section 8.2 for further details). However, there is great uncertainty as to whether there are other differences in other categories of costs (e.g. overheads, diagnostic devices, etc) between a planned trauma list and a general emergency theatre, and therefore our estimate should be considered only as an approximation of the overall cost difference between a planned trauma list and a general emergency theatre. Furthermore, there is uncertainty around the right baseline intervention as after the introduction of the BPT for hip fracture, senior staff should be performing the surgery. In particular, the GDG noted that it is not clear as to what we should consider as the usual alternative to a planned trauma list, as it is quite uncertain what could represent the “baseline” case for a hospital, and this can change depending on the type of hospital. It was also pointed out that since the introduction of the Best Practice Tariff (BpT) for hip fracture in April 2010 the hospitals that do not have planned trauma list in place on a daily basis would however have employed relevant senior staff (consultant surgeons and anaesthetist) to meet the tariff’s requirements, and therefore senior staff are already part of the comparator.

Nevertheless, the GDG thinks that these potential additional personnel costs of a planned trauma list would be at least partially off-set by savings due to lower re-operation rates and by a higher number of patients operated per hour.

Quality of evidence

No RCTs were identified evaluating a planned trauma list. There is

extrapolated evidence from surgeon seniority showed no evidence for the majority of the outcomes and only very low quality evidence from non-randomised studies for two outcomes: reoperation rate and dislocations. The recommendation is based on a consensus agreement within the GDG.

Other considerations

We have specified in the recommendation that surgery for hip fractures should occur on a planned trauma list. To establish a scheduled trauma list management and clinicians are required to provide adequate facilities and staff for it to run. For a planned list it is necessary to have a chain of responsibility to a consultant surgeon and consultant anaesthetist who have time in their programs to execute that responsibility. To run a planned trauma list requires ready access to an image intensifier and radiographer. The nursing team would need to be appropriate to the work planned for that theatre. The recommendation therefore recognises the need for adequate seniority of the surgeon but makes what we believe to be a reasonable assumption that this recognition should also apply to the rest of the operating theatre team caring for the hip fracture patient.

The GDG noted that there is high uncertainty regarding the implementation costs linked with this recommendation, as these costs will vary depending on the current set up and infrastructure of each hospital. For example, the GDG recognised that smaller hospitals may not currently provide this service at weekends.

This recommendation is in line with the British Orthopaedic Association's Advisory book on consultant trauma and orthopaedic services³⁸. The GDG consider this recommendation a key priority for implementation.

1

Recommendation	Consultants or senior staff should supervise trainee and junior members of the anaesthesia, surgical and theatre teams when they carry out hip fracture procedures.
Relative values of different outcomes	Mortality, reoperation rate, dislocations, length of stay in secondary care and wound infection were considered the main outcomes. Complications, pain and functional status were also considered.
Trade off between clinical benefits and harms	There is a significantly higher reoperation rate with unsupervised/junior orthopaedic surgeons with less than 3 years experience than senior more experienced surgeons. There was no statistically significant difference in dislocation rates. No other outcomes were reported.
Economic considerations	Higher grade surgeons or those with more experience are likely to be entitled to a higher wage than junior surgeons. However, as their rate of re-operations is statistically significantly lower, having hip fracture patients operated on by experienced surgeons will plausibly result in cost savings and improved health outcomes. In addition, the GDG believe experienced surgeons use theatre time

more efficiently allowing greater throughput of cases.

Quality of evidence

There is no evidence for the majority of the outcomes and only very low quality evidence from non-randomised studies for two outcomes: reoperation rate and dislocations.

Other considerations

The level of supervision required for a trainee or junior staff member for a particular case depends on two main factors: the junior's ability and the complexity of the case. It is therefore implicit that the senior staff responsible for the trauma list must have knowledge of both of these factors before determining the level of supervision required. Potential surgical, anaesthetic or nursing problems may be evident to an experienced surgeon, anaesthetist or nurse preoperatively. This gives the opportunity to both avoid the problem occurring and to enhance the training opportunity. An unsupervised list would therefore be one in which those responsible did not have adequate prior knowledge of the capabilities of the more junior members of the team and the specific problems they may encounter, or when they did not use this knowledge to provide adequate supervision.

1

2

1 **10 Surgical procedures**

2 **10.1 Introduction**

3 The options for hip fracture surgery depend on the type of fractures. They can be divided
4 into two main groups according to their relationship to the capsular attachment of the hip
5 joint. Those above the insertion of the capsule are termed intracapsular and those below
6 are termed extracapsular. Extracapsular fractures can be further divided into three types:
7 pertrochanteric (also called intertrochanteric), reverse oblique or subtrochanteric.
8

9 Broadly speaking there are two surgical options for treating hip fractures, replacement
10 arthroplasty or internal fixation. Replacement arthroplasty involves removing part or all of
11 the damaged bone and replacing it with a prosthesis which then functions in place of the
12 removed bone. It may describe a hemiarthroplasty or a total hip arthroplasty. Both involve
13 replacement of the femoral head with a metal implant, the stem of which is secured in the
14 femoral shaft. A total hip arthroplasty involves, in addition, replacement of the socket. Both
15 implants can be inserted with or without the use of cement. Internal fixation involves
16 returning the bone fragments to an acceptable position and then holding that position with
17 screws, plates or nails. This should allow healing of the fracture fragments in an acceptable
18 position for long term function and maintenance of patient function whilst that healing
19 occurs.
20

21 **10.2 Surgery with regard to early mobilisation**

22 This section relates to the section on early mobilisation (chapter 11) as well as surgery.
23 When embarking on any surgical procedure there should be a clear objective. In
24 orthopaedic and trauma surgery it is easy to attach a rather bland aim of "safe restoration
25 of function". Prior to any surgery commencing the surgeon should already know what his
26 planned postoperative care of that patient is to be. Given the poor reserve functional
27 capacity of many hip fracture patients any prescribed limits on mobility and weight-bearing
28 may significantly alter and restrict their postoperative care. In particular unnecessary
29 restriction of weight-bearing has the potential to compromise independence, discharge
30 destination, general health and final level of function. As a consequence of these
31 considerations, and as a result of the recommendation for early mobilisation (section
32 11.2.2) the GDG felt it appropriate to make a recommendation on postoperative weight-
33 bearing status.

1 10.2.1 Recommendations and link to evidence

Recommendation	Operate on patients with the aim to allow them to fully weight bear (without restriction) in the immediate postoperative period.
Relative values of different outcomes	The aim of surgery and rehabilitation is for patients to regain their prefracture functional status. Early mobilisation with a physiotherapist appears safe and is effective in promoting early recovery. The most important outcomes considered by the GDG were functional status, mobility, pain and quality of life.
Trade off between clinical benefits and harms	The evidence from the early mobilisation question shows that the only outcome relating to harm or safety was mortality, which showed no statistically significant difference. If safety issues were a concern it is likely that they would be reflected in the overall functional outcomes, all of which improved or had no significant effect, therefore we don't believe that harm is caused harm from this evidence.
Economic considerations	See also early mobilisation section 8.2. One of the main aims of surgery is for patients to regain their pre-fracture functional status. As the GDG has agreed to consider early mobilisation strategy as a cost-effective intervention for our population, this recommendation is unlikely to result in extra costs.
Quality of evidence	There is no direct evidence relating to this recommendation, but the evidence from the early mobilisation review question is indirectly applicable, see Chapter 8.
Other considerations	Elderly patients may be physically frail, suffering from cognitive impairment or delirium and so cannot be expected to mobilise non-weight-bearing or partially weight-bearing. Postoperative instructions requesting non-or partial weight-bearing will frequently result in the patient not mobilising at all.

2
3

4 10.3 Displaced intracapsular fractures

5 In an intracapsular fracture the proximal fragment includes the femoral head alone or the
6 femoral head with a small portion of neck. The size and shape of this fragment combined
7 with the often soft nature cancellous bone of which it is constituted makes secure fixation
8 difficult. This can potentially compromise early function. In addition, the blood supply of
9 the femoral head may be disrupted, leading to poor healing or bone death.

10
11 The displacement of an intracapsular fracture is determined on the anteroposterior and
12 lateral radiographs of the area. An undisplaced fracture may as its name suggests
13 demonstrate no change in position from that it would have occupied prior to the injury.
14 However it is also customary to include in the undisplaced group valgus impacted fractures.
15 In this impacted group the harder bone of the femoral neck has been driven into the softer
16 bone of the femoral head. In both of these these undisplaced fracture types there is

1 generally already inherent stability and little likelihood of damage to the blood supply.
2 Fixation in situ is generally accepted

3
4 In practice a displaced fracture is one in which the preoperative radiographs demonstrate
5 the fragments have moved in relation to each other to an unacceptable position for fixation
6 in situ. The implication of this is that the fragments have moved in relation to each other to
7 a greater extent. The particular anatomy of the region means that the blood supply to the
8 femoral head is at risk. There will also be less inherent stability either as a consequence of
9 fragmentation along the fracture line or difficulties in obtaining precise reduction.

10
11 In patients with these displaced intracapsular fractures a decision initially needs to be made
12 as to whether to reduce the fracture and internally fix it or to carry out some form of
13 replacement arthroplasty. Each has potential advantages and disadvantages. Internal
14 fixation retains the patient's own tissues and is often a smaller procedure. However, it may
15 require a more prescriptive postoperative regime to protect the healing bone. Should
16 replacement arthroplasty be appropriate it is necessary to determine the indications for a
17 hemiarthroplasty in which only the damaged bone of the proximal femur is replaced or a
18 total hip replacement when both the femoral head and the hip socket are replaced.

19 10.3.1 Internal fixation versus hemiarthroplasty

20 10.3.1.1 Review question

21 In patients having treatment for displaced intracapsular hip fracture what is the clinical and
22 cost effectiveness of internal fixation compared to hemiarthroplasty on mortality, number
23 of reoperations, functional status, length of stay in hospital, total time to resettlement in
24 the community, quality of life, pain and place of residence after hip fracture.

25 One systematic review²⁶⁴ was identified and one additional RCT¹⁰². Overall, there were 13
26 RCTs with 2195 participants. See evidence table 7, Appendix E and forest plots G74 to G82
27 in Appendix G.

28 10.3.1.2 Clinical evidence

29 Table 10-32: Internal fixation vs hemiarthroplasty – Clinical study characteristics

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality at 1 month ¹⁰²	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality at 3 to 6 months ^{27,102,161,174,267,276,317,324,341,343}	10	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality at 1 year ^{27,102,161,174,267,317,324,341,343}	9	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Mortality at 2 to 3 years ^{27,102,161,174,267,276,317,324,341,343}	10	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Total no. of reoperations (follow-up 1 to 5 years) ^{27,69,102,161,174,267,276,288,313,317,324,341,343}	13	RCT	serious limitations ^(a)	serious inconsistency ^(c)	no serious indirectness	no serious imprecision
Failure to return to same residence (follow-up 1 to 3 years) ^{161,267}	2	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Failure to regain mobility (follow-up 1 to 5 years) ^{27,161,267,288,317,341}	6	RCT	serious limitations ^(a)	serious inconsistency ^(f)	no serious indirectness	serious imprecision ^(b)
No. of patients reporting pain at 1 year ^{27,174,267}	3	RCT	serious limitations ^(a)	serious inconsistency ^(d)	no serious indirectness	serious imprecision ^(b)
Harris Hip Score (follow-up 1 year) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Harris Hip Score (follow-up 2 years) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Number of patients with Barthel Index Score of 95 or 100 (follow-up 1 year) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Number of patients with Barthel Index Score of 95 or 100 (follow-up 2 years) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Eq-5d (Euroqol) Index Score (follow-up 1 year) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Eq-5d (Euroqol) Index Score (follow-up 2 years) ¹⁰²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)
Length of hospital stay ^{102,174,267,341}	4	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(e)

1 (a) The studies with the most weight in the meta-analysis have inadequate or unclear allocation
2 concealment.

3 (b) The relatively few events and few patients give wide confidence intervals around the estimate of
4 effect. This makes it difficult to know the true effect size for this outcome.

- 1 (c) There is significant unexplained statistical heterogeneity between the studies. This could be due to
 2 the different types of implant or arthroplasty and different follow up periods.
 3 (d) There is significant statistical heterogeneity between the studies. This could be due to the different
 4 types of implant or arthroplasty.
 5 (e) The wide confidence intervals around the estimate make the result imprecise. Consequently, it is
 6 difficult to determine the true effect size for this outcome.
 7 (f) There is significant statistical heterogeneity between the studies. This Cochrane review reports this
 8 is likely to be due to the variation in the definition for this outcome.
 9

10 **Table 10-33: Internal fixation vs hemiarthroplasty - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality at 1 month	7/112 (6.3%)	10/110 (9.1%)	RR 0.69 (0.27 to 1.74)	28 fewer per 1,000 (from 66 fewer to 67 more)	Low
Mortality at 3 to 6 months	107/765 (14%)	122/709 (16.7%)	RR 0.81 (0.64 to 1.03)	32 fewer per 1,000 (from 60 fewer to 5 more)	Low
Mortality at 1 year	148/636 (23.3%)	143/584 (23.6%)	RR 0.93 (0.78 to 1.12)	17 fewer per 1,000 (from 52 fewer to 28 more)	Moderate
Mortality at 2 to 3 years	265/750 (35.3%)	254/683 (37.8%)	RR 0.96 (0.84 to 1.09)	15 fewer per 1,000 (from 60 fewer to 34 more)	Moderate
Total no. of reoperations (follow- up 1 to 5 years)	355/1001 (35.5%)	99/1033 (9.4%)	RR 3.59 (2.93 to 4.39)	243 more per 1,000 (from 181 more to 319 more)	Low
Failure to return to same residence (follow-up 1 to 3 years)	29/187 (15.5%)	34/185 (23.6%)	RR 0.84 (0.54 to 1.33)	38 fewer per 1,000 (from 109 fewer to 78 more)	Low
Failure to regain mobility (follow-up 1 to 5 years)	155/287 (54%)	165/306 (45.7%)	RR 1.02 (0.74 to 1.39)	9 more per 1,000 (from 119 fewer to 178 more)	Very low
No. of patients reporting pain at 1 year	126/280 (45%)	127/281 (44.2%)	RR 0.97 (0.66 to 1.44)	13 fewer per 1,000 (from 150 fewer to 194 more)	Very low
Harris Hip Score (follow-up 1 year)	87	74	N/A	MD -6.8 (-12 to -1.6)	Moderate
Harris Hip Score (follow-up 2 years)	71	68	N/A	MD -3.3 (-9.1 to 2.5)	Moderate
Number of patients with Barthel Index Score of 95 or 100 (follow-up 1 year)	31/87 (35.6%)	39/73 (53.4%)	RR 0.67 (0.47 to 0.95)	176 fewer per 1,000 (from 27 fewer to 283 more)	Moderate
Number of patients with Barthel Index Score of 95 or 100 (follow-up 2 years)	24/69 (34.8%)	26/68 (38.2%)	RR 0.91 (0.58 to 1.42)	34 fewer per 1,000 (from 160 fewer to 160 more)	Moderate
Eq-5d (Euroqol) Index Score (follow-up 1 year)	70	62	N/A	MD -0.09 (-0.2 to 0.02)	Moderate

Eq-5d (Euroqol) Index Score (follow-up 2 years)	52	52	N/A	MD -0.11 (-0.21 to -0.01)	Moderate
Length of hospital stay	486	478	N/A	MD -0.6 (-2.04 to 0.83)	Moderate

1

2 **10.3.1.3 Economic evidence**

3 Two economic studies were identified ^{173,291}. Rogmark et al (2003)²⁹¹ is a cost-consequence
4 analysis based on a RCT but it was excluded because it does not distinguish patients on the
5 basis of whether they received hemiarthroplasty or total hip replacement. Keating et al
6 (2005)¹⁷³ compare internal fixation vs. hemiarthroplasty in a cost-consequence analysis
7 based on a RCT. Please see Economic Evidence Table 14 in Appendix F for further details

8

9 **Table 10-34: Internal Fixation vs Hemiarthroplasty - Economic study characteristics**

Study	Limitations	Applicability	Other Comments
Keating 2005 ¹⁷³	Minor limitations ^(a)	Partially applicable ^(b)	Costs not discounted because mainly incurred within 1 year of injury

10 ^(a) Small number of patients.11 ^(b) UK study, but does a CUA.

12

13 **Table 10-35: Internal Fixation vs Hemiarthroplasty - Economic summary of findings**

Study	Incremental cost per patient (£)	Incremental effects	ICER	Uncertainty
Keating 2005 ¹⁷³	£2726(a)	Various (b)	N/A	Two-way sensitivity analysis showed that the direction of change in cost did not change when cost of prostheses and cost of readmission were varied over a range from -50% to +100% around the baseline values.

14 ^(a) The mean cost per patient for internal fixation was £12,623 (95% CI: 10,768 – 14,478) and for £9,897
15 (95% CI: 8,062 – 11,732) for hemiarthroplasty (2001 GBP)16 ^(b) Several outcomes were reported. Internal fixation entailed lower mortality at 4 and 12 months from the
17 operation than hemiarthroplasty (3% vs. 5%; 8% vs. 10%) and slightly higher EQ-5D scores at 24 months
18 (0.55 vs 0.53); (all effects were not statistically significant). Hemiarthroplasty involved a significantly
19 lower number of patients needing further surgery at 12 and 24 months (31% vs. 5% and 39% vs. 5%),
20 and higher EQ-5D scores at 4 and 12 months (0.56 vs. 0.61 and 0.58 vs. 0.64; difference not statistically
21 significant).
2223 **10.3.1.4 Evidence statement (s)**

Clinical There is a statistically and clinically significant decrease in patients who require reoperations with hemiarthroplasty than with internal fixation. The follow up varied between 1 and 5 years. (LOW QUALITY)

There is a statistically significant, but not clinically significant, increase in patients who have a Barthel Index Score of 95 or 100 at 1 year with

hemiarthroplasty compared to internal fixation but there is no statistically significant difference at 2 years (MODERATE QUALITY)

There is a statistically significant, but not clinically significant, increase in patients who have a higher Harris Hip Score at 1 year with hemiarthroplasty compared to internal fixation but there is no statistically significant difference at 2 years (MODERATE QUALITY)

There is a statistically significant, but not clinically significant, increase in patients who have a higher Eq-5d (Euroqol) score at 2 years with hemiarthroplasty compared to internal fixation but there is no statistically significant difference at 1 year (MODERATE QUALITY)

There is no statistically significant difference between internal fixation and hemiarthroplasty in mortality at 1 months (LOW QUALITY), 3 to 6 months (LOW QUALITY) or 1 to 2 years (MODERATE QUALITY), the number of patients reporting pain at 1 year (VERY LOW QUALITY), the number of patients failing to return to the same residence at 1 to 3 years (LOW QUALITY), failure to regain mobility at 1 to 5 years and length of hospital stay (MODERATE QUALITY).

No RCT evidence was identified reporting on total time to resettlement in the community.

Economic Hemiarthroplasty is cost saving with respect to internal fixation. This evidence has minor limitations and partial applicability.

1

2 10.3.2 Internal fixation versus total hip replacement

3 10.3.2.1 Review question

4 In patients having treatment for intracapsular hip fracture what is the clinical and cost
5 effectiveness of internal fixation compared to total hip replacement on mortality, number
6 of reoperations, functional status, length of stay in hospital, total time to resettlement in
7 the community, quality of life, pain and place of residence after hip fracture.

8 One systematic review²⁶⁴ was identified. Overall, there were 6 RCTs with 888 participants
9 were included. See evidence table 7, Appendix E and forest plots G83 to 86 in Appendix G.

10 10.3.2.2 Clinical evidence

11 **Table 10-36: Internal fixation vs. total hip replacement – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality at 2 to 4 months ^{162,174,239,327}	4	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality at 12 to 18 months ^{162,174,239}	3	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality at 2 years ^{162,166,174,327}	4	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Reoperations – any (follow-up 1 to 13 years) ^{162,166,174,239,313,327}	6	RCT	serious limitations ^(a)	serious inconsistency ^(c)	no serious indirectness	no serious imprecision
Number of patients reporting pain at 1 year ^{166,174}	2	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Length of hospital stay ¹⁷⁴	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(d)

(a) The studies with the most weight in the meta-analysis have inadequate or unclear allocation concealment.

(b) The relatively few events and few patients give wide confidence intervals around the estimate of effect. This makes it difficult to know the true effect size for this outcome.

(c) There is significant statistical heterogeneity between the studies. This could be due to the different types of implant or arthroplasty and different follow up periods. One study had a 13 year follow up whereas the others varied between 1 and 4 years.

(d) The wide confidence intervals around the estimate make it difficult to determine and effect size for this outcome.

Table 10-37: Internal fixation vs total hip replacement - Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality at 2 to 4 months	15/210 (7.1%)	6/196 (3.7%)	RR 2.21 (0.91 to 5.4)	45 more per 1,000 (from 3 fewer to 163 more)	Low
Mortality at 12 to 18 months)	25/157 (15.9%)	21/147 (10%)	RR 1.08 (0.64 to 1.82)	8 more per 1,000 (from 36 fewer to 82 more)	Low
Mortality at 2 years	44/224 (19.6%)	34/209 (11.6%)	RR 1.18 (0.79 to 1.75)	21 more per 1,000 (from 24 fewer to 87 more)	Low
Reoperations – any (follow-up 1 to 13 years)	126/325 (38.8%)	44/308 (9.4%)	RR 2.70 (1.99 to 3.67)	160 more per 1,000 (from 93 more to 251 more)	Low
Number of patients reporting pain at 1 year	47/78 (60.3%)	34/79 (37.7%)	RR 1.4 (1.02 to 1.9)	150 more per 1,000 (from 8 more to 339 more)	Moderate
Length of hospital stay	69	69	-	MD -1.7 (-4.45 to 1.05)	Moderate

10.3.2.3 Economic evidence

Three economic studies were identified^{163,173,291}. Rogmark et al (2003)²⁹¹ is a cost-consequence analysis based on a RCT which was excluded because it does not distinguish patients on the basis of whether they received hemiarthroplasty or total hip replacement.

1 Keating et al (2005)¹⁷³ compare Internal Fixation vs Total Hip Replacement in a cost-
 2 consequences analysis included in a Health Technology Assessment based on a RCT.
 3 Johansson et al (2006)¹⁶³ is a cost-consequence analysis based on a RCT. Please see
 4 Economic Evidence Tables 14 in Appendix F for further details.

6 **Table 10-38: Internal fixation vs total hip replacement - Economic study characteristics**

Study	Limitations	Applicability	Other Comments
Keating 2005 ¹⁷³	Minor limitations ^(a)	Partial applicability ^(b)	Costs not discounted because mainly incurred within 1 year of injury
Johansson 2006 ¹⁶³	Potentially serious limitations ^(c)	Partial applicability ^(d)	

7 (a) Small number of patients.

8 (b) Study set in the UK, but not a CUA.

9 (c) Costs were derived from just one hospital. No sensitivity analysis was conducted.

10 (d) Study set in Sweden.

12 **Table 10-39: Internal fixation vs total hip replacement - Economic summary of findings**

Study	Incremental cost per patient (£)	Incremental effects	ICER	Uncertainty
Keating 2005 ¹⁷³	£3224 ^(a)	THR has higher EQ-5D scores at 4, 12 and 24 months by 0.08; 0.12 and 0.14 respectively ^(b)	THR dominant	Two-way sensitivity analysis showed that the direction of change in cost did not change when cost of prostheses and cost of readmission were varied over a range from -50% to +100% around the baseline values.
Johansson 2006 ¹⁶³	£265	More patients with good/fair Harris hip score at 1 and 2 years in THR group ^(c)	THR dominant	NR

13 (a) The mean cost per patient included cost of hospital admission (inpatient and day case), theatre
 14 costs, prosthesis and profile of hardware. The mean cost per patient for internal fixation was
 15 £12,623 (95% CI: 10,768 – 14,478) and £9,399 (95% CI: 8,265-10,532) for THR.

16 (b) THR had better outcomes than internal fixation: lower number of deaths within 4, 12 and 24
 17 months from operation: (3% vs. 4%; 8% vs. 6% and 15% vs. 9%; p value not significant). Lower
 18 number of patients requiring further surgery within 4, 12 and 24 months from operation: 22% vs.
 19 7%; 31% vs. 9% and 39% vs. 9%; p value not reported). Higher mean EQ-5D scores at 4, 12 and 24
 20 months from operation: 0.56 vs 0.68 (p value not significant); 0.58 vs 0.70 (p = 0.04); 0.55 vs 0.69 (p
 21 value not significant).

22 (c) Percentage of patients with a Harris hip score excellent or good/fair or poor at 1 year: 12.5% vs.
 23 100% (p value: <0.0001); at 2 years: 14.29% vs.95.23% (p value: <0.001)

25 10.3.2.4 Evidence statement (s)

Clinical There is a statistically and clinically significant decrease in patients who
 require reoperations with total hip replacement than with internal fixation.
 The follow up varied between 1 and 13 years. (LOW QUALITY)

There is a statistically significant, but not clinically significant, increase in patients who reported pain at 1 year with internal fixation compared to total hip replacement (MODERATE QUALITY).

There is no statistically significant difference in mortality at 2 to 4 months, 12 to 18 months or 2 years (LOW QUALITY) and length of hospital stay (MODERATE QUALITY) between internal fixation and total hip replacement.

No RCT evidence was identified reporting functional status, quality of life, total time to resettlement in the community and place of residence after hip fracture.

Economic THR is the dominant strategy with respect to internal fixation (less costly and more effective). This evidence has minor limitations and partial applicability.

1

2

3 **10.3.3 Hemiarthroplasty versus total hip replacement**

4 **10.3.3.1 Review question**

5 In patients having treatment for intracapsular hip fracture what is the clinical and cost
6 effectiveness hemiarthroplasty versus total hip replacement on mortality, number of
7 reoperations, functional status, length of stay in hospital, total time to resettlement in the
8 community, quality of life, pain and place of residence after hip fracture.

9 One systematic review²⁶⁵ was identified. From this, 7 RCTs with 734 participants met the
10 inclusion criteria. See evidence table 7, Appendix E and forest plots G87 to G95 in Appendix
11 G.

12 **10.3.3.2 Clinical evidence**

13 **Table 10-40: Hemiarthroplasty vs total hip replacement – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality (follow up 3-6 months) ^{174,197,313}	3	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality (follow up 1 year) ^{26,218,313}	4	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Mortality (follow up 2-4 years) ^{11,174,197,218}	4	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Total no. of reoperations (follow-up 8 to 48 months) ^{11,26,73,174,218,313}	6	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
No. of patients reporting pain at 1 years ^{174,313}	2	RCT	no serious limitations	serious inconsistency ^(d)	no serious indirectness	serious imprecision ^(b)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Harris Hip Score for pain - 12 months ²⁶	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Failure to regain mobility (follow-up 1 to 4 years) ^{73,313}	2	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(b)
Oxford Hip Score - mean of 40 months ¹¹	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Barthel score - one year ²¹⁸	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Barthel score - four years ²¹⁸	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Hip rating questionnaire - 24 months ¹⁷⁴	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Harris Hip Score - total score - 12 months ^{26,218}	2	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Harris Hip Score - total score - four years ²¹⁸	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Harris Hip Score for function - 12 months ²⁶	1	RCT	serious limitations ^(a)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Short form 36 physical score - mean of 40 months ¹¹	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Self reported walking distance (kilometres) - mean of 40 months ¹¹	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
EuroQol (EQ-5d) questionnaire - 24 months ¹⁷⁴	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Length of hospital stay ¹⁷⁴	4	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(c)

1 (a) The studies with the most weight in the meta-analysis have inadequate or unclear allocation
2 concealment.

3 (b) The relatively few events and few patients give wide confidence intervals around the estimate of
4 effect. This makes it difficult to know the true effect size for this outcome.

5 (c) The wide confidence intervals around the measurement make the result imprecise. This makes it
6 difficult to know the true effect size for this outcome.

7 (d) There is significant heterogeneity between the studies which maybe due to the types of
8 arthroplasty used.
9

10 **Table 10-41: Hemiarthroplasty vs total hip replacement - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
---------	--------------	---------	---------------	-----------------	---------

Mortality (follow up 3-6 months)	25/192 (13%)	11/166 (6.6%)	RR 1.88 (0.96 to 3.68)	57 more per 1,000 (from 3 fewer to 174 more)	Low
Mortality (follow up 1 year)	42/272 (15.4%)	32/252 (10.3%)	RR 1.15 (0.76 to 1.74)	15 more per 1,000 (from 25 fewer to 76 more)	Low
Mortality (follow up 2-4 years)	38/176 (21.6%)	29/169 (19.1%)	RR 1.23 (0.8 to 1.87)	44 more per 1,000 (from 38 fewer to 166 more)	Low
Total no. or reoperations (follow-up 8 to 48 months)	42/350 (12%)	36/331 (10.6%)	RR 1.06 (0.7 to 1.6)	6 more per 1,000 (from 32 fewer to 64 more)	Low
No. of patients reporting pain (follow-up 1 years)	50/133 (37.6%)	29/123 (23.8%)	RR 1.68 (1.16 to 2.42)	161 more per 1,000 (from 38 more to 338 more)	Low
Harris Hip Score for pain - 12 months	55	56	N/A	MD -4 (-6.33 to -1.67)	Low
Failure to regain mobility (follow-up 1 to 4 years)	17/110 (15.5%)	20/101 (19.5%)	RR 0.78 (0.43 to 1.4)	43 fewer per 1,000 (from 111 fewer to 78 more)	Low
Oxford Hip Score - mean of 40 months	33	36	N/A	MD 3.50 (0.34 to 6.66)	Moderate
Barthel score - one year	30	33	N/A	MD -8 (-13.61 to -2.39)	Low
Barthel score - four years	20	23	N/A	MD -5.7 (-11.19 to -0.21)	Low
Hip rating questionnaire – 2 years	50	56	N/A	MD -6.1 (-12.38 to 0.18)	Moderate
Harris Hip Score - total score at 1 year	85	89	N/A	MD -5.47 (-8.39 to -2.55)	Low
Harris Hip Score - total score at 4 years	20	23	N/A	MD -4.2 (-7.66 to -0.74)	Low
Harris Hip Score for function - 12 months	55	56	N/A	MD -3.7 (-7.13 to -0.27)	Low
Short form 36 physical score - mean of 40 months	33	36	N/A	MD -2.43 (-7.56 to 2.7)	Moderate
Self reported walking distance (kilometres) - mean of 40 months	33	36	N/A	MD -1.7 (-3.28 to -0.12)	Moderate
EuroQol (EQ-5d) questionnaire at 2 years	65	66	N/A	MD -0.16 (-0.28 to -0.04)	Moderate
Length of hospital stay	69	69	N/A	MD -0.80 (-3.82 to 2.22)	Moderate

1

2 **10.3.3.3 Economic evidence**

Two studies were identified. Rogmark et al (2003)²⁹¹ is a cost-consequence analysis based on a RCT which was excluded because it does not distinguish patients on the basis of whether they received hemiarthroplasty or total hip replacement. A cost-consequence analysis comparing internal fixation vs. total hip replacement by Keating et al (2005)¹⁷³ was included. (Economic Evidence Table 14 in Appendix F)

Table 10-42: Hemiarthroplasty vs total hip replacement - Economic study characteristics

Study	Limitations	Applicability	Other Comments
Keating 2005 ¹⁷³	Minor limitations ^(a)	Partially applicability ^(b)	Costs not discounted because mainly incurred within 1 year of injury

(a) Small number of patients.

(b) UK study but did not a CUA.

Table 10-43: Hemiarthroplasty vs total hip replacement - Economic summary of findings

Study	Incremental cost per patient (£)	Incremental effects	ICER	Uncertainty
Keating 2005 ¹⁷³	£498 ^(b)	Hemiarthroplasty has lower EQ-5D scores at 4, 12 and 24 months ^(b)	NA	Two-way sensitivity analysis showed that the direction of change in cost did not change when cost of prostheses and cost of readmission were varied over a range from -50% to +100% around the baseline values.

(a) The mean cost per patient for hemiarthroplasty was 9,897 (95% CI: 8,062 – 11,732) and £9,399 (95% CI: 8,265-10,532) for THR.

(b) Hemiarthroplast had higher number of deaths within 4, 12 and 24 months from operation than THR: 5% vs.4%; 10% vs. 6% and 16% vs. 9%; (p values not significant), but lower reoperation rates at 4, 12 and 24 months: 5% vs. 7%; 5% vs 9%; and 5% vs. 9% (p value NR). THR had higher mean EQ-5D scores at 4, 12 and 24 months: 0.61 vs. 0.68 (not significant); 0.64 vs. 0.70 (not significant); 0.53 vs 0.69 (p=0.008).

10.3.3.4 Evidence statement (s)

Clinical There is a statistically significant, but not clinically significant, decrease in patients who reported pain and had a lower Harris Hip score for pain (indicating better function) at 1 year with total hip replacement compared to hemiarthroplasty (LOW QUALITY).

There is a statistically significant, but not clinically significant, increase in patients who have a lower Oxford Hip Score at 40 months (indicating better function) with total hip replacement compared to hemiarthroplasty (MODERATE QUALITY).

There is a statistically significant, but not clinically significant, increase in patients who have a higher Barthel Score (indicating better function) at 1 and 4 years (LOW QUALITY), a higher total Harris Hip Score at 1 and 4 years (LOW QUALITY), a higher Harris Hip Score for function at 1 year (LOW QUALITY) and a longer self reported walking distance at 40 months (MODERATE QUALITY)

with total hip replacement compared to hemiarthroplasty.

There is a statistically significant, but not clinically significant, increase in patients who have a higher Eq-5d (Euroqol) score at 2 years with total hip replacement compared to hemiarthroplasty (MODERATE QUALITY).

There is no statistically significant difference in mortality at 2 to 4 months (LOW QUALITY), 6 months (MODERATE QUALITY), 1 year (LOW QUALITY) or 2 to 4 years (LOW QUALITY), the number of reoperation at 8 to 48 months (LOW QUALITY), the number of patients who fail to regain mobility at 1 to 4 years (LOW QUALITY), the Hip Rating Questionnaire Score at 2 years (MODERATE QUALITY), the Short Form 36 (SF 36) score (MODERATE QUALITY) and length of hospital stay (MODERATE QUALITY) between hemiarthroplasty and total hip replacement.

No RCT evidence was identified reporting total time to resettlement or place of residence after hip fracture for studies comparing total hip replacement and hemiarthroplasty.

Economic THR is dominant compared to hemiarthroplasty. This evidence has minor limitations and partial applicability.

1

2 10.3.4 Recommendations and link to evidence

Recommendation	Perform replacement arthroplasty (hemiarthroplasty or total hip replacement) in patients with a displaced intracapsular fracture.
Relative values of different outcomes	The number of reoperations, functional status, pain and quality of life were considered the important outcomes with the number of reoperations being the most important. The interventions were not anticipated to have a significant impact on mortality so this was considered to be less important. Place of residence after hip fracture was also considered to be less important as it is a surrogate measurement for functional status.
Trade off between clinical benefits and harms	Compared to internal fixation there was a significantly lower reoperation rate with both hemiarthroplasty and total hip replacement, less patient reported pain with total hip replacement and better functional or quality of life scores with hemiarthroplasty. There was no significant difference for mortality, length of stay, failure to return to the same place of residence and failure to regain mobility. None of the reported outcomes showed any advantage of internal fixation over arthroplasty.
Economic considerations	Evidence partially applicable to the UK with only minor limitations was available on the cost-effectiveness of internal fixation vs. hemiarthroplasty and internal fixation vs. total hip replacement. The evidence shows that hemiarthroplasty is cost saving compared to internal fixation. In particular, hemiarthroplasty involved a significantly lower number of patients needing further surgery at 12 and 24 months compared to internal fixation. Similarly, THR required a lower rate of re-operation than internal

fixation, albeit not statistically significant.

Quality of evidence

The evidence was of low or moderate quality. Most outcomes were downgraded due to poor or uncertain allocation concealment. Several results were imprecise as the confidence intervals were near to one, making it difficult to determine the true effect size. Some studies were also heterogenous that could be due to the different types of arthroplasty.

Overall, the GDG felt that despite some of the results being of low quality and data not being available for some outcomes where there is a difference it shows arthroplasty being better than internal fixation. Consequently arthroplasty is recommended.

Other considerations

There maybe rare circumstances where reduction and internal fixation is appropriate for displaced intracapsular fragility fractures.

People with cognitive impairment were excluded from a lot of the studies. However, the GDG felt there is no reason for this group of patients should be excluded from equal treatment to others.

All patients should be allowed to be mobilised full weight bearing after hip fracture surgery (see section 10.2). All modern implants are designed to be load sharing devices to facilitate this.

The GDG consider this recommendation a key priority for implementation.

1

Recommendation

Offer total hip replacement to patients with a displaced intracapsular fracture who:

- **were able to walk independently out of doors with no more than the use of a stick and**
- **are not cognitively impaired and**
- **are medically fit for anaesthesia and the procedure.**

Relative values of different outcomes

The number of reoperations, functional status, pain and quality of life were considered the important outcomes with the number of reoperations being the most important. The interventions were not anticipated to have a significant impact on mortality so this was considered to be less important. Place of residence after hip fracture was also considered to be less important as it is a surrogate measurement for functional status.

Trade off between clinical benefits and harms

There was a significantly less patient reported pain and a better Oxford Hip Score, Barthel Score, Harris Hip Score, self reported walking distance and quality of life score (Eq-5d) with total hip replacement compared to hemiarthroplasty. There was no significant difference for mortality, length of stay, failure to return to the same place of residence and failure to regain mobility. None of the reported outcomes showed any advantage of

hemiarthroplasty over total hip replacement in the selected patient group.

Economic considerations

The cost-effectiveness evidence shows that THR replacement was cost-saving compared to both hemiarthroplasty and internal fixation.

Quality of evidence

The evidence was of low or moderate quality. Most outcomes were downgraded due to poor or uncertain allocation concealment. Several results were imprecise as the confidence intervals were near to one making it difficult to determine the true effect size. Some studies were also heterogenous that could be due to the different types of arthroplasty.

Overall, the GDG felt that despite some of the results being of low quality and data not being available for some outcomes where there is a difference it all shows total hip replacement being better than hemiarthroplasty in the selected patient group. Consequently total hip replacement is recommended for that group.

Other considerations

All but one of the studies excluded patients who were not medically fit, were not independently mobile before the fracture and were cognitively impaired. Consequently this recommendation does not include these groups. All the studies included in this review used a small head size for total hip replacement. Modern total hip replacements use a larger head which can reduce the risk of dislocation.

All patients should be allowed to be mobilised full weight bearing after hip fracture surgery (see section 10.2). All modern implants are designed to be load sharing devices to facilitate this.

The GDG consider this recommendation a key priority for implementation.

1

<i>Recommendation</i>	Use a proven femoral stem design rather than Austin Moore or Thompson stems for arthroplasties. Suitable designs include those with an Orthopaedic Data Evaluation Panel rating of 10A, 10B, 10C, 7A, 7B, 5A, 5B, 3A or 3B.
------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Relative values of different outcomes

The number of reoperations, functional status, pain and quality of life were considered the important outcomes. The interventions were not believed to have a significant impact on mortality so this was considered to be less important. Place of residence after hip fracture was also considered to be less important.

Trade off between clinical benefits and harms

Stem designs recommended here have a revision rate less than other stem designs. A higher failure rate would lead to a lower quality of life for patients.

Economic considerations

No economic evidence was found. Stems with a higher failure rate would require more reoperations and consequently, increased

costs and a lower quality of life for patients. Data supplied by an expert advisor reported the cost of an Exeter Trauma stem (ETS) monoblock as an example of a proven femoral stem design as £249 at 2008 prices.

Quality of evidence

No randomised evidence comparing modern stems with older stems was found.

Other considerations

There is a move towards modern style cemented stems. The Orthopaedic Data Evaluation Panel (ODEP) was set up in response to the NICE guidance on selection of prosthesis for primary total hip replacement²²⁶. The ratings used relate to the revision rate of stems and cups in arthroplasty. The results are available via the NHS Supply Chain website (<http://www.supplychain.nhs.uk/portal/page/portal/Communities/Orthopaedics/ODEP%20database>). A rating of 10A, 10B or 10C relates to devices with a failure rate of arthroplasty of 10% or less at 10 years. A rating of 7A and 7B relate to a failure rate of 7% or less at 7 years. A rating of 5A and 5B relate to a failure rate of 5% or less at 5 years. A rating of 3A and 3B relate to a failure rate of 3% or less at 3 years.

This recommendation was based on NICE guidance on selection of prosthesis for primary total hip replacement and expert opinion. In the light of such good evidence being available for the adequacy of femoral stem designs for patients with degenerative change it was thought that specific research in the fracture group would not be appropriate.

All patients should be allowed to be mobilised full weight bearing after hip fracture surgery (see section 10.2). All modern implants are designed to be load sharing devices to facilitate this.

Patients with hip fracture, particularly older patients have been treated by methods which have evolved very little over the last 50 years. This has led to a perception that they may be receiving second-class treatment. An example is the difference in the design of hip replacement implants used in patients with fractures compared with those used in patients with degenerative change. Many of those used in the fracture patients now appear archaic and their equivalents in the elective orthopaedic patients were superseded many years ago.

Long-term follow-up studies to identify function and durability of a replacement component in a fracture patient are difficult to carry out as so many of the patients are frail and their life expectancy is limited. However such studies are easier in patients with degenerative change and there is a well recognised system of assessing the adequacy of the design of a femoral stem for these patients.

1 **10.3.5 Research recommendations on displaced intracapsular fractures**

2 ***10.3.5.1 Large head total hip replacement versus hemiarthroplasty***

3 The GDG recommended the following research question:

- 4 ➤ What is the clinical and cost effectiveness of large-head total hip replacement
5 versus hemiarthroplasty on functional status, reoperations and quality of life in
6 patients with displaced intracapsular hip fracture?

7 **Why this is important**

8 Large-head total hip replacement is a development of traditional total hip replacement,
9 where a larger head makes the joint more stable and hence reduces the risks of dislocation.
10 Three small trials have shown traditional small-head total hip replacement to have better
11 outcomes and function, albeit with an increased dislocation rate in selected groups of
12 patients. The drawback with large-head arthroplasty is the additional implant cost and
13 theatre time. This cost can account for up to 20% of current NHS tariff (up to £2000) and
14 the study aims to address whether this translates to improved patient outcome. The study
15 design for the proposed research would be best addressed by a randomised controlled trial.
16 This would have two arms to compare current standard care (using hemiarthroplasty) with
17 using large-head total hip replacement for patients sustaining displaced intracapsular hip
18 fractures. The primary outcome would be patient mobility at 1 year and secondary
19 outcomes would include functional outcomes, quality of life and cost effectiveness of the
20 intervention.

21 It would be expected that a sample size of approximately 500 patients would be required to
22 show a significant difference in the mobility, hip function and quality of life (assuming 80%
23 power, $p < 0.05$). By recruiting through a trauma research network it is estimated that 10
24 centres would be able to recruit 20 patients per month (from 45 eligible patients) giving a
25 recruitment period of 25 months.

1

2 **10.4 Use of cement in arthroplasty**

3 The cement used in securing a hip replacement is not an adhesive but a grout, that is it is
 4 used to fill the gaps between the metal prosthesis and the bone. Thus, a component fixed
 5 with cement may be more secure resulting in less pain after surgery and decreased need
 6 for surgical revision due to loosening of the prosthesis. However, it has been suggested that
 7 cementing may induce side effects including cardiac arrhythmias and cardiorespiratory
 8 collapse, both of which may be fatal. NPSA data reports 26 deaths and six cases of severe
 9 harm when bone cement was used during hip surgery between October 2003 and October
 10 2008. Data from the MHRA reports 20 deaths and four cases of severe harm with bone
 11 cement between 2000 and 2008. The NPSA published advice on cementing techniques to
 12 reduce such risk. However, patients undergoing surgery for proximal femoral fractures are
 13 often elderly and frequently have multiple comorbidities, often severe. Therefore some
 14 intraoperative deaths may occur and be unrelated to the use of cement.

15 **10.4.1 Use of cement in original Thompson and Austin Moore designs of**
 16 **arthroplasty**

17 **10.4.1.1 Review question**

18 In patients having replacement arthroplasty for hip fracture what is the clinical and cost
 19 effectiveness of a cemented stem versus an uncemented stem on mortality, number of
 20 reoperations, wound healing complications, functional status, length of stay in hospital and
 21 total time to resettlement in the community, quality of life, pain and place of residence
 22 after hip fracture?

23 One systematic review²⁶⁵ including 6 RCTs with 899 participants was identified. See
 24 Evidence Table 7 and forest plots G52 to G66 in Appendix G.

25 **10.4.1.2 Clinical evidence**

26 **Table 10-44: Cemented vs. uncemented stem (original Thompson and Austin Moore designs of**
 27 **arthroplasty) – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Perioperative mortality ^{136,260}	2	RCT	no serious limitations ^(b)	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up <1 month) ^{81,260}	2	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up 3 months) ^{81,136,260,316}	4	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up 1 year) ^{34,81,136,260,298}	5	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision ^(g)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Failure to regain mobility (follow-up 12 to 17 months) ^{81,260,316}	4	RCT	no serious limitations	serious inconsistency (i,j)	no serious indirectness	serious imprecision (k)
Change in mobility score (follow-up 12 months; better indicated by less) ²⁶⁰	1	RCT	no serious limitations	no serious inconsistency	serious indirectness (a,l)	serious imprecision (l)
Length of hospital stay ^{81,136,260,298}	4	RCT	serious limitations (d,e)	no serious inconsistency	no serious indirectness	serious imprecision (b)
Failure to return home (follow up 1.5 to 5 years) ^{81,260}	2	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision (b)
Pain (follow up 3 months) ^{260,316}	2	RCT	no serious limitations	no serious inconsistency	serious indirectness (a,m)	serious imprecision (b)
Pain (follow up 1-2 years) ^{81,260,316}	3	RCT	no serious limitations	no serious inconsistency	serious indirectness (b)	no serious imprecision
Pain score (follow up 6 months) ²⁶⁰	1	RCT	no serious limitations	no serious inconsistency	serious indirectness (a,m)	serious imprecision (b)
Reoperations (follow-up 8 to 20 months) ^{34,260}	2	RCT	no serious limitations	no serious inconsistency	serious indirectness (a)	serious imprecision (b)
Deep Sepsis (follow-up 1 to 5 years) ^{136,260,298,316}	4	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision (b)
Wound haematoma (follow-up 1 to 5 years) ²⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious imprecision (b)

- 1 (a) Data only available for unipolar hemiarthroplasty
- 2 (b) The relatively few events and few patients give wide confidence intervals around the estimate of
- 3 effect. This makes it difficult to know the true effect size for this outcome.
- 4 (c) The result is calculated using only one of the two studies with no allocation concealment or
- 5 blinding of the intervention. However, the effect size is similar between the two studies, the second
- 6 study is larger and does not having any serious limitations in design. There fore the evidence has
- 7 not been downgraded on the basis of study quality.
- 8 (d) Unclear or no allocation concealment in 2 of the 4 studies which account for over 75% of the weight
- 9 of the result.
- 10 (e) Randomisation method by odd or even hospital number in 1 of the 4 studies, and by alternate days
- 11 in another of the 4 studies. These 2 of the 4 studies account for over 75% of the weight of the result.
- 12 (f) The estimate of effect is derived from the data relating to unipolar hemiarthroplasty. There is a
- 13 small study relating to bipolar arthroplasty, this has little impact on the overall result.
- 14 (g) The confidence intervals around the estimate of effect are wide enough to suggest some
- 15 uncertainty in the estimate of the effect. A larger number of patients may show a statistically
- 16 significant difference in the outcome.
- 17 (h) The estimate of effect is calculated with the better quality studies having more weight than the
- 18 lower quality studies. Consequently, the result has not been downgraded for quality.

- 1 (i) There is significant statistical heterogeneity in the results: there is no statistical for unipolar
 2 hemiarthroplasty; Significantly more patients failed to regain mobility with uncemented bipolar
 3 hemiarthroplasty than cemented bipolar hemiarthroplasty.
- 4 (j) The definition for failure to regain mobility is different in the studies. The two studies, one showing
 5 no statistical difference the other favouring cement, measure the number of people with a change
 6 in their walking status. The third study showing no statistical difference measures the number of
 7 people unable to walk properly (this includes walking without a limp) .
- 8 (k) The confidence intervals around the estimate of effect are wide enough to suggest some
 9 uncertainty in the estimate of the effect.
- 10 (l) Definition of mobility score not given. Unable to determine if it is a valid measurement for mobility
 11 or if the estimate of effect is clinically significant.
- 12 (m) How pain was measured is not reported for the study with the most weight in the meta-analysis.
 13 Unable to determine if it is a valid measurement or if the estimate of effect is clinically significant.

14
 15 **Table 10-45: Cemented vs uncemented stem (original Thompson and Austin Moore designs of**
 16 **arthroplasty) - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Perioperative mortality	1/277 (0.4%)	0/266 (0%)	RR 2.58 (0.11 to 62.21)	0 fewer per 1,000 (from 0 fewer to 0 more)	Low
Mortality (follow up <1 month)	11/227 (4.8%)	13/226 (6.6%)	RR 0.84 (0.38 to 1.84)	10 fewer per 1,000 (from 28 fewer to 54 more)	Low
Mortality (follow up 3 months)	49/359 (13.6%)	49/349 (13%)	RR 0.98 (0.68 to 1.41)	3 fewer per 1000 (from 45 fewer to 57 more)	Low
Mortality (follow up 1 year)	101/395 (25.6%)	113/398 (26.4%)	RR 0.9 (0.71 to 1.13)	28 fewer per 1000 (from 82 fewer to 37 more)	Moderate
Failure to regain mobility (follow-up 12 to 17 months)	117/196 (59.7%)	124/182 (68.1%)	RR 0.84 (0.64 to 1.11)	109 fewer per 1000 (from 245 fewer to 75 more)	Low
Change in mobility score (follow-up 12 months; better indicated by less)	150	144	N/A	MD -0.8 (-1.23 to -0.37)	Low
Length of hospital stay	354	342	N/A	MD -1.42 (-3.15 to 0.32)	Low
Failure to return home (follow up 1.5 to 5 years)	16/219 (7.3%)	26/220 (11.8%)	RR 0.62 (0.34 to 1.12)	45 fewer per 1000 (from 78 fewer to 14 more)	Moderate
Pain (follow up 3 months)	67/192 (34.9%)	84/183 (45.9%)	RR 0.77 (0.6 to 0.98)	106 fewer per 1000 (from 9 fewer to 184 fewer)	Low
Pain (follow up 1-2 years)	44/193 (22.8%)	73/176 (41.5%)	RR 0.55 (0.4 to 0.75)	187 fewer per 1000 (from 104 fewer to 249 fewer)	Moderate
Pain score (follow up 6 months)	147	142	-	MD -0.6 (-0.9 to -0.3)	Low

Reoperations (follow-up 8 to 60 months)	10/238 (4.2%)	19/253 (7.5%)	RR 0.55 (0.27 to 1.14)	34 fewer per 1000 (from 55 fewer to 11 more)	Low
Deep sepsis (follow up 1 to 5 years)	8/385 (2.1%)	6/376 (1.6%)	RR 1.25 (0.48 to 3.24)	4 more per 1000 (from 8 fewer to 36 more)	Moderate
Wound Haematoma (follow up 2 to 5 years)	2/200 (1%)	1/200 (0.5%)	RR 2.01 (0.18 to 22.35)	5 more per 1000 (from 4 fewer to 107 more)	Moderate

1

2 **10.4.1.3 Economic evidence**

3 Two economic studies were identified. Santini (2005)²⁹⁸ is a cost-consequence analysis
 4 based on a RCT included in our clinical review (see 10.3.3.2). See evidence table 15 in
 5 Appendix F for additional details. Marinelli (2008)²⁰⁶ was excluded because of poor
 6 methodology.

7 **Table 10-46: Cemented vs. uncemented hemiarthroplasty - Economic study characteristics**

Study	Limitations	Applicability	Other Comments
Santini 2005 ²⁹⁸	Potentially serious limitations (a)	Partially applicable (b)	Based on RCT included in our clinical review (see 10.3.3.2).

8 (a) Surgical time not included in cost calculation although it was significantly different (patients in the
 9 uncemented hemiarthroplasty group had shorter operating time). The only difference considered was
 10 the cost of prostheses.

11 (b) Not a cost-utility analysis. Study conducted in Italy.

12

13 **Table 10-47: Cemented vs. uncemented hemiarthroplasty - Economic summary of findings**

Study	Incremental cost per patient (£)	Incremental effects	ICER	Uncertainty
Santini 2005 ²⁹⁸	Cost saving (-£710) (b)	(b)	N/R	N/R

14 (a) Cost of medical and nursing staff, drugs, diagnostic procedures, prostheses, blood transfusion and
 15 hospital sta. Converted into GBP from 2001 euro using the Purchasing Power Parities.

16 (b) Different outcomes were reported but none of them were significantly different.

17 **10.4.1.4 Evidence statement (s)**

Clinical There is a statistically significant, but not clinically significant, increase in patients who have a lower reduction in mobility score (less loss of mobility) at 12 months (LOW QUALITY).

There is a statistically significant, but not clinically significant, decrease in patients who reported pain at 3 months (LOW QUALITY) and 1 to 2 years (MODERATE QUALITY). However, there was no significant difference in a pain score at 6 months (LOW QUALITY).

There is no statistically significant difference in perioperative mortality (LOW QUALITY), mortality at 3 months (LOW QUALITY) or 1 year (MODERATE QUALITY), failure to return home (MODERATE QUALITY), length of hospital stay (LOW QUALITY), number of patients requiring reoperations (LOW

QUALITY), number of patients failing to regain mobility (LOW QUALITY), deep sepsis (MODERATE QUALITY), wound haematoma (MODERATE QUALITY) and all medical complications combined (VERY LOW QUALITY).

No RCT evidence was identified reporting quality of life, total length of stay to community resettlement or place of residence after hip fracture

No RCT evidence was identified to suggest there is a safety issue with using cement.

Economic Cemented hemiarthroplasty is cost saving compared to uncemented hemiarthroplasty. This evidence has potentially serious limitations and partial applicability.

1

2 10.4.2 Use of cement in newer designs of arthroplasty

3 10.4.2.1 Review question

4 In patients having replacement arthroplasty for hip fracture what is the clinical and cost
5 effectiveness of a cemented stem versus an uncemented stem on mortality, number of
6 reoperations, wound healing complications, functional status, length of stay in hospital and
7 total time to resettlement in the community, quality of life, pain and place of residence
8 after hip fracture.

9 One RCT⁹⁴ including 220 participants was identified. See Evidence Table 7 and forest plots
10 G67 to G73 in Appendix G.

11 10.4.2.2 Clinical evidence

12 **Table 10-48: Cemented vs. uncemented stem (newer designs of arthroplasty) – Clinical study**
13 **characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality (follow up 30 days)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up 90 days)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up 1 year)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(b)	serious imprecision ^(b)
Mortality (follow up 2 years)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Total number of reoperations (follow up 12 months)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Need for pain medication (follow up 12 months)⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Unable to walk without aids (follow up 12 months) ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Barthel score of less than 19 (follow up 12 months) ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Harris Hip Score (follow up 12 months) ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Eq-5d index score (follow up 12 months) ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Eq-5d visual analogue score (follow up 12 months) ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(b)
Length of hospital stay ⁹⁴	1	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(a)	serious imprecision ^(c)

1
2
3
4
5
6
7

(a) Data only available for bipolar hemiarthroplasty

(b) The relatively few events and few patients give wide confidence intervals around the estimate of effect. This makes it difficult to know the true effect size for this outcome.

(c) The effect size is uncertain as the confidence intervals suggest the length of stay could be over 2 days shorter or over 1 day longer with cemented hemiarthroplasty.

Table 10-49: Cemented vs uncemented stem - Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality (follow up 30 days)	8/142 (5.6%)	10/153 (6.5%)	RR 0.47 (0.15 to 1.57)	35 fewer per 1000 (from 56 fewer to 37 more)	Low quality
Mortality (follow up 90 days)	13/108 (12%)	15/105 (14.3%)	RR 0.84 (0.42 to 1.68)	23 fewer per 1000 (from 83 fewer to 97 more)	Low quality
Mortality (follow up 1 year)	34/142 (23.9%)	46/153 (30.1%)	RR 0.65 (0.39 to 1.07)	105 fewer per 1000 (from 183 fewer to 21 more)	Low quality
Mortality (follow up 2 years)	32/108 (29.6%)	36/105 (34.3%)	RR 0.86 (0.58 to 1.28)	48 fewer per 1000 (from 144 fewer to 96 more)	Low quality
Total number of reoperations (follow up 12 months)	7/112 (6.3%)	8/108 (7.4%)	RR 0.84 (0.32 to 2.25)	12 fewer per 1000 (from 50 fewer to 93 more)	Low quality
Need for pain medication (follow up 12 months)	23/91 (25.3%)	14/77 (18.2%)	RR 1.39 (0.77 to 2.51)	71 more per 1000 (from 42 fewer to 275 more)	Low quality
Unable to walk without aids (follow up 12 months)	4/91 (4.4%)	6/77 (7.8%)	RR 0.56 (0.17 to 1.93)	34 fewer per 1000 (from 65 fewer to 72 more)	Low quality
Barthel score of less than 19 (follow up 12 months)	46/91 (50.5%)	29/77 (37.7%)	RR 1.34 (0.94 to 1.91)	128 more per 1000 (from 23 fewer to 343 more)	Low quality

Harris Hip Score (follow up 12 months)	90	77	N/A	MD 0.9 lower (6 lower to 4.2 higher)	Low quality
Eq-5d index score (follow up 12 months)	56	57	N/A	MD 0.07 higher (0.03 lower to 0.17 higher)	Low quality
Eq-5d visual analogue score (follow up 12 months)	61	60	N/A	MD 4 lower (10.75 lower to 2.75 higher)	Low quality
Length of hospital stay	109	106	N/A	MD 0.6 lower (2.48 lower to 1.28 higher)	

1

2 10.4.2.3 Economic evidence

3 No cost-effectiveness evidence was identified. A cost analysis was conducted based on
4 the resources used in the Figved study⁹⁴ and on GDG expert opinion. Please see section
5 20.8 of Appendix H of this guideline for further details.

6 **Table 10-50: Cemented stems versus uncemented stems (newer designs of arthroplasty) –**
7 **Economic study characteristics**

Study	Limitations	Applicability	Other Comments
NCGC cost analysis	Minor limitations ^(a)	Partially applicable ^(b)	Cost analysis based on resources reported in Figved (2009) ⁹⁴ and on GDG's expert opinion

8 (a) No sensitivity analysis.

9 (b) Cost analysis based on one study alone by Figved⁹⁴ and on GDG's expert opinion. The study by Figved⁹⁴ is
10 not UK based and therefore may not completely reflect current NHS practice.

11

12

13 **Table 10-51: Cemented stems versus uncemented stems (newer designs of arthroplasty) –**
14 **Economic summary of findings**

Study	Incremental cost (£)	Incremental effects	ICER	Uncertainty
NCGC cost analysis	£171.79 ^(a) (cost saving)	N/A	N/A	N/R

15 (a) The following cost categories were considered in the cost analysis: cost of implants;
16 length of hospital stay; cost of cement accessories; theatre time costs; re-operation
17 costs. The costs of length of stay and re-operation were considered in the analysis even
18 if in the RCT by Figved⁹⁴ there was not statistically significant difference between the
19 two groups for these outcomes. The total cost for the new design cemented stems was
20 estimated to correspond to £2751.64 and that for the new design uncemented stems
21 to £2923.43. The estimate for the total cost for the cemented stems could increase up
22 to £2859.75 when a more thorough set of accessories are assumed to be used in the
23 operation, in which case the incremental savings associated with using cemented
24 stems would amount to £63.68. See Appendix H section 20.8 for further details.

25 10.4.2.4 Evidence statement (s)

Clinical There is no statistically significant difference in mortality at 30 days, 90 days,
1 year or 2 years (LOW QUALITY).

There is no statistically significant difference at 1 year in the number of patients requiring reoperations, number of patients pain requiring medication, number of patients unable to walk without aids, Barthel Score of less than 19, Harris Hip Score, Eq-5d index score and visual analogue score, deep wound sepsis, any wound infection, length of hospital stay (LOW QUALITY).

No RCT evidence was identified reporting total time to resettlement in the community and place of residence after hip fracture

No RCT evidence was identified to suggest there is a safety issue with using cement.

Economic No studies were identified on the cost-effectiveness of cemented vs. uncemented stem (newer designs of arthroplasty). An NCGC cost analysis found that cemented stems are £171.79 cheaper than the newer design uncemented stems. This evidence has minor limitation and partial applicability.

1
2

3 **10.4.3 Recommendations and link to evidence**

<i>Recommendation</i>	Use cemented implants in patients undergoing surgery with arthroplasty
Relative values of different outcomes	The outcomes considered were mortality, functional status, quality of life, pain, requirement for reoperation, non-healing and requirement for surgical revision, total length of stay (i.e. the time in hospital plus any time spent in rehabilitation). Mortality was of particular importance because of reported deaths by the NPSA.
Trade off between clinical benefits and harms	<p>There is no significant difference in mortality. There is evidence of less pain at 3 months and 1 to 2 years and better mobility score at 12 months with the older designs of cemented hemiarthroplasties. There was no significant difference for length of stay, failure to return to the same place of residence and failure to regain mobility. None of the reported outcomes showed any advantage of uncemented arthroplasty over cemented.</p> <p>There is no direct evidence comparing the use of cemented and uncemented total hip replacement.</p> <p>No RCT evidence was found to raise concerns about the safety of the use of cement.</p>
Economic considerations	<p>One study with potentially serious limitations and partial applicability found that the older cemented hemiarthroplasty are cost saving compared to uncemented hemiarthroplasty.</p> <p>The NCGC cost analysis on cemented stems vs. uncemented stems for newer designs of arthroplasty has considered several cost components, such as the cost of the implants, length of stay in hospital, rate of re-operations, accessories costs for the cemented</p>

implants.

However, the GDG did not consider the higher level of blood loss reported in Figved et al (2009)⁹⁴ for patients receiving cemented implants (89mL) to be significant in terms of both patients' outcomes and costs.

As the clinical evidence did not show any advantage of uncemented over cemented arthroplasty in the newer design, and as the cost of new designs of cemented implants was shown to be lower than that of uncemented implants, the GDG agreed to consider cemented implants cost-effective for hip fracture patients.

Quality of evidence

The evidence was of low or moderate quality. All but one of the studies comparing older arthroplasty designs used a Thompson or Austin Moore hemiarthroplasty (these are the first generation of implants to be used). The other study used an unspecified bipolar hemiarthroplasty. The evidence for modern stem designs is low quality mainly due to the lack of certainty around the effect size and only evidence being identified in bipolar hemiarthroplasty.

Overall, the GDG felt there was sufficient evidence to recommend the use of cemented arthroplasties over uncemented.

Other considerations

All studies comparing the effectiveness of internal fixation with THR and hemiarthroplasty with THR used cemented THR (see section 10.3.2)

All patients should be allowed to be mobilised full weight bearing after hip fracture surgery (see section 10.2). All modern implants are designed to be load sharing devices to facilitate this.

1

2 10.5 Surgical approach to hemiarthroplasty

3 Hemiarthroplasties are usually inserted using one of two approaches, either an
4 anterolateral or a posterior approach. The choice of surgical approach for a surgeon is
5 often dictated by local custom and practice and personal experience. This review looks at
6 the evidence to see if one is better than the other. RCTs and cohorts adjusted for
7 confounders were included.

8 10.5.1.1 Review question

9 In patients having surgical treatment for intracapsular hip fracture with hemiarthroplasty
10 what is the clinical and cost effectiveness of anterolateral compared to posterior surgical
11 approach on mortality, number of reoperations, dislocation, functional status, length of
12 hospital stay, quality of life and pain.

13 One systematic review²⁶⁹ including 1 RCT with 114 participants and one cohort study
14 involving 720 participants were identified. See Evidence Table 9, Appendix E.

15 10.5.1.2 Clinical evidence

1 **Table 10-52: Posterior vs. anterolateral approach to hemiarthroplasty – Clinical study**
 2 **characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality ³⁰⁹	1	RCT	serious limitations ^(a, b)	Unable to assess this ^(f)	serious indirectness ^(c)	serious imprecision ^(d)
Number of patients with impairment of mobility at 6 months compared to prefracture ³⁰⁹	1	RCT	serious limitations ^(a, b)	no serious inconsistency	serious indirectness ^(c)	serious imprecision ^(d)
Dislocation at 0 to 2 years ³⁰⁹	1	RCT	Very serious limitations ^(a, b)	no serious inconsistency	serious indirectness ^(c)	serious imprecision ^(d)
Dislocation at 0 to 10 years ⁸⁵	1	Cohort	serious limitations ^(e)	no serious inconsistency	no serious indirectness	no serious imprecision
Pain at 1 month ³⁰⁹	1	RCT	serious limitations ^(a)	no serious inconsistency	serious indirectness ^(c)	serious imprecision ^(d)

- 3 (a) Unclear allocation concealment and randomisation method
 4 (b) Patients allocated to the posterior approach were nursed flat in bed for two weeks after surgery as
 5 a precaution against dislocation.
 6 (c) Most operations performed by surgical trainees
 7 (d) The wide confidence intervals make the estimate of effect imprecise.
 8 (e) Only a limited number of confounders were included in the analysis. No adjustment or mention of
 9 the anaesthetists experience or grade.
 10 (f) Actual event rates were not provided for this, mortality was given as percentages in a graph. The
 11 percentages were estimated using this. Mortality was significantly higher at three months, six
 12 months, 12 months and two years in the posterior group $p < 0.05$. The rate was around double for
 13 all these time points.
 14

15 **Table 10-53: Posterior vs. anterolateral approach to hemiarthroplasty - Clinical summary of**
 16 **findings**

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality at 6 months, 12 months & 2 years	Not reported	Not reported	Significantly higher in posterior group ($p < 0.05$)	Not estimable	Very low
Number of patients with impairment of mobility at 6 months compared to prefracture	5/34 (14.7%)	15/41 (36.6%)	RR 0.40 (0.16 to 0.99)	220 fewer per 1000 (from 4 fewer to 307 fewer)	Very Low
Dislocation at 0 to 2 years	1/57 (1.8%)	1/57 (1.8%)	RR 1.00 (0.06 to 15.60)	0 fewer per 1000 (from 16 fewer to 256 more)	Very Low
Dislocation at 0 to 10 years (posterior approach with posterior repair)	15/176 (8.5%)	13/431 (3%)	multivariate odds ratio 3.9 (1.6 to 9.8)	87 more per 1000 (from 18 more to 265 more)	Very Low
Dislocation at 0 to 10 years (posterior approach without posterior repair)	17/129 (13.2%)	13/431 (3%)	multivariate odds ratio 6.9 (2.6 to 19)	178 more per 1000 (from 48 more to 543 more)	Very Low

Pain at 1 month	2/55 (3.6%)	6/55 (10.9%)	RR 3.0 (0.63, 14.22)	218 more per 1000 (from 40 fewer to 1442 more)	Very low
------------------------	-------------	--------------	----------------------	------------------------------------------------	----------

1

2 **10.5.1.3 Economic evidence**

3 No cost-effectiveness evidence was identified.

4 **10.5.1.4 Evidence statement (s)**

5

Clinical Two studies of different designs showed different effects for dislocation rates. One old RCT showed no statistically significant difference in dislocation rate between approaches. (VERY LOW QUALITY). One recent cohort which adjusted for confounders showed a statistically and clinically significant higher dislocation rate with the posterior approach compared to the anterolateral approach. (VERY LOW QUALITY)

Significantly fewer patients had impaired mobility at 6 months with a posterior approach to hemiarthroplasty compared to an anterior approach when the procedure was performed by surgical trainees. (VERY LOW QUALITY)

One study reported a significantly higher mortality with a posterior approach at 6 months, 12 months and two years but did not provide the event rates. (VERY LOW QUALITY]

Economic No evidence was identified regarding the cost-effectiveness of posterior vs. anterolateral approach to hemiarthroplasty.

6

7 **10.5.2 Recommendations and link to evidence**

Recommendation

Consider an anterolateral approach in favour of a posterior approach when inserting a hemiarthroplasty.

Relative values of different outcomes

Functional status, reoperation rate, and quality of life were considered the main outcomes. Pain, wound infection, dislocations, length of stay in secondary care and mortality were also considered.

Trade off between clinical benefits and harms

The cohort study showed a significantly higher dislocation rate with a large effect size with the posterior approach compared to the anterolateral approach. This reduces the potential complications of re-operation or revision surgery. An old RCT data showed a significantly lower impaired mobility at 6 months with a posterior approach, a doubling of mortality and no difference in dislocations compared to an anterolateral approach. However, the operations had been carried out by trainees with varying degrees

	<p>of experience. Also, the group operated on with an anterolateral approach were allowed to mobilise straight away and the group operated on with a posterior approach had two weeks postoperatively bed rest.</p> <p>None of the other outcomes were reported.</p>
Economic considerations	An anterolateral approach is likely to result in cost savings because of their lower dislocation rates, and hence less revision surgery.
Quality of evidence	Both the studies available are of very low quality. The RCT is an old study where the operations were mostly carried out by surgical trainees. This RCT also treated patients differently, with those receiving a posterior approach being nursed flat in bed for two weeks after surgery as a precaution against dislocation and had a much higher mortality in the posterior group. The cohort study, which adjusted for important factors in their results, is a recent study and shows a large effect size in favour of an anterolateral approach.
Other considerations	The GDG considered this evidence along with the GDG opinions and decided the recent evidence is more relevant. They therefore recommend the anterolateral approach over the posterior. It is also recognized that the posterior approach may well be as safe in preventing dislocation in those surgeons with a large experience of using it. However, the GDG believe the majority of surgeons who perform the surgery do not regularly perform posterior approaches. It is also noted that all the RCTs comparing hemiarthroplasty and total hip replacement utilized the anterolateral approach in all of the studies.

1

2 10.6 Extracapsular fracture fixation

3 In the extracapsular fractures the femoral head blood supply is unaffected and the proximal
4 fragment large enough to allow secure fixation, therefore internal fixation is the norm. The
5 surgical decision in this group is which of the various available methods of fracture fixation
6 is most effective for each pattern. When treating the extracapsular fractures around the
7 trochanter it is necessary to stabilise the intact femoral head and neck onto the shaft of the
8 femur. The head portion is stabilised by one or more screws up the neck and into the head.
9 This screw is attached to either a plate on the outside of the bone (called extramedullary
10 fixation) or a metal rod which is inserted down the middle of the femoral shaft
11 (intramedullary fixation). The rod can either be short, spanning approximately a third of the
12 length of the femur, or long spanning the whole length of the femur. The generic term for
13 the plate and screw used for the extramedullary fixation is a sliding hip screw and the term
14 for the intramedullary fixation is the intramedullary nail.

15

16 Extracapsular fractures are split into pertrochanteric (also called intertrochanteric), reverse
17 oblique and subtrochanteric (see Introduction, Figure 1).

18

10.6.1 Intramedullary versus extramedullary implants for fixation of trochanteric extracapsular fractures

There are numerous studies comparing intramedullary and extramedullary results. The intramedullary nails can vary in size and shape, with most evolving from the initial nail design which was changed due to an increase in per- and postoperative fractures of the femur. When reviewing the evidence, the trochanteric fractures were divided into stable fractures, (those with an intact lesser trochanter – AO/OTA A1), unstable fractures (those with a fracture between the trochanters with displacement of the lesser trochanter – AO/OTA A2 fractures) and reverse oblique fractures (AO/OTA A3). Historically and presently there have been numerous implants used to treat these and we have divided them into intramedullary (those which have a rod down the shaft of the bone) and extramedullary where the device sits on the outside of the bone. Commonly these are called intramedullary nails and sliding hip screws respectively. The intramedullary nails can come in various designs from different manufacturers. Their size and shape have evolved over the last twenty years. The design of the sliding hip screw has not changed over the last thirty years and sliding hip screws are generally very similar between the different manufacturers.

10.6.1.1 Review question

In patients undergoing repair for trochanteric extracapsular hip fractures what is the clinical and cost effectiveness of extramedullary sliding hip screws compared to intramedullary nails on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?

21 studies met the inclusion criteria for this review question with a total of 4,336 patients. See evidence table 5.8, Appendix E and forest plots G96 to G106 in Appendix G.

10.6.1.2 Clinical evidence

Table 10-54: Intramedullary vs. extramedullary implants for trochanteric extracapsular fracture – Clinical study characteristics

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality – 30 days ^{14,37,128,137,191,195,244,279,337}	9	RCT	no serious limitations ^(a,b)	no serious inconsistency	no serious indirectness	no serious imprecision
Mortality – 3 months ^{128,134,251}	3	RCT	serious limitations ^(d,e)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Mortality – 1 year ^{3,14,37,77,134,191,195,251,294,300,337}	11	RCT	no serious limitations ^(f)	no serious inconsistency	no serious indirectness	no serious imprecision
Reoperation – within follow up period of study ^{9,14,77,128,134,147,191,195,214,244,251,254,279,294,300,337}	16	RCT	no serious limitations	no serious inconsistency	no serious indirectness ^(h)	no serious imprecision

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Operative or postoperative fracture - within follow up period of study 3,9,37,77,128,134,137,147,191,214,244,251,258,279,300,337,364	17	RCT	no serious limitations ^(h, j)	no serious inconsistency	serious indirectness ^(k)	no serious imprecision
Cut-out (at latest follow up) 9,14,37,77,128,134,137,147,191,195,214,244,251,254,258,279,294,300,337,364	20	RCT	no serious limitations ^(l)	no serious inconsistency	no serious indirectness	serious imprecision ^(c)
Infection (deep infection or requires reoperation) ^{128,134,147,191,195,214,244,251,254,258,279,294,300,337}	14	RCT	no serious limitations ^(m)	no serious inconsistency	no serious indirectness ⁽ⁿ⁾	serious imprecision ^(c)
Non-union (at latest follow up) ^{77,137,191,251,258,279,294,300,364}	9	RCT	no serious limitations ^(o)	no serious inconsistency	no serious indirectness ^(p)	serious imprecision ^(c)
Pain (at latest follow up) ^{134,147,191,337}	4	RCT	no serious limitations	no serious inconsistency	serious indirectness ^(q)	no serious imprecision
Length of stay in hospital ^{137,147,191,244,251,254,294,300}	8	RCT	no serious limitations	serious ^(g)	no serious indirectness	serious imprecision ^(c)
Mean mobility (Parker – Palmer score. At 1 year) ^{134,294,300,337}	4	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision

- 1 (a) Unclear allocation concealment in 4 out of 9 studies.
- 2 (b) Loss to follow up not reported or more than 5% in 4 out of 9 studies
- 3 (c) The relatively few events and few patients give wide confidence intervals around the estimate of
- 4 effect. This makes it difficult to know the true effect size for this outcome.
- 5 (d) Unclear allocation concealment in 2 out of 3 studies.
- 6 (e) Loss to follow up not reported or more than 5%, in 2 out of 3 studies.
- 7 (f) Unclear allocation concealment in 3 out of 11 studies.
- 8 (g) There is significant statistical heterogeneity in the results
- 9 (h) The definition of reoperation varies between studies to include minor or major revisions.
- 10 (i) Unclear allocation concealment in 7 out of 15 studies.
- 11 (j) Loss to follow up not reported more than 5% in 8 out of 16 studies.
- 12 (k) All fractures of the femur that were reported have been combined.
- 13 (l) Loss to follow up not reported or more than 5% in 8 out of 19 studies.
- 14 (m) Loss to follow up not reported or more than 5% in 5 out of 15 studies
- 15 (n) Inclusion of reported infection varied between studies and included deep infection and infection
- 16 that required reoperation.
- 17 (o) Loss to follow up not reported or more than 5% in 4 out of 10 cases.
- 18 (p) All cases of non-union were combined using data at latest follow up.
- 19 (q) Different definitions of patient reported pain combined.
- 20
- 21

1 **Table 10-55: Intramedullary vs. extramedullary implants for trochanteric extracapsular fracture**
 2 **- Clinical summary of findings**

Outcome	Intramedullary	Extramedullary	Relative risk	Absolute effect	Quality
Mortality – 30 days	78/712 (11%)	56/729 (7.7%)	RR 1.44 (1.04 to 1.99)	34 more per 1000 (from 3 more to 76 more)	High
Mortality – 3 months	19/173 (11%)	21/173 (10%)	RR 0.9 (0.52 to 1.59)	12 fewer per 1000 (from 58 fewer to 72 more)	Low
Mortality – 1 year	186/1005 (18.5%)	175/1021 (17.1%)	RR 1.09 (0.91 to 1.31)	15 more per 1000 (from 15 fewer to 53 more)	High
Reoperation – within follow up period of study	69/1261 (5.5%)	50/1312 (3.8%)	RR 1.39 (0.87 to 2.23)	15 more per 1000 (from 5 fewer to 47 more)	High
Operative or postoperative fracture - within follow up period of study	54/1334 (4%)	5/1380 (0%)	RR 5.61 (2.98 to 10.59)	16 more per 1000 (from 7 more to 33 more)	Low
Cut-out (at latest follow up)	39/1446 (2.7%)	42/1508 (2.8%)	RR 0.95 (0.63 to 1.45)	1 fewer per 1000 (from 10 fewer to 13 more)	Moderate
Infection (deep infection or requires reoperation)	8/922 (0.9%)	10/943 (1%)	RR 0.86 (0.38 to 1.93)	1 fewer per 1000 (from 7 fewer to 10 more)	Moderate
Non-union (at latest follow up)	3/610 (0.5%)	3/621 (0.5%)	RR 1.01 (0.3 to 3.46)	0 more per 1000 (from 3 fewer to 12 more)	Moderate
Pain (at latest follow up)	90/278 (32.4%)	90/285 (25.9%)	RR 1.03 (0.81 to 1.30)	9 more per 1000 (from 60 fewer to 95 more)	Low
Length of stay in hospital	474	482	N/A	MD 0.54 lower (1.93 lower to 0.84 higher)	Moderate
Mean mobility (Parker – Palmer score. At 1 year)	274	281	N/A	MD 0.17 higher (0.17 lower to 0.51 higher)	High

3

4 **10.6.1.3 Economic evidence**

5 Three economic studies were indentified^{110,114,179}. All these studies have been excluded.¹¹⁴
 6 is a cost-consequence analysis based on a retrospective cohort study set in the US
 7 comparing trochanteric fixation nail with sliding hip screw. This study was excluded due to
 8 poor methodological design and to the limited applicability to the UK NHS.¹⁷⁹ compared
 9 proximal femoral nail with long-stem cementless calcar-replacement prosthesis which was
 10 not an included intervention. Another study¹¹⁰ was excluded as no cost figures were
 11 reported.

1 The GDG was informed of the prices of implants produced by all major orthopaedic
 2 suppliers in the UK. At 2010 prices, the average cost for a sliding hip screw was estimated at
 3 £252.51, of a short intramedullary nail at £760.08, and of a long intramedullary nail at
 4 £1,175.40. Please see section 8.3 in Appendix H for further details.

5

6 **10.6.1.4 Evidence statement (s)**

Clinical There is a statistically significant and clinically significant increase in operative or postoperative fracture of the femur with intramedullary implants compared to extramedullary implants for fixation of trochanteric extracapsular fractures. (LOW QUALITY)

There is no statistically significant difference in mortality, reoperation, and mean mobility score with intramedullary implants compared to extramedullary implants for fixation of trochanteric extracapsular fractures. (HIGH QUALITY)

There is no statistically significant difference in cut-out, infection, non-union and length of hospital stay with intramedullary implants compared to extramedullary implants for fixation of trochanteric extracapsular fractures. (MODERATE QUALITY)

There is no statistically significant difference in pain, with intramedullary implants compared to extramedullary implants for fixation of trochanteric extracapsular fractures. (LOW QUALITY)

No studies were identified investigating reverse oblique trochanteric extracapsular fractures.

Economic -No applicable evidence was identified regarding the cost-effectiveness of Intramedullary vs. extramedullary implants.

7 **10.6.1.5 Recommendations and link to evidence**

Recommendation	Use extramedullary implants such as a sliding hip screw in preference to an intramedullary nail in patients with trochanteric fractures above and including the lesser trochanter (AO classification types A1 and A2).
Relative values of different outcomes	The most important outcomes considered by the GDG include early and late mortality, re-operation, postoperative fracture, length of hospital stay and post fracture mobility.
Trade off between clinical benefits and harms	None of the studies reported have shown any advantage of intramedullary devices over extramedullary devices. Intramedullary devices had been shown to have a higher re-operation rate due to an increased incidence of periprosthetic fracture both in the perioperative period and the postoperative period (risk ratio 5.61). This may be due to the inclusion of studies with original nail designs no longer implanted. All other outcomes have been reported as similar. An additional meta-analysis is

included in Appendix G, page 510. By grouping studies using a cut off of publication after 2000, no changes to the existing evidence statement are made.

Economic considerations

In patients with trochanteric fractures above and including the lesser trochanter (AO classification types A1 and A2) the price of intramedullary fixation devices varies but on average is three times the price of sliding hip screws for short nails and five times the price for long nails. As pointed out in the clinical evidence statement, no significant benefit has been proven of the advantages of intramedullary devices over extramedullary devices, so that the GDG agreed to consider extramedullary implants cost-effective for hip fracture patients.

Quality of evidence

The level of evidence is high with numerous studies producing very similar findings.

Other considerations

All patients should be allowed to be mobilised full weight bearing after hip fracture surgery (see section 10.2). All modern implants are designed to be load sharing devices to facilitate this. Full weight bearing allows early mobilisation and rehabilitation.

The GDG highlighted this recommendation as a key priority for implementation.

1

2 **10.6.2 Intramedullary versus extramedullary implants for fixation of reverse**
3 **oblique trochanteric extracapsular fractures**

4 In the reverse oblique fractures, which lie anatomically between the trochanteric and the
5 subtrochanteric fractures there is loss of this lateral stabilizing cortical buttress. Such
6 fractures are difficult to adequately reduce and fix at the time of the surgery. It is then the
7 more unpredictable as to whether that adequate reduction will be retained during the
8 healing process whilst allowing early mobilisation of the patient

9 **10.6.2.1 Review question**

10 In patients undergoing repair for reverse oblique trochanteric extracapsular hip fractures
11 what is the clinical and cost effectiveness of extramedullary sliding hip screws compared to
12 intramedullary nails on mortality, surgical revision, functional status, length of stay, quality
13 of life, pain and place of residence after hip fracture?

14 **10.6.2.2 Clinical evidence**

15 No RCT evidence was identified.

16 **10.6.2.3 Economic evidence**

17 No cost-effectiveness evidence was identified.

18 **10.6.2.4 Research recommendations**

19 **Intramedullary versus extramedullary fixation**

1 The GDG recommended the following research question:

- 2 ➤ What is the clinical and cost effectiveness of intramedullary versus extramedullary
3 fixation on mortality, functional status and quality of life in patients with reverse
4 oblique trochanteric hip fracture?

5 **Why this is important**

6 Reverse oblique trochanteric fractures account for approximately 5 % of all trochanteric hip
7 fractures. This means it affects approximately over 1000 patients per year in the UK.
8 Presently there is little evidence as to which is the preferable implant (which can be either
9 extramedullary – outside the bone, or intramedullary - inside the bone). The potential
10 biomechanical advantage of intramedullary advantage may be offset by increased cost
11 (which can be over £1000 more expensive). A randomised trial comparing the two implants
12 using patient mobility, function and re-operation would allow a more informed choice of
13 treatment for this injury.

14

15 **10.6.3 Intramedullary versus extramedullary implants for fixation of**
16 **subtrochanteric extracapsular fractures**

17 Subtrochanteric fractures involve the shaft of the femur somewhere between the base of
18 the lesser trochanter and a point 5 cm distal to this. They may extend proximally or distally.
19 They have been considered as a separate group for practical purposes. Many of the
20 implants available for treating a standard trochanteric fracture are not long enough to
21 reach the intact bone distal to a subtrochanteric fracture. Thus whilst the general principles
22 of extra and intramedullary fixation described earlier still apply a different inventory of
23 implants to deal with these fractures is required.

24 It is noted that subtrochanteric fractures can often occur as a result of a metastatic
25 pathological deposit affecting the strength of the bone. The presence of pathological
26 deposits may not be obvious on the initial radiographs.

27 **10.6.3.1 Review question**

28 In patients undergoing repair for subtrochanteric extracapsular hip fractures, what is the
29 clinical and cost effectiveness of extramedullary sliding hip screws compared to
30 intramedullary nails on mortality, surgical revision, functional status, length of stay, quality
31 of life, pain and place of residence after hip fracture?

32 Four studies met the inclusion criteria for this review question with a total of 149 patients.
33 See evidence table 5.8, Appendix E and forest plots G107 to G111 in Appendix G.

34

35 **10.6.3.2 Clinical evidence**

1 **Table 10-56: Intramedullary vs. extramedullary implants for subtrochanteric extracapsular**
 2 **fracture – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality – 1 year ^{77,281}	2	RCT	no serious limitations	serious ^(c)	serious ^(b)	serious ^(a)
Reoperation – within follow up period of study ^{9,77,214,281}	4	RCT	no serious limitations	serious ^(c)	no serious indirectness	serious ^(a)
Cut-out (at latest follow up) ⁷⁷	1	RCT	serious ^(d)	no serious inconsistency	no serious indirectness	serious ^(a)
Infection (deep infection or requires reoperation) ^{214,281}	2	RCT	no serious limitations	no serious inconsistency	serious ^(b)	serious ^(a)
Non-union (at latest follow up) ^{77,281}	2	RCT	no serious limitations	no serious inconsistency	serious ^(b)	no serious imprecision

3 (a) The relatively few events and few patients give wide confidence intervals around the estimate of
 4 effect. This makes it difficult to know the true effect size for this outcome.

5 (b) These studies are comparing intramedullary nailing to a Medoff sliding plate or fixed angle blade
 6 plate.

7 (c) There is significant statistical heterogeneity in the results.

8 (d) Only one study with a small sample size.

9
 10 **Table 10-57: Intramedullary vs. extramedullary implants for subtrochanteric extracapsular**
 11 **fracture - Clinical summary of findings**

Outcome	Intramedullary	Extramedullary	Relative risk	Absolute effect	Quality
Mortality – 1 year	7/48 (14.6%)	5/42 (15%)	RR 0.93 (0.08 to 11.52)	10 fewer per 1,000 (from 138 fewer to 1578 more)	Very low
Reoperation – within follow up period of study	4/78 (5.1%)	11/71 (12.5%)	RR 0.56 (0.06 to 5.47)	55 fewer per 1,000 (from 117 fewer to 559 more)	Low
Cut-out (at latest follow up)	1/19 (5.3%)	1/13 (7.7%)	RR 0.68 (0.05 to 9.98)	25 fewer per 1,000 (from 73 fewer to 691 more)	Low
Infection (deep infection or requires reoperation)	3/45 (6.7%)	2/41 (5.9%)	RR 1.27 (0.28 to 5.88)	16 more per 1,000 (from 42 fewer to 288 more)	Low
Non-union (at latest follow up)	1/48 (2.1%)	9/42 (17.6%)	RR 0.15 (0.03 to 0.82)	150 fewer per 1,000 (from 32 fewer to 171 fewer)	Moderate

1 **10.6.3.3 Economic evidence**

2 No economic evidence was identified.

3 **10.6.3.4 Evidence statement (s)**

Clinical There is a statistically significant and clinically significant decrease in non-union with intramedullary implants compared to extramedullary implants for fixation of subtrochanteric extracapsular fractures. (MODERATE QUALITY)

There is no statistically significant difference in reoperation, cut-out and infection with intramedullary implants compared to extramedullary implants for fixation of subtrochanteric extracapsular fractures. (LOW QUALITY)

There is no statistically significant difference in mortality, with intramedullary implants compared to extramedullary implants for fixation of subtrochanteric extracapsular fractures. (VERY LOW QUALITY)

Economic No economic evidence was identified.

4

5 **10.6.3.5 Recommendations and link to evidence**

Recommendation	Use an intramedullary nail to treat patients with a subtrochanteric fracture.
Relative values of different outcomes	The GDG considered the most important outcomes to be functional status, pain, requirement for reoperations and wound healing complications.
Trade off between clinical benefits and harms	There was no evidence of a difference except for non-union of fracture. It is accepted by expert opinion that the treatment of choice is intramedullary fixation which allows splinting of the whole of the femoral shaft.
Economic considerations	Although intramedullary nails are more expensive than extramedullary implants, the latter lead to more patients with non-union of fracture, which would require more re-operation.
Quality of evidence	There were few studies investigating this type of fracture. Several studies were excluded as the population was from road traffic accidents, therefore high energy trauma fractures, which were excluded from the scope. The reported outcomes were predominantly of low quality.
Other considerations	Surgeons should use a technique where they are happy for the patient to mobilise fully weight bearing (see section 10.2). When patients suffer from subtrochanteric fractures it is advised to consider whether there is a pathological process which would increase the fracture risk (such as a metastatic deposit). As noted in the introduction subtrochanteric fractures may occur as a result of a pathological process in the bone such as metastatic disease. This pre-existing pathology may not always be recognised

on the initial radiographs. It is considered to be an additional advantage of using a long intramedullary device that it provides mechanical protection to a potentially diseased bone.

1
2

1 11 Mobilisation strategies

2 11.1 Introduction

3 Mobilisation is the process of re-establishing the ability to move between postures (for
4 example sit to stand), maintain an upright posture, and to ambulate with increasing levels
5 of complexity (speed, changes of direction, dual and multi-tasking).

6 Early restoration of mobility after surgery for hip fracture has been suggested as an
7 essential part of high quality care since the early 1980s^{309,310}. The suggested benefits are
8 minimisation of hospital stay, avoiding complications of prolonged bed confinement, and
9 re-establishing people into their normal environments^{168,168}.

10 Early restoration of mobility is an aspiration of many clinical services, although guidance on
11 the optimal time to re-mobilise patients and strategies that can be used to accelerate and
12 optimise recovery of mobility are less clear. Good quality clinical care, in particular effective
13 pain management should be considered essential components of early mobilisation and a
14 rehabilitation programme, as discussed in Chapter 7.

15 Specific therapeutic procedures, such as those implemented by physiotherapists and
16 occupational therapists have the potential to accelerate the recovery of mobility. Timing of
17 the intervention examined evidence about early (within 48 hours of surgery) mobilisation
18 and physiotherapy assessment, as opposed to later mobilisation (> 48 hours). Within the
19 type of intervention the GDG considered regimes that tested protocols delivering more
20 than one short session of physiotherapy per day (the benchmark for usual care), or more
21 intensive protocols than would comprise usual care. These protocols included intensive
22 strength training regimes (characterised by prescription and progression using recognised
23 American College of Sports Medicine criteria), intensive weight bearing exercise regimes
24 (supplemented by treadmill training), and increased numbers of physiotherapy usual care
25 sessions. Usual care was taken to be prescription of walking aids, gait re-education, and bed
26 exercises^{247,247}.

27 Mobility can be measured in a range of different ways. The most simple and basic mobility
28 indicators, are the ability to transfer independently. This is usually taken to be between a
29 bed and a chair, but not all investigators report the exact definition they have used. Chair
30 rise ability and time to complete chair rises, along with timed tests of walking and balance
31 have a long established history for measuring mobility. In addition, the GDG considered
32 muscle strength, length of stay, discharge destination, independence in activity of daily
33 living (such as washing, bathing) and more complex tasks (for example, meal preparation),
34 and mortality as outcomes. Measurement of falls, and time to first fall are considered good

1 safety indicators for interventions like early mobilisation, but no studies reported these
2 outcomes.

3 11.2 Early vs. delayed mobilisation

4 11.2.1 Review question

5 In patients who have undergone surgery for hip fracture, what is the clinical and cost
6 effectiveness of early mobilisation (<48 hours after surgery) compared to late mobilisation
7 on functional status, mortality, place of residence/discharge, pain and quality of life?

8 See Evidence Table 5.10, Appendix E and forests G112 to 116).

9 11.2.1.1 Clinical evidence

10 Only one, small randomised controlled trial was identified with 60 patients.

11 **Table 11-58: Early vs. delayed mobilisation – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Independent to transfer at day 7 ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	no serious imprecision
Independent to step at day 7 ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	no serious imprecision
Discharged to home ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	Serious imprecision ^(a)
Discharged to fast stream rehab ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	Serious imprecision ^(a)
Discharged to slow stream rehab ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	Serious imprecision ^(a)
Discharged to nursing home ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	Serious imprecision ^(a)
Mortality ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	Serious imprecision ^(a)
Mean walking distance ²⁴⁷	1	RCT	Serious limitations ^(b)	no serious inconsistency	no serious indirectness	— ^(c)

12 (a) The relatively few events and few patients give wide confidence intervals around the estimate of
13 effect. This makes it difficult to know the true effect size for this outcome.

14 (b) Unclear blinding and allocation concealment, also the small sample size makes it difficult to know
15 the true effect size for this outcome.

16 (c) The data is a mean with a range and therefore no relative risk was calculated. The wide range
17 around the mean indicates that the result may be imprecise.

18
19 **Table 11-59: Early vs. delayed mobilisation - Clinical summary of findings**

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Independent to transfer at day 7	16/29 (55.2%)	4/31 (12.9%)	RR 4.28 (1.62 to 11.3)	423 more per 1000 (from 80 more to 1329 more)	Moderate

Independent to step at day 7	10/29 (34.5%)	23/31 (74.2%)	RR 0.46 (0.27 to 0.8)	401 fewer per 1000 (from 148 fewer to 542 fewer)	Moderate
Discharged to home	5/29 (17.2%)	1/31 (3.2%)	RR 5.34 (0.66 to 43.06)	140 more per 1000 (from 11 fewer to 1357 more)	Low
Discharged to fast stream rehab	8/29 (27.6%)	14/31 (45.2%)	RR 0.61 (0.3 to 1.24)	176 fewer per 1000 (from 316 fewer to 108 more)	Low
Discharged to slow stream rehab	14/29 (48.3%)	16/31 (51.6%)	RR 0.94 (0.56 to 1.55)	31 fewer per 1000 (from 227 fewer to 284 more)	Low
Discharged to nursing home	1/29 (3.4%)	0/31 (0%)	RR 3.2 (0.14 to 75.55)	0 more per 1000 (from 0 fewer to 0 more)	Low
Mortality	1/29 (3.4%)	0/31 (0%)	RR 3.2 (0.14 to 75.55)	0 more per 1000 (from 0 fewer to 0 more)	Low
Mean walking distance, metres	82.55 (0.5-400)	34.7 (5-103)	N/A	_(a)	Moderate

1 (a) An absolute effect could not be calculated as the study did not provide a mean, only a range.

2 **11.2.1.2 Economic evidence**

3 No studies were identified.

4 The GDG was informed of the hourly cost of physiotherapy in a hospital setting for England
5 and Wales, which corresponds to £23⁶¹. Physiotherapist sessions delivered during the
6 weekends and during public holidays would be paid at an enhanced rate of pay of time and
7 a third (BMA contract, 2008).

8 **11.2.1.3 Evidence statement (s)**

Clinical There is a statistically significant and clinically significant increase in independence to transfer at day 7 for patients who had early mobilisation compared to delayed mobilisation. (MODERATE QUALITY)

There is a doubling in the distance walked at day 7 for patients who had early mobilisation compared to delayed mobilisation. (MODERATE QUALITY)

There is no statistically significant difference between early versus delayed mobilisation for discharge destination or mortality. (LOW QUALITY)

There is a statistically significant and clinically significant decrease in independence to step at day 7 for patients who had early mobilisation compared to delayed mobilisation. (MODERATE QUALITY)

Economic No studies were identified on the cost-effectiveness of early vs. delayed mobilisation.

1 11.2.2 Recommendations and link to evidence

Recommendation	Offer patients a physiotherapy assessment and, unless medically or surgically contraindicated, mobilisation on the day after surgery.
Relative values of different outcomes	Early mobilisation with a physiotherapist appears safe and is effective in promoting early recovery of ability to transfer without help of a person or walking aid. These outcomes are important markers of early recovery of mobility. See also, chapter 10 section 10.2 where the recommendation is made that surgeons should operate on patients with the aim to allow them to fully weight bear (without restriction) in the immediate postoperative period.
Trade off between clinical benefits and harms	The only outcome relating to harm or safety was mortality, which showed no statistically significant difference. If safety issues were a concern it is likely that they would be reflected in the overall functional outcomes, all of which improved or had no significant effect, therefore the GDG do not believe that harm is caused in relation to this evidence. If any attempt at mobilisation is supervised by a physiotherapist it should in any case be sensitive to limitations imposed by individuals' pre-fracture abilities and postoperative pain and fatigue. Thus a policy of early mobilisation with a physiotherapist should be seen as beneficial, and delayed only when individuals' clinical circumstances indicate this as appropriate.
Economic considerations	Evidence on the cost effectiveness of early mobilisation treatments is lacking. The GDG acknowledged that early mobilisation strategies will generally involve higher personnel costs (linked to the provision of physiotherapy sessions over the entire week, thus also during weekends and public holidays). However, the GDG considered the cost-savings associated with an earlier recovery of ability to transfer and step without help of a person or walking aid, and agreed that early mobilisation strategy represent a cost-effective intervention for our population.
Quality of evidence	There is only one RCT of low to moderate quality with a relatively small sample size (n = 60) and therefore the findings were interpreted with caution by the GDG.
Other considerations	<p>Early mobilisation protocols may require new service delivery models for weekend or 7 day physiotherapy services.</p> <p>The GDG also noted that albeit the intervention should be overseen by physiotherapists it is also important for nurses to re-enforce and encourage patients' mobility at all other times, under the guidance of the physiotherapist.</p>

The GDG highlighted this recommendation as a key priority for

implementation

1

2 11.3 Intensity of physiotherapy

3 11.3.1 Review question

4 In patients who have undergone surgery for hip fracture, what is the clinical and cost
5 effectiveness of intensive physiotherapy compared to non intensive physiotherapy on
6 functional status, mortality, place of residence/discharge, pain and quality of life?

7 See evidence table 5.10, Appendix E and forest plots G116 to G128.

8 11.3.1.1 Clinical evidence

9 Three randomised studies were found with a total of 288 patients, comparing three
10 different types of intensive physiotherapy/physical medicine programme. Hauer et al
11 (2002)^{139,140} investigated intensive, progressive strength training. Moseley et al (2009)^{216,216}
12 tested an intensive weight bearing exercise programme supplemented by treadmill gait re-
13 training programme, and Karumo (1977)^{171,171} investigated twice daily physiotherapy (of
14 one hours duration) in comparison to usual care (<=30 mins, once daily).

15 **Table 11-60: Intensive exercise or physiotherapy vs. usual care – Clinical study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Intensive physiotherapy (strength training)						
Leg-press strength fractured side (kg) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)
Leg extensor strength fractured side (Newtons) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Ankle plantar flexion strength fractured side (Newtons) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)
Walking speed – 3 months ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Tinetti's POMA ^(d) – overall ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Tinetti's POMA – part 1 (balance) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Tinetti's POMA – part 2 (gait) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Timed up-and-go (seconds) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Chair rise (seconds) ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)
Barthel's ADL ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)
Lawton's IADL ¹⁴⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Intensive physiotherapy (weight bearing exercise and treadmill training)						
Knee extensor strength – 4 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Knee extensor strength – 16 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Walking speed – 4 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Walking speed – 8 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Sit-to-stand test at 4 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Sit-to-stand test at 16 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Quality of life – 4 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Quality of life – 16 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Pain – 4 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Pain – 16 weeks ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Length of hospital stay ²¹⁶	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Intensive (more frequent) physiotherapy						
Adductor muscle strength (kp) at 9 weeks ¹⁷¹	1	RCT	serious ^(c)	no serious inconsistency	no serious indirectness	serious ^(b)
Length of hospital stay ¹⁷¹	1	RCT	serious ^(c)	no serious inconsistency	no serious indirectness	serious ^(b)

(a) Low number of subjects in each arm (N = 24) therefore the study may be underpowered.

(b) The relatively few events and few patients give wide confidence intervals around the estimate of effect. This makes it difficult to know the true effect size for this outcome.

(c) Method of randomisation, blinding and allocation concealment is unclear.

(d) POMA: Tinetti's performance oriented mobility assessment

Table 11-61: Intensive exercise or physiotherapy vs. usual care - Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Intensive physiotherapy (strength training)					
Leg-press strength fractured side (kg)	12	12	N/A	MD 21 higher (2.09 lower to 44.09 higher)	Low
Leg extensor strength fractured side	12	12	N/A	MD 17 higher (2.54 to 31.46)	Moderate

1
2
3
4
5
6
7
8

<i>Outcome</i>	<i>Intervention</i>	<i>Control</i>	<i>Relative risk</i>	<i>Absolute effect</i>	<i>Quality</i>
(Newtons)				higher)	
Ankle plantar flexion strength fractured side (Newtons)	12	12	N/A	MD 23 higher (2.23 lower to 48.23 higher)	Low
Walking speed – 3 months	12	12	N/A	MD 0.23 higher (0.05 to 0.41 higher)	Moderate
Tinetti's POMA - overall	12	12	N/A	MD 3 higher (0.41 lower to 6.41 higher)	Moderate
Tinetti's POMA – part 1 (balance)	12	12	N/A	MD 1.3 higher (0.54 lower to 3.14 higher)	Moderate
Tinetti's POMA – part 2 (gait)	12	12	N/A	MD 1.7 higher (0.15 lower to 3.55 higher)	Moderate
Timed up-and-go (seconds)	12	12	N/A	MD 0.8 lower (12.3 lower to 10.7 higher)	Low
Chair rise (seconds)	12	12	N/A	MD 1.8 lower (6.61 lower to 3.01 higher)	Low
Barthel's ADL	12	12	N/A	MD 3.1 lower (9.66 lower to 3.46 higher)	Low
Lawton's IADL	12	12	N/A	MD 0.4 higher (0.68 lower to 1.48 higher)	Moderate
<u>Intensive physiotherapy (weight bearing exercise and treadmill training)</u>					
Knee extensor strength – 4 weeks	80	80	N/A	MD 0.1 higher (1.12 lower to 1.32 higher)	Moderate
Knee extensor strength – 16 weeks	80	80	N/A	MD 1 higher (0.46 lower to 2.46 higher)	Moderate
Walking speed – 4 weeks	80	80	N/A	MD 0.05 higher (0.02 lower to 0.12 higher)	High
Walking speed – 8 weeks	80	80	N/A	MD 0.03 higher (0.07 lower to 0.13 higher)	High
Sit-to-stand test at 4 weeks	80	80	N/A	MD 0.05 higher (0.01 to 0.09 higher)	High
Sit-to-stand test at 16 weeks	80	80	N/A	MD 0.04 higher (0 to 0.08 higher)	High
Quality of life – 4 weeks	80	80	N/A	MD 0 higher (0.08 lower to 0.08 higher)	High
Quality of life – 16 weeks	80	80	N/A	MD 0 higher (0.09 lower to 0.09 higher)	High
Pain – 4 weeks	44/80 (55%)	41/80	RR 1.07 (0.8 to 1.44)	36 more per	High

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
		(51.3%)		1000 (from 102 fewer to 226 more)	
Pain – 16 weeks	30/80 (37.5%)	29/80 (36.3%)	RR 1.03 (0.69 to 1.55)	11 more per 1000 (from 112 fewer to 199 more)	High
Length of hospital stay (Moseley)	80	80	N/A	MD 3 higher (1.5 lower to 7.5 higher)	Moderate
Intensive (more frequent) physiotherapy					
Adductor muscle strength (kp) at 9 weeks	38	49	N/A	MD 0.76 lower (2.42 lower to 0.9 higher)	Low
Length of hospital stay	39	39	N/A	MD 2.8 lower (12.09 lower to 6.49 higher)	Low

1

2

11.3.1.2 Economic evidence

3

No studies were identified. A cost analysis was conducted based on the resources used in the studies included in the clinical review, which is reported in section 8.4 of Appendix Hof this guideline.

5

6

Table 11-62: Intensive exercise or physiotherapy vs. usual care – Economic study characteristics

Study	Limitations	Applicability	Other Comments
NCGC cost analysis	Minor limitations ^(a)	Partially applicable ^(b)	Cost analysis based on resources used in the studies included in the clinical review ^{140,171,216}

7

(c) No sensitivity analysis.

8

(d) UK study but does not estimate QALYs. One study ¹⁷¹ quite outdated. All studies not UK based and therefore may not reflect current NHS practice.

9

10

11

12

Table 11-63: Intensive exercise or physiotherapy vs. usual care - Economic summary of findings

Study	Incremental cost (£)	Incremental effects	ICER	Uncertainty
NCGC cost analysis	<ul style="list-style-type: none"> - £12 (strength training programme vs. usual care ¹⁴⁰) ^(a) - £180.18 (more intensive physiotherapy vs usual care ¹⁷¹) ^(b) - £827.62 (inpatient-based part of the weight bearing and treadmill exercise programme ²¹⁶) ^(c) 	N/A	N/A	N/R

13

(b) Intervention group slightly more costly than the control group because of the use of ad-hoc exercise equipment.

14

15

(c) Intervention group more costly because of longer physiotherapy sessions

16

(d) It was not possible to estimate the outpatient costs of the rehabilitation programme as insufficient information was given in the study.

17

1 *Evidence statement (s)*

Clinical Strength training

Additional, progressive strength training produces a statistically significant and clinically significant increase in leg extensor power, hip flexor strength and walking speed compared to placebo motor training (control) at 3 months after surgery. (HIGH QUALITY)

There is no statistically significant difference in basic or extended activities of daily living or gait and balance as measure by the Performance Orientated Mobility Assessment with strength training compared to placebo motor training (control) at 3 months after surgery. (HIGH QUALITY)

There is no statistically significant difference in timed up and go test and chair rises with strength training compared to placebo motor training (control) at 3 months after surgery. (MODERATE QUALITY)

Weight bearing exercise and treadmill training

There is no statistically significant difference in functional performance tests, quality of life, walking speed or pain with weight bearing exercise and treadmill gait training compared to the control. (HIGH QUALITY)

There is no statistically significant difference in length of hospital stay with weight bearing exercise and treadmill gait training compared to the control. (MODERATE QUALITY)

Intensive (more frequent) physiotherapy

There is no statistically significant difference in knee extensor strength adductor muscle strength or length of stay in hospital with an increased number of physiotherapy sessions per day compared to the control. (LOW QUALITY)

Economic All intensive exercise and physiotherapy programmes are more expensive than usual care, albeit the strength programme is only slightly more costly compared to usual care.

This evidence has minor limitations and partial applicability.

2 **11.3.2 Recommendations and link to evidence**

<i>Recommendation</i>	Offer patients mobilisation at least once a day and ensure regular physiotherapy review.
Relative values of different outcomes	<p>The outcomes considered most important were mobility, functional status, pain, quality of life and length of hospital stay.</p> <p>There is evidence of training effects in muscle strength and other variables which are known to be important determinants of ability to walk, and hence live independently. Further research is needed to confirm effects on outcomes including return to independent</p>

living, quality of life, health service resource, and time to discharge.

The evidence shows that there was no difference in once a day or twice a day physiotherapy for length of hospital stay and adductor muscle strength¹⁷¹, and thus the GDG are recommending at least once a day mobilisation.

Trade off between clinical benefits and harms

GDG consensus was that mobilisation at least once a day has potential benefits of improved mobility and balance, increased independence, and reduced need for institutional and social care. The included studies failed to show improvements for these outcomes, but are all small low quality studies. There is no evidence of harm from mobilisation once a day. There is potential to exacerbate pain and induce excessive fatigue, and training should be prescribed and overseen by a physiotherapist.

There is insufficient evidence to suggest what the exact dosing of physiotherapy should be, and this will vary according to the physical capabilities of each patient. Those who are very ill will not tolerate as much physical activity as those who are progressing well. The dosing should be based on a physiotherapist assessment. Hence the issue is one of professional judgement as we have no evidence to guide us any further. However, an additional observation is that the principles of management should not be any different for people with dementia, than those without.

Economic considerations

The GDG acknowledged the lack of cost-effective evidence on this question, and agreed that intensive rehabilitation sessions are likely to be more expensive than usual care. The GDG also noted that intensive rehabilitation can bring some benefits in terms of strength and on other factors affecting the ability to walk and live independently.

The GDG agreed that daily mobilisation sessions and regular physiotherapy review represent a cost-effective intervention for our patients.

Quality of evidence

Although 3 RCTs were included, the interventions were not comparable and could not be combined in a meta-analysis. The studies were all considered individually and the evidence base is limited. The quality of the evidence ranged from low to high, but due to few studies being identified the GDG considered the overall quality to be poor.

The economic evidence is based on the resources described in the programmes in the three RCTs included in the clinical review. Only the costs of the interventions and of the usual care programme were considered. The analysis is also only partially applicable in that, even current NHS unit costs were used, the actual level of resources reported in the trials may not reflect the current practice in the UK NHS.

Other considerations

GDG expert opinion indicates that patients may benefit from more intensive protocols of rehabilitation therapy (including

occupational and physiotherapy), but that more evidence is needed.

The GDG highlighted this recommendation as a key priority for implementation

1

2 **11.3.3 Research recommendations on mobilisation**

3 **11.3.3.1 Frequency of physiotherapy**

4 The GDG recommended the following research question:

- 5 ➤ What is the clinical and cost effectiveness of additional intensive physiotherapy
6 and/or occupational therapy (for example progressive resistance training) after hip
7 fracture?

8 **Why this is important**

9 The rapid restoration of physical and self care functions is critical to recovery from hip
10 fracture, particularly where the goal is to return the patient to preoperative levels of
11 function and residence. Approaches that are worthy of future development and
12 investigation include progressive resistance training, progressive balance and gait training,
13 supported treadmill gait re-training, dual task training, and activities of daily living training.
14 The optimal time point at which these interventions should be started requires clarification.

15 The ideal study design is a randomised controlled trial. Initial studies may have to focus on
16 proof of concept and be mindful of costs. A phase III randomised controlled trial is required
17 to determine clinical effectiveness and cost-effectiveness. The ideal sample size will be
18 around, 400 to 500 patients, and the primary outcome should be physical function and
19 health related quality of life. Outcomes should also include falls. A formal sample size
20 calculation will need to be undertaken. Outcomes should be followed over a minimum of 1
21 year, and compare if possible, either the recovery curve for restoration of function or time
22 to attainment of functional goals.

23

1 12 Multidisciplinary management

2 12.1 Introduction

3 Multidisciplinary care is central to the management of frail older people with multiple
4 medical, psychological and social problems. Since these are the people who typically suffer
5 hip fracture every Trauma Unit will provide some form of multidisciplinary care. Although
6 the prevalence of comorbidity is generally lower in younger patients, the key principles of
7 multidisciplinary intervention are applicable across the adult age spectrum and the same
8 skills and organisational approaches derived within the development of a focus on the older
9 population should be provided irrespective of chronological age.

10 In this chapter the evidence for the different models of enhanced inpatient and community
11 management were considered that have evolved to meet the specific needs of patients
12 with hip fracture.

13 Secondary prevention of fracture by means of the assessment and management of both
14 osteoporosis^{234,235} and risk of falling²²⁷ are covered in separate NICE guidance. It is,
15 however, important in practice that the elements of multidisciplinary management covered
16 in this guidance relate in an organized manner closely and reliably with these secondary
17 prevention programmes to deliver all the elements of comprehensive care required by each
18 patient. The precise organizational approach to this differs amongst centres. In some there
19 is considerable overlap and/or cross-representation between the secondary prevention
20 programmes and the service models covered in this guideline.

21 Units across the UK have adopted a variety of multidisciplinary service models, but most
22 have at least some form of access to geriatrician input into the care of these patients. Local
23 circumstances and expertise have determined the precise model developed in different
24 centres, but in general these are variations on the following four approaches.

25 The traditional model of orthopaedic care - 'usual care'.

- 26 • The patient with hip fracture is admitted to a trauma ward where the orthopaedic
27 surgical team lead both their surgical care and subsequent rehabilitation.
28 Geriatrician input to such wards may be limited, with referrals and medical queries
29 being dealt with on a consultative basis by the on-call medical registrar or on
30 occasional geriatrician visits, but without a proactive geriatrician lead to the
31 multidisciplinary team.

32 A more collaborative model of trauma ward working is formal 'orthogeriatric' care - with
33 trauma patients admitted to a specialised ward under the joint care of both geriatricians

- 1 and orthopaedic surgeons. Surgical and geriatrician ward rounds may happen
2 independently, or be combined in multidisciplinary ward rounds.
- 3 • This collaborative model is particularly relevant to hip fracture patients. Such joint
4 working can thus lead to the development of a formal 'Hip Fracture Programme'
5 (HFP), with the geriatric medical team contributing to joint preoperative patient
6 assessment, and increasingly taking the lead for postoperative medical care,
7 multidisciplinary rehabilitation (MDR) and discharge planning.
- 8 Both 'traditional' and 'orthogeriatric' models of the acute trauma ward may continue to
9 care for patients throughout their recovery and rehabilitation following hip fracture, or
10 each may be followed by a transfer of some patients to other models of rehabilitation.
- 11 • In some centres, surgical care and initial mobilisation is followed by early
12 postoperative transfer to a 'Geriatric Orthopaedic Rehabilitation Unit' (GORU) - a
13 separate geriatrician-led rehabilitation ward. The extent of surgical input to the
14 GORU varies, depending on how early patients are moved from the acute trauma
15 wards.
- 16 • In other centres, similar patients would be transferred to a generic 'Mixed
17 Assessment and Rehabilitation Unit' (MARU), able to accept patients with a variety
18 of medical, surgical and orthopaedic conditions.
- 19 A further service model is some form of community rehabilitation.
- 20 • One approach is 'Early Supported Discharge' (ESD) or 'Intermediate Care' at home.
21 Patients are discharged home from the acute trauma ward, or in some cases a
22 rehabilitation ward within the hospital, with a supported 4-6 week rehabilitation
23 package. This may include patients living in care homes but in many parts of the
24 country is limited to patients returning to live independently in their own homes.
- 25 • Alternatively, patients with more complex needs may be moved for rehabilitation
26 to an Intermediate Care facility outside the hospital setting, such as a care home, or
27 a community hospital. Again this will vary depending on the provision of services
28 available locally.
- 29
- 30

1

2 12.2 Hospital-based multidisciplinary rehabilitation versus usual care

3 Multidisciplinary rehabilitation (MDR) after hip fracture has been taken by the GDG to
4 incorporate medicine, nursing, physiotherapy, occupational therapy and social care as core
5 components of assessment and management. Additional components may include
6 dietetics, pharmacy and clinical psychology.

7 The GDG also assumes:

- 8 • The required degree of relevant specialist expertise in each case.
- 9 • Formal arrangements for co-ordination/teamwork, and
- 10 • Regular on-going multidisciplinary assessment.

11 'Usual care' will be taken to imply the traditional model, with *ad hoc* or selective referral to
12 some or all of the separate MDR components listed above, but without formal
13 arrangements for co-ordinated multidisciplinary teamwork.

14 In contrast, the different models of 'orthogeriatric care' all assume the involvement of a
15 geriatrician, in addition to the orthopaedic surgical team, in the development and
16 supervision of a formal process of coordinated multidisciplinary care.

17 Such orthogeriatric models have been sub-divided into:

- 18 • Those focused predominantly or exclusively on the acute trauma ward; typified by
19 the HFP model⁴³.
- 20 • Those provided in a subsequent inpatient rehabilitation setting (with GORU and
21 MARU having been combined because no evidence has addressed a comparison of
22 these models).
- 23 • Those with a community focus (the focus of Section 12.4).

24 12.2.1 Review questions

25 In this section two review questions were combined as the evidence overlapped and could
26 not be separated in a useful way. The questions were:

27 In patients with hip fracture what is the clinical and cost effectiveness of hospital-based
28 multidisciplinary rehabilitation on functional status, length of stay in secondary care,
29 mortality, place of residence/discharge, hospital readmission and quality of life?

30 All the published studies included in the analysis of hospital-based MDR are of models that
31 include geriatrician input. The results of a collective analysis of all such studies therefore
32 reflect both the effectiveness of hospital-based MDR, and the overall value of
33 orthogeriatrician involvement in hip fracture care.

34 In addition, the benefits of different models of hospital-based MDR can be considered by
35 comparing 'usual care' with the two general sub-types of orthogeriatric care:

- 1 • Hip Fracture Programme (HFP)
- 2 • Geriatric Orthopaedic Rehabilitation Unit (GORU), or near equivalents such as a
3 Mixed Assessment and Rehabilitation Unit (MARU).
- 4 In patients with hip fracture what is the clinical and cost effectiveness of orthogeriatrician
5 involvement in the whole pathway of assessment, peri-operative care and rehabilitation on
6 functional status, length of stay in secondary care, mortality, place of residence/discharge,
7 hospital readmission and quality of life?
- 8 The geriatrician is increasingly seen as having a key role in the integration of initial
9 assessment and peri-operative care with the coordinated MDR (in whatever setting) which
10 follows it.
- 11 The usefulness of this early element of orthogeriatric input has been assessed; an element
12 that it is central to the first of the two models (HFP), but lacking from the second
13 (GORU/MARU). In the absence of trials directly comparing the two models the impact of
14 early geriatrician involvement can only be inferred from any differences that might be
15 apparent when each is compared to 'usual care'.
- 16 11 studies met the inclusion criteria for this question, with a total of 2214 patients. See
17 Evidence Table 5.10, Appendix E and forest plots G129 to 138 in Appendix G.
- 18

1 **12.2.1.1 Clinical evidence**2 **Table 12-64: Hospital based multidisciplinary rehabilitation vs. usual care – Clinical**
3 **study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality at 6 months – GORU/MARU ^{113,222}	2	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Mortality at 12 months – GORU/MARU ^{107,158,176,319}	4	RCT	serious ^(a, b, c)	no serious inconsistency	no serious indirectness ^(d)	no serious imprecision
Mortality at 12 months – HFP ^{44,305,325,344}	4	RCT	serious ^(e, f)	no serious inconsistency	no serious indirectness	no serious imprecision
Mortality (at discharge) – GORU/MARU ^{107,113,158,176,222,319}	6	RCT	serious ^(a, b, c, g)	no serious inconsistency	no serious indirectness ^(d)	no serious imprecision
Mortality (at discharge) – HFP ^{325,344}	2	RCT	no serious limitations ^(f)	serious ^(h)	no serious indirectness	serious ^(h)
Non-recovery/decline in walking at 6 months – GORU/MARU ²²²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Decline in transfers (bed to chair etc) at – GORU/MARU ²²²	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
More dependent (based on Katz index) at 1 year – GORU/MARU ^{176,319}	2	RCT	serious ^(b, g)	no serious inconsistency	no serious indirectness ^(d)	serious ^(k)
Non-recovery in activities of daily living (ADL) at 1 year – GORU/MARU ³¹⁹	1	RCT	no serious limitations ^(g)	no serious inconsistency	no serious indirectness	serious ^(k)
Non-recovery of ADL/decline in walking at 1 year – HFP ^{305,344}	2	RCT	no serious limitations ^(e, f)	no serious inconsistency	no serious indirectness	serious ^(k)
Chinese Barthel Index at 6 months - HFP ³⁰⁵	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(k)
Modified Barthel Index at 6 months – HFP ³²⁵	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(k)

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Length of hospital stay - GORU/MARU ^{107,1} 13,176,222,319	5	RCT	no serious limitations	serious ^(l,i)	no serious indirectness ^(d)	serious ^(k)
Length of hospital stay - HFP ^{44,305,325}	3	RCT	no serious limitations	serious ^(l)	no serious indirectness	serious ^(h)
Pressure sores ³⁴⁴	1	RCT	no serious limitations ^(f)	no serious inconsistency	no serious indirectness	no serious imprecision
Heart failure ³⁴⁴	1	RCT	no serious limitations ^(f)	no serious inconsistency	no serious indirectness	serious ^(k)
Pneumonia ³⁴⁴	1	RCT	no serious limitations ^(f)	no serious inconsistency	no serious indirectness	serious ^(h)
Confusion ³⁴⁴	1	RCT	no serious limitations ^(f)	no serious inconsistency	no serious indirectness	no serious imprecision
Chest infection, cardiac problem, bedsore ³²⁵	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(k)
Stroke, emboli ³²⁵	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(h)
Delirium ²⁰³	1	RCT	no serious limitations	no serious inconsistency	serious ^(m)	serious ^(k)
Severe delirium ²⁰³	1	RCT	no serious limitations	no serious inconsistency	serious ^(m)	serious ^(k)
Readmitted to hospital during follow-up – GORU/MARU ^{107,3} 19	2	RCT	serious ^(c, g)	serious ⁽ⁿ⁾	no serious indirectness	no serious imprecision
Readmitted to hospital during follow-up – HFP ^{44,305,325,344}	4	RCT	serious ^(f, g)	no serious inconsistency	no serious indirectness	no serious imprecision

- 1 (a) Intervention group in Huusko 2002^{157,158} had greater number of patients with dementia (32/120 vs. 20/123); fewer were functionally independent in ADL before hip fracture (41 vs. 66).
- 2 (b) Kennie 1988^{176,176}: difference in age mental state. Control group average age higher and with more moderate and severe impairment.
- 3 (c) In Galvard 1995^{107,107}, the intervention group were older than usual care (79.1 vs. 73.6), and there were a higher proportion of patients with subtrochanteric fractures, which often require longer rehab (12% vs. 4%).
- 4 (d) Kennie 1988^{176,176} is an all female population.
- 5 (e) In Shyu 2008³⁰⁵ the patient's insurance policy determined the number of physiotherapy sessions in the control group.
- 6 (f) In Vidan 2005^{344,344} there is potential for contamination bias given both groups were on the same ward and had the same staff.
- 7 (g) In Stenvall 2007a^{319,320}, outpatient rehabilitation was not standardised.
- 8 (h) The relatively few events and few patients give wide confidence intervals around the estimate of effect. This makes it difficult to know the true effect size for this outcome.
- 9 (i) Galvard 2002^{107,107}, author's note that geriatric department had less experience with hip fracture patients than the orthopaedic ward, which may have contributed to increased length of stay in intervention group.
- 10 (j) The intervention in Naglie 2002^{222,222} was expected to increase the length of stay in hospital.

- 1 (k) The wide confidence intervals around the estimate make it difficult to determine and effect size for
 2 this outcome.
 3 (l) There is significant statistical heterogeneity between the studies. This could be due to the variation
 4 in intervention and country of study.
 5 (m) The intervention in Marcantonio 2001^{203,203} does not examine multidisciplinary rehabilitation in the
 6 form of an HFP, but focuses on the value of early comprehensive geriatric assessment and targeted
 7 intervention.
 8 (n) There is significant statistical heterogeneity between the studies. However, this could be due to
 9 differences in access to hospital services and follow up procedures.
 10
 11

Table 12-65: Hospital based multidisciplinary rehabilitation vs. Usual care - Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality at 6 months – GORU/MARU	31/238 (13%)	44/263 (16.8%)	RR 0.79 (0.52 to 1.21)	35 fewer per 1,000 (from 80 fewer to 35 more)	High
Mortality at 12 months – GORU/MARU	89/455 (19.6%)	96/466 (19.7%)	RR 0.95 (0.74 to 1.23)	10 fewer per 1000 (from 54 fewer to 47 more)	Moderate
Mortality at 12 months – HFP	72/400 (18%)	90/404 (21%)	RR 0.81 (0.61 to 1.06)	42 fewer per 1000 (from 87 fewer to 13 more)	Moderate
Mortality (at discharge) – GORU/MARU	46/693 (6.6%)	62/729 (8.4%)	RR 0.78 (0.54 to 1.13)	19 fewer per 1000 (from 39 fewer to 11 more)	Moderate
Mortality (at discharge) – HFP	3/193 (1.6%)	11/197 (5.8%)	RR 0.27 (0.07 to 0.96)	41 fewer per 1000 (from 2 fewer to 52 fewer)	Low
Non-recovery/decline in walking at 6 months – GORU/MARU	59/124 (47.6%)	56/117 (47.9%)	RR 0.99 (0.76 to 1.29)	5 fewer per 1000 (from 115 fewer to 139 more)	Moderate
Decline in transfers (bed to chair etc) at – GORU/MARU	45/124 (36.3%)	44/117 (37.6%)	RR 0.96 (0.69 to 1.34)	15 fewer per 1000 (from 117 fewer to 128 more)	Moderate
More dependent (based on Katz index) at 1 year – GORU/MARU	57/127 (44.9%)	77/111 (72.2%)	RR 0.64 (0.51 to 0.81)	250 fewer per 1000 (from 132 fewer to 340 fewer)	Low
Non-recovery in activities of daily living (ADL) at 1 year – GORU/MARU	51/84 (60.7%)	59/76 (77.6%)	RR 0.78 (0.63 to 0.96)	171 fewer per 1000 (from 31 fewer to 287 fewer)	Moderate
Non-recovery in ADL/decline in walking at 1 year – HFP	86/207 (41.5%)	108/207 (52.2%)	RR 0.79 (0.65 to 0.97)	171 fewer per 1000 (from 31 fewer to 287 fewer)	Moderate
Chinese Barthel Index at 6 months - HFP	73	75	N/A	MD 6.17 (0.86 to 13.2)	Moderate

Modified Barthel Index at 6 months – HFP	33	27	N/A	MD 6.3 (0.53 to 13.13)	Moderate
Length of hospital stay - GORU/MARU	572	606	N/A	MD 1.32 (-12.83 to 15.47)	Low
Length of hospital stay - HFP	245	240	N/A	MD -6.06 (-14.5 to 2.38)	Low
Pressure sores	8/155 (5.2%)	27/164 (16.5%)	RR 0.31 (0.15 to 0.67)	114 fewer per 1000 (from 54 fewer to 140 fewer)	High
Heart failure	12/155 (7.7%)	5/164 (3.1%)	RR 2.54 (0.92 to 7.04)	47 more per 1000 (from 2 fewer to 184 more)	Moderate
Pneumonia	6/155 (3.9%)	6/164 (3.7%)	RR 1.06 (0.35 to 3.21)	2 more per 1000 (from 24 fewer to 81 more)	Moderate
Confusion	53/155 (34.2%)	67/164 (40.9%)	RR 0.84 (0.63 to 1.11)	65 fewer per 1000 (from 151 fewer to 45 more)	High
Chest infection, cardiac problem, bed sore	6/38 (15.8%)	13/33 (39.4%)	RR 0.4 (0.17 to 0.94)	236 fewer per 1000 (from 24 fewer to 327 fewer)	Moderate
stroke, emboli	4/38 (10.5%)	1/33 (3%)	RR 3.47 (0.41 to 29.56)	75 more per 1000 (from 18 fewer to 865 more)	Moderate
Delirium	20/62 (32.3%)	32/64 (50%)	RR 0.65 (0.42 to 1)	175 fewer per 1000 (from 290 fewer to 0 more)	Low
Severe delirium	7/62 (11.3%)	18/64 (28.1%)	RR 0.4 (0.18 to 0.89)	169 fewer per 1000 (from 31 fewer to 231 fewer)	Low
Readmitted to hospital during follow-up - GORU/MARU	74/256 (28.9%)	87/262 (33.2%)	RR 0.86 (0.67 to 1.12)	46 fewer per 1000 (from 110 fewer to 40 more)	Low
Readmitted to hospital during follow-up – HFP	86/373 (23.1%)	78/378 (17%)	RR 1.14 (0.87 to 1.48)	29 more per 1000 (from 27 fewer to 99 more)	Moderate

1 **12.2.1.2 Economic evidence**

2 The included studies for hospital-based MDR consisted of Cameron (1994)^{42,45}, Galvard
3 (1995)^{107,107}, Farnworth (1994)^{91,91} and Huusko (2002)^{157,158}. Further details on the studies
4 are available in Evidence Table 16 of Appendix F. An HTA by Cameron (2000)⁴¹ was
5 excluded because the studies were grouped in a different way to that considered for our
6 clinical review, and therefore its cost analysis was not applicable for our review question.

1 An original decision analysis has been conducted comparing the cost-effectiveness of the
2 HFP vs. GORU/MARU vs. usual care. A Markov model was developed, adopting a life-time
3 horizon.

4 An indirect comparison between the HFP and GORU/MARU models of care was made as no
5 evidence was available which compares directly the two rehabilitation programmes. The
6 usual care arms in the trials of HFP vs. usual care and of GORU/MARU vs. usual care were
7 combined for this purpose.

8 Treatment effects were based on the findings of the clinical review and applied only up to 1
9 year from follow-up. Resource use was determined from the NHS and PSS perspective.
10 Effectiveness was measured in QALYs. Costs and QALYs were discounted at a rate of 3.5%.
11 Please see section 8.6 of Appendix H for further details.

12

1 **Table 12-66: Hospital based multidisciplinary rehabilitation vs. usual care - Economic study**
 2 **characteristics**

Study	Limitations	Applicability	Other Comments
Cameron 1994 ⁴⁵ – HFP	Potentially serious limitations ^(a)	Partial applicability ^(b)	Accelerated rehab was compared to usual care. The follow up time was 4 months.
Farnworth 1994 ⁹¹ – HFP	Potentially serious limitations ^(c)	Partial applicability ^(b)	Fractured Hip Management Program (FHMP) was compared to usual care. The follow up time was 6 months.
Galvard 1995 ¹⁰⁷ - GORU	Potentially serious limitations ^(d)	Partial applicability ^(e)	Rehabilitation in a geriatric department was compared to usual care. The follow up time was 1 year.
Huusko (2002) ¹⁵⁸ - MARU	Potentially serious limitations ^(f)	Partial applicability ^(g)	Intensive multidisciplinary geriatric team rehabilitation versus usual care. Follow up was 1 year.
NCGC economic model	Minor limitations ^(h)	Direct applicability	Cost-effectiveness analysis of HFP vs. GORU/MARU vs. usual care based on the meta-analysis of the trails included in the clinical review of this guideline

- 3 (a) Patients in the intervention and control group treated in the same ward, so that results could be biased
 4 due to an underestimation of the cost effectiveness of accelerated rehab.
 5 (b) Study conducted in Australia. Not a CUA.
 6 (c) The year in which cost data were collected is not clear. The duration of follow up is not clear. HRQoL not
 7 calculated. The statistical significance of the outcome and cost measures between the two groups was
 8 not reported. Outcome at 1 year was not known for 12% of the intervention and 14% of the control
 9 group.
 10 (d) No sensitivity analysis was performed to test robustness of findings. HRQoL not calculated. The source
 11 used to estimate the unit cost of resources was unclear.
 12 (e) Study conducted in Sweden. Not a CUA.
 13 (f) Not a cost-effectiveness analysis. No sensitivity analysis was performed. 38 patients were lost during
 14 follow up. The year(s) at which cost data refer to is not clear. Imbalance of baseline characteristics.
 15 Intervention group had a more patients with dementia (32/120 vs. 20/123, and fewer who were
 16 functionally independent in ADL before hip fracture (41 vs. 66).
 17 (g) Study conducted in Finland. Not a CUA.
 18 (h) Treatment effects from meta-analysis of clinical trials available up to 1 year from follow-up.

1 ***Table 12-67: Hospital based multidisciplinary rehabilitation vs. usual care - Economic***
2 ***summary of findings***

Study	Incremental cost per patient (£)	Incremental effects	ICER	Uncertainty
Cameron 1994 – HFP	-£956 ^(a)	Several outcomes were reported ^(b)	Accelerated rehabilitation is the dominant strategy (less costly and more effective)	Threshold sensitivity analysis: results not sensitive to changes in % of patients recovering nor to the definition of recovery. Accelerated rehab becomes more costly than usual care when difference in LOS less than 1.5-2 days and when cost of treatment more than 40% per bed day.
Farnworth 1994 – HFP	£784 ^(c)	Several outcomes were reported ^(d)	N/A	Deterministic sensitivity analysis showed that results were robust to changes in the time spent to get patients to surgery more quickly; to the proportion of nursing home patients and to the average cost of the final days of a patient's stay
Galvard 1995 - GORU	-£665 ^(e)	Several outcomes were reported ^(f)	N/A	N/R
Huusko 2002 - MARU	£1310 ^(g)	Several outcomes were reported ^(h)	N/A	N/R
NCGC economic model – HFP vs. GORU/MARU vs. usual care (Appendix H)	-£ 2,000 (HFP vs. GORU/MARU) -£25,000 (HFP vs. usual care) ⁽ⁱ⁾	-0.13 QALYs (HFP vs. GORU/MARU) -1.01 QALYs (HFP vs. usual care) ⁽ⁱ⁾	HFP is the dominant strategy compared to both GORU/MARU and usual care	Deterministic sensitivity analysis showed that results were sensitive to changes in the proportion of patients discharged to their own home following rehabilitation. A probabilistic sensitivity analysis showed that there is no uncertainty that hospital MDR is better than usual care. However, there is some uncertainty over the cost-effectiveness of HFP vs. GORU/MARU. ^(k) 95% CI (HFP vs usual care and GORU/MARU vs usual care): usual care dominated. 95% CI (HFP vs. GORU/MARU): HFP dominant – GORU dominant.

1
2
3

(a) Accelerated rehab is cost saving. A\$ converted using the PPP of 1990. $p=0.186$. The cost components estimated were: inpatients hospital costs, readmissions, community support services, institutional care.

- 1 (b) No. of patients recovered at 4 months from surgery (mean Barthel index score): 63 (49.6%) vs. 52
 2 (41.6%); 95% CI (-3% to 21%). Median length of stay (days, interquartile range): 13 (7-25) vs. 15 (8-
 3 44).
- 4 (c) Fractured Hip Management Program (FHMP) is cost saving.
- 5 (d) FHMP entails lower mortality and readmission at 1 year, and lower length of stay.
- 6 (e) Swedish Krona (SEK) converted using the PPP of 1989; Rehabilitation in geriatric department more
 7 expensive than usual care (£665 per patient)
- 8 (f) The intervention had a lower level of readmissions to hospital than usual care (36 vs. 57; p value
 9 NR) but it had a higher mortality at 1 year (45 vs. 40, p value NR) and a higher mean length of stay
 10 in hospital (53.3 vs. 28 days, p value NR).
- 11 (g) The study expressed costs in Euros (values of 1999). The intervention is more costly than usual care
 12 (p value NR).
- 13 (h) Intervention did not statistically differ from usual care in terms of mortality at 12 months (15% vs.
 14 16%); mortality at discharge (5 vs. 5) and length of stay in hospital during 1 year (80 vs. 80 days),
 15 and number of patients reporting complications (51% vs. 46%, p=0.4). Patients in the intervention
 16 group regained their independency in the IADL functions faster (p=0.005) than usual care at 3
 17 months (but after 1 year there was no significant difference between the two groups).
- 18 (i) The mean costs associated with HFP were estimated to be £34,000, for GORU/MARU £36,000 and
 19 for usual care £59,000.
- 20 (j) The mean effectiveness corresponded to 3.74 QALYs for HFP, 3.61 QALYs for GORU/MARU and 2.73
 21 QALYs for usual care.
- 22 (k) Usual care was never the most cost-effective strategy. At a willingness to pay of £20k per
 23 incremental QALY, HFP was found to be the most cost-effective option in 70% of the 10,000
 24 simulations run in the PSA, while GORU/MARU was the most cost-effective option in 30% of the
 25 simulations. At a willingness to pay of £30K per incremental QALY, HFP was found to be the most
 26 cost-effective option in 80% of the 10,000 simulations run in the PSA, while GORU/MARU was the
 27 most cost-effective option in 20% of simulations.

30 12.2.2 Evidence statement (s)

Clinical

Hospital-based MDR (GORU/MARU)

There is a statistically significant and clinically significant reduction in pressure sores with hospital-based MDR (GORU/MARU) compared to usual care. (HIGH QUALITY)

There is a statistically significant, but not clinically significant improvement in recovery of activities of daily living at 1 year with hospital-based MDR (GORU/MARU) compared to usual care. (MODERATE QUALITY)

There is a statistically significant, but not clinically significant improvement in transfers (bed to chair) and being more dependent (Katz index) at 1 year with hospital-based MDR (GORU/MARU) compared to usual care. (LOW QUALITY)

There is a statistically significant, but not and clinically significant reduction in severe delirium with hospital-based MDR (GORU/MARU) compared to usual care. (LOW QUALITY)

There is no statistically significant difference in mortality at 6 months and functional outcomes at 6 months between hospital-based MDR (GORU/MARU) and usual care. (MODERATE QUALITY)

There is no statistically significant difference in mortality at 12 months and mortality at discharge between hospital-based MDR (GORU/MARU) and usual

care. (MODERATE QUALITY)

There is no statistically significant difference in length of hospital stay and readmission to hospital between hospital-based MDR (GORU/MARU) and usual care. (LOW QUALITY)

Hip fracture programme (HFP)

There is a statistically significant and clinically significant improvement in functional outcomes at 1 year with HFP compared to usual care. (MODERATE QUALITY)

There is a statistically significant and clinically significant reduction in mortality at discharge between HFP and usual care. (LOW QUALITY)

There is no statistically significant difference in mortality at 12 months and readmission to hospital, between HFP and usual care. (MODERATE QUALITY)

There is no statistically significant difference in length of hospital stay, between HFP and usual care. (LOW QUALITY)

Economic

HFP is the dominant strategy (less costly and more effective) than both GORU/MARU and usual care as a hospital based multidisciplinary rehabilitation of hip fracture patients. This evidence has minor limitations and direct applicability.

1 **12.2.3 Recommendations and link to evidence**

Recommendation	<p>From admission, offer patients a formal, acute orthogeriatric or orthopaedic ward-based Hip Fracture Programme that includes all of the following:</p> <ul style="list-style-type: none"> • orthogeriatric assessment • rapid optimisation of fitness for surgery • early identification of individual goals for multidisciplinary rehabilitation to recover mobility and independence, and to facilitate return to prefracture residence and long-term wellbeing. • continued, coordinated, orthogeriatric and multidisciplinary review • liaison or integration with related services, particularly mental health, falls prevention, bone health, primary care and social services. • clinical and service governance responsibility for all stages of the pathway of care and rehabilitation, including those delivered in the community.
-----------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Relative values of different outcomes

Patients, clinical staff and health services share the objective of safely returning patients to their original functional state and residence as quickly as possible. However, these objectives are often in conflict – for instance earlier discharge may be at the

Trade off between clinical benefits and harms

expense of functional improvement, while length of stay may increase if mortality is prevented among frailer individuals.

Therefore the most important outcomes considered by the GDG were functional status, length of stay, discharge destination and mortality. All these outcomes were incorporated into an original economic decision analysis.

Studies of MDR show no significant evidence of harm and a trend towards improved outcomes across all outcomes. There is no suggestion of harm resulting from orthogeriatric collaboration in the HFP literature.

Evidence to support the effectiveness of coordinated hospital-based orthogeriatric MDR is derived from studies of both HFP and GORU/MARU models.

Taken together these studies suggest:

- improvement in functional outcome at 1 year, though this has not been shown to lead to greater success in achieving patients' objective of returning to their original residence.
- trend toward reduced mortality at discharge, 1, 6 and 12 months, which must reflect an effect in reducing medical and/or surgical complications (problems with diagnosis, definition and ascertainment leave this issue unclear).
- reduced hospital length of stay, though some studies only examined orthopaedic ward length of stay, so the preferred measure of 'super-spell' (the total time until return home) was inconsistently characterised.

Additional evidence supporting the effectiveness of a hospital-based model incorporating continuous orthogeriatrician supervision is derived from studies of Hip Fracture Programmes which suggest:

- reduced patient mortality at discharge and follow-up
- improved functional outcomes
- reduced hospital LOS
- reduced risk of delirium²⁰³.

Both HFP and GORU/MARU proved markedly more cost-effective than usual care, although HFP emerged as the dominant strategy. The GDG took the view that HFP approach is also preferable because of its provision of a more extensive programme of multidisciplinary care that:

- supports admission assessment and peri-operative care, in addition to rehabilitation, discharge planning and follow-up
- addresses the needs of all patients, including those who might be viewed as inappropriate for a GORU/MARU (because of ongoing orthopaedic, medical or psychiatric problems)
- provides a coordinated multidisciplinary structure that will

support other recommendations in this guideline (eg. early operation).

Economic considerations

There were no published economic studies on hospital-based MDR for hip fracture patients, so an original decision analysis was developed to determine the cost-effectiveness of HFP vs. GORU/MARU vs. usual inpatient rehabilitation (usual care).

The cost-effectiveness model was based on an indirect comparison of randomised trials, but clearly showed that usual care was not the optimal approach.

The increased costs of hospital MDR were more than offset by:

- reduction in the acute hospital stay costs, including those associated with complications such as delirium and pressure sores.
- a reduction in the level of domiciliary social care costs as a result of increased probability of regaining pre-fracture independence in activities of daily living.
- reduction in costs for patients who avoid the need for long-term care in a residential or a nursing home.

HFP was the strategy with the highest incremental net benefit averaged across all the probabilistic simulations, and appeared to be the optimal strategy in the cost-effectiveness analysis both in comparison to usual care, and in comparison to GORU/MARU.

However, there remains some uncertainty about the relative cost-effectiveness of HFP and GORU/MARU. In particular, the results were sensitive to the proportion of patients returning home after completing the rehabilitation programme. Sensitivity analysis suggested that if the probability of returning home in the GORU/MARU programme was increased to 83% (instead of 79% as in the base case) then GORU/MARU would become the optimal strategy.

Quality of evidence

The GDG noted that the precision of the cost-effectiveness analysis was partially limited by the lack of clinical trials directly comparing HFP vs. GORU/MARU, and by the heterogeneous patient population in the meta-analysis of clinical trials on which the cost-effectiveness analysis is based.

However, the GDG agreed that the outcomes used in the economic analysis were overall of moderate quality and that the decision model is likely to provide a relatively unbiased estimate of cost effectiveness.

There are consistent trends towards benefit across all outcomes, but the small size of individual trials with a highly heterogeneous patient population means that statistical significance is difficult to achieve.

Inconsistency in definition of outcome (variable length of follow-up, differing functional outcome measures, and poor definition of

'super-spell') result in several similar outcomes reported separately which could not be combined in a meta-analysis.

There are no studies in which orthogeriatrician input is confined to initial assessment and peri-operative medical care, (without then leading into orthogeriatric MDR). Therefore, the value of such early orthogeriatrician involvement can only be inferred from the outcome of HFP studies.

The quality of the studies ranges from low to high, with the majority of outcomes obtaining a moderate score.

Other considerations

The orthogeriatric assessment that would be provided to individual patients by a multidisciplinary HFP team will vary according to individual circumstances, and it was not felt appropriate to specify these in detail in this Guideline.

Assumptions – all papers included an orthogeriatrician, but the outcomes are most plausibly those of coordinated hospital-based multidisciplinary team working, with orthogeriatricians playing a medical and supervisory role within the team.

An important function of the HFP is to ensure the required liaison with, or cross-coverage of, the programmes in place for the secondary prevention of fracture by means of the assessment and treatment of osteoporosis and risk of falling (see NICE Clinical Guideline 21 & Technology Appraisal 161^{227,234,235}). In some centres HFP staff (including the orthogeriatrician) have common or parallel commitments within these programmes, with the resulting potential to achieve additional economies over and above those identified in the model.

The GDG highlighted this recommendation as a key priority for implementation.

1

Recommendation

If a hip fracture complicates or precipitates a terminal illness, the multidisciplinary team should still consider the role of surgery as part of a palliative care approach that:

- **minimises pain and other symptoms and**
- **establishes patients' own priorities for rehabilitation and**
- **considers patients' wishes about their end-of-life care.**

Relative values of different outcomes

Patients with advanced, life-threatening cardiorespiratory, neurological, and malignant disease make up a substantial proportion of those presenting with hip fracture.

In addition the trauma of suffering a hip fracture, and orthopaedic and medical complications of the injury, immobility and surgery can themselves precipitate a deterioration in the health of individuals.

In these circumstances such individuals and their families may view relief of pain, restoration of function and return home as a higher

	<p>priority than survival. Taking this into consideration the GDG prioritised pain, functional status and discharge destination as the most important outcomes.</p> <p>Sometimes this may make it necessary to move from an active surgical and rehabilitative approach to a palliative focus that ensures that the patient can die with dignity, with appropriate attention pain and other symptoms, and all the support necessary to minimise their and their family's distress.</p>
Trade off between clinical benefits and harms	<p>Pain, immobility, continence, pressure ulcer risk and dignity are all improved if the hip fracture can be addressed surgically, and perioperative risk should not preclude consideration of surgical management as an integral component of palliative care.</p> <p>The prognosis for an individual patient's recovery, mobility and return home can change markedly and multidisciplinary assessment is necessary if patients, their families and carers are given information with which to make informed decisions about their priorities for care (see chapter 13 Patient and carer views and information).</p> <p>High quality palliative and terminal care requires a multi-disciplinary approach, which should be provided as a key part of the support that the Hip Fracture Programme offers. Early orthogeriatric assessment and ongoing multidisciplinary working will help in:</p> <ul style="list-style-type: none"> • avoidance of complications such as pressure sores³⁴⁴ and delirium²⁰³ • expediting discharge.
Economic considerations	<p>No cost-effectiveness evidence was identified on this sub-group of patients. Additional time spent in counseling and supporting patients and their families will clearly carry a cost. While improvements in a patient's symptoms and quality of life may be of only short duration, sensitively handled palliative care can substantially improve their relatives' distress both before and for many years after bereavement.</p>
Quality of evidence	<p>There is no evidence directly relating to this very frail sub-group. Terminally ill patients were often excluded from these papers and if included were not reported in specific sub group analysis. This recommendation was based on GDG consensus opinion.</p>
Other considerations	<p>For patients whose hip fracture occurs in the context of advanced or terminal cancer-related illness, please see NICE Clinical Guideline "Improving supportive and palliative care for adults with cancer"²²⁷.</p>

Recommendation

Healthcare professionals should deliver care that minimises the patient's risk of delirium and maximises their independence, by:

- actively looking for cognitive impairment when patients first present with hip fracture
- reassessing patients to identify delirium that may arise during their admission
- offering individualised care in line with 'Delirium' (NICE clinical guideline 103)

Relative values of different outcomes

Patients with memory problems make up a substantial proportion of admissions, and face increased risk of delirium, medical complications, mortality, prolonged length of stay, and failure to return to pre-fracture independence.

The GDG considered medical complications, mortality, length of stay and discharge destination as the most important outcomes.

Trade off between clinical benefits and harms

Patients with memory problems are known to benefit from acute comprehensive geriatric assessment and targeted intervention as a means of reducing their risk of delirium and severe delirium, which are significant contributors to increased length of stay and increased risk of mortality at 6 months^{150,150}, as well as being a source of profound distress for patients, their families and carers^{203,203}.

In addition, intensive rehabilitation has been shown to be effective in improving outcome in terms of independent living among patients with mild to moderate cognitive impairment^{157,157}.

No evidence of harm was found and the GDG would not expect harm. Although no evidence met our inclusion criteria for this area, GDG consensus is that the potential benefits include avoidance of the distress that delirium causes to patients, their family, carers, and other inpatients, along with avoidance of the persistent reduction in cognitive function that can follow an episode of delirium, and of the increased length of stay and mortality associated with delirium.

The avoidance and management of delirium in patients with hip fracture is specifically addressed in the NICE Guideline on Delirium²²⁴.

Economic considerations

The decision model from the NICE guideline on Delirium (CG103) found that the tailored multi-component intervention package was cost-effective for hip fracture patients (£8,000 per QALY gained), as this care would lead to a reduced risk of long-term institutional care placement, lower incidence of other medical complications and lower length of hospital stay for these patients.

Quality of evidence	Patients with cognitive impairment are usually a group excluded from studies. Over 60% of the papers reviewed either excluded patients with cognitive impairment and/ or dementia, or made no specific comments relating to this subgroup. The studies that specifically analysed this subgroup ^{157,203} are of moderate quality.
Other considerations	<p>For patients whose hip fracture occurs in the context of dementia, please see the NICE guidance on dementia²²⁴.</p> <p>Identification of cognitive impairment is a key part of assessment, and a number of tools have been used in patients with hip fracture. The Abbreviated Mental Test (AMT) score is often used, and forms part of the National Hip Fracture Database's dataset, but the GDG did not examine the choice of tool or approach to assessment.</p> <p>Assessment of mental state can be complex in patients who are in pain, or who have received strong analgesia at the time of presentation. Approaches to the prevention and management of delirium require much more than screening for cognitive impairment at admission, and must include a sensitivity to changes in mental state and an awareness that delirium may arise at any stage of a patient's stay.</p> <p>Delirium is not confined to patients with pre-existing cognitive problems, and its incidence will be reduced most effectively by the provision of continuous orthogeriatric support to all patients²⁰³. Evidence on the effectiveness of models to prevent and manage delirium following hip fracture were key to the recommendations made in the NICE Guideline on Delirium²²⁴, and that Guideline should be read alongside our own when developing services for patients with hip fracture.</p>

1 **12.3 Research recommendations on hospital multidisciplinary**

2 **rehabilitation**

3 **12.3.1 Hip fracture unit**

4 The GDG recommended the following research question:

- 5 ➤ What is the clinical and cost effectiveness of a designated hip fracture unit within
6 the trauma ward compared to units integrated into acute trusts on mortality,
7 quality of life and functional status in patients with hip fracture?

8 **Why this is important**

9 The increasingly structured approach to hip fracture care has led to a number of UK units
10 considering or establishing a specific 'hip fracture ward' as a specialist part of their acute
11 orthopaedic service.

12
13 Designated hip fracture wards may prove an effective means of delivering the whole programme
14 of coordinated perioperative care and multidisciplinary rehabilitation which this NICE Guidance

1 has proposed, but at present there is no high quality evidence of their clinical effectiveness when
2 compared to such care within general orthopaedic or trauma beds.
3 It may not be practical to run an RCT within a trauma unit, but there is certainly potential for
4 cohort studies to explore the effect of such units on individual patients' mobility, discharge
5 residence, mortality and length of stay. Units considering the establishment of hip fracture wards
6 should be encouraged to consider performing such trials.
7
8

1 12.4 Community-based multidisciplinary rehabilitation versus usual care

2 In addition or as an alternative to hospital based multidisciplinary rehabilitation (MDR), a
3 number of studies have evaluated the role of community based MDR.

4 Community-based MDR includes approaches that are:

- 5 ➤ based in the patient's own home - Early Supported Discharge (ESD)
- 6 ➤ based within a residential care unit or community hospital
- 7 ➤ based within a Social Care Unit (SC) - or their near equivalents.

8 The many versions of these services across the country are named differently (for example
9 'intermediate care at home', 'intermediate care residential rehabilitation'), but each
10 consists of a rehabilitation component delivered in one of the above settings.

11 12.4.1 Review question

12 In patients with hip fracture what is the clinical and cost effectiveness of community-based
13 multidisciplinary rehabilitation on functional status, length of stay in secondary care,
14 mortality, place of residence/discharge, hospital readmission and quality of life?

15 Two studies met the inclusion criteria for this review question, with a total of 168 patients.
16 See evidence table 11, Appendix E and forest plots G139 to G145 Appendix G.

17 12.4.1.1 Clinical evidence

18 **Table 12-68: Home-based multidisciplinary early supported discharge vs. usual care – Clinical**
19 **study characteristics**

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
Mortality at 12 months ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Moved to a higher level of care ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Unable to walk ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
SF-36 scores at 12 months (0: worst to 100: best) - Physical component summary scores ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision

Outcome	Number of studies	Design	Limitations	Inconsistency	Indirectness	Other considerations/ imprecision
SF-36 scores at 12 months (0: worst to 100: best) - Mental component summary scores ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(c)
Length of hospital stay ^{59,360}	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Lengths of hospital or rehabilitation stays (days) - Length of rehabilitation (hospital + home) ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	no serious imprecision
Readmission to hospital during 4 months follow-up ⁵⁹	1	RCT	serious ^(a)	no serious inconsistency	no serious indirectness	serious ^(b)
Degree of independence (Functional Independent Measure) - FIM Self-care – 1 month ³⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Degree of independence (Functional Independent Measure) - FIM Mobility ³⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Degree of independence (Functional Independent Measure) - FIM Locomotion ³⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision
Mobility and strength tests - Up and go test ³⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	serious ^(b)
Mobility and strength tests - Sit-to-stand test ³⁶⁰	1	RCT	no serious limitations	no serious inconsistency	no serious indirectness	no serious imprecision

1 (a) Baseline data for Crotty et al., 2003⁵⁹ each study arm not given.

1
2
3
4
5
6
7

- (b) The relatively few events and few patients give wide confidence intervals around the estimate of effect. This makes it difficult to know the true effect size for this outcome
- (c) The wide confidence intervals around the measurement make the result imprecise. This makes it difficult to know the true effect size for this outcome.

Table 12-69: Home-based multidisciplinary early supported discharge vs. usual care - Clinical summary of findings

Outcome	Intervention	Control	Relative risk	Absolute effect	Quality
Mortality at 12 months	3/34 (8.8%)	4/32 (12.5%)	RR 0.71 (0.17, 2.91)	36 fewer per 1000 (from 104 fewer to 239 more)	Low
Moved to a higher level of care	1/34 (2.9%)	2/32 (6.3%)	RR 0.47 (0.04 to 4.94)	33 fewer per 1000 (from 60 fewer to 246 more)	Low
Unable to walk	0/34 (0%)	2/32 (6.3%)	RR 0.19 (0.01 to 3.78)	51 fewer per 1000 (from 62 fewer to 174 more)	Low
SF-36 scores at 12 months (0: worst to 100: best) - Physical component summary scores	34	32	N/A	MD 4.7 (0.04 to 9.44)	Moderate
SF-36 scores at 12 months (0: worst to 100: best) - Mental component summary scores	34	32	N/A	MD 1.5 (2.54 to 5.54)	Low
Length of hospital stay (days)	82	86	N/A	MD -2.96 (-5.50 to -0.42)	Moderate
Lengths of hospital or rehabilitation stays (days) - Length of rehabilitation (hospital + home)	34	32	N/A	MD 2.96 (5.5 to 0.42)	Moderate
Readmission to hospital during 4 months follow-up	8/34 (23.5%)	7/32 (21.9%)	RR 1.08 (0.44, 2.62)	18 more per 1000 (from 123 fewer to 354 more)	Low
Degree of independence (Functional Independent Measure) - FIM Self-care	48	54	N/A	MD 4.90 (2.81, 6.99)	High
Degree of independence (Functional Independent Measure) - FIM Mobility – 1 month	48	54	N/A	MD 2.00 (1.02, 2.98)	High

Degree of independence (Functional Independent Measure) - FIM Locomotion	48	54	N/A	MD 2.80 (1.61, 3.99)	High
Mobility and strength tests - Up and go test	48	54	N/A	MD 5.9 lower (12 lower to 0.2 higher)	Moderate
Mobility and strength tests - Sit-to-stand test	48	54	N/A	MD 1.5 lower (2.49 to 0.51 lower)	High

1

2 12.4.2 Economic evidence

3 Our search identified five studies on community MDR versus usual care. Of these, one^{55,55}
4 was excluded as it included a mixed population with only 31% hip fracture patients. Van
5 Balen et al., 2002^{340,340} was excluded as patients in the early supported discharge scheme
6 were only discharged to a nursing home with rehabilitation facilities and not to their own
7 home.

8 The following studies were included as economic evidence on the cost-effectiveness of
9 home-based multidisciplinary early supported discharge vs. usual care: Hollingworth
10 (1993)^{148,148} O’Cathain (1994)²⁴⁵ and Parker (1991)^{270,270}. Hollingworth (1993)^{148,148} is a cost
11 analysis based on a case series. O’Cathain (1994)²⁴⁵ is a cost-consequences analysis based
12 on a non-randomised trial with concurrent controls. Parker (1991)^{270,270} is a cost-
13 consequences analysis based on a prospective observational study. For further details on
14 these studies please refer to the Evidence Table 16 in Appendix F.

15 An original decision analysis has been conducted comparing the cost-effectiveness of the
16 community MDR vs. usual care. A decision tree model with Markov states was developed,
17 adopting a life-time horizon.

18 Treatment effects and EQ-5Ds scores were based on the findings of Crotty (2002)⁶⁰ and
19 applied only up to 4 months from follow-up. Resource use was determined from the NHS
20 and PSS perspective. Effectiveness was measured in QALYs. Costs and QALYs were
21 discounted at a rate of 3.5%. Please see section 8.7 in Appendix H for further detail.

22

1 **Table 12-70: Home-based multidisciplinary early supported discharge vs. usual care -**
 2 **Economic study characteristics**

Study	Limitations	Applicability	Other Comments
Hollingworth 1993 ¹⁴⁸	Potentially serious limitations (a)	Partial applicability	A community-based MDR at home scheme was compared to usual care. The MDR at home programme consisted of: care from trained nurses, nursing auxiliaries, physiotherapists, and occupational therapists in the patient's home for up to 24 hrs a day under the medical supervision of the general practitioner
O'Cathain 1994 ²⁴⁵	Potentially serious limitations (b)	Partial applicability	MDR at home compared to usual care. MDR team consisted of district nurses, physiotherapists, occupational therapists and generic workers, all working under the clinical responsibility of a GP for a maximum of 12 days.
Parker 1991 ²⁷⁰	Potentially serious limitations (c)	Partial applicability	MDR at home scheme compared to usual care. MDR team consisted of trained nurses, nursing auxiliaries, physiotherapists, and occupational therapists.
NGC economic model	Minor limitations ^(d)	Direct applicability	Cost-effectiveness analysis of community MDR – ESD versus usual care based on the RCT by Crotty et al (2002) ⁶⁰ included in the clinical review.

- 3 (a) Unclear follow up time. HRQoL not calculated. Information on costs obtained from hospital records,
 4 not national statistics. Not an RCT.
 5 (b) The length of time during which costs are calculated is unclear. No sensitivity analysis was
 6 conducted. Not based on a RCT. Not a CUA.
 7 (c) Not based on a RCT. No sensitivity analysis. Cost data from hospital source, not national statistics.
 8 Only patients admitted from their own home were then discharged under the HAH scheme.
 9 (d) The analysis consists of a decision tree with Markov states which spans a life-time horizon.
 10 Treatment effects based on the findings of the paper by Crotty in the clinical review and applied
 11 only up to 4 months from follow-up. Resource use determined from the NHS and PSS perspective,
 12 Effectiveness measured in QALYs. QALYs discounted at a rate of 3.5%.
 13
 14

1 **Table 12-71: Home-based multidisciplinary early supported discharge vs. usual care - Economic**
 2 **summary of findings**

Study	Incremental cost (£)	Incremental effects	ICER	Uncertainty
Hollingworth 1993	-£722	LOS; readmissions ^(l)	N/A	One way sensitivity analysis: costs of MDR scheme at home would still be lower than usual care if inpatients costs 50% lower and MDR at home costs 50% higher than predicted.
O’Cathain 1994	-£370	Several outcomes reported ^(m)	N/A	N/R
Parker 1991	-£799.80 ⁽ⁿ⁾	Several outcomes reported ^(o)	N/A	N/R
NCGC economic model	£434.6 ^(p)	0.0456 QALYs ^(q)	£9533/QALYs	95% CI: Community MDR dominant –usual care dominant ^(r)

3 *(l)* LOS for MDR at home vs. usual care: 32.5 vs. 41.7 days ($p < 0.001$); readmission rates at 1 year: 6.8%
 4 (53 patients) vs. 2.7% (8 patients), $p = 0.008$

5 *(m)* Several outcomes were reported: HRQoL measured with the Nottingham Health Profile
 6 questionnaire (14 vs. 24, $p < 0.05$); Mortality (5.3% vs. 5.9%; $p = \text{NR}$); readmission rates at 3 months:
 7 (15.8% vs. 8.8%, $p = 0.187$); LOS (median no of days): 10 vs. 17, $p < 0.001$

8 *(n)* Costs based on the following resource use: hospital length of stay; sessions with hospital
 9 occupational therapist; readmission days; MDR ESD staff time; other NHS or social services (GP
 10 visits, day care, meals on wheels, community services)

11 *(o)* LOS (mean, days): 29 vs. 38 (p value: 0.035). Mortality (at 90 days): 40 (14%) vs. 14 (11%)

12 *(p)* The mean costs associated with community MDR were estimated to be £6901.20 and for usual care
 13 £6466.60

14 *(q)* The mean effectiveness corresponded to 3.1283 QALYs and 3.0827 QALYs for usual care.

15 *(r)* Deterministic sensitivity analysis showed that findings were sensitive to the length of stay spent in
 16 hospital and during rehabilitation at home. Community MDR was found to be the most cost-
 17 effective option in 50% of the 10,000 simulations run in the PSA at a willingness to pay of £20k, and
 18 in 60% of the simulations at a willingness to pay of 30k per QALY.
 19
 20

21 12.4.3 Evidence statement (s)

Clinical There is a statistically significant and clinically significant reduction in hospital length of stay, but an increase in total length of rehabilitation (hospital + home) with home-based multidisciplinary early supported discharge (ESD) compared with usual care. (MODERATE QUALITY)

There is a statistically significant and clinically significant increase in functional independence measures with home-based multidisciplinary ESD compared with usual care. (HIGH QUALITY)

There is no statistically significant difference in mortality at 12 months and readmission to hospital at 4 months with home-based multidisciplinary ESD compared with usual care. (LOW QUALITY)

Economic Home-based MDR – ESD is cost-effective in the rehabilitation of patients with hip fracture. This evidence has minor limitations and direct applicability.

1 12.4.4 Recommendations and link to evidence

Recommendation	<p>Consider early supported discharge as part of the Hip Fracture Programme, provided the Hip Fracture Programme multidisciplinary team remains involved, and the patient:</p> <ul style="list-style-type: none"> • is medically stable and • has the mental ability to participate in continued rehabilitation and • is able to transfer and mobilise short distances and • has not yet achieved their full rehabilitation potential, as discussed with the patient, carer and family.
Relative values of different outcomes	<p>Length of hospital stay, functional outcomes and re-admission rates were considered the primary outcomes of interest. All these outcomes were used in the decision analytical model.</p>
Trade off between clinical benefits and harms	<p>Multidisciplinary ESD at home in selected patients reduces hospital length of stay but may result in overall prolonged rehabilitation (hospital + home) compared to hospital MDR. Selected patients were defined from the studies as medically stable, cognitively intact, able to transfer independently, and mobilise short distances.</p>
Economic considerations	<p>Despite only a few low quality studies being identified the GDG consensus was that multidisciplinary ESD at home is beneficial to a specific patient group, as defined above. The evidence reviewed showed an increase in functional independence measures with ESD compared to usual care.</p> <p>Our decision analysis found QALYs were 0.0456 higher in the community MDR arm of the study compared to usual care.</p> <p>No cost-effectiveness studies were identified for this clinical question. An original decision analytical model was developed, which was based on the findings of an RCT included in our clinical review^{58,60}. The analysis showed that there is uncertainty as to whether MDR ESD at home is cost-effective compared to usual care. In particular, findings were sensitive to the length of hospital stay and length of the home-based rehabilitation programme.</p> <p>However, the GDG noted that the ICER of £9533/QALYs is well below the £20,000 threshold.</p> <p>It is also important to note that our model did not find community MDR to be cost saving compared to usual care. This was because patients in the community MDR branch of the model underwent rehabilitation in their own home for a relatively longer period of time than those of the other studies included in the economic evidence profile in section 8.7 in Appendix H.</p>
Quality of evidence	<p>There were few studies identified, which ranged from low to high quality with often only one study per outcome. Therefore our</p>

confidence in the results is low.

Studies were undertaken in medically stable and cognitively intact patients and there were no studies that evaluated multidisciplinary ESD at home in cognitively impaired patients or patients living in care/nursing homes. This recommendation was therefore partly based on evidence and partly GDG consensus opinion.

Other considerations

Patient selection, as defined above is very important for multidisciplinary ESD at home and may represent a very small number of eligible patients.

The benefits of MDR ESD in patient with mild to moderate cognitive impairment living at home alone or with a relative /carer are unknown. MDR ESD in this context may be beneficial and should be considered.

The benefits of MDR ESD in patients living in care /nursing homes are unknown. MDR ESD in these patients, undertaken alongside the care/nursing homes may be beneficial.

Interaction with any key carer and evaluation of his/her ability and willingness to provide support and care is in all cases an essential and normative element of the decision making process in considering the appropriateness or otherwise of early supported discharge

The GDG highlighted this recommendation as a key priority for implementation.

1

Recommendation

Only consider intermediate care (continued rehabilitation in a community hospital or residential care unit) if all of the following criteria are met:

- **intermediate care is included in the Hip Fracture Programme and**
- **the Hip Fracture Programme team retains the clinical lead, including patient selection, agreement of length of stay and ongoing objectives for intermediate care and**
- **the Hip Fracture Programme team retains the managerial lead, ensuring that intermediate care is not resourced as a substitute for an effective acute hospital Programme.**

Relative values of different outcomes

The GDG considered the most important outcomes to be length of stay in hospital (in particular superspell) and return to pre fracture residence.

Trade off between clinical benefits and harms

There are risks that transfer to intermediate care may prematurely move a co-morbid patient group from a diagnostically supported environment, impair continuity, and prolong the superspell.

	In certain settings and specific circumstances, proximity to home with access for relatives/carers visiting and a more relaxed and “homely” atmosphere for continued rehabilitation than the acute hospital might be considered advantageous.
Economic considerations	The average weekly cost of the social care received in an intermediate care setting based in residential homes varies from a minimum of £412 to a maximum of £840 for schemes run by local authorities. The average weekly cost of social and health care services in the same setting but for schemes run by the local authority in conjunction with primary care trusts amounts to £574 (source: PSSRU 2009 ⁶¹). Subject to the criteria in the recommendation above, intermediate care may be feasible for our population, but there is currently no evidence on its cost-effectiveness.
Quality of evidence	There is no evidence on the effectiveness or cost-effectiveness of rehabilitation within a community hospital or residential care unit in hip fracture rehabilitation. This recommendation was based on GDG consensus opinion.
Other considerations	Intermediate care rehabilitation for hip fracture remains ill-defined and highly variable in the UK in terms of its admission criteria, multidisciplinary composition, intervention components and mechanisms for shared outcome and resource accountability within a comprehensive hip fracture programme.

1

Recommendation	Patients admitted from care or nursing homes should not be excluded from rehabilitation programmes in the community or hospital, or as part of an early supported discharge programme.
Relative values of different outcomes	<p>The GDG considered the most important outcomes to be functional status, readmission to hospital and return to pre-fracture residence.</p> <p>Early assessment and MDR offered as part of a hip fracture programme with continued rehabilitation for patients admitted from care/nursing homes is likely to improve/maintain the patient’s functional ability with regard to mobility, transfers from bed to chair and activities of daily living. This is in the interests of both patients and care/nursing home staff. In addition patient status as a care home resident as opposed to a nursing home resident may be maintained and equality for patients in care/nursing homes is maintained with regard to access to rehabilitation.</p>
Trade off between clinical benefits and harms	<p>There is no evidence of harm accruing to care/nursing home residents from the provision of appropriately individualised rehabilitation programmes.</p> <p>For some patients admitted from care/nursing homes there may be advantages (and no particular risks) in completing their rehabilitation after hospital MDR within that home (subject to the</p>

recommended criteria above), recognising that their rehabilitation goals may be more complex and must be shared by the HFP team on a continuing basis with the care/nursing home staff.

The potential benefits of ESD for patients admitted from care/nursing homes include the possibility of functional recovery within the patient's familiar environment, shared communication, goal setting and collaboration between care/nursing home staff and HFP team resulting in improved functional outcome, and the possibility of reduced hospital stay and inappropriate hospital readmission.

This subgroup is considered at particular risk of premature discharge because of ease of access to the care/nursing home environment and the corresponding perception that functional recovery matters less. Failure to undertake adequate rehabilitation carries the subsequent risk of inappropriate functional decline and/or levels of dependency, reduced quality of life, unnecessary hospital readmission, and premature mortality.

Provision of part of a patient's continuing rehabilitation programme in the care or nursing home of origin is correctly categorised as either early supported discharge or intermediate care, and the continued involvement of the Hip Fracture Programme team in liaison with the community-based component is therefore correspondingly a requirement.

Economic considerations

There was no cost-effectiveness evidence. The GDG believe that any increase in the cost of hospital bed days from the avoidance of premature discharge should be at least partially offset by the avoidance of inappropriate readmissions and reduction in subsequent care costs resulting from optimised functional status.

Quality of evidence

No RCTs were identified regarding patients admitted from care or nursing homes undergoing community ESD, as this patient subgroup has typically been excluded from clinical trials. The recommendation is based on GDG opinion and consensus that this group of patients would benefit from ESD.

Other considerations

There is a high prevalence of cognitive impairment in this population, therefore realistic rehabilitation goals need to be defined, but not at the expense of excluding rehabilitation.

1 **12.5 Research recommendations on community multidisciplinary** 2 **rehabilitation**

3 **12.5.1 Early supported discharge**

4 The GDG recommended the following research question:

- 5 ➤ What is the clinical and cost effectiveness of early supported discharge on
6 mortality, quality of life and functional status in patients with hip fracture who are
7 admitted from a care home?

8 **Why this is important**

9 Residents of care and nursing homes account for about 30% of all patients with hip fracture
10 admitted to hospital. Two-thirds of these come from care homes and the remainder from
11 nursing homes. These patients are frailer, more functionally dependent and have a higher
12 prevalence of cognitive impairment than patients admitted from their own homes. One-
13 third of those admitted from a care home are discharged to a nursing home and one-fifth
14 are readmitted to hospital within 3 months. There are no clinical trials to define the optimal
15 rehabilitation pathway following hip fracture for these patients and therefore represent a
16 discrete cohort where the existing meta-analyses do not apply. As a consequence, many
17 patients are denied structured rehabilitation and are discharged back to their care home or
18 nursing home with very little or no rehabilitation input.

19 Given the patient frailty and comorbidities, rehabilitation may have a limited effect on
20 clinical outcomes for this group. However, the fact that they already live in a home where
21 they are supported by trained care staff, clearly provides an opportunity for a systematic
22 approach to rehabilitation. Early care/nursing home based multidisciplinary rehabilitation
23 would take advantage of the day-to-day care arrangements already in place in homes and
24 provide additional NHS support to deliver naturalistic rehabilitation, where problems are
25 tackled in the setting in which the patient lives.

26 Early supported multidisciplinary rehabilitation could reduce hospital stay, improve early
27 return to function, and affect both readmission rates and the level of NHS-funded nursing
28 care required.

29 The research would follow a two-stage design: (1) An initial feasibility study to refine the
30 selection criteria and process for reliable identification and characterisation of those
31 considered most likely to benefit, together with the intervention package and measures for
32 collaboration between the HFP team, care-home staff and other community-based
33 professionals, and (2) A cluster randomized controlled comparison (with two or more
34 intervention units and matched control units) set against agreed outcome criteria. The
35 latter should include those specified above, together with measures of the impact on care-
36 home staff activity and cost, as well as qualitative data from patients on relevant quality-of-
37 life variables.

38 **12.5.2 Care/nursing home residents**

39 The GDG recommended the following research question:

- 1 ➤ Do patients admitted to hospital with a fractured hip who live permanently in a
2 care/nursing home have equal access to multidisciplinary rehabilitation as patients
3 admitted from their own homes?

4 **Why this is important**

5 The existing literature on the effectiveness of multidisciplinary rehabilitation typically
6 excludes patients who live in care/nursing homes. From an equality perspective it
7 hypothesised that this group of people do not have access to the same multidisciplinary
8 rehabilitation as patients who are returning home as it is assumed patients returning to
9 care/nursing homes will have their care needs met by the home. The research design would
10 be a prospective observational cohort study to determine the extent and quality of
11 rehabilitation services available to this group in comparison to patients returning to their
12 own homes.

13

1 13 Patient and carer views and information

2 13.1 Introduction

3 Patient views about their hip fracture and its management, and the way patients are
 4 provided with information are important elements of the natural recovery and treatment of
 5 hip fracture. Care givers also have need for information, and can influence the recovery
 6 process. Timely and clear information could reduce stress and uncertainty for patients and
 7 potentially improve their outcome. This section examines the literature on patient views
 8 and the provision of information to patients.

9 13.2 Patient and carer views

10 A systematic literature review was conducted into the views of patients and carers about
 11 their experience of hip fracture management from hospital admission until discharge from
 12 rehabilitation. Studies examining areas not covered by the guideline scope were not
 13 included. For example, hip protectors for falls management, nutrition support or patient
 14 views relating to the time after discharge from rehabilitation programmes.

15 The aim of this review was to provide:

- 16 • Supplementary evidence to clinical questions addressed in the guideline
- 17 • A general overview of patients views’ on hip fracture and hip fracture
 18 management
- 19 • Evidence relating to the provision of information to patients and carers

20 Eleven qualitative studies are included here, only two of which are UK based studies. More
 21 details about the studies are presented in the evidence table (Evidence table 12 in
 22 Appendix E). Studies were assessed using the NICE methodology checklist for qualitative
 23 studies²³³.

24 13.2.1 Summary of studies

25
 26 **Table 13-72 Patient views study quality**

Study	Population	Methods	Analysis	Relevance to guideline population
Archibald 2003 ⁸	Adequately reported	Adequately reported	Adequately reported, credible	Community hospital in UK 4 patients interviewed during rehabilitation

Study	Population	Methods	Analysis	Relevance to guideline population
Borkan 1991 & 1992 ^{28,29}	Adequately reported	Well reported	Well reported, credible	4 hospitals in USA 80 patients interviewed during hospital stay
Bowman 1997 ³³	Adequately reported	Poorly reported	Poorly reported, credible	Teaching hospital in Canada 17 patients interviewed on day of admission
Furstenberg 1986 ¹⁰⁵	Adequately reported	Poorly reported	Poorly reported, credible	Urban hospital in USA 11 patients interviewed at one or more points during hospital stay
Olsson 2007 ²⁴⁹	Well reported	Well reported	Well reported, credible	Geriatric/ orthopaedic ward in Sweden 13 patients interviewed soon after the operation
Pownall 2004 ²⁷⁴	Well reported	Poorly reported	Adequately reported, credible	Trauma/ orthopaedic ward in UK 1 patient interviewed prior to discharge from acute trauma and orthopaedic ward
Slauenwhite 1998 ³¹⁴	Poorly reported	Adequately reported	Poorly reported, credible	Hospital in Canada 23 'caregivers' for 23 patients interviewed 4 to 6 weeks after discharge
William 1994 ³⁵⁴	Poorly reported	Poorly reported	Poorly reported, credible	Hospital in USA 120 patients interviewed before hospital discharge and followed up at 2, 8 & 14 weeks
Wykes 2009 ³⁵⁵	Well reported	Well reported	Well reported, credible	Rehabilitation hospital in Australia 5 patients interviewed during rehabilitation
Young 2009 ³⁵⁸	Adequately reported	Well reported	Well reported, credible	Rehabilitation centre in USA 62 patients interviewed after 12 month follow up meeting
Ziden 2010 ³⁶²	Well reported	Well reported	Well reported, credible	Hospital in Sweden 18 patients interviewed at 1 month follow up meeting and 15 at 1 year follow up

1

2 **Archibald et al (2003)**⁸ conducted a qualitative study of 5 hip fracture patients in a
3 community hospital in the UK. Their aim was to explore experiences of individuals who had
4 suffered a hip fracture. Interviews with open ended questions were conducted during their
5 stay in hospital.

6 Four main themes were identified: injury experience, pain experience, recovery experience,
7 disability experience. Only the pain and recovery experience relate to their time in hospital
8 and rehabilitation. Most patients described the pain they experienced, one mentioned
9 being in a lot of pain in the orthopaedic unit despite pain killers. Another mentioned they
10 thought the pain went with rest after a while, but not completely. Only 1 person was still
11 having pain at time of interview. The recovery experience was split into 3 sequential
12 categories: the operation, beginning the struggle and regaining independence. Only 1

1 person described the operation, they had a “horrendous” recollection of a noisy operating
2 theatre, like being in an engineering shop or something”. Three patients remembered
3 ‘beginning the struggle’: they reported not being able to do anything; struggling to get to
4 the toilet and into a chair; and hating using a bed pan. The comments relating to regaining
5 independence were all positive. Motivation, be it getting to the toilet, the dining room or
6 smoke room was found to be a key factor in the recovery of the patients.

7 **Borkan et al (1991 & 1992)**^{28,29} conducted a qualitative study of 80 hip fracture patients in 4
8 hospitals in the USA. Their aim was to investigate the meanings of hip fracture to older
9 patients, and to identify potentially important prognostic indicators or risk factors for
10 rehabilitation outcomes. Patients were interviewed during the first week after hip fracture
11 with a combination of open-ended and multiple choice questions.

12 The study reports how patients perceive their fracture, their perception of their disability
13 and whether they were hopeful for the future (see evidence table). Also reported were
14 patient expectations of recovery (43 expected full recovery, 14 partial recovery and the rest
15 did not know or did not give an answer) and patient expectations about their living
16 situation (61% predicted going home, 15% into a nursing home though none came from
17 one, 9% predicted being discharged to their children’s houses and 15% did not know or did
18 not respond). The actual figures showed that 43% were discharged to long term care
19 institutions, of these 38% remained in the institution at 1 year, 53% returned home and 9%
20 died.

21 **Bowman (1997)**³³ conducted a quantitative study of 43 patients undergoing surgery on the
22 hip in a hospital in Canada, 17 of these had a hip fracture. The main aim was to describe
23 sleep satisfaction, pain perceptions and psychological concerns of patients undergoing hip
24 operations. Also two open ended questions were asked at the time of admission to
25 elucidate the patient’s biggest concerns about their injury and forthcoming surgery, and
26 whether they had concerns about their ability to recover fully and quickly. The mean age of
27 hip fracture patients was 80 years old and, unlike most the other studies, it also included
28 patients with delirium (8 out of 17). Six out of 17 patients feared being unable to walk
29 again, an additional 3 out of 17 were concerned about their recovery and managing on
30 their own.

31 **Fustenberg (1986)**¹⁰⁵ conducted a qualitative study of 11 patients of hospitalised patients
32 with hip fracture in a hospital in the USA. The aim of the study was to “construct a natural
33 history of the hip fracture”, from the events surrounding the hip fracture through the
34 hospitalisation period. Ethnographic interviews were carried out at one or more points
35 during their hospital stay.

36 The findings were split into two main sets: immediate patient expectations about their
37 recovery and “contextual factors” to the evolving expectations about their recovery. The
38 immediate expectations mostly included expressions of despair and discouragement: hip
39 fracture was going to result in extended period of slow recovery of function, with attendant
40 dependency, postponement or relinquishment of plans and changed living situation with
41 the threat of permanent loss of independent living. Participants also suffered uncertainty
42 about timing and completeness of return to full recovery

43 As time progressed participants commented that although progress was slow they could
44 see improvements. They also took encouragement from other people’s recovery. The study
45 notes that while patients could focus on positive and negative points, the participants only
46 focused on encouraging examples.

1 The study also reports that healthcare professionals' cues, encouragement and feedback
2 guided the participants' perceptions about their own progress. However, some participants
3 "referred to the elusiveness of the doctors and their own unanswered questions."

4 **Olsson et al (2007)**²⁴⁹ conducted a qualitative study of 13 hospitalised patients in Sweden.
5 The aim of the study was to describe patient's own perceptions of their situation and views
6 of their responsibility in the rehabilitation process. Interviews were conducted with semi-
7 structured questions as soon after the operation as the patients felt strong enough.

8 The study categorised the findings into different conceptions: 'autonomous' – responses
9 from people who appeared confident and accustomed to managing on their own; 'modest'
10 – responses from people who gave the impression of being vulnerable and dependent on
11 others, this group worried about their future more than the others; 'heedless' – responses
12 from people who appeared to have a sense of detachment. The heedless did not doubt
13 they would recover and that people around them would care for them. This group was
14 characterised predominantly by a reluctance to reflect on their own situation, by a refusal
15 to accept responsibility and by their need for information.

16 The study also identified some common traits: a lack of awareness - most patients lacked
17 awareness about their condition, what to do and how to act, and needed more
18 information; a shocking event - although several suspected they had a fracture all were
19 distressed by the diagnosis. The period before surgery was mostly blurred and filled with
20 fear and pain. The participants worried about how they would function postoperatively;
21 zest for life - all expressed a strong desire to recuperate although, while confined to bed
22 they worried about the pain, their inability to move their leg, their forthcoming operation
23 and the fear of being unable to walk again.

24 **Pownall 2004**²⁷⁴ conducted a critical appraisal of a 60 year old women's experience with hip
25 fracture in a UK hospital. The study was undertaken in an effort to understand further the
26 nature of personal experience. Narrative was acquired as part of a routine nursing
27 evaluation and helped to illuminate nursing care issues through the eyes of the patient. The
28 participant was interviewed prior to discharge with four open-ended questions.

29 The study identified three areas for improvement within the hospital: better
30 communication skills; time management for staff so time spent with the patient is used
31 effectively; and better pain management. The participant's comments included not
32 understanding why they had to wait so long in the Emergency department after the x-ray as
33 they had already been told their hip was fractured; staff were so busy, no one had time to
34 sit and explain things to her; concern that the operation was explained to her son but not
35 her; shock at being mobilised the day after surgery.

36 **Slauenwhite and Simpson (1998)**³¹⁴ conducted a qualitative study of 23 "caregivers" for 23
37 patients who had experienced hip fracture in Canada. The purpose of the study was to
38 investigate the impact of enhanced early discharge on families experiencing a repaired hip
39 fracture in an older adult. "Caregivers" were interviewed 4 to 6 weeks after discharge.

40 The length of stay was considered too long by the patient with the fracture and too short by
41 the carer for families. 15 out of the 23 families found length of stay not an issue. 20 of the
42 families stated pain management was not a problem in hospital or at home. Several
43 families thought the transition from hospital to home was a problem as it took several
44 hours to days for all the information to be relayed to home care system. This went hand in
45 hand for those with comorbidities. Many caregivers had stories of dissatisfaction which was
46 suggested to be related to health care system and mismatched care. Mismatched care was
47 not well defined.

1 **Williams et al (1994)**³⁵⁴ conducted a study into patient recovery and views for 120 patients
2 after hospital discharge in the USA. Participants were asked what advice they would offer
3 to other patients who had just fractured their hip. Patients were interviewed at 14 weeks
4 after discharge.

5 The advice offered was grouped into categories: 94 patients emphasised the importance of
6 mental attitude with comments such as patients should “maintain hope” and “look to the
7 future”; 76 patients suggested that following experts’ advice; 34 advised mobility was key
8 with comments such as keep mobile, rest before getting up to walk, use walker to help get
9 up; 15 advised maintain healthy lifestyle; 7 said use caution and be careful not to fall; 3
10 suggested limiting stay in institution and get help to be at home if possible; and 6 gave no
11 specific advice as they commented that everyone is different.

12 **Wykes et al (2009)**³⁵⁵ conducted a qualitative pilot study to explore the impact of hip
13 fracture on the lives of previously independent women and to identify their concerns when
14 participating in inpatient rehabilitation. Five patients were interviewed during their stay in a
15 rehabilitation hospital in Australia.

16 The impact of the fracture was an issue for all five women as others had to assume
17 responsibility for things they had done previously. The study categorised the women’s
18 concerns into four categories: the behaviour of others; what was happening to them; the
19 impact of their injury on others; and other health issues. A few comments were raised
20 about the behaviour of others including things others said and did, friends and family doing
21 things without asking first, the family not being told when one woman had moved hospital,
22 concern that staff expect one woman’s daughter to look after her until rehabilitation
23 started. Concerns about what was happening to them included a possible loss of
24 independence, possible accommodation changes after discharge and money issues. The
25 women were also concerned about inconveniencing or upsetting others by telling them
26 what they were feeling or asking too many questions. Two women had pre-existing health
27 issues which, combined with their hip fracture, had adverse effects on their outcome. These
28 overshadowed specific concerns about their hip fracture.

29 **Young and Resnick (2009)**³⁵⁸ conducted a qualitative study to explore the perceptions of 62
30 older adults regarding their functional recovery 1 year after hip fracture and after
31 participating in rehabilitation programme in the USA. Participants were asked whether they
32 were satisfied with their functional recovery, what helped or hindered recovery, what
33 would improve recovery and what one piece of advice they would offer other hip fracture
34 patients. The themes identified are listed below.

35 53 participants were satisfied with their functional recovery. The main factors they listed as
36 facilitators of recovery were seeing health care professionals and their positive attitude (40
37 respondents); social support, particularly from family and friends (13 respondents); and
38 their own determination (12 respondents). Other factors mentioned included lifestyle
39 factors or an environment that encourage healthy living, individualised care & verbal
40 encouragement; spirituality and identifying goals. The nine people who were dissatisfied
41 with their recovery listed medical complications or comorbidities, unpleasant sensations
42 and age as factors that hindered their recovery.

43 The respondents also identified areas that would facilitate recovery: more direct physical &
44 occupational therapy and more education about the recovery process and ways to optimise
45 physical function (26 respondents); better follow up and care in the home setting after
46 discharge from rehabilitation (9 respondents); spirituality (3 respondents), social support (2
47 respondents); additional information (8 respondents); elimination of unpleasant sensations
48 (4 respondents) and policy (1 respondent).

1 The patients also offered the following advice on how to facilitate recovery to anyone with
2 a hip fracture: listen to healthcare instructions (19 respondents) and participate as much as
3 possible in rehabilitation activities (48 respondents); participants strongly recommended
4 that older adults who sustain hip fractures maintain a positive attitude (20 respondents)
5 and remain determined throughout the recovery experience (13 respondents); be careful
6 to avoid subsequent trauma and prevent anything that would impede recovery (8
7 respondents); push through the pain and use all medication offered (6 respondents); and
8 don't worry (4 respondents).

9 **Ziden et al (2008 & 2010)**^{362,363} conducted a qualitative study to explore and describe the
10 consequences of an acute hip fracture among home dwelling elderly people shortly after
11 discharge from hospital in Sweden. Patients, who had participated in a randomised
12 controlled trial investigating rehabilitation³⁶⁰ included in the rehabilitation chapter (Section
13 12.2), attended semi-structured interviews at 1 month and 1 year after hip fracture.

14 The study identified different responses or perceptions over time. At 1 month patients:
15 found they were limited in movement and have lost confidence in their body (18 people);
16 had become humble and grateful (7 people); respected themselves and their own needs (2
17 people); had become more dependent on others (12 people); gain more human contact
18 and are treated in a friendly way by others (2 people); were secluded and trapped at home
19 (4 people); were old, closer to death and have lost your zest for life (4 people); were taking
20 one day at a time and were uncertain about the future (7 people). At 1 year after discharge
21 patients felt: more insecure and afraid (11 patients); they had more limited ability to move
22 (12 patients); disappointed and sad that identity and life have changed (8 patients);
23 satisfied with the situation or felt even better than before their fracture (5 patients).

24 The study also identified some patient views about determinants of hip fracture recovery:
25 10 patients stated their own mind and actions influenced recovery; 4 patients stated that
26 treatment and the actions from others influenced recovery; whereas 6 patients stated you
27 cannot influence recovery.

28

29 **13.2.2 Common themes**

30 The following themes have been identified from the studies:

31 **Initial outlook in hospital**

32 Five studies with 126 participants reported views from this period^{8,28,29,33,105,249}. One of the
33 studies reported the responses varied "from stubborn optimism to despair"^{28,29}. Another
34 study also reported all 13 participants expressed a strong desire to recuperate²⁴⁹. However,
35 most of the expressions were negative with no positive comments reported in the papers.

36 The concerns covered:

- 37 • pain and the inability to move their leg while confined to bed
- 38 • the fear of being unable to walk
- 39 • not being able to do anything
- 40 • hating using a bed pan
- 41 • struggling to get out of the chair or bed
- 42 • concern about recovery and managing on their own
- 43 • return to independent living
- 44 • limitations on their functioning and consequent implications
- 45 • being burden on their "caretakers" [families and carers]

- 1 • further falls
- 2 • uncertainty about timing and completeness of return to full recovery

3 **Attitude as patients began to regain independence**

4 Two studies reported comments relating to this period.

- 5 • Archibald (2003)⁸ with 5 participants reported motivation to be key factor in recovery.
6 All comments in the study were positive about regaining independence during their
7 rehabilitation.
- 8 • In Furstenberg (1986)¹⁰⁵ (11 participants) participants commented that although
9 progress was slow they could see improvement. Participants also took encouragement
10 from others' progress.

11 **Management by health care professionals**

12 Positive and negative comments were reported about healthcare professionals:

- 13 • Encouragement and positive attitude - Furstenberg (1986)¹⁰⁵ (11 participants) reported
14 that healthcare professionals' cues, encouragement and feedback guided the
15 informants' perceptions about their own progress. 40 out of the 62 participants in
16 Young (2009)³⁵⁸ identified that communication and a positive attitude by professionals
17 were seen as a facilitator of recovery.
- 18 • Provision of information to patients - Two studies also noted some negative points,
19 some patients "referred to the elusiveness of the doctors and their own unanswered
20 questions." in Furstenberg (1986)¹⁰⁵. The woman with a hip fracture in the individual
21 patient narrative²⁷⁴ was unhappy that things were not explained to her. One of her
22 comments highlighted this where she reported that the "staff were so busy no one has
23 time to sit and explain things to you".
- 24 • Explaining directly to patients - The patient from the individual narrative²⁷⁴ was also
25 unhappy that she could hear the nurse explaining the operation to her son, but
26 nothing was explained to her.

27

28 **13.2.3 Recommendations and link to evidence**

29 Overall, little evidence was identified that provided direct comments relating to our review
30 questions. Where applicable data were identified, reference to the evidence has been
31 made in the link to evidence of the relevant recommendations. These related to:

- 32 • Several comments were identified that fed into our recommendation relating to the
33 provision of information to patients (see next section 13.3).
- 34 • Some supplementary evidence was identified relating to pain that fed into our
35 analgesia recommendations (see section 7.2.2).

36

1 **13.3 Information for patients**

2 This section covers structured health education approaches, advice, information and
3 reassurance. In addition to qualitative literature the search conducted for patient views
4 included terms relating to patient education interventions. This also aimed to identify
5 randomised controlled trials investigating the effectiveness of different ways of providing
6 information to patients with hip fracture in improving outcomes.

7 **13.3.1 Evidence**

8 No randomised evidence was identified. However, good quality advice, reassurance,
9 information and education were highlighted by patients as important to the recovery
10 process in the qualitative review presented above.

11 The evidence above suggests that

- 12 • The positive attitude of and encouragement by health professionals is important
- 13 • Patients value time spent with them, and the advice and explanation given. This seems
14 important in the recovery process
- 15 • Patients should be treated with dignity, and provided with an explanation about their
16 condition and information about recovery.

17 Two studies asked participants to suggest what advice they would offer other hip fracture
18 patients based on their experiences. The main advice by participants in the studies to other
19 patients with hip fracture was:

- 20 • Maintain a positive attitude
- 21 • Follow experts advice and participate as much as possible in the rehabilitation process
- 22 • Keep mobile

23 **13.3.2 Recommendations and link to evidence**

24

<p>Recommendation</p>	<p>Offer patients (or, as appropriate, their carer and/or family) verbal and printed information about treatment and care including:</p> <ul style="list-style-type: none"> • diagnosis • choice of anaesthesia • choice of analgesia and other medications • surgical procedures • possible complications • postoperative care • rehabilitation programme • long-term outcomes • healthcare professionals involved.
<p>Relative values of different outcomes</p>	<p>Patient views on their satisfaction with the management of their condition were the main outcomes.</p>
<p>Trade off between clinical benefits and harms</p>	<p>The data highlighted examples where information was not provided to individual patients. Patients were unhappy when things were not explained to them. Patients were also unhappy when issues about their fracture were discussed with their family members instead of directly to them.</p> <p>The themes that came out of the evidence suggest that: a positive attitude of healthcare professionals is important; patients value time spent with them, and the advice and explanation given; and patients should be treated with dignity, and provided with an explanation about their condition and information about recovery.</p> <p>The GDG were unanimous in their view that discussion with patients (and where necessary their carers) about all aspects of the management of their hip fracture in is an important contributory factor in the recovery process.</p>
<p>Economic considerations</p>	<p>Although staff time is a scarce resource, information can be passed on to patients in the course of usual care and therefore needn't increased costs. Furthermore there may be benefits from greater adherence to treatment plans.</p>
<p>Quality of evidence</p>	<p>The qualitative evidence identified was of mixed quality. Data were not identified covering all the points mentioned above.</p>
<p>Other considerations</p>	<p>No comments were identified in the studies mentioning that adequate or good information was provided. However, the studies did not specifically ask about the quality of the information provided.</p>

1

2

1 **13.4 Carer involvement**

2 In patients who have been discharged after hip fracture repair, what is the effectiveness of
3 having a non paid carer (e.g. spouse, relative or friends) on mortality, length of stay, place
4 of residence/discharge, functional status, hospital readmission and quality of life?

5
6 No published evidence was identified. The GDG recognised the often crucial and sometimes
7 major contribution made by involved relatives and other non-professional carers to
8 successful rehabilitation. Early discussion with carers of prognosis and discharge planning
9 avoids misunderstanding of rehabilitation objectives, enables those involved to prepare in
10 an informed and timely manner for a patient's return home, consequently averts
11 inappropriate delay in discharge, and may reduce both length of stay and the likelihood of
12 inappropriate readmission to hospital.

13
14 There is the potential for the delay of some decisions with this approach and it remains
15 incumbent on clinicians with the agreement of patients (and/or any nominated proxy) to
16 ensure that their best interests are correctly identified and not compromised, particularly
17 (but not exclusively) in any urgent decision-making situation.

18 **13.4.1 Clinical evidence**

19 No relevant studies were identified.

20 **13.4.2 Economic evidence**

21 No relevant studies were identified.
22

23 **13.5 Research recommendations**

24 **13.5.1 Quality of life**

25 The GDG recommended the following research question:

- 26 ➤ What quality of life value do individual patients and their carers place on different
27 mobility, independence and residence states following rehabilitation?

28 **Why this is important**

29 It is important in evaluating future priorities for intervention to determine whether the
30 perceived clinical and health economic benefits of rehabilitation outcomes in the research
31 literature are matched over the same time-frame by the quality of life judgements,
32 aspirations and expectations of patients themselves and their carers. There is currently no
33 evidence.

34 **13.5.2 Patient experience**

35 The GDG recommended the following research question:

- 36 ➤ What is the patient's experience of being admitted to hospital with a hip fracture in
37 relation to surgery, pain management, timeliness of information given, and
38 rehabilitation?

1

2 **Why this is important**

3 No studies from NHS populations were identified where patients commented specifically on
4 their surgery, their pain management and rehabilitation programme. There were comments
5 in the patient views studies about not being kept informed about the management of their
6 condition, however, there was no information identified about the appropriate time to be
7 told. It may be that different patients want the information at different times. The studies
8 suggest that patients suffer from fear, pain and delirium until after surgery and it is
9 important to learn what (if anything) can be done to alleviate this which for many will be
10 considered the worst stage in their treatment.

11

12

1 Glossary

Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.
Algorithm (in guidelines)	A flow chart of the clinical decision pathway described in the guideline, where decision points are represented with boxes, linked with arrows.
Allocation concealment	The process used to prevent advance knowledge of group assignment in a RCT. The allocation process should be impervious to any influence by the individual making the allocation, by being administered by someone who is not responsible for recruiting participants.
AO classification	Classification system used to describe stable trochanteric fractures (type A1), unstable trochanteric (type A2), and transtrochanteric which includes those fracture lines at the level of the lesser trochanter and reversed fracture lines (type A3) ²¹⁹ .
Applicability	The degree to which the results of an observation, study or review are likely to hold true in a particular clinical practice setting.
Arm (of a clinical study)	Sub-section of individuals within a study who receive one particular intervention, for example placebo arm.
Association	Statistical relationship between two or more events, characteristics or other variables. The relationship may or may not be causal.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable), with which subsequent results are compared.
Before-and-after study	A study that investigates the effects of an intervention by measuring particular characteristics of a population both before and after taking the intervention, and assessing any change that occurs.
Bias	Systematic (as opposed to random) deviation of the results of a study from the 'true' results that is caused by the way the study is designed or conducted.
Blinding	Keeping the study participants, caregivers, researchers and outcome assessors unaware about the interventions to which the participants have been allocated in a study.

Carer (caregiver)	Someone other than a health professional who is involved in caring for a person with a medical condition.
Case-control study	Comparative observational study in which the investigator selects individuals who have experienced an event (For example, developed a disease) and others who have not (controls), and then collects data to determine previous exposure to a possible cause.
Case-series	Report of a number of cases of a given disease, usually covering the course of the disease and the response to treatment. There is no comparison (control) group of patients.
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinical effectiveness	The extent to which an intervention produces an overall health benefit in routine clinical practice.
Clinician	A healthcare professional providing direct patient care, for example doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence-based medicine databases including the Cochrane Database of Systematic Reviews (reviews of randomised controlled trials prepared by the Cochrane Collaboration).
Cohort study	A retrospective or prospective follow-up study. Groups of individuals to be followed up are defined on the basis of presence or absence of exposure to a suspected risk factor or intervention. A cohort study can be comparative, in which case two or more groups are selected on the basis of differences in their exposure to the agent of interest.
Comorbidity	Co-existence of more than one disease or an additional disease (other than that being studied or treated) in an individual.
Community hospital	A local hospital, unit or centre providing an appropriate range and format of accessible health care facilities and resources. These are typically small, and provide non-emergency services.
Comparability	Similarity of the groups in characteristics likely to affect the study results (such as health status or age).
Concordance	This is a recent term whose meaning has changed. It was initially applied to the consultation process in which doctor and patient agree therapeutic decisions that incorporate their respective views, but now includes patient support in medicine taking as well as prescribing communication. Concordance reflects social values but does not address medicine-taking and may not lead to improved adherence.
Confidence interval (CI)	A range of values for an unknown population parameter with a stated 'confidence' (conventionally 95%) that it contains the true value. The interval is calculated from sample data, and generally straddles the sample estimate. The 'confidence' value means that if the method used

to calculate the interval is repeated many times, then that proportion of intervals will actually contain the true value.

Confounding	In a study, confounding occurs when the effect of an intervention on an outcome is distorted as a result of an association between the population or intervention or outcome and another factor (the 'confounding variable') that can influence the outcome independently of the intervention under study.
Consensus methods	Techniques that aim to reach an agreement on a particular issue. Consensus methods may be used when there is a lack of strong evidence on a particular topic.
Control group	A group of patients recruited into a study that receives no treatment, a treatment of known effect, or a placebo (dummy treatment) - in order to provide a comparison for a group receiving an experimental treatment, such as a new drug.
Cost benefit analysis	A type of economic evaluation where both costs and benefits of healthcare treatment are measured in the same monetary units. If benefits exceed costs, the evaluation would recommend providing the treatment.
Cost-consequences analysis (CCA)	A type of economic evaluation where various health outcomes are reported in addition to cost for each intervention, but there is no overall measure of health gain.
Cost-effectiveness analysis (CEA)	An economic study design in which consequences of different interventions are measured using a single outcome, usually in 'natural' units (For example, life-years gained, deaths avoided, heart attacks avoided, cases detected). Alternative interventions are then compared in terms of cost per unit of effectiveness.
Cost-effectiveness model	An explicit mathematical framework, which is used to represent clinical decision problems and incorporate evidence from a variety of sources in order to estimate the costs and health outcomes.
Cost-utility analysis (CUA)	A form of cost-effectiveness analysis in which the units of effectiveness are quality-adjusted life-years (QALYs).
Credible Interval	The Bayesian equivalent of a confidence interval.
Lag screw cut-out	A complication in which the implant may protrude into the surrounding tissue or penetrate into the acetabulum. Symptoms include increasing pain and impaired mobility; and treatment depends on the severity of the symptoms as well as the fitness of the patient to undergo what may be major revision surgery. It may take the form of re-fixation of the fracture, replacement arthroplasty, or simple removal of the implant.

Decision analysis	An explicit quantitative approach to decision making under uncertainty, based on evidence from research. This evidence is translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Dominance	An intervention is said to be dominated if there is an alternative intervention that is both less costly and more effective.
Drop-out	A participant who withdraws from a trial before the end.
Early Supported Discharge (ESD)	Patients are discharged home from the acute trauma ward, or in some cases a subsequent rehabilitation ward within the hospital, with a supported 4-6 week rehabilitation package.
Economic evaluation	Comparative analysis of alternative health strategies (interventions or programmes) in terms of both their costs and consequences.
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	The observed association between interventions and outcomes or a statistic to summarise the strength of the observed association.
Effectiveness	See 'Clinical effectiveness'.
Efficacy	See 'Clinical efficacy'.
Epidemiological study	The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (For example, infection, diet) and interventions.
EQ-5D (EuroQol-5D)	A standardise instrument used to measure a health outcome. It provides a single index value for health status.
Evidence	Information on which a decision or guidance is based. Evidence is obtained from a range of sources including randomised controlled trials, observational studies, expert opinion (of clinical professionals and/or patients).
Exclusion criteria (literature review)	Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.
Exclusion criteria (clinical study)	Criteria that define who is not eligible to participate in a clinical study.

Extended dominance	If Option A is both more clinically effective than Option B and has a lower cost per unit of effect, when both are compared with a do-nothing alternative then Option A is said to have extended dominance over Option B. Option A is therefore more efficient and should be preferred, other things remaining equal.
Extramedullary implant	Implants used to fix extracapsular fractures. Examples of extramedullary implants include the sliding hip screw and the Medoff plate.
Extrapolation	In data analysis, predicting the value of a parameter outside the range of observed values.
Follow-up	Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been assessed in order to observe changes in health status or health-related variables.
Generalisability	The extent to which the results of a study based on measurement in a particular patient population and/or a specific context hold true for another population and/or in a different context. In this instance, this is the degree to which the guideline recommendation is applicable across both geographical and contextual settings. For instance, guidelines that suggest substituting one form of labour for another should acknowledge that these costs might vary across the country.
Gold standard	See 'Reference standard'.
Geriatric Orthopaedic Rehabilitation Unit (GORU)	A separate geriatrician-led trauma ward. The extent of surgical input to the GORU varies, depending on how early patients are moved from the acute trauma wards.
GRADE / GRADE profile	A system developed by the GRADE Working Group to address the shortcomings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE profile.
Harms	Adverse effects of an intervention.
Health economics	The study of the allocation of scarce resources among alternative healthcare treatments. Health economists are concerned with both increasing the average level of health in the population and improving the distribution of health.
Health-related quality of life (HRQoL)	A combination of an individual's physical, mental and social well-being; not merely the absence of disease.
Heterogeneity	Or lack of homogeneity. The term is used in meta-analyses and systematic reviews when the results or estimates of effects of

treatment from separate studies seem to be very different – in terms of the size of treatment effects or even to the extent that some indicate beneficial and others suggest adverse treatment effects. Such results may occur as a result of differences between studies in terms of the patient populations, outcome measures, definition of variables or duration of follow-up.

Hip fracture programme (HFP)

Formal 'orthogeriatric' care - with the geriatric medical team contributing to joint preoperative patient assessment, and increasingly taking the lead in postoperative medical care, MDR and discharge planning.

Imprecision

Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.

Inclusion criteria (literature review)

Explicit criteria used to decide which studies should be considered as potential sources of evidence.

Incremental analysis

The analysis of additional costs and additional clinical outcomes with different interventions.

Incremental cost

The mean cost per patient associated with an intervention minus the mean cost per patient associated with a comparator intervention.

Incremental cost effectiveness ratio (ICER)

The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another.

$$ICER = \frac{Cost_A - Cost_B}{Effectiveness_A - Effectiveness_B}$$

Incremental net benefit (INB)

The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the INB is calculated as: (£20,000 x QALYs gained) – Incremental cost.

Indirectness

The available evidence is different to the review question being addressed, in terms of PICO (population, intervention, comparison and outcome).

Intention to treat analysis (ITT)

A strategy for analysing data from a randomised controlled trial. All participants are included in the arm to which they were allocated, whether or not they received (or completed) the intervention given to that arm. Intention-to-treat analysis prevents bias caused by the loss of participants, which may disrupt the baseline equivalence established by randomisation and which may reflect non-adherence to the protocol.

Intermediate care

Care provided in community hospitals or residential care units as an intermediate step between hospital care and care in a person's own home

Intervention	Healthcare action intended to benefit the patient, for example, drug treatment, surgical procedure, psychological therapy.
Intraoperative	The period of time during a surgical procedure.
Intramedullary implant	Implants used to fix extracapsular fractures. Examples of intramedullary implants are the Gamma nail, the intramedullary hip screw and the proximal femoral nail.
Kappa statistic	A statistical measure of inter-rater agreement that takes into account the agreement occurring by chance.
Length of stay	The total number of days a participant stays in hospital.
Licence	See 'Product licence'.
Life-years gained	Mean average years of life gained per person as a result of the intervention compared with an alternative intervention.
Likelihood ratio	The likelihood ratio combines information about the sensitivity and specificity. It tells you how much a positive or negative result changes the likelihood that a patient would have the disease. The likelihood ratio of a positive test result (LR+) is sensitivity divided by 1- specificity.
Long-term care	Care in a home that may include skilled nursing care and help with everyday activities. This includes nursing homes and care homes.
Loss to follow-up	Also known as attrition. The loss of participants during the course of a study. Participants that are lost during the study are often call dropouts.
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
Meta-analysis	A statistical technique for combining (pooling) the results of a number of studies that address the same question and report on the same outcomes to produce a summary result. The aim is to derive more precise and clear information from a large data pool. It is generally more reliably likely to confirm or refute a hypothesis than the individual trials.
Mixed Assessment and Rehabilitation Unit (MARU)	A rehabilitation unit able to accept patients with a variety of medical, surgical and orthopaedic conditions.
Mobilisation	Mobilisation is the process of re-establishing the ability to move between postures (for example sit to stand), maintain an upright posture, and to ambulate with increasing levels of complexity (speed, changes of direction, dual and multi-tasking).

Multidisciplinary rehabilitation (MDR)	Rehabilitation after hip fracture incorporating the following core components of assessment and management: medicine; nursing; physiotherapy; occupational therapy; social care. Additional components may include: dietetics, pharmacy, clinical psychology.
Multivariate model	A statistical model for analysis of the relationship between two or more predictor (independent) variables and the outcome (dependent) variable.
Negative predictive value (NPV)	<p>[In screening/diagnostic tests:] A measure of the usefulness of a screening/diagnostic test. It is the proportion of those with a negative test result who do not have the disease, and can be interpreted as the probability that a negative test result is correct. It is calculated as follows:</p> $NPV = \frac{(\text{specificity})(1 - \text{prevalence})}{(\text{specificity})(1 - \text{prevalence}) + (1 - \text{sensitivity})(\text{prevalence})}$
Non-union	The terms non-union, pseudarthrosis or delayed union are used for those fractures that fail to heal after a few months.
Number needed to treat (NNT)	The number of patients that who on average must be treated to prevent a single occurrence of the outcome of interest.
Observational study	Retrospective or prospective study in which the investigator observes the natural course of events with or without control groups; for example, cohort studies and case-control studies.
Odds ratio	A measure of treatment effectiveness. The odds of an event happening in the treatment group, expressed as a proportion of the odds of it happening in the control group. The 'odds' is the ratio of events to non-events.
Opportunity cost	The loss of other health care programmes displaced by investment in or introduction of another intervention. This may be best measured by the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.
Orthogeriatrician	A care of the elderly physician with an interest in fracture care.
Outcome	Measure of the possible results that may stem from exposure to a preventive or therapeutic intervention. Outcome measures may be intermediate endpoints or they can be final endpoints. See 'Intermediate outcome'.

P-value	The probability that an observed difference could have occurred by chance, assuming that there is in fact no underlying difference between the means of the observations. If the probability is less than 1 in 20, the P value is less than 0.05; a result with a P value of less than 0.05 is conventionally considered to be 'statistically significant'.
Perioperative	The period from admission through surgery until discharge, encompassing the preoperative and postoperative periods.
Placebo	An inactive and physically identical medication or procedure used as a comparator in controlled clinical trials.
Polypharmacy	The use or prescription of multiple medications.
Positive predictive value (PPV)	In screening/diagnostic tests: A measure of the usefulness of a screening/diagnostic test. It is the proportion of those with a positive test result who have the disease, and can be interpreted as the probability that a positive test result is correct. It is calculated as follows: $PPV = \frac{(\text{sensitivity})(\text{prevalence})}{(\text{sensitivity})(\text{prevalence}) + (1 - \text{specificity})(1 - \text{prevalence})}$
Postoperative	Pertaining to the period after patients leave the operating theatre, following surgery.
Post-test probability	For diagnostic tests. The proportion of patients with that particular test result who have the target disorder (post test odds/[1 + post-test odds]).
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Preoperative	The period before surgery commences.
Pre-test probability	For diagnostic tests. The proportion of people with the target disorder in the population at risk at a specific time point or time interval. Prevalence may depend on how a disorder is diagnosed.
Primary care	Healthcare delivered to patients outside hospitals. Primary care covers a range of services provided by general practitioners, nurses, dentists, pharmacists, opticians and other healthcare professionals.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Product licence	An authorisation from the MHRA to market a medicinal product.
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good prognosis is associated with low rate of undesirable outcomes; poor

prognosis is associated with a high rate of undesirable outcomes.

Prospective study	A study in which people are entered into the research and then followed up over a period of time with future events recorded as they happen. This contrasts with studies that are <i>retrospective</i> .
Publication bias	Also known as reporting bias. A bias caused by only a subset of all the relevant data being available. The publication of research can depend on the nature and direction of the study results. Studies in which an intervention is not found to be effective are sometimes not published. Because of this, systematic reviews that fail to include unpublished studies may overestimate the true effect of an intervention. In addition, a published report might present a biased set of results (e.g. only outcomes or sub-groups where a statistically significant difference was found).
Quality of life	See 'Health-related quality of life'.
Quality-adjusted life year (QALY)	An index of survival that is adjusted to account for the patient's quality of life during this time. QALYs have the advantage of incorporating changes in both quantity (longevity/mortality) and quality (morbidity, psychological, functional, social and other factors) of life. Used to measure benefits in cost-utility analysis. The QALYs gained are the mean QALYs associated with one treatment minus the mean QALYs associated with an alternative treatment.
Quick Reference Guide	An abridged version of NICE guidance, which presents the key priorities for implementation and summarises the recommendations for the core clinical audience.
Randomisation	Allocation of participants in a research study to two or more alternative groups using a chance procedure, such as computer-generated random numbers. This approach is used in an attempt to ensure there is an even distribution of participants with different characteristics between groups and thus reduce sources of bias.
Randomised controlled trial (RCT)	A comparative study in which participants are randomly allocated to intervention and control groups and followed up to examine differences in outcomes between the groups.
Residential care unit	A unit or centre where care is given outside of the patient's home. Care can be 24 hour care or partial care depending on the person's needs.
RCT	See 'Randomised controlled trial'.
Receiver operated characteristic (ROC) curve	A graphical method of assessing the accuracy of a diagnostic test. Sensitivity is plotted against 1-specificity. A perfect test will have a positive, vertical linear slope starting at the origin. A good test will be somewhere close to this ideal.
Reference standard	The test that is considered to be the best available method to establish the presence or absence of the outcome – this may not be the one that

is routinely used in practice.

Relative risk (RR)	The number of times more likely or less likely an event is to happen in one group compared with another (calculated as the risk of the event in group A/the risk of the event in group B).
Reporting bias	See publication bias.
Resource implication	The likely impact in terms of finance, workforce or other NHS resources.
Retrospective study	A retrospective study deals with the present/ past and does not involve studying future events. This contrasts with studies that are <i>prospective</i> .
Review question	In guideline development, this term refers to the questions about treatment and care that are formulated to guide the development of evidence-based recommendations.
Secondary outcome	An outcome used to evaluate additional effects of the intervention deemed a priori as being less important than the primary outcomes.
Selection bias	A systematic bias in selecting participants for study groups, so that the groups have differences in prognosis and/or therapeutic sensitivities at baseline. Randomisation (with concealed allocation) of patients protects against this bias.
Sensitivity	<p>Sensitivity or recall rate is the proportion of true positives which are correctly identified as such. For example in diagnostic testing it is the proportion of true cases that the test detects.</p> <p>See the related term 'Specificity'</p>
Sensitivity analysis	<p>A means of representing uncertainty in the results of economic evaluations. Uncertainty may arise from missing data, imprecise estimates or methodological controversy. Sensitivity analysis also allows for exploring the generalisability of results to other settings. The analysis is repeated using different assumptions to examine the effect on the results.</p> <p>One-way simple sensitivity analysis (univariate analysis): each parameter is varied individually in order to isolate the consequences of each parameter on the results of the study.</p> <p>Multi-way simple sensitivity analysis (scenario analysis): two or more parameters are varied at the same time and the overall effect on the results is evaluated.</p> <p>Threshold sensitivity analysis: the critical value of parameters above or below which the conclusions of the study will change are identified.</p> <p>Probabilistic sensitivity analysis: probability distributions are assigned to the uncertain parameters and are incorporated into evaluation models based on decision analytical techniques (For example, Monte Carlo simulation).</p>

Significance (statistical)	A result is deemed statistically significant if the probability of the result occurring by chance is less than 1 in 20 ($p < 0.05$).
Specificity	<p>The proportion of true negatives that are correctly identified as such. For example in diagnostic testing the specificity is the proportion of non-cases incorrectly diagnosed as cases.</p> <p>See related term 'Sensitivity'.</p> <p>In terms of literature searching a highly specific search is generally narrow and aimed at picking up the key papers in a field and avoiding a wide range of papers.</p>
Stakeholder	Those with an interest in the use of the guideline. Stakeholders include manufacturers, sponsors, healthcare professionals, and patient and carer groups.
Subtrochanteric extracapsular fracture	Subtrochanteric fractures are those in which the fracture is predominantly in the 5cms of bone immediately distal to the lesser trochanter.
Superspell	Total time in NHS care.
Systematic review	Research that summarises the evidence on a clearly formulated question according to a pre-defined protocol using systematic and explicit methods to identify, select and appraise relevant studies, and to extract, collate and report their findings. It may or may not use statistical meta-analysis.
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Treatment allocation	Assigning a participant to a particular arm of the trial.
Trochanteric extracapsular fracture	Extracapsular fractures occur outside or distal to the hip joint capsule and include basal, trochanteric and subtrochanteric fractures. Trochanteric fractures may be further subdivided into two part fractures, which are also termed stable fractures, and those that are comminuted or multi-fragmentary, which may be termed unstable fractures.
Univariate	Analysis which separately explores each variable in a data set.

1

2

3

4 **The management of hip fracture in**

5 **adults**

6

7 **Appendices A – J**

8

9

	APPENDICES
--	-------------------

12

13 Produced by the National Clinical Guideline Centre

1 Appendices

2	CONTENTS.....	1
3	GUIDELINE DEVELOPMENT GROUP MEMBERS	1
4	NCGC STAFF MEMBERS OF THE GUIDELINE DEVELOPMENT GROUP	2
5	EXPERT ADVISORS	2
6	ACKNOWLEDGEMENTS	3
7	ACRONYMS AND ABBREVIATIONS.....	4
8	1 INTRODUCTION.....	7
9	2 DEVELOPMENT OF THE GUIDELINE	10
10	2.1 WHAT IS A NICE CLINICAL GUIDELINE?	10
11	2.2 REMIT	11
12	2.3 WHO DEVELOPED THIS GUIDELINE?	11
13	2.4 WHAT THIS GUIDELINE COVERS.....	12
14	2.5 WHAT THIS GUIDELINE DOES NOT COVER.....	13
15	2.6 RELATIONSHIPS BETWEEN THE GUIDELINE AND OTHER NICE GUIDANCE	13
16	3 METHODS	15
17	3.1 DEVELOPING THE REVIEW QUESTIONS AND OUTCOMES	15
18	3.2 SEARCHING FOR EVIDENCE.....	20
19	3.2.1 <i>Clinical literature search</i>	20
20	3.2.2 <i>Health economic literature search</i>	21
21	3.3 EVIDENCE OF EFFECTIVENESS.....	21
22	3.3.1 <i>Inclusion/exclusion</i>	22
23	3.3.2 <i>Methods of combining clinical studies</i>	22
24	3.3.3 <i>Appraising the quality of evidence by outcomes</i>	23
25	3.3.4 <i>Grading the quality of clinical evidence</i>	24
26	3.3.5 <i>Study limitations</i>	25
27	3.3.6 <i>Inconsistency</i>	25
28	3.3.7 <i>Indirectness</i>	26
29	3.3.8 <i>Imprecision</i>	26
30	3.4 EVIDENCE OF COST-EFFECTIVENESS.....	27
31	3.4.1 <i>Literature review</i>	27
32	3.4.2 <i>NICE economic evidence profiles</i>	28
33	3.4.3 <i>Undertaking new health economic analysis</i>	29
34	3.4.4 <i>Cost-effectiveness criteria</i>	30
35	3.5 DEVELOPING RECOMMENDATIONS	30
36	3.5.1 <i>Research recommendations</i>	31
37	3.6 VALIDATION PROCESS	31
38	3.7 UPDATING THE GUIDELINE	31
39	3.8 DISCLAIMER.....	31

1	3.9 FUNDING	31
2	4 GUIDELINE SUMMARY	32
3	4.1 MAP OF RECOMMENDATIONS	32
4	AN ALGORITHM WILL BE ADDED BEFORE PUBLICATION. KEY PRIORITIES FOR IMPLEMENTATION	32
5	KEY PRIORITIES FOR IMPLEMENTATION	33
6	4.2 FULL LIST OF RECOMMENDATIONS.....	35
7	4.2.1 <i>Imaging options in occult hip fracture</i>	35
8	4.2.2 <i>Timing of surgery</i>	35
9	4.2.3 <i>Analgesia</i>	35
10	4.2.4 <i>Anaesthesia</i>	36
11	4.2.5 <i>Planning the theatre team</i>	36
12	4.2.6 <i>Surgical procedures</i>	36
13	4.2.7 <i>Mobilisation strategies</i>	37
14	4.2.8 <i>Multidisciplinary management</i>	37
15	4.2.9 <i>Patient and carer information</i>	39
16	4.3 RESEARCH RECOMMENDATIONS	39
17	4.3.1 <i>Research recommendation on imaging options in occult hip fracture</i>	39
18	4.3.2 <i>Research recommendation on anaesthesia</i>	40
19	4.3.3 <i>Research recommendation on displaced intracapsular hip fracture</i>	40
20	4.3.4 <i>Research recommendation on early supported discharge</i>	41
21	4.3.5 <i>Research recommendation on physiotherapy</i>	41
22	4.3.6 <i>Additional research recommendations</i>	42
23	5 IMAGING OPTIONS IN OCCULT HIP FRACTURE.....	46
24	5.1 INTRODUCTION	46
25	5.2 REVIEW QUESTION.....	47
26	5.3 RADIOGRAPHS	47
27	5.3.1 <i>What is the diagnostic accuracy of additional radiographs (X-Rays) after 48 hours compared</i>	
28	<i>to MRI in the diagnosis of occult hip fractures</i>	47
29	5.4 RADIONUCLIDE BONE SCAN (RNS).....	48
30	5.4.1 <i>What is the diagnostic accuracy of RNS compared to MRI in the diagnosis of occult hip</i>	
31	<i>fractures</i>	48
32	5.4.2 <i>Economic evidence</i>	48
33	5.4.3 <i>Recommendations and link to evidence</i>	49
34	5.5 ULTRASOUND (US).....	49
35	5.5.1 <i>Diagnostic accuracy of ultrasound (US) compared to MRI in the diagnosis of occult hip</i>	
36	<i>fractures</i>	49
37	5.5.2 <i>Recommendations and link to evidence</i>	50
38	5.6 COMPUTED TOMOGRAPHY (CT)	50
39	5.6.2 <i>Recommendations and link to evidence</i>	51
40	5.7 RESEARCH RECOMMENDATION ON IMAGING OPTIONS IN OCCULT HIP FRACTURE	54
41	6 TIMING OF SURGERY	55
42	6.1 INTRODUCTION	55
43	6.1.1 <i>Review question</i>	56
44	6.1.2 <i>Recommendations and link to evidence</i>	65
45	6.2 RESEARCH RECOMMENDATIONS ON TIMING OF SURGERY	69
46	6.2.1 <i>Surgery within 36 hours</i>	69
47	7 ANALGESIA	70
48	7.1 INTRODUCTION	70
49	7.2 SYSTEMIC ANALGESIA	71
50	7.2.1 <i>Review question</i>	71
51	7.2.2 <i>Recommendations and link to evidence</i>	71
52	7.3 NERVE BLOCKS COMPARED TO SYSTEMIC ANALGESIA.....	71
53	7.3.1 <i>Review question</i>	71

1	7.3.2	<i>Recommendations and link to evidence</i>	75
2	7.4	RESEARCH RECOMMENDATIONS ON ANALGESIA	82
3	8	REGIONAL (SPINAL OR EPIDURAL) VERSUS GENERAL ANAESTHESIA	83
4	8.1	INTRODUCTION	83
5	8.2	REGIONAL VERSUS GENERAL ANAESTHESIA	83
6	8.2.1	<i>Review question</i>	83
7	8.2.2	<i>Recommendations and link to evidence</i>	86
8	8.3	RESEARCH RECOMMENDATION ON ANAESTHESIA	88
9	9	SURGEON SENIORITY	90
10	9.1	INTRODUCTION	90
11	9.2	SURGEON SENIORITY	90
12	9.2.1	<i>Review question</i>	90
13	9.2.2	<i>Clinical evidence</i>	90
14	9.3	RECOMMENDATIONS AND LINK TO EVIDENCE	92
15	10	SURGICAL PROCEDURES	96
16	10.1	INTRODUCTION	96
17	10.2	SURGERY WITH REGARD TO EARLY MOBILISATION	96
18	10.2.1	<i>Recommendations and link to evidence</i>	97
19	10.3	DISPLACED INTRACAPSULAR FRACTURES	97
20	10.3.1	<i>Internal fixation versus hemiarthroplasty</i>	98
21	10.3.2	<i>Internal fixation versus total hip replacement</i>	102
22	10.3.3	<i>Hemiarthroplasty versus total hip replacement</i>	105
23	10.3.4	<i>Recommendations and link to evidence</i>	109
24	10.3.5	<i>Research recommendations on displaced intracapsular fractures</i>	113
25	10.4	USE OF CEMENT IN ARTHROPLASTY.....	114
26	10.4.1	<i>Use of cement in original Thompson and Austin Moore designs of arthroplasty</i>	114
27	10.4.2	<i>Use of cement in newer designs of arthroplasty</i>	118
28	10.4.3	<i>Recommendations and link to evidence</i>	121
29	10.5	SURGICAL APPROACH TO HEMIARTHROPLASTY	122
30	10.5.2	<i>Recommendations and link to evidence</i>	124
31	10.6	EXTRACAPSULAR FRACTURE FIXATION.....	125
32	10.6.1	<i>Intramedullary versus extramedullary implants for fixation of trochanteric extracapsular fractures</i> 126	
33	10.6.2	<i>Intramedullary versus extramedullary implants for fixation of reverse oblique trochanteric extracapsular fractures</i>	130
34	10.6.3	<i>Intramedullary versus extramedullary implants for fixation of subtrochanteric extracapsular fractures</i>	131
35			
36			
37			
38	11	MOBILISATION STRATEGIES	135
39	11.1	INTRODUCTION	135
40	11.2	EARLY VS. DELAYED MOBILISATION	136
41	11.2.1	<i>Review question</i>	136
42	11.2.2	<i>Recommendations and link to evidence</i>	138
43	11.3	INTENSITY OF PHYSIOTHERAPY	139
44	11.3.1	<i>Review question</i>	139
45	11.3.2	<i>Recommendations and link to evidence</i>	143
46	11.3.3	<i>Research recommendations on mobilisation</i>	145
47	12	MULTIDISCIPLINARY MANAGEMENT.....	146
48	12.1	INTRODUCTION	146
49	12.2	HOSPITAL-BASED MULTIDISCIPLINARY REHABILITATION VERSUS USUAL CARE	148
50	12.2.1	<i>Review questions</i>	148
51	12.2.2	<i>Evidence statement (s)</i>	158
52	12.2.3	<i>Recommendations and link to evidence</i>	159

1	12.3	RESEARCH RECOMMENDATIONS ON HOSPITAL MULTIDISCIPLINARY REHABILITATION	165
2	12.3.1	<i>Hip fracture unit</i>	165
3	12.4	COMMUNITY-BASED MULTIDISCIPLINARY REHABILITATION VERSUS USUAL CARE	167
4	12.4.1	<i>Review question</i>	167
5	12.4.2	<i>Economic evidence</i>	170
6	12.4.3	<i>Evidence statement (s)</i>	172
7	12.4.4	<i>Recommendations and link to evidence</i>	173
8	12.5	RESEARCH RECOMMENDATIONS ON COMMUNITY MULTIDISCIPLINARY REHABILITATION	177
9	12.5.1	<i>Early supported discharge</i>	177
10	12.5.2	<i>Care/nursing home residents</i>	177
11	13	PATIENT AND CARER VIEWS AND INFORMATION.....	179
12	13.1	INTRODUCTION	179
13	13.2	PATIENT AND CARER VIEWS.....	179
14	13.2.1	<i>Summary of studies</i>	179
15	13.2.2	<i>Common themes</i>	184
16	13.2.3	<i>Recommendations and link to evidence</i>	185
17	13.3	INFORMATION FOR PATIENTS	186
18	13.3.1	<i>Evidence</i>	186
19	13.3.2	<i>Recommendations and link to evidence</i>	186
20	13.4	CARER INVOLVMENT.....	188
21	13.4.1	<i>Clinical evidence</i>	188
22	13.4.2	<i>Economic evidence</i>	188
23	13.5	RESEARCH RECOMMENDATIONS	188
24	13.5.1	<i>Quality of life</i>	188
25	13.5.2	<i>Patient experience</i>	188
26		GLOSSARY	190
27		APPENDICES.....	203
28		APPENDIX A: SCOPE	210
29	13.6	GUIDELINE TITLE.....	210
30	13.6.1	<i>Short title</i>	210
31	13.7	THE REMIT	210
32	13.8	CLINICAL NEED FOR THE GUIDELINE.....	210
33	13.8.1	<i>Epidemiology</i>	210
34	13.8.2	<i>Current practice</i>	211
35	13.9	THE GUIDELINE.....	213
36	13.9.1	<i>Population</i>	213
37	13.9.2	<i>Healthcare setting</i>	214
38	13.9.3	<i>Clinical management</i>	214
39	13.9.4	<i>Main outcomes</i>	215
40	13.9.5	<i>Economic aspects</i>	216
41	13.9.6	<i>Status</i>	216
42	13.10	RELATED NICE GUIDANCE	216
43	13.10.1	<i>Published</i>	216
44	13.10.2	<i>Guidance under development</i>	217
45	13.11	FURTHER INFORMATION.....	217
46	14	APPENDIX B: DECLARATIONS OF INTEREST	219
47	14.1	INTRODUCTION	219
48	14.2	DECLARATIONS OF INTERESTS OF THE GDG MEMBERS.....	220
49	14.2.1	<i>Professor Cameron Swift</i>	220
50	14.2.2	<i>Professor Opinder Sahota</i>	221
51	14.2.3	<i>Dr Antony Johansen</i>	222
52	14.2.4	<i>Mr Tim Chesser</i>	223
53	14.2.5	<i>Mr Bob Handley</i>	224

1	14.2.6	<i>Ms Karen Hertz</i>	225
2	14.2.7	<i>Dr Richard Griffiths</i>	226
3	14.2.8	<i>Professor Sallie Lamb</i>	227
4	14.2.9	<i>Mrs Heather Towndrow</i>	228
5	14.2.10	<i>Dr Sally Hope</i>	229
6	14.2.11	<i>Ms Tessa Somerville</i>	230
7	14.2.12	<i>Mr Anthony Field</i>	231
8	14.2.13	<i>Mr Martin Wise</i>	232
9	14.3	DECLARATIONS OF INTERESTS OF THE NCGC MEMBERS	233
10	14.4	DECLARATIONS OF INTERESTS OF THE EXPERT ADVISORS	234
11	14.4.1	<i>Mr Martin Parker</i>	234
12	14.4.2	<i>Mrs Pamela Holmes</i>	234
13	14.4.3	<i>Professor Judith Adams</i>	234
14	15	APPENDIX C: REVIEW PROTOCOLS.....	235
15	15.1	REVIEW PROTOCOL – IMAGING IN OCCULT HIP FRACTURE.....	235
16	15.2	REVIEW PROTOCOL – TIMING OF SURGERY.....	237
17	15.3	REVIEW PROTOCOL – ANALGESIA- SYSTEMIC MEDICATIONS	239
18	15.4	REVIEW PROTOCOL – ANALGESIA- NERVE BLOCKS COMAPARED TO SYSTEMIC ANALGESICS	241
19	15.5	REVIEW PROTOCOL - ANAESTHESIA	243
20	15.6	REVIEW PROTOCOL – SURGEON SENIORITY	245
21	15.7	REVIEW PROTOCOL – CEMENT.....	246
22	15.8	REVIEW PROTOCOL – INTRACAPSULAR FRACTURES.....	248
23	15.9	REVIEW PROTOCOL – SURGICAL APPROACH	250
24	15.10	REVIEW PROTOCOL – HEMIARTHROPLASTY STEM DESIGN.....	252
25	15.11	REVIEW PROTOCOL – EXTRACAPSULAR FRACTURES	254
26	15.12	REVIEW PROTOCOL – MOBILISATION STRATEGIES.....	256
27	15.13	REVIEW PROTOCOL – MULTIDISCIPLINARY REHABILITATION	258
28	15.14	REVIEW PROTOCOL – CARER INVOLVEMENT	264
29	15.15	REVIEW PROTOCOL – HEALTH ECONOMICS	266
30	16	APPENDIX D: LITERATURE SEARCH STRATEGIES.....	268
31	16.1	SEARCH STRATEGIES	268
32	16.2	SEARCH TERMS.....	270
33		<i>Anaesthesia</i>	270
34		<i>Analgesia</i>	271
35		<i>Animal/publication filter</i>	272
36		<i>Carer involvement</i>	272
37		<i>Diagnostic filter</i>	273
38		<i>Early Surgery</i>	274
39		<i>Economic</i>	274
40		<i>Hip Fracture Terms</i>	276
41		<i>Orthogeriatrician</i>	277
42		<i>Patient education</i>	278
43		<i>Patient views</i>	279
44		<i>Radiological Imaging</i>	280
45		<i>RCT filter</i>	281
46		<i>Rehabilitation</i>	282
47		<i>Surgeon seniority</i>	288
48		<i>Surgical Interventions</i>	288
49		<i>Systematic review filter</i>	289
50	17	APPENDIX E: EVIDENCE TABLES - CLINICAL STUDIES	291
51	17.1	EVIDENCE TABLE 1: IMAGING OPTIONS IN OCCULT HIP FRACTURE.....	293
52	17.2	EVIDENCE TABLE 2: TIMING OF SURGERY	296
53	17.3	EVIDENCE TABLE 3: OPTIMAL ANALGESIA	314
54	17.4	EVIDENCE TABLE 4: ANAESTHESIA	317

1	17.5	EVIDENCE TABLE 5: SURGEON SENIORITY.....	320
2	17.6	EVIDENCE TABLE 6: DISPLACED INTRACAPSULAR FRACTURES.....	324
3	17.7	EVIDENCE TABLE 7: SURGERY – CEMENT VERSUS NO CEMENT.....	333
4	17.8	EVIDENCE TABLE 8: EXTRACAPSULAR FIXATION.....	335
5	17.9	EVIDENCE TABLE 9: SURGICAL APPROACH TO HEMIARTHROPLASTY.....	383
6	17.10	EVIDENCE TABLE 10: MOBILISATION STRATEGIES.....	385
7	17.11	EVIDENCE TABLE 11: MULTIDISCIPLINARY REHABILITATION.....	394
8	17.12	EVIDENCE TABLE 12: PATIENT VIEWS.....	418
9	18	APPENDIX F: EVIDENCE TABLES - ECONOMIC STUDIES.....	434
10	18.1	EVIDENCE TABLE 13: GENERAL VERSUS REGIONAL ANAESTHESIA.....	435
11	18.2	EVIDENCE TABLE 14: DISPLACED INTRACAPSULAR FRACTURES.....	437
12	18.3	EVIDENCE TABLE 15: CEMENTED ARTHROPLASTIES.....	441
13	18.4	EVIDENCE TABLE 16: MULTIDISCIPLINARY REHABILITATION.....	443
14	19	APPENDIX G: FOREST PLOTS.....	456
15	19.1	RADIOLOGY.....	459
16	19.2	TIMING OF SURGERY.....	460
17	19.3	ANALGESIA.....	466
18	19.4	ANAESTHESIA.....	473
19	19.5	SURGICAL INTERVENTIONS.....	478
20	19.5.1	<i>Surgeon seniority.....</i>	<i>478</i>
21	19.5.2	<i>Cement in older designs of arthroplasty.....</i>	<i>479</i>
22	19.5.3	<i>Cement in newer designs of arthroplasty.....</i>	<i>487</i>
23	19.5.4	<i>Internal fixation versus hemiarthroplasty.....</i>	<i>489</i>
24	19.5.5	<i>Internal fixation versus total hip replacement.....</i>	<i>494</i>
25	19.5.6	<i>Hemiarthroplasty versus total hip replacement.....</i>	<i>497</i>
26	19.5.7	<i>Trochanteric extracapsular fracture – all studies.....</i>	<i>502</i>
27	19.5.8	<i>Trochanteric extracapsular fracture – studies from 2000.....</i>	<i>510</i>
28	19.5.9	<i>Subtrochanteric extracapsular fracture.....</i>	<i>515</i>
29	19.6	MOBILISATION STRATEGIES.....	517
30	19.6.1	<i>Timing of mobilisation.....</i>	<i>517</i>
31	19.7	INTENSIVE EXERCISE OR PHYSIOTHERAPY VS. USUAL CARE.....	519
32	19.7.1	<i>Intensive physiotherapy (Strength training).....</i>	<i>519</i>
33	19.7.2	<i>Intensive physiotherapy (treadmill training).....</i>	<i>521</i>
34	19.7.3	<i>Intensive (more frequent) physiotherapy.....</i>	<i>523</i>
35	19.8	MULTIDISCIPLINARY REHABILITATION.....	524
36	19.8.1	<i>Hospital-based MDR.....</i>	<i>524</i>
37	19.9	HOME-BASED MDR VERSUS USUAL INPATIENT REHABILITATION.....	529
38	20	APPENDIX H: HEALTH ECONOMIC ANALYSIS.....	532
39	20.1	COST ANALYSIS OF NERVE BLOCKS, NON-OPIOID ANALGESICS AND OPIOID ANALGESICS.....	532
40	20.1.1	<i>Nerve block cost analysis.....</i>	<i>532</i>
41	20.1.2	<i>Non-opioid analgesics.....</i>	<i>534</i>
42	20.1.3	<i>Opioid analgesics.....</i>	<i>534</i>
43	20.2	HOURLY WAGE COSTS FOR A PLANNED TRAUMA LIST.....	535
44	20.3	PRICES FOR SLIDING HIP SCREWS AND SHORT AND LONG INTRAMEDULLARY NAILS.....	538
45	20.4	COST ANALYSIS OF THE INTERVENTIONS FOR INTENSIVE MOBILISATION STRATEGIES.....	539
46	20.5	COST-EFFECTIVENESS ANALYSIS OF HOSPITAL INVESTMENT VERSUS NO HOSPITAL INVESTMENT FOR EARLY SURGERY	
47		542	
48	20.5.1	<i>Introduction.....</i>	<i>542</i>
49	20.5.2	<i>Background.....</i>	<i>542</i>
50	20.5.3	<i>Population and time horizon.....</i>	<i>543</i>
51	20.5.4	<i>Software.....</i>	<i>543</i>
52	20.5.5	<i>Methods.....</i>	<i>543</i>
53	20.5.6	<i>Treatment effects.....</i>	<i>544</i>
54	20.5.7	<i>Quality of life.....</i>	<i>546</i>

1	20.5.8	<i>Cost analysis</i>	547
2	20.5.9	<i>Cost-effectiveness analysis</i>	550
3	20.5.10	<i>Discussion</i>	557
4	20.6	COST-EFFECTIVENESS ANALYSIS OF HOSPITAL MDR VS USUAL CARE	559
5	20.6.1	<i>Introduction</i>	559
6	20.6.2	<i>Population and time horizon</i>	560
7	20.6.3	<i>Software</i>	560
8	20.6.4	<i>Structure of the model</i>	560
9	20.6.5	<i>Evidence and treatment effects on mortality</i>	566
10	20.6.6	<i>Utilities data</i>	567
11	20.6.7	<i>Calculating QALYs gained</i>	570
12	20.6.8	<i>Cost data</i>	570
13	20.6.9	<i>Cost-effectiveness findings for base-case analysis</i>	581
14	20.6.10	<i>Discussion</i>	592
15	20.7	COST-EFFECTIVENESS ANALYSIS OF COMMUNITY MDR VS USUAL CARE	593
16	20.7.1	<i>Introduction</i>	593
17	20.7.2	<i>Population and time horizon</i>	593
18	20.7.3	<i>Software</i>	593
19	20.7.4	<i>Economic evaluation type</i>	593
20	20.7.5	<i>Time horizon, Perspective, Discount rates used</i>	594
21	20.7.6	<i>Structure of the model</i>	594
22	20.7.7	<i>Utility data</i>	594
23	20.7.8	<i>Mortality</i>	595
24	20.7.9	<i>Calculating QALYs gained</i>	595
25	20.7.10	<i>Cost analysis</i>	596
26	20.7.11	<i>Cost effectiveness findings</i>	598
27	20.7.12	<i>Discussion</i>	604
28	20.8	COST ANALYSIS OF CEMENTED VS. UNCEMENTED IMPLANTS (NEWER DESIGNS OF ARTHROPLASTY)	604
29	a)	<i>Cost of implants</i>	604
30	c)	<i>Cost of length of stay in hospital</i>	606
31	d)	<i>Re-operation costs</i>	607
32	e)	<i>Theatre time costs</i>	608
33		<i>Summary of costs components</i>	608
34	21	APPENDIX I: HIGH PRIORITY RESEARCH RECOMMENDATIONS	610
35	21.1	IMAGING OPTIONS IN OCCULT HIP FRACTURE	610
36	21.2	ANAESTHESIA	613
37	21.3	DISPLACED INTRACAPSULAR HIP FRACTURES	616
38	21.4	INTENSIVE REHABILITATION THERAPIES AFTER HIP FRACTURE	620
39	21.5	EARLY SUPPORTED DISCHARGE IN CARE HOME PATIENTS	623
40	22	APPENDIX J: EXCLUDED STUDIES	627
41		BIBLIOGRAPHY	636
42			

1

2

Appendix A: Scope

3

13.6 Guideline title

4 Hip fracture: the management of hip fracture in adults

5

13.6.1 Short title

6 Hip fracture

7

13.7 The remit

8 The Department of Health has asked NICE: "to prepare a clinical guideline on the
9 management of fractured neck of femur".

10

13.8 Clinical need for the guideline

11

13.8.1 Epidemiology

12

13 a) About 70–75,000 hip fractures (proximal femoral fractures) occur annually in
14 the UK. Hip fracture is the commonest reason for admission to an orthopaedic
15 ward, and is usually a 'fragility' fracture² caused by a fall affecting an older
16 person with osteoporosis or osteopaenia (a lesser degree of bone reduction
17 and weakness due to the same process as in osteoporosis). The average age of
a person with hip fracture is 77 years. The annual cost of medical and social

² The strict definition of a fragility fracture is one caused by a fall from standing height or less. For the purposes of this guidance, the definition will be slightly more flexible to encompass all hip fractures judged to have an osteoporotic or osteopaenic basis

- 1 care for all the hip fracture cases in the UK amounts to about £2 billion.
2 Demographic projections indicate that the UK annual incidence will rise to
3 91,500 by 2015 and 101,000 in 2020, with an associated increase in annual
4 expenditure that could reach £2.2 billion by 2020. The majority of this
5 expenditure will be accounted for by hospital bed days and a further
6 substantial contribution will come from health and social aftercare. About a
7 quarter of patients with hip fracture are admitted from institutional care.
8 About 10–20% of those admitted from home ultimately move to institutional
9 care.
- 10 b) Mortality is high – about 10% of people with a hip fracture die within
11 1 month, and about one third within 12 months. However, fewer than half of
12 deaths are attributable to the fracture. This reflects the high prevalence of
13 comorbidity in people with hip fractures; often the combination of fall and
14 fracture brings to light underlying ill health. This presents major challenges for
15 anaesthetic, surgical, postoperative and rehabilitative care.
- 16 **13.8.2 Current practice**
- 17 a) The primary and secondary prevention of fragility fractures by treating
18 osteoporosis and reducing the risk of falls are of key importance to the
19 current and future epidemiology of hip fracture. These are, or will be, covered
20 by related NICE guidance (see section 5).
- 21 b) The diagnosis and management of hip fracture itself and of any comorbidity
22 before, during and after surgery, have a profound effect on outcome, both for
23 individuals and for services.
- 24 c) Patients with hip fracture need immediate referral to hospital (other than in
25 exceptional circumstances). Their assessment and management on admission
26 commonly involve a range of specialties and disciplines, but it is not always
27 clear how and when this involvement should take place. Prompt surgery is
28 important but is sometimes delayed for administrative or clinical reasons. It is

- 1 essential that mobilisation and rehabilitation after surgery are undertaken
2 according to individual need, but this does not always happen.
- 3 d) In spite of a significant body of evidence, hip fracture management and the
4 resulting length of hospital stay vary markedly among centres across England
5 and Wales.
- 6 e) Existing UK guidance from other sources includes:
- 7 • Scottish Intercollegiate Guidelines Network (2002) Prevention and
8 management of hip fracture in older people. Available from
9 www.sign.ac.uk/guidelines/fulltext/56/index.html
 - 10 • British Orthopaedic Association (2007) The care of patients with fragility
11 fracture. Available from www.nhfd.co.uk
 - 12 • Department of Health (2001) National service framework for older people³.
13 Available from www.dh.gov.uk
- 14 f) This clinical guideline will provide guidance on the emergency, preoperative,
15 operative and postoperative management of hip fracture, including
16 rehabilitation, in adults. It will not cover those aspects of hip fracture
17 addressed by related NICE guidance, but will refer to them as appropriate.
- 18 g) At all stages of hip fracture management, and especially during rehabilitation,
19 the importance of optimal communication with, and support for, patients
20 themselves and those who provide or will provide care – including unpaid
21 care family members or others – will be a fundamental tenet of guidance
22 development.

³ Elaborates on relevant (but not specific) standards of contextual importance (intermediate care, general hospital care and falls).

1 **13.9 The guideline**

2 The guideline development process is described in detail on the NICE website (see section
3 6, 'Further information').

4 This scope defines what the guideline will (and will not) examine, and what the guideline
5 developers will consider. The scope is based on the referral from the Department of
6 Health.

7 The areas that will be addressed by the guideline are described in the following sections.

8 **13.9.1 Population**

9 ***13.9.1.1 Groups that will be covered***

10 a) Adults aged 18 years and older presenting to the health service with a clinical
11 diagnosis (firm or provisional) of fragility fracture of the hip.

12 b) People with the following types of hip fracture:⁴

- 13 • intracapsular (undisplaced and displaced)
- 14 • extracapsular (trochanteric and subtrochanteric).

15 c) Those with comorbidity strongly predictive of outcome, and those without
16 such comorbidity. The influence (if any) of advanced age or gender on clinical
17 decision-making, management and outcome will be specifically evaluated.

18 ***13.9.1.2 Groups that will not be covered***

19 People younger than 18 years.

20 People with fractures caused by specific pathologies other than osteoporosis or
21 osteopaenia (because these would require more condition-specific guidance).

⁴ These terms explain where the bone has fractured, which can be either near or within the hip joint.

1 **13.9.2 Healthcare setting**

2 a) Secondary care settings where preoperative, operative, and postoperative
3 acute and subacute care are undertaken.

4 b) Primary, secondary and social care settings, as well as an individual's own
5 home, where rehabilitation is undertaken.

6 **13.9.3 Clinical management**

7 ***13.9.3.1 Key clinical issues that will be covered***

8 a) Using alternative radiological imaging to confirm or exclude a suspected hip
9 fracture in patients with a normal X-ray.

10 b) Involving a physician or orthogeriatrician in the care of patients presenting
11 with hip fracture.

12 c) Early surgery (within 48 hours).

13 d) Optimal preoperative and postoperative analgesia (pain relief), including the
14 use of nerve blockade.

15 e) Regional (spinal – also known as 'epidural') versus general anaesthesia in
16 patients undergoing surgery for hip fracture.

17 f) Does surgeon experience reduce the incidence of mortality, the need for
18 repeat surgery, and poor outcome in terms of mobility?

19 g) For displaced intracapsular fracture:

- 20 • internal fixation versus arthroplasty (hip replacement surgery)
21 • total hip replacement versus hemiarthroplasty (replacing the head of the
22 femur only) .

23 h) Choice of surgical implants - Sliding hip screw versus intramedullary nail for
24 trochanteric extracapsular fracture.

- 1 i) Choice of surgical implants - Sliding hip screw versus intramedullary nail for
2 subtrochanteric extracapsular fracture.
- 3 j) Cemented versus non-cemented arthroplasty implants.
- 4 k) Hospital-based multidisciplinary rehabilitation for patients who have
5 undergone hip fracture surgery.
- 6 l) Early transfer to community-based multidisciplinary rehabilitation for patients
7 who have undergone hip fracture surgery.

8 **13.9.3.2 Clinical issues that will not be covered**

9 The following will not be directly covered in this guideline, but related NICE guidance will
10 be referred to if appropriate:

- 11 a) Primary and secondary prevention of fragility fracture.
- 12 b) Prevention and management of pressure sores.
- 13 c) Prophylaxis for venous thromboembolism.
- 14 d) Prevention and management of infection at the surgical site.
- 15 e) Nutritional support.
- 16 f) Selection of prostheses for hip replacement.
- 17 g) Complementary and alternative therapies.

18 **13.9.4 Main outcomes**

- 19 a) Requirement for surgical revision.
- 20 b) Short-term and long-term mortality.
- 21 c) Length of stay in secondary care.
- 22 d) Length of time before community resettlement/discharge.
- 23 e) Place of residence (compared with baseline) 12 months after fracture.

1 f) Short-, medium- and long-term functional status.

2 g) Short-, medium- and long-term quality of life.

3 **13.9.5 Economic aspects**

4 Developers will take into account both clinical and cost effectiveness when making
5 recommendations involving a choice between alternative interventions. A review of the
6 economic evidence will be conducted and analyses will be carried out as appropriate. The
7 preferred unit of effectiveness is the quality-adjusted life year (QALY), and the costs
8 considered will usually be only from an NHS and personal social services (PSS)
9 perspective. Further detail on the methods can be found in 'The guidelines manual' (see
10 'Further information').

11 **13.9.6 Status**

12 **13.9.6.1 Scope**

13 This is the final scope.

14 **13.9.6.2 Timing**

15 The development of the guideline recommendations will begin in June 2010.

16 **13.10 Related NICE guidance**

17 **13.10.1 Published**

- 18 • Surgical site infection. NICE clinical guideline 74 (2008). Available from
19 www.nice.org.uk/CG74
- 20 • Alendronate, etidronate, risedronate, raloxifene, strontium ranelate and teriparatide
21 for the secondary prevention of osteoporotic fragility fractures in postmenopausal
22 women. NICE technology appraisal guidance 161 (2008). Available from
23 www.nice.org.uk/TA161
- 24 • Alendronate, etidronate, risedronate, raloxifene and strontium ranelate for the
25 primary prevention of osteoporotic fragility fractures in postmenopausal women. NICE
26 technology appraisal guidance 160 (2008). Available from www.nice.org.uk/TA160

- 1 • Venous thromboembolism. NICE clinical guideline 46 (2007). Available from
- 2 www.nice.org.uk/CG46
- 3 • Delirium: diagnosis, prevention and management of delirium. NICE clinical guideline
- 4 103 (2010). Available from www.nice.org.uk/guidance/CG103
- 5 • Venous thromboembolism –prevention. NICE clinical guideline 92 (2010). Available
- 6 from www.nice.org.uk/guidance/CG92 Minimally invasive hip replacement. NICE
- 7 interventional procedure guidance (2010). Available from
- 8 www.nice.org.uk/guidance/IPG363
- 9 • Nutrition support in adults. NICE clinical guideline 32 (2006). www.nice.org.uk/CG32
- 10 • The management of pressure ulcers in primary and secondary care. NICE clinical
- 11 guideline 29 (2005). Available from www.nice.org.uk/CG29
- 12 • Falls. NICE clinical guideline 21 (2004). Available from www.nice.org.uk/CG21
- 13 • Preoperative tests. NICE clinical guideline 3 (2003). Available from
- 14 www.nice.org.uk/CG3
- 15 • The selection of prostheses for primary total hip replacement. NICE technology
- 16 appraisal guidance 2 (2000). Available from www.nice.org.uk/TA2

17 **13.10.2 Guidance under development**

18 NICE is currently developing the following related guidance (details available from the
19 NICE website).

- 20 • Osteoporosis. NICE clinical guideline. Publication date to be confirmed.

21 **13.11 Further information**

22 Information on the guideline development process is provided in:

- 23 • ‘How NICE clinical guidelines are developed: an overview for stakeholders, the public
- 24 and the NHS’
- 25 • ‘The guidelines manual’.

26 These are available from the NICE website (www.nice.org.uk/guidelinesmanual).

27 Information on the progress of the guideline will also be available from the NICE website
28 (www.nice.org.uk).

1
2

1 **14 Appendix B: Declarations of Interest**

2 **14.1 Introduction**

3 All members of the GDG and all members of the NCGC staff were required to make
4 formal declarations of interest at the outset of each meeting, and these were
5 updated at every subsequent meeting throughout the development process. No
6 interests were declared that required actions.

1 14.2 Declarations of interests of the GDG members

2 14.2.1 Professor Cameron Swift

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup meeting) (18th January 2010)	No interests to declare
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	Declared a non personal non pecuniary interest: has been invited to join the Department of Health Board on Fragility Fractures Programme
Twelfth GDG Meeting (29th July 2010)	No change
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

3

4

1

14.2.2 Professor Opinder Sahota

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	Did not attend
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	No interests to declare
Seventh GDG Meeting (9th March 2010)	Declared a non personal, non pecuniary interest regarding involvement in the Map of Medicine project with the Department of Health.
Eighth GDG Meeting (26th April 2010)	No change
Tenth GDG Meeting (11th June 2010)	Did not attend
Eleventh GDG Meeting (30th June 2010)	Declared a non personal non pecuniary interest: has been invited to join the Department of Health Board on Fragility Fractures Programme
Twelfth GDG Meeting (29th July 2010)	No change
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	Did not attend
Actions	None required

2

3

1 **14.2.3 Dr Antony Johansen**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup Workshop) (18th January 2010)	No interests to declare
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	Did not attend
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	No interests to declare
Thirteenth GDG Meeting (8th September 2010)	No interests to declare
Fourteenth GDG Meeting (18th January 2011)	No interests to declare
Fifteenth GDC Meeting (24th March 2011)	No interests to declare
Actions	None required

2

3

1 14.2.4 Mr Tim Chesser

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	Performed consultancy work with orthopaedic manufacturer for unrelated orthopaedic implants (locking plates for particular fractures)- compliance and worded guidelines. His Department receives research support from orthopaedic manufacturers including DePuy, Smith and Nephew, Biomet and Stryker. Department have research fellows funded by orthopaedic manufacturer. Publishing RCT on surgical treatment for peri-articular fractures which was not funded by industry.
Second GDG Meeting (17th July 2009)	Did not attend
Third GDG Meeting (15th September 2009)	Performed consultancy work with orthopaedic manufacturer for unrelated orthopaedic implants (locking plates for particular fractures)- compliance and worded guidelines. His Department receives research support from orthopaedic manufacturers including DePuy, Smith and Nephew, Biomet and Stryker. Department have research fellows funded by orthopaedic manufacturer. Publishing RCT on surgical treatment for peri-articular fractures which was not funded by industry.
Fourth GDG Meeting (8th December 2009)	Travel and accommodation funded by the Orthopaedic Trauma Association in the US to present a poster on outcomes in Pelvic Fractures at an Experts in Pelvic Trauma meeting (sponsored by Stryker Trauma).
Fifth GDG Meeting (14th December 2009)	No change
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	Did not attend
Eighth GDG Meeting (26th April 2010)	No change
Tenth GDG Meeting (11th June 2010)	No change
Eleventh GDG Meeting (30th June 2010)	Did not attend meeting
Twelfth GDG Meeting (29th July 2010)	Declared a personal non pecuniary interest- invited to teach on a hip fracture surgical techniques course organised by Stryker who paid his travel expenses. No other payment was received.
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	Declared that he has a contract with an orthopaedic company (Stryker) to design reduction clamps, instrumentation and update for pelvic ring and acetabular fractures. He was also invited to be on the NHS Map of Medicine Commissioners' toolkit.
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

2

3

1 **14.2.5 Mr Bob Handley**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	Non personal pecuniary interest: responsibility for – Synthes Fellows in the Trauma Department at the John Radcliffe hospital- 2 week fellowships usually 3-4 per year.
Second GDG Meeting (17th July 2009)	No change
Third GDG Meeting (15th September 2009)	No change
Fourth GDG Meeting (8th December 2009)	No change
Fifth GDG Meeting (14th December 2009)	No change
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	No change
Eighth GDG Meeting (26th April 2010)	No change
Tenth GDG Meeting (11th June 2010)	No change
Eleventh GDG Meeting (30th June 2010)	Did not attend meeting
Twelfth GDG Meeting (29th July 2010)	No change
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

2

3

4

1 **14.2.6 Ms Karen Hertz**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	Miss Karen Hertz- funding for flights and accommodation by a Chinese university to attend a conference in Hong Kong.
Fifth GDG Meeting (14th December 2009)	No change
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	No change
Seventh GDG Meeting (9th March 2010)	KH declared a non personal, non pecuniary interest regarding involvement in the Map of Medicine project with the Department of Health.
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	Did not attend
Eleventh GDG Meeting (30th June 2010)	No change
Twelfth GDG Meeting (29th July 2010)	Declared a non personal non pecuniary interest: has been invited to join the Department of Health Board on Fragility Fractures Programme
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

2

3

1 **14.2.7 Dr Richard Griffiths**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	Did not attend
Third GDG Meeting (15th September 2009)	Did not attend
Fourth GDG Meeting (8th December 2009)	Did not attend
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	Did not attend
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	Did not attend
Thirteenth GDG Meeting (8th September 2010)	Did not attend
Fourteenth GDG Meeting (18th January 2011)	Did not attend
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

2

3

1 **14.2.8 Professor Sallie Lamb**

GDG meeting	Declaration of Interests
First GDG meeting (1st July 2009)	Did not attend
Second GDG Meeting (17th July 2009)	Declared a non personal pecuniary interest: NIHR funded research grant. One trial is in the final stages of finding approval in primary care- using peripheral fracture (including hip fracture). The second- potential trial- ideas unclear as to whether they will be submitted. Vitamin D in Hip fracture; anaemia in hip fracture.
Third GDG Meeting (15th September 2009)	No change
Fourth GDG Meeting (8th December 2009)	No change
Fifth GDG Meeting (14th December 2009)	No change
Sixth GDG Meeting (Subgroup workshop) (28th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	Did not attend
Eighth GDG Meeting (26th April 2010)	No change
Tenth GDG Meeting (11th June 2010)	No change
Eleventh GDG Meeting (30th June 2010)	No change
Twelfth GDG Meeting (29th July 2010)	Did not attend
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	Did not attend
Fifteenth GDC Meeting (24th March 2011)	Did not attend
Actions	None required

2

3

1 **14.2.9 Mrs Heather Towndrow**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	No interests to declare
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	No interests to declare
Thirteenth GDG Meeting (8th September 2010)	No interests to declare
Fourteenth GDG Meeting (18th January 2011)	Did not attend
Fifteenth GDC Meeting (24th March 2011)	Did not attend
Actions	None required

2

1 **14.2.10 Dr Sally Hope**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	Did not attend
Second GDG Meeting (17th July 2009)	Declared a personal pecuniary interest- MSD paid for hotel in Manchester for NOS Conference (approx £200) in July 2009: in accordance with NOS policy to reduce costs for speakers.
Third GDG Meeting (15th September 2009)	Did not attend
Fourth GDG Meeting (8th December 2009)	No change
Fifth GDG Meeting (14th December 2009)	No change
Sixth GDG Meeting (Subgroup workshop) (28th January 2010)	Did not attend meeting
Seventh GDG Meeting (9th March 2010)	Declared a non personal, non pecuniary interest regarding involvement in the Map of Medicine project with the Department of Health.
Eighth GDG Meeting (26th April 2010)	Did not attend
Tenth GDG Meeting (11th June 2010)	No change
Eleventh GDG Meeting (30th June 2010)	Did not attend meeting
Twelfth GDG Meeting (29th July 2010)	No change
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	Did not attend
Actions	None required

2

1 **14.2.11 Ms Tessa Somerville**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend meeting
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	No interests to declare
Thirteenth GDG Meeting (8th September 2010)	No interests to declare
Fourteenth GDG Meeting (18th January 2011)	No interests to declare
Fifteenth GDC Meeting (24th March 2011)	No interests to declare
Actions	None required

2

1 **14.2.12 Mr Anthony Field**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	Did not attend
Fifth GDG Meeting (14th December 2009)	Did not attend
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	No interests to declare
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	No interests to declare
Thirteenth GDG Meeting (8th September 2010)	No interests to declare
Fourteenth GDG Meeting (18th January 2011)	No interests to declare
Fifteenth GDC Meeting (24th March 2011)	No interests to declare
Actions	None required

2

1 **14.2.13 Mr Martin Wise**

<i>GDG meeting</i>	<i>Declaration of Interests</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	No interests to declare
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	Did not attend
Seventh GDG Meeting (9th March 2010)	No interests to declare
Eighth GDG Meeting (26th April 2010)	Did not attend
Tenth GDG Meeting (11th June 2010)	No interests to declare
Eleventh GDG Meeting (30th June 2010)	No interests to declare
Twelfth GDG Meeting (29th July 2010)	No interests to declare
Thirteenth GDG Meeting (8th September 2010)	No interests to declare
Fourteenth GDG Meeting (18th January 2011)	No interests to declare
Fifteenth GDC Meeting (24th March 2011)	No interests to declare
Actions	None required

2

3

1 14.3 Declarations of interests of the NCGC members

<i>GDG meeting</i>	<i>Declaration of Interests of the NCGC members</i>
First GDG meeting (1st July 2009)	No interests to declare
Second GDG Meeting (17th July 2009)	No interests to declare
Third GDG Meeting (15th September 2009)	No interests to declare
Fourth GDG Meeting (8th December 2009)	No interests to declare
Fifth GDG Meeting (14th December 2009)	Antonia Morga declared her husband works for Novartis
Sixth GDG Meeting (Subgroup workshop) (18th January 2010)	No change
Seventh GDG Meeting (9th March 2010)	No change
Eighth GDG Meeting (26th April 2010)	No change
Tenth GDG Meeting (11th June 2010)	No change
Eleventh GDG Meeting (30th June 2010)	No change
Twelfth GDG Meeting (29th July 2010)	No change
Thirteenth GDG Meeting (8th September 2010)	No change
Fourteenth GDG Meeting (18th January 2011)	No change
Fifteenth GDC Meeting (24th March 2011)	No change
Actions	None required

2

3

1 **14.4 Declarations of interests of the Expert Advisors**

2 **14.4.1 Mr Martin Parker**

3 Mr Martin Parker only attended the first and second GDG meetings. He declared
4 that he had received and may in the future receive money for advising implant
5 manufacturing companies about their products and advising on implant design. He
6 has produced research papers with different conclusions and publically presented
7 the results. No actions were required as the first two meetings were introductory
8 and did not involve any discussions about the evidence or formulating
9 recommendations.

10 **14.4.2 Mrs Pamela Holmes**

11 Mrs Pamela Holmes had no interests to declare and did not attend any GDG
12 meetings

13 **14.4.3 Professor Judith Adams**

14 Professor Judith Adams only attended the twelfth GDG meeting on July 29th 2010
15 and did not have any interests to declare.

16

17

1 15 Appendix C: Review protocols

2 15.1 Review protocol – Imaging in occult hip fracture

Component	Description
Review question	In patients with a continuing clinical suspicion of hip fracture, despite negative radiographic findings, what is the clinical and cost-effectiveness of additional imaging (radiography after at least 48 hours), Radionuclide scanning (RNS), ultrasound (US) and computed tomography (CT), compared to magnetic resonance imaging (MRI), in confirming, or excluding, a hip fracture?
Objectives	To identify an alternative method of diagnosis of occult hip fractures when MRI is not available.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	<ul style="list-style-type: none"> ▪ Computed tomography ▪ Radionuclide scanning (also known as isotope scanning or scintigraphy).
Comparison	<ul style="list-style-type: none"> ▪ Magnetic resonance imaging
Outcomes	<ul style="list-style-type: none"> ▪ Sensitivity ▪ Specificity ▪ Positive and negative predictive values ▪ Positive and negative likelihood ratios

Search strategy

The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.

Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.

Studies will be restricted to English language only

No date restriction will be applied. Databases will be searched from their date of origin

The review strategy

Meta-analysis will not be conducted for diagnostic studies. Ranges of results will be reported.

If there is heterogeneity the following subgroups will be analysed separately:

- Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them)
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- *Palliative care patients*

1

2

1 15.2 Review protocol – Timing of surgery

Component	Description
Review question	In patients with hip fractures what is the clinical and cost effectiveness of early surgery (within 24, 36 or 48 hours) on the incidence of complications such as mortality, pneumonia, pressure sores, cognitive dysfunction and increased length of hospital stay?
Objectives	To investigate whether early surgery improves patient outcomes.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Early surgery (within the cut off of 24, 36 and 48 hours of admission to hospital)
Comparison	Late surgery (after the cut off of 24, 36 and 48 hours of admission)
Outcomes	<p>Mortality (30 days, 3 months, 1 year)</p> <p>Length of stay in secondary care</p> <p>Length of time before community resettlement/discharge.</p> <p>Place of residence (compared with baseline) 12 months after fracture.</p> <p>Functional status (30 days, 3 months, 1 year)</p> <p>Quality of life (30 days, 3 months, 1 year)</p> <p>Complications (including pressure ulcers)</p>
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found well conducted cohort studies and observational studies may also be considered. In particular, cohort studies using logistic regression to adjust for confounders such as comorbidity and age, which is a particular bias in this area.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>If there is heterogeneity the following subgroups will be analysed separately:</p> <ul style="list-style-type: none"> ▪ Reason for delay to surgery (administrative or medical reasons) ▪ Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them) ▪ Concurrent medication ▪ Age

- Gender
- Cognitive impairment

1

2

3

1 15.3 Review protocol – Analgesia- systemic medications

Component	Description
Review question	In patients who have or are suspected of having a hip fracture, what is the comparative effectiveness and cost effectiveness of systemic analgesics in providing adequate pain relief and reducing side effects and mortality?
Objectives	To identify the most effective systemic analgesia medication for pain relief in hip fracture patients
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	<p>Systemic:</p> <ul style="list-style-type: none"> ▪ Opioids e.g. <ul style="list-style-type: none"> ○ Buprenorphine ○ Codeine ○ Dihydrocodeine ○ Hydromorphone ○ Morphine ○ Oxycodone ○ Papaveretum (no, has been withdrawn) ○ Pentazocine ○ Pethidine (?) causes delirium in elderly ○ Tramadol (potent cause of delirium in elderly) ▪ Non Opioid e.g. <ul style="list-style-type: none"> ○ Paracetamol, iv, PR, oral ○ Non steroidal anti inflammatory (NSAIDs)
Comparison	<p>Systemic:</p> <ul style="list-style-type: none"> ▪ Opioids e.g. <ul style="list-style-type: none"> ○ Buprenorphine ○ Codeine ○ Dihydrocodeine ○ Hydromorphone ○ Morphine ○ Oxycodone ○ Papaveretum (no, has been withdrawn) ○ Pentazocine ○ Pethidine (?) causes delirium in elderly ○ Tramadol (potent cause of delirium in elderly) ▪ Non Opioid e.g.

- Paracetamol, iv, PR, oral
- Non steroidal anti inflammatory (NSAIDs)

Outcomes

- Pain (generally measured by visual analogue scale or verbal rating)
- Need for 'breakthrough' analgesia
- Mortality
- Adverse effects
 - Paracetamol
 - Virtually none but may decrease blood pressure with iv
 - Opioids
 - Itching/histamine release,
 - PONV,
 - respiratory depression,
 - decrease in blood pressure,
 - delerium

Search strategy

The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.

Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.

Studies will be restricted to English language only

No date restriction will be applied. Databases will be searched from their date of origin

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Comorbidities strongly predictive of outcome
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Type of fracture
- Type of surgery
 - THR vs. hemiarthroplasty
 - THR vs. internal fixation

1 **15.4 Review protocol – Analgesia- Nerve blocks compared to systemic**
 2 **analgesics**

Component	Description
Review question	In patients who have or are suspected of having a hip fracture, what is the clinical and cost effectiveness of nerve blocks compared to systemic analgesia in providing adequate pain relief and reducing side effects and mortality?
Objectives	To identify an optimal analgesia protocol including the use of nerve blocks which may help reduce usage of systemic analgesics with strong side effects in this patient group.
Population	<p>Patients over 18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Nerve blocks (any type: lateral cutaneous, femoral, triple, psoas, 3-in-1 [includes femoral, obturator, lateral femoral cutaneous nerves], fascia iliaca, with ultrasound guidance for localisation)
Comparison	<p>Pharmacological (systemic):</p> <ul style="list-style-type: none"> ▪ Opioids e.g. <ul style="list-style-type: none"> ○ Buprenorphine ○ Codeine ○ Dihydrocodeine ○ Hydromorphone ○ Morphine ○ Oxycodone ○ Papaveretum (no, has been withdrawn) ○ Pentazocine ○ Pethidine (?) causes delirium in elderly ○ Tramadol (potent cause of delirium in elderly) ▪ Non Opioid e.g. <ul style="list-style-type: none"> ○ Paracetamol, iv, PR, oral ▪ NSAIDs <ul style="list-style-type: none"> ○ upper gastrointestinal bleeding ○ renal, hepatic and cardiovascular side effects
Outcomes	<ul style="list-style-type: none"> ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Need for 'breakthrough' analgesia ▪ Mortality

- Adverse effects
 - Nerve Block:
 - Nerve damage
 - Pressure necrosis following motor block
 - Postoperative nausea and vomiting (PONV)
 - Paracetamol
 - Virtually none but may decrease blood pressure with iv
 - Opioids
 - Itching/histamine release,
 - PONV,
 - respiratory depression,
 - decrease in blood pressure,
 - delirium
 - NSAIDs
 - upper gastrointestinal bleeding
 - renal, hepatic and cardiovascular side effects

Search strategy

The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.

Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.

Studies will be restricted to English language only

No date restriction will be applied. Databases will be searched from their date of origin

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Comorbidities strongly predictive of outcome
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Type of fracture
- Type of surgery
 - THR vs. hemiarthroplasty
 - THR vs. internal fixation

1

2 **15.5 Review protocol - Anaesthesia**

Component	Description
Review question	In patients undergoing surgical repair for hip fractures, what is the clinical and cost effectiveness of regional (spinal/epidural) anaesthesia compared to general anaesthesia in reducing complications such as mortality, cognitive dysfunction thromboembolic events, postoperative respiratory morbidity, renal failure and length of stay in hospital?
Objectives	To identify whether regional anaesthesia confers any benefit compared to general anaesthesia with regards to reducing complications and improving patient outcomes after surgery.
Population	<p>Patients over 18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	General anaesthesia for different types of surgery
Comparison	<ul style="list-style-type: none"> • Regional anaesthesia for the same type of surgery <ul style="list-style-type: none"> ▪ Spinal/epidural without nerve block ▪ Spinal/epidural with nerve block
Outcomes	<ul style="list-style-type: none"> ▪ Patient preference ▪ Mortality at 30 days ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Adverse effects <ul style="list-style-type: none"> ○ General: <ul style="list-style-type: none"> ▪ postoperative lung complications ▪ Pulmonary emboli ▪ Pneumonia ▪ Myocardial infarction ▪ Renal failure ▪ Postoperative nausea and vomiting (PONV) ○ Regional <ul style="list-style-type: none"> ▪ Neural damage ▪ Spinal haematoma

Search strategy

The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.

Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.

Studies will be restricted to English language only

No date restriction will be applied. Databases will be searched from their date of origin

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately (where possible):

- Comorbidities strongly predictive of outcome
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Type of surgery
 - THR vs. hemiarthroplasty
 - THR vs. internal fixation
- Duration of anaesthesia

1

2 **15.6 Review protocol – surgeon seniority**

Component	Description
Review question	Does surgeon seniority (consultant or equivalent) reduce the incidence of mortality, operative revision and poor functional outcome?
Objectives	To investigate whether senior surgeons lead to better outcomes for hip fracture patients
Population	Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.
Intervention	<ul style="list-style-type: none"> ▪ Consultant grade or equivalent
Comparison	<ul style="list-style-type: none"> ▪ Below consultant grade or equivalent ▪ Trainee
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Reoperation rate ▪ Dislocations ▪ Wound infection
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library and CINAHL.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found well conducted cohort studies and observational studies may also be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>If there is heterogeneity the following subgroups will be analysed separately:</p> <ul style="list-style-type: none"> ▪ Age

3

1 15.7 Review protocol – Cement

Component	Description
Review question	In hip fracture patients undergoing total hip replacement what is the clinical and cost effectiveness of cemented total hip replacement versus uncemented total hip replacement on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?
Objectives	To examine the effectiveness of cement when inserting arthroplasty for surgical repair
Population	<p>Patients >18 years old with a hip fracture undergoing surgical repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Cemented arthroplasty
Comparison	Uncemented arthroplasty
Outcomes	<ul style="list-style-type: none"> ▪ Perioperative mortality ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture ▪ Wound healing complications
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p> <p>All questions relating to surgical repair for hip fractures will be searched together.</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>Studies will be restricted to English language articles</p> <p>If there is heterogeneity the following subgroups will be analysed</p>

separately:

- Comorbidities
- Age
- Ideally “younger and fitter” patients compared to the “older and frailer” patients. Could be a combination of age and comorbidities
- Type of arthroplasty

1 15.8 Review protocol – Intracapsular fractures

Component	Description
Review question	In patients undergoing repair for intracapsular hip fractures what is the clinical and cost effectiveness of internal fixation compared to hemiarthroplasty compared to total hip replacement on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?
Objectives	To examine the effectiveness of the 3 different techniques for fixing displaced intracapsular fractures
Population	<p>Patients >18 years old with a hip fracture undergoing surgical repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	<ul style="list-style-type: none"> • Internal fixation • Hemiarthroplasty • Total hip replacement
Comparison	All of the above are compared to each other.
Outcomes	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p> <p>All questions relating to surgical repair for hip fractures will be searched together.</p>

1

The review strategy

Meta-analyses will be conducted where possible.

Studies will be restricted to English language articles

If there is heterogeneity the following subgroups will be analysed separately:

- Ideally “younger and fitter” patients compared to the “older and frailer” patients. Could be a combination of age and comorbidities
- Type of internal fixation or arthroplasty
- Use of cement in arthroplasty

2

1

2 **15.9 Review protocol – surgical approach**

Component	Description
Review question	In patients having surgical treatment for intracapsular hip fracture with hemiarthroplasty what is the clinical and cost effectiveness of anterolateral compared to posterior surgical approach on mortality, number of reoperations, dislocation, functional status, length of hospital stay, quality of life and pain?
Objectives	To investigate whether one surgical approach is better than the other when inserting a hemiarthroplasty.
Population	<p>Patients >18 years old with a hip fracture undergoing replacement arthroplasty with a hemiarthroplasty</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	<ul style="list-style-type: none"> ▪ Anterolateral approach
Comparison	<ul style="list-style-type: none"> ▪ Posterior approach
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of hospital stay ▪ Reoperation rate ▪ Dislocations ▪ Functional status ▪ Quality of life ▪ Pain
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library and CINAHL.</p> <p>Randomised controlled trials (RCTs) and well conducted cohort studies and observational studies that adjust for confounders will be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Type of procedure

1

2

1 15.10 Review protocol – Hemiarthroplasty stem design

Component	Description
Review question	In patients undergoing surgery for hip fracture what is the clinical and cost effectiveness of 'OEDP 10A rating' designs of stems in preference to Austin Moore or Thompson stems when inserting a hemiarthroplasty on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?
Objectives	To examine the effectiveness of modern design stems ('OEDP 10A rating') compared to Austin Moore or Thompson stems.
Population	<p>Patients >18 years old with a hip fracture undergoing hemiarthroplasty</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Hemiarthroplasty with a modern design stem ('OEDP 10A rating')
Comparison	Hemiarthroplasty with an Austin Moore or Thompson
Outcomes	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Place of residence 12 months after fracture
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p> <p>All questions relating to surgical repair for hip fractures will be searched together.</p>

1

The review strategy

Meta-analyses will be conducted where possible.

Studies will be restricted to English language articles

If there is heterogeneity the following subgroups will be analysed separately:

- Ideally “younger and fitter” patients compared to the “older and frailer” patients. Could be a combination of age and comorbidities

2

3

1 **15.11 Review protocol – extracapsular fractures**

2

Component	Description
Review question	<p>In patients undergoing repair for trochanteric extracapsular hip fractures what is the clinical and cost effectiveness of extramedullary sliding hip screws compared to intramedullary nails on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?</p> <p>In patients undergoing repair for subtrochanteric extracapsular hip fractures, what is the effectiveness of extramedullary sliding hip screws compared to intramedullary nails on mortality, surgical revision, functional status, length of stay, quality of life, pain and place of residence after hip fracture?</p>
Objectives	To examine the effectiveness of extramedullary implants, including sliding hip screws, compared to intramedullary implants, including nails, in fixing trochanteric and subtrochanteric fractures.
Population	<p>Patients >18 years old with a extracapsular hip fracture undergoing surgical repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Extramedullary sliding hip screws
Comparison	Intramedullary nails
Outcomes	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Requirement for reoperation (operative or postoperative fracture of the femur, cut-out and non-union) ▪ Length of stay in hospital/acute care ▪ Length of stay in to community or resettlement (i.e. superspell) ▪ Wound healing complications
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>

All questions relating to surgical repair for hip fractures will be searched together.

The review strategy

Meta-analyses will be conducted where possible.

Studies will be restricted to English language articles

If there is heterogeneity the following subgroups will be analysed separately:

- Stability of fracture
- Comorbidities
- Age
- Previous fracture or surgery to femur

1

2 **15.12 Review protocol – Mobilisation strategies**

Component	Description
Review question	In patients who have undergone surgery for hip fracture, what is the clinical and cost effectiveness of early mobilisation (<48 hours after surgery) compared to late mobilisation on functional status, mortality, place of residence/discharge, pain and quality of life?
Objectives	To examine the effectiveness of early mobilisation on functional outcomes compared to delayed mobilisation
Population	<p>Patients >18 years old that have had surgery for a hip fracture.</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Mobilisation (physiotherapy) within 48 hours of surgery.
Comparison	Mobilisation (physiotherapy) after 48 hours of surgery.
Outcomes	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Discharge destination
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p> <p>All questions relating to surgical repair for hip fractures will be searched together.</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>Studies will be restricted to English language articles</p> <p>If there is heterogeneity the following subgroups will be analysed separately:</p> <ul style="list-style-type: none"> ▪ Comorbidities ▪ Age ▪ Previous fracture or surgery to femur

Component	Description
Review question	In patients who have undergone surgery for hip fracture, what is the clinical and cost effectiveness of intensive physiotherapy compared to non intensive physiotherapy on functional status, mortality, place of residence/discharge, pain and quality of life?
Objectives	To examine the effectiveness of intensity of mobilisation on functional outcomes.
Population	Patients >18 years old that have had surgery for a hip fracture. People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.
Intervention	Intensive physiotherapy, defined by an increased number of sessions or an increase in intensity (strength) of exercise.
Comparison	Fewer sessions of physiotherapy or usual care ad defined by the paper.
Outcomes	<ul style="list-style-type: none"> ▪ Mortality at 30 days, 3 months & 1 year or longer ▪ Functional status up to 1 year ▪ Pain (generally measured by visual analogue scale or verbal rating) ▪ Quality of life ▪ Discharge destination ▪ Mobility
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered.</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p> <p>All questions relating to surgical repair for hip fractures will be searched together.</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>Studies will be restricted to English language articles</p> <p>If there is heterogeneity the following subgroups will be analysed separately:</p> <ul style="list-style-type: none"> ▪ Type or component of exercise programme ▪ Comorbidities ▪ Age ▪ Previous fracture or surgery to femur

1

2 **15.13 Review protocol – Multidisciplinary rehabilitation**

Component	Description
Review question	In patients with hip fracture what is the clinical and cost effectiveness of 'orthogeriatrician' involvement in the whole pathway of assessment, peri-operative care and rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?
Objectives	To identify the benefit of an orthogeriatrician involved early in the care pathway to patient outcomes.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Involvement of an orthogeriatrician/physician throughout patient care, starting from admission
Comparison	No involvement of an orthogeriatrician/physician throughout the care pathway (e.g. only present in rehabilitation).
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge. ▪ Place of residence (compared with baseline) 12 months after fracture. ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them)
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Palliative care patients
- Patients from nursing homes

1

Component	Description
Review question	In patients with hip fracture what is the clinical and cost effectiveness of hospital-based multidisciplinary rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?
Objectives	To identify the effectiveness of hospital-based multidisciplinary rehabilitation compared to usual care.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	<p>Multidisciplinary hospital-based rehabilitation. Multidisciplinary rehabilitation after hip fracture will be assumed if the following core components are present: medicine; nursing; physiotherapy; occupational therapy; social care. Additional components may include: nutrition, pharmacy, clinical psychology. Additional criteria include formal arrangements for co-ordination/teamwork and regular on-going multidisciplinary assessment.</p> <p>Types of multidisciplinary hospital-based rehabilitation include Geriatric orthopaedic rehabilitation unit (GORU); mixed assessment and rehabilitation unit (MARU); geriatric hip fracture programme (GHFP).</p>
Comparison	Usual hospital-based care (not multidisciplinary)
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge. ▪ Place of residence (compared with baseline) 12 months after fracture. ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>

The review strategy

Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Type of hospital-based MDR
- Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them)
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Palliative care patients
- Patients from nursing homes

1
2
3

1

Component	Description
Review question	In patients with hip fracture what is the clinical and cost effectiveness of community-based multidisciplinary rehabilitation on functional status, length of stay in secondary care, mortality, place of residence/discharge, hospital readmission and quality of life?
Objectives	To compare community-based programmes with each other and usual care.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Community-based multidisciplinary rehabilitation, including intermediate care unit-based, home-based (early supported discharge) and social care unit-based. Any programme starting more than 1 week postoperatively will be excluded.
Comparison	Usual hospital-based care (not multidisciplinary)
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge. ▪ Place of residence (compared with baseline) 12 months after fracture. ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>

2

1 **The review strategy** Meta-analyses will be conducted where possible.

If there is heterogeneity the following subgroups will be analysed separately:

- Type of community rehabilitation programme
- Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them)
- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Palliative care patients
- Patients from nursing homes

2
3
4

1 15.14 Review protocol – Carer involvement

Component	Description
Review question	In patients who have been discharged after hip fracture repair, what is the clinical and cost effectiveness of having a non paid carer (e.g. spouse, relative, friends) on mortality, length of stay, place of residence/discharge, functional status, hospital readmission and quality of life?
Objectives	To compare the effectiveness of hospital-based multidisciplinary rehabilitation with involvement of a carer versus without a carer.
Population	<p>Patients >18 years old with a hip fracture undergoing different types of surgery for hip fracture repair</p> <p>People with fractures caused by specific pathologies other than osteoporosis or osteopaenia, and patients under 18 years old are excluded from the scope.</p>
Intervention	Hospital-based multidisciplinary rehabilitation with involvement of a non paid carer (e.g. spouse, relative, friends).
Comparison	Hospital-based multidisciplinary rehabilitation without involvement of a non paid carer (e.g. spouse, relative, friends).
Outcomes	<ul style="list-style-type: none"> ▪ Mortality (30 days, 3 months, 1 year) ▪ Length of stay in secondary care ▪ Length of time before community resettlement/discharge. ▪ Place of residence (compared with baseline) 12 months after fracture. ▪ Functional status (30 days, 3 months, 1 year) ▪ Hospital readmission ▪ Quality of life (30 days, 3 months, 1 year)
Search strategy	<p>The databases to be searched are Medline, Embase, The Cochrane Library, CINAHL and AMED.</p> <p>Randomised controlled trials (RCTs) will be considered. If no RCTs are found for certain outcomes such as adverse events, well conducted cohort studies and observational studies may also be considered.</p> <p>Studies will be restricted to English language only</p> <p>No date restriction will be applied. Databases will be searched from their date of origin</p>
The review strategy	<p>Meta-analyses will be conducted where possible.</p> <p>If there is heterogeneity the following subgroups will be analysed separately:</p> <ul style="list-style-type: none"> ▪ Comorbidities strongly predictive of outcome (as mentioned in the scope but will need the GDG to list them)

- Concurrent medication
- Age
- Gender
- Cognitive impairment
- Palliative care patients
- Patients from nursing homes

1
2

1

2 **15.15 Review protocol – Health Economics**

Objectives The aim is to identify economic studies relevant to the review questions set out above.

Criteria Populations, interventions and comparators as specified in the review protocols above. Must be a relevant economic study design (cost-utility analysis, cost-benefit analysis, cost-effectiveness analysis, cost-consequence analysis, comparative cost analysis).

Search strategy See Appendix D, section 4.2

The review strategy Each study is assessed using the NICE economic evaluation checklist – NICE (2009) Guidelines Manual, Appendix H.

Inclusion/exclusion criteria

- If a study is rated as both ‘Directly applicable’ and ‘Minor limitations’ (using the NICE economic evaluation checklist) then it should be *included* in the guideline. An evidence table should be completed and it should be included in the economic profile.
- If a study is rated as either ‘Not applicable’ or ‘Very serious limitations’ then it should be *excluded* from the guideline. It should not be included in the economic profile and there is no need to include an evidence table.
- If a study is rated as ‘Partially applicable’ and/or ‘Potentially serious limitations’ then there is *discretion* over whether it should be included. The health economist should make a decision based on the relative applicability and quality of the available evidence for that question, in discussion with the GDG if required. The ultimate aim being to include studies that are helpful for decision making in the context of the guideline. Where exclusions occur on this basis, this should be noted in the relevant section of the guideline with references.

Also exclude:

- unpublished reports unless submitted as part of the call for evidence
- abstract-only studies
- letters

- editorials
- reviews of economic evaluations⁵
- foreign language articles

Where there is discretion

The health economist should be guided by the following hierarchies.

Setting:

1. UK NHS
2. OECD countries with predominantly public health insurance systems (e.g. France, Germany, Sweden)
3. OECD countries with predominantly private health insurance systems (e.g. USA, Switzerland)
4. Non-OECD settings (always 'Not applicable')

Economic study type:

1. Cost-utility analysis
2. Other type of full economic evaluation (cost-benefit analysis, cost-effectiveness analysis, Cost-consequence analysis)
3. Comparative cost analysis
4. Non-comparative cost analyses including cost of illness studies (always 'Not applicable')

Year of analysis:

- The more recent the study, the more applicable it is

Quality of effectiveness data used in the economic analysis:

- The more closely the effectiveness data used in the economic analysis matches with the studies included for the clinical review the more useful the analysis will be to decision making for the guideline.

⁵ Recent reviews will be ordered although not reviewed. The bibliographies will be checked for relevant studies, which will then be ordered.

1 16 Appendix D: Literature search strategies

2 16.1 Search Strategies

3 Searches were constructed by using the following groups of terms. These groups are
4 expanded in full in Section 1.2 below.

5 All searches were run in Medline, Embase and the Cochrane Library. Additionally CINAHL and
6 PsychINFO were searched where this was deemed appropriate. Economic searches were
7 conducted in Medline, Embase, NHS EED and the HTA (Health Technology Reports) database
8 from the Cochrane Library.

9

10 Anaesthesia search

11

12

Hip fracture terms

13

AND

14

Anaesthesia terms

15

AND

16

RCT filter or systematic review filter

17

NOT

18

Animal/publication filter

19

Analgesia search

20

21

Hip fracture terms

22

AND

23

Analgesia terms

24

AND

25

RCT filter or systematic review filter

26

NOT

27

Animal/publication filter

28

29

Carer involvement search

30

31

Hip fracture terms

32

AND

33

Carer involvement terms

34

NOT

35

Animal/publication filter

1
2 Early surgery search
3
4 Hip fracture terms
5 AND
6 Early surgery terms
7 NOT
8 Animal/publication filter
9
10 Economic searches (Medline and Embase)
11
12 Hip fracture terms
13 AND
14 Economic filter
15 NOT
16 Animal/publication filter
17
18 Economic searches (NHS EED and HTA)
19
20 Hip fracture terms
21
22 Orthogeriatrician search
23
24 Hip fracture terms
25 AND
26 Orthogeriatrician terms
27 NOT
28 Animal/publication filter
29
30 Patient education search
31
32 Hip fracture terms
33 AND
34 Patient education terms
35 NOT
36 Animal/publication filter
37
38 Patient views search
39
40 Hip fracture terms
41 AND
42 Patient view terms
43 NOT
44 Animal/publication filter
45
46 Radiological imaging search
47

1 Hip fracture terms
 2 AND
 3 Radiological imaging terms
 4 AND
 5 RCT filter or systematic review filter or diagnostic filter
 6 NOT
 7 Animal/publication filter
 8
 9 Rehabilitation search
 10
 11 Hip fracture terms
 12 AND
 13 Rehabilitation terms
 14 NOT
 15 Animal/publication filter
 16
 17 Surgeon seniority search
 18
 19 Hip fracture terms
 20 AND
 21 Surgeon seniority terms
 22 NOT
 23 Animal/publication filter
 24
 25 Surgical interventions search
 26
 27 Hip fracture terms
 28 AND
 29 Surgical intervention terms
 30 AND
 31 RCT filter or systematic review filter
 32 NOT
 33 Animal/publication filter
 34
 35

36 16.2 Search terms

37 Anaesthesia

Anaesthesia terms – Cochrane Library

- 1 MeSH descriptor Anesthesia explode all trees
- 2 ((an?esthet* or an?esthesia) NEAR/4 (regional* or local* or general or spinal or epidural)):ti,ab,kw
- 3 #1 OR #2

38

Anaesthesia terms - OVID Embase

- 1 exp Anesthesia/
 2 ((an?esthet\$ or an?esthesia) adj4 (regional\$ or local\$ or general or spinal or
 epidural)).ti,ab.
 3 1 or 2

1

Anaesthesia terms - OVID Medline

- 1 exp Anesthesia/
 2 ((an?esthet\$ or an?esthesia) adj4 (regional\$ or local\$ or general or spinal or
 epidural)).ti,ab.
 3 1 or 2

2

3 Analgesia**Analgesia terms – Cochrane Library**

- 1 MeSH descriptor Analgesia explode all trees
 2 MeSH descriptor Analgesics explode all trees
 3 MeSH descriptor Nerve Block explode all trees
 4 (analg\$ or (pain* NEAR/3 relie*) or ((nerve* or neural*) NEAR/3 block*)):ti,ab,kw
 5 (opioid* or opiate*):ti,ab,kw
 6 (paracetamol or propacetamol or acetaminophen or co-codamol):ti,ab,kw
 7 (morphine or buprenorphine or codeine or diphenoxylate or dipipanone or
 diamorphine or dihydrocodeine or alfentanil or fentanyl or remifentanil or
 meptazinol or methadone or oxycodone or papaveretum or pentazocine or
 pethidine or tramadol):ti,ab,kw
 8 MeSH descriptor Opiate Alkaloids explode all trees
 9 MeSH descriptor Acetaminophen explode all trees
 10 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9

4

Analgesia terms - OVID Embase

- 1 exp analgesia/
 2 exp Nerve Block/
 3 (analg\$ or (pain\$ adj3 relie\$) or ((nerve\$ or neural\$) adj3 block\$)).ti,ab.
 4 exp analgesic agent/
 5 (morphine or buprenorphine or codeine or diphenoxylate or dipipanone or
 diamorphine or dihydrocodeine or alfentanil or fentanyl or remifentanil or
 meptazinol or methadone or oxycodone or papaveretum or pentazocine or
 pethidine or tramadol).ti,ab.
 6 (paracetamol or propacetamol or acetaminophen or co-codamol).ti,ab.
 7 (opioid\$ or opiate\$).ti,ab.
 8 or/1-7

5

Analgesia terms - OVID Medline

- 1 exp Analgesia/
 2 exp Nerve Block/
 3 exp Analgesics/
 4 (analg\$ or (pain\$ adj3 relie\$) or ((nerve\$ or neural\$) adj3 block\$)).ti,ab.
 5 (opioid\$ or opiate\$).ti,ab.

6 (paracetamol or propacetamol or acetaminophen or co-codamol).ti,ab.
 7 (morphine or buprenorphine or codeine or diphenoxylate or dipipanone or
 diamorphine or dihydrocodeine or alfentanil or fentanyl or remifentanil or
 meptazinol or methadone or oxycodone or papaveretum or pentazocine or
 pethidine or tramadol).ti,ab.
 8 exp Opiate Alkaloids/
 9 acetaminophen/
 10 or/1-9

1

2 Animal/publication filter

Animal/publication filter - OVID Embase

1 Case-Study/ or Abstract-Report/ or Letter/ or (case adj report).tw.
 2 (exp Animal/ or Nonhuman/ or exp Animal-Experiment/) not exp Human/
 3 or/1-2

3

Animal/publication filter - OVID Medline

1 ((Case-Reports not Randomized-Controlled-Trial) or Letter or Historical-Article or
 Review-Of-Reported-Cases).pt.
 2 exp Animal/ not Human/
 3 or/1-2

4

5 Carer involvement

Carer involvement terms – Cochrane Library

1 MeSH descriptor Family explode all trees
 2 MeSH descriptor Caregivers, this term only
 3 MeSH descriptor Friends, this term only
 4 MeSH descriptor Voluntary Workers, this term only
 5 (carer* or caregiver* or care giver* or ((care* or caring) NEAR/5 (child* or parent*
 or husband* or wife* or wives or relative* or relation* or spous* or partner* or
 offspring or son* or daughter* or famil* or brother* or sister* or sib* or friend* or
 volunteer*))) :ti,ab,kw
 6 #1 or #2 or #3 or #4 or #5

6

Carer involvement terms – EBSCO CINAHL

1 mh Family+ or mh caregivers or mh friends or mh voluntary workers
 2 carer* or caregiver* or care giver* or care* n5 child* or care* n5 parent* or care*
 n5 husband* or care* n5 wife* or care* n5 wives or care* n5 relative* or care* n5
 relation* or care* n5 spous* or care* n5 partner*
 3 care* n5 offspring or care* n5 son* or care* n5 daughter* or caring n5 child* or
 caring n5 parent* or caring n5 husband* or caring n5 wife* or caring n5 wives or
 caring n5 relative* or caring n5 relation* or caring n5 spous* or caring n5
 partner*
 4 care* n5 famil* or care* n5 brother* or care* n5 sister* or caring n5 offspring or
 caring n5 son* or caring n5 daughter* or caring n5 famil* or caring n5 brother* or
 caring n5 sister* or caring n5 sib* or caring n5 friend* or caring n5 volunteer*
 5 care* n5 sib* or care* n5 friend* or care* n5 volunteer*
 6 S1 or S2 or S3 or S4 or S5

1 Case-Study/ or Abstract-Report/ or Letter/ or (case adj report).tw. or ((exp Animal/
or Nonhuman/ or exp Animal-Experiment/) not exp Human/)

1

Carer involvement terms – Ovid Embase

1 (carer\$ or caregiver\$ or care giver\$ or ((care\$ or caring) adj5 (child\$ or parent\$ or
husband\$ or wife\$ or wives or relative\$ or relation\$ or spous\$ or partner\$ or
offspring or son\$ or daughter\$ or famil\$ or brother\$ or sister\$ or sib\$ or friend\$ or
volunteer\$ or voluntary))).ti,ab.

2 exp family/ or friend/ or caregiver/ or volunteer/

3 or/1-2

2

Carer involvement terms – Ovid Medline

1 exp Family/ or caregivers/ or friends/ or voluntary workers/

2 (carer\$ or caregiver\$ or care giver\$ or ((care\$ or caring) adj5 (child\$ or parent\$ or
husband\$ or wife\$ or wives or relative\$ or relation\$ or spous\$ or partner\$ or
offspring or son\$ or daughter\$ or famil\$ or brother\$ or sister\$ or sib\$ or friend\$ or
volunteer\$ or voluntary))).ti,ab.

3 or/1-2

3

4 Diagnostic filter

Diagnostic filter - OVID Embase

1 exp "SENSITIVITY AND SPECIFICITY"/

2 (sensitivity or specificity).tw.

3 (predictive adj3 value\$).tw.

4 ((false adj positiv\$) or (false adj negativ\$)).tw.

5 (observer adj variation\$).tw.

6 (roc adj curve\$).tw.

7 (likelihood adj3 ratio\$).tw.

8 *Diagnostic Accuracy/

9 exp *hip fracture/di

10 or/1-9

5

Diagnostic filter - OVID Medline

1 exp "Sensitivity and Specificity"/

2 (sensitivity or specificity).tw.

3 (predictive adj3 value\$).tw.

4 exp diagnostic errors/

5 ((false adj positiv\$) or (false adj negativ\$)).tw.

6 (observer adj variation\$).tw.

7 (roc adj curve\$).tw.

8 (likelihood adj3 ratio\$).tw.

9 likelihood functions/

10 exp *hip fractures/di, ra, ri, us

11 or/1-10

6

1 Early Surgery

Early surgery terms – Cochrane Library

- 1 MeSH descriptor Time Factors explode all trees
- 2 (((early or time* or delay*) NEAR/3 (surger* or operat*)) or (fast NEAR/2 track*) or (rapid NEAR/2 transit*) or (time* NEAR/2 factor*)):ti,ab,kw
- 3 #1 OR #2

2

Early surgery terms – EBSCO CINAHL

- 1 early n3 surger* or early n3 operat* or time* n3 surger* or time* n3 operat* or delay* n3 surger* or delay* n3 operat* or fast n2 track* or rapid n2 transit* or time* n2 factor*
- 2 mh time factors+ or mh treatment delay+
- 3 S1 or S2

3

Early surgery terms - OVID Embase

- 1 (((early or time\$ or delay\$) adj3 (surger\$ or operat\$)) or (fast adj2 track\$) or (rapid adj2 transit\$) or (time\$ adj2 factor\$)).ti,ab.
- 2 Therapy Delay/
- 3 1 or 2

4

Early surgery terms - OVID Medline

- 1 time factors/
- 2 (((early or time\$ or delay\$) adj3 (surger\$ or operat\$)) or (fast adj2 track\$) or (rapid adj2 transit\$) or (time\$ adj2 factor\$)).ti,ab.
- 3 1 or 2

5

6 Economic

Economic filter - OVID Embase

- 1 exp economic aspect/
- 2 cost\$.tw.
- 3 (price\$ or pricing\$).tw.
- 4 (fee or fees).tw.
- 5 (financial or finance or finances or financed).tw.
- 6 (value adj2 (money or monetary)).tw.
- 7 resourc\$ allocat\$.tw.
- 8 expenditure\$.tw.
- 9 (fund or funds or funding or fundings or funded).tw.
- 10 (ration or rations or rationing or rations or rationed).tw.
- 11 (saving or savings).tw.
- 12 or/1-11
- 13 Quality of Life/
- 14 quality of life.tw.
- 15 life quality.tw.
- 16 quality adjusted life.tw.
- 17 (qaly\$ or qald\$ or qale\$ or qtime\$).tw.
- 18 disability adjusted life.tw.
- 19 daly\$.tw.

- 20 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).tw.
- 21 (sf6 or sf 6 or short form 6 or shortform 6 or sf six or sfsix or shortform six or short form six).tw.
- 22 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).tw.
- 23 (sf16 or sf 16 or short form 16 or shortform 16 or sf sixteen or sfsixteen or shortform sixteen or short form sixteen).tw.
- 24 (sf20 or sf 20 or short form 20 or shortform 20 or sf twenty or sftwenty or shortform twenty or short form twenty).tw.
- 25 (euroqol or euro qol or eq5d or eq 5d).tw.
- 26 (hql or hqol or h qol or hrqol or hr qol).tw.
- 27 (hye or hyes).tw.
- 28 health\$ equivalent\$ year\$.tw.
- 29 (hui or hui1 or hui2 or hui3).tw.
- 30 health utilit\$.tw.
- 31 disutilit\$.tw.
- 32 rosser.tw.
- 33 (quality of wellbeing or quality of well being).tw.
- 34 qwb.tw.
- 35 willingness to pay.tw.
- 36 standard gamble\$.tw.
- 37 time trade off.tw.
- 38 time tradeoff.tw.
- 39 tto.tw.
- 40 factor analy\$.tw.
- 41 preference based.tw.
- 42 (state adj2 valu\$.tw.
- 43 Life Expectancy/
44 life expectancy\$.tw.
- 45 ((duration or length or period of time or lasting or last or lasted) adj4 symptom\$.tw.
- 46 or/13-46
- 47 exp model/
48 exp Mathematical Model/
49 markov\$.tw.
- 50 Monte Carlo Method/
51 monte carlo.tw.
- 52 exp Decision Theory/
53 (decision\$ adj2 (tree\$ or anlay\$ or model\$)).tw.
- 54 model\$.tw.
- 55 or/47-55
- 56 12 or 46 or 55

1

Economic filter - OVID Medline

- 1 exp "Costs and Cost Analysis"/
- 2 Economics/
3 Economics, Nursing/ or Economics, Medical/ or Economics, Hospital/ or Economics, Pharmaceutical/
4 exp "Fees and Charges"/
5 exp Budgets/
6 budget\$.tw.

7 cost\$.ti.
 8 (cost\$ adj2 (effective\$ or utilit\$ or benefit\$ or minimi\$)).ab.
 9 (economic\$ or pharmaco-economic\$ or pharmaco-economic\$).ti.
 10 (price\$ or pricing\$).tw.
 11 (financial or finance or finances or financed).tw.
 12 (fee or fees).tw.
 13 (value adj2 (money or monetary)).tw.
 14 Value of Life/
 15 quality adjusted life.tw.
 16 (qaly\$ or qald\$ or qale\$ or qtime\$).tw.
 17 disability adjusted life.tw.
 18 daly\$.tw.
 19 Health Status Indicators/
 20 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or
 shortform thirtysix or shortform thirty six or short form thirtysix or short form
 thirty six).tw.
 21 (sf6 or sf 6 or short form 6 or shortform 6 or sf six or sfsix or shortform six or short
 form six).tw.
 22 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or
 shortform twelve or short form twelve).tw.
 23 (sf16 or sf 16 or short form 16 or shortform 16 or sf sixteen or sfsixteen or
 shortform sixteen or short form sixteen).tw.
 24 (sf20 or sf 20 or short form 20 or shortform 20 or sf twenty or sftwenty or
 shortform twenty or short form twenty).tw.
 25 (euroqol or euro qol or eq5d or eq 5d).tw.
 26 (hql or hqol or h qol or hrqol or hr qol).tw.
 27 (hye or hyes).tw.
 28 (hui or hui1 or hui2 or hui3).tw.
 29 utilit\$.tw.
 30 disutilit\$.tw.
 31 rosser.tw.
 32 quality of wellbeing.tw.
 33 qwb.tw.
 34 willingness to pay.tw.
 35 standard gamble\$.tw.
 36 time trade off.tw.
 37 time tradeoff.tw.
 38 tto.tw.
 39 exp models, economic/
 40 models, theoretical/ or models, organizational/
 41 economic model\$.tw.
 42 markov chains/
 43 markov\$.tw.
 44 Monte Carlo Method/
 45 monte carlo.tw.
 46 exp Decision Theory/
 47 (decision\$ adj2 (tree\$ or anlay\$ or model\$)).tw.
 48 or/1-47

1

2 Hip Fracture Terms

Hip fracture terms – Cochrane Library

1 MeSH descriptor Hip Fractures explode all trees

2 ((hip* or pertrochant* or intertrochant* or trochant* or subtrochant* or
intracapsular* or extracapsular* or ((femur* or femoral*) NEAR/3 (neck or
proximal))) NEAR/4 fracture*):ti,ab,kw
3 #1 OR #2

1

Hip fracture terms – EBSCO CINAHL

1 mh hip fractures+
2 femur* n3 proximal n4 fracture* or femur* n3 neck n4 fracture* or femoral* n3
proximal n4 fracture* or femoral* n3 neck n4 fracture* or pertrochant* n4
fracture* or intertrochant* n4 fracture* or trochanteric n4 fracture* or
subtrochanteric n4 fracture* or extracapsular* n4 fracture* or hip* n4 fracture*
3 intracapsular* n4 fracture* or femur* n4 fracture* or femoral* n4 fracture*
4 S1 or S2 or S3

2

Hip fracture terms - OVID Embase

1 exp Hip Fracture/
2 ((femur\$ or femoral\$) adj3 (head or neck or proximal) adj4 fracture\$).ti,ab.
3 ((hip\$ or femur\$ or femoral\$ or trochant\$ or pertrochant\$ or intertrochant\$ or
subtrochant\$ or intracapsular\$ or extracapsular\$) adj4 fracture\$).ti,ab.
4 1 or 2 or 3

3

Hip fracture terms - OVID Medline

1 exp Hip Fractures/
2 ((femur\$ or femoral\$) adj3 (head or neck or proximal) adj4 fracture\$).ti,ab.
3 ((hip\$ or femur\$ or femoral\$ or trochant\$ or pertrochant\$ or intertrochant\$ or
subtrochant\$ or intracapsular\$ or extracapsular\$) adj4 fracture\$).ti,ab.
4 1 or 2 or 3

4

Hip fracture terms - OVID PsychInfo

1 hips/
2 ((femur\$ or femoral\$) adj3 (head or neck or proximal) adj4 fracture\$).ti,ab.
3 ((hip\$ or femur\$ or femoral\$ or trochant\$ or pertrochant\$ or intertrochant\$ or
subtrochant\$ or intracapsular\$ or extracapsular\$) adj4 fracture\$).ti,ab.
4 1 or 2 or 3

5

6 Orthogeriatrician**Orthogeriatrician terms – Cochrane Library**

1 (geriatr*-orthop* or orthop?edic-geriatr* or ortho*-geriatr* or
orthogeriatr*):ti,ab,kw
2 (orthop* NEAR/2 geriatr*):ti,ab,kw
3 MeSH descriptor Physicians, this term only
4 MeSH descriptor Geriatrics explode all trees
5 #1 or #2 or #3 or #4

7

Orthogeriatrician terms – EBSCO CINAHL

1 orthop* n2 geriatr*
 2 geriatr*-orthop* or orthogeriatr* or ortho*-geriatr* or orthop?edic-geriatr*
 3 (MH "Physicians")
 4 (MH "Geriatrics")
 5 (MH "Multidisciplinary Care Team")
 6 S1 or S2 or S3 or S4 or S5

1

Orthogeriatrician terms - OVID Embase

1 (geriatr\$-orthop\$ or orthop?edic-geriatr\$ or ortho\$-geriatr\$ or
 orthogeriatr\$).ti,ab.
 2 (orthop\$ adj2 geriatr\$).ti,ab.
 3 geriatric care/
 4 geriatrics/
 5 physician/
 6 or/1-5

2

Orthogeriatrician terms - OVID Medline

1 (geriatr\$-orthop\$ or orthop?edic-geriatr\$ or ortho\$-geriatr\$ or
 orthogeriatr\$).ti,ab.
 2 (orthop\$ adj2 geriatr\$).ti,ab.
 3 Physicians/
 4 Geriatrics/
 5 or/1-4

3

4 Patient education**Patient education – EBSCO CINAHL**

1 mh Patients or mh Inpatients or mh Outpatients
 2 mh Caregivers or mh Family+ or mh Parents+ or mh Guardianship, Legal
 3 patients or carer* or famil*
 4 S1 or S2 or S3
 5 mh Information Services+ or mh Books+ or mh Pamphlets or mh Counseling
 6 S4 and S5
 7 patient n3 education or patient n3 educate or patient n3 educating or patient n3
 information or patient n3 literature or patient n3 leaflet* or patient n3 booklet* or
 patient n3 pamphlet*
 8 patients n3 education or patients n3 educate or patients n3 educating or patients
 n3 information or patients n3 literature or patients n3 leaflet* or patients n3
 booklet* or patients n3 pamphlet*
 9 mh Patient Education+
 10 S6 or S7 or S8 or S9

5

Patient education - OVID Embase

1 Patient/ or Hospital patient/ or Outpatient/
 2 Caregiver/ or exp Family/ or exp Parent/
 3 (patients or carer\$ or famil\$).tw.
 4 or/1-3

5 Information Service/ or Information center/ or Publication/ or Book/ or
 Counseling/ or Directive counseling/
 6 4 and 5
 7 ((patient or patients) adj3 (education or educate or educating or information or
 literature or leaflet\$ or booklet\$ or pamphlet\$)).ti,ab.
 8 Patient information/ or Patient education/
 9 or/6-8

1

Patient education – OVID Medline

1 Patients/ or Inpatients/ or Outpatients/
 2 Caregivers/ or exp Family/ or exp Parents/ or exp Legal-Guardians/
 3 (patients or carer\$ or famil\$).tw.
 4 or/1-3
 5 Popular-Works-Publication-Type/ or exp Information-Services/ or Publications/ or
 Books/ or Pamphlets/ or Counseling/ or Directive-Counseling/
 6 4 and 5
 7 ((patient or patients) adj3 (education or educate or educating or information or
 literature or leaflet\$ or booklet\$ or pamphlet\$)).ti,ab.
 8 Patient-Education/ or Patient-Education-Handout-Publication-Type/
 9 or/6-8

2

Patient education – Ovid PsychInfo

1 exp patients/
 2 caregivers/ or exp family/ or exp parents/ or exp guardianship/
 3 (patients or carer\$ or famil\$).tw.
 4 or/1-3
 5 exp information services/ or exp printed communications media/ or reading
 materials/ or exp counseling/
 6 4 and 5
 7 ((patient or patients) adj3 (education or educate or educating or information or
 literature or leaflet\$ or booklet\$ or pamphlet\$)).ti,ab.
 8 client education/
 10 or/6-9

3

4 Patient views

Patient views – EBSCO CINAHL

1 mh Consumer Satisfaction+ or mh Consumer Attitudes or mh Personal Satisfaction
 or mh Consumer Participation or mh Patient Rights+ or mh Questionnaires+ or mh
 Interviews+ or mh Focus groups or mh surveys
 2 patient* n3 view* or patient* n3 opinion* or patient* n3 awareness or patient* n3
 tolerance or patient* n3 perception or patient* n3 persistenc* or patient* n3
 attitude* or patient* n3 compliance or patient* n3 satisfaction or patient* n3
 concern* or patient* n3 belief* or patient* n3 feeling*
 3 patient* n3 position or patient* n3 idea* or patient* n3 preference* or patient* n3
 choice*
 4 discomfort or comfort or inconvenience or bother* or trouble or fear* or anxiety
 or anxious or embarrass*
 5 S1 or S2 or S3 or S4

5

Patient views - OVID Embase

1 Consumer attitude/ or patient satisfaction/ or patient compliance/ or patient right/
 or health survey/ or questionnaire/ or interview/
 2 (patient\$ adj3 (view\$ or opinion\$ or awareness or tolerance or perception or
 persistenc\$ or attitude\$ or compliance or satisfaction or concern\$ or belief\$ or
 feeling\$ or position or idea\$ or preference\$ or choice\$)).tw.
 3 (Discomfort or comfort or inconvenience or bother\$4 or trouble or fear\$ or anxiety
 or anxious or embarrass\$4).tw.
 4 or/1-3

1

Patient views - OVID Medline

1 exp Consumer-Satisfaction/ or Personal-Satisfaction/ or exp Patient-Acceptance-
 Of-Health-Care/ or exp Consumer-Participation/ or exp Patient-Rights/ or Health
 Care Surveys/ or Questionnaires/ or Interview/ or Focus groups/
 2 (patient\$ adj3 (view\$ or opinion\$ or awareness or tolerance or perception or
 persistenc\$ or attitude\$ or compliance or satisfaction or concern\$ or belief\$ or
 feeling\$ or position or idea\$ or preference\$ or choice\$)).tw.
 3 (Discomfort or comfort or inconvenience or bother\$4 or trouble or fear\$ or anxiety
 or anxious or embarrass\$4).tw.
 4 or/1-3

2

Patient views - OVID PsychInfo

1 exp consumer satisfaction/ or exp client attitudes/ or client participation/ or exp
 client rights/ or treatment compliance/ or consumer surveys/ or exp
 questionnaires/ or interviews/ or expectations/
 2 (patient\$ adj3 (view\$ or opinion\$ or awareness or tolerance or perception or
 persistenc\$ or attitude\$ or compliance or satisfaction or concern\$ or belief\$ or
 feeling\$ or position or idea\$ or preference\$ or choice\$ or expect\$)).tw.
 3 ((Discomfort or comfort or inconvenience or bother\$4 or trouble or fear\$ or
 anxiety or anxious or embarrass\$4).tw..
 4 or/1-3

3

4 Radiological Imaging**Radiological imaging terms – Cochrane Library**

1 MeSH descriptor Magnetic Resonance Imaging, this term only
 2 ((MR or NMR) NEAR/2 tomograph*):ti,ab,kw
 3 (MRI):ti,ab,kw
 4 ((magnetic resonance or MR or NMR) NEAR/2 imag*):ti,ab,kw
 5 MeSH descriptor Tomography, X-Ray Computed, this term only
 6 MeSH descriptor Tomography, Spiral Computed, this term only
 7 mdct:ti,ab,kw
 8 (ct or compute* tomograph* or compute*-tomograph* or cat):ti,ab,kw
 9 MeSH descriptor Radionuclide Imaging, this term only
 10 (((radionuclide or radioisotope or isotope) NEAR (imag* or scan*)) or rns or
 scintigraph* or scintiphotograph*):ti,ab,kw
 11 MeSH descriptor Ultrasonography, this term only
 12 (ultrason* or ultrasound* or sonograph* or echograph*):ti,ab,kw
 13 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12

5

Radiological imaging terms – EBSCO CINAHL

1 mh Magnetic Resonance Imaging or magnetic resonance n2 imag* or MR n2 imag*
 or NMR n2 imag* or MR n2 tomograph\$ or NMR n2 tomograph\$ or MRI
 2 mdct or compute* tomograph* or cat or MH "Tomography, X-Ray Computed" or
 mh Tomography, Spiral Computed or compute*-tomograph* or "ct"
 3 mh Radionuclide Imaging or radionuclide n1 imag* or radioisotope n1 imag* or
 isotope n1 imag* or radionuclide n1 scan* or radioisotope n1 scan* or isotope n1
 scan* or rns or scintigraph* or scintiphotograph*
 4 mh Ultrasonography or ultrason* or sonograph* or echograph* or ultrasound*
 5 S1 or S2 or S3 or S4

1

Radiological imaging terms – OVID Embase

1 nuclear magnetic resonance imaging/
 2 ((magnetic resonance or MR or NMR) adj2 imag\$).ti,ab.
 3 (((MR or NMR) adj2 tomograph\$) or MRI).ti,ab.
 4 computer assisted tomography/
 5 spiral computer assisted tomography/
 6 mdct.ti,ab.
 7 (ct or compute\$ tomograph\$ or compute\$-tomograph\$ or cat).ti,ab.
 8 scintiscanning/ or scintigraphy/
 9 (((radionuclide or radioisotope or isotope) adj1 (imag\$ or scan\$)) or rns or
 scintigraph\$ or scintiphotograph\$).ti,ab.
 10 (ultrason\$ or ultrasound\$ or sonograph\$ or echograph\$).ti,ab.
 11 echography/
 12 or/1-11

2

Radiological imaging terms – OVID Medline

1 Magnetic Resonance Imaging/
 2 ((magnetic resonance or MR or NMR) adj2 imag\$).ti,ab.
 3 (((MR or NMR) adj2 tomograph\$) or MRI).ti,ab.
 4 Tomography, X-Ray Computed/
 5 Tomography, Spiral Computed/
 6 mdct.ti,ab.
 7 (ct or compute\$ tomograph\$ or compute\$-tomograph\$ or cat).ti,ab.
 8 Radionuclide Imaging/
 9 (((radionuclide or radioisotope or isotope) adj1 (imag\$ or scan\$)) or rns or
 scintigraph\$ or scintiphotograph\$).ti,ab.
 10 Ultrasonography/
 11 (ultrason\$ or ultrasound\$ or sonograph\$ or echograph\$).ti,ab.
 12 or/1-11

3

4 RCT filter**RCT filter Embase**

1 Clinical-Trial/ or Randomized-Controlled-Trial/ or Randomization/ or Single-Blind-
 Procedure/ or Double-Blind-Procedure/ or Crossover-Procedure/ or Prospective-
 Study/ or Placebo/

2 (((clinical or control or controlled) adj (study or trial)) or ((single or double or triple) adj (blind\$3 or mask\$3)) or (random\$ adj (assign\$ or allocat\$ or group or grouped or patients or study or trial or distribut\$)) or (crossover adj (design or study or trial)) or placebo or placebos).ti,ab.

3 1 or 2

1

RCT filter Medline

1 Randomized-Controlled-Trials/ or Random-Allocation/ or Double-Blind-Method/ or Single-Blind-Method/ or exp Clinical-Trials as topic/ or Cross-Over-Studies/ or Prospective-Studies/ or Placebos/

2 (Randomized-Controlled-Trial or Clinical-Trial or Controlled-Clinical-Trial).pt.

3 (((clinical or control or controlled) adj (study or trial)) or ((single or double or triple) adj (blind\$3 or mask\$3)) or (random\$ adj (assign\$ or allocat\$ or group or grouped or patients or study or trial or distribut\$)) or (crossover adj (design or study or trial)) or placebo or placebos).ti,ab.

4 or/1-3

2

3 Rehabilitation

Rehabilitation terms - Cochrane Library

- 1 MeSH descriptor Rehabilitation explode all trees
- 2 MeSH descriptor Rehabilitation Centers explode all trees
- 3 MeSH descriptor Rehabilitation Nursing explode all trees
- 4 MeSH descriptor Patient Care Team explode all trees
- 5 MeSH descriptor Patient Care Management explode all trees
- 6 MeSH descriptor Occupational Therapy explode all trees
- 7 MeSH descriptor Physical Therapy Modalities explode all trees
- 8 MeSH descriptor Physical Therapy Department, Hospital explode all trees
- 9 MeSH descriptor Physical Therapy (Specialty) explode all trees
- 10 MeSH descriptor Critical Pathways explode all trees
- 11 MeSH descriptor Therapy, Computer-Assisted explode all trees
- 12 MeSH descriptor Exercise Therapy explode all trees
- 13 MeSH descriptor Social Work explode all trees
- 14 MeSH descriptor Social Support explode all trees
- 15 MeSH descriptor Pain Clinics explode all trees
- 16 MeSH descriptor Patient Education as Topic explode all trees
- 17 MeSH descriptor Health Education explode all trees
- 18 MeSH descriptor Recovery of Function, this term only
- 19 MeSH descriptor Subacute Care, this term only
- 20 MeSH descriptor Residential Facilities explode all trees
- 21 MeSH descriptor Day Care, this term only
- 22 MeSH descriptor Home Care Services, this term only
- 23 MeSH descriptor Home Care Services, Hospital-Based, this term only
- 24 MeSH descriptor Home Nursing, this term only
- 25 MeSH descriptor Hospital Units, this term only
- 26 MeSH descriptor Nursing Homes explode all trees
- 27 MeSH descriptor Walking explode all trees
- 28 MeSH descriptor Caregivers, this term only
- 29 (rehab* or habilitat* or recover*):ti,ab,kw
- 30 (multidisciplinary* or interdisciplinary* or multiprofessional* or multimodal* or mdt or mdr):ti,ab,kw
- 31 (social NEAR (work* or support or care)):ti,ab,kw

- 32 (pain clinic* or pain service* or pain relief unit* or (pain center* or pain
centre*)):ti,ab,kw
- 33 ((treatment* or therap* or training or education* or healthcare) NEAR/10
(program* or intervention* or approach*)):ti,ab,kw
- 34 (early NEAR (mobil* or discharg* or ambulat*)):ti,ab,kw
- 35 (occupational therap* or physical therap* or physiotherap* or physio):ti,ab,kw
- 36 (exercis* NEAR/3 therap*):ti,ab,kw
- 37 ((early or earli* or immediat* or initial* or begin* or first* or first-line or first line
or first choice or primar* or precede* or original*) NEAR/3 (interven* or treat* or
therap* or care or medicine* or technique* or strateg* or activit* or
mobili*)):ti,ab,kw
- 38 (walk or walks or walking):ti,ab,kw
- 39 mobili?ation strateg*:ti,ab,kw
- 40 (ambulate* or ambulation* or ambulating*):ti,ab,kw
- 41 (exerci* NEAR/3 (rehab* or habilitat* or recover* or therap* or treat* or
medicine* or intervention* or technique* or strateg*)):ti,ab,kw
- 42 ((walk* or mobil* or mov* or motor* or physi*) NEAR/3 (rehab* or habilitat* or
recover* or therap* or treat* or medicine* or intervention* or technique* or
strateg*)):ti,ab,kw
- 43 (extend* NEAR/2 care* NEAR/3 (facilit* or service* or unit* or center* or clinic* or
program* or residen* or home* or hous*)):ti,ab,kw
- 44 ((residen* or intermediate* or assist* liv*) NEAR/3 (facilit* or care* or service* or
unit* or center* or clinic* or program* or residen* or home* or hous*)):ti,ab,kw
- 45 ((halfway or transition*) NEAR/3 (home* or hous* or facilit* or care* or residen*
or service* or unit* or center* or clinic* or program*)):ti,ab,kw
- 46 (nurs* NEAR/2 home*):ti,ab,kw
- 47 (geriatr*-orthop* or orthop?edic-geriatr* or ortho*-geriatr* or orthogeriatr* or
goru):ti,ab,kw
- 48 (orthop* NEAR/2 geriatr*):ti,ab,kw
- 49 rehabilitation unit*:ti,ab,kw
- 50 (mixed assessment or maru):ti,ab,kw
- 51 (geriatric hip fracture program* or ghfp):ti,ab,kw
- 52 (day NEAR (hospital* or care or unit*)):ti,ab,kw
- 53 ((home-based or home based) NEAR care):ti,ab,kw
- 54 carer* involve*:ti,ab,kw
- 55 (esd or early supported discharge):ti,ab,kw
- 56 sequential care:ti,ab,kw
- 57 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14
or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26
or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38
or #39 or #40 or #41 or #42 or #43 or #44 or #45 or #46 or #47 or #48 or #49 or #50
or #51 or #52 or #53 or #54 or #55 or #56

1

Rehabilitation terms – EBSCO CINAHL

- 1 (MH "Rehabilitation+")
- 2 (MH "Rehabilitation Nursing")
- 3 (MH "Recovery")
- 4 (MH "Subacute Care")
- 5 (MH "Rehabilitation Centers+")
- 6 mh residential facilities or mh Assisted Living Facilities or mh Halfway Houses
- 7 mh Day Care or mh home care services or mh home care services, hospital-based
or mh home nursing or mh Hospital Units
- 8 mh Nursing Homes+ or mh Patient Care Team+ or mh Patient Care Management+
or mh Physical Therapy Techniques+ or mh Physical Therapy Department,

- Hospital+
- 9 mh Critical Pathways+ or mh Therapy, Computer-Assisted+ or mh Exercise Therapy+ or mh Walking+
- 10 mh Social Work+ or mh Social Support+ or mh Pain Clinics+ or mh Patient Education+ or mh Health Education+ or mh Caregivers
- 11 (MH "Multidisciplinary Care Team+")
- 12 rehab* or habilitat* or recover*
- 13 multidisciplinary* or mdr or mdt or multimodal* or multiprofessional* or interdisciplinary*
- 14 social n1 work* or social n1 support or social n1 care
- 15 pain clinic* or pain service* or pain relief unit* or pain center* or pain centre*
- 16 treatment* n10 program* or treatment* n10 intervention* or treatment* n10 approach* or therap* n10 program* or therap* n10 intervention* or therap* n10 approach* or training n10 program* or training n10 intervention* or training n10 approach* or education* n10 program* or education* n10 intervention* or education* n10 approach*
- 17 healthcare n10 program* or healthcare n10 intervention* or healthcare n10 approach*
- 18 early n1 mobil* or early n1 discharg* or early n1 ambulat*
- 19 occupational therap* or physical therap* or physiotherap* or physio
- 20 exercis* n3 therap*
- 21 early n3 interven* or early n3 treat* or early n3 therap* or early n3 care or early n3 medicine* or early n3 technique* or early n3 strateg* or early n3 activit* or early n3 mobili*
- 22 earli* n3 interven* or earli* n3 treat* or earli* n3 therap* or earli* n3 care or earli* n3 medicine* or earli* n3 technique* or earli* n3 strateg* or earli* n3 activit* or earli* n3 mobili*
- 23 immediat* n3 interven* or immediat* n3 treat* or immediat* n3 therap* or immediat* n3 care or immediat* n3 medicine* or immediat* n3 technique* or immediat* n3 strateg* or immediat* n3 activit* or immediat* n3 mobili*
- 24 initial* n3 interven* or initial* n3 treat* or initial* n3 therap* or initial* n3 care or initial* n3 medicine* or initial* n3 activit* or initial* n3 technique* or initial* n3 strateg* or initial* n3 mobili*
- 25 begin* n3 interven* or begin* n3 treat* or begin* n3 therap* or begin* n3 care or begin* n3 medicine* or begin* n3 technique* or begin* n3 strateg* or begin* n3 activit* or begin* n3 mobili*
- 26 first* n3 interven* or first* n3 treat* or first* n3 therap* or first* n3 care or first* n3 medicine* or first* n3 technique* or first* n3 strateg* or first* n3 activit* or first* n3 mobili*
- 27 first-line n3 interven* or first-line n3 treat* or first-line n3 therap* or first-line n3 care or first-line n3 medicine* or first-line n3 technique* or first-line n3 strateg* or first-line n3 activit* or first-line n3 mobili*
- 28 primar* n3 interven* or primar* n3 treat* or primar* n3 therap* or primar* n3 care or primar* n3 medicine* or primar* n3 technique* or primar* n3 strateg* or primar* n3 activit* or primar* n3 mobili*
- 29 original* n3 interven* or original* n3 treat* or original* n3 therap* or original* n3 care or original* n3 medicine* or original* n3 technique* or original* n3 strateg* or original* n3 activit* or original* n3 mobili*
- 30 precede* n3 interven* or precede* n3 treat* or precede* n3 therap* or precede* n3 care or precede* n3 medicine* or precede* n3 technique* or precede* n3 strateg* or precede* n3 activit* or precede* n3 mobili*
- 31 walk or walks or walking
- 323 mobili?ation strateg*
- 33 ambulate* or ambulation* or ambulating*
- 34 exerci* n3 rehab* or exerci* n3 habilitat* or exerci* n3 recover* or exerci* n3

- therap* or exerci* n3 treat* or exerci* n3 medicine* or exerci* n3 intervention* or exerci* n3 technique* or exerci* n3 strateg*
- 35 walk* n3 rehab* or walk* n3 habilitat* or walk* n3 recover* or walk* n3 therap* or walk* n3 treat* or walk* n3 medicine* or walk* n3 intervention* or walk* n3 technique* or walk* n3 strateg*
- 36 mov* n3 rehab* or mov* n3 habilitat* or mov* n3 recover* or mov* n3 therap* or mov* n3 treat* or mov* n3 medicine* or mov* n3 intervention* or mov* n3 technique* or mov* n3 strateg*
- 37 motor* n3 rehab* or motor* n3 habilitat* or motor* n3 recover* or motor* n3 therap* or motor* n3 treat* or motor* n3 medicine* or motor* n3 intervention* or motor* n3 technique* or motor* n3 strateg*
- 38 physi* n3 rehab* or physi* n3 habilitat* or physi* n3 recover* or physi* n3 therap* or physi* n3 treat* or physi* n3 medicine* or physi* n3 intervention* or physi* n3 technique* or physi* n3 strateg*
- 39 extend* n2 care* n3 facilit* or extend* n2 care* n3 service* or extend* n2 care* n3 unit* or extend* n2 care* n3 center* or extend* n2 care* n3 clinic* or extend* n2 care* n3 program* or extend* n2 care* n3 residen* or extend* n2 care* n3 home* or extend* n2 care* n3 hous*
- 40 residen* n3 facilit* or residen* n3 care* or residen* n3 service* or residen* n3 unit* or residen* n3 center* or residen* n3 clinic* or residen* n3 program* or residen* n3 residen* or residen* n3 home* or residen* n3 hous*
- 41 intermediate* n3 facilit* or intermediate* n3 care* or intermediate* n3 service* or intermediate* n3 unit* or intermediate* n3 center* or intermediate* n3 clinic* or intermediate* n3 program* or intermediate* n3 residen* or intermediate* n3 home* or intermediate* n3 hous*
- 42 assist* liv* n3 facilit* or assist* liv* n3 care* or assist* liv* n3 service* or assist* liv* n3 unit* or assist* liv* n3 center* or assist* liv* n3 clinic* or assist* liv* n3 program* or assist* liv* n3 residen* or assist* liv* n3 home* or assist* liv* n3 hous*
- 43 halfway n3 home* or halfway n3 hous* or halfway n3 facilit* or halfway n3 care* or halfway n3 residen* or halfway n3 service* or halfway n3 unit* or halfway n3 center* or halfway n3 clinic* or halfway n3 program*
- 44 transition* n3 home* or transition* n3 hous* or transition* n3 facilit* or transition* n3 care* or transition* n3 residen* or transition* n3 service* or transition* n3 unit* or transition* n3 center* or transition* n3 clinic* or transition* n3 program*
- 45 nurs* n2 home* or geriatr*-orthop* or orthop?edic-geriatr* or ortho*-geriatr* or orthogeriatr* or goru or orthop* n2 geriatr* or rehabilitation unit* or mixed assessment or maru
- 46 geriatric hip fracture program* or ghfp or day n1 hospital* or day n1 care or day n1 unit* or home-based n1 care or home based n1 care or carer* involve* or esd or early supported discharge or sequential care
- 47 S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11 or S12 or S13 or S14 or S15 or S16 or S17 or S18 or S19 or S20 or S21 or S22 or S23 or S24 or S25 or S26 or S27 or S28 or S29 or S30 or S31 or S32 or S33 or S34 or S35 or S36 or S37 or S38 or S39 or S40 or S41 or S42 or S43 or S44 or S45 or S46

1

Rehabilitation terms - OVID Embase

- 1 exp Rehabilitation/ or exp Rehabilitation Nursing/ or exp daily life activity/
 2 assisted living facility/ or nursing home/ or pain clinic/ or rehabilitation center/ or residential home/ or halfway house/
 3 day hospital/ or home care/ or home health agency/ or home physiotherapy/ or home rehabilitation/ or patient care/ or patient care planning/ or rehabilitation care/
 4 exp mobilization/ or exp Occupational Therapy/ or exp Physiotherapy/ or exp

- kinesiotherapy/ or walking/
 5 exp clinical pathway/ or social care/ or caregiver support/ or social support/ or
 caregiver/
 6 (rehab\$ or habilitat\$ or recover\$).ti,ab.
 7 (multidisciplinar\$ or interdisciplinar\$ or multiprofessional\$ or multimodal\$ or mdt
 or mdr).ti,ab.
 8 (social adj1 (work\$ or support or care)).ti,ab.
 9 (pain clinic\$ or pain service\$ or pain relief unit\$ or (pain center\$ or pain
 centre\$)).ti,ab.
 10 ((treatment\$ or therap\$ or training or education\$ or healthcare) adj10 (program\$
 or intervention\$ or approach\$)).ti,ab.
 11 (early adj1 (mobil\$ or discharg\$ or ambulat\$)).ti,ab.
 12 (occupational therap\$ or physical therap\$ or physiotherap\$ or physio).ti,ab.
 13 (exercis\$ adj3 therap\$).ti,ab.
 14 ((early or earli\$ or immediat\$ or initial\$ or begin\$ or first\$ or first-line or first line
 or first choice or primar\$ or precede\$ or original\$) adj3 (interven\$ or treat\$ or
 therap\$ or care or medicine\$ or technique\$ or strateg\$ or activit\$ or
 mobili\$)).ti,ab.
 15 (walk or walks or walking).ti,ab.
 16 mobili?ation strateg\$.ti,ab.
 17 (ambulate\$ or ambulation\$ or ambulating\$).ti,ab.
 18 (exerci\$ adj3 (rehab\$ or habilitat\$ or recover\$ or therap\$ or treat\$ or medicine\$ or
 intervention\$ or technique\$ or strateg\$)).ti,ab.
 19 ((walk\$ or mobil\$ or mov\$ or motor\$ or physi\$) adj3 (rehab\$ or habilitat\$ or
 recover\$ or therap\$ or treat\$ or medicine\$ or intervention\$ or technique\$ or
 strateg\$)).ti,ab.
 20 (extend\$ adj2 care\$ adj3 (facilit\$ or service\$ or unit\$ or center\$ or clinic\$ or
 program\$ or residen\$ or home\$ or hous\$)).ti,ab.
 21 ((residen\$ or intermediate\$ or assist\$ liv\$) adj3 (facilit\$ or care\$ or service\$ or
 unit\$ or center\$ or clinic\$ or program\$ or residen\$ or home\$ or hous\$)).ti,ab.
 22 ((halfway or transition\$) adj3 (home\$ or hous\$ or facilit\$ or care\$ or residen\$ or
 service\$ or unit\$ or center\$ or clinic\$ or program\$)).ti,ab.
 23 (nurs\$ adj2 home\$).ti,ab.
 24 (geriatr\$-orthop\$ or orthop?edic-geriatr\$ or ortho\$-geriatr\$ or orthogeriatr\$ or
 goru).ti,ab.
 25 (orthop\$ adj2 geriatr\$).ti,ab.
 26 rehabilitation unit\$.ti,ab.
 27 (mixed assessment or maru).ti,ab.
 28 (geriatric hip fracture program\$ or ghfp).ti,ab.
 29 (day adj (hospital\$ or care or unit\$)).ti,ab.
 30 ((home-based or home based) adj care).ti,ab.
 31 carer\$ involve\$.ti,ab.
 32 (esd or early supported discharge).ti,ab.
 33 sequential care.ti,ab.
 34 or/1-33

1

Rehabilitation terms - OVID Medline

- 1 exp rehabilitation/ or exp rehabilitation nursing/ or "Recovery of Function"/ or
 Subacute Care/
 2 exp rehabilitation centers/ or Residential Facilities/ or Assisted Living Facilities/ or
 Halfway Houses/
 3 Day Care/ or home care services/ or home care services, hospital-based/ or home

- nursing/ or Hospital Units/
 4 exp Nursing Homes/ or exp Patient Care Team/ or exp Patient Care Management/
 or exp Occupational Therapy/ or exp Physical Therapy Techniques/ or exp Physical
 Therapy Department, Hospital/
 5 exp "Physical Therapy (Specialty)"/ or exp Critical Pathways/ or exp Therapy,
 Computer-Assisted/ or exp Exercise Therapy/ or exp Walking/
 6 exp Social Work/ or exp Social Support/ or exp Pain Clinics/ or exp Patient
 Education/ or exp Health Education/ or Caregivers/
 7 (rehab\$ or habilitat\$ or recover\$).ti,ab.
 8 (multidisciplinar\$ or interdisciplinar\$ or multiprofessional\$ or multimodal\$ or mdt
 or mdr).ti,ab.
 9 (social adj1 (work\$ or support or care)).ti,ab.
 10 (pain clinic\$ or pain service\$ or pain relief unit\$ or (pain center\$ or pain
 centre\$)).ti,ab.
 11 ((treatment\$ or therap\$ or training or education\$ or healthcare) adj10 (program\$
 or intervention\$ or approach\$)).ti,ab.
 12 (early adj1 (mobil\$ or discharg\$ or ambulat\$)).ti,ab.
 13 (occupational therap\$ or physical therap\$ or physiotherap\$ or physio).ti,ab.
 14 (exercis\$ adj3 therap\$).ti,ab.
 15 ((early or earli\$ or immediat\$ or initial\$ or begin\$ or first\$ or first-line or first line
 or first choice or primar\$ or precede\$ or original\$) adj3 (interven\$ or treat\$ or
 therap\$ or care or medicine\$ or technique\$ or strateg\$ or activit\$ or
 mobili\$)).ti,ab.
 16 (walk or walks or walking).ti,ab.
 17 mobilization strateg\$.ti,ab.
 18 (ambulate\$ or ambulation\$ or ambulating\$).ti,ab.
 19 (exerci\$ adj3 (rehab\$ or habilitat\$ or recover\$ or therap\$ or treat\$ or medicine\$ or
 intervention\$ or technique\$ or strateg\$)).ti,ab.
 20 ((walk\$ or mobil\$ or mov\$ or motor\$ or physi\$) adj3 (rehab\$ or habilitat\$ or
 recover\$ or therap\$ or treat\$ or medicine\$ or intervention\$ or technique\$ or
 strateg\$)).ti,ab.
 21 (extend\$ adj2 care\$ adj3 (facilit\$ or service\$ or unit\$ or center\$ or clinic\$ or
 program\$ or residen\$ or home\$ or hous\$)).ti,ab.
 22 ((residen\$ or intermediate\$ or assist\$ liv\$) adj3 (facilit\$ or care\$ or service\$ or
 unit\$ or center\$ or clinic\$ or program\$ or residen\$ or home\$ or hous\$)).ti,ab.
 23 ((halfway or transition\$) adj3 (home\$ or hous\$ or facilit\$ or care\$ or residen\$ or
 service\$ or unit\$ or center\$ or clinic\$ or program\$)).ti,ab.
 24 (nurs\$ adj2 home\$).ti,ab.
 25 (geriatr\$-orthop\$ or orthop?edic-geriatr\$ or ortho\$-geriatr\$ or orthogeriatr\$ or
 goru).ti,ab.
 26 (orthop\$ adj2 geriatr\$).ti,ab.
 27 rehabilitation unit\$.ti,ab.
 28 (mixed assessment or maru).ti,ab.
 29 (geriatric hip fracture program\$ or ghfp).ti,ab.
 30 (day adj (hospital\$ or care or unit\$)).ti,ab.
 31 ((home-based or home based) adj care).ti,ab.
 32 carer\$ involve\$.ti,ab.
 33 (esd or early supported discharge).ti,ab.
 34 sequential care.ti,ab.
 35 or/1-34

1 Surgeon seniority

Surgeon seniority terms – Cochrane Library

- 1 MeSH descriptor Clinical Competence explode all trees
- 2 (surgeon* NEAR/3 (senior* or experience* or supervision* or volume* or grade*)):ti,ab,kw
- 3 (consultant* or registrar* or spr or staff grade or trust grade or associate specialist*):ti,ab,kw
- 4 (surg* NEAR (team* or list*)):ti,ab,kw
- 5 (list* NEAR (organise* or organize* or consultant-led or consultant led)):ti,ab,kw
- 6 #1 or #2 or #3 or #4 or #5

2

Surgeon seniority terms – EBSCO CINAHL

- 1 surgeon* n3 senior* or surgeon* n3 volume* or surgeon* n3 supervision* or surgeon* n3 experience* or surgeon* n3 grade* or surg* n1 team* or surg* n1 list* or list* n1 organise* or list* n1 organize* or list* n1 consultant-led or list* n1 consultant led
- 2 consultant* or spr or registrar* or staff grade or trust grade or associate specialist* or mh clinical competence+
- 3 S1 or S2

3

Surgeon seniority terms - OVID Embase

- 1 exp clinical competence/
- 2 (surgeon\$ adj3 (senior\$ or experience\$ or supervision\$ or volume\$ or grade\$)).ti,ab.
- 3 (consultant\$ or registrar\$ or spr or staff grade or trust grade or associate specialist\$).ti,ab.
- 4 (surg\$ adj1 (team\$ or list\$)).ti,ab.
- 5 (list\$ adj1 (organise\$ or organize\$ or consultant-led or consultant led)).ti,ab.
- 6 or/1-5

4

Surgeon seniority terms - OVID Medline

- 1 Clinical Competence/
- 2 (surgeon\$ adj3 (senior\$ or experience\$ or supervision\$ or volume\$ or grade\$)).ti,ab.
- 3 (consultant\$ or registrar\$ or spr or staff grade or trust grade or associate specialist\$).ti,ab.
- 4 (surg\$ adj1 (team\$ or list\$)).ti,ab.
- 5 (list\$ adj1 (organise\$ or organize\$ or consultant-led or consultant led)).ti,ab.
- 6 or/1-5

5

6 Surgical Interventions

Surgical Interventions terms – Cochrane Library

- 1 MeSH descriptor Fracture Fixation, Internal explode all trees
- 2 MeSH descriptor Internal Fixators explode all trees
- 3 MeSH descriptor Bone Nails explode all trees
- 4 MeSH descriptor Bone Screws explode all trees

5 MeSH descriptor Bone Plates explode all trees
 6 MeSH descriptor Bone Cements explode all trees
 7 MeSH descriptor Arthroplasty explode all trees
 8 (pin* or nail* or screw* or plate* or arthroplast* or fix* or prosthes* or ((cement*
 or glue* or paste*) NEAR/3 bone*)):ti,ab,kw
 9 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8

1

Surgical interventions terms - OVID Embase

1 (pin\$ or nail\$ or screw\$ or plate\$ or arthroplast\$ or hemiarthroplast\$ or fix\$ or
 prosthes\$).ti,ab.
 2 arthroplasty/ or hip arthroplasty/
 3 ((cement\$ or glue\$ or paste\$) adj3 bone\$).ti,ab.
 4 Fracture Treatment/ or Hip Surgery/ or Femur Intertrochanteric Osteotomy/ or Hip
 Osteotomy/ or exp Fracture Fixation/ or Bone Screw/ or Bone Plate/ or Bone Nail/
 or ender Nail/ or Interlocking Nail/ or Osteosynthesis Material/ or external fixator/
 or exp bone cement/
 5 or/1-4

2

Surgical interventions terms - OVID Medline

1 (pin\$1 or nail\$ or screw\$1 or plate\$1 or arthroplast\$ or fix\$ or prosthes\$).ti,ab.
 2 Internal Fixators/ or Bone Screws/ or Fracture Fixation, Internal/ or Bone Plates/ or
 Bone Nails/ or Bone Cements/
 3 ((cement\$ or glue\$ or paste\$) adj3 bone\$).ti,ab.
 4 Arthroplasty/ or Arthroplasty, Replacement, Hip/
 5 or/1-4

3

4 Systematic review filter

Systematic review filter - OVID Medline

1 meta-analysis/
 2 (metaanalys\$ or meta-analys\$ or meta analys\$).tw.
 3 exp "review literature"/
 4 (systematic\$ adj3 (review\$ or overview\$)).tw.
 5 (selection criteria or data extraction).ab. and review.pt.
 6 (cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or
 cinhal or science citation index or bids or cancerlit).ab.
 7 (reference list\$ or bibliograph\$ or hand search\$ or hand-search\$ or manual
 search\$ or relevant journals).ab.
 8 or/1-7

5

Systematic review filter - OVID Embase

1 meta analysis/
 2 (metaanalys\$ or meta-analys\$ or meta analys\$).tw.
 3 systematic review/
 4 (systematic\$ adj3 (review\$ or overview\$)).tw.
 5 (selection criteria or data extraction).ab. and Review.pt.
 6 (cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or
 cinhal or science citation index or bids or cancerlit).ab.
 7 (reference list\$ or bibliograph\$ or hand search\$ or manual search\$ or relevant
 journals).ab.

8

or/1-7

17 Appendix E: Evidence tables - Clinical studies

17.1	Evidence Table 1: Imaging options in occult hip fracture.....	293
17.2	Evidence Table 2: Timing of surgery	296
17.3	Evidence Table 3: Optimal analgesia	314
17.4	Evidence Table 4: Anaesthesia	317
17.5	Evidence Table 5: Surgeon seniority.....	320
17.6	Evidence Table 6: Displaced intracapsular fractures	324
17.7	Evidence Table 7: Surgery – Cement versus no cement	333
17.8	Evidence Table 8: Extracapsular fixation	335
17.9	Evidence Table 9: Surgical approach to hemiarthroplasty.....	383
17.10	Evidence Table 10: Mobilisation strategies	385
17.11	Evidence Table 11: Multidisciplinary rehabilitation.....	394
17.12	Evidence Table 12: Patient views	418
18.1	Evidence Table 13: General versus regional anaesthesia.....	435
18.2	Evidence Table 14: Displaced intracapsular fractures.....	437
18.3	Evidence Table 15: Cemented arthroplasties.....	441
18.4	Evidence Table 16: Multidisciplinary rehabilitation.....	443

Abbreviations

CI	Confidence interval
IQR	Interquartile range
ITT	Intention to treat analysis
LOS	Length Of Stay
LR+	Positive likelihood ratio
LR-	Negative likelihood ratio
M/F	Male/female
N	Total number of patients randomised
NA	Not Applicable
NPV	Negative predictive value
NR	Not reported
PPV	Positive predictive value
QALY	Quality-Adjusted Life Years
QoL	Quality of life
RCT	Randomised controlled trial
RR	Relative risk
SD	Standard Deviation
SE	Standard Error
Sig	Statistically significant at 5%

1 17.1 Evidence Table 1: Imaging options in occult hip fracture

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
<p>Study name: Safran et al., 2009^{296,297}</p> <p>Study design: Prospective cross-sectional study</p> <p>Duration of follow up: Not reported</p>	<p>Patient group: Patients with painful hips after low energy trauma (e.g. fall from a sitting or standing position)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Difficulty or inability to bear weight after a fall • Tenderness around the hip with painful hip motion • Negative pelvic and hip radiographic finding <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Prior ipsilateral hip fractures or surgery • Contraindications to MRI <p>All patients N: 30 Mean age (range): 73 (26-94) M/F: 6/24</p> <p>Drop outs: 0</p>	<p>Assessment tool under investigation: Sonography (HDI 5000 ultrasound device) Bilateral hips were examined and saggital, axial and coronal planes and particular attention was paid to the hip joint and greater trochanteric regions searching for fracture lines, joint and bursal effusions and peritrochaneric oedema The findings were recorded before the MRI examination</p> <p>Reference standard: MRI within 72 hours of admission on a 1.5-T Sigma scanner or a 1.5-T Avanto scanner. Scans were performed in the axial and coronal planes with a T1 weighted fast spin echo sequence and with Short Tau inversion recovery with magnitude display sequence. The scans were performed in the axial plane from the level of the anterior superior iliac spine to 5 cm below the level of the lesser trochanter. In the coronal plane, the scans were performed from the symphysis pubis to the sacrum.</p> <p>The MRI scans were read by a radiologist with 15 years experience in musculoskeletal MRI, who was blinded to the sonographic findings</p>	Sensitivity	100%	<p>Funding: Not reported</p> <p>Limitations: Sonographic examinations performed by 2 musculoskeletal radiologists who may not always be available at community hospitals</p> <p>72 hours delay before MRI was given</p> <p>The time from injury to admission ranged from 0 to 14 days (average 1.7 days)</p> <p>Notes: An overall well conducted and well reported study with low risk of bias</p>
			Specificity	65%	
			PPV	59%	
			NPV	100%	
			LR+	2.85	
			LR-	0	
			Prevalence	33%	

1 Evidence tables – imaging

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
Study name: Rizzo et al., 1993 ^{286,286} Study design: Prospective Cross sectional Duration of follow-up: 6 months	Patient group: Patients whose history and clinical examination suggestive of a hip fracture but whose radiographs were negative Inclusion/exclusion criteria: Not reported <u>All patients</u> N: 62 Mean age (range): 73 (26-93) M/F: 23/39 Drop outs: 0	Assessment tool under investigation: bone scanning 72 hours after admission using a technetium-99m bone scan Reference standard: MRI within 24 hours after admission. Only T1-weighted coronal spin-echo pulse sequences were obtained	Sensitivity	97.3%	Funding: None Limitations: Patients had MRI within 24 hours of admission whereas bone scanning was carried out 72 hours after admission Notes: 1 patient had an initial negative CT scan bit a positive MRI scan. CT scanning after 6 days showed a positive result. This patient has been considered as a false negative in this analysis
			Specificity	100%	
			PPV	100	
			NPV	95.8	
			LR+	0	
			LR-	0.02	
			Prevalence	60	

2
3

1 Evidence tables – imaging

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
Study name: Evans et al., 1994 ^{87,88} Study design: Prospective cross sectional study Duration of follow-up: 3 months	Patient group: Elderly patients admitted to hospital with hip pain after a fall and whose radiographs were normal or showed a fracture of the greater trochanter Inclusion/exclusion criteria: Not reported <u>All patients</u> N: 37 Mean age (range): not reported Drop outs: 0	Assessment tool under investigation: Isotope scanning Technetium 99m, 48 hours after MRI scan Reference standard: MRI, 5 minute sequence of T1-weighted coronal images. Where necessary Short tau inversion recovery and/or T2 weighted images were also obtained	Sensitivity	75%	Funding: None Limitations: Relatively small patient numbers Isotope scans given 48 hours after the fall to avoid false positives Not clear who interpreted the results and whether they were blind to the results of the reference standard test Authors did not report any information on patient demographics Notes:
			Specificity	100%	
			PPV	100	
			NPV	93	
			LR+	0	
			LR-	0.25	
			Prevalence	22	

2
3

1 17.2 Evidence Table 2: Timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Alani et al., 2008⁴</p> <p>Country of study: Sweden</p> <p>Study design: <i>Prospective cohort</i></p> <p>Duration of follow-up: Hospital stay</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Danderd and Huddinge hospitals, Stockholm, Sweden.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients with acute hip fracture aged 50 years or older <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients with a pathological fracture and patients who arrived at the hospital one calendar day after the time of injury. <p>All patients N: 744 Lost to follow up: 22 patients (missing data for return to independent living) Age (mean \pmSD): 81 M/F: 200/544 Diagnosis of dementia: 209 (28%) N for time to surgery: ≤ 24h = 359 > 24h = 385 ≤ 36h = 550 > 36h = 194</p>	<p>Group 1 Early surgery. ≤ 48 hours</p> <p>Group 2 Late surgery. > 48 hours</p>	<p>Return to independent living Adjusted odds ratio adjusted for age, sex, prefracture walking ability, whether patient was living with someone, ASA score, treatment modality, reoperation, and reason for delay of surgery.</p> <p>Pressure ulcers Adjusted odds ratio adjusted for age, prefracture walking ability, dementia, ASA score, and duration of surgery.</p>	<p>Unadjusted (patients without dementia): Group 1: 320/375 Group 2: 43/59 Missing data: 22 (5%)</p> <p><24 hours: 178/209 ≥ 24 hours: 185/225 Missing data: 22 (5%)</p> <p><36 hours: 282/329 ≥ 36 hours: 81/105 Missing data: 22 (5%)</p> <p>Adjusted odds ratio: Delay > 24h: 0.86 (0.45 to 1.65) NS Delay > 36h: 0.44 (0.21 to 0.90) $P < 0.05$ Delay > 48h: 0.33 (0.14 to 0.78) $P < 0.01$</p> <p>Unadjusted: Group 1: 41/646 Group 2: 20/98</p> <p><24 hours: 53/354 ≥ 24 hours: 60/345 $p < 0.05$</p> <p><36 hours: 31/550 ≥ 36 hours: 30/194 $p < 0.0001$</p>	<p>Funding: One or more authors received, in any one year, outside funding or grants in excess of \$10,000 from the Stockholm County Council Research Fund for clinical studies. No benefits received from commercial entities.</p> <p>Limitations: Impact of comorbidity on mortality (unadjusted data).</p> <p>Additional outcomes reported: None</p> <p>Notes: None</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>≤48 = 646 >48 = 98</p> <p>Group 1 Early No.: 646 No. of dropouts: not stated Age (mean): 81 M/F: 166/480 Other factors: Diagnosis of dementia: 181 (28%)</p> <p>Group 2 Late No. : 98 No. of dropouts: not stated Age (mean): 81 M/F: 34/64 Other factors: Diagnosis of dementia: 28 (29%) Delay due to: Patient related (e.g. medical): 57 (58%) System related (e.g. no available operating room): 41 (42%)</p>			<p>Adjusted odds ratio: Delay >24h: 2.19 (1.21 to 3.96) P<0.01 Delay >36 hours: 3.42 (1.94 to 6.04) P<0.001 Delay >48 hours: 4.34 (2.34 to 8.04) P<0.001</p>	
			Length of hospital stay – median (including rehab)	<p>Unadjusted: Group 1: 15 Group 2: 21</p> <p><24 hours: 14 ≥24 hours: 18 p <0.001</p> <p><36 hours: 15 ≥36 hours: 19 p <0.001</p>	
			Length of hospital stay – median (including rehab), excluding days prior to surgery	<p>Unadjusted: Group 1: 13 Group 2: 16 p <0.01</p> <p><24 hours: 14 ≥24 hours: 17 p <0.05</p> <p><36 hours: 15 ≥36 hours: 18 p <0.05</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mortality rate – 4 months Adjusted odds ratio adjusted for age, sex, prefracture walking ability, dementia and ASA score.	Adjusted odds ratio: Delay >24h: 1.07 (0.67 to 1.70) NS Delay >36h: 1.05 (0.63 to 1.74) NS Delay >48h: 0.86 (0.44 to 1.69) NS	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Diabetes: 16.4% Anticoagulation: 6.6% Chronic renal dialysis: 2.1% Active cancer: 2.1% Cirrhosis: 0.3% Fall occurred at: Home: 58.2% Nursing home: 21.5% Outdoor: 19% In-hospital: 1.2% Time of surgery: <24h: 523 24-48h: 325 >48h: 129 <u>Group 1 Early</u> No.: 848 No. of dropouts: not stated Age (mean): <24 hrs: 79 24-48 hrs: 80 M/F: <24 hrs: 25%/75% 24-48 hrs: 21.5%/78.5% Sever complications: 17.2% Dementia: 308/848 <u>Group 2 Late</u> No. : 129 No. of dropouts: not stated Age (mean): 80 M/F: 24%/76% Sever complications: 24.8% Dementia: 49/129			Group 1: <24 hrs: 16 24-48 hrs: 15 Group 2: 20 All Group 1: 147/848 Group 2: 40/129 <24 hours: 88/523 ≥24 hours: 90/454 Adjusted Odds ratio: 24-48hs (vs. 24h): 0.87 (0.58-1.29) >48 hours (vs. 24h): 1.32 (0.79-2.20)	

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bottle et al., 2006^{30,31}</p> <p>Country of study: England</p> <p>Study design: <i>Retrospective cohort</i></p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: NHS hospital trusts in England with at least 100 admissions for fractured neck of femur</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients aged ≥65 admitted with a primary diagnosis of fractured neck of femur admitted from their own home. Patients with a first hip fracture only were included. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients admitted from nursing and residential homes <p>All patients N: 114,942</p> <p>Group 1 Early No.: 90551 No. of dropouts: not stated Age (mean ±SD): not stated</p> <p>Group 2 Late No. : 24391 No. of dropouts: not stated Age (mean ±SD): not stated</p>	<p>Patients underwent one of 4 types of surgery: fixation, prosthetic replacement of head of femur, other procedure (including non-orthopaedic) and no procedure recorded (medical management).</p> <p>Group 1 Early surgery. < 2days</p> <p>Group 2 Late surgery. > 2 days</p>	<p>30 day mortality</p> <p>30 day mortality Adjusted Odds ratios (adjusted for age, sex, deprivation fifth and comorbidity)</p> <p>Emergency readmission within 28 days (adjusted for age, sex, deprivation fifth and comorbidity)</p>	<p>Group 1: 6366/90551 Group 2: 2625/24391</p> <p>>1 day vs. ≤1 day: 1.25 (1.19 to 1.31) >2 day vs. ≤2 day: 1.36 (1.29 to 1.43)</p> <p>>1 day vs. ≤1 day: 1.04 (0.99 to 1.08) >2 day vs. ≤2 day: 1.04 (0.99 to 1.10)</p>	<p>Funding: The unit is funded by a grant from Dr Foster Ltd (an independent health service research organisation).</p> <p>Limitations: Baseline characteristics given for entire cohort, which includes patients who did not receive surgery.</p> <p>Additional outcomes reported: Adjusted effect of operative delay on mortality, excess risk of death</p>

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Grimes et al.,2002 ¹²⁵ Country of study: USA Study design: <i>Retrospective cohort</i> Duration of follow-up: 5 – 10 years	Patient group: Patients with hip fracture Setting: 20 hospitals in New Brunswick, New Jersey; San Antonio, Texas; Philadelphia, Pennsylvania; and Richmond, Virginia – and represented university, community, and Veterans Affairs medical centers. Inclusion criteria: <ul style="list-style-type: none"> Consecutive patients with hip fracture who were aged 60 years or older and who underwent surgical repair between 1983 and 1993. Exclusion criteria: <ul style="list-style-type: none"> Patients were excluded if they had metastatic cancer, trauma resulting in multiple injuries requiring surgery, or declined blood transfusion for religious reasons. Patients with a fracture occurring >48 hours before admission to the hospital. All patients N: 8383 Lost to follow up: Not stated	Time from admission to surgery. Group 1 Early surgery Group 2 Late surgery	30 day mortality	Group 1: 175 Group 2: Active medical problems: 56 No medical problems:166	Funding: Not stated Limitations: No baseline data provided
			30 day mortality (adjusted odds ratio)	>48-72h: 0.71 (0.45-1.10) n = 3805	
			Decubitus Ulcer (adjusted odds ratio)	>48-72h: 1.2 (0.9-1.6) n = 3579	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean \pmSD): 80.4 \pm8.6 M/F: 1751/6632</p> <p><u>Group 1 Early (\leq 24 hours)</u> No.: 4578 No. of dropouts: not stated Age (mean \pmSD): 60-69: 590 70-79: 1356 80-89: 1972 \geq90: 3683 M/F: 895/3683 Other factors: ASA class: 1 or 2: 1341 3: 2852 4 or 5: 385</p> <p><u>Group 2 Late (\geq 24 hours)</u> No. : 3805 No. of dropouts: not stated Age (mean \pmSD): 60-69: 485 70-79: 1089 80-89: 1683 \geq90: 549 M/F: 858/2949 Other factors: ASA class: 1 or 2: 974 3: 2279 4 or 5: 552</p>				

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Lefaiivre et al., 2009^{189,189}</p> <p>Country of study: Canada</p> <p>Study design: <i>Retrospective cohort</i></p> <p>Duration of follow-up: In hospital</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Vancouver General Hospital</p> <p>Inclusion criteria: All patients over the age of 65 who had been admitted with an isolated fracture of the proximal femur between 1998 and 2001.</p> <p>All patients N: 607 M/F: 125/482</p> <p>Delay to surgery <24h: 245 24 to 48: 264 >48: 98</p> <p>Age: <75: 102, 76 – 85: 262 86 – 95: 212, 96 – 105: 30 106 – 115: 1</p> <p>Medical comorbidities: 0: 141 1 to 2: 405 ≥3: 61</p>	<p>Pre-existing medical comorbidity was quantified by listing the pre-injury medical diagnoses by a body system such as cardiac, pulmonary, autoimmune, substance dependence etc. Patients were categorised into no major comorbidity, those with one to two body systems with major comorbidity and those with ≥3 body systems with major comorbidities.</p>	<p>Logistic regression model (adjusted for medical comorbidity age, gender and fracture type)</p> <p>24 to 48h</p> <p>Odds ratio (95% CI)</p>	<p>Death 0.82 (0.42 to 1.62) p = 0.5713</p> <p>Major medical complication 0.96 (0.52 to 1.75) p = 0.8868</p> <p>Minor medical complication 1.53 (1.05 to 2.22) p = 0.0257</p> <p>Pressure sores 1.23 (0.71 to 2.12) p = 0.4700</p>	<p>Funding: None</p> <p>Limitations:</p> <p>Notes: 690 patients added to the database, of these they were only able to review the complete medical records of 607 patients.</p>
			<p>Logistic regression model (adjusted for medical comorbidity age, gender and fracture type)</p> <p>> 48h</p>	<p>Death 0.93 (0.38 to 2.33) p = 0.8840</p> <p>Major medical complication 2.21 (1.01 to 4.34) p = 0.0260</p> <p>Minor medical complication 2.27 (1.38 to 3.72) p = 0.0012</p> <p>Pressure sores 2.29 (1.19 to 4.40) p = 0.0128</p>	

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Majumdar et al., 2006^{200,200}</p> <p>Country of study: Canada</p> <p>Study design: <i>Retrospective cohort</i></p> <p>Duration of follow-up: 30 days</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Tertiary care hospitals in Edmonton, Alberta, Canada</p> <p>Inclusion criteria: Consecutive patients with hip fracture during March 1994 to February 2000 Patients aged 60 years or older Hip fracture patients included femoral neck, intertrochanteric, subtrochanteric or subcapital fractures.</p> <p>Exclusion criteria: Patients with multiple traumatic fractures, pathologic hip fractures, or bilateral hip fractures.</p> <p>All patients N: 3981 (3846 – had surgery) Age (mean ±SD): 82 (±8.52) M/F: 1154/2827 Time of surgery: <24h: 1048 24 – 48h: 2152 >48h: 664</p> <p>Group 1 Early No.: 3200</p>	<p>Timing of surgery was based on the calendar date of hospital admission and calendar date of surgical repair.</p> <p>Group 1 Early surgery. Within 48 hours of admission</p> <p>Group 2 Late surgery. After 48 hours of admission</p>	In hospital mortality	<p>Group 1: 160/3200 Group 2: 66/664</p> <p><24 hours: 5/1046 ≥24 hours: 36/2933</p> <p>Adjusted odds ratio: 24 -48hr vs. <24: 0.90 (0.85-1.99) P = 0.59</p> <p>>48hr vs. <24h: 1.30 (0.86-2) p = 0.21</p>	<p>Funding: None</p> <p>Limitations: Adjusted odds ratios compare <24h to >48h time to surgery.</p> <p>Additional outcomes reported: Type of fracture, % with dementia, prefracture comorbidities</p> <p>Notes:</p>
			1 year mortality	<p>Group 1: 970/3200 Group 2: 219/664</p> <p><24 hours: 5/1046 ≥24 hours: 35/497</p>	
			Length of stay (after surgery) (in days, median, with interquartile range)	<p>Group 1: <24h: 7 (1-13) 24-48h: 8 (2-14) Group 2: 11 (0-24)</p>	
			Complications (Myocardial infarction, heart failure, cardiac arrhythmia, electrolytes abnormal, anaemia, pneumonia, urinary tract infection).	<p>Group 1: 614/3200 Group 2: 130/664</p> <p><24 hours: 235/1046 ≥24 hours: 509/497</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>No. of dropouts: not stated Age (mean \pmSD): 82 M/F: 892/2308</p> <p><u>Group 2 Late</u> No. : 664 No. of dropouts: not stated Age (mean \pmSD): 81 M/F: 214/450</p>				

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Moran et al., 2005^{215,215}</p> <p>Country of study: UK</p> <p>Study design: <i>Prospective cohort</i></p> <p>Duration of follow-up: 30 days</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: University hospital Nottingham</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> All adult patients with a fracture of the femoral neck. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Isolated femoral head fractures and acetabular fractures 140 patients who did not have surgery were excluded <p>All patients N: 2148</p> <p>Lost to follow up: Age (mean \pmSD): 80 M/F: 684/2219</p> <p>Group 1 Early No.: 982 No. of dropouts: not stated Age (mean \pmSD): not stated</p> <p>Group 2 Late No. : 1166 No. of dropouts: not stated Age (mean \pmSD): not stated</p>	<p>Group 1 Early surgery. No delay, surgery performed in less than one day of admission</p> <p>Group 2 Late surgery. Surgery after 1 day or more from admission</p>	<p>30 day mortality of patients fit for surgery:</p>	<p>No delay: 85/982 Delay 1 day: 85/1166 $p = 0.51$</p> <p>No delay: 134/1651 Delay 2 day: 36/497</p> <p>No delay: 158/1978 Delay 3 day: 12/170</p> <p>No delay: 166/2092 Delay 4 day: 4/56</p>	<p>Funding: Not stated</p> <p>Limitations: No protocol for determining which patients were unfit for surgery and anaesthesia, therefore variation between clinicians.</p> <p>Notes: Delay to surgery was most frequently due to acute medical comorbidity (206 patients). The subgroup of patients who were fit for surgery is given; any delay here is due to logistical reasons.</p>

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Orosz et al., 2004^{250,250}</p> <p>Country of study: USA</p> <p>Study design: Prospective cohort</p> <p>List who was masked to interventions: Nurses identifying complications were not aware of the study hypothesis, but physicians categorising complications were not blinded.</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: 4 hospitals in the New York City metropolitan area (an academic medical centre, an urban teaching hospital, and a suburban hospital)</p> <p>Inclusion criteria: Patients with hip fracture aged 50 and over.</p> <p>Exclusion criteria: Patients aged younger than 50 years, fractures that occurred as an inpatient, transfers from another hospital, multiple trauma, pathological fractures, distal and femoral shaft fractures, bilateral hip fractures, or previous fracture or surgery on the currently fractured site.</p> <p>All patients N: 1203 Age (mean \pmSD): M/F:</p> <p>Group 1 Early No.: 398 No. of dropouts: not stated</p>	<p>Patients enrolled as early in the admission as possible (69% on or before the day of surgery).</p> <p>Group 1 Surgery within 24 hours</p> <p>Group 2 Surgery after 24 hours</p> <p>Adjustments to odd ratios were based on age, sex, nursing home residence, needing a proxy for consent, delirium on admission, prefracture FIM locomotion score, fracture type, history of diabetes, COPD, stroke syndrome, dementia, cardiac disease,</p>	<p>Major postoperative complications (those that pose a threat to life or bodily functions and that typically are treated with parenteral medications, procedures, or intensive monitoring e.g. pneumonia or arrhythmias. Data for patients enrolled in 1st 12 months only.</p>	<p>Adjusted OR = 0.26 (0.07to 0.95) p = 0.04</p>	<p>Funding: Grants were received from the Agency for Healthcare Research and Quality</p> <p>Limitations: Baseline data given for study arms, but not for reported separately for the restricted cohort.</p> <p>Additional outcomes reported:</p> <p>Notes: Restricted cohort excluded patients who might not be candidates for early surgery because of markedly abnormal clinical findings or the need for additional time for preoperative evaluation. This, the restricted cohort excludes patients admitted with</p>
			<p>Mean pain scores over the first 5 hospital days. Data for patients enrolled in 1st 12 months only. Score from 1 (none) - 5 (very severe pain).</p>	<p>Group 1: 2.52 Group 2: 2.90</p> <p>Difference (95% CI) = -0.38 (-0.61 to -0.16) p = 0.001</p>	
			<p>Number of days of severe pain over hospital days 1-5 (assessed by asking if they were experiencing no pain, or mild, moderate or severe pain). Data for patients enrolled in 1st 12 months only.</p>	<p>Group 1: 0.50 Group 2: 0.80</p> <p>Difference (95% CI) = -0.30 (-0.50 to -0.08) p = 0.007</p>	
			<p>Length of stay, mean stay in days and adjusted odds ratio</p>	<p>Group 1: 6.94 Group 2: 7.85</p> <p>Difference (95% CI) = -0.91 (-1.81 to -</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean \pmSD): 82 (9.2) M/F: 82/316 Delirium at admission: 10 Admitted from nursing home : 63</p> <p>Group 2 Late No. : 780 No. of dropouts: Age (mean \pmSD): 82 (8.6) M/F: 147/633 Delirium at admission: 20 Admitted from nursing home : 90 The restricted cohort is a subset of the groups shown above, which is described in the notes section.</p>	hypertension, hospitalisation within 6 months, hospital site, day and time of admission and abnormal clinical findings.		0.01) p = 0.05	abnormal clinical findings, aortic stenosis, dementia, and endstage renal disease on dialysis.
			FIM locomotion score at 6 months (2-item subscale focusing on walking and climbing stairs)	<p>Group 1: 9.94 Group 2: 9.97</p> <p>Difference (95% CI) = -0.03 (-0.60 to 0.54) p = 0.91</p>	
			FIM self care (6 item scale of self-care activities including bathing and dressing)	<p>Group 1: 34.8 Group 2: 35.4</p> <p>Difference (95% CI) = -0.60 (-1.98 to 0.65) p = 0.32</p>	
			FIM transferring (3 item scale focusing on transfers from the bed, toilet and bath tub)	<p>Group 1: 15.7 Group 2: 15.7</p> <p>Difference (95% CI) = 0 (-0.64 to 0.77) p = 0.85</p>	
			Dead or needing total assistance in locomotion at 6 months	Adjusted OR = 0.62 (0.35 to 1.08) p = 0.09	

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Siegmeth et al., 2005A ^{308,308} Country of study: England Study design: <i>Prospective cohort</i> Duration of follow-up: 1 year	Patient group: Patients with hip fracture Setting: Peterborough District Hospital Inclusion criteria: Patients with hip fracture admitted to the Peterborough Hip fracture service Exclusion criteria: <ul style="list-style-type: none"> Patients aged younger than 60 years, those treated conservatively and those with a pathological fracture or a fracture of the shaft or distal femur. Patients who were delayed for any medical reason when orthopaedic or anaesthetic staff felt that operation should be delayed in order to improve the patient's fitness for surgery All patients N: 3628 Lost to follow up: 2 Age (mean \pmSD): 81 (8.06) Group 1 Early (\leq 48 hours) No.: 3454 Age (mean \pmSD): M/F: 656/2798	Surgical treatment involved either internal fixation with cannulated screws or hemiarthroplasty for intracapsular fixation. Those with extracapsular fractures were operated on with a dynamic hip screw or an intramedullary nail device. Group 1 Early surgery Group 2 Late surgery	Mean hospital stay in days (95% CI) (includes time spent on orthopaedic ward and any other hospital wards or convalescent units until eventual discharge to a permanent place of residence)	Group 1: 21.6 Group 2: 36.5 (5.7-16) P value(s): <0.0001	Funding: No benefits in any form were/will be received from a commercial party related directly or indirectly to the subject of the article. Limitations: Baseline data reported for 6 individual groups, but not split according to <48 or >48 hours delay. Outcomes not reported: <i>List the outcomes in which we are interested that are not reported here</i> Additional outcomes reported: N/A Notes: Delay for non-medical reasons was because of lack of operating theatre space, equipment or available staff.
			Return to original residence (%)	Group 1: 2974 (86.1%) Group 2: 128 (73.6%) P value(s): <0.0001	
			Change in residence (admitted to a more dependent accommodation)	Group 1: 240 (6.9%) Group 2: 22 (12.6%) P value(s): <0.0007	
			Mortality at 1 year	Group 1: 238 (6.9%) Group 2: 24 (13.8%) P value(s): <0.001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<u>Group 2 Late (> 48 hours)</u> No. : 174 Age (mean \pm SD): M/F: 39/135				

1 Evidence tables – timing of surgery

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Weller et al., 2005^{351,351}</p> <p>Country of study: Canada</p> <p>Study design: <i>Retrospective cohort</i></p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Inclusion criteria:</p> <ul style="list-style-type: none"> Patients aged over 50 years who were admitted to hospital in Ontario, Canada between 1993 and 1999 for surgical treatment of a hip fracture from the Canadian Institute for Health Information Discharge Abstracts Database <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Delay to surgery ≥ 7 days. <p>All patients N: 57,315</p> <p>Lost to follow up: Not stated</p> <p>Age (mean \pmSD): Men: 77.7 \pm10.2 Woman: : 81.4 \pm8.8</p> <p>M/F: 14,329/42,986</p> <p>Group 1 Early (≤ 2 days) No.: 52,937</p> <p>No. of dropouts: not stated</p> <p>Age (mean \pmSD): not stated</p> <p>M/F: not stated</p>	<p>Group 1 Early surgery < 2 days</p> <p>Group 2 Late surgery >2 days</p>	In-hospital mortality	<p>Group 1: 3509 (6.6%) Group 2: 433 (10%)</p> <p><24hr: 1177/20303 \geq24hr: 2765/37012</p> <p>Adjusted Odds Ratio: 1 day: 1.17 (1.08-1.26) 2 days: 1.36 (1.23 – 1.52) >2 days: 1.60 (1.42 to 1.80)</p>	<p>Funding: N/R</p> <p>Limitations: One aim of the study was to determine whether mortality after hip fracture is related to type of hospital (teaching or non teaching and urban or rural) in which the patient is treated.</p> <p>Notes: A modified Charlson-Deyo index was used to adjust for comorbidity. An algorithm was used in order to identify any major complications after hip fracture surgery, including infection deep vein thrombosis, intra-operative surgical complications and</p>
			3 -month mortality	<p>Group 1: 7277 (13.7%) Group 2: 790 (18%)</p> <p><24hr: 2552/20303 \geq24hr: 5515/37012</p> <p>Adjusted Odds Ratio: 1 day: 1.11 (1.05 – 1.17) 2 days: 1.27 (1.17 – 1.37) >2 days: 1.40 (1.27 to 1.53)</p>	
			6-month mortality	<p>Group 1: 9441 (17.8%) Group 2: 1038 (24%)</p> <p><24hr: 3361/20303 \geq24hr: 7118/37012</p> <p>Adjusted Odds Ratio: 1 day: 1.09 (1.04 – 1.15) 2 days: 1.20 (1.12 – 1.29) >2 days: 1.42 (1.31 to 1.55)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Other factors:</p> <p><u>Group 2 Late (> 2 days)</u> No. : 4378 No. of dropouts: not stated Age (mean \pmSD): not stated M/F: not stated</p> <p>Data given by type of hospital, not by delay to surgery.</p>		1-Year mortality	<p>Group 1: 12233 (23.1%) Group 2: 1313 (30%)</p> <p><24hr: 4366/20303 \geq24hr: 9180/37012</p> <p>Adjusted Odds Ratio: 1 day: 1.13 (1.05 – 1.22) 2 days: 1.26 (1.11 – 1.44) >2 days: 1.58 (1.26 to 1.99)</p>	significant medical complications.

1 17.3 Evidence Table 3: Optimal analgesia

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Parker et al., 2002^{262,270}</p> <p>Study design: Cochrane systematic review. The review includes 17 randomised and quasi randomised studies</p> <p>Setting: Hospitals in Europe, Turkey, South Africa and Israel.</p> <p>Duration of follow-up: Range: 24 hours-6 months. Also includes: length of</p>	<p>Patient group: Hip fracture</p> <p>Inclusion criteria: Skeletally mature patients with a proximal femoral fracture undergoing nerve blocks (including epidurals) versus no nerve blocks.</p> <p>Exclusion criteria: Not stated</p> <p>All patients N (range): 888 (19-100) Age range: 59-86 M/F: 70-95%</p> <p>Drop outs: Most trials report 0%. 1 trial reported 2% and 3 did not state the number lost to follow up.</p>	<p>Group 1 Nerve blocks (any type, subcostal, lateral cutaneous, femoral, triple, psoas)</p> <p>Group 2 no block (either systemic analgesics or placebo)</p>	Pain	<p>Group 1: 106 Group 2: 104 SMD -0.52 (-0.8 to -0.25) p value: p = 0.0002</p>	<p>Funding: Supported internally by Peterborough and Stamford NHS Foundation Trust, UK and externally by Scottish Home and Health Department, UK.</p> <p>Additional outcomes: Length of operation, operative hypotension, intra-operative blood gases, complications specific to methods of treatment, allergic reactions, cerebrovascular accident, congestive cardiac failure, renal failure</p> <p>Notes:</p>
			Unsatisfactory pain control preoperatively or need for 'breakthrough' analgesia	<p>Group 1: 18/150 (12%) Group 2: 47/148 (31.8%) Relative risk: 0.37 95% CI: (0.23-0.61) p value: p<0.0001</p>	
			Nausea and/or vomiting	<p>Group 1: 18/141 (12.8%) Group 2: 25/159 (15.7%) Relative risk: 1.05 95% CI: (0.63-1.75) p value: 0.84</p>	
			Need for anti-emetics	<p>Group 1: 0/20 (0%) Group 2: 5/20 (25%) Relative risk: 0.09 95% CI: (0.01-1.54) p value: not reported</p>	
			Wound infection	<p>Group 1: 0/28 (0%) Group 2: 2/27(7.4%) Relative risk: 0.019 95% CI: (0.01-3.85) p value: p= 0.14</p>	
			Pneumonia	<p>Group 1: 12/129 (9.3%) Group 2: 25/130 (19.2%) Relative risk: 0.49 95% CI: (0.26-0.94)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
hospital stay and duration of time in emergency department				p value: 0.03	
			Any cardiac complication	Group 1: 3/62 (4.8%) Group 2: 12/62 (19.4%) Relative risk: 0.25 95% CI: (0.07-0.84) p value: 0.02	
			Myocardial infarction	Group 1: 1/34 Group 2: 4/34 Relative risk: 0.25 95% CI: (0.03-2.12) p value: Not significant	
			Puritis	Group 1: 0/20 Group 2: 5/20 Relative risk: 0.09 95% CI: (0.01-1.54) p value:	
			Pulmonary embolism	Group 1: 1/53 (1.9%) Group 2: 2/52 (3.8%) Relative risk: 0.66 95% CI: (0.11-3.86) p value: 0.64	
			Deep vein thrombosis	Group 1: 7/116 (6%) Group 2: 7/137 (5.1%) Relative risk: 1.12 95% CI: (0.43-2.93) p value: 0.82	
			Mortality	Group 1: 9/189 (4.8%) Group 2: 19/205 (9.3%) Relative risk: 0.59 d 95% CI: (0.29-1.21)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				p value: 0.15	
			Pressure sores	Group 1: 3/86 (3.5%) Group 2: 9/106 (8.5%) Relative risk: 0.51 95% CI: (0.11-2.39) p value: 0.39	
			Confusional state	Group 1: 15/77 (19.5%) Group 2: 34/101 (33.7%) Relative risk: 0.63 95% CI: (0.37-1.06) p value: 0.08	

1 17.4 Evidence Table 4: Anaesthesia

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Parker et al., 2004^{266,270}</p> <p>Study design: Cochrane systematic review. Includes 22 randomised and quasi randomised controlled trials</p> <p>Duration of follow-up: Range: 2 days to 30 months</p>	<p>Patient group: Hip fracture patients</p> <p>Inclusion criteria Skeletally mature patients undergoing hip fracture surgery</p> <p>Exclusion criteria Not stated</p> <p>All patients N (range): 2567 Age range: 60-91</p> <p>Drop outs: 0-7%. Not stated</p> <p>Setting: Hospitals in Europe, Hong Kong, New Zealand, Japan</p>	<p>Group 1 Regional (spinal or epidural) anaesthesia</p> <p>Group 2 General anaesthesia</p>	Mortality (early up to 1 month)	<p>Group 1: 64/912 (7%) Group 2: 93/966 (9.6%) Relative risk: RR 0.73 95% CI: (0.54-0.99) p value: 0.04</p>	<p>Funding: Supported internally University of Teesside, Middlesbrough, UK and Peterborough and Stamford Hospitals NHS Foundation Trust, Peterborough, UK.</p> <p>Limitations:</p> <p>Additional outcomes: Length of operation, operative hypotension, operative blood loss, patients receiving blood transfusion, transfusion requirements, postoperative hypoxia, cerebrovascular accident, congestive cardiac failure, renal failure, urine retention.</p> <p>Notes: All results reported in this table have been obtained using a fixed</p>
			Mortality at 1 month	<p>Group 1: 56/811 (6.9%) Group 2: 86/857 (10%) Relative risk: 0.69 95% CI: (0.50-0.95) p value: 0.02</p>	
			Mortality at 3 months	<p>Group 1: 86/726 (12%) Group 2: 98/765 (13%) Relative risk: 0.92 95% CI: (0.92-1.21) p value: 0.55</p>	
			Mortality at 6 months	<p>Group 1: 103/613 (17%) Group 2: 105/651 (16%) Relative risk: 1.04 95% CI: (0.81-1.33) p value: 0.76</p>	
			Mortality at 12 months	<p>Group 1: 80/354 Group 2: 78/372 Relative risk: 1.07 95% CI: (0.82-1.33) p value: 0.61</p>	
			Length of stay in hospital	<p>Group 1: n=108 Group 2: n=110 Mean Difference: -0.21</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				95% CI: -5.21-4.78 p value: (If no p-value: Sig/Not sig/NR)	effect model. Where there was heterogeneity a random effects model was used the results of which have not been reported here (please refer to forest plots).
			Vomiting	Group 1: 2/46 (4.3%) Group 2: 3/49 (6.1%) Relative risk: 0.7 95% CI: (0.12-3.94) p value: 0.68	
			Acute confusional state	Group 1: 11/117 (9.4%) Group 2: 23/120 (19.2%) Relative risk: 0.5 95% CI: (0.26-0.95) p value: 0.03	
			Pneumonia	Group 1: 21/574 (3.7%) Group 2: 29/612 (4.7%) Relative risk: 0.76 95% CI: (0.44-1.3) p value:0.32	
			Myocardial infarction	Group 1: 5/502 (1%) Group 2: 11/531 (2.1%) Relative risk: 0.55 95% CI: (0.22-1.37) p value: 0.2	
			Pulmonary embolism	Group 1: 9/605 (1.5%) Group 2: 13/640 (2%) Relative risk: 0.88 95% CI: (0.32-2.39) p value: 0.8	
			Deep vein thrombosis	Group 1: 39/129 (30.2%) Group 2: 61/130 (36.9%) Relative risk: 0.64 95% CI: (0.48-0.86)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				p value: 0.003	

1

1 **17.5 Evidence Table 5: Surgeon seniority**

Study details	Patients	Exposure	Outcome measures	Effect size	Comments
Enocson et al., 2008 ^{85,85}	<p>Patient group: Consecutive patients who had a hemiarthroplasty for non-pathological displaced femoral neck fracture</p> <p>Setting: Orthopaedics department</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Not reported <p>Exclusion criteria:</p> <ul style="list-style-type: none"> None reported <p>All patients N: 739 hips in 720 patients No. of dropouts: not reported Age (mean \pmSD): women: 84 (54-103), men 82 (55-97) years M/F: 147/592</p>	<p>Surgeon experience</p> <p><u>Group 1</u> Post registrar: 604 operations</p> <p><u>Group 2</u> Registrar: 135 operations</p> <p>59 surgeons in total - number of surgeons by grade not reported</p>	Number of dislocations	<p>Group 1: 37/404 (9.2%) Group 2: 8/135 (5.9%)</p>	<p>Funding: None reported</p> <p>Limitations: No details about surgeons and the number in each group.</p> <p>Not reported how patients were allocated to surgeons, no mention of anaesthetists grade/experience involved in operations.</p> <p>Outcomes not reported: Mortality, length of stay in secondary care, reoperations, quality of life, functional status, wound infection.</p>
<p>Country of study: Sweden</p> <p>Study design: Historical cohort</p> <p>List who was masked to interventions: Not applicable</p> <p>Duration of follow-up: Median 2.3 (0-10) years</p>			<p>Dislocation by 'post registrars' compared to 'Registrars'. Logistic regression univariate analysis</p>	Odds ratio: 1.0 (0.4, 2.2) P=0.9	
			<p>Dislocation by 'post registrars' compared to 'Registrars'. Logistic regression multivariate analysis adjusted for age, sex, indication for surgery, surgical approach and type of hemiarthroplasty</p>	Odds ratio: 1.3 (0.6, 3.0) P=0.5	

2

1 Evidence tables – surgeon seniority

Study details	Patients	Exposure	Outcome measures	Effect size	Comments
Enocson et al., 2009 ^{83,85}	<p>Patient group: Consecutive patients who had a primary total hip replacement for non-pathological displaced femoral neck fracture (Garden III or IV) or secondary total hip replacement due to a fracture healing complication (non-union or avascular necrosis) after internal fixation.</p> <p>Setting: Orthopaedics department</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Not reported <p>Exclusion criteria:</p> <ul style="list-style-type: none"> None reported <p>All patients N: 713 hips in 698 patients No. of dropouts: not reported Age (mean \pmSD): women: 78 \pm8.6 (46-96) , men 74 \pm9.8 (45-90) years M/F: 140/573</p>	<p>Surgeon experience</p> <p><u>Group 1</u> Post registrar: 636 operations</p> <p><u>Group 2</u> Registrar: 77 operations</p> <p>54 surgeons in total - number of surgeons by grade not reported</p>	Number of dislocations	<p>Group 1: 38*/636 (6%) Group 2: 3*/77 (3.9%)</p>	<p>Funding: None reported</p> <p>Limitations: No details about surgeons and the number in each group. Not reported how patients were allocated to surgeons, no mention of anaesthetists grade/experience involved in operations.</p> <p>Outcomes not reported: Mortality, length of stay in secondary care, reoperations, quality of life, functional status, wound infection.</p>
<p>Country of study: Sweden</p> <p>Study design: Historical cohort</p> <p>List who was masked to interventions: Not applicable</p> <p>Duration of follow-up: Median 4.3 (0-11) years</p>			<p>Dislocation by 'post registrars' compared to 'Registrars'. Cox regression univariate analysis</p>	Hazard ratio: 1.4 (0.4, 4.5) P=0.6	
			<p>Dislocation by 'post registrars' compared to 'Registrars'. Cox regression multivariate analysis adjusted for age, sex, indication for surgery, surgical approach and femoral head size</p>	Hazard ratio: 0.9 (0.3, 2.8) P=0.8	
				* number calculated by NCGC	

2

1 Evidence tables – surgeon seniority

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Palm et al., 2007 ^{256,257} Country of study: Denmark Study design: Prospective cohort List who was masked to interventions: None Duration of follow-up: 6 months	Patient group: Consecutive patients with proximal fracture of the femur. Various classifications of fracture. All patients N: 600 No. of dropouts: none Group 1 No.: 137 No. of dropouts: 0 Age (mean \pmSD): 81 (72-87) M/F: 12/44 Types of fracture:: Technically demanding fractures <ul style="list-style-type: none"> ○ Posterior angulated Garden I-II (n=8) ○ Garden III-IV (n=23) ○ Petrotrochanteric (Evans type 5) (n=23) ○ Per-/subtrochanteric (n=2) ○ Subtrochanteric (n=0) ○ Pathological (n=0) Technically undemanding fractures <ul style="list-style-type: none"> ○ Garden I-II (n=13) ○ Basocervical (n=4) ○ Petrotrochanteric (Evans type 1-4) 	Surgeon experience. Number of surgeons not reported. Group 1 Unsupervised orthopaedic junior surgeon (<3 years orthopaedic surgical experience) 137 operations (56 classified as technically demanding). Group 2 Experienced surgeon (> 3 years orthopaedic surgical experience) 463 operations (309 classified as technically demanding).	Reoperation at 6 months for technical demanding fractures (unadjusted for other factors)	Group 1: 16/56 (29%) Group 2: 47/309 (15%) P=0.015	Funding: Supported by grant from IMK Fonden Limitations: Not stated how patients were allocated to surgeons, no mention of anaesthetists grade/experience involved in operations. Senior surgeons operated on significantly more patients with a poor prefracture mobility score Outcomes not reported: Mortality, length of stay in secondary care, requirement for surgical revision, wound infection. Additional outcomes reported:
			Reoperation at 6 months for technical demanding fractures (multivariate analysis combining age >85, female gender, ASA score III-IV, Pre fracture New Mobility score 0-5 (poor score), time to surgery >1 day from admission & type of implant (arthroplasty or osteosynthesis)).	Odds ratio 2.01 (1.01, 4.02) P=0.048	
			Prefracture New Mobility Score of 0-5 (scale 0f 0-9, score of 0 means patient is unable do any of the following: to get around the house, get out of the house or go shopping. Score of 9 means the patient can do all 3 with no difficulty)	Group 1: 173/309 (56%) Group 2: 21/56 (38%) P=0.011	
			Number of patients receiving arthroplasty	Group 1: 166/309 (54%) Group 2: 12/56 (21%) P<0.0001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>(n=64)</p> <p>Group 2 No.: 463 No. of dropouts: 0 Age (mean \pmSD): 83 (77-88) M/F: 63/246</p> <p>Types of fracture:</p> <p>Technically demanding fractures</p> <ul style="list-style-type: none"> ○ Posterior angulated Garden I-II (n=18) ○ Garden III-IV (n=176) ○ Petrotrochanteric (Evans type 5) (n=73) ○ Per-/subtrochanteric (n=18) ○ Subtrochanteric (n=20) ○ Pathological (n=4) <p>Technically undemanding fractures</p> <ul style="list-style-type: none"> ○ Garden I-II (n=43) ○ Basocervical (n=11) ○ Petrotrochanteric (Evans type 1-4) (n=100) 				<p>multivariate analysis for age >85, female gender, ASA score III-IV, Pre fracture New Mobility score 0-5 (poor score), time to surgery >1 day from admission & type of implant.</p> <p>Notes: Only technically demanding fractures were analysed by logistic regression.</p>

1 **17.6 Evidence Table 6: Displaced intracapsular fractures**

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Parker et al., 2010^{263,265,270}</p> <p>Country of study:</p> <p>Study design: Systematic review including 6 out of the 19 RCTs from the review with 734 participants. The remaining RCTs were not relevant to this comparison.</p> <p>Duration of follow-up: Average ranged from 6 months to 4 years</p>	<p>Patient group: Skeletally mature patients with a proximal femoral fracture.</p> <p>Setting: Hospital</p>	<p>Group 1 Hemiarthroplasty (cemented or uncemented)</p> <p>Group 2 Total hip replacement</p> <p>Additional non-comparative prophylaxis: Not applicable</p>	<p>Outcomes extracted</p>	<p>Results reported in forest plots for:</p> <ul style="list-style-type: none"> - Mortality at 3 to 4 months, 1 year & 2 to 4 years - Number of reoperations - Pain – residual pain and Harris Hip Score for pain at 1 year - Failure to regain mobility at final follow up - Functional scores: Oxford Hip Score, Harris Hip Score, Barthel Score, Hip Rating Questionnaire, Short Form 36 physical function score - Self reported walking distance at end of study. - Quality of Life – Eq-5d index score - All medical complications - Length of hospital stay 	<p>Funding: supported internally at Peterborough and Stamford Hospitals NHS Trust, UK. No external source of funding.</p> <p>Limitations:</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: length of surgery, hypotension during surgery, operative blood loss, postoperative blood transfusion, cost of treatment, leg shortening, external rotation deformity</p> <p>Notes:</p>

1 Evidence tables – displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Parker et al., 2006^{264,270}</p> <p>Country of study:</p> <p>Study design: Systematic review including 17 RCTs with 2694 participants.</p> <p>Duration of follow-up: Average ranged from 1 to 13 years</p>	<p>Patient group: Skeletally mature patients with a intracapsular proximal femoral fracture.</p> <p>Setting: Hospital</p> <p>12 trials involving 1973 participants compared internal fixation to hemiarthroplasty.</p> <p>6 trials involving 881 participants compared internal fixation to total hip replacement.</p> <p>The numbers do not add up to 17 trials and 2694 participants as: 1 trial of 409 patients was not included in our analysis as it did not distinguish between hemiarthroplasty and total hip replacement; and two trials investigated a three way comparison of internal fixation, hemiarthroplasty and total hip replacement .</p>	<p>Group 1 Internal fixation</p> <p>Group 2 a. Hemi-arthroplasty b. total hip replacement</p> <p>Additional non-comparative prophylaxis: Not applicable</p>	<p>Outcomes extracted</p>	<p>Results in forest plots for:</p> <ul style="list-style-type: none"> - Mortality at 1 month, 3 months, 1 year & 2 to 4 years - Number of reoperations split into major, moderate, minor and total number of reoperations - Pain at 1 year and 2 to 3 years - Failure to return to same place of residence by final follow up - Failure to regain mobility at final follow up - All medical complications - Length of hospital stay 	<p>Funding: supported internally at Peterborough and Stamford Hospitals NHS Trust, UK. No external source of funding.</p> <p>Limitations:</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: length of surgery, hypotension during surgery, operative blood loss, postoperative blood transfusion, cost of treatment, leg shortening, external rotation deformity</p> <p>Notes:</p>

2
3

1 Evidence tables – displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Frihagen et al., 2007^{102,103}</p> <p>Country of study: Norway</p> <p>Study design: RCT</p> <p>List who was masked to interventions: Investigators of functional outcomes were blinded to interventions. Unclear if anyone else was masked to the intervention after randomisation.</p> <p>Duration of follow-up: 24 months</p>	<p>Patient group: Patients with a intracapsular femoral neck fracture with angular displacement in either radiographic plane.</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> - age \geq60 - ability for independent ambulation before fracture - displaced femoral neck fracture <p>Exclusion criteria</p> <ul style="list-style-type: none"> - unfit for arthroplasty according to anaesthesiologist - previous symptomatic hip pathology such as arthritis - pathological fracture - delay of more than 96 hours from injury to treatment - living outside hospital's designated area <p>Setting: Hospital</p> <p>All patients N: 222 No. of dropouts: 0</p>	<p>Group 1 Closed reduction and internal fixation with two parallel cannulated screws (Olmed, DePuy/Johnson and Johnson, Sweden)</p> <p>Group 2 Charnley-Hastings bipolar cemented hemiarthroplasty (DePuy/Johnson and Johnson, Sweden).</p>	Mortality at 30 days	Group 1: 7/112 Group 2: 10/110 P value(s): 0.42	<p>Funding: Norwegian Foundation for Health and Rehabilitation through the Norwegian Osteoporosis Society and the Norwegian Research Council, Nycomed, Smith and Nephew, and OrtoMedic</p> <p>Limitations: Functional outcome data not available for all patients.</p> <p>Outcomes not reported: length of superspell, place of residence 12 months after fracture, pain</p> <p>Additional outcomes reported: time from admission to surgery, time in operation theatre, time of surgery,</p>
			Mortality at 90 days	Group 1: 16/112 Group 2: 20/110 P value(s): 0.43	
			Mortality at 12 months	Group 1: 24/112 Group 2: 29/110 P value(s): 0.39	
			Mortality at two years	Group 1: 39/112 Group 2: 39/110 P value(s): 0.92	
			Any medical complication	Group 1: 28/111 Group 2: 30/109 P value(s): 0.70	
			Total number of reoperations at 24 months	Group 1: 70/111 Group 2: 13/108 P value(s): <0.001	
			Total number of hips with any reoperation at 24 months	Group 1: 47/111 Group 2: 11/108 P value(s): <0.001	
			Total number of hips with major reoperation at 24 months	Group 1: 44/111 Group 2: 11/108 P value(s): <0.001	
			Length of hospital stay (mean \pmSD)	Group 1: 8.2 \pm 7.35 (n= 111) Group 2: 10.2 \pm 11.95 (n= 109) P value(s): 0.14	
Harris hip score (mean \pmSD) at 4 months	Group 1: 59.6 \pm 19.5 (n= 89) Group 2: 67.7 \pm 15.8 (n= 84) P value(s): 0.003				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 1: internal fixation No. randomised: 112 No. of dropouts: 0 Mean age (SD): 83.2 (7.65) M/F: 25/87 Other factors: Concurrent symptomatic medical disease: 52 Previously recognised cognitive failure: 40 Ability to walk without any aid: 67 Mean time from injury to admission: 8 hours</p> <p>Group 2: hemiarthroplasty No. randomised: 110 No. of dropouts: 0 Mean age (SD): 82.5 (7.32) M/F: 32/78 Other factors: Concurrent symptomatic medical disease: 64 Previously recognised cognitive failure: 29 Ability to walk without any aid: 60 Mean time from injury to admission: 5.5 hours</p>		Harris hip score (mean \pm SD) at 12 months	Group 1: 65.8 \pm 15.9 (n= 87) Group 2: 72.6 \pm 17.5 (n= 74) P value(s): 0.01	<p>intraoperative blood loss, main surgeons with >3 years experience with procedure, spinal anaesthesia, no. receiving blood transfusion while admitted, postoperative confusion, cognitive failure at 4 months, type of reoperation</p> <p>Notes:</p>
			Harris hip score (mean \pm SD) at 24 months	Group 1: 67.3 \pm 15.5 (n= 71) Group 2: 70.6 \pm 19.1 (n= 68) P value(s): 0.26	
			Eq-5d index score (mean \pm SD) at 4 months	Group 1: 0.53 \pm 0.29 (n= 79) Group 2: 0.61 \pm 0.30 (n= 70) P value(s): 0.06	
			Eq-5d index score (mean \pm SD) at 12 months	Group 1: 0.56 \pm 0.33 (n= 70) Group 2: 0.65 \pm 0.30 (n= 62) P value(s): 0.07	
			Eq-5d index score (mean \pm SD) at 24 months	Group 1: 0.61 \pm 0.31 (n= 52) Group 2: 0.72 \pm 0.23 (n= 52) P value(s): 0.03	
			Eq-5d visual analogue scale (mean \pm SD) at 4 months	Group 1: 53 \pm 18.5 (n= 69) Group 2: 62 \pm 21.0 (n= 60) P value(s): 0.01	
			Eq-5d visual analogue scale (mean \pm SD) at 12 months	Group 1: 57 \pm 21.6 (n= 59) Group 2: 63 \pm 24.3 (n= 54) P value(s): 0.16	
			Eq-5d visual analogue scale (mean \pm SD) at 24 months	Group 1: 60 \pm 18.0 (n= 45) Group 2: 60 \pm 21.0 (n= 43) P value(s): 0.84	
			No. patients with Barthel Index Score of 95 or 100 at 4 months	Group 1: 41/88 Group 2: 40/80 P value(s): 0.66	
			No. patients with Barthel Index Score of 95 or 100 at 12 months	Group 1: 31/87 Group 2: 39/73 P value(s): 0.02	
			No. patients with Barthel	Group 1: 24/69	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Index Score of 95 or 100 at 24 months	Group 2: 26/68 P value(s): 0.02	
			Total number of complications at 24 months	Group 1: 70/111 Group 2: 16/108 P value(s): <0.001	
			Total number of hips with any complication at 24 months	Group 1: 56/111 Group 2: 16/108 P value(s): <0.001	
			Total number of hips with major complication at 24 months	Group 1: 47/111 Group 2: 11/108 P value(s): <0.001	
			Complications at 24 months – deep infection	Group 1: 7/111 Group 2: 7/108 P value(s):	
			Complications at 24 months – mechanical failure of internal fixation/non-union	Group 1: 40/111 Group 2: 3/108 P value(s):	
			Complications at 24 months – dislocation of hemiarthroplasty	Group 1: 6/111 Group 2: 1/108 P value(s):	
			Complications at 24 months – avascular necrosis	Group 1: 6/111 Group 2: 0/108 P value(s):	
			Median (range) time to complication	Group 1: 137.5 (8-730) days (n= 111) Group 2: 18 (6-730) days (n= 109) P value(s): 0.01	

1 Evidence tables – displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Macauley et al., 2008^{197,197}</p> <p>Country of study: USA</p> <p>Study design: RCT</p> <p>List who was masked to interventions: Unclear if anyone was masked to the intervention after randomisation.</p> <p>Duration of follow-up: 24 months</p>	<p>Patient group: Patients with a displaced intracapsular proximal femoral fracture.</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> - age \geq50 - ability for independent ambulation before fracture - displaced femoral neck fracture (Garden III or IV which the surgeon considered not amenable to treatment with open reduction internal fixation (ORIF)) - ability to comprehend either English or Spanish <p>Exclusion criteria</p> <ul style="list-style-type: none"> - chronic severe dementia (defined as $<$23 of 30 on Folstein Mini Mental State Examination (MMSE)) - pathological fracture - other concomitant long bone fractures or fractures requiring surgical repair - preexisting arthritis of the ipsilateral hip <p>Setting: Hospital</p>	<p>Group 1 Hemiarthroplasty (unipolar or bipolar, cemented or uncemented stem).</p> <p>Group 2 Total hip replacement with a femoral head of 28mm or more (cemented or uncemented stem).</p>	Mortality at 6 months after surgery	Group 1: 5/23 Group 2: 1/17 P value(s): 0.21	<p>Funding: American Association of Hip and Knee Surgeons, Orthopaedic Research and Education Foundation</p> <p>Limitations:</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: duration of operation</p> <p>Notes: study designed to demonstrate the feasibility of a large randomised, multicentre trial with multiple surgeons treating subjects with displaced intracapsular femoral neck fractures.</p>
			Mortality at mean follow up of 34 months (29 to 42 months)	Group 1: 9/23 Group 2: 5/17 P value(s): 0.53	
			Bodily pain at 12 months (SF-36 subscales 1-100) (mean \pmSD)	Group 1: 42.4 \pm 11.5 (n= 23) Group 2: 53.2 \pm 10.2 (n= 17) P value(s): 0.02	
			Pain on injured side at 12 months (WOMAC 1-100) (mean \pmSD)	Group 1: 88.5 \pm 13.6 (n= 23) Group 2: 92.5 \pm 14.6 (n= 17) P value(s): 0.50	
			Bodily pain at 24 months (SF-36 subscales 1-100) (mean \pmSD)	Group 1: 44.7 \pm 10.5 (n= 23) Group 2: 54.8 \pm 7.9 (n= 17) P value(s): 0.03	
			Pain on injured side at 24 months (WOMAC 1-100) (mean \pmSD)	Group 1: 77.8 \pm 20.9 (n= 23) Group 2: 94.4 \pm 6.8 (n= 17) P value(s): 0.05	
			Physical function at 12 months (SF-36 subscales 1-100) (mean \pmSD)	Group 1: 32.8 \pm 10.0 (n= 23) Group 2: 33.5 \pm 12.0 (n= 17) P value(s): 0.87	
			Function at 12 months (WOMAC 1-100) (mean \pmSD)	Group 1: 78.7 \pm 16.8 (n= 23) Group 2: 75.9 \pm 19.8 (n= 17) P value(s): 0.71	
			Physical function at 24 months (SF-36 subscales 1-100) (mean \pmSD)	Group 1: 35.1 \pm 12.9 (n= 23) Group 2: 38.6 \pm 8.9 (n= 17) P value(s): 0.52	
Function at 24 months (WOMAC 1-100) (mean	Group 1: 65.1 \pm 18.1 (n= 23) Group 2: 81.8 \pm 10.2 (n= 17)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients N: 41 No. of dropouts: 1 (2.5%)</p> <p>Group 1: hemiarthroplasty No. randomised: 23 No. of dropouts: 0 Mean age (SD): 77 (9) M/F: 9/14 Other factors: Average no. comorbid conditions: 4.2 (1-11)</p> <p>Group 2: total hip replacement No. randomised: 18 No. of dropouts: 1 Mean age (SD): 82 (7) M/F: 10/7 Other factors: Average no. comorbid conditions: 3.5 (0-7)</p>		±SD)	P value(s): 0.66	
			Physical component summary score at 12 months (SF-36 subscales 1-100) (mean ±SD)	Group 1: 36.4 ±9.2 (n= 23) Group 2: 40.2 ±9.9 (n= 17) P value(s): 0.35	
			Physical component summary score at 24 months (SF-36 subscales 1-100) (mean ±SD)	Group 1: 40.9 ±12.3 (n= 23) Group 2: 43.0 ±7.5 (n= 17) P value(s): <0.68	
			Harris Hip Score on injured side at 12 months (1-100) (mean ±SD)	Group 1: 80.6 ±14.3 (n= 23) Group 2: 84.2 ±12.0 (n= 17) P value(s): 0.55	
			Harris Hip Score on injured side at 24 months (1-100) (mean ±SD)	Group 1: 81.1 ±11.7 (n= 23) Group 2: 84.0 ±12.2 (n= 17) P value(s): 0.64	
			TUG score (Take "Up and Go" score at 12 months (mean ±SD)	Group 1: 16.5 ±10.1 (n= 23) Group 2: 17.2 ±13.5 (n= 17) P value(s): 0.89	
			TUG score (Take "Up and Go") score at 24 months (mean ±SD)	Group 1: 16.9 ±10.1 (n= 23) Group 2: 14.7 ±7.2 (n= 17) P value(s): 0.64	
			Length of stay in hospital (mean ±SD days)	Group 1: 7.7 ±5.5 (n= 23) Group 2: 5.4 ±2.8 (n= 17) P value(s): 0.18	
			Length of stay in hospital (median days)	Group 1: 7 (n= 23) Group 2: 6 (n= 17)	

1 Evidence tables – displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Mouzopoulos et al., 2008^{218,218}</p> <p>Country of study: Greece</p> <p>Study design: RCT</p> <p>List who was masked to interventions:</p> <p>Duration of follow-up: 4 years</p>	<p>Patient group: Patients with a displaced subcapital hip fractures (Garden III or IV)</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> - displaced femoral neck fracture (Garden III or IV) <p>Exclusion criteria</p> <ul style="list-style-type: none"> - previous hip surgery - history of cancer or Paget's disease - rheumatic arthritis <p>Setting: Hospital</p> <p>All patients N: 129 No. of dropouts: 34 at 1 year, 67 at 4 years</p> <p>Group 1: internal fixation No. randomised: 43 No. of dropouts: 11 at 1 year, 24 at 4 years Mean age (SD): 75.38 (4.62)* M/F: 12/26* Other factors: Average no. comorbid conditions: 4.2 (1-11)</p>	<p>Group 1 Internal fixation (Richards plate screw, Smith & Nephew, Memphis, TN, USA)</p> <p>Group 2 Hemiarthroplasty (Merete, Berlin, Germany).</p> <p>Group 3 Total hip replacement (Plus; DePuy, Warsaw, IN, USA).</p>	Mortality at 1 year	<p>Group 1: 6/43 Group 2: 6/43 Group 3: 5/43 P value(s):</p>	<p>Funding: not reported</p> <p>Limitations: method of randomisation unclear study: patients assigned in order of type of fixation: hemiarthroplasty, total hip replacement, internal fixation. No indication that anyone was masked to the intervention.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: mentions but provides no figures for range of passive motion, and walking speed. Barthel Index score prefracture</p> <p>Notes:</p>
			Mortality at 4 years	<p>Group 1: 15/43 Group 2: 13/43 Group 3: 11/43 P value(s):</p>	
			Prefracture function according to the Barthel Index Score	<p>Group 1: 85.2 ±4.8 (n= 43) Group 2: 81.05 ±8.95 (n= 43) Group 3: 87.4 ±17.4 (n= 43)</p>	
			Function according to the Barthel Index Score at 1 year	<p>Group 1: 77.1 ±7.1 (n= 32) Group 2: 76.8 ±6.8 (n= 30) Group 3: 84.8 ±14.8 (n= 33)</p>	
			Function according to the Barthel Index Score at 4 years	<p>Group 1: 80.1 ±5.3 (n= 19) Group 2: 79.6 ±6.3 (n= 20) Group 3: 85.3 ±11.6 (n= 23)</p>	
			Harris Hip Score at 1 year	<p>Group 1: 71.3 ±5.3 (n= 32) Group 2: 77.81 ±9.6 (n= 30) Group 3: 83.7 ±4.8 (n= 33) P value <0.05 for comparison between group 1 and 3</p>	
			Harris Hip Score at 4 years	<p>Group 1: 73.6 ±6.7 (n= 19) Group 2: 79.5 ±6.5 (n= 20) Group 3: 83.7 ±4.8 (n= 23) P value <0.05 for comparison between group 1 and 3</p>	
			Number of revisions	<p>Group 1: 12/43 Group 2: 5/43 Group 3: 1/43</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2: hemiarthroplasty No. randomised: 43 No. of dropouts: 13 at 1 year, 23 at 4 years Mean age (SD): 74.24 (3.77)* M/F: 10/24* Other factors: Average no. comorbid conditions: 3.5 (0-7)</p> <p>Group 3: total hip replacement No. randomised: 43 No. of dropouts: 10 at 1 year, 20 at 4 years Mean age (SD): 73.07 (4.93)* M/F: 9/28* Other factors: Average no. comorbid conditions: 3.5 (0-7)</p> <p>* data not provided for all patients</p>			<p>P value(s):</p>	

1 17.7 Evidence Table 7: Surgery – Cement versus no cement

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Parker et al., 2010 ²⁶⁵ Country of study: UK Study design: Systematic review including 19 RCTs, 6 relating to cemented stems in old designs of hemiarthroplasty with 899 participants, 1 relating to new styles of stems with 220 participants Duration of follow-up: Average ranged from 6 months to 4 years	Patient group: Skeletally mature patients with a proximal femoral fracture. Setting: Hospital	Group 1 Cemented prostheses Group 2 Uncemented prostheses Additional non-comparative prophylaxis: Not applicable	Outcomes extracted for older designs of hemiarthroplasty	Results reported in forest plots for: <ul style="list-style-type: none"> - Mortality at up to 1 month, 1 to 3 months, 1 year & 3 years - Number of reoperations at 8 to 20 months - Failure to regain mobility at 12 to 17 months - Change in mobility score at 12 months - Length of hospital stay - Number of patients failing to return home at 1.5 to 5 years - Pain at 3 months and 1 to 2 years - Pain score at 6 months - Number of reoperations at 8 to 20 months - Deep sepsis at 1 to 5 years - Wound haematoma at 1 to 5 years - All medical complications 	Funding: Not reported Limitations: Outcomes not reported: Additional outcomes reported: length of surgery, hypotension during surgery, operative blood loss, postoperative blood transfusion, cost of treatment, leg shortening, external rotation deformity Notes: Review also compares: different types of unipolar or bipolar hemiarthroplasties, unipolar vs. bipolar hemiarthroplasty, uncemented
			Outcomes extracted for new designs of hemiarthroplasty	Results reported in forest plots for: <ul style="list-style-type: none"> - Mortality at 30 days, 9 days, 1 year & 2 years - Number of reoperations at 12 months - Need for pain medication at 12 months - Unable to walk without aids at 12 to months - Functional scores: Barthel Index, 	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Harris Hip Score and Eq-5d at 12 months - Length of hospital stay	hemiarthroplasty vs. total hip replacement, cemented hemiarthroplasty vs. total hip replacement, different types of total hip replacement.

1 17.8 Evidence Table 8: Extracapsular fixation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ahrengart et al., 2002^{3,3}</p> <p>Country of study: Sweden and Finland</p> <p>Study design: RCT</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients with intertrochanteric fractured femur.</p> <p>Setting: 5 hospitals.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Fracture types 1 to 5 of the Evans' classification of intertrochanteric fractures, as modified by Jensen and Michaelsen. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Subtrochanteric and pathologic fractures, earlier fractures or operations on the same hip, or if the surgeon was unfamiliar with the Gamma nail technique. <p>All patients N: 492 No. of dropouts: 66 (13%) Group 1: Gamma nail No. randomised: 210 No. of dropouts: Mean age (range): F: 82 (48-96) M: 77 (44-90) M/F: 63/147 Other factors:</p>	<p>96% of patients were operated on within 2 days.</p> <p>Group 1 Gamma nail The 12mm diameter Gamma nail was used in 73%, the 14mm nail in 20% and the 16mm nail in 7% of patients. The proximal femur was reamed to a 2mm larger diameter than the diameter of the nail. In patients with stable fractures, distal locking was used in 68% of patients, and in unstable fractures 74% of patients.</p> <p>Group 2 Compression hip screw The Richard's classic or the Dynamic hip screw was used. 2 hole plates were used in 5%, 4 hole in 67%, 5 hole in 20%, 6 hole in 7%, and 8 or 10 in 2% of patients in whom a compression screw was used.</p>	Additional fissure/fracture perioperatively	Group 1: 5 Group 2: 2	<p>Funding: The Karolinska Institute Foundation, Lund University, Skane County Council and Stryker-Howmedica.</p> <p>Outcomes not reported: Place of residence</p> <p>Additional outcomes reported: Radiological parameters, operation time, blood loss, % of fractures healed in preoperative position, hip rotation</p> <p>Notes: Of the 5 hospitals participating in the study, 1 centre was active for 3 years, whereas the others participated for 2</p>
			Other technical/surgical problems	Group 1: 5 Group 2: 2	
			Duration of hospital stay, mean (range)	Group 1: 10 (1 – 100) Group 2: 10 (1 – 100)	
			Wound infection	Group 1: 10 (1 – 100) Group 2: 10 (1 – 100)	
			Cut out of lag screw	Group 1: 14 Group 2: 4	
			Mortality of 6 months	Group 1: 41 Group 2: 37	
			Healed fracture at 6 months	Group 1: 89% Group 2: 88%	
			Lateral pain over the femoral head screw at 6 months	Group 1: 27% Group 2: 26%	
			Pain at the top of the greater trochanter at 6 months	Group 1: 20% Group 2: 6% p<0.001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	ASA score: 1: 16% 2: 42% 3: 34% 4: 8% 5: 0 Group 2: Compression hip screw No. randomised: 216 No. of dropouts: Age (mean \pmSD): F: 81 (54-99) M: 74 (32-98) M/F: 60/156 Other factors: ASA score: 1: 20% 2: 39% 3: 36% 4: 6% 5: 0	Additional non-comparative prophylaxis: 81% of patients received antibiotic prophylaxis. Prophylactic anticoagulants were given to 75% of the patients; 56% received dextran and 18% received heparin preparations or warfarin.	Needs walking aid Lives at home Internal hip rotation of the fractured leg External hip rotation of the fractured leg	Group 1: 72% Group 2: 70% Group 1: 65% Group 2: 62% Group 1: 15° (0 – 50°) Group 2: 15° (0 – 45°) Group 1: 20° (0 – 70°) Group 2: 30° (0 – 60°) p = p<0.001	years each. In most cases patient who were lost to follow up were absent at the final exam due to advanced age, other physical illness, or dementia. Spinal anaesthesia used in 90% of patients.

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Aune et al., 1994^{9,9}</p> <p>Country of study: Norway</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: Median follow-up was 17 months (10-27)</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospitals, Norway</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Trochanteric or subtrochanteric femoral fractures <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • None stated <p>All patients N: 378</p> <p>No. of dropouts: 0</p> <p>Group 1: Gamma nail No. randomised: 177 Mean age (range): 82 (49-96) M/F: 66/109 Other factors: Stable trochanteric = 84 Unstable trochanteric = 76 Subtrochanteric = 14</p> <p>Group 2: Hip compression screw (HCS) No. randomised: 201 Age (mean ±SD): M/F: 89/114 Other factors: Stable trochanteric = 89 Unstable trochanteric = 98 Subtrochanteric = 17</p>	<p>Group 1 All the Gamma nails (Howmedica) were modified to a 6 degree valgus angle, 4 degrees less than in the standard nail. The slot for the lag screw had a 131 degree angle in relation to the shaft. The diameters of the nails used were 12 or 14mm. The medullary canal was over-reamed 2mm. In 119 of 177 nailings distal locking screws were inserted through a jig.</p> <p>Group 2 Hip compression screw (Smith and Nephew)</p>	<p>Requirement for reoperation</p>	<p>Group 1: 13/177</p> <ul style="list-style-type: none"> – Stable trochanteric =5 femoral shaft fractures and 2 cut out of the lag screw – Unstable trochanteric =4 femoral shaft fractures and 1 cut out of the lag screw – Subtrochanteric = 1 femoral shaft fracture <p>Group 2: 2/201</p> <ul style="list-style-type: none"> – Stable trochanteric = 1 cut out of the lag screw – Unstable trochanteric = 1 cut out of the lag screw – Subtrochanteric = 0 <p>P value(s): P < 0.003</p>	<p>Funding: Not reported</p> <p>Limitations: Small study, little detail about randomization and few outcomes reported e.g. mortality etc.</p> <p>Outcomes not reported: Mortality, length of stay in hospital, place of residence, functional status.</p> <p>Additional outcomes reported: Further details of the 15 patients requiring reoperation, including time from operation to reoperation.</p> <p>Notes: Fracture type assessed by methods of Jenson and Zickel.</p>

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Barton et al., 2010 ^{14,14} Country of study: UK Study design: RCT List who was masked to interventions: No blinding of assessor or patients. Duration of follow-up: 1 year	Patient group: Patients with fracture of the proximal femur Setting: Dept Trauma and Orthopaedics, Frenchay Hospital, Bristol. Inclusion criteria: Patients aged over 18 with AO/OTA 31-A2 fracture of the proximal femur. Exclusion criteria: Pathological fractures, previous proximal femoral fracture, reverse oblique fractures, and a decision by the surgeon responsible for the patient's care not to include the patient in the study. All patients N: 210 No. of dropouts: 2 Mean age (range): 83.2 (42 to 99) M/F: 44/166 Group 1: No. randomised: 110 No. of dropouts: 0 Mean age (range): 83.3 (56 to 97) M/F: 25/85 Other factors: ASA score	All surgeons performing the operations had experience with the 2 implants. Following surgery, patients were mobilized bearing full weight under the supervision of a physiotherapist. Following discharge, patients were evaluated both clinically and radiographically at 3, 6 and 12 months. Group 1 Sliding hip screw (Omega 2; Stryker, Newbury, UK) A four-hole, 135° plate was inserted. Group 2 Long gamma nail (Dyax; Stryker) The femur was reamed to 1mm greater than the	Reoperation (screw cut-out, implant failure, late fracture, and deep infection)	Group 1: 2 Group 2: 3 P value(s): 0.67 (all were screw cut out)	Funding: No external funding Limitations: Initial power calculation produced a sample requirement of 220 patients. Outcomes not reported: Additional outcomes reported: Requirement for transfusion, demographic characteristics (side of fracture, minimal score), tip-apex distance >25mm Notes:
			Mortality	30 days Group 1: 11 Group 2: 21 P value(s): 0.13 1 year Group 1: 24 Group 2: 32 P value(s): 0.26	
			Length of hospital stay	Group 1: 31 (1 to 154) Group 2: 32 (1 to 164) P value(s): 0.17	
			Mobility (change in score – points) (1 – unaided, 2 – one cane or crutch, 3 – two canes or crutches, 4 – walker, 5 – wheelchair)	Group 1: 1.49 Group 2: 1.86 P value(s): 0.26	
			Change in residence (change in score – points) (1 – own home, 2 – sheltered housing, 3 – residential home, 4 – nursing home, 5 – hospital)	Group 1: 1.23 Group 2: 1.16 P value(s): 0.79	
			EuroQol 5D	QUALY Group 1: 0.46	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	1: 2, 2: 46, 3: 59, 4: 3 Group 2: No. randomised: 100 No. of dropouts: 2 died before surgery Mean age (range): 83.1 (42 to 99) M/F: 19/81 Other factors: ASA score 1: 0, 2: 47, 3: 49, 4: 4	diameter of the nail, and a 130° nail of the appropriate length was inserted; all nails were locked distally with 2 screws.		Group 2: 0.37	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																					
Bridle et al., 1991 ^{37,37} Country of study: UK, London Study design: Randomised prospective comparison List who was masked to interventions: Not reported. Duration of follow-up: At least 6 months	Patient group: Patients with intertrochanteric fractured femur. Inclusion criteria: <ul style="list-style-type: none"> Patients diagnosed with intertrochanteric fractured femur Exclusion criteria: <ul style="list-style-type: none"> Not reported All patients N: 100 No. lost to follow up: 6 Group 1: Gamma nail No. randomised: 49 Age: 81.0 M/F: 9/40 Other factors: ASA score: I = 2 II = 23 III = 20 IV = 4 Fracture type: Stable: 18 Unstable: 31	Group 1 The Gamma nail was inserted using a 'closed' technique under image intensifier control. The patient is positioned on the traction table, and the fracture is reduced with the leg adducted. A 6 cm incision is made just proximal to the greater trochanter, which is entered using a curved awl. The entry point is just lateral to the tip of the trochanter. A guide wire is introduced into the femoral shaft, and flexible reamers are used to the appropriate size. A nail, 1 to 1.5 mm smaller than the final reamer, is selected. No attempt is made to ream the shaft to accept a large nail. The angle of the nail ranges from 125 to 140 degrees. Group 2 Dynamic hip screws were inserted using the standard technique.	Mortality	Before discharge Group 1: 10 Group 2: 9 6 months post op Group 1: 15 Group 2: 19	Funding: Not reported Limitations: Allocation concealment unclear. Outcomes not reported: Length of stay in hospital, reoperation. Additional outcomes reported: Operative details Notes: Treatment was randomised at the time of anaesthesia.																					
			Complications	<table border="1"> <thead> <tr> <th>Group</th> <th>1</th> <th>2</th> </tr> </thead> <tbody> <tr> <td>CVA</td> <td>4</td> <td>0</td> </tr> <tr> <td>Bronchopneumonia</td> <td>1</td> <td>3</td> </tr> <tr> <td>Pulmonary embolism</td> <td>1</td> <td>0</td> </tr> <tr> <td>Pressure score</td> <td>4</td> <td>1</td> </tr> <tr> <td>Wound infection</td> <td>1</td> <td>2</td> </tr> <tr> <td>Wound haematoma</td> <td>0</td> <td>2</td> </tr> </tbody> </table>		Group	1	2	CVA	4	0	Bronchopneumonia	1	3	Pulmonary embolism	1	0	Pressure score	4	1	Wound infection	1	2	Wound haematoma	0	2
			Group	1		2																				
CVA	4	0																								
Bronchopneumonia	1	3																								
Pulmonary embolism	1	0																								
Pressure score	4	1																								
Wound infection	1	2																								
Wound haematoma	0	2																								
Accommodation Before injury	Home Group 1: 32 Group 2: 24 Non-institution Group 1: 3 Group 2: 8 Non-hospital institution Group 1: 9 Group 2: 13 Hospital Group 1: 5 Group 2: 6																									
Accommodation Latest review (at least 6 months post op)	Home Group 1: 24 Group 2: 18 Non-institution Group 1: 2 Group 2: 4																									

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Anaesthesia: Spinal = 6 General = 43 <u>Group 2: Dynamic hip screw (DHS)</u> No. randomised: 51 Mean age: 82.7 M/F: 7/44 Other factors: ASA score: I = 2 II = 22 III = 16 IV = 11 Fracture type: Stable: 23 Unstable: 28 Anaesthesia: Spinal = 7 General = 44			Non-hospital institution Group 1: 3 Group 2: 15 Hospital Group 1: 11 Group 2: 3	
			Mobility (before injury)	Unaided Group 1: 31 Group 2: 25 Sticks Group 1: 6 Group 2: 16 Frame Group 1: 17 Group 2: 9 Non-walker Group 1: 5 Group 2: 1	
			Mobility (final review)	Unaided Group 1: 7 Group 2: 11 Sticks Group 1: 24 Group 2: 9 Frame Group 1: 13 Group 2: 14 Non-walker Group 1: 13 Group 2: 3	
			Cut out	Group 1: 2 Group 2: 3	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Femoral shaft fracture	Group 1: 4 Group 2: 0	

1

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Ekstrom et al., 2007 ^{77,77}	<p>Patient group: Unstable trochanteric and subtrochanteric fractures</p> <p>Inclusion criteria: Unstable intertrochanteric proximal femoral fractures and subtrochanteric fractures.</p> <p>Exclusion criteria: Stable trochanteric fractures, high energy trauma, pathological fractures, previous surgery to the proximal femur, daily steroids of more than 10mg of prednisolone, ongoing chemotherapy, irradiation treatment, presence of degenerative osteoarthritis of the injured hip.</p> <p>Setting: 2 orthopaedic hospitals, Sweden</p> <p>All patients N: 210 No. of dropouts: 25% (7 exclusions made: 5 wrong fracture and 2 wrong treatment).</p> <p>Group 1: Proximal femoral nail No. randomised: 105</p>	<p>Spinal anaesthesia was used, although 13 patients had general anaesthesia and 1 had a combination of both.</p> <p>Group 1: Proximal femoral nail (Stratec) The nail used was a 240mm long nail with a 130 degree shaft angle. The nail was inserted according to the surgical technique recommended by the manufacturer</p> <p>Group 2: Medoff sliding plate (Medpac) Both 4 hole and 2 hole were used for trochanteric fractures, whereas only the 6 hole plates were used for subtrochanteric fractures. The locking screw set was used in all subtrochanteric fractures to prevent compression along the femoral neck. No locking set screw was used in the trochanteric fractures.</p>	<p>Mortality at 1 year</p> <p>Functional outcome – able to walk the 15m test at 6 weeks</p> <p>Functional outcome – rise from a chair without arm support (6 weeks)</p> <p>Functional outcome – rise from a chair without arm support (4 months)</p> <p>Functional outcome – rise from a chair without arm support (12 months)</p> <p>Pain while walking</p>	<p>Trochanteric Group 1: 14 Group 2: 15 Subtrochanteric Group 1: 1 Group 2: 3</p> <p>Trochanteric Group 1: 86% Group 2: 72% Subtrochanteric Group 1: 94% Group 2: 77%</p> <p>Trochanteric Group 1: 25% Group 2: 19% Subtrochanteric Group 1: 35% Group 2: 31%</p> <p>Trochanteric Group 1: 46% Group 2: 40% Subtrochanteric Group 1: 56% Group 2: 23%</p> <p>Trochanteric Group 1: 50% Group 2: 53% Subtrochanteric Group 1: 60% Group 2: 50%</p> <p>Trochanteric</p>	<p>Funding: Not reported</p> <p>Outcomes not reported: Length of stay in hospital</p> <p>Additional outcomes reported: Operative details, ability to climb a curb, living conditions, union, minor complications.</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>No. of dropouts: 0 Age (SD): 82 (48-96) M/F: 24/76 Other factors: Trochanteric =86 Jensen-Michaelsen (JM) JM 3: 16% JM 4: 10% JM 5: 56% Subtrochanteric = 19 Seinsheimer (S) S3: 1% S4: 8% S5: 9%</p> <p><u>Group 2: Medoff sliding plate</u> No. randomised: 98 No. of dropouts: 0 Mean age (SD): 82 (52-97) M/F: 25/75 Other factors: Trochanteric = 85 Jensen-Michaelsen (JM) JM 3: 11% JM 4: 19% JM 5: 57% Subtrochanteric = 13 Seinsheimer (S) S3: 5% S4: 1%</p>	<p>All patients received preoperative iv antibiotics with 2g of cloxacillin. Subcutaneous low molecular weight heparin was used as thromboembolic prophylaxis for 7 days.</p>	<p>at 6 weeks (assesses using a visual analogue scale – VAS 0-100)</p> <p>Pain while walking at 4 months (assesses using a visual analogue scale – VAS 0-100)</p> <p>Pain while walking at 12 months (assesses using a visual analogue scale – VAS 0-100)</p> <p>Complications: Femoral fracture</p> <p>Complications: cut out</p>	<p>Group 1: 30 Group 2: 30 <u>Subtrochanteric</u> Group 1: 30 Group 2: 25</p> <p><u>Trochanteric</u> Group 1: 20 Group 2: 20 <u>Subtrochanteric</u> Group 1: 0 Group 2: 20</p> <p><u>Trochanteric</u> Group 1: 0 Group 2: 0 <u>Subtrochanteric</u> Group 1: 0 Group 2: 0.5</p> <p><u>Trochanteric</u> Group 1: 1 Group 2: 0 <u>Subtrochanteric</u> Group 1: 0 Group 2: 0</p> <p><u>Trochanteric</u> Group 1: 5 Group 2: 1 <u>Subtrochanteric</u> Group 1: 1 Group 2: 1</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	S5: 7%		Complications: femoral neck fracture	<u>Trochanteric</u> Group 1: 0 Group 2: 0 <u>Subtrochanteric</u> Group 1: 1 Group 2: 0	
			Complications: Non union	<u>Trochanteric</u> Group 1: 0 Group 2: 1 <u>Subtrochanteric</u> Group 1: 0 Group 2: 1	
			reoperations	<u>Trochanteric</u> Group 1: 6 Group 2: 1 <u>Subtrochanteric</u> Group 1: 3 Group 2: 0	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Guyer et al., 1993A^{127,128}</p> <p>Country of study: Switzerland</p> <p>Study design: Randomised prospective comparison</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: Petrochanteric and intertrochanteric fractures</p> <p>Exclusion criteria: Not reported</p> <p>Setting: Orthopaedic hospital, Switzerland</p> <p>All patients N: 100 No. of dropouts: 0</p> <p>Group 1: Gamma nail No. randomised: 50 No. of dropouts: 10 lost to follow up Age (SD): 79.5 M/F: 82% women Other factors: Fracture stability: Petrochanteric: Stable: 23 Unstable: 24 Intertrochanteric: 3</p> <p>Group 2: Dynamic hip screw No. randomised: 50 No. of dropouts: 14 lost to follow up Mean age (SD): 80.3 M/F: 88% women Other factors:</p>	<p>All patients were operated on within 24 hours where possible.</p> <p>Group 1: Gamma nail The greater trochanter was exposed after standard intramedullary technique and the entry point was holed with the awl. 12 mm diameter nails used in 44 cases and 14mm in 6 cases.</p> <p>Group 2: Dynamic hip screw 135° 4 to 12 hole plates were used.</p> <p>All patients received prophylactic cephalosporin and low dose heparin.</p>	Mortality (termed lethality in the study)	<p>30 days Group 1: 4 Group 2: 2 Late lethality (not defined) Group 1: 4 Group 2: 5</p>	<p>Funding: not reported</p> <p>Limitations: Allocation concealment unclear.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Operative details including blood loss and length of surgery, leg shortening, social situation</p>
			Length of stay in hospital (excluding those who died in hospital)	Group 1: 30.9 Group 2: 30.9	
			Reoperation	Group 1: 5/50 Group 2: 6/50	
			Complications	<p>Cranial screw perforation (cut out) Group 1: 1 Group 2: 3 Intra op femoral fragmentation Group 1: 1 Group 2: 0 Wound haematoma Group 1: 2 Group 2: 2 Deep wound infection Group 1: 0 Group 2: 1</p>	
			Pain during walking (12 weeks)	Group 1: 19/28 Group 2: 18/32	
Walking capacity (12 weeks)	<p>Full Group 1: 4/28 Group 2: 6/32 More than 1 hr Group 1: 13/28</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Fracture stability: Pertrochanteric: Stable: 19 Unstable: 26 Intertrochanteric: 5			Group 2: 16/32 Less than 1 hr Group 1: 11/28 Group 2: 10/32	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Hardy et al., 1998^{134,134}</p> <p>Country of study: Belgium, Brussels</p> <p>Study design: Randomised prospective comparison</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: At least 6 months</p>	<p>Patient group: Trochanteric proximal femoral fractures</p> <p>Inclusion criteria:</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients aged <60, pathological fractures, incorrect anatomy, history of fracture or operation involving same limb. <p>All patients N: 100 No. of dropouts: 0</p> <p>Group 1: Compression hip-screw No. randomised: 50 No. of dropouts: 0 Age (SD): 79.5 (±10.7) M/F: 15/35 Other factors: Fracture stability: Stable: 16 Unstable: 34</p> <p>ASA score: I = 5 II = 13 III = 18</p>	<p>Group 1: Compression hip-screw The compression hip-screw with a plate was inserted with a standard technique by means of a straight lateral incision on the lateral aspect of the thigh, as described by Clawson*. The barrel of the plate was at a 135 degree angle in each patient.</p> <p>Group 2: Intramedullary hip screw A cannulated intramedullary nail with a 4 degree mediolateral bend to allow insertion through the greater trochanter. The nail is 21 cm long and available in 3 diameters (12, 14 and 16 mm). The opening for the lag-screw is available in 2 angles (130 and 135 degrees). It can be locked with one or 2 4.5 mm diameter interlocking screws. A keyed centering sleeve, which is held by a set-screw, passes through</p>	<p>Mobility score (Parker and Palmer) Ability to walk indoors (SD)</p> <p>Mobility score (Parker and Palmer) Ability to walk outdoors (SD)</p>	<p>Pre op Group 1: 2.3 (0.8) Group 2: 2.4 (0.8)</p> <p>1 month Group 1: 0.9 (0.6)*** Group 2: 1.9 (0.7)***</p> <p>6 month Group 1: 1.5 (1.1) Group 2: 1.9 (1.0)</p> <p>12 month Group 1: 1.6 (1.2) Group 2: 1.9 (1.0) *** p<0.01</p> <p>Pre op Group 1: 2.1 (2.3) Group 2: 3.0 (2.6)</p> <p>1 month Group 1: 0.3 (0.7)** Group 2: 0.7 (0.9)**</p> <p>6 month Group 1: 1.7 (2.2)* Group 2: 2.7 (2.1)*</p> <p>12 month Group 1: 1.7 (2.2)* Group 2: 2.8 (2.2)* * p=0.05</p>	<p>Funding: Smith and Nephew Richards, Memphis, Tennessee</p> <p>Limitations: Allocation concealment unclear.</p> <p>Outcomes not reported: Reoperations, length of stay in hospital.</p> <p>Additional outcomes reported: Operative data e.g. time, blood loss. Sliding of lag screw.</p> <p>Notes: The fractures healed in all but one of the seventy patients who were still alive at 12 months. The one non-union was in a patient who had a compression hip</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	IV = 13 V = 1 Jenson index 1 = 10 2 = 7 3 = 7 4 = 26 Anaesthesia: Spinal: 36 General: 14 <u>Group 2: Intramedullary hip screw</u> No. randomised: 50 No. of dropouts: 0 Mean age (SD): 81.7 (± 11.8) M/F: 8/42 Other factors: Fracture stability: Stable: 13 Unstable: 37 ASA score: I = 5 II = 12 III = 23 IV = 10 V = 0 Jenson index 1 = 11 2 = 10	the intramedullary nail and over the lag-screw. The sleeve helps to prevent rotation while allowing the lag-screw to slide freely. Piritramide was administered postoperatively and paracetamol given in the recovery period. Patients were permitted to get out of bed and sit in a chair on the second day after operation and bear full weight on the fourth day.	** 0.01<p<0.05 *** p<0.01 Perioperative complications Bronchpneumonia: Group 1: 6 Group 2: 4 Cardiac failure Group 1: 5 Group 2: 7 Mortality 3 months Group 1: 12/50 Group 2: 13/50 6 months Group 1: 13/50 Group 2: 13/50 1 year Group 1: 15/50 Group 2: 15/50 Pain in hip whilst walking (4 point scale, 1 = no pain, 2 = slight pain that does not effect ability tp walk, 3 = moderate pain that that effects ability to walk, 4 – severe intractable pain even in bed) 3 months Group 1: 7/40 Group 2: 4/37 1 year Group 1: 2/35 Group 2: 2/35 Pain in mid portion of thigh while 1 year Group 1: 2/35	screw.	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	3 = 5 4 = 24 Anaesthesia: Spinal: 36 General: 14		walking, resulting in inability to walk (4 point scale, 1 = no pain, 2 = slight pain that does not effect ability tp walk, 3 = moderate pain that that effects ability to walk, 4 – severe intractable pain even in bed)	Group 2: 7/35	
			Cut-out	Group 1: 0 Group 2: 1	

*Clawson DK. Trochanteric fractures treated by the sliding screw plate fixation method. J. Trauma, 4:737-752, 1964.

1
2

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Harrington et al., 2002^{137,137}</p> <p>Country of study: UK</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Unstable trochanteric proximal femoral fractures <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients aged <65 years, pathological fractures, previous fractures, other fracture. Patients with dementia who were unable to give informed consent were excluded <p>All patients N: 102 No. lost to follow up: not reported</p> <p>Group 1: Compression hip screw No. randomised:52 No. of dropouts: 0 Mean age (SD): 82.1 (8.6) M/F: 11/41 Other factors: ASA score I: 4 II: 20</p>	<p>Group 1: Compression hip screw</p> <p>Group 2 Intramedullary hip screw</p> <p>The nail is 21cm long with a 4 degree valgus angulation and distal locking screws measuring 4.5mm in diameter. A 12 mm diameter nail and 2 locking screws were used for distal locking were used in all patients.</p> <p>Additional non-comparative prophylaxis: n/a</p>	<p>Post-op stay, days (SD)</p>	<p>Group 1: 16.3 (7.5) Group 2:16.5 (8.8)</p>	<p>Funding: Not reported</p> <p>Limitations: Reference made to some surgeons who had only used the IMHS on bone model sessions.</p> <p>Outcomes not reported: Reoperation, length of stay in hospital, functional status, pain.</p> <p>Additional outcomes reported: Operative details, ambulatory status</p> <p>Notes:</p>
			<p>Mortality in hospital</p>	<p>Group 1: 2/52 Group 2: 4/50</p>	
			<p>Ambulatory status at 1 year (retained pre injury living status)</p>	<p>Group 1: 22/33 Group 2: 19/30</p>	
			<p>Technical complications</p>	<p>Group 1: Screw cut out = 1 Barrel-plate pulled off femur = 1 Group 2: Screw cut out = 1 Intraoperative fracture propagation= 1 Late fracture of femoral shaft = 1</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>III: 17 IV: 11 V: 0</p> <p>Anaesthesia: Spinal: 34 General: 18</p> <p><u>Group 2: Intramedullary hip screw</u> No. randomised: 50 No. of dropouts: 0 Mean age (SD): 83.8 (8.5) M/F: 10/40 Other factors: ASA score I: 3 II: 22 III: 16 IV: 9 V: 0</p> <p>Anaesthesia: Spinal: 35 General: 15</p>				

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hoffman et al., 1996 ^{147,147}	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, New Zealand</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Trochanteric proximal femoral fractures • Patients aged >50 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Pathological fractures excluded <p>All patients N: 69</p> <p>No. lost to follow up: none</p> <p>Died before surgery: 2</p> <p>Mean age: 81 years</p> <p>Group 1: Ambi hip screw No. randomised:36 Mean age (SD): 79.0 (10.4) M/F: 12/24</p> <p>Other factors:</p> <p>ASA score: II: 18 III: 15 IV: 3 V: 0</p>	<p>The selected device was inserted following a detailed operative protocol based on the manufacturer's guidelines.</p> <p>Group 1: Ambi hip screw</p> <p>Group 2 Gamma nail</p> <p>The Gamma nail was interlocked in all cases initially, as recommended, but after the first 5 cases locking was reserved for unstable fractures and in line with manufacturer's updated recommendation.</p> <p>No cases were locked after patient number 50.</p> <p>Antibiotic prophylaxis (IV</p>	<p>Delay to surgery (SD)</p> <p>Total hospital stay (SD)</p> <p>Postoperative stay (SD)</p> <p>Postoperative complications</p> <p>Fracture union (% united)</p>	<p>Group 1: 1.9 (± 1.4) Group 2: 1.6 (± 1.1)</p> <p>Group 1: 30.3 (±18.9) Group 2: 31.4 (± 19.7)</p> <p>Group 1: 28.5 (±18.9) Group 2: 29.8 (±20.1)</p> <p>CVA Group 1: 1 Group 2: 1</p> <p>Cardiac Group 1: 3 Group 2: 2</p> <p>Pressure areas Group 1: 1 Group 2: 0</p> <p>Pneumonia Group 1: 1 Group 2: 1</p> <p>DVT Group 1: 0 Group 2: 1</p> <p>6 weeks Group 1: 38 Group 2: 32</p> <p>12 weeks Group 1: 79</p>	<p>Funding: Not reported</p> <p>Limitations: The manufacturer's guidelines were modified during the course of the study for the Gamma nail.</p> <p>Outcomes not reported: Reoperations, functional status.</p> <p>Additional outcomes reported: Intraoperative complications</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Anaesthesia: Spinal: 11 General: 25 Fracture stability: Unstable: 12 Stable: 24 Group 2: Gamma nail No. randomised: 31 Mean age (SD): 83.2 (8.1) M/F: 4/27 Other factors: ASA score: II: 10 III: 15 IV: 5 V: 1 Anaesthesia: Spinal: 6 General: 25 Fracture stability: Unstable: 10 Stable: 21	cephradine – 1g) prior to induction of anaesthesia.		Group 2: 79 26 weeks Group 1: 96 Group 2: 96	
			Resolution of hip pain (% without pain)	2 weeks Group 1: 52 Group 2: 48 6 weeks Group 1: 55 Group 2: 67 12 weeks Group 1: 75 Group 2: 37 26 weeks Group 1: 71 Group 2: 60	
			intra-operative fracture	Group 1: 0 Group 2: 3	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Leung et al., 1992^{191,191}</p> <p>Country of study: Hong Kong</p> <p>Study design: Randomised prospective comparison</p> <p>Duration of follow-up: 7 months</p>	<p>Patient group: Trochanteric proximal femoral fractures</p> <p>Inclusion criteria: Patients over 65 years with pertrochanteric fractures (including subtrochanteric extensions).</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Pure subtrochanteric fractures were excluded. <p>Setting: Orthopaedic hospitals</p> <p>All patients N: 225 (226 fractures) No. of dropouts: 0</p> <p>Group 1: Gamma nail No. randomised: 93 No. of dropouts: 0 Age (SD): 80.9 (±8.41) M/F: 25/68</p> <p>Other factors: ASA grade 1:15 2:47 3:23 4:8</p> <p>Fracture stability:</p>	<p>Group 1: Gamma nail 2mm Kirschner wire passed percutaneously, anterior to the femoral shaft and parallel to the femoral neck. 6 to 8cm incision made above the tip of the greater trochanter and then the medullary canal is entered. The cavity is reamed 1mm larger than the diameter of the intended nail. The nail is passed into the canal, without hammering, and the corresponding device is assembled on the nail mount and the lateral cortex of the femur is perforated by the awl and the lag screw guide wire is inserted. Distal locking is indicated for unstable fractures.</p> <p>Group 2: Dynamic hip screw Inserted using the standard technique.</p>	<p>Mortality</p> <p>Mean duration of hospital stay (acute hospital) in days (SD)</p> <p>Mean duration of hospital stay (convalescent hospital) in days (SD)</p> <p>mean time to full weight bearing (SD)</p>	<p>4 weeks Group 1: 7 Group 2: 5</p> <p>6 months Group 1: 13 Group 2: 15</p> <p>Group 1 n = 93 (30 stable, 63 unstable) Group 2 n = 93 (20 stable, 73 unstable)</p> <p>Stable Group 1: 9.2 (6.43) Group 2: 10.7 (6.27)</p> <p>Unstable Group 1: 9.5 (3.38) Group 2: 9.6 (4.46)</p> <p>Stable Group 1: 17.7 (11.97) Group 2: 15.4 (10.86)</p> <p>Unstable Group 1: 15.9 (8.2) Group 2: 19.1 (10.34)</p> <p>Stable Group 1: 1.3 (0.88) Group 2: 1.9 (0.89)</p> <p>Unstable Group 1: 1.2 (0.64) Group 2: 1.7 (0.76)</p>	<p>Funding: Not reported</p> <p>Limitations: Allocation concealment unclear.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Operative details intra-operative complications, mean sliding of lag screws, shortening, external rotation.</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Stable: 30 Unstable: 63 <u>Group 2: Dynamic hip screw</u> No. randomised: 93 No. of dropouts: 0 Mean age (SD): 78.3 (±9.46) M/F: 30/63 ASA grade 1:10 2:42 3:38 4:3 Fracture stability: Stable: 20 Unstable: 73		Postoperative mobility	Stable Independent Group 1: 12 (40%) Group 2: 8 (40%) Aided Group 1: 11 (36.7%) Group 2: 11 (55%) Chair/bed bound Group 1: 7 (23.3%) Group 2: 1 (5%) Unstable Independent Group 1: 22 (34.9%) Group 2: 23 (31.5%) Aided Group 1: 36 (57.1%) Group 2: 42 (57.5%) Chair/bed bound Group 1: 5 (8%) Group 2: 8 (11%)	
			Pain in hip	Stable Group 1: 8 (26.7%) Group 2: 5 (25%) Unstable Group 1: 14 (22.2%) Group 2: 27 (40%)	
			Pain in thigh	Stable Group 1: 4 (13.4%) Group 2: 5 (25%) Unstable	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 7 (11.1%) Group 2: 3 (4.1%)	
			Non union	Stable Group 1: 1 Group 2: 0 Unstable Group 1: 0 Group 2: 0	
			Postoperative complications	Infection Group 1: 1 Group 2: 3 Superior cutting out Group 1: 2 Group 2: 3 Fracture of shaft Group 1: 2 Group 2: 0	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Little et al., 2008^{195,195}</p> <p>Country of study: England</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Inclusion criteria: Patients presenting to the Accident and Emergency department with an extracapsular intertrochanteric fracture</p> <p>Exclusion criteria: Patients with subtrochanteric extensions of the fracture were excluded.</p> <p>All patients N: 190 No. lost to follow up: 0 Mean age: 83.4 (50 to 102)</p> <p>Group 1: Holland nail No. randomised: 92 Mean age (range): 82.6 (54 to 102) M/F: 8/84 ASA score: 1= 2 (2.2%), 2= 57 (62.0%) 3= 33 (35.8%), 4= 0</p> <p>Group 2: Dynamic hip screw No. randomised: 98 Mean age (range): 84.2 (50 to 98) M/F: 20/78 ASA score: 1= 3 (3.1%), 2= 55 (56.1%) 3= 37 (37.7%), 4= 3 (3.1%)</p>	<p>A standard operative technique either recommended by manufacturer's guidelines or as detailed in previous studies was used.</p> <p>Group 1: Holland nail (long trochanteric-entry intramedullary nail)</p> <p>Group 2 Gamma nail</p> <p>Each patient was given a single-dose antibiotic teicoplanin and gentamicin induction.</p>	Mortality	<p>30 day Group 1: 7/92 (7.6%) Group 2: 6/98 (6.1%)</p> <p>1 year Group 1: 16/92 (17.4%) Group 2: 17/98 (17.3%)</p>	<p>Funding: Not reported</p> <p>Limitations:</p> <p>Outcomes not reported: Reoperation, length of stay in hospital, pain</p> <p>Additional outcomes reported: Intra-operative variables</p> <p>Notes: 2 implant failures in group II. The proximal screws migrated laterally in 4 patients in group I.</p>
			Time to frame in days (95% CI)	<p>Group 1: 3.6 (3.3 to 3.9) Group 2: 4.23 (3.9 to 4.8) p = 0.012</p>	
			Patients with wound infections (%) None were reopened and all healed within 6 weeks	<p>Group 1: 5 (5.4) Group 2: 10 (10.2) p = 0.286</p>	
			Mobility at 1 year (95% CI)	<p>Group 1: 5.9 (5.3 to 6.5) Group 2: 3.8 (3.3 to 4.3) p <0.001</p>	
			Patients with mobility restored at 1year (%)	<p>Group 1: 49 (64) Group 2: 30 (37) p <0.001</p>	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size			Comments																																					
<p>Miedel et al., 2005^{214,214}</p> <p>Country of study: Sweden</p> <p>Study design: Randomised prospective comparison</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Unstable trochanteric and subtrochanteric proximal femoral fractures</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Acute unstable trochanteric (J-M type 3-5) or subtrochanteric fractures after a simple fall. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Pathological fractures, rheumatoid arthritis or osteoarthritis were excluded. Fractures extending more than 5cm distal to the lesser trochanter were excluded. <p>All patients N: 217 Lost to follow up: 3</p> <p>Group 1: Standard gamma nail No. randomised: 109 No. of dropouts: 0 Age (SEM): 84.6 (±0.6) M/F: 17/92 Other factors: Fracture type: Trochanteric 93</p>	<p>Group 1: Standard gamma nail</p> <p>Diameter 11mm, length 200mm, valgus bend 10°, neck angle 125 or 130° (Stryker Howmedica, Malmo, Sweden). Nails were inserted by hand and not by hammering and not to use the awl before drilling for the distal locking screw.</p> <p>Group 2: Medoff sliding plate</p> <p>Neck angle 135°, 6 hole plate (Swemac, Linkoping, Sweden). Used in the biaxial dynamisation mode, which allows sliding along both the femoral neck and shaft.</p> <p>All patients were given low-molecular weight heparin before and for approximately 10 to 14 days before operation and one dose of cefuroxim before operation.</p>	<p>Technical failures</p> <p>Trochanteric</p> <p>No complication 87 Penetration of lag screw 3 Redisplacement/medialisation 0 intra-operative femoral fracture 3 Deep infection 0</p> <p>Subtrochanteric</p> <p>No complication 16 Penetration of lag screw 0 Redisplacement/medialisation 0 intra-operative femoral fracture 0 Deep infection 0</p> <p>Reoperation</p> <p>Trochanteric Group 1: 3 Group 2: 6</p> <p>Subtrochanteric Group 1: 0 Group 2: 3</p>	<table border="1"> <thead> <tr> <th></th> <th>Grp1</th> <th>Grp2</th> </tr> </thead> <tbody> <tr> <td>Trochanteric</td> <td></td> <td></td> </tr> <tr> <td>No complication</td> <td>87</td> <td>91</td> </tr> <tr> <td>Penetration of lag screw</td> <td>3</td> <td>4</td> </tr> <tr> <td>Redisplacement/medialisation</td> <td>0</td> <td>1</td> </tr> <tr> <td>intra-operative femoral fracture</td> <td>3</td> <td>0</td> </tr> <tr> <td>Deep infection</td> <td>0</td> <td>1</td> </tr> <tr> <td>Subtrochanteric</td> <td>Grp1</td> <td>Grp2</td> </tr> <tr> <td>No complication</td> <td>16</td> <td>10</td> </tr> <tr> <td>Penetration of lag screw</td> <td>0</td> <td>0</td> </tr> <tr> <td>Redisplacement/medialisation</td> <td>0</td> <td>2</td> </tr> <tr> <td>intra-operative femoral fracture</td> <td>0</td> <td>0</td> </tr> <tr> <td>Deep infection</td> <td>0</td> <td>1</td> </tr> </tbody> </table>		Grp1	Grp2	Trochanteric			No complication	87	91	Penetration of lag screw	3	4	Redisplacement/medialisation	0	1	intra-operative femoral fracture	3	0	Deep infection	0	1	Subtrochanteric	Grp1	Grp2	No complication	16	10	Penetration of lag screw	0	0	Redisplacement/medialisation	0	2	intra-operative femoral fracture	0	0	Deep infection	0	1	<p>Funding: Grants received from the Trygg-Hansa Insurance company, the Swedish Orthopaedic Association and, in equal parts from Stryker Howmedica and Swemac.</p> <p>Outcomes not reported: Mortality, length of stay in hospital, place of residence, pain.</p> <p>Additional outcomes reported: Some outcomes grouped together (e.g. not reported separately for trochanteric and subtrochanteric) such as length of stay in hospital, HRQOL (EQO-5D), operative data, pain</p> <p>Notes:</p>
	Grp1	Grp2																																										
Trochanteric																																												
No complication	87	91																																										
Penetration of lag screw	3	4																																										
Redisplacement/medialisation	0	1																																										
intra-operative femoral fracture	3	0																																										
Deep infection	0	1																																										
Subtrochanteric	Grp1	Grp2																																										
No complication	16	10																																										
Penetration of lag screw	0	0																																										
Redisplacement/medialisation	0	2																																										
intra-operative femoral fracture	0	0																																										
Deep infection	0	1																																										

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>J-M 3: 12 J-M 4: 28 J-M 5: 53</p> <p>Subtrochanteric 16</p> <p>S2B: 1 S2C: 11 S3A: 3 S3B: 1 S4: 0 S5: 0</p> <p><u>Group 2: Medoff sliding plate</u></p> <p>No. randomised: 108 No. of dropouts: 0 Mean age (SEM): 82.7 (± 0.6) M/F: 24/84 Other factors: Fracture type: Trochanteric 96</p> <p>J-M 3: 11 J-M 4: 24 J-M 5: 61</p> <p>Subtrochanteric 12</p> <p>S2B: 0 S2C: 6 S3A: 2 S3B: 1 S4: 1 S5: 2</p>				

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>O'Brien et al., 1995²⁴⁴</p> <p>Country of study: Canada</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 52 weeks</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting:</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients with intertrochanteric fractures of the femur <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Fractures more than 1 week old Pathological fractures Subtrochanteric fractures. <p>All patients N: 101 (102 fractures) No. lost to follow up: 18%</p> <p>Group 1: Dynamic hip screw No. randomised: 49 Mean age (range): 77 (39 to 94) M/F: 17/32 Other factors: Fracture stability: Unstable: 21 Stable: 28</p> <p>Group 2: Gamma nail No. randomised: 53 Mean age (range): 83 (57 to 95))</p>	<p>The standard operative technique for fracture fixation was followed.</p> <p>Group 1: Dynamic hip screw The 135 degree four hole DHS was used more than 80% of the time in this group.</p> <p>Group 2 Gamma nail 130 or 135 degree nails were used 86% of the time. 88% of nails were distally locked.</p> <p>All but 4 patients received prophylactic antibiotic coverage with cefazolin intravenously.</p>	<p>Length of hospital stay, range (median), days</p>	<p>Orthopaedic ward Group 1: 4 – 102 (16) Group 2: 3 – 52 (14)</p> <p>Total hospital stay Group 1: 4 – 108 (18) Group 2: 3 – 92 (16)</p>	<p>Funding: Not reported</p> <p>Limitations: Mortality rate could be higher as the number of people lost to follow up is unclear</p> <p>Outcomes not reported: Functional status, place of residence,</p> <p>Additional outcomes reported: Blood loss and fluid replacement., length of surgery, early (in hospital) general complications</p> <p>Notes:</p>
			<p>Early (in hospital) local complications</p>	<p>Superficial wound infection Group 1: 1 Group 2: 0</p> <p>Wound haematoma Group 1: 0 Group 2: 1</p> <p>Malalignment Group 1: 0 Group 2: 1</p> <p>Early failure of fixation Group 1: 0 Group 2: 2</p> <p>Intraoperative fracture Group 1: 0 Group 2: 2</p> <p>Neuropraxia Group 1: 2 Group 2: 0</p>	
			<p>Late local complications</p>	<p>Failure of fixation Group 1: 1</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>M/F: 9/43 Other factors: Fracture stability: Unstable: 23 Stable: 30</p>		<p>Complications requiring reoperation</p>	<p>Group 2: 1</p> <p>Femoral shaft fracture Group 1: 0 Group 2: 1</p> <p>Varus malunion Group 1: 3 Group 2: 5</p> <p>Varus collapse with pain Group 1: 0 Group 2: 2</p> <p>Varus collapse with malunion Group 1: 1 Group 2: 0</p> <p>Failure of fixation (cut-out) Group 1: 1 Group 2: 2</p> <p>Femoral shaft fracture Group 1: 0 Group 2: 1</p>	
			<p>Mortality (early postoperative)</p>	<p>Group 1: 1 Group 2: 6</p>	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ovesen et al., 1996^{251,251}</p> <p>Country of study: Denmark</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, Odense, Denmark</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients with Intertrochanteric fractures having given informed consent. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Subtrochanteric or pathological fractures Secondary exclusions included wrong diagnosis and transfer to hospitals outside the inclusion area. <p>All patients N: 150 (101 fractures) No. lost to follow up: 17%</p> <p>Group 1: Dynamic hip screw No. randomised: 73 Mean age (sd): 78.5 (±11.7) M/F: 21/52 Other factors: lost to follow up = 4</p>	<p>Group 1: Dynamic hip screw (DHS) The use of a trochanteric stabilizing plate in combination with the DHS was allowed, but only used in 2 patients.</p> <p>Group 2 Gamma nail The distal femur was reamed 13 mm and the proximal femur to 18 mm. The use of a hammer during insertion was avoided.</p> <p>Additional non-comparative prophylaxis: Prophylaxis against DVT and pulmonary embolism consisting of Enoxaparine 40 mg once daily starting at admission until mobilisation,</p>	Mortality	<p>4 months Group 1: 3/66 Group 2: 3/67</p> <p>12 months Group 1: 3/56 Group 2: 3/59</p>	<p>Funding: Not reported</p> <p>Limitations: Surgeon experience may cause bias as operations were by surgical team on call – 49 surgeons participated in the trial.</p> <p>Outcomes not reported: Place of residence, pain</p> <p>Additional outcomes reported: Intraoperative details</p> <p>Notes:</p>
			Reoperation by 12 months	<p>Group 1: 6 Group 2: 12</p>	
			Walking aids pre fracture	<p>Sticks, crutches or no walking aid Group 1: 50 Group 2: 50</p> <p>Walking frame or wheelchair Group 1: 22 Group 2: 22</p> <p>Missing or deceased Group 1: 1 Group 2: 1</p> <p>p = 0.41</p>	
			Walking aids at discharge	<p>Sticks, crutches or no walking aid Group 1: 22 Group 2: 13</p> <p>Walking frame or wheelchair Group 1: 47 Group 2: 59</p> <p>Missing or deceased</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	ASA score: 1 = 19 2 = 18 3 = 26 4 = 10 Group 2: Trochanteric gamma nail No. randomised: 73 Mean age (sd): 79.9 (±10) M/F: 20/53 Other factors: lost to follow up = 11 ASA score: 1 = 20 2 = 21 3 = 25 4 = 7	discharge or for 7 days. Antibiotic prophylaxis was also given.		Group 1: 4 Group 2: 1 p = 0.03	
			Walking aids at 4 months	Sticks, crutches or no walking aid Group 1: 43 Group 2: 37 Walking frame or wheelchair Group 1: 23 Group 2: 30 Missing or deceased Group 1: 7 Group 2: 6 p = 0.14	
			Complications requiring reoperation	Group 1: Cut- out = 2 Redislocation = 3 Femoral fracture = 0 Infection = 1 Haematoma = 0 Group 2: Cut- out = 7 Redislocation = 0 Femoral fracture = 2 Infection = 2 Haematoma = 1	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments	
Pajarinen et al., 2005 (Also Pajarinen 2004) ^{254,255} Country of study: Finland Study design: Prospective randomized study List who was masked to interventions: Not reported. Duration of follow-up: 4 months	Patient group: Patients with hip fracture Setting: Orthopaedic hospital, Helsinki, Finland Inclusion criteria: <ul style="list-style-type: none"> Low energy extracapsular pertrochanteric femoral fractures Exclusion criteria: <ul style="list-style-type: none"> Pathological fractures, multiple injuries, and those unable to give informed consent were excluded. All patients N: 108 No. lost to follow up: 15 (14%) Group 1: Dynamic hip screw No. randomised: 54 Mean age (sd): 80.3 (±10.8) M/F: 14/40 Other factors: ASA score: 2 = 8 3 = 32 4 = 14 Anaesthetic: General = 2	All operations were performed within 2 days of admission, in most cases by a senior orthopaedic resident. Standard operative techniques, which are recommended by the manufacturers and have been described in detail in instruction manuals or earlier studies were used. Group 1: Dynamic hip screw (DHS) Group 2 Proximal femoral nail Intravenous antibiotic prophylaxis was given. Patients were also treated with a low-molecular weight	Mean hospitalisation time in days (sd) Group 1: 5.4 (3) Group 2: 6.1 (3.3) $p = 0.251$	Group 1: 5.4 (3) Group 2: 6.1 (3.3) $p = 0.251$ Own home Group 1: 4 (7.4) Group 2: 6 (11.1) Nursing home Group 1: 2 (3.7) Group 2: 1 (1.9) Rehabilitation hospital Group 1: 48 (88.0) Group 2: 45 (83.3) Died at our hospital Group 1: 0 Group 2: 3.7 (0.495)	Funding: Not reported Limitations: Outcomes not reported: Pain Additional outcomes reported: Intraoperative details, radiographic findings at 4 months post-op. Notes:	
			Discharged to (%)			Place of residence at 4 months (%)
			Own home Group 1: 4 (7.4) Group 2: 6 (11.1) Nursing home Group 1: 2 (3.7) Group 2: 1 (1.9) Rehabilitation hospital Group 1: 48 (88.0) Group 2: 45 (83.3) Died at our hospital Group 1: 0 Group 2: 3.7 (0.495)			Own home Group 1: 22 (53.7) Group 2: 24 (57.1) $p = 0.827$ Nursing home Group 1: 6 (14.6) Group 2: 10 (23.8) $p = 0.405$ Institution Group 1: 13 (31.7) Group 2: 8 (19.0) $p = 0.214$

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Spinal = 52 Group 2: Proximal femoral nail No. randomised: 54 Mean age (sd): 80.9 (±9.1) M/F: 13/41 Other factors: ASA score: 2 = 6 3 = 28 4 = 20 Anaesthetic: General = 3 Spinal = 51	heparin during their stay in hospital	Recovery of abilities to pre-op status (%)	Yes Group 1: 32 (78) Group 2: 34 (81) No Group 1: 9 (22) Group 2: 8 (19) p = 0.791	
			Walking ability (%)	No aids needed Group 1: 12 (29.3) Group 2: 15 (35.7) p = 0.641 In need of aids, but independent Group 1: 22 (53.7) Group 2: 24 (57.1) p = 0.827 In need of assistance Group 1: 7 (17.1) Group 2: 3 (7.1) p = 0.194	
			Recovery of walking ability to pre-op status (%)	Yes Group 1: 22 (53.7) Group 2: 32 (76.2) No Group 1: 19 (46.3) Group 2: 10 (23.8) p = 0.040	
			Drop out patients	Fracture redisplacement (reoperation) Group 1: 2 Group 2: 2 p = 1.00 Died before follow up was complete	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 2 Group 2: 4 $p = 0.678$ Did not attend final review Group 1: 9 Group 2: 6 $p = 0.578$	

1

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Park et al., 1998 ^{258,258} Country of study: Korea Study design: Prospective randomized study List who was masked to interventions: Not reported. Duration of follow-up: 1 year	Patient group: Patients with hip fracture Setting: University Hospital, Korea Inclusion criteria: Intertrochanteric fractures of the femur. Patients aged 60 and over Exclusion criteria: Not reported All patients N: 60 No. lost to follow up: 0 Group 1: Gamma Asia Pacific nail (GAPN) No. randomised: 30 Mean age: 73.7 M/F: 10/20 Other factors: ASA score: 1 = 3 2 = 19 3 = 8 4 = 0 Fracture pattern (Tronzo) Stable (II): 14 (47%)	Group 1: Gamma Asia Pacific nail (GAPN) These were inserted using a closed technique under image intensifier control. Group 2: Compression hip screw (CHS) CHS (135°) were inserted using the standard technique.	Mean time to union (weeks)	Systemic Group 1: 14.3 Group 2: 15.1 $p = 0.06$ Stable Group 1: 14.28 Group 2: 14.55 $p = 0.73$ Unstable Group 1: 14.31 Group 2: 15.42 $p = 0.03$	Funding: Not reported Limitations: Unclear allocation concealment. Outcomes not reported: Pain, place of residence Additional outcomes reported: Operative details, decrease of neck shaft angle, length of sliding of the lag screw. Notes:
			Mobility assessment (Ceder et al)	Mean Group 1: 5.1 Group 2: 4.7 $p > 0.05$	
			Complications	Fracture of the shaft of the femur Group 1: 0 Group 2: 0 Greater trochanter fracture Group 1: 1 Group 2: 0 Fracture displaced by nail insertion Group 1: 2 Group 2: 0 Cut out	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Unstable (III & IV): 16 (53%) <u>Group 2: Compression hip screw (CHS)</u> No. randomised: 30 Mean age: 72.2 M/F: 14/16 Other factors: ASA score: 1 = 4 2 = 16 3 = 9 4 = 1 Fracture pattern (Tronzo) Stable (II): 11 (37%) Unstable (III & IV): 19 (63%)			Group 1: 1 Group 2: 1 Deep infection Group 1: 1 Group 2: 1 Non union Group 1: 0 Group 2: 1	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Radford et al., 1993^{279,279}</p> <p>Country of study: England</p> <p>Study design: Prospective randomized study</p> <p>List who was masked to interventions: Not reported.</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients aged over 60, with a pertrochanteric femoral fracture <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Not reported <p>All patients N: 200 No. lost to follow up: not stated</p> <p>Group 1: Dynamic hip screw No. randomised: 100 Mean age: 78 (60 to 90) M/F: 76/24 Other factors: Number with diabetes: 4 Unstable: 43%</p> <p>Group 2: Gamma nail No. randomised: 100 Mean age: 72.2 M/F: 14/16 Other factors:</p>	<p>The operations were performed using image intensification. For both implants they aimed to have a central position of the screw in the femoral head on both anteroposterior and lateral views, with its tip 5 to 10 mm from the subchondral bone.</p> <p>Group 1: Dynamic hip screw 4 hole 135° plate with a screw of appropriate length</p> <p>Group 2: Gamma nail A preoperative radiograph was taken of the other hip to compare with the implant template to decide the angle of the chosen nail.</p>	<p>Mortality</p> <p>Delayed wound healing or persistent discharge leading to another course of antibiotics to be given</p> <p>Infection (bacteriologically proven)</p> <p>Thromboembolism during hospital stay</p> <p>Fixation failure requiring surgical revision</p> <p>Fracture of the femoral shaft</p> <p>Fracture of the femoral shaft – requiring surgical revision</p> <p>Reoperation</p> <p>Cut-out</p> <p>Non-union</p>	<p>3 months Group 1: 10 Group 2: 12</p> <p>Group 1: 8 Group 2: 3</p> <p>3 months Group 1: 4 Group 2: 0</p> <p>Group 1: 6 Group 2: 8</p> <p>Group 1: 3 Group 2: 2</p> <p>Group 1: 1 Group 2: 11</p> <p>Group 1: 1 Group 2: 3</p> <p>Group 1: 3 Group 2: 6</p> <p>Group 1: 3 Group 2: 2</p> <p>Group 1: 0 Group 2: 0</p>	<p>Funding: Not reported</p> <p>Limitations: Includes diabetic patients. Unclear allocation concealment.</p> <p>Outcomes not reported: Pain, place of residence</p> <p>Additional outcomes reported: Prefracture mobility and housing score, femoral shaft fracture details of patients treated with gamma nails, preoperative blood loss.</p> <p>Notes: Only surgeons of registrar grade and above took part in the trial and were already experienced</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Number with diabetes: 6 Unstable: 38%	Distal locking of the nail in the femoral shaft was performed only when indicated for longitudinal instability.			in the use of the DHS. The first 2 Gamma nail operations performed by each surgeon were not included in the trial. Perioperative fractures were caused by too forceful insertion of the nail into the femoral shaft – often by hammer

1
2

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Rahme et al., 2007 ^{281,281} Country of study: Australia Study design: Randomised prospective comparison List who was masked to interventions: Not reported. Duration of follow-up: 9 months	Patient group: Subtrochanteric femoral fractures Inclusion criteria: All skeletally mature patients presenting with acute subtrochanteric fractures Exclusion criteria: Ipsilateral femoral shaft or neck fractures. All patients N: 58 No. of dropouts: 0 Group 1: Blade plate No. randomised: 29 No. of dropouts: 0 Mean age : 67 M/F: 12/17 Seinsheimer classification: Type 1: 0, Type 2: 8 Type 3: 8, Type 4: 4 Type 5: 9 Group 2: Proximal femoral nail No. randomised: 29 No. of dropouts: 0 Mean age: 73 M/F: 13/16 Seinsheimer classification: Type 1: 1, Type 2: 7 Type 3: 10, Type 4: 1 Type 5: 10	Group 1: Proximal femoral nail Treated with closed reduction using a traction table and percutaneous insertion of the nail (Synthes AG, Chur, Switzerland) without anatomic reduction. Group 2: Intramedullary hip screw Treated with open anatomic reduction, Internal fixation was achieved using a 95° angled blade plate (Synthes AG, Chur, Switzerland).	Length of stay in hospital	Group 1: 22 Group 2: 25 p=0.7	Funding: not reported Limitations: Allocation concealment unclear. Underpowered. Outcomes not reported: Pain, mobility, functional status Additional outcomes reported: mean operating time, blood transfusion, infection (all, including whether an organism was confirmed as present or not) Notes: Intention to treat analysis performed
			Non-union (absence of bridging callus on 2 radiographic views 9 months after injury)	Group 1: 8 Group 2: 1 p=0.025	
			Revision	Group 1: 8 Group 2: 0 p=0.005	
			Mortality	Group 1: 2 Group 2: 6 p=0.25	
			Infection	Group 1: 1 Group 2: 3 p=0.6	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Sadowski et al., 2002 ^{294,294} Country of study: Switzerland Study design: Prospective randomized study Duration of follow-up: 12 months	Patient group: Patients with hip fracture Setting: Orthopaedic hospital, Geneva, Switzerland. Inclusion criteria: <ul style="list-style-type: none"> Patients aged over 55, with AO/OTA 31-A3 fractures (trochanteric proximal femoral fractures) Low energy fractures Exclusion criteria: <ul style="list-style-type: none"> Patients with pathological fractures, fractures associated with polytrauma, fractures associated with polytrauma, a preexisting femoral deformity preventing hip screw osteosynthesis or intramedullary nailing, previous surgery on the ipsilateral hip or femur, and a fractures extending 5cm distal to the inferior border of the lesser trochanter. All patients N: 39	All procedures were performed by staff surgeons. Group 1: Dynamic hip screw Operative technique described by Blatter and Janssen Group 2: Gamma nail Operative technique as described in Simmermacher et al. The fracture was not exposed for nailing unless it could not be reduced with closed techniques. A 10 or 11mm diameter nail was used 18/20 patients and the lag screw measured 100 or 105mm in 10/20 patients. The proximal fragment was reamed in all	Residence - Preoperative (chi square with Yates correction)	Home Group 1: 15 Group 2: 13 Nursing home Group 1: 4 Group 2: 7 p = 0.54	Funding: Not reported Limitations: Includes diabetic patients Additional outcomes reported: Operative time, blood transfusion, difficulty of operation, type of reduction, conversion from static to dynamic construct, consolidation time. Notes:
			Postoperative data – complications (chi square)	Pneumonia Group 1: 3 Group 2: 2 Cardiac failure or infarction Group 1: 1 Group 2: 1 Cerebrovascular accident Group 1: 0 Group 2: 1 p = 0.83	
			Wound complications (chi square with Yates correction)	Group 1: 2 Group 2: 3 p = 0.95	
			Hospital stay (days) (student t test)	Group 1: 18 ± 7 Group 2: 13 ± 4 p = 0.01	
			Discharge to: (chi square)	Home Group 1: 15 Group 2: 13	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>No. lost to follow up: 1</p> <p><u>Group 1: Dynamic condylar screw</u> No. randomised: 19 Mean age: 77 (± 14) M/F: 5/14 Other factors: ASA score: 1 = 1 2 = 9 3 = 9 4 = 0 Anaesthesia: General = 10 Regional = 9</p> <p><u>Group 2: Proximal femoral nail</u> No. randomised: 20 Mean age: 80 (± 13) M/F: 7/13 Other factors: ASA score: 1 = 0 2 = 6 3 = 11 4 = 3 Anaesthesia: General = 11 Regional = 9</p>	<p>20 patients, but distal reaming was only performed on 1 patient. All of the nails were interlocked distally with 2 screws.</p> <p>All patients were given one dose of prophylactic intravenous antibiotic. In addition all patients were treated with low-molecular weight heparin</p>	<p>Status of patient at 1 year (chi square)</p> <p>Orthopaedic complications at 1 year (chi square)</p>	<p>Nursing home Group 1: 4 Group 2: 7</p> <p>Home Group 1: 15 Group 2: 13</p> <p>Nursing home Group 1: 4 Group 2: 7 p = 0.26</p> <p>Mortality Group 1: 1 Group 2: 2</p> <p>Lost to follow-up Group 1: 1 Group 2: 0</p> <p>Available for review Group 1: 17 Group 2: 18</p> <p>Implant failure Group 1: 6 Group 2: 0</p> <p>Non-union Group 1: 1 Group 2: 1</p> <p>Infection Group 1: 1</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 0 $p = 0.007$ cut-out Group 1: 5 Group 2: 0	
			Major reoperations at 1 year (chi square)	Group 1: 6 * Group 2: 0 *(1 hip prosthesis, 1 change of implant, 4 change of implant and bone graft) $p = 0.008$	
			Hip/thigh pain score at 1 year (student t test)	Group 1: 1.77 ±0.73 Group 2: 1.44 ±0.86 $p = 0.2$	
			Jenson social-function score at 1 year (student t test)	Group 1: 2.5 ±1.3 Group 2: 2.6 ±1.0 $p = 0.9$	
			Parker-and-palmer score at 1 year (student t test)	Group 1: 6.0 ±3.5 Group 2: 5.0 ±2.6 $p = 0.39$	
			Residence at 1 year (chi square)	Home Group 1: 15 Group 2: 13 Nursing home Group 1: 4 Group 2: 7	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Saudan et al., 2002^{300,300}</p> <p>Country of study: Switzerland</p> <p>Study design: Prospective randomized study</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, Geneva, Switzerland.</p> <p>Inclusion criteria: All fractures of the trochanteric region (in persons over the age of 55 years) caused by a low energy injury. Included classifications were AO/OTA Type 31-A1 or A2.</p> <p>Exclusion criteria: Pathologic fractures, fractures associated with polytrauma, a patient with previous ipsilateral hip or femur surgery, or any fractures with extension 5 cm distal to the inferior border of the lesser trochanter.</p> <p>All patients N: 206 No. lost to follow up: 4%</p> <p>Group 1: Dynamic hip screw No. randomised: 106 Mean age: 83.7 (±10.1) M/F: 22/84 Other factors: ASA score:</p>	<p>Group 1: Dynamic hip screw In 50% of patients the length of the screw was 90 or 95mm, and in almost all cases the side plate was 135° with 4 holes</p> <p>Group 2: Proximal femoral nail Operative technique as described by Simmermacher.</p> <p>All patients were given one dose of antibiotic prophylaxis preoperatively, and treated with a low-molecular weight heparin followed by Coumadin as prophylactic anticoagulation, begun after surgery and continued for 6 weeks.</p>	<p>Postoperative data – complications (chi square)</p> <p>Wound complications</p> <p>Hospital stay (days)</p> <p>Discharge to:</p>	<p>Respiratory Group 1: 7 Group 2: 7</p> <p>Cardiovascular Group 1: 9 Group 2: 5</p> <p>Pulmonary embolism Group 1: 1 Group 2: 1</p> <p>Deep vein thrombosis Group 1: 1 Group 2: 1</p> <p>Gastrointestinal Group 1: 2 Group 2: 1</p> <p>Neurologic Group 1: 1 Group 2: 2 p = 0.24</p> <p>Group 1: 10 Group 2: 11 p = 0.71</p> <p>Group 1: 14 ±10 Group 2: 13 ±4 p = 0.71</p> <p>Home</p>	<p>Funding: Not reported</p> <p>Limitations:</p> <p>Additional outcomes reported: Intraoperative data.</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	1 = 3 2 = 30 3 = 66 4 = 7 Anaesthesia: General = 37 Regional = 69 Group 2: Proximal femoral nail No. randomised: 100 Mean age: 83 (± 9.7) M/F: 24/76 Other factors: ASA score: 1 = 1 2 = 30 3 = 63 4 = 6 Anaesthesia: General = 38 Regional = 62			Group 1: 24 Group 2: 22 Nursing home/rehabilitation hospital Group 1: 78 Group 2: 74 Died in hospital Group 1: 4 Group 2: 4 $p = 0.99$	
			Status of patient at 1 year	Died Group 1: 13 Group 2: 16 Lost to follow up Group 1: 4 Group 2: 5 Available for review Group 1: 89 Group 2: 79	
			Complications at 1 year	Fixation failure (cut-out_ Group 1: 1 Group 2: 3 Non-union Group 1: 0 Group 2: 0 Infection Group 1: 1 Group 2: 3	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				p = 0.15	
			Reoperation at 1 year	<p>Hip prosthesis Group 1: 1 Group 2: 3</p> <p>Removal of implant and/or debridement Group 1: 1 Group 2: 3 p = 0.15</p>	
			Habitation	<p>Home Group 1: 50 Group 2: 37</p> <p>Nursing home Group 1: 39 Group 2: 42 p = 0.22</p>	
			Pain (score)	<p>Group 1: 1.31 ±0.63 Group 2: 1.36 ±0.63 p = 0.59</p>	
			Social function – Jensen (mean)	<p>Group 1: 2.65 ±1.14 Group 2: 2.88 ±1.16 p = 0.2</p>	
			Mobility score – Palmer/Parker (mean)	<p>Group 1: 5.07 ±2.97 Group 2: 4.94 ±3.33 p = 0.8</p>	

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Utrilla et al., 2005 ^{337,337}	<p>Patient group: Patients with hip fracture</p> <p>Setting: Orthopaedic hospital, Alicante, Spain</p> <p>Inclusion criteria: Patients aged over 65 years who sustained a trochanteric fracture of the femur.</p> <p>Exclusion criteria: Patients with subtrochanteric fractures or subtrochanteric fracture extension, pathologic fractures, history of a previous injury involving the lower limbs, and patients who had a severe concomitant medical condition (grade V ASA score).</p> <p>All patients N: 210 No. lost to follow up: 7 (3.3%)</p> <p>Group 1: Trochanteric Gamma Nail (TGN) No. randomised: 106 Mean age: 80.6 (±7.5) M/F: 38/66 Other factors: ASA score: 1 = 13</p>	<p>Fracture fixation was performed within 4 days.</p> <p>4 surgeons experienced in the standard gamma nail did all the operations, but the first 3 TGN operations performed by each surgeon were not included in the study.</p> <p>Spinal anaesthesia was performed in all but 3 patients.</p> <p>Group 1: Trochanteric Gamma Nail (TGN) This was a modification of the standard implant: shorter in length (180mm), with a lower mediolateral curvature (4°) and available only in</p>	<p>Mortality</p> <p>Walking ability (Parker/Palmer score) at 12 months</p> <p>Hip flexion (°)</p> <p>Hip pain (no.)</p>	<p>0 – 30 days Group 1: 7 Group 2: 10</p> <p>31 – 90 days Group 1: 1 Group 2: 5</p> <p>91 – 180 days Group 1: 3 Group 2: 0</p> <p>181 – 365 days Group 1: 8 Group 2: 6</p> <p>Total Group 1: 6.4 ±2.8 n= 82 Group 2: 6.2 ±2.8 n= 81 p = 0.74</p> <p>Stable Group 1: 7.6 ±2.2 Group 2: 7.3 ±2.4 p = 0.92</p> <p>Unstable Group 1: 7.0 ±2.1 Group 2: 5.8 ±2.7 p = 0.017</p> <p>Group 1: 97.9 ±10.3 Group 2: 95.6 ± 9.5 p = 0.15</p> <p>Group 1: 41</p>	<p>Funding: Not reported</p> <p>Limitations:</p> <p>Outcomes not reported: <i>List the outcomes in which we are interested that are not reported here</i></p> <p>Additional outcomes reported: Perioperative data, leg shortening</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																											
	2 = 39 3 = 41 4 = 11 Group 2: Compression hip screw No. randomised: 106 Mean age: 79.8 (± 7.3) M/F: 28/78 Other factors: ASA score: 1 = 14 2 = 35 3 = 54 4 = 3	proximal and distal diameters of 17 and 11 mm. The neck shaft angle was 130° and was inserted by a percutaneous technique. Distal locking with 1 screw only was performed on those fractures with rotational instability of the diaphyseal fragment. Group 2: Compression hip screw (CHS) The CHS was inserted using the standard technique, the implant was a 135° plate with 4 holes. All patients received antibiotic and thromboembolic prophylaxis.		Group 2: 44 $p = 0.75$																												
			Thigh pain (no.)	Group 1: 50 Group 2: 45 $p = 0.52$																												
			Postoperative complications	<table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp2</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>82</td> <td>81</td> </tr> <tr> <td>DVT</td> <td>4</td> <td>3</td> </tr> <tr> <td>Local wound</td> <td>6</td> <td>7</td> </tr> <tr> <td>Deep infection</td> <td>0</td> <td>1</td> </tr> <tr> <td>Trochanter fracture</td> <td>4</td> <td>2</td> </tr> <tr> <td>Fixation failure</td> <td>5</td> <td>6</td> </tr> <tr> <td>Cut out</td> <td>1</td> <td>2</td> </tr> <tr> <td>Reoperation</td> <td>1</td> <td>4</td> </tr> </tbody> </table>		Grp 1	Grp2	n	82	81	DVT	4	3	Local wound	6	7	Deep infection	0	1	Trochanter fracture	4	2	Fixation failure	5	6	Cut out	1	2	Reoperation	1	4	
	Grp 1	Grp2																														
n	82	81																														
DVT	4	3																														
Local wound	6	7																														
Deep infection	0	1																														
Trochanter fracture	4	2																														
Fixation failure	5	6																														
Cut out	1	2																														
Reoperation	1	4																														

1 Evidence tables – extracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Zou et al., 2009 ^{364,364} Country of study: China Study design: RCT Duration of follow-up: 1 year	Patient group: Consecutive patients with low-energy trochanteric femoral fractures Setting: Dept orthopaedic surgery, The first affiliated hospital of Soochow University, Suzhou, Jiangsu, China Inclusion criteria: Patients with 31-A1 stable trochanteric or 31-A2/31-A3 unstable trochanteric fractures. Exclusion criteria: Patients with a pathological fracture or multiple injuries were excluded. <u>All patients</u> N: 121 <u>Group 1</u> No. randomised: 63 Stable: 52 Unstable: 11 Age (mean \pmSD): 65 (34-89) M/F: 24%/76% Operative time: 93 +/- 13 mins <u>Group 2</u> No. randomised: 58 Stable: 42 Unstable: 16 Age (mean \pmSD): 65 (37-91) M/F: 21%/79%	Surgery was performed with the patient in the supine position on a fracture table, with the injured extremity slight adducted to facilitate insertion of the implant. After surgery the patients were mobilised and given standard rehabilitation instructions by a physiotherapist. <u>Group 1</u> Dynamic hip screw <u>Group 2</u> Proximal femoral nail antirotation	Femoral shaft fracture	Group 1: 0 Group 2: 0 P value(s): not significant	Funding: Not stated Limitations: Outcomes not reported: <i>The Salvati and Wilson scoring system for hip function</i> Additional outcomes reported: list additional outcomes reported in the study that we are not interested in Notes:
			Cut-out	Group 1: 0 Group 2: 0 P value(s): not significant	
			Non-union	Group 1: 1 (unstable) Group 2: 0	
			Breakage of implant	Group 1: 0 Group 2: 2 (1 unstable, 1 stable, of which 1 required reoperation) P value(s): not significant	
			Wound infection	Group 1: 1 (stable) Group 2: 1 (unstable) P value(s): not significant	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Operative time: 52 +/- 10 mins				

1

1 **17.9 Evidence Table 9: Surgical approach to hemiarthroplasty**

Study details	Patients	Exposure	Outcome measures	Effect size	Comments
Enocson et al., 2008 ^{85,85}	Patient group: Consecutive patients who had a hemiarthroplasty for non-pathological displaced femoral neck fracture Setting: Orthopaedics department Inclusion criteria: <ul style="list-style-type: none"> Not reported Exclusion criteria: <ul style="list-style-type: none"> None reported All patients N: 739 hips in 720 patients No. of dropouts: not reported Age (mean ±SD): women: 84 (54-103), men 82 (55-97) years M/F: 147/592	Surgical approach Group 1 431 operations performed by an anterolateral approach. Group 2 176 operations performed by a posterolateral approach with posterior repair. Group 3 129 operations performed by a posterolateral approach without posterior repair.	Number of dislocations	Group 1: 13/431 (3%) Group 2: 15/176 (9%) Group 2: 17/129 (13%)	Funding: None reported Limitations: Not stated how patients allocated to a surgeon. Surgical approach based on surgeon's own preference Outcomes not reported: Mortality, length of stay in secondary care, requirement for surgical revision, wound infection. Operations performed by registrars or post-registrars.
Country of study: Sweden Study design: Historical cohort List who was masked to interventions: Not applicable Duration of follow-up: Median 2.3 (0-10) years			Dislocation for posterior lateral approach with posterior repair compared to anterolateral approach. Logistic regression univariate analysis Odds ratio: 3.0 (1.4, 6.4) P=0.005 Logistic regression multivariate analysis adjusted for age, sex, indication for surgery, surgeon seniority and femoral head size Odds ratio: 3.9 (1.6, 9.8) P=0.003		
Dislocation for posterior lateral approach without posterior repair compared to anterolateral approach. Logistic regression univariate analysis Odds ratio: 4.9 (2.3, 10) P<0.001 Logistic regression multivariate analysis adjusted for age, sex, indication for surgery, surgeon seniority and femoral head size Odds ratio: 6.9 (2.6, 19) P<0.001					

2

1 Evidence tables – surgical approach to hemiarthroplasty

Study details	Patients	Exposure	Outcome measures	Effect size	Comments
<p>Parker et al., year²⁶⁹</p> <p>Country of study: UK</p> <p>Study design: Systematic review including 1 RCT</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Patients with displaced intracapsular hip fracture</p> <p>Setting: Hospital</p> <p>All patients N: 114 patients No. of dropouts: not reported</p>	<p>Surgical approach</p> <p><u>Group 1</u> 57 cemented Thompson hemiarthroplasties by an anterolateral approach.</p> <p><u>Group 2</u> 57 cemented Thompson hemiarthroplasties by posterior approach</p>	<p>Outcomes extracted</p>	<p>Results reported in forest plots for:</p> <ul style="list-style-type: none"> -Number of dislocations -Pain at 1 month -Impairment of mobility at 6 months 	<p>Funding: None reported</p> <p>Limitations: Most operations were performed by trainees with different levels of experience. No blinding of anyone reported. Unclear allocation concealment.</p> <p>Outcomes not reported: Mortality (only presented in graphs), length of stay in secondary care, reoperations (unable to work out numbers), quality of life.</p>

2

1 17.10 Evidence Table 10: Mobilisation strategies

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hauer et al., 2002 ^{139,140} Country of study: Germany Study design: <i>RCT</i> Duration of follow-up: 3 month	Patient group: Patients with hip fracture Setting: Inclusion criteria: <ul style="list-style-type: none"> Hip surgery, recent history of injurious falls, age over 75 years, female, consent of orthopaedic surgeon, patient willingness to participate in the study. Exclusion criteria: <ul style="list-style-type: none"> Acute neurological impairment, severe cardio-vascular disease, unstable chronic or terminal illness, major depression, severe cognitive impairment or severe musculo-skeletal impairment. All patients N: 28 No. of dropouts: Age (mean \pmSD): 81 (\pm 3.9) M/F: All female Group 1 No. randomised: 15	Group 1 High intensity progressive resistance training of functionally relevant muscle groups and a progressive functional training for 3 days a week for 12 weeks. Intensity of strength training was adjusted to 70-90% of the individual maximal workload. Basic functions such as walking, stepping or balancing were trained progressively with increasing complexity. Group 2 Patients in the control group met 3 times a week for 1 hour for motor placebo activities. Typical activities, which were not supposed to be relevant for the study purpose, were calisthenics, games and memory tasks whilst seated	Barthel/Mahoney activities of daily living (ADL)	Group 1: 93.0 (8.2) Group 2: 96.1 (8.2) $p = 0.636$	Funding: A grant received from the Ministerium fur Wissenschaft, Forschung und Kunst Baden-Wuerttemberg and the University of Heidelberg. Limitations: Small study size Additional outcomes reported: Further baseline characteristics. Balance score, functional reach, total activity, 'sports' activity. Household activities, emotional state Notes:
			Lawton/Brody Instrumental activities of daily living index	Group 1: 7.3 (1.4) Group 2: 6.9 (1.3) $p = 0.416$	
			Maximal dynamic and isometric muscle strength, at 3 months mean, (\pmSD)	Leg-press, fractured side 1RM (kg) Group 1: 71 (35) Group 2: 50 (21) $p = 0.021$ Leg-press, non-affected side 1RM (kg) Group 1: 88 (39) Group 2: 67 (17) $p = 0.018$ Leg-extensor, fractured side, Newton Group 1: 68 (13) Group 2: 51 (22) $p = 0.011$ Leg-extensor, non affected side, Newton Group 1: 80 (11) Group 2: 60 (20) $p = 0.006$ Leg flexor, fractured side, Newton Group 1: 37 (7) Group 2: 34 (13) $p = 0.036$	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	No. of dropouts: Age (mean \pmSD): 81.7 (\pm 7.6) M/F: All female Group adherence: 93.1 (\pm 13.5%) Group 2 No. randomised: 13 No. of dropouts: Age (mean \pmSD): 80.8 (\pm 7.0) M/F: All female Group adherence: 96.7 (\pm 6.1%)	Both groups received identical physiotherapy two times a week for 25 mins. Strength and balance training were excluded during physiotherapy and control group sessions. Physiotherapy consisted of massage, stretching and application of heat or ice.		Leg flexor, non affected side, Newton Group 1: 39 (11) Group 2: 37 (13) $p = 0.113$ Ankle plantar flexion, fractured side, Newton Group 1: 88 (30) Group 2: 65 (33) $p = 0.944$ Ankle plantar flexion, non affected side, Newton Group 1: 98 (32) Group 2: 78 (32) $p = 0.968$	
			Handgrip strength, both hands, Kilopascal	Group 1: 121 (29) Group 2: 108 (28) $p = 0.270$	
			Maximal gait speed, m/sec	Group 1: 0.72 (0.28) Group 2: 0.49 (0.15) $p = 0.121$	
			Timed up and go, (sec)	Group 1: 26.1 (17.8) Group 2: 26.9 (9.8) $p = 0.731$	
			Tinetti's performance oriented mobility assessment (POMA)	Overall Group 1: 23.5 (4.5) Group 2: 20.5 (4) $p = 0.505$ Part 1 Group 1: 12.7 (2.2) Group 2: 11.4 (2.4) $p = 0.747$ Part 2 Group 1: 10.8 (2.5) Group 2: 9.1 (2.1) $p = 0.249$	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Box step, cm	Fractured leg Group 1: 34.5 (6.4) Group 2: 30.6 (9.8) p = 0.482 Unaffected leg Group 1: 38.5 (7.8) Group 2: 34.4 (5.8) p = 0.420	

1

1 Evidence tables – mobilisation strategies

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Karumo et al., 1977^{171,171}</p> <p>Country of study: Finland</p> <p>Study design: RCT</p> <p>List who was masked to interventions: Not reported</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Patients with hip fracture</p> <p>Inclusion criteria: Consecutive patients aged over 50 with dislocated fractures of the femoral neck.</p> <p>Exclusion criteria: Inadequate follow up examination.</p> <p>All patients N: 100 Lost to follow up: 13</p> <p>Group 1 No. randomised: 23 treated with prosthesis 26 with internal fixation No. of dropouts: Age (mean \pmSD): M/F: 13/26 Subgroup category numbers: Other factors:</p> <p>Group 2 No. randomised: 16 treated with prosthesis 22 with internal fixation</p> <p>No. of dropouts: Age (mean \pmSD): M/F: 9/29</p>	<p>Group 1 – usual care Average of 30mins physiotherapy per day.</p> <p>Group 2 – Intensive Physiotherapy performed twice daily – average of 1 hour.</p> <p>Physiotherapy shame: Walking on crutches on first postoperative day with almost all allowed full weight bearing from the beginning. From first post op day training in sitting in a chair with the hip and knee joint in 90° flexion. In second postoperative week training in walking up and down stairs. Patients urged to perform extension-flexion movements of the knee joint.</p>	<p>Length of hospital stay</p> <p>Strength of the adductor muscle (9 weeks post op) – operated leg</p>	<p>Prosthesis Group 1: 33.9 (\pm20.1) Group 2: 31.8 (\pm19.6)</p> <p>Internal fixation Group 1: 36.0 (\pm23.2) Group 2: 32.5 (\pm23.6)</p> <p>Cochrane report: Group 1: 35.01 (21.8) Group 2: 32.21 (22.03)</p> <p>Prosthesis Group 1: 5.6 (3.3) Group 2: 6.3 (5.7)</p> <p>Internal fixation Group 1: 6.4 (4.0) Group 2: 4.5 (2.3)</p> <p>Cochrane report: Group 1: 5.26 (4.08) Group 2: 6.02 (3.69)</p>	<p>Funding: Not stated</p> <p>Limitations: Most data presented for overall trial population or split by surgical treatment rather than rehab type.</p> <p>Additional outcomes reported: Ability to move/sit up/stand/walking ability/social management – all split by surgical treatment. No difference reported.</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments

1

1 Evidence tables – mobilisation strategies

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Moseley et al., 2009^{216,216}</p> <p>Country of study: Australia</p> <p>Study design: <i>RCT</i></p> <p>List who was masked to interventions: Assessor-blinded</p> <p>Duration of follow-up: 16 weeks</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Inpatient rehab units of 3 teaching hospitals in Sydney</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients with surgical fixation for hip fracture admitted to inpatient rehab units who had approval to weight bear or partial weight bear; able to tolerate the exercise programmes; able to take 4 plus steps with a forearm support walking frame and the assistance of one person; no medical contraindications that would limit ability to exercise; living at home or low care residential facility prior to the hip fracture, with the plan to return to this accommodation at discharge. Subjects with cognitive impairment were included if a carer who was able to supervise the exercise programme was available. Middle band of people with hip 	<p>Group 1 High group. Weight bearing exercise twice daily for a total of 60 minutes per day for 16 weeks. 5 weight bearing exercises were prescribed in addition to walking on a tread mill with partial body weight support using a harness (for inpatients) or a walking programme (after hospital discharge). The 5 weight bearing exercises used for both legs included stepping in different directions, standing up and sitting down, tapping the foot and stepping onto and off a block. Hand support could be used if necessary. The exercises were progressed by reducing support from the hands, increasing block height, decreasing chair height and increasing the number of repetitions. This started</p>	<p>Knee extensor strength (isometric knee extensor strength at 90° measured using a spring balance.)kg, mean (SD)</p> <p>Walking speed (measured over a 6 m distance using a stop watch) m/sec. Mean (SD)</p> <p>Pain (7 item ordinal scale) – some. Moderate or severe.</p> <p>Quality of life (EQ 5D and expressed as a utility score)</p> <p>Length of stay in hospital</p>	<p>4 week Group 1: 7.8 (3.9) Group 2: 7.7 (4.0)</p> <p>16 week Group 1: 10.3 (5.0) Group 2: 9.3 (4.4)</p> <p>4 week Group 1: 0.53 (0.25) Group 2: 0.48 (0.22)</p> <p>16 week Group 1: 0.63 (0.32) Group 2: 0.60 (0.31)</p> <p>4 week Group 1: 44 Group 2: 41</p> <p>16 week Group 1: 30 Group 2: 29</p> <p>4 weeks Group 1: 0.53 (0.27) Group 2: 0.53 (0.27)</p> <p>16 week Group 1: 0.62 (0.30) Group 2: 0.62 (0.26)</p> <p>16 week Group 1: 28 (15) Group 2: 25 (14)</p>	<p>Funding: Project grant from the National Health and Medical Research Council , Australia.</p> <p>Limitations:</p> <p>Additional outcomes reported: Fear of falling, balance, step test, body sway, stability test, falls efficiency scale. Further participant characteristics.</p> <p>Notes:</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	fracture Exclusion criteria: <ul style="list-style-type: none"> High functioning patients who are discharged directly to home and low functioning patients who are discharged to a residential aged care facility from the acute orthopaedic ward were excluded. All patients N: 160 No. of dropouts: Group 1 No. randomised: 80 No. of dropouts: 2 withdrew by 16 weeks Age (mean \pmSD): 84 (8) M/F: 15:65 Subgroup category numbers: Other factors: Group 2 No. randomised: 80 No. of dropouts: 1 withdrew at 16 weeks Age (mean \pmSD): 84 (7) M/F: 15:65	as an inpatient programme, followed by home visits and a structured home exercise programme. Group 2 Low group. Patients undertook 5 exercises in sitting or lying plus a small amount of walking using parallel bars or walking aids for a total of 30 mins each day for 4 weeks. The exercises were progressed by increasing the repetitions and resistance. This type of exercise programme is regarded as usual care. All patients received usual post-op mobilisation, and the rehab programme usually provided by other health professionals and any gait aids were progressed as per usual protocols. No physiotherapy treatments were administered during the trial.	Total exercise time with a physiotherapist or physiotherapy assistant as an inpatient, (min) mean (IQR) sit-to-stand (stand-ups per sec) mean (SD) Barthel index	Group 1: 545 (463) Group 2: 363 (318) P value(s): 0.001 4 week Group 1: 0.24 (0.15) Group 2: 0.19 (0.09) 16 week Group 1: 0.26 (0.14) Group 2: 0.22 (0.11) 4 week Group 1: 93 (85-100) Group 2: 90 (85-95) 16 week Group 1: 95 (90-100) Group 2: 95 (85-100)	

1 Evidence tables – mobilisation strategies

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Oldmeadow et al., 2006^{247,247}</p> <p>Country of study: Australia</p> <p>Study design: RCT</p> <p>Duration of follow-up: 1 week post surgery</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: The Alfred Hospital, Victoria, Australia</p> <p>Inclusion criteria: Consecutive patients admitted through the emergency department for surgical fixation of an acute neck of femur fracture (by sliding screw, gamma nail or a hemiarthroplasty) were considered for inclusion in the study.</p> <p>Exclusion criteria: Pathological fractures, if postoperative orders were for non-weight bearing on the operated hip, the patient admitted from a nursing home or the patient was non-ambulant preoperatively.</p> <p>All patients N: 60 Mean age: 79.4 years (53-95) M/F: 68% women</p> <p>Group 1 No. randomised: 29 No. of dropouts: 10 patients failed to achieve their first walk within the 48h. Age (mean \pmSD): 78.8 (2.14)</p>	<p>Group 1 Early ambulation (within 48 h/postoperative day) with a physiotherapist during standard working hours.</p> <p>Group 2 delayed (longer than 48 h/postoperative day 3 or 4)</p> <p>All patients received routine postoperative medical and nursing clinical care, as currently practiced at The Alfred.</p> <p>The physiotherapy ambulation re-education program was implemented once per day over 7 days. This program was the same for all and included</p>	<p>Function – Assistance required to transfer from supine to sit, sit to stand</p>	<p>independent Group 1: 16 Group 2: 4</p> <p>assistance Group 1: 10 Group 2: 21</p> <p>P value(s): 0.009</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <p>Additional outcomes reported: Further baseline characteristics, Troponin, subgroup analysis of true early ambulation and failed early ambulation.</p> <p>Notes:</p>
			<p>Function – Mean walking metres</p>	<p>Group 1: 58.63 (0.05 – 400) Group 2: 29.71 (0 – 150) P value(s): 0.03</p>	
			<p>Assistance required to negotiate one step on day 7 post-surgery.</p>	<p>Independent Group 1: 10 Group 2: 23</p> <p>Failed/unable Group 1: 13 Group 2: 1</p> <p>P value(s): 0.32</p>	
			<p>Discharge destination</p>	<p>Group 1: Home: 5 Fast stream rehab: 8 Slow stream rehab: 14 Nursing home: 1 Death: 1</p> <p>Group 2: Home: 1 Fast stream rehab: 14</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>M/F: 8/21</p> <p>Group 2</p> <p>No. randomised: 31</p> <p>Age (mean \pmSD): 80.0 (2.08)</p> <p>M/F: 11/20</p>	walking re-education, bed exercises and chest physiotherapy as indicated. Only the time to first walk differed between groups		<p>Slow stream rehab: 16</p> <p>Nursing home: 0</p> <p>Death: 0</p> <p>P value(s): 0.19</p>	
			Length of stay, mean (range)	<p>Group 1: 9.27 (4-33) – outlier removed n =18.</p> <p>17.90 (5-33) – failed early ambulation n = 10</p> <p>Group 2: 11.39 (5-24)</p> <p>P value(s): 0.59</p>	

1 **17.11 Evidence Table 11: Multidisciplinary rehabilitation**

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Cameron 1993 ^{42,44} Country of study: Australia Study design: RCT Duration of follow-up: 4 months	Patient group: Patients with proximal femoral hip fracture Setting: General hospital serving an outer urban area of Sydney, Australia. Inclusion criteria: Patients aged over 50 with an uncomplicated proximal femoral fracture (non-pathological, no additional fractures), surgical intervention within 7 days of injury and residence in the district. Exclusion criteria: Fractures sustained whilst in hospital or who were transferred to another hospital for surgical treatment. All patients N: 252 Lost to follow up:	Group 1 A nursing care plan starts immediately post-op that supports early mobility and self-care. A physician sees the patient the same day or the next day of the operation to identify and treat concurrent illness, review previous level of disability and assess social support needs. The Physician also liaises with the orthopaedic surgeon regarding likely complications or precautions (e.g. limitations of weight bearing). The physician leads on planning the rehab according to the patient's pre fracture condition. Patients from nursing homes are returned there as soon as feasible to undergo supervised mobilization and physiotherapy. Patients not from nursing homes are discharged once they can walk (with an aid) and go to the toilet independently. The patient received	Median length of hospital stay, days (interquartile range)	Group 1: 13 (7-25) Group 2: 15 (8-44) $p=0.034$	Funding: Australian Department of Health, Housing and Community Services. Limitations: Mean age significantly lower in accelerated rehab group ($p = 0.0042$) No assessor blinding Outcomes not reported: Additional outcomes reported: Additional baseline characteristics such as pre-injury situation, injury details. Notes:
			Mortality (obtained from the Cochrane review- Handoll 2009) 12 months	Group 1: 32 Group 2: 38	
			Mean Barthel index	2 weeks after injury Group 1: 32 Group 2: 38 1 month after injury Group 1: 32 Group 2: 38	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean \pmSD): 84 M/F: 17% male Cognitively impaired: 122</p> <p><u>Group 1 Accelerated rehab</u> No.: 127 No. of dropouts: Age (mean): Nursing home: 84.2 (n = 48) Non nursing home+moderate to severe disability: 87.2 (n = 21) Non-nursing home+limited disability: 79.2 (n = 58) M/F:</p> <p><u>Group 2 Usual care</u> No. : 125 No. of dropouts: Age (mean): Nursing home: 88.5 (n=46) Non nursing home+moderate to severe disability: 89.3 (n = 22) Non-nursing home+limited disability: 81.4 (n = 57) M/F: Other factors: Living alone</p>	<p>physiotherapy on each weekday (ideally 2 sessions per day). The orthopaedic surgeon and rehab physician review the patient 3 or 4 times weekly. After discharge the patient's rehab continues either at home (physiotherapist home visit) or at a day hospital until they reach their pre-fracture level of function or plateau at a lower level.</p> <p><u>Group 2</u> Conventional care</p>			<p>Patients stratified into 3 groups: Nursing home, non nursing home + moderate to severe disability and non-nursing home + limited disability.</p> <p>Key difference in accelerated rehab was concentrated input of an experienced physician with training in geriatric and rehab medicine.</p>

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Crotty 2002 ^{58,60}	<p>Patient group: Patients with hip fracture</p> <p>Setting: 2 Australian teaching hospitals in Adelaide (Flinders Medical centre, Repatriation General Hospital)</p> <p>Inclusion criteria: Aged 65 or over, medically stable, needed a formal rehabilitation program, had adequate physical and mental capacity to participate in rehabilitation, were expected to return home after discharge from the hospital, and had a home environment suitable for rehabilitation.</p> <p>Exclusion criteria: If patients had inadequate social support in the community, no telephone at home, or did not live in Adelaide's southern metropolitan region.</p> <p>All patients N: 66</p>	<p>Randomisation was undertaken by the hospital pharmacy department (computer generated allocation sequence in sealed opaque envelopes).</p> <p>Group 1 Patients were discharged within 48 hours of randomisation and were visited by physiotherapists, occupational therapists, speech pathologists, social workers, and therapy aides, who negotiated a set of realistic, short-term, and measureable treatment goals with both participants and their care-givers. Standard therapy services podiatry, nursing care, and assistance with light domestic tasks, were provided as required.</p> <p>Group 2 Conventional care in routine hospital interdisciplinary rehabilitation.</p>	Mortality at 12 months	Group 1: 3 Group 2: 4	<p>Funding: Supported by the South Australian Department of Human Services</p> <p>Limitations: Baseline data not given for male/female ration or mean age in each arm.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported:</p> <p>Notes:</p>
Country of study: Australia			Moved to higher level of care	Group 1: 1 Group 2: 2	
Study design: RCT			Unable to walk	Group 1: 0 Group 2: 2	
Duration of follow-up: 12 months			SF-36 physical component score at one year, mean (95% CI)	Group 1: 38 (34.0-41.9) Group 2: 33.3 (27.6-39.1)	
			SF-36 mental component score at one year, mean (95% CI)	Group 1: 53.8 (49.2-58.3) Group 2: 52.3 (47.3-57.3)	
			Length of hospital stay, mean (SD) – from Cochrane review, Handoll 2009	Group 1: 7.8 (9.3) Group 2: 14.3 (10.6)	
			Length of rehab, mean (SD) – from Cochrane review, Handoll 2009	Group 1: 28.3 (14.5) Group 2: 14.3 (10.6)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Lost to follow up: 3 Age (mean \pmSD): 82.5 M/F: 33% male</p> <p><u>Group 1 Early discharge + home rehab</u> No.: 34 Age (mean): not stated M/F: not stated</p> <p><u>Group 2 Usual care</u> No. : 32 Age (mean): not stated M/F: not stated</p>		<p>Hospital readmissions during 4 month follow up – from Cochrane review, Handoll et al., 2009^{132,133}</p>	<p>Group 1: 8 Group 2: 7</p>	

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Galvard 1995 ^{107,107}	Patient group: Patients with hip fracture	All patients were treated at the orthopaedic department and then randomization took place immediately after the operation, using a random number generator.	length of stay in hospital, days (mean, SD)	Group 1: 53.3 (47.7) Group 2: 28 (24.2)	Funding: Not stated Limitations: Higher number of subtrochanteric fractures and higher mean age of men in the geriatric MDR group. Unclear allocation concealment. Outcomes not reported: Additional outcomes reported: Baseline data – distribution of fracture types. Destination at discharge from hospital. Causes for hospital readmissions. Hip pain and walking ability one year postoperatively. Indoor walking speed. Notes: Study states that longer length of hospital stay in geriatric MDR group may relate to lack of experience in geriatric department at the time and that the orthopaedic (usual care) group had over 25 years of experience with these patients.
Country of study: Sweden	Setting: Vaernhem Hospital, Malmo, Sweden			Median Group 1: 40 Group 2: 21	
Study design: RCT	Inclusion criteria: Independently living hip fracture patients in the municipality of Malmo Exclusion criteria: People resident in nursing homes or waiting for a nursing bed, or already in hospital		Mortality at 1 year	Group 1: 45 Group 2: 40	
List who was masked to interventions: :	All patients N: 371 Age (mean + range): 79 (52-102) M/F: 26% male Group 1 Geriatric (MDR) No.: 179 Age (mean ±SD): men: 79.1 (8.6) women: 80.9 (9.2) M/F: 50/129 Group 2 Usual care No. : 192 Age (mean ±SD): men: 73.6 (10) women: 79.6 (8.2) M/F: 45/147		Group 1 Patients were transferred on the second postoperative day, and once weekly a visiting orthopaedic surgeon would decide on further treatment of the fracture Group 2 Usual care – stayed on the orthopaedic ward.	Total no. of patients readmitted to hospital	

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gilchrist et al., 1988^{113,113}</p> <p>Country of study: Glasgow, UK</p> <p>Study design: RCT</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients with femoral neck fractures</p> <p>Setting: Orthopaedic unit, Western Infirmary</p> <p>Inclusion criteria: Women aged over 65</p> <p>Exclusion criteria: Patients referred from nearby hospitals, patients who made a rapid recovery and were sent directly home.</p> <p>All patients N: 222 Age (mean ±SD):</p> <p>Group 1 Orthopaedic geriatric unit No.: 97 Age (mean): 82 Length of stay before transfer (days): 10.2</p> <p>Group 2 Usual care (orthopaedic ward) No.: 125 Age (mean): 80.6 Length of stay before transfer (days): 9.8</p>	<p>Patients were admitted to the orthopaedic unit and had standard preoperative medical assessment. After surgery were transferred to orthopaedic wards at Gartnavel General Hospital for rehab. Randomisation occurred at time of transfer.</p> <p>Group 1 Patients were under overall care of the orthopaedic surgical staff. A weekly combined ward round was performed by a geriatrician (consultant or senior registrar), an orthopaedic senior registrar, and the senior ward nurse. A physiotherapist, occupational therapist, and a social worker participated in the case conference that followed. Advice was given on medical problems that arose between ward rounds by consultation with the geriatrician. Patients were seen on average, 4 times by a geriatrician.</p> <p>Group 2 Similar nursing cover and paramedical services as group 1, but no case conference. Referral for any medical problem to the geriatric service was made by letter, and patients were seen by a different geriatrician than in group 1.</p>	<p>Mortality</p> <p>Length of stay in hospital (mean, SE)</p>	<p>Inpatient Group 1: 4 Group 2: 13</p> <p>3 month Group 1: 10 Group 2: 18</p> <p>6 month Group 1: 14 Group 2: 23</p> <p>Group 1: 44 (5.7) Group 2: 47.7 (7.7)</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Type of fracture, placement of patients admitted from home, conditions in patients at discharge</p> <p>Notes:</p>

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Huusko et al., 2002^{157,158} (Huusko et al., 2000^{157,157} gives subgroup data for patients with dementia)</p> <p>Country of study: Finland</p> <p>Study design: RCT</p> <p>List who was masked to interventions: No assessor blinding</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Patients with proximal femoral fracture</p> <p>Setting: Specialist district hospital in Jyväskylä, Finland</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Community-dwelling patients with acute hip fractures over 64 years of age. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Pathological fracture, multiple fractures, terminally ill, serious early complication, receiving calcitonin, unable to communicate <p>All patients N: 243</p> <p>Lost to follow up: Age (mean and range): 80 (66-97) M/F: 28% male</p> <p>Group 1 Geriatric rehab No.: 84 Age (mean + range): 80 (67-92) M/F: 36/84 Living alone: 62 Dementia: 32</p>	<p>Group 1 Intensive geriatric rehab within hospital: multidisciplinary geriatric team (geriatrician, specialist GP and nurses, occupational therapist, physiotherapist, social worker, neuropsychiatrist). Twice daily physiotherapy; ADL practice; daily schedule; counselling; information; discharge plan; home visits, treatment at home after discharge based in geriatric ward in same hospital as surgery.</p> <p>Group 2 Discharge to local community hospitals, treatment by GP with physiotherapists usually available. Transfer 2 to 5 days after surgery.</p>	Mortality at 12 months	Group 1: 18 Group 2: 20	<p>Funding: Study was supported by grants from Central Finland Health Care District, Kuopio University Hospital, Emil Aaltonen Foundation, Uulo Arthio Foundation and Novartis Finland Ltd</p> <p>Limitations: Imbalance of baseline characteristics. Intervention group had a greater number with Dementia 32/120 vs. 20/123); fewer were functionally independent in ADL before hip fracture (41 vs. 66)</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: IADL and ADL change</p>
			Mortality at discharge	Group 1: 5 Group 2: 5	
			Total days in hospital (during 1 year)	Group 1: 80 Group 2: 80	
			Length of hospital stay (median + range) – severe dementia (mini mental state examination score 0-11)	Group 1: 85 (13-365) N = 19 Group 2: 67 (15-365) N = 9 P=0.902	
			Length of hospital stay (median + range) – moderate dementia (mini mental state examination score 12-17)	Group 1: 47 (10-365) N = 24 Group 2: 147 (18-365) N = 12	
			Place of residence and mortality – severe dementia (mini mental state examination score 0-11)	1 year Independent living Group 1: 7 Group 2: 3 Nursing home Group 1: 5 Group 2: 0 Hospital Group 1: 2 Group 2: 3	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Group 2 Usual care No. : 90 Age (mean + range): 80 (66-97) M/F: 33/90 Living alone: 70 Dementia: 20			Dead Group 1: 5 Group 2: 3 Group 1: n = 19 Group 2: n = 9	from baseline. Notes: Patients were mobilised on the first postoperative day.
			Place of residence and mortality – moderate dementia (mini mental state examination score 12-17)	1 year Independent living Group 1: 15 Group 2: 4 Nursing home Group 1: 1 Group 2: 2 Hospital Group 1: 4 Group 2: 4 Dead Group 1: 4 Group 2: 2 Group 1: n = 24 Group 2: n = 12	

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Kennie et al., 1988 ^{176,176} Country of study: Stirling, UK Study design: RCT Duration of follow-up: 1 year	Patient group: Women with proximal femoral fracture Setting: Orthopaedic ward and geriatric rehab ward, Stirling Inclusion criteria: Women aged over 65 Exclusion criteria: Mortality prior to randomisation, pathological fractures, those likely to be discharged within 7 days of entering the trial, those remaining unfit for transfer by ambulance to a peripheral hospital. All patients N: 108 Lost to follow up: Age (mean ±SD): M/F: All female Group 1 Geriatric rehab No.: 54 Age (median + range): 79 (65-94) M/F: All female Group 2 Usual care No. : 54	Patients were randomised to geriatric rehab or usual care once the orthopaedic surgeon judged them fit to be moved to a rehab ward. Both treatment and control groups received physiotherapy, occupational therapy, and orthotic and other services. Group 1 Transferred by ambulance 5km to orthopaedic beds in a peripheral hospital. The median delay between entry into the study and transfer was one day (range 0-7). A GP provided day-to-day medical attention, and a consultant physician in geriatric medicine attended 2 ward round and 1 conference of the multidisciplinary team each week. Orthopaedic advice was available on demand.	Length of hospital stay	Mean +/- SD(from Cochrane review, Handoll 2009) Group 1: 37 (33) Group 2: 56 (54) Median Group 1: 24 (8-197) Group 2: 41 (9-365)	Funding: Forth Valley Health Board Limitations: No blinding of staff or patients. Outcomes not reported: Additional outcomes reported: Additional baseline data including residence, independence and mental state before admission, details of fracture. Notes: Similar baseline characteristics across groups, apart from age and difference in mental state, with more moderate and severe impairment in
			More dependent based on Katz score at 1 year (from Cochrane review, Handoll 2009)	Group 1: 22/43 Group 2: 28/35	
			Type of residence after discharge	NHS or private nursing home Group 1: 5 Group 2: 16 Own home Group 1: 31 Group 2: 19	
			Mortality (taken from Reid 1989)	At discharge Group 1: 5 Group 2: 4 At 1 year Group 1: 10 Group 2: 18	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Age (median + range): 84 (66-94) M/F: All female	Group 2 The control group remained in the orthopaedic admission ward. A few of these patients were moved into other short stay wards at the discretion of the consultant orthopaedic surgeon.			the control group (p=0.06)

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Naglie et al., 2002^{222,222}</p> <p>Country of study: Toronto, Canada</p> <p>Study design: RCT</p> <p>List who was masked to interventions: Assessors</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Teaching hospital in Toronto</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients aged over 70 from the community and from nursing homes <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Fractures occurring in an acute care hospital, pathologic fractures, multiple traumas, previous surgery on the fractured hip, expected survival less than 6 months, residence in a nursing home and dependence and at least one person for ambulation before the fracture, or residence outside metropolitan Toronto. 	<p>Within 48h of randomisation the research coordinator reviewed each case for compliance with the inclusion criteria and a panel then reviewed eligibility.</p> <p>Separate staff provided care in each group to prevent containment bias.</p> <p>Group 1 Protocols and standardized orders were used, early mobilisation, early participation in self care and individualised discharge planning. All nursing staff on the ward received specialised education about the care of elderly with hip fracture. A physiotherapist, occupational therapist or a clinical nurse specialist and</p>	<p>Mortality</p>	<p>At discharge Group 1: 7 (5%) Group 2: 13 (9.4%)</p> <p>3 months Group 1: 10 (7.1%) Group 2: 12 (8.7%)</p> <p>6 months Group 1: 17 (12.1%) Group 2: 21 (15.2%)</p>	<p>Funding: Supported by a grant from Ontario Ministry of Health and the Research Institute of the Queen Elizabeth Hospital, Toronto.</p> <p>Limitations: Anticipated that the intervention would increase length of hospital stay.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Baseline characteristics such as: Functional and cognitive scores,</p>
			<p>Decline in ambulation- data missing for 3 patients in group 1 and 8 patients in group 2 at 3 months.</p>	<p>3 months Group 1: 73 (57%) Group 2: 72 (61%)</p> <p>6 months Group 1: 59 (47.6%) Group 2: 56 (47.9%)</p>	
			<p>Decline in transfers- data missing for 3 patients in group 1 and 8 patients in group 2 at 3 months.</p>	<p>3 months Group 1: 57 (44.5%) Group 2: 48 (40.7%)</p> <p>6 months Group 1: 45 (36.3%) Group 2: 44 (37.6%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Patients were excluded postoperatively if the surgery failed for technical reasons, if they required care in an intensive care unit or if there was no bed available on the interdisciplinary care ward.</p> <p>All patients N: Lost to follow up: Age (mean \pmSD): M/F:</p> <p>Group 1 interdisciplinary care No.: 141 No. of dropouts: 0 Age (mean): 83.8 (6.9) M/F: 32/109 Other factors: Living alone: 23.4% Mean time to surgery: 1.3 days Subcapital fractures: 46.8%</p> <p>Group 2 Usual care No. : 138 Withdrawal: 1 Age (mean): 84.6 (7.3) M/F: 24/114 Other factors: Living alone: 23.2% Mean time to surgery: 1.4 days Subcapital fractures: 39.1%</p>	<p>social worker assigned to the ward routinely assessed all study patients within 72 hours. Daily medical care from a senior internal medicine resident supervised by an internist-geriatrician.</p> <p>Group 2</p> <p>Patients had access to allied health professionals if a consultation was requested, but had limited access to an occupational therapist or a clinical nurse specialist.</p>	<p>Change in residence</p>	<p>3 months Group 1: 31 (23.7%) Group 2: 32 (25.4%)</p> <p>6 months Group 1: 22 (17.7%) Group 2: 23 (19.7%)</p>	<p>medical indicators, surgical procedure. Care by allied health professional. Place of residence at discharge,</p> <p>Notes: Intervention group received more physiotherapy hours than the control $p < 0.001$</p> <p>A subgroup analysis in the paper shows a trend towards benefit in patients with mild to moderate cognitive impairment.</p> <p>NB Intensive intervention during hospital stay.</p>
			<p>Length of stay in hospital, days (SD)</p>	<p>Group 1: 29.2 (22.6) Group 2: 20.9 (18.8)</p>	

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Marcantonio et al., 2001^{203,203}</p> <p>Country of study: USA</p> <p>Study design: RCT</p> <p>Duration of follow-up:</p>	<p>Patient group: Patients with proximal hip fracture</p> <p>Inclusion criteria: All patients aged 65 and older admitted to an academic tertiary medical center for primary surgical repair of hip fracture.</p> <p>Exclusion criteria: Presence of metastatic cancer or other comorbid illnesses likely to reduce life expectancy to less than 6 months, or inability to obtain informed consent within 24h of surgery or 48h of admission</p> <p>All patients N: 126 Lost to follow up: Age (mean \pmSD): M/F:</p> <p>Group 1 Geriatric consultation No.: 62 No. of dropouts: Age (mean): 78\pm8 M/F: 79% female Other factors: Pre fracture dementia: (Blessed score \geq4) :21 Prefracture ADL impairment (Katz ADL score <5: 11</p>	<p>Group 1: Intervention Geriatric consultation preoperatively or within 24h postoperatively. A geriatrician performed daily visits for the duration of hospitalisation and made targeted recommendations based on a structured protocol. The protocol included 10 modules each containing 2 to 5 specific recommendations. Detailed fully in the paper, includes adequate CNS oxygen delivery, fluid/electrolyte balance, treatment of severe pain, elimination of unnecessary medications, regulation of bowel/bladder function, adequate nutritional intake, early mobilization and rehab, management of postop complications, appropriate environmental stimuli, treatment of agitated delirium.</p> <p>Group 2: usual care Management by orthopaedic team, including internal medicine or geriatric consults</p>	<p>Delirium: Total cumulative incidence during acute hospitalisation</p>	<p>Group 1: 20 Group 2: 32</p>	<p>Funding: Part funded by a pilot project grant from the Older Americans Independence Centre and a grant from the Charles Farnsworth Trust.</p> <p>Limitations: Not MDR rehab, focus on impact of geriatric consultation.</p> <p>Notes: Recommendations made, and adherence to them varied. Full data given in paper.</p>
			<p>Severe delirium: cumulative incidence during acute hospitalisation</p>	<p>Group 1: 7 Group 2: 18</p>	
			<p>Hospital days of delirium per episode (mean \pmSD)</p>	<p>Group 1: 2.9\pm2 Group 2: 3.1\pm2.3</p>	
			<p>Hospital length of stay (median +IQR)</p>	<p>Group 1: 5\pm2 Group 2: 5\pm2</p>	
			<p>Discharged to institutional setting (nursing home, rehab hospital)</p>	<p>Group 1: 92% Group 2: 88%</p>	
			<p>Delirium at hospital discharge</p>	<p>Group 1: 8 Group 2: 12</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 Usual care No. : 64 No. of dropouts: Age (mean): 80±8 M/F: 78% female Other factors: Pre fracture dementia: (Blessed score ≥4) :29 Prefracture ADL impairment (Katz ADL score <5: 18</p>	<p>on a reactive rather than proactive basis.</p>			

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Shyu et al., 2008^{305,306}</p> <p>Country of study: Taiwan</p> <p>Study design: RCT</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Patients with hip fracture</p> <p>Setting: Teaching hospital in Taiwan</p> <p>Inclusion criteria: Aged 60 or over, admitted to hospital for an accidental single-side hip fracture, receiving hip arthroplasty or internal fixation, able to perform full range of motion against gravity and against some or full resistance and had a prefracture Chinese Barthel Index score >70, and living in northern Taiwan</p> <p>Exclusion criteria: Severely cognitively impaired, making them unable to follow orders or terminally ill.</p> <p>All patients N: 162 Age (mean \pmSD): 78 M/F: 31.5% male</p> <p>Group 1 Intervention No.: 80 Age (mean): 77.36 (8.19) M/F: 25/55</p>	<p>Patients recruited from the emergency room by research assistants.</p> <p>Group 1 Interdisciplinary programme of geriatric consultation, continuous rehabilitation and discharge planning. Geriatrician and geriatric nurses provided geriatric assessment/consultation; physiotherapist, geriatric nurses and rehab physician were responsible for rehab programme, Early mobilisation, home visit and follow-up services provided. 4x 30min physical therapy sessions per patient, 2 assessments from a physical therapist and one visit from rehab physician. 4 home visits during first month and 4 during second and third month from a geriatric nurse.</p> <p>Group 2 On trauma or orthopaedic ward. Occasional consultation with other disciplines depending on patient's</p>	<p>Length of hospital stay, mean days (SD)</p> <p>Recovery of walking ability</p> <p>Mortality</p> <p>Non-recovery/decline in walking ability – long-term at 12 months (additional info from Cochrane review Handoll 2009)</p>	<p>Group 1: 10.1 (3.7) Group 2: 9.72 (4.96)</p> <p>at 6 months Group 1: 62 Group 2: 44</p> <p>at 12 months Group 1: 61 Group 2: 49</p> <p>at 6 months Group 1: 6 Group 2: 8</p> <p>at 12 months Group 1: 13 Group 2: 15</p> <p>Group 1: 59 Group 2: 56</p>	<p>Funding: Supported by grants from the National Health Research Institute, Taiwan.</p> <p>Limitations:</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Marital status, educational background. Occurrence of falls, self-care ability, depressive symptoms</p> <p>Notes: Includes early mobilisation and intensive rehab.</p> <p>Intervention resembles a geriatric hip fracture rehab</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Independent walking ability: 68</p> <p><u>Group 2 Usual care</u></p> <p>No. : 82</p> <p>Age (mean): 78.94 (7.28)</p> <p>M/F: 26/56</p> <p>Independent walking ability: 69</p>	<p>condition. Exercises taught by nurses in first 2 to 3 days. Physical therapy sessions varied according to insurance policy.</p>			<p>programme and early supportive discharge.</p>

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Stenvall et al., 2007^{320,320}</p> <p>Country of study: Sweden</p> <p>Study design: RCT</p> <p>List who was masked to interventions:</p> <p>Duration of follow-up: 12 month</p>	<p>Patient group: Patients with femoral neck fracture</p> <p>Setting: Umea University Hospital, Sweden</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients aged 70 years or older <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients with severe rheumatoid arthritis, severe hip osteoarthritis or a pathological fracture, or severe renal failure. Patients who were bed bound prior to the fracture. <p>All patients N: 199</p> <p>Lost to follow up:</p> <p>Age (mean ±SD): M/F: 26% male</p> <p>Group 1 Intervention No.: 102</p> <p>No. of dropouts:</p> <p>Age (mean): 82.3 (6.6)</p> <p>M/F: 28/74</p> <p>Other factors:</p>	<p>Group 1 Geriatric unit specializing in geriatric orthopaedic patients. Active prevention, detection and treatment of post op complications implemented daily. Early mobilisation, with daily training was provided by physiotherapists, occupational therapists and care staff during hospital stay. Assessment at 4 months by geriatric team.</p> <p>Group 2 Specialist orthopaedic unit following conventional postoperative routines. A geriatric unit was used for those needing longer rehab n = 40, but this was not the</p>	<p>Living independently</p> <p>Independent walking ability</p> <p>Independent walking without walking aid indoors</p> <p>Independent in P-ADL (poorer personal activities of daily living)</p> <p>Length of stay in hospital</p>	<p>At 4 months Group 1: 54 Group 2: 46</p> <p>At 12 months Group 1: 47 Group 2: 36</p> <p>At 4 months Group 1: 59 Group 2: 52</p> <p>At 12 months Group 1: 55 Group 2: 45</p> <p>At 4 months Group 1: 31 Group 2: 19</p> <p>At 12 months Group 1: 35 Group 2: 22</p> <p>At 4 months Group 1: 35 Group 2: 23</p> <p>At 12 months Group 1: 33 Group 2: 17</p> <p>At 12 months Group 1: 30 (18.1) Group 2: 40 (40.6)</p>	<p>Funding: Supported by the Vardal Foundation, the Joint Committee of the Northern Health Region of Sweden, the JC Kempe Memorial Foundation, the Dementia Fund, and the Foundation of the Medical Faculty, the Borgerskapet of Ulmea Research Foundation, the Erik and Anne-Marie Detlof's Foundation, University of Ulmea and the County Council of Vasterbotten and the Swedish Research Council.</p> <p>Limitations: Not blinded, but independent assessors.</p> <p>Intensity and quality of outpatient rehab is unknown.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Living alone: Group 2 Usual care No. : 97 No. of dropouts: Age (mean): 82 (5.9) M/F: 23/74 Other factors: Living alone	same ward as the intervention.		p=0.028	Outcomes not reported: Baseline data such as health and medical problems, functional performance prior to fracture. Additional outcomes reported: Notes: Paper contains a detailed description of the intervention and control group.
			Mortality	At discharge Group 1: 6 Group 2: 7 At 12 months Group 1: 16 Group 2: 18	
			Hospital readmissions	At 12 months Group 1: 38 Group 2: 30	
			More dependent based on Katz index at 1 year	Group 1: 35 Group 2: 49	
			Non recovery in ADL at 1 year	Group 1: 51 Group 2: 59	

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Swanson et al., 1998 ^{325,325} Country of study: Australia Study design: RCT Duration of follow-up: 12months	Patient group: Patients with femoral fractures. Setting: Royal Brisbane Hospital, teaching hospital. Inclusion criteria: Patients aged 55 or over; non-pathological fractures; residing at home or in a hostel; independently mobile (with or without a walking aid); able to give informed consent; accessible for follow up (i.e., residing in the Brisbane area); and public patients. Exclusion criteria: Patients with dementia, with inadequate English to give informed consent or residing in a nursing home. All patients N: 71	Patients were identified by the trial coordinator in the Accident and Emergency Department Group 1 Multidisciplinary team: full time physiotherapist, occupational therapist, clinical nurse consultant, half time social worker, geriatrician, orthopaedic surgeon. Early mobilisation (1 st day after surgery if possible), twice daily intense sessions by physiotherapist, daily assessment, treatment or counselling by the occupational therapist and social worker. Review by geriatrician	Length of stay (discharge criteria used e.g. when medically stable and able to transfer and walk independently with or without aids) Mortality Modified Barthel Index at discharge (95% CI)	Mean Group 1: 21 (17.2-24.4) Group 2: 32.5 (24.2-41.1) p<0.01 Median Group 1: 17 Group 2: 24 p<0.01 In hospital Group 1: 2 Group 2: 2 12 months Group 1: 5 Group 2: 6 6 months Group 1: 92.8 (90.0-95.6) Group 2: 85.6 (81.3-89.8) 12 months Group 1: 95.3 (SD 9.8) Group 2: 89 (SD 15.8)	Funding: Medicare Incentives Hospital Access Program. Limitations: Underpowered – initial power analysis determined that 120 patients (60 in each arm) would have the power to detect a reduction in mean length of stay of 7 days at 0.05 level of significance. However the difference in length of stay was larger than anticipated. No assessor blinding Outcomes not reported:

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Lost to follow up: 0 Age (mean \pmSD): M/F: 22% male</p> <p><u>Group 1 Early intervention</u> No.: 38 Age (mean): 78.5 (75.3-81.7) M/F: 11/27 Living at home: 35 (92.1%)</p> <p><u>Group 2 Usual care</u> No. : 33 No. of dropouts: Age (mean): 77.8 (74.0-81.6) M/F: 5/28 Living at home: 29 (87.9%)</p>	<p>on next working day after surgery, 2 additional ward rounds attended by all staff, weekly case conference attended by all staff, coordination of care by trial coordinator, home assessment visit before discharge.</p> <p><u>Group 2</u> Standard orthopaedic management including daily visits from a physiotherapist, and social worker or occupational therapist visits as requested by hospital staff. Weekly discharge planning, home visits as requested by social worker.</p>	<p>Complications (additional from Cochrane review)</p>	<p>Chest infection, cardiac problem bedsore Group 1: 6 Group 2: 13</p> <p>Stroke emboli Group 1: 4 Group 2: 1</p>	<p>Additional outcomes reported:</p> <p>Notes: Surgery was carried out within 48 hours of admission for 90% of intervention and 80% of standard care. 12 month data from Day 2001.</p>

1

2

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Vidan et al., 2005 ^{344,344}	<p>Patient group: Patients with hip fracture</p> <p>Setting: Hospital General Universitario “Gregorio Marañon”.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Consecutive patients aged 65 and older between February 1 and December 15, 1997 for acute hip fracture surgery. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Inability to walk before the fracture and dependency in all basic activities of daily living; pathological hip fracture; known terminal illnesses, defined as those associated with a life expectancy of less than 12 months. <p>All patients N: 319 Lost to follow up: Age (mean ±SD): M/F: 18.5% male</p> <p>Group 1 Usual care No.: 164 No. of dropouts: not stated</p>	<p>All patients had an orthopaedic surgeon and a nurse assigned when they were admitted to hospital. The intervention and control group shared the same orthopaedic wards and used the same hospital-wide support services, including physical therapy and social work. The orthopaedic surgeon made the decision of discharge moment in both groups.</p> <p>Group 1 The surgeon and orthopaedic nurses managed patients, with counselling from different specialists as needed.</p> <p>Group 2 A geriatrician visited the patients daily and was responsible for medical care. After initial</p>	Median total length of hospital stay (25th to 75th percentile)	Group 1: 18 (13 – 24) Group 2: 16 (13 – 19) p = 0.06	<p>Funding: Not stated</p> <p>Limitations: Usual care group have a higher percentage of coexisting conditions</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Additional baseline data: coexisting conditions, type of fracture, type of surgery. Also medical complications: heart failure, DVT, myocardial infarction, arrhythmia.</p> <p>Notes: ADL = activities of daily living. (Bathing, dressing, using the toilet, getting from bed to chair, and continence) FAC = Functional</p>
Country of study: Spain			In hospital mortality	Group 1: 9 (5.5%) Group 2: 1 (0.6%) p = 0.03	
Study design: RCT			Mortality – end of scheduled follow up (from Cochrane review)	Group 1: 39 Group 2: 28	
Duration of follow-up: 12 months			Major medical complications	<p>Confusion Group 1: 67 (44.1%) Group 2: 53 (34.2%) p = 0.07</p> <p>Pressure sores Group 1: 27 (16.9%) Group 2: 8 (5.2%) p = 0.001</p> <p>Pneumonia Group 1: 6 (3.7%) Group 2: 6 (3.9%) p = 0.95</p> <p>Heart failure Group 1: 5 Group 2: 12</p>	
			Time from surgery to rehabilitation, days, mean (SD)	Group 1: 10.2 (6) Group 2: 8.3 (3.9) p = 0.007	
			Recovery of ADL or FAC	Group 1: 3 (2%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Age (mean): 82.6 (± 7.4) M/F: 35/129 Living at home before admission: 134 (82%) Type of surgery: Internal fixation: 101 (61.6%) Prosthetic replacement: 53 (32.3%) Others: 10 (6.1%) Mean time to surgery, hours (SD): 78.5 ± 53.2 Group 2 Intervention No. : 155 No. of dropouts: not stated Age (mean): 81.1 (± 7.8) M/F: 24/131 Living at home before admission: 135 (87%) Internal fixation: 91 (58.7%) Prosthetic replacement: 58 (37.4%) Others: 6 (3.9%) Mean time to surgery, hours (SD): 75.8 ± 43.2	assessment and within 72 hours after admission, there was an interdisciplinary meeting, including the orthopaedic and geriatric teams, to discuss the patient's medical, functional, and social problems and to elaborate a comprehensive therapeutic plan. The meeting was repeated weekly.	at time of hospital discharge Recovery of ADL or FAC at time of 3 months Incomplete recovery of ADL and mobility at 1 year (from Cochrane review)	Group 2: 5 (3%) Group 1: 59/134 (44%) Group 2: 82/144 (57%) p = 0.03 Group 1: 75 Group 2: 67	Ambulation Classification. This consists of 6 different functional levels.

1 Evidence tables – multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ziden et al., 2008 and Ziden et al., 2010^{360,361}</p> <p>Country of study: Sweden</p> <p>Study design: RCT</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Community-dwelling patients with hip fracture</p> <p>Setting: Patients admitted to the emergency unit at the Sahlgrenska University Hospital</p> <p>Inclusion criteria: Acute hip fracture surgery, medically approved by the responsible geriatric doctor as being in need of geriatric care and rehab, aged 65 or over and able to speak and understand Swedish.</p> <p>Exclusion criteria: Severe mental illness with expected survival of less than one year, severe drug or alcohol abuse, mental illness or documented severe cognitive impairment.</p> <p>All patients N: 102 Total of 212 randomised: Excluded: 99</p>	<p>A geriatric nurse who performed the randomisation using sealed envelopes.</p> <p>Patients with hip fracture were referred from the emergency unit to a geriatric ward with home rehab (group 1) or with conventional care.</p> <p>Both groups performed early mobilization, preferably within 48 h. When needed an occupational therapist or physiotherapist made a home visit with the patient to assess if they could manage and what aids they needed.</p> <p>Group 1 Conventional care and rehab as in group 2, plus supported discharge. An initial meeting with the patient aimed to establish individual goals. Close contact with social home services and relatives to plan discharge and cooperation during 1st few</p>	<p>Balance confidence – Swedish version of the Falls Efficacy Scale – 1 month (SD) . 0-10 scale where 0 indicates very confident, no fear of falling, 10 is not confident, very afraid of falling. Swedish. Includes 13 items covering activities of daily living.</p> <p>Activities of daily living and leisure activities – degree of independence assessed by Functional Independent Measure (FIM) motor scale (mean, SD). 13 items with a 7 point grading scale (0 = totally dependent and 7 = totally independent) max score 91 points</p>	<p>1 month Group 1: 117.4 (12.0) Group 2: 85.5 (30.5) p<0.0001</p> <p>1 month Self-care Group 1: 38.4 (2.9) Group 2: 33.5 (7.2)</p> <p>Mobility Group 1: 18.3 (1.5) Group 2: 16.3 (3.3)</p> <p>Locomotion Group 1: 10.4 (2.5) Group 2: 7.6 (3.6)</p> <p>6 months – median with range Self-care Group 1: 40 (33-42) Group 2: 37 (6-42)</p> <p>Locomotion Group 1: 31 (15-34) Group 2: 30 (5-35)</p>	<p>Funding: Supported by the Vardal Institute, the Hjalmar Svensson's Foundation and the Geriatric Section of the Swedish Association of Registered Physiotherapists.</p> <p>Limitations: No length of hospital stay or total length of rehab in control group.</p> <p>Outcomes not reported:</p> <p>Additional outcomes reported: Other baseline characteristics such as walking ability, number of medical diagnosis, functional independence and instrumental activity.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Declined to participate: 11 Lost to follow up: Age (mean \pmSD): 81.9 (6.8) M/F:</p> <p><u>Group 1 Home rehab</u> No.: 48 No. of dropouts: Age (mean): 81.2 (5.9) M/F: 19/29 Other factors: Living alone: 26</p> <p><u>Group 2 Usual care</u> No. : 54 No. of dropouts: Age (mean): 82.5 (7.6) M/F: 12/42 Other factors: Living alone 39</p>	<p>weeks at home. Home rehab consisted of a 3 week intervention period.</p> <p><u>Group 2</u> Participation in standard rehab including daily training in basic activities: transfer techniques, technical aids, indoor and stair walking. Also physiotherapy and occupational therapy group sessions. Prior to discharge the home service officer and patient's next of kin was contacted to make plans for the future. All rehab measures were adapted to the patient's individual medical and functional status and personal goals.</p>	<p></p> <p>Basic physical mobility – timed “up and go” test. Assess total time for standing up from a chair, walking 3 m, turning 180° returning and sitting down (performed twice, one trial and one timed)</p> <p>Functional lower extremity muscle strength. Ability to rise from a chair was measured by sit-to-stand. Best of 3 trials was recorded. Made in participant's home with ordinary chairs, preferable with armrests, and ordinary walking aids were used, if needed (secs)</p> <p>Length of hospital stay (mean +/- SD)</p>	<p>1 year – median with range Self-care Group 1: 40 (23-42) Group 2: 38 (12-42)</p> <p>Locomotion Group 1: 32 (11-35) Group 2: 29 (9-35)</p> <p>1 month Group 1: 24.9 (15.4) Group 2: 30.8 (16.0)</p> <p>1 month Group 1: 1.8 (0.8) Group 2: 3.3 (3.6)</p> <p>Group 1: 18.4 (8.4) Group 2: 20.0 (6.8)</p>	<p>Subsequent falls, frequency of activities, balance confidence.</p> <p>Notes:</p>

1 **17.12 Evidence Table 12: Patient views**

Study	Archibald 2003 ⁸ . Country: UK. Setting: community hospital in Bradford
Aim	To explore experiences of individuals who had suffered a hip fracture. Not to produce generalisable findings but to generate "rich description" of the experience of incurring and recovering from hip fracture to inform nursing practice.
Population	5 patients with hip fracture Age >65; 4 women and 1 man; all were cognitively intact
Method of gaining views	In depth audio-recorded interviews with open ended questions, ranging between 25 and 50 minutes duration were conducted during stay in the hospital.
Data analysis	<p>“Colaizzi's analysis framework. 6 step methodological interpretation</p> <ul style="list-style-type: none"> • Interviews transcribed verbatim and read to get a feel for responses • Significant statements and phrases extracted • Meanings formulated from significant statements • Organised into clusters of themes • Themes used to provide full description of experience • Researcher returns description to participants for confirmation of validity
Findings	<p>4 main themes: injury experience, pain experience, recovery experience, disability experience.</p> <p>Injury – relates to falling and breaking their hip</p> <p>Pain - Most participants described the pain they had. One mentioned being in a lot of pain in orthopaedic unit despite pain killers. Another mentioned they thought the pain went with rest after a while, but not completely. Only 1 person was still having pain at time of interview. One said "...I have not suffered, not what I call real pain, at all",</p> <p>Recovery - operation: varied comments - some did not remember anything or much, one had a “horrendous” recollection of operating theatre: “The operation was pretty horrendous. I had the injection in the spinal cord, [an] epidural... There was no pain, but the noises [laughs] – it was like being in an engineering shop or something. The noise was terrible. I thought ‘What are they doing me?’ Anyway, it came to an en (it took quite a long time)..and before I knew it I was back on the ward.”</p> <p>Recovery - beginning struggle: 3 patients discussed this, 1: not being able to do anything, 2: struggling to get to toilet & into the chair, 3: hated using bed pan.</p> <p>Recovery - regaining independence: Motivation found to be key factor in recovery, all comments in study positive comments about regaining independence during their rehabilitation.</p> <p>Disability: comments about reduced functional status, dependence on others, being house bound.</p>
Comments	Not stated how patients were selected for the study. No baseline data provided about patients. The role of the researcher is not described.

1 Evidence tables – patient views

Study	Borkan 1991 & 1992 ^{28,29} . Country: USA. Setting: 4 hospitals (no more detail)
Aim	Two research questions addressed: <ul style="list-style-type: none"> • What are the meanings present in the narratives of elderly hip fracture patients? • What is the importance of narrative elements as prognostic indicators or 'risk factors' for predicting rehabilitation outcomes?
Population	80 patients with hip fracture (from a pool of 174) "functionally hardy elderly, intact mental status, independent or lightly-supervised residence outside long-term care facilities, full pre-fracture ambulation; >65 years; 65 women and 15 men; diagnosed within 48 hours of fracture; treated surgically within 1 week. Excluded open pathological or multiple fractures.
Method of gaining views	Interviewed during first week after hip fracture , generally 1 or 2 days after surgery, in participant's hospital room. In depth initial interviews included demographics, open ended questions and standardised scales. Combination of open-ended and multiple choice questions. Interview content validated through pretesting with 10 subjects, and reviewed by a panel of experts. Inconsistencies and ambiguities revised or deleted from study. Follow up interviews at 3 and 6 months post-fracture generally conducted in participants' current residence, except where movement to distant states or particular patient preferences precluded face to face contact. These attempted to match some of the patient's perceptions to what actually happened. In addition, observations carried out on main orthopaedic floors over the course of 2 years in order to familiarise research team with the treatment and rehabilitation as well as to confirm information drawn from interviews and uncover unexpected associations.
Data analysis	Quantitative analysis & qualitative narrative. Names coded and interview transcripts sent to independent expert panel to identify emergent or recurrent themes. 13 dimensions identified and grouped into 3 composite. Subjects' narrative accounts rated on a 7 point bipolar scale.
Findings	Gives themes around the patient perception of hip fracture, how it happened, how they perceive their injury, what the future holds, their subsequent level of ability and their future. Categories derived from narratives are rated on a bipolar scale and presented in 3 groups. The remaining percentage not given for each category relates to patients either not giving a view or indicating an equal rating for both polar elements. <ol style="list-style-type: none"> 1. Explanation of fracture: described as disease (1%) or fracture (49%); fall as secondary (4%) or primary (82%); etiology, internal degeneration – primary (6%) or secondary (11%); broke and fell (10%) or fell and broke (64%); course of rehabilitation described as chronic (19%) or acute (49%); functional severity – total impairment (14%) or complete recovery (70%); range of severity – whole body (11%) or affected leg or hip (15%). 2. Perception of disability: vulnerable (41%) or not vulnerable (34%); dependency increased (21%) or not increased (30%); sense of alienation from the world – alienated (20%) or integrated (29%); objectification of body part – alienation (4%) or wholeness (7%). 3. Futurity: hopefulness (54%) or hopelessness (19%). <p>Expectations of recovery during initial hospitalisation: 43 (53.7%) expected full recovery; 14 (17.5%) partial recovery; the rest did not know or did not give an answer. Narrative responses varied "from stubborn optimism to despair".</p> <p>Expectations of living situation: 61% predicted going home, 15% predicted going into a nursing home (none came from nursing home), 9% going to children's house, 15% did not know or did not respond. Actual figures: 34 (43%) discharged to long-term care institutions, 13 (38%) of these remained in institution at 1 year, 18 (53%) returned home, 3 (9%) died.</p>
Comments	The role of the researcher is not well described.

1 **Evidence tables – patient views**

Study	Bowman 1997 ³³ . Country: Canada. Setting: hospital
Aim	To describe sleep satisfaction, pain perceptions & psychological concerns of patients undergoing planned & emergency hip operations. Two additional questions on perceptions of how they would manage.
Population	43 out of 50 consecutively admitted patients: 17 with hip fracture & 26 undergoing elective hip replacement. Gender for overall study 29 women and 14 men. Characteristics of hip fracture patients: mean age 80 (+7.5); 8/17 had delirium; 11/17 patients claimed to be active or very active prior to fracture
Method of gaining views	Pain assessment was conducted using a visual analogue scale. Sleep satisfaction was conducted using a 'Likert' scale. Not much detail on methods for qualitative part of study. Interviewed on day of admission . Two structured questions but no details on how or by whom they were delivered. 1. What are your biggest concerns at this time as a result of this injury and your upcoming surgery? 2. Do you have any concerns about your ability to recover fully and quickly?
Data analysis	Numerical analysis of responses to two questions.
Findings	6/17 feared being unable to walk again; additional 3/17 concerned about recovery and managing on their own; 5/17 put their trust in God.
Comments	Little detail about methods used for the qualitative part of this review. Little baseline data provided about patients. The role of the researcher is not described.

2

1 **Evidence tables – patient views**

Study	Furstenberg 1986 ¹⁰⁵ . Country: USA. Setting: Large urban teaching hospital
Aim	2 parts to study: 1. community residents without hip fracture, 2. hospitalised patients with hip fracture. "The purpose of hospital study was to construct a natural history of the hip fracture, from the events surrounding the fracture through the hospitalisation period.
Population	11 patients hospitalised for hip fracture. Patient characteristics: age 59 to 85 years; 4 men & 7 women; cognitively intact, fracture that had not resulted from malignancy or its treatment.
Method of gaining views	Interviewed at one or more points during their hospital stay. "Ethnographic interviews" recorded and transcribed in full. Interviews took place in physical therapy hospital rooms or in rehabilitation centre for 3 who were transferred. During interview informants requested to talk about the fracture, their reactions to it, their pre-fracture functioning, their experiences during hospitalisation and the process for planning for discharge.
Data analysis	"Analysis consisted of identifying salient and recurrent issues and themes and grouping the portions of the interviews dealing with each theme. The variations on each theme were described, and correlates of these variations were identified".
Findings	<p>Split into two main sets: (1) immediate expectations about recovery explicitly or implicitly expressed by patients; (2) contextual factors to the evolving expectations about recovery.</p> <ol style="list-style-type: none"> 1. Immediate or early expectations of recovery - most expressions of despair and discouragement. Only 1 patient feared "it was over". First reactions "varied from shock to a focus on immediate problems, and for some, immediate concern about the consequences of their way of life. As the situation progressed, patients' concerns focused more exclusively on limitations on their functioning and the implications these would have". Most expressed worry about the degree to which they would recover, and when. Several talked repeatedly about the slowness of the process of recovery of physical function. Some worried about being burdens on their caretakers, some worried about further falls. Those who went temporarily to a home of an adult child worried about being able to return to independent living. Summary - hip fracture was going to result in extended period of slow recovery of function, with attendant dependency, postponement or relinquishment of cherished plans and changed living situation with the threat of permanent loss of independent living. Also suffered uncertainty about timing & completeness of return to full recovery. 2. Contextual factors - as time progressed. Only positive points, not negative ones, came out in this section. Patients observing their own progress sometime after surgery commented that although progress slow they could see improvement. Participants also took encouragement from others progress. The study notes that while patients could focus on positive and negative points, the informants only focused on encouraging examples. 3. Contextual factors - health professionals influence on patients' perceptions. Healthcare professionals' cues, encouragement and feedback guided the informants' perceptions about their own progress. Quotes of the healthcare professionals were scattered throughout participants' responses. Some patients "referred to the elusiveness of the doctors and their own unanswered questions." 4. Contextual factors - other health issues. Also reports a few comments by patients on other health issues.
Comments	Little baseline data provided about patients. The role of the researcher is not described.

2

1 Evidence tables – patient views

Study	Olsson 2007 ²⁴⁹ . Country: Sweden. Setting: geriatric/orthopaedic ward
Aim	To describe patients' own perceptions of their situation and views of their responsibility in the rehabilitation process.
Population	13 hip fracture patients from a geriatric/orthopaedic ward, non-institutional residence pre-fracture, median age 81 (range 71 to 93) years, 2 men & 11 women. Excluded patients with severe illness, cognitive impairment, dementia or pathological fracture.
Method of gaining views	30-45 minute interviews conducted in informant's room or in a secluded area of the ward as soon after the operation as the informants felt strong enough. Semi-structured questions were used "such that the main questions, related to the informant's perception of the transitional properties, were included in all interviews." Deliberate efforts were made to encourage informants to reveal and comment freely on their personal experiences of and reflections on their situation, without imposing the interviewer's own values on what was being said. The interviewees all talked freely and appeared to be grateful for the attention and for having someone to listen to their reflections. All interviews were recorded and transcribed.
Data analysis	Transcripts read several times. 5 transitional properties & 542 meaning units identified & pooled. A "saturation" was observed when 9 interviews had been conducted "...meaning units describing qualitatively similar conceptions were grouped together and the nature of this similarity was articulated." Categories were labelled and exemplified with representative quotations from interviews. To test the reliability of the categories the second author evaluated the categories in relation to the interviews.
Findings	<p>Participant's responses were categorised into different conceptions:</p> <ul style="list-style-type: none"> • autonomous – appeared confident and accustomed to managing for themselves and being in control of their lives. Willing to listen to staff, but made their own decisions. Even if they appeared strong they felt just as vulnerable as the other groups. However, they were aware of the importance of information, personal support and their own responsibility. One informant commented that more information given preoperatively could have made a great difference: <ul style="list-style-type: none"> ○ "Of course, if someone had come and sat down for a little while and talked. If they had said something like, this is what it will be like and so on and after a while you will be able to walk and maybe manage on your own again. That would have been reassuring, it really would. Because I really must say, at moments like that, you get a feeling of being small and insignificant." • modest – gave the impression of being vulnerable and dependent on others and they expressed themselves cautiously. Instead of demanding community aftercare like the Autonomous, they were willing to go along with what was offered. These informants appreciated information offered to them but for some reason they did not request more, even though they seemed to want to. They worried more about their future ability to walk and maintain their former lifestyles than the other two groups. They feared being discharged, saw only problems and appeared unaware of the progress they had made. They were reluctant to talk about their hopes for the future and did not see their responsibility as clearly as the autonomous. • heedless – "appeared to view their situation with some detachment, almost as if it did not concern them. The Heedless did not doubt they would recover and they were confident that people around them would care for them." "The Heedless were characterized predominantly by a reluctance to reflect on their own situation, by a refusal to accept responsibility and by their need for information.They did not appear to have reached a stage where planning for the future was relevant." <p>Also identifies some common traits:</p> <ul style="list-style-type: none"> • [lack of] awareness - most lacked adequate awareness about their condition, what to do and how to act and needed more information. Only

	<p>1 patient knew someone who had undergone rehabilitation for hip fracture.</p> <ul style="list-style-type: none">• shocking event - although several suspected they had a fracture all were distressed by the diagnosis. Period before surgery was mostly blurred and filled with fear and pain. They worried about how they would function postoperatively;• zest for life - all expressed a strong desire to recuperate. While confined to bed they were worried remembering the pain and inability to move their leg. The suffering experienced in anticipation and preparation for the operation led them to believe they might not be able to walk.
Comments	Not stated how patients were 'strategically selected' for the study. Little baseline data provided about patients. The role of the researcher is not described.

1 Evidence tables – patient views

Study	Pownall 2004 ²⁷⁴ . Country: UK. Setting: trauma and orthopaedic ward
Aim	Critical appraisal of an individual patient narrative of their experience with hip fracture. Undertaken in an effort to understand further the nature of personal experience. Narrative was acquired as part of a routine nursing evaluation and helped to illuminate nursing care issues through they eyes of the patient.
Population	A 60 year old woman with an intracapsular fracture in Nottingham hospital. She stated she was fully independent prior to fracture.
Method of gaining views	Interviewed prior to discharge from acute trauma and orthopaedic ward, exact time point unclear. A list of structured questions were devised but not rigidly adhered to: <ul style="list-style-type: none"> • What did you feel about requiring hospitalisation? • What were the good aspects of your hospitalisation? • What were the bad aspects of your hospitalisation? • What do you feel could be improved?
Data analysis	Narrative assessment of patient's views
Findings	<p>A few areas for potential improvement for the hospital/department were identified:</p> <ul style="list-style-type: none"> • communication skills • time management for staff so time spent with patient is used effectively • pain management <p>Ann's comments that were included in the study:</p> <ul style="list-style-type: none"> • I could not understand why I had to wait so long in A & E, they had done the X-ray, it was broken the X-ray person told me that. So why did I have to wait? • The pain was unbearable; I didn't care what happened or what was said I just wanted to get rid of the pain. • The staff were so kind, they could not do enough for me. • Initially, I could not understand why they (the staff) wanted to keep checking my bottom, I was comfortable why keep moving me? • It was terrible to be kept nil by mouth the first day, I didn't feel like eating but I really wanted a drink. • It was such a disappointment to be told my operation was cancelled; I just wanted to be fixed. • When I came back from theatre I really needed a drink, but I could not reach my glass. I didn't want to bother the staff they looked so busy. • It was a relief to come back from theatre and be able to press a button and get pain relief, but it was taken away the next day when the physiotherapist came. So I had to keep asking for pain killers. • The staff are so busy no one has time to sit and explain things to you. • I could hear the nurse explaining the operation to my son, but what about me I needed to know. • It was frightening to wake up from the operation and see that I was having a blood transfusion, no-one said that I might need a blood transfusion. It makes you feel something has gone terribly wrong. • I couldn't believe it when they wanted to mobilise me the day after the operation, even my son was shocked to see me out of bed.

Comments	Almost no methodology described so results could be unreliable. It is unclear how this patient was chosen. The role of the researcher is not described.
-----------------	---------------------------------------------------------------------------------------------------------------------------------------------------------

1

1 **Evidence tables – patient views**

Study	Slauenwhite 1998 ³¹⁴ . Country: Canada. Setting: interviewed at home after discharge from hospital
Aim	Purpose of this study was to investigate the impact of enhanced early discharge on families experiencing repaired hip fracture in an older adult.
Population	Convenience sample of 23 caregivers for 23 patients who had experienced hip fracture . Patient characteristics: age 75.9 (range 56-97) years; 19 women 4 men. Care giver characteristics: 16 women, 7 men.
Method of gaining views	Interviewed 4 to 6 weeks after discharge with questions adapted from Canadian Patient Centered Hospital Care Survey, a “validated tool”. The developers of the tool determined the adaptations would not affect the validity or reliability of the questionnaire. Questionnaires mailed to caregiver 1 week before interview (questions reported in study). "Data were considered qualitative if the interviewee elaborated on an answer, expressing strong beliefs about a topic. At the end of the interview, specific questions were asked to elicit how the patient and family experienced the illness episode." One interviewer was used. Interviews were taped then transcribed.
Data analysis	2 investigators separately analysed the transcripts and developed themes that emerged from the data. Themes were then compared, contrasted and collapsed until only main themes remained.
Findings	<ul style="list-style-type: none"> • Length of stay not a major issue for 15/23 families, care-recipient thought too long while patient-carer thought too short for 3 families, 4 families said people heal better in own homes. • 20/23 families stated pain management not a problem in hospital or at home • several families thought transition from house to home a problem as took several hours to days for all info to be relayed to home care system. This went hand in hand for those with comorbidities. • “Instrumental functioning” not a concern when patients were allowed to manipulate their own resources in their own home. • Older people and men more capable of role flexibility while younger people and women talked more about role strain. • Many caregivers had stories of dissatisfaction which was suggested to be related to health care system and mismatched care. Mismatched care not well defined.
Comments	No description of ‘early supported discharge’. No baseline description of patients and no indication of how patients were selected.

2

1 **Evidence tables – patient views**

Study	Williams 1994 ³⁵⁴ . Country: USA. Setting: At home after discharge from 4 hospitals
Aim	Aim to gain information on: (1) the recovery pattern in functional status & mood in first 14 weeks after hospital discharge; (2) factors most associated with the extent of assistance required in specific mobility activities & patient assessment of their problems; (3) problems patients identified as most important; (4) advice those patients would give to others.
Population	120 consecutive patients meeting inclusion criteria with hip fracture. Older patients who were relatively healthy & home dwelling before fracture. Mean 79.9 (+9.7) range 60 to 100). Included intracapsular (68), extracapsular (52), internal fixation (76), femoral head replacement (44). Sample included only white women as a result of low number in region of study.
Method of gaining views	Interviewed before hospital discharge and followed up at 2, 8 and 14 weeks. At 14 weeks participants asked what advice they would give to other persons who fractured their hip. Also assessed functional status, perceived return to normal mobility, mood states, and other factors including urinary problems according to scales.
Data analysis	Coding of responses to advice to give to other hip fracture patients done by the two "co-principal investigators" with recategorisation occurring until 100% agreement was reached.
Findings	Advice to patients with newly fractured hips from women with a personal experience of hip fracture Number of comments by category: <ul style="list-style-type: none"> • 94 importance of mental attitude - maintain hope & look to the future • 76 follow experts advice • 34 mobility – keep mobile, rest before getting up to walk, use walker to help get up • 15 maintain healthy lifestyle • 7 use caution & be careful not to fall • 3 limit stay in institution and get help to be at home if possible; • 6 gave no specific advice as they commented that everyone is different.
Comments	

2

1 **Evidence tables – patient views**

Study	Wykes 2009 ³⁵⁵ . Country: Australia. Setting: rehabilitation hospital
Aim	Pilot study to explore "the impact of fractured neck of femur on the lives of previously independent women and identifies their concerns when participating in inpatient rehabilitation".
Population	5 patients undergoing inpatient rehabilitation for hip fracture at 2 rehabilitation hospitals, aged 60-85 years, living alone and independently before fracture, cognitively intact and able to converse fluently in English
Method of gaining views	Interviewed in a private room during stay in rehabilitation hospital . Interviews were shared by two researchers previously unknown to the patients. Interviews taped and transcribed verbatim. Each woman invited to tell her story with as few interruptions as possible. Main questions were "Can you tell me how you came to be here in this rehabilitation ward?" and "What do you think about while you are in hospital?" To ensure in depth coverage patients were frequently asked "Could you please tell me more about that?"
Data analysis	Thematic analysis using stages set out by Burnard 1991. Two researchers independently made notes of themes apparent in the data as a whole. Transcript lines were coded. Similar codes combined into higher order categories. Same two researchers carried out the analysis. Researchers engaged in "reflexive self-awareness". This included a conscious awareness of previous experiences of and with patients who had fractured a neck of the femur.
Findings	Two major findings: <ol style="list-style-type: none"> 1. Impact of fracture for previously independent women was an issue for all. Primarily, others had to assume responsibility for things they had done previously. 2. Concerns following fracture listed in 4 sections: <ol style="list-style-type: none"> a. behaviour of others (22 instances identified) - these included: <ul style="list-style-type: none"> what others do - things staff said or did (1 women upset when she overheard staff talking about the possibility of limb shortening if they stayed in a wheelchair too long, 1 women upset about being put on a ward with people "completely of the planet" as a result of dementia; 1 patient commented that staff don't understand because they encouraged her to walk when she felt she could not because of Parkinson's Disease interfering with her mobility), friends & family doing things without consulting her what others do not do -; family not told by staff when patient moved hospital; not enough information about complications what others expect – 1 women concerned by staff expecting her daughter to look after her before rehabilitation started; family and friends expectations upset participants. b. what was happening to them - possible accommodation changes after discharge; possible loss of independence; money issues c. impact of their injury on others - inconveniencing and upsetting others d. other health issues – 2 women had pre-existing conditions that overshadowed their concerns about hip fracture and had adverse effects on their rehabilitation outcomes. – 1 severely disabled with Parkinson's Disease; 1 had recent cardiac surgery and a long-standing vertebral disc prolapse.
Comments	Study notes: only 5 patients included so it only reveals some of the concerns of older women with hip fractures; not enough data to explore the differences between hospitals; analysis only at 1 point in time.

2

1 Evidence tables – patient views

Study	Young 2009 ³⁵⁸ , Country: USA. Setting: rehabilitation programme
Aim	To explore the perceptions of older adults regarding their functional recovery 1 year after hip fracture.
Population	62 hip fracture patients ('convenience sample' from a longitudinal study of rehabilitation and functional recovery after hip fracture involving 280 patients). Age 65 or older (average: 78, range 65-91), 47 women, 15 men, cognitively intact, community dwelling, admitted to one of the five predetermined rehabilitation sites with a primary diagnosis of acute hip fracture, receiving a surgical procedure, non pathological fracture, no evidence of metastatic cancer.
Method of gaining views	<p>Participants invited and completed an exit interview immediately after the 12-month post hip fracture follow up data collection. The exit interview was a thematic survey with open-ended questions that explored areas of functional recovery and participants' willingness to engage in rehabilitation activities. Questions:</p> <ol style="list-style-type: none"> 1. Have you been satisfied with your functional recovery since your hip fracture surgery? – YES, NO <ol style="list-style-type: none"> a. If "YES" what do you think has helped the most with regards to your recovery process? b. If "NO" what do you think has hindered your recovery process most? c. If "NO" what things would you have liked to see differently regarding you recovery process? 2. What do you think needs to be done to help improve the functional recovery process for future hip fracture patients? 3. What one piece of advice would you give a hip fracture patient to help them with their recovery? <p>Responses were transcribed verbatim by a physical therapist and a physician assistant, both of whom were familiar with hip fracture care and received three sessions of interview training at the Center on Aging and Health at John Hopkins University in Baltimore.</p>
Data analysis	<p>Data analysis conducted using basic content analysis. "Although the interview guide used in this study contained specific themes and directed participants to address things that facilitated their recovery process, response analysis was conducted using participants' own words to capture their particular responses and ideas about thematic areas." A list including a definition of each code was developed and continually revised as new codes were added. "</p> <p>"Confirmability" Data were initially coded by first reviewer, a geriatric nurse practitioner, and researcher familiar with the hip fracture trajectory. The coded data were then given to a second researcher, an epidemiologist and gerontologist who had studied patients post-hip fracture across the entire recovery period. The second interviewer independently coded the transcripts, compared her coding to the coding of the first reviewer, and then discussed the findings with the first reviewer. As the discrepancies were identified, the reviewers went back to the data to clarify their interpretations. This process repeated until consensus was reached. Codes were then grouped based on similarities and differences.</p> <p>Data credibility was addressed by presenting the findings to an interdisciplinary group of clinicians and researchers (one physician, four epidemiologists, three exercise trainers, one physical therapist, and one occupational therapist) familiar with the hip fracture trajectory to establish if the findings made sense and were consistent with the current understanding of the recovery process post hip-fracture. The findings were presented in a small group and one on one in the clinical setting. Participants were asked to verbally confirm or refute the findings.</p>
Findings	<p>53 participants were satisfied with their functional recovery, 9 were not satisfied. 25 codes were identified and collapsed into four main themes.</p> <ol style="list-style-type: none"> 1. Facilitators of recovery (identified by 53 participants satisfied with their recovery): <ul style="list-style-type: none"> • professionals (40) – comments covered being buoyed by seeing physician frequently, having good doctors or surgeons, getting "correct" or "professional" care. "They evaluated professionals as a team and did not single out one provider over another in terms

- of help and support received". Communication and a positive attitude by professionals also important;
- **social support** (13) – from family and friends essential to their recovery. Specifically mentioned verbal encouragement helped them maintain a positive attitude
 - **determination** (12) – own determination to exercise and be involved
 - **lifestyle factors** (4) & **environment** (1) – eating healthy food, taking appropriate medications and vitamins, and engaging in physical activity. "an environment that encouraged healthy behaviors (i.e. facilitated physical activity) was important to promote exercise"
 - **individualised care** – verbal encouragement (4);
 - **spirituality** (4) – spirituality and belief in a supreme being helped them maintain their optimism throughout the process
 - identifying **goals** (3) – returning home, regaining independence and being able to walk like they could prefracture
2. **Factors that hinder recovery** (identified by 9 participants dissatisfied with their recovery):
- medical complications/comorbidities (4)
 - **unpleasant sensations** (3) – pain reported as a limiting factor
 - age (1)
3. **System recommendations to facilitate recovery:**
- **more care** (26) – more direct physical & occupational therapy and more education about the recovery process and ways to optimise physical function
 - **better care** (9) – follow up and care in the home setting after discharge from rehabilitation
 - **spirituality** (3), **social support** (2) – some participants said they would have like exposure to spiritual support options throughout the course of their rehabilitation programme. Some participants also felt that additional social and spiritual supports were needed from family and friends.
 - additional information (8)
 - elimination of unpleasant sensations (4)
 - policy (1)
4. **Peer advice to facilitate recovery:**
- **participate** (48) & **listen to providers** (19) – listen to healthcare instructions and participate as much as possible in rehabilitation activities. Comments included "listen to the advice from medical staff such as doctors, therapists, and nurses" and "Do a lot of physical and occupational therapy even if it's painful"
 - **positive attitude** (20) & **determination** (13) – participants strongly recommended that older adults who sustain hip fractures maintain a positive attitude, avoid worry and remain determined throughout the recovery experience
 - **be careful** (8) – avoid subsequent trauma, prevent anything that would impede recovery, prevent falls
 - **push through pain** (6), relieve pain – "do your physical therapy even though it may hurt" & "use all offered medications that could alleviate pain and relax muscles"
 - don't worry (4).

Numbers in brackets relate to the number of times noted

Comments	Paper reports the study used to as the basis to recruit participants for this paper had stringent eligibility criteria because it was designed to evaluate rehabilitation. Therefore, the findings of this study may only be applicable to a similar patient group. Although the findings were found to be credible with rehabilitation clinicians and researchers they were not verified with patients who had sustained hip fracture. Themes were determined by the interview guide.
-----------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

1

2

1 Evidence tables – patient views

Study	Ziden 2008 & Ziden 2010 ^{362,363} . Country: Sweden. Setting: hospital
Aim	Aim to explore & describe the experienced consequences of an acute hip fracture among home dwelling elderly people shortly after discharge. "The ambition was to let the subjects concretize their experiences, for instance by describing in as great details as possible their ordinary daily activities before and after the fracture."
Population	Patients selected from a larger sample of 102 participants (ZIDEN2008 RCT) with acute hip fracture, >65 years old, living in own home, no cognitive impairment, and able to understand Swedish. Participants asked if they were willing to participate a few days after surgery. At 1 month: 18 participants, 16 women and 2 men At 1 year: 15 participants, 13 women and 2 men
Method of gaining views	Semi-structured interviews using the phenomenographic method. Interviews held in patients own homes 1 month & 1 year after hospital discharge . Interviews conducted in a conversational manner that allowed interviewees to speak freely and to express their own experiences of the consequences of the hip fracture. As an introduction, the subject was asked to narrate what had happened when he or she broke their hip. Follow up questions and prompts were used, such as "Tell me more about it", "What does this mean to you?" and "Can you clarify?" Interviews were taped and were transcribed verbatim.
Data analysis	Phenomenographic method described by Dahlgren & Fallsberg: interviews read through repeatedly to obtain a total concurrent overview; statements extracted that dealt with consequences of hip fracture to achieve a concentrated and representative version of entire dialogues; quotes from previous step were compared in order to uncover sources of variation or agreement; similar quotes were grouped together, an attempt was made to "describe the essence of similarity within each group" (stage called articulating); these groups were then labelled/categorised and compared to ensure categories did not overlap. The grouping and describing stages were revised several times before the analysis was judged to be satisfactory. Sequence of steps in the analysis made separately by authors before joint discussions leading finally to consensus.
Findings	At 1 month 8 categories in 3 focused areas were identified: In relation to your body and yourself: <ul style="list-style-type: none"> • You are limited to move and have lost confidence in your body (18 people) • You become humble and grateful (7 people) • You respect yourself and your own needs (2 people) In relation to others: <ul style="list-style-type: none"> • You become more dependent on others (12 people) • You gain more human contact and are treated in a friendly way by others (2 people) In relation to the life situation: <ul style="list-style-type: none"> • You are secluded and trapped at home (4 people) • You are old, closer to death and have lost your zest for life (4 people) • You take one day at a time and are uncertain about the future (7 people) At 1 year 6 categories in 2 focused areas were identified: Experienced consequences of a hip fracture 1 year after discharge

	<ul style="list-style-type: none">• Isolated life with more restricted activity and fewer social contacts<ul style="list-style-type: none">a. more insecure and afraid (11 patients)b. more limited ability to move (12 patients)• Disappointed and sad that identity and life have changed (8 patients)• Satisfied with the situation or feeling even better than before fracture (5 patients) <p>Conceptions of what influences hip fracture recovery</p> <ul style="list-style-type: none">• Own mind and actions influence recovery (10 patients)• Treatment and actions from others influences recovery (4 patients)• You cannot influence recovery (6 patients)
Comments	

1

2 **18 Appendix F: Evidence tables - Economic**

3 **studies**

4 **Abbreviations**

5

CI	Confidence interval
IQR	Interquartile range
ITT	Intention to treat analysis
Int	Intervention
LOS	Length Of Stay
LR+	Positive likelihood ratio
LR-	Negative likelihood ratio
M/F	Male/female
N	Total number of patients randomised
NA	Not Applicable
NPV	Negative predictive value
NR	Not reported
PPV	Positive predictive value
QALY	Quality-Adjusted Life Years
QoL	Quality of life
RCT	Randomised controlled trial
RR	Relative risk
SA	Sensitivity analysis
SD	Standard Deviation
SE	Standard Error
Sig	Statistically significant at 5%

1 18.1 Evidence Table 13: General versus regional anaesthesia

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Chakladar 2010 UK Economic analysis: Cost analysis Study design Survey Duration of follow-up: NA Perspective: UK NHS Discount rates: Costs: NA Effects: NA	Patient group: Hypothetical patients undergoing uncomplicated anaesthetic for hip fracture repair.	Group 1: Spinal anaesthesia Group 2: General anaesthesia	Mean (SD) anaesthetic time (minutes)	Group 1: 31 (15) Group 2: 27 (16) p value: p<0.0001	Funding/conflict of interest: The authors declared there were no competing interests or external funding. Limitations: Partial economic evaluation. Survey on hypothetical patients, not on real cohorts. Spinal anaesthesia after failure of regional was not included in the analysis. Anaesthetists from one hospital only were interviewed. Overall quality and applicability Potentially serious limitations and partial applicability. Data sources: Anaesthetic time from Brighton Hip Fracture Database. Notes: * 20 anaesthetic consultants
			Mean (SD) cost of anaesthesia equipment per patient (2010 GBP)	Group 1: £66.73 (30.05) Group 2: £108.15 (38.53) p value: NR	
			Mean (SD) cost of airway equipment per patient (2010 GBP)	Group 1: £1.81 (0) Group 2: £25.68 (2.28) p value: NR	
			Mean (SD) cost of personnel per patient (2010 GBP)	Group 1: £105.90 (0) Group 2: £106.76 (0) p value: NR	
			Mean (SD) cost of drugs per patient (2010 GBP)	Group 1: £19.03 (11.00) Group 2: £25.17 (11.04) p value: NR	
			Mean (SD) cost of gases/inhalational agents per patient (2010 GBP)	Group 1: £0.43 (0.13) Group 2: £6.26 (3.94) p value: NR	
			Mean total cost per patient (SD) 2010 GBP, sum of previous categories of costs.	Group 1: £193.81 (37.49) Group 2: £270.58 (44.68) p value: p<0.0001	
			Cost-effectiveness	NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Sensitivity analysis	NR	

Abbreviations: NR=not reported, NA=not applicable

1
2

1 18.2 Evidence Table 14: Displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Johansson2006 ¹⁶³ Sweden	<p>Patient group: Patients 75 years or older who were admitted to the Linköping University Hospital with displaced femoral neck fractures with walking ability prior to the trauma, no contraindications to major surgery, no rheumatic joint disease.</p> <p>All patients N: 143* Mean age (range): 84 (75–101) M/F: 34/109 Drop outs: 16 patients</p> <p>Group 1 N: 78* Age (mean): M/F: Drop outs: 9 patients</p> <p>Group 2 N: 68* Age (mean): M/F: Drop outs: 7 patients</p>	<p>Group 1: Internal fixation performed with two parallel and percutaneously inserted screws after closed reduction.</p> <p>Group 2: Total hip replacement performed with a cemented prosthesis using a poster-lateral approach.</p> <p>All patients had postoperative physiotherapy.</p>	Number of hips that required reoperation (%)	Group 1: 34 (44%) Group 2: 11 (16%) p value: NR	<p>Funding/conflict of interest: NR</p> <p>Limitations: Costs derived only from one hospital.</p> <p>Overall quality and applicability Potentially serious limitations and partial applicability.</p> <p>Additional outcomes: There was no difference in the change of average cost of community services/place of residency between the two groups. Pain was significantly higher in Group 1.</p> <p>Notes: *143 patients were followed up but two patients in Group 1 and one patient in Group 2 were randomised twice in the same group because they had bilateral fractures. ** Data for 7 patients in Group 1 and 4 in Group 2 were missing at 1 year, and data for 9 patients in Group 1 and 7 in Group 2 were missing at 2 years. *** Once a patient scored as poor due to a failure they remained in this group despite reoperation.</p>
Economic analysis: cost-consequences analysis			Number of patients with a Harris hip score excellent or good/fair or poor at 1 year**	Group 1: 6/48*** Group 2: 24/24 p value: <0.0001	
Study design RCT			Number of patients with a Harris hip score excellent or good/fair or poor at 2 years**	Group 1: 6/42*** Group 2: 20/21 p value: <0.001	
Duration of follow-up: 2 years			Mean cost per patient 2000 Euros, cost of surgical procedures, hospital stay, radiographic examination, home rehabilitation, emergency and outpatient visits, hospital overheads, complications and reoperations.	Group 1: 13,100 (£11,575) Group 2: 12,800 (£11,310) p value: NR	
Perspective: Hospital			Cost-effectiveness	NR	
Discount rates: Costs: NR Effects: NA			Sensitivity analysis	NR	

1 **Abbreviations:** NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis
2

1 Evidence table: displaced intracapsular fractures

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Keating 2005 ¹⁷³ UK Economic analysis: Cost-utility analysis Study design RCT Duration of follow-up: 2 years Perspective: NHS Discount rates: Costs: 0%* Effects: 0%	Patient group: previously fit patients of 60 years or older with displaced subcapital hip fractures. All patients N: 298 Age (range): 60 - 93 M/F: 65/233 Drop outs: Group 1 N: 118 Age (mean): 74.9 M/F: 29/89 Drop outs: 19 (18/19 died) Group 2 N: 111 Age (mean): 75.4 M/F: 19/92 Drop outs: 20 (19/20 died) Group 3 N: 69 Age (mean): 75.2 M/F: 17/52 Drop outs: 7 (7/7 died)	Group 1: Internal fixation Group 2: Bipolar hemiarthroplasty Group 3: Total hip replacement	Number of deaths within 4 months of operation (%)	Group 1: 3 (3%) Group 2: 6 (5%) Group 3: 2 (4%) p value: Not sig	Funding/conflict of interest: grant from the National Health Service Health Technology Assessment programme. Limitations: Small number of patients. Overall quality and applicability Minor limitations and partial applicability. Additional outcomes: Place of discharge, adverse events Notes: * Costs were not discounted because most of the costs were incurred within 1 year of injury. ** Group1 vs 3 was sig after adjusting for age and gender
			Number of patients with further surgery within 4 months of first operation (%)	Group 1: 26 (22%) Group 2: 6 (5%) Group 3: 5 (7%) p value: NR	
			Number of deaths within 12 months of operation (%)	Group 1: 10 (8%) Group 2: 11 (10%) Group 3: 4 (6%) p value: Not sig	
			Number of patients with further surgery within 12 months of first operation (%)	Group 1: 37 (31%) Group 2: 6 (5%) Group 3: 6 (9%) p value: NR	
			Number of deaths within 24 months of operation (%)	Group 1: 18 (15%) Group 2: 18 (16%) Group 3: 6 (9%) p value: Not sig	
			Number of patients with further surgery within 24 months of first operation (%)	Group 1: 46 (39%) Group 2: 6 (5%) Group 3: 6 (9%) p value: <0.001 (Group 1 vs 2 and 3) Not sig (Group 2 vs 3)	
			EQ-5D utility scores at 4 months – mean (SD)	Group 1: 0.56 (0.29) Group 2: 0.61 (0.29) Group 3: 0.68 (0.24) p value: Not sig**	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			EQ-5D utility scores at 12 months – mean (SD)	Group 1: 0.58 (0.34) Group 2: 0.64 (0.33) Group 3: 0.70 (0.29) p value: 0.04 (Group 1 vs 3) Other groups not sig	
			EQ-5D utility scores at 24 months – mean (SD)	Group 1: 0.55 (0.38) Group 2: 0.53 (0.35) Group 3: 0.69 (0.32) p value: 0.008 (Group 2 vs 3) Other groups not sig	
			Mean cost per patient over 2 years (95% CI) 2001 GBP, cost of hospital admission (inpatient and day case), theatre costs, prosthesis and profile of hardware, excluding non-hip-related admissions.	Group 1: 12,623 (10,768 – 14,478) Group 2: 9,897 (8,062 – 11,732) Group 3: 9,399 (8,265 – 10,532) p value: Sig (Group 1 vs 3) Other groups not sig	
			Cost-effectiveness Cost per utility gained	Total Hip Replacement is dominant.	
			Sensitivity analysis Two-way SA	Results did not change when cost of prostheses and cost of readmission were varied over a range from -50% to +100% around the baseline values.	

Abbreviations: NR=not reported, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, SA=sensitivity analysis

1 18.3 Evidence Table 15: Cemented arthroplasties

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Santini 2005 ²⁹⁸ Italy Economic analysis: Cost-consequences analysis Study design RCT* Duration of follow-up: One year Perspective: Provider Discount rates: Costs: NA Effects: NA	Patient group: at least 65 years old, with life expectancy of at least 3 months, low-energy trauma. All patients N: 106 Age (mean): NR M/F: 24/82 Drop outs: 0 Group 1 N: 53 Age (mean): 82 M/F: 13/40 Drop outs: 0 Group 2 N: 53 Age (mean): 80 M/F: 11/42 Drop outs: 0	Group 1: Cemented bipolar hemiarthroplasty Group 2: Uncemented bipolar hemiarthroplasty	VELCA functional score	Group 1: 9.13 Group 2: 8.95 p value: Not sig	Funding: The authors declared no conflict of interest. Limitations: Surgical time not included in cost calculation although it was significantly different (group 2 had shorter operating time). The only difference considered was the cost of prostheses. Overall quality and applicability: Potentially serious limitations and partial applicability. Additional outcomes: Social environment at 1 year was similar in the two groups. Notes: * included in our clinical review **only cost of prostheses was different between the two groups.
			Peri-operative mortality – number of patients (%)	Group 1: 3 (24.5%) Group 2: 2 (26.4%) p value: Not sig	
			Mortality at 1year – number of patients (%)	Group 1: 13 (24.5%) Group 2: 14 (26.4%) p value: Not sig	
			Number of patients with complications	Group 1: 21 Group 2: 21 p value: NR	
			Mean cost per patient** 2001 Euros, cost of medical and nursing staff, drugs, diagnostic procedures, prostheses, blood transfusion and hospital stay.	Group 1: 3,093 (£2,400) Group 2: 4,008 (£3,110) p value: NR	
			Cost-effectiveness	NR	
			Sensitivity analysis	NR	

1 **Abbreviations:** NR=not reported, NA=not applicable, M/F=male/female, N=total number of patients randomised, VELCA=Verona Elderly Care, Sig=statistically significant at 5%

2

3

1 18.4 Evidence Table 16: Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Cameron 1994 ⁴⁵ Country: Australia Economic analysis: CEA Study design RCT Duration of follow-up 4 months Perspective: Health care provider Discount rates NA	Patient group: Patients with proximal femoral hip fracture All patients N: 252 Age (mean): 84 M/F: 14/70 Drop outs: 0 Group 1: Accelerated Rehab N: 127 Age (mean): Nursing home: 84.2 (n=48) Non-nursing home + moderate to severe disability: 87.2 (n=21) Non-nursing home +limited disability: 79.2 (n=58) M/F: NR Group 2 N: 125 Age (mean): Nursing home: 88.5 (n=46) Non-nursing home +moderate to severe disability: 89.3 (n=22) Non-nursing home +limited	Group 1: Accelerated rehab (involving: early mobilization after surgery, comprehensive rehabilitation program, early discharge from hospital, community-based rehabilitation). Group 2: Conventional care	Median length of stay, days (interquartile range)	Group 1: 13 (7-25) Group 2: 15 (8-44) p value = 0.034	Funding Australian Department of Health, Housing, and Community Services. Conflict of interest: NR Limitations A longer follow up could have better reflected differences in costs and outcomes. Health-related QoL were not calculated. Overall quality and applicability The study has potentially serious limitations and partial applicability. Notes: (1)Calculated using the Power Purchasing Parity (PPP) of 1990
			Mean Barthel index: No. of patients recovered at 4 months from surgery	Group 1: 63 (49.6%) Group 2: 52 (41.6%) 95% CI (-3% to 21%) p value = Not significant	
			Mean Barthel index: No. of patients worse at 4 months from surgery	Group 1: 31 (24%) Group 2: 39 (31%) p value= NR	
			Mean Barthel index: No. of patients death at 4 months from surgery	Group 1: 19 Group 2: 20 p value = NR	
			Mean cost per patient Year: 1990 Currency: Australian dollars Cost components: inpatient hospital (surgical, post	Group 1: A\$ 10,620 (£ 4678.9 – 1990 PPP)(1) Group 2: A\$ 12,790 (£ 5635.01 – 1990 PPP)(1)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	disability: 81.4 (n=57) M/F: NR		surgical), readmissions, community support services, institutional care.	p value = 0.186	
			Cost-effectiveness Incremental cost per additional recovered patient	The accelerated rehab program is the dominant strategy (more effective, less costly)	
			Sensitivity analysis Threshold sensitivity analysis	Accelerated rehab is more costly than usual care when: (1) The difference in LOS between the 2 strategies is less than 1.5 – 2 days (2) Cost of treatment is more than 40% per bed day compared to conventional care. These results were not sensitive to the % of patients recovering nor to the definition of recovery.	

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Farnworth 1994 ⁹¹ Australia Economic analysis: CCA Study design Case study with historical control Duration of follow-up 6 months Perspective: Health care provider Discount rates NA	Patient group: Patients with hip fracture All patients N: 138 Age (mean): NR M/F: 23/115 Drop outs: 0 Group 1 N: 67 Age (mean), (SD): 78.4 (8.8) M/F 10/57 Group 2 N: 71 Age (mean): 79.8 (10.7) M/F 13/58	Group 1: Fractured Hip Management Program (FHMP) comprising: orthopaedic surgeon, geriatric physician, nurses, occupational therapist, physiotherapist. Rehabilitation took place in the patient's normal environment. Group 2: Usual care	In-hospital mortality at 1 year Length of Stay – days (nursing home patients) Length of Stay – days (non-nursing home patient) Readmission within 1 year Mean cost per patient Year: 1990 Currency: \$Aus Cost components: -Hospital costs - FHMP costs (staff time, use of medical goods, office space and travel time for home visits). Cost-effectiveness Sensitivity analysis	Group 1: 16 (24%) Group 2: 19 (27%) p value = NR Group 1: 7.3 Group 2: 10.2 p value = NR Group 1: 21.5 Group 2: 28.2 p value = NR Group 1: 4 (6%) Group 2: 6 (8%) p value = NR Group 1: \$Aus11 060 (£4872) (1) Group 2: \$Aus 9280 (£4088) (1) p value = NR NR NR	Funding/conflict of interest: NR Limitations: The year at which cost data refer is not clear. The duration of follow up is not clear. No sensitivity analysis was conducted. Health related QoL outcomes were not calculated. No incremental analysis was conducted. Overall quality and applicability The study has potentially serious limitations and partial applicability Additional outcomes: Changes in living arrangements at discharge from hospital and 1 year after hip fracture. Notes:

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
					(1) The costs were expressed in GBP using the Power Purchasing Parity for 1990.

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

1
2

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Galvard 1995 ¹⁰⁷ Sweden Economic analysis: CCA Study design RCT Duration of follow-up: 1 year Perspective: NHS and PPS Discount rates NA	<p>Patient group: Patients with hip fracture</p> <p>All patients N: 371 Age (mean): NR M/F: 95/276 Drop outs: 0</p> <p>Group 1 N: 192 Age (mean) - female: 79.6 years (SD 8.2) Age (mean) - male: 73.6 years (SD 10.0) M/F 45/147 Drop outs: 0</p> <p>Group 2 N: 179 Age (mean) - male: 79.1 (SD 8.6) Age (mean) - female: 80.9 (SD 9.2)</p> <p>M/F 50/129 Drop outs: 0</p>	<p>Group 1: Rehabilitation in geriatric department (Patients transferred on second postoperative day. Orthopaedic surgeon would visit them once weekly)</p> <p>Group 2: Usual care (rehabilitation in orthopaedic department)</p>	<p>Readmissions to hospital</p> <p>Mortality at 1 year</p> <p>Mean length of stay in hospital, days (SD)</p> <p>Mean cost per patient Year: 1989</p> <p>Currency: Swedish Krona (SEK)</p> <p>Cost components: Technical aids, home adjustment costs, stay at convalescent home, new hospital admission, daily costs at orthopaedic and geriatric department.</p> <p>Cost-effectiveness</p>	<p>Group 1: 36 Group 2: 57 p value = NR</p> <p>Group 1: 45 Group 2: 40 p value = NR</p> <p>Group 1: 53.3 (47.7) Group 2: 28 (24.2) p value = NR</p> <p>Group 1: SEK 94,026.05 (£6590.82) (1) Group 2: SEK 84,536.81 (£5925.67) (1) p value = NR</p> <p>NR</p>	<p>Funding/ Conflict of interest: NS</p> <p>Limitations: No sensitivity analysis was performed.</p> <p>Health related QoL outcomes are not calculated.</p> <p>No incremental analysis was conducted.</p> <p>The source used to estimate the unit cost of resources was unclear.</p> <p>Overall quality and applicability The study has potentially serious limitations and partial applicability</p> <p>Additional outcomes: Destination at discharge: 72.4% of patients from group 1 and 72.0% of patients in group 2 returned to their previous living arrangements (NS).</p> <p>Notes: (1) Values in GBP obtained using the Power Purchasing Parity (PPP) for</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Sensitivity analysis	NR	1989.

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

1
2

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hollingworth 1993 ¹⁴⁸ UK Economic analysis: Cost analysis Study design Case series Duration of follow-up: Until discharge (1) Perspective: NHS Discount rates NR	Patient group: Hip fracture patients. All patients N: 1080 Age (mean): NR M/F: 198/882 Drop outs: NA Group 1 N= 779 (2) Age (mean): 78.7 (SD 11.2) M/F: 143/636 Drop outs: NA Group 2 N: 301 Age (mean): 79.8 (SD 10.9) M/F: 55/246 Drop outs: NA	Group 1: Community rehabilitation - Hospital at home (HAH) scheme. The scheme provides care from trained nurses, nursing auxiliaries, physiotherapists, and occupational therapists in the patient's home for up to 24 hours a day under the medical supervision of the general practitioner. The scheme lasts for up to two weeks – after then, other community services take over. Group 2: Usual care	LOS (mean inpatients days) Readmission rates at 1 year (for patients with access to HAH scheme and for usual care patients) Mean cost per patient Year: 1992 Currency: UK sterling Cost components: Ward, Hospital at home, hotel, overheads, medical, theatre, other treatment. Cost-effectiveness Sensitivity analysis One-way deterministic sensitivity analysis.	Group 1: 32.5 Group 2: 41.7 p value: <0.001 Group 1: 53 (6.8%) Group 2: 8 (2.7%) p value =0.008 Group 1: £4884 Group 2: £5606 p value = 0.048 NR The costs in the HAH scheme would still be lower than in the usual care case even if inpatients costs were 50% lower than predicted and the HAH costs were 50% higher.	Funding/conflict of interest: NR Limitations: Unclear follow up time Parameters' uncertainty has not been subjected to appropriate probabilistic sensitivity analysis. No incremental analysis was conducted. Health-related QoL were not determined. Information on costs obtained from the hospital finance department, not from official statistics. Overall quality and applicability The study has potentially serious limitations and partial applicability Notes: (1) The duration of follow up was unclear from the paper (2) These were patients with access to the HAH scheme. Of these 779 patients, 292 patients were actually

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
					discharged to the scheme.

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

1
2

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Huusko 2002 ¹⁵⁸ Finland Economic analysis: CCA Study design RCT Duration of follow-up: One year Perspective: NHS Discount rates: Costs: NA Effects: NA	Patient group: Patients with acute hip fracture All patients N: 243 Age (mean): 80 M: 69 F: 174 Drop outs: Group 1 N: 120 Age (mean, range): 80 (66-97) M: 36 F: 84 Drop outs: Group 2 N: 123 Age (mean, range): 80 (67-92) M: 33 F: 90 Drop outs:	Group 1: 2 weeks intensive rehabilitation on the geriatric ward Group 2: Standard care in a local hospital	Length of stay, days	Group 1: 34 (95% CI 28-38) Group 2: 42 (95% CI 35-48) p value: 0.05	Funding/conflict of interest: Study was supported by grants from Central Finland Health Care District, Kuopio University Hospital, Emil Aaltonen Foundation, Uulo Arthio Foundation and Novartis Finland Ltd Limitations: No sensitivity analysis No HRQoL Overall quality and applicability The study has limited applicability and potentially serious limitations Additional outcomes: Pre-fracture instrumental activities of daily living – IADL (median) – baseline to 3 months and baseline to
			Mortality (at discharge)	Group 1: 5 (4%) Group 2: 5 (4%)	
			Mortality (at 1 year)	Group 1: 18 (15%) Group 2: 20 (16%)	
			Patients regaining their independency in ADL – median, baseline to 3 months	Group 1: 5 Group 2: 6 p value: 0.004	
			Patients regaining their independency in ADL – median, baseline to 1 year	Group 1: 5 Group 2: 6 p value: 0.008	
			Mean cost per patient (includes hospital care, nursing home care, and outpatient services) PPP = 0.667223 (of 2002)	Group 1: € 17,900 (£11,723) Group 2: € 15,900 (£10,414) p value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Cost-effectiveness	NR	1 year Data sources: Notes:
			Sensitivity analysis	NR	

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

1
2

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>O’Cathain 1994²⁴⁵</p> <p>UK</p> <p>Economic analysis: CCA</p> <p>Study design Non randomised trial with concurrent controls</p> <p>Duration of follow-up: 3 months</p> <p>Perspective: NHS</p> <p>Discount rates NA</p>	<p>Patient group: Patients with fracture neck of femur</p> <p>All patients N: 110 Age (mean): NR M/F: 16/94 Drop outs: 14</p> <p>Group 1 N: 76 Age (mean): 76.4 (SD 10.0) M/F: 11/65 Drop outs: 8</p> <p>Group 2 N: 34 Age (mean): 77.6 (SD 9.7) M/F: 5/29 Drop outs: 6</p>	<p>Group 1: Hospital at home scheme (patients discharged to their own homes and cared for by a community HAH team under the clinical responsibility of the GP for a maximum of 12 days. The HAH team consisted of district nurses, community physiotherapists, occupational therapists and generic workers.)</p> <p>Group 2: Usual care</p>	Health-related QoL: Emotional reaction at discharge (from the Nottingham Health Profile questionnaire)(1)	<p>Group 1: 14</p> <p>Group 2: 24</p> <p>p value <0.05</p>	<p>Funding: The study was funded by Trent Regional Health Authority, Southern Derbyshire Community Health Services and Southern Derbyshire Department of Public Health.</p> <p>Conflict of interest: NR</p> <p>Limitations: The length of period during which costs are calculated is unclear.</p> <p>A longer follow up would have better reflected differences in costs and outcomes.</p> <p>No sensitivity analysis was conducted.</p> <p>No incremental analysis was conducted.</p> <p>Overall quality and applicability The study has potentially serious limitations and limited applicability</p> <p>Notes: (1) The other dimensions of the NHP (Physical mobility, pain, sleep, energy and social isolation) were not statistically significant.</p>
			Mortality	<p>Group 1: 5.3%</p> <p>Group 2: 5.9%</p> <p>p value: NR</p>	
			Readmission rate (at three months)	<p>Group 1: 15.8%</p> <p>Group 2: 8.8%</p> <p>p value: 0.187 (NS)</p>	
			Hospital LOS, median number of days (interquartile range)	<p>Group 1: 10</p> <p>Group 2: 17</p> <p>p value: <0.001</p>	
			Mean cost per patient	<p>Group 1: £1500</p> <p>Group 2: £1870</p> <p>p value: NR</p>	
			Year: 1992 Currency: UK sterling		
			Cost components: staff costs, orthopaedic bed cost.		
			Cost-effectiveness	NR	
			Sensitivity analysis	NR	

1 **Abbreviations:** NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis
2

1 Evidence table - Multidisciplinary rehabilitation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Parker 1991 ²⁷⁰ Economic analysis: CCA Study design Prospective observational study Duration of follow-up: 3 years Perspective: NHS Discount rates: Costs: NR Effects: NR	Patient group: Patients with acute hip fracture <u>All patients</u> N: 410 Age (mean): 77 F: 80% Drop outs: <u>Group 1</u> N: 284 Age (mean, range): 77 F: 79% Drop outs: 113 <u>Group 2</u> N: 126 Age (mean, range): 77 F: 83% Drop outs: NA	Group 1: early supported discharge scheme – hospital at home scheme Group 2: usual inpatient rehabilitation	LOS (mean, days)	Group 1: 29 Group 2: 38 p value: 0.035	Funding/conflict of interest: Limitations: No sensitivity analysis Costs were not discounted Overall quality and applicability The study has limited applicability and potentially serious limitations Additional outcomes: Data sources: Hospital records Notes: *HAH cost saving (-£799.80). Only 171 patients (60% of 284) were discharged using the HAH scheme, and the mean cost of the scheme refers to this group only.
			Mortality (at 90 days)	Group 1: 40 (14%) Group 2: 14 (11%)	
			Mean cost per patient	Group 1: £1165.30 Group 2: £365.50* p value: NR	
			Cost-effectiveness	NR	
			Sensitivity analysis	NR	

Abbreviations: NR=not reported, NA=not applicable, M/F=male/female, Sig=statistically significant at 5%, N=total number of patients randomised, Int=intervention, SA=sensitivity analysis

2
3
4

1 19 Appendix G: Forest plots

2	Figure 3-1: Illustration of precise and imprecise outcomes based on the confidence interval of.....	26
3	Figure G-2. Sensitivity and specificity: Sonography and isotope scanning (reference standard: MRI)	459
4	Figure G-3. Mortality: Early (≤ 24 hours) vs. late surgery.....	460
5	Figure G-4. Return to independent living: late (>24 hours) vs. early surgery	460
6	Figure G-5. Pressure ulcers: late (>24 hours) vs. early surgery.....	460
7	Figure G-6. Major complications: late (>24 hours) vs. early surgery.....	461
8	Figure G-7. Mortality – in hospital: late (24-48 hours) vs. early surgery.....	461
9	Figure G-8. Complications: late (24-48 hours) vs. early surgery.....	461
10	Figure G-9. Pressure ulcers: late (24-48 hours) vs. early surgery.....	461
11	Figure G-10. Mortality – at 4 months: late (>36 hours) vs. early surgery	462
12	Figure G-11. Pressure ulcers: late (>36 hours) vs. early surgery	462
13	Figure G-12. Return to independent living: late (>36 hours) vs. early surgery.....	462
14	Figure G-13. Mortality: late (>48 hours) vs. early surgery	462
15	Figure G-14. Return to independent living: late (>48 hours) vs. early surgery.....	463
16	Figure G-15. Pressure ulcers: late (>48 hours) vs. early surgery	463
17	Figure G-16. Major complications: late (>48 hours) vs. early surgery	463
18	Figure G-17. Minor complications: late (>48 hours) vs. early surgery	463
19	Figure G-18. Mortality – 30 days: late (>24 hours) vs. early surgery with the exclusion of patients unfit for surgery ..	464
20	Figure G-19. Combined mortality and needing total assistance in locomotion at 6 months: late (>24 hours) vs. early	
21	surgery with the exclusion of patients unfit for surgery	464
22	Figure G-20. Major post operative complications: late (>24 hours) vs. early surgery with the exclusion of patients unfit	
23	for surgery.....	464
24	Figure G-21. Mortality: late (>48 hours) vs. early surgery with the exclusion of patients unfit for surgery	464
25	Figure G-22. Change in residence (more dependent): late (>48 hours) vs. early surgery with the exclusion of patients	
26	unfit for surgery.....	465
27	Figure G-23. Return to original residence: late (>48 hours) vs. early surgery with the exclusion of patients unfit for	
28	surgery	465
29	Figure G-24. Pain: Nerve blocks vs. no block (systemic drugs).....	466
30	Figure G-25. Unsatisfactory pain control pre-operatively or 'need for breakthrough analgesia': Nerve blocks vs. no	
31	block (systemic drugs).....	466
32	Figure G-26. Unsatisfactory pain control postoperatively: Nerve blocks vs. (systemic drugs).....	467
33	Figure G-27. Nausea and/ or vomiting: Nerve blocks vs. (systemic drugs)	467
34	Figure G-28. Need for anti-emetics: Nerve blocks vs. (systemic drugs)	467
35	Figure G-29. Wound infection: Nerve blocks vs. (systemic drugs)	468
36	Figure G-30. Pneumonia: Nerve blocks vs. (systemic drugs).....	468
37	Figure G-31. Any cardiac complication: Nerve blocks vs. no block (systemic drugs)	469
38	Figure G-32. Myocardial infarction: Nerve blocks vs. no block (systemic drugs).....	469
39	Figure G-33. Puritis: Nerve blocks vs. no block (systemic drugs)	469
40	Figure G-34. Pulmonary embolism: Nerve blocks vs. no block (systemic drugs)	470
41	Figure G-35. Deep vein thrombosis: Nerve blocks vs. no block (systemic drugs).....	470
42	Figure G-36. Mortality: Nerve blocks vs. no block (systemic drugs).....	471
43	Figure G-37. Pressure sores: Nerve blocks vs. no block (systemic drugs).....	472
44	Figure G-38. Confusional state: Nerve blocks vs. no block (systemic drugs)	472

1	Figure G-40. Mortality at 1 month (random effects model): Regional (spinal or epidural) versus general anaesthesia	473
2	473
3	Figure G-41. Mortality- early up to 1 month: Regional (spinal or epidural) versus general anaesthesia	473
4	Figure G-42. Length of stay in hospital: Regional (spinal or epidural) versus general anaesthesia	474
5	Figure G-43. Vomiting: Regional (spinal or epidural) versus general anaesthesia	474
6	Figure G-44. Acute confusional state: Regional (spinal or epidural) versus general anaesthesia	474
7	Figure G-45. Pneumonia: Regional (spinal or epidural) versus general anaesthesia	475
8	Figure G-46. Myocardial infarction: Regional (spinal or epidural) versus general anaesthesia	475
9	Figure G-47. Pulmonary embolism (Peto odds ratio): Regional (spinal or epidural) versus general anaesthesia	476
10	Figure G-48. Pulmonary embolism (random effects model): Regional (spinal or epidural) versus general anaesthesia	476
11	476
12	Figure G-49. Pulmonary embolism (fatal and non fatal): Regional (spinal or epidural) versus general anaesthesia	477
13	Figure G-50. Deep vein thrombosis: Regional (spinal or epidural) versus general anaesthesia	477
14	Figure G-51. Reoperation rate for technically demanding hip fractures at 6 months: Senior/higher grade surgeon	478
15	versus junior/lower grade surgeon.....	478
16	Figure G-52. Dislocation rate for arthroplasty: Senior/higher grade surgeon versus junior/lower grade surgeon	478
17	Figure G-53. Perioperative mortality - older designs of arthroplasty: cemented vs. uncemented.	479
18	Figure G-54. Mortality – at up to 1 month - older designs of arthroplasty: cemented vs. uncemented.	479
19	Figure G-55. Mortality at between 1 and 3 months - older designs of arthroplasty: cemented vs. uncemented.	480
20	Figure G-56. Mortality at 1 year - older designs of arthroplasty: cemented vs. uncemented.	481
21	Figure G-57. Mortality at 3 years - older designs of arthroplasty: cemented vs. uncemented.	481
22	Figure G-58. Number of patients failing to regain mobility - older designs of arthroplasty: cemented vs. uncemented.	482
23	482
24	Figure G-59. Change in mobility score - older designs of arthroplasty: cemented vs. uncemented.	482
25	Figure G-60. Length of hospital stay - older designs of arthroplasty: cemented vs. uncemented.....	483
26	Figure G-61. Number of patients failing to return home - older designs of arthroplasty: cemented vs. uncemented.	483
27	Figure G-62. Number of patients reporting pain at 3 months - older designs of arthroplasty: cemented vs.	484
28	uncemented.	484
29	Figure G-63. Number of patients reporting pain at 1 to 2 years - older designs of arthroplasty: cemented vs.	484
30	uncemented.	484
31	Figure G-64. Pain score at 6 months - older designs of arthroplasty: cemented vs. uncemented.	485
32	Figure G-65. Reoperations - older designs of arthroplasty: cemented vs. uncemented.	485
33	Figure G-66. Deep sepsis - older designs of arthroplasty: cemented vs. uncemented.	486
34	Figure G-67. Wound haematoma - older designs of arthroplasty: cemented vs. uncemented.....	486
35	Figure G-68. Mortality - newer designs of arthroplasty: cemented vs. uncemented.	487
36	Figure G-69. Reoperations - newer designs of arthroplasty: cemented vs. uncemented.	487
37	Figure G-70. Pain – need for pain medication - newer designs of arthroplasty: cemented vs. uncemented.	487
38	Figure G-71. Unable to walk without aids at 12 months –newer designs of arthroplasty: cemented vs. uncemented.	487
39	487
40	Figure G-72. Barthel Index –newer designs of arthroplasty: cemented vs. uncemented.	488
41	Figure G-73. Harris Hip Score and Eq-5d scores –newer designs of arthroplasty: cemented vs. uncemented.	488
42	Figure G-74. Length of hospital stay –newer designs of arthroplasty: cemented vs. uncemented.....	488
43	Figure G-75. Mortality: Internal fixation versus hemiarthroplasty.....	489
44	Figure G-76. Reoperations: Internal fixation versus hemiarthroplasty.....	490
45	Figure G-77. Failure to return to same residence by final follow up: Internal fixation versus hemiarthroplasty.....	491
46	Figure G-78. Failure to regain mobility: Internal fixation versus hemiarthroplasty.....	491
47	Figure G-79. Patients reporting pain at 1 year: Internal fixation versus hemiarthroplasty	492
48	Figure G-80. Harris Hip Score: Internal fixation versus hemiarthroplasty	492
49	Figure G-81. Number of patients with Barthel Index Score of 95 or 100: Internal fixation versus hemiarthroplasty ...	493
50	Figure G-82. Euroqol Eq-5d score: Internal fixation versus hemiarthroplasty.....	493
51	Figure G-83. Length of hospital stay: Internal fixation versus hemiarthroplasty.....	493
52	Figure G-84. Mortality: Internal fixation versus total hip replacement	494
53	Figure G-85. Reoperations – all – at final follow up of study: Internal fixation versus total hip replacement.....	495
54	Figure G-86. Number of patients reporting pain at 1 year: Internal fixation versus total hip replacement.....	496
55	Figure G-87. Length of hospital stay: Internal fixation versus total hip replacement	496
56	Figure G-88. Mortality: Hemiarthroplasty versus total hip replacement	497
57	Figure G-89. Reoperations - all: Hemiarthroplasty versus total hip replacement.....	498

1	Figure G-90. Number of patients reporting pain at 1 year: Hemiarthroplasty versus total hip replacement.....	498
2	Figure G-91. Pain scores: Hemiarthroplasty versus total hip replacement.....	499
3	Figure G-92. Failure to regain mobility at end of study: Hemiarthroplasty versus total hip replacement	499
4	Figure G-93. Functional scores (lower scores advantageous): Hemiarthroplasty versus total hip replacement	499
5	Figure G-94. Functional status (higher scores advantageous): Hemiarthroplasty versus total hip replacement.....	500
6	Figure G-95. Quality of life scores: Hemiarthroplasty versus total hip replacement.....	501
7	Figure G-96. Length of hospital stay: Hemiarthroplasty versus total hip replacement	501
8	Figure G-97. 30 days mortality: Intramedullary implants versus extramedullary implants.....	502
9	Figure G-98. 3 months mortality: Intramedullary implants versus extramedullary implants	502
10	Figure G-99. 12 months mortality: Intramedullary implants versus extramedullary implants.....	503
11	Figure G-100. Reoperation – within the follow up period of the study: Intramedullary implants versus extramedullary	
12	implants.....	504
13	Figure G-101. Operative or postoperative fracture of femur - within the follow up period of the study: Intramedullary	
14	implants versus extramedullary implants.....	505
15	Figure G-102. Cut-out (at latest follow up): Intramedullary implants versus extramedullary implants.....	506
16	Figure G-103. Infection (deep infection or requires reoperation – at latest follow up): Intramedullary implants versus	
17	extramedullary implants	507
18	Figure G-104. Non-union (at latest follow-up): Intramedullary implants versus extramedullary implants.....	508
19	Figure G-105. Pain – patient reported outcomes: Intramedullary implants versus extramedullary implants.....	508
20	Figure G-106. Length of stay in hospital (in days): Intramedullary implants versus extramedullary implants.....	509
21	Figure G-107. Mean mobility score (Parker Palmer score): Intramedullary implants versus extramedullary implants	
22	509	
23	Figure G-96. 30 days mortality: Intramedullary implants versus extramedullary implants.....	510
24	Figure G-97. 3 months mortality: Intramedullary implants versus extramedullary implants	510
25	Figure G-98. 12 months mortality: Intramedullary implants versus extramedullary implants.....	510
26	Figure G-99. Reoperation – within the follow up period of the study: Intramedullary implants versus extramedullary	
27	implants.....	511
28	Figure G-100. Operative or postoperative fracture of femur - within the follow up period of the study: Intramedullary	
29	implants versus extramedullary implants.....	511
30	Figure G-101. Cut-out (at latest follow up): Intramedullary implants versus extramedullary implants.....	513
31	Figure G-102. Infection (deep infection or requires reoperation – at latest follow up): Intramedullary implants versus	
32	extramedullary implants	513
33	Figure G-103. Non-union (at latest follow-up): Intramedullary implants versus extramedullary implants.....	514
34	Figure G-104. Pain – patient reported outcomes: Intramedullary implants versus extramedullary implants.....	514
35	Figure G-105. Length of stay in hospital (in days): Intramedullary implants versus extramedullary implants	514
36	Figure G-106. Mean mobility score (Parker Palmer score): Intramedullary implants versus extramedullary implants	
37	515	
38	Figure G-108. Mortality at 12 months: Intramedullary implants versus extramedullary implants.....	515
39	Figure G-109. Reoperation within follow up period of the study: Intramedullary implants versus extramedullary	
40	implants.....	515
41	Figure G-110. Infection (deep infection or requires reoperation – at latest follow up): Intramedullary implants versus	
42	extramedullary implants	515
43	Figure G-111. Cut-out (at latest follow up): Intramedullary implants versus extramedullary implants.....	516
44	Figure G-112. Non-union (at latest follow up): Intramedullary implants versus extramedullary implants	516
45	Figure G-113. Independent to transfer at day 7: Early versus delayed mobilisation.....	517
46	Figure G-114. Independent to step at day 7: Early versus delayed mobilisation	517
47	Figure G-115. Discharge to home or rehabilitation programme: Early versus delayed mobilisation.....	517
48	Figure G-116. Discharge to nursing home or died: Early versus delayed mobilisation	518
49	Figure G-117. Strength measures: intensive physiotherapy versus usual care.....	519
50	Figure G-118. Tinetti's POMA (Performance Orientated Mobility Assessment): intensive physiotherapy versus usual	
51	care	519
52	Figure G-119. Functional performance measures: intensive physiotherapy versus usual care.....	520
53	Figure G-120. Functional performance tests: intensive physiotherapy versus usual care.....	520
54	Figure G-121. Walking speed: intensive physiotherapy versus usual care.....	520
55	Figure G-122. Knee extensor strength: intensive physiotherapy versus usual care	521
56	Figure G-123. Functional performance tests: intensive physiotherapy versus usual care.....	521
57	Figure G-124. Quality of life: intensive physiotherapy versus usual care.....	522
	Figure G-125. Walking speed: intensive physiotherapy versus usual care.....	522
	Figure G-126. Pain: intensive physiotherapy versus usual care	522


1 Figure G-127. Length of hospital stay: intensive physiotherapy versus usual care..... 523
 2 Figure G-128. Adductor muscle strength (kp) at 9 weeks: intensive physiotherapy versus usual care 523
 3 Figure G-129. Length of hospital stay: intensive physiotherapy versus usual care..... 523
 4 Figure G-130. Mortality at 6 months: hospital MDR versus usual care 524
 5 Figure G-131. Mortality at 12 months: hospital MDR versus usual care 524
 6 Figure G-132. Mortality (at discharge): hospital MDR versus usual care..... 525
 7 Figure G-133. Functional outcomes at 6 months: orthogeriatric hospital MDR versus usual care 525
 8 Figure G-134. Functional outcomes at 1 year: orthogeriatric hospital MDR versus usual care 526
 9 Figure G-135. Functional outcomes at 1 year: hip fracture programme versus usual care 526
 10 Figure G-136. : Functional outcomes: Barthel scores at long-term follow-up: hip fracture programme versus usual care
 11 526
 12 Figure G-137. Complications: hospital MDR versus usual care 527
 13 Figure G-138. Length of hospital stay: hospital MDR versus usual care 528
 14 Figure G-139. Readmitted to hospital during follow up: hospital MDR versus usual care 528
 15 Figure G-140. Mortality: Home-based MDR versus usual care 529
 16 Figure G-141. "Poor outcome" – institutional care and unable to walk: Home-based MDR versus usual care 529
 17 Figure G-142. SF-36 scores at 12 months (0: worst to 100: best): Home-based MDR versus usual care 529
 18 Figure G-143. Lengths of hospital or rehabilitation stays (days): Home-based MDR versus usual care..... 530
 19 Figure G-144. Readmission to hospital during 4 month follow-up: Home-based MDR versus usual care..... 530
 20 Figure G-145. Degree of independence (Functional Independent Measure): Home-based MDR versus usual care..... 530
 21 Figure G-146. Mobility and strength tests: Home-based MDR versus usual care 531
 22 Figure 147: Decision tree with Markov states - investment for early surgery vs. no hospital investment for early surgery
 23 545
 24 Figure 148: Cycle 0 Markov model 561
 25 Figure 149: Cycles 1 to 3 of the Markov model 563
 26 Figure 150: Cycle 4 - onwards of the Markov model 564
 27 Figure 151: Place of residence at discharge 565
 28 Figure 152: Incremental cost-effectiveness scatter plot: HFP vs GORU/MARU 589
 29 Figure 153: Incremental cost-effectiveness scatter plot: GORU/MARU vs. usual care 590
 30 Figure 154: Incremental cost-effectiveness scatter plot - HFP vs usual care 591
 31 Figure 155: Model structure - community MDR vs usual care 594
 32 Figure 156: Two-way sensitivity analysis on length of stay at home and in hospital 600
 33 Figure 157: Incremental cost-effectiveness scatter plot - Community MDR vs usual care 604
 34

19.1 Radiology

37 Figure G-2. Sensitivity and specificity: Sonography and isotope scanning (reference standard: MRI)


Sonography

Study	TP	FP	FN	TN	Sensitivity	Specificity
Safran 2009	10	7	0	13	1.00 [0.69, 1.00]	0.65 [0.41, 0.85]



isotope scanning

Study	TP	FP	FN	TN	Sensitivity	Specificity
Evans 1994	6	0	2	29	0.75 [0.35, 0.97]	1.00 [0.88, 1.00]
Rizzo 1993	36	0	1	23	0.97 [0.86, 1.00]	1.00 [0.85, 1.00]



19.2 Timing of surgery

Figure G-3. Mortality: Early (≤ 24 hours) vs. late surgery




Figure G-4. Return to independent living: late (> 24 hours) vs. early surgery






Figure G-5. Pressure ulcers: late (> 24 hours) vs. early surgery




1 Figure G-6. Major complications: late (>24 hours) vs. early surgery



2
3

4

5 Figure G-7. Mortality – in hospital: late (24-48 hours) vs. early surgery




6

7

8


9 Figure G-8. Complications: late (24-48 hours) vs. early surgery



10

11


12 Figure G-9. Pressure ulcers: late (24-48 hours) vs. early surgery



13

14


1 Figure G-10. Mortality – at 4 months: late (>36 hours) vs. early surgery



2

3


4 Figure G-11. Pressure ulcers: late (>36 hours) vs. early surgery



5

6


7 Figure G-12. Return to independent living: late (>36 hours) vs. early surgery



8

9

10 Figure G-13. Mortality: late (>48 hours) vs. early surgery




11

12

1 Figure G-14. Return to independent living: late (>48 hours) vs. early surgery

2




3

4

5

6 Figure G-15. Pressure ulcers: late (>48 hours) vs. early surgery

7




8

9

10 Figure G-16. Major complications: late (>48 hours) vs. early surgery

11



12


13

14 Figure G-17. Minor complications: late (>48 hours) vs. early surgery


15

16


17




1 Figure G-18. Mortality – 30 days: late (>24 hours) vs. early surgery with the exclusion of patients
 2 unfit for surgery




3
 4
 5 Figure G-19. Combined mortality and needing total assistance in locomotion at 6 months: late (>24
 6 hours) vs. early surgery with the exclusion of patients unfit for surgery



7
 8
 9 Figure G-20. Major postoperative complications: late (>24 hours) vs. early surgery with the
 10 exclusion of patients unfit for surgery




11
 12
 13
 14 Figure G-21. Mortality: late (>48 hours) vs. early surgery with the exclusion of patients unfit for
 15 surgery




1
2
3

Figure G-22. Change in residence (more dependent): late (>48 hours) vs. early surgery with the exclusion of patients unfit for surgery



4
5

6 Figure G-23. Return to original residence: late (>48 hours) vs. early surgery with the exclusion of
7 patients unfit for surgery




8
9
10
11

1

2 **19.3 Analgesia**

3 Figure G-24. Pain: Nerve blocks vs. no block (systemic drugs)


4



5

6

7 Figure G-25. Unsatisfactory pain control preoperatively or 'need for breakthrough analgesia':
8 Nerve blocks vs. no block (systemic drugs)




9

1

2 Figure G-26. Unsatisfactory pain control postoperatively: Nerve blocks vs. no block (systemic
3 drugs)


4



5

6 Figure G-27. Nausea and/ or vomiting: Nerve blocks vs. no block (systemic drugs)

7




8

9

10 Figure G-28. Need for anti-emetics: Nerve blocks vs. no nerve block(systemic drugs)

11




12

13

14

1 Figure G-29. Wound infection: Nerve blocks vs. no nerve block (systemic drugs)


2



3

4 Figure G-30. Pneumonia: Nerve blocks vs. no nerve block (systemic drugs)

5




6

7

8

1 Figure G-31. Any cardiac complication: Nerve blocks vs. no nerve block (systemic drugs)

2




3

4

5 Figure G-32. Myocardial infarction: Nerve blocks vs. no nerve block (systemic drugs)

6




7

8

9 Figure G-33. Puritis: Nerve blocks vs. no nerve block (systemic drugs)

10



11


12

13

14

1 Figure G-34. Pulmonary embolism: Nerve blocks vs. no nerve block (systemic drugs)

2




3

4

5 Figure G-35. Deep vein thrombosis: Nerve blocks vs. no nerve block (systemic drugs)

6




7

8

9

1 Figure G-36. Mortality: Nerve blocks vs. no nerve block (systemic drugs)


2



3
4
5
6

1 Figure G-37. Pressure sores: Nerve blocks vs. no nerve block (systemic drugs)

2




3

4

5 Figure G-38. Confusional state: Nerve blocks vs. no nerve block (systemic drugs)

6




7

8

1 **19.4 Anaesthesia**

2 **Figure G-39. Mortality at 1 month (random effects model): Regional (spinal or epidural)**
 3 **versus general anaesthesia**

4




5
6

7 **Figure G-40. Mortality- early up to 1 month: Regional (spinal or epidural) versus general**
 8 **anaesthesia**

9


10 Additional analysis: The authors pooled mortality data from Adams 1990 and Bigler 1985 which reported
 11 early mortality during hospital stay and Ungemach 1987 which reported mortality at 2 weeks with data
 12 from the mortality at one month analysis.

13




14
15
16


1 Figure G-41. Length of stay in hospital: Regional (spinal or epidural) versus general
2 anaesthesia



3
4
5 Figure G-42. Vomiting: Regional (spinal or epidural) versus general anaesthesia




6
7
8 Figure G-43. Acute confusional state: Regional (spinal or epidural) versus general
9 anaesthesia



10
11


12

1 Figure G-44. Pneumonia: Regional (spinal or epidural) versus general anaesthesia



2
3
4
5


4 Figure G-45. Myocardial infarction: Regional (spinal or epidural) versus general
5 anaesthesia



6
7
8

1 Figure G-46. Pulmonary embolism (Peto odds ratio): Regional (spinal or epidural) versus
 2 general anaesthesia


3



4
5

6 Figure G-47. Pulmonary embolism (random effects model): Regional (spinal or epidural)
 7 versus general anaesthesia


8



9
10
11

1 Figure G-48. Pulmonary embolism (fatal and non fatal): Regional (spinal or epidural)
 2 versus general anaesthesia


3



4

5 Figure G-49. Deep vein thrombosis: Regional (spinal or epidural) versus general
 6 anaesthesia

7



8


9

1 19.5 Surgical interventions

2 19.5.1 Surgeon seniority

3 Figure G-50. Reoperation rate for technically demanding hip fractures at 6 months:
4 Senior/higher grade surgeon versus junior/lower grade surgeon


5



6

7

8 Figure G-51. Dislocation rate for arthroplasty: Senior/higher grade surgeon versus
9 junior/lower grade surgeon




10

11

1 **19.5.2 Cement in older designs of arthroplasty**

2


3 Figure G-52. Perioperative mortality - older designs of arthroplasty: cemented vs.
4 uncemented.



5

6


7 Figure G-53. Mortality – at up to 1 month - older designs of arthroplasty: cemented vs.
8 uncemented.



9

10


1 Figure G-54. Mortality at between 1 and 3 months - older designs of arthroplasty:
 2 cemented vs. uncemented.




3

4

1 Figure G-55. Mortality at 1 year - older designs of arthroplasty: cemented vs.
 2 uncemented.




3
 4
 5 Figure G-56. Mortality at 3 years - older designs of arthroplasty: cemented vs.
 6 uncemented.



8
 9
 10
 11


1 Figure G-57. Number of patients failing to regain mobility - older designs of arthroplasty:
 2 cemented vs. uncemented.

3



4

5
 6 Figure G-58. Change in mobility score - older designs of arthroplasty: cemented vs.
 7 uncemented.




8

9

10


1 Figure G-59. Length of hospital stay - older designs of arthroplasty: cemented vs.
 2 uncemented.

3



4
5
6


7 Figure G-60. Number of patients failing to return home - older designs of arthroplasty:
 8 cemented vs. uncemented.



9
10
11


1 Figure G-61. Number of patients reporting pain at 3 months - older designs of
 2 arthroplasty: cemented vs. uncemented.

3



4
5
6


7 Figure G-62. Number of patients reporting pain at 1 to 2 years - older designs of
 8 arthroplasty: cemented vs. uncemented.



10
11
12


1 Figure G-63. Pain score at 6 months - older designs of arthroplasty: cemented vs.
 2 uncemented.

3




4

5 Figure G-64. Reoperations - older designs of arthroplasty: cemented vs. uncemented.




6
7

1 Figure G-65. Deep sepsis - older designs of arthroplasty: cemented vs. uncemented.



2

3 Figure G-66. Wound haematoma - older designs of arthroplasty: cemented vs.
4 uncemented.




5

6


1 **19.5.3 Cement in newer designs of arthroplasty**

2 **Figure G-67. Mortality - newer designs of arthroplasty: cemented vs. uncemented.**



3

4 **Figure G-68. Reoperations - newer designs of arthroplasty: cemented vs. uncemented.**




5

6

7 **Figure G-69. Pain – need for pain medication - newer designs of arthroplasty: cemented**

8 **vs. uncemented.**




9


10

11 **Figure G-70. Unable to walk without aids at 12 months –newer designs of arthroplasty:**


12 **cemented vs. uncemented.**




1
2
3 Figure G-71. Barthel Index –newer designs of arthroplasty: cemented vs. uncemented.



4
5
6 Figure G-72. Harris Hip Score and Eq-5d scores –newer designs of arthroplasty: cemented
7 vs. uncemented.




8
9
10 Figure G-73. Length of hospital stay –newer designs of arthroplasty: cemented vs.
11 uncemented.



1 **19.5.4 Internal fixation versus hemiarthroplasty**

2 **Figure G-74. Mortality: Internal fixation versus hemiarthroplasty**


3



4
5

1 Figure G-75. Reoperations: Internal fixation versus hemiarthroplasty


2



34567

1 Figure G-76. Failure to return to same residence by final follow up: Internal fixation versus
 2 hemiarthroplasty


3



4
5

6 Figure G-77. Failure to regain mobility: Internal fixation versus hemiarthroplasty


7



8
9


1 Figure G-78. Patients reporting pain at 1 year: Internal fixation versus hemiarthroplasty

2



3
4


5 Figure G-79. Harris Hip Score: Internal fixation versus hemiarthroplasty



6
7
8


1 Figure G-80. Number of patients with Barthel Index Score of 95 or 100: Internal fixation
 2 versus hemiarthroplasty

3




4
5

6 Figure G-81. Euroqol Eq-5d score: Internal fixation versus hemiarthroplasty



7
8
9


10 Figure G-82. Length of hospital stay: Internal fixation versus hemiarthroplasty



11
12


1 **19.5.5 Internal fixation versus total hip replacement**

2 **Figure G-83. Mortality: Internal fixation versus total hip replacement**




3
4

1 Figure G-84. Reoperations – all – at final follow up of study: Internal fixation versus total
 2 hip replacement




3
4
5
6

1 Figure G-85. Number of patients reporting pain at 1 year: Internal fixation versus total hip
 2 replacement




3
 4
 5 Figure G-86. Length of hospital stay: Internal fixation versus total hip replacement



6
 7
 8


1 **19.5.6 Hemiarthroplasty versus total hip replacement**

2 **Figure G-87. Mortality: Hemiarthroplasty versus total hip replacement**




3
4
5

1 Figure G-88. Reoperations - all: Hemiarthroplasty versus total hip replacement




2
3
4 Figure G-89. Number of patients reporting pain at 1 year: Hemiarthroplasty versus total
5 hip replacement




6
7
8

1 Figure G-90. Pain scores: Hemiarthroplasty versus total hip replacement




2
3

4 Figure G-91. Failure to regain mobility at end of study: Hemiarthroplasty versus total hip replacement




6
7

8 Figure G-92. Functional scores (lower scores advantageous): Hemiarthroplasty versus total hip replacement




10
11
12

1 Figure G-93. Functional status (higher scores advantageous): Hemiarthroplasty versus
 2 total hip replacement




3
4
5

1 Figure G-94. Quality of life scores: Hemiarthroplasty versus total hip replacement



2
3


4 Figure G-95. Length of hospital stay: Hemiarthroplasty versus total hip replacement



5
6
7


1 **19.5.7 Trochanteric extracapsular fracture – all studies**

2 **Figure G-96. 30 days mortality: Intramedullary implants versus extramedullary implants**




3
4

5 **Figure G-97. 3 months mortality: Intramedullary implants versus extramedullary implants**



7
8


1 Figure G-98. 12 months mortality: Intramedullary implants versus extramedullary
 2 implants



3
4


5

1 Figure G-99. Reoperation – within the follow up period of the study: Intramedullary
 2 implants versus extramedullary implants




1 Figure G-100. Operative or postoperative fracture of femur - within the follow up period
 2 of the study: Intramedullary implants versus extramedullary implants

3



4


1 Figure G-101. Cut-out (at latest follow up): Intramedullary implants versus extramedullary
 2 implants



3
4


5

- 1 Figure G-102. Infection (deep infection or requires reoperation – at latest follow up):
- 2 Intramedullary implants versus extramedullary implants




- 3
- 4

1 Figure G-103. Non-union (at latest follow-up): Intramedullary implants versus
 2 extramedullary implants




3
4
5
6
7

6 Figure G-104. Pain – patient reported outcomes: Intramedullary implants versus
 7 extramedullary implants




8
9

1 Figure G-105. Length of stay in hospital (in days): Intramedullary implants versus
 2 extramedullary implants



3
4


5 Figure G-106. Mean mobility score (Parker Palmer score): Intramedullary implants versus
 6 extramedullary implants




7
8
9

1 **19.5.8 Trochanteric extracapsular fracture – studies from 2000**


2 Figure G-107. 30 days mortality: Intramedullary implants versus extramedullary implants



4
5
6 Figure G-108. 3 months mortality: Intramedullary implants versus extramedullary
7 implants




10 Figure G-109. 12 months mortality: Intramedullary implants versus extramedullary
11 implants




1
2
3

4 Figure G-110. Reoperation – within the follow up period of the study: Intramedullary
5 implants versus extramedullary implants




6

7 Figure G-111. Operative or postoperative fracture of femur - within the follow up period
8 of the study: Intramedullary implants versus extramedullary implants




1 Figure G-112. Cut-out (at latest follow up): Intramedullary implants versus extramedullary
 2 implants



3


4 Figure G-113. Infection (deep infection or requires reoperation – at latest follow up):
 5 Intramedullary implants versus extramedullary implants

6




7


1 Figure G-114. Non-union (at latest follow-up): Intramedullary implants versus
 2 extramedullary implants




3
 4
 5
 6 Figure G-115. Pain – patient reported outcomes: Intramedullary implants versus
 7 extramedullary implants



8
 9 Figure G-116. Length of stay in hospital (in days): Intramedullary implants versus
 10 extramedullary implants




1 Figure G-117. Mean mobility score (Parker Palmer score): Intramedullary implants versus
 2 extramedullary implants



3

4 **19.5.9 Subtrochanteric extracapsular fracture.**


5 Figure G-118. Mortality at 12 months: Intramedullary implants versus extramedullary
 6 implants



7

8


9 Figure G-119. Reoperation within follow up period of the study: Intramedullary implants
 10 versus extramedullary implants



11

12


13 Figure G-120. Infection (deep infection or requires reoperation – at latest follow up):
 14 Intramedullary implants versus extramedullary implants



15


16

1 Figure G-121. Cut-out (at latest follow up): Intramedullary implants versus extramedullary
2 implants



3
4

5 Figure G-122. Non-union (at latest follow up): Intramedullary implants versus
6 extramedullary implants



7

8


1

2 **19.6 Mobilisation strategies**

3 **19.6.1 Timing of mobilisation**

4 Figure G-123. Independent to transfer at day 7: Early versus delayed mobilisation

5




6

7

8 Figure G-124. Independent to step at day 7: Early versus delayed mobilisation

9




10

11

12 Figure G-125. Discharge to home or rehabilitation programme: Early versus delayed
13 mobilisation

14




15

16

1 Figure G-126. Discharge to nursing home or died: Early versus delayed mobilisation

2




3

4

1 **19.7 Intensive exercise or physiotherapy vs. usual care**

2 **19.7.1 Intensive physiotherapy (Strength training)**

3 Figure G-127. Strength measures: intensive physiotherapy versus usual care




4

5

6 Figure G-128. Tinetti's POMA (Performance Orientated Mobility Assessment): intensive
7 physiotherapy versus usual care


8



9
10


1 Figure G-129. Functional performance measures: intensive physiotherapy versus usual
2 care

3



4

5 Figure G-130. Functional performance tests: intensive physiotherapy versus usual care




6

7

8 Figure G-131. Walking speed: intensive physiotherapy versus usual care

9



10


11

12

1 **19.7.2 Intensive physiotherapy (treadmill training)**

2 Figure G-132. Knee extensor strength: intensive physiotherapy versus usual care


3



4

5


6 Figure G-133. Functional performance tests: intensive physiotherapy versus usual care



7


8

1 Figure G-134. Quality of life: intensive physiotherapy versus usual care




2
3

4 Figure G-135. Walking speed: intensive physiotherapy versus usual care




5
6

7 Figure G-136. Pain: intensive physiotherapy versus usual care



8
9
10


1 Figure G-137. Length of hospital stay: intensive physiotherapy versus usual care



2

3 **19.7.3 Intensive (more frequent) physiotherapy**


4 Figure G-138. Adductor muscle strength (kp) at 9 weeks: intensive physiotherapy versus
5 usual care



7

8

9 Figure G-139. Length of hospital stay: intensive physiotherapy versus usual care



11

12

13

14


15

1 **19.8 Multidisciplinary rehabilitation**

2 **19.8.1 Hospital-based MDR**


3 Hospital based MDR has been split into orthogeriatric hospital MDR (including GORU and
4 MARU) and hip fracture programmes.

5 Figure G-140. Mortality at 6 months: hospital MDR versus usual care



7
8


9 Figure G-141. Mortality at 12 months: hospital MDR versus usual care



11
12

1 Figure G-142. Mortality (at discharge): hospital MDR versus usual care

2




3

4

5 Figure G-143. Functional outcomes at 6 months: orthogeriatric hospital MDR versus usual care

6

7




8

9

1 Figure G-144. Functional outcomes at 1 year: orthogeriatric hospital MDR versus usual
2 care


3



4
5

6 Figure G-145. Functional outcomes at 1 year: hip fracture programme versus usual care


7



8
9


10 Figure G-146. : Functional outcomes: Barthel scores at long-term follow-up: hip fracture
programme versus usual care

11



12
13


1 Figure G-147. Complications: hospital MDR versus usual care



2

1 Figure G-148. Length of hospital stay: hospital MDR versus usual care

2




3

4

5 Figure G-149. Readmitted to hospital during follow up: hospital MDR versus usual care

6




7

8

1 **19.9 Home-based MDR versus usual inpatient rehabilitation**


2 Figure G-150. Mortality: Home-based MDR versus usual care



3

4


5 Figure G-151. "Poor outcome" – institutional care and unable to walk: Home-based MDR
6 versus usual care



8

9

10 Figure G-152. SF-36 scores at 12 months (0: worst to 100: best): Home-based MDR versus
11 usual care




13

14

15

1 Figure G-153. Lengths of hospital or rehabilitation stays (days): Home-based MDR versus
 2 usual care


3



4

5 Figure G-154. Readmission to hospital during 4 month follow-up: Home-based MDR
 6 versus usual care

7




8

9

10 Figure G-155. Degree of independence (Functional Independent Measure): Home-based
 11 MDR versus usual care

12




13

14

1 Figure G-156. Mobility and strength tests: Home-based MDR versus usual care

2



3
4

1

2 20 Appendix H: Health economic analysis

3 20.1 Cost analysis of nerve blocks, non-opioid analgesics and 4 opioid analgesics

5 20.1.1 Nerve block cost analysis

6

7 No studies were identified on the cost-effectiveness of nerve blocks compared to systemic
8 analgesia in providing adequate pain relief and reducing side effects and mortality.

9

10 As a consequence, we conducted a cost analysis where the different types and level of resources
11 used to administer a nerve block to a patient with a suspected hip fracture are based on the
12 GDG's opinion, summarised in Table 73 below.

13

14 **Table 73: Cost analysis for nerve block**

Resources	Unit price	Source of unit price
Spinal pack (gown and drape) *	£4.50	NHS hospital***
Biogel glove	£1.07	NHS hospital***
Chlorhexidine**	£1.08	NHS hospital***
Vial with Lidocaine 1%	£0.38 (10-mL am)	BNF 58
Vial of 0.5% Levobupivacaine	£3.88 (5mg/mL)	BNF 58
Syringes (10ml)	£0.06	NHS hospital***
Filter needle	£0.23	NHS hospital***
Regional block needle	£5.78	NHS hospital***
Hypodermic needle	£1.35	NHS hospital***
Personnel costs (consultant anaesthetist)	£36.00	PSSRU 2009; GDG estimate (£1.8 per minute*20 minutes)
Total cost	£54.33	

15 * Most anaesthetists use full aseptic precautions, with a gown and gloves plus a dressing pack

16 ** Chlorhexidine built into swabs are standard practice.

17 *** Peterborough and Stamford Hospital NHS Foundation Trust

18

1 The **personnel costs** can vary depending on the time required to administer a nerve block,
2 which in turn depends on the technique used (nerve stimulator, ultrasound- guided,
3 landmark only) and the block used (3-in-1, femoral nerve only or fascia iliaca block). If a
4 **fascia iliaca** block is administered using a landmark technique only, then the following
5 sequence would be observed:

- 6 - Obtaining equipment (needle, disinfectant, gloves, local anaesthetic etc)
- 7 - Estimating patient's weight
- 8 - Obtaining patient's consent
- 9 - Identifying landmark
- 10 - Disinfecting skin
- 11 - Anaesthetising skin
- 12 - Passing needle
- 13 - Injecting local anaesthetic
- 14 - Maintaining manual pressure distal to injecting site for a minute after injection

15
16 The GDG estimates that the whole process would require about 15 - 20min, and that the time
17 required would not change substantially if the block is administered by a consultant anaesthetist
18 or a SAS (staff and associate specialist).

19
20 At present, in most emergency departments that do advocate nerve blocks for hip fracture
21 patients, the block would be performed by 'middle grade doctors', i.e. specialist registrars (SpR),
22 senior specialist trainees (ST3-6) or senior clinical fellows. In some departments, junior doctors
23 can also administer the procedure. In operating departments and if asked to do elsewhere
24 anaesthetists will always have a trained assistant with them, usually an ODP, which would
25 increase the total cost for a nerve block to £63.33 (assuming an ODP wage of £27 per hour as that
26 of a senior nurse)

27
28 The GDG recognises that there is likely to be a wide variation in practice as far as the
29 administration of nerve blocks is concerned.

- 30
31 1) The nerve block may be administered with a ultrasound-guided technique, which would
32 require the use of ultrasound anaesthetic machines. An average cost of these machines
33 has been estimated at around £34,000 from hospital records supplied by the
34 Peterborough and Stamford Hospital NHS Foundation Trust. The equivalent annual cost
35 would be £5,313, assuming a life expectancy of 7 years and discount rate of 3.5%.

36
37 If we assume that the ultrasound machine would be used solely for nerve blocks in the
38 anaesthetic department and that it would be used 7 hours per day every day, including
39 weekends with 4 scans per hour, then the machine costs 52p per scan.

- 40
41 2) Bupivacaine can be used as local anaesthetic instead of Levobupivacaine, but the
42 difference in price would be minimal.
- 43
44 3) A nerve locator could be used when performing the nerve block, but its cost would be
45 minimal (GDG expert's opinion)
- 46
47

1 **20.1.2 Non-opioid analgesics**

2 We assume that patients will take a simple analgesic, such as paracetamol, continuously
 3 throughout their inpatient stay. The GDG noted that aspirin would not generally be used as an
 4 analgesic for our population, unless it is used as a low dose to prevent strokes. The average cost
 5 of these drugs is less than £0.1p per dose (BNF 58).

6 **20.1.3 Opioid analgesics**

7
 8 Table 74: Opioids controlled drugs

Category	Dose cost (source: BNF 58)
Diamorphine hydrochloride	£2.69
Morphine salts	£0.36
Oxycodone hydrochloride	£1.60
Buprenorphine	£0.72
Average cost	£1.34

9
 10 The opioids reported in Table 743 are non-controlled drugs and can be administered within
 11 existing nurse drug rounds, and therefore there is little extra cost associated with their
 12 administration.

13
 14 Table

15 2 summarises the opioids controlled drugs that could be administered to hip fracture
 16 patients. This category of analgesics requires an additional round of two trained nurses to
 17 administer. The GDG estimates that this would involve approximately 15 minutes per
 18 dose, with an extra cost of £10.50 (considering that the cost per hour of a staff nurse is
 19 £21 (PSSRU 2009)). Hence, the cost of administering these controlled drugs is £11.84
 20 (nurse time plus drug cost).

21

22 **Table 74: Opioids controlled drugs**

Category	Dose cost (source: BNF 58)
Diamorphine hydrochloride	£2.69
Morphine salts	£0.36
Oxycodone hydrochloride	£1.60
Buprenorphine	£0.72
Average cost	£1.34

1

2 The opioids reported in Table 743 are non-controlled drugs and can be administered within
3 existing nurse drug rounds, and therefore there is little extra cost associated with their
4 administration.
5

6

Table 3: Opioids non-controlled drugs

Category	Dose cost (source: BNF 58)
Codeine phosphate	£1.83
Dihydrocodeine Tartrate	£2.58
Tramadol Hydrochloride	£1.47
Average cost	£1.96

7 The remaining opioid drugs (dipipanone hydrochloride, hydromorphone hydrochloride,
8 meptazinol, methadone hydrochloride, paperetum, pentazocine, pethidine hydrochloride) are
9 very rarely used in our population, as they are highly specialist analgesics for palliative care.
10 Fentanyl is rarely used in acute care, and is therefore not included in the dose cost.
11

12

13

14 **20.2 Hourly wage costs for a planned trauma list**

15 The GDG suggested to consider a general emergency theatre as a likely alternative to a
16 planned trauma list.
17

18

18 A general emergency theatre is one to which multiple specialities have access for
19 unplanned operations. Under these circumstances there will be necessary discussions
20 between the various specialties as to whose patient should go first. With an emergency
21 theatre, there is no start and finish time that can be forecasted in advance and great
22 variation in the professional grade of the personnel involved.
23

24

24 When the hip fracture patient does go to theatre, he will clearly need the same supporting
25 staff of surgeon, anaesthetist, nursing staff, radiographer etc. as for a planned trauma list.
26 Thus, some costs will be common across the two types of lists with the exception that an
27 emergency trauma list is more likely involve more junior staff.
28

29

29 Overall, the GDG has identified the following differences between an emergency and a
30 planned trauma list:
31

32

33

- a) Senior responsible staff involved

34

34 With a general emergency theatre the involvement of senior staff may be regarded as a
35 covering on-call commitment. With a trauma list it becomes a regular work commitment to
36 which there needs to be programmed activities allocated for both senior responsible
37 anaesthetic and surgical staff. Since the nature of the work is known appropriate scrub
38 staff can be allocated
39

40

- b) Where necessary a new operating room

1
2
3 Providing trauma cases with the same level of care enjoyed by elective cases may require
4 extra operating theatre space. There have been attempts in many hospitals to use
5 operating theatres for a greater proportion of the 24 hour day to better use that resource.
6 This has in general proved to be difficult; largely because trained staff prefer to have their
7 regular commitments in what would be regarded as normal working hours. Genuine
8 emergency procedures are a small proportion of any theatre workload and these need to
9 be carried out at the necessary time whenever that may be. However, the bulk of
10 procedures are urgent or elective, these should all be given the same advantages of a
11 properly staffed theatre. Should it be necessary for best use of theatre space to utilise
12 evening operating lists it may be preferable that these are occupied by the well prepared
13 elective patients rather than the rapidly prepared often unwell urgent patient. Since this is
14 unlikely to occur more operating space may well be required for daytime lists.

15
16 The advantage of a general emergency list is it uses the resources already available, and
17 may run from early in the morning till late in the evening (therefore many operations can
18 be performed sequentially). On the other hand, a planned trauma list needs to be run in
19 parallel with other lists, preferably in the morning. It may be difficult to find a physical
20 space for a planned trauma list to be carried out, in which case a new operating room may
21 be required.

22
23
24 c) Ad-hoc technical resources

25
26 A planned trauma list needs a dedicated image intensifier, so it depends upon the other
27 lists running as to whether its availability may be a problem.

28
29
30 d) Type of patients operated

31
32 A planned trauma list would only operate trauma patients whereas in a general
33 emergency theatre there would be operations on different types of patients

34
35
36 The table below estimates the cost of one hour of personnel input for a planned trauma
37 list during weekly normal working hours (that is, excluding weekends and public holidays
38 personnel costs).

39
40 **Personnel input cost for a planned trauma list – weekly normal working hours**

Categories of personnel	Cost of hourly wage (source: PSSRU 2009)
Consultant surgeon	£108
Consultant anaesthetist	£108
Scrub nurse (senior staff nurse)	£27
Unscrub nurse (runner – staff nurse)	£21

Radiographer	£25
Anaesthesia assistant [ODP] (as senior staff nurse)	£27
Recovery nurse (staff nurse)	£21
Total personnel costs	£337

1
2 As for the personnel costs of a general emergency theatre, we assume that it mainly relies
3 on registrars (both surgeons and anaesthetists) rather than consultants, and use a hourly
4 cost for registrars of £38 (per 48 hour week; source: PSSRU 2009⁶¹). Any emergency
5 theatre also relies on having consultant surgeons and anaesthetist on call, and this cost
6 would also have to be considered in the overall costs for an emergency theatre. Once
7 again we consider the personnel costs during weekly normal working hours, and thus
8 exclude weekends and public holidays personnel costs nor additional personnel costs for
9 out-of-hours operations, which are quite common with a general emergency theatre.

10
11
12

Personnel input cost for an general emergency theatre – weekly normal working hours

Categories of personnel	Cost of hourly wage (source: PSSRU 2009)
Registrar surgeon	£38
Registrar anaesthetist	£38
Consultant surgeon on call*	£23
Consultant anaesthetist on call*	£23
Scrub nurse (senior staff nurse)	£27
Unscrub nurse (runner – staff nurse)	£21
Radiographer	£25
Anaesthesia assistant [ODP] (as senior staff nurse)	£27
Recovery nurse (staff nurse)	£21
Total personnel costs	£243

13 *= Assumes that the average emergency work undertaken per week for on-call duty is 3
14 hours. If the amount of this emergency on-call work raises to 6 hours per week, the hourly
15 rate paid to the consultant would be £39. Source: hourly on call salary costs provided by
16 the NICE costing implementation team.

17
18
19

Thus, a planned trauma list has additional personnel cost compared to a general
emergency theatre of £94 per hour. It is very important to stress that this estimate does

1 not consider the additional salary costs linked with operations taking place during
 2 weekends or public holidays and outside normal working hours.
 3

4 **20.3 Prices for sliding hip screws and short and long**

Manufacturer	Price for Sliding Hip Screw (for extramedullary fixation) IMP	Price for Short intramedullary nail (for intramedullary fixation)	Price for Long intramedullary nail (for intramedullary fixation)
Stryker	£357	£854	£1384
Biomet	£260.70	£745	£1,090
Zimmer (1)	£175	£826	£1,177
Synthes	£260.35	£796.05	£1,142.85
Smith & Nephew (2)	£245	£823.45	£1,083.16
DePuy	£217	£516	NA
Average price	£252.51	£760.08	£1,175.40

5 **intramedullary nails**

6
 7 In the table above we report the prices for sliding hip screws, short intramedullary and long intramedullary
 8 nails from quotations received by some of the major manufacturers of implants. All quotations are 2010
 9 prices. All prices include VAT.
 10
 11

12

1 **20.4 Cost analysis of the interventions for intensive mobilisation strategies.**

2 Cost analysis of the interventions for intensive mobilisation strategies.

Study	Intervention	Control	Other resources	Unit costs	Incremental cost of intervention over usual care
Hauer et al 2002 ¹⁴⁰	1 hour of physiotherapist for 3 weeks	1 hour of physiotherapist for 3 weeks	Using data provided from a GDG member, the cost of the equipment that would be used in the intervention group was estimated at £49.00 per patient. This estimate is based on a study currently under way, where the costs per person for the exercise equipment was estimated to be £49.00. This cost assumes no re-use of equipment and does not include overhead costs. When appropriately cleaned, the equipment could be re-used, in which case, assuming that it is re-used up to four times, the relevant cost per person would be approximately £12.	£23 per hour for physiotherapist input Other costs (for stepping and strength training) are considered as negligible and have not been included in the cost analysis	£12
Karumo 1977A ¹⁷¹	Physiotherapy performed twice daily – average of 1 hour for 14 days	Average of 30mins physiotherapy per day for 14 days	Crutches	£23 per hour for physiotherapist input (£161 control; £322 intervention)	£180.18

				Cost of crutches: £19.18 (a)	
Moseley et al., 2009²¹⁶	<p><u>Weight bearing</u> exercise twice daily for a total of 60 minutes per day for 16 weeks.</p> <p>Walking on a treadmill with partial body weight support using a harness (for inpatients) or a walking programme (after hospital discharge).</p> <p>LOS in hospital: 28 days (4 weeks)</p> <p>For 84 days: walking programme with home visits and exercise programme</p> <p>This started as an inpatient programme, followed by home visits and a structured home exercise programme.</p>	<p>Exercise for 30 mins each day for 4 weeks.</p> <p>LOS in hospital: 25 days.</p>	<p>For inpatients: additional inpatients costs</p> <p>Treadmill with partial weight-support</p>	<p>£23 per hour for physiotherapist input</p> <p>£13,029 for Biodex treadmill with body-weight support.</p> <p>Cost per day of a bed (elderly person care: £152 (b))</p> <p>Assumption: a physiotherapist is present for all the duration of treatment when inpatient</p> <p>Treadmill costs – assumptions:</p> <ul style="list-style-type: none"> - Treadmill life is 5 years - Treadmill overall use: 4 hours per day for 5 days of the week - Discount rate: 3.5% - Treadmill used for 20 minutes per session - Cost one session of treadmill imputable to the intervention: £0.54. - Cost of treadmill sessions over 4 weeks: $7*4*£0.54=£15.12$ <p>INTERVENTION COSTS:</p> <p>Bed days cost: $£152*28=£4256$</p> <p>Attributable treadmill costs: £15.12 per patient</p>	<p>£827.62 (for the inpatient part of the rehabilitation programme)</p> <p>The costs of the outpatient part of the programme was not calculated as it was not clear from the study what types of resources were used in that part of the rehab programme.</p>

				Physiotherapist costs (intervention): $\text{£}23 * 28 = \text{£}644$	
				Total inpatient costs of intervention: £4915.12	
				CONTROL COSTS: Physiotherapist costs: $\text{£}23 * 0.5 * 25 = \text{£}287.5$	
				Bed day costs: $\text{£}152 * 25 = \text{£}3800$	
				Total cost for control: £4087.5	

1

2 (a) Average cost obtained from the NHS Supply Catalogue 2010 for the following manufacturers: Sunrise Medical Ltd, NHS Supply Chain and Days
 3 Healthcare UK Limited

4 (b): We have estimated the hospital stay using the unit cost per excess day associated with complex elderly patients (that is, the unit cost per day for days
 5 exceeding the trim point). Using all the HRG unit costs reported for all Complex Elderly patients (Hospital Episode Statistics for England, Inpatient Statistics,
 6 2007-08) we found a weighted mean of £152.

7

1 **20.5 Cost-effectiveness analysis of hospital investment versus no** 2 **hospital investment for early surgery**

3 **20.5.1 Introduction**

4 The GDG assigned a high priority in the economic plan for an original economic analysis to the
5 question:

6 “*In patients with hip fractures what is the clinical and cost effectiveness of early surgery (within*
7 *24, 36 or 48 hours) on the incidence of complications such as mortality, pneumonia, pressure*
8 *sores, cognitive dysfunction and increased length of hospital stay?”*

9 A review of the literature was conducted. The literature search and review methods can be found
10 in Chapter 3. No cost-effectiveness analysis was found which addressed our clinical question. As a
11 consequence, the GDG felt that an original decision model was essential in order to inform their
12 recommendations.

13 The following general principles were adhered to:

- 14 • The GDG was consulted during the construction and interpretation of the model.
 - 15 • When published data was not available, we used hospital records and experts’ opinion to
16 populate the model.
 - 17 • Model assumptions were reported fully and transparently.
 - 18 • The results were subject to sensitivity analysis and limitations were discussed.
 - 19 • We followed the methods of the NICE reference case. Therefore costs were calculated
20 from the NHS and PSS perspective. Health gain was measured in terms of quality-
21 adjusted life-years (QALYs) gained. Both future costs and QALYs were discounted at 3.5%.
 - 22 • The model employed a cost-effectiveness threshold of £20,000 per QALY gained.
 - 23 • The model was peer-reviewed by another health economist at the NCGC.
- 24

25 **20.5.2 Background**

26 There are fundamentally two reasons why a patient with a diagnosed hip fracture is delayed in
27 receiving surgery. First, the patient may be considered to be *unfit* for surgery for medical reasons,
28 and therefore made to wait until the medical team optimises her status. Alternatively, a patient
29 may be deemed to be fit for surgery at the time of admission, but will still incur delays linked with
30 *administrative reasons*, such as lack of space on theatre lists and/or problems with theatre,
31 surgical and anaesthetic staff cover.

32 In our economic analysis, we focus exclusively on the *administrative reasons* for surgical delay.

33 This is because, albeit all studies in the clinical review were initially considered for inclusion in the
34 economic model, the GDG concluded that only the subgroup of papers with a population that
35 excluded patients unfit for surgery was appropriate for basing the economic model upon.

36 In particular, the GDG considered that by removing patients unfit for surgery (defined as those for
37 whom: ‘*any medical reason when orthopaedic or anaesthetic staff felt that operation should be*
38 *delayed in order to improve the patient’s fitness for surgery*³⁰⁸) from our model, we would be
39 excluding confounding factors from the decision model, thus allowing more confidence in the
40 cost-effectiveness findings.

1 Those studies that had not excluded patients unfit for surgery from their population would
2 potentially have an imbalance in baseline characteristics which could result in skewing the data in
3 favour of the early surgery group. Even though these studies had used logistic regression to adjust
4 for confounding factors (such as ASA score, sex, age and comorbidities like cardiac problems), the
5 GDG still felt that the subgroup of papers that excluded patients unfit for surgery were more
6 robust.

7 Overall, three studies which excluded patients unfit for surgery from their population were
8 included in our clinical review: Moran (2005), Siegmeth (2005) and Orosz (2004)^{215,250,308}. Of these,
9 only Siegmeth³⁰⁸ reports data regarding whether patients returned to their original place of
10 residence or whether they changed residence (at 1 year follow up) and this was considered
11 essential information for modelling the different health states in our analysis.

12 Siegmeth (2005)³⁰⁸ excluded patients who were delayed for any medical reason when orthopaedic
13 or anaesthetic staff felt that operation should have been delayed in order to improve the patient's
14 fitness for surgery. Reasons for delays included anaemia requiring transfusion, correction of
15 electrolyte imbalance, uncontrolled diabetes and untreated heart failure. The GDG agreed that
16 the study adopted a set of diagnostically objective criteria in deciding which patients were
17 considered fit for surgery, and that no selection bias had been introduced in this process.

18 Furthermore, Siegmeth³⁰⁸ is a study set in the UK, and as such was considered to be more
19 applicable to our question than studies set in different countries. As the paper interprets "early
20 surgery" as surgery that took place within 48 hours from admission, we adopt this specific cut-off
21 point in our model.

22 **20.5.3 Population and time horizon**

23 The population for the cost-effectiveness analysis consists of hip fracture patients (male and
24 female) hospitalised for surgery and considered to be fit for surgery. The model spans over a life-
25 time horizon.

26 **20.5.4 Software**

27 The cost-effectiveness analyses were conducted using TreeAge Pro 2008.

28 **20.5.5 Methods**

29 We built a decision tree with Markov states where the expected costs and effectiveness of two
30 alternatives are evaluated and compared: "investment for early surgery" vs. "no investment for
31 early surgery". As discussed in section 20.5.8, this investment consists of the addition of extra
32 operating lists to the existing weekly number of theatre lists.

33 As mentioned in section 20.5.2, the health states of the model reflect the outcomes of Siegmeth
34 (2005)³⁰⁸: at one year after surgery, patients can be "living in their own home", "living in a
35 residential home", "living in a nursing home", or "dead".

36 Since patients were followed at 1 year from surgery in Siegmeth (2005)³⁰⁸, the *cycle length* of the
37 Markov model is supposed to last one year. At the end of each cycle, patients can either stay in
38 the same health state or can transit to the "dead" state (the "absorbing" health state in the
39 model). This is because no data were available from Siegmeth (2005)³⁰⁸ over the possible
40 transitions of patients between the other health state ("living in own home", "living in residential
41 home" or "living in nursing home"). Hence, we assume that patients' place of residence at 1 year
42 stays the same for the rest of their lifetime. Although this is obviously a simplification, it is unlikely

1 that the impact of the intervention (“investment for early surgery”) will have an effect after 1 year
2 from surgery.

3 The model starts with a simple decision node, which represent the decision to invest or not in
4 providing extra operating theatre lists. Following the investment, surgery takes place. However,
5 whether surgery will indeed take place “early” (within 48 hours from admission) or “late” is an
6 *uncertain* event. As a consequence, in our decision model we are able to address the question of
7 whether it is cost-effective to invest in extra operating lists (and therefore in extra personnel and
8 all the required resources) in order to increase the *probability* that those patients deemed “fit for
9 surgery” at admission are indeed operated within a certain time target. The probabilities of a
10 patient being in one of the four possible health states in the first cycle depend on whether they
11 have been operated within 48 hours or after 48 hours.

12 **20.5.6 Treatment effects**

13 The proportion of patients in each health state depends on the effectiveness of the treatment
14 (that is, of investment for early surgery), and on the proportion of patients still alive, which falls as
15 the number of cycles and therefore age increases.

16 Primary data were obtained from a GDG expert advisor regarding the proportion of patients in
17 each health state at 1 year follow up. These data (reported in Table 75 below) have been
18 extracted from the same database used in the Siegmeth³⁰⁸ study included in our clinical review,
19 and therefore refer to patients who were delayed for surgery not for medical reason but only for
20 administrative reasons.

21




Figure 157: Decision tree with Markov states - investment for early surgery vs. no hospital investment for early surgery

1 **Table 75: Place of residence and mortality at 1 year**

	Patients who had surgery ≤ 48 hours	Patients who had surgery > 48 hours	RR (surgery ≤ 48 hours vs. surgery > 48 hours)
Total number of admissions	3445 (0.952%)	175 (0.048%)	
No. patients living in own home at 1 year	1734 (0.503%)	76 (0.434%)	1.16
No. patients living in residential home at 1 year	489 (0.142%)	22 (0.126%)	1.13
No. patients living in nursing home at 1 year	307 (0.089%)	16 (0.091%)	0.97
No. patients dead at 1 year	915 (0.266%)	61 (0.349%)	0.76

2

3 It is important to point out that, for the first cycle in our model, the mortality data are based on
4 the information obtained from the database reported in Table 75.

5 For the long-term mortality, we considered a mean age of 81 for our cohort of patient, as this was
6 the mean age of patients in Siegmeth³⁰⁸. Following Parker(1992)²⁶⁸, the life expectancy after the
7 first cycle was assumed to be the same as that of the general population, and was obtained from
8 the Life Tables for the general population of England and Wales in the year 2005-2007 from the
9 Government Actuary Department:

10 (<http://www.gad.gov.uk/Documents/Demography/EOL/ILT%202005-07/wltewm0507.xls>).

11

12 This value was then adjusted for the ratio male/female corresponding to the patients
13 characteristics in the study as follows:

14 $\text{Total LE} = \text{LE}_{\text{female}} * \% \text{female} + \text{LE}_{\text{male}} * \% \text{male}$

15 **20.5.7 Quality of life**

16 The EQ-5D utility weights for patients living in their own home, in a residential or nursing home
17 used in our model are based on the findings of the paper by Tidermark (2002)³²⁸ and are
18 summarised in Table 76 below.

19 **Table 76: EQ-5D scores for health states**

Health state	Utility score
Living in own home (at 1 year from the fracture)	0.64
Living in own home (after 1 year from the fracture)	0.56
Living in an institution	0.35

1 (a) Source: Tidermark (2002)

2 We have assumed that patients living in their own home correspond to those “living
3 independently” in Tidermark (2002)³²⁸.

4 For each strategy, the expected QALYs in each cycle are calculated as follows:

5
$$\text{Expected QALYs} = \sum (U_i \times P_i)$$

6 Where:

7 U_i = the utility score for health state i

8 P_i = the proportion of patients in health state i

9 and where health state i could be any of the health states reported in table 1.

10 The overall *lifetime expected QALYs* are given by the sum of QALYs calculated for each cycle. The
11 *incremental QALYs gained* associated with a treatment strategy (“investment for early surgery” in
12 our case) are calculated as the difference between the expected QALYs with that strategy and the
13 expected QALYs with the comparator (that is, “no investment for early surgery”).

14 **20.5.8 Cost analysis**

15 **20.5.8.1 Early surgery implementation costs**

16 The “investments for early surgery” in our model consists of adding extra operating lists aimed at
17 increasing the theatre capacity as a way of reducing the time hip fracture patients have to wait
18 before they receive surgery. The evidence for this strategy refers to hospital records supplied by a
19 GDG member. In 2008, the John Radcliffe hospital in Oxford implemented a policy aimed at
20 increasing the number of patients operated with 48 hours from admissions. This was achieved by
21 adding an extra five half-day operating lists to the weekly number of lists. All the extra lists were
22 added during a normal working week, not during the weekend. Each extra theatre list consisted of
23 four hours of operating time. Table 77 below describes the extra personnel that had to be
24 employed to run these extra lists and the associated costs incurred by the hospital.

25 **Table 77: Personnel costs for extra operating lists**

Categories of personnel	Hours per additional list	Cost of hourly wage (source: PSSRU 2009)	Additional personnel costs for the 5 extra lists
Consultant surgeon	4	£108	£2,160
Consultant anaesthetist	4	£108	£2,160
Orthogeriatrician	1	£108	£540
Scrub nurse (as senior staff nurse)	4	£27	£540
Unscrub nurse (runner)	4	£21	£420
Radiographer	4	£25	£500
Anaesthesia assistant [ODP] (as senior staff nurse)	4	£27	£540

Recovery nurse (as staff nurse)	4	£21	£420
Total personnel cost for 5 additional weekly lists			£7,280
Total personnel costs for 5 additional lists over 1 year			£378,560

1

2 In addition to the extra personnel costs, we have to consider the *overhead costs* involved with
 3 running the operating theatre for the extra five half-day lists. These costs have been estimated on
 4 the basis of hospital records obtained from the Peterborough and Stamford District Hospital, and
 5 are summarised in the Table 78 below.

6

7 **Table 78: Overhead costs for the additional operating lists**

Resource	Cost per minute (£)
Energy	0.18
Premises maintenance	0.09
Staff uniforms and clothing	0.01
Medical and surgical equipment (including instruments)	0.82
Dressings	0.06
Total overhead costs per minute	£1.16
Total overhead costs for 5 additional weekly lists	£1,392
Total overhead costs for 5 additional over 1 year	£72,384

8

9 It follows that the overall total implementation cost for early surgery amounts to £450,944.

10 ***Probability of early surgery after hospital investment***

11 The following table summarises the number of patients operated within 48 hours from
 12 admissions before the extra operating lists were added (i.e. at baseline, year 2007-08) and for the
 13 years following the investment in extra operating lists. These data are also based on hospital
 14 records supplied by the John Radcliffe Hospital in Oxford.

15 **Table 79: Patients operated within and after 48 hours from admission - before and after** 16 **investments in extra operating lists**

	2007-8 (baseline)	2008-9 (intervention)	2009-10	2010-2011*
Total cases operated during the year	431	434	441	123
Number of patients fit for surgery within 48 hours during the year	363	347	374	114

Number of patients delayed over 48hrs because unfit for surgery	68	87	67	9
Number of patients fit for surgery and operated within 48 hrs (%)	192 (52.89%)	233 (67.15%)	316 (84.49%)	109 (95.61%)
Number of patients fit for surgery but delayed >48hrs	171 (47.11%)	114 (32.85%)	58 (15.51%)	5 (4.39%)

1 *data collected up to July 2010.

2 As Table 79 shows, the addition of the extra operating lists affected the *probability* that patients
3 fit for surgery are operated “early” (in our case, within 48 hours from admission). However, even
4 following this investment, early surgery is still a *random* event which is affected by many other
5 factors beyond the number of operating sessions available. Still, the data in Table 79 shows that
6 there is a clear trend in the increase in the number of patients fit for surgery that are operated
7 within 48 hours. There are several possible reasons for this trend, but they can mostly be seen as
8 the result of a learning process (by all the health care professionals involved in the care of the
9 patients) that produced positive spillover effects and efficiency gains in the years following the
10 implementation of the extra operating lists.

11 We use the data for 2008-09 as our intervention in the base case analysis. Data referring to other
12 years (2009-10 and 2010-11) are used in a sensitivity analysis.

13 ***Incremental cost per patient of implementation costs for extra theatre lists***

14 The extra cost per patient of implementing an early surgery strategy for the first year following
15 the investment (that is, for 2008-09) correspond to £450,944/434 = £1039.04 (where 434 is the
16 total number of patients operated for hip fracture – whether within or after 48 hours from
17 admission – in the intervention year).

18 ***20.5.8.2 Costing hospital length of stay***

19 In addition to the costs linked with the extra operating lists, we have consider the costs for the
20 length of hospital stay. We assume that the daily cost of a hospital bed in an orthopaedic ward
21 corresponds to £241.69 (which is obtained from a weighted average of the costs of the excess bed
22 days for hip all hip fracture procedures (major, intermediate and minor) with all types of
23 complications). This cost is then multiplied by the length of stay for each group of patients,
24 summarised in Table 80 below and based on the findings of ³⁰⁸

25 **Table 80: Mean length of hospital stay**

	Surgery ≤ 48 hours	Surgery > 48 hours	CI
Mean hospital stay in days (95% CI)	21.6	36.5	(5.7 – 16) p<0.0001

26

27 ***20.5.8.3 Health state costs***

28 **Health and social care costs for patients in the “living at own home” health state**

1 We acknowledge that even if a patient is discharged to his own home and returns to an
 2 independent living status, he will still incur in a higher level of use of health and domiciliary social
 3 care compared to his pre-fracture status, as it is unlikely that he will completely regain his pre-
 4 fracture level of independence. The PSSRU (2009)⁶¹ describes five possible “community care
 5 packages” for individuals who live in their own home and consume a level of health and
 6 domiciliary social care resources that varies according to their specific level of independence in
 7 functional status. For our model, we assume that the health and domiciliary social care costs for
 8 the patients in the “living in their own home” health state is an average of the cost of the “very
 9 low”, “low” and “medium” community care packages stated in the report. It follows that the
 10 weekly average health care costs for patients living in their own home after the fracture amounts
 11 to £9.9, and the weekly domiciliary social care costs to £98.1. While the health care costs are fully
 12 funded by the NHS, the domiciliary social care costs will only be partially met by the local
 13 authority. We found no published evidence regarding a national average of the percentage of
 14 domiciliary social care funded by local authorities^{71, 348, 72, 144}. In our base case analysis, we
 15 assume that 60% of these costs would be funded by the local authorities, and then test this
 16 assumption in a sensitivity analysis.

17 **Health and social care costs for patients in the “living in residential home” and “living in nursing** 18 **home” health states**

19 For patients living in a residential or in a nursing home, we need to consider the cost of long term
 20 care. This is estimated from the unit cost of stay in private nursing homes and in private
 21 residential care reported in the PSSRU 2009. The health care costs and fees per permanent
 22 residential week are described in Table 81.

23 **Table 81: Weekly health and social care costs for patients living in residential or nursing homes**

Place of residence	Weekly health care costs	Weekly fees
Private nursing home	£30.80, of which: <ul style="list-style-type: none"> • £30 (GP weekly home visit) • £0.80 (community nursing) 	£ 678
Private residential care	£26.3, of which: <ul style="list-style-type: none"> • £19.30 (GP weekly home visit) • £7.00 (community nursing) 	£467

24

25 Once again, while the NHS fully funds the health care costs, it does not pay towards long-term
 26 care for all patients. Moreover, only a proportion of the weekly fees will be met by the local
 27 authorities. We found no published evidence regarding a national average of the percentage of
 28 long-term care costs funded by local authorities, and as a consequence we assumed that the
 29 proportion of the costs of long-term care borne by the NHS and PSS is equal to 60% in the base
 30 case analysis, and changed it afterwards in a sensitivity analysis.

31 **20.5.9 Cost-effectiveness analysis**

32 Table 82 below summarised the findings of the cost-effectiveness analysis for the determinist
 33 case. We found that, for the first year following the investment in extra operating lists, the

1 strategy “investment for early surgery” is not cost-effective at a willingness to pay of £20k per
2 QALYs gained.

3 **Table 82: Cost-effectiveness results - deterministic analysis – first year following investment in**
4 **extra lists**

Strategy	Cost	Incremental Cost	Effectiveness	Incremental Effectiveness	Incremental cost-effectiveness (ICER)
No hospital investment for early surgery	£46.4K		2.32		
Hospital investment for early surgery (with probability of early surgery =67.15%)	£47.4K	£1.0K	2.3622	0.0421	£/QALY 22776

5

6 Table 83: Costs breakdown for "investment" and "no investment" in early surgery reports a
7 breakdown of all the cost categories included in the model for the first year in which the extra
8 operating lists were introduced.

9 **Table 83: Costs breakdown for "investment" and "no investment" in early surgery**

Resource item	Investment in extra operating lists	No investment in extra operating lists
Rehab cost	NA	NA
Hospital-related costs (for length of stay and investment in extra operating lists)	7442	6917
Readmission	NA	NA
Community health care (own home)	1664	1630
Community social care (own home)	9892	9690
Community health care (residential and nursing home)	2224	2206
Community social care (residential and nursing home)	26200	26000
Total cost	£47422	£46443

10

11 In order to ascertain how robust the findings of Table 82 are, we ran a series of sensitivity
12 analyses. Deterministic sensitivity analysis showed that the findings of our model are not sensitive
13 to the hospital bed day cost. However, threshold sensitivity analyses found that “investing for
14 early surgery” is the strategy with the highest net benefit in correspondence to a range of values
15 for different variables of the model, as summarised in Table 84 below.

16 **Table 84: Threshold sensitivity analyses**

Variable	Threshold values	Strategy with highest net benefit
Probability of being operated within 48 hours when investing for early surgery	>0.68	Investment for early surgery
Probability of living at home at 1 year for early surgery	>0.53	Investment for early surgery
Probability of living in nursing	<0.10	Investment for early surgery

home at 1 year – early surgery		
Probability of living in residential home at 1 year – early surgery	<0.15	Investment for early surgery
Mean length of hospital stay for early surgery patients	<18.47 days	Investment for early surgery
Number of extra operating lists (of 4 hours each)	>4.38	No investment for early surgery
Proportion of social care costs paid by the NHS and local authorities	>0.43	No investment for early surgery
Cost per day in hospital	>£292.50	Investment in early surgery

1

2 20.5.9.1 Probabilistic sensitivity analysis

3 A probabilistic sensitivity analysis was performed to assess the robustness of the model results to
 4 plausible variations in the model parameters. Probability distributions were assigned to each
 5 model parameter, where there was some measure of parameter variability. We then re-calculated
 6 the main results 10000 times, and each time all the model parameters were set simultaneously,
 7 selecting from the respective parameter distribution at random. Table 85 summarises the type
 8 and properties of distributions used in the probabilistic sensitivity analysis.

9

10 **Table 85: Description of the type and properties of distributions used in the probabilistic**
 11 **sensitivity analysis**

Parameter	Type of distribution	Properties of distribution
Baseline risk	Beta	Bounded on 0 – 1 interval. Derived from sample size, number of patients experiencing events
Cost	Gamma	Bounded at 0, positively skewed. Derived from mean and standard error
Utility	Beta	Bounded on 0 – 1 interval. Derived from mean and sample size
Risk ratio	Lognormal	Bounded at 0. Derived from log (RR) and standard error of log (RR)

12

13 Table 86 reports the distribution, parameters and expected values for each variable of the model.

14 **Table 86: Distributions, parameters and expected values for probabilistic sensitivity analysis**

Name	Baseline value (deterministic analysis)	Distributions and parameters	Expected value
------	-----------------------------------------	------------------------------	----------------

EQ- 5D “living own home”	0.64	Beta, Real-numbered parameters, alpha = 37.12, beta = 20.88	0.64
EQ - 5D – “living in nursing home” and “living in residential home”	0.35	Beta, Real-numbered parameters, alpha = 2.45, beta = 4.55	0.35
EQ- 5D “living in own home” after 1 year	0.56	Beta, Real-numbered parameters, alpha = 31.92, beta = 25.08	0.56
Cost per hour – consultant (surgeon and anaesthetist)	108	Gamma, alpha = 15.36583528, lambda = 0.142276253	108
Cost per hour (staff nurse)	21	Gamma, alpha = 15.36583528, lambda = 0.731706442	21
Cost per hour - ODP	27	Gamma, alpha = 15.36583528, lambda = 0.56910501	27
Cost per hour -radiographer	25	Gamma, alpha = 15.36583528, lambda = 0.614633411	25
Cost per hour – senior nurse	27	Gamma, alpha = 15.36583528, lambda = 0.56910501	27
Operating time per each extra list (hours)	4	Triangular, Min = 1, Likeliest = 4, Max = 7	4
Initial age	81	None	
Length of hospital stay – early surgery	21.6	Log-Normal, u (mean of logs) = 3.038030773, sigma (std dev of logs) = 0.2632965680	21.6
Length of hospital stay – late surgery	36.5	Log-Normal, u (mean of logs) = 3.562649719, sigma (std dev of logs) = 0.263296568	36.5
No of patients operated in the intervention year (2008-09)	434	Poisson, lambda = 434	434
No of weekly extra operating lists added	5	Triangular, Min = 3, Likeliest = 5, Max = 7	5
Overhead cost per minute	1.16	Gamma, alpha = 15.36583528, lambda = 13.24640973	1.16
Probability of surgery within 48 hours without investments in extra lists	0.5289	Beta, Integer parameters only, n = 363, r = 192	0.5289
Probability of surgery within 48 hours after investments in extra	0.6715	Beta, Integer parameters only, n = 347, r = 233	0.6715

lists			
Proportion of social care costs borne by local authorities	0.6	Triangular, Min = 0.20, Likeliest = 0.60, Max = 1; Expected value: 0.6	0.6
Probability of dead – late surgery	0.349	Beta, Integer parameters only, n = 175, r = 61	0.349
Probability of living in own home – late surgery	0.434	Beta, Integer parameters only, n = 175, r = 76	0.434
Probability of living in nursing home – late surgery	0.092	Beta, Integer parameters only, n = 175, r = 16	0.092
Probability of living in residential home – late surgery	0.125714	Beta, Integer parameters only, n = 175, r = 22	0.125714286
Relative risk of living in nursing home	0.97	Log-Normal, u (mean of logs) = -0.060565609, sigma (std dev of logs) = 0.24538297	0.97
Relative risk of living in own home	1.16	Log-Normal, u (mean of logs) = 0.144607796, sigma (std dev of logs) = 0.08731791	1.16
Relative risk of living in residential home	1.13	Log-Normal, u (mean of logs) = 0.101743909, sigma (std dev of logs) = 0.202354755	1.13
Relative risk mortality	0.76	Log-Normal, u (mean of logs) = -0.280072176, sigma (std dev of logs) = 0.106163367	0.76
Weekly health care costs for patients living in a nursing home	30.8	Gamma, alpha = 15.36583528, lambda = 0.498890756	30.8
Weekly health care costs for patients living in their own home	9.9	Gamma, alpha = 15.36583528, lambda = 1.552104574	9.9
Weekly health care costs for patients living in a retirement home	26.3	Gamma, alpha = 15.36583528, lambda = 0.584252292	26.3
Weekly social care costs for patients living in their own home	98.1	Gamma, alpha = 15.36583528, lambda = 0.156634407	98.1
Weekly social care costs for patients living in a residential home	467	Gamma, alpha = 15.36583528, lambda = 0.032903288	467
Weekly social care costs for patients living in a nursing home	678	Gamma, alpha = 15.36583528, lambda = 0.022663474	678
Daily cost of hospital stay	241.68	Gamma, alpha = 15.36583528,	241.68

lambda = 0.063579259

1

2 The conventional way to interpret a cost-effectiveness analysis is to look at the option that is
3 optimal based on mean results from the probabilistic sensitivity analysis. These findings are
4 summarised in Table 87 below:

5 **Table 87: Cost-effectiveness findings from probabilistic sensitivity analysis – first year following**
6 **investment in extra lists**


Strategy	Cost	Incremental Cost	Effectiveness	Incremental Effectiveness	Incremental C/E ratio (ICER)	95% CI
No hospital investment for early surgery	£46.4K		2.3212			
Hospital investment for early surgery (<48 hours)	£47.4K	£1.0K	2.3637	0.0425	£/QALY 22542	Cost saving - dominated

7

8 The PSA shows that there is a high uncertainty as to whether “investment for early surgery” is
9 cost-effective compared to “no investment for early surgery”. This uncertainty can be graphically
10 represented by plotting the results of the incremental analysis for all the 10,000 simulations into a
11 cost-effectiveness plane. Each point on the scatter plot represents the ICER of investment for
12 early surgery versus no investment for early surgery for each simulation. The dotted line
13 represents the £20,000/QALY threshold while the ellipse delimits the 95% confidence interval.

14

ICE Scatterplot of Hospital investment for early surgery (<48 hours) vs. No hospital investment for early surgery



1

2 We found that the strategy of “investment in extra operating lists” was cost-effective in 50% of
 3 the simulations, both at a willingness to pay of £20,000 per QALY and of 30,000 per QALY.

4 **20.5.9.2 Scenario analysis: second year following implementation**

5 We now compare the non-investment strategy versus the investment strategy, where for the
 6 latter we use data referring to the second year following the introduction of the additional
 7 operating lists. The findings of the deterministic and of the probabilistic cost-effectiveness
 8 analysis are summarised in Table 88 and Table 89 below.

9 **Table 88: Cost-effectiveness results - deterministic analysis – second year following investment**
 10 **in extra lists**

Strategy	Cost	Incremental Cost	Effectiveness	Incremental effectiveness	Incremental Cost-effectiveness ratio (ICER)
No hospital investment for early surgery	£46.4K		2.32		
Hospital investment for early surgery (<48 hours) (with probability of early surgery from second year of investment=84.49% and with total number of patients operated in that year = 441)	£47.3K	£0.8K	2.413	0.093	£/QALY 9070

1
2 **Table 89: Cost-effectiveness findings from probabilistic sensitivity analysis – first year following**
3 **investment in extra lists**

Strategy	Cost	Incremental Cost	Effectiveness	Incremental effectiveness	Incremental Cost-effectiveness ratio (ICER)
No hospital investment for early surgery	£46.4K		2.321		
Hospital investment for early surgery (<48 hours) <i>(with probability of early surgery from second year of investment=84.49% and with total number of patients operated in that year = 441)</i>	£47.3K	£0.8K	2.415	0.094	£/QALY 8933

4
5 The strategy of introducing extra theatre list is therefore cost-effective from the second year of
6 implementing the change aimed at reducing the waiting time to surgery for hip fracture patients.
7

8 **20.5.10 Discussion**

9 Our analysis showed that adding extra operating lists as a way of undertaking surgery within 48
10 hours from admission is slightly above the threshold of 20K/QALYs in the first year of
11 implementation, but becomes clearly cost-effective from the second year onwards.

12 However, our cost-effectiveness estimates are likely to be conservative in that we did not look at
13 the impact of early surgery on the presence of complications. This was because no information on
14 complications was available from Siegmeth (2005)³⁰⁸, and the other studies from the clinical
15 review that did report data on complications could not be used since they did not exclude
16 patients unfit for surgery from their population.

17 As resources and treatment effects data are based on information received from two specific
18 hospital settings (John Radcliffe hospital in Oxford and the Peterborough and Stamford Hospital
19 Foundation Trust), our findings may not be generalised to the whole NHS. For example, for some
20 hospitals the addition of extra operating lists may not be feasible if no spare theatre capacity is
21 available for this purpose.

22 In non-linear models, such as Markov models, there is often a difference between the deterministic and
23 probabilistic results and in such cases the probabilistic results should take precedence. The findings of the
24 PSA reported in section 20.5.9.1 show that there is a high uncertainty as to whether “investment for early
25 surgery” is cost-effective compared to “no investment for early surgery”. If we consider a 95% confidence
26 interval the base case results did not reach statistical significance (as reported in table 85). Moreover, we
27 found that the strategy of “investment in extra operating lists” was cost-effective in only 50% of the
28 simulations, both at a willingness to pay of £20,000 per QALY and of 30,000 per QALY.

29 A possible extension of the model could look at the possibility of introducing extra operating lists
30 during the weekend, which would be more expensive than weekdays, as personnel would have to
31 be paid up to a time and a third more in salary (BMA contract 2003). Patients admitted at
32 weekends or public holidays tend to do worse (Foss 2006)⁹⁷). However, most large hospitals have
33 trauma lists at the weekend, with planned trauma lists built into job plans. The reason why extra
34 lists were introduced during weekdays in the model that we have developed is because it was
35 acknowledged that there are more competing patients for planned trauma lists in those days, for

- 1 example patients requiring specialist reconstructions such as pelvic fractures or complex joint
- 2 injuries.
- 3

1

2 **20.6 Cost-effectiveness analysis of Hospital MDR vs Usual care**3 **20.6.1 Introduction**

4 The GDG identified as a high priority area for economic analysis the multidisciplinary management
5 in hospital for hip fracture patients.

6 In the economic plan, the clinical question (number 13) linked to this high priority area is the
7 following:

8 *“What is the clinical effectiveness and cost-effectiveness of the following hospital-based
9 multidisciplinary rehabilitation programmes:*

- 10 • *Hip Fracture Programme (HFP),*
- 11 • *Geriatric Orthopaedic Rehabilitation Unit (GORU), and*
- 12 • *Mixed Assessment and Rehabilitation Unit (MARU)*

13 *versus each other and versus usual inpatient rehabilitation for hip fracture patients?”*

14 The GDG felt that there were sufficient similarities between the GORU and MARU rehabilitation
15 programmes, and therefore decided to group the evidence for these interventions under the
16 same category of “GORU/MARU”. A detailed discussion of the main characteristics of each
17 rehabilitation programme is presented in Chapter 12 of this Guideline, especially in sections 12.1
18 and 12.2.

19 A review of the literature was conducted. The literature search and review methods can be found
20 in section 3. Despite some cost-effectiveness studies were identified, none represented a full cost-
21 utility analysis which addressed our clinical question. As a consequence, the GDG felt that an
22 original economic model of the listed interventions was essential in order to inform their
23 recommendations.

24 The following general principles were adhered to:

- 25 • The GDG was consulted during the construction and interpretation of the model.
- 26 • When published data was not available we used expert opinion to populate the
27 model.
- 28 • Model assumptions were reported fully and transparently.
- 29 • The results were subject to sensitivity analysis and limitations were discussed.
- 30 • We followed the methods of the NICE reference case. Therefore costs were
31 calculated from a NHS and personal social services perspective. Health gain was
32 measured in terms of quality-adjusted life-years (QALYs) gained. Both future costs
33 and QALYs were discounted at 3.5%.
- 34 • The model employed a cost-effectiveness threshold of £20,000 per QALY gained.

- 1 • The model was peer-reviewed by another health economist at the NCGC.

2

3 **20.6.2 Population and time horizon**

4 The population for the cost-effectiveness analysis consists of hip fracture patients (male and
5 female) hospitalised for surgery. The model spans over a life-time horizon.

6 **20.6.3 Software**

7 The cost-effectiveness analyses were conducted using TreeAge Pro 2008.


8 **20.6.4 Structure of the model**

9 **20.6.4.1 Model cycles at time 0**

10 We develop a Markov model with a cycle length of 3 months. Thus, all events are calculated on a
11 3 month basis at the end of which patients are in one of the possible health states. As the time
12 horizon in our model is lifetime, these cycles will keep repeating for the duration of the life
13 expectancy of the population in the studies.

14 The specific health states of our Markov model have been determined on the basis of the findings
15 of the clinical review. During cycle 0 the health states are determined by the *types of*
16 *complications* experienced while in hospital (and while undergoing their rehabilitation
17 programme). Using evidence from the clinical review, we assume that during cycle 0, patients can
18 occupy one of the following health states: “not recovered and with no complications”, “not
19 recovered and with pressure sores”, “not recovered and with moderate delirium”, “not recovered
20 and with severe delirium”, and “dead”.

21 This is a graphic representation of cycle 0 of the Markov model:



1
2
3

Figure 158: Cycle 0 Markov model

4 The above diagram illustrates that throughout their hospital stay (and hence, while still
5 undergoing their rehabilitation programme) patients will be considered as “not recovered”. Some
6 of these “not recovered” patients will not develop any complications, but others will experience
7 delirium (moderate or severe), or pressure sores.

8 ***Evidence and treatment effects on complications – Cycle 0 of the Markov model***

9 The clinical review found evidence of complications only from RCTs of HFP vs usual care. The
10 following complications were identified:

11

Table 90: Types of complications identified in the clinical review

Type of complication as reported in the clinical review	Source
Pressure sores	Vidan (2005) ³⁴⁴
Heart failure	Vidan (2005) ³⁴⁴
Pneumonia	Vidan (2005) ³⁴⁴
Confusion	Vidan (2005) ³⁴⁴
Chest infection, cardiac problem, bedsore	Swanson (1998) ³²⁵
Stroke, emboli	Swanson (1998) ³²⁵
Delirium	Marcantonio (2001) ²⁰³

Severe delirium	Marcantonio (2001) ²⁰³
-----------------	-----------------------------------

1

2 The GDG decided to include the evidence on pressure sores from Vidan (2005)³⁴⁴ and on delirium
3 from Marcantonio (2001)²⁰³. This was because of the good quality of the evidence; the reliable
4 ascertainment of these complications, and their well recognised impact on costs of hospital stay.

5 The findings of Vidan (2005)³⁴⁴ on “confusion” were not considered in the economic model since
6 they were not statistically significant and because they did not distinguish between “moderate”
7 and “severe” confusion, so it was not possible to use these findings alongside those of
8 Marcantonio (2001)²⁰³ on delirium.

9 The evidence on complications from Swanson (1998)³²⁵ was not included in the economic model
10 since the paper only provided a composite figure for chest infections, cardiac problems and
11 bedsores and did not distinguish among the different types of complications. As a consequence, it
12 was not possible to determine the loss in health-related Quality of Life (QoL) due to each
13 complication and the associated costs.

14 The evidence on pneumonia (Vidan 2005)³⁴⁴ was also not included in the economic model,
15 because it showed no difference between the intervention and control group.

16 The GDG decided to exclude the remaining complications (heart failure, and stroke) due to the
17 weaker evidence of effectiveness in prevention and the unreliable ascertainment of the
18 conditions. In particular, it was pointed out that ‘heart failure’ is very difficult to define and
19 diagnose clinically, and that ‘stroke’ is a whole series of different conditions with hugely differing
20 origins and outcomes. It should also be noted that it is unlikely that we have introduced a bias in
21 our model because of the exclusion of these specific outcomes. In fact, despite the clinical review
22 reported that the relative risk for heart failure and stroke was large and in favour of usual care, it
23 was also true that they had wide confidence intervals, which meant that the difference was not
24 statistically significant. Moreover, the GDG agreed that the lower event numbers associated with
25 usual care was due to the fact that people had been less intensively monitored compared to the
26 intervention arms of the studies, so that some events may have been missed in the control arm.

27 As a consequence, the model only looked at the following complications: pressure sores (from
28 Vidan 2005)³⁴⁴, moderate delirium and severe delirium (Marcantonio 2001)²⁰³.

29 The clinical review did not find evidence of complications for GORU/MARU vs usual care. The GDG
30 decided to consider the sample complications from the HFP (pressure sores, moderate and severe
31 delirium) and assume that there was no difference between the intervention and usual care (and
32 hence to consider a RR equal to 1). This assumption was subject to a sensitivity analysis. Table 19
33 below reports the transition probabilities for cycle 0 of the Markov model.

34 **Table 91: Transition probabilities - cycle 0 of the Markov model**

Transition Probability	Usual care	HFP	GORU
Probability moderate delirium*	22.0%	20.9% (RR 0.95)	22% (RR 1.00)
Probability severe delirium*	28.12%	11.25% (RR 0.4)	28.12% (RR 1.00)

Probability pressure sores**	16.46%	5.10 % (RR 0.31)	16.46% (RR 1.00)
------------------------------	--------	------------------	------------------

1 *= source: Marcantonio (2001)²⁰³.

2 **= source: Vidan (2005)³⁴⁴

3


4

20.6.4.2 Cycles 1 – onwards

5 As for the health states for cycle 1 – onwards, we again used the findings of the clinical review and
6 assume that, after their hospital discharge (and therefore, after their hospital-based MDR or their
7 usual care has been completed), patients can transit between the following health states:
8 “recovered”, “not recovered”, and “dead”.

9 Vidan (2005)³⁴⁴, Stenvall (2007)³²⁰ and Shyu (2008)³⁰⁵ report findings regarding the effectiveness
10 of hospital MDR programmes versus usual care to help patients recover their pre-fracture
11 Activities of Daily Living (ADL) levels. The “recovered” health state in our model refers therefore
12 to the case in which patients have gone back to their pre-fracture ADL levels.

13 This is a graphic representation of cycles 1 to 3 of Markov model, following hospital discharge:



14

15


16

Figure 159: Cycles 1 to 3 of the Markov model

17 The above diagram illustrates that, up until 12 months, patients who are in the “recovered”
18 health state can stay in the same state in the following cycles, or can transit to the “dead”
19 health state.

20 However, patients in the “not recovered” health state can stay in the same state at the end of
21 each cycle, or transit to the “recovered” or “dead” states. This is because, from the clinical review,
22 we only have data regarding the transition of patients from the “not recovered” to the

- 1 “recovered” health state, and these data are only available up until 12 months follow up period.
 2 No clinical data are available regarding the possible transition of the “recovered” patients to the
 3 “not recovered health state”.
- 4 From 12 months onwards, we assume that patients will no longer transit from the “not
 5 recovered” to the “recovered” health state, and that patients can only remain in the state they
 6 are in or transit to the “dead” state. This is because no clinical data are available from the clinical
 7 review after that point. Hence, the relevant transitions between health states after 12 months will
 8 be:



9
 10 **Figure 160: Cycle 4 - onwards of the Markov model**
 11

- 12 That is, from cycle 4 onwards, patients who are in the “recovered” health state will stay in that
 13 state or transit to the “dead” state. Similarly, patients in the “not recovered” health state will
 14 remain in that state or transit to the “dead” state. The GDG noted that the assumption that
 15 people remain in the same health state from 12 months onwards is clinically reasonable, as from
 16 that time patients’ health state will no longer be influenced by their hip fracture. All possible
 17 events after this time (e.g. death, falls, needs for care home etc) will take place at rates that are
 18 consistent and in line with those of the general population and that therefore will no longer be a
 19 consequence of the hip fracture nor of the specific rehabilitation programme received.

- 20 Whether they are “recovered” or “not recovered”, the place of residence at hospital discharge for
 21 patients will also be affected by whether they received usual care, HFP or GORU/MARU as a form
 22 of rehabilitation programme. This circumstance is represented in Figure 150 below:

23




Figure 161: Place of residence at discharge

1
2
3

4 No evidence is available from the clinical review regarding whether patients discharged to their
5 own home would then transit to the “living in long term care” setting in subsequent cycles of the
6 model, and vice versa. Hence, we make the assumption that patients will keep living in the same
7 place of residence they had when they were discharged from hospital, and that they can only
8 transit to the “dead” state in the following cycles.

9 **Evidence and treatment effects on recovery of ADL levels and on place of residence at discharge**

10 Table 92 reports the levels of the transition probabilities used in the model

11
12

Table 92: Transition probability of Not Recovery of ADL pre-fracture levels

Transition probability of Not Recovery of ADL pre-fracture levels	Usual care	HFP	GORU/MARU
At 3 months ⁽¹⁾	0.73	0.5767 (RR=0.79)	0.5694 (RR=0.78)
At 6 months ⁽¹⁾	0.67	0.5293 (RR=0.79)	0.5226 (RR=0.78)
At 9 months ⁽²⁾	0.63	0.4977 (RR=0.79)	0.4914 (RR=0.78)
At 12 months ⁽³⁾	0.59	0.4661 (RR=0.79)	0.4602 (RR=0.78)

13

⁽¹⁾Data at 3 and 6 months from Vidan³⁴⁴

14
15

⁽²⁾Data at 9 months obtained with a linear extrapolation from the transition probabilities in Vidan³⁴⁴

16

⁽³⁾Data at 12 months pooled from Vidan, Shyu and Stenvall^{305,320,344}

17

1 As for the place of residence following hospital discharge, we use the following treatment effects
2 in our model:

	Usual care	HFP	GORU/MARU
Probability of returning to own home*	0.71	0.8094 (RR=1.14)	0.7881 (RR=1.11)

3 *source: NCGC meta-analysis of clinical trials
4

5 20.6.5 Evidence and treatment effects on mortality

6 In our model we distinguished two types of mortality: short-term mortality (within 12 months
7 from the start of the rehab programme) and long-term mortality (after 12 months).

8 SHORT-TERM MORTALITY

9 In order to take into account the difference in mortality due to the intervention, we used the data
10 from the RCTs included in our meta-analysis to estimate mortality. The data available from the
11 RCTs can be found in Table 21.

12 **Table 93: Proportion of patients dead at different time points**

	6 months	12 months
Usual care¹	16.73%	21.38%
HFP²	NA	17.32% (RR 0.81)
GORU/MARU²	13.22% (RR 0.79)	20.31% (RR 0.95)

13 1 Data pooled from the usual care arms of RCTs in the clinical review

14 2 RR calculated compared to usual care
15

16 Data were available for usual care and GORU at 6 and 12 months from randomisation. Only 12
17 month data were available for the HFP intervention.

18 When more than one time points was available (i.e. for the usual care and GORU/MARU arms),
19 the probability of dying was calculated from the data reported in Table 4 as follows:

$$20 \quad \text{Prob_die_y to x} = (\% \text{ dead time x} - \% \text{ dead time y}) / (1 - \% \text{ dead time y})$$

21 Where:

22 Prob_die_y to x is the probability of dying from time y to the following time x

23• “% dead time x” is the proportion of patients dead at time x

24• “% dead time y” is the proportion of patients dead at time y

25 To convert probabilities into a 3-month transition probability, which is the cycle length of the
26 model, we used the formula:

$$27 \quad 1 - \exp((\ln(1 - \text{Prob_die_y to x})) / ((x-y)/3))$$

28

1 Where x and y are the initial and final time points of the interval considered, exp(a)=exponential
2 of a; and ln(a)=natural log of a.

3 LONG-TERM MORTALITY

4 The mean age of the patients when entering the model was 81 as this was the mean age of
5 patients in the RCTs.

6 Life expectancy in people who were alive one year after a hip fracture was assumed to be the
7 same as the general population in England and Wales, as reported in a study (Parker1992, citing
8 Elmerson1988)²⁶⁸. The remaining life expectancy for the participants of the RCTs was obtained
9 from the Life Tables for the general population of England and Wales in the year 2005-2007 from
10 the Government Actuary Department
11 (<http://www.gad.gov.uk/Documents/Demography/EOL/ILT%202005-07/wltewm0507.xls>).

12 The value was adjusted for the ratio male/female corresponding to the patients characteristics in
13 the RCTs as follows:

14 Total LE = LE_{female} * %female + LE_{male} * %male

15

16 **20.6.6 Utilities data**

17 **20.6.6.1 Utilities for cycle 0 (0-3 months)**

18 Utilities indicate the preference for health states on a scale from 0 (death) to 1 (perfect health).
19 Quality of life values are attached to all health states.

20 Stage 0 of the model refers to the first three months of the Markov model. They capture the time
21 that the patients spend in hospital, during which they undergo a surgical treatment of the
22 fracture, following which the rehabilitation process starts.

23 The utility weights for the health states in cycle 0 are summarised in table 5.

24 **Table 94: Utility weights for cycle 0**

Table 5: Utility weights for cycle 0 Health state	Base case value	Source
"Not recovered, no complications"	0.314	ADL levels from Kennie (1988) ¹⁷⁶ ; EQ-5D scores from Tidermark (2002) ³²⁸
"Not recovered and with pressure sores"	0.19	Essex (2009) ⁸⁶
"Not recovered and with moderate delirium"	0.314	ADL levels from Kennie (1988) ¹⁷⁶ ; EQ-5D scores from Tidermark (2002) ³²⁸
"Not recovered and with severe delirium"	0.25	NICE clinical guideline on Delirium ²²⁴

25

1 We assume that the utility for the “not recovered, no complication” health state in the first three
 2 months is the same as that of the “Not recovered” health state after the hospital discharge (i.e.
 3 after the first cycle). The following paragraph explains how the utility for the “not recovered, no
 4 complication” health state is obtained.

5 The NICE guideline on Delirium²²⁴ reports utility weights for patients with moderate and severe
 6 delirium using the finding of Ekman (2007)⁷⁶ on patients with dementia. Ekman (2007)⁷⁶ estimates
 7 that the mean utility score for mild, moderate and severe dementia correspond to 0.62, 0.40 and
 8 0.25 respectively. As for pressure sores, Essex (2009)⁸⁶ reports an EQ-5D score of 0.19 for patients
 9 experiencing this complication. EQ-5D scores were obtained from a survey of a sample of 6
 10 patients with pressure ulcers.

11 We proceeded by selecting the *lowest* EQ-5D score between the “not recovered with no
 12 complication” health state and the EQ-5D linked with that particular complication (moderate
 13 delirium, severe delirium or pressure sores). Thus, being the utility for “moderate delirium” 0.4,
 14 and being this utility higher than the one of the “not recovered with no complication” health state
 15 (0.4 vs 0.314), we selected the latter also for the “not recovered and with moderate delirium”
 16 health state.

17 However, the utility score for patients with severe delirium identified in the literature was lower
 18 than then the score for the “not recovered, no complications” health state (0.25 vs 0.314).
 19 Similarly, the utility score for pressure sores identified in the literature was lower than the one of
 20 the “not recovered, no complications” health state (0.19 vs 0.314). Hence, we used the EQ-5D
 21 score for those specific complications (severe delirium, pressure sores) in our model.

22 **20.6.6.2 Utilities for cycles 1 – onwards (3 months – onwards)**

23 In order to assign an utility level to each of the health states for the model in cycles 1-onwards
 24 (that is, “recovered” and “not recovered”), we proceeded by using the RCT included in our clinical
 25 review by Kennie et al (1988)¹⁷⁶ which reports the number of patients (in the treatment and
 26 control group) classified according to their level of independence in activities of daily living *before*
 27 *admission* (i.e. before the hip fracture) and *at entry into study* (i.e. before the rehabilitation
 28 program has started). This information is summarised in tables 23 and 24 below.

29

30 **Table 95: ADL levels before admission for treatment and control group (source: Kennie et al**
 31 **1988)¹⁷⁶**

<i>Independence in activities of daily living before admission (Katz index)</i>	<i>Treatment group (n=54)</i>	<i>Control group (n=54)</i>
A	21	28
B	14	11
C	6	6
D	3	3
E	2	1
F	2	1
G	1	1
Not classified	5	3

1

2 **Table 96: ADL levels at entry into study for treatment and control group (source: Kennie et al**
 3 **1988)¹⁷⁶**

Independence in activities of daily living at entry into study (Katz index)	Treatment group (n=54)	Control group (n=54)
A	0	0
B	1	0
C	1	0
D	2	3
E	18	19
F	23	16
G	7	15
Not classified	2	1

4 Source: Kennie et al (1988)¹⁷⁶

5

6 We use the data for the “independence in ADL before admission” to calculate the proportion of
 7 independent and dependent patients that are in the “recovered” health state. Similarly, we use
 8 the information on ADL for patients at entry into study to calculate the proportion of independent
 9 and dependent patients that are in the “not recovered” health state.

10 As a consequence, we have:

- 11 • % of patients with A-B score in the “recovered” state:
 12 $(21+14)$ (from the treatment group) + $(28+11)$ (from the control group)/100 = 74%
- 13 • % of patients with A-B score in the “not recovered” state:
 14 $1/100 = 1\%$

15 Hence, in the “recovered” health state, 74% of patients have an ADL score of A-B, and 26% of
 16 patients in the same state have an ADL score of C-G. On the other hand, in the “not recovered”
 17 health state, only 1% of patients have ADL score of A-B, the rest having an ADL score of C-F.

18 For each of these two states we calculated the *composite utility*, that is the utility for the
 19 “independent” and for the “dependent” patients. Tidermark (2002)³²⁸ reports EQ-5D scores
 20 associated with ADL scores of A-B and C-F for hip fracture patients at 4 months after the fracture.
 21 These weights correspond to: 0.68 for ADLs of A-B, and to 0.31 for ADLs of C-F.

22 Using the proportion of patients who were reported as independent and as dependent before
 23 admission for the “recovered” health state we have:

24 $74\% * 0.68 = 0.053$

25 $26\% * 0.31 = 0.081$

26 Thus, the utility weight for “recovered” health state corresponds to 0.584

27 As for the “Not recovered” health state we have:

1

2 1% *0.68 = 0.0068

3 99%*0.31 = 0.307

4 Thus, the utility weight for “not recovered” health state is: 0.314. We summarise these findings in
5 table 8:

6

7 **Table 97: Utility weights for health states in cycles 1 -onwards**

Health state	EQ-5D	Source
“Recovered”	0.584	ADL levels from Kennie (1988) ¹⁷⁶ ; EQ-5D scores from Tidermark (2002) ³²⁸
“Not recovered”	0.314	ADL levels from Kennie (1988) ¹⁷⁶ ; EQ-5D scores from Tidermark (2002) ³²⁸

8

9 **20.6.7 Calculating QALYs gained**10 For each strategy (HFP, GORU/MARU and usual inpatient rehabilitation), the expected QALYs in
11 each cycle are calculated as follows:

12
$$\text{Expected QALYs} = \sum (U_i \times P_i)$$

13 where

14 U_i = the utility score for health state i 15 P_i = the proportion of patients in health state i 16 and where health state i could be any of the health states reported in the Figures 147 and 148.17 The proportion of patients in each health state depends on the effectiveness of the treatment,
18 and on the proportion of patients still alive, which falls as the number of cycles and therefore age
19 increases.20 The overall *lifetime expected QALYs* are given by the sum of QALYs calculated for each cycle. The
21 *incremental QALYs gained* associated with a treatment strategy are calculated as the difference
22 between the expected QALYs with that strategy and the expected QALYs with the comparator.23 **20.6.8 Cost data**24 **20.6.8.1 Cost data: cycle 0 (hospital stay)**25 During hospital stay, the costs will depend on the rehabilitation programme, the length of hospital
26 stay and health state related costs. We analyse each category in turn.

1 **Cost of the rehabilitation programme**

2 The NICE “Guide to the methods of technology appraisal” points out that national data based on
3 healthcare resource groups (HRGs), such as the Payment by Results tariff, are a valuable source of
4 information for resource use and costs and should be considered for use when they are
5 appropriate and available (“Guide to the methods of technology appraisal”, 2008, page 40).
6 However, data based on HRGs may not be appropriate in all circumstances, especially when the
7 definition of the HRG is broad or the mean cost probably does not reflect resource use in relation
8 to the interventions we are evaluating.

9 In our case, we would be using the HRG4 as the source to cost our rehab programmes. In the
10 document: “Casemix Service HRG4 - Guide to unbundling” it is pointed out that the HRG4 refers
11 to cases of **Discrete Rehab** services:

12 “[...] only discrete rehabilitation activity and costs should be reported using the rehabilitation HRG4
13 categories, for the reference costs collection.”

14 And the 2007 document on Collection Guidance on Reference Costs for 2006-07 specifies that:

15 “Rehabilitation HRGs are only generated where care is identified as taking place under a specialist
16 rehabilitation consultation or within a discrete rehabilitation ward or unit. [...] Where a patient is
17 not admitted specifically to a rehabilitation unit or where rehabilitation treatment is undertaken
18 without transfer to a specialist consultant, or without transfer to a rehabilitation unit, this should
19 not be reported as discrete rehabilitation”.

20 It would therefore seem that whilst this definition could apply to the GORU/MARU model (where
21 a patient is discharged from the orthopaedic unit and admitted to a separate geriatric
22 orthopaedic unit to receive the rehabilitation), it could not reflect the case of a HFP, where a
23 patient is not usually discharged to the care of a specialist rehabilitation consultant.

24 Thus, whilst we could use the HRG4 to cost a GORU and a MARU programme, we would not be
25 able to use it to cost a HFP.

26 As a consequence, the GDG decided to evaluate the cost of the different rehabilitation
27 programmes using the level of resources specified in the different RCTs included in the clinical
28 review. When necessary, such levels have been adjusted by expert opinion to reflect a pattern of
29 care closer to the UK health care setting (see below).

30 The resources used in the different RCTs have been reported as *incremental* resources used with
31 respect to the usual care arm of the study. Using information on unit costs for NHS personnel
32 provided by the PSSRU 2009, we were then able to estimate the *incremental cost* of both HFP and
33 GORU/MARU with respect to usual care.

34 Moreover, it is important to note that the level of resources used in the two hospital-based MDR
35 programmes are calculated in such a way to reflect the length of hospital stay of the patients in
36 our model. Thus, we use the length of stay for the HFP to calculate the incremental resources and
37 costs for that programme, as follows. Similarly, we use the length of stay for GORU/MARU to
38 calculate the incremental resources and costs for that rehab programme.

39 Tables 9 – 11 summarise the incremental resources used in the HFP and the GORU/MARU
40 programme, compared to usual care.

1 **Table 98: Incremental resource use for GORU/MARU programme versus usual care**

Staff resources	Incremental resources used, based on a LOS of 32.88 days	Source	Unit cost (source: PSSRU 2008/09), £ per hour	Incremental cost
Orthogeriatrician	Two consultant ward rounds (0.25/hour per patient each) and one weekly conference (0.25/hour = 0.75 hour per week per patient 0.75*4.6 weeks = 3.45 hours per patients	Kennie et al (1988) ¹⁷⁶	£108	£372.6
Physiotherapist	8.5 hours per patient	Naglie 2002 ²²²	£23	£195.5
Occupational therapist	5 hr/patient	GDG adjustment from the 7.5 hr/pt reported in Naglie ²²²	£23	£115
Nurse	Initial assessment within 72 hours (0.5 hour per patient) and twice weekly assessment afterwards (0.25*2)/hour per patient 0.5+0.5*4.6 weeks= 2.8 hours per patient	Naglie 2002 ²²²	Nurse team leader: £27 Nurse day ward: £21	£75.6 £58.8
Social worker	-0.4 hour per patient	Naglie 2002 ²²²	£29 (from community data)	-£11.6
Dietician	-0.4 hour per patient	Naglie 2002 ²²²	£23/	-£9.2
Total incremental cost for GORU/MARU over usual care:				£721 (with generic nurse, Band 5); £738 (with team leader nurse, Band 6)

2

3

4

1 **Table 99: Incremental resource use and incremental cost for HFP over usual care**

Staff resources	Incremental resources used based on a LOS of 25.5 days	Source	Incremental cost (using PSSRU 2008/09 unit costs)
Orthogeriatrician	Initial assessment 0.5 hour per patient, and subsequently 0.25 hour per day: $0.50 + 0.25 \times 24.5 = 6.625$ hour per patient	Cameron (1993) ⁴⁴ ; Shyu (2008) ³⁰⁵ ; Marcantonio ²⁰³	$\pounds 108 \times 6.625 = \pounds 715.50$
Physiotherapist or nurse	0.5 hour per patient per day: $0.50 \times 25.5 = 12.75$ hours	Cameron (1993) ⁴⁴	$\pounds 23 \times 12.75 = \pounds 293.25$
Total incremental cost of HFP over usual care: $\pounds 1009$			

2

3 Hence, the incremental cost for HFP over usual care is $\pounds 1009$, while for the GORU/MARU
4 programme it is $\pounds 721$ (with generic nurse) or $\pounds 738$ (with team leader nurse).

5 **Health state related costs in cycle 0**

6 To calculate the health state costs during the hospital stay, we used the NHS reference cost for
7 excess bed days reported in table 28 below. The excess bed day cost is the cost per day for days
8 exceeding the tripoint, a cut-off that determines patients with exceptionally long stay, and as
9 such usually estimates the cost of care without the cost of procedures (i.e. without the cost of the
10 surgery. These costs reflect the presence of complications experienced by hip fracture patients
11 during their entire hospital stay. Moreover, they distinguish between "major" and "intermediate"
12 complications, thus allowing users to take into account the different degrees of resource use.

13 **Table 100: National Schedule of Reference Costs Year : '2008-09' - NHS Trusts and PCTs**
14 **combined Non-Elective Inpatient (Long Stay) Excess Bed Day HRG Data for hip procedures**

Currency Code	Currency Description	Activity	National Average Unit Cost
HA11A	Major Hip Procedures Category 2 for Trauma with Major CC	360	$\pounds 243$
HA11B	Major Hip Procedures Category 2 for Trauma with Intermediate CC	620	$\pounds 242$
HA11C	Major Hip Procedures Category 2 for Trauma without CC	162	$\pounds 220$
HA12B	Major Hip Procedures Category 1 for Trauma with CC	9,760	$\pounds 237$
HA12C	Major Hip Procedures Category 1 for Trauma without CC	1,230	$\pounds 226$
HA13A	Intermediate Hip Procedures for Trauma with	14,891	$\pounds 240$

	Major CC		
HA13B	Intermediate Hip Procedures for Trauma with Intermediate CC	12,856	£249
HA13C	Intermediate Hip Procedures for Trauma without CC	2,972	£223
HA14A	Minor Hip Procedures for Trauma with Major CC	5,195	£234
HA14B	Minor Hip Procedures for Trauma with Intermediate CC	5,808	£245

1

2 The GDG decided to calculate a weighted average cost of the different categories of hip fractures
3 taking into account the level of activity associated with each procedure.

4 To cost the health state “not recovered with pressure sores” we use evidence from Bennett
5 (2004)¹⁷ regarding the cost of pressure ulcer treatment in the UK. The paper calculates the daily
6 cost of treating pressure ulcers looking at resources such as nurse time (dressing changes, patient
7 repositioning and risk assessment) dressings, antibiotics, diagnostic tests, and support surfaces.
8 These costs do not include inpatient costs, but assume that the patients are cared for in an
9 institutional setting (hospital or long-term care).

10 Pressure ulcers can have a different “grade”, ranging from 1 to 4 as their complexity increases.
11 However, the GDG emphasised that the published evidence on the incidence of the different
12 types of pressure sores in hip fracture patients reports many contradictory findings from which it
13 is difficult to draw definitive conclusions when it comes to costs. We followed the evidence in
14 Rademakers (2007)²⁷⁸ and assumed that 97% of the pressure ulcers were of grade 2, and 3% of
15 grade 3 or 4.

16 Bennett (2004)¹⁷ reports a daily cost for grade 2 pressure sores of £42, and of £50 for grade 3 and
17 4. These daily costs refer to patients who do not develop any further complications linked to the
18 pressure sores (such as critical colonisation, cellulites, or osteomyelitis), as no evidence on such
19 conditions was available from the RCTs included in our clinical review. Table 101 reports the total
20 daily cost for the “not recovered with pressure sores” health state.

21 **Table 101: Total daily hospital cost for patients with pressure sores**

Category of cost	Level of cost
Daily inpatient hospital cost without complications	£220.07
Daily cost of grade 2 pressure sore	0.97*£45
Daily cost for grade 3 and 4 pressure sore	0.03*£50
Total daily cost for patients with pressure sores	£265.22

22

23 For the cost of the health state “not recovered with moderate delirium” we used the mean
24 weighted average cost for minor complications (£237), and for the cost of the health state “not
25 recovered with severe delirium”, we used the mean weighted average cost for major and
26 intermediate complications (£242.89). One limit with this approach is that all patients with
27 moderate delirium are assumed to have undergone a Major Hip Procedures Category 1 for

1 Trauma. Even if the difference between the two cost figures is quite low (£5.89) we test the
2 impact of this assumption on the base case findings in a sensitivity analysis.

3 It has to be emphasised that this approach to calculate the health state costs in cycle 0 is
4 necessary in that only figures regarding the *total* length of hospital stay are available from the
5 evidence included in our clinical review. Ideally, we would have needed information regarding the
6 *additional* length of hospital stay for the patients experiencing a particular complication, both for
7 the control and for the intervention groups, but this information was not available from the
8 clinical review. Moreover, even if Marcantonio (2001)²⁰³ reports the hospital days of delirium per
9 episode, it does not distinguish between the two types of delirium (moderate and severe) that
10 correspond to our health states in cycle 0 of the Markov model, and only gives an overall figure
11 for all types of delirium.

12

13 **Table 102: Daily inpatient average cost for health states in cycle 0**

Health state	Average daily cost	Source
Not recovered and with no complications	£220.07	Mean weighted average of excess bed days costs – NHS reference costs 2008-08 Major, Intermediate and Minor Hip procedures <i>with no complications</i>
Not recovered and with pressure sores	£265.22	See Table 29
Not recovered and with moderate delirium	£237	Mean weighted average of excess bed days costs – NHS reference costs 2008-08. Major, Intermediate and Minor Hip procedures <i>with minor complications</i>
Not recovered and with severe delirium	£242.89	Mean weighted average of excess bed days costs – NHS reference costs 2008-08. Major, Intermediate and minor hip procedures <i>with intermediate and major complications</i>

14

15 ***Evidence and treatment effects on length of hospital stay***

16 The studies included in the clinical review comparing the GORU/MARU programme vs usual care
17 only reported the *total* length of hospital stay for patients in the intervention arm of the study.
18 Hence, no information was available to evaluate the number of days patients spent in the
19 orthopaedic ward and the number of days they spent in the orthogeriatric rehabilitation hospital
20 ward.

21 To calculate the length of stay at baseline (i.e. the usual care arm of the model), we pooled the
22 data for the usual care arm from all RCTs included in the clinical review. Table 103 reports the
23 relevant values for hospital length of stay used in the model:

24 **Table 103: Mean length of hospital stay**

Mean length of stay - usual care (days)	31.56
-----------------------------------------	-------

Mean difference length of stay - HFP (days)	-6.06
Mean difference length of stay - GORU/MARU (days)	1.32

1

2

20.6.8.2 Cost data: cycle 1 - onwards

3 From cycle 1 – onwards, the costs for our model will depend on the place of discharge (whether
4 own home or residential or nursing home), which in turn will affect the level of health care
5 services and social care used, and on the probability of hospital readmissions.

6 **Hospital readmissions**

7

8 The RCTs on HFP versus usual care included in the clinical review did not report any information
9 over the reasons for hospital readmissions nor the associated length of stay.

10 Two RCTs on GORU/MARU versus usual care (Galvard 1995 and Stenvall 2007)^{107,320} reported data
11 on length of stay following readmission available from two RCTs on GORU/MARU. However, the
12 reasons for readmissions (whether orthopaedic-related or any other medical reason) were only
13 given in Galvard (1995)¹⁰⁷.

14 Given the lack of data from the clinical review, the GDG decided to assume that readmissions
15 were composed by an equal proportion of patients are readmitted for surgery, medicine and
16 rehabilitation reasons. This assumption was also supported by unpublished data on readmissions
17 following hip fracture obtained from a GDG member and based on hospital records from
18 Peterborough and Stamford NHS Foundation Trust.

19 As for the length of stay following a hospital readmission, we followed the most recent clinical
20 paper (Stenvall 2007)³²⁰ and assumed a LOS for readmission for usual care is 11 days and in the
21 intervention (whether GORU/MARU or HFP) is 7 days.

22 The cost data for the hospital readmissions were obtained from Czoski-Murray (2007)⁶³, which
23 reports the unit costs for inpatient stay (at 2002 prices) for surgery (£381), medicine (£282) and
24 rehabilitation (£188). These costs are based on Netten et al (2002)²⁴¹. The mean unit cost for
25 inpatient stay for readmissions (at 2009 prices) was estimated at £367.00. This price has been
26 obtained using the annual percentage increases for prices of hospital and community health
27 services (HCHS) for 2002/03 – 2008/09 reported in the PSSRU 2009 report⁶¹.

28 **Community care costs for the “recovered” and “not recovered” health states when discharged to 29 own home**

30 To analyse the costs associated with the “recovered” and the “not recovered” health states we
31 need to take in to consideration whether patients are discharged to a long-term care setting or to
32 their own home.

33 The GDG decided that in determining the level of community (that is, health care and social care)
34 resources used after the hip fracture and after the rehabilitation programme it was important to
35 reflect the level of “dependency” and “independency” in activities of daily living of patients in each
36 of the health state.

37 The PSSRU 2009 identifies five different “packages” of community care provided in the home
38 setting of the patient (also known as “domiciliary care”), according to the different level of

1 dependency in the activities of daily living of the recipients. These packages of care are
2 summarised in Table 104 below.

3 **Table 104: Weekly costs of community care packages – excluding accommodation and living**
4 **expenses. Source PSSRU 2009.**

Community care package	Description of the level of functional ability of the recipient of care	Weekly cost (excluding accommodation, living expenses and independently provided home care)	Average weekly cost of social care services	Average weekly cost of health care services
“Very low cost”	Mrs A. had problems with three activities of daily living: stairs, getting around outside, and bathing. Her problems stemmed from a previous stroke.	£49	£41.3 (£18.10 of home care (one hour of weekly local authority-organised home care)) and £23.20 of meals on wheels)	£7.70 for a 11.7 minutes of GP surgery visit (one every four weeks)
“Low cost”	Mrs B. had problems with three activities of daily living: stairs, getting around outside and bathing. Her problems stemmed from arthritic conditions and cardiovascular disease.	£87(1)	£72 of home care (4 hours of local authority-organised home care)	£14.3 (of which £6.60 of community nurse (one visit per month) and £7.70 of one GP visit (one every four weeks))
“Median cost”	Mrs C. had problems with four activities of daily living: stairs, getting around outside, dressing and bathing.	£188	£181 of home care (10 hours of weekly local authority-organised home care)	£7.70 for a 11.7 minutes of GP surgery visit (one every four weeks)
“High cost”	Mr D. had problems with seven activities of daily living: stairs, getting around outside and inside the house, using the toilet, transferring between	£273	£216 (of which £181 of home care (10 hours of weekly local authority-organised home care) and £35 for a	£58 £26 of community nurse (once a week); £24 for two monthly OT visits; £7.70 for a 11.7 minutes of GP

	chair and bed, dressing and bathing. His problems stemmed from arthritic conditions and a previous stroke.		day centre attended once a week)	surgery visit (one every four weeks)
“Very high cost”	Mrs E suffered from dementia and needed help with nine activities of daily living: stairs, getting around outside and inside the house, using the toilet, transferring between chair and bed, dressing, bathing, washing and feeding.	£576	£542 of home care (30 hours of weekly local authority-organised home care)	£34 £26 of community nurse (once a week); £7.70 for a 11.7 minutes of GP surgery visit (one every four weeks)

1 (1) Please note that the cost figure reported in the PSSRU 2009 for “low cost” is not correct
 2 (£129) as the cost for the independently provided health care has not been subtracted (£42).
 3 The correct figure should be £87.

4 We used the data from Kennie (1988)¹⁷⁶ to determine the proportion of patients with level of
 5 independence from A to G to attribute the community care costs to the “recovered” and “not
 6 recovered” health state.

7 For both health states (“recovered” and “not recovered”), we assume that patients with ADL
 8 score A or B do not incur in any domiciliary care cost. However, we assume that the same type of
 9 patients will each visit the GP once weekly.

10 The weekly health and social care costs are calculated by multiplying the weekly unit cost of the
 11 different type of care (as obtained from the PSSRU 2009) times the proportion of patients with
 12 the corresponding ADL score in the specific health state and times the level of resources used
 13 (which depend on the level of dependency). The health and social care costs for the “recovered”
 14 and “not recovered” health states are described in Table 105 and in Table 106 below.

15 **Table 105: Health and social care costs for patients in the “recovered” health state discharged at**
 16 **their own home**

ADL	% ADL in recovered state	Unit health care costs	Health cost for recovered state	Unit social care costs	Social care costs for recovered state
A	0.454	7.7	3.4958	N/A	N/A
B	0.231	7.7	1.7787	N/A	N/A
C	0.112	7.7	0.8624	41.3	4.6256
D	0.056	7.7	0.4312	41.3	2.3128
E	0.028	14.3	0.4004	72	2.016
F	0.028	7.7	0.2156	181	5.068

G	0.018	58	1.044	216	3.888
NC	0.073	34	2.482	542	39.566
		Total	10.7101	Total	57.4764
		Annual health care cost	556.925	Annual social care cost	2988.77

1

2 **Table 106: Health and social care costs for patients in the “not recovered” health state**
3 **discharged at their own home**

ADL	% ADL in not recovered state	Unit health care costs (£)	Health cost for recovered state (£)	Unit social care costs (£)	Social care costs for recovered state (£)
A	0	0	0	N/A	N/A
B	0.009	7.7	0.0693	N/A	N/A
C	0.009	7.7	0.0693	41.3	0.3717
D	0.046	7.7	0.3542	41.3	1.8998
E	0.342	14.3	4.8906	72	24.624
F	0.362	7.7	2.7874	181	65.522
G	0.204	58	11.832	216	44.064
NC	0.028	34	0.952	542	15.176
			20.9548		151.658
		Annual health care cost	1089.65	Annual social care cost	7886.19

4

5 Hence, the annual health and social care costs for the “recovered” and the “not recovered” health
6 state are:

7 **Table 107: Annual health and social care costs for the “recovered” and the “not recovered”**
8 **health state**

Annual health care costs	£557	£2989
Annual social care costs	£1090	£7886
Total community care costs	£1647	£10875

9

10 While the health care costs will be fully funded by the NHS, the social care costs will only be
11 generally partially funded by the local councils^{71, 348, 72, 144}. It was not possible to identify a
12 national average for the social care costs funded by local authorities in the published literature,
13 and as a consequence an assumption had to be made regarding the proportion of this care that
14 was publicly funded. In the base case analysis, we assume that 60% of social care costs are funded
15 by the local authorities, and are therefore includable in the model, and we then test this
16 assumption in a sensitivity analysis.

1 **Community care costs for the “recovered” and “not recovered” health states when discharged to**
 2 **long term care**

3 The cost of long term care used in the model was estimated from the unit cost of stay in private
 4 nursing homes, private residential care, voluntary residential care and local authority residential
 5 care facility for older people. The care package costs per permanent residential week are
 6 described in Table 108.

7 **Table 108: Weekly long term care costs for patients not discharged to their own home.**
 8 **(Source: PSSRU 2009).**

Type of long term care	Weekly health care costs	Weekly fees (minus living expenses)
Private nursing home	£30.80 £30.00 (GP weekly home visit) £0.80 (community nursing)	£678
Private residential care	£26.3 £19.30 (GP weekly home visit) £7.00 (community nursing)	£467
Voluntary residential care	£28.7 £19.30 (GP weekly home visit) £9.40 (community nursing)	£470
Local authority residential care	£20.9 £10.60 (GP weekly surgery visit) £10.30 (community nursing)	£902

9

10 These unit costs include the cost of external services such as community nursing, GP services as
 11 well as personal living expenses. They also include capital costs for the local authority residential
 12 care, and fees for the private and voluntary residential care. We subtracted £9.20, the cost of
 13 personal living expenses per week, from each unit cost and estimated **£717.05**, the weighted
 14 average of £708.80, £493.80, £489.80 and £913.80, to be the weekly unit cost of long term care.
 15 By also subtracting the health care costs, we get: **£557.64 as the weekly fees** for long term care
 16 (£28997 per year). The (weighted) **health cost per week is £27** (£1404 per year).

17 The weighting is based on the distribution of residents, 65 years and older, in care homes in 1996.
 18 It was reported that in nursing homes, local authority, private and voluntary residential homes the
 19 number of residents were 5746, 5476, 2791 and 3664 respectively (Netten et al 1998)²⁴⁰. A
 20 similar approach is also followed in the cost-effectiveness analysis conducted in the NICE Delirium
 21 Guideline²²⁴.

22 It is important to note that, contrary to the community care packages for domiciliary care, we
 23 could not distinguish the level of long-term residential care according to the level of
 24 “dependency” in ADL of the patients in the different health state. Hence, the same figure for
 25 community costs had to be used both for the “recovered” and “not recovered” health states if not
 26 discharged at their own home.

27 As with the domiciliary care, the health care costs in table 18 will be fully funded by the NHS, but
 28 the residential fees for long term care will only be generally partially funded by the local councils.
 29 Moreover, only a very small proportion of patients (about 2%) qualifies for fully funded NHS care

1 (the so called “continuing care”)^{71, 348, 72, 144}. It was not possible to identify a national average for
 2 this figure in the published literature, and as a consequence an assumption had to be made
 3 regarding the proportion of residential costs in long term care paid by local authorities. In the
 4 base case analysis, we assume that 60% of residential fees costs are funded by the local
 5 authorities, and then change this assumption in a sensitivity analysis.

6 20.6.9 Cost-effectiveness findings for base-case analysis

7 In the base case analysis, HFP is the dominant strategy (more effective, less costly) than both
 8 GORU/MARU and usual care.

9

10 **Table 109: Cost-effectiveness findings from the deterministic base case analysis**

Strategy	Cost (£000)	Incremental Cost* (£000)	Effectiveness (QALYs)	Incremental Effectiveness* (QALYs)	Incremental cost-effectiveness
HFP	£34		3.75		
GORU/MARU	£36	£2	3.62	-0.13	(Dominated by HFP)
Usual care	£59	£26	2.73	-1.02	(Dominated by HFP)

11 *Compared with HFP

12

13 Table 38 below shows the breakdown of the different cost categories for the three strategies of
 14 the deterministic base case

15

16 **Table 110: Cost breakdown for usual care, HFP and GORU/MARU**

Resource item	Usual Care	HFP	GORU
Rehab cost (initial costs)*	-	1009	729
Complications*	-	-548	217
Readmission	969.5	762.2	535.3
Health care costs – living in own home	9178	4032	3738
Social care costs – living in own home	14,000	5,000	5,000
Health care costs – residential and nursing home	2,615	1,801	1930
Social care costs (fees) - residential and nursing home	32,000	22,000	24,000
Total cost	58762.50	33595.2	35203.3

17 * calculated incrementally vs usual care

18

19 20.6.9.1 Sensitivity analyses

1 In order to check how robust the findings in the deterministic base case analysis reported in
2 table 20 are, we ran a series of sensitivity analyses.


3 The results were not sensitive to changes in several parameters (length of hospitals stay, cost
4 of long-term care, proportion of long-term care borne by the NHS and PSS).

5 However, the results were sensitive to changes in the probability of returning home for both
6 HFP and GORU/MARU. In the base case analysis, the probability of returning home for the
7 HFP is 0.81 (RR of HFP vs usual care: 1.14), and for GORU/MARU it is 0.79 (RR of GORU/MARU
8 vs usual care: 1.11). The findings of a two-way sensitivity analysis on such probabilities are
9 reported in the graph below.

10

11

Net Monetary Benefit (wtp=20000.) Sensitivity Analysis: p_ReturnHome_GORU and p_ReturnHome_HFP



12

13

14

15 A threshold sensitivity analysis shows that:

- 16 a) If the probability of returning home for HFP < 0.77 (it is 0.81 in the base case scenario),
17 then GORU/MARU is the most cost-effective option at a willingness to pay threshold of
18 £20,000 per QALY.
- 19 b) If probability of returning home for GORU/MARU < 0.83 (it is 0.79 in the base case
20 scenario), then HFP is the most cost-effective option at a willingness to pay threshold of
21 £20,000 per QALY.

1 A two-way sensitivity analysis on a) the proportion of social care costs borne by the NHS and
 2 PSS for patients living in their own home and b) the proportion of social care costs borne by
 3 the NHS and PSS for patients living in a residential or nursing accommodation found that HFP
 4 is always the most cost-effective option.

5 **20.6.9.2 Probabilistic sensitivity analysis**

6 A probabilistic sensitivity analysis was performed to assess the robustness of the model
 7 results to plausible variations in the model parameters.

8 Probability distributions were assigned to each model parameter, where there was some
 9 measure of parameter variability. We then re-calculated the main results 10,000 times, and
 10 each time all the model parameters were set simultaneously, selecting from the respective
 11 parameter distribution at random. Table 111 describes the type and properties of the
 12 distributions used in the probabilistic sensitivity analysis.

13

14 **Table 111: Description of the type and properties of distributions used in the probabilistic**
 15 **sensitivity analysis**

Parameter	Type of distribution	Properties of distribution
Baseline risk	Beta	Bounded on 0 – 1 interval. Derived from sample size, number of patients experiencing events
Cost	Gamma	Bounded at 0, positively skewed. Derived from mean and standard error
Utility	Beta	Bounded on 0 – 1 interval. Derived from mean and sample size
Risk ratio, length of stay	Lognormal	Bounded at 0. Derived from log and standard error of log
Mean differences (e.g. in length of stay, time of therapies, etc.)	Normal	Derived from mean and standard deviation

16

17 Table 112 summarises the distribution, parameters and expected values for each variable of
 18 the model.

19

20 **Table 112: Probabilistic sensitivity analysis: formulas and expected value**

Variable name	Formula	Expected value	Deterministic value
Cost per hospital bed day (patients with moderate delirium)	Gamma alpha = 15.366, lambda = 0.0648;	237.13	237
Cost per hospital bed day (patients with no complications)	Gamma alpha = 15.366, lambda =	220.14	220.07

	0.0698;		
Cost per hospital bed day (patients with pressure sores)	Gamma alpha = 15.366, lambda = 0.057984;	265.00	265.22
Cost per hospital bed day (patients with severe delirium)	Gamma alpha = 15.366, lambda = 0.0633;	242.75	242.89
Annual health care costs – “not recovered” patients living in their own home	Gamma alpha = 15.366, lambda = 0.005141;	2988.91	2989
Annual health care costs for “recovered” patients living in their own home	Gamma alpha = 15.366, lambda = 0.0275;	558.76	557
Annual social care costs for “not recovered” patients living in their own home	Gamma alpha = 15.366, lambda = 0.001948;	7888.09	7886
Annual social care costs for “recovered” patients living in their own home	Gamma alpha = 15.366, lambda = 0.014;	1097.57	1090
Annual cost for fees in long term care – “not recovered” patients	Gamma alpha = 15.366, lambda = 0.00053;	28992.45	28997
Annual cost for fees in long term care – “recovered” patients	Gamma alpha = 15.366, lambda = 0.00053;	28992.45	28997
Annual health care costs for “not recovered” patients in long term care	Gamma alpha = 15.366, lambda = 0.0109;	1409.72	1404
Annual health care costs for “recovered” patients in long term care	Gamma alpha = 15.366, lambda = 0.0109;	1409.72	1404
Cost of hospital bed day for readmissions	Gamma alpha = 15.366, lambda = 0.0474;	324.18	324.01
Cost per hour of day ward nurse	Gamma alpha = 15.366, lambda = 0.7317;	21	21
Cost per hour of a dietician	Gamma alpha = 15.366, lambda = 0.668;	23	23
Cost per hour of a geriatrician	Gamma alpha = 15.366, lambda = 0.1423;	108	108
Cost per hour of an occupational therapist	Gamma alpha = 15.366, lambda = 0.668;	23	23
Cost per hour of a physiotherapist	Gamma alpha = 15.366, lambda = 0.668;	23	23

Cost per hour of a social worker	Gamma alpha = 15.366, lambda = 0.5298;	29	29
Cost per hour of a team lead nurse	Gamma alpha = 15.366, lambda = 0.5691;	27	27
Initial age	None – from meta analysis of RCTs	81	81
Length of stay (days) – usual care	Log-Normal, u (mean of logs) = 3.439942259 sigma (std dev of logs) = 0.154584841;	31.56	31.56
Length of stay (days) – mean difference – GORU/MARU	Normal, Mean = 1.32, Std Dev = 0.03322; Expected value: 1.32	1.32	1.32
Length of stay (days) – mean difference – HFP	Normal, Mean = -6.06, Std Dev = 0.3593	-6.06	-6.06
Length of stay for hospital readmissions – GORU/MARU	Triangular, Min = 4, Likeliest = 7, Max = 10;	7	7
Length of stay for hospital readmissions – HFP	Triangular, Min = 4, Likeliest = 7, Max = 10;	7	7
Length of stay for hospital readmissions – usual care	Triangular, Min = 7, Likeliest = 11, Max = 15;	11	11
Proportion of patients with ADL scores C-G in the “not recovered” health state	Beta Integer parameters only, n = 108, r = 107;	0.99	0.99
Proportion of patients with ADL scores C-G in the “recovered” health state	Beta Integer parameters only, n = 108, r = 34;	0.31	0.31
Probability moderate delirium – usual care	Beta Integer parameters only, n = 64, r = 14;	0.22	0.22
Probability pressure sores –usual care	Beta Integer parameters only, n = 164, r = 27;	0.1646	0.1646
Probability die at 12 months – usual care	Beta Integer parameters only, n = 870, r = 186;	0.2138	0.2138
Probability die at 6 months – usual care	Beta Integer parameters only, n = 263, r = 44 ;	0.1673	0.1673
Probability die 6 to 12 months – usual care	Beta Integer parameters only, n = 219, r = 12;	0.0548	0.0558
Probability of hospital readmission at 12 months – usual care	Beta Integer parameters only, n = 640, r = 165;	0.2578	0.26

Probability of not recovery of pre-fracture ADL levels at 12 months – usual care	Beta Integer parameters only, n = 283 , r = 167;	0.59	0.59
Probability of not recovery of pre-fracture ADL levels at 3 months – usual care	Beta Integer parameters only, n = 125, r = 91;	0.728	0.73
Probability of not recovery of pre-fracture ADL levels at 6 months – usual care	Beta, Integer parameters only, n = 125, r = 84;	0.672	0.67
Proportion of social care costs funded by the NHS or local authorities – patients living in their own home	Triangular, Min = 0.3, Likeliest = 0.6, Max = 0.9;	0.6	0.6
Proportion of long term fee costs funded by the NHS or local authorities – patients living in long term care	Triangular, Min = 0.3, Likeliest = 0.6, Max = 0.9;	0.6	0.6
Probability severe delirium – usual care	Beta Integer parameters only, n = 64 , r = 18;	0.28125	0.28125
Proportion of men - HFR and GORU/MARU	None – from meta analysis of RCTs		0.76
Proportion of men - usual care	None – from meta analysis of RCTs		0.79
Relative risk of die – 12 months – GORU/MARU	Log-Normal, u (mean of logs) = -0.05969, sigma (std dev of logs) = 0.129622261;	0.95	0.95
Relative risk of die – 12 months – HFP	Log-Normal, u (mean of logs) = -0.22022, sigma (std dev of logs) = 0.140960518;	0.81	0.81
Relative risk of die – 6 months – GORU/MARU	Log-Normal, u (mean of logs) = -0.26001, sigma (std dev of logs) = 0.220399212;	0.79	0.79
Relative risk – moderate delirium - HFP	Log-Normal, u (mean of logs) = -0.10966, sigma (std dev of logs) = 0.341655183;	0.95	0.95
Relative risk – not recovery – GORU/MARU	Log-Normal, u (mean of logs) = -0.25423, sigma (std dev of logs) = 0.107452415;	0.78	0.78
Relative risk – not recovery – HFP	Log-Normal, u (mean of logs) = -0.24094, sigma (std dev of logs) = 0.102123395;	0.79	0.78
Relative risk pressure sores – HFP	Log-Normal u (mean of logs) = -	0.31	0.31

	1.24407, sigma (std dev of logs) = 0.381796535;		
Relative risk of readmissions – GORU/MARU	Log-Normal u (mean of logs) = -0.15941, sigma (std dev of logs) = 0.131073023;	0.86	0.86
Relative risk of readmissions - HFP	Log-Normal u (mean of logs) = 0.121843, sigma (std dev of logs) = 0.135536774;	1.14	1.14
Relative risk of returning to own home – GORU/MARU	Log-Normal u (mean of logs) = 0.103769, sigma (std dev of logs) = 0.0347056;	1.11	1.11
Relative risk of returning to own home – HFP	Log-Normal u (mean of logs) = 0.129321, sigma (std dev of logs) = 0.058435349;	1.14	1.14
Relative risk – severe delirium – HFP	Log-Normal u (mean of logs) = -0.99941, sigma (std dev of logs) = 0.407720564;	0.4	0.4
Time input of dietician (incremental over usual care) – GORU/MARU	Normal Mean = -0.4, Std Dev = 0.0332;	-0.4	-0.4
Time input of nurse - (incremental over usual care) – GORU/MARU	Normal Mean = 2.8, Std Dev = 0.358;	2.8	2.8
Time input of occupational therapist - (incremental over usual care) – GORU/MARU	Normal Mean = 5, Std Dev = 0.64;	5	5
Time input of physiotherapist (incremental over usual care) – GORU/MARU	Normal Mean = 8.5, Std Dev = 1.09;	8.5	8.5
Time input of social worker - (incremental over usual care) – GORU/MARU	Normal Mean = -0.4, Std Dev = 0.32;	-0.4	-0.4
Transition probability from “not recovered” to “recovered” health state – 0 to 3 months	Beta Integer parameters only, n = 125, r = 34;	0.272	0.27
Transition probability from “not recovered” to “recovered” health state – 3 to 6 months	Beta Integer parameters only, n = 91, r = 7;	0.07692307 7	0.082192
Transition probability from “not recovered” to “recovered” health state – 6 to 12 months	Beta Integer parameters only, n = 499212, r = 60837	0.12186606 1	0.119403
EQ-5D score for “Recovered” health state	Beta Real-numbered	0.68	0.68

	parameters, alpha = 35.36, beta = 16.64;		
EQ-5D score for “Not recovered with no complications” health state	Beta Real-numbered parameters, alpha = 3.72, beta = 8.28;	0.31	0.31
EQ-5D score for “Not recovered with pressure sores” health state	Beta Real-numbered parameters, alpha = 0.952227, beta = 4.059492;	0.19	0.19
EQ-5D score for “Not recovered with severe delirium” health state	Beta Real-numbered parameters, alpha = 293, beta = 880;	0.25	0.25

1

2 The conventional way to identify the most cost-effective strategy is to look at the option that is
3 optimal based on the mean costs and mean QALYs averaged across all of the probabilistic
4 simulations. These findings are summarised in Table 113.

5

6 **Table 113: Cost-effectiveness analysis from probabilistic analysis**


Strategy	Cost	Incremental Cost	Effectiveness	Incremental effectiveness	Incremental C/E (ICER)
HFP	£34K		3.74		
GORU/MARU	£36K	£2K	3.61	-0.13	(Dominated)
Usual care	£59K	£25K	2.73	-1.01	(Dominated)

7

8 The probabilistic results are very similar to the deterministic ones indicating that HFP is dominant
9 (has lower cost and more QALYs) compared with the two alternatives.

10 These findings are described in Figures 151, 152 and 153. Each point on the second scatter plot
11 represents the incremental cost and QALYs gained for HFP vs GORU for one simulation. The
12 dotted line represents the £20,000/QALY threshold and the ellipse delimits the 95% confidence
13 space.

ICE Scatterplot of HFP vs. GORU/MARU




1
2
3

Figure 162: Incremental cost-effectiveness scatter plot: HFP vs GORU/MARU

4 The scatter plot of HFP vs usual care shows the high certainty of HFP being cost-effective as all the
5 dots in the 95% confidence ellipse are below the £20,000/QALY threshold and more than 95% are
6 cost saving.


7



1
2

3 **Figure 163: Incremental cost-effectiveness scatter plot: GORU/MARU vs. usual care**

ICE Scatterplot of HFP vs. Usual care



1
2 **Figure 164: Incremental cost-effectiveness scatter plot - HFP vs usual care**
3

4 However, when we compared HFP with GORU the 95% CI showed a greater uncertainty as HFP
5 was dominant in the lower bound and GORU was dominant in the upper bound. The uncertainty
6 can be graphically represented by plotting the results of the incremental analysis for all the
7 10,000 simulations into a cost-effectiveness plane.

8 We also found that, at a willingness to pay equal to £20,000 per QALY, HFP was the optimal
9 strategy in 70% of the simulations; GORU/MARU was the most cost-effective intervention in 30%
10 of simulations, and usual care was never the optimal strategy. These findings are summarised in
11 table 42 below:

12 **Table 42: Probability most cost-effective intervention at a willingness to pay of £20,000 and**
13 **£30,000 per QALY**

Strategy	Probability most cost-effective intervention at a WTP of £20,000 per QALY	Probability most cost-effective intervention at a WTP of £30,000 per QALY
HFP	0.70	0.80
GORU/MARU	0.30	0.20
Usual care	0	0

14
15
16
17

1 20.6.10 Discussion

2 The optimal strategy in a cost-effectiveness analysis is the one with the highest incremental net
3 benefit averaged across all the probabilistic simulations. This was HFP.

4 The model showed that usual care was clearly not the optimal strategy.

5 However, there was some uncertainty about which strategy was the most cost-effective between
6 HFP and GORU/MARU. In particular the results were sensitive to the proportion of patients
7 returning home after their rehabilitation: if the probability of returning home after undergoing a
8 GORU/MARU programme was 83% (instead of 79% in the base case) then GORU is the optimal
9 strategy.

10

11 Our analysis had to rely on several assumptions.

12 Firstly, no evidence was available which compared directly HFP vs GORU/MARU. As a
13 consequence, only an *indirect comparison* between the two hospital MDR programmes was
14 possible. This meant that findings had to be pooled in the usual care arm of the different RCTs
15 included in our clinical review, thus assuming that “usual care arms” in all such studies were
16 sufficiently similar. However, the GDG agreed that the population included in the RCTs on HFP
17 and the population included in the RCTs on GORU were sufficiently similar and that therefore our
18 findings were not affected by confounding factors.

19 Secondly, no data were available regarding the presence and incidence of complications in the
20 GORU/MARU programme versus usual care. The assumption that in this case the relative risk for
21 that rehab programme was equal to 1 implies that we may have underestimated the efficacy of
22 GORU/MARU in reducing the presence of postoperative complications, and as a consequence,
23 that we may have overestimated its costs and decrement in quality of life compared to HFP.
24 However, when we changed the probabilities of complications for GORU/MARU in a one-way
25 sensitivity analysis, the findings of the cost-effectiveness analysis did not change, and HFP was still
26 the dominant strategy.

27 Finally, the finding of the meta-analysis of clinical trials regarding the length of stay showed a
28 longer length of stay for the GORU/MARU programme versus usual care (mean difference (days):
29 1.32). However, the inclusion of the study by Galvard (1995)¹⁰⁷ in the meta-analysis may have
30 biased this finding. This is because Galvard (1995)¹⁰⁷ reports a mean length of stay of 53.3 days for
31 the intervention (GORU) group and of 28 days for usual care. This finding, according to the
32 authors, was due to the fact that GORU was a new rehabilitation programme that had just been
33 implemented in their hospital, and the hospital staff was not yet experienced in the management
34 of the programme, which could have resulted in a longer length of stay for patients in the
35 intervention group. As a consequence, we may have overestimated the costs of hospital stay for
36 GORU/MARU. However, when we changed the length of hospital stay for the GORU/MARU
37 programme in a one-way sensitivity analysis, the findings of the cost-effectiveness analysis did not
38 change, and HFP was still the dominant strategy.

39

1

2 **20.7 Cost-effectiveness analysis of Community MDR vs Usual care**

3 **20.7.1 Introduction**

4 The GDG identified the multidisciplinary management in the community for hip fracture patients
5 as a high priority area for economic analysis.

6 The clinical question linked to this high priority area is the following:

7 *What is the comparative effectiveness of community -based multidisciplinary rehabilitation*
8 *models versus usual care?*

9 A review of the literature was conducted followed by economic modelling of the cost-
10 effectiveness of the listed interventions in England and Wales. The literature search and review
11 methods can be Chapter 3. Despite some cost-effectiveness studies were identified, none
12 represented a full cost-utility analysis which addressed our clinical question. As a consequence,
13 the GDG felt that an original economic model was essential in order to support their
14 recommendations.

15 The following general principles were adhered to:

- 16 • The GDG was consulted during the construction and interpretation of the model.
- 17 • When published data was not available we used expert opinion to populate the
18 model.
- 19 • Model assumptions were reported fully and transparently.
- 20 • The results were subject to sensitivity analysis and limitations were discussed.
- 21 • We followed the methods of the NICE reference case. Therefore costs were
22 calculated from the UK NHS and PSS perspective. Health gains were measured in
23 terms of quality-adjusted life-years (QALYs) gained.
- 24 • The model employed a cost-effectiveness threshold of £20,000 per QALY gained.
- 25 • The model was peer-reviewed by another health economist at the NCGC.

26 **20.7.2 Population and time horizon**

27 The population for the cost-effectiveness analysis consists of hip fracture patients (male and
28 female) hospitalised for surgery. The model spans over a life-time horizon.

29 **20.7.3 Software**

30 The cost-effectiveness analyses were conducted using TreeAge Pro 2008.

31 **20.7.4 Economic evaluation type**

32 We conduct a cost-utility analysis, where health outcomes are measured as Quality-Adjusted Life-
33 Years (QALYs). The cost effectiveness outcome of the model is measured as cost per QALY gained.

20.7.5 Time horizon and discount rates used

The model spans over a life-time horizon. All costs considered in the model were calculated at on the basis of a four-months follow-up time and hence were not discounted. However, we used a discount rate of 3.5% for the health gains, as these were calculated throughout the remaining life of the cohort of patients.

20.7.6 Structure of the model

The structure of our model reflects the findings of the RCT by Crotty et al (2002)⁶⁰. The paper reports SF-36 scores for surviving patients, both in the community MDR and in the usual care arm of the study, at a 4 months follow up.

We develop a decision tree with Markov states, where a hip fracture patients can either receive a community based MDR programme or usual inpatient rehabilitation. Following this decision node, a chance node determines whether patients survive or die following their specific rehabilitation programme. The probability associated to this chance node is derived from Crotty et al (2002)⁶⁰ at a 4 months follow up. Subsequently, patients who are alive after the 4-months follow up period transit in a Markov state, "alive after follow up". Patients will then either stay in that state or transit to the "dead" state in the following cycles.

The structure of the model is the following:




Figure 165: Model structure - community MDR vs usual care

20.7.7 Utility data

Utility weights are calculated using SF-36 scores obtained from Crotty et al (2002)⁶⁰. The paper only reports total scores for the physical and mental components. Following personal communications with the authors, we were able to access individual SF-36 scores, reported in Table 114 below:

1 **Table 114: SF-36 scores based on Crotty et al (2002)⁶⁰.**

SF-36 domain, Mean (SD)	Conventional Care, n=29	Early Discharge, n=30
Physical functioning	28.8 (25.2)	41.2 (26.6)
Social functioning	62.1 (40.0)	72.5 (32.4)
Role-physical	61.2 (41.0)	53.3 (40.9)
Role-emotional	83.9 (31.6)	77.8 (38.5)
Mental health	77.9 (14.2)	80.1 (19.8)
Vitality	45.0 (21.9)	54.2 (24.3)
Bodily pain	61.4 (30.9)	65.1 (24.4)
General health	61.8 (30.1)	69.3 (24.1)

2 Source: primary data supplied by the authors of Crotty et al (2002)⁶⁰

3 Using the Ara-Brazier method⁷, we mapped the individual SF-36 scores in EQ-5D utility weights.
 4 We found that the EQ-5D weight for patients undergoing community MDR is 0.732, and for
 5 patients undergoing usual inpatient rehabilitation is 0.643. As the effectiveness data refer to
 6 findings at 4 months⁶⁰, we used these utility weights for cycle 0 only. For cycle 1-onwards we
 7 assume that there is no difference in the utility score of the two groups of patients, and use the
 8 EQ-5D score of the control group also for patients in the community MDR arm of the model.

9 **20.7.8 Mortality**

10 The mortality rates for the community MDR and usual care patients have been adjusted to take
 11 into account the baseline characteristics of the two groups, which were very different in the two
 12 arms of Crotty et al (2002)⁶⁰, since 62% of patients were female in the COMMUNITY MDR versus
 13 75% in the usual care group, and the median age for COMMUNITY MDR patients was 81.6 versus
 14 83.5 years in the usual care arm.

15 First, we have calculated the age and gender-adjusted mortality rate (AMR) for the general UK
 16 population as per characteristics in usual care arm and the same for community MDR arm. Then,
 17 we have calculated the Standardised Mortality Rate (SMR) as = MR/AMR, both the usual care and
 18 the community MDR arm. We have then assumed that the average age for the overall population
 19 in the model was 80 years of age, and we have determined the probability of death using the
 20 formula: SMR*pDeath[80].

21 We have found that that probability of death at 4 months for the patients in the usual care arm
 22 corresponds to 0.07239, and for patients in the community MDR group is equal to 0.067. The
 23 relative risk of the mortality rate for community MDR compared to usual care is 0.925.

24 **20.7.9 Calculating QALYs gained**

25 For each strategy (community MDR and usual inpatient rehabilitation), the expected QALYs in the
 26 "survived" health state at each cycle are calculated as follows:

$$27 \quad \text{Expected QALYs} = \Sigma (U_{\text{survived}} \times P_{\text{survived}})$$

28 where: U_{survived} = the utility score for the patients who are still alive and P_{survived} = the proportion of
 29 alive patients

1 The proportion of patients in the “alive” health state depends on the effectiveness of the
2 treatment, and on the proportion of patients still alive, which falls as the number of cycles and
3 therefore age increases.

4 The overall *lifetime expected QALYs* are given by the sum of QALYs calculated for each cycle. The
5 *incremental QALYs gained* associated with a treatment strategy are calculated as the difference
6 between the expected QALYs with that strategy and the expected QALYs with the comparator.

7 **20.7.10 Cost analysis**

8 **20.7.10.1 Cost for the community MDR and inpatient rehabilitation programmes.**

9 While in hospital, we assume that there is no difference in the level and type of resources used by
10 patients in the two groups, as no evidence of the contrary was found in the literature. Moreover,
11 as patients receive their inpatients rehabilitation services without being discharged to a different
12 ward, they will still be under the same HRG recorded at admission. Thus, the rehabilitation that
13 patients receive while in hospital is not a type of discrete rehabilitation service, that is, a service
14 that can be cost using its own HRG, since: *“rehabilitation HRGs are only generated where care is
15 identified as taking place under a specialist rehabilitation consultation or within a discrete
16 rehabilitation ward or unit. [...] Where a patient is not admitted specifically to a rehabilitation unit
17 or where rehabilitation treatment is undertaken without transfer to a specialist consultant, or
18 without transfer to a rehabilitation unit, this should not be reported as discrete rehabilitation”*
19 (Collection Guidance on Reference Costs for 2006-07⁷⁰).

20 As a consequence, we use the reference cost for excess bed days reported in the National
21 Schedule of Reference Costs Year: '2008-09' - NHS Trusts and PCTs combined Non-Elective
22 Inpatient (Long Stay).

23 Crotty et al (2002)⁶⁰ report evidence on the presence of complications experienced by hip fracture
24 patients in the two groups while in acute care. None of these complications were statistically
25 significant different between usual care and community MDR (the complications were:
26 pneumonia, pressure sores, confusion, wound infection and urinary tract infection). Moreover, no
27 additional information was provided in the paper as to whether those complications resulted in a
28 prolonged length of hospital stay for patients in the community MDR scheme. Thus, we used the
29 weighted average NHS reference cost for excess bed days for major, intermediate and minor hip
30 procedures with all types of complications, amounting to £241.68 per day.

31 As for the daily cost of the community MDR scheme, we use the NHS reference cost (2008-09)
32 reported for “Hospital at Home/ Early Discharge Schemes - Fractured Neck of Femur”, which
33 corresponds to £94 per day.

34 We conduct a sensitivity analysis on these values in section 32.1 of this chapter.

35 **20.7.10.2 Length of stay**

36 Crotty et al (2002)⁶⁰ reports the following findings for the length of stay for the community MDR
37 and the usual inpatient rehabilitation:

38 Table 115: Length of stay in hospital and in own home

Length of stay community MDR (at home stay)	20.3 (mean, days)
Length of stay community MDR (at home stay) (in	7.8 (mean, days)

hospital stay)	
Length of stay usual care (in hospital stay)	14.3 (mean, days)

1

2 **20.7.10.3 Hospital readmissions and related length of stay**

3 Crotty et al (2002)⁶⁰ gives information about the levels of readmissions during the four months
 4 follow up of the study. The paper distinguishes between related readmissions and unrelated
 5 readmissions, and gives the length of stay for both cases. However what these related and
 6 unrelated admissions were was not clear in the paper. We consider surgery and the rehabilitation
 7 admissions to be the “related” readmissions, and we consider the cost of a bed day in medicine
 8 for the cost of not-related admissions.

9 These unit bed day costs are based on Czoski-Murray (2007)⁶³, which reports the cost per day for
 10 hospital stay in an orthopaedic, rehabilitation or general medicine ward at 2002 prices. We
 11 assume that the “related readmissions” take place either for orthopaedic or for rehabilitation
 12 reasons, and that the “unrelated readmission” are those in the generic medicine ward.

13 Taking into account of the inflation index, the cost per day of hospital stay for a related
 14 readmission corresponds to £367.85 (assuming that half of these readmissions took place for
 15 surgery and half for rehabilitation reasons) and to £364.61 for unrelated readmission.

16

17 **Table 116: Evidence on readmissions (Crotty et al, 2002)⁶⁰**

Number of not related readmission for usual care	0.43
Mean difference for unrelated readmissions	0.38
Number of related readmission for usual care	0.27
Mean difference for related readmissions	-0.05
Length of hospital stay for not related readmissions (usual care)	4.9
Mean difference for length of hospital stay for unrelated readmissions	-0.3
Length of hospital stay for related readmissions (usual care)	3.6
Mean difference for length of hospital stay for related readmissions	0.1

18

19 **20.7.10.4 Social services costs**

20 For community services, Crotty et al (2002)⁶⁰ intended any of the following: outpatient
 21 rehabilitation; private therapy, district nursing, day care, respite care, employment rehabilitation
 22 training, carer time off work and Meals on Wheels. As we do not have data regarding the exact
 23 amount of resources for each of the above categories that were actually used by patients in the

1 two arms of the study, we assume that the weekly cost of social care is given by a weighted
 2 average of the five categories of packages of care reported in the PSSRU 2009⁶¹ and discussed in
 3 section 18.2.2 of the hospital MDR model. We assume that an equal proportion of patients used
 4 each type of social care package. However, in a sensitivity analysis we look at the case in which all
 5 patients used a “very low cost” type of social care package and when all of them used a “very high
 6 cost” package of care.

7 Only a proportion of the social care costs will generally be funded by local authorities^{71, 348, 72, 144}.
 8 It was not possible to identify a national average for the social care costs funded by local
 9 authorities in the published literature, and as a consequence an assumption had to be made
 10 regarding the proportion of this care that was publicly funded. In the base case analysis, we
 11 assume that 60% of social care costs are borne by local authorities, and are therefore includable
 12 in the model, and we then test this assumption in a sensitivity analysis.

13 As no further data were given regarding the use of social care services after the 4 months follow
 14 up, we adopted a conservative approach and assumed that after that period there was no
 15 difference in the use of social services that could be due to the different rehabilitation scheme
 16 used.

17 **20.7.10.5 Primary care costs**

18 Crotty et al (2002)⁶⁰ point out that: “[...] patients [in the community MDR scheme] tended to call
 19 the GPs if problems arose and this invariably meant a visit to the home for the GP” (Crotty et al
 20 2002, page 11⁶⁰). On the other hand, no details were provided regarding whether *all* GP visits to
 21 patients in the community MDR scheme took in fact place in the patients’ own home. Similarly,
 22 no information was given regarding where GP visit took place for patients in the usual care arm.
 23 As a consequence, we have assumed that the unit cost for a GP visit for patients in the usual care
 24 scheme is the average between the cost of a GP visit at the patient’s own home (£117) and a GP
 25 surgery visit (£76) as reported in the PSSRU 2009⁶¹, and corresponds to £96.5.

26 As no further data were given regarding the use of primary care services after the 4 months follow
 27 up, we adopted a conservative approach and assumed that after that period there was no
 28 difference in the use of GP services that could be ascribed to the different rehabilitation scheme
 29 used.

30

31 **Table 117: Evidence on GP visits (Crotty et al, 2002)**

Number of GP visits (usual care)	4.5
Number of GP visits (community MDR)	3.3
	(mean difference: -1.2)

32

33 **20.7.11 Cost effectiveness findings**

34 The cost-effectiveness findings for the deterministic base case analysis is presented in Table 118:
 35 Cost-effectiveness analysis - deterministic base case below:

36 **Table 118: Cost-effectiveness analysis - deterministic base case**

Strategy	Cost	Incremental	Effectiveness	Incremental	Incremental cost-
----------	------	-------------	---------------	-------------	-------------------

		Cost		Effectiveness	effectiveness ratio
Usual care	£6469.1		3.0827 QALYs		
Community MDR	£6903.2	£434.1	3.1283 QALYs	0.0456 QALYs	9521 £/QALYs

1

2 Hence, the community MDR scheme is a cost-effective treatment for the rehabilitation of hip
3 fracture patients in the deterministic case scenario. **Table 119** reports a breakdown of costs for
4 the relevant resources used in the community MDR and in the usual care group.

5 **Table 119: Cost breakdown for community MDR and usual care**

Resource item	Usual Care	Community MDR
Rehab cost	3456	3793
Readmission	1124	1657
Domiciliary social care	1453	1133
GP visits	434	318
Total cost	£6467	£6901

6

7 **20.7.11.1 Sensitivity analysis**

8 We now proceed by investigating how robust the findings of the deterministic analysis are by
9 conducting a series of sensitivity analysis.

10 To begin with we note that the model is not sensitive to changes in the level of social services paid
11 by the NHS (from 0 to 100%), as community MDR is still cost-effective.

12 Moreover, when the cost per week of social services is varied between the minimum (£41 per
13 week) and the maximum (£542) the option with the highest net benefit is still community MDR.

14 However, our findings are sensitive to the length of hospital stay (both for community MDR and
15 for usual care patients) and on the length of rehabilitation programme at home, as well as on the
16 daily cost of hospital stay following surgery and on the daily cost of the community MDR
17 programme. These findings are summarized in Table 120 below.

18

19 **Table 120: Threshold sensitivity analysis**

Variable	Values	Strategy with the highest net benefit
Length of stay in hospital for community MDR patients	≥ 9.78 (days)	Usual care
	< 9.78 (days)	Community MDR
Length of stay at home for community MDR patients	≥ 25.38 (days)	Usual care
	< 25.38 (days)	Community MDR
Length of stay in hospital for usual care patients	≥ 12.32 (days)	Community MDR
	< 12.32 (days)	Usual care
Daily cost for hospital stay	> £168.18	Community MDR


Daily cost for community MDR rehab programme	< £ 117.53	Community MDR
-----------------------------------------------------	------------	---------------

1

2 Our cost-effectiveness findings are not sensitive to changes in the cost per day in hospital of the
 3 related readmissions and to changes in the proportion of social care costs borne by local
 4 authorities.

5 The following figure summarise the findings of a two-ways sensitivity analysis on the length of
 6 stay in hospital and at home (the vertical axe reports the length of stay at home for community
 7 MDR patients and the horizontal axe the length of stay in hospital for usual care).

**Net Monetary Benefit (wtp=20000.) Sensitivity Analysis on
 LOS_community_MDR_at_home and LOS_in_hospital_usual_c**



8

Figure 166: Two-way sensitivity analysis on length of stay at home and in hospital

9

20.7.11.2 Probabilistic sensitivity analysis

12 A probabilistic sensitivity analysis was performed to assess the robustness of the model results to
 13 plausible variations in the model parameters.

14 Probability distributions were assigned to each model parameter, where there was some measure
 15 of parameter variability. We then re-calculated the main results 10000 times, and each time all
 16 the model parameters were set simultaneously, selecting from the respective parameter
 17 distribution at random.

**Table 121: Description of the type and properties of distributions used in the probabilistic
 19 sensitivity analysis**

Parameter	Type of distribution	Properties of distribution
-----------	----------------------	----------------------------

Baseline risk	Beta	Bounded on 0 – 1 interval. Derived from sample size, number of patients experiencing events
Cost	Gamma	Bounded at 0, positively skewed. Derived from mean and standard error
Utility	Beta	Bounded on 0 – 1 interval. Derived from mean and sample size
Risk ratio, length of stay	Lognormal	Bounded at 0. Derived from log and standard error of log
Mean differences (e.g. in length of stay, time of therapies, etc.)	Normal	Derived from mean and standard deviation

1

2 Table 122 summarises the expected values of the variables in our model from the different
3 distributions used in the PSA.

4 **Table 122: Distribution and parameters - probabilistic sensitivity analysis**

Name	Baseline value	Expected value	Distribution and parameters
Probability use of community services for usual care	0.72	0.7187	Beta, Integer parameters only, n = 32, r = 23
Weekly social care unit cost for “high need” patients (£)	216	216	Gamma, alpha = 15.36583528, lambda = 0.071138126
Weekly social care unit cost for “low need” patients	72	72	Gamma, alpha = 15.36583528, lambda = 0.213414379
Weekly social care unit cost for “median need” patients (£)	180	180	Gamma, alpha = 15.36583528, lambda = 0.085365752
Weekly social care unit cost for “very high need” patients (£)	542	542	Gamma, alpha = 15.36583528, lambda = 0.02835025
Weekly social care unit cost for “very low need” patients (£)	41	41	Gamma, alpha = 15.36583528, lambda = 0.37477647
NHS reference costs for community MDR (£; daily)	94	94	Gamma, alpha = 15.36583528, lambda = 0.163466333
NHS reference cost for usual care	240	240	Gamma, alpha = 15.36583528, lambda = 0.064024314
Mean difference in GP visits for community MDR	-1.2	-1.2	Normal, Mean = -1.2, Std Dev = 0.0957

Number of GP visits for usual care	4.5	4.5	Normal, Mean = 4.5, Std Dev = 0.646
Length of stay (days) at own home for community MDR programme	20.3	20.3	Log-Normal, μ (mean of logs) = 3.006198781, σ (std dev of logs) = 0.094043657
Length of stay (days) in hospital for community MDR patients	7.8	7.8	Log-Normal, μ (mean of logs) = 2.028128367, σ (std dev of logs) = 0.228014764
Length of stay (days) unrelated readmissions	4.9	4.9	Log-Normal, μ (mean of logs) = 1.264935055, σ (std dev of logs) = 0.80535725
Length of stay (days) related readmissions	3.6	3.6	Log-Normal, μ (mean of logs) = 1.061775585, σ (std dev of logs) = 0.662054772
Length of stay (days) in hospital – usual care	14.3	14.3	Log-Normal, μ (mean of logs) = 2.650611207, σ (std dev of logs) = 0.138912419
Probability mortality – community MDR	0.067	0.067	Beta, Real-numbered parameters, α = 2.278, β = 31.722
Probability mortality usual care	0.0724	0.0724	Beta, Real-numbered parameters, α = 2.31648, β = 29.68352
Proportion of patients with “very low”/”low”/”median”/”high”/”very high” social care costs	0.2	0.2	Dirichlet; Alpha list (proportion of patients with very low social care costs; proportion of patients with low social care costs; proportion of patients with median social care costs; proportion of patients with high social care costs; proportion of patients with very high social care costs)
Proportion of social care costs funded by the NHS	0.6	0.6	Triangular, Min = 0.30, Likeliest = 0.60, Max = 0.90;
EQ-5D score (community MDR)	0.732	0.732	Beta, Real-numbered parameters, α = 24.888, β = 9.112
EQ-5D score (usual care)	0.643	0.643	Beta, Real-numbered parameters, α = 20.576, β = 11.424
Number of not related	0.43	0.43	Normal, Mean = 0.43, Std Dev =

readmission for usual care			0.0617
Number of related readmission for usual care	0.27	0.27	Normal, Mean = 0.27, Std Dev = 0.387
Mean difference for length of hospital stay for related readmissions	0.1	0.1	Normal, Mean = 0.1, Std Dev = 0.0145
Mean difference for related readmissions	-0.05	-0.05	Normal, Mean = -0.05, Std Dev = 0.04
Mean difference for length of hospital stay for unrelated readmissions	-0.3	-0.3	Normal, Mean = -0.3, Std Dev = 0.03442
Mean difference for unrelated readmissions	0.38	0.38	Normal, Mean = 0.38, Std Dev = 0.545;
Relative Risk use of community service	0.78	0.78	Log-Normal, μ (mean of logs) = -0.265778098, σ (std dev of logs) = 0.186100721
Unit cost for a GP visit	76	76	Gamma, α = 15.36583528, λ = 0.202182043
Unit cost for related readmissions	352	352	Gamma, α = 15.36583528, λ = 0.043652941
Unit cost for unrelated readmissions	249	249	Gamma, α = 15.36583528, λ = 0.061710182

1

2 The cost-effectiveness findings of the PSA are summarized in Table 123 below:

3

4 **Table 123: cost-effectiveness finding from probabilistic sensitivity analysis**

Strategy	Cost	Incr Cost	Eff	Incr Eff	Incremental cost-effectiveness ratio	95% CI on ICERs
Usual care	£6466.6		3.0827 QALYs			
Community Rehab	£6901.2	£434.6	3.1283 QALYs	0.0456 QALYs	9533 £/QALYs	Cost saving - dominated

5

6 The PSA shows that there is a high uncertainty as to whether community MDR is cost-effective
7 compared to usual care. This uncertainty can be graphically represented by plotting the results of
8 the incremental analysis for all the 10,000 simulations into a cost-effectiveness plane. Each point
9 on the scatter plot represents the ICER of community MDR versus usual care for each simulation.

1 The dotted line represents the £20,000/QALY threshold while the ellipse delimits the 95%
2 confidence interval.

3

4 **Figure 167: Incremental cost-effectiveness scatter plot - Community MDR vs usual care**

5

6 From the simulations conducted for the PSA, we found that at a willingness to pay equal to
7 £20,000 per QALY, community MDR was the optimal strategy in 50% of the simulations. At a
8 willingness to pay of £30,000 per QALY, community MDR was the optimal strategy in 60% of the
9 simulations.

10

11 **Table 124: Probability most cost-effective intervention at a willingness to pay of £20,000 and**
12 **30,000 per QALY**

Strategy	Probability most cost-effective intervention at a WTP of £20,000 per QALY	Probability most cost-effective intervention at a WTP of £30,000 per QALY
Community MDR	0.50	0.60
Usual care	0.50	0.40

13

14 **20.7.12 Discussion**

15 The model shows that community MDR is cost-effective in the rehabilitation of patients after a
16 hip fracture. However, this finding is rather sensitive to variations in the length of stay, both in
17 hospital and at home. Moreover, a PSA has shown that there is high uncertainty over the cost-
18 effectiveness of community MDR compared to usual care.

19 The model has several limitations, such as the fact that it is based on the clinical evidence derived
20 from only one RCT⁶⁰ based in Australia. Moreover, the evidence on treatment effects⁶⁰ was
21 available only up to 4 months follow up. No information was available regarding the impact of
22 community MDR after that time point.

23

24 **20.8 Cost analysis of cemented vs. uncemented implants (newer designs of arthroplasty)**

25

26 In order to conduct a cost analysis for the cemented and uncemented implants, we need to consider the
27 following cost components: implants cost; accessories costs; length of stay; re-operation and theatre costs.
28

29 **a) Cost of implants**

30

1 The National Joint Registry (NJRv7) was accessed on February 11th 2011, in order to find out the five most
 2 commonly used types of both cemented (stems with no head) and uncemented implants in the UK.
 3 Furthermore, the NHS Supply catalogue 2011 was searched in order to obtain the most recent price for
 4 each of these items. All this information is reported in **Table 125** below:

5
6

Table 125: Price of new design cemented and uncemented stems most commonly used in the UK

Cemented implants – stems with no head (from most common to less common)	Price per item (£)	Uncemented implants (from most common to less common)	Price per item (£)
Exeter	410.53	Corail	893.47
C-Stem	347.37	Taperloc	Could not retrieve price on NHS supply catalogue
CPT	393.68	JRI Furlong	789.47
Stanmore	Could not retrieve price on NHS supply catalogue	Accolade	684.21
Charnely	Could not retrieve price on NHS supply catalogue	SL-Plus	789.47
Average price for most common cemented implants	383.86	Average price for most common uncemented implants	789.15

7
8
9
10
11
12
13

b) Accessories costs for the cemented group of patients

The price for accessories used when new design cemented stems are implanted are presented in **Table 126** below.

14 **Table 126: Prices for accessories used with new design cemented stems**

Resources	Unit price	Source of unit price
Pulse Lavage	£27.86	NHS Supply Chain catalogue 2011
Biogel glove	£1.07	NHS hospital*
Cement	£45	Data supplied by GDG member
Cement mixing kit	£35	Data supplied by GDG member
Cement Restrictor (size 16/18/20mm)	£35.88	Smith & Nephews**
Femoral canal brush (NW 12.5/19MM)	£64.48	Smith & Nephews**
MIXOR femoral pressurizer (large/medium/small)	£14.70	Smith & Nephews**

Sterilized tray	£25	NHS hospital record***
Total cost for accessories	£248.99	

- 1
- 2 *Peterborough and Stamford Hospital NHS Trust
- 3 **Price list available online at: [http://browse.uk-](http://browse.uk-plc.net/Companies/SMITH_NEPHEW/products/CEMENT_ACCESSORIES.htm)
- 4 [plc.net/Companies/SMITH_NEPHEW/products/CEMENT_ACCESSORIES.htm](http://browse.uk-plc.net/Companies/SMITH_NEPHEW/products/CEMENT_ACCESSORIES.htm)
- 5 (accessed 21st February 2011)
- 6 ***Data from John Radcliffe NHS Hospital Trust.

7 It is important to note that the cost of the sterilized tray could vary from £25 to £50 (source: John
 8 Radcliffe NHS Hospital Trust) depending on the number of instruments contained in the tray. It is
 9 also relevant to point out that our cost calculation for the accessories used with the new design
 10 of cemented stems is very similar to the cost reported in Unnanuntana et al (2009)³³⁵, which
 11 calculates the cost of accessories used for a third-generation cement technique. They considered
 12 two 40-g bags of bone cement without antibiotics, a vacuum mixing cartridge, cement pressurizer,
 13 canal plug and distal cement centralizer, canal brush and cement scrapers. The average total cost
 14 for the two 40-g batches of bone cement and all accessories used to achieve a third-generation
 15 cement technique was estimated to be \$386 (range, \$351-\$407) (January 2008 prices), which
 16 correspond to £252 (range, £229 - £266) (converted using 2008 purchasing power parity).
 17 The GDG noted that the accessories costs determined in table 126 represent a through end of
 18 the spectrum; at the “lower end” of the spectrum, the only accessories costs to consider would be
 19 those for the cement, cement mixing kit, restrictor and sterilized tray, for an overall cost of
 20 **£140.88**.

21

22 c) Cost of length of stay in hospital

23 To calculate the health state costs during the hospital stay, we use the NHS reference cost for excess bed
 24 days reported in Table 127 below. The excess bed day cost is the cost per day for days exceeding the
 25 trimpoint, a cut-off that determines patients with exceptionally long stay, and as such usually estimates the
 26 cost of care without the cost of procedures (i.e. without the cost of the surgery). These costs reflect the
 27 presence of complications experienced by hip fracture patients during their entire hospital stay. Moreover,
 28 they distinguish between “major” and “intermediate” complications, thus allowing users to take into
 29 account the different degrees of resource use.

30

31 **Table 127: National Schedule of Reference Costs Year : '2008-09' - NHS Trusts and PCTs combined Non-**
 32 **Elective Inpatient (Long Stay) Excess Bed Day HRG Data for hip procedures**

Currency Code	Currency Description	Activity	National Average Unit Cost
HA11A	Major Hip Procedures Category 2 for Trauma with Major CC	360	£243
HA11B	Major Hip Procedures Category 2 for Trauma with Intermediate CC	620	£242

HA11C	Major Hip Procedures Category 2 for Trauma without CC	162	£220
HA12B	Major Hip Procedures Category 1 for Trauma with CC	9,760	£237
HA12C	Major Hip Procedures Category 1 for Trauma without CC	1,230	£226
HA13A	Intermediate Hip Procedures for Trauma with Major CC	14,891	£240
HA13B	Intermediate Hip Procedures for Trauma with Intermediate CC	12,856	£249
HA13C	Intermediate Hip Procedures for Trauma without CC	2,972	£223
HA14A	Minor Hip Procedures for Trauma with Major CC	5,195	£234
HA14B	Minor Hip Procedures for Trauma with Intermediate CC	5,808	£245
Mean weighted average of excess bed days costs – NHS reference costs 2008-08 Major, Intermediate and Minor Hip procedures with all types of complications			£240

1

2 Using the evidence reported in Figved (2009)⁹⁴, the mean LOS in hospital for patients in the cemented
3 group was 7.8 days and in the uncemented group 8.4 ($p < 0.52$). This implies that the LOS costs for the
4 cemented group correspond to £1872 and for the uncemented group to £2016.

5

6 d) Re-operation costs

7 The cost of the re-operations in the two groups of patients is calculated by using the weighted average of
8 the NHS reference cost for non-elective inpatient short stay data for NHS Trusts and PCTs combined. The
9 different HRGs and unit costs associated with each type of surgical procedure and possible presence of
10 complications are summarised in **Table 128** below.

11 **Table 128: National Schedule of Reference Costs Year : '2008-09' - NHS Trusts and PCTs combined Non-**
12 **Elective Inpatient (Short Stay) HRG Data**
13

Currency Code	Currency Description	Activity	National Average Unit Cost
HB11A	Major Hip Procedures for non Trauma Category 2 with Major CC	4	£1,793
HB11B	Major Hip Procedures for non Trauma Category 2 with CC	5	£2,001
HB11C	Major Hip Procedures for non Trauma Category 2 without CC	3	£1,765
HB12A	Major Hip Procedures for non Trauma Category 1 with Major CC	16	£1,811
HB12B	Major Hip Procedures for non Trauma Category 1 with CC	89	£2,097
HB12C	Major Hip Procedures for non Trauma Category 1 without CC	88	£1,611
HB13Z	Intermediate Hip Procedures for non Trauma Category 2	51	£2,771
HB14B	Intermediate Hip Procedures for non Trauma Category 1 with CC	746	£1,671

HB14C	Intermediate Hip Procedures for non Trauma Category 1 without CC	529	£1,785
HB15D	Minor Hip Procedures for non Trauma Category 2 19 years and over with CC	213	£912
HB15E	Minor Hip Procedures for non Trauma Category 2 19 years and over without CC	124	£1,028
HB15F	Minor Hip Procedures for non Trauma Category 2 18 years and under with CC	8	£779
HB15G	Minor Hip Procedures for non Trauma Category 2 18 years and under without CC	73	£1,304
HB16B	Minor Hip Procedures for non Trauma Category 1 with CC	141	£2,057
HB16C	Minor Hip Procedures for non Trauma Category 1 without CC	111	£833
Mean weighted average of short-stay non-elective NHS reference costs 2009-10 Major, Intermediate and Minor Hip procedures with all types of complications			£1,598.39

1

2 Figved (2009)⁹⁴ reports a re-operation rate of 6.3% for the cemented group and of 7.4% for the uncemented
3 group. The re-operation costs therefore correspond to £100.70 in the cemented group and to £118.28 for
4 the uncemented group.

5 e) Theatre time costs

6 Figved (2009)⁹⁴ reports the duration of the operation for the cemented group which was 12.4 minutes
7 longer than for the uncemented group. Using a cost per minute for the theatre use of £20.50 (from
8 Peterborough and Stamford NHS Trust accountant data), the higher theatre costs for the cemented group
9 correspond to: £254.2

10

11 Summary of costs components

	Patients who received cemented implants	Patients who received uncemented implants
Cost categories:		
a) Implants	£383.86	£789.15
b) Accessories costs for cemented implants	£248.99	-
c) LOS	£1872	£2016
d) Re-operations	£100.70	£118.28
e) Incremental theatre costs for cemented group	£254.2	-
Total costs	£2859.75	£2923.43

12

13 It follows that the overall incremental cost of the newer design of uncemented implants over the cemented
14 ones £63.68. When the lower estimate for accessories costs is used (£140.88) the total costs for the

1 cemented group corresponds to £2751.64, and the incremental cost of the uncemented implants to
2 £171.79. These cost does not include the additional pain relief required by patients in the uncemented
3 group. However, the unit costs for analgesics is relatively low, as showed in Appendix H section 20.1.

4

5

6

21 Appendix I: High Priority Research Recommendations

21.1 Imaging options in occult hip fracture

Research question: In patients with a continuing suspicion of a hip fracture but whose radiographs are normal, what is the clinical and cost effectiveness of computed tomography (CT) compared to magnetic resonance imaging (MRI), in confirming or excluding the fracture?

Why this is important:

The GDG's consensus decision to recommend CT over a radionuclide bone scan as an alternative to MRI to detect occult hip fractures reflects current NHS practice but assumes that advances in technology have made the reliability of CT comparable with that of MRI. If modern CT can be shown to have similar reliability and accuracy to MRI, then this has considerable implications because of its widespread availability out of hours and lower cost. It is therefore a high priority to confirm or refute this assumption by direct randomised comparison. The study design would need to retain MRI as the 'gold standard' for cases of uncertainty and to standardise the criteria, expertise and procedures for radiological assessment. Numbers required would depend on the degree of sensitivity/specificity (the key outcome criteria) set as target requirement for comparability, but need not necessarily be very large.

Criteria for selecting high-priority research recommendations:

<p><u>PICO question</u></p> <p>Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>In patients with a continuing suspicion of a hip fracture but whose radiographs are normal, what is the clinical and cost effectiveness of computed tomography compared to magnetic resonance imaging, in confirming or excluding the fracture?</p> <p>Patient: patients with a continuing suspicion of a hip fracture but whose radiographs are normal</p> <p>Intervention: Modern Computed Tomography techniques e.g. 64-slice scanners with three dimensional capabilities and spiral multidetector CT (MDCT)</p> <p>Comparison: Magnetic resonance imaging</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Outcomes: Diagnostic accuracy including sensitivity and specificity
<u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality)?	The altered guidance would ensure the availability of accurate diagnosis out of hours and thus promote the benefits of prompt, accurate surgery to all patients in this group – prompt pain relief, lower mortality, enhanced return to independent living, fewer complications and shorter hospital stay.
<u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?	Demonstration of comparable sensitivity and specificity with MRI would enable CT techniques to be recommended as investigation of first choice in these circumstances.
<u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?	Avoiding delay to surgery in hip fracture is cost-effective. The altered guidance would support this objective. CT is in addition available at lower NHS cost than MRI.
<u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.	The question has a direct bearing on the Department of Health Best Practice Tariff initiative to achieve time-to-surgery not exceeding 36 hours.
<u>Current evidence base</u> What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.	There have been no studies comparing the sensitivity and specificity of modern multi-detector CT techniques with the current gold standard (MRI) in the diagnosis of hip fracture. See Section 5.5.1 of the Full Guideline.
<u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?	No specific equality issues.
<u>Study design</u> It should also specify the most appropriate	The research design of choice would be a two-stage design comprising (1) an initial small-scale prospective randomised trial to test an

<p>study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>agreed minimum percentage variability between methods followed (subject to outcome) by (2) a prospective cohort study using CT alone.</p>
<p><u>Feasibility</u></p> <p>Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>It should be possible to undertake both elements in a realistic timescale and at reasonable cost. This would not be the case if a full-scale Phase 3 trial (as distinct from a prospective cohort) were considered essential.</p> <p>It would be ethically necessary to retain the availability of MRI as opt-out gold standard throughout both studies.</p>
<p><u>Other comments</u></p> <p>Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>The ideal study would compare both CT and MRI in the same patients. This is, however, impractical. The proposed research design has some limitations, but does have the potential to provide useful evidence. The alternative of awaiting an “evolutionary” approach to progress in this area is less acceptable.</p>
<p><u>Importance</u></p> <p>How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline • Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates • Low: the research is of interest and will fill existing evidence gaps. 	<p>The research is of high priority, since its findings have the potential to alter future guidance on the diagnosis of occult hip fracture.</p>

1 21.2 **Anaesthesia**

2 Research question: What is the clinical and cost effectiveness of regional versus general
3 anaesthesia on postoperative morbidity in patients with hip fracture?

4 **Why this is important**

5 No recent randomised controlled trials were identified that fully address this question. The
6 evidence is old and does not reflect current practice. In addition, in most of the studies the
7 patients are sedated before regional anaesthesia is administered, and this is not taken into
8 account when analysing the results. The study design for the proposed research would be
9 best addressed by a randomised controlled trial. This would ideally be a multi-centre trial
10 including 3000 participants in each arm. This is achievable given that there are about
11 70,000 to 75,000 hip fractures a year in the UK. The study should have three arms that look
12 at spinal anaesthesia versus spinal anaesthesia plus sedation versus general anaesthesia;
13 this would separate those with regional anaesthesia from those with regional anaesthesia
14 plus sedation. The study would also need to control for surgery, especially type of fracture,
15 prosthesis and grade of surgeon. A qualitative research component would also be helpful to
16 study on patient preference for type of anaesthesia.

17 This needs to be multicentre and could be conducted in one year in the U.K. Sample size,
18 may need to have 3,000 in each limb which is achievable if one considers that there are 80,
19 000 hip fractures a year in the UK.

20 Criteria for selecting high-priority research recommendations:

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>What is the clinical and cost effectiveness of regional versus general anaesthesia on postoperative morbidity in patients with hip fracture? Patient: patients undergoing surgical repair for hip fractures Intervention: regional anaesthesia Comparison: general anaesthesia Outcome: postoperative morbidity</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>Improved survival following hip fracture. Improved analgesia following surgery. Reduced complications such as acute delirium, nausea and vomiting.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>The study may give the evidence to give better guidance to anaesthetists. There have been no studies comparing modern anaesthesia techniques in this group of patients. The current evidence is old and unreliable. The hip fracture population is now older and has more comorbidities than the population in which the historical studies were conducted.</p>

	<p>The studies are also important to help patients and their carers make informed decisions about the form of anaesthesia most appropriate for them.</p> <p>Importance : High</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>There may be a reduction in length of stay in patients receiving spinal anaesthesia, without sedation. Postoperative recovery should be quicker.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>SIGN recommend spinal but without any evidence base. The evidence for benefit is weak and was conducted over 30 years ago.</p>
<p><u>Current evidence base</u> What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>No trial evidence was identified</p>
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>This recommendation does not exclude any patient group. However, special consideration should be given to very frail older people with a high prevalence of cognitive impairment.</p>
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>The study design for the proposed research would be best addressed by an RCT. This would ideally have three arms (3000 participants each) which looks at spinal versus spinal plus sedation versus general anaesthesia, this would separate those with regional anaesthesia from those with regional anaesthesia plus sedation. The study would also need to control for surgery, especially type of fracture, prosthesis and grade of surgeon.</p> <p>A qualitative research component would also be helpful to study on patient preference for</p>

	type of anaesthesia.
<p><u>Feasibility</u></p> <p>Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>Although the number of participants suggested is relatively high, it is worth considering that there are over 80,000 patients admitted with hip fractures each year. This should be feasible by conducting a multi-centre RCT.</p>
<p><u>Other comments</u></p> <p>Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>Potential funders include : The National Institute for Health Research (NIHR), ASTRA foundation.</p>
<p><u>Importance</u></p> <p>How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline • Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates • Low: the research is of interest and will fill existing evidence gaps. 	<p>High. The research is essential to inform future updates of key recommendations in the guideline.</p>

1

2

21.3 Displaced intracapsular hip fractures

Research question:

What is the clinical and cost effectiveness of large head total hip replacement versus hemiarthroplasty on functional status, reoperations and quality of life in patients with displaced intracapsular hip fracture?

Why this is important:

Large-head total hip replacement is a development of traditional total hip replacement, where a larger head makes the joint more stable and hence reduces the risks of dislocation. Three small trials have shown traditional small-head total hip replacement to have better outcomes and function, albeit with an increased dislocation rate in selected groups of patients. The drawback with large-head arthroplasty is the additional implant cost and theatre time. This cost can account for up to 20% of current NHS tariff (up to £2000) and the study aims to address whether this translates to improved patient outcome. The study design for the proposed research would be best addressed by a randomised controlled trial. This would have two arms to compare current standard care (using hemiarthroplasty) with using large-head total hip replacement for patients sustaining displaced intracapsular hip fractures. The primary outcome would be patient mobility at 1 year and secondary outcomes would include functional outcomes, quality of life and cost effectiveness of the intervention.

It would be expected that a sample size of approximately 500 patients would be required to show a significant difference in the mobility, hip function and quality of life (assuming 80% power, $p < 0.05$). By recruiting through a trauma research network it is estimated that 10 centres would be able to recruit 20 patients per month (from 45 eligible patients) giving a recruitment period of 25 months.

Criteria for selecting high-priority research recommendations:

<p><u>PICO question</u></p> <p>Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>Question: What is the clinical and cost effectiveness of large head total hip replacement versus hemiarthroplasty on functional status, reoperations and quality of life in patients with displaced intracapsular hip fracture?</p> <p><i>Patients:</i> Patients sustaining displaced intracapsular hip fractures</p> <p><i>Intervention:</i> Arthroplasty</p> <p><i>Comparison:</i> Either hemiarthroplasty (half a hip replacement) or total hip replacement with a large head</p> <p><i>Outcome:</i> Timely functional status, cost effectiveness, re-operations and quality of life at one year</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>Presently there are over 30,000 who sustain a displaced intracapsular hip fracture per year in the United Kingdom. Whilst there is evidence that total hip replacement with a small femoral head gives some advantages in specific groups (3 small RCTs) the concern has been the risk of dislocations. The technology has advanced and it is now possible to perform large head (>36mm) total hip replacement which significantly reduces the risk of dislocation and may improve function. The drawback is the increased cost (between £1000 - £2000 or >10-20% of the tariff)</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>Presently the NICE recommendations recommend replacement arthroplasty and is only specific about a defined group of cognitively unimpaired, previously mobile and with no significant comorbidities. There is currently widespread practice in arthroplasty and there is an increase use of more expensive prosthesis. Surgeons are beginning to adopt large head technology without evidence of effectiveness, cost benefit or consideration of complication rates.</p> <p>This recommendation is considered high by the NICE Hip Fracture Development Group as the results of this study would advise NICE on future recommendations for the large and vulnerable group of patients</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>The NHS would be in a better position to focus resources on those in most need. Better function of the large head total hip replacement may reduce care costs in both the acute setting and rehabilitation.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>Improving the care of those suffering fragility fractures is a NHS priority. Hip fractures are the largest cost of this group and account for two thirds of all hospital days due to fractures and 87% of the costs (£385million 2007).</p>
<p><u>Current evidence base</u> What is the current evidence base? What are</p>	<p>One cohort study has been presented on large head total hip replacement and three previous RCTs on small head total hip replacements have</p>

<p>the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>been published</p>
<p><u>Equality</u></p> <p>Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>Yes, very frail older people with a high prevalence of cognitive impairment.</p>
<p><u>Study design</u></p> <p>It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p><i>Design:</i> A randomised controlled trial of displaced intracapsular fractures in previously mobile patients between hemiarthroplasty and large head total hip replacement.</p> <p><i>Outcome:</i> Does large head arthroplasty improve recovery of mobility one year after surgical management of displaced intracapsular hip fracture</p>
<p><u>Feasibility</u></p> <p>Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The research would be ethically and technically feasible.</p> <p>The research costs would need to be considered in the context that participants would still need treatment if outside a trial which would set the research costs into proper context and perspective.</p>
<p><u>Other comments</u></p> <p>Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>The National Institute for Health Research (NIHR) would be an appropriate funding source. Industry support would off lay excess implant costs</p>
<p><u>Importance</u></p> <p>How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline 	<p>High. The research is essential to inform future updates of key recommendations in the guideline.</p>

- | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| <ul style="list-style-type: none">• Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates• Low: the research is of interest and will fill existing evidence gaps. | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|

1

1 **21.4 Intensive rehabilitation therapies after hip fracture**

2 Research question:

3 What is the clinical and cost effectiveness of additional intensive physiotherapy and/or
4 occupational therapy (for example progressive resistance training) after hip fracture?

5 Why this is important:

6 The rapid restoration of physical and self care functions is critical to recovery from hip
7 fracture, particularly where the goal is to return to the patient to preoperative levels of
8 function and residence. Approaches that are worthy of future development and
9 investigation include progressive resistance training, progressive balance and gait training,
10 supported treadmill gait re-training, dual task training, and activities of daily living training.
11 The optimal time point at which these interventions should be started requires clarification.

12 The ideal study design is a randomised controlled trial. Initial studies may have to focus on
13 proof of concept and be mindful of costs. A phase III randomised controlled trial is required
14 to determine clinical effectiveness and cost-effectiveness. The ideal sample size will be
15 around, 400 to 500 patients, and the primary outcome should be physical function and
16 health related quality of life. Outcomes should also include falls. A formal sample size
17 calculation will need to be undertaken. Outcomes should be followed over a minimum of 1
18 year, and compare if possible, either the recovery curve for restoration of function or time
19 to attainment of functional goals.

20 Criteria for selecting high-priority research recommendations:

<p><u>PICO question</u></p> <p>Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p><i>Question:</i> What is the clinical and cost effectiveness of additional intensive physiotherapy and/or occupational therapy (for example progressive, resistance training) after hip fracture?</p> <p><i>Patients:</i> All patients who have a fracture, studies should consider all forms of surgical treatment. Separate studies maybe needed for those with severe cognitive impairment and those without (depending on specifics of the intervention)</p> <p><i>Intervention:</i> Progressive therapy protocols</p> <p><i>Comparison:</i> Usual care therapy</p> <p><i>Outcome:</i> Restoration of mobility, health related quality of life, falls, residence, ADL /IADL abilities, linked geriatric syndromes, resource use.</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for</p>	<p>Patients and their families value mobility very highly. The ability to walk even short distances, can mean the difference between being able</p>

<p>example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>to live at home, or not. The step between being able to walk outside and inside is greater still. The same can be said for key skills like dressing and bathing. The impact of improved mobility, strength, balance and function would have a substantial impact on the patient and their family, as well as the requirement for long term residential or at home care.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>It would enable NICE to come to a decision on whether to recommend more intensive physiotherapy and/or occupational therapy, by generating new evidence on clinical and cost effectiveness. This is very important to the guideline – at the moment we have made several statements about volume (frequency of therapy) but not of content. The guideline would be strengthened considerably with this additional information.</p> <p>The level would be high.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>There would possibly be an increase in the amount of therapy time that is needed, and this would incur impacts on strategic planning and service delivery.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>Yes the national service framework for older people.</p>
<p><u>Current evidence base</u> What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>Very limited trial evidence, and no trial evidence for some interventions.</p>
<p><u>Equality</u> Does the research recommendation address</p>	<p>Yes these are vulnerable older adults who need special consideration, particularly if they have cognitive impairment or frailty. These</p>

<p>equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>types of services are not currently provided to many hip fracture patients, and certainly not those with cognitive impairments</p>
<p><u>Study design</u></p> <p>It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p><i>Design:</i> A randomised controlled trial of intensive therapy (to be specified) versus usual care therapy</p> <p><i>Outcome:</i> Mobility, function, health related quality of life, resource use, and costs (health and social care)</p>
<p><u>Feasibility</u></p> <p>Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The research would be ethically and technically feasible.</p> <p>The outcome and research question is sufficiently important to merit a large scale randomised controlled trial</p>
<p><u>Other comments</u></p> <p>Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>The National Institute for Health Research (NIHR) HTA would be an appropriate funding source.</p>
<p><u>Importance</u></p> <p>How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline • Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates • Low: the research is of interest and will fill existing evidence gaps. 	<p>High. The research is essential to inform future updates of key recommendations in the guideline.</p>

1 **21.5 Early Supported Discharge in Care Home patients**

2 Research question:

3 What is the clinical and cost effectiveness of early supported discharge on mortality, quality
4 of life and functional status in patients with hip fracture who are admitted from a care
5 home?

6 Why this is important:

7 Residents of care and nursing homes account for about 30% of all patients with hip fracture
8 admitted to hospital. Two-thirds of these come from care homes and the remainder from
9 nursing homes. These patients are frailer, more functionally dependent and have a higher
10 prevalence of cognitive impairment than patients admitted from their own homes. One-
11 third of those admitted from a care home are discharged to a nursing home and one-fifth
12 are readmitted to hospital within 3 months. There are no clinical trials to define the optimal
13 rehabilitation pathway following hip fracture for these patients and therefore represent a
14 discrete cohort where the existing meta-analyses do not apply. As a consequence, many
15 patients are denied structured rehabilitation and are discharged back to their care home or
16 nursing home with very little or no rehabilitation input.

17 Given the patient frailty and comorbidities, rehabilitation may have no effect on clinical
18 outcomes for this group. However, the fact that they already live in a home where they are
19 supported by trained care staff, clearly provides an opportunity for a systematic approach
20 to rehabilitation. Early multidisciplinary rehabilitation based in care homes or nursing
21 homes would take advantage of the day-to-day care arrangements already in place and
22 provide additional NHS support to deliver naturalistic rehabilitation, where problems are
23 tackled in the patient's residential.

24 Early supported multidisciplinary rehabilitation could reduce hospital stay, improve early
25 return to function, and affect both readmission rates and the level of NHS-funded nursing
26 care required.

27 The research would follow a two-stage design: (1) an initial feasibility study to refine the
28 selection criteria and process for reliable identification and characterisation of those
29 considered most likely to benefit, together with the intervention package and measures for
30 collaboration between the Hip Fracture Programme team, care-home staff and other
31 community-based professionals, and (2) a cluster randomized controlled comparison (with
32 two or more intervention units and matched control units) set against agreed outcome
33 criteria. The latter should include those specified above, together with measures of the
34 impact on care-home staff activity and cost, as well as qualitative data from patients on
35 relevant quality-of-life variables.

36

1 Criteria for selecting high-priority research recommendations:

<p><u>PICO question</u></p> <p>Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome).</p>	<p><i>Patients:</i> Elderly hip fracture patients admitted from a care/nursing home</p> <p><i>Intervention:</i> Structured multidisciplinary rehabilitation</p> <p><i>Comparison:</i> Standard care</p> <p><i>Outcome:</i> Reduction in hospital LOS, short and long-term functional improvement, reduction in readmission to hospital, reduction in upgrade from care to nursing home dependency.</p>
<p><u>Importance to patients or the population.</u></p> <p>What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>Reduced dependency</p>
<p><u>Relevance to NICE guidance</u></p> <p>How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>The answer to this question is key to guidance on early supported discharge in hip fracture patients admitted from care home, who represent a significant proportion of patients</p> <p>With this information available NICE would be in a position to recommend early supported discharge in this group of patients.</p> <p>Importance : High</p>
<p><u>Relevance to the NHS</u></p> <p>What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>Reduction in hospital LOS will allow greater efficiency with respect to usage of trauma beds.</p> <p>Reduction in re-admissions, upgraded dependency to nursing homes represent significant cost savings</p>
<p><u>National priorities</u></p> <p>Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>A number of national guidelines now recommend the need for research in care home patients following hip fracture (SIGN, Orthopaedic Blue Book, NIHR HTA review, Cochrane review).</p>
<p><u>Current evidence base</u></p> <p>What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research</p>	<p>No trial evidence was identified.</p>

<p>required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	
<p><u>Equality</u></p> <p>Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>Yes, very frail older people with a high prevalence of cognitive impairment.</p>
<p><u>Study design</u></p> <p>It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>This will comprise: a systematic literature review, focusing on rehabilitation in care homes; a qualitative interview study with care home residents, their families, care home staff, allied health professionals and inpatient orthopaedic staff regarding discharge planning and rehabilitation for these patients; and an evaluation of a pilot early supported multidisciplinary rehabilitation service compared to usual care.</p>
<p><u>Feasibility</u></p> <p>Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The research would be ethically and technically feasible, at an acceptable level of cost.</p>
<p><u>Other comments</u></p> <p>Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>Potential funders include :The National Institute for Health Research (NIHR), BUPA, Alzheimer's Society</p>
<p><u>Importance</u></p> <p>How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in 	<p>High. The research is essential to inform future updates of key recommendations in the guideline.</p>

the guideline

- Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates
- Low: the research is of interest and will fill existing evidence gaps.

1 22 Appendix J: Excluded studies

Chapter	Study ID	Reasons for exclusion
Diagnosis	Lubovsky et al (2005) ¹⁹⁶	The trial was excluded because of the very small sample size. Only 13 patients included and only 6 patients received CT and MRI. The results were reported in a way that did not allow calculations of sensitivity and specificity.
Timing of surgery	Davis et al (1988) ⁶⁸	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Franzo et al (2005) ¹⁰¹	No clear explanation of adjustment and no baseline characteristics for each group.
Timing of surgery	Gdalevich et al (2004) ¹⁰⁸	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Hoening et al (1997) ¹⁴⁶	Not only surgical delay investigated, unable to extract raw data.
Timing of surgery	Kenzora et al (1984) ¹⁷⁷	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Mackenzie wt al (2006) ¹⁹⁸	Letter/short correspondence.
Timing of surgery	McGuire et al (2004) ²⁰⁹	The aim of the study is on day of the week of admission.
Timing of surgery	Moran et al (2005) ²¹⁵	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Novack et al (2007) ²⁴³	Adjusted hazard ratios given.
Timing of surgery	Rae et al (2007) ²⁸⁰	Baseline characteristics not given for each group.
Timing of surgery	Rogers et al (1995) ²⁹⁰	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Sebestyen et al (2008) ³⁰²	No adjustment for comorbidity.
Timing of surgery	Shabat et al (2003) ³⁰⁴	Inadequate methodology.
Timing of surgery	Sircar et al (2007) ³¹²	No baseline characteristics, no adjustment for comorbidity.
Timing of surgery	Sund & Liski (2005) ³²²	Adjusted odds ratios for provider characteristics.
Analgesia	Gorodetskyi et al (2007) ¹²²	Not a study of nerve blocks.

Analgesia	Mannion et al (2005) ²⁰²	No 'control' group without the nerve block.
Analgesia	Marhofer et al (1998) ²⁰⁵	No 'control' group without the nerve block.
Analgesia	Mutty et al (2007) ²²¹	No proximal femoral fractures included.
Analgesia	Piangatelli et al (2004) ²⁷²	No 'control' group without the nerve block.
Analgesia	Schiferer et al (2007) ³⁰¹	Inclusion of participants with other conditions. The trialists were unable to provide separate results for only the hip fracture participants.
Analgesia	Turker et al (2003) ³³²	No 'control' group without the nerve block.
Analgesia	Van Leeuwen et al (2000) ³⁴²	No 'control' group without the nerve block.
Anaesthesia	Alonso Chico et al (2003) ⁶	Not a trial of different types of anaesthesia but a comparison of different drugs within one form of anaesthesia.
Anaesthesia	Barna (1981) ¹³	No randomisation of patients.
Anaesthesia	Ben-David et al (2000) ¹⁶	Not a trial of different types of anaesthesia but a comparison of different drugs within one form of anaesthesia.
Anaesthesia	Coleman et al (1988) ⁵⁶	The study was excluded as it involved a change in the types of drugs used only, not a change in the method of anaesthesia.
Anaesthesia	Critchley et al (1995) ⁵⁷	Not a trial of different types of anaesthesia but a comparison of different drugs within one form of anaesthesia.
Anaesthesia	Darling et al (1994) ⁶⁴	The study was excluded as it was not felt relevant to this review as no clinical outcomes were reported.
Anaesthesia	Dyson et al (1988) ⁷⁵	Lack of outcome data for the anaesthesia comparison.
Anaesthesia	El-Zahaar et al (1995) ^{79,334}	This trial was excluded because separate results for patients having surgery for a hip fracture were not presented.
Anaesthesia	Favarel-Garrigues et al (1996) ⁹²	The trial was excluded as it was not considered a comparison of different forms of anaesthesia, only of a modification of anaesthetic technique.
Anaesthesia	Hemmingsen & Nielsen (1991) ¹⁴³	Not a trial of different types of anaesthesia but a comparison of different drugs within one form of anaesthesia.
Anaesthesia	Marhofer et al (1999) ²⁰⁴	Not a comparison of anaesthetic methods.
Anaesthesia	Matot et al (2003) ²⁰⁷	Compared techniques outside the scope of this review.
Anaesthesia	Maurette et al (1993) ²⁰⁸	The trial was excluded as it was a trial of different drugs with the same anaesthetic technique, not a comparison of different types of anaesthesia.
Anaesthesia	Naja et al (2000) ²²³	No randomisation of patients.
Anaesthesia	Nishikawa et al (2002) ²⁴²	Not a comparison of different types of anaesthesia.

Anaesthesia	Owen & Hutton (1982) ²⁵²	Not a comparison of anaesthetic techniques.
Anaesthesia	Sinclair et al (1997) ³¹¹	Not a comparison of different types of anaesthesia.
Anaesthesia	Sutcliffe & Parker (1994) ³²³	No randomisation of patients.
Anaesthesia	Tonczar & Hammerle (1981) ³²⁹	The study was excluded as it involved a neuroleptic anaesthesia and the only outcome measures were plasma catecholamines, cortisol, blood pressure and changes in heart rate.
Anaesthesia	Ungemach (1987) ³³³	The trial was excluded as it was a comparison of different drugs within one type of anaesthesia (general anaesthesia) and not a comparison of different anaesthetic techniques.
Surgeon Seniority	Claque et al (2002) ⁵²	Retrospective study, unclear if adjusted for confounders. Not stated how patients were allocated to surgeons.
Surgeon Seniority	Englesbe et al (2009) ⁸²	Compares outcomes at time when new trainees start compared to other times of the year. Not about surgeon seniority.
Surgeon Seniority	Evans et al (1979) ⁸⁷	No results or data for surgeon seniority analysis.
Surgeon Seniority	Faraj & Drakau (2007) ⁹⁰	No adjustment for confounders and no indication of how patients were allocated to surgeons.
Surgeon Seniority	Fung et al (2007) ¹⁰⁴	No outcome of interest.
Surgeon Seniority	Giannoudis et al (1998) ¹¹¹	No outcome of interest.
Surgeon Seniority	Grimley et al (1980) ¹²⁶	Compares hospitals outcomes rather than surgeon seniority. Unclear if retrospective or prospective. No indication of how patients were allocated to surgeons.
Surgeon Seniority	Harper & Walsh (1985) ¹³⁵	Unclear if retrospective or prospective, no adjustment for confounders.
Surgeon Seniority	Holmberg et al (1987) ¹⁴⁹	Unclear if retrospective or prospective, no adjustment for confounders.
Surgeon Seniority	Holt et al (1994) ¹⁵¹	No adjustment for confounders.
Surgeon Seniority	Levi & Gebuhr (2000) ¹⁹²	Unclear if retrospective or prospective, no adjustment for confounders, no outcomes measured by surgeon seniority only reports in words there was no difference between registrars and consultants.
Surgeon Seniority	Kukla et al (2001) ¹⁸¹	Unclear if retrospective or prospective. Examines years of experience but inexperienced surgeons were supervised. Results presented as a continuous variable.
Surgeon Seniority	Parker et al (1994) ²⁷¹	Not surgeon seniority, investigates the use of a special "Hip Fracture Team".
Surgeon Seniority	Sarvilinna et al (2002) ²⁹⁹	Retrospective study, no adjustment for

		confounders.
Surgeon Seniority	Sehat et al (2006) ³⁰³	Not about surgeon seniority.
Surgeon Seniority	Weinrauch (2006) ³⁵⁰	Not stated how patients were assigned to surgeons. Not stated the total number of surgeons involved nor how many involved in each category. Does not adjust for any confounders.
Internal fixation vs arthroplasty	Bhandari et al (2003) ²¹	Systematic review, used Cochrane review instead.
Internal fixation vs arthroplasty	Bjorgul et al (2006) ²⁵	Non-randomised study.
Internal fixation vs arthroplasty	Bray et al (1988) ³⁵	Excluded from Cochrane review due to inadequate randomisation. Patients were allocated according to day of week and surgeon preference. In addition to the low numbers recruited five were lost to follow-up.
Internal fixation vs arthroplasty	El-Abed et al (2005) ⁷⁸	Excluded from Cochrane review as non-randomised study, type of procedure used was by the preference of the attending surgeon on the day of admission.
Internal fixation vs arthroplasty	Gjertsen et al (2010) ¹¹⁸	Non-randomised study.
Internal fixation vs arthroplasty	Haentjens et al (2005) ¹³⁰	Non-randomised study.
Internal fixation vs arthroplasty	Heetveld et al (2009) ¹⁴²	Non-randomised study.
Internal fixation vs arthroplasty	Hunter (1974) ¹⁵⁶	Excluded from Cochrane review as non-randomised study.
Internal fixation vs arthroplasty	Hunter (1969) ¹⁵⁵	Excluded from Cochrane review as non-randomised study.
Internal fixation vs arthroplasty	Neander (2000) ²³⁸	Excluded from Cochrane review due to inadequate randomisation procedure. The first 20 patients were randomised with closed envelopes but the last 80 were allocated according to the day of week they were admitted (Monday to Thursday total hip replacement, Friday to Sunday reduction and fixation).
Internal fixation vs arthroplasty	Parker (1992) ²⁶¹	Excluded from Cochrane review as non-randomised study.
Internal fixation vs arthroplasty	Riley (1978) ²⁸⁵	Excluded from Cochrane review as study provided no adequate data.
Internal fixation vs arthroplasty	Rodriguez et al (1987) ²⁸⁹	Excluded from Cochrane review as non-randomised study.
Internal fixation vs arthroplasty	Rogmark & Johnell (2006) ²⁹²	Systematic review, used Cochrane review instead.
Internal fixation vs arthroplasty	Sikorski & Barrington (1981) ³⁰⁹	This comparison excluded from Cochrane review due to poor methodological quality.
Internal fixation vs arthroplasty	Stewart (1984) ³²¹	Excluded from Cochrane review as non-randomised study.

Internal fixation vs arthroplasty	Wang et al (2009) ³⁴⁷	Systematic review, used Cochrane review instead.
Hemiarthroplasty vs total hip replacement	Goh et al (2009) ¹²⁰	Systematic review, used Cochrane review instead.
Hemiarthroplasty vs total hip replacement	Haentjens et al (2005) ¹³⁰	Non-randomised study.
Hemiarthroplasty vs total hip replacement	Heetveld et al (2009) ¹⁴²	Non-randomised study.
Hemiarthroplasty vs total hip replacement	Kavcic et al (2006) ¹⁷²	Methodology not reported. Only mentions patients were randomly selected. No indication of allocation concealment, method of randomisation, blinding, or inclusion/exclusion criteria.
Cement	Ahn et al (2008) ²	Systematic review that includes randomised and non-randomised studies. Used Cochrane review.
Cement	Bajammal et al (2008) ¹⁰	Systematic review of cement use in appendicular fractures, not just hip fractures. Used Cochrane review.
Cement	Christie et al (1994) ⁵⁰	Excluded from Cochrane review as biometric study with no clinical outcome measures. No methods given for RCT, no outcomes from our protocol.
Cement	Clark et al (2001) ⁵³	Excluded from Cochrane review as non-randomised study.
Cement	Dorr et al (1986) ⁷³	Cemented vs uncemented hemiarthroplasty not a randomised comparison
Cement	Faraj & Branfoot (1999) ⁸⁹	Excluded from Cochrane review as non-randomised study, use of cement was at operating surgeon's preference.
Cement	Field & Rushton (2005) ⁹³	Excluded from Cochrane review because of a limited number of cases using what is at present an experimental new cup.
Cement	Georgescu et al (2004) ¹⁰⁹	Excluded from Cochrane review because of a lack of reported results within the conference abstract
Cement	Gierer et al (2002) ¹¹²	Excluded from Cochrane review as non-randomised study, use of cement was at operating surgeon's preference.
Cement	Graf et al (2000) ¹²³	Excluded from Cochrane review as non-randomised study.
Cement	Johnson et al (2001) ¹⁶⁴	Excluded from Cochrane review as non-randomised study.
Cement	Karpmann et al (1992) ¹⁷⁰	Excluded from Cochrane review as there was inadequate reporting of the trial. Attempts were made to contact the trialists for further information, without

		success.
Cement	Khan et al (2002) ¹⁷⁸	Systematic review, excluded as used Cochrane review instead.
Cement	Lachiewicz et al (2008) ¹⁸⁴	Elective hip replacement patients, not hip fracture patients.
Cement	Leidinger et al (2002) ¹⁹⁰	Excluded from Cochrane review as variations of cementing technique are not part of the protocol
Cement	Pitto et al (2000) ²⁷³	Excluded from Cochrane review as small numbers and only outcome measure is transoesophagal echocardiography shown embolism. No methods given for RCT, no outcomes from our protocol.
Cement	Sadr & Arden (1977) ²⁹⁵	Excluded from Cochrane review as unclear whether randomised, the use of Proplast coated prosthesis is no longer prevalent, small study of 40 patients with limited reporting of outcomes for the 25 assessed patients at follow up.
Cement	Vochteloo et al (2009) ³⁴⁵	Protocol for a randomised study, study not completed.
Surgical approach to hemiarthroplasty	Barden et al (2001) ¹²	Excluded from Cochrane review as not a comparison of different surgical approaches.
Surgical approach to hemiarthroplasty	Cashman & Cashman (2008) ⁴⁷	Elective hip replacement patients.
Surgical approach to hemiarthroplasty	Chan & Hoskin (1975) ⁴⁹	No adjustment for confounders.
Surgical approach to hemiarthroplasty	Enocson et al (2009) ⁸³	About total hip replacement.
Surgical approach to hemiarthroplasty	Enocson et al (2010) ⁸⁴	Possible double counting of the included study ENOCSON 2008. No adjustment for confounders.
Surgical approach to hemiarthroplasty	Keene et al (1993) ¹⁷⁵	Not about surgical approach.
Surgical approach to hemiarthroplasty	Lafosse et al (2007) ¹⁸⁷	About a minimally invasive approach.
Surgical approach to hemiarthroplasty	Lafosse et al (2007) ¹⁸⁶	About a minimally invasive approach.
Surgical approach to hemiarthroplasty	Lafosse et al (2008) ¹⁸⁵	About a minimally invasive approach.
Surgical approach to hemiarthroplasty	Unwin & Thomas (1994) ³³⁶	No adjustment for confounders.
Surgical approach to hemiarthroplasty	Wang et al (2010) ³⁴⁶	About a minimally invasive approach.
Surgical approach to hemiarthroplasty	Yang et al (2010) ³⁵⁶	About a minimally invasive approach.
Surgical approach to hemiarthroplasty	Widman & Isacson (2001) ³⁵³	Excluded from Cochrane review as not a comparison of different surgical approaches.
Screws/nails	Baumgaertner et al	No relevant outcomes.

	(1998) ¹⁵	
Screws/nails	Benum et al (1994) ¹⁸	Abstract only.
Screws/nails	Butt et al (1995) ⁴⁰	Does not meet our inclusion criteria: includes trochanteric and subtrochanteric combined.
Screws/nails	Davis et al (1988) ⁶⁷	Does not meet our inclusion criteria: includes trochanteric and associated subtrochanteric combined.
Screws/nails	Dujardin et al (2001) ⁷⁴	Experimental nail not used commercially.
Screws/nails	Kuwabara et al (1998) ¹⁸³	Unable to obtain paper.
Screws/nails	Lee et al (2007) ¹⁸⁸	Does not meet our inclusion criteria: all high energy trauma (subtrochanteric fractures).
Screws/nails	Mehdi et al (2000) ²¹²	Abstract only.
Screws/nails	Michos et al (2001) ²¹³	Abstract only.
Screws/nails	Mott et al (1993) ²¹⁷	Abstract only.
Screws/nails	Pahlpatz & Langius (1993) ²⁵³	Does not meet our inclusion criteria: Includes trochanteric and subtrochanteric fractures combined.
Screws/nails	Rahme & Harris (2007) ²⁸¹	Does not meet our inclusion criteria: all high energy trauma (subtrochanteric).
Surgical procedures (economic evidence)	Giancola et al (2008) ¹¹⁰	No cost figures were reported.
Surgical procedures (economic evidence)	Gill & Ursic (2007) ¹¹⁴	Inadequate methodological design and limited applicability to the UK NHS.
Surgical procedures (economic evidence)	Kim et al (2005) ¹⁷⁹	Proximal femoral nail compared to long-stem cementless calcar-replacement prosthesis (not an included intervention).
Surgical procedures (economic evidence)	Marinelli et al (2008) ²⁰⁶	Inadequate methodology.
Surgical procedures (economic evidence)	Rogmark et al (2003) ²⁹¹	The study does not distinguish patients on the basis of whether they received hemiarthroplasty or total hip replacement.
Mobilisation	Binder et al (2004) ²⁴	The comparison is not versus usual care.
Mobilisation	Galea et al (2008) ¹⁰⁶	The comparison is not versus usual care, both have a targeted plan.
Mobilisation	Graham (1968) ¹²⁴	The intervention is weight bearing at 2 weeks or 12 weeks. Not relevant to our review question.
Mobilisation	Mangione et al (2005) ²⁰¹	The comparison is not versus usual care.
Mobilisation	Resnick et al (2007) ²⁸³	Does not answer our review question: augmented mobilisation vs. usual care.
Mobilisation	Tsauo et al (2005) ³³⁰	Does not answer our review question: community mobilisation vs. usual care.
Mobilisation	Yu-yahiro et al (2009) ³⁵⁹	Does not answer our review question: community mobilisation vs usual care.
MDR	Fordham et al (1986) ⁹⁶	Discussion paper with a cost benefit analysis
MDR	Giusti et al (2006) ¹¹⁷	Does not meet our inclusion criteria for MDR team: medicine; nursing;

		physiotherapy; occupational therapy; and social care. Additional components may include: nutrition; pharmacy; and clinical psychology.
MDR	Gonzalez-Montalvo et al (2010) ¹²¹	Mixed intervention, acute orthogeriatric unit model, plus early surgery.
MDR	Ho et al (2009) ¹⁴⁵	Letter to editor.
MDR	Holt et al (2010) ¹⁵²	Does not meet our inclusion criteria: no outcomes reported that were prioritised in our protocol. Survival analysis rather than mortality.
MDR	Iliffe et al (2010) ¹⁵⁹	Protocol only, not full results.
MDR	Kuisma (2002) ¹⁸⁰	Does not meet our inclusion criteria for MDR team: medicine; nursing; physiotherapy; occupational therapy; and social care. Additional components may include: nutrition; pharmacy; and clinical psychology.
MDR	O’Cathain (1994) ²⁴⁵	Observational study.
MDR	Olsson et al (2007) ²⁴⁸	Does not meet our inclusion criteria for MDR team: medicine; nursing; physiotherapy; occupational therapy; and social care. Additional components may include: nutrition; pharmacy; and clinical psychology.
MDR	Pryor & Williams (1989) ²⁷⁵	Observational study.
MDR	Richards et al (1998) ²⁸⁴	Mixed population, only 31% hip fracture patients.
MDR	Ryan et al (2006) ²⁹³	Does not answer our review question. Intervention is intensity of multidisciplinary rehab (intensive: 6 or more face-to-face sessions per week from MDR team vs. less intensive: 3 or less face-to-face sessions per week).
MDR	Shyu et al (2010) ³⁰⁷	Reports 2 year follow up. 1 year data already included, which is the longest time point stated in our protocol.
MDR	Uy et al (2008) ³³⁸	Very low number of patients. N = 11
Hospital MDR (economic evidence)	Cameron et al (2000) ⁴¹	The studies included in the HTA were grouped in a different way to that considered for our clinical review, and therefore its cost analysis was not applicable for our review question.
Community MDR (economic evidence)	Coast et al (1998) ⁵⁵	Mixed population with only 31% hip fracture patients.
Community MDR (economic evidence)	Van Balen et al (2002) ³⁴⁰	Patients in the early supported discharge scheme were only discharged to a nursing home with rehabilitation facilities and not to their own home.

Patient views	Boutin-Lester & Gibson (2002) ³²	Only 1 / 5 of the patients had HF. This patient also had osteoporosis.
Patient views	Closs & Briggs (2002) ⁵⁴	Words used by patients to describe pain, not hip fracture patients only.
Patient views	Franchignoni (2002) ¹⁰⁰	Only 5/55 patients had hip fracture.
Patient views	Gjertsen et al (2008) ¹¹⁹	Not qualitative research into patient views.
Patient views	Hallstrom et al (2000) ¹³¹	7/9 patients had cervical fractures.
Patient views	Harrison (2006) ¹³⁸	Very brief summary of MSc thesis, unable to obtain a copy of thesis.
Patient views	Hedman et al (2008) ¹⁴¹	Compares level of care received between cognitively impaired and cognitively intact hip fracture patients in two Swedish hospitals.
Patient views	Huang & Acton (2009) ¹⁵⁴	Patient views about the period after discharge from rehabilitation in Taiwan.
Patient views	Lin & Lu (2005) ¹⁹⁴	Caregivers views after discharge from hospital not patient views.
Patient views	Lin (2006) ¹⁹³	Not a patient view study.
Patient views	Magasi et al (2009) ¹⁹⁹	About choice of a rehabilitation facility in the US, not applicable to UK.
Patient views	Resnick et al (2005) ²⁸²	Patient views on a specific exercise programme adopted at a centre in the USA.
Patient views	Robinson (1999) ²⁸⁷	Patient views about adapting to life after rehabilitation.
Patient views	Smith et al (1997) ³¹⁵	Review of article on report about patient views on discharge information. Unable to obtain a copy of full report with qualitative research.
Patient views	Webster (1976) ³⁴⁹	Not qualitative research of patient views.
Patient education	Allegrante et al (2007) ⁵	Not patient education intervention alone.
Patient education	Bhandari & Tornetta (2004) ²²	About which way of communicating risk ratios to patients.
Patient education	Elinge et al (2003) ⁸⁰	Group learning programme started 3 months after fracture.
Patient education	Gill & Ursic (1994) ¹¹⁵	Education for nurses not patients.
Patient education	Jackson (2010) ¹⁶⁰	Education intervention for healthcare professionals not patients.
Patient education	Tappen et al (2003) ³²⁶	Effect of video intervention of recovery from hip surgery. Unclear how patients were allocated to interventions.
Patient education	Yoon et al (2008) ³⁵⁷	Non-randomised study.

1 Bibliography

2 Bibliography

- 3
4 1. Adams HA, Wolf C, Michaelis G, Hempelmann G. Postoperative course and endocrine
5 stress reaction of geriatric patients with para-articular hip fractures. Prospective
6 randomized study comparing spinal anesthesia and halothane intubation narcosis.
7 *Anästhesie, Intensivtherapie, Notfallmedizin* 1990, **25**(4):263-70. (Guideline Ref ID:
8 ADAMS1990)
- 9 2. Ahn J, Man LX, Park S, Södl JF, Esterhai JL. Systematic review of cemented and
10 uncemented hemiarthroplasty outcomes for femoral neck fractures. *Clinical*
11 *Orthopaedics and Related Research* 2008, **466**(10):2513-8. (Guideline Ref ID: AHN2008)
- 12 3. Ahrengart L, Tornkvist H, Fornander P, Thorngren KG, Pasanen L, Wahlstrom P *et al.* A
13 randomized study of the compression hip screw and Gamma nail in 426 fractures.
14 *Clinical Orthopaedics and Related Research* 2002,(401):209-22. (Guideline Ref ID:
15 AHRENGART2002)
- 16 4. Al-Ani AN, Samuelsson B, Tidermark J, Norling A, Ekstrom W, Cederholm T *et al.* Early
17 operation on patients with a hip fracture improved the ability to return to independent
18 living. A prospective study of 850 patients. *Journal of Bone & Joint Surgery - American*
19 *Volume* 2008, **90A**(7):1436-42. (Guideline Ref ID: ALANI2008)
- 20 5. Allegrante JP, Peterson MG, Cornell CN, Mackenzie CR, Robbins L, Horton R *et al.*
21 Methodological challenges of multiple-component intervention: lessons learned from a
22 randomized controlled trial of functional recovery after hip fracture. *Hss Journal* 2007,
23 **3**(1):63-70. (Guideline Ref ID: ALLEGRANTE2007)
- 24 6. Alonso Chico A, Cruz Pardos P, Alvarez Grau J, Pachoco Jimenez A, Arregui Martinez de
25 Lejarza M, Sanchez Garcia ML *et al.* Comparison of the hemodynamic response in
26 subarachnoid anesthesia with bupivacaine versus bupivacaine with fentanyl in
27 traumatology surgery in elderly patients. *Revista Espanola de Anestesiologia y*
28 *Reanimacion* 2003, **50**(1):17-22. (Guideline Ref ID: ALONSOCHICO2003)
- 29 7. Ara R, Brazier J. Deriving an algorithm to convert the eight mean SF-36 dimension scores
30 into a mean EQ-5D preference-based score from published studies (where patient level
31 data are not available). *Value in Health* 2008, **11**(7):1131-43. (Guideline Ref ID:
32 ARA2008)

- 1 8. Archibald G. Patients' experiences of hip fracture. *Journal of Advanced Nursing* 2003,
2 **44**(4):385-92. (Guideline Ref ID: ARCHIBALD2003)
- 3 9. Aune AK, Ekeland A, Odegaard B, Groggaard B, Alho A. Gamma nail vs compression screw
4 for trochanteric femoral fractures. 15 reoperations in a prospective, randomized study
5 of 378 patients. *Acta Orthopaedica Scandinavica* 1994, **65**(2):127-30. (Guideline Ref ID:
6 AUNE1994)
- 7 10. Bajammal SS, Zlowodzki M, Lelwica A, Tornetta P, III, Einhorn TA, Buckley R *et al.* The
8 use of calcium phosphate bone cement in fracture treatment. A meta-analysis of
9 randomized trials. *Journal of Bone & Joint Surgery - American Volume* 2008, **90**(6):1186-
10 96. (Guideline Ref ID: BAJAMMAL2008)
- 11 11. Baker RP, Squires B, Gargan MF, Bannister GC. Total hip arthroplasty and
12 hemiarthroplasty in mobile, independent patients with a displaced intracapsular
13 fracture of the femoral neck. A randomized, controlled trial. *Journal of Bone & Joint*
14 *Surgery - American Volume* 2006, **88**(12):2583-9. (Guideline Ref ID: BAKER2006)
- 15 12. Barden B, Kroger K, Loer F. [Intraoperative Doppler ultrasound of the femoral vein for
16 maintaining venous patency in hip joint prosthesis implantation]. *Unfallchirurg* 2001,
17 **104**(2):138-42. (Guideline Ref ID: BARDEN2001)
- 18 13. Barna B. Comparison of spinal and general anesthesia in the surgical treatment of hip
19 fractures. *Orvosi Hetilap* 1981, **122**(19):1135-8. (Guideline Ref ID: BARNA1981)
- 20 14. Barton TM, Gleeson R, Topliss C, Greenwood R, Harries WJ, Chesser TJ. A comparison of
21 the long gamma nail with the sliding hip screw for the treatment of AO/OTA 31-A2
22 fractures of the proximal part of the femur: a prospective randomized trial. *Journal of*
23 *Bone & Joint Surgery - American Volume* 2010, **92**(4):792-8. (Guideline Ref ID:
24 BARTON2010)
- 25 15. Baumgaertner MR, Curtin SL, Lindskog DM. Intramedullary versus extramedullary
26 fixation for the treatment of intertrochanteric hip fractures. *Clinical Orthopaedics and*
27 *Related Research* 1998,(348):87-94. (Guideline Ref ID: BAUMGAERTNER1998)
- 28 16. Ben-David B, Frankel R, Arzumov T, Marchevsky Y, Volpin G. Minidose bupivacaine-
29 fentanyl spinal anesthesia for surgical repair of hip fracture in the aged. *Anesthesiology*
30 2000, **92**(1):6-10. (Guideline Ref ID: BENDAVID2000)
- 31 17. Bennett G, Dealey C, Posnett J. The cost of pressure ulcers in the UK. *Age and Ageing*
32 2004, **33**(3):230-5. (Guideline Ref ID: BENNETT2004)
- 33 18. Benum P, Grontvedt T, Braten M, Rossvoll I, Walloe A, Ekeland A *et al.* Gamma nailing
34 versus CHS in intertrochanteric and subtrochanteric femoral fractures: a prospective
35 randomised multicentre study. *Acta Orthopaedica Scandinavica Supplementum* 1994,
36 **260**:33-4. (Guideline Ref ID: BENUM1994)
- 37 19. Bergeron E, Lavoie A, Moore L, Bamvita JM, Ratte S, Gravel C *et al.* Is the delay to
38 surgery for isolated hip fracture predictive of outcome in efficient systems? *Journal of*
39 *Trauma-Injury Infection & Critical Care* 2006, **60**(4):753-7. (Guideline Ref ID:
40 BERGERON2006)

- 1 20. Berggren D, Gustafson Y, Eriksson B, Bucht G, Hansson LI, Reiz S *et al.* Postoperative
2 confusion after anesthesia in elderly patients with femoral neck fractures. *Anesthesia*
3 *and Analgesia* 1987, **66**(6):497-504. (Guideline Ref ID: BERGGREN1987)
- 4 21. Bhandari M, Devereaux PJ, Swiontkowski MF, Tornetta P, III, Obremskey W, Koval KJ *et*
5 *al.* Internal fixation compared with arthroplasty for displaced fractures of the femoral
6 neck. A meta-analysis. *Journal of Bone & Joint Surgery - American Volume* 2003, **85-**
7 **A**(9):1673-81. (Guideline Ref ID: BHANDARI2003)
- 8 22. Bhandari M, Tornetta P. Communicating the risks of surgery to patients. *European*
9 *Journal of Trauma* 2004, **30**(3):177-81. (Guideline Ref ID: BHANDARI2004)
- 10 23. Bigler D, Adelhoj B, Petring OU, Pederson NO, Busch P, Kalhke P. Mental function and
11 morbidity after acute hip surgery during spinal and general anaesthesia. *Anaesthesia*
12 1985, **40**(7):672-6. (Guideline Ref ID: BIGLER1985)
- 13 24. Binder EF, Brown M, Sinacore DR, Steger-May K, Yarasheski KE, Schechtman KB. Effects
14 of extended outpatient rehabilitation after hip fracture: a randomized controlled trial.
15 *JAMA* 2004, **292**(7):837-46. (Guideline Ref ID: BINDER2004)
- 16 25. Bjorgul K, Reikeras O. Hemiarthroplasty in worst cases is better than internal fixation in
17 best cases of displaced femoral neck fractures: a prospective study of 683 patients
18 treated with hemiarthroplasty or internal fixation. *Acta Orthopaedica* 2006, **77**(3):368-
19 74. (Guideline Ref ID: BJORGUL2006)
- 20 26. Blomfeldt R, Tornkvist H, Eriksson K, Soderqvist A, Ponzer S, Tidermark J. A randomised
21 controlled trial comparing bipolar hemiarthroplasty with total hip replacement for
22 displaced intracapsular fractures of the femoral neck in elderly patients. *Journal of Bone*
23 *& Joint Surgery - British Volume* 2007, **89**(2):160-5. (Guideline Ref ID: BLOMFELDT2007)
- 24 27. Blomfeldt R, Tornkvist H, Ponzer S, Soderqvist A, Tidermark J. Internal fixation versus
25 hemiarthroplasty for displaced fractures of the femoral neck in elderly patients with
26 severe cognitive impairment. *Journal of Bone & Joint Surgery - British Volume* 2005,
27 **87**(4):523-9. (Guideline Ref ID: BLOMFELDT2005)
- 28 28. Borkan JM, Quirk M. Expectations and outcomes after hip fracture among the elderly.
29 *International Journal of Aging & Human Development* 1992, **34**(4):1992-350. (Guideline
30 Ref ID: BORKAN1992)
- 31 29. Borkan JM, Quirk M, Sullivan M. Finding meaning after the fall: injury narratives from
32 elderly hip fracture patients. *Social Science & Medicine* 1991, **33**(8):947-57. (Guideline
33 Ref ID: BORKAN1991)
- 34 30. Bottle A, Aylin P. Mortality associated with delay in operation after hip fracture:
35 observational study. *British Medical Journal* 2006, **332**(7547):947-51. (Guideline Ref ID:
36 BOTTLE2006)
- 37 31. Bottle A, Jarman B, Aylin P, Taylor R. Some way to go for consistent implementation of
38 guidance on hip fracture. *British Medical Journal* 2004, **328**(7448):1097. (Guideline Ref
39 ID: BOTTLE2004)

- 1 32. Boutin-Lester P, Gibson RW. Patients' perceptions of home health occupational therapy.
2 *Australian Occupational Therapy Journal* 2002, **49**(3):146-54. (Guideline Ref ID:
3 BOUTINLESTER2002)
- 4 33. Bowman AM. Sleep satisfaction, perceived pain and acute confusion in elderly clients
5 undergoing orthopaedic procedures. *Journal of Advanced Nursing* 1997, **26**(3):550-64.
6 (Guideline Ref ID: BOWMAN1997)
- 7 34. Branfoot T, Faraj AA, Porter P. Cemented versus uncemented Thompson's prosthesis: a
8 randomised prospective functional outcome study. *Injury* 2000, **31**:280-1. (Guideline Ref
9 ID: BRANFOOT2000)
- 10 35. Bray TJ, Smith-Hoefer E, Hooper A, Timmerman L. The displaced femoral neck fracture.
11 Internal fixation versus bipolar endoprosthesis. Results of a prospective, randomized
12 comparison. *Clin Orthop Relat Res* 1988,(230):127-40. (Guideline Ref ID: BRAY1988)
- 13 36. Brichant JF, Blom-Peters L, Buffels R, Lamy M. Central neural blockage failed to decrease
14 deep venous thrombosis in patients undergoing hip surgery and receiving low molecular
15 weight heparin. *British Journal of Anaesthesia* 1995, **74**(Suppl 1):75. (Guideline Ref ID:
16 BRICHANT1995)
- 17 37. Bridle SH, Patel AD, Bircher M, Calvert PT. Fixation of intertrochanteric fractures of the
18 femur. A randomised prospective comparison of the gamma nail and the dynamic hip
19 screw. *Journal of Bone & Joint Surgery - British Volume* 1991, **73**(2):330-4. (Guideline Ref
20 ID: BRIDLE1991)
- 21 38. British Orthopaedic Association. (2007) Advisory book on consultant trauma and
22 orthopaedic services. London: British Orthopaedic Association. (Guideline Ref ID:
23 BOA2007A)
- 24 39. British Orthopaedic Association. (2007) The care of patients with fragility fractures.
25 (Guideline Ref ID: BOA2007)
- 26 40. Butt MS, Krikler SJ, Nafie S, Ali MS. Comparison of dynamic hip screw and gamma nail: a
27 prospective, randomized, controlled trial. *Injury* 1995, **26**(9):615-8. (Guideline Ref ID:
28 BUTT1995)
- 29 41. Cameron I, Crotty M, Currie C, Finnegan T, Gillespie L, Gillespie W *et al.* Geriatric
30 rehabilitation following fractures in older people: A systematic review. *Health
31 Technology Assessment* 2000, **4**(2):i-102. (Guideline Ref ID: CAMERON2000C)
- 32 42. Cameron I, Lyle D, Quine S. Accelerated rehabilitation after proximal femoral fracture: a
33 randomised controlled trial. *Proceedings of the Annual Conference of the Australian
34 Association of Gerontology* 1991, **26**:87-90. (Guideline Ref ID: CAMERON1991)
- 35 43. Cameron ID. Coordinated multidisciplinary rehabilitation after hip fracture. *Disability
36 and Rehabilitation* 2005, **27**(18-19):1081-90. (Guideline Ref ID: CAMERON2005)
- 37 44. Cameron ID, Lyle DM, Quine S. Accelerated rehabilitation after proximal femoral
38 fracture: a randomized controlled trial. *Disability and Rehabilitation* 1993, **15**(1):29-34.
39 (Guideline Ref ID: CAMERON1993)

- 1 45. Cameron ID, Lyle DM, Quine S. Cost effectiveness of accelerated rehabilitation after
2 proximal femoral fracture. *Journal of Clinical Epidemiology* 1994, **47**(11):1307-13.
3 (Guideline Ref ID: CAMERON1994A)
- 4 46. Casati A, Aldegheri G, Vinciguerra E, Marsan A, Frascini G, Torri G. Randomized
5 comparison between sevoflurane anaesthesia and unilateral spinal anaesthesia in
6 elderly patients undergoing orthopaedic surgery. *European Journal of Anaesthesiology*
7 2003, **20**(8):640-6. (Guideline Ref ID: CASATI2003)
- 8 47. Cashman JP, Cashman WF. Comparison of complications in transtrochanteric and
9 anterolateral approaches in primary total hip arthroplasty. *Orthopedics* 2008,
10 **31**(11):1085. (Guideline Ref ID: CASHMAN2008)
- 11 48. Chakladar A, White SM. Cost estimates of spinal versus general anaesthesia for
12 fractured neck of femur surgery. *Anaesthesia* 2010, **65**(8):810-4. (Guideline Ref ID:
13 CHAKLADAR2010)
- 14 49. Chan RN, Hoskinson J. Thompson prosthesis for fractured neck of femur. A comparison
15 of surgical approaches. *Journal of Bone & Joint Surgery - British Volume* 1975, **57**(4):437-
16 43. (Guideline Ref ID: CHAN1975)
- 17 50. Christie J, Burnett R, Potts HR, Pell AC. Echocardiography of transatrial embolism during
18 cemented and uncemented hemiarthroplasty of the hip. *J Bone Joint Surg Br* 1994,
19 **76**(3):409-12. (Guideline Ref ID: CHRISTIE1994)
- 20 51. Chudinov A, Berkenstadt H, Salai M, Cahana A, Perel A. Continuous psoas compartment
21 block for anesthesia and perioperative analgesia in patients with hip fractures. *Regional*
22 *Anesthesia and Pain Medicine* 1999, **24**(6):563-8. (Guideline Ref ID: CHUDINOV1999)
- 23 52. Clague JE, Craddock E, Andrew G, Horan MA, Pendleton N. Predictors of outcome
24 following hip fracture. Admission time predicts length of stay and in-hospital mortality.
25 *Injury* 2002, **33**(1):1-6. (Guideline Ref ID: CLAGUE2002)
- 26 53. Clark DI, Ahmed AB, Baxendale BR, Moran CG. Cardiac output during hemiarthroplasty
27 of the hip. A prospective, controlled trial of cemented and uncemented prostheses. *J*
28 *Bone Joint Surg Br* 2001, **83**(3):414-8. (Guideline Ref ID: CLARK2001)
- 29 54. Closs SJ, Briggs M. Patients' verbal descriptions of pain and discomfort following
30 orthopaedic surgery. *International Journal of Nursing Studies* 2002, **39**(5):563-72.
31 (Guideline Ref ID: CLOSS2002)
- 32 55. Coast J, Richards SH, Peters TJ, Gunnell DJ, Darlow MA, Pounsford J. Hospital at home or
33 acute hospital care? A cost minimisation analysis. *British Medical Journal* 1998,
34 **316**(7147):1802-6. (Guideline Ref ID: COAST1998)
- 35 56. Coleman SA, Boyce WJ, Cosh PH, McKenzie PJ. Outcome after general anaesthesia for
36 repair of fractured neck of femur. A randomized trial of spontaneous v. controlled
37 ventilation. *British Journal of Anaesthesia* 1988, **60**(1):43-7. (Guideline Ref ID:
38 COLEMAN1988)
- 39 57. Critchley LA, Stuart JC, Conway F, Short TG. Hypotension during subarachnoid
40 anaesthesia: haemodynamic effects of ephedrine. *British Journal of Anaesthesia* 1995,
41 **74**(4):373-8. (Guideline Ref ID: CRITCHLEY1995)

- 1 58. Crotty M, Kittel A, Hayball N. Home rehabilitation for older adults with fractured hips:
2 how many will take part? *Journal of Quality in Clinical Practice* 2000, **20**(2-3):65-8.
3 (Guideline Ref ID: CROTTY2000A)
- 4 59. Crotty M, Whitehead C, Miller M, Gray S. Patient and caregiver outcomes 12 months
5 after home-based therapy for hip fracture: a randomized controlled trial. *Archives of*
6 *Physical Medicine and Rehabilitation* 2003, **84**(8):1237-9. (Guideline Ref ID:
7 CROTTY2003)
- 8 60. Crotty M, Whitehead CH, Gray S, Finucane PM. Early discharge and home rehabilitation
9 after hip fracture achieves functional improvements: a randomized controlled trial.
10 *Clinical Rehabilitation* 2002, **16**(4):406-13. (Guideline Ref ID: CROTTY2002)
- 11 61. Curtis L. (2009) Unit costs of health & social care 2009. Canterbury, Kent: Personal
12 Social Services Research Unit. (Guideline Ref ID: CURTIS2009)
- 13 62. Cuvillon P, Ripart J, Debureaux S, Boisson C, Veyrat E, Mahamat A *et al.* Analgesia after
14 hip fracture repair in elderly patients: the effect of a continuous femoral nerve block: a
15 prospective and randomised study. *Annales Françaises d'Anesthésie et de Réanimation*
16 2007, **26**(1):2-9. (Guideline Ref ID: CUVILLON2007)
- 17 63. Czoski-Murray C, De-Nigris E, Brazier JE, Walters S. (2007) A prospective controlled
18 study of the costs and health related quality of life following hip fracture. School of
19 Health and Related Research, University of Sheffield. (Guideline Ref ID:
20 CZOSKIMURRAY2007)
- 21 64. Darling JR, Murray JM, Hainsworth AM, Trinick TR. The effect of isoflurane or spinal
22 anesthesia on indocyanine green disappearance rate in the elderly. *Anesthesia &*
23 *Analgesia* 1994, **78**(4):706-9. (Guideline Ref ID: DARLING1994)
- 24 65. Davis FM, Laurenson VG. Spinal anaesthesia or general anaesthesia for emergency hip
25 surgery in elderly patients. *Anaesthesia & Intensive Care* 1981, **9**(4):352-8. (Guideline
26 Ref ID: DAVIS1981)
- 27 66. Davis FM, Woolner DF, Frampton C, Wilkinson A, Grant A, Harrison RT *et al.* Prospective,
28 multi-centre trial of mortality following general or spinal anaesthesia for hip fracture
29 surgery in the elderly. *British Journal of Anaesthesia* 1987, **59**(9):1080-8. (Guideline Ref
30 ID: DAVIS1987)
- 31 67. Davis TR, Sher JL, Checketts RG, Porter BB. Intertrochanteric fractures of the femur: a
32 prospective study comparing the use of the Kuntscher-Y nail and a sliding hip screw.
33 *Injury* 1988, **19**(6):421-6. (Guideline Ref ID: DAVIS1988)
- 34 68. Davis TR, Sher JL, Porter BB, Checketts RG. The timing of surgery for intertrochanteric
35 femoral fractures. *Injury* 1988, **19**(4):244-6. (Guideline Ref ID: DAVIS1988A)
- 36 69. Davison JN, Calder SJ, Anderson GH, Ward G, Jagger C, Harper WM *et al.* Treatment for
37 displaced intracapsular fracture of the proximal femur. A prospective, randomised trial
38 in patients aged 65 to 79 years. *Journal of Bone & Joint Surgery - British Volume* 2001,
39 **83**(2):206-12. (Guideline Ref ID: DAVISON2001)
- 40 70. Department of Health. (2007) Reference Costs 2006-07 Collection Guidance. London:
41 (Guideline Ref ID: DH2007)

- 1 71. Department of Health. (2009) The national framework for NHS continuing healthcare
2 and NHS-funded nursing care. (*Guideline Ref ID: DH2009*)
- 3 72. Department of Health. (2010) Prioritising need in the context of Putting People First: a
4 whole system approach to eligibility for social care - guidance on eligibility criteria for
5 adult social care, England 2010. (*Guideline Ref ID: DH2010A*)
- 6 73. Dorr LD, Glousman R, Hoy AL, Vanis R, Chandler R. Treatment of femoral neck fractures
7 with total hip replacement versus cemented and noncemented hemiarthroplasty.
8 *Journal of Arthroplasty* 1986, **1**(1):21-8. (*Guideline Ref ID: DORR1986*)
- 9 74. Dujardin FH, Benez C, Polle G, Alain J, Biga N, Thomine JM. Prospective randomized
10 comparison between a dynamic hip screw and a mini-invasive static nail in fractures of
11 the trochanteric area: preliminary results. *Journal of Orthopaedic Trauma* 2001,
12 **15**(6):401-6. (*Guideline Ref ID: DUJARDIN2001*)
- 13 75. Dyson A, Henderson AM, Chamley D, Campbell ID. An assessment of postoperative
14 oxygen therapy in patients with fractured neck of femur. *Anaesthesia & Intensive Care*
15 1988, **16**(4):405-10. (*Guideline Ref ID: DYSON1988*)
- 16 76. Ekman M, Berg J, Wimo A, Jonsson L, McBurney C. Health utilities in mild cognitive
17 impairment and dementia: a population study in Sweden. *International Journal of*
18 *Geriatric Psychiatry* 2007, **22**(7):649-55. (*Guideline Ref ID: EKMAN2007*)
- 19 77. Ekstrom W, Karlsson-Thur C, Larsson S, Ragnarsson B, Alberts KA. Functional outcome in
20 treatment of unstable trochanteric and subtrochanteric fractures with the proximal
21 femoral nail and the Medoff sliding plate. *Journal of Orthopaedic Trauma* 2007,
22 **21**(1):18-25. (*Guideline Ref ID: EKSTROM2007*)
- 23 78. El-Abed K, McGuinness A, Brunner J, Dallovedova P, O'Connor P, Kennedy JG.
24 Comparison of outcomes following uncemented hemiarthroplasty and dynamic hip
25 screw in the treatment of displaced subcapital hip fractures in patients aged greater
26 than 70 years. *Acta Orthopaedica Belgica* 2005, **71**(1):48-54. (*Guideline Ref ID:*
27 *ELABED2005*)
- 28 79. El-Zahaar MS, Al-Kawally HM, Said AS. A double-blind randomized study of the effects of
29 tourniquet use and type of anesthetic techniques on the incidence of deep vein
30 thrombosis (DVT) in orthopedic surgery. *Journal of Neurological & Orthopaedic*
31 *Medicine & Surgery* 1995, **16**(2):70-4. (*Guideline Ref ID: ELZAHAAR1995*)
- 32 80. Elinge E, Lofgren B, Gagerman E, Nyberg L. A group learning programme for old people
33 with hip fracture: A randomized study. *Scandinavian Journal of Occupational Therapy*
34 2003, **10**(1):27-33. (*Guideline Ref ID: ELINGE2003*)
- 35 81. Emery RJ, Broughton NS, Desai K, Bulstrode CJ, Thomas TL. Bipolar hemiarthroplasty for
36 subcapital fracture of the femoral neck. A prospective randomised trial of cemented
37 Thompson and uncemented Moore stems. *Journal of Bone & Joint Surgery - British*
38 *Volume* 1991, **73**(2):322-4. (*Guideline Ref ID: EMERY1991*)
- 39 82. Englesbe MJ, Fan Z, Baser O, Birkmeyer JD. Mortality in medicare patients undergoing
40 surgery in July in teaching hospitals. *Annals of Surgery* 2009, **249**(6):871-6. (*Guideline*
41 *Ref ID: ENGLLESBE2009*)

- 1 83. Enocson A, Hedbeck CJ, Tidermark J, Pettersson H, Ponzer S, Lapidus LJ. Dislocation of
2 total hip replacement in patients with fractures of the femoral neck. *Acta Orthopaedica*
3 2009, **80**(2):184-9. (Guideline Ref ID: ENOCSON2009)
- 4 84. Enocson A, Lapidus G, Tornkvist H, Tidermark J, Lapidus LJ. Direction of hip arthroplasty
5 dislocation in patients with femoral neck fractures. *International Orthopaedics* 2010,
6 **34**(5):641-7. (Guideline Ref ID: ENOCSON2010)
- 7 85. Enocson A, Tidermark J, Tornkvist H, Lapidus LJ. Dislocation of hemiarthroplasty after
8 femoral neck fracture: better outcome after the anterolateral approach in a prospective
9 cohort study on 739 consecutive hips. *Acta Orthopaedica* 2008, **79**(2):211-7. (Guideline
10 Ref ID: ENOCSON2008)
- 11 86. Essex HN, Clark M, Sims J, Warriner A, Cullum N. Health-related quality of life in hospital
12 inpatients with pressure ulceration: assessment using generic health-related quality of
13 life measures. *Wound Repair & Regeneration* 2009, **17**(6):797-805. (Guideline Ref ID:
14 ESSEX2009)
- 15 87. Evans JG, Prudham D, Wandless I. A prospective study of fractured proximal femur:
16 factors predisposing to survival. *Age and Ageing* 1979, **8**(4):246-50. (Guideline Ref ID:
17 EVANS1979)
- 18 88. Evans PD, Wilson C, Lyons K. Comparison of MRI with bone scanning for suspected hip
19 fracture in elderly patients. *Journal of Bone & Joint Surgery - British Volume* 1994,
20 **76**(1):158-9. (Guideline Ref ID: EVANS1994)
- 21 89. Faraj AA, Branfoot T. Cemented versus uncemented Thompson's prostheses: A
22 functional outcome study. *Injury* 1999, **30**(10):671-5. (Guideline Ref ID: FARAJ1999)
- 23 90. Faraj AA, Drakau NN. Press-fit hemiarthroplasty for elderly with femoral neck fracture:
24 High complication rates in operations performed by younger surgeons. *European*
25 *Journal of Orthopaedic Surgery and Traumatology* 2007, **17**(3):267-71. (Guideline Ref ID:
26 FARAJ2007)
- 27 91. Farnworth MG, Kenny P, Shiell A. The costs and effects of early discharge in the
28 management of fractured hip. *Age and Ageing* 1994, **23**(3):190-4. (Guideline Ref ID:
29 FARNWORTH1994)
- 30 92. Favarel-Garrigues JF, Sztark F, Petitjean ME, Thicoipe M, Lassie P, Dabadie P.
31 Hemodynamic effects of spinal anesthesia in the elderly: single dose versus titration
32 through a catheter. *Anesthesia & Analgesia* 1996, **82**(2):312-6. (Guideline Ref ID:
33 FAVARELGARRIGUES1996)
- 34 93. Field RE, Rushton N. Five-year clinical, radiological and post-mortem results of the
35 Cambridge Cup in patients with displaced fractures of the neck of the femur. *Journal of*
36 *Bone and Joint Surgery - Series B* 2005, **87**(10):1344-51. (Guideline Ref ID: FIELD2005)
- 37 94. Figved W, Opland V, Frihagen F, Jervidallo T, Madsen JE, Nordsletten L. Cemented versus
38 uncemented hemiarthroplasty for displaced femoral neck fractures. *Clin Orthop Relat*
39 *Res* 2009, **467**(9):2426-35. (Guideline Ref ID: FIGVED2009)

- 1 95. Fletcher AK, Rigby AS, Heyes FL. Three-in-one femoral nerve block as analgesia for
2 fractured neck of femur in the emergency department: a randomized, controlled trial.
3 *Annals of Emergency Medicine* 2003, **41**(2):227-33. (Guideline Ref ID: FLETCHER2003)
- 4 96. Fordham R, Thompson R, Holmes J, Hodgkinson C. (1986) A cost-benefit study of
5 geriatric-orthopaedic management of patients with fractured neck of femur. Discussion
6 paper 14. York: University of York: Centre for Health Economics. (Guideline Ref ID:
7 FORDHAM1986)
- 8 97. Foss NB, Kehlet H. Short-term mortality in hip fracture patients admitted during
9 weekends and holidays. *British Journal of Anaesthesia* 2006, **96**(4):450-4. (Guideline Ref
10 ID: FOSS2006)
- 11 98. Foss NB, Kristensen BB, Bundgaard M, Bak M, Heiring C, Virkelyst C *et al.* Fascia iliaca
12 compartment blockade for acute pain control in hip fracture patients: a randomized,
13 placebo-controlled trial. *Anesthesiology* 2007, **106**(4):773-8. (Guideline Ref ID:
14 FOSS2007)
- 15 99. Foss NB, Kristensen MT, Kristensen BB, Jensen PS, Kehlet H. Effect of postoperative
16 epidural analgesia on rehabilitation and pain after hip fracture surgery: a randomized,
17 double-blind, placebo-controlled trial. *Anesthesiology* 2005, **102**(6):1197-204.
18 (Guideline Ref ID: FOSS2005A)
- 19 100. Franchignoni F, Ottonello M, Benevolo E, Tesio L. Satisfaction with hospital
20 rehabilitation: Is it related to life satisfaction, functional status, age or education?
21 *Journal of Rehabilitation Medicine* 2002, **34**(3):105-8. (Guideline Ref ID:
22 FRANCHIGNONI2002)
- 23 101. Franzo A, Francescutti C, Simon G. Risk factors correlated with postoperative mortality
24 for hip fracture surgery in the elderly: a population-based approach. *European Journal*
25 *of Epidemiology* 2005, **20**(12):985-91. (Guideline Ref ID: FRANZO2005)
- 26 102. Frihagen F, Nordsletten L, Madsen JE. Hemiarthroplasty or internal fixation for
27 intracapsular displaced femoral neck fractures: randomised controlled trial. *British*
28 *Medical Journal* 2007, **335**(7632):1251-4. (Guideline Ref ID: FRIHAGEN2007)
- 29 103. Frihagen F, Nordsletten L, Tariq R, Madsen JE. MRI diagnosis of occult hip fractures.
30 *Acta Orthopaedica* 2005, **76**(4):524-30. (Guideline Ref ID: FRIHAGEN2005)
- 31 104. Fung W, Jonsson A, Buhren V, Bhandari M. Classifying intertrochanteric fractures of the
32 proximal femur: does experience matter? *Medical Principles & Practice* 2007, **16**(3):198-
33 202. (Guideline Ref ID: FUNG2007)
- 34 105. Furstenberg AL. Expectations about outcome following hip fracture among older
35 people. *Social Work in Health Care* 1986, **11**(4):33-47. (Guideline Ref ID:
36 FURSTENBERG1986)
- 37 106. Galea MP, Levinger P, Lythgo N, Cimoli C, Weller R, Tully E *et al.* A targeted home- and
38 center-based exercise program for people after total hip replacement: a randomized
39 clinical trial. *Archives of Physical Medicine and Rehabilitation* 2008, **89**(8):1442-7.
40 (Guideline Ref ID: GALEA2008)

- 1 107. Galvard H, Samuelsson SM. Orthopedic or geriatric rehabilitation of hip fracture
2 patients: a prospective, randomized, clinically controlled study in Malmo, Sweden.
3 *Aging-Clinical & Experimental Research* 1995, **7**(1):11-6. (Guideline Ref ID:
4 GALVARD1995)
- 5 108. Gdalevich M, Cohen D, Yosef D, Tauber C. Morbidity and mortality after hip fracture:
6 the impact of operative delay. *Archives of Orthopaedic and Trauma Surgery* 2004,
7 **124**(5):334-40. (Guideline Ref ID: GDALEVICH2004A)
- 8 109. Georgescu N, Iancu C, Trandabat C, Sirbu P, Alexa O. (2004) A prospective
9 randomised study comparing two types of hemiarthroplasty for displaced fractures of the
10 femoral neck in elderly patients *Journal of Bone & Joint Surgery - British Volume*
11 **86-B** (Suppl III): 338-Abstract (Guideline Reference ID: GEORGESCU2004)
- 12 110. Giancola R, Antonini G, Delle Rose G, Crippa C. Percutaneous compression plating
13 versus gamma nail for the treatment of pertrochanteric hip fractures. *Strategies in*
14 *Trauma & Limb Reconstruction* 2008, **3**(1):9-14. (Guideline Ref ID: GIANCOLA2008)
- 15 111. Giannoudis PV, McGuigan J, Shaw DL. Ionising radiation during internal fixation of
16 extracapsular neck of femur fractures. *Injury* 1998, **29**(6):469-72. (Guideline Ref ID:
17 GIANNOUDIS1998)
- 18 112. Gierer P, Landes J, Grubwinkler M, Gradl G, Lob G, Andress HJ. [The femoral neck
19 fracture in the elderly patient - cemented or cementless hip arthroplasty?]. *Zentralbl*
20 *Chir* 2002, **127**(6):514-8. (Guideline Ref ID: GIERER2002)
- 21 113. Gilchrist WJ, Newman RJ, Hamblen DL, Williams BO. Prospective randomised study of an
22 orthopaedic geriatric inpatient service. *British Medical Journal* 1988, **297**(6656):1116-8.
23 (Guideline Ref ID: GILCHRIST1988)
- 24 114. Gill JB, Jensen L, Chin PC, Rafiei P, Reddy K, Schutt RC, Jr. Intertrochanteric hip fractures
25 treated with the trochanteric fixation nail and sliding hip screw. *Journal of Surgical*
26 *Orthopaedic Advances* 2007, **16**(2):62-6. (Guideline Ref ID: GILL2007)
- 27 115. Gill KP, Ursic P. The impact of continuing education on patient outcomes in the elderly
28 hip fracture population. *Journal of Continuing Education in Nursing* 1994, **25**(4):181-5.
29 (Guideline Ref ID: GILL1994)
- 30 116. Gille J, Gille M, Gahr R, Wiedemann B. Acute pain management in proximal femoral
31 fractures: femoral nerve block (catheter technique) vs. systemic pain therapy using a
32 clinic internal organisation model. *Anaesthetist* 2006, **55**(4):414-22. (Guideline Ref ID:
33 GILLE2006)
- 34 117. Giusti A, Barone A, Oliveri M, Pizzonia M, Razzano M, Palummeri E *et al.* An analysis of
35 the feasibility of home rehabilitation among elderly people with proximal femoral
36 fractures. *Archives of Physical Medicine and Rehabilitation* 2006, **87**(6):826-31.
37 (Guideline Ref ID: GIUSTI2006A)
- 38 118. Gjertsen JE, Vinje T, Engesaeter LB, Lie SA, Havelin LI, Furnes O *et al.* Internal screw
39 fixation compared with bipolar hemiarthroplasty for treatment of displaced femoral
40 neck fractures in elderly patients. *Journal of Bone & Joint Surgery - American Volume*
41 2010, **92**(3):619-28. (Guideline Ref ID: GJERTSEN2010)

- 1 119. Gjertsen JE, Vinje T, Lie SA, Engesaeter LB, Havelin LI, Furnes O *et al.* Patient
2 satisfaction, pain, and quality of life 4 months after displaced femoral neck fractures: a
3 comparison of 663 fractures treated with internal fixation and 906 with bipolar
4 hemiarthroplasty reported to the Norwegian Hip Fracture Register. *Acta Orthopaedica*
5 2008, **79**(5):594-601. (Guideline Ref ID: GJERTSEN2008)
- 6 120. Goh SK, Samuel M, Su DH, Chan ES, Yeo SJ. Meta-analysis comparing total hip
7 arthroplasty with hemiarthroplasty in the treatment of displaced neck of femur
8 fracture. *Journal of Arthroplasty* 2009, **24**(3):400-6. (Guideline Ref ID: GOH2009)
- 9 121. Gonzalez-Montalvo JI, Alarcon T, Mauleon JL, Gil-Garay E, Gotor P, Martin-Vega A. The
10 orthogeriatric unit for acute patients: a new model of care that improves efficiency in
11 the management of patients with hip fracture. *Hip International* 2010, **20**(2):229-35.
12 (Guideline Ref ID: GONZALEZMONTALVO2010)
- 13 122. Gorodetskiy IG, Gorodnichenko AI, Tursin PS, Reshetnyak VK, Uskov ON. Non-invasive
14 interactive neurostimulation in the postoperative recovery of patients with a
15 trochanteric fracture of the femur. A randomised, controlled trial. *Journal of Bone &*
16 *Joint Surgery - British Volume* 2007, **89**(11):1488-94. (Guideline Ref ID:
17 GORODETSKIY2007)
- 18 123. Graf C, Scherer MA, Gumpfenberg S. Cemented vs.cement-free endoprothetics in the
19 traumatology of older people. *Hefte zur der Unfallchirurg* 2000, **282**:343. (Guideline Ref
20 ID: GRAF2000)
- 21 124. Graham J. Early or delayed weight-bearing after internal fixation of transcervical
22 fracture of the femur. A clinical trial. *Journal of Bone & Joint Surgery - British Volume*
23 1968, **50**(3):562-9. (Guideline Ref ID: GRAHAM1968)
- 24 125. Grimes JP, Gregory PM, Noveck H, Butler MS, Carson JL. The effects of time-to-surgery
25 on mortality and morbidity in patients following hip fracture. *American Journal of*
26 *Medicine* 2002, **112**(9):702-9. (Guideline Ref ID: GRIMES2002A)
- 27 126. Grimley EJ, Wandless I, Prudham D. A prospective study of fractured proximal femur;
28 hospital differences. *Public Health* 1980, **94**(3):149-54. (Guideline Ref ID: GRIMLEY1980)
- 29 127. Guyer P, Landolt M, Keller H, Eberle C. The Gamma Nail in per- and intertrochanteric
30 femoral fractures--alternative or supplement to the dynamic hip screw? A prospective
31 randomized study of 100 patients with per- and intertrochanteric femoral fractures in
32 the surgical clinic of the City Hospital of Triemli, Zurich, September 1. *Aktuelle*
33 *Traumatologie* 1991, **21**(6):242-9. (Guideline Ref ID: GUYER1991)
- 34 128. Guyer P, Landolt M, Keller H, Eberle C. The Gamma nail in per- and intertrochanteric
35 femoral fractures - alternative or complimentary to the DHS? A prospective randomised
36 study. In: Marti RK, Dunki Jacobs PB, eds. *Proximal femoral fractures. Operative*
37 *technique and complications. Volume 2*, 1993. pp 481-98. London: Medical Press
38 Limited. (Guideline Reference ID: Ref ID: GUYER1993A)
- 39 129. Haddad FS, Williams RL. Femoral nerve block in extracapsular femoral neck fractures.
40 *Journal of Bone & Joint Surgery - British Volume* 1995, **77**(6):922-3. (Guideline Ref ID:
41 HADDAD1995)

- 1 130. Haentjens P, Autier P, Barette M, Boonen S, Belgian Hip Fracture Study Group.
2 Predictors of functional outcome following intracapsular hip fracture in elderly women.
3 A one-year prospective cohort study. *Injury* 2005, **36**(7):842-50. (Guideline Ref ID:
4 HAENTJENS2005)
- 5 131. Hallstrom I, Elander G, Rooke L. Pain and nutrition as experienced by patients with hip
6 fracture. *Journal of Clinical Nursing* 2000, **9**(4):639-46. (Guideline Ref ID:
7 HALLSTROM2000)
- 8 132. Handoll HHG, Cameron ID, Mak JCS, Finnegan TP. Multidisciplinary rehabilitation for
9 older people with hip fractures. *Cochrane Database of Systematic Reviews* 2009, **Issue**
10 **4**:CD007125. (Guideline Ref ID: HANDOLL2009)
- 11 133. Handoll HHG, Sherrington C. Mobilisation strategies after hip fracture surgery in adults.
12 *Cochrane Database of Systematic Reviews* 2007, **Issue 1**:CD001704. (Guideline Ref ID:
13 HANDOLL2007)
- 14 134. Hardy DC, Descamps PY, Krallis P, Fabeck L, Smets P, Bertens CL *et al.* Use of an
15 intramedullary hip-screw compared with a compression hip-screw with a plate for
16 intertrochanteric femoral fractures. A prospective, randomized study of one hundred
17 patients. *Journal of Bone & Joint Surgery - American Volume* 1998, **80**(5):618-30.
18 (Guideline Ref ID: HARDY1998)
- 19 135. Harper MC, Walsh T. Ender nailing for peritrochanteric fractures of the femur. An
20 analysis of indications, factors related to mechanical failure, and postoperative results.
21 *Journal of Bone & Joint Surgery - American Volume* 1985, **67**(1):79-88. (Guideline Ref ID:
22 HARPER1985)
- 23 136. Harper WM. Treatment of intracapsular proximal femoral fractures. A prospective
24 randomised trial comparing cemented and uncemented Thompson hemiarthroplasty in
25 the treatment of displaced intracapsular proximal femoral fractures. 1994. University
26 of Leicester. (Guideline Reference ID: HARPER1994)¶
- 27 137. Harrington P, Nihal A, Singhanian AK, Howell FR. Intramedullary hip screw versus sliding
28 hip screw for unstable intertrochanteric femoral fractures in the elderly. *Injury* 2002,
29 **33**(1):23-8. (Guideline Ref ID: HARRINGTON2002)
- 30 138. Harrison SA. Focus on research... Patients' perspectives of written information supplied
31 in an acute trauma setting following fractured neck of femur. *British Journal of*
32 *Occupational Therapy* 2006, **69**(8):385. (Guideline Ref ID: HARRISON2006)
- 33 139. Hauer K, Rost B, Rutschle K, Opitz H, Specht N, Bartsch P *et al.* Exercise training for
34 rehabilitation and secondary prevention of falls in geriatric patients with a history of
35 injurious falls. *Journal of the American Geriatrics Society* 2001, **49**(1):10-20. (Guideline
36 Ref ID: HAUER2001)
- 37 140. Hauer K, Specht N, Schuler M, Bartsch P, Oster P. Intensive physical training in geriatric
38 patients after severe falls and hip surgery. *Age and Ageing* 2002, **31**(1):49-57. (Guideline
39 Ref ID: HAUER2002)
- 40 141. Hedman AR, Heikkila K, Grafstrom M, Stromberg L. Hip fractures and cognitive state:
41 patient outcomes and proxies' perceptions of the rehabilitation period. *International*
42 *Journal of Older People Nursing* 2008, **3**(3):178-86. (Guideline Ref ID: HEDMAN2008)

- 1 142. Heetveld MJ, Rogmark C, Frihagen F, Keating J. Internal fixation versus arthroplasty for
2 displaced femoral neck fractures: what is the evidence? *Journal of Orthopaedic Trauma*
3 2009, **23**(6):395-402. (Guideline Ref ID: HEETVELD2009)
- 4 143. Hemmingsen C, Nielsen JE. Intravenous ketamine for prevention of severe hypotension
5 during spinal anaesthesia. *Acta Anaesthesiologica Scandinavica* 1991, **35**(8):755-7.
6 (Guideline Ref ID: HEMMINGSEN1991)
- 7 144. HM Government. (2009) Shaping the future of care together. (Guideline Ref ID:
8 HMG2009)
- 9 145. Ho WWS, Kwan Dai DL, Liu KW, Chow KM, Lau E, Woo J *et al.* To investigate the effect
10 and cost-effectiveness of implementing an orthogeriatric intervention for elderly
11 patients with acute hip fracture: the experience in Hong Kong. *Journal of the American*
12 *Geriatrics Society* 2009, **57**(11):2153-4. (Guideline Ref ID: HO2009)
- 13 146. Hoenig H, Rubenstein LV, Sloane R, Horner R, Kahn K. What is the role of timing in the
14 surgical and rehabilitative care of community-dwelling older persons with acute hip
15 fracture? *Archives of Internal Medicine* 1997, **157**(5):513-20. (Guideline Ref ID:
16 HOENIG1997A)
- 17 147. Hoffman CW, Lynskey TG. Intertrochanteric fractures of the femur: a randomized
18 prospective comparison of the Gamma nail and the Ambi hip screw. *Australian and New*
19 *Zealand Journal of Surgery* 1996, **66**(3):151-5. (Guideline Ref ID: HOFFMAN1996)
- 20 148. Hollingworth W, Todd C, Parker M, Roberts JA, Williams R. Cost analysis of early
21 discharge after hip fracture. *British Medical Journal* 1993, **307**(6909):903-6. (Guideline
22 Ref ID: HOLLINGWORTH1993)
- 23 149. Holmberg S, Kalen R, Thorngren KG. Treatment and outcome of femoral neck fractures.
24 An analysis of 2418 patients admitted from their own homes. *Clinical Orthopaedics and*
25 *Related Research* 1987,(218):42-52. (Guideline Ref ID: HOLMBERG1987)
- 26 150. Holmes J, House A. Psychiatric illness predicts poor outcome after surgery for hip
27 fracture: a prospective cohort study. *Psychological Medicine* 2000, **30**(4):921-9.
28 (Guideline Ref ID: HOLMES2000)
- 29 151. Holt EM, Evans RA, Hindley CJ, Metcalfe JW. 1000 femoral neck fractures: the effect of
30 pre-injury mobility and surgical experience on outcome. *Injury* 1994, **25**(2):91-5.
31 (Guideline Ref ID: HOLT1994)
- 32 152. Holt G, Smith R, Duncan K, McKeown DW. Does delay to theatre for medical reasons
33 affect the peri-operative mortality in patients with a fracture of the hip? *Journal of Bone*
34 *& Joint Surgery - British Volume* 2010, **92**(6):835-41. (Guideline Ref ID: HOLT2010)
- 35 153. Hood G, Edbrooke DL, Gerrish SP. Postoperative analgesia after triple nerve block for
36 fractured neck of femur. *Anaesthesia* 1991, **46**(2):138-40. (Guideline Ref ID: HOOD1991)
- 37 154. Huang TT, Acton GJ. Ways to maintain independence among Taiwanese elderly adults
38 with hip fractures: a qualitative study. *Geriatric Nursing* 2009, **30**(1):28-35. (Guideline
39 Ref ID: HUANG2009)

- 1 155. Hunter GA. A comparison of the use of internal fixation and prosthetic replacement for
2 fresh fractures of the neck of the femur. *Br J Surg* 1969, **56**(3):229-32. (Guideline Ref ID:
3 HUNTER1969)
- 4 156. Hunter GA. A further comparison of the use of internal fixation and prosthetic
5 replacement for fresh fractures of the neck of the femur. *Br J Surg* 1974, **61**(5):382-4.
6 (Guideline Ref ID: HUNTER1974)
- 7 157. Huusko TM, Karppi P, Avikainen V, Kautiainen H, Sulkava R. Randomised, clinically
8 controlled trial of intensive geriatric rehabilitation in patients with hip fracture:
9 subgroup analysis of patients with dementia. *British Medical Journal* 2000,
10 **321**(7269):1107-11. (Guideline Ref ID: HUUSKO2000)
- 11 158. Huusko TM, Karppi P, Avikainen V, Kautiainen H, Sulkava R. Intensive geriatric
12 rehabilitation of hip fracture patients: a randomized, controlled trial. *Acta Orthopaedica*
13 *Scandinavica* 2002, **73**(4):425-31. (Guideline Ref ID: HUUSKO2002A)
- 14 159. Iliffe S, Kendrick D, Morris R, Skelton D, Gage H, Dinan S *et al.* Multi-centre cluster
15 randomised trial comparing a community group exercise programme with home based
16 exercise with usual care for people aged 65 and over in primary care: protocol of the
17 ProAct 65+ trial. *Trials [Electronic Resource]* 2010, **11**(1):6. (Guideline Ref ID: ILIFFE2010)
- 18 160. Jackson SE. The efficacy of an educational intervention on documentation of pain
19 management for the elderly patient with a hip fracture in the emergency department.
20 *JEN: Journal of Emergency Nursing* 2010, **36**(1):10-5. (Guideline Ref ID: JACKSON2010)
- 21 161. Jensen J, Rasmussen TB, Christensen S, Holm-Moller S, Lauritzen J. Internal fixation or
22 prosthetic replacement in fresh femoral neck fractures. *Acta Orthopaedica Scandinavica*
23 1984, **55**(6):712. (Guideline Ref ID: JENSEN1984)
- 24 162. Johansson T. Displaced femoral neck fractures: a prospective randomized study of
25 clinical outcome, nutrition and costs. 2002. Linkoping University. (Guideline Reference
26 ID: JOHANSSON2002)¶
- 27 163. Johansson T, Bachrach-Lindstrom M, Aspenberg P, Jonsson D, Wahlstrom O. The total
28 costs of a displaced femoral neck fracture: comparison of internal fixation and total hip
29 replacement. A randomised study of 146 hips. *International Orthopaedics* 2006, **30**(1):1-
30 6. (Guideline Ref ID: JOHANSSON2006)
- 31 164. Johnson MM, Thompson B, Hartford JM. Complications of cemented versus
32 noncemented hemiarthroplasty for femoral neck fractures [abstract]. Mid-America
33 Orthopaedic Association 19th annual meeting; 2001 April 18-22; Amelia Island (FL).
34 <http://www.jbjs.org/abstracts/maoa01/absy001.htm> [accessed 24-11-2003]. (Guideline
35 Ref ID: JOHNSON2001)
- 36 165. Jones SF, White A. Analgesia following femoral neck surgery. Lateral cutaneous nerve
37 block as an alternative to narcotics in the elderly. *Anaesthesia* 1985, **40**(7):682-5.
38 (Guideline Ref ID: JONES1985)
- 39 166. Jonsson B, Sernbo I, Carlsson A, Fredin H, Johnell O. Social function after cervical hip
40 fracture. A comparison of hook-pins and total hip replacement in 47 patients. *Acta*
41 *Orthopaedica Scandinavica* 1996, **67**(5):431-4. (Guideline Ref ID: JONSSON1996)

- 1 167. Juelsgaard P, Sand NP, Felsby S, Dalsgaard J, Jakobsen KB, Brink O *et al.* Perioperative
2 myocardial ischaemia in patients undergoing surgery for fractured hip randomized to
3 incremental spinal, single-dose spinal or general anaesthesia. *European Journal of*
4 *Anaesthesiology* 1998, **15**(6):656-63. (Guideline Ref ID: JUELSGAARD1998)
- 5 168. Kamel HK, Iqbal MA, Mogallapu R, Maas D, Hoffmann RG. Time to ambulation after hip
6 fracture surgery: relation to hospitalization outcomes. *Journals of Gerontology Series A-*
7 *Biological Sciences & Medical Sciences* 2003, **58**(11):1042-5. (Guideline Ref ID:
8 KAMEL2003)
- 9 169. Kamitani K, Higuchi A, Asahi T, Yoshida H. Postoperative delirium after general
10 anesthesia vs. spinal anesthesia in geriatric patients. *Masui* 2003, **52**(9):972-5.
11 (Guideline Ref ID: KAMITANI2003)
- 12 170. Karpman RR, Lee TK, Moore BM. Austin-Moore versus bipolar hemi-arthroplasty for
13 displaced femoral neck fractures: a randomized prospective study. *Orthopaedic*
14 *Transactions* 1992, **16**:749-50. (Guideline Ref ID: KARPMAN1992)
- 15 171. Karumo I. Recovery and rehabilitation of elderly subjects with femoral neck fractures.
16 *Annales Chirurgiae et Gynaecologiae* 1977, **66**(3):170-6. (Guideline Ref ID:
17 KARUMO1977A)
- 18 172. Kavcic G, Hudoklin P, Mikek M, Hussein M. Hemiarthroplasty versus total arthroplasty
19 for treatment of femoral neck fractures. *European Journal of Trauma* 2006, **32**(Suppl
20 1):24. (Guideline Ref ID: KAVCIC2006)
- 21 173. Keating JF, Grant A, Masson M, Scott NW, Forbes JF. Displaced intracapsular hip
22 fractures in fit, older people: a randomised comparison of reduction and fixation,
23 bipolar hemiarthroplasty and total hip arthroplasty. *Health Technology Assessment*
24 2005, **9**(41):iii-x, 1. (Guideline Ref ID: KEATING2005)
- 25 174. Keating JF, Grant A, Masson M, Scott NW, Forbes JF. Randomized comparison of
26 reduction and fixation, bipolar hemiarthroplasty, and total hip arthroplasty. Treatment
27 of displaced intracapsular hip fractures in healthy older patients. *Journal of Bone & Joint*
28 *Surgery - American Volume* 2006, **88**(2):249-60. (Guideline Ref ID: KEATING2006)
- 29 175. Keene GS, Parker MJ, Pryor GA. Mortality and morbidity after hip fractures. *British*
30 *Medical Journal* 1993, **307**(6914):1248-50. (Guideline Ref ID: KEENE1993)
- 31 176. Kennie DC, Reid J, Richardson IR, Kiamari AA, Kelt C. Effectiveness of geriatric
32 rehabilitative care after fractures of the proximal femur in elderly women: a
33 randomised clinical trial. *British Medical Journal* 1988, **297**(6656):1083-6. (Guideline Ref
34 ID: KENNIE1988)
- 35 177. Kenzora JE, McCarthy RE, Lowell JD, Sledge CB. Hip fracture mortality. Relation to age,
36 treatment, preoperative illness, time of surgery, and complications. *Clinical*
37 *Orthopaedics and Related Research* 1984,(186):45-56. (Guideline Ref ID:
38 KENZORA1984A)
- 39 178. Khan RJK, MacDowell A, Crossman P, Keene GS. Cemented or uncemented
40 hemiarthroplasty for displaced intracapsular fractures of the hip - A systematic review.
41 *Injury* 2002, **33**(1):13-7. (Guideline Ref ID: KHAN2002B)

- 1 179. Kim SY, Kim YG, Hwang JK. Cementless calcar-replacement hemiarthroplasty compared
2 with intramedullary fixation of unstable intertrochanteric fractures. A prospective,
3 randomized study. *Journal of Bone & Joint Surgery - American Volume* 2005,
4 **87**(10):2186-92. (Guideline Ref ID: KIM2005)
- 5 180. Kuisma R. A randomized, controlled comparison of home versus institutional
6 rehabilitation of patients with hip fracture. *Clinical Rehabilitation* 2002, **16**(5):553-61.
7 (Guideline Ref ID: KUISMA2002)
- 8 181. Kukla C, Heinz T, Gaebler C, Heinze G, Vecsei V. The standard Gamma nail: a critical
9 analysis of 1,000 cases. *Journal of Trauma-Injury Infection & Critical Care* 2001,
10 **51**(1):77-83. (Guideline Ref ID: KUKLA2001)
- 11 182. Kullenberg B, Ysberg B, Heilman M, Resch S. Femoral nerve block as pain relief in hip
12 fracture. A good alternative in perioperative treatment proved by a prospective study.
13 *Lakartidningen* 2004, **101**(24):2104-7. (Guideline Ref ID: KULLENBERG2004)
- 14 183. Kuwabara H, Wada T, Minagi Y, Iwasaki T, Tsuji H. Compression hip screw and gamma
15 nail for intertrochanteric fractures - randomized prospective study. *Hokkaido Journal of*
16 *Orthopaedics & Traumatology* 1998, **40**(2):29-33. (Guideline Ref ID: KUWABARA1998)
- 17 184. Lachiewicz PF, Kelley SS, Soileau ES. Survival of polished compared with precoated
18 roughened cemented femoral components. A prospective, randomized study. *Journal of*
19 *Bone & Joint Surgery - American Volume* 2008, **90**(7):1457-63. (Guideline Ref ID:
20 LACHIEWICZ2008)
- 21 185. Laffosse JM, Accadbled F, Molinier F, Chiron P, Hocine B, Puget J. Anterolateral mini-
22 invasive versus posterior mini-invasive approach for primary total hip replacement.
23 Comparison of exposure and implant positioning. *Archives of Orthopaedic and Trauma*
24 *Surgery* 2008, **128**(4):363-9. (Guideline Ref ID: LAFFOSSE2008)
- 25 186. Laffosse JM, Chiron P, Molinier F, Bensafi H, Puget J. Prospective and comparative study
26 of the anterolateral mini-invasive approach versus minimally invasive posterior
27 approach for primary total hip replacement. Early results. *International Orthopaedics*
28 2007, **31**(5):597-603. (Guideline Ref ID: LAFFOSSE2007)
- 29 187. Laffosse JM, Chiron P, Tricoire JL, Giordano G, Molinier F, Puget J. Prospective and
30 comparative study of minimally invasive posterior approach versus standard posterior
31 approach in total hip replacement. *Revue de Chirurgie Orthopedique et Reparatrice de l*
32 *Appareil Moteur* 2007, **93**(3):228-37. (Guideline Ref ID: LAFFOSSE2007A)
- 33 188. Lee PC, Hsieh PH, Yu SW, Shiao CW, Kao HK, Wu CC. Biologic plating versus
34 intramedullary nailing for comminuted subtrochanteric fractures in young adults: a
35 prospective, randomized study of 66 cases. *Journal of Trauma-Injury Infection & Critical*
36 *Care* 2007, **63**(6):1283-91. (Guideline Ref ID: LEE2007)
- 37 189. Lefaivre KA, Macadam SA, Davidson DJ, Gandhi R, Chan H, Broekhuysse HM. Length of
38 stay, mortality, morbidity and delay to surgery in hip fractures. *Journal of Bone & Joint*
39 *Surgery - British Volume* 2009, **91**(7):922-7. (Guideline Ref ID: LEFAIVRE2009)
- 40 190. Leidinger W, Hoffmann G, Meierhofer JN, Wolfel R. Reduction of severe cardiac
41 complications during surgical repair of femoral neck fractures with cemented hip

- 1 arthroplasty. [German]. *Unfallchirurg* 2002, **105**(8):675-9. (Guideline Ref ID:
2 LEIDINGER2002)
- 3 191. Leung KS, So WS, Shen WY, Hui PW. Gamma nails and dynamic hip screws for
4 peritrochanteric fractures. A randomised prospective study in elderly patients. *Journal*
5 *of Bone & Joint Surgery - British Volume* 1992, **74**(3):345-51. (Guideline Ref ID:
6 LEUNG1992)
- 7 192. Levi N, Gebuhr P. Early failure and mortality following intramedullary fixation of
8 peritrochanteric fractures. *International Journal of Risk and Safety in Medicine* 2000,
9 **13**(2-3):99-101. (Guideline Ref ID: LEVI2000)
- 10 193. Lin PC, Hung SH, Liao MH, Sheen SY, Jong SY. Care needs and level of care difficulty
11 related to hip fractures in geriatric populations during the post-discharge transition
12 period. *Journal of Nursing Research* 2006, **14**(4):251-60. (Guideline Ref ID: LIN2006)
- 13 194. Lin PC, Lu CM. Hip fracture: family caregivers' burden and related factors for older
14 people in Taiwan. *Journal of Clinical Nursing* 2005, **14**(6):719-26. (Guideline Ref ID:
15 LIN2005)
- 16 195. Little NJ, Verma V, Fernando C, Elliott DS, Khaleel A. A prospective trial comparing the
17 Holland nail with the dynamic hip screw in the treatment of intertrochanteric fractures
18 of the hip. *Journal of Bone & Joint Surgery - British Volume* 2008, **90**(8):1073-8.
19 (Guideline Ref ID: LITTLE2008)
- 20 196. Lubovsky O, Liebergall M, Mattan Y, Weil Y, Mosheiff R. Early diagnosis of occult hip
21 fractures: MRI versus CT scan. *Injury* 2005, **36**(6):788-92. (Guideline Ref ID:
22 LUBOVSKY2005)
- 23 197. Macaulay W, Nellans KW, Garvin KL, Iorio R, Healy WL, Rosenwasser MP *et al.*
24 Prospective randomized clinical trial comparing hemiarthroplasty to total hip
25 arthroplasty in the treatment of displaced femoral neck fractures: winner of the Dorr
26 Award. *Journal of Arthroplasty* 2008, **23**(6 Suppl 1):2-8. (Guideline Ref ID:
27 MACAULAY2008)
- 28 198. Mackenzie DG, Wild S, Muir R. Mortality associated with delay in operation after hip
29 fracture: Scottish data provide additional information... *British Medical Journal* 2006,
30 **332**(7549):1093. (Guideline Ref ID: MACKENZIE2006B)
- 31 199. Magasi S, Durkin E, Wolf MS, Deutsch A. Rehabilitation consumers' use and
32 understanding of quality information: a health literacy perspective. *Archives of Physical*
33 *Medicine and Rehabilitation* 2009, **90**(2):206-12. (Guideline Ref ID: MAGASI2009)
- 34 200. Majumdar SR, Beaupre LA, Johnston DWC, Dick DA, Cinats JG, Jiang HX. Lack of
35 association between mortality and timing of surgical fixation in elderly patients with hip
36 fracture: results of a retrospective population-based cohort study. *Medical Care* 2006,
37 **44**(6):552-9. (Guideline Ref ID: MAJUMDAR2006)
- 38 201. Mangione KK, Craik RL, Tomlinson SS, Palombaro KM. Can elderly patients who have
39 had a hip fracture perform moderate- to high-intensity exercise at home? *Physical*
40 *Therapy* 2005, **85**(8):727-39. (Guideline Ref ID: MANGIONE2005)

- 1 202. Mannion S, Hayes I, Loughnane F, Murphy DB, Shorten GD. Intravenous but not
2 perineural clonidine prolongs postoperative analgesia after psoas compartment block
3 with 0.5% levobupivacaine for hip fracture surgery. *Anesthesia and Analgesia* 2005,
4 **100**(3):873-8. (Guideline Ref ID: MANNION2005)
- 5 203. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip
6 fracture: a randomized trial. *Journal of the American Geriatrics Society* 2001, **49**(5):516-
7 22. (Guideline Ref ID: MARCANTONIO2001)
- 8 204. Marhofer P, Faryniak B, Iler C, Koinig H, Kapral S, Mayer N. Cardiovascular effects of 6%
9 hetastarch and lactated Ringer's solution during spinal anesthesia. *Regional Anesthesia*
10 *& Pain Medicine* 1999, **24**(5):399-404. (Guideline Ref ID: MARHOFER1999)
- 11 205. Marhofer P, Schrogendorfer K, Wallner T, Koinig H, Mayer N, Kapral S. Ultrasonographic
12 guidance reduces the amount of local anesthetic for 3-in-1 blocks. *Regional Anesthesia*
13 *and Pain Medicine* 1998, **23**(6):584-8. (Guideline Ref ID: MARHOFER1998)
- 14 206. Marinelli M, Soccetti A, Panfoli N, de Palma L. Cost-effectiveness of cemented versus
15 cementless total hip arthroplasty. A Markov decision analysis based on implant cost.
16 *Journal of Orthopaedics and Traumatology* 2008, **9**(1):23-8. (Guideline Ref ID:
17 MARINELLI2008)
- 18 207. Matot I, Oppenheim-Eden A, Ratrot R, Baranova J, Davidson E, Eylon S *et al.*
19 Preoperative cardiac events in elderly patients with hip fracture randomized to epidural
20 or conventional analgesia. *Anesthesiology* 2003, **98**(1):156-63. (Guideline Ref ID:
21 MATOT2003)
- 22 208. Maurette P, Bonada G, Djiane V, Erny P. A comparison between lidocaine alone and
23 lidocaine with meperidine for continuous spinal anesthesia. *Regional Anesthesia* 1993,
24 **18**(5):290-5. (Guideline Ref ID: MAURETTE1993)
- 25 209. McGuire KJ, Bernstein J, Polsky D, Silber JH. The 2004 Marshall Urist Award: delays until
26 surgery after hip fracture increases mortality. *Clinical Orthopaedics and Related*
27 *Research* 2004,(428):294-301. (Guideline Ref ID: MCGUIRE2004)
- 28 210. McKenzie PJ, Wishart HY, Smith G. Long-term outcome after repair of fractured neck of
29 femur. Comparison of subarachnoid and general anaesthesia. *British Journal of*
30 *Anaesthesia* 1984, **56**(6):581-5. (Guideline Ref ID: MCKENZIE1984)
- 31 211. McLaren AD, Stockwell MC, Reid VT. Anaesthetic techniques for surgical correction of
32 fractured neck of femur. A comparative study of spinal and general anaesthesia in the
33 elderly. *Anaesthesia* 1978, **33**(1):10-4. (Guideline Ref ID: MCLAREN1978)
- 34 212. Mehdi SA, Kinninmonth AWG, MacLeod C, McKenzie E, James PJ. Extracapsular hip
35 fracture fixation: a prospective randomised comparison of the intramedullary hip
36 screw with the sliding hip screw. *Injury* 2000, **31**:287. (Guideline Ref ID: MEHDI2000)
- 37 213. Michos I, Brakoulakis E, Pastroudis A, Loutriotis A, Adamopoulos G. The Gamma nail
38 system compared to sliding nail and plate for peritrochanteric fractures. *Journal of Bone*
39 *& Joint Surgery - British Volume* 2001, **83**(Suppl 2):193. (Guideline Ref ID: MICHOS2001)
- 40 214. Miedel R, Ponzer S, Tornkvist H, Soderqvist A, Tidermark J. The standard Gamma nail or
41 the Medoff sliding plate for unstable trochanteric and subtrochanteric fractures. A

- 1 randomised, controlled trial. *Journal of Bone & Joint Surgery - British Volume* 2005,
2 **87**(1):68-75. (Guideline Ref ID: MIEDEL2005)
- 3 215. Moran CG, Wenn RT, Sikand M, Taylor AM. Early mortality after hip fracture: is delay
4 before surgery important? *Journal of Bone & Joint Surgery - American Volume* 2005,
5 **87A**(3):483-9. (Guideline Ref ID: MORAN2005)
- 6 216. Moseley AM, Sherrington C, Lord SR, Barraclough E, St George RJ, Cameron ID. Mobility
7 training after hip fracture: a randomised controlled trial. *Age and Ageing* 2009,
8 **38**(1):74-80. (Guideline Ref ID: MOSELEY2009)
- 9 217. Mott MP, Kronik JL, Fitzgerald RHJr, Morawa LG, Georgiadis GM, Salot WH. Gamma nail
10 versus the sliding hip screw: a prospective randomized comparison. *Orthopaedic*
11 *Transactions* 1993, **17**:1049. (Guideline Ref ID: MOTT1993)
- 12 218. Mouzopoulos G, Stamatakos M, Arabatzi H, Vasiliadis G, Batanis G, Tsembeli A *et al.* The
13 four-year functional result after a displaced subcapital hip fracture treated with three
14 different surgical options. *International Orthopaedics* 2008, **32**(3):367-73. (Guideline Ref
15 ID: MOUZOPOULOS2008)
- 16 219. Muller ME, Allgower N, Schneider R, Willenegger H. The comprehensive classification of
17 fractures of long bones. In: Allgower M, ed. *Manual of internal fixation*, 1991. Berlin:
18 Springer-Verlag. (Guideline Reference ID: Ref ID: MULLER1991)
- 19 220. Murgue D, Ehret B, Massacrier-Imbert S, Durand O, Gibaud F, Maakel A *et al.* Equimolar
20 nitrous oxide/oxygen combined with femoral nerve block for emergency analgesia of
21 femoral neck fractures. *Journal Europeen Des Urgences (Jeur)* 2006, **19**(1):9-14.
22 (Guideline Ref ID: MURGUE2006)
- 23 221. Muttu CE, Jensen EJ, Manka MA, Jr., Anders MJ, Bone LB. Femoral nerve block for
24 diaphyseal and distal femoral fractures in the emergency department. *Journal of Bone &*
25 *Joint Surgery - American Volume* 2007, **89**(12):2599-603. (Guideline Ref ID: MUTTY2007)
- 26 222. Naglie G, Tansey C, Kirkland JL, Ogilvie-Harris DJ, Detsky AS, Etchells E *et al.*
27 Interdisciplinary inpatient care for elderly people with hip fracture: A randomized
28 controlled trial. *Canadian Medical Association Journal* 2002, **167**(1):25-32. (Guideline
29 Ref ID: NAGLIE2002)
- 30 223. Naja Z, el Hassan MJ, Khatib H, Ziade MF, Lonqvist PA. Combined sciatic-paravertebral
31 nerve block vs. general anaesthesia for fractured hip of the elderly. *Middle East Journal*
32 *of Anesthesiology* 2000, **15**(5):559-68. (Guideline Ref ID: NAJA2000)
- 33 224. National Clinical Guideline Centre. (2010) Delirium: diagnosis, prevention and
34 management. (Guideline Ref ID: NCGC2010)
- 35 225. National Clinical Guideline Centre. (2010) Reducing the risk of venous
36 thromboembolism (deep vein thrombosis and pulmonary embolism) in patients
37 admitted to hospital. London: (Guideline Ref ID: NCGC2010A)
- 38 226. National Institute for Clinical Excellence. Guidance on the selection of prostheses for
39 primary total hip replacement <http://www.nice.org.uk/ta2> [accessed 6-10-2008].
40 (Guideline Ref ID: TA022000)

- 1 227. National Institute for Health and Clinical Excellence. (2004) The assessment and
2 prevention of falls in older people. London: Royal College of Nursing. (*Guideline Ref ID:*
3 *NICE2004*)
- 4 228. National Institute for Health and Clinical Excellence. (2005) The management of
5 pressure ulcers in primary and secondary care. London: Royal College of Nursing.
6 (*Guideline Ref ID: NICE2005*)
- 7 229. National Institute for Health and Clinical Excellence. (2006) Nutrition support in adults:
8 oral nutrition support, enteral tube feeding and parenteral nutrition. London: National
9 Institute for Health and Clinical Excellence. (*Guideline Ref ID: NICE2006A*)
- 10 230. National Institute for Health and Clinical Excellence. (2006) Supporting people with
11 dementia and their carers in health and social care. (*Guideline Ref ID: NICE2006*)
- 12 231. National Institute for Health and Clinical Excellence. (2008) Prevention and treatment of
13 surgical site infection. London: National Institute for Health and Clinical Excellence.
14 (*Guideline Ref ID: NICE2008B*)
- 15 232. National Institute for Health and Clinical Excellence. Social value judgements: principles
16 for the development of NICE guidance
17 <http://www.nice.org.uk/media/C18/30/SVJ2PUBLICATION2008.pdf> [accessed 13-10-
18 2009]. (*Guideline Ref ID: NICE2008A*)
- 19 233. National Institute for Health and Clinical Excellence. The guidelines manual 2009
20 <http://www.nice.org.uk> [accessed 13-1-2009]. (*Guideline Ref ID: NICE2009*)
- 21 234. National Institute for Health and Clinical Excellence. (2010) Alendronate, etidronate,
22 risedronate, raloxifene and strontium ranelate for the primary prevention of
23 osteoporotic fragility fractures in postmenopausal women. London: National Institute
24 for Health and Clinical Excellence. (*Guideline Ref ID: NICE2010*)
- 25 235. National Institute for Health and Clinical Excellence. (2010) Alendronate, etidronate,
26 risedronate, raloxifene, strontium ranelate and teriparatide for the secondary
27 prevention of osteoporotic fragility fractures in postmenopausal women. London:
28 National Institute for Health and Clinical Excellence. (*Guideline Ref ID: NICE2010A*)
- 29 236. National Institute for Health and Clinical Excellence. (2010) Denosumab for the
30 prevention of osteoporotic fractures in postmenopausal women. London: National
31 Institute for Health and Clinical Excellence. (*Guideline Ref ID: NICE2010B*)
- 32 237. National Institute for Health and Clinical Excellence. (2010) Reducing the risk of venous
33 thromboembolism (deep vein thrombosis and pulmonary embolism) in patients
34 admitted to hospital. London: National Institute for Health and Clinical Excellence.
35 (*Guideline Ref ID: NICE2010C*)
- 36 238. Neander G. (2000) Reduction and fixation versus total hip arthroplasty in the treatment
37 of displaced femoral neck fractures. Results after four years of a prospective
38 randomised study in 100 patients. *Displaced femoral neck fractures. Studies on*
39 *osteosynthesis and total hip arthroplasty*. Stockholm: Division of Orthopaedics,
40 Karolinska Institutet, Danderyds Hospital. (*Guideline Ref ID: NEANDER2000*)

- 1 239. Neander G, Adolphson P, von Sivers K, Dahlborn M, Dalen N. Bone and muscle mass
2 after femoral neck fracture. A controlled quantitative computed tomography study of
3 osteosynthesis versus primary total hip arthroplasty. *Archives of Orthopaedic and*
4 *Trauma Surgery* 1997, **116**(8):470-4. (Guideline Ref ID: NEANDER1997)
- 5 240. Netten A, Bebbington A, Darton R, Forder J, Miles K. (1998) Survey of Care Homes for
6 Elderly People. Final Report. Canterbury, Kent: Personal Social Services Research
7 Unit. (Guideline Ref ID: NETTEN1998)
- 8 241. Netten A, Curtis L. (2002) Unit costs of health and social care 2001. Personal and Social
9 Services Research Unit (PSSRU). (Guideline Ref ID: NETTEN2002)
- 10 242. Nishikawa K, Yamakage M, Omote K, Namiki A. Prophylactic IM small-dose
11 phenylephrine blunts spinal anesthesia-induced hypotensive response during surgical
12 repair of hip fracture in the elderly. *Anesthesia & Analgesia* 2002, **95**(3):751-6.
13 (Guideline Ref ID: NISHIKAWA2002)
- 14 243. Novack V, Jotkowitz A, Etzion O, Porath A. Does delay in surgery after hip fracture lead
15 to worse outcomes? A multicenter survey. *International Journal for Quality in Health*
16 *Care* 2007, **19**(3):170-6. (Guideline Ref ID: NOVACK2007)
- 17 244. O'Brien PJ, Meek RN, Blachut PA, Broekhuysen HM, Sabharwal S. Fixation of
18 intertrochanteric hip fractures: gamma nail versus dynamic hip screw. A randomized,
19 prospective study. *Canadian Journal of Surgery* 1995, **38**(6):516-20. (Guideline Ref ID:
20 OBRIEN1995)
- 21 245. O'Cathain A. Evaluation of a Hospital at Home scheme for the early discharge of
22 patients with fractured neck of femur. *Journal of Public Health Medicine* 1994,
23 **16**(2):205-10. (Guideline Ref ID: OCATHAIN1994)
- 24 246. Office for Economic Cooperation and Development (OECD). OECD PPPs and exchange
25 rates http://stats.oecd.org/Index.aspx?DatasetCode=SNA_TABLE4 [accessed 11-1-
26 2009]. (Guideline Ref ID: OECD2009)
- 27 247. Oldmeadow LB, Edwards ER, Kimmel LA, Kippen E, Robertson VJ, Bailey MJ. No rest for
28 the wounded: Early ambulation after hip surgery accelerates recovery. *ANZ Journal of*
29 *Surgery* 2006, **76**(7):607-11. (Guideline Ref ID: OLDMEADOW2006)
- 30 248. Olsson LE, Karlsson J, Ekman I. Effects of nursing interventions within an integrated care
31 pathway for patients with hip fracture. *Journal of Advanced Nursing* 2007, **58**(2):116-25.
32 (Guideline Ref ID: OLSSON2007A)
- 33 249. Olsson LE, Nystrom AE, Karlsson J, Ekman I. Admitted with a hip fracture: patient
34 perceptions of rehabilitation. *Journal of Clinical Nursing* 2007, **16**(5):853-9. (Guideline
35 Ref ID: OLSSON2007)
- 36 250. Orosz GM, Magaziner J, Hannan EL, Morrison RS, Koval K, Gilbert M *et al.* Association of
37 timing of surgery for hip fracture and patient outcomes. *JAMA* 2004, **291**(14):1738-43.
38 (Guideline Ref ID: OROSZ2004)
- 39 251. Ovesen O, Andersen M, Poulsen T, Nymark T, Overgaard S, Rock ND. The trochanteric
40 gamma nail versus the dynamic hip screw: a prospective randomised study. One-year

- 1 follow-up of 146 intertrochanteric fractures. *Hip International* 2006, **16**(4):293-8.
2 (Guideline Ref ID: OVESEN2006)
- 3 252. Owen H, Hutton P. Doxapram and the fractured femur. *Anaesthesia* 1982, **37**(3):301-4.
4 (Guideline Ref ID: OWEN1982)
- 5 253. Pahlpatz PVM, Langius FB. Comparing the Gamma nail and the Dynamic Hip Screw in
6 the treatment of pertrochanteric fractures. Preliminary results of a prospective
7 randomised study. In: Marti RK, Dunki Jacobs PB, eds. *Proximal femoral fractures.*
8 *Operative techniques and complications. Volume 2.*, 1993. pp 475-80. London: Medical
9 Press Limited. (Guideline Reference ID: Ref ID: PAHLPATZ1993)
- 10 254. Pajarinen J, Lindahl J, Michelsson O, Savolainen V, Hirvensalo E. Pertrochanteric femoral
11 fractures treated with a dynamic hip screw or a proximal femoral nail. A randomised
12 study comparing postoperative rehabilitation. *Journal of Bone & Joint Surgery - British*
13 *Volume* 2005, **87**(1):76-81. (Guideline Ref ID: PAJARINEN2005)
- 14 255. Pajarinen J, Lindahl J, Savolainen V, Michelsson O, Hirvensalo E. Femoral shaft
15 medialisation and neck-shaft angle in unstable pertrochanteric femoral fractures.
16 *International Orthopaedics* 2004, **28**(6):347-53. (Guideline Ref ID: PAJARINEN2004)
- 17 256. Palm H, Jacobsen S, Krasheninnikoff M, Foss NB, Kehlet H, Gebuhr P *et al.* Influence of
18 surgeon's experience and supervision on re-operation rate after hip fracture surgery.
19 *Injury* 2007, **38**(7):775-9. (Guideline Ref ID: PALM2007)
- 20 257. Palm H, Krasheninnikoff M, Jacobsen S. Surgical treatment of proximal femoral fracture.
21 *Ugeskrift for Laeger* 2006, **168**(35):2891-6. (Guideline Ref ID: PALM2006)
- 22 258. Park SR, Kang JS, Kim HS, Lee WH, Kim YH. Treatment of intertrochanteric fracture with
23 the Gamma AP locking nail or by a compression hip screw--a randomised prospective
24 trial. *International Orthopaedics* 1998, **22**(3):157-60. (Guideline Ref ID: PARK1998)
- 25 259. Parker M, Johansen A. Hip fracture. *British Medical Journal* 2006, **333**(7557):27-30.
26 (Guideline Ref ID: PARKER2006)
- 27 260. Parker M, Pryor G, Gurusamy K. Cemented versus uncemented hemiarthroplasty for
28 intracapsular hip fractures: a randomised controlled trial in 400 patients. *Journal of*
29 *Bone & Joint Surgery - British Volume* 2010, **92B**(1):116-22. (Guideline Ref ID:
30 PARKER2010)
- 31 261. Parker MJ. Internal fixation or arthroplasty for displaced subcapital fractures in the
32 elderly? *Injury* 1992, **23**(8):521-4. (Guideline Ref ID: PARKER1992B)
- 33 262. Parker MJ, Griffiths R, Appadu B. Nerve blocks (subcostal, lateral cutaneous, femoral,
34 triple, psoas) for hip fractures. *Cochrane Database of Systematic Reviews* 2002, **Issue**
35 **1**:CD001159. (Guideline Ref ID: PARKER2002A)
- 36 263. Parker MJ, Gurusamy K. Arthroplasties (with and without bone cement) for proximal
37 femoral fractures in adults. *Cochrane Database of Systematic Reviews* 2006, **Issue**
38 **3**:CD001706. (Guideline Ref ID: PARKER2006A)

- 1 264. Parker MJ, Gurusamy K. Internal fixation versus arthroplasty for intracapsular proximal
2 femoral fractures in adults. *Cochrane Database of Systematic Reviews* 2006, **Issue**
3 **4**:CD001708. (Guideline Ref ID: PARKER2006C)
- 4 265. Parker MJ, Gurusamy KS, Azegami S. Arthroplasties (with and without bone cement) for
5 proximal femoral fractures in adults. *Cochrane Database of Systematic Reviews* 2010,
6 **6**:CD001706. (Guideline Ref ID: PARKER2010A)
- 7 266. Parker MJ, Handoll HHG, Griffiths R. Anaesthesia for hip fracture surgery in adults.
8 *Cochrane Database of Systematic Reviews* 2004, **Issue 4**:CD000521. (Guideline Ref ID:
9 PARKER2004B)
- 10 267. Parker MJ, Khan RJ, Crawford J, Pryor GA. Hemiarthroplasty versus internal fixation for
11 displaced intracapsular hip fractures in the elderly. A randomised trial of 455 patients.
12 *Journal of Bone & Joint Surgery - British Volume* 2002, **84**(8):1150-5. (Guideline Ref ID:
13 PARKER2002)
- 14 268. Parker MJ, Myles JW, Anand JK, Drewett R. Cost-benefit analysis of hip fracture
15 treatment. *Journal of Bone & Joint Surgery - British Volume* 1992, **74**(2):261-4.
16 (Guideline Ref ID: PARKER1992)
- 17 269. Parker MJ, Pervez H. Surgical approaches for inserting hemiarthroplasty of the hip.
18 *Cochrane Database of Systematic Reviews* 2002, **Issue 3**:CD001707. (Guideline Ref ID:
19 PARKER2002B)
- 20 270. Parker MJ, Pryor GA, Myles JW. Early discharge after hip fracture. Prospective 3-year
21 study of 645 patients. *Acta Orthopaedica Scandinavica* 1991, **62**(6):563-6. (Guideline Ref
22 ID: PARKER1991)
- 23 271. Parker MJ, Pryor GA, Myles JW. The value of a special surgical team in preventing
24 complications in the treatment of hip fractures. *International Orthopaedics* 1994,
25 **18**(3):184-8. (Guideline Ref ID: PARKER1994)
- 26 272. Piangatelli C, De AC, Pecora L, Recanatini F, Testasecca D. Levobupivacaine versus
27 ropivacaine in psoas compartment block and sciatic nerve block in orthopedic surgery
28 of the lower extremity. *Minerva Anestesiologica* 2004, **70**(12):801-7. (Guideline Ref ID:
29 PIANGATELLI2004)
- 30 273. Pitto RP, Blunk J, Kossler M. Transesophageal echocardiography and clinical features of
31 fat embolism during cemented total hip arthroplasty. A randomized study in patients
32 with a femoral neck fracture. *Archives of Orthopaedic and Trauma Surgery* 2000, **120**(1-
33 2):53-8. (Guideline Ref ID: PITTO2000)
- 34 274. Pownall E. Using a patient narrative to influence orthopaedic nursing care in fractured
35 hips. *Journal of Orthopaedic Nursing* 2004, **8**(3):151-9. (Guideline Ref ID:
36 POWNALL2004)
- 37 275. Pryor GA, Williams DR. Rehabilitation after hip fractures. Home and hospital
38 management compared. *Journal of Bone & Joint Surgery - British Volume* 1989,
39 **71**(3):471-4. (Guideline Ref ID: PRYOR1989)
- 40 276. Puolakka TJ, Laine HJ, Tarvainen T, Aho H. Thompson hemiarthroplasty is superior to
41 Ullevaal screws in treating displaced femoral neck fractures in patients over 75 years. A

- 1 prospective randomized study with two-year follow-up. *Annales Chirurgiae et*
2 *Gynaecologiae* 2001, **90**(3):225-8. (Guideline Ref ID: PUOLAKKA2001)
- 3 277. Racle JP, Benkhadra A, Poy JY, Gleizal B, Gaudray A. Comparative study of general and
4 spinal anesthesia in elderly women in hip surgery. *Annales Françaises d'Anesthésie et de*
5 *Réanimation* 1986, **5**(1):24-30. (Guideline Ref ID: RACLE1986)
- 6 278. Rademakers L, Vainas T, van Zutphen S, Brink P, van Helden S. Pressure ulcers and
7 prolonged hospital stay in hip fracture patients affected by time-to-surgery. *European*
8 *Journal of Trauma and Emergency Surgery* 2007, **33**(3):238-44. (Guideline Ref ID:
9 RADEMAKERS2007)
- 10 279. Radford PJ, Needoff M, Webb JK. A prospective randomised comparison of the dynamic
11 hip screw and the gamma locking nail. *Journal of Bone & Joint Surgery - British Volume*
12 1993, **75**(5):789-93. (Guideline Ref ID: RADFORD1993)
- 13 280. Rae HC, Harris IA, McEvoy L, Todorova T. Delay to surgery and mortality after hip
14 fracture. *ANZ Journal of Surgery* 2007, **77**(10):889-91. (Guideline Ref ID: RAE2007A)
- 15 281. Rahme DM, Harris IA. Intramedullary nailing versus fixed angle blade plating for
16 subtrochanteric femoral fractures: a prospective randomised controlled trial. *Journal of*
17 *Orthopaedic Surgery* 2007, **15**(3):278-81. (Guideline Ref ID: RAHME2007)
- 18 282. Resnick B, Orwig D, Wehren L, Zimmerman S, Simpson M, Magaziner J. The Exercise Plus
19 Program for older women post hip fracture: participant perspectives. *Gerontologist*
20 2005, **45**(4):539-44. (Guideline Ref ID: RESNICK2005)
- 21 283. Resnick B, Orwig D, Yu-Yahiro J, Hawkes W, Shardell M, Hebel JR *et al.* Testing the
22 effectiveness of the exercise plus program in older women post-hip fracture. *Annals of*
23 *Behavioral Medicine* 2007, **34**(1):67-76. (Guideline Ref ID: RESNICK2007A)
- 24 284. Richards SH, Coast J, Gunnell DJ, Peters TJ, Pounsford J, Darlow MA. Randomised
25 controlled trial comparing effectiveness and acceptability of an early discharge, hospital
26 at home scheme with acute hospital care. *British Medical Journal* 1998,
27 **316**(7147):1796-801. (Guideline Ref ID: RICHARDS1998)
- 28 285. Riley TBH. Knobs or screws? - a prospective trial of prosthetic replacement against
29 internal fixation of subcapital fractures [Abstract]. *Journal of Bone & Joint Surgery -*
30 *British Volume* 1978, **60**(1):136. (Guideline Ref ID: RILEY1978)
- 31 286. Rizzo PF, Gould ES, Lyden JP, Asnis SE. Diagnosis of occult fractures about the hip.
32 Magnetic resonance imaging compared with bone-scanning. *Journal of Bone & Joint*
33 *Surgery - American Volume* 1993, **75**(3):395-401. (Guideline Ref ID: RIZZO1993)
- 34 287. Robinson SB. Transitions in the lives of elderly women who have sustained hip fractures.
35 *Journal of Advanced Nursing* 1999, **30**(6):1341-8. (Guideline Ref ID: ROBINSON1999)
- 36 288. Roden M, Schon M, Fredin H. Treatment of displaced femoral neck fractures: a
37 randomized minimum 5-year follow-up study of screws and bipolar hemiprostheses in
38 100 patients. *Acta Orthopaedica Scandinavica* 2003, **74**(1):42-4. (Guideline Ref ID:
39 RODEN2003)

- 1 289. Rodriguez J, Herrera A, Canales V, Serrano S. Epidemiologic factors, morbidity and
2 mortality after femoral neck fractures in the elderly. A comparative study: internal
3 fixation vs. hemiarthroplasty. *Acta Orthopaedica Belgica* 1987, **53**(4):472-9. (Guideline
4 Ref ID: RODRIGUEZ1987)
- 5 290. Rogers FB, Shackford SR, Keller MS. Early fixation reduces morbidity and mortality in
6 elderly patients with hip fractures from low-impact falls. *Journal of Trauma-Injury*
7 *Infection & Critical Care* 1995, **39**(2):261-5. (Guideline Ref ID: ROGERS1995)
- 8 291. Rogmark C, Carlsson A, Johnell O, Sembo I. Costs of internal fixation and arthroplasty for
9 displaced femoral neck fractures: a randomized study of 68 patients. *Acta Orthopaedica*
10 *Scandinavica* 2003, **74**(3):293-8. (Guideline Ref ID: ROGMARK2003)
- 11 292. Rogmark C, Johnell O. Primary arthroplasty is better than internal fixation of displaced
12 femoral neck fractures: a meta-analysis of 14 randomized studies with 2,289 patients.
13 *Acta Orthopaedica* 2006, **77**(3):359-67. (Guideline Ref ID: ROGMARK2006)
- 14 293. Ryan T, Enderby P, Rigby AS. A randomized controlled trial to evaluate intensity of
15 community-based rehabilitation provision following stroke or hip fracture in old age.
16 *Clinical Rehabilitation* 2006, **20**(2):123-31. (Guideline Ref ID: RYAN2006)
- 17 294. Sadowski C, Lubbeke A, Saudan M, Riand N, Stern R, Hoffmeyer P. Treatment of reverse
18 oblique and transverse intertrochanteric fractures with use of an intramedullary nail or
19 a 95 degrees screw-plate: a prospective, randomized study. *Journal of Bone & Joint*
20 *Surgery - American Volume* 2002, **84-A**(3):372-81. (Guideline Ref ID: SADOWSKI2002)
- 21 295. Sadr B, Arden GP. A comparison of the stability of proplast-coated and cemented
22 Thompson prostheses in the treatment of subcapital femoral fractures. *Injury* 1977,
23 **8**(3):234-7. (Guideline Ref ID: SADR1977)
- 24 296. Safran DG, Graham JD, Osberg JS. Social supports as a determinant of community-based
25 care utilization among rehabilitation patients. *Health Services Research* 1994, **28**(6):729-
26 50. (Guideline Ref ID: SAFRAN1994)
- 27 297. Safran O, Goldman V, Applbaum Y, Milgrom C, Bloom R, Peyser A *et al.* Posttraumatic
28 painful hip: sonography as a screening test for occult hip fractures. *Journal of*
29 *Ultrasound in Medicine* 2009, **28**(11):1447-52. (Guideline Ref ID: SAFRAN2009)
- 30 298. Santini S, Rebeccato A, Bolgan I, Turi G. Hip fractures in elderly patients treated with
31 bipolar hemiarthroplasty: comparison between cemented and cementless implants.
32 *Journal of Orthopaedics and Traumatology* 2005, **6**(2):80-7. (Guideline Ref ID:
33 SANTINI2005)
- 34 299. Sarvilinna R, Pajamaki J, Sovelius R, Puolakka T, Huhtala H, Jarvinen M. Bipolar
35 hemiarthroplasty in the treatment of hip fracture: are special centres necessary for the
36 optimal outcome? *Scandinavian Journal of Surgery: SJS* 2002, **91**(2):182-5. (Guideline
37 Ref ID: SARVILINNA2002)
- 38 300. Saudan M, Lubbeke A, Sadowski C, Riand N, Stern R, Hoffmeyer P. Pertrochanteric
39 fractures: is there an advantage to an intramedullary nail?: a randomized, prospective
40 study of 206 patients comparing the dynamic hip screw and proximal femoral nail.
41 *Journal of Orthopaedic Trauma* 2002, **16**(6):386-93. (Guideline Ref ID: SAUDAN2002)

- 1 301. Schiferer A, Gore C, Gorove L, Lang T, Steinlechner B, Zimpfer M *et al*. A randomized
2 controlled trial of femoral nerve blockade administered preclinically for pain relief in
3 femoral trauma. *Anesthesia and Analgesia* 2007, **105**(6):1852-4. (Guideline Ref ID:
4 SCHIFERER2007)
- 5 302. Sebestyen A, Boncz I, Sandor J, Nyarady J. Effect of surgical delay on early mortality in
6 patients with femoral neck fracture. *International Orthopaedics* 2008, **32**(3):375-9.
7 (Guideline Ref ID: SEBESTYEN2008B)
- 8 303. Sehat K, Baker RP, Pattison G, Price R, Harries WJ, Chesser TJ. The use of the long
9 gamma nail in proximal femoral fractures. *Injury* 2005, **36**(11):1350-4. (Guideline Ref ID:
10 SEHAT2005)
- 11 304. Shabat S, Heller E, Mann G, Gepstein R, Fredman B, Nyska M. Economic consequences
12 of operative delay for hip fractures in a non-profit institution. *Orthopedics* 2003,
13 **26**(12):1197-9. (Guideline Ref ID: SHABAT2003)
- 14 305. Shyu YI, Liang J, Wu CC, Su JY, Cheng HS, Chou SW *et al*. Interdisciplinary intervention
15 for hip fracture in older Taiwanese: benefits last for 1 year. *Journals of Gerontology*
16 *Series A-Biological Sciences & Medical Sciences* 2008, **63**(1):92-7. (Guideline Ref ID:
17 SHYU2008A)
- 18 306. Shyu YI, Liang J, Wu CC, Su JY, Cheng HS, Chou SW *et al*. A pilot investigation of the
19 short-term effects of an interdisciplinary intervention program on elderly patients with
20 hip fracture in Taiwan. *Journal of the American Geriatrics Society* 2005, **53**(5):811-8.
21 (Guideline Ref ID: SHYU2005)
- 22 307. Shyu Y-I, Liang J, Wu C-C, Su J-Y, Cheng H-S, Chou S-W *et al*. Two-year effects of
23 interdisciplinary intervention for hip fracture in older Taiwanese. *Journal of the*
24 *American Geriatrics Society* 2010, **58**(6):1081-9. (Guideline Ref ID: SHYU2010)
- 25 308. Siegmeth AW, Gurusamy K, Parker MJ. Delay to surgery prolongs hospital stay in
26 patients with fractures of the proximal femur. *Journal of Bone & Joint Surgery - British*
27 *Volume* 2005, **87**(8):1123-6. (Guideline Ref ID: SIEGMETH2005A)
- 28 309. Sikorski JM, Barrington R. Internal fixation versus hemiarthroplasty for the displaced
29 subcapital fracture of the femur. A prospective randomised study. *Journal of Bone &*
30 *Joint Surgery - British Volume* 1981, **63-B**(3):357-61. (Guideline Ref ID: SIKORSKI1981)
- 31 310. Sikorski JM, Davis NJ, Senior J. The rapid transit system for patients with fractures of
32 proximal femur. *British Medical Journal Clinical Research Ed* 1985, **290**(6466):439-43.
33 (Guideline Ref ID: SIKORSKI1985)
- 34 311. Sinclair S, James S, Singer M. Intraoperative intravascular volume optimisation and
35 length of hospital stay after repair of proximal femoral fracture: randomised controlled
36 trial. *BMJ* 1997, **315**(7113):909-12. (Guideline Ref ID: SINCLAIR1997)
- 37 312. Sircar P, Godkar D, Mahgerefteh S, Chambers K, Niranjana S, Cucco R. Morbidity and
38 mortality among patients with hip fractures surgically repaired within and after 48
39 hours. *American Journal of Therapeutics* 2007, **14**(6):508-13. (Guideline Ref ID:
40 SIRCAR2007)

- 1 313. Skinner P, Riley D, Ellery J, Beaumont A, Coumine R, Shafighian B. Displaced subcapital
2 fractures of the femur: a prospective randomized comparison of internal fixation,
3 hemiarthroplasty and total hip replacement. *Injury* 1989, **20**(5):291-3. (Guideline Ref ID:
4 SKINNER1989)
- 5 314. Slauenwhite CA, Simpson P. Patient and family perspectives on early discharge and care
6 of the older adult undergoing fractured hip rehabilitation. *Orthopaedic Nursing* 1998,
7 **17**(1):30-6. (Guideline Ref ID: SLAUENWHITE1998)
- 8 315. Smith M, Rousseau N, Lecouturier J, Gregson B, Bond J, Rodgers H. Are older people
9 satisfied with discharge information? *Nursing Times* 1997, **93**(43):52-3. (Guideline Ref
10 ID: SMITH1997)
- 11 316. Sonne-Holm S, Walter S, Jensen JS. Moore hemi-arthroplasty with and without bone
12 cement in femoral neck fractures. A clinical controlled trial. *Acta Orthopaedica*
13 *Scandinavica* 1982, **53**(6):953-6. (Guideline Ref ID: SONNEHOLM1982)
- 14 317. Soreide O, Molster A, Raugstad TS. Internal fixation versus primary prosthetic
15 replacement in acute femoral neck fractures: a prospective, randomized clinical study.
16 *British Journal of Surgery* 1979, **66**(1):56-60. (Guideline Ref ID: SOREIDE1979)
- 17 318. Spansberg NL, Anker-Moller E, Dahl JB, Schultz P, Christensen EF. The value of
18 continuous blockade of the lumbar plexus as an adjunct to acetylsalicylic acid for pain
19 relief after surgery for femoral neck fractures. *European Journal of Anaesthesiology*
20 1996, **13**(4):410-2. (Guideline Ref ID: SPANSBERG1996)
- 21 319. Stenvall M, Olofsson B, Lundstrom M, Englund U, Borssen B, Svensson O *et al.* A
22 multidisciplinary, multifactorial intervention program reduces postoperative falls and
23 injuries after femoral neck fracture. *Osteoporosis International* 2007, **18**(2):167-75.
24 (Guideline Ref ID: STENVALL2007A)
- 25 320. Stenvall M, Olofsson B, Nyberg L, Lundstrom M, Gustafson Y. Improved performance in
26 activities of daily living and mobility after a multidisciplinary postoperative
27 rehabilitation in older people with femoral neck fracture: a randomized controlled trial
28 with 1-year follow-up. *Journal of Rehabilitation Medicine* 2007, **39**(3):232-8. (Guideline
29 Ref ID: STENVALL2007)
- 30 321. Stewart HD. Pugh's nail fixation versus Thompson's prosthesis for displaced subcapital
31 fractures of the femur. *Injury* 1984, **15**(4):227-31. (Guideline Ref ID: STEWART1984)
- 32 322. Sund R, Liski A. Quality effects of operative delay on mortality in hip fracture treatment.
33 *Quality and Safety in Health Care* 2005, **14**(5):371-7. (Guideline Ref ID: SUND2005)
- 34 323. Sutcliffe AJ, Parker M. Mortality after spinal and general anaesthesia for surgical fixation
35 of hip fractures. *Anaesthesia* 1994, **49**(3):237-40. (Guideline Ref ID: SUTCLIFFE1994)
- 36 324. Svenningsen S, Benum P, Nesse O, Furset OI. Femoral neck fractures in the elderly--a
37 comparison of 3 treatment methods. *Nordisk Medicin* 1985, **100**(10):256-9. (Guideline
38 Ref ID: SVENNINGSEN1985)
- 39 325. Swanson CE, Day GA, Yelland CE, Broome JR, Massey L, Richardson HR *et al.* The
40 management of elderly patients with femoral fractures. A randomised controlled trial of

- 1 early intervention versus standard care. *Medical Journal of Australia* 1998, **169**(10):515-
2 8. (Guideline Ref ID: SWANSON1998)
- 3 326. Tappen RM, Whitehead D, Folden SL, Hall R. Effect of a video intervention on functional
4 recovery following hip replacement and hip fracture repair. *Rehabilitation Nursing*
5 2003, **28**(5):148-53. (Guideline Ref ID: TAPPEN2003)
- 6 327. Tidermark J, Ponzer S, Svensson O, Soderqvist A, Tornkvist H. Internal fixation compared
7 with total hip replacement for displaced femoral neck fractures in the elderly. A
8 randomised, controlled trial. *Journal of Bone & Joint Surgery - British Volume* 2003,
9 **85**(3):380-8. (Guideline Ref ID: TIDERMARK2003B)
- 10 328. Tidermark J, Zethraeus N, Svensson O, Tornkvist H, Ponzer S. Femoral neck fractures in
11 the elderly: functional outcome and quality of life according to EuroQol. *Quality of Life*
12 *Research* 2002, **11**(5):473-81. (Guideline Ref ID: TIDERMARK2002)
- 13 329. Tonczar L, Hammerle AF. The impairment of stress parameters by hip joint close
14 operations and the influence of anaesthesia. Preliminary results of a prospective study.
15 *Unfallchirurgie* 1981, **7**(3):138-41. (Guideline Ref ID: TONCZAR1981)
- 16 330. Tsauo JY, Leu WS, Chen YT, Yang RS. Effects on function and quality of life of
17 postoperative home-based physical therapy for patients with hip fracture. *Archives of*
18 *Physical Medicine and Rehabilitation* 2005, **86**(10):1953-7. (Guideline Ref ID:
19 TSAUO2005)
- 20 331. Tuncer S, Sert OA, Yosunkaya A, Mutlu M, Celik J, Okesli S. Patient-controlled femoral
21 nerve analgesia versus patient-controlled intravenous analgesia for postoperative
22 analgesia after trochanteric fracture repair. *Acute Pain* 2003, **4**(3-4):105-8. (Guideline
23 Ref ID: TUNCER2003)
- 24 332. Türker G, Uçkunkaya N, Yavaşoğlu B, Yılmazlar A, Özçelik S. Comparison of the
25 catheter-technique psoas compartment block and the epidural block for analgesia in
26 partial hip replacement surgery. *Acta Anaesthesiologica Scandinavica* 2003, **47**(1):30-6.
27 (Guideline Ref ID: TÜRKER2003)
- 28 333. Ungemach J. Inhalation anesthesia or "balanced anesthesia"? A comparative
29 perioperative study in geriatric patients. *Anaesthesist* 1987, **36**(6):288-91. (Guideline Ref
30 ID: UNGEMACH1987)
- 31 334. Ungemach JW, Andres FJ, Eggert E, Schoder K. The role of anaesthesia in geriatric
32 patients with hip fractures: a prospective study. *European Journal of Anaesthesiology*
33 1993, **10**(5):380. (Guideline Ref ID: UNGEMACH1993)
- 34 335. Unnanuntana A, Dimitroulias A, Bolognesi MP, Hwang KL, Goodman SB, Marcus RE.
35 Cementless femoral prostheses cost more to implant than cemented femoral
36 prostheses. *Clin Orthop Relat Res* 2009, **467**(6):1546-51. (Guideline Ref ID:
37 UNNANUNTANA2009)
- 38 336. Unwin AJ, Thomas M. Dislocation after hemiarthroplasty of the hip: a comparison of the
39 dislocation rate after posterior and lateral approaches to the hip. *Annals of the Royal*
40 *College of Surgeons of England* 1994, **76**(5):327-9. (Guideline Ref ID: UNWIN1994)

- 1 337. Utrilla AL, Reig JS, Munoz FM, Tufanisco CB. Trochanteric gamma nail and compression
2 hip screw for trochanteric fractures: a randomized, prospective, comparative study in
3 210 elderly patients with a new design of the gamma nail. *Journal of Orthopaedic*
4 *Trauma* 2005, **19**(4):229-33. (Guideline Ref ID: UTRILLA2005)
- 5 338. Uy C, Kurrle SE, Cameron ID. Inpatient multidisciplinary rehabilitation after hip fracture
6 for residents of nursing homes: a randomised trial. *Australasian Journal on Ageing* 2008,
7 **27**(1):43-4. (Guideline Ref ID: UY2008)
- 8 339. Valentin N, Lomholt B, Jensen JS, Hejgaard N, Kreiner S. Spinal or general anaesthesia
9 for surgery of the fractured hip? A prospective study of mortality in 578 patients. *British*
10 *Journal of Anaesthesia* 1986, **58**(3):284-91. (Guideline Ref ID: VALENTIN1986)
- 11 340. van Balen R, Steyerberg EW, Cools HJ, Polder JJ, Habbema JD. Early discharge of hip
12 fracture patients from hospital: transfer of costs from hospital to nursing home. *Acta*
13 *Orthopaedica Scandinavica* 2002, **73**(5):491-5. (Guideline Ref ID: VAN2002)
- 14 341. van Dortmont LM, Douw CM, van Breukelen AM, Laurens DR, Mulder PG, Wereldsma JC
15 *et al.* Cannulated screws versus hemiarthroplasty for displaced intracapsular femoral
16 neck fractures in demented patients. *Annales Chirurgiae et Gynaecologiae* 2000,
17 **89**(2):132-7. (Guideline Ref ID: VANDORTMONT2000)
- 18 342. Van Leeuwen FL, Bronselaer K, Gilles M, Sabbe MB, Delooz HH. The 'three in one' block
19 as locoregional analgesia in an emergency department. *European Journal of Emergency*
20 *Medicine* 2000, **7**(1):35-8. (Guideline Ref ID: VANLEEUWEN2000)
- 21 343. van Vugt AB, Oosterwijk WM, Goris RJ. Osteosynthesis versus endoprosthesis in the
22 treatment of unstable intracapsular hip fractures in the elderly. A randomised clinical
23 trial. *Archives of Orthopaedic and Trauma Surgery* 1993, **113**(1):39-45. (Guideline Ref ID:
24 VANVUGT1993)
- 25 344. Vidan M, Serra JA, Moreno C, Riquelme G, Ortiz J. Efficacy of a comprehensive geriatric
26 intervention in older patients hospitalized for hip fracture: a randomized, controlled
27 trial. *Journal of the American Geriatrics Society* 2005, **53**(9):1476-82. (Guideline Ref ID:
28 VIDAN2005)
- 29 345. Vochteloo AJ, Niesten D, Riedijk R, Rijnberg WJ, Bolder SB, Koeter S *et al.* Cemented
30 versus non-cemented hemiarthroplasty of the hip as a treatment for a displaced
31 femoral neck fracture: design of a randomised controlled trial. *BMC Musculoskeletal*
32 *Disorders* 2009, **10**:56. (Guideline Ref ID: VOCHTELOO2009)
- 33 346. Wang G, Gu GS, Li D, Sun DH, Zhang W, Wang TJ. Comparative study of anterolateral
34 approach versus posterior approach for total hip replacement in the treatment of
35 femoral neck fractures in elderly patients. *Chinese Journal of Traumatology* 2010,
36 **13**(4):234-9. (Guideline Ref ID: WANG2010)
- 37 347. Wang J, Jiang B, Marshall RJ, Zhang P. Arthroplasty or internal fixation for displaced
38 femoral neck fractures: which is the optimal alternative for elderly patients? A meta-
39 analysis. *International Orthopaedics* 2009, **33**(5):1179-87. (Guideline Ref ID:
40 WANG2009)
- 41 348. Wanless D. (2006) *Securing good care for older people: taking a long-term view.* London:
42 King's Fund. (Guideline Ref ID: WANLESS2006)

- 1 349. Webster R. Fractured femur: a patient's viewpoint. *New Zealand Nursing Journal* 1976,
2 **69**(12):6-8. (Guideline Ref ID: WEBSTER1976)
- 3 350. Weinrauch P. Intra-operative error during Austin Moore hemiarthroplasty. *Journal of*
4 *Orthopaedic Surgery* 2006, **14**(3):249-52. (Guideline Ref ID: WEINRAUCH2006)
- 5 351. Weller I, Wai EK, Jaglal S, Kreder HJ. The effect of hospital type and surgical delay on
6 mortality after surgery for hip fracture. *Journal of Bone & Joint Surgery - British Volume*
7 2005, **87B**(3):361-6. (Guideline Ref ID: WELLER2005)
- 8 352. White IW, Chappell WA. Anaesthesia for surgical correction of fractured femoral neck. A
9 comparison of three techniques. *Anaesthesia* 1980, **35**(11):1107-10. (Guideline Ref ID:
10 WHITE1980)
- 11 353. Widman J, Isacson J. Lateral position reduces blood loss in hip replacement surgery: a
12 prospective randomized study of 74 patients. *International Orthopaedics* 2001,
13 **25**(4):226-7. (Guideline Ref ID: WIDMAN2001)
- 14 354. Williams MA, Oberst MT, Bjorklund BC. Posthospital convalescence in older women
15 with hip fracture. *Orthopaedic Nursing* 1994, **13**(4):55-64. (Guideline Ref ID:
16 WILLIAMS1994)
- 17 355. Wykes C, Pryor J, Jeeawody B. The concerns of older women during inpatient
18 rehabilitation after fractured neck of femur. *International Journal of Therapy &*
19 *Rehabilitation* 2009, **16**(5):261-70. (Guideline Ref ID: WYKES2009)
- 20 356. Yang C, Zhu Q, Han Y, Zhu J, Wang H, Cong R *et al.* Minimally-invasive total hip
21 arthroplasty will improve early postoperative outcomes: A prospective, randomized,
22 controlled trial. *Irish Journal of Medical Science* 2010, **179**(2):285-90. (Guideline Ref ID:
23 YANG2010)
- 24 357. Yoon RS, Macaulay W, Torres G, Nellans KW, Siris ES, Bigliani LU *et al.* Assessment of
25 inpatient fragility fracture education and outpatient follow-up at an urban tertiary care
26 institution. *Endocrine Practice* 2008, **14**(1):58-68. (Guideline Ref ID: YOON2008)
- 27 358. Young Y, Resnick B. Don't worry, be positive: improving functional recovery 1 year after
28 hip fracture. *Rehabilitation Nursing* 2009, **34**(3):110-7. (Guideline Ref ID: YOUNG2009A)
- 29 359. Yu-Yahiro JA, Resnick B, Orwig D, Hicks G, Magaziner J. Design and implementation of a
30 home-based exercise program post-hip fracture: the Baltimore hip studies experience.
31 *PM&R* 2009, **1**(4):308-18. (Guideline Ref ID: YUYAHIRO2009)
- 32 360. Ziden L, Frandin K, Kreuter M. Home rehabilitation after hip fracture. A randomized
33 controlled study on balance confidence, physical function and everyday activities.
34 *Clinical Rehabilitation* 2008, **22**(12):1019-33. (Guideline Ref ID: ZIDEN2008)
- 35 361. Ziden L, Kreuter M, Frandin K. Long-term effects of home rehabilitation after hip
36 fracture - 1-year follow-up of functioning, balance confidence, and health-related
37 quality of life in elderly people. *Disability and Rehabilitation* 2010, **32**(1):18-32.
38 (Guideline Ref ID: ZIDEN2010A)

- 1 362. Ziden L, Scherman MH, Wenestam CG. The break remains--Elderly people's experiences
2 of a hip fracture 1 year after discharge. *Disability and Rehabilitation* 2010, **32**(2):103-13.
3 (Guideline Ref ID: ZIDEN2010)
- 4 363. Ziden L, Wenestam CG, Hansson-Scherman M. A life-breaking event: early experiences
5 of the consequences of a hip fracture for elderly people. *Clinical Rehabilitation* 2008,
6 **22**(9):801-11. (Guideline Ref ID: ZIDEN2008A)
- 7 364. Zou J, Xu Y, Yang H. A comparison of proximal femoral nail antirotation and dynamic hip
8 screw devices in trochanteric fractures. *Journal of International Medical Research* 2009,
9 **37**(4):1057-64. (Guideline Ref ID: ZOU2009)
- 10
11