National Clinical Guideline Centre

Draft

Appendix O

Pressure ulcer prevention and management

Risk assessment

.

.

Draft for Consultation

Commissioned by the National Institute for Health and Care Excellence

Royal College of Physicians

The Royal College of Surgeons of England

RCGP Royal College of General Practitioners

Royal College of Nursing

NICE accredited www.nice.org.uk/accreditation

Disclaimer
Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and/or their guardian or carer.

Copyright
National Clinical Guideline Centre 2013.

Funding
<Please insert your statement here

Contents

Appendix O: Risk assessment and skin assessment 9
O.1 Summary of risk assessment tools 9
O.2 Area under the ROC curve 11
O.2.1 Median AUC across studies for additional scales 11
O.2.2 AUC within study comparisons 14
O.3 Predictive ability (sensitivity and specificity) 17
O.4 Quality of the studies 23
O.5 Incidence and predictive ability of risk assessment scales - all thresholds 25
O.6 Forest plots area under the receiver operating characteristics curve (AUC) 39
O.6.1 Modified Braden scale 41
O.6.2 Braden-Q scale 42
O.6.3 Norton scale 42
O.6.4 Waterlow scale 42
O.6.5 Cubbin-Jackson scale 43
O.6.6 Douglas scale 44
O.6.7 Fragmment scale 44
O.6.8 Song and choi scale 44
O.6.9 The Northern Hospital Pressure Ulcer Prevention Plan 45
O.6.10 Schoonhoven 2002 (general patient group, mean age 60.1 years, 1229 patients, 135 events (11%), 5% received preventative treatment) 45
O.6.11 Perneger 2002 (general patient group, 1190 patients, 170 events (14%), 24% received preventative interventions) 45
O.6.12 Seongsook 2004 (ICU, mean age 62 years, 112 patients, 35 events (31%), all patients received preventative measures) 46
O.6.13 Chan 2009 (Orthopaedic patients, mean age 79 years, 197 patients, 18 events (9%), preventative measures applied ‘as normal practice’) 46
O.6.14 Kim 2009 (Surgical ICU patients, mean age 58 years, 219 patients, 40 events (18%), all patients received preventative measures) 46
O.6.15 Serpa 2009 (ICU patients, 98 patients, 7 events (7%), selected group (no low risk patients), mean age 71 years, preventative measures not mentioned) 47
O.6.16 Serpa 2011 (ICU patients, 72 patients, 8 events (11%), selected group (no low risk patients), mean age 61 years, preventative measures according to the institution) 47
O.7 Forest plots and sensitivity 47
O.7.1 Braden scale cut-off score 17 – follow-up < 1 week – general population – all grades 47
O.7.2 Braden scale cut-off score 18 – follow-up < 1 week – general population – all grades 48

O.7.3 : Braden scale cut-off score 19 – follow-up < 1 week – general population – all grades 48
O.7.4 Braden scale cut-off score 12 – follow-up 48 hours – ICU – all grades 48
O.7.5 Braden scale cut-off score 13 – follow-up 4 and 6 days – ICU – all grades 49
O.7.6 Braden scale cut-off score 16 – follow-up < 1 week – ICU – all grades 49
O.7.7 Braden scale cut-off score 18 – follow-up > 1 week – general population – all grades 49
O.7.8 Braden scale cut-off score 19 – follow-up > 1 week – general population – all grades 50
O.7.9 Braden scale cut-off score 20 – follow-up > 1 week – general population – all grades 50
O.7.10 Braden scale cut-off score 15 – follow-up > 1 week – ICU – all grades 51
O.7.11 Figure 29: Braden scale cut-off score 16 – follow-up > 1 week – ICU – all grades 51
O.7.12 Braden scale cut-off score 17 – follow-up > 1 week – ICU – all grades 51
O.7.13 Braden scale cut-off score 17 – follow-up > 1 week – general population – stage 2+ 51
O.7.14 Braden scale cut-off score 18 – follow-up > 1 week – general population – stage 2+ 51
O.7.15 Braden scale cut-off score 19 – follow-up > 1 week – general population – stage 2+ 52
O.7.16 Braden-Q scale cut-off score 16 – follow-up > 1 week – paediatric ICU – all stages 52
O.7.17 Braden-Q scale cut-off score 17 – follow-up > 1 week – paediatric ICU – all stages 52
O.7.18 Norton scale cut-off score 14 – follow-up > 1 week – general population – all stages 52
O.7.19 Norton scale cut-off score 15 – follow-up > 1 week – general population – all stages 53
O.7.20 Norton scale cut-off score 16 – follow-up > 1 week – general population – all stages 53
O.7.21 Waterlow scale cut-off score 17 – follow-up 48 hours – general population – all stages 53
O.7.22 Waterlow scale cut-off score 20 – follow-up 4 days and 6 days – general population – all stages 53
O.7.23 Waterlow scale cut-off score 10 – follow-up > 1 week – general population – all stages 54
O.7.24 Waterlow scale cut-off score 15 – follow-up > 1 week – general population – all stages 54
O.7.25 Waterlow scale cut-off score 16 – follow-up > 1 week – general population – all stages 54
O.7.26 Waterlow scale cut-off score 15 – follow-up > 1 week – ICU – stage 2+ 54
O.7.27 Cubbin-Jackson scale cut-off score 24 – follow-up > 1 week – ICU – all

stages ... 55
O.7.28 Cubbin-Jackson scale cut-off score 28 – follow-up > 1 week – ICU – all stages ... 55
O.7.29 Fragmment scale cut-off score 1 – follow-up > 1 week – general population– all stages ... 55
O.7.30 Fragmment scale cut-off score 2 – follow-up > 1 week – general population– all stages ... 55
O.7.31 Fragmment scale cut-off score 3 – follow-up > 1 week – general population– all stages ... 55
O.7.32 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 2 – follow-up > 1 week – general population– all stages ... 56
O.7.33 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 3 – follow-up > 1 week – general population– all stages ... 56
O.7.34 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 4 – follow-up > 1 week – general population– all stages ... 56
O.7.35 Douglas scale cut-off score 18 – follow-up > 1 week – ICU – all stages ... 56
O.7.36 Song and Choi scale cut-off score 2 – follow-up > 1 week – ICU – all stages ... 57
O.7.37 Suriadi and Sanada scale cut-off score 3 – follow-up > 1 week – ICU – all stages ... 57
O.7.38 Suriadi and Sanada scale cut-off score 4 – follow-up > 1 week – ICU – all stages ... 57
O.7.39 Suriadi and Sanada scale cut-off score 5 – follow-up > 1 week – ICU – all stages ... 57
O.7.40 Clinical judgement – follow-up > 1 week – general population – all stages ... 57
O.8 Skin assessment ... 60
O.8.1 Forest plots and sensitivity ... 60
O.9 References ... 66

Appendix O: Risk assessment and skin assessment

O.1 Summary of risk assessment tools

Table 1: Summary of existing risk assessment tools

Risk assessment tool	Risk factors	Scores
Braden scale (Bergstrom 1987a[5])	Sensory perception (completely limited to no impairment) Moisture (constantly to rarely) Activity (bedfast to walks frequently) Mobility (completely immobile to no limitation) Nutrition (very poor to excellent) Friction and shear (problem to no apparent problem)	Score ranges from 6 to 23*
Norton scale (Norton 1962[29])	Physical condition (very bad to good) Mental condition (stupor to alert) Activity (bedfast to ambulant) Mobility (immobile – full) Incontinent (urinary and faecal to not)	Score ranges from 5 to 20*
Waterlow scale (Waterlow 1985[48]; revised Waterlow, 2005[49])	Build/weight for height (average to below average) Skin type visual risk area (healthy to broken/spots grade 2-4) Sex (male or female) Age (14 to 81+) Continence (complete/catheterised to urinary and faecal incontinence) Mobility (fully to chair bound) Malnutrition screening tool (MST) (nutrition score) Special risk: tissue malnutrition (terminal cachexia, multiple/single organ failure, peripheral vascular disease, anaemia, smoking); neurological deficit (diabetes, MS, CVA, motor/sensory, paraplegia); major surgery/trauma (orthopaedic/spinal, on table ≥2hrs/6hrs); medication (cytotoxic, long term/high dose steroids, anti-inflammatory)	Score ranges from 2 to 20+**
Cubbin-Jackson scale (Cubbin 1991[11]; revised Jackson 1999[19])	Age Weight Skin condition of the whole body Mental state Mobility Nutrition Respiration Incontinence Hygiene Hemodynamic state	Score ranges from 10 to 40*

Risk assessment tool	Risk factors	Scores
Braden-Q scale (Quigley 1996[34])	Mobility (completely immobile to no limitations) Activity (bedfast to all patients too young to ambulate or walks frequently) Sensory perception (completely limited to no impairment) Moisture (constantly to rarely) Friction and shear (problem to no apparent problem) Nutrition (very poor to excellent) Tissue perfusion and oxygenation (extremely compromised to excellent)	Scores ranges from 7 to 28*

O.2 Area under the ROC curve

O.2.1 Median AUC across studies for additional scales

Table 2: Modified Braden scale

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95%CI)	Acceptability of values*	Quality
General population									
Chan 2009[9]	Serious1	No serious inconsistency	No serious indirectness	Serious2	197	18	74.0 (95% CI: 63.0 to 84.0)	Fair discrimination	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 Study had high risks of bias (see quality table)
2 Confidence interval around the value is consistent with two decisions

Table 3: Braden-Q scale

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95%CI)	Acceptability of values*	Quality
General population									
Curley 2003[12]	Serious1	No serious inconsistency	No serious indirectness	Serious2	322	86	83.0 (95% CI: 76.0-91.0)	Good discrimination	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 Study had high risks of bias (see quality table)
2 Confidence interval around the median is consistent with two decisions

Table 4: Waterlow scale – subgroups

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients (range)	Number of events (range)	Median AUC (%) (95%CI) range	Acceptability of values*	Quality
General population									
Schoonhoven 2002[37]; Anthony 2003[2]; Serpa 2009[39]	Very serious1	Serious2	No serious indirectness	Serious imprecision	98-45735	7-203	61 (56 to 66), range54– 90)	Poor discrimination	VERY LOW
Intensive care patients									
Compton 2008[10]	Very serious1	No serious inconsistency	No serious indirectness	No serious imprecision	698	121	59.0 (95% CI: 54.0 to 65.0)	Fail discrimination	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The studies had high to very high risks of bias (see quality table)
2 Wide variation in AUC across the studies
3 Confidence interval around the median is consistent with two decisions
4 Confidence interval around the median is consistent with three decisions

Table 5: Douglas scale

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%)	Acceptability of values*	Quality
Intensive care patients									
Seongsook 2004[38]**	Serious1	No serious inconsistency	No serious indirectness	Serious2	112	170	79	Fair discrimination	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
*** Unclear if patients with a PU at start of the study were included.*
1 Study had high risks of bias (see quality table)
2 Low event rates (< 100); no confidence interval

Table 6: Fragmment scale

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (95% CI)	Acceptability of values*	Quality
General population and intensive care patients									
Perneger 2002[33]	Serious1	No serious inconsistency	No serious indirectness	No serious imprecision	1190	170	79 (95% CI: 75.0 to 82.0)	Fair discrimination	MODERATE

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 Study had high risks of bias (see quality table)

Table 7: Song and Choi scale

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%)	Acceptability of values*	Quality
Intensive care patients									
Kim 2009[21]	Serious1	No serious inconsistency	No serious indirectness	Serious2	219	40	89	Good discrimination	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 Study had high risks of bias (see quality table)
2 Low event rates (< 100); no confidence interval

Table 8: The Northern Hospital Pressure Ulcer Prevention Plan (TNH-PUPP)

Study	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (95% CI) (range)	Acceptability of values*	Quality
Intensive care patients									
Page 2011[31]	Very serious1	No serious inconsistency	No serious indirectness	Serious2	165	7	90.0 (95% CI: 82 to 99)	Perfect discrimination	VERY LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*

1 Study had very high risks of bias (see quality table)
2 Very low events rates (< 100)

O.2.2 AUC within study comparisons

Table 9: Schoonhoven 2002[37]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
General population									
Braden scale Norton scale Waterlow scale	Serious1	No serious inconsistency	No serious indirectness	No serious imprecision	1129	135	55 (95% CI 49 to-60) 56 (95% CI 51 to 61) 61 (95% CI 56 to 66)	Fail Fail Poor	MODERATE

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The study had high risks of bias (see quality table)

Table 10: Perneger 2002[33]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
General population and intensive care patients									
Braden scale Norton scale Fragmment scale	Serious1	No serious inconsistency	No serious indirectness	No serious imprecision	1190	170	74 (95% CI 70 to 78) 74 (95% CI 70 to 78) 79 (95% CI 75 to 82)	Fair Fair Fair	MODERATE

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The study had high risks of bias (see quality table)

Table 11: Seongsook 2004[38]**

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%)	Acceptability of values*	Quality
Intensive care patients									
Braden scale Cubbin-Jackson scale Douglas scale	Serious 1	No serious inconsistency	No serious indirectness	Serious2	112	35	71 83 79	Fair Good Fair	LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
*** Unclear if patients with a PU at start of the study were included.*
1 The study had high risks of bias (see quality table)
2 Low event rates (< 100); no confidence interval

Table 12: Chan 2009[9]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
General population									
Braden scale Modified Braden scale	Serious 1	No serious inconsistency	No serious indirectness	Very serious2	197	18	73 (95% CI 63 to 84) 68 (95% CI 51 to 79)	Fair Poor	VERY LOW

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The study had high risks of bias (see quality table)
2 Low event rates (< 100); wide confidence interval

Table 13: Kim 2009[21]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%)	Acceptability of values*	Quality
Intensive care patients									
Braden scale	Serious 1	No serious inconsistency	No serious indirectness	Serious2	219	40	88	Good	LOW

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%)	Acceptability of values*	Quality
Cubbin-Jackson							91	Excellent	
Song and Choi							89	Good	

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The study had high risks of bias (see quality table)
2 Low event rates (< 100); no confidence interval

Table 14: Serpa 2009[39]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
General population									
Waterlow scale (48 hours)	Very serious1	No serious inconsistency	Serious indirectness3	Very serious2	98	7	64 (95% CI 35 to 93)	Poor	VERY LOW
Waterlow scale (4 days)							59 (95% CI 34 to 83)	Fail	
Waterlow scale (6 days)							54 (95% CI 35 to 74)	Poor	

** 90.0-100.0: perfect discrimination; 80.0-89.0: good discrimination; 70.0-79.0: fair discrimination; 60.0-69.0: poor discrimination;50.0-59.0: fail to discriminate*
1 The study had very high risks of bias (see quality table)
2 Very low event rates (< 100); very wide confidence intervals
3 Low risk patients excluded

Table 15: Serpa 2011[40]

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
Intensive care patients									
Braden scale (48 hours)	Very serious1	No serious inconsistency	Serious indirectness3	Very serious2	72	8	79 (95% CI 29 to 100)	Fair	VERY LOW
Braden scale							79(95% CI 27 to 100)	Fair Good	

Scale	Risk of bias	Inconsistency	Indirectness	Imprecision	Number of patients	Number of events	AUC (%) (95% CI)	Acceptability of values*	Quality
(4 days)									
Braden scale (6 days)							80 (95% CI 28 to 100)		

** 90-100: perfect discrimination; 80-89: good discrimination; 70-79: fair discrimination; 60-69: poor discrimination;50-59: fail to discriminate*
1 The study had very high risks of bias (see quality table)
2 Very low event rates (< 100); very wide confidence intervals
3 Low risk patients excluded

O.3 Predictive ability (sensitivity and specificity)

Table 16: Braden scale

Study	Cut-off score*	Median sensitivity** (range)	Specificity**‡ (range)
Follow-up < 1 week – all stages – general population			
Bergstrom 1998a; Braden 1994a	≤ 17	59 (range 50-78)	80 (range 76-85)
Bergstrom 1998[4]a; Braden 1994a	≤ 18	75 (range 60-88)	68 (range 68-81)
Bergstrom 1998a; Braden 1994a	≤ 19	86.5 (range 67-100)	62.5 (range 40-73)
Follow-up < 1 week – all stages – ICU			
Serpa 2011[40] (48 hours)	≤ 12	88 (95%CI 47 to 100)	64 (95%CI 51 to 76)
Serpa 2011 (6 days)	≤ 13	75 (95%CI 35to 97)	81 (95%CI 70 to 90)
Feuchtinger 2007[15]	≤ 16	77 (95%CI 56 to 91)	30 (95%CI 14 to 50)
Follow-up > 1 week – all stages – general population			
Bergstrom 1987a; Bergstrom 1998[4]a; Braden 1994a; Capobianco 1996; Chan	≤ 18	80 (95%CI 68 to 89b	73 (95%CI 66 to 79 b

Study	Cut-off score*	Median sensitivity** (range)	Specificity**‡ (range)
2009; Goodridge 1998; Langemo 1991[23]; Lyder 1999; Pang 1998; Salvadalena 1992		(range 46.2-100)	(range 14-100)
Bergstrom 1987a; Bergstrom 1998a; Braden 1994a; Capobianco 1996[8]; Salvadalena 1992	≤ 19	86 (95%CI 57 to 98) (range 71.4-100)	78 (95%CI 61 to 90) c (range 42.9-77.8)
Bergstrom 1987a; Bergstrom 1998a; Braden 1994a; Capobianco 1996; Salvadalena 1992	≤ 20	93.2 (95%CI 76 to 99) d (range 65-100)	43 (95%CI 32 to 55) d (range 31.6-66.7)
Follow-up > 1 week – all stages – ICU			
Braden 1994[7]	≤ 15	32 (95%CI 16 to 52)	95 (95%CI 87 to 99)
Braden 1994[7]a; Seongsook 2004[38]a***	≤ 16	50 (95%CI 31 to 69) and 97 (95%CI 85 to 100)	89 (95%CI 80 to 95) 26 (95%CI 17 to 37)
Braden 1994[7]	≤ 17	87.5	50.0
Follow-up > 1 week – stage 2+ – general population			
Ramundo 1995[35] e	≤ 17	42.9 (95%CI 10 to 82)	63.4 (95%CI 47 to 78)
Ramundo 1995[35] e	≤ 18	100.0 (95%CI 59 to 100)	34.1 (95%CI 20 to 51)
Ramundo 1995[35] e	≤ 19	100.0 (95%CI 59 to 100)	22.0 (95%CI 11 to 38)

** The reported thresholds are these with the highest values for median sensitivity and specificity*
*** Percentage*
**** Unclear if patients with a PU at start of the study were included.*
‡ Specificity corresponding to the median sensitivity
a Study of which sensitivity and specificity are presented.
b Sensitivity analysis without studies with < 10 events (Bergstrom 1987a and Langemo 1991) revealed a median sensitivity of 78.6 (range: 46.2-90.5) and a corresponding specificity of 74.3 (range: 14.0-100.0)
c Sensitivity analysis without study with < 10 events (Bergstrom 1987a) revealed a median sensitivity of 85.7 (range: 71.4-100.0) and a corresponding specificity of 59.1 (range: 43.0-77.8)
d Sensitivity analysis without study with < 10 events (Bergstrom 1987a) revealed a median sensitivity of 91.7 (range: 43.2-100.0) and a corresponding specificity of 40.1 (range: 31.6-66.7)
e The study of Ramundo 1995 had 7 events

Table 17: Braden-Q scale

Study	Cut-off score*	Sensitivity	Specificity
Follow-up > 1 week – all stages – paediatric ICU patients			
Curley 2003[12]	≤ 15	75.6	67.8
Curley 2003[12]	≤ 16	88.4	58.1
Curley 2003[12]	≤ 17	91.9	44.1

Table 18: Norton scale

Study	Cut-off score*	Median sensitivity**	Specificity**‡
Follow-up > 1 week – all stages – general population			
Kwong 2005[22]; Lincoln 1986[25]; Stotts 1998[42]a***; Wai-Han 1997[47]a	≤ 14	16 (95%CI 8 to 27) b range 0.0-88.9	94 (95%CI 91 to 97) b Range 61.0-94.4
Schoonhoven 2002[37] c	≤ 15	45.9 (95%CI 37 to 55)	60.3 (95%CI 57 to 63)
Pang 1998[32]a; Smith 1989[41]a	≤ 16	60 (95%CI 41 to 77) and 81 (95%CI 58 to 95)	31 (95%CI 21 to 43) and 59 (95%CI 48 to 69)

** The reported thresholds are these with the highest values for median sensitivity and specificity*
*** Percentage*
**** Unclear if patients with a PU at start of the study were included.*
‡ Specificity corresponding to the median sensitivity
a Study of which sensitivity and specificity are presented.
b Sensitivity analysis without studies with < 10 events (Kwong 2005 and Lincoln 1986) revealed a median sensitivity of 45.7 (range: 16.4-75.0) and a corresponding specificity of 80.6 (range: 66.7-94.4)
c The study of Schoonhoven 2002 had 135 events

Table 19: Waterlow scale

Study	Cut-off score*	Median sensitivity**	Specificity**‡
Follow-up < 1 week – all stages – general population			
Serpa 2009[39] (48 hours) b	≥17	71.4 (95%CI 29 to 96)	67.0 (95%CI 56 to 77)
Serpa 2009[40] (4 days) b	≥ 20	85.7 (95%CI 42 to 100)	41.0 (95%CI 30 to 51)
Follow-up > 1 week – all stages – general population			

Study	Cut-off score*	Median sensitivity**	Specificity**‡
Anthony 2003[2]a; Schoonhoven 2002[37]; Wai-Han 1997[47]	≥ 10	87.5 (95%CI 47 to 100)c range 82.3-89.6	28.2 (95%CI 22 to 35) c range 22.4-85.2
Anthony 2003[2] d	≥ 15	48.8 (95%CI 42 to 56)	94.4 (95%CI 94 to 95)
Pang 1998[32]a; Smith 1989[41]a***	≥ 16	95 (95%CI 76 to 100) and 73 (95%CI 54 to 88)	44 (95%CI 33 to 55) and 38 (95%CI 27 to 50)
Follow-up < 1 week – stage 2+ – ICU			
Weststrate 1998[50]	≥ 15	80.9 (95%CI 67 to 91)	28.5 (95%CI 25 to 33)

** The reported thresholds are these with the highest values for median sensitivity and specificity*
*** Percentage*
**** Unclear if patients with a PU at start of the study were included.*
‡ Specificity corresponding to the median sensitivity
a Study of which sensitivity and specificity are presented.
b The study of Serpa 2009 had 7 events and low risk patients were excluded
c Sensitivity analysis with only studies with > 100 events (Antony 2003 and Schoonhoven 2002) revealed a median sensitivity of 86.0 (range: 82.3-89.6) and a corresponding 53.8 (range: 22.4-85.2)
d The study of Antony 2003 had 203 events

Table 20: Cubbin-Jackson scale

Study	Cut-off score*	Sensitivity	Specificity*
Follow-up > 1 week – all stages – ICU			
Seongsook 2004[38]***	≤ 24	88.6 (95%CI 73 to 97)	61.0 (95%CI 49 to 72)
Kim 2009[21]	≤ 28	95.0 (95%CI 83 to 99)	81.6 (95%CI 75 to 87)

*** Percentage*
**** Unclear if patients with a PU at start of the study were included.*

Table 21: Fragmment scale

Study	Cut-off score*	Sensitivity	Specificity
Follow-up > 1 week – all stages – general population			
Perneger 2002[33] a	≤ 1	78.7	53.5
Perneger 2002[33] a	≤ 2	76.7	71.9

Study	Cut-off score*	Sensitivity	Specificity
Perneger 2002[33] a	≤ 3	62.1	85.0

*** Percentage*
a The study of Perneger 2002 had 170 events

Table 26: Douglas scale

Study	Cut-off score*	Sensitivity	Specificity*
Follow-up > 1 week – all stages – ICU			
Seongsook 2004[38]***	≤ 18	100.0	18.2

*** Percentage*
**** Unclear if patients with a PU at start of the study were included.*

Table 27: The Northern Hospital Pressure Ulcer Prevention Plan

Study	Cut-off score*	Sensitivity	Specificity
Follow-up > 1 week – all stages – general population			
Page 2011[31] a	≥ 2	85.7	62.0
Page 2011[31] a	≥ 3	71.4	81.0
Page 2011[31] a	≥ 4	71.4	88.0

a The study of Page 2011 had a 7 events

Table 22: Song and Choi scale

Study	Cut-off score*	Sensitivity	Specificity
Follow-up > 1 week – all stages – ICU			
Kim 2009[21]	≤ 21	95.0	69.3

*

Table 23: Suriadi and Sanada scale

Study	Cut-off score*	Sensitivity	Specificity
Follow-up > 1 week – all stages – ICU			
Suriadi 2008[43]	≥ 3	97.2	53.0
Suriadi 2008[43]	≥ 4	80.6	82.9
Suriadi 2008[43]	≥ 5	72.2	86.7

‡ Specificity corresponding to the median sensitivity

Table 24: Clinical judgement

Study	Cut-off score	Sensitivity**	Specificity**‡
Follow-up > 1 week – all stages – ICU			
Salvadalena 1992[36]a; VandenBosch 1996[46]a	Yes/no	50 (95%CI 27 to 73) and 52 (95%CI 33 to 70)	80 (95%CI 69 to 88) and 59 (95%CI 47 to 70)

*** Percentage*
a Study of which sensitivity and specificity are presented.

O.4 Quality of the studies

Table 25: Quality of the studies

Study	Selection bias*	Risk tool bias**	Outcome bias***	Analysis bias****
Andersen 1982[1]	Low	High	Low	High
Anthony 2003[2]	High	High	Low	High
Barnes 1993[3]	High	High	Low	High
Bergstrom 1987a[5]	Low	High	Low	Very high
Bergstrom 1987b[6]	Low	High	Low	Very High
Bergstrom 1998[4]	Low	High	Low	High
Braden 1994[7]	Low	High	Low	High
Capobianco 1996[8]	Low	High	Low	High
Chan 2009[9]	High	High	Low	High
Compton 2008[10]	Very high	High	Low	High
Curley 2003[12]	Low	High	Low	High
de Souza 2010[13]	Very high	High	Low	High
Edwards 1995[14]	Low	High	Low	Very high
Feuchtinger 2007[15]	Low	High	Low	High
Goodridge 1998[16]	High	High	Low	High
Halfens 2000[17]	High	High	Low	High
Hatanaka 2008[18]	High	High	High	High
Jalali 2005[20]	High	Very high	Low	High
Kim 2009[21]	High	High	Low	High
Kwong 2005[22]	High	High	Low	Very high
Langemo 1991[23]	High	High	Low	Very high
Lewicki 2000[24]	High	High	Low	Very high
Lincoln 1986[25]	High	High	Low	Very high

Study	Selection bias*	Risk tool bias**	Outcome bias***	Analysis bias****
Lindgren 2002[26]	High	High	Low	High
Lothian 1989[27]	Very high	Very high	Very high	High
Lyder 1999[28]	High	High	High	High
Ongoma 2006[30]	High	Very high	Very high	High
Page 2011[31]	Very high	High	High	Very high
Pang 1998[32]	High	High	Low	High
Perneger 2002[33]	High	High	Low	High
Ramundo 1995[35]	High	High	Low	Very high
Salvadalena 1992[36]	High	High	Low	High
Schoonhoven 2002[37]	High	High	Low	Low
Seongsook 2004[38]	Very high	High	Low	High
Serpa 2009[39]	High	High	High	Very high
Serpa 2011[40]	High	High	High	Very high
Smith 1989[41]	Very high	High	High	High
Stotts 1988[42]	Very high	High	Low	High
Suriadi 2006[44]	High	High	Low	High
Suriadi 2008[43]	High	High	Low	High
Towey 1988[45]	Low	High	High	High
VandenBosch 1996[46]	High	High	Low	High
Wai-Han 1997[47]	High	High	High	Very high
Weststrate 1998[50]	High	High	High	High

** inappropriate patient enrolment, inappropriate study design, not representative population*
*** unclear definition and measurement of predictive test, absence of imputation technique or unclear description of exclusion, inadequate threshold*
**** unclear definition and measurement of reference test, inappropriate duration*
***** no use of time to event analysis, number of events < 100, reason for missing data not reported*

O.5 Incidence and predictive ability of risk assessment scales - all thresholds

Table 26: Braden scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Barnes 1993[3]	2 weeks	6.1	≤ 16	72.7	90.6
Braden 1994[7]	48-72 hours‡	NR	≤ 14	29.0	97.0
			≤ 15	43.0	95.0
			≤ 16	46.0	84.0
			≤ 17	61.0	78.0
			≤ 18	79.0	68.0
			≤ 19	93.0	51.0
			≤ 20	96.0	35.0
	4 weeks	27.5	≤ 14	21.4	95.9
			≤ 15	32.1	94.6
			≤ 16	50.0	89.2
			≤ 17	57.1	85.1
			≤ 18	78.6	74.3
			≤ 19	85.7	59.3
			≤ 20	92.9	43.2
Bergstrom 1987a[5] (a)	6 weeks	7.1	≤ 9	14.3	100.0
			≤ 10	14.3	98.9
			≤ 13	28.6	98.9
			≤ 14	42.9	98.9
			≤ 15	71.4	94.6
			≤ 16	100.0	90.2
			≤ 17	100.0	88.0
			≤ 18	100.0	82.6
			≤ 19	100.0	73.9
			≤ 20	100.0	65.2
			≤ 21	100.0	50.0

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 22	100.0	35.9
			≤ 23	100.0	0.0
(b)	12 weeks	9.0	≤ 8	11.1	95.6
			≤ 9	11.1	91.2
			≤ 11	22.2	89.0
			≤ 12	44.4	86.8
			≤ 13	55.6	83.5
			≤ 14	66.7	78.0
			≤ 15	77.8	73.
			≤ 16	100.0	63.7
			≤ 17	100.0	60.4
			≤ 18	100.0	50.5
			≤ 19	100.0	42.9
			≤ 20	100.0	31.9
			≤ 21	100.0	26.4
			≤ 22	100.0	9.9
			≤ 23	100.0	0.0
Bergstrom 1987b[6]	2 weeks	40.0	≤ 9	8.3	100.0
			≤ 10	8.3	97.2
			≤ 11	16.7	91.7
			≤ 12	33.3	88.9
			≤ 13	58.3	77.8
			≤ 14	70.8	75.0
			≤ 15	75.0	66.7
			≤ 16	83.3	63.9
			≤ 17	87.5	50.0
			≤ 18	91.7	38.9
			≤ 19	91.7	25.0
			≤ 20	95.8	13.9
			≤ 21	95.8	5.6

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 22	100.0	0.0
Bergstrom 1998[4] (c)	48-72 hours‡	NR	≤ 9	4.0	100.0
			≤ 10	12.0	100.0
			≤ 11	19.0	99.0
			≤ 12	31.0	99.0
			≤ 13	38.0	98.0
			≤ 14	38.0	95.0
			≤ 15	46.0	90.0
			≤ 16	58.0	84.0
			≤ 17	62.0	76.0
			≤ 18	88.0	68.0
			≤ 19	100.0	59.0
			≤ 20	100.0	40.0
			≤ 21	100.0	23.0
	11 days	8.5	≤ 9	7.7	100.0
			≤ 10	11.5	100.0
			≤ 11	11.5	98.9
			≤ 12	15.4	98.9
			≤ 13	15.4	97.9
			≤ 14	15.4	97.1
			≤ 15	23.1	92.9
			≤ 16	30.8	88.9
			≤ 17	38.5	83.9
			≤ 18	46.2	68.9
			≤ 19	100.0	58.9
			≤ 20	100.0	40.0
			≤ 21	100.0	22.9
(d)	48-72 hours‡	NR	≤ 9	0.0	99.0
			≤ 10	10.0	99.0
			≤ 11	10.0	98.0

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 12	10.0	98.0
			≤ 13	10.0	97.0
			≤ 14	20.0	96.0
			≤ 15	20.0	94.0
			≤ 16	30.0	90.0
			≤ 17	50.0	85.0
			≤ 18	60.0	81.0
			≤ 19	80.0	73.0
			≤ 20	80.0	69.0
			≤ 21	90.0	41.0
	11 days	7.4	≤ 9	0.0	99.2
			≤ 10	0.0	99.2
			≤ 11	0.0	99.2
			≤ 12	0.0	98.1
			≤ 13	28.6	98.1
			≤ 14	28.6	96.9
			≤ 15	52.4	93.9
			≤ 16	52.4	92.0
			≤ 17	61.9	87.0
			≤ 18	71.4	78.9
			≤ 19	71.4	70.9
			≤ 20	90.5	50.2
			≤ 21	90.5	32.2
(e)	48-72 hours‡	NR	≤ 9	0.0	99.0
			≤ 10	2.0	99.0
			≤ 11	2.0	99.0
			≤ 12	5.0	99.0
			≤ 13	13.0	99.0
			≤ 14	23.0	97.0
			≤ 15	33.0	93.0

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 16	41.0	88.0
			≤ 17	56.0	81.0
			≤ 18	72.0	68.0
			≤ 19	67.0	48.0
			≤ 20	83.0	34.0
			≤ 21	97.0	17.0
	11 days	23.9	≤ 9	0.0	99.0
			≤ 10	19.7	99.0
			≤ 11	29.5	97.9
			≤ 12	8.2	97.9
			≤ 13	13.1	97.9
			≤ 14	19.7	95.9
			≤ 15	31.1	94.8
			≤ 16	49.2	90.2
			≤ 17	60.7	86.1
			≤ 18	80.3	73.2
			≤ 19	86.9	57.2
			≤ 20	93.4	40.2
			≤ 21	98.4	25.3
Capobianco 1996[8]	2 weeks	28.0	≤ 12	28.6	97.2
			≤ 13	28.6	97.2
			≤ 14	28.6	97.2
			≤ 15	35.7	94.4
			≤ 16	42.9	91.7
			≤ 17	57.1	91.7
			≤ 18	71.4	83.3
			≤ 19	85.7	77.8
			≤ 20	92.9	66.7
Chan 2009[9]	9 days	9.1	≤ 16	66.7	64.2
			≤ 17	72.2	40.8

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 18	88.9	21.2
de Souza 2010[13] (f)	3 months	3.9	≤ 13	56.8	71.9
(g)	3 months	3.9	≤ 17	71.4	75.8
Feuchtinger 2007[15]	4 days	62.3	≤ 9	19.2	100.0
			≤ 10	23.1	100.0
			≤ 11	30.8	100.0
			≤ 16	76.9	29.6
			≤ 20	96.2	3.7
Goodridge 1998[16]	3 months	9.7	≤ 11	12.5	97.3
			≤ 16	25.0	85.6
			≤ 18	50.0	52.3
Halfens 2000[17]	NR	58.1	≤ 10	1.1	100.0
			≤ 11	3.2	100.0
			≤ 12	5.4	99.3
			≤ 13	11.8	97.8
			≤ 14	17.7	97.0
			≤ 15	22.0	94.8
			≤ 16	32.3	91.8
			≤ 17	40.9	90.3
			≤ 18	51.1	85.8
			≤ 19	61.3	79.9
			≤ 20	73.7	70.1
			≤ 21	78.5	56.7
			≤ 22	88.2	42.5
			≤ 23	100.0	29.9
Jalali 2005[20]	14 days	9.1	NR	52.7	100.0
Kim 2009[21]	90 days	18.3	≤ 14	92.5	69.8
Kwong 2005[22]	21 days	2.1	≤ 14	88.9	71.9
Langemo 1991[23] (h)	16 days	14.9	≤ 15	54.5	93.7

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 16	63.6	87.3
(i)	31 days	28.0	≤ 18	57.1	61.1
Pang 1998[32]	2 weeks	19.8	≤ 18	90.5	62.4
Lyder 1999[28] (j)	NR‡	NR	≤ 18	81.0	100.0
(k)	NR‡	NR	≤ 16	77.0	50.0
(l)	NR‡	NR	≤ 18	90.0	14.0
Ramundo 1995[35]	4 weeks	14.6	≤ 11	14.3	97.6
			≤ 12	14.3	95.1
			≤ 13	14.3	95.1
			≤ 14	14.3	90.2
			≤ 15	14.3	82.9
			≤ 16	28.6	80.5
			≤ 17	42.9	63.4
			≤ 18	100.0	34.1
			≤ 19	100.0	22.0
			≤ 20	100.0	12.2
			≤ 21	100.0	4.9
			≤ 22	100.0	0.0
Salavadalena 1992[36]	6 months	20.2	≤ 9	0.0	98.7
			≤ 10	5.0	97.5
			≤ 11	5.0	91.1
			≤ 12	15.0	89.9
			≤ 13	20.0	86.1
			≤ 14	30.0	79.
			≤ 15	30.0	77.2
			≤ 16	40.0	69.6
			≤ 17	45.0	63.3
			≤ 18	60.0	54.4
			≤ 19	80.0	43.0

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
			≤ 20	85.0	31.6
			≤ 21	95.0	13.9
			≤ 22	100.0	1.3
			≤ 23	100.0	0.0
Schoonhoven 2002[37]	12 weeks	11.0	≤ 17	43.7	67.8
Seongsook 2004[38]	NR	31.3	≤ 16	97.1	26.0
Serpa 2011[40]	48 hours	11.1	≤ 12	87.5	64.1
	4 days	11.1	≤ 13	75.0	81.3
	6 days	11.1	≤ 13	75.0	82.8
Suriadi 2006[44]	21 days	33.3	≤ 14	80.0	54.3
VandenBosch 1996[46]	2 weeks	28.8	≤ 17	59.0	NR
			≤ 18	NR	79.0

** Percentage*
‡ No raw data was available to recalculate the sensitivity and specificity
NR: not reported
(a) ward one in Bergstrom 1987a study
(b) ward two in Bergstrom 1987a study
(c) tertiary care hospitals
(d) veteran medical centres
(e) skilled nursing facilities
(f) group of patients with a Braden score < 18 on admission
(g) group of patients with a Braden score < 19 on admission
(h) hospitalized patients
(i) long-term care patients
(j) black elders ≥ 75 yrs
(k) black elders < 75 yrs

(l) Latino/Hispanic < 75 yrs

Table 27: Extended Braden scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Halfens 2000[17]	NR	58.1	≤ 11	0.5	100.0
			≤ 12	1.6	100.0
			≤ 13	2.2	100.0
			≤ 14	3.8	99.3
			≤ 15	6.5	98.5
			≤ 16	12.4	97.9
			≤ 17	17.7	96.3
			≤ 18	24.2	94.8
			≤ 19	32.8	91.0
			≤ 20	40.9	88.8
			≤ 21	51.1	85.1
			≤ 22	62.9	79.1
			≤ 23	73.7	69.4
			≤ 24	78.5	55.2
			≤ 25	88.2	42.5
			≤ 26	100.0	29.1

** Percentage*
NR: not reported

Table 28: Modified Braden scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Chan 2009[9]	9 days	9.1	≤ 17	38.9	79.9
			≤ 18	55.6	72.6
			≤ 19	88.9	62.0
Kwong 2005[22]	21 days	2.1	≤ 16	88.9	75.0

** Percentage*

Table 29: Braden-Q scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Curley 2003[12]	10 days	26.7	≤ 10	3.5	100.0
			≤ 11	16.3	97.0
			≤ 12	47.7	92.8
			≤ 13	67.4	89.0
			≤ 14	72.1	78.8
			≤ 15	75.6	67.8
			≤ 16	88.4	58.1
			≤ 17	91.9	44.1
			≤ 18	100.0	30.1
			≤ 19	100.0	19.9
			≤ 20	100.0	8.1

** Percentage*

Table 30: Norton scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Kwong 2005[22]	21 days	2.1	≤ 14	88.9	61.0
Lincoln 1986[25]	26 days	13.9	≤ 14	0.0	93.5
Pang 1998[32]	2 weeks	19.8	≤ 16	81.0	58.8
Schoonhoven 2002[37]	12 weeks	11.0	≤ 15	45.9	60.3
Smith 1989[41]	NR	29.7	≤ 16	60.0	31.0
Stotts 1988[42]	3 weeks	17.3	≤ 14	16.4	94.4
Wai-Hang 1997[47]	4 weeks	4.3	≤ 14	75.0	66.7

** Percentage*
NR: Not reported

Table 31: Modified Norton scale (ICU)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Feuchtinger 2007[15]	4 days	62.3	≤ 19	26.9	100.0
			≤ 21	34.6	92.6
			≤ 23	42.3	88.9
			≤ 25	57.7	48.1

** Percentage*

Table 32: Modified Norton scale (South African Hospital)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Ongoma 2005[30]	1 week	37.9	≤ 20	92.0	29.3

** Percentage*

Table 33: Waterlow scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Anthony 2003[2]	NR	0.4	≥ 10	82.3	85.2
			≥ 15	48.8	94.5
			≥ 20	16.7	98.1
Compton 2008[10]	13 days	17.3	NR	37.2	94.6
Edwards 1995[14]	8 weeks	6.5	NR	100.0	10.3
Jalali 2005[20]	14 days	9.1	NR	63.5	83.3
Pang 1998[32]	2 weeks	19.8	≥ 16	95.2	43.5
Serpa 2009[39]	48 hours	7.1	≥ 17	71.4	67.0
	4 days	7.1	≥ 20	85.7	40.7
	6 days	7.1	≥ 20	85.7	33.0
Schoonhoven 2002[37]	12 weeks	11.0	≥ 10	89.6	22.4
Smith 1989[41]	NR	29.7	≥ 16	73.3	38.0
Wai-Han 1997[47]	4 weeks	4.3	≥ 10	87.5	28.2
Weststrate 1998[50]	183 days	7.9	≥ 15	80.9	28.5

** Percentage*

NR: not reported

Table 34: Andersen scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Andersen 1982[1]	3 months	1.2	≥ 2	87.5	86.7

** Percentage*

Table 35: Pressure Sore Prediction Score scale (PSPS)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Lothian 1989[27]	3 weeks	4.3	> 6	88.7	76.0

** Percentage*

Table 36: Knoll scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Towey 1988[45]	28 days	46.7	≥ 12	85.7	56.3

** Percentage*

Table 37: <Insert Table Title here>

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Kim 2009[21]	90 days	18.3	≤ 28	95.0	81.6
Seongsook 2004[38]	NR	31.3	≤ 24	88.6	61.0

** Percentage*
NR: not reported

Table 38: Sunderland Pressure Sore Risk Calculator (modified Cubbin-Jackson)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Ongoma 2005[30]	1 week	37.9	≤ 34	80.0	70.7

Table 39: Risk Assessment Pressure Sore scale (RAPS)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Lindgren 2002[26]	12 weeks	11.7	≤ 31	31.5	84.6
			≤ 32	33.3	80.2
			≤ 33	38.9	75.3
			≤ 34	46.3	69.4
			≤ 35	50.0	64.3
			≤ 36	57.4	57.6
			≤ 37	70.4	46.5
			≤ 38	77.8	34.8

** Percentage*

Table 40: Fragmment scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Perneger 2002[33]	3 weeks	29.9	= 0	91.6	34.2
			≤ 1	78.7	53.5
			≤ 2	76.7	71.9
			≤ 3	62.1	85.0
			≤ 4	49.7	91.0
			≤ 5	40.2	94.2
			≤ 6	27.0	97.6
			≤ 7	17.7	98.9
			≤ 8	2.2	99.5

** Percentage*

Table 41: Douglas scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Seongsook 2004[38]	NR	31.3	≤ 18	100.0	18.2

** Percentage*
NR: Not reported

Table 42: Grosnell scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Jalali 2005[20]	2 weeks	9.1	NR	85.1	83.3

** Percentage*
NR: not reported

Table 43: Song and Choi scale

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Kim 2009[21]	90 days	18.3	≤ 21	95.0	69.3

** Percentage*

Table 44: 4-factor model

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Feuchtinger 2007[15]	4 days	62.3	≥ 2	84.6	29.6

** Percentage*

Table 45: Suriadi and Sanada scale (SS)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Suriadi 2008[43]	NR	28.5	≥ 0	100.0	0.0
			≥ 2	97.2	42.0
			≥ 3	97.2	53.0
			≥ 4	80.6	82.9
			≥ 5	72.2	86.7
			≥ 6	61.1	92.3
			≥ 7	58.3	95.0
			≥ 9	6.9	100.0

** Percentage*
NR: not reported

Table 46: The Northern Hospital Pressure Ulcer Prevention Plan (TNH-PUPP)

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Page 2011[31]	NR	4.2	≥ 1	100.0	34.2
			≥ 2	85.7	62.0
			≥ 3	71.4	81.0
			≥ 4	71.4	88.0
			≥ 5	42.9	96.2
			≥ 6	57.1	99.4

** Percentage*
NR: not reported

Table 47: Clinical judgement

Study	Time point	Incidence*	Cut-off score	Sensitivity*	Specificity*
Salvadalena 1992[36]	6 months	20.2	Yes/no	50.0	79.7
VandenBosch 1996[46]	2 weeks	28.2	Yes/no	51.7	58.1

O.6 Forest plots area under the receiver operating characteristics curve (AUC)

- Perneger 2002 and Schoonhoven 2002 were the only studies with more than 100 events; Serpa 2011 had fewer than 10 events.
- The proportion of pressure ulcers was less than 10% for Chan 2009;10-20% for de Souza 2010, Kim 2009, Perneger 2002, Schoonhoven 2002, Serpa 2011; 20-50% for Hatanaka 2008, Seongsook 2004, Suriadi 2006
- De Souza 2010 and Suriadi 2006 did not give preventative treatment; Schoonhoven 2002 and Perneger 2002 both gave less than half the patients preventative treatment; Hatanaka 2007, Kim 2009 and Seongsook 2004 gave preventative treatment to all patients.
- Suriadi 2006, Seongsook 2004 and Serpa 2011 were conducted in intensive care; Hatanaka 2007 had bedridden hospitalised patients and the patients in de Souza 2010 were in long term care.
- Serpa 2011 included a selected risk group (low risk patients missing)

- Kim 2009 and Suriadi 2006 had a mean age of between 50 and 60 years; Schoonhoven 2002, Seongsook 2004, Serpa 2011, had a mean patient age of 60-70 years; Chan 2009, de Souza 2010 and Hatanake 2007 had a mean age of 70-80 years;

Table 48: Braden scale

Sample sizes: Chan 2009 (n=197); de Souza 2010 (n=233); Hatanaka 2007 (n=149); Kim 2008 (219); Perneger 2002 (n=1190); Schoonhoven 2002 (n=1229); Seongsook 2004 (n=112); Serpa 2011 (n=72); Suriadi 2006 (n=105)







O.6.1 Modified Braden scale



O.6.2 Braden-Q scale



O.6.3 Norton scale

Sample sizes: Perneger 2002 (n=1190); Schoonhoven 2002 (n=1229);



O.6.4 Waterlow scale







O.6.5 Cubbin-Jackson scale

Both studies reported that all patients received preventative treatment. Confidence intervals were not reported in either study, but were estimated (using the number of patients) to be: Seongsook 2004 AUC 83% (95%CI 75 to 89) and Kim 2009 AUC 90% (95%CI 85 to 94).



O.6.6 Douglas scale



O.6.7 Fragmment scale



O.6.8 Song and choi scale



O.6.9 The Northern Hospital Pressure Ulcer Prevention Plan



Comparison of scales in the same study

O.6.10 Schoonhoven 2002 (general patient group, mean age 60.1 years, 1229 patients, 135 events (11%), 5% received preventative treatment)



O.6.11 Perneger 2002 (general patient group, 1190 patients, 170 events (14%), 24% received preventative interventions)



O.6.12 Seongsook 2004 (ICU, mean age 62 years, 112 patients, 35 events (31%), all patients received preventative measures)

Confidence intervals were not reported, but were estimated (using the number of patients) to be: Braden AUC 71% (95%CI 62 to 79); Cubbin Jackson AUC 83% (95%CI 75 to 89) and Douglas scale AUC 79% (95%CI 70 to 86%).



O.6.13 Chan 2009 (Orthopaedic patients, mean age 79 years, 197 patients, 18 events (9%), preventative measures applied 'as normal practice')



O.6.14 Kim 2009 (Surgical ICU patients, mean age 58 years, 219 patients, 40 events (18%), all patients received preventative measures)



O.6.15 Serpa 2009 (ICU patients, 98 patients, 7 events (7%), selected group (no low risk patients), mean age 71 years, preventative measures not mentioned)



O.6.16 Serpa 2011 (ICU patients, 72 patients, 8 events (11%), selected group (no low risk patients), mean age 61 years, preventative measures according to the institution)



O.7 Forest plots and sensitivity

O.7.1 Braden scale cut-off score 17 – follow-up < 1 week – general population – all grades

Sample sizes: Bergstrom 1998 (1): 306; Bergstrom 1998 (2): 282; Bergstrom 1998 (3): 61; Braden 1994: 102

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1998 (1)	0	0	0	0	0.62	0.76
Bergstrom 1998 (2)	0	0	0	0	0.50	0.85
Bergstrom 1998 (3)	0	0	0	0	0.56	0.81
Braden 1994	0	0	0	0	0.61	0.78



No raw data available
Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2): veteran medical centre; Bergstrom 1998 (3): skilled nursing facility

O.7.2 Braden scale cut-off score 18 – follow-up < 1 week – general population – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1998 (1)	0	0	0	0	0.88	0.68
Bergstrom 1998 (2)	0	0	0	0	0.60	0.81
Bergstrom 1998 (3)	0	0	0	0	0.72	0.68
Braden 1994	0	0	0	0	0.79	0.68



No raw data available
Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2): veteran medical centre; Bergstrom 1998 (3): skilled nursing facility

O.7.3 : Braden scale cut-off score 19 – follow-up < 1 week – general population – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1998 (1)	0	0	0	0	1.00	0.40
Bergstrom 1998 (2)	0	0	0	0	0.80	0.73
Bergstrom 1998 (3)	0	0	0	0	0.67	0.48
Braden 1994	0	0	0	0	0.93	0.51



No raw data available
Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2): veteran medical centre; Bergstrom 1998 (3): skilled nursing facility

O.7.4 Braden scale cut-off score 12 – follow-up 48 hours – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Serpa 2011	7	23	1	41	0.88 [0.47, 1.00]	0.64 [0.51, 0.76]



O.7.5 Braden scale cut-off score 13 – follow-up 4 and 6 days – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Serpa 2011 [1]	6	11	2	53	0.75 [0.35, 0.97]	0.83 [0.71, 0.91]
Serpa 2011 [2]	6	12	2	52	0.75 [0.35, 0.97]	0.81 [0.70, 0.90]



Serpa 2011 1: 4 days; Serpa 2011 2: 6 days

O.7.6 Braden scale cut-off score 16 – follow-up < 1 week – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Feuchtinger 2007	20	19	6	8	0.77 [0.56, 0.91]	0.30 [0.14, 0.50]



O.7.7 Braden scale cut-off score 18 – follow-up > 1 week – general population – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1987a (1)	7	16	0	76	1.00 [0.59, 1.00]	0.83 [0.73, 0.90]
Bergstrom 1987a (2)	9	45	0	46	1.00 [0.66, 1.00]	0.51 [0.40, 0.61]
Bergstrom 1998 (1)	12	62	14	218	0.46 [0.27, 0.67]	0.78 [0.73, 0.83]
Bergstrom 1998 (2)	15	55	6	206	0.71 [0.48, 0.89]	0.79 [0.73, 0.84]
Bergstrom 1998 (3)	49	52	12	142	0.80 [0.68, 0.89]	0.73 [0.66, 0.79]
Braden 1994	22	19	5	55	0.81 [0.62, 0.94]	0.74 [0.63, 0.84]
Capobianco 1996	10	6	4	30	0.71 [0.42, 0.92]	0.83 [0.67, 0.94]
Chan 2009	16	141	2	38	0.89 [0.65, 0.99]	0.21 [0.15, 0.28]
Goodridge 1998	16	142	16	156	0.50 [0.32, 0.68]	0.52 [0.47, 0.58]
Langemo 1991 (2)	4	7	3	11	0.57 [0.18, 0.90]	0.61 [0.36, 0.83]
Lyder 1999 (1)	0	0	0	0	0.81	1.00
Lyder 1999 (3)	0	0	0	0	0.90	0.14
Pang 1998	19	32	2	53	0.90 [0.70, 0.99]	0.62 [0.51, 0.73]
Salvadalena 1992	12	36	8	43	0.60 [0.36, 0.81]	0.54 [0.43, 0.66]



No raw data for Lyder 1991
Bergstrom 1987a (1): ward one; Bergstrom 1987a (2): ward two; Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2):

veteran medical centre; Bergstrom 1998 (3): skilled nursing facility; Langemo 1991 (2): skilled nursing facility; Lyder 1999 (1): black elders ≥ 75 yrs; Lyder 1999 (2): Latino/Hispanic < 75 yrs

O.7.8 Braden scale cut-off score 19 – follow-up > 1 week – general population – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1987a (1)	7	24	0	68	1.00 [0.59, 1.00]	0.74 [0.64, 0.83]
Bergstrom 1987a (2)	9	52	0	39	1.00 [0.66, 1.00]	0.43 [0.33, 0.54]
Bergstrom 1998 (1)	12	87	14	193	0.46 [0.27, 0.67]	0.69 [0.63, 0.74]
Bergstrom 1998 (2)	15	76	6	185	0.71 [0.48, 0.89]	0.71 [0.65, 0.76]
Bergstrom 1998 (3)	53	83	8	111	0.87 [0.76, 0.94]	0.57 [0.50, 0.64]
Braden 1994	24	30	4	44	0.86 [0.67, 0.96]	0.59 [0.47, 0.71]
Capobianco 1996	12	8	2	28	0.86 [0.57, 0.98]	0.78 [0.61, 0.90]
Salvadalena 1992	16	45	4	34	0.80 [0.56, 0.94]	0.43 [0.32, 0.55]



Bergstrom 1987a (1): ward one; Bergstrom 1987a (2): ward two; Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2): veteran medical centre; Bergstrom 1998 (3): skilled nursing facility

O.7.9 Braden scale cut-off score 20 – follow-up > 1 week – general population – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Bergstrom 1987a (1)	7	32	0	60	1.00 [0.59, 1.00]	0.65 [0.55, 0.75]
Bergstrom 1987a (2)	9	62	0	29	1.00 [0.66, 1.00]	0.32 [0.22, 0.42]
Bergstrom 1998 (1)	17	126	9	154	0.65 [0.44, 0.83]	0.55 [0.49, 0.61]
Bergstrom 1998 (2)	19	130	2	131	0.90 [0.70, 0.99]	0.50 [0.44, 0.56]
Bergstrom 1998 (3)	57	116	4	78	0.93 [0.84, 0.98]	0.40 [0.33, 0.47]
Braden 1994	26	42	2	32	0.93 [0.76, 0.99]	0.43 [0.32, 0.55]
Capobianco 1996	13	12	1	24	0.93 [0.66, 1.00]	0.67 [0.49, 0.81]
Salvadalena 1992	17	54	3	25	0.85 [0.62, 0.97]	0.32 [0.22, 0.43]



Bergstrom 1987a (1): ward one; Bergstrom 1987a (2): ward two; Bergstrom 1998 (1): tertiary hospital; Bergstrom 1998 (2): veteran medical centre; Bergstrom 1998 (3): skilled nursing facility

O.7.10 Braden scale cut-off score 15 – follow-up > 1 week – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Braden 1994	9	4	19	70	0.32 [0.16, 0.52]	0.95 [0.87, 0.99]



O.7.11 Figure 29: Braden scale cut-off score 16 – follow-up > 1 week – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Braden 1994	14	8	14	66	0.50 [0.31, 0.69]	0.89 [0.80, 0.95]
Seongsook 2004	34	57	1	20	0.97 [0.85, 1.00]	0.26 [0.17, 0.37]



O.7.12 Braden scale cut-off score 17 – follow-up > 1 week – ICU – all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Braden 1994	16	11	12	63	0.57 [0.37, 0.76]	0.85 [0.75, 0.92]



O.7.13 Braden scale cut-off score 17 – follow-up > 1 week – general population – stage 2+

Study	TP	FP	FN	TN	Sensitivity	Specificity
Ramundo 1995	3	15	4	26	0.43 [0.10, 0.82]	0.63 [0.47, 0.78]



O.7.14 Braden scale cut-off score 18 – follow-up > 1 week – general population – stage 2+

Study	TP	FP	FN	TN	Sensitivity	Specificity
Ramundo 1995	7	27	0	14	1.00 [0.59, 1.00]	0.34 [0.20, 0.51]



O.7.15 Braden scale cut-off score 19 – follow-up > 1 week – general population – stage 2+

Study	TP	FP	FN	TN	Sensitivity	Specificity
Ramundo 1995	7	32	0	9	1.00 [0.59, 1.00]	0.22 [0.11, 0.38]



O.7.16 Braden-Q scale cut-off score 16 – follow-up > 1 week – paediatric ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Curley 2003	76	99	10	137	0.88 [0.80, 0.94]	0.58 [0.51, 0.64]



O.7.17 Braden-Q scale cut-off score 17 – follow-up > 1 week – paediatric ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Curley 2003	79	132	7	104	0.92 [0.84, 0.97]	0.44 [0.38, 0.51]



O.7.18 Norton scale cut-off score 14 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Kwong 2005	8	164	1	256	0.89 [0.52, 1.00]	0.61 [0.56, 0.66]
Lincoln 1986	0	2	5	29	0.00 [0.00, 0.52]	0.94 [0.79, 0.99]
Stotts 1988	11	18	56	302	0.16 [0.08, 0.27]	0.94 [0.91, 0.97]
Wai-Han 1997	6	59	2	118	0.75 [0.35, 0.97]	0.67 [0.59, 0.74]



O.7.19 Norton scale cut-off score 15 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Schoonhoven 2002	62	434	73	660	0.46 [0.37, 0.55]	0.60 [0.57, 0.63]



O.7.20 Norton scale cut-off score 16 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Pang 1998	17	35	4	50	0.81 [0.58, 0.95]	0.59 [0.48, 0.69]
Smith 1989	18	49	12	22	0.60 [0.41, 0.77]	0.31 [0.21, 0.43]



O.7.21 Waterlow scale cut-off score 17 – follow-up 48 hours – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Serpa 2009	5	30	2	61	0.71 [0.29, 0.96]	0.67 [0.56, 0.77]



O.7.22 Waterlow scale cut-off score 20 – follow-up 4 days and 6 days – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Serpa 2009 [1]	6	54	1	37	0.86 [0.42, 1.00]	0.41 [0.30, 0.51]
Serpa 2009 [2]	6	61	1	30	0.86 [0.42, 1.00]	0.33 [0.23, 0.44]



Serpa 2009 1: 4 days; Serpa 2009 2: 6 days

O.7.23 Waterlow scale cut-off score 10 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Anthony 2003	167	6757	36	38775	0.82 [0.76, 0.87]	0.85 [0.85, 0.85]
Schoonhoven 2002	121	849	14	245	0.90 [0.83, 0.94]	0.22 [0.20, 0.25]
Wai-Han 1997	7	127	1	50	0.88 [0.47, 1.00]	0.28 [0.22, 0.35]



O.7.24 Waterlow scale cut-off score 15 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Anthony 2003	99	2519	104	43013	0.49 [0.42, 0.56]	0.94 [0.94, 0.95]



O.7.25 Waterlow scale cut-off score 16 – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Pang 1998	20	48	1	37	0.95 [0.76, 1.00]	0.44 [0.33, 0.55]
Smith 1989	22	44	8	27	0.73 [0.54, 0.88]	0.38 [0.27, 0.50]



O.7.26 Waterlow scale cut-off score 15 – follow-up > 1 week – ICU – stage 2+

Study	TP	FP	FN	TN	Sensitivity	Specificity
Weststrate 1998	38	391	9	156	0.81 [0.67, 0.91]	0.29 [0.25, 0.33]



O.7.27 Cubbin-Jackson scale cut-off score 24 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Seongsook 2004	31	30	4	47	0.89 [0.73, 0.97]	0.61 [0.49, 0.72]



O.7.28 Cubbin-Jackson scale cut-off score 28 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Kim 2009	38	33	2	146	0.95 [0.83, 0.99]	0.82 [0.75, 0.87]



O.7.29 Fragmment scale cut-off score 1 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Perneger 2002	280	388	76	446	0.79 [0.74, 0.83]	0.53 [0.50, 0.57]



O.7.30 Fragmment scale cut-off score 2 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Perneger 2002	273	234	83	600	0.77 [0.72, 0.81]	0.72 [0.69, 0.75]



O.7.31 Fragmment scale cut-off score 3 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Perneger 2002	221	125	135	709	0.62 [0.57, 0.67]	0.85 [0.82, 0.87]



O.7.32 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 2 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Page 2011	6	60	1	98	0.86 [0.42, 1.00]	0.62 [0.54, 0.70]



O.7.33 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 3 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Page 2011	5	30	2	128	0.71 [0.29, 0.96]	0.81 [0.74, 0.87]



O.7.34 The Northern Hospital Pressure Ulcer Prevention plan cut-off score 4 – follow-up > 1 week – general population– all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Page 2011	5	19	2	139	0.71 [0.29, 0.96]	0.88 [0.82, 0.93]



O.7.35 Douglas scale cut-off score 18 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Seongsook 2004	35	63	0	14	1.00 [0.90, 1.00]	0.18 [0.10, 0.29]



O.7.36 Song and Choi scale cut-off score 2 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Kim 2009	38	55	2	124	0.95 [0.83, 0.99]	0.69 [0.62, 0.76]



O.7.37 Suriadi and Sanada scale cut-off score 3 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Suriadi 2008	70	85	2	96	0.97 [0.90, 1.00]	0.53 [0.45, 0.60]



O.7.38 Suriadi and Sanada scale cut-off score 4 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Suriadi 2008	58	31	14	150	0.81 [0.70, 0.89]	0.83 [0.77, 0.88]



O.7.39 Suriadi and Sanada scale cut-off score 5 – follow-up > 1 week – ICU – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Suriadi 2008	52	24	20	157	0.72 [0.60, 0.82]	0.87 [0.81, 0.91]



O.7.40 Clinical judgement – follow-up > 1 week – general population – all stages

Study	TP	FP	FN	TN	Sensitivity	Specificity
Salvadalena 1992	10	16	10	63	0.50 [0.27, 0.73]	0.80 [0.69, 0.88]
VandeBosch 1996	15	30	14	43	0.52 [0.33, 0.71]	0.59 [0.47, 0.70]



Figure 1: Norton, Waterlow, Braden scales and clinical judgement – all stages, general population – for studies reporting more than one scale

Norton

Study	TP	FP	FN	TN	Threshold	Sensitivity	Specificity
Jalali 2005_Norton	36	0	38	156	16.0	0.49 [0.37, 0.61]	1.00 [0.98, 1.00]
Pang 1998_Norton	17	35	4	50	16.0	0.81 [0.58, 0.95]	0.59 [0.48, 0.69]
Schoonhoven 2002_Norton	62	434	73	660	15.0	0.46 [0.37, 0.55]	0.60 [0.57, 0.63]
Waihan 1997_Norton	6	59	2	118	14.0	0.75 [0.35, 0.97]	0.67 [0.59, 0.74]



Waterlow

Study	TP	FP	FN	TN	Threshold	Sensitivity	Specificity
Jalali 2005_Waterlow	47	27	27	129	16.0	0.64 [0.52, 0.74]	0.83 [0.76, 0.88]
Pang 1998_Waterlow	20	48	1	37	16.0	0.95 [0.76, 1.00]	0.44 [0.33, 0.55]
Schoonhoven 2002_Waterlow	121	849	14	245	10.0	0.90 [0.83, 0.94]	0.22 [0.20, 0.25]
Waihan 1997_Waterlow	7	127	1	50	10.0	0.88 [0.47, 1.00]	0.28 [0.22, 0.35]



Clinical judgement

Study	TP	FP	FN	TN	Threshold	Sensitivity	Specificity
Salvadalena 1992_clinjudg	10	16	10	63	100.0	0.50 [0.27, 0.73]	0.80 [0.69, 0.88]
Vandenbosch 1996_clinjudg	15	30	14	43	100.0	0.52 [0.33, 0.71]	0.59 [0.47, 0.70]



Braden

Study	TP	FP	FN	TN	Threshold	Sensitivity	Specificity
Jalali 2005_Braden	39	0	35	156	18.0	0.53 [0.41, 0.64]	1.00 [0.98, 1.00]
Pang 1998_Braden	19	32	2	53	18.0	0.90 [0.70, 0.99]	0.62 [0.51, 0.73]
Salvadalena 1992_braden	12	36	8	43	18.0	0.60 [0.36, 0.81]	0.54 [0.43, 0.66]
Schoonhoven 2002_Braden	59	353	76	741	17.0	0.44 [0.35, 0.53]	0.68 [0.65, 0.70]
Vandenbosch 1996_Braden	17	30	12	44	17.0	0.59 [0.39, 0.76]	0.59 [0.47, 0.71]



Figure 2: ROC curve for scales Norton, Waterlow, Braden and clinical judgement – all stages, general population – for studies reporting more than one scale



O.8 Skin assessment

O.8.1 Forest plots and sensitivity

Figure 3: Erythema / redness - grades 2-4 – from multivariable analyses (unless otherwise stated)

Study or Subgroup	log[Odds Ratio]	SE	Odds Ratio IV, Fixed, 95% CI
1.1.1 Unadjusted - blanchable vs no erythma			
Konishi 2008	2.2468	1.1643	9.46 [0.97, 92.65]
Nixon 2007	-1.117	1.2321	0.33 [0.03, 3.66]
1.1.2 Multivariable analysis - redness vs no redness			
Compton 2008	0.8351	0.2538	2.31 [1.40, 3.79]
1.1.3 Multivariable analysis - non-blanchable vs blanchable+none			
Nixon 2007	1.9488	0.7326	7.02 [1.67, 29.51]
1.1.4 Unadjusted - non-blanchable vs not non-blanchable			
Vanderwee 2007	1.7362	0.2883	5.68 [3.23, 9.99]



Figure 4: Subjective nursing assessment of skin assessment features - grades 2-4 – from multivariable analyses

Study or Subgroup	log[Odds Ratio]	SE	Odds Ratio IV, Fixed, 95% CI
2.1.1 Moist skin vs non-moist skin			
Compton 2008	0.8544	0.2597	2.35 [1.41, 3.91]
2.1.2 Oedematous skin vs not oedematous skin			
Compton 2008	0.8087	0.2617	2.24 [1.34, 3.75]
2.1.3 Mottled skin vs not mottled skin			
Compton 2008	0.7036	0.2921	2.02 [1.14, 3.58]
2.1.4 Livid skin vs not livid skin (not significant, no data)			
Compton 2008	0	0	Not estimable
2.1.5 Centralised circulation vs not centralised circulation			
Compton 2008	0.8738	0.2656	2.40 [1.42, 4.03]
2.1.6 Cyanosis vs not cyanosis (not significant, no data)			
Compton 2008	0	0	Not estimable
2.1.7 Hyperaemic skin vs not hyperaemic skin (not significant, no data)			
Compton 2008	0	0	Not estimable
2.1.8 Redness vs no redness			
Compton 2008	0.8351	0.2538	2.31 [1.40, 3.79]



Figure 5: Subjective nursing assessment of moist skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	92	202	29	375	0.76 [0.67, 0.83]	0.65 [0.61, 0.69]



Figure 6: Subjective nursing assessment of oedematous skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	77	135	44	442	0.64 [0.54, 0.72]	0.77 [0.73, 0.80]



Figure 7: Subjective nursing assessment of mottled skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	40	48	81	529	0.33 [0.25, 0.42]	0.92 [0.89, 0.94]



Figure 8: Subjective nursing assessment of centralised circulation – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	86	171	35	406	0.71 [0.62, 0.79]	0.70 [0.66, 0.74]



Figure 9: Subjective nursing assessment of livid skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	38	48	83	529	0.31 [0.23, 0.40]	0.92 [0.89, 0.94]



Figure 10: Subjective nursing assessment of cyanosis – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	77	135	44	442	0.64 [0.54, 0.72]	0.77 [0.73, 0.80]



Figure 11: Subjective nursing assessment of reddened skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	83	172	38	405	0.69 [0.60, 0.77]	0.70 [0.66, 0.74]



Figure 12: Subjective nursing assessment of hyperaemic skin – ICU- grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Compton 2008	26	50	95	527	0.21 [0.15, 0.30]	0.91 [0.89, 0.93]



Figure 13: Presence of blanchable erythema assessed by finger test – hospitalized patients- all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Konishi 2008	6	56	2	185	0.75 [0.35, 0.97]	0.77 [0.71, 0.82]



Figure 14: Presence of thermal anomaly (an area of the skin at least 1°C warmer than the surrounding skin) – follow-up 10 days- geriatric inpatients- all grades

Study	TP	FP	FN	TN	Sensitivity	Specificity
Newman 1981	6	22	0	63	1.00 [0.54, 1.00]	0.74 [0.63, 0.83]



Figure 15: Presence of blanchable erythema assessed by finger test – hospitalized patients – grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Konishi 2008	3	59	1	186	0.75 [0.19, 0.99]	0.76 [0.70, 0.81]
Nixon 2006	3	55	1	6	0.75 [0.19, 0.99]	0.10 [0.04, 0.20]



Figure 16: Presence of non-blanchable erythema assessed by finger test – surgical inpatients – grades 2-4

Study	TP	FP	FN	TN	Sensitivity	Specificity
Nixon 2007	11	21	4	61	0.73 [0.45, 0.92]	0.74 [0.64, 0.83]
Vanderwee 2007	26	102	30	668	0.46 [0.33, 0.60]	0.87 [0.84, 0.89]



Figure 17: Presence of non-blanchable erythema assessed by transparent disk and Braden score then non-blanchable erythema (in comparison with non-blanchable erythema alone)

NBE

Study	TP	FP	FN	TN	Sensitivity	Specificity
Vanderwee 2007	26	102	30	668	0.46 [0.33, 0.60]	0.87 [0.84, 0.89]



Braden then NBE

Study	TP	FP	FN	TN	Sensitivity	Specificity
Vanderwee 2007	43	208	10	530	0.81 [0.68, 0.91]	0.72 [0.68, 0.75]



Figure 18: Unadjusted odds ratios for presence of non-blanchable erythema assessed by transparent disk and Braden score then non-blanchable erythema

Study or Subgroup	Positive on test Events	Total	Negative on test Events	Total	Odds Ratio M-H, Fixed, 95% CI
4.1.1 NBE alone					
Vanderwee 2007	26	128	30	698	5.68 [3.23, 9.99]
4.1.2 NBE plus Braden					
Vanderwee 2007	43	251	10	540	10.96 [5.41, 22.21]



O.9 References

1 Andersen KE, Jensen O, Kvorning SA, Bach E. Prevention of pressure sores by identifying patients at risk. BMJ. 1982; 284(6326):1370-1371

2 Anthony D, Reynolds T, Russell L, Anthony D, Reynolds T, Russell L. A regression analysis of the Waterlow score in pressure ulcer risk assessment. Clinical Rehabilitation. England 2003; 17(2):216-223

3 Barnes D, Payton RG. Clinical application of the Braden Scale in the acute-care setting. Dermatology Nursing. 1993; 5(5):386-388

4 Bergstrom N, Braden B, Kemp M, Champagne M, Ruby E. Predicting pressure ulcer risk: a multisite study of the predictive validity of the Braden Scale. Nursing Research. 1998; 47(5):261-269

5 Bergstrom N, Braden BJ, Laguzza A, Holman V. The Braden Scale for Predicting Pressure Sore Risk. Nursing Research. 1987; 36(4):205-210

6 Bergstrom N, Demuth PJ, Braden BJ. A clinical trial of the Braden Scale for Predicting Pressure Sore Risk. Nursing Clinics of North America. 1987; 22(2):417-429

7 Braden BJ, Bergstrom N. Predictive validity of the Braden Scale for pressure sore risk in a nursing home population. Research in Nursing and Health. 1994; 17(6):459-470

8 Capobianco ML, McDonald DD. Factors affecting the predictive validity of the Braden Scale. Advances in Wound Care. 1996; 9(6):32-36

9 Chan WS, Pang SM, Kwong EW. Assessing predictive validity of the modified Braden scale for prediction of pressure ulcer risk of orthopaedic patients in an acute care setting. Journal of Clinical Nursing. England 2009; 18(11):1565-1573

10 Compton F, Hoffmann F, Hortig T, Strauss M, Frey J, Zidek W et al. Pressure ulcer predictors in ICU patients: nursing skin assessment versus objective parameters. Journal of Wound Care. 2008; 17(10):417-424

11 Cubbin B, Jackson C. Trial of a pressure area risk calculator for intensive therapy patients. Intensive Care Nursing. 1991; 7(1):40-44

12 Curley MAQ, Razmus IS, Roberts KE, Wypij D. Predicting pressure ulcer risk in pediatric patients: the Braden Q Scale. Nursing Research. 2003; 52(1):22-33

13 de Souza DM, Santos VL, Iri HK, Sadasue Oguri MY. Predictive validity of the Braden Scale for Pressure Ulcer Risk in elderly residents of long-term care facilities. Geriatric Nursing. United States 2010; 31(2):95-104

14 Edwards M. The levels of reliability and validity of the Waterlow pressure sore risk calculator. Journal of Wound Care. 1995; 4(8):373-378

15 Feuchtinger J, Halfens R, Dassen T. Pressure ulcer risk assessment immediately after cardiac surgery--does it make a difference? A comparison of three pressure ulcer risk assessment instruments within a cardiac surgery population. Nursing in Critical Care. England 2007; 12(1):42-49

16 Goodridge DM, Sloan JA, LeDoyen YM, McKenzie JA, Knight WE, Gayari M. Risk-assessment scores, prevention strategies, and the incidence of pressure ulcers among the elderly in four Canadian health-care facilities. Canadian Journal of Nursing. 1998; 30(2):23-34

17 Halfens RJG, Van Achterberg T, Bal RM. Validity and reliability of the Braden scale and the influence of other risk factors: a multi-centre prospective study. International Journal of Nursing Studies. 2000; 37(4):313-319

18 Hatanaka N, Yamamoto Y, Ichihara K, Mastuo S, Nakamura Y, Watanabe M et al. A new predictive indicator for development of pressure ulcers in bedridden patients based on common laboratory tests results. Journal of Clinical Pathology. England 2008; 61(4):514-518

19 Jackson C. The revised Jackson/Cubbin Pressure Area Risk Calculator. Intensive & Critical Care Nursing. 1999; 15(3):169-175

20 Jalali R, Rezaie M. Predicting pressure ulcer risk: comparing the predictive validity of 4 scales. Advances in Skin and Wound Care. United States 2005; 18(2):92-97

21 Kim E, Lee S, Lee E, Eom M. Comparison of the predictive validity among pressure ulcer risk assessment scales for surgical ICU patients. Australian Journal of Advanced Nursing. 2009; 26(4):87-94

22 Kwong E, Pang S, Wong T, Ho J, Shao-ling X, Li-jun T. Predicting pressure ulcer risk with the modified Braden, Braden, and Norton scales in acute care hospitals in Mainland China. Applied Nursing Research. 2005; 18(2):122-128

23 Langemo DK, Olson B, Hunter S, Hanson D, Burd C, Cathcart-Silberberg T. Incidence and prediction of pressure ulcers in five patient care settings. Decubitus. 1991; 4(3):25-36

24 Lewicki LJ, Mion LC, Secic M. Sensitivity and specificity of the Braden Scale in the cardiac surgical population. Journal of Wound, Ostomy and Continence Nursing. 2000; 27(1):36-41

25 Lincoln R, Roberts R, Maddox A, Levine S, Patterson C. Use of the norton pressure sore risk assessment scoring system with elderly patients in acute care. Journal of Enterostomal Therapy. 1986; 13:132-138

26 Lindgren M, Unosson M, Krantz AM, Ek AC. A risk assessment scale for the prediction of pressure sore development: reliability and validity. Journal of Advanced Nursing. 2002; 38(2):190-199

27 Lothian P. Identifying and protecting patients who may get pressure sores. Nursing Standard. 1989; 18(4):26-29

28 Lyder CH, Yu C, Emerling J, Mangat R, Stevenson D, Empleo-Frazier O et al. The Braden Scale for pressure ulcer risk: evaluating the predictive validity in Black and Latino/Hispanic elders. Applied Nursing Research. 1999; 12(2):60-68

29 Norton D, McLaren R, Exton-Smith AN. An Investigation of Geriatric Nursing Problems in Hospital. Edinburgh: Churchill-Livingstone; 1962

30 Ongoma C, Schmollgruber S, Langley GC. Predictive validity of pressure risk assessment scales in a private sector trauma intensive care unit. Southern African Journal of Critical Care. 2005; 21(2):78-86

31 Page KN, Barker AL, Kamar J. Development and validation of a pressure ulcer risk assessment tool for acute hospital patients. Wound Repair and Regeneration. United States 2011; 19(1):31-37

32 Pang SM, Wong TK. Predicting pressure sore risk with the Norton, Braden, and Waterlow scales in a Hong Kong rehabilitation hospital. Nursing Research. 1998; 47(3):147-153

33 Perneger TV, Rae AC, Gaspoz JM, Borst F, Vitek O, Heliot C. Screening for pressure ulcer risk in an acute care hospital: development of a brief bedside scale. Journal of Clinical Epidemiology. 2002; 55(5):498-504

34 Quigley SM, Curley MAQ. Skin integrity in the pediatric population: preventing and managing pressure ulcers. Journal of the Society of Pediatric Nurses. 1996; 1(1):7-18

35 Ramundo JM. Reliability and validity of the Braden Scale in the home care setting. Journal of Wound, Ostomy and Continence Nursing. 1995; 22(3):128-134

36 Salvadalena GD, Snyder ML, Brogdon KE. Clinical trial of the Braden Scale on an acute care medical unit. Journal of Enterostomal Therapy Nursing. 1992; 19(5):160-165

37 Schoonhoven L, Haalboom JRE, Bousema MT, Algra A, Grobbee DE, Grypdonck MH et al. Prospective cohort study of routine use of risk assessment scales for prediction of pressure ulcers. BMJ. 2002; 325(7368):797

38 Seongsook RNJ, Ihnsook RNJ, Younghee RNL. Validity of pressure ulcer risk assessment scales; Cubbin and Jackson, Braden, and Douglas scale. International Journal of Nursing Studies. 2004; 41(2):199-204

39 Serpa LF, de Gouveia Santos VL, Gomboski G, Rosado SM. Predictive validity of Waterlow Scale for pressure ulcer development risk in hospitalized patients. Journal of Wound, Ostomy and Continence Nursing. United States 2009; 36(6):640-646

40 Serpa LF, Santos VL, Campanili TC, Queiroz M. Predictive validity of the Braden scale for pressure ulcer risk in critical care patients. Revista Latino-Americana De Enfermagem. Brazil 2011; 19(1):50-57

41 Smith I. Waterlow/Norton scoring system: a ward view. Care: Science and Practice. 1989; 7(4):93-95

42 Stotts NA. Predicting pressure ulcer development in surgical patients. Heart and Lung. 1988; 17(6 Pt 1):641-647

43 Suriadi, Sanada H, Sugama J, Thigpen B, Subuh M. Development of a new risk assessment scale for predicting pressure ulcers in an intensive care unit. Nursing in Critical Care. England 2008; 13(1):34-43

44 Suriadi, Sanada H, Sugama J, Thigpen B, Kitagawa A, Kinosita S et al. A new instrument for predicting pressure ulcer risk in an intensive care unit. Journal of Tissue Viability. 2006; 16(3):21-26

45 Towey AP, Erland SM. Validity and reliability of an assessment tool for pressure ulcer risk. Decubitus. 1988; 1(2):40-48

46 VandenBosch T, Montoye C, Satwicz M, Durkee-Leonard K, Boylan-Lewis B. Predictive validity of the Braden Scale and nurse perception in identifying pressure ulcer risk. Applied Nursing Research. 1996; 9(2):80-86

47 Wai-Han C, Kit-Wai C, French P, Yim-Sheung L, Lai-Kwan T. Which pressure sore risk calculator? A study of the effectiveness of the Norton scale in Hong Kong. International Journal of Nursing Studies. 1997; 34(2):165-169

48 Waterlow J. Pressure sore: a risk assessment card. Nursing Times. 1985; 81:49-55

49 Waterlow J. The Waterlow Pressure Ulcer prevention manual (revised 2005). 2005. Available from: www.judy-waterlow.co.uk [Last accessed: 27 September 2012]

50 Weststrate JT, Hop WC, Aalbers AG, Vreeling AW, Bruining HA. The clinical relevance of the Waterlow pressure sore risk scale in the ICU. Intensive Care Medicine. 1998; 24(8):815-820