

# Evidence tables

---

## 4. Physical features

### 4.1 Injuries

#### 4.1.1 Bruises

Bibliographic details	Study type and evidence level	Study details	Patient characteristics	Intervention and comparisons	Comments
Maguire S;Mann MK; 2005  10	Study Type: Systematic Review/Meta-Analysis  Evidence Level: 2+	23 studies	All papers that defined patterns of bruising in non-abused or abused children aged less than 18 year of age.	Papers about bruising in non-abused children: 2 case-control studies 4 cross sectional studies 3 case series  Papers about bruising in abused children: 2 case-control studies 1 cross sectional study 13 case-series	Source of Funding: Supported by the NSPCC (funding?)  11 study populations were located in the USA, 7 in the UK, 2 in Australia, 1 in Canada, 1 in South Africa and 1 Germany.

#### 4.1.2 Bites

Bibliographic details	Study type	Study details	Patient characteristics	Intervention and comparisons	Comments
11;12	Systematic review  Evidence level: 2+	Systematic review of inflicted bites in children.	5 case studies: 4 children <30 months, 1 in her teens.	n/a	

#### 4.1.3 Cuts and abrasions

When to suspect child maltreatment: full guideline DRAFT (December 2008)

No literature identified.

#### 4.1.4 Thermal injuries

Bibliographic details	Study type	Study details	Patient characteristics	Comparisons	Comments
Maguire S;Moynihan S;Mann M;Potokar T;Kemp AM;  2007 Jan 12  166  Country: USA and UK	Study Type: Systematic Review/Meta-Analysis  Evidence Level: 2+	26 studies 587 children	One case-control study, eight cross sectional studies and 17 case series and case studies	Whether a scald was intentional or accidental. (neglectful scalds excluded) <b>Intentional scalds:</b> immersion injuries, caused by hot tap water, affecting the extremities, buttocks or perineum or both symmetrical with clear upper margins, associated with old fractures and unrelated injuries. <b>Unintentional scalds:</b> due to spill injuries of other hot liquids, affecting the upper body with irregular margins and depth.	Source of Funding: NSPCC Narrative review
CORE-INFO;  2006  14	Study Type: Systematic Review/Meta-Analysis  Evidence Level: 2+	28 studies (1 case-control, 27 case series) 255 children of which 76 were abused	Children < 14 years of age	Contact burns commonest non-scald injury. Injuries with demarcated edges in shape of implement (e.g. cigarette, iron) Age not a factor in intentional non-scald burns.	Limitations of the review: Small numbers of children No comparative studies of cigarette burns Lack of comparative data for contact burns

#### 4.1.5 Cold injury

No literature identified.

#### 4.1.6 Hair loss

No literature identified.

#### 4.1.7 Fractures

When to suspect child maltreatment: full guideline DRAFT (December 2008)

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Hui C;Joughin E;Goldstein S;Cooper N;Harder J;Kiefer G;Parsons D;Howard J;  2008 Apr  21	Study Type: Other  Evidence Level: 4	Comparison: Non-accidentally injured children vs accidentally injured children	127	Children under 3 years with femoral fracture	Abuse categorised as:  definite - multiple recent fractures, fractures of various ages, eyewitness, multiple internal injuries, physical findings, abuse of sibling, definite act causing physical harm to child, suspicious injury with definite later abuse  likely abuse - previous injury diagnosed as abuse AND inconsistent history  questionable abuse - inconsistent history	14 categorised as non-accidental injury  mechanism of injury was unwitnessed or with an inconsistent or absent explanation in 10/14 compared to 3/113.  No specific fracture type or location.  Multiple injuries in 6/14 abused children compared to 13/113 in accidental injuries group.		Possible neglect cases are included in 'accidental' injury cases.

Kemp et al, 2008<sup>15</sup>, Day et al, 2006<sup>17</sup>, Carty, 2002<sup>18, 19, 20</sup> to be added.

4.1.8 Intra-cranial injuries

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Reviewer Comments
CORE-INFO <sup>24</sup>	Study type: Systematic review  Evidence level: 2+	12 studies USA, 2 UK studies, 2 France, 1 New Zealand	971 abused 1252 non-abused (215 motor vehicle accident, 905 other trauma, 132 other medical causes)				

When to suspect child maltreatment: full guideline DRAFT (December 2008)

#### 4.1.9 Eye trauma

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Reviewer Comments
Vinchon M;foort-Dhellemmes S;Desurmont M;Dhellemmes P;  2005 May  27	Study Type: Cohort  Evidence level: 2+	150 head-injured children, of which 129 were assessed for retinal haemorrhage.	69% male  mean age 5.3 months, median age 3.6 months  children <24 months hospitalised for craniocerebral traumatic lesions.			42/56 abused children had RH 5/73 accidental trauma children had RH  sens=75%, spec=93%.	High level of confirmation of abuse.

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Adams G;Ainsworth J;Butler L;Bonshek R;Clarke M;Doran R;Dutton G;Green M;Hodgkinson P;Leitch J;Lloyd C;Luthert P;Parsons A;Punt J;Taylor D;Tehrani N;Willshaw H;  2004  26	Study Type: Other Consensus statement  Evidence Level: 4	Comparison:	NA	NA	NA		In children under 2 years, retinal haemorrhage is highly unlikely to be caused by rough play or an attempt to arouse an apparently unconscious child	

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Taylor et al 1999 25	Consensus statement EL=4							

#### 4.1.10 Spinal injuries

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Reviewer Comments
24	Study Type: Systematic review - meta-analysis  Evidence level: 2+	15 studies, 33 abused children	median age 6 months (range 1.2 to 48 months). >50% younger than 6 months.	n/a	26 children died due to injuries, 2 survivors had quadriplegia.  Diagnosis delayed in 7 cases.  25/33 had cervical injuries  17/25 had significant head trauma  23/25 had retinal haemorrhage  Other presenting features: focal neurological signs, apnoea, signs of raised intra-cranial pressure, general neurological deterioration.  17/33 had thoracolumbar injuries	Abusive spinal injury is rare. Major accidental trauma must be excluded.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Reviewer Comments
					(median age 14m, range 9-16m) → 3 thoracic, one lumbar and 3 thoraco-lumbar injuries.  Presenting features: focal neurological signs and orthopaedic deformity,	

4.1.11 Visceral injuries

Bibliographic Details	Study Type & Evidence Level	No. of Patients	Patient Characteristics	Comparison	Outcome Measures, Follow-Up & Effect Size	Comments
Wood J; 2005 Nov <sup>30</sup> Country: USA	Study Type: Cohort Study  Evidence Level: 2-	Total number of patients = 121	Children less than 6y, Abbreviated Injury Scale score >= 2, sustained abdominal injuries, Exclusions: associated neurological injury, abdominal injury secondary to severe thoracic injury, abused children because of trauma denial, also where injuries could be not be classified as accidental or inflicted.			Source of Funding: Not stated  Non-accidental trauma was defined as suspicion being established by clinician and formal evaluation completed by child protection service. No significant differences between number of abdominal injuries due to non-accidental trauma and accidental trauma. Significantly more thoracic injuries due to accidental trauma.

DRAFT FOR CONSULTATION

DiScala C; 2000 Jan <sup>28</sup> Country: USA	Study Type: Cohort Study  Evidence Level: 2-	Total number of patients = 18828 Child abuse N = 1997 Unintentional injury N = 16831	Children aged <5 y hospitalised between 1988 and 1997. Data from National Pediatric Trauma Registry.			Source of Funding: Not stated  Patient databases often produce biased results. Unclear whether public and private hospitals covered. Children were identified as abused in the treating hospital. It is not clear how these decisions were reached. Cases of 'suspected abuse' were excluded.
Roaten JB; 2006 Dec <sup>29</sup> Country: USA	Study Type: Cohort Study  Evidence Level: 2+	Total number of patients = 6186 Accidental trauma patients N = 5733 Non-accidental trauma patients as identified by admitting clinician and evaluation by child advocacy and protection service. N = 453	Children aged under 18y attending trauma centre.			Source of Funding: Not stated
Trokel, M 2004 <sup>31</sup> Country: USA	Study type: Cohort  Evidence Level: 2-	927 cases of blunt abdominal trauma	Children aged <5 y between 1995 and 2001. Data from National Pediatric Trauma Registry. 46% female Median age 34 months  MVA 63% Abuse 16% Fall 14% Other 8%		Excluding MVAs, abuse accounted for 79% of injuries in children <12 mo 61% in 13mo - 24mo 39% in 25mo – 36mo 25% in 37mo – 48mo	No description of differences in presentation between unintentional and abusive injuries.

4.1.12 Oral injury

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
------------------------------	-----------------------------------	-----------------------	----------------------------	------------	------------------------------------	-------------	---------------	----------------------

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Maguire S;Hunter B;Hunter L;Sibert JR;Mann M;Kemp AM;  2007  32	Study Type: Systematic review - meta-analysis  Evidence level: 2+	19 studies (case studies, case series and one case-control study) 603 children  27 abused children with torn labial frenum	children aged 0-18 with intra-oral injuries due to physical abuse and torn labial frenum with any cause	Are intra-oral injuries indicative of maltreatment?  Comparison: n/a	Follow-up period: n/a  Outcome Measures:	27 abused children had torn labial frenum (7 case series, 1 case-control study, 1 case study) 22 were <5y  Non-abusive torn frenum found in two children. Torn frenum not regularly documented as it is considered a trivial injury.  Other abusive intra-oral injuries were found in 580 children, namely:  lacerations or bruising to the lips, mucosal lacerations, dental trauma, tongue injuries, gingival lesions		Oral injuries not specific to abuse.

4.2.1 Genital and anal symptoms

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
DeLago C;Deblinger E;Schroeder C;Finkel MA;  2008 Aug 1  33	Study Type: OtherRetrospective chart review of girls who disclosed sexual abuse.  Evidence Level: 3	n/a  Comparison: n/a	161 girls who had disclosed sexual abuse by direct genital contact.	Median age 10.5 y (range 3.1-17.8y)		% of girls reporting genital symptoms: genital pain/soreness 53 dysuria 37 genital bleeding 11		Study took place at child abuse referral centre. Only charts reviewed were those written by a doctor with a standard method for eliciting information.



Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Klevan JL;De Jong AR;  1990 Feb  34	Study Type: Other  Evidence Level: 3	NA  Comparison:	428 CSA victims seen at sexual assault health centre based in a hospital.	mean age 8.6 years (range 1-16), 84% female		85 (20%) of sample had GU symptoms at 1-3 weeks after first report of CSA.  Recent onset of enuresis in 24 (6% of total series) Vaginal pain 43 (10%) Dysuria 21 (5%) Increased urinary frequency 20 (5%)		Cohort of sexually abused children. No data on non-abused children.

#### 4.2.2 Genital and anal signs

<sup>3</sup> to be added

#### 4.2.3 Sexually transmitted infection

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Royal College of Paediatrics and Child Health;  2008  3	Study Type: Systematic review - meta-analysis  Evidence level: 1+	Neisseria gonorrhoeae n=17 studies Chlamydia trachomatis n=10 studies Bacterial vaginosis n=6 studies Genital Mycoplasmas n=6 studies Syphilis n=9 studies Anogenital warts n=10 studies Oral warts n=1 study Genital herpes	Newborn and children aged 0 to 18 years with an STI for which child sexual abuse had been confirmed or actively excluded.	Observational studies  Comparison:	Follow-up period:  Outcome Measures: Prevalence of sexual abuse in children with the STI  Prevalence of the STI in sexually abused children	Neisseria gonorrhoea: Gonorrhoea is not often seen in sexually abused prepubertal and pubertal children. A significant number of children with gonorrhoea who have been evaluated for sexual abuse were found to have been abused. This suggests that sexual contact was the mode of transmission. Sexual abuse is the most likely mode of transmission in pubertal and prepubertal children. The evidence does not	Child sexual abuse should be strongly considered in children with Neisseria gonorrhoeae, Chlamydia trachomatis and anogenital warts. A high prevalence of abuse was also found in studies on Trichomonas vaginalis, genital herpes and HIV, although population numbers were small.  For syphilis, Hepatitis B and C was too limited to offer information on the association between the presence of the infection and sexual abuse.	The review suffers from a lack in evidence but is methodologically rigorous. The limitations of the study relate to the very thin evidence base. This results of this review are mainly expert consensus based.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
		simplex n=5 studies Hepatitis B n=4 studies Hepatitis C n=2 studies HIV n=4 studies Trichomonas vaginalis n=10 studies				<p>helped to establish the age at which the possibility of verital transmission can be excluded.</p> <p>Chlamydia trachomatis: Chlamyida infection is not often seen in sexually abused children. A significant number of children with chlamydia trachomatis who have been evaluated for sexual abuse were found to have been abused. This suggests that sexual contact was the mode of transmission. Chlamydia is more frequent in pubertal than prepubertal sexually abused girls. This result may have been confounded by consensual sexual activity and/or younger children less likely to disclose abuse. The evidence does not helped to establish the age at which the possibility of verital transmission can be excluded.</p> <p>Bacterial vaginosis: There is insufficient data in children to determine the significance of bacterial vaginosis in relation to CSA.</p> <p>Genital Mycoplasmas: The evidence does not</p>	<p>Attributing infection to perinatal transmission or sexual abuse is difficult in very young children who are preverbal and cannot disclose abuse.</p> <p>For Neisseria gonorrhoea, Chlamydia trachomatis, anogenital warts and Trichomonas vaginalis, the likelihood of sexually transmitted infection in sexually abused children increased with the child's age. This is complicated by a lack of consideration of consensual sexual activity in adolescents, difficulties in obtaining disclosure of abuse in young children and incomplete information about how other modes of transmission were excluded.</p> <p>Penetrative sexual contact is associated with an increased risk of infection by Neisseria gonorrhoea, Chlamydia trachomatis, Trichomonas vaginalis and HIV.</p>	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						<p>help establish whether or not genital mycoplasmas are sexually transmitted children.</p> <p>Syphilis: The literature cannot help in establishing whether sexual contact is a likely route of transmission in children with syphilis.</p> <p>Anogenital warts: A significant proportion of children with anogenital warts have been sexually abused. Sexual abuse is more likely to be confirmed in older prepubertal children. The evidence does not help to establish the age at which the possibility of verital transmission can be excluded.</p> <p>Oral warts: There is insufficient evidence to determine the significance of oral warts in relation to child sexual abuse at the current time.</p> <p>Genital herpes simplex: There are very few published studies to inform whether sexual abuse is likely to be the mode of transmission. Where infected children had been evaluated 1/2 and 6/8 were found to</p>		

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						<p>have been abused.</p> <p>Hepatitis B: There is insufficient evidence to determine the significance of Hepatitis B in relation to sexual abuse in children. Despite the lack of evidence, in view of the fact that Hepatitis B can be sexually transmitted in adults, sexual abuse should be considered in a child with Hepatitis B if vertical, perinatal or blood contamination has been excluded. A positive diagnosis in the mother does not exclude child sexual abuse.</p> <p>Hepatitis C: There is insufficient evidence to determine the significance of Hepatitis C in relation to sexual abuse in children. Despite the lack of evidence, in view of the fact that Hepatitis C can be sexually transmitted in adults, sexual abuse should be considered in a child with Hepatitis C if vertical, perinatal or blood contamination has been excluded. A positive diagnosis in the mother does not exclude child sexual abuse.</p> <p>HIV:</p>		

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						<p>Published studies suggests that sexual abuse is a likely source of infection in children with HIV in whom the possibility of mother-child transmission or blood contamination has been excluded.</p> <p>In a child with HIV with an uninfected mother, the possibility of sexual abuse is highly likely. HIV infection in the mother of a child with HIV does not exclude the possibility of sexual transmission.</p> <p>Trichomonas vaginalis: Published studies suggests that sexual abuse is a likely source of infection in girls. The evidence does not help to establish the age at which the possibility of vertical transmission can be excluded. Consensual sexual activity should be considered.</p>		

4.2.4 Pregnancy

No literature identified.

5.1 Neglect

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
<p>Strathearn L;Gray PH;O'Callaghan F;Wood DO;</p> <p>2001 Jul</p> <p>35</p>	<p>Study Type: Other</p> <p>Evidence Level: 2+</p>	<p>NA</p> <p>Comparison: Children referred for maltreatment vs all others</p> <p>Substantiated maltreatment cases vs all others</p>	<p>353 children - 52 referred for maltreatment - 21 substantiated for neglect</p>	<p>Extremely low birthweight children followed from birth to 4 years.</p>	<p>Child General Cognitive Index (normative mean=100 sd 15)</p>	<p>GCI at 4 years (n=269) Referred children vs non referred children 83 vs 98 (p&lt;0.001)</p> <p>Substantiated referrals vs all others 82 vs 98 (p&lt;0.001)</p> <p>Physical abuse referrals vs all others 83 vs 97 (p=0.004) not significant when using only substantiated cases</p> <p>Emotional abuse referrals vs all others 81 vs 98 (p&lt;0.001) - similar for substantiated referrals vs all others</p> <p>Neglect referrals vs all others 77 vs 98 (p&lt;0.001) - similar for substantiated referrals vs all others</p> <p>Difference in GCI score at 4 years accounting for risk factors: Physical abuse referrals: 5 (95%CI -6.9-16.8)</p> <p>Emotional abuse referrals: -3 (95%CI -16.8-10.9)</p> <p>Neglect referrals: -17.6 (95%CI -31.9 - -3.3)</p> <p>Substantiated neglect referrals had increasingly worse outcomes over time than all other study participants.</p>		<p>Some participants lost to follow up. After four years, only 269 children in study, of these 21 substantiated referrals of maltreatment.</p> <p>Wide confidence intervals (because of small numbers in groups of interest)</p>

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Chester DL;Jose RM;Aldiyami E;  2006 Mar  39	Study Type: Other  Evidence Level: 3	Comparison: Burns in neglect cases (n=41) vs accidental burns (n=395)	440 patients including 41 with burns due to neglect and 4 with burns due to inflicted injury.	Admissions to burns unit <16 yrs.		<p>% of cases presenting after 24hours: neglect 49% vs accidental 14% (p&lt;0.0001)</p> <p>% of cases where first aid performed neglect 22% vs accidental 70% (p&lt;0.0001)</p> <p>% with presence of deep burns: neglect 71% vs accidental 54% (p=0.49)</p> <p>% with skin grafting performed: neglect 76% vs accidental 41% (p&lt;0.0001)</p> <p>Mean age neglect 4.2 vs accidental 4.0 (not significant)</p> <p>Gender - not significant</p> <p>Anatomical site - not significant</p> <p>Mechanism of injury - not significant.</p> <p>Mean body surface area affected neglect 7.1% vs accidental 6.4% (not significant)</p>		<p>Concern about circumstances of injury in 178 children. These children were investigated by family support team including home assessment. Definition of neglect based on these investigations.</p> <p>Study in burns unit, so more severe cases?</p>

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Whitaker RC;Phillips SM;Orzol SM;Burdette HL;  2007  37	Study Type: Other  Evidence Level: 2-	NA  Comparison: Ever neglected in last year vs never neglected (as measured by Parent-Child Conflicts Tactics Scales and items)  Corporal punishment (frequency per year)  Psychological aggression (frequency per year)	2412 children and their mothers	Age of children 3.2 years.		OR for obesity if neglected adjusted for maternal BMI and other covariates including birth weight.  1.56 (1.14 - 2.14):  OR for obesity if experienced corporal punishment adjusted for maternal BMI and other covariates including birth weight.  0.94 (0.72 - 1.24)  OR for obesity if experienced psychological aggression adjusted for maternal BMI and other covariates including birth weight.  0.90 (0.70 - 1.18)		11% of mothers reported ever doing one of the itmes on the negelct subscale  18% of children in the study were obese as measured by having BMI>=95th percentile on Centers for Desease Control and Prevention 2000 growth reference.



DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Stockwell MS;Brown J;Chen S;Vaughan RD;Irigoyen M;  2008  38	Study Type: OtherComparative case series  Evidence Level: 2-	Immunisation status at 3 months and 7 months of age.  Comparison: confirmed abuse cases vs ruled out abuse cases	285 children evaluated at a child advocacy centre with reliable immunisation data.	mean age 24.4m, SD 14m 63% female  confirmed abuse 17% suspected 11% ruled out 66% indeterminate 6%	Does underimmunisation predict maltreatment status?	Odds ratio of being a confirmed maltreatment case if:  underimmunised at 3 months 4.0 (95% CI 1.7-9.5)  underimmunised at 7 months 4.8 (95% CI 1.5- 15.7)  (comparison is underimmunised vs not)  Neglect not addressed separately.		Only immunisations that happened before first contact with child advocacy centre used in analysis.  Biased sample because non-abuse cases come from original 'suspected abuse', therefore not general population.

5.2 Over- and under-nutrition

Ref	Year	Country	Study Type	Population	Outcome	Conclusion	Evidence Level
-----	------	---------	------------	------------	---------	------------	----------------

DRAFT FOR CONSULTATION

Ref	Year	Country	Study Type	Population	Outcome	Conclusion	Evidence Level
<sup>41</sup>	1988	UK	cohort study	n = 260. Growth patterns of maltreated children (diagnosis based on case conference and social services intervention) based on remaining at home or entering foster care.	The study found that of the 260 children 39 had height greater than 2 Standard Deviations (SD) below mean for the cohort, and 21 had weight greater than 2 SD below mean for cohort. The study reported that 10 of 11 children in foster care compared to 4 of 28 children who remained at home showed 0.5 SD increase in height (p = 0.001). However, 8 of 16 who remained at home compared to 4 of 4 who were in foster care showed a 0.5 SD increase in weight (ns).	Study concluded maltreated children should not be rehabilitated at home.	EL = 3
<sup>42</sup>	1989	USA	case-control study	n = 196. Growth patterns of children who had been maltreated (n = 53 – 64.2% female, 86.5% non-Caucasian, 84% less than 5 years old) or not (n = 143 – 51% female, 59.3% non-Caucasian, 87% less than 5 years old).	The study reported low weight for height in 16.35% of abused and 0.7% of non-maltreated (OR 16.6, 95% CI 1.9 to 145.0, p < 0.05). The study found a low height for age in 11.6% of abused and 5.6% of non-maltreated (OR = 2.2, 95% CI 0.61 to 7.9). All the figures were adjusted for age, sex, and ethnicity.	The study concluded that malnutrition was found more amongst abused children than amongst non-abused.	EL = 2+
<sup>37</sup>	2007	USA	cohort study	n = 2412. Assessed the association between obesity (BMI > 30) and maltreatment (based on parent-child conflict tactics scale – neglect, physical punishment, psychological aggression) in children (aged 3 years, 48.2% female, 19.4% Caucasian).	The study found that 23.6% of neglected children were obese compared to 17.5% of children who were not neglected (OR = 1.56, 95% CI 1.14 to 2.14, adjusted for maternal BMI and other covariates). For physical punishment the study found that 19.8% of children whose parents reported 0 to 2 incidences per year were obese, 19.8% for those that reported 2 to 6, 18.4% for those that reported 7 to 14, 15% for those that reported 15 to 30, 17.8% for those that reported 31 to 104 (OR = 0.94, 95% CI 0.72 to 1.24). For psychological	The study concluded that neglect was associated with obesity.	EL = 3

DRAFT FOR CONSULTATION

Ref	Year	Country	Study Type	Population	Outcome	Conclusion	Evidence Level
					aggression the study found that 19.7% of children whose parents reported 0 to 5 incidences per year were obese, 18% for those that reported 6 to 16, 17.5% for those that reported 7 to 29, 17.4% for those that reported 30 to 49, 18% for those that reported 50 to 125 (OR = 0.90 to 1.18).		
<sup>43</sup>	2007	USA	case-control study	n = 173. Examined the link between childhood sexual abuse (based on child protection services, n = 84, 39% minority) or not (n = 89, 51% minority) and obesity (BMI > 30) from childhood to adulthood in females.	The study found that as children (aged 6 to 14) 25.42% of abused compared to 21.88% of non-abused were obese (OR = 1.25, 95% CI -0.05 to 3.00, p = 0.52). As adolescents (aged 15 to 19) the figures were 27.87% vs. 15.49% (OR 2.03, 95% CI 0.54 to 4.60, p = 0.09). As adults (aged 20 to 27) the figures were 42.25% vs. 28.4% (OR = 2.85, 95% CI 1.06 to 4.64, p = 0.009).	The study concluded that identification of high-risk growth trajectories may improve health outcomes for victims.	EL = 2+
<sup>44</sup>	2002	USA	community-based prospective cohort study	n = 782 mothers and off-spring. Examined link between childhood adversity (abuse based on referral to child protection services) and weight problems during adolescence and early adulthood. Children were interviewed three times over a ten year period. The study was 91% white and 385 of 782 were female.	The study found that 5 of 24 who reported neglect were obese compared to 36 of 711 who did not report neglect (OR = 4.66, 95%CI 1.65 to 13.16). The figures for recurrent weight change and physical abuse were 10 of 24 compared to 117 of 711 (OR = 3.63, 95% CI 1.58 to 8.36). For recurrent weight change and sexual abuse the figures were 9 of 22 compared to 120 of 644 (OR = 3.02, 95% CI 1.26 to 7.24). The figures for strict dieting and physical abuse were 9 of 24 compared to 120 of 711 (OR = 2.96, 95% CI 1.26 to 6.91). The study also undertook sub-group analysis on females. For females the study found that low body weight and physical abuse 4 of 24	The study reported that parental relationship factors were the most significant for eating disorders and weight problems.	EL = 2+

Ref	Year	Country	Study Type	Population	Outcome	Conclusion	Evidence Level
					compared to 13 of 319 (OR = 4.71, 95% CI 1.41 to 15.76). The figures for obesity and physical neglect were 3 of 14 compared to 14 of 356 (OR = 6.66, 95% CI 1.67 to 26.59).		

### 5.3 Oral health

Bibliographic Information	Study Type & Evidence Level	Study Aims/Objectives	No. of Patients	Patient Characteristics	Outcomes	Comments
Greene P; 1995 Jun <sup>46</sup>  Country: USA	Study Type: Case-Control Study  Evidence Level: 2+	To identify the role of child abuse/neglect on the oral health status in the primary dentition of children.	Total No. of Patients = 864, Cases = 42, Controls = 822	Age range 3yr - 11yr. Recruited from military bases. Controls matched on age, parental education, sponsor's military rank. No other descriptive statistics reported.	Presence of lifetime caries (treated or untreated) in child's primary teeth. No odds ratio reported for cases vs. controls.  Presence of untreated dental decay in primary teeth. Abused/neglected with noncombatant sponsor vs. noncombatant sponsor control OR 5.19 (95% CI 2.04, 13.2) Combatant but non-abused vs. noncombatant sponsor control OR 1.33 (95% CI 0.95, 1.87) Cases with combatant sponsor vs. non-combatant sponsor control OR 1.04 (95% CI 0.38, 2.85)	Funding: Not stated Maltreated cases stated as 'confirmed cases on the social services registry', controls recruited from general oral health survey from schools in the military bases.  The significant findings of this study relate to very specific circumstances that are not applicable to a general UK population. It should be noted that there were no significant differences between cases and controls in presence of lifetime caries.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Study Aims/Objectives	No. of Patients	Patient Characteristics	Outcomes	Comments
Greene PE; 1994 Jan <sup>47</sup>  Country: USA	Study Type: Case-Control Study  Evidence Level: 2+	To assess relationship between child abuse/neglect and oral health status.	Total No. of Patients = 903, Cases = 30, Controls = 873	Age 5yr - 13yr Cases and controls from military bases. Controls matched on age, parental education, sponsor's military rank.No other descriptive statistics reported.	Presence of lifetime caries (treated or untreated) in child's permanent teeth. Cases vs. controls: OR 2.20 (95% CI 0.90, 5.42)  Presence of untreated dental decay in primary teeth. Cases vs. controls: OR 8.00 (95% CI 3.90, 17.7)	Funding: Not stated  Significant relationship between presence of untreated decay and abuse/neglect, although confidence intervals are wide.  Maltreated cases stated as 'confirmed cases on the social services registry', controls recruited from general oral health survey from schools in the military bases. Military setting of study lends a bias to the results.

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
---------------------------	-----------------------------	--------------	--	----------------------------	------------------	--------------------	---------------	------------------

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Valencia-Rojas N;Lawrence HP;Goodman D;  2008  48	Study Type: OtherRetrospective chart review  Evidence Level: 3	a) prevalence of early childhood caries in abused/neglected children b)compare prevalence in maltreated with general c)Is untreated decay associated with different types of maltreatment  Comparison: dmft index in abused vs neglected children  comparisons between study children and general population data (3185 5- year old school children in city of Toronto)	66 children in care of Children's Aid Society of Toronto	80.3% neglected 19.7% physical or sexual abuse  mean age 4.1y (sd 1.16)	dmft index (decayed, missing, filled teeth)  Early Childhood Caries (ECC) = dmft index>=1 Severe Early Childhood Caries (S-ECC) =dmft >=4 Dental trauma Mean dmft and components (dt, mt, ft)	No children had missing or filled teeth at first dental visit so dmft=dt in these children.  58% of maltreated group had ECC vs 30% of general population  No significant differences between physical/sexual abuse group and neglect group  Mean dt 3.78 (se 0.73) in abused/neglected 4-6y olds 0.42 (se 0.02) in general population		Dental records not independent of knowledge of maltreatment status.

### 6.1 Repeat attendance at A&E

Bibliographic details	Study type and evidence level	Study details	Patient characteristics	comparisons	Comments
-----------------------	-------------------------------	---------------	-------------------------	-------------	----------

Bibliographic details	Study type and evidence level	Study details	Patient characteristics	comparisons	Comments
Spivey  2008  49	Cohort  Evidence level = 2+	Emergency department data linked to child protective services (CPS) data	All children aged <5y who attended emergency department with an injury in year 2000 Sexual abuse cases excluded. Repeat visits for same injury within 7 days excluded. Children who died excluded. n=50068 children accounting for 56364 injury visits	12% of children had a CPS report  2% of total had substantiated report  RR of being reported to CPS after 2 injury visits: 1.9 (95% CI 1.8-2.0)  RR of being reported to CPS after 4 injury visits: 3.8 95% CI 3.0 – 4.7)  RR of substantiated CPS report after 2 injury visits: 2.5 (95% CI 2.1-2.9)  RR of being reported to CPS after 4 injury visits: 4.7 (95% CI 2.4 – 9.2)  Unadjusted RRs used. Sensitivity analysis suggested age, sex, race or insurance status do not affect results.	Time of CPS report in relation to injury visit taken into account.

Woodman, 2008<sup>4</sup> to be added.

## 6.2 Dehydration

No literature identified.

## 6.3 Strangulation and suffocation

<sup>50</sup> to be added.

## 6.4 Apparent life threatening events

Ref	Author	Year	Study Design	Population	Outcome	Evidence Level	

DRAFT FOR CONSULTATION

Ref	Author	Year	Study Design	Population	Outcome	Evidence Level	
<sup>57</sup>	Altman	2003	Prospective case-series	243	6 had head injuries	3	A prospective case-series (n = 243) of infants under 12 months of age admitted to one tertiary unit in the USA examined the diagnosis given to cases of ALTE. The study found that a total 35 diagnosis were made: 80 were infection, 69 were gastrointestinal, 32 were neurologic (including 6 (2.5% of total) abusive head injuries within this group), 7 were airway obstruction, 6 were congenital or birth related problems, 39 were unknown, 6 were normal or benign. The study concluded that a wide spectrum of diseases and disorders can precipitate an ALTE. In relation to maltreatment the study concluded that "Among them, abusive head injury, a recently recognized cause, occurs frequently enough to obligate its inclusion in the differential diagnosis." (EL = 3) 2003 40750
<sup>59</sup>	Cote	1997	Retrospective case-series	73	abuse not reported	3	A retrospective case-series (n = 73) of infants (mean age 7.4 weeks) who were seen at a single apnoea program in USA. The results were that 47 had negative investigation, 17 recurrent events but no diagnosis, 5 respiratory infection, 2 had GORT, 1 had pallid syncope and 1 had tracheal stenosis. (EL = 3) 1998 40752
<sup>61</sup>	Ward	1986	Survey	11 of 31 apnea programmes and 4 of 10 vendors	13 sids, 4 non-accidental trauma, 6 sudden unexpected death at home, 1 subarachnoid hemorrhage, 1 cardiac disease	3	A survey of 11 apnoea monitoring programs and 4 apnoea monitoring device vendors in USA was undertaken examined reports of infant deaths. Over a 5-year period 1841 children were monitored. There were 25 reported deaths in this group: 13 due to SIDs, 4 due to non-accidental trauma (0.2% of total), 6 due to sudden unexpected death at home, 1 due to subarachnoid haemorrhage, and 1 caused by cardiac disease. The study no specific conclusions relating to maltreatment (EL = 3) 40791 1986
<sup>167</sup>	Johnson	1995	Retrospective case-series	28 infants with non-accidental head injury	57% had history of apnea	3	A retrospective case-series (n = 28) from the USA of children who suffered proven non-accidental head injury examined their presentation and outcome. Of the children examined only 3 older than 1 year of age. The results showed 16 of 28 presented with apnoea. Of those who presented with apnoea 57% had history of apnoea and 71% had previous seizures within 24 hours. The study found that 12 were left with severe disability, 4 died, 1 was in vegetative state and 7 survived. The authors concluded that trauma induced apnoeas is more important to outcome than mechanism of injury. (EL= 3) 38448 1995



DRAFT FOR CONSULTATION

Ref	Author	Year	Study Design	Population	Outcome	Evidence Level	
<sup>168</sup>	Light	1989	Survey of 127 apnea monitoring programs	20090	54 had mbpa	3	A survey of 51 of 127 (n = 20090) apnoea monitoring programs in USA investigated prevalence of fabricated and induced illness. The results showed that 54 (0.25% of total) cases of fabricated or induced illness were reported. The average age of infants with this diagnosis was 8.2 weeks. Detailed information on 32 of these 54 showed that 18 were re-hospitalised between 1 and 4 times, 13 were re-hospitalised 5 or more and 1 was unknown. The study concluded that fabricated or induced illness presents as unexplained multiple, serious apnoea events occurring in the presence of only one person (not witnessed). (EL = 3) 40767 1989.
<sup>52</sup>	McGovern	2004	Systematic review	Diagnoses when infants first present with an ALTE	2912 assessed, 8 studies -643 - 227 were GORD, 83 were seizures, 58 were LRTI, 2 were factitious	2+	One systematic review (n = 8 papers; search undertaken in 2002) assessed the initial diagnosis given when infants presented with an ALTE. The review included 8 studies involving 643 infants seen in Emergency Departments or paediatric units. The study calculated that 0.6% to 0.8% of emergency admissions for infants were for ALTE. A total of 728 diagnosis covering 50 conditions were reported, of these: 227 were gastro-oesophageal reflux disease (GORD), 169 were unknown, 83 were seizures, 58 were Lower Respiratory Tract Infection (LRTI), 26 were Ears, Nose and Throat (ENT) problems, 17 were breath holding, 11 were metabolic disease, 11 were ingestion of toxin or drugs, 6 were cardiac problems, 8 were Urinary Tract Infection (UTIs), 5 were benign cause, and 2 were fabricated illness (0.3% of children). The study concluded that careful investigation needed of ALTE due to variation in cause. (EL = 2+)(40770) 2004
<sup>55</sup>	Piteeti	2002	Prospective case-series	128	51 had gerd, 38 apnea, 11 choking episode, 6 infection, 5 bronchiolitis, 5 URI, 4 seizure, 3 abuse, 3 swallowing disorder, 2	3	A prospective case-series (n = 128) from the USA of children aged less than 24 months presenting at a single Emergency Department examined the diagnosis applied to cases of ALTE. Of the 128 cases of ALTE: 51 were GORD, 38 were apnoea, 11 were choking episode, 6 were infection, 5 were bronchiolitis, 5 were URI, 4 were seizures, 3 were abuse (2.3% of total), 3 were swallowing disorder, and 2 breathing spell. The study concluded that abuse diagnosed in 2.3% of cases of ALTE and this should be considered in patients who present with ALTE. (EL = 3) 39794 2002

DRAFT FOR CONSULTATION

Ref	Author	Year	Study Design	Population	Outcome	Evidence Level	
					breathing spell		
<sup>56</sup>	Samuels	1993	Prospective case-series	157	80 had no diagnosis; 77 had diagnosis - 2 disturbances in skin perfusion 7 fabricated, 18 suffocation, 40 hypoxaemic.	3	A prospective case-series (n = 157) from the UK of children (aged 1 week to 96 months) presenting once or more in one hospital setting examined the diagnosis applied to cases of ALTE. The study reported that of the 157 reported cases: 80 had no diagnosis; 77 had diagnosis. Of those diagnosed: 2 had disturbances in skin perfusion, 7 had fabricated illness (9% of those diagnosed and 4% of total), 18 suffocation (23% of those diagnosed and 11.5% of total), 40 had hypoxaemic, and 10 had hypoxaemia induced by epilepsy. The study concluded that identification of mechanisms is essential to the appropriate management of infants with apparent life threatening events. (EL = 3), 26532 1993
<sup>58</sup>	Stratton	2003	Retrospective case-series	60	abuse not reported	3	A retrospective case-series (n = 60) from the USA examined the diagnosis applied to infants with ALTE. The study setting was a single emergency medical service over a 12 month period. The study found that 60 (7.5%) out of 804 infants encountered by met criteria for ALTE (absence of breath, colour change, change in muscle tone). The diagnosis applied to these cases were: 20 (33%) had no diagnosis, 7 (12%) were pneumonia or bronchiolitis, 6 (10%) were GORD, 5 (8%) were seizures, 4 (7%) were sepsis, 4 (7%) were Upper Respiratory infection, 3 (5%) were apnoea episodes 2 (3%) were intracranial haemorrhage, 2 (3%) left against advice, 1 (2%) was bacterial meningitis, 1 (2%) was dehydration, and 1 (2%) was severe anaemia. Furthermore, of the 60 infants 35% had diagnosed underlying conditions. The study reported 1 case of intracranial injury caused by maltreatment, but highlighted the in 20 cases no diagnosis was made and in 2 the parents left against medical advice. The study concluded that "An apparent life-threatening event in an infant can present without signs of acute illness and is commonly encountered in the EMS setting. It is often associated with significant medical conditions, and EMS personnel should be aware of the clinical importance of an apparent life-threatening event. Infants

DRAFT FOR CONSULTATION

Ref	Author	Year	Study Design	Population	Outcome	Evidence Level	
							meeting criteria for an apparent life-threatening event should receive a timely and thorough medical evaluation" (EL = 3) 2003 40788
<sup>53</sup>	Kiechl-Kohlendorfer	2004	Prospective case-series	44184		3	A prospective cohort study (n = 44184) undertaken in Austria investigated the epidemiology of ALTE. The study identified 164 cases of ALTE or 2.46 per 1000 live births. An underlying cause was identified 91 of 164 cases (55%) and of these: 29% were respiratory, 22% were digestive tract, 2% were congenital cardiac malformation, 1% were inborn metabolic errors, and 1% were convulsions. The study made no conclusions in relation to child maltreatment. (EL = 3) 41232 2004
<sup>54</sup>	rahilly	1991	Prospective case-series	340		3	A prospective cohort study (n = 340) undertaken in Australia examined the diagnosis of ALTEs 289 of 340 had a diagnosis: 211 were GORT, 17 airway pathology, 25 fits/seizures, 2 brain-stem tumours, 2 hypoglycaemia, 8 respiratory syncytial virus, 5 fabricated or induced illness (1.7% of those diagnosed, 1.5% of total), 27 abnormal pneumograms (11 with reflux). 51 had no abnormal finding. The study made no conclusions in relation to child maltreatment. (EL = 3) 41230 1991

6.5 Poisoning

No literature identified.

6.6 Near drowning

Bibliographic Information	Study Type & Evidence Level	Study Aims/Objectives	No. of Patients	Patient Characteristics	Outcomes	Comments
Gillenwater JM; 1996 <sup>63</sup> Country: USA	Study Type: Other  Evidence Level: 3	To improve characterization and recognition of inflicted pediatric submersions.	Total No. of Patients = 205 were judged as being inflicted N = 16 were classified as unintentional submersions N = 186	Children younger than 19 years who sustained submersion injury and were hospitalised or autopsied. All children in the inflicted submersion group were below the age of 5years.		The study population is located in the USA.

6.7 Fabricated or induced illness

When to suspect child maltreatment: full guideline DRAFT (December 2008)

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Sheridan MS; 2003 Apr 64	Study Type: Systematic review - meta-analysis  Evidence level: 2++	451 cases of MSBP	52% male (n=415) Mean age at diagnosis: 48.6m (s.d. 49m), range 0-204m (n=404)	Comparison:	Follow-up period:  Outcome Measures: Length of time from onset to diagnosis Perpetrator Reported symptoms	Length of time from onset to diagnosis Estimated mean 21.8m, range (0-195m) (n=201)  Perpetrator 76.5% mothers 6.7% fathers  Reported symptoms (% of case reports) Apnoea (26.8%) Diarrhoea (24.6%) Seizures (17.5%) Behaviour [not defined] (10.4%) Asthma (9.5%) Allergy (9.3%) Fever (8.6%) Unspecified pain (8.0%) Unspecified infection (7.5%) Injury (6.4%) Unspecified bleeding (6.4%) Developmental delay (5.7%) Lethargy, fatigue (5.7%) Otitis (5.1%) Respiratory tract infection (5.1%) ...64 other symptoms occurring in fewer than 5% of the case reports.  Mean number of medical problems per victim: 3.25 (range 0-19)  Symptoms were produced in 57.2% of cases; of these, 48.8% produced while in hospital.		Update of review by Rosenberg (1987). Comparisons made between reviews. Includes cases from articles published 1972-1999.  Reporting bias from case reports and case series.  Possible replication of cases as anonymity to be preserved.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						74 cases of suffocation, 43 cases of giving drugs and 22 cases of poisoning.		
Feldman MD; Brown RM; 2002 May <sup>66</sup>	Study Type: Systematic review - meta-analysis  Evidence level: 2++	122 cases of MSBP		NA Comparison: NA	Follow-up period: NA  Outcome Measures:	Perpetrator (n=93) Mother in 86% of cases Father in 4% Spouses unrelated to cases in 4% Grandmother in 2%  Age (n=76) 26% <3y 52% between 3y and 13y 12% >13y 9% involved adult  54% male (n=81)  Data not given on presentations but said to be similar to presentations in UK and USA data; comment made that induced apnoea more infrequent than in other data.		Some cases appear in <sup>64</sup> .

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
<p>Awadallah N;Vaughan A;Franco K;Munir F;Sharaby N;Goldfarb J;</p> <p>2005 Aug</p> <p>70</p>	<p>Study Type: Systematic review - meta-analysis</p> <p>Evidence level: 2-</p>	<p>42 children from literature review (1966-2002)</p> <p>9 children from local clinic (2001-2003)</p>	<p>Children older than 6 years with MSBP.</p> <p>Mean age in lit review cases (n=32): 8.8 years (range 6-17years)</p> <p>Mean age in clinic cases (n=9): 11.3 years (range 9-16years)</p>	<p>NA</p> <p>Comparison: NA</p>	<p>Follow-up period: NA</p> <p>Outcome Measures:</p>	<p>Lit review cases 48% male False reporting in 62% Falsification of records or sample in 14% Induced illness in 57%</p> <p>Chronic subjective pain in 31%.</p> <p>Induced illnesses included poisoning, inappropriate medication, injection-related infections, starvation, seizures from medication overdoses, suffocation and induced ALTE.</p> <p>Clinic case series 2 males False reporting in 9 Fabrication of records or samples in 1 Induced illness in 2 Diagnoses: rash, aches and fever after surgery, chronic pain, juvenile rheumatoid arthritis, neurological complaints, seizures, corneal abrasions.</p>		<p>Overlap of many cases in <sup>64</sup>.</p>

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Korpershoek M;Flisher AJ;  2004  67	Study Type: Systematic review - meta-analysis  Evidence level: 2++	Literature review	NA	NA  Comparison: NA	Follow-up period: NA  Outcome Measures:		Section on presentations of MSBP details results from <sup>64</sup> , <sup>169</sup> and <sup>65</sup>  Folks(1995) found two patterns of presentation: apnoea, seizures and cyanosis or d&v, nausea, bone and joint problems.  Most common forms of assault were suffocaation, giving drugs and poisoning..  Noted wide variety in fabricateed illnesses. History of multiple hospitalisations and repeated medical investigations also mentioned.	
de Ridder L.;Hoekstra JH;  2000 Aug  69	Study Type: Systematic review - meta-analysis  Evidence level: 2++	NA	Cases of MSBP involving gastroenterology.	NA  Comparison: NA	Follow-up period: NA  Outcome Measures:		Manifestations of MSBP in paediatric gastroenterology.  Chronic diarrhoea, failure to thrive, vomiting, abdominal pain, hematemesis, gastric erosions, allory-Weis tears, colitis, hematochezia, constipation, cystic fibrosis, central line complications, ingestion of foreign bodies, creating aphthous ulcers in the mouth.	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
<p>Feldman KW;Stout JW;Inglis AF; 2002 May 71</p>	<p>Study Type: OtherCase series  Evidence Level: 3</p>	<p>NA  Comparison: PCF children with history of allergy, asthma, sinopulmonary infections, ENT surgery or drug sensitivity compared to PCF children without these attributes.</p>	<p>104 children identified with paediatric condition falsification</p>	<p>71 children with PCF history of allergy, asthma, sinopulmonary infections, ENT surgery or drug sensitivity compared  33 PCF children without these attributes.</p>		<p>Of 99 children with data, presenting complaints (reason that PCF was brought to HCP's attention) were:  Asthma: 4 Sinopulmonary disease: 8 Hearing loss: 2 CNS disease/seizure: 23 Apnoea: 17 GI symptoms: 15 Other infections:8 Failure to thrive: 5 Sexual abuse: 2 Immune dysfunction: 1 Other: 3  Mother was perpetrator in all cases.  Associated falsified or induced conditions. PCF children with history of allergy, asthma, sinopulmonary infections, ENT surgery or drug sensitivity vs other PCF patients Haematologic bleeding 27% vs 27% Undifferentiated sepsis 14% vs 3% Catheter sepsis: 13% vs 0% UTI 11% vs 6% Vomiting 46% vs 36% Past pH probe study for reflux: 28% vs 18% Diarrhoea: 34% vs 27% Failure to thrive: 35% vs 27% Apnoea: 46% vs 39% Past apnoea study: 21% vs 24% Seizures: 45% vs 24%</p>		<p>Cases identified over 24 years (1974-1998) through author's practice.  Evidence that carer intentionally falsified history of nonexistent illness, exaggerated history of legitimate illness, fabricated medical signs and symptoms or induced illness in the child.  Siblings included in study so correlations present.</p>



DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						ADHD 29% vs 6% etc  Median age of onset: 4 months vs 1 month Median age at diagnosis: 48m vs 7m		
Light MJ; Sheridan MS;  1990 Mar  168	Study Type: OtherSurvey  Evidence Level: 3	NA  Comparison: NA	20090	Infants in apnoea monitoring programs		54 children with MSBP and apnoea detailed information available for 32 children; average age at presentation 8.4 weeks presenting diagnoses: prematurity (n=4), ALTE/infantile apnoea (n=25), subsequent sibling of a SIDS case (n=2) family history of apnoea (n=1)		Survey of apnoea monitoring programs

7.1 Emotional, behavioural and interpersonal/social functioning

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
---------------------------	-----------------------------	--------------------	-------------------------	------------	------------------------------	-------------	---------------	-------------------

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Bolger KE;Patterson CJ;  2001 Mar  75	Study Type: Cross-sectional  Evidence level:	107 maltreated children (maltreatment identified before initial participation in study) 107 non-maltreated children matched on age, gender, ethnicity, school and SES.	51% male, 60% white, 40% African-American	Comparison:	Follow-up period:  Outcome Measures:			
Holmes WC;  1998 Dec  77	Study Type: Systematic review - meta-analysis  Evidence level:	NA		Comparison:	Follow-up period:  Outcome Measures:		Compared to nonabused males, abused males were reported to have greater difficulty controlling sexual feelings, were hypersexual and more likely to perpetrate coercive sexual acts against others.	Narrative review. Search dates were 1985-1997
Inderbitzen-Pisaruk H;Shawchuck CR;Hoier TS;  1992 Mar  140	Study Type: Case-control  Evidence level: 2-	17 CSA cases 17 controls	CSA cases validated by Child Protection Department, child protected from perpetrator, age 5 - 15  Controls matched on age, sex, socioeconomic status and current living situation recruited from community. Sexual abuse excluded.	Number of validated questionnaires administered, including CBCL. The 6 sex behaviour problems were combined to give a sex-problem score.  Comparison: CSA vs controls.	Follow-up period:  Outcome Measures: Sex-problem score	Mean (SD) CSA: 1.47 (1.84) Controls: 0.41 (1.23) p=0.05		Small study. Large span of ages in this context.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Kendall-Tackett KA;Williams LM;Finkelhor D; 1993 Jan 74	Study Type: Systematic review - meta-analysis  Evidence level: 1+			Comparison:	Follow-up period:  Outcome Measures:		Nonclinical samples Demeanour or behaviour and the number of studies in which CSA children were more symptomatic than non-cases.  Anxiety: 5/8 studies  Fear: 5/5 studies  Depressed: 10/11 studies  Withdrawn: 11/11 studies  Poor self-esteem: 3/6 studies  Aggressive antisocial: 10/11 studies  Cruel: 2/2 studies  Delinquent 6/6 studies  School/learning problems: 5/6 studies  Regression/immaturity: 2/2 studies  Running away: 1/1 studies  Percentage of CSA cases who were displayed each symptom ranged from 0% to 70%	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Kitzmann KM;Gaylord NK;Holt AR;Kenny ED;  2003 Apr  72	Study Type: Systematic review - meta-analysis  Evidence level:			Comparison:	Follow-up period:  Outcome Measures:			
Wolfe DA;Crooks CV;Lee V;Intyre-Smith A;Jaffe PG;  2003  73	Study Type: Systematic review - meta-analysis  Evidence level: 1-	41 studies	Children exposed to domestic violence	Comparison: Differences in negative emotional and behavioural outcomes between children exposed to domestic violence and children not exposed to domestic violence.	Follow-up period:  Outcome Measures:	No significant differences between males and females in terms of negative outcomes due to domestic violence.  Small and significant effect of domestic violence on negative outcomes (internalising, externalising and PTSD pooled).  Lack of clarity on the effect of age on outcome.		

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
<p>Evans SE;Davies C;DiLillo D;</p> <p>2008</p> <p><sup>76</sup></p>	<p>Study Type: Systematic review - meta-analysis</p> <p>Evidence level: 2-</p>	61 studies		<p>Effects of exposure to domestic violence on internalising and externalising behaviours.</p> <p>Comparison:</p>	<p>Follow-up period: n/a</p> <p>Outcome Measures: Internalising behaviour and externalising behaviour as measured by a validated tool (e.g. CBCL).</p>	<p>Internalising</p> <p>All children: 58 studies, n=7602, weighted sample mean= 0.48 (95%CI 0.39 to 0.57), no significant heterogeneity.</p> <p>Boys: 15 studies, n=1697, weighted sample mean=0.44 (p&lt;0.05), no significant heterogeneity.</p> <p>Girls: 14 studies, n=1758, weighted sample mean=0.39 (p&lt;0.05).</p> <p>Externalising</p> <p>All children: 53 studies, n=7200, weighted sample mean=0.47, 95% CI 0.38 to 0.56, significant heterogeneity.</p> <p>Boys: 16 studies, n=1787, weighted sample mean=0.46 (p&lt;0.05), no significant heterogeneity.</p> <p>Girls: 13 studies, n=1570, weighted sample mean=0.23 (p&lt;0.05).</p> <p>Significant difference between boys and girls' externalising behaviour.</p>		<p>Some crossover in studies with <sup>72</sup>. Studies 1990-2006 used here and unpublished data included.</p> <p>Unclear whether included studies were all comparative. Difficulty in interpreting effect size.</p>

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
<p>Finzi R;Har-Even D;Weizman A; 2003 Sep 86</p>	<p>Study Type: OtherDescriptive study Evidence Level: 3</p>	<p>Child Suicidal Potential Scales (CSPS) A clinician administered interview schedule consisting of nine sections  Comparison: Group 1 (study group) physically abused children (PA) Group 2 (first control group) children neglected by their parents (NEG) Group 3 (second control group) children who were neither abused nor neglected (N-M)</p>	<p>Sample of 114 children divided into three groups on the basis of maltreatment Group 1 (study group) consisted of 41 physically abused children (PA) Group 2 (first control group) consisted of 38 children neglected by their parents (NEG) Group 3 (second control group) consisted of 35 children who were neither abused nor neglected (N-M)</p>	<p>Children aged 6 to 12 years</p>	<p>Ego Defenses-Regression Denial Projection Introjection Reaction-Formation Undoing Displacement Intellectualization Compensation Sublimation Repression</p>	<p>Significant differences between physically abused and the non abused/ non neglected (N-M) for all ego defenses except displacement.  Significant differences between physically abused (PA) and neglected children (NEG) for regression, denial and splitting, projection, and introjection (high scores for physically abused (PA) children) for compensation and undoing (higher scores for the neglected (NEG) children)</p>	<p>Physically abused children should be distinguished as a high-risk population for future personality disorders.</p>	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Wells RD;McCann J;Adams J;Voriss J;Ensign J; 1995 Feb 87	Study Type: OtherDescriptive study Evidence Level: 3	Structured Interview for Signs Associated with Sexual Abuse (SASA)  Comparison: - Nonabuse group (NA group) -Sexual abuse with perpetrator confession (SA group) -Sexual abuse without perpetrator confession (AA group)	3 matched samples -68 in Nonabuse group (NA group) -68 in Sexual abuse with perpetrator confession (SA group) -68 in Sexual abuse without perpetrator confession (AA group)	Parents of prepubescent females	Generalized signs of disturbance-sleep problems, withdrawal, concentration problems Specific sexualized symptoms-increased masturbation, sexual aggressiveness, increased knowledge	Both SA and AA groups reported increased sleep problems, fearfulness, emotional and behavioral changes, concentration problems, and sexual curiosity and knowledge.  Self consciousness, nightmares, and fearfulness of being left alone emerged significantly more frequently in SA group as compared to AA group	The presence of signs and symptoms in sexually abused children should be considered suggestive of abuse although the lack of symptoms does not necessarily confer safety form victimization.	
Dubowitz H;Black M;Harrington D;Verschoore A; 1993 Nov 170	Study Type: Other Evidence Level:	Comparison:						Parents or guardians of children suspected of being sexually abused completed the CBCL.  Comparisons recruited from primary care clinics.  Likelihood of abuse rated by interdisciplinary team.
Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Slusser MM; 1995 171	Study Type: Systematic review - meta-analysis  Evidence level: 2-	NA		Comparison:	Follow-up period:  Outcome Measures:		Overt sexual behaviour, inappropriate for age, is an indication of sexual abuse.	Narrative review of 6 studies. Study included if used comparison group, validated assessment tool, structured clinical interviews and systematic clinical record review.
Zolotor A;Kotch J;Dufort V;Winsor J;Catellier D;Bou-Saada I; 1999 Mar 81	Study Type: Cohort  Evidence level: 2-	842 primary caregivers predominantly mothers with their infants	Primary caregivers, predominantly mothers eligibility criteria low birth weight (below 2,500 g) low maternal age (less than 18 years of age) significant medical problems such as intracranial hemorrhage, meningitis, seizures or respiratory distress syndrome Significant social problems such as single parent without family support problem, or any parent with incapacitating medical or mental handicap or alcohol or substance abuse	Achenbach Teacher Report form and project developed questions regarding peer status.  Comparison: for every 4 eligible infants whose mothers agreed to participate, the next mother to deliver an infant without any risk criteria was recruited	Follow-up period: At 6 years (n=217) and 8 years (n=181)  Outcome Measures: School performance measured by -academic success -peer status -adaptive functioning	Mean academic performance (100-500) at age 6 was 260 (SD=85) at age 8 was 263 (SD=95) Mean Peer status (1-5) at age 6 was 3.5 (SD=0.85) at age 8 was 3.3 (SD=0.96) Total adaptive functioning (4-28) at age 6 was 14.6 (SD=5.16) at age 8 was 14.6 (SD=5.28) Maltreatment associated significantly associated with poorer academic performance (p<0.01) and poorer adaptive functioning (p<0.001) but not with peer status	The study concluded that maltreatment is associated significantly with poorer academic performance (p<0.01) and poorer adaptive functioning (p<0.001) but not with peer status.	



DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Antao V;Maddocks A;Street E;Sibert JR;  1996 <sup>80</sup>	Study Type: Case-control  Evidence level: 2-	107 school boys sexually abused (cases) 107 school boys not sexually abused (controls)	Majority of boys aged 8 to 10 years	Information from general practice records, school health records, hospital records and the records of consultant child psychiatrists  Comparison: Boys sexually abused (cases) and boys not abused (controls)	Follow-up period:  Outcome Measures: Somatic and behavioural symptoms	Somatic and behavioural symptoms uncommon in both cases and controls 83.6% of cases and 76.7% controls did not have symptoms.  No significant difference between the numbers of cases and controls who had presented with somatic and behavioral complaints (18 cases vs 25 controls).  Significant difference between cases and controls with symptoms lasting over a year (p<0.05).	Where somatic symptoms are long standing should be considered as a possible diagnosis for abuse.	
Baer J;Martinez CD;  2006 <sup>88</sup>	Study Type: Systematic review - meta-analysis  Evidence level: 1+	791 children from 8 studies	Studies selected if maltreated children were <48 months old, studies used 'Strange Situation' tool, included comparison groups, reported sufficient data.	Comparison: Insecure vs secure attachment style in maltreated vs. nonmaltreated children.	Follow-up period:  Outcome Measures: Odds of showing insecure attachment and having been maltreated vs. not being maltreated.	Odds of having insecure attachment in abused group compared to nonabused group 6.5 (95% CI 3.7-11.6)	Maltreated children under 2 years of age are more likely to have insecure or disorganized attachment than their non-maltreated peers.	The results of the study reflect a composite score and do not provide key information for healthcare professionals who may observe one-off interactions.
van Ijzendoorn MH;Schuengel C;Bakermans-Kranenburg MJ;  1999 <sup>89</sup>	Study Type: Systematic review - meta-analysis  Evidence level: 1+	For question of interest 323 children from 5 studies		Attachment disorganisation as measured by a validated tool.  Comparison: Attachment styles in maltreated and nonmaltreated children under 2 years of age.	Follow-up period: NA  Outcome Measures:	Across all studies, 48% of maltreated children had disorganised attachment compared to 17% of nonmaltreated children.  Pooled effect size (standardised correlation coefficient): 0.41	Maltreatment (including failure to provide adequate nourishment) is associated with disorganised attachment.	At least 3 of the samples used in the meta-analysis are used in <sup>88</sup> .

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Eckenrode J;Laird M;Doris J;  1993 Jan  83	Study Type: OtherComparative study  Evidence Level: 3	Iowa test scores  Comparison: Maltreated children compared with nonmaltreated children	420 maltreated children 420 nonmaltreated children The 2 groups matched on the following variables: gender, school, grade level, residential neighborhood, and classroom.	Children in kindergarten through grade 12	Academic performance- Test scores in reading and math- Iowa tests Final grades Grade repetitions Discipline referrals and suspensions	Comparable percentile scores in math 49.5 for maltreated children and 60.03 for nonmaltreated children $t(411)=4.4, p<0.001$ Reading/ english mean grade 2.0 for maltreated children 2.3 for nonmaltreated children  Overall percentage of repeating a grade in entire sample 28.6% Likelihood of repeating much higher in maltreated children $X^2(1, N=773)=32.3, p<0.001$  Maltreated children are 2.5 times more likely to repeat a grade than nonmaltreated children $OR= 2.53 (.606/.239)$ 95% CI 1.8-3.4  Discipline Referrals for entire sample 29.2% had atleast one discipline referral Among maltreated children 34.3% had one or more referrals Among nonmaltreated children 24.1% had one or more referrals $p<0.01$  Suspensions for entire sample 25.1% had been suspended atleast once Among maltreated children 1.2	The study concluded that maltreated children performed significantly below nonmaltreated children in standardized tests and grades and were more likely to repeat a grade. Maltreated children also had significantly more discipline referrals and suspensions.	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
						Among nonmaltreated children 0.4 p<0.01		
Rowe E;Eckenrode J;  1999  82	Study Type: OtherComparative study  Evidence Level: 3	Survival analysis used to investigate the timing of risk of experiencing an academic difficulty.  Comparison: Maltreated children compared with non-maltreated children.	300 maltreated children 300 non-maltreated children	Children (aged 5-18 years) in grades K-12 enrolled in public schools	Grade repetitions Poor English grades Poor math grades	Maltreated children at higher risk than non-maltreated children of repeating kindergarten and first grade.  No difference in the risk of repeating grade for the first time.  Absolute risk of receiving a poor english or mathematics grade changed across elementary years whereas the relative risk by maltreatment status did not.	The risk changed across time for grade repetitions but not for the first occurrence of a poor English or mathematics grade.	
Quas JA;Goodman GS;Jones D;  2003 Jul  84	Study Type: OtherDescriptive study  Evidence Level: 3	Sexual Assault profile Child behavior checklist Social adjustment scale  Comparison: NA	218 victims of sexual abuse	Victims of sexual abuse aged 4 to 17 years	Child characteristics Characteristics of the abuse Maternal support following disclosure of the abuse	Child having these factors had increase attributions of self blame -close relationship with the perpetrator -severe sexual abuse -perceiving sexual abuse as disgusting -coping with abuse by pretending it never happened  These factors did not predict internalizing behavior problems.	Self-blame attributions and behavior problems need to be distinguished as separate outcomes in children who are sexually abused.	
Reyome ND;  1994 Oct  85	Study Type: OtherDescriptive study  Evidence Level: 3	Class room behaviors linked with school achievement Hahnemann Elementary school behavior rating scale used	33 maltreated children 33 non-maltreated children receiving public assistance 33 non-maltreated lower middle class children	Maltreated and non-maltreated (on public assistance and lower middle class) children from 5 to 12 years of age	Achievement related classroom behaviors.	Maltreated children exhibited less classroom behavior positively linked with academic achievement as compared to non-maltreated children (receiving public assistance).	Maltreated children exhibited less classroom behavior positively linked with academic achievement as compared to non-maltreated children (receiving public assistance) and non-maltreated children of	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
		Comparison: Maltreated compared with non-maltreated (receiving public assistance and lower middle class)				<p>Maltreated children did not differ significantly from children receiving public assistance in most behaviors negatively linked with academic achievement (except withdrawn behavior) but clearly engage in less academically oriented classroom behaviors.</p> <p>Maltreated children significantly engaged in less classroom behavior positively linked to academic achievement and significantly more classroom behavior negatively related to academic achievement as compared to non-maltreated children of lower middle class.</p>	lower middle class.	
<p>Wells RD;McCann J;Adams J;Voriss J;Ensign J;</p> <p>1995 Feb</p> <p>87</p>	<p>Study Type: OtherDescriptive study</p> <p>Evidence Level: 3</p>	<p>Structured Interview for Signs Associated with Sexual Abuse (SASA)</p> <p>Comparison: - Nonabuse group (NA group) -Sexual abuse with perpetrator confession (SA group) -Sexual abuse without perpetrator confession (AA group)</p>	<p>3 matched samples</p> <p>-68 in Nonabuse group (NA group)</p> <p>-68 in Sexual abuse with perpetrator confession (SA group)</p> <p>-68 in Sexual abuse without perpetrator confession (AA group)</p>	Parents of prepubescent females	<p>Generalized signs of disturbance-sleep problems, withdrawal, concentration problems</p> <p>Specific sexualized symptoms-increased masturbation, sexual aggressiveness, increased knowledge</p>	<p>Both SA and AA groups reported increased sleep problems, fearfulness, emotional and behavioral changes, concentration problems, and sexual curiosity and knowledge.</p> <p>Self consciousness, nightmares, and fearfulness of being left alone emerged significantly more frequently in SA group as compared to AA group</p>	<p>The presence of signs and symptoms in sexually abused children should be considered suggestive of abuse although the lack of symptoms does not necessarily confer safety form victimization.</p>	

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Maas C;Herrenkohl TI;Sousa C;  2008  78	Study Type: Systematic review - meta-analysis  Evidence level: 2+	8 studies		Comparison:	Follow-up period:  Outcome Measures:			Authors looked for longitudinal studies on association of child maltreatment with youth violence in 12-21 year olds.  Authors conclude: physical abuse more likely to lead to youth violence than other forms of abuse; extreme and/or repeated abuse increases risk for youth violence.

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Root C;MacKay S;Henderson J;Del B;Warling D; 2008 79	Study Type: Other  Evidence Level: 3	Comparison: maltreated vs non-maltreated firesetters  Children defined as maltreated if caregiver confirmed that the child had ever experienced at least one of physical abuse, physical neglect or sexual abuse.  'suspected abuse' cases were excluded	205 caregivers and their children	87% male mean age 11.2y, sd 3.1y) children referred to assessment and treatment programme for juvenile firesetters.	fire-setting frequency, versatility, interest ascertained by Fire Involvement Interview. - semi-structured interview asking. Frequency = total number of episodes coded 1-10 where 10 represents all numbers above 10  versatility=sum of different ignition materials and target types (out of a possible 7 and 6, respectively)  externalising behaviour (CBCL) internalising behaviour (CBCL)	48% maltreated  frequency (maltreated vs non-maltreated) 6 vs 5 (p<0.05)  versatility 4 vs 3 (p<0.05)  fire interest 15 vs 12 (ns)  externalising 71.6 vs 64.7 (p<0.05) internalising 65.8 vs 57.6 (p<0.05)		Maltreatment status validated by cross-referring to child protective services data.  maltreatment was as common as non-maltreatment in fire-setting children. frequency and versatility were greater in maltreated children.

### 7.2.1 Self-harm

90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 115, 121 to be added.

### 7.2.2 Repeated nightmares

No literature identified.

### 7.2.3 Abdominal pain

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Rimsza ME; 1988 <sup>123</sup>	Study Type: Case-control  Evidence level: 2-	67 girls who reported sexual abuse and 5 who had abuse confirmed on clinical examination  68 controls of similar age, sex and initial clinic visit date	Cases: female children who were evaluated at a medical centre because of sexual abuse Mean age 10 years (range 2 years to 17 years)  Controls: females identified from the general clinic admission records.	Comparison: Sexual abuse vs controls	Follow-up period: mean = 24 months (range 9 mo - 48 mo)  Outcome Measures: Skeletal muscle tension (including sleep problems), gastrointestinal irritability (including chronic abdominal pain), Genitourinary symptoms (including dysuria and vaginal discharge), "Emotional reactions" (including suicide attempt), Runaway behaviour, School problems, "Other behaviour problems", Early pregnancy	Sexually abused vs controls Muscle tension: 26% vs 7%, p<0.01  Gastrointestinal irritability: 31% vs 10%, p<0.01 [Chronic abdominal pain: 26% vs 7%, no analysis] Genitourinary symptoms: 40% vs 10%, p<0.001 Emotional reactions: 18% vs 3%, p<0.001 Runaway behaviour: 8% vs 0%, p<0.05 School problems: 10% vs 10%, p=NS Other behavioural problems: 28% vs 4%, p<0.001 Early pregnancy: 8% vs 12%, p=NS	Cases are more likely than controls to display somatic or behavioural difficulties after a period of sexual abuse has ended.	This study is a review of medical records, confirmation was ascertained by patient history or medical examination (in 5 cases).  This study reports somatic and behavioural difficulties after abusive period. It is not clear how long this is.

#### 7.2.4 Disturbances in eating and feeding behaviour

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
---------------------------	-----------------------------	--------------------	-------------------------	------------	------------------------------	-------------	---------------	-------------------

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Chandy JM;Blum RW;Resnick MD;  1996 Dec  128	Study Type: Cross-sectional  Evidence level: 2-	370 males, 2681 females	adoescents who reported sexual abuse in a large school survey	NA  Comparison:	Follow-up period:  Outcome Measures:	Evaluates self as overweight f=52.3%, m=21.3% (p<0.001)  V satisfied with present weight: f=8.1%, m=26.0% (p<0.001)  V proud of body f=6.9%, m=28.4% (p<0.001)  Binge-eating: f=40.8%, m=22.3% (p<0.001)  Non-stop eating: f=22.7%, m=8.1% (p<0.001)  Dieting during last year: f=69.6%, m=26.8% (p<0.001)  Self-induced vomiting: f=19.9%, m=10.4% (p<0.001)  Use of diuretics: f=3.7%, m=1.4% (p<0.05)  Use of laxatives: f=3.0%, m=1.6% (NS)  Use of ipecac: f=1.4%, m=1.1%		Sexual abuse=someone in your family, or someone else, touches you in a place you did not want to be touched, or does something to you sexually which they shouldn't have done.
Hernandez J;  1995  127	Study Type: Case-control  Evidence level: 3	6224 children (10% subsample of Minnesota study)	3238 males, 2986 females  318 females who reported eating disorder and abuse 84 males who reported eating disorder and abuse	Comparison:	Follow-up period:  Outcome Measures:	Eating disorders more commonly reported in abused than non-abused whether abuse was physical or sexual.		Eating disorder defined as two of the following true: out-of-control eating, using laxatives and vomiting  10% randomly selected subsample of Minnesota study.  9th and 12th graders only



DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Perkins DF;Luster T;  1999 Apr  129	Study Type: Other  Evidence Level: 3	Comparison:	7903 female adolescents	Mean age 14.5y (sd 1.6)		Physically abused girls more likely to purge than non-physically abused girls: OR=1.81 (p=0.0014) (from logistic regression model)  No signif relationship found between sexual abuse and purging behaviour.		Surveyquestionnaire. Stat modelling fairly simple. Opportunity for multilevel modelling (with school as one of the levels).  Abuse = Have you ever been sexually abused? Have you ever been physically abused by an adult (that is, where an adult caused you to have a scar, black and blue marks, welts, bleeding or a broken bone?  Purging = How often do you vomit on purpose after eating?  Study suggests physical abuse is associated with purging behaviour but sexual abuse is not.  No info on past or current abuse.

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Chandy JM;Blum RW;Resnick MD;  1996 Dec  126	Study Type: Other  Evidence Level: 3	Comparison:	1011 girls with history of CSA, 1011 comparison subjects	Female adolescents mean age 15.28 vs 14.92		Abuse vs no abuse Thinks of self as overweight 55.6 vs 43.7%  Binge-eating 40.3% vs 31.7%  Nonstop eating 24.6% vs 16.7%  10+ dieting episodes in last year 17.9% vs 12.3%  Use of diuretics 4.4% vs 2.7%  Self-induced vomiting 1+ times per week 4.4% vs 2.7%  Use of laxatives 3.7% vs 2.2%  All statistically significant		Simple proportions presented. No accounting for confounders. Multiple comparisons.  Time between 'abuse' and eating behaviours unclear.

Wonderlich SA;Crosby RD;Mitchell JE;Roberts JA;Haseltine B;Demuth G;Thompson KM; 2000<sup>125</sup> to be added.

### 7.2.5 Selective mutism

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Intervention & Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
---------------------------	-----------------------------	--------------------	-------------------------	---------------------------	------------------------------	-------------	---------------	-------------------

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Intervention & Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
MacGregor R; Pullar A; Cundall D; 1994 130	Study Type: Case-control  Evidence level: 2-	52 (18 each of selectively mute, speech or language problems, no speech or language problem)	Children >=6years, mute in school for >1y but able to speak normally in other circumstances.  Age 6-14years, controls matched by age and sex from same class at school.	Intervention: Does maltreatment lead to selective mutism?  Comparison:	Follow-up period: n/a  Outcome Measures: Abuse status derived from community paediatrics database.	Selectively mute children - 5 definitely abused, 3. possibly  Speech or language problems - 1 possibly abuse  Controls - no abuse		

### 7.2.6 Head-banging and body rocking

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Troster H; 1994 Feb 131	Study Type: Cross-sectional  Evidence level: 2-	142 140 included in analysis of interest including 45 with history of suspected abuse	Children in residential care homes, aged 10 mo - 11y,	Comparison: Suspected abuse vs non-abuse.	Follow-up period: Cross-sectional study but data are observations of weekly or daily behaviours.  Outcome Measures: Percentage of children in whom behaviours were observed at least once day.	Suspected abuse n=45 Other n=95  Body rocking 11.1% (susp. abuse) vs. 6.3% (other) Not significant  Head nodding/shaking 4.4% (susp. abuse) vs. 4.2% (other) Not significant  Head banging 4.4% (susp. abuse) vs 1.1% (other) Not significant		Data collected via caregivers' ratings.  Children not living with parents so hard to disentangle cause of stereotyped behaviour.  Similarly, non-significance of proportions between suspected abuse and other children is biased by the reason that the children are in residential care.  Suspicion of abuse is not described.

### 7.2.7 Wetting and soiling

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Klevan JL;De Jong AR;  1990 Feb  34	Study Type: Other  Evidence Level: 3	NA  Comparison:	428 CSA victims seen at sexual assault health centre based in a hospital.	mean age 8.6 years (range 1-16), 84% female		85 (20%) of sample had GU symptoms at 1-3 weeks after first report of CSA.  Recent onset of enuresis in 24 (6% of total series) Vaginal pain 43 (10%) Dysuria 21 (5%) Increased urinary frequency 20 (5%)		Cohort of sexually abused children. No data on non-abused children.
Mellon MW;Whiteside SP;Friedrich WN;  2006 Feb  134	Study Type: Other  Evidence Level: 2+	Comparison: NA	1114 normative children, 577 psychiatric patients, 620 children with confirmed history of CSA mostly in the last 12 months	Children aged 2-12 years.		Prevalence of encopresis  Normative vs psych vs abused 2% vs 10.5% vs 10.3%  Does encopresis=abuse? Sensitivity 10%, PPP 45%  Broken down by age and sex, lowest		PPV depends on prevalence of abuse in the sample.  Encopresis defined as a rating of "sometimes true" or "often true" on the "bowel movements outside the toilet" item on the CBCL.

### 7.2.8 Sexualised behaviour

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Cosentino CE;Meyer-Bahlburg HF;Alpert JL;Weinberg SL;Gaines R;  1995 Aug  138	Study Type: Case-control  Evidence level: 2+	Sexually abused = 20 Psychiatric controls = 20 Nonpsychiatric controls = 20	Girls aged 6-12 years	Child sexual behaviour inventory  Comparison: CSA vs psych. controls and nonpsych controls	Follow-up period: NA  Outcome Measures: Child sexual behaviour checklist	Total CSBI mean (sd) CSA 30.6 (20.3) Psych. controls 15.2 (9.9) Non-psych controls 10.8 (9.6)  Significant difference found between groups.	Sexually abused girls exhibited more sexualised behaviours than girls who had not been sexually abused.	Cases attended a treatment centre for child abuse, had been referred there within the preceding 2 years and were undergoing treatment.

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Slusser MM; 1995 171	Study Type: Systematic review - meta-analysis  Evidence level: 2-	NA		Comparison:	Follow-up period:  Outcome Measures:		Overt sexual behaviour, inappropriate for age, is an indication of sexual abuse.	Narrative review of 6 studies. Study included if used comparison group, validated assessment tool, structured clinical interviews and systematic clinical record review.
Gordon BN;Schroeder CS;Abrams JM; 1990 Apr 139	Study Type: Cross-sectional  Evidence level: 2-	Abused children: n=22 Nonabused children: n=22	Abused children recruited at first contact with psychology clinic. Nonabused children recruited at paediatric clinic and public health clinic. Age not stated but cases and controls matched on age and socioeconomic status.	Comparison:	Follow-up period:  Outcome Measures: Areas of knowledge: Gender identity Sexual body parts Nonsexual body parts Sexual behaviour Pregnancy Private parts Total knowledge	Abuse vs. non-abuse Mean score (s.d)  Gender identity 16.6 (2.5) vs. 17.4 (4.0) n.s  Sexual body parts 9.5 (4.1) vs. 9.3 (4.8) n.s  Nonsexual body parts 12.6 (2.1) vs. 11.4 (2.9) n.s  Sexual behaviour 4.9 (2.6) vs. 4.0 (2.3) n.s  Pregnancy 4.9 (4.3) vs. 5.6 (4.2) n.s  Private parts 8.1 (5.6) vs. 7.1 (6.4) n.s.  Total knowledge 57.0 (15.6) vs. 55.0 (19.5) n.s	No observed differences in sexual knowledge between abused and non-abused children.	Confounders (such as parents' marital status, parental attitude, sex education) not accounted for in analysis.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Holmes WC; 1998 Dec 77	Study Type: Systematic review - meta-analysis  Evidence level:	NA		Comparison:	Follow-up period:  Outcome Measures:		Compared to nonabused males, abused males were reported to have greater difficulty controlling sexual feelings, were hypersexual and more likely to perpetrate coercive sexual acts against others.	Narrative review. Search dates were 1985-1997
Inderbitzen-Pisaruk H; Shawchuck CR; Hoier TS; 1992 Mar 140	Study Type: Case-control  Evidence level: 2-	17 CSA cases 17 controls	CSA cases validated by Child Protection Department, child protected from perpetrator, age 5 - 15  Controls matched on age, sex, socioeconomic status and current living situation recruited from community. Sexual abuse excluded.	Number of validated questionnaires administered, including CBCL. The 6 sex behaviour problems were combined to give a sex-problem score.  Comparison: CSA vs controls.	Follow-up period:  Outcome Measures: Sex-problem score	Mean (SD) CSA: 1.47 (1.84) Controls: 0.41 (1.23) p=0.05		Small study. Large span of ages in this context.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Kendall-Tackett KA;Williams LM;Finkelhor D; 1993 Jan 74	Study Type: Systematic review - meta-analysis  Evidence level: 1+			Comparison:	Follow-up period:  Outcome Measures:		Nonclinical samples Demeanour or behaviour and the number of studies in which CSA children were more symptomatic than non-cases.  Anxiety: 5/8 studies  Fear: 5/5 studies  Depressed: 10/11 studies  Withdrawn: 11/11 studies  Poor self-esteem: 3/6 studies  Aggressive antisocial: 10/11 studies  Cruel: 2/2 studies  Delinquent 6/6 studies  School/learning problems: 5/6 studies  Regression/immaturity: 2/2 studies  Running away: 1/1 studies  Percentage of CSA cases who were displayed each symptom ranged from 0% to 70%	

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Merrick MT; Lutrownik AJ; Everson MD; Cox CE; 2008 <sup>141</sup>	Study Type: Other  Evidence Level: 2-	Are sexualised behaviours related to early (before age 4y) or late (between 4y and 8y) maltreatment. Sexual abuse cases were excluded.  Comparison: Binary variables of: early physical abuse y/n early emotional abuse y/n early neglect y/n late physical abuse y/n late emotional abuse y/n late neglect y/n	690	maltreated children and children at risk of maltreatment, age ~8 years at time of interview 47% female	predictors of sexualised behaviours: boundary problems displaying private parts sexual interest sexual intrusiveness sexual knowledge	Boundary problems:  Late physical abuse OR: 1.94 95% CI 1.1-3.5  Displaying private parts:  Early physical abuse OR 2.4 95% CI 1.1-5.4  Early emotional abuse: OR 0.3 95% CI 0.1-0.8  Late physical abuse OR: 2.4 95% CI 1.0-5.6  Sexual interest No significant predictors  Sexual intrusiveness: Early neglect OR: 0.4 95% CI 0.2-0.9  Late physical abuse OR: 2.6 95% CI 1.3-5.2  Sexual knowledge: Late emotional abuse OR 2.0 95% CI 1.2-3.4		No normative data. Predictors of sexualised behaviours misleading because of the mix in the 'no' groups

Hall T; Hogben M; Carlton AL; Liddon N; Koumans EH; 2008 <sup>142</sup> to be added.

### 7.2.9 Runaway behaviour

<sup>154</sup>, <sup>153</sup>, <sup>152</sup>, Powers (1990)<sup>143</sup> and Powers (1988)<sup>144</sup> Stiffman(1989)<sup>145</sup> Gary (1996)<sup>146</sup>, Warren (1997)<sup>147</sup> and Warren (1994)<sup>148</sup>, Kufeldt (1987)<sup>150</sup> to be added.

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Intervention & Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
---------------------------	-----------------------------	--------------------	-------------------------	---------------------------	------------------------------	-------------	---------------	-------------------



DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Number of Patients	Patient Characteristics	Intervention & Comparison	Follow-up & Outcome Measures	Effect Size	Study Summary	Reviewer Comments
Thompson SJ; Zittel-Palamara KM; Maccio EM; 2004 Dec 149	Study Type: Cross-sectional Evidence level: 2-	156	44% male 79% African-American 40% European-American mean age 16y (sd 1.5y)	Intervention: Comparison:	Follow-up period: Maltreatment status ascertained up to 48 hours after admission to shelter for runaway youth. Outcome Measures:	Physical: 35% Sexual: 12% Emotional: 30% Neglect: 29%		
Craig TK; Hodson S; 1998 Nov 172	Study Type: Cohort Evidence level: 2-	161 homeless 104 domiciled	Young people 16-21 years modal age 17 in both groups	Intervention: Comparison:	Follow-up period: Outcome Measures:	Reasons for running away: conflict with parents (most common reason, numbers not stated) 11 people in domiciled group had run away before. Total of 20 cited physical violence or sexual abuse as reason for running.		Runaway=leaving formal care arrangement for at least one night before age 16 without carer's consent - recruited from inner London accommodation facilities for young homeless people.  Domiciled=recruited from inner city GP practice lists.

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients & Patient Characteristics	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Feitel B; Margetson N; Chamas J; Lipman C; 1992 Feb 151	Study Type: Other Evidence Level:	Intervention: NA Comparison:	372	Homeless and runaway adolescents 55% male, median age 17 years (range 13-21), 48% white, 22% gay, lesbian or bisexual		47% reported physical abuse as a child		

7.2.10 Dissociation

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Lansford JE;Dodge KA;Pettit GS;Bates JE;Crozier J;Kaplow J;  2002  USA  155	Study Type: Prospective longitudinal study  Evidence Level: 2+	To determine whether child physical maltreatment early in life has long-terms effects on psychological (including dissociation), behavioural and academic problems	n=585 children	Participants were randomly recruited from two cohorts of children in 1987 and 1988 from kindergarten in public schools. The sample was 52% male and 48% female. The ethnic profile was 82% European American, 16% African American and 2% other. No ages given	Detailed interviews of developmental history with mothers in home prior to entrance to kindergarten including details on child misbehaviour, discipline practices and whether child has ever been harmed by an adult. At this point, Investigators also rated whether maltreatment had occurred as 0= definitely no physical maltreatment and 1= physical maltreatment probably having occurred. This interview also used the Retrospective Infant Characteristics Questionnaire.  Children's official school records were available from 9th to 11th grades  At 11th grade, mothers completed 113-item Child Behaviour Checklist (CDC)  On completion of 11th grade	CDC dissociation subscales results: (mother reports)  Unadjusted analysis *not maltreated (n=392) maltreated (n=52)  Not maltreated 1.68 SD 0.11 Maltreated 3.38 SD 0.43 F= 28.63 p<0.001  Covariates adjusted analysis (socioeconomic status, single-parent status, family stress, maternal social support, child's exposure to violence, child temperament, child health and harsh parental treatment during adolescence)  Not maltreated 1.58 SD 0.16 Maltreated 2.8 SD 0.37 F=10.01 p<0.01  *presence or absence of maltreatment was determined at the initial interview by two interviewers independently. Agreement was 90% (kappa 0.56)	There was a significant association between suspected child physical maltreatment in pre-kindergarten children and the presence of dissociation later in school life (11th grade).	This study was rated EL=2-  The strengths of this study were that it was it was conducted over a 12 year period in a normative setting with a suitable control group.  The weaknesses were that the presence of child maltreatment at the start of the study was assessed in the home by an interviewer. No details given but presumably a psychology professional  Random sampling was not described  Only physical abuse was considered  The number of abused children was small n=52.  Details of dropouts were not described

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
					adolescents completed Youth Self-report form of the CDC			This study was funded by the Child Development Project from the National Institute of Mental Health Bethesda, USA
Macfie J;Cicchetti D;Toth SL;  2001 Sep  USA  156	Study Type: Case control series  Evidence Level: 2-	To investigate a) the link between child maltreatment and pre-school children and b) to examine which sub-groups of maltreated preschoolers are likely to present with dissociation.	n= 198 pre-school children	Pre-school children mean age 5 yrs SD 6mths (range 3 yrs 7 mths to 6 years). 62% ethnic minorities: 35% African American, 7% Hispanic, 20% other. Recruited from families referred to social services and classified as physically abused, sexually abused, neglected or non-maltreated  no numbers given per group	Child Dissociative Checklist (CDC)	Using ANOVA there was a significant overall effect for maltreatment subtypes on dissociation $F(3, 194)=21.05$ $p<0.00001$  mean dissociation scores per group (no SD given, p value compared with non-maltreated group)  Physical abuse 8.91 Sexual abuse 7.27 Neglected group 5.52 Non-maltreated group 1.88  All clinical groups demonstrated greater dissociation than the non-maltreated group, $p<0.001$ for all.  Further analysis (Pearson r correlations) between the three maltreatment subgroups showed physical abuse and neglect were significantly related to dissociation ( $P<0.001$ for both) but sexual abuse was not ( $p>0.1$ ).	Child maltreatment is a factor in pre-school aged children. Sexually abused, physically abused and neglected groups each demonstrated more dissociation than the non-maltreated group. Dissociation within the three clinical groups was associated greatest with physical abuse.	This study was rated EL=2-  Small study  control group comprised of children who were referred to social services i.e. not general population  62% were of minority status  Therefore external validity is limited.  The funding of this study was undeclared.

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
<p>Kisiel CL;Lyons JS;  2001 Jul USA  157</p>	<p>Study Type: Case-control series  Evidence level: 2-</p>	<p>To investigate the role of dissociation as a mediator of mental-health outcomes in children with a history of sexual abuse</p>	<p>n=114 children and adolescents</p>	<p>Children aged 10 to 18 years old, 52% male and 48% female. The majority were African American (69%), 24% Caucasian and 5% Hispanic who were recruited from a group who were wards of the social services Inclusion criteria were 1) removed from family 2) placed in residential treatment 3) suitable age 4) from the city area 5) agreed to participate They were classified into no abuse, sexual abuse, physical abuse, sexual and physical abuse</p>	<p>Outcome Measures: Adolescent Dissociative Experiences (ADE) scale  or  Child Dissociative Checklist (CDC)  depending on age</p>	<p>8/114 had incomplete data as the children/caregiver was unwilling to complete certain items.  97% of the study group had a history of some type of abuse and 84% had an abuse history that was considered moderate to severe.  Results from ADE and CDC reported by type of abuse: no abuse (n=27),sexual abuse (n=25), physical abuse (n=18) and sexual &amp; physical abuse (n=44)  ADE No abuse 2.4 SD 4.7 Sexual abuse 3.4 SD 2.6 Physical abuse 2.4 SD 1.8 Sexual &amp; physical abuse 3.7 SD 2.1  Children with sexual abuse reported significantly higher levels of dissociation p&lt;0.01  CDC No abuse 4.7 SD 2.0 Sexual abuse 6.0 SD 4.8 Physical abuse 6.2 SD 6.1 Sexual &amp; physical abuse 10.4 SD 6.9  Children with a history of sexual and physical abuse had higher levels of 'perceived' dissociation( p&lt;0.05 for each)</p>	<p>The authors conclude that the findings of this study suggest a unique relationship between sexual abuse and dissociation.</p>	<p>This study is rated EL=2-  Study is small  All participants recruited from social services including the control group.  This study was supported in part by a grant from the Philanthropic Education Organization</p>
<p>Eisen ML;Qin J;Goodman GS;Davis SL;  2002 Nov USA</p>	<p>Study Type: Case-control series  Evidence level: 2-</p>	<p>To assess children's memory and suggestibility in the context of ongoing maltreatment investigations</p>	<p>n=189 children who were stratified into three age groups: 3-5 years n=51 6-10 years n=64 11-17 years n=31</p>	<p>Children (107 females and 82 males), predominantly low socioeconomic status (77% African American , 9%</p>	<p>Relevant outcome Measures:  Child Dissociative Checklist (CDC)  Children's</p>	<p>Results given for CDC, CPAS and A-DES (measures of dissociation).  Results presented by age : CDC Age 3-5: 9.7 SD5.6 (n=31)</p>	<p>Dissociation findings extracted from text.  There was no significant</p>	<p>This study was rated EL=2-  A case-control study although the controls were also potentially</p>

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
158			<p>they were also divided into three abuse-status groups:</p> <p>1. Abused physical n=101 sexual n=55 both physical and sexual n=22</p> <p>2. Neglected neglected n=27 children with documented evidence of parental addiction but not abuse n=16</p> <p>3. Non-abused control group n=40 who had no history of abuse, neglect or maltreatment but who were referred due to suspicions.</p> <p>For n=5 children there was no abuse status information</p>	<p>Hispanic American, 10% European American, 4% other mean age 7.3 years (range 3-17) years old. Approximately 38% were siblings. Referred to a child abuse evaluation unit in a hospital for a five day physical and psychological assessment</p>	<p>Perceptual Alteration Scale (CPAS)</p> <p>The Adolescent Dissociation Experiences (ADE) Scale</p>	<p>Age 6-10: 6.8 SD6.2 (n=35) Age 11-15: 4.6 SD3.8 (n=14)</p> <p>CPAS Age 3-5: n/a Age 6-10: 54.8 SD 11.4 (n=84) Age 11-15: 51.2 SD9.8 (n=22)</p> <p>ADE Age 3-5:n/a Age 6-10: n/a Age 11-15: 76.5 SD 52.5 (n=41)</p> <p>Results presented by abuse status:</p> <p>CDC Abused 7.6 SD 5.5 (n=40) Neglect 7.9 SD 6.9 (n=18) Control 6.8 SD 5.7 (n=20)</p> <p>CPAS Abused 54.1 SD 10.3 (n=68) Neglect 51.6 SD 11.2 (n=31) Control 55.2 SD 11.8 (n=22)</p> <p>ADE Abused 79.1 SD 51.2 (n=2) Neglect 86.1 SD 63.4(n=9) Control 63.3 SD 50.1(n=9)</p> <p>No statistical analysis given.</p>	<p>association between prior history of abuse in any of the groups with any of the dissociation measures.</p>	<p>abused children.</p> <p>The aim of the study was to assess memory and suggestibility in maltreated children as opposed to finding a relationship between dissociation and abuse.</p> <p>Assessment was undertaken in the artificial (intense) environment.</p> <p>The funding of this study was undeclared</p>

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Collin-Vezina D;Hebert M;  2005 Jan  Canada  159	Study Type: Case-control series  Evidence level: 2-	To assess and contrast dissociation and posttraumatic stress disorder symptoms in a group of sexually abused school aged girls compared with a matched group	n=67 girls with a history of sexual abuse n=67 girls as a comparison	Sexually abused (SA) group French -speaking, Canadian girls referred for evaluation to child protection clinic after alleged sexual episode within the previous six months. (Age mean =9.0 SD 1.4 range 7-12 yrs) 65.6% cases were classed as very serious 68.7% cases involved family or extended family 46.9% abuse was chronic over months or years  Comparison group was girls recruited from three public schools. There were no sexual abuse cases. Mean age 9.2 SD 1.7	Relevant outcome Measures: Child Dissociative Checklist (CDC) in French	CDC scores SA group (no penetration) 7.14 SD 6.36  SA group (penetration) 8.48 SD 5.34  SA group (no intrafamilial abuse) 9.11 SD 7.01  SA group (Intrafamilial abuse) 7.63 SD 5.45  SA group (no chronic abuse) 7.51 SD 6.37  SA group (chronic abuse) 8.67 SD 5.68  Comparative group 3.43 SD 3.95  20/67 of the SA (29.9%) and 3/67 of the comparative group (4.5%) presented with clinical levels of dissociation.  After correcting for confounding factors e.g. parent's education, the odds of presenting with a dissociative tendencies was 8-fold in the SA group compared to the control group	These results indicate that sexual victimisation of school aged girls significantly increases the odds (eight-fold) of presenting with a clinical level of dissociation.  Degree/type of sexual abuse did not prove to be predictive of dissociation symptoms.	This case control study was given an EL=2-  Small study  SA population was narrow (attending clinic).  Comparison group was suitable but although similar on age and socioeconomic factors differed in terms of family structure and parental level of education  The funding of this study was undeclared

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
Atlas JA;Hiott J; 1994 Feb USA 160	Study Type: Case-control series  Evidence level: 2-	To assess the extent of dissociative experience and possible differences in amount of dissociation related to type of abuse (physical , sexual and both)	n=57 of which n=34 girls n=23 boys	Adolescents with a history of abuse admitted into an acute adolescent inpatient unit Age range (11yrs 3mths to 17yrs 8mths) Minority backgrounds (African-American and Hispanic) lower middle-class homes mean IQ 70 (Peabody picture vocabulary test) n=23 physical abuse (PA) n=24 sexual abuse (SA) n=10 physical and sexual abuse (PA & SA)	Outcome Measures: Adolescent Dissociative Experiences Scale (ADE)	Mean ADE score PA = 28.1 SD 25.1 SA = 34.7 SD 31.7 PA & SA= 34.9 SD 22.9  No statistically significant differences between groups but combined group mean of 32 reflects moderate to severe dissociation.  Authors quote a 'control' group from a separate paper of adolescents 13-17 yrs with a variety of diagnoses and abuse backgrounds with a mean ADE 19.2 SD 15.00 Two-tailed t-test between the two group p<0.005	In this study adolescent inpatients with histories of abuse showed moderate to severe dissociation as measured by the ADE scale. Adolescents with a history of sexual abuse had a higher ADE score than physical abuse but this was not statistically significant	This study was rated EL=2-  A small case series with historical control group. This control group was not a 'normal' population  High chance of bias as no confounding factors was considered.  Select population of inpatients  Population was of low socioeconomic class  i.e. these factors affect the external validity of the results  The funding of this study was undeclared
Friedrich WN;Jaworski TM;Huxsahl JE;Bengtson BS; 1997 Apr USA 161	Study Type: Case-controlled series  Evidence level: 2-	Authors hypothesised that longer duration and greater severity , earlier age of onset of abuse and a history physical abuse would result in greater level of dissociative and sexual symptoms	n=350 of which n=75 nonpsychiatric n=165 psychiatric nonabused n=72 psychiatric abused n=38 psychiatric suspected abuse	Four groups of children aged 7-18 years  Non-psychiatric comparative sample recruited from two local mainstream protestant religious education groups mean age 11.96 SD 2.25 56% female  Psychiatric	Relevant outcome measures:  Child completed the Trauma Symptom Checklist -Children (TSC-C)  Parents completed the CDC	Dissociation Subscale total score from the TSC-C  Psychiatric abused 7-11 years 8.83 SD 4.47 12-14 years 9.94 SD 4.86 15-18 years 14.85 SD 6.56  Psychiatric suspected abused 7-11 years 7.57 SD 4.88 12-14 years 9.69 SD 4.87 15-18 years 10.14 SD 6.04  Psychiatric non-abused	The authors concluded that a combination of child and parent related reports were very useful in understanding symptoms of sexually abused children. A correlation was noted between	This study is rated as EL=2-  However the aim of the study was not primarily to demonstrate the association of sexual abuse and dissociation.  The funding of this study was undeclared

DRAFT FOR CONSULTATION

Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
				<p>nonabused recruited from consecutive inpatient admissions to a child psychiatric inpatient unit or consecutive outpatient evaluations (by author) mean age 12.56 SD 2.74 42% female</p> <p>Psychiatric abused children with a documented history of sexual abuse from in and out patients settings mean age 12.05 SD 2.84 80% female</p> <p>Psychiatric suspected abuse children with suspected sexual abuse from in and out patient settings mean age 12.05 SD 2.84 68% female</p>		<p>7-11 years 9.18 SD 6.39 12-14 years 7.96 SD 5.76 15-18 years 10.49 SD 6.51</p> <p>Non-psychiatric 7-11 years 1.11 SD 1.73 12-14 years 1.2 SD 1.58 15-18 years 3.43 SD 3.26</p> <p>'Significant differences' were found between all three clinical groups and the nonpsychiatric group for dissociation but there were no differences between the three clinical groups for dissociation although the psychiatric abused group scored highest.</p> <p>No details of tests were given</p> <p>In a foot note post-hoc analysis using the TSC-C subscale DIS1 Overt Dissociation showed there was a significant difference between the psychiatric abused and psychiatric nonabused for the 12-14 and 15-18 year old groups.</p> <p>CDC reporting was brief. The authors stated that post hoc analysis for the CDC found that the clinical groups differed significantly from the non psychiatric comparison group but not from each other (no further details given)</p>	<p>all the clinical groups and dissociation but no statistical testing was reported</p>	



Bibliographic Information	Study Type & Evidence Level	Aim of Study	Number of Patients	Population Characteristics	Outcome measures	Results & Comments	Study Summary	Reviewer Comment
<p>Nilsson D;Svedin CG; 2006 Sweden 162</p>	<p>Study Type: Retrospective questionnaire validation study  Evidence Level: 2-</p>	<p>To investigate a) the psychometric properties of the Dissociation Questionnaire (DIS-Q) in Swedish in a normative adolescent population and b) dissociative symptoms associated with trauma including sexual and physical abuse.</p>	<p>n=74 adolescents with known experiences of trauma (clinical group)  n=449 normal adolescents (control group)</p>	<p>Clinical group Adolescents who had been patients at a child and adolescent psychiatric clinic. All had a history of sexual and or physical abuse self reported and confirmed by authorities. Mean age 16.03 (range 12-19 )years n=64 girls, n=10 boys  Control group Adolescents recruited from within schools in the same city and of different socioeconomic backgrounds mean age 15.07 SD 1.92 no range given n=210 boys n=190 girls (remainder were dropouts)</p>	<p>DIS-Q in Swedish</p>	<p>Prevalence of dissociation  2.3% in the control group  50% in the clinical group  Chi-square test P&lt;0.001  Total scores of DIS-Q 1.42 SD 0.43 control group 2.52 SD 0.8 clinical group  p&lt;0.001</p>	<p>The prevalence of dissociative symptoms was greater in a group of adolescents with a history of abuse compared with a control group of adolescents</p>	<p>This study was rated EL=2-  This was a retrospective study.  The main aim of it was to investigate the validity of the DIS-Q-Sweden and not to look at the prevalence of dissociation with sexual and physical abuse.  Although they endeavoured to match control and clinical group there was no analysis to detect any differences in confounding factors  There were significant dropouts from the control group.</p>

8 Parent-child interactions  
No literature identified.

# References

---

1. HM Government. Working Together to Safeguard Children. London: The Stationery Office; 2006.
2. National Institute for Health and Clinical Excellence. The guidelines manual 2007. London: NICE; 2007.
3. Royal College of Paediatrics and Child Health. The physical signs of child sexual abuse. An evidence-based review and guidance for best practice. 2008.
4. Woodman J, Pitt M, Wentz R *et al.* Accuracy of screening procedures for non-accidental injury in children: systematic review and modelling of clinical effectiveness. *Health Technology Assessment* . 2008.
5. National Institute for Clinical Excellence. Guideline Development Methods: Information for National Collaborating Centres and Guideline Developers. London: National Institute for Clinical Evidence; 2005.
6. Oxman AD, Sackett DL, and Guyatt GH. Users' guide to the medical literature. I. How to get started. *JAMA: the journal of the American Medical Association* 1993; 270:(17)2093-5.
7. Sackett DL, Straus SE, Richardson WS, Rosenberg W, and Haynes RB. Evidence-based medicine. How to practice and teach EBM. 3rd ed. Edinburgh: Churchill Livingstone; 2005.
8. Scottish Intercollegiate Guidelines Network. A guideline developers' handbook. No. 50. Edinburgh: SIGN; 2001.
9. HM Government. What to do if you're worried a child is being abused. London: Department for Children, Schools and Families; 2006.
10. Maguire S and Mann MK. Are there patterns of bruising in childhood which are diagnostic or suggestive of abuse? A systematic review. *Archives of Disease in Childhood* 2005; 90:(2)182-6.
11. CORE-INFO. Bites review. Is this an abusive bite? Welsh Child Protection Systematic Review Group [online] 2008 [cited 2008 Jul 1]; Available from: URL:[http://www.core-info.cf.ac.uk/oral\\_inj/bites/bites\\_key.htm](http://www.core-info.cf.ac.uk/oral_inj/bites/bites_key.htm)
12. Kemp A, Maguire SA, Sibert J *et al.* Can we identify abusive bites on children? *Archives of Disease in Childhood* 2006; 91:(11)951.
13. Maguire S, Moynihan S, Mann M *et al.* A systematic review of the features that indicate intentional scalds in children. *Burns* 2008; 34:(8)1072-81.
14. CORE-INFO. Thermal Injuries Review. Welsh Child Protection Systematic Review Group [online] 2006 Available from: URL:<http://medweb.uwcm.ac.uk/editcore-info/thermal/index.htm>
15. Kemp A, Dunstan F, Harrison S *et al.* Patterns of skeletal fractures in child abuse: systematic review. *British Medical Journal* 2008; 337:(a1518).
16. Gilks WR, Thomas A, and Spiegelhalter DJ. A language and program for complex Bayesian modelling. *The Statistician* 1994; 43:169-78.
17. Day F, Clegg S, McPhillips M *et al.* A retrospective case series of skeletal surveys in children with suspected non-accidental injury. *Journal of Clinical Forensic Medicine* 2006; 13:(2)55-9.
18. Loder RT. Orthopaedic injuries in children with nonaccidental trauma: demographics and incidence from the 2000 kids' inpatient database. *Journal of Pediatric Orthopedics* 2007; 27:(4)421-6.

## DRAFT FOR CONSULTATION

19. Carty H. Non-accidental injury: a retrospective analysis of a large cohort. *European Radiology* 2002; 12:(12)2919-25.
20. Krishnan J. Patterns of osseous injuries and psychosocial factors affecting victims of child abuse. *Australian and New Zealand Journal of Surgery* 1990; 60:(6)447-50.
21. Hui C, Joughin E, Goldstein S *et al.* Femoral fractures in children younger than three years: the role of nonaccidental injury. *Journal of Pediatric Orthopedics* 2008; 28:(3)297-302.
22. Jayawant S, Rawlinson A, Gibbon F *et al.* Subdural haemorrhages in infants: Population based study. *British Medical Journal* 1998; 317:(7172)1558-61.
23. Hobbs C, Childs AM, Wynne J *et al.* Subdural haematoma and effusion in infancy: An epidemiological study. *Archives of Disease in Childhood* 2005; 90:(9)952-5.
24. Welsh Child Protection Systematic Review Group. Welsh Child Protection Systematic Review Group. Cardiff University [online] 2008 [cited 2008 Mar 11]; Available from: URL:<http://www.core-info.cardiff.ac.uk/>
25. Taylor D, Bonshek R, Brosnahan D *et al.* Child abuse and the eye. *Eye* 1999; 13:(1)3-10.
26. Adams G, Ainsworth J, Butler L *et al.* Update from the ophthalmology child abuse working party: Royal College Ophthalmologists. *Eye* 2004; 18:(8)795-8.
27. Vinchon M, foort-Dhellemmes S, Desurmont M *et al.* Accidental and nonaccidental head injuries in infants: a prospective study. *Journal of Neurosurgery* 2005; 102:(4 Suppl)380-4.
28. DiScala C. Child abuse and unintentional injuries: a 10-year retrospective. *Archives of Pediatrics and Adolescent Medicine* 2000; 154:(1)16-22.
29. Roaten JB, Partrick DA, Nydam TL *et al.* Nonaccidental trauma is a major cause of morbidity and mortality among patients at a regional level 1 pediatric trauma center. *Journal of Pediatric Surgery* 2006; 41:(12)2013-5.
30. Wood J. Distinguishing inflicted versus accidental abdominal injuries in young children. *Journal of Trauma-Injury Infection and Critical Care* 2005; 59:(5)1203-8.
31. Trokel M. Blunt abdominal injury in the young pediatric patient: child abuse and patient outcomes. *Child Maltreatment* 2004; 9:(1)111-7.
32. Maguire S, Hunter B, Hunter L *et al.* Diagnosing abuse: a systematic review of torn frenum and other intra-oral injuries. *Archives of Disease in Childhood* 2007; 92:(12)1113-7.
33. DeLago C, Deblinger E, Schroeder C *et al.* Girls Who Disclose Sexual Abuse: Urogenital Symptoms and Signs After Genital Contact. *Pediatrics* 2008; 122:(2)e281-e286.
34. Klevan JL and De Jong AR. Urinary tract symptoms and urinary tract infection following sexual abuse. *American Journal of Diseases of Children* 1990; 144:(2)242-4.
35. Strathearn L, Gray PH, O'Callaghan F *et al.* Childhood neglect and cognitive development in extremely low birth weight infants: a prospective study. *Pediatrics* 2001; 108:(1)142-51.
36. Wright C, Loughridge J, and Moore G. Failure to thrive in a population context: two contrasting studies of feeding and nutritional status. *Proceedings of the Nutrition Society* 2000; 59:(1)37-45.
37. Whitaker RC, Phillips SM, Orzol SM *et al.* The association between maltreatment and obesity among preschool children. *Child Abuse and Neglect* 2007; 31:(11-12)1187-12.
38. Stockwell MS, Brown J, Chen S *et al.* Is Underimmunization Associated With Child Maltreatment? *Ambulatory Pediatrics* 2008; 8:(3)210-3.

## DRAFT FOR CONSULTATION

39. Chester DL, Jose RM, and Aldlyami E. Non-accidental burns in children--are we neglecting neglect? *Burns* 2006; 32:(2)222-8.
40. Maguire S, Moynihan S, Mann M, Potokar T, and Kemp AM. A systematic review of the features that indicate intentional scalds in children. 1-12-2007. [Unpublished]
41. Taitz LS and King JM. Growth patterns in child abuse. *Acta Paediatrica Scandinavica* 1988; 77:(SUPPL. 343)62-72.
42. Karp RJ, Scholl TO, Decker E *et al.* Growth of abused children. Contrasted with the non-abused in an urban poor community. *Clinical Pediatrics* 1989; 28:(7)317-20.
43. Noll JG, Zeller MH, Trickett PK *et al.* Obesity risk for female victims of childhood sexual abuse: a prospective study. *Pediatrics* 2007; 120:(1)e61-e67.
44. Johnson JG, Cohen P, Kasen S *et al.* Childhood adversities associated with risk for eating disorders or weight problems during adolescence or early adulthood. *American Journal of Psychiatry* 2002; 159:(3)394-400.
45. World Health Organization. Oral Health. World Health Organization [online] 2008 [cited 2008 Mar 11]; Available from: URL:[http://www.who.int/topics/oral\\_health/en/](http://www.who.int/topics/oral_health/en/)
46. Greene P. Child abuse/neglect and the oral health of children's primary dentition. *Military Medicine* 1995; 160:(6)290-3.
47. Greene PE, Chisick MC, and Aaron GR. A comparison of oral health status and need for dental care between abused/neglected children and nonabused/non-neglected children. *Pediatric Dentistry* 1994; 16:(1)41-5.
48. Valencia-Rojas N, Lawrence HP, and Goodman D. Prevalence of early childhood caries in a population of children with history of maltreatment. *Journal of Public Health Dentistry* 2008; 68:(2)94-101.
49. Spivey MI, Schnitzer PG, Kruse RL *et al.* Association of Injury Visits in Children and Child Maltreatment Reports [Epub ahead of print]. *Journal of Emergency Medicine* 2008.
50. Paranothy S, Fone D, Mann M *et al.* The incidence and aetiology of epistaxis in infants: a population-based study. *Archives of Disease in Childhood* . 2008.
51. National Institutes of Health Consensus Development Conference on Infantile Apnea and Home Monitoring, Sept 29 to Oct 1, 1986. *Pediatrics* 1987; 79:(2)292-9.
52. McGovern MC and Smith MBH. Causes of apparent life threatening events in infants: A systematic review. *Archives of Disease in Childhood* 2004; 89:(11)1043-8.
53. Kiechl-Kohlendorfer U, Hof D, Peglow UP *et al.* Epidemiology of apparent life threatening events. *Archives of Disease in Childhood* 2005; 90:(3)297-300.
54. Rahilly PM. The pneumographic and medical investigation of infants suffering apparent life threatening episodes. *Journal of Paediatrics and Child Health* 1991; 27:(6)349-53.
55. Pitetti RD, Maffei F, Chang K *et al.* Prevalence of retinal hemorrhages and child abuse in children who present with an apparent life-threatening event. *Pediatrics* 2002; 110:(3)557-62.
56. Samuels MP, Poets CF, Noyes JP *et al.* Diagnosis and management after life threatening events in infants and young children who received cardiopulmonary resuscitation. *British Medical Journal* 1993; 306:(6876)489-92.
57. Altman RL, Brand DA, Forman S *et al.* Abusive head injury as a cause of apparent life-threatening events in infancy. *Archives of Pediatrics and Adolescent Medicine* 2003; 157:(10)1011-5.
58. Stratton SJ, Taves A, Lewis RJ *et al.* Apparent life-threatening events in infants: High risk in the out-of-hospital environment. *Annals of Emergency Medicine* 2004; 43:(6)711-7.

59. Cote A, Hum C, Brouillette RT *et al.* Frequency and timing of recurrent events in infants using home cardiorespiratory monitors. *Journal of Pediatrics* 1998; 132:(5)783-9.
60. Southall DP, Plunkett MC, Banks MW *et al.* Covert video recordings of life-threatening child abuse: lessons for child protection. *Pediatrics* 1997; 100:(5)735-60.
61. Ward SLD, Keens TG, and Chan LS. Sudden infant death syndrome in infants evaluated by apnea programs in California. *Pediatrics* 1986; 77:(4)451-8.
62. Dine MS and McGovern ME. Intentional poisoning of children - An overlooked category of child abuse: Report of seven cases and review of the literature. *Pediatrics* 1982; 70:(1)32-5.
63. Gillenwater JM. Inflicted submersion in childhood. *Archives of Pediatrics and Adolescent Medicine* 1996; 150:(3)298-303.
64. Sheridan MS. The deceit continues: an updated literature review of Munchausen Syndrome by Proxy. *Child Abuse and Neglect* 2003; 27:(4)431-51.
65. Rosenberg DA. Web of deceit: A literature review of Munchausen syndrome by proxy. *Child Abuse and Neglect* 1987; 11:(4)547-63.
66. Feldman MD and Brown RM. Munchausen by Proxy in an international context. *Child Abuse and Neglect* 2002; 26:(5)509-24.
67. Korpershoek M and Flisher AJ. Diagnosis and management of Munchausen's Syndrome by Proxy. *Journal of Child and Adolescent Mental Health* 2004; 16:(1)1-9.
68. Folks DG. Munchausen's syndrome and other factitious disorders. *Neurologic Clinics* 1995; 13:(2)267-81.
69. de Ridder L. and Hoekstra JH. Manifestations of Munchausen syndrome by proxy in pediatric gastroenterology. *Journal of Pediatric Gastroenterology and Nutrition* 2000; 31:(2)208-11.
70. Awadallah N, Vaughan A, Franco K *et al.* Munchausen by proxy: A case, chart series, and literature review of older victims. *Child Abuse and Neglect* 2005; 29:(8)931-41.
71. Feldman KW, Stout JW, and Inglis AF, Jr. Asthma, allergy, and sinopulmonary disease in pediatric condition falsification. *Child Maltreatment* 2002; 7:(2)125-31.
72. Kitzmann KM, Gaylord NK, Holt AR *et al.* Child witnesses to domestic violence: a meta-analytic review. *Journal of Consulting and Clinical Psychology* 2003; 71:(2)339-52.
73. Wolfe DA, Crooks CV, Lee V *et al.* The effects of children's exposure to domestic violence: a meta-analysis and critique. *Clinical Child and Family Psychology Review* 2003; 6:(3)171-87.
74. Kendall-Tackett KA, Williams LM, and Finkelhor D. Impact of sexual abuse on children: a review and synthesis of recent empirical studies. *Psychological Bulletin* 1993; 113:(1)164-80.
75. Bolger KE and Patterson CJ. Developmental pathways from child maltreatment to peer rejection. *Child Development* 2001; 72:(2)549-68.
76. Evans SE, Davies C, and DiLillo D. Exposure to domestic violence: A meta-analysis of child and adolescent outcomes. *Aggression and Violent Behavior* 2008; 13:(2)131-40.
77. Holmes WC. Sexual abuse of boys: Definition, prevalence, correlates, sequelae, and management. *JAMA: the journal of the American Medical Association* 1998; 280:(21)1855-62.
78. Maas C, Herrenkohl TI, and Sousa C. Review of research on child maltreatment and violence in youth. *Trauma, Violence, and Abuse: A Review Journal* 2008; 9:(1)56-67.
79. Root C, MacKay S, Henderson J *et al.* The link between maltreatment and juvenile firesetting: Correlates and underlying mechanisms. *Child Abuse and Neglect* 2008; 32:(2)161-76.

80. Antao V, Maddocks A, Street E *et al.* A case-control study of somatic and behavioural symptoms in sexually abused boys. *Archives of Disease in Childhood* 1996; 75:(3)237-8.
81. Zolotor A, Kotch J, Dufort V *et al.* School performance in a longitudinal cohort of children at risk of maltreatment. *Maternal and Child Health Journal* 1999; 3:(1)19-27.
82. Rowe E and Eckenrode J. The timing of academic difficulties among maltreated and nonmaltreated children. *Child Abuse and Neglect* 1999; 23:(8)813-32.
83. Eckenrode J, Laird M, and Doris J. School performance and disciplinary problems among abused and neglected children. *Developmental Psychology* 1993; 29:(1)53-62.
84. Quas JA, Goodman GS, and Jones D. Predictors of attributions of self-blame and internalizing behavior problems in sexually abused children. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 2003; 44:(5)723-36.
85. Reyome ND. Teacher ratings of the achievement-related classroom behaviors of maltreated and non-maltreated children. *Psychology in the Schools* 1994; 31:(4)253-60.
86. Finzi R, Har-Even D, and Weizman A. Comparison of ego defenses among physically abused children, neglected, and non-maltreated children. *Comprehensive Psychiatry* 2003; 44:(5)388-95.
87. Wells RD, McCann J, Adams J *et al.* Emotional, behavioral, and physical symptoms reported by parents of sexually abused, nonabused, and allegedly abused prepubescent females. *Child Abuse and Neglect* 1995; 19:(2)155-63.
88. Baer J and Martinez CD. Child maltreatment and insecure attachment: A meta-analysis. *Journal of Reproductive and Infant Psychology* 2006; 24:(3)187-97.
89. van Ijzendoorn MH, Schuengel C, and Bakermans-Kranenburg MJ. Disorganized attachment in early childhood: meta-analysis of precursors, concomitants, and sequelae. *Development and Psychopathology* 1999; 11:(2)225-49.
90. Gore-Felton C, Koopman C, McGarvey E *et al.* Relationships of sexual, physical, and emotional abuse to emotional and behavioral problems among incarcerated adolescents. *Journal of Child Sexual Abuse* 2001; 10:(1)73-88.
91. Glassman LH, Weierich MR, Hooley JM *et al.* Child maltreatment, non-suicidal self-injury, and the mediating role of self-criticism. *Behaviour Research and Therapy* 2007; 45:(10)2483-90.
92. Weierich MR and Nock MK. Posttraumatic stress symptoms mediate the relation between childhood sexual abuse and nonsuicidal self-injury. *Journal of Consulting and Clinical Psychology* 2008; 76:(1)39-44.
93. Martin G, Bergen HA, Richardson AS *et al.* Sexual abuse and suicidality: gender differences in a large community sample of adolescents. *Child Abuse and Neglect* 2004; 28:(5)491-503.
94. Lau JT, Chan KK, Lam PK *et al.* Psychological correlates of physical abuse in Hong Kong Chinese adolescents. *Child Abuse and Neglect* 2003; 27:(1)63-75.
95. Goldstein TR, Birmaher B, Axelson D *et al.* History of suicide attempts in pediatric bipolar disorder: Factors associated with increased risk. *Bipolar Disorders* 2005; 7:(6)525-35.
96. Cohen Y, Spirito A, Sterling C *et al.* Physical and sexual abuse and their relation to psychiatric disorder and suicidal behavior among adolescents who are psychiatrically hospitalized. *Journal of Child Psychology and Psychiatry and Allied Disciplines* 1996; 37:(8)989-93.
97. Noll JG, Horowitz LA, Bonanno GA *et al.* Revictimization and self-harm in females who experienced childhood sexual abuse: results from a prospective study. *Journal of Interpersonal Violence* 2003; 18:(12)1452-71.

98. Kaplan SJ, Pelcovitz D, Salzinger S *et al.* Adolescent physical abuse and risk for suicidal behaviors. *Journal of Interpersonal Violence* 1999; 14:(9)976-88.
99. Lipschitz DS, Winegar RK, Nicolaou AL *et al.* Perceived abuse and neglect as risk factors for suicidal behavior in adolescent inpatients. *Journal of Nervous and Mental Disease* 1999; 187:(1)32-9.
100. Glowinski AL, Bucholz KK, Nelson EC *et al.* Suicide attempts in an adolescent female twin sample. *Journal of the American Academy of Child and Adolescent Psychiatry* 2001; 40:(11)1300-7.
101. Beautrais AL, Joyce PR, and Mulder RT. Risk factors for serious suicide attempts among youths aged 13 through 24 years. *Journal of the American Academy of Child and Adolescent Psychiatry* 1996; 35:(9)1174-82.
102. Tulloch AL, Blizzard L, Hornsby H *et al.* Suicide and self-harm in Tasmanian children and adolescents. *Medical Journal of Australia* 1994; 160:(12)775-80.
103. Bayatpour M, Wells RD, and Holford S. Physical and sexual abuse as predictors of substance use and suicide among pregnant teenagers. *Journal of Adolescent Health* 1992; 13:(2)128-32.
104. Finzi R, Ram A, Shnit D *et al.* Depressive symptoms and suicidality in physically abused children. *American Journal of Orthopsychiatry* 2001; 71:(1)98-107.
105. Shaunesey K, Cohen JL, Plummer B *et al.* Suicidality in hospitalized adolescents: relationship to prior abuse. *American Journal of Orthopsychiatry* 1993; 63:(1)113-9.
106. De Wilde EJ, Kienhorst IC, Diekstra RF *et al.* The relationship between adolescent suicidal behavior and life events in childhood and adolescence. *American Journal of Psychiatry* 1992; 149:(1)45-51.
107. Edwall GE, Hoffmann NG, and Harrison PA. Psychological correlates of sexual abuse in adolescent girls in chemical dependency treatment. *Adolescence* 1989; 24:(94)279-88.
108. Garnefski N, Diekstra RF, and de HP. A population-based survey of the characteristics of high school students with and without a history of suicidal behavior. *Acta Psychiatrica Scandinavica* 1992; 86:(3)189-96.
109. Molnar BE, Shade SB, Kral AH *et al.* Suicidal behavior and sexual/physical abuse among street youth. *Child Abuse and Neglect* 1998; 22:(3)213-22.
110. Thompson R, Briggs E, English DJ *et al.* Suicidal ideation among 8-year-olds who are maltreated and at risk: findings from the LONGSCAN studies. *Child Maltreatment* 2005; 10:(1)26-36.
111. Zoroglu SS, Tuzun U, Sar V *et al.* Suicide attempt and self-mutilation among Turkish high school students in relation with abuse, neglect and dissociation. *Psychiatry and Clinical Neurosciences* 2003; 57:(1)119-26.
112. Martin G. Reported family dynamics, sexual abuse, and suicidal behaviors in community adolescents. *Archives of Suicide Research* 1996; 2:(3)183-95.
113. Grilo CM, Sanislow CA, Fehon DC *et al.* Correlates of suicide risk in adolescent inpatients who report a history of childhood abuse. *Comprehensive Psychiatry* 1999; 40:(6)422-8.
114. Calam R, Horne L, Glasgow D *et al.* Psychological disturbance and child sexual abuse: a follow-up study. *Child Abuse and Neglect* 1998; 22:(9)901-13.
115. Vajda J and Steinbeck K. Factors associated with repeat suicide attempts among adolescents. *Australian and New Zealand Journal of Psychiatry* 2000; 34:(3)437-45.
116. Grossman DC, Milligan BC, and Deyo RA. Risk factors for suicide attempts among Navajo adolescents. *American Journal of Public Health* 1991; 81:(7)870-4.
117. Johnson JG, Cohen P, Gould MS *et al.* Childhood adversities, interpersonal difficulties, and risk for suicide attempts during late adolescence and early adulthood. *Archives of General Psychiatry* 2002; 59:(8)741-9.

## DRAFT FOR CONSULTATION

118. Wright J, Friedrich W, Cinq-Mars C *et al.* Self-destructive and delinquent behaviors of adolescent female victims of child sexual abuse: rates and covariates in clinical and nonclinical samples. *Violence and Victims* 2004; 19:(6)627-43.
119. Borowsky IW, Resnick MD, Ireland M *et al.* Suicide attempts among American Indian and Alaska native youth: Risk and protective factors. *Archives of Pediatrics and Adolescent Medicine* 1999; 153:(6)573-80.
120. Brezo J, Paris J, Tremblay R *et al.* Identifying correlates of suicide attempts in suicidal ideators: A population-based study. *Psychological Medicine* 2007; 37:(11)1551-62.
121. Brent DA, Perper JA, Moritz G *et al.* Familial risk factors for adolescent suicide: a case-control study. *Acta Psychiatrica Scandinavica* 1994; 89:(1)52-8.
122. Berger MY, Gieteling MJ, and Benninga MA. Chronic abdominal pain in children. *British Medical Journal* 2007; 334:(7601)997-1002.
123. Rimsza ME, Berg RA, and Locke. Sexual abuse: Somatic and emotional reactions. *Child Abuse and Neglect* 1988; 12:(2)201-8.
124. National Collaborating Centre for Mental Health. Eating Disorders: Core interventions in the treatment and management of anorexia nervosa, bulimia nervosa and related eating disorders. No. CG9. London: National Institute for Clinical Excellence; 2004.
125. Wonderlich SA, Crosby RD, Mitchell JE *et al.* Relationship of childhood sexual abuse and eating disturbance in children. *Journal of the American Academy of Child and Adolescent Psychiatry* 2000; 39:(10)1277-83.
126. Chandy JM, Blum RW, and Resnick MD. Female adolescents with a history of sexual abuse: Risk outcome and protective factors. *Journal of Interpersonal Violence* 1996; 11:(4)503-18.
127. Hernandez J. The concurrence of eating disorders with histories of child abuse among adolescents. *Journal of Child Sexual Abuse* 1995; 4:(3)73-85.
128. Chandy JM, Blum RW, and Resnick MD. Gender-specific outcomes for sexually abused adolescents. *Child Abuse and Neglect* 1996; 20:(12)1219-31.
129. Perkins DF and Luster T. The relationship between sexual abuse and purging: findings from community-wide surveys of female adolescents. *Child Abuse and Neglect* 1999; 23:(4)371-82.
130. MacGregor R, Pullar A, and Cundall D. Silent at school -- elective mutism and abuse. *Archives of Disease in Childhood* 1994; 70:(6)540-1.
131. Troster H. Prevalence and functions of stereotyped behaviors in nonhandicapped children in residential care. *Journal of Abnormal Child Psychology* 1994; 22:(1)79-97.
132. Butler RJ, Golding J, Northstone K *et al.* Nocturnal enuresis at 7.5 years old: prevalence and analysis of clinical signs. *BJU International* 2005; 96:(3)404-10.
133. Hjalmas K, Arnold T, Bower W *et al.* Nocturnal enuresis: an international evidence based management strategy. *Journal of Urology* 2004; 171:(6 Pt 2)2545-61.
134. Mellon MW, Whiteside SP, and Friedrich WN. The relevance of fecal soiling as an indicator of child sexual abuse: a preliminary analysis. *Journal of Developmental and Behavioral Pediatrics* 2006; 27:(1)25-32.
135. Davies SL, Glaser D, and Kossoff R. Children's sexual play and behavior in pre-school settings: staff's perceptions, reports, and responses. *Child Abuse and Neglect* 2000; 24:(10)1329-43.
136. Lindblad F, Gustafsson P, Larson I *et al.* Preschoolers' sexual behavior at day care centers: An epidemiological study. *Child Abuse and Neglect* 2005; 19:569-77.
137. Friedrich WN, Grambsch P, Damon L *et al.* Child Sexual Behavior Inventory: Normative and clinical comparisons. *Psychological Assessment* 1992; 4:(3)303-11.



## DRAFT FOR CONSULTATION

138. Cosentino CE, Meyer-Bahlburg HF, Alpert JL *et al.* Sexual behavior problems and psychopathology symptoms in sexually abused girls. *Journal of the American Academy of Child and Adolescent Psychiatry* 1995; 34:(8)1033-42.
139. Gordon BN, Schroeder CS, and Abrams JM. Children's knowledge of sexuality: a comparison of sexually abused and nonabused children. *American Journal of Orthopsychiatry* 1990; 60:(2)250-7.
140. Inderbitzen-Pisaruk H, Shawchuck CR, and Hoier TS. Behavioral characteristics of child victims of sexual abuse: A comparison study. *Journal of Clinical Child Psychology* 1992; 21:(1)14-9.
141. Merrick MT, Lutrownik AJ, Everson MD *et al.* Beyond sexual abuse: the impact of other maltreatment experiences on sexualized behaviors. *Child Maltreatment* 2008; 13:(2)122-32.
142. Hall T, Hogben M, Carlton AL *et al.* Attitudes toward using condoms and condom use: difference between sexually abused and nonabused African American female adolescents. *Behavioral Medicine* 2008; 34:(2)45-52.
143. Powers JL, Eckenrode J, and Jaklitsch B. Maltreatment among runaway and homeless youth. *Child Abuse and Neglect* 1990; 14:(1)87-98.
144. Powers JL, Jaklitsch B, and Eckenrode J. Behavioral characteristics of maltreatment among runaway and homeless youth. *Early Child Development and Care* 1989; 42:127-39.
145. Stiffman AR. Physical and sexual abuse in runaway youths. *Child Abuse and Neglect* 1989; 13:(3)417-26.
146. Gary F, Moorhead J, and Warren J. Characteristics of troubled youths in a shelter. *Archives of Psychiatric Nursing* 1996; 10:(1)41-8.
147. Warren JK, Gary FA, and Moorhead J. Runaway youths in a southern community: four critical areas of inquiry. *Journal of Child and Adolescent Psychiatric Nursing* 1997; 10:(2)26-35.
148. Warren JK, Gary F, and Moorhead J. Self-reported experiences of physical and sexual abuse among runaway youths. *Perspectives in Psychiatric Care* 1994; 30:(1)23-8.
149. Thompson SJ, Zittel-Palamara KM, and Maccio EM. Runaway Youth Utilizing Crisis Shelter Services: Predictors of Presenting Problems. *Child and Youth Care Forum* 2004; 33:(6)387-404.
150. Kufeldt K and Nimmo M. Youth on the street: abuse and neglect in the eighties. *Child Abuse and Neglect* 1987; 11:(4)531-43.
151. Feitel B, Margetson N, Chamas J *et al.* Psychosocial background and behavioral and emotional disorders of homeless and runaway youth. *Hospital and Community Psychiatry* 1992; 43:(2)155-9.
152. Noell J, Rohde P, Seeley J *et al.* Childhood sexual abuse, adolescent sexual coercion and sexually transmitted infection acquisition among homeless female adolescents. *Child Abuse and Neglect* 2001; 25:(1)137-48.
153. Tyler KA and Cauce AM. Perpetrators of early physical and sexual abuse among homeless and runaway adolescents. *Child Abuse and Neglect* 2002; 26:(12)1261-74.
154. Janus MD, Archambault FX, Brown SW *et al.* Physical abuse in Canadian runaway adolescents. *Child Abuse and Neglect* 1995; 19:(4)433-47.
155. Lansford JE, Dodge KA, Pettit GS *et al.* A 12-year prospective study of the long-term effects of early child physical maltreatment on psychological, behavioral, and academic problems in adolescence. *Archives of Pediatrics and Adolescent Medicine* 2002; 156:(8)824-30.
156. Macfie J, Cicchetti D, and Toth SL. Dissociation in maltreated versus nonmaltreated preschool-aged children. *Child Abuse and Neglect* 2001; 25:(9)1253-67.
157. Kisiel CL and Lyons JS. Dissociation as a mediator of psychopathology among sexually abused children and adolescents. *American Journal of Psychiatry* 2001; 158:(7)1034-9.

## DRAFT FOR CONSULTATION

158. Eisen ML, Qin J, Goodman GS *et al.* Memory and suggestibility in maltreated children: age, stress arousal, dissociation, and psychopathology. *Journal of Experimental Child Psychology* 2002; 83:(3)167-212.
159. Collin-Vezina D and Hebert M. Comparing dissociation and PTSD in sexually abused school-aged girls. *Journal of Nervous and Mental Disease* 2005; 193:(1)47-52.
160. Atlas JA and Hiott J. Dissociative experience in a group of adolescents with history of abuse. *Perceptual and Motor Skills* 1994; 78:(1)121-2.
161. Friedrich WN, Jaworski TM, Huxsahl JE *et al.* Dissociative and sexual behaviors in children and adolescents with sexual abuse and psychiatric histories. *Journal of Interpersonal Violence* 1997; 12:(2)155-71.
162. Nilsson D and Svedin CG. Evaluation of the Swedish version of Dissociation Questionnaire (DIS-Q), Dis-Q-Sweden, among adolescents. *Journal of Trauma and Dissociation* 2006; 7:(3)65-89.
163. Murphy MK, Black NA, Lamping DL *et al.* Consensus development methods, and their use in clinical guideline development. *Health Technology Assessment* 1998; 2:(3).
164. Fitch K, Bernstein S, Aguilar MS, Burnand B, LaCalle JR, Van het Loo M, McDonnell J, Vader JP, and Kahan JP. The RAND/UCLA Appropriateness Method User's Manual. No. MR-1269-DG-XII/RE. Rand Corporation; 2000.
165. Elwyn G, Greenhalgh T, and Macfarlane F. *Groups: a guide to small group work in healthcare, management, education and research.* Abingdon: Radcliffe Medical Press; 2001.
166. Maguire S, Moynihan S, Mann M, Potokar T, and Kemp AM. A systematic review of the features that indicate intentional scalds in children. 1-12-2007. [Unpublished]
167. Johnson DL. Role of apnea in nonaccidental head injury. *Pediatric Neurosurgery* 1995; 23:(6)305-10.
168. Light MJ and Sheridan MS. Munchausen syndrome by proxy and apnea (MBPA). A survey of apnea programs. *Clinical Pediatrics* 1990; 29:(3)162-8.
169. Herman-Staab B. Antecedents to nonorganic failure-to-thrive. *Pediatric Nursing* 1992; 18:(6)579-90.
170. Dubowitz H, Black M, Harrington D *et al.* A follow-up study of behavior problems associated with child sexual abuse. *Child Abuse and Neglect* 1993; 17:(6)743-54.
171. Slusser MM. Manifestations of sexual abuse in preschool-aged children. *Issues in Mental Health Nursing* 1995; 16:(5)481-91.
172. Craig TK and Hodson S. Homeless youth in London: I. Childhood antecedents and psychiatric disorder. *Psychological Medicine* 1998; 28:(6)1379-88.