

The management of lower urinary tract symptoms in men

Appendices A – H

	<p>APPENDICES</p>
--	-------------------

Contents

APPENDIX A - SCOPE	3
APPENDIX B – DECLARATIONS OF INTEREST	11
APPENDIX C – SEARCH STRATEGIES	25
APPENDIX D – EVIDENCE TABLES	44
APPENDIX E – FOREST PLOTS	514
APPENDIX F - COST-EFFECTIVENESS ANALYSIS	658
APPENDIX G - RECOMMENDATIONS FOR RESEARCH	695
APPENDIX H – INTERNATIONAL PROSTATE SYMPTOMS SCORE (IPSS)	709
BIBLIOGRAPHY	711

Appendix A - Scope

NATIONAL INSTITUTE FOR HEALTH AND CLINICAL EXCELLENCE

SCOPE

1 Guideline title

The management of lower urinary tract symptoms in men

1.1 Short title

Lower urinary tract symptoms in men

2 Background

- a) The National Institute for Health and Clinical Excellence ('NICE' or 'the Institute') has commissioned the National Collaborating Centre for Acute Care to develop a clinical guideline on the management of lower urinary tract symptoms (LUTS) in men for use in the NHS in England and Wales. This follows referral of the topic by the Department of Health (see appendix). The guideline will provide recommendations for good practice that are based on the best available evidence of clinical and cost effectiveness.
- b) The Institute's clinical guidelines support the implementation of National Service Frameworks (NSFs) in those aspects of care for which a Framework has been published. The statements in each NSF reflect the evidence that was used at the time the Framework was prepared. The clinical guidelines and technology appraisals published by the Institute after an NSF has been issued will have the effect of updating the Framework.
- c) NICE clinical guidelines support the role of healthcare professionals in providing care in partnership with patients, taking account of their individual needs and preferences, and ensuring that patients (and their carers and families, where appropriate) can make informed decisions about their care and treatment.

3 Clinical need for the guideline

- a) Lower urinary tract symptoms (LUTS) are a collection of symptoms related to problems with the voiding, storage and post-micturition of urine. They generally arise as a result of abnormalities or inadequate functioning of the prostate, urethra, bladder or sphincters. The pathophysiology of LUTS are diverse. In men, benign prostate enlargement, which is secondary to benign prostatic hyperplasia and causes bladder outlet obstruction, is frequently considered to be the major cause of LUTS. However, many other conditions can cause LUTS, including detrusor muscle weakness or overactivity, prostatitis, urinary tract infection, malignancy and neurological disease. In acknowledgement of the non-specific nature of many male LUTS, this clinical guideline will advise on the effective evidence-based management of male LUTS in general, with a specific focus on LUTS associated with benign prostatic disease (presumed benign prostatic hyperplasia).
- b) LUTS in men are best categorised into voiding, storage or post-micturition symptoms to help define the source of the problem. Voiding symptoms (previously known as obstructive symptoms) include weak or intermittent urinary stream, straining, hesitancy, terminal dribbling and incomplete emptying. Storage symptoms (previously known as irritative symptoms, and currently often considered as a symptom complex known as 'overactive bladder') include urgency, frequency, urgency incontinence and nocturia. The major post-micturition symptom is dribbling, which is common and bothersome. Although LUTS do not usually cause severe illness, they can considerably reduce patients' quality of life, and may point to serious pathology of the urogenital tract.
- c) LUTS are a major burden for the ageing male population. Approximately 30% of men aged 50 and older have moderate to severe LUTS. This is a very large group potentially requiring treatment. Age is an important risk factor for LUTS and the prevalence of LUTS increases as men get older. Other risk factors include hormonal status (presence of androgens), increased size of the prostate gland and bladder decompensation. Ethnicity may also be a risk factor: men of black origin seem to be more likely to need surgery for prostate enlargement than men of white origin. Men of Asian origin seem to be less likely than men of white origin to need surgery.
- d) Because prevalence increases with age, the figure above will continue to rise with increasing life expectancy and the resulting growth of the elderly population. This will place increasing demands on health service resources in the coming years. The past 25 years have seen an increase in the use of pharmacotherapy for LUTS, with a considerable decline in surgical rates. Nevertheless, in England, for the year 2003–2004, there were almost 30,000 endoscopic resections of the male bladder outlet, accounting for more than 138,000 bed days. Although transurethral resection of the prostate is often effective in reducing symptoms in men, it is associated with considerable morbidity and a significant overall annual cost. In addition, a significant

proportion of men (25–30%) do not benefit from prostatectomy and have poor post-surgical outcome with no improvement of symptoms. Some failures can be attributed to poor surgical technique, whereas others may be due to incorrect diagnosis of the cause of LUTS. Therefore, to minimise the number of unnecessary operations, predicting the outcome of transurethral resection of the prostate is important.

- e) The British Association of Urological Surgeons primary care guidelines (2004) include recommendations on management and referral to secondary care. There are no specific recommendations on urodynamic studies. The European Association of Urology guidelines (2004) recommend the routine use of uroflowmetry before prostatectomy, and that pressure-flow studies should be used in certain circumstances (but not routinely). According to expert opinion, most UK clinicians carry out uroflowmetry and, in appropriate patients in secondary care, pressure-flow studies are done before surgical intervention in units with access to the equipment. However, experts agree that there is wide variation in clinical practice in the UK. This is due to individual clinicians' belief in the value of urodynamic studies, and also due to staffing issues and access to the technology. There are many national and international guidelines concerned with the management of men with LUTS; however, these vary in quality.
- f) This NICE clinical guideline will address the variations in practice to allow equitable and appropriate treatment for all affected men. There may be cost savings in defining the appropriate use of suitable investigational modalities and existing pharmacotherapy, and by potentially preventing unnecessary surgical treatment and the costs of failed prostatectomy. However, costs incurred would include the cost of equipment, carrying out the tests and associated staff time. Uncertainty over the effectiveness of urodynamic studies makes it impossible to estimate resource impact.

4 The guideline

- a) The guideline development process is described in detail in two publications that are available from the NICE website (see 'Further information'). 'The guideline development process: an overview for stakeholders, the public and the NHS' describes how organisations can become involved in the development of a guideline. 'The guidelines manual' provides advice on the technical aspects of guideline development.
- b) This document is the scope. It defines exactly what this guideline will (and will not) examine, and what the guideline developers will consider. The scope is based on the referral from the Department of Health (see appendix).
- c) The areas that will be addressed by the guideline are described in the following sections.

4.1 *Population*

4.1.1 **Groups that will be covered**

- a) Adult men (18 years or older) with a clinical working diagnosis of LUTS.
- b) Men who have a higher prevalence of LUTS or may be at higher risk including:
 - older men
 - men who are of black origin.

4.1.2 **Groups that will not be covered**

- a) Women.
- b) Men younger than 18 years.

4.2 **Healthcare setting**

Primary, secondary and tertiary care settings.

4.3 **Clinical management**

- a) The clinical and cost effectiveness, and possibly morbidity, of intervention in the management of LUTS.
- b) Initial diagnostic assessments of LUTS, including:

- digital rectal examination (DRE)
 - symptom scores assessments
 - prostate-specific antigen
 - urinary flow rate
 - post-void residual
 - appropriate use of pressure/flow urodynamics
 - cystoscopy.
- c) Monitoring of chronic LUTS.
- d) Non-pharmacological interventions:
- active observation ('watchful waiting')
 - devices (such as catheters, pads and clamps)
 - lifestyle and behavioural changes (such as diet, bladder retraining and pelvic floor exercises).
- e) Pharmacological interventions as first- and/or second-line treatment:
- 5-alpha reductase inhibitors
 - alpha blockers
 - anticholinergics
 - other pharmacotherapeutic agents (such as phytotherapy and phosphodiesterase inhibitors)
 - combination therapy.
- f) Note that guideline recommendations will normally fall within licensed indications; exceptionally, and only if clearly supported by evidence, use outside a licensed indication may be recommended. The guideline will assume that prescribers will use a drug's summary of product characteristics to inform their decisions for individual patients.
- g) Surgical interventions or minimally invasive alternatives:
- transurethral electrovaporisation of the prostate
 - transurethral radiofrequency needle ablation of the prostate
 - all forms of laser therapy directed at the prostate, including enucleation and vaporisation

- transurethral resection of the prostate, including newer forms of therapy such as bipolar excision
 - transurethral incision of the prostate
 - open prostatectomy.
- h) Combinations of the above interventions.
- i) Condition-specific information, support and communication needs of patients, carers and families with LUTS.
- j) General advice on the appropriate evaluation and management of LUTS in men.
- k) The Guideline Development Group will consider making recommendations on the principal complementary and alternative interventions or approaches to care relevant to male LUTS. This will include phytotherapy.
- l) The Guideline Development Group will take reasonable steps to identify ineffective interventions and approaches to care. If robust and credible recommendations for re-positioning the intervention for optimal use, or changing the approach to care to make more efficient use of resources can be made, they will be clearly stated. If the resources released are substantial, consideration will be given to listing such recommendations in the 'Key priorities for implementation' section of the guideline.

4.4 Status

4.4.1 Scope

This is the final version of the scope.

The NICE has published the following related guidance:

- Urinary incontinence: the management of urinary incontinence in women. NICE clinical guideline 40 (2006)
- Referral guidelines for suspected cancer. NICE clinical guideline 27 (2005)
- Potassium-titanyl-phosphate (KTP) laser vapourisation of the prostate for benign prostatic obstruction. NICE interventional procedure guidance 120 (2005)
- Holmium laser prostatectomy. NICE interventional procedure guidance 17 (2003)
- Transurethral radiofrequency needle ablation of the prostate. NICE interventional procedure guidance 15 (2003)
- Transurethral electrovaporisation of the prostate. NICE interventional procedure guidance 14 (2003).

NICE is in the process of producing the following related guidance:

- Prostate cancer: diagnosis and treatment. NICE clinical guideline (publication expected February 2008).

4.4.2 Guideline

The development of the guideline recommendations will begin on 12 December 2007.

5 Further information

Information on the guideline development process is provided in:

- 'The guideline development process: an overview for stakeholders, the public and the NHS'
- 'The guidelines manual'.

These booklets are available as PDF files from the NICE website (www.nice.org.uk/guidelinesmanual). Information on the progress of the guideline will also be available from the website.

6 Referrals from the Department of Health

The Department of Health asked the Institute:

'To prepare a clinical guideline on the management of benign prostatic hyperplasia.'

'To prepare a guideline on the assessment, investigation, management and onward referral of men with lower urinary tract symptoms (including male incontinence) within primary care.'

Appendix B – Declarations of interests

1.1 Introduction

All members of the GDG and all members of the NCGC-ACC staff were required to make formal declarations of interest at the outset, and these were updated at every subsequent meeting throughout the development process.

1.2 Declarations of interests of the GDG members

1.2.1 Chris Chapple (Chair)

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	CC declared a personal pecuniary interest, his attendance in National and International conferences for BAUS, EAU and AUA. He declared a personal pecuniary interest in private practice. He declared that he knew of no personal family interest. He declared his non-personal pecuniary interest, consultancy and research honoraria up to 6 months ago from Allergan, AMS, Astellas, Novartis, Pfizer and UCB – this was put into the department to provide funding for a researcher. He declared a personal non-pecuniary interest as principal investigator and author on pharmaceutical sponsored papers. He is a member of the committee of the BAUS section of female and functional urology and the Adjunct Secretary General of EAU- responsible for their educational activities. He has written books on the subject of BPH/LUTS. He is editor in chief of the Neurourology and Urodynamics journal (official journal of ICS and SUFU).
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	CC declared a personal pecuniary interest, his attendance in National and International conferences for ICS.
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change

GDG meeting	Declaration of Interests
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change
Eleventh GDG Meeting (23 rd February 2009)	CC declared a non-personal pecuniary interest as a consultant for Astellas, Pfizer, Allergan, Xention, Ono, Recordati and Ranbaxy. He declared a personal non-pecuniary interest that any concerns over his views should be expressed at any stage. He declared that he knew of no personal pecuniary interest or personal family interest, above those declared at the previous meeting.
Twelfth GDG Meeting (25 th March 2009)	No change
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	CC declared a personal non-pecuniary interest; he spoke as invited speaker at Astellas symposium at the British Association of Urological Surgeons meeting. He was a speaker at a symposium provided by the European Association of Urology on behalf of Astellas. He was a speaker at a symposium organised by Allergan at the American urology Association meeting. He declared that he had no personal pecuniary interest, personal family interest or non-personal pecuniary interest above those previously declared.
Actions	None required.

1.2.2 Angela Billington

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	She did not attend this meeting.
Second GDG Meeting (13 th December 2007)	AB declared a personal pecuniary interest, Pfizer education support committee. AB did not declare a personal family interest. AB did not declare a non-personal pecuniary interest. She did not declare a personal non-pecuniary interest.
Third GDG Meeting (17 th March 2008)	She did not attend this meeting.
Fourth GDG Meeting (30 th April 2008)	AB declared a personal pecuniary interest, attended conferences for Pfizer, Coloplast, Rochester Medical and Bard. Faculty for Pfizer sense of leadership conference and CARE program for nurses. She did not declare a personal family interest, non-personal pecuniary interests or personal non-pecuniary interest.
Fifth GDG Meeting (6 th June 2008)	She did not attend this meeting.
Sixth GDG Meeting (14 th July 2008)	AB declared a personal pecuniary interest; she is involved in an educational package for Pfizer and educational symposium for Coloplast. Articles for nursing press on catheters. She had dinner courtesy of Pfizer at the ICI meeting. She did not declare a personal family interest, non-personal pecuniary interest or personal non-pecuniary interest.
Seventh GDG Meeting (8 th September 2008)	No change

GDG meeting	Declaration of Interests
Eighth GDG Meeting (15 th October 2008)	She did not attend this meeting
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	She did not attend this meeting
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	During both the 14 th GDG on the 8 June 2009 and the 15 th GDG on the 29 June 2009, The Chair noted that AB had personal pecuniary interests and required AB to be present in an observatory role during the discussion of the pharmacologic recommendations.

1.2.3 Paul Joachim

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	PJ did not declare a personal pecuniary interest or personal family interest. He declared a non-personal pecuniary interest, trustee of Incontact, a charity that benefits from grants from the industry. He declared a personal non-pecuniary interest, trustee of Incontact (as above) Chair of the patient advisory board. He declared that he has had personal and family experience of symptoms.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	No change
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change

GDG meeting	Declaration of Interests
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	No change
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	PJ declared that his interests have not changed, but he informed the group that 'Incontact' had changed its name to 'The Bladder and Bowel Foundation' in September 2008.
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	None required

1.2.4 Malcolm Lucas

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	He did not attend this meeting
Second GDG Meeting (13 th December 2007)	He did not attend this meeting
Third GDG Meeting (17 th March 2008)	ML declared a personal pecuniary interest; I have received lecture fees from Pfizer, UCB Pharma and Astellas within the last 12 months and sponsorship to attend national and international meetings also from Pfizer, Gynecare and AMS. I am not involved in private practice and I am not now accepting invitations to serve on advisory boards. Any current income from lecturing will be payable to a research fund which pays expenses for research fellow and nurses. He did not declare a personal family interest. He declared a non-personal pecuniary interest, I am Principle local investigator for trials with Astellas, Plethora and Bioxell and Lead investigator for trials with Astra. All income goes to Clinical Research Unit, Swansea NHS Trust. He declared a personal non-pecuniary interest, current chairman of Section of Female and Reconstructive Urology, BAUS.
Fourth GDG Meeting (30 th April 2008)	No change
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	He did not attend this meeting

GDG meeting	Declaration of Interests
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	ML declared a non-personal pecuniary interest of departmental research fund receiving income from the UK Continence Society Conference April 2009. The primary source of income in this conference derives from healthcare companies (pharmaceutical and device manufactures). He declared that he knew of no personal pecuniary interest, personal non-pecuniary interest or personal family interest, above those declared at the previous meeting.
Thirteenth GDG Meeting (1 st May 2009)	ML declared a non-personal pecuniary interest of the clinical research unit receiving research income from Astra tech, Pfizer and Astellas. He declared that he knew of no personal pecuniary interest, personal non-pecuniary interest or personal family interest, above those declared at the previous meeting.
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	He did not attend this meeting.
Actions	None required

1.2.5 Roy Latham

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	RL declared a personal pecuniary interest, he acted as a Lay Member on an Invited Service Review carried out by the Royal College of Physicians (July 07). He received a fee for this. He did not declare a personal family interest or non-personal pecuniary interest. He declared a personal non-pecuniary interest, he is personally affected by BPH/LUTS as a patient and as the relative/friend of affected people.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	No change
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change
Ninth GDG Meeting (27 th November 2008)	No change

GDG meeting	Declaration of Interests
Tenth GDG Meeting (16 th January 2009)	He did not attend this meeting
Eleventh GDG Meeting (23 rd February 2009)	He did not attend this meeting
Twelfth GDG Meeting (25 th March 2009)	No change
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	None required

1.2.6 Thomas Ladds

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	He did not attend meeting
Second GDG Meeting (13 th December 2007)	He did not attend meeting
Third GDG Meeting (17 th March 2008)	TL declared a personal pecuniary interest, regular attendance at national and international conferences. BAUS, BAUN, EAU and AUA. Advisory board member for Bard UK Ltd – January 2008. He did not declare a personal family interest or non-personal pecuniary interest. He declared a personal non-pecuniary interest, member and current president of British Association of Urological Nurses (BAUN). Ex officio member BAUS Council Editorial Board member of International Journal of Urological Nursing and Urology News.
Fourth GDG Meeting (30 th April 2008)	TL declared a personal pecuniary interest, sponsorship to attend EAU from Bayer. Lecture fee from Astra Zenecu Marhcin in 2008.
Fifth GDG Meeting (6 th June 2008)	He did not attend this meeting
Sixth GDG Meeting (14 th July 2008)	TL declared that he knew of no personal pecuniary interest, personal family interest or personal non-pecuniary interest. He declared a non-personal pecuniary interest, lecture fees for Astrazeneca and Coloplast Ltd, which were paid to departmental charitable research fund.
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change
Ninth GDG Meeting (27 th November 2008)	He did not attend this meeting

GDG meeting	Declaration of Interests
Tenth GDG Meeting (16 th January 2009)	TL declared a personal pecuniary interest, that he has notified his NHS employer, Central Manchester University Hospitals NHS Foundation Trust that he wished to terminate his contract with them on 27 th March 2009. He is in the process of setting up a limited company, TL Consulting Ltd, of which he will be the director and sole shareholder; he will be employed there from April 1 2009. TL Consulting Ltd. has entered into a contract with ProstaLund Operations AB of Sweden to supply services, including advising them on clinical issues and potential business activities in the UK and overseas. This contract will be operational from April 1 2009. ProstaLund AB currently develops, manufacture and supply equipment, consumables and software in the field of microwave thermotherapy for BPH. TL Consulting may also negotiate and enter into contracts with other suppliers in urology pharmaceutical and medical technical sectors in the future. He declared that he knew of no non-personal pecuniary interest, personal non-pecuniary interest or personal family interest, above those declared at the previous meeting.
Eleventh GDG Meeting (23 rd February 2009)	TL withdrew from the GDG due to new interests declared in the 10 th GDG meeting.
Actions	None required

1.2.7 James N'Dow

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	JN declared a personal pecuniary interest, principle investigator (PI) on a clinical trial with payment per patient going to the urology department. Involved in private practice. He is a member of BAUS Academic Section. He did not declare a personal family interest. He declared a non-personal pecuniary interest, PI of commissioned research with University of Aberdeen by CYTOSYSTEMS on evaluation of a urinary diagnostic marker for bladder cancer. He declared a personal non-pecuniary interest; he led HTA commissioned research on systematic review of surgical treatments of BPH (in press).
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	JN declared a non-personal pecuniary interest, principle investigator (PI) on a clinical trial with payment per patient going to the urology department.
Fifth GDG Meeting (6 th June 2008)	He did not attend this meeting
Sixth GDG Meeting (14 th July 2008)	He did not attend this meeting
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	He did not attend this meeting
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change

GDG meeting	Declaration of Interests
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	He did not attend this meeting
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	None required

1.2.8 Jon Rees

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	JR declared a personal pecuniary interest, involved in private urological practice. He declared that he knew of no personal family interest, non-personal pecuniary interest or personal non-pecuniary interest.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	No change
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	He did not attend this meeting
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	He did not attend this meeting
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	He did not attend this meeting

GDG meeting	Declaration of Interests
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	None required

1.2.9 Mark Speakman

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	MS declared a personal pecuniary interest, he is involved in giving lectures for drug companies at national and international meetings in last 12 months (Astellas, GSK, Boehringer Ingelheim, Pfizer). No new consulting work and new projects declined for duration of guideline. Involved in private practice. He did not declare a personal family interest. He declared a non-personal pecuniary interest, investigator in BPH trials (Astellas, Bayer, GSK, Pfizer, MSD, Allergan). None in last 12 months (sponsorship). Previous research sponsorship from Yamanouchi and MSD in last 5 years. He declared a personal non-pecuniary interest, his clear opinion - author of BAUS BPH Guideline 2004. Author of a number of peer-reviewed LUTS/BPH papers.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	MS declared a personal non-pecuniary interest, he is a member of the editorial board for European Urology.
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	MS declared a personal pecuniary interest, single lecture (debate) on anticholinergics for Astellas. He declared that he knew of no personal family interest, non-personal pecuniary interest or personal non-pecuniary interest, above those declared at the previous meeting.
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	He did not attend this meeting
Eleventh GDG Meeting (23 rd February 2009)	No change

GDG meeting	Declaration of Interests
Twelfth GDG Meeting (25 th March 2009)	MS declared a non-personal pecuniary interest of future research studies planned with Allergan and GSK. He declared a personal non-pecuniary interest as national investigator for new LUTS/BPH Registry for the European Association of Urology. He declared that he knew of no personal pecuniary interest or personal family interest, above those declared at the previous meeting.
Thirteenth GDG Meeting (1 st May 2009)	He did not attend this meeting.
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	MS declared a non-personal pecuniary interest, new supported research studies with Allergan, Astellas and GSK. He declared participation in EAU LUTS/BPH database. He declared that he knew of no personal pecuniary interest, personal family interest or personal non-pecuniary interest, above those declared at the previous meeting.
Actions	None required

1.2.10 Julian Spinks

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	JS declared a personal pecuniary interest, he is a member of advisory boards on LUTS and received honoraria from Boehringer Ingelheim (March 07). He has attended advisory boards on Restless legs syndrome organised by RLS UK with payment from Boehringer Ingelheim. He has been paid for attendance at a focus group on faecal incontinence by Continence UK (Nov 07). He has been paid to speak and chair meetings by Astellas, BMS and ALK. He is a paid member of the editorial boards of Continence UK. He has received payment for attending focus meetings on child growth hormone. He did not declare a personal family interest of non-personal pecuniary interest. He declared a personal non-pecuniary interest, member of the strategy board of Incontact, Chairman of the local division of the BMA and board member of RLS UK.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	JS declared a personal pecuniary interest, I have received sponsorship to attend the EAU congress in Milan from Pfizer. I have received speaker fees to speak at a conference from Pfizer on GPs and OAB. He is a member of advisory boards on LUTS and received honoraria from Boehringer Ingelheim (March 07). He has attended advisory boards on Restless legs syndrome organised by RLS UK with payment from Boehringer Ingelheim. He has been paid for attendance at a focus group on faecal incontinence by Continence UK (Nov 07). He has been paid to speak and chair meetings by Astellas, BMS and ALK. He is a paid member of the editorial boards of Continence UK. He has received payment of attending focus meetings on child growth hormone. He did not declare a personal family interest of non-personal pecuniary interest. He declared a personal non-pecuniary interest, member of the strategy board of Incontact, Chairman of the local division of the BMA and board member of RLS UK.
Fifth GDG Meeting (6 th June 2008)	No change
Sixth GDG Meeting (14 th July 2008)	No change

GDG meeting	Declaration of Interests
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	JS declared a personal non-pecuniary interest, he attended a planning meeting for the “Sense of Leadership” organised by Pfizer. He declared that he knew of no personal pecuniary interest, personal family interest or non-personal pecuniary interest, above those declared at the previous meeting.
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	No change
Thirteenth GDG Meeting (1 st May 2009)	JS declared that he had no current personal pecuniary interests. He declared that he knew of no non-personal family interest, personal non-pecuniary interest or personal family interest, above those declared at the previous meeting.
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	No change
Actions	During the 12 th GDG on the 25 th March 2009, JS was only present as an observer for the presentations on medical interventions and did not participate in discussion due to previously declared interest.

1.2.11 William Turner

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	WT declared a personal pecuniary interest, private practice in urology. He did not declare a personal family interest. He declared a non-personal pecuniary interest, he is the principal local investigator in clinical trials with Allergan (not yet opened), Dianippo Sumuto, Yamanouchi (now Astellas), Schwarz Pharma. He is the principal local investigator in clinical trial with Novartis 2005-6. He declared a personal non-pecuniary interest, executive committee member section of female and reconstructive urology, British Association of Urological Surgeons. Author of papers, chapters and books on urology. Member of NICE Topic Selection Panel and Technology Appraisal Committee.
Second GDG Meeting (13 th December 2007)	No change
Third GDG Meeting (17 th March 2008)	No change
Fourth GDG Meeting (30 th April 2008)	No change
Fifth GDG Meeting (6 th June 2008)	No change

GDG meeting	Declaration of Interests
Sixth GDG Meeting (14 th July 2008)	No change
Seventh GDG Meeting (8 th September 2008)	No change
Eighth GDG Meeting (15 th October 2008)	No change
Ninth GDG Meeting (27 th November 2008)	No change
Tenth GDG Meeting (16 th January 2009)	No change
Eleventh GDG Meeting (23 rd February 2009)	No change
Twelfth GDG Meeting (25 th March 2009)	No change
Thirteenth GDG Meeting (1 st May 2009)	No change
Fourteenth GDG Meeting (8 th June 2009)	No change
Fifteenth GDG Meeting (29 th June 2009)	He declared a non-personal pecuniary interest; he stated that his participation in the clinical trial with Allergan never materialised. He declared that he knew of no personal pecuniary interest, personal family interest or personal non-pecuniary interest above those declared at the previous meeting.
Actions	None required.

1.2.12 Adrian Wagg

GDG meeting	Declaration of Interests
First GDG meeting (12 th December 2007)	He did not attend this meeting
Second GDG Meeting (13 th December 2007)	AW declared a personal pecuniary interest, Astellas pharmaceutical – consultant. Pfizer – occasional consultant. He did not declare a personal family interest. He declared a non-personal pecuniary interest, fees for lectures/writing to research healthcare commission – research fund for Pfizer, Astellas, UCB. He declared a personal non-pecuniary interest, Chairman of trustees of the Continence Foundation and Vice Chairman trustees of Incontact. Researcher for Astellas. Plethora, Boehringer Ingelheim –Lilly. Associate Director CEEU, Royal College of Physicians. He is the National leader for audit of the Continence care.
Third GDG Meeting (17 th March 2008)	He declared a non-personal pecuniary interest, he declared a Pfizer research study, European CI and UK PI.
Fourth GDG Meeting (30 th April 2008)	AW declared a personal pecuniary interest, Astellas pharmaceutical – consultant. Pfizer – occasional consultant. Pfizer pharmaceutical advisory board. Sense of leadership course for Pfizer. SCA conference. Lecture fees from Astellas and telephone symposium on LUTS on geriatric medicine. He did not declare a personal family interest. He declared a non-personal pecuniary interest, fees for lectures/writing to research healthcare commission – research fund for Pfizer, Astellas, and UCB. Pfizer research study, European C.I. and UK principal investigator. BUPA grant for research £13K. Sponsorship to EAU by Astellas. He

GDG meeting	Declaration of Interests
	declared a personal non-pecuniary interest, Vice-chairman of the Continence Foundation and Incontact (merged). Researcher for Astellas. Plethora, Boehringer Ingelheim –Lilly. Associate Director CEEU, Royal College of Physicians. He is the National leader for audit of the Continence care. Papers for Pharma funded studies.
Fifth GDG Meeting (6th June 2008)	AW declared a personal pecuniary interest, since last declaration, speaker for Pfizer at launch meeting for Fesoterodine. Astellas pharmaceutical – consultant. Pfizer – occasional consultant. Pfizer pharmaceutical advisory board. Sense of leadership course for Pfizer. SCA conference. Lecture fees from Astellas and telephone symposium on LUTS on geriatric medicine. He did not declare a personal family interest. He declared a non-personal pecuniary interest, fees for lectures/writing to research healthcare commission – research fund for Pfizer, Astellas, and UCB. Pfizer research study, European C.I. and UK principal investigator. BUPA grant for research £13K. Sponsorship to EAU by Astellas. He declared a personal non-pecuniary interest, Vice-chairman of the Continence Foundation and Incontact (merged). Researcher for Astellas. Plethora, Boehringer Ingelheim –Lilly. Associate Director CEEU, Royal College of Physicians. He is the National leader for audit of the Continence care. Papers for Pharma funded studies.
Sixth GDG Meeting (14th July 2008)	AW declared a non-personal pecuniary interest, Chairman of Bladder Master class for Astellas Pharma. He declared a personal non-pecuniary interest; he had dinner courtesy of Pfizer at the ICI meeting in Paris and BAUS. He declared that he knew of no personal pecuniary interest or personal family interest, above those declared at the previous meeting.
Seventh GDG Meeting (8th September 2008)	No change
Eighth GDG Meeting (15th October 2008)	No change
Ninth GDG Meeting (27th November 2008)	No change
Tenth GDG Meeting (16th January 2009)	AW declared a non personal pecuniary interest, donation to fellows research fund from Astellas. He declared that he knew of no personal pecuniary interest, personal family interest or personal non-pecuniary interest, above those declared at the previous meeting.
Eleventh GDG Meeting (23rd February 2009)	He did not attend this meeting
Twelfth GDG Meeting (25th March 2009)	He did not attend this meeting
Thirteenth GDG Meeting (1st May 2009)	No change
Fourteenth GDG Meeting (8th June 2009)	AW declared a personal pecuniary interest and had received fees for a talk from Glaxo, he did not declare a personal family interest. He declared a non-personal pecuniary interest for research from Pfizer. He declared a personal non-pecuniary interest that a donation from Astellas for filming.
Fifteenth GDG Meeting (29th June 2009)	AW declared a non-personal pecuniary interest, Pfizer talk at BAUS – payment into the department. He declared that he had no personal pecuniary interest, personal family interest or personal non-pecuniary interest above those previously declared.
Actions	During both the 14 th GDG on the 8 June 2009 and the 15 th GDG on the 29 June 2009, The Chair noted that AW had personal pecuniary interests and required AW to be present in an observatory role during the discussion of the pharmacologic recommendations.

1.3 Personal pecuniary interests

ML, MS and CC personal pecuniary interests that were deemed significant conflicts of interest had expired before medical intervention recommendations were discussed in the 10th GDG meeting on the 16th January 2009. Further details of the GDG meetings can be found in the minutes on the [NICE website](#).

Appendix C – Search Strategies

Overview of Search Strategies

2.1 Search Strategies

Searches were constructed by using the following groups of terms. These groups are expanded in full in Section 1.2 below.

All searches were run in Medline, Embase and Cochrane Library. Additionally Cinahl and PsychINFO were searched where this was deemed appropriate. Economic searches were conducted in Medline, Embase, NHS EED and the HTA (Health Technology Reports) database from the Cochrane Library. Additionally in HEED (Health Economic Evaluations Database).

Medications search

BPH/LUTS terms
AND
Medication terms
AND
RCT filter or systematic review filter
NOT
Animal/publication filter

Surgery search

BPH/LUTS terms
AND
Surgery terms
AND
RCT filter or systematic review filter
NOT
Animal/publication filter

Laser search

BPH/LUTS terms
AND
Laser terms
AND
RCT filter or systematic review filter
NOT
Animal/publication filter

Conservative treatment search

BPH/LUTS terms
AND
Conservative treatment terms
AND
RCT filter or systematic review filter
NOT
Animal/publication filter

Diagnosis search

BPH/LUTS terms
AND
Diagnosis terms
NOT
Animal/publication filter

Monitoring search

BPH/LUTS terms
AND
Monitoring terms
NOT
Animal/publication filter

Economic searches (Medline and Embase)

BPH/LUTS terms
AND
Economic filter
NOT
Animal/publication filter

Economic searches (NHS EED and HEED)

BPH/LUTS terms

Patient education search

BPH/LUTS terms
AND
Patient education terms
NOT
Animal/publication filter

Patient views search

BPH/LUTS terms
AND
Patient view terms

2.2 Search terms

Animal/publication filter

Animal/publication filter - OVID Embase

- 1 Case-Study/ or Abstract-Report/ or Letter/ or (case adj report).tw. or ((exp Animal/ or Nonhuman/ or exp Animal-Experiment/) not exp Human/)

Animal/publication filter - OVID Medline

- 1 (Case-Reports NOT Randomized-Controlled-Trial OR Letter OR Historical-Article OR Review-Of-Reported-Cases).PT. OR (exp Animals/ NOT Humans/)

Benign Prostatic Hyperplasia (BPH) / Lower Urinary Tract Infection (LUTS) Terms

BPH/LUTS terms – Cochrane Library

- 1 MeSH descriptor Prostatic Hyperplasia, this term only
- 2 (Benign prostat* disease or prostatism or benign prostat* hyperplasia or benign prostat* enlargement or prostat* hypertrophy or prostat* obstruct* or enlarged prostate):ti,ab
- 3 (Lower urinary tract symptom* or urinary symptom* or LUTS or irritable bladder syndrome):ti,ab
- 4 MeSH descriptor Urinary Retention, this term only
- 5 (Bladder obstruct* or incomplete bladder emptying or impaired bladder emptying or storage symptom* or (retention adj5 (chronic or urinary or acute)) or residual urine):ti,ab
- 6 MeSH descriptor Urinary Bladder, Overactive, this term only
- 7 MeSH descriptor Urinary Incontinence, this term only
- 8 MeSH descriptor Enuresis explode all trees
- 9 ((micturition or urin* or bladder or voiding) near (disorder or dysfunction or symptom* or urgency or incontinen*)):ti,ab
- 10 (post micturition dribble or enuresis or nocturia or pollakisuria or weak bladder or overactive bladder or bedwetting):ti,ab
- 11 (haematuria or hematuria):ti,ab
- 12 male or man or men
- 13 ((#3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11) AND #12)
- 14 #1 OR #2 OR #13

BPH/LUTS terms - OVID Embase

- 1 Prostate Hypertrophy/
- 2 (Benign prostat\$ disease or prostatism or benign prostat\$ hyperplasia or benign prostat\$ enlargement or prostat\$ hypertrophy or prostat\$ obstruct\$ or enlarged prostate).tw.
- 3 (Lower urinary tract symptom\$ or urinary symptom\$ or LUTS or irritable bladder syndrome).tw.
- 4 exp Micturition Disorder/
- 5 (Bladder obstruct\$ or incomplete bladder emptying or impaired bladder emptying or storage symptom\$ or (retention adj5 (chronic or urinary or acute)) or residual urine).tw.
- 6 Urinary Frequency/
- 7 ((micturition or urin\$ or bladder or voiding) adj2 (disorder or dysfunction or symptom\$ or urgency or incontinen\$)).tw.
- 8 (post micturition dribble or enuresis or nocturia or pollakisuria or weak bladder or overactive bladder or bedwetting).tw.

- 9 (haematuria or hematuria).tw.
- 10 (male or man or men).mp.
- 11 ((or/3-9) and 10)
- 12 1 or 2 or 11

BPH/LUTS terms - OVID Medline

- 1 prostatic hyperplasia/
- 2 (Benign prostat\$ disease or prostatism or benign prostat\$ hyperplasia or benign prostat\$ enlargement or prostat\$ hypertrophy or prostat\$ enlargement or enlarged prostate).tw.
- 3 (Lower urinary tract symptom\$ or urinary symptom\$ or LUTS or irritable bladder syndrome).tw.
- 4 urinary retention/
- 5 (Bladder obstruct\$ or incomplete bladder emptying or impaired bladder emptying or storage symptom\$ or (retention adj5 (chronic or urinary or acute)) or residual urine).tw.
- 6 urinary bladder, overactive/ or urinary incontinence/ or exp enuresis/
- 7 ((micturition or urin\$ or bladder or voiding) adj2 (disorder or dysfunction or symptom\$ or urgency or incontinen\$)).tw.
- 8 (post micturition dribble or enuresis or nocturia or pollakisuria or weak bladder or overactive bladder or bedwetting).tw.
- 9 (haematuria or hematuria).tw.
- 10 (male or man or men).mp.
- 11 ((or/3-9) and 10)
- 12 1 or 2 or 11

Conservative

Conservative terms – Cochrane Library

- 1 (conservative next (management or treatment* or therap*))
- 2 MeSH descriptor Pelvic Floor, this term only
- 3 MeSH descriptor Exercise Therapy, this term only
- 4 ((Pelvic floor or pelvic muscle) next (exercise or training))
- 5 MeSH descriptor Behavior Therapy, this term only
- 6 (bladder next (training or education or exercise*))
- 7 Post void milking or post-void milking
- 8 MeSH descriptor Drinking Behavior, this term only
- 9 MeSH descriptor Drinking, this term only
- 10 MeSH descriptor Beverages, this term only
- 11 (Fluid* or water) near (consumption or intake)
- 12 MeSH descriptor Caffeine, this term only
- 13 MeSH descriptor Sweetening Agents, this term only
- 14 MeSH descriptor Carbonated Beverages, this term only
- 15 alcohol* or caffeine or tea or coffee or artificial sweetener* or carbonated drink* or fizzy drink* or beverage*
- 16 MeSH descriptor Catheterization, this term only
- 17 MeSH descriptor Catheters, Indwelling, this term only
- 18 MeSH descriptor Absorbent Pads, this term only
- 19 MeSH descriptor Incontinence Pads, this term only
- 20 Catheter*
- 21 Sheath* or penile clamp*

- 22 (Absorbent or incontinence or continence or protective or bed) near (pad* or pants or product*)
 23 (bed or seat or chair) near (protection or pad* or sheet*)
 24 MeSH descriptor Biofeedback (Psychology), this term only
 25 (biofeedback or bio feedback or bio-feedback)
 26 MeSH descriptor Electric Stimulation, this term only
 27 Electric stimulation
 28 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27

Conservative terms - OVID Embase

- 1 (conservative adj (management or treatment\$ or therap\$)).tw.
 2 Pelvic floor muscle training/
 3 ((Pelvic floor or pelvic muscle) adj (exercise or training)).tw.
 4 Bladder training/
 5 (bladder adj (training or education or exercise\$)).tw.
 6 (Post void milking or post-void milking).tw.
 7 Fluid intake/ or exp beverage/ or drinking behavior/
 8 ((Fluid\$ or water) adj (consumption or intake)).tw.
 9 Alcohol consumption/ or caffeine/ or sweetening agent/ or carbonated beverage/
 10 (alcohol\$ or caffeine or tea or coffee or artificial sweetener\$ or carbonated drink\$ or fizzy drink\$ or beverage\$).tw.
 11 Catheter/
 12 Catheter\$.tw.
 13 (Sheath\$ or penile clamp\$).tw.
 14 ((Absorbent or incontinence or continence or protective or bed) adj (pad\$ or pants or product\$)).tw.
 15 ((bed or seat or chair) adj2 (protection or pad\$ or sheet\$)).tw.
 16 Feedback system/
 17 (Biofeedback or bio feedback or bio-feedback).tw.
 18 Electrostimulation/
 19 Electrical stimulation.tw
 20 or/1-19

Conservative terms - OVID Medline

- 1 (conservative adj (management or treatment\$ or therap\$)).tw.
 2 Pelvic floor/ or exercise therapy/
 3 ((Pelvic floor or pelvic muscle) adj (exercise or training)).tw.
 4 behavior therapy/
 5 (bladder adj (training or education or exercise\$)).tw.
 6 (Post void milking or post-void milking).tw.
 7 Drinking behavior/ or Drinking/ or Beverages/
 8 ((Fluid\$ or water) adj (consumption or intake)).tw.
 9 Caffeine/ or sweetening agents/ or carbonated beverages/
 10 (alcohol\$ or caffeine or tea or coffee or artificial sweetener\$ or carbonated drink\$ or fizzy drink\$ or beverage\$).tw.
 11 Catheterization/ or catheters, indwelling/ or absorbent pads/ or incontinence pads/
 12 Catheter\$.tw.
 13 (Sheath\$ or penile clamp\$).tw.

- 14 ((Absorbent or incontinence or continence or protective or bed) adj (pad\$ or pants or product\$)).tw.
 15 ((bed or seat or chair) adj2 (protection or pad\$ or sheet\$)).tw.
 16 "Biofeedback (Psychology) /"
 17 (biofeedback or bio feedback or bio-feedback).tw
 18 Electric stimulation/
 19 Electrical stimulation.tw.
 20 or/1-19

Diagnosis

Diagnosis terms - Central

- 1 (IPSS or I-PSS or (symptom near score))
 2 ((American Urological Association or AUA*) near (symptom or score or index or questionnaire)).tw.
 3 MeSH descriptor Urinalysis, this term only
 4 MeSH descriptor Kidney Function Tests explode all trees
 5 kidney function test* or renal function test* or serum creatinine or eGFR or urea or serum biochemistry or blood test* or dipstick test* or urine analys* or urinalys*
 6 MeSH descriptor Digital Rectal Examination, this term only
 7 rectal exam*
 8 MeSH descriptor Prostate-Specific Antigen, this term only
 9 (prostate specific antigen or PSA) and (test* or assess*)
 10 MeSH descriptor Urodynamics, this term only
 11 urinary flow rate* or urodynamics or pressure flow studies or post void residual measurement* or uroflowmetry
 12 (Frequency volume chart* or ((bladder or volume or void* or urine or urinary or incontinence) adj (diar* or record*)))
 13 MeSH descriptor Cystoscopy, this term only
 14 Cystoscopy or cystometry or cystourethroscopy or videocystogram or cystometrogram
 15 MeSH descriptor Ultrasonography, this term only
 16 ultrasound or non-invasive test*
 17 pad test*
 18 MeSH descriptor X-Rays, this term only
 19 abdominal x-ray*
 20 KUB
 21 MeSH descriptor Urography, this term only
 22 IVU or IVP
 23 (intravenous or intra-venous) near (urogram* or pyelogram* or urography)
 24 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23

Diagnosis terms - OVID Embase

- 1 international prostate symptom score/
 2 (IPSS or I-PSS or (symptom adj3 score)).tw.
 3 ((American Urological Association or \$AUA\$) adj3 (symptom or score or index or questionnaire)).tw.
 4 urinalysis/ or kidney function test/
 5 (kidney function test\$ or renal function test\$ or serum creatinine or eGFR or urea or serum biochemistry or blood test\$ or dipstick test\$ or urine analys\$ or urinalys\$).tw.
 6 digital rectal examination/

7 rectal exam\$.tw.
 8 Prostate Specific Antigen/
 9 ((prostate specific antigen or PSA) and (test\$ or assess\$)).tw.
 10 urodynamics/
 11 (urinary flow rate\$ or urodynamics or pressure flow studies or post void residual
 measurement\$ or uroflowmetry).tw.
 12 (Frequency volume chart\$ or ((bladder or volume or void\$ or urine or urinary or
 incontinence) adj (diar\$ or record\$))).tw.
 13 cystoscopy/ or urethrocytometry/
 14 (Cystoscopy or cystometry or cystourethroscopy or videocystogram or
 cystometrogram).tw.
 15 (ultrasound or ultrasonography or non-invasive test\$).tw.
 16 pad test\$.tw.
 17 X Ray/
 18 abdominal x-ray\$.tw.
 19 KUB.tw.
 20 Intravenous Urography/ or Intravenous Pyelography/
 21 (IVU or IVP).tw.
 22 ((intravenous or intra-venous) adj (urogram\$ or pyelogram\$ or urography)).tw.
 23 or/1-22

Diagnosis terms - OVID Medline

1 (IPSS or I-PSS or (symptom adj3 score)).tw.
 2 ((American Urological Association or \$AUA\$) adj3 (symptom or score or index or
 questionnaire)).tw.
 3 urinalysis/ or exp kidney function tests/
 4 (kidney function test\$ or renal function test\$ or serum creatinine or eGFR or urea or
 serum biochemistry or blood test\$ or dipstick test\$ or urine analys\$ or urinalys\$).tw.
 5 digital rectal examination/
 6 rectal exam\$.tw.
 7 prostate specific antigen/
 8 ((prostate specific antigen or PSA) and (test\$ or assess\$)).tw.
 9 urodynamics/
 10 (urinary flow rate\$ or urodynamics or pressure flow studies or post void residual
 measurement\$ or uroflowmetry).tw.
 11 (Frequency volume chart\$ or ((bladder or volume or void\$ or urine or urinary or
 incontinence) adj (diar\$ or record\$))).tw.
 12 cystoscopy/
 13 (Cystoscopy or cystometry or cystourethroscopy or videocystogram or
 cystometrogram).tw.
 14 ultrasonography/
 15 (ultrasound or non-invasive test\$).tw.
 16 pad test\$.tw.
 17 X-Rays/
 18 abdominal x-ray\$.tw.
 19 KUB.tw.
 20 Urography/
 21 (IVU or IVP).tw.
 22 ((intravenous or intra-venous) adj (urogram\$ or pyelogram\$ or urography)).tw.
 23 or/1-22

Economic**Economic filter - OVID Embase**

- 1 exp economic aspect/
- 2 cost\$.tw.
- 3 (price\$ or pricing\$).tw.
- 4 (fee or fees).tw.
- 5 (financial or finance or finances or financed).tw.
- 6 (value adj2 (money or monetary)).tw.
- 7 resourc\$ allocat\$.tw.
- 8 expenditure\$.tw.
- 9 (fund or funds or funding or fundings or funded).tw.
- 10 (ration or rations or rationing or rations or rationed).tw.
- 11 (saving or savings).tw.
- 12 or/1-11
- 13 Quality of Life/
- 14 quality of life.tw.
- 15 life quality.tw.
- 16 quality adjusted life.tw.
- 17 (qaly\$ or qald\$ or qale\$ or qtime\$).tw.
- 18 disability adjusted life.tw.
- 19 daly\$.tw.
- 20 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty six).tw.
- 21 (sf6 or sf 6 or short form 6 or shortform 6 or sf six or sfsix or shortform six or short form six).tw.
- 22 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform twelve or short form twelve).tw.
- 23 (sf16 or sf 16 or short form 16 or shortform 16 or sf sixteen or sfsixteen or shortform sixteen or short form sixteen).tw.
- 24 (sf20 or sf 20 or short form 20 or shortform 20 or sf twenty or sftwenty or shortform twenty or short form twenty).tw.
- 25 (euroqol or euro qol or eq5d or eq 5d).tw.
- 26 (hql or hqol or h qol or hrqol or hr qol).tw.
- 27 (hye or hyes).tw.
- 28 health\$ equivalent\$ year\$.tw.
- 29 (hui or hui1 or hui2 or hui3).tw.
- 30 health utilit\$.tw.
- 31 disutilit\$.tw.
- 32 rosser.tw.
- 33 (quality of wellbeing or quality of well being).tw.
- 34 qwb.tw.
- 35 willingness to pay.tw.
- 36 standard gamble\$.tw.
- 37 time trade off.tw.
- 38 time tradeoff.tw.
- 39 tto.tw.
- 40 factor analy\$.tw.
- 41 preference based.tw.
- 42 (state adj2 valu\$).tw.
- 43 Life Expectancy/
- 44 life expectancy\$.tw.
- 45 ((duration or length or period of time or lasting or last or lasted) adj4 symptom\$).tw.

46 or/13-46
 47 exp model/
 48 exp Mathematical Model/
 49 markov\$.tw.
 50 Monte Carlo Method/
 51 monte carlo.tw.
 52 exp Decision Theory/
 53 (decision\$ adj2 (tree\$ or anlay\$ or model\$)).tw.
 54 model\$.tw.
 55 or/47-55
 56 12 or 46 or 55

Economic filter - OVID Medline

1 exp "Costs and Cost Analysis"/
 2 Economics/
 3 Economics, Nursing/ or Economics, Medical/ or Economics, Hospital/ or Economics,
 Pharmaceutical/
 4 exp "Fees and Charges"/
 5 exp Budgets/
 6 budget\$.tw.
 7 cost\$.ti.
 8 (cost\$ adj2 (effective\$ or utilit\$ or benefit\$ or minimi\$)).ab.
 9 (economic\$ or pharmacoeconomic\$ or pharmaco-economic\$).ti.
 10 (price\$ or pricing\$).tw.
 11 (financial or finance or finances or financed).tw.
 12 (fee or fees).tw.
 13 (value adj2 (money or monetary)).tw.
 14 Value of Life/
 15 quality adjusted life.tw.
 16 (qaly\$ or qald\$ or qale\$ or qtime\$).tw.
 17 disability adjusted life.tw.
 18 daly\$.tw.
 19 Health Status Indicators/
 20 (sf36 or sf 36 or short form 36 or shortform 36 or sf thirtysix or sf thirty six or
 shortform thirtysix or shortform thirty six or short form thirtysix or short form thirty
 six).tw.
 21 (sf6 or sf 6 or short form 6 or shortform 6 or sf six or sfsix or shortform six or short
 form six).tw.
 22 (sf12 or sf 12 or short form 12 or shortform 12 or sf twelve or sftwelve or shortform
 twelve or short form twelve).tw.
 23 (sf16 or sf 16 or short form 16 or shortform 16 or sf sixteen or sfsixteen or shortform
 sixteen or short form sixteen).tw.
 24 (sf20 or sf 20 or short form 20 or shortform 20 or sf twenty or sftwenty or shortform
 twenty or short form twenty).tw.
 25 (euroqol or euro qol or eq5d or eq 5d).tw.
 26 (hql or hqol or h qol or hrqol or hr qol).tw.
 27 (hye or hyes).tw.
 28 (hui or hui1 or hui2 or hui3).tw.
 29 utilit\$.tw.
 30 disutilit\$.tw.
 31 rosser.tw.
 32 quality of wellbeing.tw.
 33 qwb.tw.
 34 willingness to pay.tw.

- 35 standard gamble\$.tw.
 36 time trade off.tw.
 37 time tradeoff.tw.
 38 tto.tw.
 39 exp models, economic/
 40 models, theoretical/ or models, organizational/
 41 economic model\$.tw.
 42 markov chains/
 43 markov\$.tw.
 44 Monte Carlo Method/
 45 monte carlo.tw.
 46 exp Decision Theory/
 47 (decision\$ adj2 (tree\$ or anlay\$ or model\$)).tw.
 48 or/1-47

Laser

Laser terms - Central

- 1 MeSH descriptor Prostatic Hyperplasia, this term only with qualifier: SU
 2 MeSH descriptor Prostatic Hyperplasia, this term only
 3 MeSH descriptor Urinary Bladder Neck Obstruction, this term only
 4 benign prostat* near (hyperplas* or hypertroph* or obstruct* or enlarge* or disease)
 5 bph or bpo or bpe
 6 (bladder neck or bladder outlet or bladder outflow) near obstruct*
 7 #2 or #3 or #4 or #5 or #6
 8 MeSH descriptor Prostatectomy explode all trees
 9 MeSH descriptor Transurethral Resection of Prostate, this term only
 10 Transurethral near (resect* or electroresect* or incision* or diatherm* or vaporiz* or electrovaporiz* or evaporiz* or ablat* or thermo* or inject* or coagulat*)
 11 MeSH descriptor Electrosurgery explode all trees
 12 MeSH descriptor Laser Therapy, this term only
 13 MeSH descriptor Laser Coagulation, this term only
 14 laser near (resect* or ablat* or coagulat* or incision* or vaporiz*)
 15 laser near (enucleat* or prostatect*)
 16 laser near (holmium or yag or nd or ktp or green light)
 17 photoselectiv* near vaporiz*
 18 needle near ablat*
 19 microwave near thermo*
 20 coretherm or prostatron or targis or thermatrix or prolieve
 21 ethanol near inject*
 22 (water or cooled) near thermotherapy
 23 MeSH descriptor Ultrasound, High-Intensity Focused, Transrectal, this term only
 24 high intensity near ultrasound
 25 MeSH descriptor Stents, this term only
 26 prostat* near (stent* or spiral*)
 27 turp or tvap or tevap or tvp or tuevap
 28 tuip or vlap or holrp or holep or tuna or tumt
 29 ilc or tulip or hifu
 30 #11 or #12 or #13 or #14 or #16 or #17 or #18 or #19 or #21 or #22 or #23 or #24 or #25 or #29

- 31 #7 AND #30
 32 #1 or #8 or #9 or #10 or #15 or #20 or #26 or #27 or #28 or #31

Laser terms - OVID Embase

- 1 Prostate hypertrophy/su
 2 Prostate hypertrophy/
 3 bladder obstruction/
 4 (benign prostat\$ adj1 (hyperplas\$ or hypertroph\$ or obstruct\$ or enlarge\$ or disease)).tw.
 5 (bph or bpo or bpe).tw.
 6 ((bladder neck or bladder outlet or bladder outflow) adj1 obstruct\$).tw.
 7 or/2-6
 8 exp prostate surgery/
 9 (Transurethral adj3 (resect\$ or electroresect\$ or incision\$ or diatherm\$ or vaporis\$ or electrovaporis\$ or evaporis\$ or ablat\$ or thermo\$ or inject\$ or coagulat\$)).tw.
 10 exp laser/
 11 laser prostatectomy/
 12 laser surgery/
 13 Laser Coagulation/
 14 (laser adj3 (resect\$ or ablat\$ or coagulat\$ or incision\$ or vaporis\$)).tw.
 15 (laser adj3 (enucleat\$ or prostatect\$)).tw.
 16 (laser adj3 (holmium or yag or ktp or nd or green light)).tw.
 17 (photoselectiv\$ adj1 vaporis\$).tw.
 18 (needle adj3 ablat\$).tw.
 19 (microwave adj3 thermo\$).tw.
 20 (coretherm or prostatron or targis or thermatrix or prolieve).tw.
 21 (ethanol adj3 inject\$).tw.
 22 Laser thermotherapy/
 23 ((water or cooled) adj3 thermotherapy).tw.
 24 high intensity focused ultrasound/
 25 (high intensity adj3 ultrasound).tw.
 26 stents/
 27 (prostat\$ adj3 (stent\$ or spiral\$)).tw.
 28 (turp or tvvp or tevap or tvp or tuevap).tw.
 29 (tuip or vlap or holrp or holep or tuna or tumt).tw.
 30 (ilc or tulip or hifu).tw.
 31 or/10-14,16-19,21-26,30
 32 7 and 31
 33 or/1,8-9,15,20,27-29,32
 34 prostate cancer/ or bladder cancer/
 35 (cancer\$ or carcinoma\$ or neoplasm\$).tw.
 36 34 or 35
 37 36 not 7
 38 33 not 37

Laser terms - OVID Medline

- 1 Prostatic hyperplasia/su
 2 Prostatic hyperplasia/

- 3 Bladder neck obstruction/
 4 (benign prostat\$ adj1 (hyperplas\$ or hypertroph\$ or obstruct\$ or enlarge\$ or
 disease)).tw.
 5 (bph or bpo or bpe).tw.
 6 ((bladder neck or bladder outlet or bladder outflow) adj1 obstruct\$).tw.
 7 or/2-6
 8 exp prostatectomy/
 9 Transurethral resection of prostate/
 10 (Transurethral adj3 (resect\$ or electroresect\$ or incision\$ or diatherm\$ or vaporis\$ or
 electrovaporis\$ or evaporis\$ or ablat\$ or thermo\$ or inject\$ or coagulat\$)).tw.
 11 exp electrosurgery/
 12 laser therapy/
 13 laser coagulation/
 14 (laser adj3 (resect\$ or ablat\$ or coagulat\$ or incision\$ or vaporis\$)).tw.
 15 (laser adj3 (enucleat\$ or prostatect\$)).tw.
 16 (laser adj3 (holmium or yag or nd or ktp or green light)).tw.
 17 (photoselectiv\$ adj1 vaporis\$).tw.
 18 (needle adj3 ablat\$).tw.
 19 (microwave adj3 thermo\$).tw.
 20 (coretherm or prostatron or targis or thermatrix or prolieve).tw.
 21 (ethanol adj3 inject\$).tw.
 22 ((water or cooled) adj3 thermotherapy).tw.
 23 ultrasound, high-intensity focused, transrectal/
 24 (high intensity adj3 ultrasound).tw.
 25 stents/
 26 (prostat\$ adj3 (stent\$ or spiral\$)).tw.
 27 (turp or tvap or tevap or tvp or tuevap).tw.
 28 (tuip or vlap or holrp or holep or tuna or tumt).tw.
 29 (ilc or tulip or hifu).tw.
 30 or/11-14,16-19,21-25,29
 31 7 and 30
 32 or/1,8-10,15,20,26-28,31
 33 prostatic neoplasms/ or bladder neoplasms/
 34 (cancer\$ or carcinoma\$ or neoplasm\$).tw.
 35 33 or 34
 36 35 not 7
 37 32 not 36

Medications

Medication terms - Central

- 1 MeSH descriptor Adrenergic alpha-Antagonists, this term only
 2 (Alpha near (blocker or blocking agent or antagonist)):ti,ab
 3 MeSH descriptor Doxazosin, this term only
 4 MeSH descriptor Indoramin, this term only
 5 MeSH descriptor Prazosin, this term only
 6 (Doxazosin or Tamsulosin or Alfuzozin or Terazosin or Indoramin or Prazosin or
 Cardura or Stronazon or Flomaxtra or Flomax or Xaltral or Hytrin or Doralese or
 Hypovase):ti,ab
 7 (5-Alpha reductase inhibitor* or Alpha V reductase inhibitor*):ti,ab

- 8 MeSH descriptor Finasteride, this term only
 9 (Finasteride or Dutasteride or Avodart or Proscar):ti,ab
 10 MeSH descriptor Cholinergic Antagonists, this term only
 11 (Anticholinergic* or cholinergic antagonist* or antimuscarinic*):ti,ab
 12 (Oxybutynin or Tolterodine or Darifenacin or Propiverine or Solifenacin or Trosipium
 or Cystrin or Ditropan or Lyrinel or Detrusitol or Emselex or Detrunorm or Vesicare or
 Regurin):ti,ab
 13 MeSH descriptor Cyclic Nucleotide Phosphodiesterases, Type 5, this term only
 14 (Phosphodiesterase 5 inhibitor* or Phosphodiesterase V inhibitor*):ti,ab
 15 (PDE5 or sildenafil or viagra or vardenafil or levitra or tadalafil or cialis):ti,ab
 16 MeSH descriptor Phytotherapy, this term only
 17 MeSH descriptor Plant Extracts, this term only
 18 MeSH descriptor Plants, Medicinal, this term only
 19 (Phytotherapy or plant extract*):ti,ab
 20 MeSH descriptor Serenoa, this term only
 21 MeSH descriptor Sterols, this term only
 22 MeSH descriptor Sitosterols, this term only
 23 (Saw palmetto or serenoa or sabal or s repens or sitosterol* or b-sitosterol* or
 sitosteryl* or phytosterol*):ti,ab
 24 MeSH descriptor Secale cereale, this term only
 25 (pollen or secale cereale or rye or cernitin or cernilton):ti,ab
 26 MeSH descriptor Cucurbita, this term only
 27 (pumpkin seed\$ or cucurbita or pepita):ti,ab
 28 MeSH descriptor Urtica dioica, this term only
 29 (nettle or urtica):ti,ab
 30 MeSH descriptor Pygeum, this term only
 31 (pygeum africanum or prunus or tadenan or docosonal or pigenil):ti,ab
 32 (cranberry AND (juice or extract)):ti,ab
 33 MeSH descriptor Diuretics, this term only
 34 Diuretic*:ti,ab
 35 MeSH descriptor Furosemide, this term only
 36 MeSH descriptor Bumetanide, this term only
 37 (Frusemide or furosemide or bumetanide or burinex):ti,ab
 38 (Desmopressin or DDAVP or desmotabs or desmomelt or desmospray or octim):ti,ab
 39 MeSH descriptor Anti-Inflammatory Agents, Non-Steroidal, this term only
 40 (Aceclofenac or acemetacin or azapropazone or celecoxib or dexibuprofen or
 dexketoprofen or diclofenac or etodolac or etoricoxib or fenbufen or fenobufen or
 flurbiprofen or ibuprofen or indometacin or ketoprofen or mefenamic acid or
 meloxicam or nabumetone or naproxen or piroxicam or sulindac or tenoxicam or
 tiaprofenic acid or aspirin):ti,ab
 41 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or
 #13 or #14 or #15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23
 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or
 #34 or #35 or #36 or #37 or #38 or #39 or #40

Medication terms - OVID Embase

- 1 Alpha Adrenergic Receptor Blocking Agent/
 2 (Alpha ad[3 (blocker or blocking agent or antagonist)):ti,ab.
 3 Doxazosin/ or Tamsulosin/ or Alfuzosin/ or Terazosin/ or Indoramin/ or Prazosin/
 4 (Doxazosin or Tamsulosin or Alfuzosin or Terazosin or Indoramin or Prazosin or
 Cardura or Stronazon or Flomaxtra or Flomax or Xaltral or Hytrin or Doralase or
 Hypovase).ti,ab.

- 5 Steroid 5alpha Reductase Inhibitor/
6 (5-Alpha reductase inhibitor\$ or Alpha V reductase inhibitor\$).ti,ab.
7 Dutasteride/ or Finasteride/
8 (Finasteride or Dutasteride or Avodart or Proscar).ti,ab.
9 (Anticholinergic\$ or cholinergic antagonist\$ or antimuscarinic\$).ti,ab.
10 Oxybutynin/ or Tolterodine/ or Darifenacin/ or Propiverine/ or Solifenacin/ or
Tropium/
11 (Oxybutynin or Tolterodine or Darifenacin or Propiverine or Solifenacin or Tropium
or Cystrin or Ditropan or Lyrinel or Detrusitol or Emselex or Detrunorm or Vesicare or
Regurin).ti,ab.
12 Phosphodiesterase V Inhibitor/
13 (Phosphodiesterase 5 inhibitor\$ or Phosphodiesterase V inhibitor\$).ti,ab.
14 Sildenafil/ or Vardenafil/ or Tadalafil/
15 (PDE5 or sildenafil or viagra or vardenafil or levitra or tadalafil or cialis).ti,ab.
16 Phytotherapy/ or Plant extract/ or Medicinal plant/
17 (Phytotherapy or plant extract\$).ti,ab.
18 Sabal/ or Sterol/ or Sitosterol derivative/
19 (Saw palmetto or serenoa or sabal or s repens or sitosterol\$ or b-sitosterol\$ or
sitosteryl\$ or phytosterol\$).ti,ab.
20 Rye/ or Grass pollen extract/
21 (pollen or secale cereale or rye or cernitin or cernilton).ti,ab.
22 (pumpkin seed\$ or cucurbita or pepita).ti,ab.
23 Urtica extract/
24 (nettle or urtica).ti,ab.
25 Pygeum Africanum extract/
26 (pygeum africanum or prunus or tadenan or docosonal or pigenil).ti,ab.
27 Cranberry extract/ or Cranberry juice/
28 (cranberry adj1 (juice or extract)).ti,ab.
29 Diuretic Agent/
30 Diuretic\$.ti,ab.
31 Furosemide/ or Bumetanide/
32 (Frusemide or furosemide or bumetanide or burinex).ti,ab.
33 Desmopressin Acetate/ Or Desmopressin/
34 (Desmopressin or DDAVP or desmotabs or desmomelt or desmospray or octim).ti,ab.
35 Nonsteroid Antiinflammatory Agent/
36 (Non steroidal anti inflammator\$3 or NSAID\$).ti,ab.
37 Aceclofenac/ or acemetacin/ or azapropazone/ or celecoxib/ or dexibuprofen/ or
dexketoprofen/ or diclofenac/ or etodolac/ or etoricoxib/ or fenbufen/ or
fenobufen/ or flurbiprofen/ or ibuprofen/ or indometacin/ or ketoprofen/ or
mefenamic acid/ or meloxicam/ or nabumetone/ or naproxen/ or piroxicam/ or
sulindac/ or tenoxicam/ or tiaprofenic acid/ or aspirin/
38 (Aceclofenac or acemetacin or azapropazone or celecoxib or dexibuprofen or
dexketoprofen or diclofenac or etodolac or etoricoxib or fenbufen or fenobufen or
flurbiprofen or ibuprofen or indometacin or ketoprofen or mefenamic acid or
meloxicam or nabumetone or naproxen or piroxicam or sulindac or tenoxicam or
tiaprofenic acid or aspirin).ti,ab.
39 or/1-38

Medication terms - OVID Medline

- 1 Adrenergic alpha-Antagonists/
2 (Alpha adj3 (blocker or blocking agent or antagonist)).ti,ab.
3 Doxazosin/ or Indoramin/ or Prazosin/

- 4 (Doxazosin or Tamsulosin or Alfuzosin or Terazosin or Indoramin or Prazosin or Cardura or Stronazon or Flomaxtra or Flomax or Xaltral or Hytrin or Doralese or Hypovase).ti,ab.
- 5 (5-Alpha reductase inhibitor\$ or Alpha V reductase inhibitor\$).ti,ab.
- 6 Finasteride/
- 7 (Finasteride or Dutasteride or Avodart or Proscar).ti,ab.
- 8 Cholinergic Antagonists/
- 9 (Anticholinergic\$ or cholinergic antagonist\$ or antimuscarinic\$).ti,ab.
- 10 (Oxybutynin or Tolterodine or Darifenacin or Propiverine or Solifenacin or Trospium or Cystrin or Ditropan or Lyrinel or Detrusitol or Emselex or Detrunorm or Vesicare or Regurin).ti,ab.
- 11 Cyclic Nucleotide Phosphodiesterases, Type 5/
- 12 (Phosphodiesterase 5 inhibitor\$ or Phosphodiesterase V inhibitor\$).ti,ab.
- 13 (PDE5 or sildenafil or viagra or vardenafil or levitra or tadalafil or cialis).ti,ab.
- 14 Phytotherapy/ or Plant extracts/ or Plants, medicinal/ or serenoa/
- 15 (Phytotherapy or plant extract\$).ti,ab.
- 16 Serenoa/ or Sterols/ or Sitosterols/
- 17 (Saw palmetto or serenoa or sabal or s repens or sitosterol\$ or b-sitosterol\$ or sitosteryl\$ or phytosterol\$).ti,ab.
- 18 Secale Cereale/
- 19 (pollen or secale cereale or rye or cernitin or cernilton).ti,ab.
- 20 Cucurbita/
- 21 (pumpkin seed\$ or cucurbita or pepita).ti,ab.
- 22 Urtica dioica/
- 23 (nettle or urtica).ti,ab.
- 24 Pygeum/
- 25 (pygeum africanum or prunus or tadenan or docosonal or pigenil).ti,ab.
- 26 (cranberry adj1 (juice or extract)).ti,ab.
- 27 Diuretics/
- 28 Diuretic\$.ti,ab.
- 29 Furosemide/ or Bumetanide/
- 30 (Frusemide or furosemide or bumetanide or burinex).ti,ab.
- 31 (Desmopressin or DDAVP or desmotabs or desmomelt or desmospray or octim).ti,ab.
- 32 Anti-Inflammatory Agents, Non-Steroidal/
- 33 (Non steroidal anti inflammator\$3 or NSAID\$).ti,ab.
- 34 (Aceclofenac or acemetacin or azapropazone or celecoxib or dexibuprofen or dexketoprofen or diclofenac or etodolac or etoricoxib or fenbufen or fenobufen or flurbiprofen or ibuprofen or indometacin or ketoprofen or mefenamic acid or meloxicam or nabumetone or naproxen or piroxicam or sulindac or tenoxicam or tiaprofenic acid or aspirin).ti,ab.
- 35 or/1-34

Monitoring

Monitoring terms – Cochrane Library

- 1 (review* near (interval* or visit* or inspect* or examin* or attend* or check-up* or recall*))
- 2 (routine* near (interval* or visit* or inspect* or examin* or attend* or check-up* or recall*))
- 3 (periodic* near (interval* or visit* or inspect* or examin* or attend* or check-up* or recall*))
- 4 (regular near (visit* or inspect* or examin* or attend* or check-up*))
- 5 recall* near interval*

- 6 visit* near clinic*
7 #1 or #2 or #3 or #4 or #5 or #6

Monitoring terms – OVID Embase and Medline

- 1 (review\$ adj (interval\$ or visit\$ or inspect\$ or examin\$ or attend\$ or check-up\$ or recall\$)).tw.
2 (routine\$ adj (interval\$ or visit\$ or inspect\$ or examin\$ or attend\$ or check-up\$ or recall\$)).tw.
3 (periodic\$ adj (interval\$ or visit\$ or inspect\$ or examin\$ or attend\$ or check-up\$ or recall\$)).tw.
4 (regular adj (visit\$ or inspect\$ or examin\$ or attend\$ or check-up\$)).tw.
5 (recall\$ adj interval\$).tw.
6 (visit\$ adj5 clinic\$).tw.
7 or/1-6

Patient education

Patient education - OVID Embase

- 1 Patient/ or Hospital patient/ or Outpatient/
2 Caregiver/ or exp Family/ or exp Parent/
3 (patients or carer\$ or famil\$).tw.
4 or/1-3
5 Information Service/ or Information center/ or Publication/ or Book/ or Counseling/ or Directive counseling/
6 4 or 5
7 ((patient or patients) adj3 (education or educate or educating or information or literature or leaflet\$ or booklet\$ or pamphlet\$)).ti,ab.
8 Patient information/ or Patient education/
9 or/6-8

Patient education OVID Medline

- 1 Patients/ or Inpatients/ or Outpatients/
2 Caregivers/ or exp Family/ or exp Parents/ or exp Legal-Guardians/
3 (patients or carer\$ or famil\$).tw.
4 or/1-3
5 Popular-Works-Publication-Type/ or exp Information-Services/ or Publications/ or Books/ or Pamphlets/ or Counseling/ or Directive-Counseling/
6 4 or 5
7 ((patient or patients) adj3 (education or educate or educating or information or literature or leaflet\$ or booklet\$ or pamphlet\$)).ti,ab.
8 Patient-Education/ or Patient-Education-Handout-Publication-Type/
9 or/6-8

Patient views

Patient views - OVID Embase

- 1 Consumer attitude/ or patient satisfaction/ or patient compliance/ or patient right/ or health survey/ or questionnaire/ or interview/
2 (patient\$ adj3 (view\$ or opinion\$ or awareness or tolerance or perception or persistenc\$ or attitude\$ or compliance or satisfaction or concern\$ or belief\$ or feeling\$ or position or idea\$ or preference\$ or choice\$)).tw.

- 3 (Discomfort or comfort or inconvenience or bother\$4 or trouble or fear\$ or anxiety
or anxious or embarrass\$4).tw.
- 4 or/1-3

Patient views - OVID Medline

- 1 exp Consumer-Satisfaction/ or Personal-Satisfaction/ or exp Patient-Acceptance-Of-
Health-Care/ or exp Consumer-Participation/ or exp Patient-Rights/ or Health Care
Surveys/ or Questionnaires/ or Interview/ or Focus groups/
- 2 (patient\$ adj3 (view\$ or opinion\$ or awareness or tolerance or perception or
persistenc\$ or attitude\$ or compliance or satisfaction or concern\$ or belief\$ or
feeling\$ or position or idea\$ or preference\$ or choice\$)).tw.
- 3 (Discomfort or comfort or inconvenience or bother\$4 or trouble or fear\$ or anxiety
or anxious or embarrass\$4).tw.
- 4 or/1-3

RCT filter

RCT filter Embase

- 1 Clinical-Trial/ or Randomized-Controlled-Trial/ or Randomization/ or Single-Blind-
Procedure/ or Double-Blind-Procedure/ or Crossover-Procedure/ or Prospective-
Study/ or Placebo/
- 2 (((((((clinical or control or controlled) adj (study or trial)) or (single or double or
triple)) adj (blind\$3 or mask\$3)) or randomised or randomized or random\$) adj
(assign\$ or allocat\$ or group or grouped or patients or study or trial or distribut\$))
or crossover) adj (design or study or trial)) or placebo or placebos).ti,ab.
- 3 1 or 2

RCT filter Medline

- 1 Randomized-Controlled-Trials/ or Random-Allocation/ or Double-Blind-Method/ or
Single-Blind-Method/ or exp Clinical-Trials as topic/ or Cross-Over-Studies/ or
Prospective-Studies/ or Placebos/
- 2 (Randomized-Controlled-Trial or Clinical-Trial or Controlled-Clinical-Trial).pt.
- 3 (((((((clinical or control or controlled) adj (study or trial)) or (single or double or
triple)) adj (blind\$3 or mask\$3)) or randomised or randomized or random\$) adj
(assign\$ or allocat\$ or group or grouped or patients or study or trial or distribut\$))
or crossover) adj (design or study or trial)) or placebo or placebos).ti,ab.
- 4 or/1-3

Surgery

Surgery terms – Cochrane Library

- 1 MeSH descriptor Surgery, this term only
- 2 MeSH descriptor Urologic Surgical Procedures, this term only
- 3 MeSH descriptor Botulinum Toxins, this term only
- 4 botulinum or botox
- 5 Cystoplasty or bladder neck incision
- 6 Neuromodulation
- 7 Sacral nerve stimulation
- 8 Myectomy
- 9 MeSH descriptor Suburethral Slings, this term only
- 10 sling

- 11 injectable
- 12 MeSH descriptor Urinary Diversion, this term only
- 13 (Continent or incontinent) and diversion
- 14 MeSH descriptor Urinary Sphincter, Artificial, this term only
- 15 Artificial sphincter
- 16 Compression device
- 17 MeSH descriptor Catheterization, this term only
- 18 Suprapubic catheter*
- 19 Sphincterotomy
- 20 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18 or #19

Surgery terms - OVID Embase

- 1 Urologic Surgery/ or Male Genital System Surgery/ or Surgery/ or Bladder Surgery/ or Prostate Surgery/
- 2 Botulinum Toxin/
- 3 (botulinum or botox).tw.
- 4 Bladder Reconstruction/
- 5 (Bladder neck incision or cystoplasty).tw.
- 6 Neuromodulation/
- 7 neuromodulation.tw.
- 8 sacral nerve stimulation/
- 9 Sacral nerve stimulation.tw.
- 10 muscle resection/
- 11 Myectomy.tw.
- 12 sling.tw.
- 13 injectable.tw.
- 14 Urinary Diversion/
- 15 ((Continent or incontinent) and diversion).tw.
- 16 Bladder Sphincter Prosthesis/
- 17 Artificial sphincter.tw.
- 18 Compression device.tw.
- 19 Ureter Catheterization/ or Catheterization/
- 20 Suprapubic Catheter/
- 21 Suprapubic catheter\$.tw.
- 22 Sphincterotomy/
- 23 Sphincterotomy.tw.
- 24 or/1-23

Surgery terms - OVID Medline

- 1 Surgery/
- 2 Urologic Surgical Procedures/
- 3 Botulinum Toxins/
- 4 (botulinum or botox).tw.
- 5 (Cystoplasty or bladder neck incision).tw.
- 6 Neuromodulation.tw.
- 7 Sacral nerve stimulation.tw.
- 8 Myectomy.tw.

- 9 Suburethral Slings/
- 10 sling.tw.
- 11 injectable.tw.
- 12 Urinary Diversion/
- 13 ((Continent or incontinent) and diversion).tw.
- 14 Urinary Sphincter, Artificial/
- 15 Artificial sphincter.tw.
- 16 Compression device.tw.
- 17 Catheterization/
- 18 Suprapubic catheter\$.tw.
- 19 Sphincterotomy.tw.
- 20 or/1-19

Systematic review filter

Systematic review filter - OVID Medline

- 1 meta-analysis/
- 2 (metaanalys\$ or meta-analys\$ or meta analys\$).tw.
- 3 exp "review literature"/
- 4 (systematic\$ adj3 (review\$ or overview\$)).tw.
- 5 (selection criteria or data extraction).ab. and review.pt.
- 6 (cochrane or embase or psychlit or psychlit or psychinfo or psycinfo or cinahl or cinhal or science citation index or bids or cancerlit).ab.
- 7 (reference list\$ or bibliograph\$ or hand search\$ or hand-search\$ or manual search\$ or relevant journals).ab.
- 8 or/1-7

Systematic review filter - OVID Embase

- 1 meta analysis/
- 2 (metaanalys\$ or meta-analys\$ or meta analys\$).tw.
- 3 systematic review/
- 4 (systematic\$ adj3 (review\$ or overview\$)).tw.
- 5 (selection criteria or data extraction).ab. and Review.pt.
- 6 (cochrane or embase or psychlit or psychlit or psychinfo or psycinfo or cinahl or cinhal or science citation index or bids or cancerlit).ab.
- 7 (reference list\$ or bibliograph\$ or hand search\$ or manual search\$ or relevant journals).ab.
- 8 or/1-7

Appendix D – Evidence Tables

Evidence Table 1: Diagnostic accuracy for urinalysis	48
Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?	50
Evidence Table 3: Diagnostic accuracy of uroflowmetry.....	60
Evidence Table 4: Diagnostic accuracy of post void residual.....	63
Evidence Table 5: Pelvic floor exercises (with or without electrical stimulation or biofeedback)	64
Evidence Table 6: Post void milking vs. no intervention or other conservative intervention.....	87
Evidence Table 7: Product vs. no product or other conservative intervention	88
Evidence Table 8: Catheters vs. no catheters	97
Evidence Table 9: Alpha-blockers vs. placebo	103
Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors	152
Evidence Table 11: Alpha-blockers vs. anticholinergics	165
Evidence Table 12: Alpha-blockers vs. phosphodiesterase-5 inhibitors.....	166
Evidence Table 13: 5-alpha reductase inhibitors vs. placebo	168
Evidence Table 14: Anticholinergics vs. placebo	192
Evidence Table 15: Phosphodiesterase-5 inhibitors vs. placebo.....	193
Evidence Table 16: Diuretics vs. placebo	204
Evidence Table 17: Desmopressin vs. placebo	206
Evidence Table 18: Non steroidal anti-inflammatory drugs (NSAIDS) vs. placebo	207
Evidence Table 19: Combination therapy: 5-Alpha reductase inhibitor and alpha-blocker.....	209
Evidence Table 20: Combination therapy: Anticholinergic added to alpha-blocker.....	210
Evidence Table 21: Combination therapy: phosphodiesterase-5-inhibitor added to alpha-blocker	212
Evidence Table 22: Holmium laser enucleation (or resection) of the prostate HoLEP (HoLRP) vs. transurethral resection of the prostate.....	216
Evidence Table 23: Thulium laser resection vs. transurethral resection of the prostate.....	232
Evidence Table 24: Holmium laser enucleation of the prostate (HoLEP) vs. transurethral incision of the prostate (HoBNI)	235
Evidence Table 25: Holmium laser enucleation of the prostate (HoLEP) vs. open prostatectomy (OP)	239
Evidence Table 26: Laser coagulation vs. transurethral resection of the prostate (TURP).....	246
Evidence Table 27: Laser vaporisation vs. transurethral resection of the prostate (TURP).....	270
Evidence Table 28: Laser vs. open prostatectomy	292
Evidence Table 29: Laser vs. transurethral microwave thermotherapy (TUMT)	296
Evidence Table 30: Laser vs. transurethral vaporisation of the prostate (TUVV)	298
Evidence Table 31: Laser coagulation vs. laser vaporisation	306
Evidence Table 32: Holmium laser resection of the prostate (HoLRP) vs. laser coagulation	311
Evidence Table 33: Holmium laser enucleation of the prostate (HoLEP) vs. laser vaporisation.....	314
Evidence Table 34: Transurethral microwave thermotherapy (TUMT) vs. no treatment.....	317
Evidence Table 35: Transurethral microwave thermotherapy (TUMT) vs. transurethral resection of the prostate (TURP)	333
Evidence Table 36: Transurethral vaporisation of the prostate (TUVV) vs. transurethral resection of the prostate (TURP)	347
Evidence Table 37: Bipolar transurethral vaporisation of the prostate (TUVV) vs. transurethral resection of the prostate (TURP)	375
Evidence Table 38: Transurethral needle ablation (TUNA) vs. transurethral resection of the prostate (TURP)	381
Evidence Table 39: Transurethral incision of the prostate (TUIP) vs. transurethral resection of the prostate (TURP).....	391
Evidence Table 40: Botulinum toxin vs. placebo	411
Evidence Table 41: Transurethral vaporesction of the prostate (TUVRP) vs. transurethral resection of the prostate (TURP)	412
Evidence Table 42: Bipolar TUVRP vs. transurethral resection of the prostate (TURP)	426

Evidence Table 43: Transurethral ethanol ablation of the prostate (TEAP) vs. transurethral resection of the prostate (TURP)	428
Evidence Table 44: Transurethral resection of the prostate (TURP) vs. watchful waiting	429
Evidence Table 45: Bipolar transurethral resection of the prostate (TURP) vs. TURP	433
Evidence Table 46: Conservative vs. surgery	454
Evidence Table 47: What is the effectiveness of alpha-blockers in treating men after acute urinary retention?	460
Evidence Table 48: Phytotherapy vs. placebo	466
Evidence Table 49: Phytotherapy combinations vs. placebo	474
Evidence Table 50: Phytotherapy vs. Alpha-blockers	478
Evidence Table 51: Phytotherapy vs. 5-Alpha Reductase inhibitors	484
Evidence Table 52: Provision of information.....	488
Evidence Table 53: Economic evidence	494

Abbreviations

5-ARI	5-Alpha-Reductase Inhibitors
AB	Alpha-Blockers
AUA	American Urological Association
AUASS	American Urological Association Symptom Score
AUR	Acute Urinary Retention
BOO	Bladder outlet obstruction
BPE	Benign prostatic enlargement
BPH	Benign prostatic hyperlasia
BPO	Benign prostatic obstruction
CI 95%	95% Confidence interval
DRE	Digital rectal examination
ED	Erectile dysfunction
GP	General Practitioner
HIFU	High Intensity Focused Ultrasound
HoLAP	Holmium Laser Ablation of the Prostate
HoLEP	Holmium Laser Enucleation of the prostate
HoLRP	Holmium Laser Resection of the Prostate
ICER	Incremental Cost-Effectiveness Ratio
ICS	International Continence Society
ILC	Interstitial Laser Coagulation
Int	Intervention
IPSS	International prostate symptom score
IQR	Interquartile range
ITT	Intention to treat analysis
KTP	Potassium-Titanyl-Phosphate
LOS	Length Of Stay
LUTS	Lower urinary tract symptoms
M/F	Male/female
N	Total number of patients randomised
NA	Not Applicable
NR	Not reported
OAB	Overactive bladder
PFMT	Pelvic floor muscle training
PMD	Post micturition dribble
PPP	Purchasing Power Parities
PSA	Prostate specific antigen
PVM	Post-void milking
PVP	Photoselective vaporisation of the prostate
PVR	Post voidal residual
QALY	Quality-Adjusted Life Years
Q_{max}	Maximum urinary flow rate
QoL	Quality of life
RBC	Red blood cells
RCT	Randomised controlled trial
RR	Relative risk

SA	Sensitivity Analysis
SD	Standard Deviation
SE	Standard Error
Sig	Statistically significant at 5%
TEAP	Transurethral ethanol ablation of the prostate
TUIP	Transurethral incision of the prostate
TUMT	Transurethral microwave thermotherapy
TUNA	Transurethral needle ablation
TURP	Transurethral resection of the prostate
TUVP	Transurethral vaporisation of the prostate
TUVRP	Transurethral vaporisation resection of the prostate
TVP	Transurethral electroVaporisation of the Prostate
TWOC	Trial Without Catheter
UI	Urinary incontinence
UTI	Urinary Tract Infection
Vs	Versus
WW	Watchful Waiting

Evidence Table 1: Diagnostic accuracy for urinalysis

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
<p>Ezz et al., 1996⁸⁵</p> <p>Study design: Cross sectional study</p> <p>Evidence level: Level-2 study (II)</p> <p>Duration of follow-up: NR. Tests carried out over 2 visits.</p>	<p>Patient group: Consecutive men at one outpatient department (Department of Urology, Nijmegen, The Netherlands) with BPE and LUTS, either irritative or obstructive.</p> <p>Exclusion criteria: Patients excluded from further assessment for BPH once a prostate carcinoma suspected.</p> <p>All patients N: 750 Av Age (range): 64 years (40-85) Drop outs: 0</p>	<p>Assessment tool under investigation: Urinalysis by dipstick readings from clean mid-stream specimen, If revealed erythrocytes urine sediment microscopy was completed.</p> <p>Sediment grading completed by number of red blood cells (RBC): Grade 1 = 0 RBC Grade 2 = 1-5 RBC Grade 3 = 6-10 RBC Grade 4 = 10+ RBC</p> <p>Results: Grade 1: 516 (68.8%) Grade 2: 207 (27.2%) Grade 3: 15 (2%) Grade 4: 12 (1.6%)</p> <p>Gold standard: Cystoscopy and histology.</p> <p>Additional tests: All patients underwent: History, IPSS, physical examination with Digital rectal examination, biochemistry (PSA and serum creatinine), urine culture and cytology, trans rectal ultrasonography, plain abdominal X-ray, renal ultrasound, flexible cystoscopy, flow, post void residual (PVR) and urodynamic investigations.</p>	<p>Bladder tumours</p> <p>Sensitivity Specificity PPV NPV Prevalence Positive LR Negative LR Pre-test Odds (CI 95%) Post-Test Odds +ve result Post-Test Odds -ve result</p> <p>Urinary tract infection by urine culture</p> <p>Sensitivity Specificity PPV NPV Prevalence Positive LR Negative LR Pre-test Odds (CI 95%) Post-Test Odds +ve result Post-Test Odds -ve result</p> <p>Urinary calculi (Stones) by abdominal X-ray</p> <p>Sensitivity Specificity PPV NPV</p>	<p>Grade 1: 1/516 (0.2%) Grade 2, 3 & 4: 2/234 (0.9%) Grade 2: 2/207 Grade 3: 0/15 Grade 4: 0/12 66.7% 68.9% 0.9% 99.8% 3/750 (0.4%) 2.15 0.48 0.004(0-0.01) 0.01 0.01</p> <p>Grade 1: 7/516 (1.4%) Grade 2, 3 & 4: 10/234 (4.3%) Grade 2: 9/207 Grade 3: 0/15 Grade 4: 1/12 58.8% 69.4% 4.3% 98.6% 17/750 (2.3%) 1.9 0.6 0.02 (0.01-0.03) 0.04 0.03</p> <p>Grade 1: 35/516 (6.8%) Grade 2, 3 & 4: 14/234 (6.0%) Grade 2: 12/207 Grade 3: 1/15 Grade 4: 1/12 28.6% 68.6% 6.0% 93.2%</p>	<p>Funding: NR.</p> <p>Limitations: Cystoscopy performed on second visit after initial tests.</p> <p>Additional tests: Correlation of grades of RBC to age, prostate volume, IPSS, residual urine and outlet obstruction. Papillary lesion and dilatation were reported. One renal tumour was reported.</p> <p>Notes: All patients with positive dipstick readings were found to have red cells on microscopy.</p> <p>Sensitivity and specificity values calculated by NCGC using no RBC found (negative) compared to any RBC (positive).</p> <p>All values calculated to 1d.p.</p>

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
			Prevalence Positive LR Negative LR Pre-test Odds (CI 95%) Post-Test Odds +ve result Post-Test Odds -ve result	49/750 (6.5%) 0.91 1.04 0.07 (0.05-0.09) 0.06 0.07	
			Cyst by renal ultrasound Sensitivity Specificity PPV NPV Prevalence Positive LR Negative LR Pre-test Odds (CI 95%) Post-Test Odds +ve result Post-Test Odds -ve result	Grade 1: 39/516 (7.6%) Grade 2, 3, & 4: 22/234 (9.4%) Grade 2: 11/207 Grade 3: 10/15 Grade 4: 1/12 36.1% 69.2% 9.4% 92.4% 61/750 (8.1%) 1.17 0.92 0.09 (0.07-0.11) 0.10 0.10	

Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?

Study details	Patients	Outcome measures & Analysis	Effect size	Comments
<p>Carter et al., 2005⁴⁶</p> <p>Study design: Longitudinal Cohort</p> <p>Duration of follow-up: Long-term from 1959</p>	<p>Patient group: cohort of men from the Baltimore Longitudinal Study of Aging (BLSA).</p> <p>Setting: USA</p> <p>Interventions: Not applicable</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • < 70 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Medical or surgical treatment of BPH • Development of prostate cancer <p>All patients N: 704</p> <p>Drop outs:</p> <p>Group 1 (age <50) N: 370 Age (median + range): 37.4 (22.5 – 49.9) 25th percentile PSA (ng/mL): 0.3 50th percentile PSA (ng/mL): 0.5 75th percentile PSA (ng/mL): 0.8 Median symptom evaluation (range): 6 (1-18)</p> <p>Group 2 (age 50 – 69.9) N: 334 Age (median + range): 59.3 (50.1 – 69.9) 25th percentile PSA (ng/mL): 0.5 50th percentile PSA (ng/mL): 0.9 75th percentile PSA (ng/mL): 2.0 Median symptom evaluation (range): 10.5 (0-28)</p>	<p>Change in IPSS over time with PSA</p> <p>Mixed effect Poisson model (because of repeated measures between subjects) used to test whether there was a significant relationship between PSA percentile grouping and symptom score with time</p>	<p>No correlation – analysis not shown</p>	<p>Funding: National Institute on Aging Intramural Research Program and gift from GSK.</p> <p>Limitations: No results for regression analysis of IPSS score and PSA</p> <p>Additional outcomes:</p> <ul style="list-style-type: none"> • Symptom score distribution by percentile against PSA percentile grouped by age • Correlation plot of medical history symptom score with IPSS. • Plot of symptom score vs. age for each PSA percentile <p>Notes: Baseline PSA was divided into percentiles: <25th 25th – 50th >75th Patients also divided into age groups at the time of 1st PSA measurement</p> <p>PSA measurements at visits started in 1991 otherwise measured retrospectively from serum samples</p> <p>Medical history questionnaire used from 1959 – 1991 and IPSS also used from 1991 – 2000. Questions relating to lower urinary tract score from medical history were used to devise score 0 - 13</p>

Study details	Patients	Outcomes			Analysis conducted	Results	Comments	
<p>Laguna et al. 2002¹⁵⁷</p> <p>Study design: Cohort</p> <p>Duration of follow-up: Minimum of 1 year. Evaluated every 3 months during year 1 and every 6 months in year 2 and thereafter</p>	<p>Patient group: Consecutive patients treated with transurethral thermotherapy</p> <p>Setting: Secondary care, Netherlands</p> <p>Interventions: transurethral thermotherapy</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Treated with transurethral thermotherapy between February 1992 to June 1999, when data were available on pre-treatment determination of PSA, free uroflowmetry, voided and post-void residual urine, ultrasound measurement of prostate volume, and IPSS scores. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previously treated with transurethral thermotherapy, medical therapy or manipulation of the lower urinary tract interfering with baseline PSA. Neurogenic or systemic disorder that may have impaired bladder function. <p>All patients N: 404 M/F: 404/0 Age (mean, range): 66.3 (44.8-89.7) Drop outs: 16/404, 388 analysed</p>						<p>Funding: not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> Patients received surgical treatment (TUMT) “Retreated patients”, analysed as having unchanged values at 12 months Report: “no relevant linear correlation was noted for baseline PSA with changes in IPSS, QoL or Qmax.” <p>Additional outcomes:</p> <ul style="list-style-type: none"> Values for a subgroup of patients, who have similar inclusion criteria for Djavan 2004 was reported. <p>Notes:</p> <ul style="list-style-type: none"> Seems to address the question of “does baseline PSA predict TUMT surgery outcomes”? Retrospective study, on “prospectively collected data”. 	
		Age (years):	66.3 (44.8-89.7)	-		Linear regression: Change in QoL vs. pretreatment PSA		Spearman r: -0.135 “linear regression coefficient”: -0.04 P value: 0.01
		PSA (ng/ml):	5.3 (0.1-45)	-				
		IPSS:	19.1 (3-35)	9.4(0-32)		Linear regression: Change in Qmax vs. pretreatment PSA		Spearman r: 0.105 , “linear regression coefficient”: 0.105 P value: 0.1
		QoL (IPSS)	3.9(0-6)	1.9(0-5)				
		Prostate volume, PV (cm3)	57.7(25-178) 18 (11-31)	-		Mann Whitney test: Baseline PSA vs. these outcomes at 1 year		Box and whisker plots shown, reported as “no association”
		Qmax (mL/s):	9.4 (2-19.9)	14.6(2.4-50.3)		- IPSS>7 vs. less		
		Voided vol (ml)	226(22-763)			- Qmax >12 vs. less		
		Post-void vol (ml)	86(0-755)			- QoL 1 or 2 (or 1 or 0)		
		All values reported were mean (range), unless otherwise specified						

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>McConnell et al., 2003¹⁹¹</p> <p>MTOPS research group NCT00021814</p> <p>Setting: multi-centre, 17 centres USA</p> <p>Study design: RCT double blinded (4 arms)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Mean follow up 4.5 years</p> <p>Study also reported in Bautista et al., 2003²⁵</p>	<p>Patient group: Men with BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ≥ 50 years Qmax between 4 - 15 mL/sec; and voided volume ≥ 125 mL. AUA-7 Symptom Score 8 - 30. Voluntarily signed the informed consent agreement prior to the performance of any study procedures. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Serum PSA > 10 ng/mL. Supine blood pressure < 90/70 mmHg Orthostatic hypotension. Prior medical/surgical intervention for BPH. Received prior experimental intervention (either medical or surgical) for prostate disease or enrolled in any other study protocol. <p>All patients N: 3047 out of 4391 screened Mean age: 62.6 ± 7.3 Drop outs: ?</p> <p>Group 1 (Doxazosin) N: 756 Age Mean (± SD): 62.7 ± 7.2 White race (%): 82.5 AUA-7 (± SD): 17.0 ± 5.8 Qmax (± SD), mL/s: 10.3 ± 2.5 Prostate volume (± SD), mL: 36.9 ± 21.6</p>	<p>Group 1: Doxazosin 10 mg (+ placebo) Single daily dose at bedtime. Dose doubled at 1 week intervals starting at 1 mg/day for the 1st week until final dose of 8 mg/day. Men who could not tolerate 8mg were given 4 mg. Those who could not tolerate 4 or 8 mg were discontinued.</p> <p>Group 2: Finasteride 5mg (+ placebo) Single daily dose at bedtime</p> <p>Group 3: Doxazosin 10 mg + finasteride 5 mg Single daily dose at bedtime</p> <p>Group 4: placebo for Doxazosin and placebo for finasteride Single daily dose at bedtime</p> <p>Examination methods: Vital signs, AUA symptom score, Qmax, compliance, adverse events</p>	<p>Cumulative incidence of clinical progression defined as first occurrence of increase of ≥ 4 points AUA-7 score over baseline at 4 years log rank test</p> <p>Cumulative incidence of clinical progression defined as incidence of acute urinary retention at 4 years log rank test</p> <p>Mean change in AUA ± SD at 4 years</p> <p>Mean change in Qmax ± SD at 4 years</p> <p>Change of prostate volume compared to baseline, mean±sd (ml)</p>	<p>Grp 1: 55/756 Grp 2: 65/768 Grp 3: 36/786 Grp 4: 97/737 P value: grp 1 v grp 4 <0.001, P value: grp 2 v grp 4 <0.016 P value: grp 3 v grp 4 <0.001 No significant differences between grps 1, 2 or 3</p> <p>Grp 1: 9/756 Grp 2: 6/768 Grp 3: 4/786 Grp 4: 18/737 P value: grp 1 v grp 4 =0.23 P value: grp 2 v grp 4 =0.009 P value: grp 3 v grp 4 <0.001</p> <p>Grp 1: 6.6 ± 5.8** Grp 2: 5.6 ± 5.0** Grp 3: 7.4 ± 5.7* Grp 4: 4.9 ± 4.1* P value: grp 1 v grp 4 <0.001 P value: grp 2 v grp 4 =0.001* P value: grp 3 v grp 4 <0.001 P value: grp 1 v grp 3 =0.006* P value: grp 2 v grp 3 <0.001 P value: grp 1 v grp 2 =0.001*</p> <p>Grp 1: 4.0 ± NR Grp 2: 3.2 ± NR Grp 3: 5.1 ± NR Grp 4: NR P values were only available for median change from baseline</p> <p>Group 1: 8.00±16.07 Group 2: -2.76±14.42 Group 3: -1.91±13.63</p>	<p>Funding: National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) National Institutes of Health, National Centre for Minority Health & Health Disparities, Merck and Pfizer.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Standard deviations were not reported for mean changes from baseline for secondary outcomes Number of patients discontinuing in the placebo group were not reported. <p>Additional outcomes: Median changes from baseline for symptom score, Qmax and serum PSA at 1 year and 4 years.</p> <p>Percentage discontinued therapy (most of them due to adverse events) Doxazosin: 27% Finasteride: 24% Combination: 18% (discontinued both)</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																																											
	<p>PVR (± SD), mL: 69.2 ± 88.2 PSA serum(± SD), ng/mL: 2.4 ± 2.1 Dropouts: 204/756 (27%)</p> <p>Group 2 (Finasteride) N: 768 Age Mean (± SD): 62.67 ± 7.3 White race (%): 83.7 AUA-7 (± SD): 17.6 ± 5.9 Qmax (± SD), mL/s:10.5 ± 2.5 Prostate volume (± SD), mL: 36.9 ± 20.6 PVR (± SD), mL: 66.2 ± 80.0 PSA serum(± SD), ng/mL: 2.4 ± 2.1 Dropouts: 174/768 (24%)</p> <p>Group 3: (Doxazosin + finasteride 5 mg) N: 786 Age Mean (± SD): 62.7 ± 7.1 White race (%): 80.8 AUA-7 (± SD): 16.8 ± 5.8 Qmax (± SD), mL/s:10.6 ± 2.5 Prostate volume (± SD), mL: 36.4 ± 19.2 PVR (± SD), mL: 67.5 ± 81.1 PSA serum(± SD), ng/mL: 2.3 ± 1.9 Dropouts: 141/786 (18%)</p> <p>Group 4: (placebo for Doxazosin and placebo for Finasteride) N: 737 Age Mean (± SD): 62.5 ± 7.5 White race (%): 82.4 AUA-7 (± SD): 16.8 ± 5.9 Qmax (± SD), mL/s:10.5 ± 2.6 Prostate volume (± SD), mL: 35.2 ± 18.8 PVR (± SD), mL: 69.6 ± 82.1</p>	<p>measured every 3 months. DRE, Serum PSA and urinalysis performed annually. Prostate volume assessed by TRUS at baseline and 5 year follow up.</p>	<p>[Calculated by NCC-AC from Kaplan2008B¹³⁵]</p> <p>Adverse events\$</p> <table border="1"> <thead> <tr> <th>Total no. of person-year</th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp3</th> <th>Grp4</th> </tr> </thead> <tbody> <tr> <td>Erectile Dysfunction</td> <td>3489</td> <td>3600</td> <td>3832</td> <td>3489</td> </tr> <tr> <td>Libido decrease</td> <td>3.56</td> <td>4.53</td> <td>5.11</td> <td>3.32</td> </tr> <tr> <td>Ejaculation disorder</td> <td>1.56</td> <td>2.36</td> <td>2.51</td> <td>1.40</td> </tr> <tr> <td>Postural hypotension</td> <td>1.10</td> <td>1.78</td> <td>3.05</td> <td>0.83</td> </tr> <tr> <td>Asthenia</td> <td>4.03</td> <td>2.56</td> <td>4.33</td> <td>2.29</td> </tr> <tr> <td>Dizziness</td> <td>4.08</td> <td>1.56</td> <td>4.20</td> <td>2.06</td> </tr> <tr> <td>Peripheral oedema</td> <td>4.41</td> <td>2.33</td> <td>5.35</td> <td>2.29</td> </tr> <tr> <td>Dyspnea</td> <td>0.88</td> <td>0.72</td> <td>1.25</td> <td>0.66</td> </tr> <tr> <td>Allergic reaction</td> <td>0.93</td> <td>0.56</td> <td>1.20</td> <td></td> </tr> <tr> <td>Somnolence</td> <td>0.57</td> <td></td> <td></td> <td></td> </tr> <tr> <td>\$ 10 most frequently reported adverse expressed as rate per 100 person-year of follow up.</td> <td>0.85</td> <td>0.58</td> <td>0.73</td> <td></td> </tr> <tr> <td></td> <td>0.46</td> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td>0.82</td> <td>0.39</td> <td>0.78</td> <td></td> </tr> <tr> <td></td> <td>0.37</td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p>Prognosis value of PSA, based on placebo arm</p> <p>[Data from Crawford2006,⁵⁷]</p> <p>Overall BPH progression was defines as the first occurrence of an increase of at least 4 points in the AUASS, AUR, urinary incontinence or renal insufficiency or recurrent UTI</p>	Total no. of person-year	Grp 1	Grp 2	Grp3	Grp4	Erectile Dysfunction	3489	3600	3832	3489	Libido decrease	3.56	4.53	5.11	3.32	Ejaculation disorder	1.56	2.36	2.51	1.40	Postural hypotension	1.10	1.78	3.05	0.83	Asthenia	4.03	2.56	4.33	2.29	Dizziness	4.08	1.56	4.20	2.06	Peripheral oedema	4.41	2.33	5.35	2.29	Dyspnea	0.88	0.72	1.25	0.66	Allergic reaction	0.93	0.56	1.20		Somnolence	0.57				\$ 10 most frequently reported adverse expressed as rate per 100 person-year of follow up.	0.85	0.58	0.73			0.46					0.82	0.39	0.78			0.37				<p>Group 4: 6.67±15.98</p>	<p>Notes: Urn method of randomisation and stratified according to centre. Merck and Pfizer supplied active drugs and placebo designed to look and taste like Doxazosin and Finasteride. Allocation concealment preserved by coded medications distributed by drug company.</p> <p>Eligible patients entered 2 week single blind placebo run-in.</p> <p>Patients discontinued were followed for primary and secondary outcomes</p> <p>* P values between comparisons were used along with mean differences to estimate standard deviations for groups. Where possible exact p values were used. As numbers of patients as each follow up point not clear the ITT numbers were used. Methods were following Cochrane Handbook. **Where >1 possible standard deviations were calculated for a group the mean was used</p>
Total no. of person-year	Grp 1	Grp 2	Grp3	Grp4																																																																												
Erectile Dysfunction	3489	3600	3832	3489																																																																												
Libido decrease	3.56	4.53	5.11	3.32																																																																												
Ejaculation disorder	1.56	2.36	2.51	1.40																																																																												
Postural hypotension	1.10	1.78	3.05	0.83																																																																												
Asthenia	4.03	2.56	4.33	2.29																																																																												
Dizziness	4.08	1.56	4.20	2.06																																																																												
Peripheral oedema	4.41	2.33	5.35	2.29																																																																												
Dyspnea	0.88	0.72	1.25	0.66																																																																												
Allergic reaction	0.93	0.56	1.20																																																																													
Somnolence	0.57																																																																															
\$ 10 most frequently reported adverse expressed as rate per 100 person-year of follow up.	0.85	0.58	0.73																																																																													
	0.46																																																																															
	0.82	0.39	0.78																																																																													
	0.37																																																																															

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	PSA serum(\pm SD), ng/mL: 2.3 ± 2.0 Dropouts: Not reported			P=0.0029 Incidence rate of invasive therapy (events/100 person year) PSA \geq 1.6ng/ml: 1.8 PSA<1.6ng/ml: 0.8 P=0.018	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Roehrborn et al., 2006²⁵⁵</p> <p>Study design: RCT</p> <p>Setting: multi-centre in US, Europe, Australia, Middle-east and South Africa.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Men at risk of having progression events from LUTS/BPH enrolled between May 2001 and March 2005.</p> <p>Inclusion criteria: ≥55 years with a ≥6 month history of LUTS related to BPH, an IPSS of ≥13, a Qmax of 5-12mL/s for a voided volume of ≥150mL, a PVR of ≥350mL, a prostate of ≥30g estimated by DRE, and a PSA level of 1.4-10ng/mL.</p> <p>Exclusion criteria: previous occurrence of AUR or prostatic surgery; concomitant urological diseases; diagnosed or suspected prostate carcinoma; previous x-ray therapy of the pelvic region; history of postural hypotension or syncope; concomitant use of medications that may alter the voiding pattern; and clinically relevant biochemical abnormalities.</p> <p>All patients N: 1522</p> <p>Group 1 N: 759 (ITT analysis N: 749) Mean (±SD) Age: 66.4 (6.7) Dropouts: 230 (Lack of efficacy or disease progression 75; adverse events 71; patients request=39; poor compliance with protocol=8, lost to follow-up=6; other 31)</p> <p>Group 2 N: 763 (ITT analysis N: 757)</p>	<p>Group 1: alpha-blocker Alfuzosin 10mg once daily</p> <p>Group 2: Placebo</p>	Number (%) progressed to AUR	Group1: 16 (2.1%) Group 2: 14 (1.8%) P=0.82	<p>Funding: Sanofi-Aventis</p> <p>Limitations: Method of randomisation and allocation concealment unclear.</p> <p>Additional outcomes: Haematological or biochemical measurement s-reported that there were no significant changes.</p> <p>Notes: Baseline variables analysed as predictors of IPSS worsening, AUR or BPH related surgery.</p>
			Number (%) men with BPH-related surgery	Group1: 38 (5.1%) Group 2: 49 (6.5%) P=0.18 RR: 22 (-18 to 48)%	
			Number (%) patients with symptom progression of ≥ 4points	Group1: 88 (11.7%) Group 2: 127 (16.8%) P=0.0013 RR with alfuzosin: 30 (10-46)%	
			Number (%) of men having any LUTS/BPH progression event (AUR and/or surgery and/or IPSS deterioration of ≥4 points)	Group1: 122 (16.3%) Group 2: 167 (22.1%) P<0.001 RR with alfuzosin: 26 (9-40)%	
			Mean (SD) decrease from baseline in IPSS	Group1: -5.9 (6.9) Group 2: -4.7 (6.9)	
			Mean (SD) decrease from baseline in bother score	Group1: -1.3 (1.5) Group 2: -0.9 (1.6) P<0.001	
			Mean (SD) decrease from baseline in Qmax, mL/s at 12 months	Group1: 2.0 (3.8) Group 2: 1.3 (3.6) P=0.001	
			Median change in serum PSA levels	Group 1: -0.6% Group 2: 3.6%; P=0.07	
			Treatment emergent adverse events	Group 1: 400 (53.1%) Group 2: 390 (51.2%)	
			Discontinuation after TEAE	Group 1: 69 (9.2%) Group 2: 58 (7.6%)	
Adverse events	<p>Dizziness Group 1: 45 (6.0%) Group 2: 35 (4.6%)</p> <p>Headache Group 1: 25 (3.3%) Group 2: 17 (2.2%)</p> <p>Hypotension</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean (\pmSD) Age: 66.5 (7.0)</p> <p>Dropouts: 283 (Lack of efficacy or disease progression=111; adverse events=62; patients request=58; poor compliance with protocol=13, lost to follow-up=12; other=27)</p>			<p>Group 1: 9 (1.2%) Group 2: 4 (0.5%)</p> <p>Syncope Group 1: 5 (0.7%) Group 2: 2 (0.3%)</p> <p>Malaise Group 1: 1 (0.1%) Group 2: 0</p> <p>Ejaculatory dysfunction Group 1: 15 (2.0%) Group 2: 14 (1.8%)</p> <p>Ejaculatory disorders Group 1: 3 (0.4%) Group 2: 0</p> <p>Asthenia/fatigue Group 1: 16 (2.1%) Group 2: 8 (1.1%)</p> <p>Somnolence Group 1: 0 Group 2: 3 (0.4%)</p>	
			<p>Mean (SD) changes in SBP/DBP, mmHg</p>	<p>Supine Group 1: -3.2 (15.6)/-2.9 (10.1) Group 2: -0.1 (15.3)/-0.8 (9.3)</p> <p>Standing Group 1: -3.8 (15.5)/ -2.8 (10.3) Group 2: -0.2 (15.5)/-0.5 (10.0)</p>	
			<p>Number (%) symptom worsening (IPSS worse \geq4 points) by baseline PSA</p>	<p>Group 1: PSA <2.3: 22/248 (8.9%) PSA 2.3-3.9: 33/261 (12.6%) PSA >3.9: 32/228 (14.8%) P=NS</p> <p>Group 2: PSA <2.3: 36/242 (14.9%) PSA 2.3-3.9: 49/237 (20.7%); PSA >3.9: 39/264 (14.0%) P=NS</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Roehrborn et al., 1999²⁵⁶</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 years</p>	<p>Patient group: men with clinical BPH diagnosed on the basis of moderate to severe symptoms.</p> <p>Setting: 95 centres (Finasteride Long-Term Efficacy & Safety Study Group)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Moderate to severe symptoms Peak flow rate <15 mL/s with voided volume \geq 150 mL Enlarged prostate by digital rectal examination Serum PSA 4 -9.9 ng/mL with negative biopsy <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Current therapy of α-blocking agents or anti-androgens History of chronic prostatitis Recurrent urinary tract infections Surgery for prostate or bladder cancer Serum PSA >10ng/mL <p>All patients N: 3040 Drop outs: 1157</p> <p>Group 1 N: 1524 Age (mean \pm SD): 64 \pm 7 Quasi-AUA: 15 \pm 6 Serum PSA (ng/mL): 2.8 \pm 2.1 (n=1512)* 1st tertile PSA (ng/mL): 0.83 \pm 0.3 (n=472) 2nd tertile PSA (ng/mL): 2.21 \pm 0.6 (n=</p>	<p>Group 1 Finasteride (Proscar) 5mg 1/day</p> <p>Group 2 Placebo</p> <p>Assessment: 1 month single blind placebo run in after which randomisation and baseline measurements performed</p> <p>Quasi AUA symptom score (1-34), adverse events, urinary flow were assessed every 4 month. PSA was measured at baseline and every 4 months in year 1 and every 8 months thereafter. Physical examinations and routine haematological and serum chemistry tests performed yearly. MRI to determine prostate volume performed at baseline and yearly in a subset of 10% of patients</p>	<p>Mean Change in Quasi-AUA Symptom Score (\pm SE) v baseline PSA at 4 years</p> <p>Within tertile group and between treatment group analysis of variance performed to compare effect of baseline PSA and prostate volume on symptom changes over time</p> <p>Mean Change in Quasi-AUA Symptom Score (\pm SE) over time (years 1-4) for each PSA tertile in placebo patients (group 2)</p> <p>Mean Change in Quasi-AUA Symptom Score (\pm SE) over time (years 1-4) for each PSA tertile group 1 v group 2</p>	<p>1st Tertile Group 1: -3.2 \pm 0.4 Group 2: -2.4 \pm 0.3 Group1 v Group 2 p=0.128 Not sig. (ANOVA)</p> <p>2nd Tertile Group 1: -3.4 \pm 0.3 Group 2: -0.4 \pm 0.4 Group1 v Group 2 p<0.001 (ANOVA)</p> <p>3rd Tertile Group 1: -3.4 \pm 0.3 Group 2: -0.2 \pm 0.4 P Group1 v Group 2 p<0.001 (ANOVA)</p> <p>1st tertile had a significantly better long-term symptom improvement than those in other tertiles p < 0.001 There was no significant difference between long term symptom improvement between 2nd and 3rd tertiles p=0.65</p> <p>1st tertile Not sig. 2nd tertile (p=0.004) 3rd tertile (p=0.001)</p>	<p>Funding: Merck & Co., Inc.</p> <p>Limitations: No adjustment mentioned and no regression analysis</p> <p>Additional outcomes:</p> <ul style="list-style-type: none"> Mean Change in Quasi-AUA Symptom Score (\pm SE) v baseline prostate volume tertile at 4 years Mean Change in Quasi-AUA Symptom Score (\pm SE) v PSA tertile over time Mean Change in Quasi-AUA Symptom Score (\pm SE) v prostate volume tertile over time Mean Change in Qmax (\pm SE) v PSA tertile over time Mean Change in Qmax (\pm SE) v prostate volume tertile over time <p>Notes: Baseline PSA was divided into 3 tertiles: First (0.2 - 1.3) Second (1.4 - 3.2) Third (3.3 - 12.0)</p> <p>Quasi AUA symptom score: Had all components of the</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>536) 3rd tertile PSA (ng/mL): 5.39 ± 1.7 (n=504) Qmax (mL/s): 11 ± 4 Prostate Volume (mL): 54 ± 25 (n=157) Drop outs: 524</p> <p>Group 2 N: 1516 Age (mean \pm SD): 64 ± 6 Quasi-AUA: 15 ± 6 Serum PSA (ng/mL): 2.8 ± 2.1 (n=1498)* 1st tertile PSA (ng/mL): 0.86 ± 0.3 (n=511) 2nd tertile PSA (ng/mL): 2.24 ± 0.6 (n=514) 3rd tertile PSA (ng/mL): 5.36 ± 1.7 (n=473) Qmax (mL/s): 11 ± 4 Prostate Volume (mL): 55 ± 26 (n=155) Drop outs: 633</p>				<p>AUA score but the score differed from AUA per question: 0-5 for six questions and 0-4 for one question. Total 0-34</p> <p>*Patients numbers quoted for baseline characteristics were different in Roehborn 1999 paper from original study report McDonnell et al 1998 (NEJM).</p>

Study details	Patients	Outcomes	Analysis conducted	Results	Comments																								
<p>Tubaro et al., 2004²⁹⁸</p> <p>Study design: Cross sectional, observational</p> <p>Duration of follow-up: Nil</p>	<p>Patient group: Men with LUTS, ambulatory</p> <p>Setting: 45 urological centres in Italy between Feb 1998 and Jan 1999</p> <p>Interventions: Not applicable</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> - Age: 50-80 years - Persistent LUTS/BPH and BPE (as estimated by DRE) - Minimal voided volume (VV) of 150ml <p>Exclusion criteria:</p> <ul style="list-style-type: none"> - Associated urological diseases, psychiatric or mental illness, previous surgical or minimally invasive treatments of BPH, indwelling catheter, - Pharmacological treatments (e.g. tricyclic antidepressants, anticholinergic and sympathomimetic drugs) - Current or previous treatment for LUTS/BPH (e.g. alpha adrenoreceptor antagonists, finasteride, plant extracts) <p>All patients N: 866 M/F: 866/0 Age (mean, range): 64(50-80) Drop outs: 64/866, 802 analysed, dropouts are due to missing data Mean duration of LUTS: 30.2 months, median 24 months</p>	<table border="1"> <tr> <td>Age (range) (years):</td> <td>66.3 (44.8-89.7)</td> </tr> <tr> <td>PSA (ng/ml):</td> <td>2.23±2.36</td> </tr> <tr> <td>IPSS:</td> <td>13.4 ±6.1</td> </tr> <tr> <td>- Voiding</td> <td>7.6±4.4</td> </tr> <tr> <td>- Storage</td> <td>5.8±2.9</td> </tr> <tr> <td>Prostate volume, PV (cm³)</td> <td>34.5±18.8</td> </tr> <tr> <td>Uroflowmetry</td> <td></td> </tr> <tr> <td>Qmax (ml/s)</td> <td>13.6±6.6</td> </tr> <tr> <td>Qave (ml/s)</td> <td>6.8±3.7</td> </tr> <tr> <td>Flow time(s)</td> <td>46.3±27.3</td> </tr> <tr> <td>VV(ml)</td> <td>265.9±123.4</td> </tr> <tr> <td>Post void volume, PVR (ml)</td> <td>58.3±72.6</td> </tr> </table>	Age (range) (years):	66.3 (44.8-89.7)	PSA (ng/ml):	2.23±2.36	IPSS:	13.4 ±6.1	- Voiding	7.6±4.4	- Storage	5.8±2.9	Prostate volume, PV (cm³)	34.5±18.8	Uroflowmetry		Qmax (ml/s)	13.6±6.6	Qave (ml/s)	6.8±3.7	Flow time(s)	46.3±27.3	VV(ml)	265.9±123.4	Post void volume, PVR (ml)	58.3±72.6	<p>Multiple logistic regressions: IPSS >7 vs. PSA (ng/ml), IPSS <7 is the reference</p>	<p>Odds ratio (95%CI) PSA ≤2: 1.0 PSA >2-4: 1.62(1.2-2.2) PSA >4-10: 2.64 (1.5-4.7) PSA >10: 4.28 (1.8-10.3) ≤2</p>	<p>Funding: not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> - Cross sectional study - Answers the questions of association of PSA vs. IPSS, rather than ability of PSA to predict IPSS over time (prognosis) <p>Additional outcomes: Logistic regression of IPSS vs. prostate related variables- PVR, PV, Qmax, Abrams-Griffiths number etc..</p> <p>Notes: - All values reported were mean ± standard deviation unless otherwise specified</p>
Age (range) (years):	66.3 (44.8-89.7)																												
PSA (ng/ml):	2.23±2.36																												
IPSS:	13.4 ±6.1																												
- Voiding	7.6±4.4																												
- Storage	5.8±2.9																												
Prostate volume, PV (cm³)	34.5±18.8																												
Uroflowmetry																													
Qmax (ml/s)	13.6±6.6																												
Qave (ml/s)	6.8±3.7																												
Flow time(s)	46.3±27.3																												
VV(ml)	265.9±123.4																												
Post void volume, PVR (ml)	58.3±72.6																												

Evidence Table 3 Diagnostic accuracy of uroflowmetry

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
<p>Ref ID: Oelke et al., 2007²³¹</p> <p>Study design: Cross-sectional study</p> <p>Evidence level: Level-2 study (II)</p> <p>Duration of follow-up: 1-3 weeks duration between the index test and the gold standard</p>	<p>Patient group: Men with LUTS, clinical BPH and/or prostate volume >25ml</p> <p>Setting: single centre – urologic outpatient clinic - Germany</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> > 40 years with LUTS, clinical BPH and/or prostate volume >25ml <p>Exclusion criteria: Patients with:</p> <ul style="list-style-type: none"> Prostate cancer Acute urinary retention Neurological disease Previous prostatic or urethral surgery Medication treating BPH α- blockers, α-reductase inhibitors <p>All patients N: 160 Age median (range): 62 (40-89) Drop outs: 0</p>	<p>Assessment tool under investigation: Uroflowmetry – number of voids not specified.</p> <p>Gold standard: Pressure flow studies (PFS) performed using Ellipse (Andromeda) machine with CHESSE used to classify obstruction</p>	<p>Qmax threshold < 10 mL/s</p> <p>Sensitivity 68% (51/75) CI95% 57 - 79 Specificity 73% (62/85) CI95% 64 - 82 Positive predictive value 69% (51/74) Negative predictive value 72% (62/86) Prevalence 47% 75/160 Positive Likelihood Ratio 2.51 Negative Likelihood Ratio 0.44 Pre-test Odds (CI 95%) 0.88 (CI95%: 0.81-0.96) Post-Test Odds +ve result 2.22 Post-Test Odds -ve result 0.39</p> <p>Qmax threshold < 15 mL/s</p> <p>Sensitivity 99% (74/75) CI95% 97 - 100 Specificity 39% (33/85) CI95% 29 - 49 Positive predictive value 59% (74/126) Negative predictive value 97% (33/34) Prevalence 47% 75/160 Positive Likelihood Ratio 1.61 Negative Likelihood Ratio 0.03 Pre-test Odds (CI 95%) 0.88 (CI95%: 0.81-0.96) Post-Test Odds +ve result 1.42 Post-Test Odds -ve result 0.03</p>	<p>Funding: NR</p> <p>Limitations: Details of Uroflowmetry methods not reported</p> <p>1-3 week delay between</p> <p>Uroflowmetry as index test and PFS</p> <p>No mention whether the procedures tested were conducted by the same investigator(s)</p> <p>Additional outcomes: This study also reports Detrusor Wall Thickness measured by 7.5 MHz ultrasound, Post Void Residual measured with 3.5 MHz ultrasound. Prostate Volume measured with TRUS</p> <p>Notes: None</p>	

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
Ref ID: Poulsen et al., 1994 ²⁴¹ Study design: Cross-sectional study Evidence level: Level-2 study (II) Duration of follow-up: NA	Patient group: Men with symptomatic BPH (94% uncomplicated), 5% also with recurrent urinary tract infection and 1% with previous AUR Setting: single centre Denmark Exclusion criteria: NR All patients N: 188 Age median (range): 68 (32-90) Drop outs: Free flow missing for 35/188 (19%) and PFS data missing for 5/188 (3%)	Assessment tool under investigation: Void into Dantec Urodyn 1000 uroflowmeter. Number of voids not reported Gold standard: Pressure flow studies (PFS) performed using Dantec Urodyn 1000 uroflowmeter after filling with Foley 14F catheter. Patients characterised for BOO using Abrams-Griffiths nomogram.	Qmax threshold < 10 mL/s Sensitivity 69% (68/99) CI95%: 59 - 78 Specificity 57% (31/54) CI95%: 44 - 70 Positive predictive value 75% (68/91) Negative predictive value 50% (31/62) Prevalence 65% (99/153) Positive Likelihood Ratio 1.61 Negative Likelihood Ratio 0.55 Pre-test Odds (CI 95%) 1.83 (CI95%: 1.76 -1.91) Post-Test Odds +ve result 2.96 Post-Test Odds -ve result 1.00	90% (89/99) CI95%: 84 - 96 Specificity 31% (17/54) CI95%: 19 - 43 Positive predictive value 71% (68/91) Negative predictive value 63% (31/62) Prevalence 65% (99/153) Positive Likelihood Ratio 1.31 Negative Likelihood Ratio 0.32 Pre-test Odds (CI 95%) 1.83 (CI95%: 1.76 -1.91) Post-Test Odds +ve result 2.41 Post-Test Odds -ve result 0.59	Funding: NR Limitations: Masking of assessors to test results NR Not clear whether tests were independent (implies PFS before entry into study) Number of voids NR Additional outcomes: DAN-PSS Symptom Score also recorded Notes: None
			Qmax threshold < 15 mL/s Sensitivity 90% (89/99) CI95%: 84 - 96 Specificity 31% (17/54) CI95%: 19 - 43 Positive predictive value 71% (68/91) Negative predictive value 63% (31/62) Prevalence 65% (99/153) Positive Likelihood Ratio 1.31 Negative Likelihood Ratio 0.32 Pre-test Odds (CI 95%) 1.83 (CI95%: 1.76 -1.91) Post-Test Odds +ve result 2.41 Post-Test Odds -ve result 0.59		

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
<p>Ref ID: Reynard et al., 1996²⁴⁸</p> <p>Study design: Cross-sectional study</p> <p>Evidence level: Level-2 study (II)</p> <p>Duration of follow-up: NA</p>	<p>Patient group: Men > 45 years with) LUTS suggestive of benign prostatic obstruction (BPO)</p> <p>Setting: 2 centres UK</p> <p>Exclusion criteria: Patients with:</p> <ul style="list-style-type: none"> Prostate cancer (DRE + TRUS) Diabetes Lower urinary tract infection Previous prostatic or urethral surgery Medication affecting lower urinary tract <p>All patients N: 165</p> <p>Age median (range): 68 (50-84)</p> <p>Drop outs: PFS data missing for 8/165 (5%) patients</p>	<p>Assessment tool under investigation: Uroflowmetry 4 voids into Dantec Urolyn 1000 uroflowmeter. Qmax below threshold indicates BOO</p> <p>3 voids: 17 (10%) 4 voids: 148 (90%)</p> <p>Gold standard: Pressure flow studies (PFS) performed using Dantec Menuet or Dantec 5500 multichannel recorder. Patients characterised for BOO using Abrams-Griffiths nomogram as obstructed or equivocal/unobstructed.</p>	<p>*Qmax threshold < 10 mL/s</p> <p>Sensitivity 49% (47/95) CI95% 39 - 59</p> <p>Specificity 87% (54/62) CI95% 79 - 95</p> <p>Positive predictive value 85%</p> <p>Negative predictive value 53%</p> <p>Prevalence 61% (95/157)</p> <p>Positive Likelihood Ratio 3.83</p> <p>Negative Likelihood Ratio 0.58</p> <p>Pre-test Odds (CI 95%) 1.53 (CI95%:1.46 -1.61)</p> <p>Post-Test Odds +ve result 5.88</p> <p>Post-Test Odds -ve result 0.89</p>	<p>Funding: NR</p> <p>Limitations: No indication of who carried out the tests-whether by the same people, or whether the investigator or patients were masked to the results of other tests.</p> <p>Results of individual centres not compared, and inter-rater agreement (presumably tests in different tests done by different people) was not addressed</p> <p>Notes: *Qmax taken as highest value on voids 1 & 2. Also reported < 8 mL/s</p> <p>Study suggests increasing specificity and decreasing specificity with increasing number of voids</p>	
			<p>*Qmax threshold < 12 mL/s</p> <p>Sensitivity 65% (62/95) CI95% 55 - 75</p> <p>Specificity 74% (46/62) CI95% 79 - 95</p> <p>Positive predictive value 79%</p> <p>Negative predictive value 58%</p> <p>Prevalence 61% (95/157)</p> <p>Positive Likelihood Ratio 2.53</p> <p>Negative Likelihood Ratio 0.47</p> <p>Pre-test Odds (CI 95%) 1.53 (CI95%:1.46 -1.61)</p> <p>Post-Test Odds +ve result 3.88</p> <p>Post-Test Odds -ve result 0.72</p>		
			<p>*Qmax threshold < 15 mL/s</p> <p>Sensitivity 85% (81/95) CI95% 78 - 92</p> <p>Specificity 53% (33/62) CI95% 63 - 85</p> <p>Positive predictive value 74%</p> <p>Negative predictive value 70%</p> <p>Prevalence 61% (95/157)</p> <p>Positive Likelihood Ratio 1.82</p> <p>Negative Likelihood Ratio 0.38</p> <p>Pre-test Odds (CI 95%) 1.53 (CI95%:1.46 -1.61)</p> <p>Post-Test Odds +ve result 2.79</p> <p>Post-Test Odds -ve result 0.42</p>		

Evidence Table 4: Diagnostic accuracy of post void residual

Study details	Patients	Diagnostic tools	Measure of Disorders	Results	Comments
<p>Ref ID: REYNARD1998 (ICS- 'BPH' study)</p> <p>Study design: Cross-sectional study</p> <p>Evidence level: Level-2 study (II)</p> <p>Duration of follow-up: NA</p>	<p>Patient group: Men with LUTS and benign prostatic enlargement (BPE)</p> <p>Setting: multi-centre 12 centres in Europe, Australia, Canada, Taiwan & Japan</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> > 45 years Symptoms of BOO secondary to BPH <p>Exclusion criteria: Patients with:</p> <ul style="list-style-type: none"> Prostate cancer Neurological disease Previous prostatic or urethral surgery Medication affecting lower urinary tract <p>All patients N: 1271</p> <p>Age mean (range): 66.5 (45-88)</p> <p>Drop outs: Uroflowmetry data missing for 81/1271 (6%) PFS data missing for 338/1271 (27%)</p>	<p>Assessment tool under investigation: Uroflowmetry 3 voids 1 void: 211 (17%) 2 voids: 443 (35%) 3 voids: 537 (42%) Details of technique not reported</p> <p>Gold standard: Pressure flow studies (PFS) performed according to International Continence Society guidelines with diagnosis of BOO using Schafer classification Ratings 0-2 categorised as non-obstructive while 3-6 were obstructed. Definition of Schaefer method: 0 no obstruction, 1 slightly obstructed, 2-6 obstructed with increasing severity</p>	<p>*Qmax threshold < 10 mL/s</p> <p>Sensitivity 47% (252/540) CI95% 42 - 51 Specificity 70% (250/357) CI95% 65 - 75 Positive predictive value 70% (252/359) Negative predictive value 46% (250/538) Prevalence 60% 540/897 Positive Likelihood Ratio 1.56 Negative Likelihood Ratio 0.76 Pre-test Odds (CI 95%) 1.51 (CI95%:1.48 -1.54) Post-Test Odds +ve result 2.36 Post-Test Odds -ve result 1.15</p> <p>*Qmax threshold < 15 mL/s</p> <p>Sensitivity 81% (440/540) CI95% 78 - 85 Specificity 38% (136/357) CI95% 33 - 43 Positive predictive value 67% (440/661) Negative predictive value 58% (136/236) Prevalence 60% 540/897 Positive Likelihood Ratio 1.32 Negative Likelihood Ratio 0.49 Pre-test Odds (CI 95%) 1.51 (CI95%:1.48 -1.54) Post-Test Odds +ve result 1.99 Post-Test Odds -ve result 0.74</p>	<p>Funding: International Continence Society (ICS)</p> <p>Limitations: No information provided about the specific protocol followed in carrying out tests, who carried them out, whether they were blinded and also interval between the tests.</p> <p>Notes: *Qmax taken as highest value for each patient from voids</p>	

See Evidence Table 3 Diagnostic accuracy of uroflowmetry for Oelke et al., 2007²³¹.

Evidence Table 5: Pelvic floor exercises (with or without electrical stimulation or biofeedback)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bales et al., 2000²²</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months after surgery</p> <p>Outcome assessment was masked</p>	<p>Patient group: Men with stages T1c-T2c prostate cancer who were to undergo radical retropubic prostatectomy by a single surgeon</p> <p>Inclusion criteria: Men with stages T1c-T2c prostate cancer who were to undergo radical retropubic prostatectomy by a single surgeon. None of the men had undergone transurethral resection of the prostate or had pre-existing neurologic disease.</p> <p>Exclusion criteria: See above, exclusion criteria not specifically stated.</p> <p>All patients N: 100 Drop outs: 3</p> <p>Group 1: N: 50 Age (mean): 59.3 Drop outs: 3</p> <p>Group 2: N: 50 Age (mean): 60.9 Drop outs: 0</p>	<p>Group 1: Biofeedback 45-minute session with a nurse trained in biofeedback techniques 2 to 4 weeks prior to radical prostatectomy. Patients instructed how to perform graded PFMT using biofeedback. Surface electrodes were used to assess muscle strength and contractions of 5 to 10 seconds, and 10 to 15 repetitions were performed. Patients advised to practice these exercises 4/day until their surgery.</p> <p>Group 2: Control Patients underwent radical prostatectomy without any biofeedback training. These patients received only written and brief verbal instructions on how to perform PFMT to isolate the muscle that starts and stops urine flow and to practice contractions 4/day with 10 to 15 repetitions. Patients were given written instructions and briefly reviewed these instructions with a nurse.</p> <p>All patients: Postoperatively, the urethral catheter was removed approximately 2 weeks following surgery in both groups. Patients in both groups were encouraged to perform pelvic muscle strengthening exercises 4/day after catheter removal. No patient in either group received adjuvant radiation therapy or hormonal therapy within 6 months following surgery.</p>	<p>Incidence of urinary continence at 6 months post op.</p>	<p>Group 1: 44/47 (94%) Group 2: 48/50 (96%) p value: 0.60</p>	<p>Funding: NR</p> <p>Limitations: This study is poorly reported: Method of randomisation and allocation concealment not described, there is insufficient information about patients' baseline characteristics, no description of sample size calculation. Assessments methods could be unreliable. Other limitations stated by authors:</p> <ul style="list-style-type: none"> - no effort was made to assess pelvic muscle floor strength prior to surgery - incidence of incontinence in Group 2: was very low - patients received only one preoperative biofeedback session. - subtle differences in results might have been detected if more rigorous measures of incontinence had been used, such as weighted pad testing. No objective measurement of continence was used. <p>Notes: Patients wearing one pad or less per day were considered to be continent. Those using two or more pads per day were considered incontinent.</p>
			<p>Incidence of urinary continence at 3 months post op</p>	<p>Group 1: 27/47 Group 2: 31/50 p value: 0.64</p>	
			<p>Proportion of still incontinent at 3 months (ITT analysis)</p>	<p>Group 1: 23/50 Group 2: 19/50 p value: NR</p>	
			<p>Proportion of still incontinent at 6 months (ITT analysis)</p>	<p>Group 1: 6/50 Group 2: 2/50 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Burgio et al., 2006³⁹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months post surgery</p>	<p>Patient group: Men elected for radical prostatectomy for prostate cancer</p> <p>Setting: single centre university urology clinic(USA)</p> <p>Inclusion criteria: Ambulatory and continent</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • If reporting > 2 episodes of urinary incontinence in past 6 months • Had documented incontinence in a bladder diary • Previous prostatectomy • Mental impaired status (<20 on the Mini-Mental State Examination) • <1 week before scheduled surgery <p>All patients N: 112 Age (mean ± SD): 60.9 ± 6.9 Drop outs: 0</p> <p>Group 1 N: 57* Age (mean ± SD): 60.7 ± 6.6 M: 57 Black: 13 Previous TURP: 2 Drop outs: 0</p> <p>Group 2 N: 55* Age (mean ± SD): 61.1 ± 7.2</p>	<p>Group 1 Single session of preoperative biofeedback enhanced behavioural training on pelvic floor muscle control and instructions on daily PMFT. Rectal probe used to provide feedback of rectal pressure. Daily practice 3 x 15 exercises. Also instructed to interrupt stream when voiding. Postoperatively patients were reminded to resume exercise regimen</p> <p>Group 2 Brief instructions on how to interrupt stream when voiding and usual care.</p> <p>All patients Instructed on use of bladder diaries and use of pads to record incontinence. Patients sent a weekly bladder diary to investigators during follow up.</p> <p>Patients were contacted for follow-up at 6 weeks, 3 and 6 months after surgery. They completed patient questionnaire on bladder control, 7-day bladder diary, QoL score, and</p>	<p>Proportion of patients with severe/continual leakage at 6 months</p>	<p>Group 1: 3/50 (6%) Group 2: 9/47 (19%) p value: 0.04 (Chi squared) not ITT NCGC Chi-squared calculation p=0.058 using ITT</p>	<p>Funding: National Institute for Diabetes and Digestive Kidney Diseases, National Institute of Health</p> <p>Limitations: There were significantly more men in the control group with preserved urethral length. P=0.03 favouring continence. At 6 months data was not presented as an ITT analysis</p> <p>Notes: Bladder diaries were scored by an individual kept blind to group assignment. Those performing intervention were blinded to next group assignment. Randomisation by computer. Kaplan-Meier data extraction by Hunter et al., 2007¹²³ et al Cochrane review</p>
			<p>Number of patients wearing pads at 6 months</p>	<p>Group 1: 16/50 (32%) Group 2: 24/46 (52%) p value: <0.05 not ITT NCGC Chi-squared calculation p=0.086 using ITT</p>	
			<p>Mean days ± SD with no leakage at 6 months</p>	<p>Group 1: 72.6 ± 0.39 Group 2: 54.2 ± 0.47 p value: 0.04 not ITT NCGC t-test with equal variance test calculation p<0.00001 using ITT</p>	
			<p>Kaplan-Meier survival curve of proportion of still incontinent at < 3 months (data from Hunter et al., 2007¹²³)</p>	<p>Group 1: 49/54 Group 2: 51/53 p value: 0.25 (NCGC Chi-squared calculation – not ITT)</p>	
			<p>Kaplan-Meier survival curve of proportion of still incontinent at 3 - 6 months (data from Hunter et al., 2007¹²³)</p>	<p>Group 1: 32/53 Group 2: 40/51 p value: 0.046 (NCGC Chi-squared calculation – not ITT)</p>	
			<p>Kaplan-Meier survival curve of proportion of still incontinent at 6 - 12 months (data from Hunter et al., 2007¹²³)</p>	<p>Group 1: 22/51 Group 2: 30/50 p value: 0.09 (NCGC Chi-squared calculation – not ITT)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>M: 55 Black: 18 Previous TURP: 1 Drop outs: 0</p> <p>* excludes patients with cancelled operations</p>	<p>Incontinence Impact Questionnaire modified for men.</p> <p><i>Continence defined as 3 consecutive weekly bladder diaries returned with no leakage.</i></p>			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Filocamo et al., 2005⁹¹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: men undergoing retropubic radical prostatectomy for localised prostate cancer</p> <p>Setting: urology clinic, University of Florence, Italy</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Prior bladder or prostate surgery • Prior urinary or faecal incontinence • Neurogenic dysfunction of lower urinary tract • Preoperative history of overactive bladder <p>All patients N: 300 Age (mean ± SD): NR Drop outs: 0</p> <p>Group 1 N: 150 Age (mean ± SD): 65 ± 4.79 (51-75) M: 150 Mean preop PSA (ng/ml): 8.13 Drop outs: 0</p> <p>Group 2 N: 150 Age (mean ± SD): 66.8 ± 5.33 (45-75) M: 150 Mean preop PSA (ng/ml): 8.11 Drop outs: 0</p>	<p>Group 1 In 1st treatment session PFMT was taught using verbal and visual feedback. Strength of muscles evaluated by digital anal control. Patients instructed to perform 3x10 sets/day at home for 6 months. In 2nd treatment session PMFT taught in all positions and patients asked to identify movements causing incontinence. Patients asked to practice new exercises at home for 7 days. At 3rd treatment session patients asked to practise PFMT before any activity that may cause incontinence.</p> <p>Group 2 No treatment</p> <p>All patients Asked to complete a bladder diary and counselled to prevent leakage by increasing frequency of micturation.</p> <p>All patients were assessed at 1,3,6 and 12 months. Incontinence was assessed objectively using 1h and</p>	<p>Proportion of patients still incontinent at 1 month (using subjective ICS male questionnaire)</p>	<p>Group 1: 121/150 (81%) Group 2: 138/150 (92%) p value: NR NCGC Chi-squared calculation p=0.004 using ITT analysis signif.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method not described • Masking of outcome assessment not mentioned • Proportion of patients still incontinent reported as subjective measurement using ICS questionnaire <p>Additional outcomes: Correlation between patient age and continence at each time interval</p> <p>Notes: Study reports numbers of patients continent at time intervals but data are presented as number of patients <i>still incontinent</i></p>
			<p>Proportion of patients still incontinent at 3 months (using subjective ICS male questionnaire)</p>	<p>Group 1: 39/150 (26%) Group 2: 105/150 (70%) p value: NR NCGC Chi-squared calculation p<0.00001 using ITT analysis signif.</p>	
			<p>Proportion of patients still incontinent at 6 months (using subjective ICS male questionnaire)</p>	<p>Group 1: 6/150 (4%) Group 2: 53/150 (35%) p value: NR NCGC Chi-squared calculation p<0.00001 using ITT analysis signif.</p>	
			<p>Proportion of patients still incontinent at 12 months (using subjective ICS male questionnaire)</p>	<p>Group 1: 2/150 (1%) Group 2: 18/150 (12%) p value: NR NCGC Chi-squared calculation p=0.0002 using ITT analysis signif.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
		<p>24h pad test – number of pads used daily. Subjective assessment by completion of International Continence Society (ICS) questionnaire. All patients still incontinent at 6 months underwent urodynamic evaluation</p> <p><i>Continence defined as 1 precautionary pad</i></p>			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Floratos et al., 2002⁹³</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients undergoing radical retropubic prostatectomy for localised prostate cancer.</p> <p>Setting: multi-centre. Greece and Netherlands</p> <p>Inclusion criteria: Patients with objectively confirmed urinary incontinence, no significant perioperative complications (ureteric or rectal injury, urine leakage from anastomosis, thrombo-embolism), no history of preoperative incontinence and pelvic or lower urinary tract operations, no psychiatric history, a recognised ability to participate in a learning programme, good general condition and willingness to participate in the study.</p> <p>All patients N: 42 Age (mean ± SD): Drop outs: Unclear</p> <p>Group 1: N: 28 Age (mean ± SD): 63.1 +/- 4 Received Oxybutynin: n=3</p> <p>Group 2: N: 14 Age (mean ± SD): 65.8 +/- 4.3 Received Oxybutynin: n=2</p>	<p>Group 1: Biofeedback Patients referred to a specialist in physical therapy and rehabilitation to have 15 sessions of electromyographic (EMG) biofeedback (2 channel Totem Biofeedback, BEAC, Italy) 3/week of 30 min duration each. During the initial 2/3 sessions, a strong emphasis was placed on the specificity of muscle contraction. During the sessions the exercises were designed to increase the power, endurance and coordination of the pelvic floor muscles. In parallel, patients practised 50-100 exercises daily at home.</p> <p>Group 2: Control Patients were taught how to contract their pelvic muscles without contracting abdominal muscles simultaneously. Patient was placed in the lateral decubitus position and the instructor inserted index finger into patient's rectum to check for simultaneous contraction whilst palpating the abdominal muscles. Verbal feedback used to instruct the patient how to correctly and selectively contract the anal sphincter while. Patients received an informative leaflet with these instructions. Home practise comprised 80-100 exercises daily, divided in four sessions of 20-25 exercises each. The duration of each constriction was 3-5 s with submaximal strength (70%) and relaxation period of 6-10 s between the exercises. Initially patients practised these exercises while supine but later when sitting and standing. After the first month patients were encouraged to practise the exercises during normal daily activities, including movements that provoked incontinence.</p>	<p>Mean urine loss as assessed by the 1-h pad test Patients were evaluated at 1,2, 3 and 6 months of treatment using 1-h pad test. For the best intra- and inter-patient estimates in the pad test, a special type of 'pocket pad' was used which covered only the penis, thus reducing the interference from sweat on the pad weight gained during the test.</p> <p>Mean no. pads/ day Patients were evaluated subjectively with a questionnaire (to determine the number and extent of incontinence episodes, number of pads used per day, and any LUTS).</p> <p>Number of men still incontinent at 3-6 months (data from Hunter et al., 2007¹²³)</p>	<p>Group 1: Baseline: 39 g 1st month: 18 g 2nd month: 7 g 3rd month: 4 g 6th month: 3 g</p> <p>Group 2: Baseline: 31 g 1st month: 11 g 2nd month: 3 g 3rd month: 1 g 6th month: 0 g</p> <p>P value > 0.05</p> <p>Group 1: Baseline: 3.9 1st month: 3.4 2nd month: 1.2 3rd month: 0.8 6th month: 0.4</p> <p>Group 2: Baseline: 3.6 1st month: 1.8 2nd month: 0.9 3rd month: 0.4 6th month: 0.2</p> <p>P value > 0.05</p> <p>Group 1: 4/28 Group 2: 0/14</p>	<p>Funding: NR</p> <p>Limitations: Randomisation and allocation concealment is not described. There is insufficient information about patients' baseline characteristics, no description of sample size calculation. Masking of outcome assessment is not reported.</p> <p>Additional outcomes: No additional outcomes reported</p> <p>Notes: All patients: During the study, patients with irritative symptoms and a negative urine culture received empirical anticholinergic medication (oxybutynin).</p> <p>Continence defined as <1 g loss / 1 hour pad test or < 2 pads per day</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Franke at al., 2000 ⁶ Study design: RCT Evidence level: 1+ Duration of follow-up: 24 weeks (6 months)	Patient group: Incontinent men after radical prostatectomy Setting: Urology department, Vanderbilt Medical Centre, Tennessee, USA Inclusion criteria: 2 weeks post prostatectomy Exclusion criteria: <ul style="list-style-type: none"> • Previous TURP • Neurological condition affecting the urinary tract. • Men with residual urine greater than 50ml or urinary tract infection were excluded at 6 week visit. All patients N: 30 Drop outs: 5 withdrew after randomisation Group 1 N: 15 Age (mean): 62.3 Dropouts: At 3 months= 2, 6 months= 8 Group 2 N: 15 Age (mean): 60.7 Drop outs: 3 months: 5, 6 months: 7	Group 1 45 minute biofeedback behavioural therapy session 6, 7, 9, 11 and 16 weeks postoperatively. Perineal patch electromyography biofeedback was performed using abdominal electromyography leads to ensure proper isolation. Patients instructed to continue pelvic floor muscle exercises at home (20 contractions 3 times a day). A timed voiding schedule was encouraged and patients instructed in techniques to decrease urgency and urge incontinence. Group 2 No instruction and asked to return voiding diary and 48 hour pad test at the routine follow-up visits. All patients: Urinalysis and post void residual urine volume tests at 6 week visit. Completed voiding diary and 48 hour pad test at 6, 12 and 24 weeks postoperatively.	Number still incontinent at 3 months	Group 1: 6/13 (46%) Group 2: 3/10 (30%) P value: NR NCGC Chi-squared calculation p=0.23 using ITT analysis Not sig.	Funding: NR Limitations: <ul style="list-style-type: none"> • Randomisation method not described • Masking of outcome assessment not mentioned • Not an ITT analysis Additional outcomes: Improvement in pelvic muscle work using electromyography training effect (only assessed in intervention group). Notes: Study reports number of patients continent at time intervals but data are presented as number of patients <i>still incontinent</i> . Incontinent defined as still using pads in the study.
			Number still incontinent at 6 months	Group 1: 1/7 (14%) Group 2: 1/8 (12%) P value: NR NCGC Chi-squared calculation p=1.00 using ITT analysis Not sig.	
			Mean incontinence (gm/24hours) using pad tests	At 6 weeks Group 1: 162 Group 2: 152, p value: 0.91(CI95%: 193-214) At 3 months: Group 1: 58 Group 2: 93, p value: 0.67(CI95%: 199-128) At 6 months: Group 1: 8 Group 2: 62, p value: 0.41(CI95%: 200-90)	
			Mean incontinent episodes/day (mean voiding diary differences)	At 6 weeks Group 1: 7.2 Group 2: 5.2, p value: 0.48 (-3.7-7.7) At 3 months: Group 1: 1.3 Group 2: 0.8, p value: 0.38 (-0.7-1.6) At 6 months: Group 1: 0.3 Group 2: 0.1, p value: 0.45 (-0.3-0.6)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Manassero et al., 2007¹⁸⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p> <p>Masked outcome assessment and computer generated random numbers</p>	<p>Patient group: men undergoing retropubic radical prostatectomy for localised prostate cancer</p> <p>Setting: urology clinic, University of Pisa, Italy</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Compliance with protocol clinic attendance Objectively confirmed urinary incontinence (>2g urine on 24h pad test) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> History of preoperative incontinence Significant perioperative complications Active rectal lesions or infections Psychiatric or neurological disorders Inability to contract pelvic floor muscles or weak contraction Detrusor over activity <p>All patients N: 107 Age (mean): M: 107 Drop outs: 13</p> <p>Group 1 N: 54 Age (mean ± SD): 66.8 ± 6.3 M: 54 Mean urine leakage/day: 247 ± 505g</p>	<p>Group 1 Pelvic floor muscle training programme by trained urologists with verbal feedback and measurement of muscle strength using digital anal control. Patients with weak muscles had additional electrical stimulation. Home practice 3x15 sessions/day increasing to 3x30 sessions in supine, sitting and standing positions. After 1 month patients were encourage to integrate exercise into daily life.</p> <p>Group 2 No treatment.</p> <p>All patients Assessed at 1 week and 1,3,6,9 and 12 months after catheter removal including a physical examination and IPSS score. At home patients weighed pads and residual incontinence assessed subjectively using visual analogue score (VAS) where 0=completely continent, 10=completely incontinent. Patients also filled out frequency volume charts</p> <p>Continence defined as <2g urine lost per day on 24h</p>	<p>Proportion of patients still incontinent at 1 month</p>	<p>Group 1: 45/54 (83%) Group 2: 39/40 (98%) p value: 0.04 (Fishers exact test) signif. NCGC Chi-squared calculation p=0.21 using ITT analysis Not sig.</p>	<p>Funding: NR</p> <p>Limitations: High drop out rate 13/53 (28%) in control group and results for control group are not presented as intention to treat (ITT) analysis</p> <p>Additional outcomes: Correlation between VAS score subjective assessment and 24h pad test at each time interval.</p> <p>Multivariate logistic regression to find variables that predict incontinence at 12 months (adjusting for age, IPSS score, blood loss, baseline QoL, incontinence at 1 week, tumour stage & nerve preservation)</p> <p>Notes: None</p>
			<p>Proportion of patients still incontinent at 3 months</p>	<p>Group 1: 29/54 (54%) Group 2: 31/40 (76%) p value: 0.03 (Fishers exact test) signif NCGC Chi-squared calculation p=0.61 using ITT analysis Not sig.</p>	
			<p>Proportion of patients still incontinent at 6 months</p>	<p>Group 1: 18/54 (33%) Group 2: 24/40 (60%) p value: 0.01 (Fishers exact test) signif NCGC Chi-squared calculation p=0.21 using ITT analysis Not sig.</p>	
			<p>Proportion of patients still incontinent at 12 months</p>	<p>Group 1: 9/54 (17%) Group 2: 21/40 (53%) p value: 0.0003 (Fishers exact test) signif NCGC Chi-squared calculation p=0.008 using ITT analysis signif.</p>	
			<p>Proportion of patients still incontinent at 12 months (incontinence severity)</p>	<p>Group 1: 1 mild (2-9g), 1 moderate (10-49g), 7 severe (≥50g) Group 2: 7 mild (2-9g), 10 moderate (10-49g), 4 severe (≥50g)</p>	
			<p>Subjective comparison of incontinence at 12 months using VAS score</p>	<p>Group 1: NR Group 2: NR p value: 0.01 (Wilcoxon Rank Sum Tets) signif</p>	
			<p>Subjective comparison of incontinence at 12 months</p>	<p>Group 1: NR Group 2: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Drop outs: 0</p> <p>Group 2</p> <p>N: 53</p> <p>Age (mean ± SD): 67.9 ± 5.5 (n=40)</p> <p>M: 53</p> <p>Mean urine leakage/day: 97 ± 138g</p> <p>Drop outs: 13 (social reasons and refusal to complete follow-up)</p> <p>Baseline data only available for 40 patients</p>		<p>months using Quality of Life (QoL) question from IPSS symptom score.</p>	<p>p value: 0.03 (Wilcoxon Rank Sum Tets) signif</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Mathewson-Chapman 1997¹⁸⁵</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Men with a radical retropubic prostatectomy (RP) for localised prostate cancer</p> <p>Setting: University of Florida College of Nursing</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Incontinent on day 15 after surgery after catheter removal Able to regularly attend hospital appointments <p>All patients N: 53 Age (mean): 62 (range 47-75) M: 53 Drop outs: 2 (unaccounted for in report)</p> <p>Group 1 N: 27 Age (mean): NR M: 27 Drop outs: NR</p> <p>Group 2 N: 24 Age (mean): NR M: 24 Drop outs: NR</p>	<p>Group 1 Preoperative education and instruction*</p> <p>Then postoperative Pelvic Muscle Exercise protocol (PME) practiced 3/week for 36 sessions starting at week 3. 15 repetitions performed at home, increasing by 10 every 4 weeks to a maximum of 35</p> <p>Biofeedback using an anal probe (PRS 8900 Incare). Evaluations were done at baseline, weeks 5, 12 and any other times requested by the patient.</p> <p>Group 2 Preoperative education and instruction*</p> <p>Postoperatively no intervention.</p> <p>Examination methods: Bladder diary was used to measure the number of pads used, number of episodes of incontinence /day over a 3 day period and frequency of episodes of urine loss. 24h pad test measured</p>	<p>Mean ± SD number of episodes of incontinence at week 2</p>	<p>Group 1: 25.1 ± 39.5 Group 2: 12.5 ± 26.3 p value: 0.17 (t test) Not sig.</p>	<p>Funding: In part by a Geriatric Nurse Fellowship from Dept. Veteran Affairs, USA</p> <p>Limitations:</p> <ul style="list-style-type: none"> The results from the intervention arm are potentially confounded by the preoperative instruction on pelvic floor muscle contraction given to both groups No allocation concealment No blinding Not an ITT analysis – report says 53 randomised but only 51 in patient groups. Drop outs not explained. <p>Notes: *Both groups were taught preoperatively how to contract perineal muscle prior to lifting, standing, coughing or sneezing and also to limit tea, coffee, chocolate and alcohol uptake.</p>
			<p>Mean ± SD number of episodes of incontinence at week 5</p>	<p>Group 1: 13.4 ± 31.1 Group 2: 10.4 ± 26.8 p value: 0.71 (t test) Not sig.</p>	
			<p>Mean ± SD number of episodes of incontinence at week 9</p>	<p>Group 1: 1.5 ± 3.2 Group 2: 5.6 ± 26.3 p value: 0.34 (t test) Not sig.</p>	
			<p>Mean ± SD number of episodes of incontinence at week 12</p>	<p>Group 1: 0.84 ± 1.99 Group 2: 1.00 ± 0.27 p value: 0.68 (t test) Not sig.</p>	
			<p>Mean ± SD number of pads used at week 2</p>	<p>Group 1: 3.88 ± 3.15 Group 2: 3.84 ± 3.3 p value: 0.95 (t test) Not sig.</p>	
			<p>Mean ± SD number of pads used at week 5</p>	<p>Group 1: 2.35 ± 2.97 Group 2: 2.84 ± 3.1 p value: 0.56 (t test) Not sig.</p>	
			<p>Mean ± SD number of pads used at week 9</p>	<p>Group 1: 1.1 ± 2.1 Group 2: 2.04 ± 2.7 p value: 0.2 (t test) Not sig.</p>	
			<p>Mean ± SD number of pads used at week 12</p>	<p>Group 1: 0.6 ± 1.6 Group 2: 1.8 ± 2.7 p value: 0.07 (t test) Not sig.</p>	
			<p>Mean ± SD time to continence - no pad needed (days)</p>	<p>Group 1: 51 ± 28.9 Group 2: 56 ± 30.47 p value: 0.59 (t test) Not sig.</p>	
			<p>Mean amount of urine (ounces ± SD) lost in 24h at week 5</p>	<p>Group 1: 4.3 ± 8.9 (4.3 oz = 121g) Group 2: 4.5 ± 7.7 (4.5 oz = 128g) p value: 0.95 (t test) Not sig.</p>	
			<p>Mean amount of urine (ounces ± SD) lost in 24h at week 12</p>	<p>Group 1: 0.0 ± 80.0 Group 2: 0.5 ± 1.7 (1.7 oz = 48g) p value: 0.22 (t test) Not sig.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
		<p>amount of urine lost.</p> <p>Volume of urine lost (ounces), number of pads used, number of episodes of urine loss, number of episodes of incontinence and length of time urine loss was experienced were all evaluated at weeks 2, 5, 9 and 12.</p>	<p>Proportion of still incontinent at 0 – 3 months (60-79 days)</p> <p>Data from Hunter et al., 2007¹²³</p>	<p>Group 1: 8/27 Group 2: 10/24 p value: NR</p>	<p>.</p> <p>Included study in SR by Hunter et al., 2007¹²³.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Moore et al, 1999²⁰³</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24 weeks</p> <p>Computer generated randomisation sequence and allocation concealment</p>	<p>Patient group: Patients who had undergone radical retropubic prostatectomy</p> <p>Setting: University-affiliated hospitals in Edmonton, Canada</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 4 weeks after radical prostatectomy (RP) • (>2 g of urine loss on pad test) • Neurologically normal • Within 2 h drive of study centre • Able to speak and read English • Willing to comply with protocol • No current treatment • Not seeking other treatment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Demand pacemaker • Previous pelvic muscle stimulation • Active rectal lesions or infections • Known detrusor instability <p>All patients N: 63</p> <p>Drop outs: 5 3 because of bladder neck contractures</p>	<p>Group 1 (PFMT) Pre and postoperative verbal + written instructions about PFMT by nurses in preadmission clinic and follow-up visits to urologist.</p> <p>Also Intensive physiotherapy 30 min 2/week for 12 weeks. Initial contractions were of 5-10 s + a 10-20 s rest, with 12-20 repetitions. For endurance exercises the 'hold' time was 20-30 s + equal rest time, with 8-10 repetitions. Speed was achieved by sets of quick repetitive contractions in a 10 s span with a 20-s rest. Finally, purposeful control occurred in 3 stages, with a 5-s hold each stage and a slow release, with a rest period of 15-30s.</p> <p>Group 2 (PFMT+ ES) Pre and postoperative verbal + written instructions about PFMT by nurses in preadmission clinic and follow-up visits to urologist</p> <p>Also patients met with the same physiotherapist 2/week for 30 min. Electrical stimulation (ES) with a surface anal electrode (InCare) was alternated with PMFT as for Group 1. Stimulation parameters were 50 Hz, a biphasic pulse shape with 1-s bursts, a 1 s pulse width and 1</p>	<p>Mean (median) [SD, range] urinary loss (g) in 24 h at baseline*</p> <p>Mean (median) [SD, range] urinary loss (g) in 24 h at 3 months*</p> <p>Mean (median) [SD, range] urinary loss (g) in 24 h at 4 months*</p> <p>Mean (median) [SD, range] urinary loss (g) in 24 h at 6 months*</p> <p>QOL Objective QoL measures (IIQ-7 and EORTC QLQ C-30)</p>	<p>Group 1 (PFMT): n=18: 565.6 (513.9) [403.3, 21.5-1538.6]</p> <p>Group 2 (PFMT+ ES) n= 19: 452.5 (492.1) [385.1, 5.3-1344.8]</p> <p>Group 3 (Control) n=21: 385.9 (395.5) [256.9, 6.3-921.5]</p> <p>Total n=58: 463.5 (419.8) [352.2, 5.3-1538.6] p value: Not sig</p> <p>Group 1 (PFMT): n=18: 86.9 (32.50) [123.0, 2.2-385.9]</p> <p>Group 2 (PFMT+ ES) n= 19: 155.5 (87.5) [168.1, 1.0-509.3]</p> <p>Group 3 (Control) n=21: 103.8 (23.8) [176.3, 1.0-702.4]</p> <p>Total n=58: 115.5 (27.2) [158.7, 1.0-702.4] p value: Not sig</p> <p>Group 1 (PFMT): n=18: 73.5 (10.35) [131.4, 1.0-494.6]</p> <p>Group 2 (PFMT+ ES) n= 19: 202.2 (85.7) [242.23, 1.0-753.4]</p> <p>Group 3 (Control) n=21: 67.3 (11.5) [137.4, 2.0-530.3]</p> <p>Total n=58: 114.2 (14.1) [185.6, 1.0-595.7] p value: Not sig</p> <p>Group 1 (PME): n=18: 69.9 (8.7) [113.5, 1.0-362.8]</p> <p>Group 2 (PME+ ES) n= 19: 98.2 (8.95) [132.1, 1.0-424.2]</p> <p>Group 3 (Control) n=21: 54.1 (6.9) [103.1, 1.0-277.3]</p> <p>Total n=58: 72.5 (7.5) [115.7, 1.0-424.2] p value: Not sig</p> <p>There were no significant group differences in either IIQ-7 or the QLQ C30 P NR</p>	<p>Funding: Oncology Nurses' Society, Canadian Nurses' Foundation, Caritas Health, Alberta Physiotherapy Association, Edna Minton Foundation, and the University of Alberta, Edmonton, Canada.</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Masking of outcome assessment was not reported • The results from the intervention arm are potentially confounded by the preoperative instruction on pelvic floor muscle contraction given to all groups <p>Notes: *Data from text for median urinary loss: A one-way repeated-measures ANOVA using a general linear model was computed to test the difference between and within groups, as well as the change over time at 12, 16 and 24 weeks. There were no differences among the</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>1 because of rectal pain when he did the exercises 1 because he went on vacation for 4 months and could not continue therapy Age (mean): 67 (range 49-77)</p> <p>Group 1 (PFMT) N: 20 Age (mean): 67.4 Drop outs: 2</p> <p>Group 2 (PFMT+ ES) N: 22 Age (mean): 65.7 Drop outs: 3</p> <p>Group 3 (Standard treatment) N: 21 Age (mean): 66.8 Drop outs: 0</p>	<p>s pulse trains.</p> <p>Group 3(Standard treatment) Pre and postoperative verbal + written instructions about PFMT by nurses in preadmission clinic and follow-up visits to urologist</p> <p><i>Continence was defined as a loss of <= 2 g of urine; socially acceptable continence was considered as <= 10 g</i></p>	<p>Proportion of still incontinent at 0 – 3 months (data from Hunter et al., 2007¹²³)</p> <p>Proportion of still incontinent at 3 – 6 months (data from Hunter et al., 2007¹²³)</p>	<p>Other data for QoL is reported in text for the whole population and not per group.</p> <p>Group 1: 12/20 Group 2: 11/22 Group 3: 14/21 p value: NR</p> <p>Group 1: 8/20 Group 2: NR Group 3: 7/21 p value: NR</p>	<p>groups (F=0.23, P=0.80) at any of the measurements</p> <p>Data for proportion of patients still incontinent was taken from Hunter et al., 2007¹²³</p> <p>Cochrane Review though it is unclear how this data was extracted from the paper.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Parekh et al., 2003 ²³³ Study design: RCT Evidence level: 1+ Duration of follow-up: 12 months	Patient group: men scheduled to undergo radical prostatectomy for localised prostate cancer Setting: Urology clinic, USA Exclusion criteria: Prior bowel or bladder incontinence All patients N: 38 Age (mean ± SD): NR Drop outs: 0 Group 1 N: 19 Age (mean ± SD): 61.6 M: 19 Mean preop PSA (ng/ml): 8.3 Drop outs: 0 Group 2 N: 19 Age (mean ± SD): 55.5 M: 19 Mean preop PSA (ng/ml): 8.1 Drop outs: 0	Group 1 PMFT using verbal and visualisation techniques and biofeedback using rectal probe was delivered by a physiotherapist comprising initial evaluation and 2 treatment sessions prior to surgery and then every 3 weeks for 3 months postoperatively. Home exercise programme was followed for 6 months or longer. Group 2 No treatment. All patients Completed urinary incontinence questionnaire by telephone or when questioned by medical students at weeks 6, 12, 16, 20, 28 and 52. Incontinence measured by number of pads used daily with continence defined as 0-1 precautionary pad	Median time to regain continence	Group 1: 12 weeks Group 2: 16 weeks p value: <0.05 (2 tailed <i>t</i> -test)	Funding: NR Limitations: <ul style="list-style-type: none"> • Randomisation method not described • Masking of outcome assessment not mentioned Notes: Study reports numbers of patients continent at time intervals but data are presented as number of patients still incontinent
			Proportion of patients still incontinent at 3 months	Group 1: 6/19 (32%) Group 2: 12/19 (63%) p value: NR NCGC Chi-squared calculation p=0.051 using ITT analysis Not sig.	
			Proportion of patients still incontinent at 6.5 months	Group 1: 4/19 (21%) Group 2: 7/19 (37%) p value: NR NCGC Chi-squared calculation p=0.28 using ITT analysis Not sig.	
			Proportion of patients still incontinent at 13 months	Group 1: 3/19 (16%) Group 2: 4/19 (21%) p value: NR NCGC Chi-squared calculation p=0.68 using ITT analysis Not sig.	
			Severe incontinence (>3 pads) at 12 months	Group 1: 2/19 (11%) Group 2: 3/19 (16%) p value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Paterson et al., 1997²³⁷</p> <p>Study design: RCT Observer masked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 13 weeks</p>	<p>Patient group: Men with post-micturation dribbling (PMD)</p> <p>Setting: Repatriation General Hospital, South Australia</p> <p>Inclusion criteria: Patients with an history of post-micturation dribbling (PMD)</p> <p>Exclusion criteria: No history of surgery on the bladder, prostate or urethra, or had a history of urgency or stress incontinence. All were able to comply with instructions</p> <p>All patients N: 49 Drop outs: 6</p> <p>Group 1 (counselling) N: 15 Age (mean [SEM]): 69.5 [2.4] Initial pad weight gain (g) (mean [SEM]): 7.56 [1.27] Initial pelvic muscle (mean [SEM]): 2.5 [0.21]</p> <p>Group 2 (milking) N: 15 Age (mean [SEM]): 69.3 [3.1] Initial pad weight gain (g) (mean [SEM]): 10.43 [2.99] Initial pelvic muscle (mean [SEM]): 2.6 [0.30]</p> <p>Group 3 (PFMT)</p>	<p>Group 1 (counselling) Advice on drinking patterns, types of beverages, aperient use, toileting habits, hints to alleviate oedema, dietary advice and relaxation therapy</p> <p>Group 2 (milking) Patients were given insights into the anatomy of the urethra and where the urine pools. They performed the procedure in the clinic to ensure that they did so correctly. An education sheet based on the technique outlined by Millard was issued to this group to reinforce their understanding of the procedure.</p> <p>Group 3 (PFMT) Pelvic muscle exercise: Patients were given simple education on the anatomy and physiology of the act of micturition. Time and effort were taken to enable correct identification of the pelvic muscles. Participants were taught to tighten and lift these muscles as if they were controlling flatus or interrupting the flow of urine mid-stream. They were encouraged to do them in front of the mirror to observe penile and scrotal lift and to recognize inappropriate tightening of abdominal and gluteal muscles. The fast-twitch muscle fibres were exercised by a series of 1-second contractions (usually five) and gradually extending the number of repetitions, depending on the individual ability of each participant. The slow-twitch fibres were exercised by repeating the maximum contraction as many times as possible without weakening of the length</p>	<p>Urinary loss measured by difference in mean pad weight gain</p> <p>Urinary loss was measured at baseline and at 5, 7, and 13 weeks using pad weighing method. Participants were given instruction on how to wear the pads, seal them in plastic bags and how to complete a bladder chart. The weighing and coding of the pads was the responsibility of the research assistant who was unaware of the participant's group allocation.</p> <p>Crude and adjusted mean (SEM) improvement in pad weight gain (g) Adjusted for initial pad weight gain</p>	<p>Data is reported in figures. The mean pad weight initially decreased rapidly in the exercise group and less so in the milking group but did not change dramatically in the counselling group (p values not reported).</p> <p>Counselling: n=15 Crude 0.019 (1.04) Adjusted: -1.387</p> <p>Milking: n=15 Crude 3.97 (2.07) Adjusted: 2.877 p<0.01 compared to counselling</p> <p>Exercise: n=13 Crude 4.28 (2.47) Adjusted: 4.707 p<0.001 compared to counselling</p>	<p>Funding: Cello Paper Pty donated weighing scales. Sancella Pty Ltd supplied the male incontinent pads</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method and allocation concealment were not reported. • Standard deviations were not available for adjusted improvement in pad weight again. • Sample size calculation is not reported. <p>Notes: Authors report compliance of participants was excellent, with all patients completing pad wearing and bladder charts, and 99.6% attendance of the required number of clinic visits.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 14 (1 patient completed 9 of the 13 weeks of the study) Age (mean [SEM]): 70.8 (2.7) Initial pad weight gain (g) (mean [SEM]): 11.68 [5.43] Initial pelvic muscle (mean [SEM]): 2.5 [0.23]</p> <p>Height and weight reported not included in this table.</p> <p>Differences in initial pad weight gain was Not sig.</p>	<p>and strength of the contraction. Participants were instructed to spread exercise sessions throughout the day and to vary the positions from lying to sitting and standing.</p>		<p>Improvement in pad weight gain was strongly influenced by initial pad weight gain, or degree of urine loss at the start of the study. After allowing for the effects of initial pad weight gain, the counselling group showed no improvement, the urethral milking group showed an adjusted mean improvement in urine loss of 2.9 g after 13 weeks, compared with 4.7 in the exercise group.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Porru et al., 2001²⁴⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 month</p> <p>Blinded outcome assessment for pelvic muscle strength</p>	<p>Patient group: diagnosis of symptomatic BPH selected to undergo TURP</p> <p>Setting: single centre, university urology clinic, Italy</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> > 80 years History of urethral or pelvic surgery Neurogenic bladder Prostate carcinoma <p>All patients N: 58 Age (mean): NR M: 58 Drop outs: 5</p> <p>Group 1: N: 30 Age (mean): 66 (range 53-71) M: 30 Drop outs: 2</p> <p>Group 2 N: 28 Age (mean): 67.5 (range 55-73) M: 28 Drop outs: 3</p>	<p>Group 1 Pelvic floor muscle training through verbal instructions and feedback on contractions. Patients received verbal and written instructions for home PFMT with a regimen of 3x15 exercises/day</p> <p>Group 2 No treatment</p> <p>All patients Pelvic floor muscle strength was measured using digital examination and graded from 0 (none) to 4 (strong) preoperatively and at follow up visits on week 1, 2, 3 and 4.</p> <p>Patients began voiding diaries immediately post TURP over 48 hour periods</p> <p>The AUA symptom score was administered preoperatively and at 30 days postoperatively. ICS male questionnaire was used to assess Quality of Life</p> <p>Uroflowmetry was performed pre and 30 days post TURP and pressure flow studies confirmed existence of BOO Incontinence assessed by voiding diary.</p>	<p>Proportion of patients still incontinent at 4 weeks</p>	<p>Group 1: 1/30 (3%) Group 2: 3/28 (11%) p value: NR NCGC Fishers exact test calculation p=0.34 using ITT analysis Not sig.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method not described Masking of outcome assessment not mentioned Incontinence was not clearly defined <p>Notes: Urologist measuring pelvic floor muscle strength was masked to treatment allocation</p>
			<p>Change in AUA symptom score at 30 days</p>	<p>Group 1: from 22 to 9 Group 2: from 24 to 10 p value: reported as Not sig. ANOVA</p>	
			<p>Change in ICS-Male Quality of Life score at 30 days</p>	<p>Group 1: from 5.8 to 1.5 Group 2: from 5.5 to 3.2 p value: <0.001 signif. ANOVA</p>	
			<p>Mean muscle contraction strength (grade 0-4) ± SD at 4 weeks</p>	<p>Group 1: 3.8 ± 0.3 Group 2: 2.4 ± 0.2 p value: NR. NCGC calculation using a two-sample t test with unequal variances p <0.00001 signif.</p>	
			<p>Mean voiding interval at 4 weeks (± SD)</p>	<p>Group 1: 110 ± 23 Group 2: 118.5 ± 24 p value: reported as Not sig.</p>	
			<p>Proportion of patients with post micturation dribbling and incontinence episodes at 4 weeks</p>	<p>Group 1: NR Group 2: NR p value: reported as Not sig.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size				Comments				
<p>Tibaek et al., 2007²⁹⁴</p> <p>Study design: RCT single blinded</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months after TURP</p>	<p>Patient group: Men with uncomplicated BPO (benign prostatic obstruction) scheduled for TURP (transurethral resection of the prostate).</p> <p>Setting: single centre, university hospital, Denmark</p> <p>Inclusion criteria: Fit, ambulatory, uncomplicated BPO scheduled for TURP</p> <p>Exclusion criteria: Prostate cancer, previous lower urinary tract surgery and neurological disease</p> <p>All patients N: 58 Drop outs: 9/58 (before intervention – group not specified)</p> <p>Group 1 N: 26 Age, median (range): 70(58-77) DAN-PSS-1 - Symptom score: 15(7-24) - Bother score: 17 (8-28) - Total Score: 28 (10-61) Urine output per 24 h (ml): 1827(1023-3187) Voided volume (ml): 165(50-350) Frequency (no. of voidings/24hr): 12(5-21) Max flow (ml/s): 7(3-15) Residual urine (ml): 116(0-877) 1st sensation (ml): 64(10-270) Max cystometric bladder capacity (ml): 131(38-406)</p>	<p>Group 1 (PFMT) Pre-TURP pelvic floor muscle training (digital-anal guided) lasting 4 consecutive weeks Program consisted of :</p> <p>- Individual information: 1 hour session including symptoms, anatomy and instructions on PFMT</p> <p>- 3 group treatments 1 hour of isolated PFM contractions, strength exercises, endurance exercises repeated 4-8x in the supine, standing and sitting positions and PFM contractions before and during rising from sitting position and walking</p> <p>- Home exercises: PFM strength and endurance exercises repeated gradually 6 - 10 x in the supine, standing and sitting positions, 1 or 2/day. Patients received new progressive programme after</p>	<p>DansPSS-1 total score (values range from 0-108) Results presented as median (range).</p> <p>Leakage in pad test (g/24 hours)</p> <p>Patients who used pads per 24hours, n(%)</p> <p>Urine output/24hours (ml)</p> <p>Voiding volume (diary) (ml)</p>	<p>2 weeks 4 weeks 3 months</p>				<p>Funding: Prof Jens C Christoffersen's Memory Fund, Danish Physiotherapist Research Fund, SCA Hygiene Products A/s. Astra Tech Denmark and Coloplast</p> <p>Limitations:</p> <ul style="list-style-type: none"> Physiotherapists assessing the PFM outcomes were masked. However, no mention on whether urological nurses who measured the subjective and objective voided parameters were blinded. No mention whether urologists performing the TURP were blinded Both groups received information about PMFT 				
				<p>Group 1:</p>	15(3-61)	11(0-52)	3 (0-24)					
				<p>Group 2:</p>	13.5(0-51)	6 (0-37)	4.5(0-51)					
				<p>P value:</p>	0.927	0.452	0.754					
				<p>Leakage in pad test (g/24 hours)</p>					<p>N#</p>	12/26	12/23	
				<p>Group 1:</p>	1(0-188)	12(0-374)	-					
				<p>Group 2:</p>	0(0-23)	4(0-56)	-					
				<p>P value:</p>	0.656	0.755						
				<p>#The others were continent and refused to do the test</p>					<p>2 weeks 4 weeks 3 months</p>			
				<p>Group 1:</p>	9/25 (36)	4/26(15)	3/26(12)					
				<p>Group 2:</p>	6/21(29)	4/21(19)	5/22(23)					
				<p>Relative risk: (95%CI)</p>	()	()	()					
<p>p value:</p>												
<p>Urine output/24hours (ml)</p>				<p>2 weeks</p>	<p>4 weeks</p>	<p>3 months</p>						
<p>Group 1:</p>	1985(105-3415)	1694(923-3003)	1875(775-3387)									
<p>Group 2:</p>	1887(583-3557)	1903(617-3803)	1820(367-2716)									
<p>p value:</p>	0.638	0.412	0.640									
<p>Voiding volume (diary) (ml)</p>				<p>2 weeks</p>	<p>4 weeks</p>	<p>3 months</p>						
<p>Group 1:</p>	165.5(40-250)	150(30-250)	200(50-300)									
<p>Group 2:</p>	127.5(50-360)	150(50-350)	155(50-360)									
<p>P value:</p>	0.563	0.599	0.510									

Study details	Patients	Interventions	Outcome measures	Effect size				Comments	
	<p>Unstable detrusor; n(%): 22/26(85) Pressure flow AG number (ml/s): 79.5(33-170) Weight of prostate specimen (g): 22(4-61) Histology; no with prostate cancer: 2 Time from randomisation to TURP (days): 42(18-140)</p> <p>Group 2 N: 23 Age, median (range): 68(52-79) DAN-PSS-1 - Symptom score: 15(6-22) - Bother score: 15(3-28) - Total Score: 26(3-64) Urine output per 24 h (ml): 1650 (418-3180) Voided volume (ml): 140 (50-350) Frequency (no. of voidings per 24 hour): 11.7(5-21) Max flow (ml/s): 7(1.5-17) Residual urine (ml): 108(0-875) First sensation (ml): 97(13-238) Max cystometric bladder capacity (ml): 174(42-338) Unstable detrusor; n(%): 19/23(83) Pressure flow AG number (ml/s): 76(22-228) Weight of prostate specimen (g): 24(10-58) Histology; no with prostate cancer: 2 Time from randomisation to TURP (days): 35(5-162)</p>	<p>the weekly lessons and motivated to continue until at least 4 weeks after surgery.</p> <p>Group 2 (control) -no preoperative physiotherapy treatment</p> <p>Both groups received brief information regarding the anatomy and physiology of the bladder and PFM, and were given verbal, instructions about PFMT in the ward 2-3 days after TURP</p>	<p>Frequency of voiding, times/24 hours</p> <p>Maximal Urine Flow (ml/s)</p> <p>Residual urine (ml)</p>		<p>2 weeks</p>	<p>4 weeks</p>	<p>3 months</p>	<p>after TURP. Confounding</p> <p>Additional outcomes: Attendance was 100% for 24/26 and 75% for 2/26</p> <p>All men had good initial PFM function (minimum rating 2), but did not improve to optimum function post-test.</p> <p>At 2 weeks, 41 men "improved", and 8 "worse". At 3 months, 3 patients still had higher DAN-PSS-1 score than before surgery</p> <p>Significant difference ($p=0.049$) between groups on dynamic muscle endurance.</p> <p>Notes: None.</p>	
					<p>Group 1:</p>	<p>11.85(7.5-28.3)</p>	<p>10.3(4.3-26.3)</p>		<p>10.0(6.0-17.3)</p>
					<p>Group 2:</p>	<p>13.2(5.7-20.7)</p>	<p>11.3(6.7-17.3)</p>		<p>10.7(4.3-19.0)</p>
					<p>P value:</p>	<p>0.657</p>	<p>0.499</p>		<p>0.794</p>
					<p>Group 1:</p>	<p>-</p>	<p>-</p>		<p>16.6(4.1-47)</p>
					<p>Group 2:</p>	<p>-</p>	<p>-</p>		<p>16.8(5.3-36.5)</p>
					<p>P value:</p>	<p>-</p>	<p>-</p>		<p>0.726</p>
					<p>Group 1:</p>	<p>-</p>	<p>-</p>		<p>22(0-661)</p>
					<p>Group 2:</p>	<p>-</p>	<p>-</p>		<p>1(0-56)</p>
					<p>P value:</p>	<p>-</p>	<p>-</p>		<p>0.127</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Van Kampen et al., 2000 ³⁰⁴ Study design: RCT Evidence level: 1+ Duration of follow-up: 12 months Blinded outcome assessment and allocation concealment	<p>Patient group: Men with a radical retropubic prostatectomy (RP) for localised prostate cancer</p> <p>Setting: Department of Urology, Leuven University Hospital, Belgium</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Incontinent on day 15 after surgery after catheter removal Able to regularly attend hospital appointments <p>Exclusion criteria: NR</p> <p>All patients N: 102 Age (mean): 65 range (52-76) M: 102 Drop outs: 4</p> <p>Group 1 N: 50 Age (mean): 64.4 ± 0.8 M: 50 Drop outs: 2 Previous TURP: 2 (4%) Preoperative micturation (IPSS): <10: 37 (74%) 10-20: 9 (18%) >20: 4 (8%)</p> <p>Group 2 N: 52</p>	<p>Group 1 Pelvic floor re-education programme extending for as long as incontinence persisted within time limit of 1 year. Programme comprised anatomical education pelvic floor and function, active pelvic floor muscle training (PFMT) with biofeedback. Strength of pelvic-floor muscles assessed using digital anal control and scored. 7 patients who could not contract were given electrical stimulation by anal probe. Patients were required to do 90 home exercises/day supine, sitting or standing. Each patient received treatment at weekly outpatient clinic</p> <p>Group 2 Attendance of weekly outpatient clinic receiving education on aetiology of UI and placebo electrotherapy that couldn't affect muscle function.</p> <p>Examination methods: Continence measured by 24h weighed pad test after catheter removal and everyday until patient was continent. **Continence defined as <2g urine lost per day on 24h and 1</p>	<p>Number of men achieving continence at 3 months</p> <p>Number of incontinent* patients at 12 months</p> <p>Duration of incontinence (Kaplan-Meier Survival Analysis)</p> <p>Number of patients with VAS score=0 completely dry at 1 month</p> <p>Number of patients with VAS score=0 completely dry at 6 months</p> <p>Number of patients with VAS score=0 completely dry at 12 months</p> <p>Proportion of still incontinent at 0 – 3 months</p> <p>Proportion of still incontinent at 3 - 6 months</p> <p>Proportion of still incontinent at 6 - 12 months</p>	<p>Group 1: 43/48 (not ITT) Group 2: 29/52 p value: 0.001 (Fishers Exact test) NCGC check using ITT analysis p=0.0008 (Chi-squared) signif.</p> <p>Group 1: 2/50 Group 2: 9/52 p value: 0.001 (Wald test) NCGC check using ITT analysis p=0.03 (Chi-squared) Not sig.</p> <p>Group 1: NR Group 2: NR p value: 0.0001 (log rank test)</p> <p>Group 1: 15/50 Group 2: 8/52 p value: NR NCGC check using ITT analysis p=0.08 (Chi-squared) Not sig.</p> <p>Group 1: 29/50 Group 2: 27/52 p value: NR NCGC check using ITT analysis p=0.5 (Chi-squared) Not sig.</p> <p>Group 1: 26/50 Group 2: 22/52 p value: NR NCGC check using ITT analysis p=0.3 (Chi-squared) Not sig.</p> <p>Group 1: 5/48 Group 2: 23/52 p value: NR</p> <p>Group 1: 2/48 Group 2: 12/52 p value: NR</p> <p>Group 1: 2/48 Group 2: 9/49 p value: NR</p>	<p>Funding: Grant from Fund of Scientific Research, Flanders, Belgium</p> <p>Limitations: No IPSS change data. No QoL score</p> <p>Notes: Patients placed in 6 subgroups according to amount of initial urine loss (>50g, <250g, >250g) and whether they had had a previous TURP. They were then randomised using permuted blocks by an independent person. Sealed envelopes but no statement of opacity. All patients treated by same physiotherapist. All continence assessments done by therapist who was not involved in the study.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean): 66.6 ± 0.8 M: 52 Drop outs: 2 Previous TURP: 5 (10%) Preoperative micturation (IPSS): <10: 41 (81%) 10-20: 9 (17%) >20: 2 (2%)</p>	<p><i>h pad test as well as patients indicating no incontinence in past 3 days</i></p> <p>Confirmation was by 1h pad test in hospital with additional assessment.</p> <p>Continence was also assessed subjectively by visual analogue scale (0=completely continent, 10=completely incontinent)</p> <p>Continence assessed preoperatively and at 1, 6, 12 months</p>			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Willie et al., 2003 ³²¹ Study design: RCT Evidence level: 1 + Duration of follow-up: 12 months post.op	Patient group: Men with clinically localized prostate cancer who were scheduled for radical prostatectomy. Setting: Department of urology Inclusion criteria: Patient willingness to make 2 visits 3 and 12 months postoperatively. Patients who underwent previous transurethral prostatic resection were not excluded from the study. Exclusion criteria: NR <u>All patients</u> N: 139 Drop outs: see outcomes <u>Group 1: PFMT</u> N: 47 Age (no units reported): 65.9 Prostate wt (gm): 58.5 % pathological tumor stage: pT1a-2b: 71.7 pT3a-3b: 28.3 pT4: 0 patients continent at baseline according to questionnaire: 20.5% Patients continent at baseline according to pad test: 29% Drop outs: see outcomes <u>Group 2: PFMT + Electrical Stimulation</u>	Group 1: PFMT: Patients received verbal and written instructions about postoperative PFMT from a physiotherapist. After this introduction each patient received intensive physiotherapy for 20 to 30 minutes for 3 days. All patients encouraged to perform the exercises twice daily for 3 months after discharge. Group 2: PFMT + Electrical Stimulation (ES) Patients received PFMT and ES and shown how to use the device by a dedicated nurse. ES was provided with a bioimpulser (Haynl Elektronik, Schonebeck, Germany) surface anal electrode. Therapy time was set for 15 minutes in the device. After this time the device was automatically downloaded to ensure that each patient had same therapy duration. Stimulation parameters were 27 Hz, biphasic pulse shape with 1-second bursts, a 5-second pulse width and 2-second pulse trains. Intensity was controlled by each patient from 10% to 100%. Group 3: PFMT +ES and Biofeedback: These patients were additionally treated with biofeedback (BFB) 15 minutes twice daily for 3	% patients continent at 3 months according to questionnaires to determine number of pads daily Results available at 3 months for questionnaires: n= 120	Group 1: PFMT: 3 months: 60% Group 2: PFMT + ES: 3 months: 65% Group 3: PMFT + ES + Biofeedback: 3 months: 53% p= 0.8	Funding: NR Limitations: Method of randomisation, allocation concealment and sample size calculation not described. Additional outcomes: Compliance to treatment Measured by asking the patients how long they had done the recommended treatment. Notes: Subjective continence was defined as no or 1 pad used daily. Objective continence <1 g/20 minute pad test
			% patients continent at 12 months according to questionnaires to determine number of pads daily Results available at 12 months for questionnaires: n= 129	Group 1: PFMT: 12 months: 88% Group 2: PFMT + ES: 12 months: 81% Group 3: PMFT + ES + Biofeedback: 12 months: 88.6% p= 0.50	
			% patients continent at 3 months according to 20 minute pad test Results available at 3 months for pad test: n= 79	Group 1: PFMT: 3 months: 64% Group 2: PFMT + ES: 3 months: 78% Group 3: PMFT + ES + Biofeedback: 3 months: 73% p= 0.5	
			% patients continent at 12 months according to 20 minute pad test Results available at 12 months for pad test: n= 124	Group 1: PFMT: 3 months: 76% Group 2: PFMT + ES: 3 months: 82% Group 3: PMFT + ES + Biofeedback: 3 months: 90.5% p= 0.24	
			Number of men still incontinent at 3 months (ITT analysis)	Group 1: PFMT: 17/47 (36%) Group 2: PFMT + ES: 10/46 (22%) Group 3: PMFT + ES + Biofeedback:	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 46 Age (no units reported): 64.6 Prostate wt (gm): 53.7 % pathological tumor stage: pT1a-2b: 70.4 pT3a-3b: 27.3 pT4: 2.3 Patients continent at baseline according to questionnaire: 22.9% Patients continent at baseline according to pad test: 36.4% Drop outs: see outcomes</p> <p>Group 3: PFMT +ES and Biofeedback N: 46 Age (no units reported): 64.6 Prostate wt (gm): 55.4 % pathological tumor stage: pT1a-2b: 55.6 pT3a-3b: 42.2 pT4: 2.2 Patients continent at baseline according to questionnaire: 20.7% Patients continent at baseline according to pad test: 33% Drop outs: see outcomes</p>	<p>months using the same device and the same anal probe. Each contraction of the anal sphincter and pelvic floor led to a corresponding signal in the device display to ensure that the patient had control over training. The combined ES and BFB programme consisted of a stimulation time of 5 seconds, and a contracting the relaxing time of 5 and 15 seconds, respectively.</p> <p>All patients: Patients were encouraged to perform the treatment they were randomised to for 3 months. There was regular personal interaction between the patient and a health professional during the 6 weeks of surgery. After that time they had no further support.</p>	<p>Number of men still incontinent at 12 months (ITT analysis)</p>	<p>12/46 (27%)</p> <p>Group 1: PFMT: 11/47 (24%) Group 2: PFMT + ES: 8/46 (18%) Group 3: PMFT + ES + Biofeedback: 5/46 (10%)</p>	

Evidence Table 6: Post void milking vs. no intervention or other conservative intervention

See Evidence Table 5: Pelvic floor exercises (with or without electrical stimulation or biofeedback) for Paterson et al., 1997²³⁷

Evidence Table 7: Product vs. no product or other conservative intervention

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Fader et al, 2006⁸⁷</p> <p>Study design: Cross over RCT</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: 4 weeks, 1 week for each design</p>	<p>Patient group: Men with light urinary incontinence</p> <p>Setting: United Kingdom</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> - ≥18 years old - usually use an absorbent product for light urinary incontinence or had been accessed by a health care professional to as suitable to use such products <p>All patients N: 74 Age: median 70 years (range 23-92) Dropouts: 6 (did not return any data) Type of incontinence:</p> <ul style="list-style-type: none"> - 50% did not know type - 21% stress, 16% urge, 13% mixed <p>Output type: 90% described as “dribbled”, 7% as “gush” and 3% as constant flow Time of incontinence:</p> <ul style="list-style-type: none"> - 31(46%) both day and night - 37(54%) during the day only <p>Usual products: <u>Leaf:</u> 38%</p>	<p>Products: All products available for leaf (6 types) and pouch (6 types) design. The best product for pads and pants with inserts were chosen.</p> <p>Products in random order for up to 1 week. Total test time was 14 weeks.</p> <p>Product performance: Rated using product performance questionnaire (developed from earlier study)</p> <p>Wet product weights Measured and recorded using pad leakage diaries.</p>	<p>Prioritisation of product performance characteristics (% of patients rated it as top 5):</p> <ul style="list-style-type: none"> - Ability to hold urine (Absorbance without leakage-82%) - Comfort (88%) – leaf design allowed the scrotum to stay wet, and this can cause skin irritation and discomfort. - Fit (71%) – designs which are flatter preferred - Discreteness and ability to stay in place (23%) -- elastics help product to stay in place. If a product fall off (ie down the trouser leg), it can be very embarrassing. <p>Other issues: Ease of use and practical issues</p> <ul style="list-style-type: none"> - Absorbent products can be difficult to manage away from home when wet. <ul style="list-style-type: none"> o Men’s toilet cubicles may not have the equivalent of sanitary disposal unit. Discrete disposal difficult o For washables, need to bring home for washing. Washing and drying can be problematic and embarrassing o Pouches fiddly to apply, especially through a trouser fly, and difficult to reinsert when it is swollen with absorbent gel. Some men may need a cubicle instead of urinal. <p>Design performance results*: <u>Very good/good:</u> Leaf : 59% Pouch: 24% Pantegral: 50% Small pad: 51%</p> <p><u>Okay:</u> Leaf : 25% Pouch: 21% Pantegral: 12% Small pad: 31%</p> <p><u>Poor/very poor:</u> Leaf : 16% Pouch: 55%</p>		<p>Funding: The products were provided from manufacturers.</p> <p>Limitations: - Not a blinded study. - Method of qualitative analysis not well described</p> <p>Additional outcomes: Specific product performance measured by product performance questionnaire provided for each brand of leaf or pouches tested.</p> <p>Related outcomes Fader et al 2008 ⁸⁶ reported that men and women have different preferences of products. The suitability of products may depend on time of use (day vs. night) due to the position of the penis and whether when going out or staying at home. For overall acceptability, men preferred pull ups or diapers to pads. Washable diapers were most popular among men for use at night.</p> <p>Notes:</p>

	<p><u>Small disposable pads</u> : 35% <u>Other methods (including pouches or Pantegral)</u>: 27%</p> <p>Most use 1-2 products during the day (66%), and during the night (87%).</p> <p>Other characteristics: 76% walked independently, 21% use walking aids routinely, 3% use occasionally. 32% reported penile retraction</p>		<p>Pantegral: 38% Small pad: 18%</p> <p><u>Leakage performance (10g)</u> 96(90-98)% 88(78-94)% 57(43-70)% 93(84-97)%</p> <p><u>Leakage performance (50g)</u> 87(76-93)% 85(75-91)% 7(0-56)% 87(76-93)%</p> <p>*Results from best products in each design category.</p> <p>Leaf products:</p> <ul style="list-style-type: none"> - Varied in performance within group. Tena Level 2 significantly better (score of 79% in overall opinion) compared to others brands (19-40%) in the same leaf design group - Leakage performance was generally better for disposables compared to washables (88-96% vs. 59% do not leak when holding 10g of urine) <p>Pouches:</p> <ul style="list-style-type: none"> - Least successful design - More homogenous in performance (range of 15-28%). Generally lower score than leaves. - 74-88% do not leak when holding 10g of urine. 	<p>None</p>
--	---	--	---	-------------

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Jakobsson et al, 2002¹²⁶</p> <p>Study design: qualitative study</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: Questionnaire</p>	<p>Patient group: sample selected from men with prostate cancer and BPH that were part of larger questionnaire study.</p> <p>Setting: They were randomly selected from 2 urological clinic registers in Sweden.</p> <p>Inclusion criteria: Men with experience of indwelling urinary catheter treatment.</p> <p>All patients N: 108 Group 1: n=37 Group 2: n=71</p> <p>Treatment duration: Group 1: Men with BPH <1 week=48.6 2-4 weeks=18.9 1-2 months=27.0 >3 months=5.4 Group 2: Men with prostate cancer <1 week=11.3 2-4 weeks=54.9 1-2 months=24.0 >3 months=8.5</p>	<p>Questionnaire – questions on experiences of indwelling catheter installation, wearing and handling and background data. Response format was on nominal (no-yes) and ordinal (ranging from 'not at all' to 'much') scale levels.</p> <p>Assessment of health related quality of life with the QLQ-C30 questionnaire – which includes five functional scales (physical, role, emotional, social and cognitive functioning), three symptoms scales (fatigue, pain, and nausea and vomiting) a global health status and additional single items. Response format comprised yes-no questions and assessment ranging from 'very bad' to 'excellent' (1-7). All scores linearly transformed to a 0-100 scale.</p> <p>Sense of Coherence Questionnaire, 13 item format used in the study (1-7 score to disagree completely to agree completely).</p>	<p>Information about wearing a catheter:</p> <p>Information about handling a catheter</p> <p>Mean (SD) functional scales: higher score better function):</p> <p>Feelings of discomfort, tagging, smarting and pain at catheter instalment, resting, moving and problems related to indwelling catheter treatment:</p>	<p><u>Little or less than wanted:</u> Group 1: 23.9% Group 2: 29.9%</p> <p><u>Satisfaction with information:</u> Group 1: 24.3% Group 2: 52.1%</p> <p><u>Question not applicable:</u> Group 1: 35.1% Group 2: 16.9%</p> <p><u>Little or less than wanted:</u> Group 1: 22.6% Group 2: 23.9%</p> <p><u>Satisfaction:</u> Group 1: 24.3% Group 2: 56.3%</p> <p><u>Not applicable:</u> Group 1: 40.5% Group 2: 14.1%</p> <p>Physical: 85.5 (22) / 84.3 (24.1) Role: 83.3 (28) / 83.3 (29) Emotional: 85.4 (19.5) / 86.0 (17.8) Cognitive: 85.1 (15) / 85.2 (18.3) Social: 85.0 (14.6) / 85.2 (18.3) QoL: 69.0 (26) / 72.0 (23.0)</p> <p><u>Discomfort: % Rather much / much</u> Instalment: 38 / 5.6% Resting: 32.4 / 1.9% Moving:40.8 / 7.4% <u>Tagging: % Rather much / much</u> Instalment: 25.9 / 0.9% Resting: 19.4 / 2.8% Moving:38.9 / 5.6%</p> <p><u>Smarting: % Rather much / much</u></p>	<p>Funding: Supported by the medical faculty, Lund University, the Swedish Foundation for Health Care Science sand Allergy Research, the County Council of Kristianstad, and Kristianstad University college.</p> <p>Limitations: - Aim of study to compare results from men with BPH to men with prostate cancer. - QLQ C-30 score is cancer specific. - study only looked at negative views of catheters.</p> <p>Additional outcomes: Factor solution of indwelling catheter treatment and mean values. Single items on health related quality of life scores.</p> <p>Notes: None</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Instalment: 25 / 2.8% Resting: 15.7 / 1.9% Moving:23.2 / 1.9% <u>Pain: % Rather much / much</u> Instalment: 26.9 / 2.8% Resting: 14.8 / 1.9% Moving:20.3 / 2.8% Infections % Rather often / often: 18.5 / 7.4% Smeary urethra: 25 / 6.5% Difficulties attaching catheter comfortably: 30.5 / 1.9% Difficulties attaching drainage bag comfortably: 31.5 / 0.9% Difficulties changing drainage bag: 13.9 / 0.9% Fear of leaking urine: 25.9 / 4.6% Fear of drainage bag rupture: 16.7 / 3.7% Difficulties finding comfortable resting/sleeping position: 46.3 / 1.9%	
			Bivariate significant relationship between health related quality of life and sense of coherence	Global quality of life had a moderate correlation to sense of coherence: $r=.0.52$	
			Multiple logistic regression test:	No association between global quality of life, QOL, and the independent variables under study in any of the groups.	

Study details	Patients	Intervention (Methodology)	Outcomes	Comments
<p>Macaulay et al, 2004^{177,177}</p> <p>Study design: 2 interviews (pre and post tests), and a survey (questionnaire)</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: Not stated. Up to 8 washes for each product</p>	<p>Patient group: Men/Women who had moderate/ eavy incontinence. Fully mobile.</p> <p>Participants recruited from advertisement in a consumer journal (Incontact)</p> <p>Cause of incontinence: Varied, not specified.</p> <p>Setting: UK</p> <p>All participants N: 14 Age (mean): 43.6 , range 28-67 years M/F: 10/4</p>	<p>Purpose: To evaluate all the reusable products for moderate/heavy incontinence and compare them with disposable alternatives.</p> <p>Methods: Order of product testing was randomized. Subjects tests products one after another based on randomization order, and repeat the process until each product tested a maximum of 8 times.</p> <p>Sequence of follow up: <u>Pretests interview</u> – to determine attributes of products considered to be important</p> <p><u>Testing period:</u> Completion of product performance questionnaire and pad leakage diary. Questionnaire was designed based on the pretest interview.</p> <p><u>Post test interview</u> Feedback regarding reusables</p>	<p>Difference in men vs. women in fitting of pads.</p> <ul style="list-style-type: none"> - Men were not always happy with a product they perceived to be designed for women. - Fitting of insert pads (for pants with integral pads), shaping of pads did not reflect anatomy. - Some reversed the inset pads thereby having their larger end situated to their front. This left the smaller end feeling uncomfortable around the buttocks. <p>Problems with washing</p> <ul style="list-style-type: none"> - A man who had to use a launderette found it difficult. Even when washed at home, this could lead to some embarrassment when they are part of the family laundry, in a bucket or on a drying line. <p>Most important product attributes:</p> <ul style="list-style-type: none"> - Leakage/absorbency, discreteness, comfort and fit. - More details about the specific performance attributed were reported. 	<p>Funding: conducted by Continence Product Evaluation (CPE) Network , funded by MHRA</p> <p>Limitations:</p> <ul style="list-style-type: none"> - Selection of participants from specialized consumer journal – not certain how this is representative of men with LUTS. Patients noted to be relatively young. - This was a pilot study with small sample size. - Feedback from men and women were not reported separately. - Method of qualitative analysis not well described <p>Additional outcomes: More details about the specific performance attributed were reported</p> <p>Notes: A full report on the product performances are detailed in a report to MHRA: MHRA. A pilot study to evaluate reusable absorbent body- word products for adults with moderate/heavy urinary incontinence. Med healthcare Prod Reg Agency. 2003:IN11</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Moore et al., 2004²⁰⁴</p> <p>Study design: Cross over randomised</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 days, 1 day for each product/control</p>	<p>Patient group: Men with radical prostatectomy ≤ 6 months ago</p> <p>Setting: Canada</p> <p>Inclusion/Exclusion criteria:</p> <ul style="list-style-type: none"> - Men with stress incontinence who required continuous incontinence pad protection after radical prostatectomy - Normal perineal and penile sensation, intact penile skin, no neurologic disorders that could affect sensation or peripheral circulation, sufficient manual dexterity to manage the penile compression device - No overactive bladder - No cognitive impairment that could affect their ability to follow instructions or perceive penile discomfort (Mini-Mental State Examination score ≥27), ability to read and speak English <p>All patients N: 12 Mini Mental State Score (Mean 29.6 ± 1.2) No other baseline data provided</p>	<p>Group 1: Control-no device</p> <p>Group 2: Timms C-3 penile compression device</p> <p>Group 3: Cunningham Clamp</p> <p>Group 4: U-TeX Male Adjustable Tension Band</p> <p>All these interventions were randomly carried out on 4 sequential days. Subjects were instructed to standardise their activities, time of day for wearing the devices and the amount of fluid intake.</p>	<p>Mean urine loss (grams loss in 4 hour pad test)</p>	<p>Group 1 (No device): 122.8 ± 130.8</p> <p>Group 2 (C-3): 32.3 ± 24.3</p> <p>Group 3 (Cunningham): 17.1 ± 21.3</p> <p>Group 4 (U-TeX): 53.3 ± 65.7</p> <p>p value: <0.05 for all groups vs. Group 1</p> <p>Note: The standard deviation sizes were larger than the mean values, indicating that the data was potentially skewed and not normally distributed.</p>	<p>Funding: University of Alberta: Internal Allocations Fund and Department of Radiology. One investigator was supported by the Ministry of Health of the Province of British Columbia.</p> <p>Limitations:</p> <ul style="list-style-type: none"> - Data analysis – Data was potentially not normally distributed, but a parametric test (analysis of variance, Dunnett’s procedure for post hoc) was used. Interpretation of results need to be treated with caution since n=12. - The duration of intervention was only 4 hours or each product, or the control (1 pad test each). - The value for Doppler tests for Cunningham clamp was reported for the loosest setting, but setting for others was not reported. - The outcome for patient satisfaction was measured using Male Continence Device Satisfaction Questionnaire, which was adapted from another product testing questionnaire. It is unclear whether this is a fully validated instrument. The criteria for determining “rated positively” were not stated. <p>Additional outcomes: None of the clamps completely eliminated urine loss.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			<p>Patient satisfaction (rating device positively, using Male Continenace Device Satisfaction Questionnaire)</p>	<p>Group 1 (Control): NR Group 2 (C-3): 2/12 Group 3 (Cunningham): 10/12 Group 4 (U-Tex): 0/12 p value: NR For U-Tex, none reported it “positively” because it was difficult to apply, did not stay on with activity and did not control urine leakage satisfactorily. The patient satisfaction for no control was not reported.</p>	<p>Safety data: Blood flow (Systolic velocity)- measured using Doppler Ultrasound. <u>Right:</u> Group 1 (Control): 12.4±2.8 Group 2 (C-3): 12.4±5.5 Group 3 (Cunningham): 9.5±2.3* Group 4 (U-Tex): 11.9±4.4 p value: * <0.05 vs. control <u>Left:</u> Group 1 (Control): 12.3±3.0 Group 2 (C-3): 11.7±4.7 Group 3 (Cunningham): 7.3±3.0* Group 4 (U-Tex): 13.8±7.3 p value: * 0.05 vs. control</p> <p>Resistance Index- measured using Doppler Ultrasound. <u>Right:</u> Group 1 (Control): 0.90±0.10 Group 2 (C-3): 0.92±0.10 Group 3 (Cunningham): 0.92±0.13 Group 4 (U-Tex): 0.93±0.08 p value: * 0.05 vs. control) <u>Left:</u> Group 1 (Control): 0.87±0.10 Group 2 (C-3): 0.92±0.11 Group 3 (Cunningham): 0.86±0.29 Group 4 (U-Tex): 0.91±0.11 p value: * 0.05 vs. control</p> <p>Notes: Information from author: Patient satisfaction data was based on the reply to a single question "What is your overall opinion of the penile compression device?" Response choices for this question was not provided.</p>

Study details	Patients	Methodology	Outcomes	Comments
<p>Paterson et al, 2003²³⁶</p> <p>Study design: Qualitative Study Semi structured interviews and focus groups</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: NR</p>	<p>Patient group: Participants included people who had incontinence or cared for someone with incontinence, or were part of an advocacy group that had significant numbers of people with incontinence in its membership, from metropolitan, rural and remote Australia. Included people of minority backgrounds and indigenous Australians.</p> <p>Purposive and snowballed sampling. Participant recruitment ceased once no new themes emerged.</p> <p>Cause of incontinence: Varied widely and included congenital malformations, chronic debilitating diseases, sever spinal cord injuries and degenerative diseases.</p> <p>All participants N: 82 NR Age (mean): NR M/F: NR Dropouts: NR</p>	<p>Purpose: To understand issues, needs and concerns of people with incontinence to inform development of comprehensive Australian consumer guide to continence products.</p> <p>Analysis method: Key issues transcribed from audio tapes. Constant comparison, thematic data analysis was commenced concurrently with data collection enabling the opportunity to follow up an emerging theme. (grounded theory)</p> <p>Transcriptions and notes taken during sessions Integrated into common themes, shared meanings, similarities and difference.</p> <p>3 researchers conducted analysis, cross- validated with another.</p> <p>Analysis focused on the similarities in experiences and concerns of consumers across the group.</p>	<p>Overall: Striking similarities in experiences and concerns about selection of consumer products.</p> <p>Seeking information:</p> <ul style="list-style-type: none"> - <u>Did not know how to begin to search for information and had problems finding it:</u> Most gathered information themselves, and these are usually not all available in one place. - <u>Feeling vulnerable:</u> Most felt discussing about incontinence management and shopping for products very personal and embarrassing. Some reluctant to speak to professionals. - <u>Lack of confidence in healthcare professional's knowledge:</u> Although dependent on healthcare professionals for assessment and referral, they had not received much helpful advice on products or directed to sources of advice. The most satisfactory help was from specialist continence nurse advisers. Local doctors knew little about assessment and management and many participants were dissatisfied. There was a pervasive "grin and bear with it" attitude and participants were expected to purchase a supermarket product and learn to live with it. - <u>Assessment and management:</u> Participants expressed a need for these to be standardised and coordinated. <p>Finding a suitable product:</p> <ul style="list-style-type: none"> - <u>Tried different products</u> to find one which enable them to remain socially continent. - <u>Advice for product selection:</u> Most had limited product knowledge in early stages and selected from limited range accessible to them in shops, hospital suppliers and recommendations of professionals. However, participants in support networks benefited from exchange of information. - <u>Key factors influencing selection of continence products</u> were quality, comfort and design balanced against availability and cost. Specific product features of concern including noise, allergy, trouble of keeping on, leakage around the seams 	<p>Funding: National Continence Management Strategy, an initiative of the Commonwealth of Australia Department of Health and Aged Care</p> <p>Limitations: Possible selection bias as details of demography, disease, disease severity and role of participants not reported. Not clear whether their target group of 'incontinent' patients is for urinary or faecal incontinence or both.</p> <p>Notes: Analysis did not use verbatim transcripts.</p>

Study details	Patients	Methodology	Outcomes	Comments
			<p>Information about product use and disposal required:</p> <ul style="list-style-type: none"> - Instructions for use and wear - Best methods for care and disposal of products <hr/> <p>Suggestions for content and format of the consumer guide to products:</p> <ul style="list-style-type: none"> - Detailed product description - More information in general about incontinence (causes, treatments and sources of help) and - Use simple layman’s language throughout guide. - Make available a variety of formats and a wide distribution throughout the community 	

Evidence Table 8: Catheters vs. no catheters

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Logan et al, 2008¹⁷⁰</p> <p>Study design: Qualitative study</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: NR</p>	<p>Patient group: selected from case lists of a continence and urology service. Patients with experiences of learning clean intermittent self catheterisation (CISC).</p> <p>Patients selected to include maximum variation of characteristics likely to impact on views, attitudes and access to services.</p> <p>Setting: Continence and urology service in Wales.</p> <p>All patients N: 15 M/F: 8/7 Median age (range): 65 (33-81) Duration of use: 6m to >2y Frequency: weekly to four times per day.</p> <p>Reasons for catheterisation: MS, urethral stricture, urine retention.</p>	<p>In depth interviews from January to June 2006 in the UK by two of authors and by a continence nurse. Interview guide developed based on the literature and experience and expertise of the research team. Topics helped guide the interviewer to explore reasons for CISC duration and frequency of CISC, experience of being taught, location, teaching aids, information, ongoing support and follow-up. Guide covered all relevant areas but allowed interviews to pursue themes emerging during the interview.</p>	<p>Psychological-embarrassment and privacy: Views not separated out for men and women.</p> <p>Physical: Technical difficulties were expressed by both sexes. Men's difficulties were related to negotiating the penile anatomy and handling the lengthy catheters. Generally men had no problem in visualising the urethra. One man experienced muscle spasms and urethral 'clamping', causing difficult insertion and frustration in the first few months.</p> <p>The entire sample used coated catheters, commonly describing them as 'slippery'. To overcome this, some men developed practical handling strategies; another recruited his wife to help. Men with urethral strictures described complications such as discomfort, bleeding and problems negotiating the stricture: 'Sometimes you (have) got to twiddle, twirl it in around it and just sort of ease it in the best way I can'.</p> <p>Both sexes avoided touching the catheter tip for fear of contamination and infection, illustrating concerns about hygiene and the development of a good technique.</p> <p>In the beginning, respondents found CISC emotionally and technically difficult. Gaining confidence was related to pace of skill acquisition. Men were squeamish at the thought of inserting a catheter for the first time, because of psychological issues and fear of causing internal damage. Q: You were going weak at the knees were you? A: Yes, definitely yes, and the perspiration... I was afraid to blink, I wouldn't see... you know, from a man's point of view to think you got something that long to push into yourself!</p> <p>Only two men in study felt confident immediately while the majority took considerably longer to accept CISC as part of their lives.</p> <p>Service interaction: Information-giving: Participants were unfamiliar with CISC, and on hearing the word catheter feared it would involve a permanent 'catheter</p>	<p>Effect size</p>	<p>Funding: Gwent Health Care Trust research and development small grant scheme.</p> <p>Limitations: Mix of views from men and women.</p> <p>Additional outcomes: Service interaction was also covered.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
					<p>and bag': 'I didn't know enough about it – I was just told that I had to start using a catheter and I didn't know any thing at the point...I didn't know that there was a much simpler, straight forward version that you could use yourself and that point I was not at all happy about it' .</p> <p>Practical demonstration was an important component of learning CISC, and a few participants felt that their demonstrations had been insufficient: 'I would have liked more than one demonstration or more time spent...I was only shown once and I had to get on with it then.'</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Saint et al, 1999²⁶⁶</p> <p>Study design: Qualitative study</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: NR</p>	<p>Patient group: Consecutive male patients between May and November 1998 who were using an indwelling or condom urinary catheter.</p> <p>Setting: Patients housed on the medical, rehabilitation and nursing home units of Puget Sound VA health Care System.</p> <p>Inclusion criteria: patients with a urinary catheter in use for at least 24 hours were eligible to participate.</p> <p>All patients N: 116 Mean age (SD): 71 (12) Drop outs: 12 90% response rate.</p> <p>Group 1: n = 21 Group 2: n = 83</p> <p>Location: Hospitalised on an acute care ward: 72% Other ward (nursing home, surgery, neurology, rehabilitation): 28%</p>	<p>Face to face interviews with a simple instrument requiring only yes or no answers for each of the 5 questions.</p> <p>Group 1: men using a condom catheter</p> <p>Group 2: men using an indwelling catheter</p>	<p>% of men reporting yes to questions at interview: Question: Is the current urinary catheter...</p> <ol style="list-style-type: none"> 1. Comfortable? 2. Painful? 3. Convenient? 4. Restricting your daily activity? 5. Causing you embarrassment? <p>Logistic regression: Condom catheters compared to indwelling were found to be: More comfortable: Less painful: Less restrictive: Convenience or embarrassment: Patients were also asked if they remembered having another type of urinary collection device in the past (alternative catheter or disposable diaper). If yes, we</p>	<p>Group 1: 86% Group 2: 58%, p=0.04</p> <p>Group 1: 14% Group 2: 48%, p=0.008</p> <p>Group 1: 86% Group 2: 75%, p=0.40</p> <p>Group 1: 24% Group 2: 61%, p=0.002</p> <p>Group 1: 24% Group 2: 30%, p=0.50</p> <p>OR=4.2; 95% CI: 1.1 to 15.6, p=0.03</p> <p>OR=0.17; 95% CI: 0.05 to 0.64, p=0.008</p> <p>OR=0.23; 95% CI: 0.07 to 0.75, p=0.01</p> <p>Catheter type not significantly related. N=36 Preferred condom: 17 (47%) Preferred indwelling: 14 (39%) No preference: 5 (14%)</p>	<p>Funding: Supported, in part, by the Department of Veterans Affairs and the Robert Wood Johnson Clinical Scholars Program.</p> <p>Limitations: Not population of interest.</p> <p>Additional outcomes: Nurses views by questionnaire.</p> <p>Notes: Logistic regression analysis using each 'yes' or 'no' answer as the dependent variable with patient age, hospital service and current catheter type as independent variables.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			<p>asked whether they preferred current or previous device.</p> <p>Previous experience of disposable diapers, n=27</p> <p>Men with experience of condom catheter (n=43)</p>	<p>Group 1: n=10 preferred current catheter Group 2: n=17; 9 preferred current catheter, four preferred diapers and four had no preference.</p> <p>N=7 (16%) offered spontaneously that main drawback was the associated leaking.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Shaw et al, 2008²⁷³</p> <p>Same trial as Logan, et al (see evidence table above) reporting more outcomes on QOL</p> <p>Study design: Qualitative study</p> <p>Evidence level: 3+</p> <p>Duration of follow-up: NR</p>	<p>Patient group: selected from case lists of a continence and urology service. Patients with experiences of learning clean intermittent self catheterisation (CISC).</p> <p>Patients selected to include maximum variation of characteristics likely to impact on views, attitudes and access to services.</p> <p>Setting: Continenence and urology service in Wales.</p> <p>All patients N: 15 M/F: 8/7 Median age (range): 65 (33-81) Duration of use: 6m to >2y Frequency: weekly to four times per day.</p> <p>Reasons for catheterisation: MS, urethral stricture, urine retention.</p>	<p>In depth interviews from January to June 2006 in the UK by two of authors and by a continence nurse. Interview guide developed based on the literature and experience and expertise of the research team. Topics helped guide the interviewer to explore reasons for CISC duration and frequency of CISC, experience of being taught, location, teaching aids, information, ongoing support and follow-up. Guide covered all relevant areas but allowed interviews to pursue themes emerging during the interview.</p>	<p>Impact on QoL:</p> <p>Positive impacts <u>Specific comments from men:</u> There were reports of relief from symptoms such as recurrent urinary tract infections. “I would rather do this than put up with the symptoms of infection.”</p> <p>CISC was also deemed to be a preferable option compared to other management strategies, such as permanent catheters with leg bags. “I said, ‘I don’t want a catheter fixed to me permanent, this bag on the leg or whatever they use’”.</p> <p>Negative impacts <u>Specific comments from men:</u> “..if I found a disabled toilet where you can go into the room and wash your hands and whatever, and in a normal toilet you can’t do that”</p> <p>“I have a problem when I am out...Finding water... If you go to a public toilet you have to fill it and then go into the toilet.”</p> <p>Difficulty experienced in travelling Carrying the necessary equipment was a particular problem:</p> <p>“Yes. I can’t travel light. Where I would much prefer to get on the train and go over and come back again, I now drive”</p> <p>Physical impacts <u>Specific comments from men:</u> Some reported occasional bleeding, or ongoing discomfort: “Oh it still gets sore now...especially with the withdrawal, insertion and withdrawal. And, of course, when you empty your bladder for the first time after the procedure, it’s grit your teeth..”</p> <p>Carrying out CISC <u>Specific comments from men:</u> One man had a common problem of muscle spasm preventing insertion of the catheter. Whilst he had learned how to manage this,</p>		<p>Funding: Gwent Health Care Trust research and development small grant scheme.</p> <p>Limitations: Mix of views from men and women.</p> <p>Additional outcomes: Same trial as Logan, et al (see evidence table above) reporting more outcomes on QOL</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			<p>he found it an inconvenience as he had to wait before trying to catheterize again.</p> <p>Factors explaining variation in QoL impacts</p> <p>Reasons for carrying out CISC and sex issues: More men found CISC to be a nuisance and time-consuming. This was related to the reasons carrying out CISC. More women carried is out to relieve previously severe urinary tract symptoms, whereas men tended to have problems with urethral stricture or voiding difficulties in the absence of severe symptoms. Because of differences in physiology and the longer urethra, men were more likely to be anxious about the catheter causing discomfort or pain, or about inadvertent damage because of poor technique.</p> <p>Type of catheter and sex issues There were sex differences related to type of catheter as male catheters are longer and more unwieldy. This had implications for carrying catheters discreetly. Women easily carried catheters in their handbags, whereas men were less likely to carry a bag and had difficulty carrying catheters in their pockets.</p>		

See Evidence Table 7: Product vs. no product or other conservative intervention

for Jakobsson et al., 2002¹²⁶.

Evidence Table 9: Alpha-blockers vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Andersen et al., 2000 ¹⁶	Patient group: Men between 50-80 years with evidence of BPH.	Phase 1: 2 week wash out Phase 2: Run-in period 2-week single blind placebo run-in period Phase 3: Treatment period: 13 weeks double blind	Mean (SE) adjusted change from baseline to final visit for total IPSS score (per-protocol analysis)	Group 1 (n=310): -8.0 ± 0.3 ; $p < 0.01$ Group 2 (n=311): -8.4 ± 0.3 ; $p < 0.01$ Group 3 (n=151): -6.0 ± 0.4	Funding: Pfizer Inc. Limitations: Method of randomisation and allocation concealment was NR. Additional outcomes: Mean changes from baseline in individual symptom IPSS score. Graphical presentation of IPSS and Qmax over each visit. Blood pressure and heart rate, pharmacokinetics.
Study design: RCT	Inclusion criteria: Maximum urinary flow rate ≥ 5 ml/s and ≤ 15 ml/s in a total voided volume of ≥ 150 ml and IPSS score of 12 or more.		IPSS Mean difference \pmSEM (95% CI) in change from baseline at the final visit for Group 1-Group 2 [least squares difference]	0.39 \pm 0.39 (-0.38, 1.15)	
Setting: Multi-centre, Scandinavia.	Exclusion criteria: Patients who had undergone prostate surgery, had a prostatic stent, or had undergone microwave thermotherapy were excluded, as were those who had had balloon dilation within the previous 6 months. Suspected or known malignancy and or PSA > 10 ng/ml; any known cause of urinary symptoms or reduced flow rate other than BPH; known acute urinary retention within the year, major residual urine, bladder stones, recurrent urinary tract infections, or large bladder diverticulum. Hepatic, renal, cardiac and gastrointestinal dysfunction or disease; uncontrolled diabetes, hypotension; and known allergy to study drugs. Use of prespecified drugs that might interfere with treatment or of an investigational drug or donation of blood 4 weeks prior to or during the study and conditions precluding good compliance were also cause for exclusion.	Group 1: Doxazosin Gastrointestinal therapeutic system (GITS) 4mg or 8mg once daily with a doxazosin standard placebo tablet. Initially 4mg dose given for at least 7 weeks. At week 7 the dose was increased to 8mg once daily if subjects had not experienced an increase in the maximum urinary flow are of at least 3ml/s and a 30% reduction in IPSS.	Mean (SE) adjusted change from baseline to final visit for Qmax (per-protocol analysis)	Group 1 (n=300): 2.6 ± 0.2 Group 2 (n=303): 2.2 ± 0.2 Group 3 (n=151): 0.8 ± 0.3	
Evidence level: 1+			Mean (SD) adjusted change from baseline to final visit for urinary flow (per-protocol analysis)	Group 1 (n=300): 1.2 ± 2.4 ; $p < 0.04$ Group 2 (n=303): 1.1 ± 2.0 ; $p < 0.05$ Group 3 (n=151): 0.6 ± 2.1	
Duration of follow-up: 13 weeks			Mean (SD) adjusted change from baseline to final visit for total quality of life IPSS question (per-protocol analysis) – least squares difference	Group 1 (n=310): -1.3 ± 0.1 Group 2 (n=311): -1.4 ± 0.1 Group 3 (n=151): -0.9 ± 0.1 $P < 0.001$	Notes: Mean changes are adjusted and can not be combined for meta-analysis.
	All patients N: 795	Group 2: Doxazosin standard 1 to 8mg once daily Initial dose 1 mg that was increased at the end of 1 week to 2mg, at week to	Adverse events	Dizziness Group 1: 18/317 (5.7%) Group 2: 27/322 (8.4%) Group 3: 3/156 (1.9%) Headache Group 1: 18/317 (5.7%) Group 2: 13/322 (4.0%) Group 3: 7/156 (4.5%) Asthenia Group 1: 10/317 (3.2%) Group 2: 16/322 5.0%) Group 3: 2/156 (1.3%) Vertigo Group 1: 8/317 (2.5%) Group 2: 24/322 (7.5%)	Per protocol analysis: Group 1 GITS: 44.2% remained at the 4mg and 55.8% received 8mg at the final visit. Group 2: doxazosin standard group 14.9% were receiving 2mg/day, 34% were on 4mg/day and 51.1% were receiving 8mg/day. Mean final dose for Group 1: 6.2mg/day

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>ITT analysis: 784 Per protocol analysis: 780 Mean age: 65.2 years Drop outs:</p> <p>Group 1 N: 317 ITT analysis =311 Mean (\pmSD) Age: 64.9 Baseline IPSS: 17.7\pm4.3 Race: White=311 Dropouts:22 (treatment related adverse events=11)</p> <p>Group 2 N: 322 (ITT analysis =318) Mean (\pmSD) Age: 65.3 Baseline IPSS: 17.8\pm4.5 Race: White=318 Dropouts:38 (treatment related adverse events=20; insufficient clinical response=1)</p> <p>Group 3 N: 156 (ITT analysis =155) Mean (\pmSD) Age: 65.4 Baseline IPSS: 18.0\pm4.3 Race: White=153; Asian=1; Other=1 Dropouts: 8 (treatment related adverse events=1)</p>	<p>4mg and at week 7 the dose was increased to 8mg once daily if required to achieve the target increasing urinary flow and decrease in IPSS.</p> <p>Group 3: Placebo once daily Received double-dummy matching placebo</p> <p>Study medications taken once daily at breakfast, except on study visit days, when medication was administered after study assessments.</p>	<p>Reduction from baseline IPSS of \geq30%</p> <p>Increase in maximum urinary flow rate \geq3ml/s</p> <p>Investigator s assessment of efficacy (intention to treat analysis)</p>	<p>Group 3: 1/156 (0.6%) Flu syndrome Group1: 4/317 (1.3%) Group 2: 6/322 (1.9%) Group 3: 7/156 (4.5%) Back pain Group1: 4/317 (1.3%) Group 2: 4/322 (1.2%) Group 3: 4/156 (2.6%) Postural hypotension Group1: 4/317 (1.3%) Group 2: 7/322 (2.2%) Group 3: 1/156 (0.6%) Nausea Group1: 3/317 (0.9%) Group 2: 8/322 (2.5%) Group 3: 1/156 (0.6%) Discontinuation - adverse events Group 1: 11 (3.5%) Group 2:20 (6.2%) Group 3: 1 (0.6%)</p> <p>Group1: 73.5% Group 2: 74.7% Group 3: 53.5%</p> <p>Group1: 38.8% Group 2: 38.7% Group 3: 21.4%</p> <p>Excellent or good rating Group 1: 193 (62.3%) Group 2:207 (65.5%) Group 3: 57 (37.5%) Poor rating Group 1: 39 (12.6%) Group 2:48 (15.2%) Group 3: 47 (30.9%)</p>	Group 2: 5.7mg/day

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Carbin et al., 1991⁴²</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: NR</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 8 weeks</p>	<p>Patient group: Males from 50 to 76 years of age with a known diagnosis of BPH.</p> <p>All patients N: 33 Drop outs: 3 (1 did not enter trial due to pneumonia, 2 discontinued treatment due to palpitations and tachycardia)</p> <p>Group 1 N: 16 Mean (\pmSD) Age: 68.7 (5.0) Prostatic size, g: 41 (15) Dropouts: 1</p> <p>Group 2 N: 16 Mean (\pmSD) Age: 64.6 (6.4) Prostatic size, g: 61 (40) Dropouts: 1</p>	<p>Group 1: Alpha-blocker Alfzozin 2.5mg X 3 If no effect of therapy noticed by the patient after 3 weeks of treatment and body weight more than 80kg the dose was increased to 4 tablets daily (e.g. 10mg).</p> <p>Group 2: Placebo</p>	<p>Mean urinary flow rate, ml/sec</p>	<p>Baseline Group 1: 8.1 (2.2) Group 2: 8.4 (3.0) 3 weeks Group 1: 9.2 (3.3) Group 2: 8.2 (3.8) 8 weeks Group 1: 8.9 (2.8) Group 2: 8.9 (3.4) P=NS</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, allocation concealment and blinding were unclear.</p> <p>Additional outcomes: Serum concentration, heart rate and blood pressure reported.</p> <p>Notes: Baseline number in each group not reported in methods. The table for adverse events reports that 15 in the intervention group.</p>
			<p>Timed micturition seconds</p>	<p>Baseline Group 1: 19.6 (13.1) Group 2: 23.9 (15.4) 3 weeks Group 1: 14.7 (10.4) Group 2: 22.6 (13.2) 5 weeks Group 1: 14.3 (9.8) Group 2: 23.9 (17.8) 8 weeks Group 1: 15.8 (11.7) Group 2: 21.8 (10.6) P=0.023</p>	
			<p>Residual urine</p>	<p>Baseline Group 1: 97.9 (115) Group 2: 92.7 (86) 3 weeks Group 1: 30.9 (32) Group 2: 114 (167) 8 weeks Group 1: 42.8 (51) Group 2: 94.2 (121) P=0.02</p>	
			<p>Frequency number</p>	<p>Baseline Group 1: 8.9 (3) Group 2: 10.7 (3.0) 3 weeks Group 1: 7.1 (2) Group 2: 10.4 (3)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				5 weeks Group 1: 8.6 (3) Group 2: 9.5 (3) 8 weeks Group 1: 7.4 (2) Group 2: 9.4 (3) P=NS	
			Boyarsky score	Baseline Group 1: 11.3 (3.0) Group 2: 11.7 (3.7) 3 weeks Group 1: 7.3 (3.0) Group 2: 8.9 (2.6) 5 weeks Group 1: 6.3 (3.2) Group 2: 7.9 (2.6) 8 weeks Group 1: 5.9 (3.6) Group 2: 7.1 (2.2) P=NS	
			% of patients that had the dose increased	Group 1: 27% Group 2: 47%	
			Patients/physicians correct guess of treatment given	Group 1: 60% / 60% Group 2: 67% / 58%	
			Adverse events	Vertigo Group 1: 3/15 Group 2: 2/15 Headache Group 1: 1/15 Group 2: 1/15 Weakness Group 1: 1/15 Group 2: 0/15 Weight gain Group 1: 1/15 Group 2: 0/15 Indigestion	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 2/15 Group 2: 0/15 Diarrhoea Group 1: 1/15 Group 2: 2/15 Constipation Group 1: 1/15 Group 2: 0/15 Dry mouth Group 1: 0/15 Group 2: 1/15 Dry hands Group 1: 1/15 Group 2: 0/15 Herpes simplex Group 1: 1/15 Group 2: 0/15 Conjunctivitis Group 1: 1/15 Group 2: 0/15	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Chapple et al., 1994⁵⁰</p> <p>Study design: Randomised controlled study</p> <p>Setting: Multi-centre, UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group:</p> <p>Inclusion criteria: Maximum urinary flow rate <15ml/s accompanied by symptoms of bladder outflow obstruction and in whom outflow obstruction at the level of the prostate was confirmed by means of videocystometry. Only patients with a functioning detrusor muscle were included (residual urine <200ml).</p> <p>Exclusion criteria: Patients with other conditions giving rise to urinary symptoms and reduced urine flow rates, such as carcinoma of the prostate. Previous prostatic surgery, serum creatinine >200mmol/l, poorly controlled diabetes, a history of myocardial infarction or a cerebrovascular accident within the preceding 6 months.</p> <p>All patients N: 135</p> <p>Group 1 N: 67 Mean (\pmSD) Age: 67 (7.3) Race: Caucasian=55, other=12 Dropouts: 7 (drop out during 2 week run-in=2, withdrew due to concomitant or associated illness=3; adverse events=2) Data for efficacy=60 [Evaluable in 2 of 12 that withdrew; inevaluable in 1 due to protocol violations]</p>	<p>Baseline evaluation: Lasting 2 weeks during which patients received one doxazosin or placebo tablet each morning.</p> <p>Group 1: Alpha-blocker Doxazosin commenced with daily dose 1mg, increased to 2mg after 2 weeks and to maximum of 4mg after 4 weeks</p> <p>Group 2: Placebo</p>	<p>Mean (SEM) maximum flow rate, ml/s</p>	<p>Baseline Group 1: 9.1 (0.5) Group 2: 9.1 (0.5) Change Group 1: 2.6 (0.7) Group 2: 1.1 (0.6) P=0.09</p>	<p>Funding: Pfizer provided medications and material support for study.</p> <p>Limitations: Method of randomisation and allocation concealment unclear.</p> <p>Additional outcomes: Maximum bladder capacity, volume of first unstable contraction, end filling pressure reported. Modified Boyarsky scale used to report obstructive and irritative symptoms but figures not provided.</p> <p>Notes: Headache and dizziness reported as most frequent side effects but actual figures not reported.</p>
			<p>Mean (SEM) maximum detrusor voiding pressure, cmH2O</p>	<p>Baseline Group 1: 78.5 (2.7) Group 2: 74.2 (4.6) Change Group 1: -4.6 (3.2) Group 2: 7.9 (3.0) P=0.007</p>	
			<p>Mean flow rate, ml/s</p>	<p>Baseline Group 1: 4.4 (0.3) Group 2: 4.3 (0.3) Change Group 1: 1.0 (0.3) Group 2: 0.2 (0.3) P=0.04</p>	
			<p>Number of reported adverse events in number of patients with adverse events</p>	<p>Group 1: 44/25 Group 2: 12/11</p>	
			<p>Withdrawn due to adverse events</p>	<p>Group 1: 2 Group 2: 0</p>	
			<p>% Improvement in symptoms (evaluation in response to questioning at tend of study)</p>	<p>Hesitancy Group 1: 59% Group 2: 26% P=0.003 Nocturia Group 1: 39% Group 2: 19% P=0.017 Urgency Group 1: 60%</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 N: 68 Mean (\pmSD) Age: 67 (7.5) Race: Caucasian=64, other4 Dropouts: 5 (drop out during 2 week run-in=1, withdrew due to concomitant or associated illness=4) Data for efficacy=62 [inevaluable in 2 due to protocol violations]</p>			<p>Group 2: 38% P=0.041 Impaired urinary stream Group 1: 56% Group 2: 33% P=0.019 Frequency Group 1: 44% Group 2: 27% P=0.062</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Chapple et al., 2005 ⁴⁹ Study design: RCT Setting: Multi national (18 countries), multi-centre (138 mainly European) Evidence level: 1+ Duration of follow-up: 12 weeks	<p>Patient group: Men with lower urinary tract symptoms suggestive of BPH.</p> <p>Inclusion criteria: Men aged 45 years or over with voiding and storage symptoms diagnosed as LUTS/BPH with a total IPSS ≥ 13 and a maximum flow rate ≥ 4ml/s and ≤ 12ml/s.</p> <p>Exclusion criteria: any other urological procedures or conditions what may cause LUTS ; patients with hepatic or renal insufficiency, clinically significant cardiovascular or cerebrovascular diseases within 6 months prior to enrolment, central nervous system conditions or life-threatening diseases. Patients taking or had taken other drugs for LUTS or were hypersensitive to $\alpha 1$ AR antagonists or their recipients, were taking drugs which could interfere with the pharmacodynamics of tamsulosin OCAS or were taking or had taken other investigational drugs within the previous 3 months.</p> <p>All patients N: 2152 Mean age: 65 years Mean IPSS: 18.5 Mean prostate volume: 43-45ml Drop outs: 107 (5%) due to treatment emergent adverse events=57, insufficient response=18, lost to follow-up=9, protocol violations=3, adverse</p>	<p>Group 1: Tamsulosin: Oral controlled absorption system 0.4mg once daily</p> <p>Group 2: Tamsulosin: Old modified release tamsulosin: 0.4mg once daily</p> <p>Group 3: Tamsulosin: Oral controlled absorption system 0.8mg once daily</p> <p>Group 4: placebo Placebo once daily</p>	<p>Mean (SD) IPSS at baseline</p> <p>IPSS reduction at endpoint</p> <p>Mean (SD) change at endpoint IPSS- QOL</p> <p>Investigator reported as slightly improved</p> <p>Investigator reported as much improved</p> <p>Treatment-emergent Adverse events attributable to alpha-blocker</p>	<p>Baseline: Group 1: 18.5 (4.4) Group 2: 18.5 (4.5) Group 3: 18.6 (4.5) Group 4: 18.3 (4.5)</p> <p>End point: Group 1 (n=355): 10.8 (6.2) Group 2 (n=703): 10.6 (5.9) Group 3 (n=709): 10.6 (5.9) Group 4 (n=351): 12.4 (6.4)</p> <p>Group 1 (n=354): -7.7 (5.8); p<0.001 Group 2 (n=700): -8.0 (5.6); p<0.001 Group 3 (n=707): -8.0 (5.9) Group 4 (n=350): -5.8 (5.6)</p> <p>Baseline: Group 1 (n=354): 3.8 (1.1) Group 2 (n=699): 3.8 (1.1) Group 3 (n=706): 3.8 (1.1) Group 4 (n=350): 3.8 (1.0)</p> <p>Change at endpoint: Group 1 (n=354): -1.4 (1.3) Group 2 (n=699): -1.4 (1.3) Group 3 (n=706): -1.4 (1.4) Group 4 (n=350): -1.1 (1.3)</p> <p>Group 1: 33.1% Group 2: 33.5% Group 3: 33.0% Group 4: 35.7%</p> <p>Group 1: 46.5% Group 2: 48.7% Group 3: 48.4% Group 4: 35.7%</p> <p>Non cardiovascular Group 1: 16 (4.4%) Group 2: 36 (5.1%) Group 3: 57 (7.9%) Group 4: 7 (2.0%) Cardiovascular</p>	<p>Funding: NR.</p> <p>Limitations: None.</p> <p>Additional outcomes: Blood pressure was reported.</p> <p>Notes: Additional information retrieved from the authors.</p> <p>Outcomes reported for group 1 and 2 combined for meta-analysis by NCGC.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>events starting during the placebo run in =3, death=3, abnormal laboratory values=1, non-specified reasons=13</p> <p>Group 1 N: 361 Dropouts:18</p> <p>Group 2 N: 710 Dropouts: 25</p> <p>Group 3 N: 724 Dropouts: 45</p> <p>Group 4 N: 357 Dropouts: 19</p>			<p>Group 1: 9 (2.5%) Group 2: 23 (3.2%) Group 3: 28 (3.9%) Group 4: 8 (2.2%) All: Group 1: 25 (6.9%) Group 2: 55 (7.8%) Group 3: 80 (11.1%) Group 4: 13 (3.7%)</p> <p>Number (%) Dizziness Group 1: 5/360 (1.4%) Group 2: 9/709 (1.3%) Group 3: 17/722 (2.4%) Group 4: 5/356 (1.4%)</p> <p>Number (%) Retrograde ejaculation Group 1: 6/360 (1.7%) Group 2: 10/709 (1.4%) Group 3: 18/722 (2.5%) Group 4: 1/356 (0.3%)</p> <p>Number (%) of at least one Treatment-emergent adverse events Group 1: 93/360 (26.0%) Group 2: 168/709 (24.0%) Group 3: 192/722 (27.0%) Group 4: 71/356 (20.0%)</p> <p>Number (%) at least one treatment-related adverse events Group 1: 40/360 (11.0%) Group 2: 82/709 (12.0%) Group 3: 103/722 (14.0%) Group 4: 25/356 (7.0%)</p> <p>% Responders (defined as patients who had at least a 25% improvement in total IPSS vs. baseline) Group 1: 71.2% Group 2: 75.4% Group 3: 73.8% Group 4: 60.9%</p> <p>Serious adverse events Group 1: 7/360 Group 2: 9/709 Group 3: 12/722 Group 4: 3/356</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Discontinuation due to adverse events	Group 1: 14/360 Group 2: 11/709 Group 3: 28/722 Group 4: 6/356	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Christensen et al., 1993⁵³</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Denmark</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 9 weeks</p>	<p>Patient group: consecutive patients from Feb 1988-May 1989 referred to the out patient clinics of the 2 participating surgical departments for BPH.</p> <p>Inclusion criteria: All had moderate or severe symptoms resulting from infravesical obstruction, an obstructive flow curve pattern as determined by uroflowmetry and were candidates for TURP.</p> <p>Exclusion criteria: previous prostatic/bladder neck surgery, suspicion of prostatic cancer on DRE, non-prostatic obstruction on the urethra, overflow incontinence, renal dysfunction, positive urine cytology, hematuria, urinary infection, symptomatic hypotension, previous or present cerebrovascular disease, history of intolerance to doxazosin, prazosin or other quinazolines, current treatment with alpha adrenoceptor blocking agents, severe psychiatric or neurologic disease.</p> <p>All patients N: 100 Drop outs: 9</p> <p>Group 1 N: 52 Mean (\pmSD) Age: 66.7 (7.9) Dropouts: 4 (diabetes=1, withdrew consent=2, urinary tract infection=1)</p>	<p>Run-in period One week</p> <p>Group 1: alpha-blocker Doxazosin once daily at bed time. 1mg week 1, 2mg week 2-5 and 4mg week 6-9.</p> <p>Group 2: Placebo Once daily at bedtime</p>	<p>Mean (SEM) maximum urinary flow rate (estimated from graph)</p>	<p>Baseline Group 1 (n=52): 7.6 (SD 3.7) Group 2 (n=48): 7.5 (SD 3.5)</p> <p>0 weeks Group 1 (n=46): 7.4 Group 2 (n=43): 8.0</p> <p>5 weeks Group 1 (n=47): 9.5 (0.7) Group 2 (n=42): 9.1 (0.8)</p> <p>9 weeks Group 1 (n=46): 9.4 (0.7) Median improvement: 1.5 (range: -9.0, 22.0) Group 2 (n=42): 8.0 (0.5) Median improvement: -0.3 (-7.0 to 7.2)</p>	<p>Funding: NR</p> <p>Limitations: Method of allocation concealment unclear.</p> <p>Additional outcomes: Mean urinary flow rate – reported but actual figures not provided. Changes in blood pressure and weight were reported.</p> <p>Notes: Maximum urinary flow rates were estimated from a graph.</p>
			<p>Median reduction in voiding frequency chart (3 days average 24-hour voiding frequencies)</p>	<p>9 weeks Group 1: 2.3 Group 2: 1.2 P=0.005</p>	
			<p>Median (range) baseline and change in frequency (daytime)</p>	<p>Baseline Group 1 (n=52): 8 (3/18) Group 2 (n=48): 7 (3/16)</p> <p>Week 9 Group 1 (n=48): -1.5 (-9/3) Group 2 (n=43): 0.3 (-7/7) P=0.001</p>	
			<p>Median (range) baseline and change in nocturia</p>	<p>Baseline Group 1 (n=52): 2.5 (0/6) Group 2 (n=48): 2.5 (0/7)</p> <p>Week 9 Group 1 (n=48): -1.1 (-4/1) Group 2 (n=43): -1.0 (-4/1) P=0.12</p>	
			<p>Baseline and change in residual urine</p>	<p>Baseline Group 1 (n=52): 100 (10/450) Group 2 (n=48): 85 (10/340)</p> <p>Week 9</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 N: 48 Mean (\pmSD) Age: 68.1 (7.4) Dropouts: 5 (S-creatinine > 130 micromoles/l, withdrawn due to side effects=2, urinary retention=1, lost to follow-up=1).</p>			Group 1 (n=48): -15.0 (-430/150) Group 2 (n=43): -1.0 (-305/355) P=0.56	
			Median (range) Bladder capacity (ml)	Baseline Group 1 (n=52): 288 (134/490) Group 2 (n=48): 271 (124/660) Week 9 Group 1 (n=48): 0.0 (-228/197) Group 2 (n=43): 3.0 (-297/159) P=0.34	
			Number of symptoms improved (%) - all symptoms pooled for each group	Baseline: Group 1: 239 Group 2: 270 Week 9: Group 1: 159 (67) Group 2: 95 (35) P=0.023	
			Number of obstructive symptoms improved (%) - all symptoms pooled for each group	Baseline: Group 1: 177 Group 2: 196 Week 9: Group 1: 112 (63) Group 2: 62 (32) P=0.015	
			Number of irritative symptoms improved (%) - all symptoms pooled for each group	Baseline: Group 1: 62 Group 2: 74 Week 9: Group 1: 47 (76) Group 2: 33 (45) P=0.12	
					Group 1: 11 patients reported 13 events Group 2: 10 patients reported 11 events P=Not sign Dizziness/vertigo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1:5 Group 2: 5 (2 withdrew due to dizziness)	
			Patients subjective overall assessment at 9 weeks	Group 1 Much worse: 0/48 Worse: 1/48 Unchanged: 9/48 Better: 28/48 Much better: 10/48 Group 2 Much worse: 1/43 Worse: 0/43 Unchanged: 23/43 Better: 12/28 Much better: 7/43	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Djavan et al., 2005D⁷³</p> <p>Study design: RCT</p> <p>Setting: European multi-centre (3 countries)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 8 week</p>	<p>Patient group: Men aged 45 years or over with voiding and storage symptoms diagnosed as LTUS/BPH.</p> <p>Inclusion criteria: After a 2 week placebo run in, men 45 years or older, with lower urinary tract symptoms (IPSS: 13 or above suggestive of BPH (maximum flow rate 4-12ml/s and 2 or more nocturnal voids per night.</p> <p>Exclusion criteria: any other urological procedures or conditions, which may cause LUTS; hepatic or renal insufficiency, clinically significant cardiovascular or cerebrovascular diseases within six months prior to enrolment, central nervous system conditions or life-threatening diseases. Alcohol consumption of more than 15 units per week; post voiding residual volume of >250ml in at least two assessment over the last 3 months. Patient taking or had taken other drugs for BPH; hypersensitive to alpha-blockers, were taking drugs with could interfere with the pharmacodynamics of tamsulosin or were taking or had taken over investigational drugs within previous 3 months.</p> <p>All patients N: 117 Mean age: 67</p>	<p>Group 1: Alpha-blocker Tamsulosin oral controlled absorption system 0.4mg once daily</p> <p>Group 2: Placebo</p>	<p>Mean (SD) IPSS symptom scores</p>	<p>Baseline Group 1: 18.2 (4.0) Group 2: 18.1 (3.3) Change at endpoint Group 1: -8.0 (5.2) Group 2: -5.6 (4.7) Difference: 2.4; p=0.0099</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation and allocation concealment was unclear.</p> <p>Additional outcomes: Analysis of IPSS by subgroup of voiding and storage symptoms.</p> <p>Notes: None.</p>
			<p>Mean change in nocturia question on IPSS questionnaire</p>	<p>Group 1: 1.1 Group 2: 0.7 Difference: 0.4; p=0.028</p>	
			<p>Mean IPSS quality of life question reduction at endpoint</p>	<p>Group 1: 2.0 Group 2: 1.3 OR: 2.4; p=0.0087</p>	
			<p>Adverse events</p>	<p>Treatment-emergent adverse events (TEAE) Group 1 (n=61): 10 Group 2 (n=56): 8 At least one TEAE Group 1: 5 (8.2%) Group 2: 7 (12.5%) Dizziness Group 1: 2 (3.3%) Group 2: 0 Nasopharyngitis Group 1: 0 Group 2: 2 (3.4%) Orthostatic hypotension Group 1: 0 Group 2: 0 Discontinuations due to AE Group 1: 0 Group 2: 0</p>	
			<p>Mean change in total hours of undisturbed sleep (defined as time between falling asleep</p>	<p>Group 1: 81 minutes (60%) Group 2: 60 minutes (40%) Difference: 21 minutes; p=0.198</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 1 N: 61 Mean (\pmSD) Age: 66.8 (8.5) Baseline IPSS: 19.0 (5.1) Dropouts: 1 (discontinued due to non compliance)</p> <p>Group 2 N: 56 Mean (\pmSD) Age: 67.6 (7.6) Baseline IPSS: 18.1 (3.5) Dropouts: 0</p>		<p>and first awakening to void)</p> <p>Mean decrease in nocturnal voids as measured by means of voiding diary (defined as time between falling asleep and first awakening to void)</p> <p>Questionnaire to assess level of tiredness or alertness during the day (not validated)</p> <p>Correlation between number of nocturnal void and the hours undisturbed sleep</p> <p>Correlation between IPSS nocturia and IPSS QoL domains</p>	<p>Group 1: 1.0 Group 2: 0.7 OR: 0.56; p=0.099</p> <p>Group 1: 0.49 Group 2: 0.32 OR: 0.672; p=.27</p> <p>Spearman's rank coefficient: -0.63</p> <p>Spearman's rank coefficient: 0.64</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Fawzy et al., 1995⁸⁹</p> <p>Study design: RCT</p> <p>Setting: Multi-centre, US.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 16 week</p>	<p>Patient group: normotensive patients (sitting diastolic blood pressure <90mm.Hg) with BPH.</p> <p>Inclusion criteria: AUA of 10 or greater , maximum urinary flow rate of 5-15ml/s in a voided volume of 125-500ml and post void residual volume of 250ml or less on 2 consecutive weeks of the placebo run in period. aged 45 years or over</p> <p>Exclusion criteria: recent urinary retention, sever outflow obstruction, or non BPH conditions that caused obstruction or symptoms. Patients who had serious concurrent disease, history of clinically significant cardiovascular, hepatic or renal dysfunction, poorly controlled diabetes, urinary calculi or intolerance/sensitivity to quinazoline derivatives.</p> <p>All patients N: 100 Race: 96% white, 2% Asian, 1% Hispanic and 1% Black. Drop outs: 2 (did not undergo any efficacy measurement). Patient withdrawal: 22</p> <p>Group 1 N: 50 Mean (\pmSD) Age: 62.1 (7.8) Withdrawals: 11 (adverse events – related and unrelated=7; other=4)</p>	<p>Placebo run-in: 2 weeks</p> <p>Group 1: Alpha-blocker Doxazosin: 8 week dose titration phase the initial dose of doxazosin was 1mg, increasing to 2mg, 4mg, or 8mg at 2-week intervals until the optimum dose was attained. During the final 6-week phase of the study the dose was held constant at the optimum level.</p> <p>41 patients in the study dosage was titrated to a maximally efficacious s and/or tolerated, stable level of doxazosin; 36 reached dose of 8mg, 1 reached a daily dose of 4mg and 4 reached a daily dose of 2mg.</p> <p>Group 2: Placebo</p>	<p>Mean change in AUA6 symptom score</p> <p>Mean change from baseline in Qmax, ml/s</p> <p>Mean change from baseline in average urinary flow rate, ml/s</p> <p>Percent improvement in patient assessed symptoms (AUA)</p> <p>Adverse events</p>	<p>Group 1: -5.7 Group 2: -2.5 P<0.001</p> <p>Group 1: 2.9 Group 2: 0.7 P<0.01</p> <p>Group 1: 1.4 Group 2: 0.3 P<0.01</p> <p>Total symptoms Group 1: 39 Group 2: 17</p> <p>Obstructive symptoms Group 1: 43 Group 20</p> <p>Irritative symptoms Group 1: 35 Group 2: 15</p> <p>Total Group 1: 44% Group 2: 30%</p> <p>Events in patients over 65 years Group 1: 28% Group 2: 37%</p> <p>Discontinuation due to adverse events Group 1: 1 Group 2: 0</p> <p>Dizziness Group 1: 15/50 Group 2: 2/50</p> <p>Fatigue Group 1: 6/50 Group 2: 2/50</p> <p>Headache Group 1: 6/50 Group 2: 2/50</p> <p>Somnolence</p>	<p>Funding: Pfizer</p> <p>Limitations: Method of randomisation and allocation concealment unclear.</p> <p>Frequency of nocturia significantly greater in placebo arm.</p> <p>Additional outcomes: Graphical presentation of Qmax by week. Intervention arm significantly improved compared to placebo by 2 weeks. Boyarsky modified score also reported.</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 N: 48 Mean (\pmSD) Age: 61.6 (8.7) Withdrawals: 11 (adverse events – related and unrelated=1; patient request=3; protocol violation=4; entry criteria not met=1; other=2)</p>			<p>Group 1: 5/50 Group 2: 2/50 Hypotension Group 1: 4/50 Group 2: 0 Nausea Group 1: 4/50 Group 2: 0</p>	
			<p>Mean sitting blood pressure change, mmHg</p>	<p>Group 1: -5.6/-4.1 Group 2: 0.7/-0.4 P<0.05</p>	
			<p>Mean standing blood pressure change, mmHg</p>	<p>Group 1: -6.0/-4.5 Group 2: 1.9/-0.4 P<0.05</p>	
			<p>Mean change in daytime micturition frequency from patient daily diary</p>	<p>Group 1: -1.3 Group 2: -0.7 P=0.043</p>	
			<p>Mean change in nocturia frequency from patient daily diary</p>	<p>Group 1: -0.5 Group 2: -0.5 P=0.470</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gillenwater et al., 1995¹⁰¹</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Multi-centre, USA</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 16 weeks</p>	<p>Patient group: men 45 years or older with BPH and mild to moderate essential hypertension.</p> <p>Inclusion criteria: maximum urinary flow rate of 5-15ml/s in a voided volume of 150-500ml, post void residual volume of less than 200ml, daytime micturition frequency of 4 or more, nocturia of more than 2 times per night and a sitting diastolic blood pressure of 90-114 mm.Hg.</p> <p>Exclusion criteria: Any other conditions causing urinary symptoms or decreased flow rate, previous or imminent prostatic surgery, prostate specific antigen level greater than 10ng/ml, acute urinary retention, recent catheterisation for outflow obstruction or prostate malignancy were excluded from the study. Insulin-dependent or poorly controlled noninsulin-dependent diabetes, significant hepatic, renal or cardiovascular dysfunction; secondary hypertension, concurrent serious disease or malignancy, or significant psychiatric disorders. Intolerance/sensitivity to quinazoline derivatives, substance abuse, recent blood donation, obesity, antihypertensive drug therapy or any treatment known to affect vesicourethral function, and recent therapy with any other investigational drug or any prior doxazosin therapy.</p>	<p>Screening: 0-4 week period allowed for the discontinuation and wash out of excluded medication, including any other antihypertensive agents.</p> <p>Placebo- run in phase: 2 weeks.</p> <p>Group 1: Alpha-blocker Doxazosin 2, 4, 8 or 12mg once daily in the morning. The initial dose was 1mg, increasing sequentially at weekly intervals during a 5-week titration phase to the randomised, fixed dose level. The dose then remained constant during the 9-week efficacy phase.</p> <p>Group 2: Placebo</p>	<p>Mean (SD) Qmax at trough and peak measurements, ml/s</p> <p>Trough defined as assessment approximately 24 hours following the previous morning dose. Peak defined as assessment 2 -6 hours following administration of medication</p>	<p>Trough</p> <p>Group 1: 2mg (n=39): 10.5 (2.1) 4mg (n=46): 9.8 (2.0) 8mg (n=45): 10.7 (2.1) 12mg (n=45): 10.5 (2.2) Group 2 (n=41): 10.3 (2.3)</p> <p>Peak</p> <p>Group 1: 2mg (n=39): 10.1 (2.7) 4mg (n=46): 9.4 (2.9) 8mg (n=45): 10.3 (2.6) 12mg (n=45): 9.7 (2.4) Group 2 (n=41): 10.5 (2.6)</p>	<p>Funding: Gillenwater, Conn, Chrysant and Roy, and the Multicenter Study Group have participated in clinical studies sponsored by Pfizer Central Research, New York.</p> <p>Limitations: Method of randomisation and allocation concealment unclear. Method states that compliance assessed by tablet count of returned medication – results not reported.</p> <p>Additional outcomes: Obstructive and irritative sub-groups results for Boyarsky score. Qmax also reported as adjusted mean change.</p> <p>Notes: Boyarsky score was reversed so that lower scores indicated improvement, as with other commonly used symptom scores.</p> <p>Treatment effect tested for significance after adjusting for the</p>
			<p>Patients with ≥ 3ml/s increase in Qmax</p>	<p>Trough</p> <p>Group 1: 8mg: 37% 2mg: 39% Group 2: 13%</p> <p>Peak</p> <p>Group 1: 8mg: 42% 2mg: 51% Group 2: 17% * 2mg and 4mg Not sig.ly different from placebo group</p>	
			<p>Mean (adjusted) change in average flow rate (* significantly different from placebo p<0.05, ** p<0.01)</p>	<p>Trough</p> <p>Group 1: 2mg: 0.6 4mg: 0.6 8mg: 1.5** 12mg: 1.3* Group 2: 0.2</p> <p>Peak</p> <p>Group 1: 2mg: 0.9</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients N: 248 Efficacy analysis Group 1: 175 Efficacy analysis Group 2: 41 Drop outs: 32 (no efficacy follow-up measurements=7; not meet inclusion criterion for maximum urinary flow rate=25).</p> <p>Group 1 N: 199 Efficacy analysis: 175 2mg: 39 4mg: 46 8mg: 45 12mg: 45 Mean (±SD) Age: Dropouts: 69 (adverse events 11%, lack of blood pressure efficacy 7%, and protocol violations 9%)</p> <p>Group 2 N: 49 Efficacy analysis: 41 Mean (±SD) Age: 64.5 (7.7) Dropouts: 18 (adverse events 4%, lack of blood pressure efficacy 12%, lack of BPH efficacy 4% and protocol violations 10%)</p>		<p>4mg: 1.1 8mg: 1.6** 12mg: 2.1** Group 2: 0.2</p> <p>BPH symptom questionnaire (modified Boyarsky) mean change from baseline (adjusted for baseline effect) Key: * significantly different from placebo mean changes, p<0.01; \$significantly different from placebo mean changes, p<0.05</p> <p>% of patients with adverse events</p>	<p>2mg: 0.2 4mg: 0.2 8mg: 0.2 12mg: 0.2 Group 2: 0.2</p> <p>End point analysis of severity Group 1 2mg (n=34): -2.8 4mg(n=38): -5.0* 8mg(n=42): -4.2\$ 12mg(n=39): -3.6 Group 2 (n=37): -0.25</p> <p>End point analysis of bothersomeness Group 1 2mg (n=34): -3.4 4mg (n=38):-5.3\$ 8mg (n=42): -4.7 12mg (n=39): -4.9 Group 2 (n=37): -3.0</p> <p>Total Group 1 (n=199): 48% Group 2 (n=49): 35%</p> <p>Dizziness Group 1 (n=199): 19% Group 2 (n=49): 4%</p> <p>Headache Group 1 (n=199): 14% Group 2 (n=49): 18%</p> <p>Fatigue Group 1 (n=199): 10% Group 2 (n=49): 0%</p> <p>Hypotension Group 1 (n=199): 2.5% Group 2 (n=49): NR</p> <p>Withdrawal due to adverse events Group 1 (n=199): 11.1% Group 2 (n=49): 4.1%</p>	<p>baseline effect.</p> <p>Intervention at 1 week of treatment with 1mg dose - Qmax +0.8ml/s.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Hansen et al., 1994¹¹²</p> <p>Study design: RCT</p> <p>Setting: Multi-centre, Denmark and Netherlands</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: Men with BPH enrolled from November 1991 to March 1993.</p> <p>Inclusion criteria: Madsen-Iversen symptom score >6; urinary peak flow rate <10ml/s with a voided volume of at least 100ml. Men with very low urinary flow rates were included.</p> <p>Exclusion criteria: patients whose digital rectal examination suggested presence of prostatic cancer, or patients suffering from other urological diseases such as neurogenic bladder, urethral stricture, current urinary tract infection, macroscopic or microscopic hematuria, prostatitis or previous prostatectomy were excluded. Incidence of total urinary retention, history of bladders tones, repeated urinary tract infections, overflow incontinence, azotemia, abnormal acid phosphatase, a history of orthostatic hypotension or known hypersensitivity to alpha-blockers.</p> <p>All patients: N: 205 Mean age: 45-81</p> <p>Group 1</p>	<p>Run-in phase: All patients entered a four week placebo run-in phase. Single blind.</p> <p>Group 1: Alpha-blocker Alfuzosin 2.5mg TID</p> <p>Group 2: Placebo Three times a day</p>	<p>Median (25% and 75% quartiles) Madsen-Iversen symptom score</p>	<p>Baseline Group 1: 7 (6-8.5) Group 2: 7 (6-9) 12 weeks Group 1: 5 (3.5-7) Group 2: 6 (5-7.5)</p>	<p>Funding: Research grant from Synthelabo International.</p> <p>Limitations: Method of randomisation and allocation concealment was not reported.</p> <p>Additional outcomes: Blood pressure reported. Small but significant decrease in diastolic blood pressure in alfuzosin group compared to placebo.</p> <p>Notes: None</p>
			<p>Median (25% and 75% quartiles) peak flow rate, ml/s</p>	<p>Baseline Group 1: 9 (7-11) Group 2: 9 (7-11) 12 weeks Group 1: 11 (7.6-13.5) Group 2: 10 (8-11)</p>	
			<p>Median (25% and 75% quartiles) residual urinary volume, ml</p>	<p>Baseline Group 1: 50 (20-89) Group 2: 42 (20-100) 12 weeks Group 1: 30 (15-80) Group 2: 45 (15-80)</p>	
			<p>Adverse events – vasodilatory events</p>	<p>Dizziness Group 1: 3 Group 2: 0 Headache Group 1: 2 Group 2: 2 Postural hypotension Group 1: 1 Group 2: 0 Fatigue Group 1: 1 Group 2: 1 Syncope Group 1: 2 Group 2: 0</p>	
			<p>Adverse events – gastro-intestinal disorders</p>	<p>Nausea Group 1: 2 Group 2: 1 Diarrhoea Group 1: 4</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 104 (91 completed study) Median (\pmSD) Age: 65 (47-81) Withdrawals: 5 (lost to follow-up=1; adverse event=1; other=3)</p> <p>Group 2 N: 101 (87 completed study) Median (\pmSD) Age: 64 (45-81) Withdrawals: 12 (lack of efficacy=4; lost to follow-up=2; adverse events=1; other=5)</p>		<p>Adverse events – urinary tract disorders</p> <p>Other adverse events (including pain in arm, lymph disease, pneumonia, hypertension)</p> <p>Discontinuation due to adverse events</p>	<p>Group 2: 1 Vomiting Group 1: 0 Group 2: 0 Pyrosis Group 1: 1 Group 2: 0 Abdominal pain Group 1: 5 Group 2: 0 Obstipation Group 1: 0 Group 2: 1 Flatulence Group 1: 1 Group 2: 0 Haematemesis Group 1: 1 Group 2: 0</p> <p>Cystitis Group 1: 1 Group 2: 0 Urinary tract infection Group 1: 0 Group 2: 0 Hameatura Group 1: 0 Group 2: 0</p> <p>Group 1: 2 Group 2: 9</p> <p>Group 1: 1 Group 2: 1</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kaplan et al., 2006¹³⁶</p> <p>Also reported in Kaplan2008¹³⁴ and Rovner2008A²⁶⁴</p> <p>Study identifier: NCT00147654</p> <p>Study design: RCT, Double blind Patients, investigators and researchers masked to treatment allocation</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Men with overactive bladder or other LUTS recruited between Nov 2004 – Feb 2006</p> <p>Setting: multi-centre, USA</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • ≥ 40 years • IPSS ≥ 12 • Self-rated bladder condition of 'some moderate problems', 'severe problems' or 'many severe problems' based on the validated Patient Perception of Bladder Condition questionnaire. • Micturition frequency ≥ 8/24 hrs and urgency ≥ 3/24 hrs for ≥ 3 months <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Clinically significant bladder outlet obstruction defined as PVR ≥ 200 mL and Qmax < 5 mL/s • Serum PSA > 10 ng/mL with risk of prostate cancer • History of postural hypotension or syncope • Significant hepatic or renal disease • Neurological conditions such as MS, spinal cord injury and Parkinson disease • Prostate cancer • Prostate surgery or other intervention • History of acute urinary retention requiring catheterisation 	<p>Group 1: Tolterodine ER 4mg/day in evening</p> <p>Group 2: Tamsulosin 0.4 mg/day in evening</p> <p>Group 3: Tolterodine ER 4mg + Tamsulosin 0.4 mg/day in evening</p> <p>Group 4: Placebo in evening</p> <p>Examination methods: A Perception of Treatment Benefit question was posed at weeks 1, 6 and 12. "Have you had any benefit from your treatment? – YES/NO" and if so "How much benefit (little/a lot)?" Bladder diaries for 5 days were assessed prior to each visit at baseline and weeks 1, 6 and 12. IPSS measured at baseline and weeks 1, 6 and 12. PVR and Qmax measured at baseline and at week 12.</p>	<p>Change in IPSS from baseline at 12 weeks (estimated from graph) Analysis of covariance with covariates – smoking status, age, baseline score, duration of OAB, centre</p>	<p>Grp 1: -6.7 ± NR, n=206 Grp 2: -7.6 ± NR, n=197 Grp 3: -8.0 ± NR, n=203 Grp 4: -6.2 ± NR, n=213 P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: =0.007 Grp 3 vs Grp 4: =0.003</p>	<p>Funding: Pfizer</p> <p>Limitations: Incomplete reporting of outcomes:</p> <ul style="list-style-type: none"> ▪ Only the statistical significance of Combination vs placebo was reported. The statistical significance of difference between active arms unknown ▪ There were inconsistencies in the results reported within the paper ▪ Standard deviations were not reported. <p>Additional outcomes: Number of patients reporting treatment benefit from Perception of Treatment Benefit Question: Grp 1: 136/217 Grp 2: 146/215 Grp 3: 172/225 Grp 4: 132/222 Not sig, except : Grp 1 v Grp 3 p value 0.02, Grp 3 v Grp 4 p value 0.01 Pair wise analysis using Fishers 2 sided test</p>
			<p>Change in IPSS QoL from baseline at 12 weeks (estimated from graph) Analysis of covariance with covariates – smoking status, age, baseline score, duration of OAB, centre</p>	<p>Grp 1: -1.4 ± NR, n=206 Grp 2: -1.4 ± NR, n=198 Grp 3: -1.6 ± NR, n=205 Grp 4: -1.2 ± NR, n=213 P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: not sig Grp 3 vs Grp 4 =0.003</p>	
			<p>Change in Qmax from baseline at 12 weeks Analysis of covariance with covariates – centre, treatment, baseline value.</p>	<p>Grp 1: -0.60 ± NR Grp 2: -0.22 ± NR Grp 3: 0.07 ± NR Grp 4: -0.53 ± NR P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: not sig Grp 3 vs Grp 4: not sig.</p>	
			<p>Change in urgency incontinence/24h from baseline at 12 weeks (estimated from graph) Analysis of covariance with covariates – treatment, centre, PVR, Qmax and baseline value</p>	<p>Grp 1: -0.7. ± NR [n=48] Grp 2: -0.8 ± NR [n=46] Grp 3: -0.9 ± NR [n=47] Grp 4: -0.3 ± NR [n=43] P values: Grp 1 vs Grp 4= 0.008 Grp 2 vs Grp 4: Not sig Grp 3 vs Grp 4 p value =0.005</p>	
<p>Change in urgency episodes/24h from baseline at 12 weeks</p>	<p>Grp 1: -2.9 ± NR, n=209 Grp 2: -2.4 ± NR, n=205 Grp 3: -3.3 ± NR, n=211</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																																																					
	<ul style="list-style-type: none"> BOO due to diseases other than BPH Any condition for which antimuscarinics are contraindicated Men treated with alpha-blockers with 2 weeks or antimuscarinics, phytotherapy or electrical stimulation within 1 month, any investigational drug within 2 months or 5-alpha reductase within 3 months <p>All patients N: 879 out of 1531 evaluated Mean age: 61.8±9.9 Drop outs: 851/879 included in efficacy analysis, 754 /879 completed the study IPSS ± SD: 19.9±5.3 IPSS QoL ± SD: 4.57 ± 0.93 Qmax ± SD, mL/s: 12.9 ± 7.2</p> <p>Group 1 (Tolterodine ER) N: 217 (baseline data/efficacy analysis for N=210) Mean (± SD) Age: 61.8 ± 9.6 (range 41-91) Urge urinary incontinence : 53/217 Urgency episodes/24h: 7.58 ± 3.49 Micturitions/24h: 11.79 ± 2.83 Micturitions/night: 1.97 ± 1.27 IPSS ± SD: 19.53 ± 5.15 IPSS QoL ± SD: 4.57 ± 0.94 Qmax ± SD, mL/s: 13.3 ± 7.8 PVR ± SD, mL: 50.5 ± 55.8 Dropouts: 28/217 (12.9%) 1 patient did not receive study medication</p>		<p>(estimated from graph) Analysis of covariance with covariates – treatment, centre, PVR, Qmax and baseline value</p> <p>Change in micturitions/24h from baseline at 12 weeks (estimated from graph) Analysis of covariance with covariates – treatment, centre, PVR, Qmax and baseline value</p> <p>Change in micturitions/night from baseline at 12 weeks (estimated from graph) Analysis of covariance with covariates – treatment, centre, PVR, Qmax and baseline value</p>	<p>Grp 4: -2.5 ± NR, n=210 P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: not sig Grp 3 vs Grp 4: = 0.03</p> <p>Grp 1: -1.7 ± NR, n=209 Grp 2: -1.8 ± NR, n=205 Grp 3: -2.5 ± NR, n=211 Grp 4: -1.4 ± NR, n=212 P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: not sig Grp 3 vs Grp 4: <0.001</p> <p>Grp 1: -0.36 ± NR, n=209 Grp 2: -0.54 ± NR, n=205 Grp 3: -0.59 ± NR, n=209 Grp 4: -0.39 ± NR, n=212 P values: Grp 1 vs Grp 4: not sig. Grp 2 vs Grp 4: not sig Grp 3 vs Grp 4: =0.02</p>	<p>Notes: The study reported the adverse events based on the safety population, ie patients who had received at least one dose of the allocated treatment.</p> <p>The average IPSS score puts the patients in the study in the severely symptomatic category</p> <p>Sample size based on projected treatment difference of 15% between Tolterodine ER + Tamsulosin group compared to placebo for number of patients reporting treatment benefit at week 12.</p>																																																																																					
			<p>Reasons for discontinuation</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> <th>Grp 4</th> </tr> </thead> <tbody> <tr> <td>Adverse event</td> <td>5</td> <td>7</td> <td>20</td> <td>7</td> </tr> <tr> <td>Lack of efficacy</td> <td>8</td> <td>0</td> <td>4</td> <td>7</td> </tr> <tr> <td>Withdrew consent</td> <td>9</td> <td>9</td> <td>2</td> <td>5</td> </tr> <tr> <td>Protocol deviation</td> <td>2</td> <td>4</td> <td>0</td> <td>4</td> </tr> <tr> <td>Lost to follow up</td> <td>1</td> <td>4</td> <td>6</td> <td>4</td> </tr> <tr> <td>Death</td> <td>1</td> <td>0</td> <td>0</td> <td>0</td> </tr> <tr> <td>Other</td> <td>1</td> <td>5</td> <td>2</td> <td>5</td> </tr> </tbody> </table>		Grp 1	Grp 2	Grp 3	Grp 4	Adverse event	5	7	20	7	Lack of efficacy	8	0	4	7	Withdrew consent	9	9	2	5	Protocol deviation	2	4	0	4	Lost to follow up	1	4	6	4	Death	1	0	0	0	Other	1	5	2	5	<table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> <th>Grp 4</th> </tr> </thead> <tbody> <tr> <td>All cause adverse events</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>N</td> <td>216</td> <td>215</td> <td>225</td> <td>220</td> </tr> <tr> <td>Constipation</td> <td>9</td> <td>2</td> <td>8</td> <td>5</td> </tr> <tr> <td>Diarrhoea</td> <td>7</td> <td>6</td> <td>5</td> <td>3</td> </tr> <tr> <td>Dizziness</td> <td>3</td> <td>12</td> <td>6</td> <td>2</td> </tr> <tr> <td>Dry mouth</td> <td>16</td> <td>15</td> <td>47</td> <td>5</td> </tr> <tr> <td>Dyspepsia</td> <td>2</td> <td>1</td> <td>3</td> <td>5</td> </tr> <tr> <td>Ejaculation failure</td> <td>0</td> <td>4</td> <td>7</td> <td>0</td> </tr> </tbody> </table>		Grp 1	Grp 2	Grp 3	Grp 4	All cause adverse events					N	216	215	225	220	Constipation	9	2	8	5	Diarrhoea	7	6	5	3	Dizziness	3	12	6	2	Dry mouth	16	15	47	5	Dyspepsia	2	1	3	5	Ejaculation failure	0	4	7	0	<p>Randomisation sequence using block method prepared by statistician. Study medication kits were identical in appearance and smell.</p> <p>Missing data imputed for treatment benefit question (YES/NO), bladder diary variables, IPSS and IPSS QoL using Last</p>
	Grp 1	Grp 2	Grp 3	Grp 4																																																																																						
Adverse event	5	7	20	7																																																																																						
Lack of efficacy	8	0	4	7																																																																																						
Withdrew consent	9	9	2	5																																																																																						
Protocol deviation	2	4	0	4																																																																																						
Lost to follow up	1	4	6	4																																																																																						
Death	1	0	0	0																																																																																						
Other	1	5	2	5																																																																																						
	Grp 1	Grp 2	Grp 3	Grp 4																																																																																						
All cause adverse events																																																																																										
N	216	215	225	220																																																																																						
Constipation	9	2	8	5																																																																																						
Diarrhoea	7	6	5	3																																																																																						
Dizziness	3	12	6	2																																																																																						
Dry mouth	16	15	47	5																																																																																						
Dyspepsia	2	1	3	5																																																																																						
Ejaculation failure	0	4	7	0																																																																																						

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 (Tamsulosin) N: 215 (baseline data/efficacy analysis for N=209) Mean (\pm SD) Age: 61.7 \pm 10.5 (range 40-90) Urge urinary incontinence : 50/215 Urgency episodes/24h: 7.10 \pm 3.83 Micturitions/24h: 12.10 \pm 3.51 Micturitions/night: 1.74 \pm 1.20 IPSS \pm SD: 20.04 \pm 5.02 IPSS QoL \pm SD: 4.57 \pm 0.86 Qmax \pm SD, mL/s: 13.4 \pm 7.6 PVR \pm SD, mL: 56.5 \pm 55.0 Dropouts: 29/215 (13.5%)</p> <p>Group 3 (Tolterodine ER + Tamsulosin) N: 225 (baseline data/efficacy analysis for N=217) Mean (\pm SD) Age: 61.0 \pm 9.6 (range 40-92) Urge urinary incontinence : 52/225 Urgency episodes/24h: 6.72 \pm 3.95 Micturitions/24h: 11.92 \pm 3.35 Micturitions/night: 2.07 \pm 1.32 IPSS \pm SD: 20.10 \pm 5.49 IPSS QoL \pm SD: 4.55 \pm 0.93 Qmax \pm SD, mL/s: 12.7 \pm 6.8 PVR \pm SD, mL: 58.8 \pm 53.8 Dropouts: 34/225 (15.1%)</p> <p>Group 4 (Placebo) N: 222 (baseline data/efficacy analysis for N=215) Mean (\pm SD) Age: 62.8 \pm 9.7 (range 40-88) Urge urinary incontinence : 48/220 Urgency episodes/24h: 7.33 \pm 3.82 Micturitions/24h: 11.86 \pm 3.24</p>		<p>Fatigue 2 3 2 6 Headache 2 9 14 7 Nasal congestion 0 3 10 2 Somnolence 2 5 4 2 Urinary retention 2 0 2 3</p> <p>*See Notes</p> <p>Number of patients reporting treatment benefit at 12 weeks (ITT post hoc figures with imputed data) Pair wise analysis using Fishers 2 sided test</p> <p>PPBC at week 12 (%)</p> <p>Willingness to continue at week 12 (%)</p>	<p>Grp 1: 136/217 Grp 2: 146/215 Grp 3: 172/225 Grp 4: 132/222</p> <p>Grp 1 v Grp 4 p value 0.49 Grp 1 v Grp 2 p value 0.27 Grp 1 v Grp 3 p value 0.002 Grp 2 v Grp 4 p value 0.07 Grp 2 v Grp 3 p value 0.06 Grp 3 v Grp 4 p value <0.001</p> <p>Grp 1 Grp 2 Grp 3 Grp 4</p> <p>Major improvement 32 35 35 27 Minor improvement 32 27 32 30 No change 28 30 27 38 Deterioration 8 8 5 5</p> <p>Grp 1 Grp 2 Grp 3 Grp 4</p> <p>Very willing 44 39 51 38 Little bit willing 21 19 15 21 Little bit unwilling 12 20 12 12 Very unwilling 23 21 23 30</p>	<p>observation carried forward (LOCF)</p> <p>PPBC is a single item global measure questionnaire with sex options to the question of " which of the following statements described your bladder condition best at the moment"?</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Micturitions/night: 2.02 ± 1.19 IPSS \pm SD: 20.00 ± 5.42 IPSS QoL \pm SD: 4.58 ± 0.95 Qmax \pm SD, mL/s: 12.2 ± 6.6 PVR \pm SD, mL: 47.1 ± 47.7 Dropouts: 34/222 (15.3%) 2 patients did not receive study medication</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																			
Kirby et al., ¹⁴⁷ Study design: RCT double blinded(4 arms) Setting: 90 European centres Evidence level: 1+ Duration of follow-up: 1 year(52 weeks)	Patient group: Symptomatic BPH Inclusion criteria: <ul style="list-style-type: none"> ▪ Aged 50 to 80 years ▪ IPSS ≥ 12 ▪ Qmax of ≥5 mL/s but ≤15 mL/s in a total voided volume of ≥150 mL ▪ Enlarged prostate as determined by DRE. Exclusion criteria: <ul style="list-style-type: none"> ▪ Previous prostate surgery or other invasive procedures for treating BPH ▪ Prostate cancer or a PSA level exceeding 10 ng/mL. If PSA was between 4.1 to 10 ng/mL, need to have ≥2 of the following : negative DRE or transrectal ultrasound findings(within the past 3 months) or negative biopsy findings(within the past 4 weeks) ▪ lower urinary tract symptoms or reduced urinary flow rates resulting from a condition other than BPH ▪ large bladder diverticulum, bladder stones, recurrent urinary tract infection, or two or more episodes of AUR requiring catheterization within the year before study entry ▪ residual urine volumes greater 	Group 1: Doxazosin 4 mg(+ placebo) Initiated on 1 mg/day, titrated to 2 mg at end of week 2 and, 4 mg from end of week 6. At the end of week 10, the 4-mg dose was maintained in subjects who met the following two criteria: (a) total IPSS had decreased by 30% or more from baseline, and(b) Qmax had increased by 3 mL/s or more from baseline. For subjects who did not meet these goals, the doxazosin dose was increased to 8 mg/day and maintained for the remaining 42 weeks. Doses were reduced to the next lower dose if the SBP/diastolic BP(DBP) fell to less than 90/60 mm Hg or tolerability was limited. Subjects unable to tolerate a 2-mg/day dose of doxazosin were withdrawn.	IPSS, mean ±SD at 1 year	Group 1: 8.7 ± 5.8 Group 2: 10.9 ± 6.2 Group 3: 8.7 ± 6.2 Group 4: 11.8 ± 6.9	Funding: Grant provided by Pfizer Ltd. Finasteride & placebo provided by Merck & Co Limitations: Randomisation allocation and concealment methods not stated. Additional outcomes: Mean change in sitting and SBP and DBP: Normotensive subjects: Not sig Hypertensive subjects (sitting DBP ≥ 90mmHg, SBP ≥ 140mmHg): LS mean change (sitting SBP/DBP, mmHg) for doxazosin: -11.8/-5.7 Doxazosin + finasteride: -9.2/-5.6 (P<0.05, clinically sig) For Finasteride: -5.7/-																																																			
			IPSS LS mean change ±SEM at 1 year	<u>Compared to baseline value</u> Group 1: -8.3 ± 0.4## Group 2: -6.6 ± 0.4 Group 3: -8.5 ± 0.4## Group 4: -5.7 ± 0.4 ##P<0.0001 compared to placebo, <0.01 compared to finasteride																																																				
			Qmax, ml/s mean ±sd at 1 year	Group 1: 14.0 ± 4.9 Group 2: 12.1 ± 4.7 Group 3: 14.5 ± 5.1 Group 4: 12.1 ± 4.2																																																				
			Qmax, ml/s change from baseline at endpoint, LS mean change ±sem	Group 1: 3.6 ± 0.3 ## Group 2: 1.8 ± 0.3 Group 3: 3.8 ± 0.3 ## Group 4: 1.4 ± 0.3 **P<0.0001 compared to placebo or finasteride																																																				
			Reason for withdrawal Total withdrawals Reasons	<table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> <th>Grp 4</th> </tr> </thead> <tbody> <tr> <td>Total withdrawals</td> <td>78(28.4)</td> <td>81(30.7)</td> <td>89(31.1)</td> <td>76(28.1)</td> </tr> <tr> <td>Adverse Events</td> <td>32(11.6)</td> <td>34(12.9)</td> <td>35(12.2)</td> <td>30(11.1)</td> </tr> <tr> <td> Death**</td> <td>0(0.0)</td> <td>2(0.8)</td> <td>1(0.3)</td> <td>2(0.7)</td> </tr> <tr> <td>Inadequate response</td> <td>3(1.1)</td> <td>6(2.3)</td> <td>3(1.0)</td> <td>9(3.3)</td> </tr> <tr> <td>Noncompliance</td> <td>7(2.5)</td> <td>12(4.2)</td> <td>6(2.1)</td> <td>9(3.3)</td> </tr> <tr> <td>Protocol violation</td> <td>5(1.8)</td> <td>4(1.5)</td> <td>6(2.1)</td> <td>3(1.1)</td> </tr> <tr> <td>Failed screening guidelines</td> <td>3(1.1)</td> <td>2(0.8)</td> <td>1(0.3)</td> <td>1(0.4)</td> </tr> <tr> <td>Other therapy indicated</td> <td>5(1.8)</td> <td>3(1.1)</td> <td>6(2.1)</td> <td>5(1.9)</td> </tr> <tr> <td>Lost to follow-up</td> <td>4(1.5)</td> <td>15(5.7)</td> <td>5(1.7)</td> <td>4(1.5)</td> </tr> <tr> <td>Other</td> <td>19(6.9)</td> <td>15(5.7)</td> <td>26(9.1)</td> <td>13(4.8)</td> </tr> </tbody> </table>			Grp 1	Grp 2	Grp 3	Grp 4	Total withdrawals	78(28.4)	81(30.7)	89(31.1)	76(28.1)	Adverse Events	32(11.6)	34(12.9)	35(12.2)	30(11.1)	Death**	0(0.0)	2(0.8)	1(0.3)	2(0.7)	Inadequate response	3(1.1)	6(2.3)	3(1.0)	9(3.3)	Noncompliance	7(2.5)	12(4.2)	6(2.1)	9(3.3)	Protocol violation	5(1.8)	4(1.5)	6(2.1)	3(1.1)	Failed screening guidelines	3(1.1)	2(0.8)	1(0.3)	1(0.4)	Other therapy indicated	5(1.8)	3(1.1)	6(2.1)	5(1.9)	Lost to follow-up	4(1.5)	15(5.7)	5(1.7)	4(1.5)	Other
	Grp 1	Grp 2	Grp 3	Grp 4																																																				
Total withdrawals	78(28.4)	81(30.7)	89(31.1)	76(28.1)																																																				
Adverse Events	32(11.6)	34(12.9)	35(12.2)	30(11.1)																																																				
Death**	0(0.0)	2(0.8)	1(0.3)	2(0.7)																																																				
Inadequate response	3(1.1)	6(2.3)	3(1.0)	9(3.3)																																																				
Noncompliance	7(2.5)	12(4.2)	6(2.1)	9(3.3)																																																				
Protocol violation	5(1.8)	4(1.5)	6(2.1)	3(1.1)																																																				
Failed screening guidelines	3(1.1)	2(0.8)	1(0.3)	1(0.4)																																																				
Other therapy indicated	5(1.8)	3(1.1)	6(2.1)	5(1.9)																																																				
Lost to follow-up	4(1.5)	15(5.7)	5(1.7)	4(1.5)																																																				
Other	19(6.9)	15(5.7)	26(9.1)	13(4.8)																																																				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																				
	<p>than 200 ml</p> <ul style="list-style-type: none"> ▪ Active urinary tract infection. ▪ Serious diseases ▪ History of drug or alcohol abuse ▪ History of sensitivity to alpha-adrenergic blocking agents, quinazolines, or finasteride. ▪ Hypotension(sitting BP less than 95/60 mm Hg) or orthostatic hypotension(greater than a 20-mm Hg decrease in systolic BP [SBP] when changing from a supine to standing position ▪ Concomitant therapy with anticholinergics, cholinergics, other alpha-blockers, calcium channel blockers, antiandrogens, other 5-alpha-reductase inhibitors, and plant extract preparations was prohibited during the study. <p>All patients N: 1095(79.5%) out of 1378 screened Age, mean ±sd,(yr): 64 IPSS mean ± sd: 17.2 Qmax, ml/s mean±sd: 10.5 Mean PSA, ng/ml, mean= 2.6 Prostate volume, g, mean= 36.3 Drop outs:</p> <p>Group 1(Doxazosin) N: 250 Dropouts: Age, mean ±sd,(yr): 63 ±7</p>	<p>Mean final dose: 6.4mg/day 8mg: 63.2% 4mg: 31.2% 2 mg: 4.8% 1 mg: 0.8%</p> <p>Group 2: Finasteride 5mg(+ placebo)</p> <p>Group 3: Doxazosin 4 mg + finasteride 5 mg Mean final dose: 6.1mg/day 8mg: 57.0% 4mg: 35.5% 2 mg:6.0% 1 mg:1.5%</p> <p>Group 4: placebo for terazosin and placebo for finasteride All subjects advised to take medications at about 8am</p> <p>Concomitant treatment: Diuretic and beta-blocker dosages which were stable for 4 weeks before the initial screening and were maintained during the study.</p>	<p>AUR TURP Either AUR or TURP</p> <p>Dizziness</p> <p>Postural hypotension</p> <p>Hypertension</p> <p>Hypotension</p> <p>Syncope</p> <p>Asthenia</p>	<table border="0"> <tr> <td>Grp 1</td> <td>Grp 2</td> <td>Grp 3</td> <td>Grp4</td> </tr> <tr> <td>N=275</td> <td>N=264</td> <td>N=286</td> <td>N=269</td> </tr> <tr> <td>0(0)</td> <td>3(1.1)</td> <td>0(0)</td> <td>4(1.5)</td> </tr> <tr> <td>1(0.4)</td> <td>3(1.1)</td> <td>0(0)</td> <td>7(2.6)</td> </tr> <tr> <td>1(0.4)</td> <td>5(1.9)</td> <td>0(0)</td> <td>7(2.6)</td> </tr> </table> <p>Group 1: 43/275(15.6%)# Group 2: 21/264(8.0%) Group 3: 39/286(13.6%)# Group 4: 20/269(7.4%) P<0.01 vs. finasteride and placebo</p> <p>Group 1: 16/275(5.8%)# Group 2: 2/264(0.8%) Group 3: 8/286(2.8%) Group 4: 4/269(1.5%) P<0.01 vs. finasteride and placebo</p> <p>Group 1: 5/275(1.8%)# Group 2: 11/264(4.2%) Group 3: 4/286(1.4%)# Group 4: 15/269(5.6%) P=0.02 vs. placebo.</p> <p>Group 1: 14/275(5.1%)# Group 2: 2/264(0.8%) Group 3: 8/286(2.8%) Group 4: 4/269(1.5%) P=0.01 vs. finasteride & placebo</p> <p>Group 1: 2/275(0.7%) Group 2: 0/264(0.0%) Group 3: 6/286(2.1%)# Group 4: 1/269(0.4%) P=0.04 vs. finasteride</p> <p>Group 1: 29/275(10.5%) # Group 2: 11/264(4.2%) Group 3: 26/286(9.1%) # Group 4: 11/269(4.1%) P<0.01 vs. finasteride & placebo</p>	Grp 1	Grp 2	Grp 3	Grp4	N=275	N=264	N=286	N=269	0(0)	3(1.1)	0(0)	4(1.5)	1(0.4)	3(1.1)	0(0)	7(2.6)	1(0.4)	5(1.9)	0(0)	7(2.6)	<p>2.7 Placebo: -4.0/-2.1 Not sig</p> <p>Notes: Analysis of covariance was used for efficacy data, which included effects of treatment, centre(pooled by country), and treatment by centre interaction</p> <p>Last observed carried forward algorithm was used for subjects who discontinued early.</p> <p>*No overall baseline differences were found except for Qmax. †P <0.0001 vs. placebo. ‡P <0.09 vs. finasteride. §Estimated by DRE(in increments of 5 g). ** Excludes one post therapy death, which occurred approximately 35 days after discontinuation of</p>
Grp 1	Grp 2	Grp 3	Grp4																						
N=275	N=264	N=286	N=269																						
0(0)	3(1.1)	0(0)	4(1.5)																						
1(0.4)	3(1.1)	0(0)	7(2.6)																						
1(0.4)	5(1.9)	0(0)	7(2.6)																						

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Dropouts: Duration of BPH at baseline, mean(yr): 1.7 ± 2.9 Prostate Vol by DRE,(g)§: 36 ± 14 IPSS mean ± sd: 17.1 ± 4.2 Qmax(ml/s): 10.4 ± 2.5†‡ PSA serum, mean(ng/ml): 2.5 ± 2.0</p> <p>Group 2(Finasteride) N: 239 Dropouts: Age, mean ±sd,(yr): 63 ±7 Duration of BPH at baseline, mean(yr) = 1.4 ± 2.2 Prostate Vol by DRE,(g)§: 36 ± 14 IPSS mean ± sd: 17.1 ± 4.4 Qmax(ml/s): 10.2 ± 2.5† PSA serum, mean(ng/ml): 2.6 ± 2.1</p> <p>Group 3: Terazosin 10 mg + finasteride 5 mg N: 265 Dropouts: Age, mean ±sd,(yr): 64 ±7 Duration of BPH at baseline, mean(yr) = 1.8 ± 2.9 Prostate Vol by DRE,(g)§: 37 ± 14 IPSS mean ± sd 17.3 ± 4.7 Qmax(ml/s): 10.4 ± 2.7† PSA serum, mean(ng/ml): 2.7 ± 2.3</p> <p>Group 4: placebo for terazosin and placebo for finasteride N: 253 Dropouts: Age Mean(±SD): 64±7 Duration of BPH at baseline, mean(yr) = 1.6 ± 3.0</p>		Somnolence	Group 1: 11/275(4.0%) Group 2: 8/264(3.0%) Group 3: 9/286(3.1%) Group 4: 6/269(2.2%) Not sig	doxazosin therapy
			Vertigo	Group 1: 8/275(2.9%) Group 2: 6/264(2.3%) Group 3: 8/286(2.8%) Group 4: 3/269(1.1%) Not sig	
			Impotence	Group 1: 16/275(5.8%) Group 2: 13/264(4.9%) Group 3: 30/286(10.5%)# ‡ Group 4: 9/269(3.3%) P<0.01 vs. finasteride, finasteride and doxazosin	
			Decreased libido	Group 1: 10/275(3.6%) Group 2: 9/264(3.4%) Group 3: 6/286(2.1%) Group 4: 5/269(1.9%) Not sig	
			Ejaculatory abnormality	Group 1: 1/275(0.4%) Group 2: 6/264(2.3%) Group 3: 7/286(2.4%) Group 4: 4/269(1.5%) Not sig	
			PSA at end point , mean±sd ng/ml	Group 1: 2.8 ± 2.3 Group 2: 1.5 ± 1.0 Group 3: 1.4 ± 1.2 Group 4: 2.9 ± 2.6	
			PSA change from baseline at endpoint , mean ±sd ng/ml	Group 1: 0.3 ± 1.0 Group 2: 1.2 ± 1.4 Group 3: 1.3 ± 1.6 Group 4: 0.3 ± 1.3	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Prostate Vol by DRE,(g)§: 36 ± 15 IPSS mean ± sd: 17.2 ± 4.5 Qmax (ml/s): 10.8 ± 2.5 PSA serum, mean(ng/ml): 2.6 ± 2.1				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Martorana et al., 1997¹⁸⁴</p> <p>Study design: RCT</p> <p>Setting: Multi-centre</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 weeks</p>	<p>Patient group: Men with clinical diagnosis of BPH.</p> <p>Inclusion criteria: Men aged 50-80 years with a clinical diagnosis of BPH confirmed by digital rectal examination and transrectal ultrasound examination showing prostate enlargement,; at least a 6 month history of BPH related symptoms with a 9-item Boyarsky score>6 before entry and after placebo run-in; peak flow rate between 5-12ml/s with a voided volume>150ml.</p> <p>Exclusion criteria: concomitant urological diseases, had undergone prostatectomy or were scheduled to have prostatectomy within 6 months had systolic blood pressure<100,,Hg or history off orthostatic hypotension, had either renal or severe hepatic insufficiency, a psychiatric disorder, insulin dependent diabetes mellitus, history of sever heart disease, myocardial infarction or cerebrovascular accident within 6 months, had hypersensitivity to afluzosin, had treatment with other drugs for BPH during the 2 weeks prior to inclusion, or concomitant treatment with other alpha-blockers, calcium antagonists, monoamine oxidase inhibitors or anticholinergic drugs.</p> <p>All patients N: 94</p> <p>Group 1 N: 47 Evaluable for efficacy analysis: 26</p>	<p>Group 1: alpha-blocker Alfuzosin2.5mg t.i.d.</p> <p>Group 2: Placebo</p>	<p>Mean (±SEM) Qmax, ml/s</p>	<p>Baseline Group 1: 10.55 (0.43) Group 2: 10.4 (0.50)</p> <p>4 weeks Group 1 (n=25): 13.16 (0.80) Group 2 (n=25): 11.75 (0.62) P=NS</p>	<p>Funding: NR</p> <p>Limitations: ITT analysis completed but only the per-protocol analysis reported in the study. This is the patient population that complied with the selection criteria and with the complete urodynamic evaluation at baseline and end point.</p> <p>Additional outcomes: Detrusor opening pressure and maximum detrusor pressure reported. Reported that blood pressure and heart rate measurement found no statistically significant changes.</p> <p>Notes: 2 week placebo run-in phase before trial. After double blind study there was an 8 week single blind treatment extension study.</p>
			<p>Mean (±SEM) flow, ml/s</p>	<p>Baseline Group 1: 5.92 (0.34) Group 2: 6.30 (0.43)</p> <p>4 weeks Group 1 (n=25): 7.80 (0.70) Group 2 (n=24): 6.90 (0.47) P=NS</p>	
			<p>Mean (±SEM) maximum flow rates, ml/s (from pressure/flow study)</p>	<p>Baseline Group 1: 7.76 (0.44) Group 2: 8.52 (0.57)</p> <p>4 weeks Group 1 (n=25): 10.01 (0.91) Group 2 (n=26): 10.26 (0.92) P=NS</p>	
			<p>Mean (±SEM) detrsor pressure at maximum flow, cmH2O (pressure/flow study)</p>	<p>Baseline Group 1: 77.88 (5.61) Group 2: 82.27 (5.91)</p> <p>4 weeks Group 1 (n=25): 54.36 (4.97) Group 2 (n=26): 76.84 (7.78) P<0.05</p>	
			<p>Mean (SEM) Boyarsky score</p>	<p>Baseline Group 1: 10.7 (0.7) Group 2: 10.5 (0.5)</p> <p>4 weeks Group 1 (n=25): 8.0 (0.4) Group 2 (n=26): 8.0 (0.5) P=NS</p>	
			<p>Adverse events</p>	<p>Total Group 1: 4/47 (8.5%) Group 2: 1/47 (2.1%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean (\pmSD) Age: 62.5 (1.0) Dropouts: 21 (10 lack of complete urodynamic evaluation; 6 lack of compliance with selection criteria at baseline; 5 lack of compliance with protocol treatment requirements; 1 lack of correspondence between treatment drug and blood detection; 2 lost to follow up; 1 lack of uroflowmetric evaluation.</p> <p>Group 2 N: 47 Evaluable for efficacy analysis: 26 Mean (\pmSEM) Age: 63.1 (1.1) Dropouts: 21 (9 lack of complete urodynamic evaluation; 8 lack of compliance with selection criteria at baseline; 2 lack of compliance with protocol treatment requirements; 3 lack of correspondence between treatment drug and blood detection, 2 lost to follow up.</p> <p>Note: 5 patients had two reasons and 1 had three reasons of non evaluability.</p>			<p>Hypertension Group 1: 1 (2.1%) Group 2: 1 (2.1%)</p> <p>arthralgia Group 1: 1 (2.1%) Group 2: 0</p> <p>Vertigo Group 1: 1 (2.1%) Group 2: 0</p> <p>Pathological fracture Group 1: 1 (2.1%) Group 2: 0</p>	

See Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?

for McConnell et al., 2003¹⁷⁰.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Mohanty et al., 2003²⁰¹</p> <p>Study design: RCT</p> <p>Setting: India</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 months</p>	<p>Patient group: male patients between 40-80years having lower urinary tract obstructive symptoms suggestive of BPH were recruited.</p> <p>Inclusion criteria: IPSS>10, maximum flow rate 5-13mL/s and average flow rate<6mL/s with post residual urine volume >100mL and PSA<4ng/mL</p> <p>Exclusion criteria: patients with renal or hepatic failure, carcinoma prostate, stricture urethra, neurogenic bladder, bladder neck stenosis, previous surgery on prostate</p> <p>All patients N: 72 Mean age: 61years Drop outs: 3</p> <p>Group 1 N: 38 Mean (±SD) Age: 61.3 (8.5) Dropouts:2</p> <p>Group 2 N: 34 Mean (±SD) Age: 62.7 (13.8) Dropouts:1</p>	<p>Group 1: ALPHA-BLOCKER Tamsulosin 0.4mg daily (sustained capsules)</p> <p>Group 2: PLACEBO Identical capsules once daily</p>	<p>Mean (SD) IPSS</p>	<p>Baseline Group1: 19.53 (3.2) Group 2: 18.52 (5)</p> <p>2 weeks Group1: 12.67 (4.3) Group 2: 15.3 (4.7)</p> <p>4 weeks Group1: 9.8 (4.4) Group 2: 13.8 (4.8)</p> <p>8 weeks Group1 (n=36): 6.9 (4.4) Group 2 (n=33): 12.7 (4.0)</p>	<p>Funding: NR</p> <p>Additional outcomes: Vital signs reported.</p> <p>Notes: Adverse events reported at end point but study included figures for each time interval.</p>
			<p>Mean (SD) Qmax, mL/s</p>	<p>Baseline Group1: 10.5 (2.1) Group 2: 11.6 (2.3)</p> <p>8 weeks Group1 (n=36): 15.7 (4.6) Group 2 (n=33): 12.5 (2.6)</p>	
			<p>Average urinary flow rate, mL/s</p>	<p>Baseline Group1: 4.5 (1.5) Group 2: 5.3 (1.7)</p> <p>8 weeks Group1 (n=36): 7.7 (2.1) Group 2 (n=33): 5.8 (1.7)</p>	
			<p>Maximum voided volume, mL</p>	<p>Baseline Group1: 341.7 (137.6) Group 2: 310.3 (105.4)</p> <p>8 weeks Group1 (n=36): 353.1 (154.3) Group 2 (n=33): 336.9 (149.4)</p>	
			<p>Mean (SD) post voided residual volume, mL</p>	<p>Baseline Group1: 100.6 (46) Group 2: 97.6 (46.4)</p> <p>8 weeks Group1 (n=36): 53.1 (19.2) Group 2 (n=33): 91.8 (40.1)</p>	

			<p>Adverse events at end point</p>	<p>Dizziness Group 1: 9 Group 2: 11 Headache Group 1: 8 Group 2: 9 Fatigue Group 1: 14 Group 2: 14 Postural hypotension Group 1: 2 Group 2: 0 Syncope Group 1: 1 Group 2: 0 Somnolence Group 1: 1 Group 2: 1 Abdominal pain Group 1: 2 Group 2: 1 Dyspnea Group 1: 0 Group 2: 3 Retrograde ejaculation Group 1: 0 Group 2: 0 Constipation Group 1: 7 Group 2: 0 Withdrawn due to adverse events Group 1: 0 Group 2: 0</p>	
--	--	--	---	--	--

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nordling et al., 2005²²⁵</p> <p>Study design: RCT</p> <p>Setting: Multi-centre, Europe and Israel</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: Men were recruited between Feb 1998 and August 1999.</p> <p>Inclusion criteria: men aged ≥ 50 years with a clinical diagnosis of symptomatic BPH and at least a 6 month history of LUTS, with all the following criteria met only a the beginning of the placebo run-in period: an IPSS of ≥ 13, nocturia twice or more, a peak flow rate of 5-12ml/s for a voided volume of 150mL or more, and a residual urine volume of 350mL or less. Patients were not required to these criteria again at the time of randomisation, simulating real-life practice.</p> <p>Exclusion criteria: concomitant urological diseases; diagnosed or suspected carcinoma of the prostate; previous prostate surgery; invasive BPH treatments; previous x-ray therapy of the pelvic region; patients previously showing no improvement with treatment with an alpha-blocker; patients with Parkinson's disease, insulin-dependent diabetes, diagnosed or suspected MS, unstable angina or sever heart failure, history of stroke or myocardial infarction within 5 months of day -28 of day 0, known hypersensitivity to alpha blockers or patients taking concomitant medications that might alter voiding patterns.</p>	<p>Run in period: 28 day single blind, placebo run in period. One placebo tablet matching Alfuzosin 10mg and one matching Tamsulosin 0.4mg at the end of the evening meal.</p> <p>Group 1: Alpha-blocker Alfuzosin 10mg once daily (one tablet plus one placebo tamsulosin capsule)</p> <p>Group 2: Alpha-blocker Alfuzosin 15mg once daily (one tablet plus one placebo tamsulosin capsule)</p> <p>Group 3: Alpha-blocker Tamsulosin 0.4mg once daily (one capsule plus one placebo alfuzosin tablet)</p> <p>Group 4: Placebo One placebo alfuzosin tablet plus one placebo tamsulosin capsule. At the end of the evening meal</p>	<p>Mean (SD) IPSS</p> <p>% of patients with a total IPSS improvement (defined as 3 or more points)</p> <p>Mean (SD) Qmax, mL/s</p> <p>Number (%) adverse events (AE)</p>	<p>Baseline Group 1: 18.0 (5.4) Group 2: 17.4 (5.6) Group 3: 17.4 (6.2) Group 4: 17.7 (5.0)</p> <p>Change from baseline Group 1: -6.5 (5.2); p=0.007 Group 2: -6.0 (5.6); p=0.050 Group 3: -6.5 (6.2); p=0.014 Group 4: -4.6 (5.8)</p> <p>Group 1: 81 Group 2: 69 Group 3: 77 Group 4: 64</p> <p>Baseline Group 1: 9.2 Group 2: 8.9 Group 3: 9.4 Group 4: 9.0</p> <p>Change from baseline Group 1: 1.5 (3.3) ; p=0.22 Group 2: 1.6; (3.8) p=0.09 Group 3: 2.4 (4.3); p=0.02 Group 4: 0.9 (3.0)</p> <p>Treatment emergent (TE) AE \geq one Group 1: 58 (38) Group 2: 61 (39) Group 3: 58 (37) Group 4: 52 (34)</p> <p>TEAE \geq one serious Group 1: 3 (2) Group 2: 7 (4) Group 3: 6 (4) Group 4: 3 (2)</p> <p>Discontinuation because of TEAE Group 1: 4 (3) Group 2: 13 (8)</p>	<p>Funding: NR.</p> <p>Limitations: Method of randomisation and allocation concealment not reported.</p> <p>Additional outcomes: Blood pressure changes were reported. Standard laboratory test results were taken but the study did not report figures but stated no significant changes.</p> <p>Notes: Alfuzosin 10mg improvement of IPSS was apparent at the first assessment at 4 weeks. Not reported for other groups.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients N: 625 Patients in ITT analysis: 611 Dropouts: 47</p> <p>Group 1 N: 154 Mean (\pmSD) Age: 65 (51-85) Dropouts: 9 (adverse events=4; other=5)</p> <p>Group 2 N: 159 Mean (\pmSD) Age: 65 (50-84) Dropouts: 17 (adverse events=14; other=3)</p> <p>Group3 N: 158 Mean (\pmSD) Age: 64 (50-87) Dropouts: 9 (adverse events=6, other=3)</p> <p>Group 4 N: 154 Mean (\pmSD) Age: 64 (50-82) Dropouts:12 (adverse events=5; lack of efficacy=2; other=5)</p>			<p>Group 3: 6 (4) Group 4: 5 (3) Discontinuation because of serious vasodilatory TEAE Group 1: 0 Group 2: 1(1) Group 3: 1 (1) Group 4: 0 Dizziness Group 1: 9 (6) Group 2: 11 (7) Group 3: 3 (2) Group 4: 6 (4) Headache Group 1: 3 (2) Group 2: 4 (3) Group 3: 7 (4) Group 4: 5 (3) Syncope Group 1: 0 Group 2: 2 (1) Group 3: 1 (1) Group 4: 0 Hypotension Group 1: 0 Group 2: 1 (1) Group 3: 1(1) Group 4: 0 Malise Group 1: 0 Group 2: 1 (1) Group 3: 0 Group 4: 0 Impotence Group 1: 2 (1) Group 2: 2 (1) Group 3: 7 (4) Group 4: 0 Ejaculation disorder Group 1: 2 (1) Group 2: 0</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 3: 5 (3) Group 4: 0 Abnormal semen Group 1: 0 Group 2: 0 Group 3: 1 (1) Group 4: 0 Asthenia/ Fatigue Group 1: 4 (3) Group 2: 10 (6) Group 3: 6 (4) Group 4: 3 (2) Somnolence Group 1: 0 Group 2: 1 (1) Group 3: 0 Group 4: 2 (1)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Resnick et al., 2007²⁴⁵</p> <p>Study design: RCT</p> <p>Setting: Multi-centre, US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 29 days</p>	<p>Patient group: Men aged ≥50 years with LUTS suggestive of BPH, including a history of storage and/or voiding symptoms.</p> <p>Inclusion criteria: IPSS of ≥13 points and IPSS bother score of ≥3 points; Qmax between 5 and 12ml/s with a voided volume ≥150ml and post void residual ≤350ml.</p> <p>Exclusion criteria: Conditions that affect urinary functioning, such as Parkinson's disease, MS, poorly controlled diabetes, severe heart failure, stroke recent myocardial infarction or concomitant lower urinary tract disease. Previous prostatic surgery or radiation therapy, an endoscopic procedure within 1 month of screening, spontaneous urinary retention during the preceding 12 months, an ongoing episode of urinary retention requiring an indwelling catheter, postural hypotension, syncope or non-responders to previous alpha blocker therapy. Concomitant use of medications. Evidence of clinically relevant biochemical abnormalities or a PSA>10ng/ml.</p> <p>All patients N: 372</p> <p>Group 1 N: 186 Mean (±SD) Age: 63.5 (8.4) Ethnicity: Black/African: 161 American:</p>	<p>Run-in phase: 28 days patients received one tablet of placebo.</p> <p>Group 1: Alpha-blocker Alfuzosin 10mg One tablet taken once daily after the evening meal, at approximately 0700 h or as late as possible.</p> <p>Group 2: Placebo One tablet taken once daily</p>	<p>Mean improvement in Qmax, ml/s</p>	<p>24 hours Group 1: 1.58 Group 2: 0.71; p<0.021</p> <p>Day 8 Group 1: 1.92 Group 2: 0.39; p<0.001</p> <p>Day 29 Group 1: 1.76 Group 2: 0.36; p<0.001</p>	<p>Funding: Sanofi-Aventis</p> <p>Limitations: Adverse events figures reported differently in text and table.</p> <p>Additional outcomes: BPH impact score reported. Method of randomisation and allocation concealment unclear.</p> <p>Notes: No clinically significant changes in blood pressure were observed (figures not provided). One serious adverse event (non-insulin dependent diabetes mellitus) in intervention group. Considered not to be due to treatment.</p>
			<p>Mean change in IPSS (acute version of IPSS: to allow evaluation of symptom relief after one week)</p>	<p>Day 8 Group 1: -3.4 Group 2: -2.7; p=0.071</p> <p>Day 29 Group 1: -4.5 Group 2: -3.1; p=0.003</p>	
			<p>Mean change in IPSS quality of life score</p>	<p>Day 29 Group 1: -0.7 Group 2: -0.6 P=0.125</p>	
			<p>Treatment emergent adverse events (with > 1% incidence in either group)</p>	<p>Total Group 1: 46/185 (24.9%) Group 2: 43/185 (23.2%)</p> <p>Dizziness Group 1: 11/185 (5.9%) Group 2: 0</p> <p>Headache Group 1: 5/185 (2.7%) Group 2: 2/185 (1.1%)</p> <p>Upper respiratory tract infection Group 1: 4/185 (2.2%) Group 2: 2/185 (1.1%)</p> <p>Orthostatic hypotension Group 1: 3/185 (1.6%) Group 2: 4/185 (2.2%)</p> <p>Fatigue Group 1: 2/185 (1.1%) Group 2: 1/185 (0.5%)</p> <p>Insomnia Group 1: 2/185 (1.1%)</p>	

	<p>White/Caucasian: 10 Other: 14 Dropouts: 10</p> <p>Group 2 N: 186 Mean (\pmSD) Age: 64.4 (8.0) Ethnicity: Black/African: 166 American: White/Caucasian: 6 Other: 13 Dropouts: 7</p>			<p>Group 2: 0 Erectile dysfunction Group 1: 1/185 (0.5%) Group 2: 2/185 (1.1%) Cough Group 1: 0 Group 2: 2/185 (1.1%) Dry mouth Group 1: 0 Group 2: 2/185 (1.1%) Gastroesophageal reflux disease Group 1: 0 Group 2: 2/185 (1.1%) Discontinuation due to adverse events Group 1: 3/185 (24.9%) Group 2: 1/185</p>	
--	--	--	--	--	--

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Roehrborn et al., 2001a²⁵⁴</p> <p>Study design: RCT</p> <p>Setting: Multi-centre, US and Canada.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Men with LUTS/BPH recruited between Jan 1998-Aug 1999.</p> <p>Inclusion criteria: men aged 50 years or older with a history of lower urinary tract symptoms consistent with clinical BPH for 6 months or longer, an IPSS of at least 13, a Qmax between 5-12mL/s with a voided volume of 150mL or more, a residual urine volume of 350mL or less, and a quality of life of at least 3 points. Patients had to meet inclusion criteria on day 1 of placebo run-in period (4 weeks) and did not need to re-qualify on randomisation.</p> <p>Exclusion criteria: Concomitant lower urinary tract disease; previous prostate surgery; history of postural hypotension or syncope; concomitant use of medications that may alter the voiding pattern; and clinically relevant biochemical abnormalities. Serum PSA >10ng/mL were excluded and those with an elevated serum PSA 4-10 had to have prostate cancer excluded to the satisfaction for the investigator.</p> <p>All patients N: 536 Mean age: 63.6 (49-92) Drop outs: 72 (13%)</p> <p>Group 1 N: 177</p>	<p>Group 1: Alpha-blocker Alfuzosin 10mg once daily without initial dose titration.</p> <p>Group 2: Alpha-blocker Alfuzosin 15mg once daily without initial dose titration.</p> <p>Group 3: Placebo</p>	<p>Mean (SD) IPSS</p> <p>[Note: * adjusted p-value compared to placebo]</p> <p>% of patients showing an improvement in IPSS of 3 or more points</p> <p>Mean (SD) quality of life</p> <p>% of patients showing an improvement in IPSS quality of life question of 2 or more points</p> <p>Mean (SD) Qmax, mL</p>	<p>Baseline Group 1: 18.2 (6.3) Group 2: 17.7 (5.7) Group 3: 18.2 (6.4)</p> <p>Change Group 1 (n=170): -3.6 (4.8); p=0.001* Group 2 (n=165): -3.4 (5.7); p=0.004 Group 3 (n=167): -1.6 (5.8)</p> <p>Group 1: 56% Group 2: 52% Group 3: 39%</p> <p>Baseline Group 1: 3.8 (1.1) Group 2: 3.7 (1.1) Group 3: 3.7 (1.1)</p> <p>Change Group 1 (n=170): -0.7 (1.1); p=0.002 Group 2 (n=165): -0.7 (1.2); p=0.002 Group 3 (n=167): -0.3 (1.1)</p> <p>Group 1: 21%; p=0.004 Group 2: 21%; p=0.003 Group 3: 12%</p> <p>Baseline Group 1: 9.9 (3.9) Group 2: 10.0 (3.2) Group 3: 10.2 (4.0)</p> <p>Mean change Group 1 (n=170): 1.7 (4.2); p=0.0004 Group 2 (n=165): 0.9 (3.6); p=0.12 Group 3 (n=167): 0.2 (3.5)</p> <p>Optimal mean change Group 1 (n=170): 1.7; p=0.0004 Group 2 (n=165): 1.2; p=0.03 Group 3 (n=167): 0.3</p>	<p>Funding: Sanofi-Synthelabo</p> <p>Limitations: Method of randomisation or allocation concealment unclear. Prostate volume in alfuzosin 10mg significantly larger than other 2 groups.</p> <p>Additional outcomes: IPSS voiding and filling sub-scores were reported. Reported that there were no significant changes in the hematologic or biochemical measurement were observed. Blood pressure changes reported (reported that no patient experienced clinically relevant changes).</p> <p>Notes: Significant improvement in IPSS for treatment groups by first post treatment assessment (day 28) and maintained throughout study.</p> <p>Qmax was not normally</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean [range] Age: 64.3 (50-92)</p> <p>Prostate volume: 40.2</p> <p>Dropouts: 11% (adverse events=8;</p> <p>Group 2</p> <p>N: 181</p> <p>Mean [range] Age: 63.9 (50-81)</p> <p>Prostate volume: 38.3</p> <p>Dropouts: 18% (adverse events=8; insufficient efficacy=2</p> <p>Group 3</p> <p>N: 178</p> <p>Mean [range] Age: 62.7 (49-85)</p> <p>Prostate volume: 36.8</p> <p>Dropouts: 11% (adverse events=4; insufficient efficacy=2</p>			<p>Median change</p> <p>Group 1 (n=170): 1.1 (4.2); p=0.0006</p> <p>Group 2 (n=165): 1.0 (3.6); p=0.0006</p> <p>Group 3 (n=167):</p> <p>Median optimal change</p> <p>Group 1 (n=170): 1.3</p> <p>Group 2 (n=165): 1.1</p> <p>Group 3 (n=167): 0.3</p>	<p>distributed so median values were also reported.</p> <p>Men over 65 years who received alfuzosin 15mg reported more adverse events potentially related to vasodilation (dizziness, malaise, hypotension) than younger patients (17% v 5%). This was not observed in the 10mg group.</p>
			<p>% of patients showing an improvement in Qmax of 2mL/s or more</p> <p>Group 1: 40%</p> <p>Group 2: 41%</p> <p>Group 3: 26%</p>		
			<p>Number (%) treatment emergent adverse events (≥2%) of the exposed population</p> <p>Total</p> <p>Group 1: 52%</p> <p>Group 2: 43%</p> <p>Group 3: 43%</p> <p>Dizziness</p> <p>Group 1: 13 (7.4)</p> <p>Group 2: 16 (9.0)</p> <p>Group 3: 5 (2.9)</p> <p>Headache</p> <p>Group 1: 9 (5.1)</p> <p>Group 2: 4 (2.3)</p> <p>Group 3: 4 (2.3)</p> <p>Respiratory tract infection</p> <p>Group 1: 6 (3.4)</p> <p>Group 2: 5 (2.8)</p> <p>Group 3: 4 (2.3)</p> <p>Back pain</p> <p>Group 1: 2 (1.1)</p> <p>Group 2: 6 (3.4)</p> <p>Group 3: 4 (2.3)</p> <p>Rhinitis</p> <p>Group 1: 3 (1.7)</p> <p>Group 2: 4 (2.3)</p> <p>Group 3: 4 (2.3)</p> <p>Fatigue</p> <p>Group 1: 4 (2.3)</p> <p>Group 2: 3 (1.7)</p> <p>Group 3: 4 (2.3)</p>		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>Inflicted injury Group 1: 4 (2.3) Group 2: 3 (1.7) Group 3: 1 (0.6)</p> <p>Impotence Group 1: 5 (2.8) Group 2: 2 (1.1) Group 3: 2 (1.1)</p> <p>Somnolence Group 1: 4 (2.3) Group 2: 3 (1.7) Group 3: 0</p> <p>Sinusitis Group 1: 5 (2.8) Group 2: 1 (0.6) Group 3: 4 (2.3)</p> <p>Constipation Group 1: 4 (2.3) Group 2: 1 (0.6) Group 3: 1 (0.6)</p> <p>Pain Group 1: 5 (2.8) Group 2: 0 Group 3: 1 (1.1)</p> <p>Nausea Group 1: 4 (2.3) Group 2: 1 (0.6) Group 3: 1 (0.6)</p> <p>Abdominal pain Group 1: 2 (1.1) Group 2: 2 (1.1) Group 3: 4 (2.3)</p> <p>Arthralgia Group 1: 2 (1.1) Group 2: 1 (0.6) Group 3: 4 (2.3)</p> <p>Dyspepsia Group 1: 3 (1.7) Group 2: 0 Group 3: 4 (2.3)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Orthostatic hypotension (decrease in systolic BP of 20mmHg or more when standing) Group 1: 3.4% Group 2: 2.3% Group 3: 3.4%	

See Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?

for Roehborn et al., 2006²⁵⁵

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Schulman et al., 1994²⁶⁹</p> <p>Study design: Randomised cross over trial</p> <p>Setting: Multi-centre</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 weeks</p>	<p>Patient group: men with clinical symptoms of BPH</p> <p>Inclusion criteria: urinary peak flow of <12.5ml/sec; prostate volume >20ml.</p> <p>Exclusion criteria: men suffering from urogenital diseases other than BPH or from neurological diseases that might influence the parameters measured during the trial were excluded.</p> <p>All patients N: 161 Mean age: 31-79 Drop outs: 19 (lost to follow-up=6; intercurrent disease=2; patient withdrawal=2; adverse event=8; lack of efficacy=1)</p> <p>Group 1 (alfuzosin-placebo) N: 79 Mean Age: 63.5</p> <p>Group 2 (placebo-alfuzosin) N: 82 Mean Age: 61.9</p>	<p>Group 1: Alpha-blockers Alfuzosin 2.5mg three times daily</p> <p>Group 2: Placebo Three times daily</p>	<p>Peak flow, ml/sec</p>	<p>Baseline Group 1: 9.06 (2.9) Group 2: 9.14 (2.8) 4 weeks Group 1(n=68): 13.95 (6.3) Group 2(n=73): 11.69 (5.5)</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation and allocation concealment unclear. No washout period between cross over of treatments.</p> <p>Additional outcomes: Results after the cross over period. Adverse events – not reported as unclear whether in phase 1 before cross over of treatments.</p> <p>Notes: After 4 weeks of treatment each group then had 4 more weeks on the opposite treatment. There was no wash out period and the effect of the initial treatment could not be distinguished from any new effects. Therefore, only the first 4 weeks of this trial are reported to limit bias.</p>
			<p>Mean flow, ml/sec</p>	<p>Baseline Group 1: 4.72 (1.9) Group 2: 5.00 (1.9) 4 weeks Group 1(n=68): 6.85 (3.4) Group 2(n=73): 6.01 (2.5)</p>	
			<p>Post voiding volume, ml</p>	<p>Baseline Group 1: 90.65 (82.2) Group 2: 83.86 (67.4) 4 weeks Group 1 (n=61): 50.88 (47.76) Group 2 (n=68): 71.13 (77.0)</p>	
			<p>Boyarsky symptoms score</p>	<p>Baseline Group 1: 12.33 (2.55) Group 2: 12.42 (2.36) 4 weeks Group 1 (n=61): 50.88 (47.76) Group 2 (n=69): 7.65 (3.58)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>VanKerrebroeck et al., 2000³⁰⁵</p> <p>Study design: RCT</p> <p>Setting: 48 Urology centres, Europe</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Men over 50 years with micturition disorders related to BPH from April 1997 to July 1998.</p> <p>Inclusion criteria: IPSS \geq13 and a maximum urinary flow rate between 5 and 12ml/s for a voided volume of at least 150ml and a residual urine volume of \leq350ml.</p> <p>Exclusion criteria: concomitant urinary tract disease, previous prostatic surgery or other invasive procedures for the treatment of BPH, associated severe visceral disease, history of postural hypotension or syncope, clinically relevant biological abnormalities, alpha blockers in the month preceding the selection, androgen, antiandrogens, 5 alpha reductase inhibitors and LHRH analogues in the 3 months preceding the selection.</p> <p>All patients N: 447 Drop outs: 40 (8.9%)</p> <p>Group 1 N: 143 Mean (\pmSD) Age: 64.9 (7.4) Dropouts: 16</p> <p>Group 2 N: 150 Mean (\pmSD) Age: 64.7 (7.5) Dropouts: 14</p> <p>Group 3</p>	<p>Run-in period: One month, placebo controlled period'</p> <p>Group 1: Alpha-blockers Alfuzosin 10mg once daily at the end of the evening meal</p> <p>Group 2: Alpha-blockers Alfuzosin 7.5mg (2.5mg thrice daily)</p> <p>Group 3: Placebo</p>	<p>Mean (SD) IPSS</p> <p>Mean (SD) IPSS quality of life question</p> <p>Mean (SD) Qmax</p> <p>Adverse events</p>	<p>Baseline Group 1: 17.3 (3.5) Group 2: 16.8 (3.7) Group 3: 17.7 (4.1)</p> <p>3 months Group 1: 10.4 (4.7) Group 2: 10.5 (6.1) Group 3: 12.8 (6.7)</p> <p>Baseline Group 1: 3.3 (0.9) Group 2: 3.3 (1.0) Group 3: 3.3 (1.0)</p> <p>3 months Group 1: 2.2 (1.1) Group 2: 2.2 (1.1) Group 3: 2.6 (1.3)</p> <p>Baseline Group 1: 9.4 (1.9) Group 2: 8.7 (1.9) Group 3: 9.2 (2.0)</p> <p>3 months Group 1: 11.7 (3.9) Group 2: 11.9 (4.3) Group 3: 10.6 (3.3)</p> <p>Vasodilatory events Group 1: 9/143 (6.3%) Group 2: 14/149 (9.4%) Group 3: 4/154 (2.6%)</p> <p>Drop outs due to Vasodilatory events (syncope) Group 1: 0 Group 2: 1/149 (0.7%) Group 3: 0</p> <p>Dizziness Group 1: 3/143 (2.1%) Group 2: 7/149 (4.7%) Group 3: 2/154 (1.3%)</p> <p>Headache Group 1: 2/143 (1.4%)</p>	<p>Funding: NR</p> <p>Limitations: Qmax was significantly lower in alfuzosin 2.5mg group at baseline. Method of randomisation and allocation concealment unclear.</p> <p>Additional outcomes: IPSS sub-scores for filling and voiding symptoms. Changes in haemodynamic parameters in normotensive and hypertensive patients (no significant differences reported).</p> <p>Notes: NCGC calculated means for Group 1 and 2 for the meta-analysis.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 154 Mean (\pmSD) Age: 64.2 (7.8) Dropouts: 10</p>			<p>Group 2: 3/149 (2%) Group 3: 1/154 (0.6%) Hypotension/postural hypotension Group 1: 1/143 (0.7%) Group 2: 2/149 (1.3%) Group 3: 0/154 Malaise Group 1: 2/143 (1.4%) Group 2: 1/149 (0.7%) Group 3: 0/154 Asthenia/fatigue Group 1: 5/143 (3.5%) Group 2: 1/149 (0.7%) Group 3: 4/154 (2.6%) Sexual dysfunction Group 1: 0 Group 2: 1/149 (0.7%) Group 3: 2/154 (1.3%) Acute urinary retention Group 1: 0 Group 2: 0 Group 3: 1/154</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Wilt et al., 2000a³²⁵</p> <p>Study design: Systematic Review – Cochrane. This comparison includes 10 randomised controlled trials.</p> <p>Setting: Europe, Canada and US.</p> <p>Evidence level: 1++</p> <p>Duration of follow-up: Range 4-52 weeks</p>	<p>Patient group: Men with symptomatic benign prostatic hyperplasia.</p> <p>Inclusion criteria: treatment duration of at least 4 weeks.</p> <p>Exclusion criteria: NR.</p> <p>All patients N: 5151 Mean age: 65 (45-94) Racial characteristics (reported in 6 trials): White: 82%, Asian: 10%, Black 6%, Other : 2% Discontinuation: 26% (5-42%) Mean symptoms score (7 trials)= 18.8 Drop outs: 23 (lost to follow-up, reported as erroneously randomised or unaccounted for and not included in outcome analysis)</p> <p>Group 1 N: 2438</p> <p>Group 2 N: 1821</p> <p>Group 3 N: 990</p>	<p>Group 1: Alpha-blocker Terazosin (hytrin) – non-uroselective alpha-blocker</p> <p>Group 2: Placebo</p> <p>Group 3: Active controls Includes phytotherapy, pharmacological or surgical therapies</p>	<p>AUA symptoms score (0-35) * extrapolated from graphs</p>	<p>Group1 (n=275): 10.1 (6.35) Group 2 (n=265): 13.2 (6.3) Mean difference: -3.10 [-4.17, -2.03]; 1 study P<0.00001</p>	<p>Funding: Minneapolis/VISN-13 Centre for Chronic Diseases Outcomes Research (CCDOR), USA. Department of Veterans Affairs Health Services Research and Development Program, USA.</p> <p>Limitations: Only 3 of 10 studies described their method of allocation concealment (unclear in remaining 7)</p> <p>Additional outcomes: Boyarsky symptom score was reported.</p> <p>Notes: Baseline values for symptoms scores, peak urine flow did not differ by treatment group. * NCGC used fixed effect meta-analysis model rather than</p>
			<p>Mean change in AUA symptom score (fixed dose studies, 10mg only)</p>	<p>Group1 (n=976): -7.6 (7.17) Group 2 (n=973): -3.7 (7.16) Mean difference: -3.90 [-4.54, -3.26]; 1 study P<0.00001</p>	
			<p>Mean change in peak flow rate (10mg), mL/s</p>	<p>Flexible dose studies: MD: 1.40 [0.56, 2.24]; n=424; 2 studies Fixed dose: 10mg MD: 1.53 [0.35, 2.70]; n=148; 2 studies Total: MD: 1.44 [0.76, 2.13]; 4 studies; p<0.0001</p>	
			<p>Mean change in Peak flow rate (5mg), mL/s</p>	<p>Flexible dose studies: MD: 1.40 [0.56, 2.24]; n=424; 2 studies Fixed dose: 5mg MD: 0.46 [-0.76, 1.69]; n=153; 2 studies Total: MD: 1.10 [0.41, 1.79]; 4 studies; p=0.002</p>	
			<p>Mean peak flow rate (up to 10mg), mL/s</p>	<p>Dose escalation/Flexible dose studies: MD: 1.75 [1.09, 2.41]; n=424; 2 studies Fixed dose: MD: 0.90 [-1.06, 2.86]; n=153; 1 study Total: MD: 1.66 [1.03, 2.29]; 3 studies; p<0.00001</p>	
			<p>Discontinuations, all causes*</p>	<p>Dose escalation/flexible-dose studies RR: 0.86 [0.78, 0.95]; 4 studies Fixed doses: all doses RR: 0.93 [0.55, 1.55]; 3 studies Total: Group 1: 521/1904 (27.4%) Group 2: 555/1621 (34.2%) RR: 0.87 [0.79, 0.95]; p=0.003; 7 studies</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Discontinuations, due to adverse events	Dose escalation/flexible-dose studies RR: 1.51 [1.24, 1.85]; 4 studies Fixed doses: all doses RR: 1.77 [0.58, 5.40]; 2 studies Total: Group 1: 229/1817 (12.6%) Group 2: 140/1607 (8.7%) RR: 1.52 [1.25, 1.86]; p<0.00001	random effect used by Cochrane. Fixed model used as there was no heterogeneity present. Cochrane model detected no significant difference between the interventions.
			Dizziness	Group 1: 252/1802 (14.0%) Group 2: 98/1586 (6.2%) RR: 2.40 [1.92, 3.00]; 6 studies; p=<0.00001	
			Asthenia	Group 1: 153/1736 (8.8%) Group 2: 62/1566 (4.0%) RR: 2.42 [1.78, 3.28]; 5 studies; p=<0.00001	
			Headache	Group 1: 40/749 (5.3%) Group 2: 25/555 (4.5%) RR: 1.24 [0.76, 2.01]; 5 studies; p=0.39	
			Postural hypotension	Group 1: 57/1655 (3.4%) Group 2: 8/1487 (%) RR: 5.52 [2.71, 11.24]; 4 studies; p=<0.00001	
			Impotence/erectile dysfunction	Group 1: 24/386 (6.2 %) Group 2: 15/384 (3.9%) RR: 1.59 [0.85, 2.99]; 2 studies; p=0.15	
			Flu syndrome	RR: 1.22 [0.49, 3.06]; 3 studies; p=0.67	
			Abnormal ejaculation	RR: 1.50 [0.05, 40.91]; 2 studies; p=0.81	
			Rhinitis	RR: 1.34 [0.77, 2.31]; 2 studies; p=0.30	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Wilt et al., 2002³²⁴</p> <p>Study design: Systematic Review – Cochrane. 14 RCTs identified; 6 included in this comparison.</p> <p>Setting: Europe, Japan and US.</p> <p>Evidence level: 1++</p> <p>Duration of follow-up: Range 4-26 weeks.</p>	<p>Patient group: Men with symptomatic benign prostatic hyperplasia.</p> <p>Inclusion criteria: treatment duration at least 30 days.</p> <p>Exclusion criteria: NR.</p> <p>All patients N: 3418</p> <p>Mean age: 64 (45 to 85)</p> <p>Drop outs: 395 (lost to follow-up, reported as erroneously randomised or unaccounted for and not included in outcome analysis)</p> <p>Mean IPSS/AUA: 19.5 (6 studies)</p> <p>Mean discontinuation rate: 12%</p> <p>Racial characteristics from one study: White > 99%</p> <p>Group 1 N: 2486</p> <p>Group 2 N: 781</p> <p>Group 3 N: 851</p>	<p>Group 1: Alpha-blockers Tamsulosin</p> <p>Group 2: Placebo</p> <p>Group 3: Active control Medical, phytotherapeutic or surgical therapies.</p>	IPSS/AUA final score by dose	<p>Tamsulosin 0.4mg: MD: -2.55[-3.46, -1.63]; p<0.00001; 2 studies</p> <p>Tamsulosin 0.8mg: MD: -3.42 [-4.32, -2.52]; p<0.00001; 2 studies</p>	<p>Funding: internal sources: Minneapolis/VISN-23 centre for chronic Disease Outcomes Research, USA. Dept of Veterans Affairs Health Service research and Development Program, USA.</p> <p>Limitations: Allocation concealment unclear in all of the studies.</p> <p>Additional outcomes: Boyarsky scores. Mean urine flow. Comparisons by dose for adverse events.</p> <p>Notes: Converted pooled analysis to fixed model rather than random effect model reported in Cochrane review – expect when there was heterogeneity.</p>
			Mean change in IPSS/AUA	<p>Tamsulosin 0.4mg: MD: -2.14[-3.42, -0.87]; p=0.001; 2 studies</p> <p>Tamsulosin 0.8mg: MD: -3.15 [-5.01, -1.28]; p=0.0009; 2 studies</p>	
			Qmax	<p>Tamsulosin 0.4mg: MD: 0.91 [0.51, 1.32]; p<0.00001; 5 studies</p> <p>Tamsulosin 0.8mg: MD: 0.96 [0.50, 1.43]; p<0.00001; 2 studies</p>	
			Mean change in Qmax	<p>Tamsulosin 0.4mg: MD: 1.02 [0.68, 1.35]; p<0.00001; 4 studies</p> <p>Tamsulosin 0.8mg: MD: 1.07 [0.65, 1.48]; p<0.00001; 2 studies</p>	
			Discontinuation due to adverse events	RR: 1.08 [0.73, 1.62]; p=0.69; 3 studies	
			Discontinuation – all men	RR: 1.02 [0.80, 1.31]; p=0.85; 3 studies	
			Serious adverse events	RR: 1.18 [0.57, 2.43]; p=0.65; 3 studies	
			Adverse events – cardiovascular	RR: 0.78 [0.40, 1.53]; p=0.47; 1 study	
			Adverse events – digestive system	RR: 0.86 [0.65, 1.12]; p=0.27; 2 studies	
			Adverse events – nervous system	RR: 1.55 [1.24, 1.95]; p=0.0002; 3 studies	
			Adverse events – urogenital system	RR: 2.67 [0.89, 7.96]; p=0.08; 3 studies	
			Adverse events - drug related	RR: 1.07 [0.71, 1.62]; p=0.75; 2 studies	
Dizziness	Group 1: 176/1473 (11.9%) Group 2: 56/714 (7.8%)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				RR: 1.53 [1.15, 2.02]; p=0.003; 4 studies	
			Headache	Group 1: 211/1473 (14.3%) Group 2: 104/714 (14.6%) RR: 1.00 [0.81, 1.24]; p=1.00; 4 studies	
			Abnormal ejaculation	Group 1: 148/1375 (10.8%) Group 2: 3/686 (0.4%) RR: 21.13 [7.33, 60.87]; p<0.00001; 3 studies	
			Rhinitis	Group 1: 154/1375 (11.2%) Group 2: 41/686 (6.0%) RR: 1.86 [1.34, 2.57]; p=0.0002; 3 studies	
			Asthenia	Group 1: 89/1473 (6.0%) Group 2: 31/714 (4.3%) RR: 1.38 [0.93, 2.04]; p=0.11; 4 studies	
			AUA bother score	Tamsulosin 0.4mg: MD: -1.60 [-2.44, -0.76]; 0.00018; 1 study Tamsulosin 0.8mg: MD: -2.00 [-2.83, -1.17]; p<0.00001; 1 study	

Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Debruyne et al, 1998 ⁶⁹ ALFIN study Study design: RCT double blinded(3 arms) Setting: European, multicenter (104 centres). Conducted from Sept 1994 to Dec1996 Evidence level: 1+ Duration of follow-up: 6 months	Patient group: Lower urinary tract symptoms related to BPH Inclusion criteria: <ul style="list-style-type: none"> ▪ Men 50-75 years ▪ IPSS≥7 ▪ Qmax of ≥5 mL/s but ≤15 mL/s in a total voided volume of >150 mL (no threshold for prostate size was specified, patients with hypertension included) Exclusion criteria: <ul style="list-style-type: none"> ▪ Other concomitant urinary tract disease (prostate cancer, neurogenic bladder dysfunction, bladder stones, chronic bacterial prostatitis, untreated urinary tract infection) ▪ Previous invasive procedure to treat BPH ▪ Associated severe visceral disease ▪ Postural hypotension ▪ Any concomitant medication affecting voiding pattern ▪ Clinically relevant biological abnormalities (aspartate aminotransferase and alanine aminotransferase > 2 times the upper limit of normal, blood creatinine ≥160 micromol/l) ▪ Serum PSA>20ng/ml All patients N: 1051 Dropouts: 133(13%) Age, mean ±sd,(yr): 63.3±6.5 IPSS mean ± sd: 15.4±5.5	All patients received placebo during a 2-week, single blinded run in period Group 1: Alfuzosin SR 5mg twice daily Group 2: finasteride 5mg once daily Group 3: Alfuzosin SR 5mg twice daily + finasteride 5 mg once daily <u>Duration:</u> 6 months	IPSS change, at 6 months (mean ±SD)	Group 1: -6.3±5.8 Group 2: -5.2±5.7 Group 3: -6.1±5.6 P values: Group 1 vs. 2: 0.01 Group 2 vs. 3: 0.03 Group 1 vs. 3: NR	Funding: Synthelabo Recherche, France Limitations: <ul style="list-style-type: none"> ▪ Method of randomisation allocation and concealment was not reported ▪ No report of placebos being used to mask the different number of pills and treatment regimens Additional outcomes: Supine blood pressure (systolic and diastolic), change compared to baseline. There were no sig. difference between groups Notes: None.
			IPSS improved by >50% at 6 months (% of patients)	Group 1: 43 Group 2: 33 Group 3: 42 P values: Group 1 vs. 2: 0.008 Group 2 vs. 3: 0.009 Group 1 vs. 3: NR	
			Qmax change, at 6 months (mean ±SD), ml/s	Group 1: 1.8±3.8 Group 2: 1.8±4.5 Group 3: 2.3±4.7 P values: Not sig	
			Qmax increase >30% compared to baseline, % (Subgroup analysis in 497/1051 men who had Qmax <10ml/s at baseline (most likely to be obstructed))	Group 1: 51 Group 2: 38 Group 3: 49 P values: Group 1 vs. 2: 0.02 Group 2 vs. 3: 0.06 Group 1 vs. 3: NR	
		Prostate volume change, at 6 months (mean ±SD), ml	Group 1: -0.2±14.3 Group 2: -4.3±15.0 Group 3: -4.9±12.4 P values: Group 1 vs. 2: <0.001 Group 2 vs. 3: Not sig Group 1 vs. 3: <0.001		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments															
	<p>Duration of symptoms, mean ± sd, (yr): 3.4±3.2 Prostate vol, mean ± SD (ml): 41.2±24.0 PSA serum, mean ± sd:(ng/ml): 4.0 ± 2.08 Qmax mean±sd (ml/sec): 9.9±3.0</p> <p>Group 1 (Alfuzosin SR) N: 358 Dropouts: 40(11%) Age, mean ±sd,(yr): 63.2±6.4 IPSS, mean ± sd: 15.3±5.5 Duration since first LUTS, mean ± sd, (yr): 3.5±3.0 Prostate vol, mean ± SD (ml):41.4±25.7 PSA serum, mean ± sd:(ng/ml): 3.0±2.5 Qmax mean±sd (ml/sec): 9.7±2.8</p> <p>Group 2 (Finasteride) N: 344 Dropouts: 39(11%) Age, mean ±sd,(yr): 63.0±6.4 IPSS, mean ± sd: 15.5±5.2 Duration since first LUTS, mean ± sd, (yr): 3.3±3.2 Prostate vol, mean ± SD (ml): 40.9±23.5 PSA serum, mean ± sd:(ng/ml): 3.4±2.5 Qmax mean±sd (ml/sec): 9.8±2.6</p> <p>Group 3: Alfuxosin SR + finasteride N: 349 Dropouts: 54(15%) Age, mean ±sd,(yr): 63.7±6.7 IPSS, mean ± sd: 15.6±5.7 Duration since first LUTS, mean ± sd, (yr): 3.4±3.3 Prostate vol, mean ± SD (ml):41.1±22.6 PSA serum, mean ± sd:(ng/ml): 3.1±2.7 Qmax mean±sd (ml/sec): 10.1±3.5</p>		<p>PSA change, at 6 months (mean ±SD), ng/ml</p>	<p>Group 1: 0.1±2.7 Group 2: -1.7±1.9 Group 3: -1.4±1.7</p> <p>P values: Group 1 vs. 2: <0.001 Group 2 vs. 3: Not sig Group 1 vs. 3: <0.001</p>																
			<p>Withdrawals</p> <p>Withdrawal due to adverse events</p> <p>Lack of efficacy</p>	<table border="1"> <thead> <tr> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> </tr> </thead> <tbody> <tr> <td>N=358</td> <td>N= 344</td> <td>N=349</td> </tr> <tr> <td>40</td> <td>54</td> <td>39</td> </tr> <tr> <td>25</td> <td>24</td> <td>18</td> </tr> <tr> <td>3</td> <td>2</td> <td>2</td> </tr> </tbody> </table>	Grp 1	Grp 2	Grp 3	N=358	N= 344	N=349	40	54	39	25	24	18	3	2	2	
Grp 1	Grp 2	Grp 3																		
N=358	N= 344	N=349																		
40	54	39																		
25	24	18																		
3	2	2																		
			<p>Adverse events:</p> <p>Vasodilatory events (%)</p> <p>Vertigo/dizziness</p> <p>Headache</p> <p>Postural hypotension/ hypotension</p> <p>Malaise</p>	<table border="1"> <tbody> <tr> <td>6(1.7)</td> <td>4(1.2)</td> <td>8(2.3)</td> </tr> <tr> <td>7(2.0)</td> <td>4(1.2)</td> <td>5(1.4)</td> </tr> <tr> <td>2(0.6)</td> <td>3(0.9)</td> <td>2(0.6)</td> </tr> <tr> <td>1(0.3)</td> <td>1(0.3)</td> <td>1(0.3)</td> </tr> </tbody> </table>	6(1.7)	4(1.2)	8(2.3)	7(2.0)	4(1.2)	5(1.4)	2(0.6)	3(0.9)	2(0.6)	1(0.3)	1(0.3)	1(0.3)				
6(1.7)	4(1.2)	8(2.3)																		
7(2.0)	4(1.2)	5(1.4)																		
2(0.6)	3(0.9)	2(0.6)																		
1(0.3)	1(0.3)	1(0.3)																		
			<p>Sexual disorders (%)</p> <p>Impotence</p> <p>Ejaculatory failure</p> <p>Decreased libido</p>	<table border="1"> <tbody> <tr> <td>8(2.2)</td> <td>23(6.7)</td> <td>26(7.4) #</td> </tr> <tr> <td>- (-)</td> <td>5(1.5)</td> <td>3(0.9)</td> </tr> <tr> <td>2(0.6)</td> <td>6(1.7)</td> <td>7(2.0)</td> </tr> </tbody> </table>	8(2.2)	23(6.7)	26(7.4) #	- (-)	5(1.5)	3(0.9)	2(0.6)	6(1.7)	7(2.0)							
8(2.2)	23(6.7)	26(7.4) #																		
- (-)	5(1.5)	3(0.9)																		
2(0.6)	6(1.7)	7(2.0)																		
			<p>Others (%)</p> <p>Somnolence</p> <p>Asthenia/fatigue</p> <p>Myocardial infarction</p> <p>Acute urine retention</p>	<table border="1"> <tbody> <tr> <td>-(-)</td> <td>2(0.6)</td> <td>1(0.3)</td> </tr> <tr> <td>4(1.1)</td> <td>-(-)</td> <td>2(0.6)</td> </tr> <tr> <td>-(-)</td> <td>1(0.3)</td> <td>1(0.3)</td> </tr> <tr> <td>2(0.6)</td> <td>1(0.3)</td> <td>1(0.3)</td> </tr> </tbody> </table> <p># p>0.002</p>	-(-)	2(0.6)	1(0.3)	4(1.1)	-(-)	2(0.6)	-(-)	1(0.3)	1(0.3)	2(0.6)	1(0.3)	1(0.3)				
-(-)	2(0.6)	1(0.3)																		
4(1.1)	-(-)	2(0.6)																		
-(-)	1(0.3)	1(0.3)																		
2(0.6)	1(0.3)	1(0.3)																		
			<p>Asymptomatic orthostatic hypotension during at least one visit</p>	<table border="1"> <thead> <tr> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> </tr> </thead> <tbody> <tr> <td>(9)/358</td> <td>(8)/344</td> <td>(8)/349</td> </tr> <tr> <td>(13)/112</td> <td>(13)/109</td> <td>(12)/115</td> </tr> <tr> <td>(10)/165</td> <td>(10)/147</td> <td>(10)/169</td> </tr> </tbody> </table>	Grp 1	Grp 2	Grp 3	(9)/358	(8)/344	(8)/349	(13)/112	(13)/109	(12)/115	(10)/165	(10)/147	(10)/169				
Grp 1	Grp 2	Grp 3																		
(9)/358	(8)/344	(8)/349																		
(13)/112	(13)/109	(12)/115																		
(10)/165	(10)/147	(10)/169																		
			<p>All</p> <p>Hypertensive</p> <p>≥65 years</p>																	

Study details	Patients	Interventions	Outcome measures	Effect size			Comments
			Study withdrawals Withdrawals Adverse events Lost to follow up Lack of efficacy Other reasons	Grp 1 N=358 40(11%)	Grp 2 N= 344 39(11%)	Grp 3 N=349 54(15%) 24 6 2 22	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Lepor et al., 1996¹⁶³</p> <p>Also reported in Lepor 1998¹⁶⁴ and Lepor 2000¹⁶²</p> <p>Study design: RCT double blinded (4 arms)</p> <p>Setting: US, outpatient clinics, multicentre (Dec 1992 to March 1995)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Symptomatic BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ▪ Age 45 to 80 years old ▪ Mean AUA symptom score ≥8 ▪ Mean Qmax ≥4ml/s, ≤15 ml/s, with a minimal voided volume 125ml and a mean residual volume after voiding <300ml <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ Taken the following drugs within the specified time periods: experimental drug < 4 weeks before screening; alpha adrenergic agonist, cholinergic agonist or antagonist, topical beta adrenergic antagonist drug for glaucoma, or any hypertensive drug other than a diuretic or angiotensin converting enzyme inhibitor within 2 weeks before lead in period; estrogens, androgens or androgen inhibitors within 3 months. ▪ Unstable angina, myocardial infarction, transient ischaemic attack, stroke within past 6 months, insulin dependent diabetes mellitus, orthostatic hypotension 	<p>Group 1: Terazosin 10 mg (+ placebo) (Titrated from 1 mg from days 1 to 3, 2 mg from days 4 to 7, 5 mg from days 8 to 14 and 10 mg from day 15 to end of study. Patients allowed to reduce to 5 mg in the event of adverse events observed)</p> <p>Group 2: Finasteride 5mg (+ placebo) Single daily dose at bedtime</p> <p>Group 3: Terazosin 10 mg + finasteride 5 mg</p> <p>Group 4: placebo for terazosin and placebo for finasteride</p>	<p>IPSS/AUASS mean ±SD at 1 year (SD calculated from SEM presented in Lepor1998^{164*}</p> <p>IPSS/AUASS mean change (95% CI) at 1 year * [calculated by NCGC team from baseline and 1 year follow up values]</p> <p>Difference in IPSS/AUA mean change (95% CI) at 1 year, between groups [calculated by NCGC team]</p> <p>Qmax, ml/s mean ±SD at 1 year (SD calculated from SEM presented in Lepor1998^{164*}</p> <p>Qmax, ml/s mean change (95% CI) at 1 year compared to baseline* [calculated by NCGCAC team from baseline and 1 year follow up values]</p> <p>Difference in Qmax mean change (95% CI) at 1 year, between groups* [calculated by NCGC team]</p> <p>Discontinuation due to adverse events</p>	<p>Group 1: 10.2 ± 4.97, n=275 Group 2: 13.0 ±4.84, n=260 Group 3: 9.80 ±5.00, n=278 Group 4: 13.2±4.88, n=265</p> <p><u>Compared to baseline value</u> Group 1: -6.00 [-6.85, -5.15] Group 2: -3.20 [-4.04, -2.36] Group 3: -6.10 [-3.97, -5.23] Group 4: -2.60 [-3.45, -1.75]</p> <p>MD Gp1-2: -2.80 [-3.99, -1.61]** MD Gp1-3: 0.10 [-1.31, 1.11] MD Gp1-4:-3.40 [-4.60, -2.20]** MD Gp2-3: 2.90 [1.70, 4.10]** MD Gp2-4:-0.60 [-1.79, 0.59] MD Gp3-4: -3.50 [-4.71, -2.29]** **p value:<0.001</p> <p>Group 1: 13.2±4.97, n=275 Group 2: 12.1±4.76, n=252 Group 3: 13.6±1.66, n=277 Group 4: 11.8±4.87, n=264</p> <p><u>Compared to baseline value</u> Group 1: 2.70[2.04, 3.36] Group 2: 1.50[0.85, 2.15] Group 3: 3.20[2.54, 3.86] Group 4: 1.40[0.74, 2.06]</p> <p>MD Gp1-2: 1.20 [0.28, 2.12]** MD Gp1-3: -0.50 [-1.43, 0.43] MD Gp1-4: 1.30 [0.37, 2.23]** MD Gp2-3: -1.70 [-2.62, -0.78]** MD Gp2-4: 0.10 [-0.82, 1.02] MD Gp3-4: 1.80 [0.87, 2.73]** **p value:<0.001</p> <p>Group 1: 18/305 (5.9%) Group 2: 15/310 (4.8%) Group 3: 24/309 (7.8%) Group 4: 5/305 (1.6%) P<0.05</p>	<p>Funding: Veterans Affairs Medical Research Service, Merck and Abbott</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Values for Qmax and AUA/IPSS had to be extrapolated from graphs, no actual values reported. <p>Additional outcomes: AUA symptoms scores started to be significantly different between arms containing terazosin vs. finasteride only or placebo at week 2, reached nadir at week 13 and maintained until week 52. There were no significant differences between terazosin only vs. terazosin + finasteride arm through out study period.</p> <p>The Qmax outcomes had a similar trend, expect that statistical significance between terazosin containing arms vs. finasteride only</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																													
	<ul style="list-style-type: none"> ▪ Previous BPH, obstruction or pelvic surgery ▪ Prostate carcinoma ▪ Urinary tract infections ▪ Renal or hepatic impairment ▪ <p>All patients N: 1229 (73%) out of 1686 screened Age Mean (±SD): Drop outs:</p> <p>Group 1 (Terazosin) N: 305 Age Mean (±SD): 65±6 Dropouts:49/305 Prostate volume (cm³): 37.5±1.1 White race (%): 81 AUASS: 16.2±5.5 Qmax (ml/s):10.5±2.6 PSA serum (ng/ml): 2.2±1.9</p> <p>Group 2 (Finasteride) N: 310 Age Mean (±SD): 65±7 Dropouts:67 Prostate volume (cm³): 36.2±1.0 White race (%): 79 AUASS:16.2±5.4 Qmax (ml/s):10.6±2.5 PSA serum (ng/ml): 2.2±1.8</p> <p>Group 3: Terazosin 10 mg + finasteride 5 mg N: 309 Age Mean (±SD): 65±7 Dropouts:55</p>		<p>Discontinuation – all men</p> <p>Reason for withdrawal *</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> <th>Grp 4</th> </tr> </thead> <tbody> <tr> <td>Total withdrawals</td> <td>49</td> <td>67</td> <td>55</td> <td>51</td> </tr> <tr> <td>Reasons</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td> Adverse Events</td> <td>18</td> <td>15</td> <td>24</td> <td>5</td> </tr> <tr> <td> Absolute indication for surgery</td> <td>2</td> <td>5</td> <td>2</td> <td>4</td> </tr> <tr> <td> Unrelated medical problem</td> <td>4</td> <td>10</td> <td>8</td> <td>10</td> </tr> <tr> <td> Death</td> <td>2</td> <td>7</td> <td>2</td> <td>3</td> </tr> <tr> <td> Lost to follow up</td> <td>9</td> <td>9</td> <td>5</td> <td>3</td> </tr> <tr> <td> Other</td> <td>14</td> <td>21</td> <td>14</td> <td>26</td> </tr> </tbody> </table>		Grp 1	Grp 2	Grp 3	Grp 4	Total withdrawals	49	67	55	51	Reasons					Adverse Events	18	15	24	5	Absolute indication for surgery	2	5	2	4	Unrelated medical problem	4	10	8	10	Death	2	7	2	3	Lost to follow up	9	9	5	3	Other	14	21	14	26	<p>Group 1: 49/305 (16%) Group 2: 67/310 (22%) Group 3: 55/309 (18%) Group 4: 51/305 (17%)</p>	<p>and placebo arms started at week 4. (based on graph, no actual values reported)</p> <p>Notes: Slight differences in values of differences between baseline and 1 year values between Lepor1996 and Lepor1998. Postural hypotension and other adverse events values reported in Lepor1996 was slightly different from 1998</p> <p>† P values for overall difference among all 4 groups</p> <p>* Values for Qmax and AUASS was obtained from Lepor1998¹⁶⁴. There are some discrepancies in differences between baseline and 1 year follow up. Values in Lepor 1998 were used.</p>
	Grp 1	Grp 2	Grp 3	Grp 4																																														
Total withdrawals	49	67	55	51																																														
Reasons																																																		
Adverse Events	18	15	24	5																																														
Absolute indication for surgery	2	5	2	4																																														
Unrelated medical problem	4	10	8	10																																														
Death	2	7	2	3																																														
Lost to follow up	9	9	5	3																																														
Other	14	21	14	26																																														
			Dizziness	<p>Group 1: 79/305 (26%) Group 2: 26/310 (8%) Group 3:66/309 (21%) Group 4: 22/305 (7%) P<0.001†</p>																																														
			Postural hypotension (determined by principal investigator, involving light headedness when standing and not measurable change in blood pressure)	<p>Group 1: 23/305 (8%) Group 2: 7/310 (2%) Group 3: 27/309 (9%) Group 4: 3/305 (1%) P<0.001†, Gp 1 +- 2: P=0.004</p>																																														
			Orthostatic hypotension , at least once during study (A fall of more than 20 mmHg in the systolic blood pressure when patient changed from supine to upright position)	<p>Group 1: 45% Group 2: 26% Group 3: 39% Group 4: 30% (Information was provided in replies and correction section NEJM1997; 336:293)</p>																																														
			Syncope	<p>Group 1: 3/305 (1%) Group 2: 3/310 (1%) Group 3: 5/309 (2.3%) Group 4: 0/305 (0%) Not sig</p>																																														
			Asthenia	<p>Group 1: 42/305 (14%) Group 2: 23/310 (7%) Group 3: 43/309 (14%)</p>																																														

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Prostate volume (cm³): 37.2±1.1 White race (%): 80 AUASS: 15.9±5.7 Qmax (ml/s): 10.4±2.7 PSA serum (ng/ml): 2.3±2.0 Group 4: placebo for terazosin and placebo for finasteride N: 305 Age Mean (±SD): 65±7 Dropouts: 51 Prostate volume (cm³): 38.4±1.3 White race (%): 79 AUASS: 15.8±5.5 Qmax (ml/s): 10.4±2.6 PSA serum (ng/ml): 2.4±2.1			Group 4: 21/305 (7%) P<0.002†, Gp 1 +- 2: P= 0.01	
			Headache	Group 1: 18/305 (6%) Group 2: 19/310 (6%) Group 3: 16/309 (5%) Group 4: 10/305 (3%) Not sig	
			Decreased libido	Group 1: 8/305 (3%) Group 2: 14/310 (5%) Group 3: 15/309 (5%) Group 4: 4/305 (1%) P=0.05†, Grp 1 vs. 2: Not sig	
			Ejaculatory abnormality	Group 1: 1/305 (0.3%) Group 2: 6/310 (2%) Group 3: 21/309 (7%) Group 4: 4 /305 (1%) P<0.001†, Grp 1 vs. 2: Not sig	
			Rhinitis	Group 1: 20/305 (7%) Group 2: 8/310 (3%) Group 3: 24/309 (8%) Group 4: 14/305 (5%) P=0.02† Grp 1 vs. 2: Not sig	
			Sinusitis	Group 1: 6/305 (2%) Group 2: 4/310 (1%) Group 3: 7/309 (2%) Group 4: 4/305 (1%) Grp 1 vs. 2: 0.02	
			BPH impact index (BII) mean ±SD at 1 year (SD calculated from SEM presented in Lepor1998¹⁶⁴*	Group 1: 2.4±1.66 n=276 Group 2: 3.0±1.61 n=259 Group 3: 2.0±1.67 n=279 Group 4: 3.0±1.63 n=265	
			BPH impact index (BII) mean change (95% CI) at 1 year * [calculated by NCGC team from baseline and 1 year follow up values]	<u>Compared to baseline value</u> Group 1: -1.2±2.4 Group 2: -0.5±2.4 Group 3: -1.7±2.4 Group 4: -0.5±2.4	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			BPH impact index (BII) mean change \pm SD(95% CI) at 1 year, between groups [calculated by NCGC team]	MD Gp1-2: -0.7 ± 3.4 (-1.0,-0.4)** MD Gp1-3: 0.5 ± 3.4 (0.2,0.8)** MD Gp1-4: -0.5 ± 3.4 (-1.0,-0.4)** MD Gp2-3: 1.2 ± 3.4 (0.9,1.5)** MD Gp2-4: 0.0 ± 3.4 (-0.3,0.3) MD Gp3-4: -1.2 ± 3.0 (-1.5,-0.9)** **P<0.001	
			Prostate volume, ml, \pmSD at 1 year (SD calculated from SEM presented in Lepor1998 ^{164*}	Group 1: 38.0 ± 21.5 n=271 Group 2: 30.1 ± 20.8 , n=252 Group 3: 30.2 ± 21.7 , n=275 Group 4: 38.9 ± 25.2 , n=258	
			Prostate volume, ml, mean change (95% CI) at 1 year * [calculated by NCGC team from baseline and 1 year follow up values]	<u>Compared to baseline value</u> Group 1: 0.5 ± 21.57 Group 2: -6.1 ± 20.80 Group 3: -7.0 ± 21.72 Group 4: 0.5 ± 25.20	
			Difference in prostate volume mean change (95% CI) at 1 year, between groups [calculated by NCGC team]	<u>Change in AUA between groups, at 1 year</u> MD Gp1-2: 6.6 (3.0, 10.2) ** MD Gp3-1: -7.5 (-11.1,-3.9) ** MD Gp1-4: 0 (-4.0, 4.0) MD Gp3-2: -0.9 (-4.5, 2.7)** MD Gp2-4: -6.6 (-10.6, -2.6) ** MD Gp3-4: -7.5 (-11.5,-3.5) ** **p value:<0.001	

See Evidence Table 9: Alpha-blockers vs. placebo

for Kirby et al., 2003¹⁴⁷

See Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?

for McConnell et al., 2003¹⁷⁰.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments				
Rigatti et al, 2003 ²⁵² MICTUS study Study design: RCT double blinded Setting: Italian, multicenter (50 centres) Evidence level: 1+ Duration of follow-up: 52 weeks	Patient group: Lower urinary tract symptoms related to benign prostatic hyperplasia Inclusion criteria: <ul style="list-style-type: none"> ▪ men between 50 and 80 y with symptomatic LUTS/BPH ▪ I-PSS ≥ 13 ▪ Qmax between 4 and 15 ml/s Total Symptom Problem Index (SPI) score ≥ 7. ▪ Post-void residual volume (PVR: evaluated by ultrasonography) < 400 ml ▪ PSA level < 3 or 3–10 ng/ml (provided that prostate cancer was ruled out by the investigator according to the usual procedure in the centre). Exclusion criteria: <ul style="list-style-type: none"> ▪ Known history or a diagnosis of urological disturbances, cardiovascular diseases, neurological diseases, hepatic or renal insufficiency ▪ Clinically significant abnormalities in haematological and biochemical tests ▪ Took an alpha-1-adrenoreceptor antagonist (A-1-ARA) or phytotherapy in the 6 weeks prior to the study or finasteride in the 6 months prior to the study. ▪ Required concomitant medications influencing pharmacodynamic or pharmacokinetic properties of tamsulosin, in particular A-1-ARA, mixed alpha- beta-antagonists, alpha-agonists and anticholinergics. 	During the 2-week, single-blind, placebo run-in period, patients took one capsule of tamsulosin-matching placebo and one tablet of finasteride-matching placebo once daily. Group 1: Tamsulosin One capsule of tamsulosin 0.4 mg + one tablet of finasteride-matching placebo once daily Group 2: Finasteride One tablet of finasteride 5 mg + one capsule of tamsulosin-matching placebo once daily. Patients were assessed at visit 1 (screening visit) and 2 weeks later (randomisation/base line visit) during the placebo run-in period. Treatment period:	IPSS change from baseline at 26 weeks (mean \pm SD)	Group 1: -6.3 ± 5.5 (-32.0%) Group 2: -5.7 ± 5.7 (-37.3%) P value: 0.080	Funding: Boehringer Ingelheim Italy SpA Limitations: Method of randomisation allocation and concealment was not reported Notes: None.				
			IPSS improved by $\geq 50\%$ at 26 weeks compared to baseline (% of patients)	Group 1: 42.5% Group 2: 35.6% P value: Not sig					
			I-PSS-QoL change from baseline at 26 weeks , (mean \pm sd)	Group 1: -1.1 ± 1.2 (-31.2%) Group 2: -1.0 ± 1.2 (-25.8%) P value: 0.163					
			Qmax change from baseline at 26 weeks , (mean \pm sd) ,ml/s	Group 1: 2.4 ± 5.9 (30.7%) Group 2: 1.9 ± 5.1 (21.7%) P value: 0.271					
			Voided volume, change from baseline at 26 weeks , (mean \pm sd), ml	Group 1: 21.3 ± 152.4 (29.9%) Group 2: 5.2 ± 141.0 (16.4%) P value: 0.043					
			Number of patients treated	<table border="0"> <tr> <td>Grp 1</td> <td>Grp 2</td> </tr> <tr> <td>N=196</td> <td>N= 204</td> </tr> </table>		Grp 1	Grp 2	N=196	N= 204
			Grp 1	Grp 2					
			N=196	N= 204					
			Any AE	63 (32.1)					
			Serious AE	15 (7.6)					
Discontinued due to AE	19 (9.7)								
Adverse events reported in more than 3% patients)									
Influenza-like symptoms	12 (6.1)								
Impotence	6 (3.1)								
Abdominal pain	6 (3.1)								
Ejaculation disorder	6 (3.1)								
Study withdrawals	<table border="0"> <tr> <td>Grp 1</td> <td>Grp 2</td> </tr> <tr> <td>N=199</td> <td>N= 204</td> </tr> </table>	Grp 1	Grp 2	N=199	N= 204				
Grp 1	Grp 2								
N=199	N= 204								
Adverse events	19(9.7%)								
Lost to follow up	13(6.6)								
Lack of efficacy	4(2.0%)								
Non compliance to protocol	4(2.0%)								
Withdrawal of consent	1(0.5%)								
Other reasons	16(8.2%)								
	7(3.6%)								

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients N: 403 randomised from 441 enrolled Dropouts: see study withdrawals Age, mean \pmsd,(yr): 63\pm7.1 Prostate vol ,mean \pm SD (ml): 39\pm18.9</p> <p>Group 1(Tamsulosin) N: 199 Dropouts: 34(17%) at week 26, 63 (31%) at week 52 IPSS, mean \pm sd: 16.3\pm5.1 IPSS-QoL, mean \pm sd: 3.2 (1.0) *Prostate vol < 50 ml): 68% Qmax mean\pmsd (ml/sec):10.8\pm3.7 Voided volume, mean\pmsd, ml 239.5 (118.4)</p> <p>Group 2(Finasteride) N: 204 Dropouts: 24(11.8%) at 26 weeks, 45 (22%) at 52 weeks IPSS, mean \pm sd: 16.9\pm5.0 IPSS-QoL, mean \pm sd: 3.1 (1.1) *Prostate vol < 50 ml): 75% Qmax mean\pmsd (ml/sec): 10.8\pm3.4 Voided volume, mean\pmsd,ml:226.5 \pm93.1</p> <p>* Not statistically significant, calculated by NCGC team using Fisher's exact test</p>	26 weeks + 26 weeks	Symptom Problem Index (SPI) ITT population	<u>Baseline</u> Group 1: 13.6 \pm 4.4, n=193 Group 2: 14.0 \pm 4.2, n=202 <u>Change at week-26</u> Group 1: -5.2 \pm 5.0 (-37.4%), n=193 Group 2: -4.5 \pm 5.0 (-31.5%), n=202 P value: 0.055	
			Symptom Problem Index (SPI): Per protocol population	<u>Baseline</u> Group 1: 13.6 \pm 4.4, n=130 Group 2: 14.1 \pm 4.2, n=152 <u>Change at week-26</u> Group 1: -5.5 \pm 5.0 (-39.6%) Group 2: -4.5 \pm 4.9 (-31.5%) P value: 0.032	
			% Symptom Problem Index (SPI) responders (50% improvement from baseline)	<u>% Patients at week-26</u> Group 1: 43.5%, n=193 Group 2: 35.1%, n=202	
			Symptom Problem Index (SPI) -storage	<u>Baseline</u> Group 1: 6.1 \pm 2.4 Group 2: 6.2 \pm 2.2 <u>Change at week-26</u> Group 1: -2.3 \pm 2.5 (-34.3%), n=193 Group 2: -1.9 \pm 2.7 (-22.0%), n=202 P value: 0.09	
			Symptom Problem Index (SPI) -voiding	<u>Baseline</u> Group 1: 7.5 \pm 3.0, n=193 Group 2: 7.8 \pm 2.7, n=202 <u>Change at week-26</u> Group 1: -3.0 \pm 3.2(-35.0%) Group 2: -2.6 \pm 3.1(-27.3%) P value: 0.069	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Roehrborn et al 2008 ²⁶³ for the 2 year results</p> <p>Study design: RCT double blinded(3 arms)</p> <p>Setting: International, multicenter (446 investigators in 35 countries)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: This is the results from the 2-year interim results</p> <p>Total: 208 weeks treatment + 16 weeks additional safety follow up(224 total)</p>	<p>Patient group: Clinical diagnosis of BPH, prostate size ≥ 30cc</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Men 50 years or older Clinical diagnosis of BPH by medical history and physical examination, including digital rectal examination IPSS ≥ 12 Qmax of ≥ 5 mL/s but ≤ 15 mL/s in a total voided volume of ≥ 125 mL Prostate volume ≥ 30 cc on TRUS Total serum PSA ≥ 1.5 ng/ml <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Total serum PSA > 10.0 ng/ml A history or evidence of prostate cancer Previous surgery to treat BPH History of AUR within 3 months before study entry. Postvoid volume > 250 mL (suprapubic ultrasound) Use of phytotherapy for BPH within 2 weeks of screening visit or /and predicted need for phytotherapy Use of any alpha adrenoceptor blockers within 2 weeks of screening visit and/or predicted need to any alpha blocker other than tamsulosin during study History of postural hypotension, dizziness, vertigo or any other signs and symptoms or orthostasis, which in the opinion of the investigators, could be exacerbated by tamsulosin and putting the subject at risk 	<p>All patients received placebo run in the 4 weeks run in period.</p> <p>Group 1: Tamsulosin 0.4mg (+ placebo dutasteride)</p> <p>Group 2: dutasteride 0.5mg(+ placebo tamsulosin)</p> <p>Group 3: Tamsulosin 0.4 mg + dutasteride 0.5 mg</p> <p><u>Duration:</u> 4 years (208 weeks) All administered once daily</p>	<p>IPSS, at 24 months (mean \pmSD) SE</p> <p>IPSS, change from baseline at 24 months (mean \pmSD) SE</p> <p>IPSS, adjusted** mean difference between groups at 24 months</p> <p>IPSS-QoL, change from baseline at 24 months (mean \pmSD) SE</p> <p>Patients who improved by more than 3 points on the IPSS at 24 months compared to baseline (%)</p> <p>Qmax, ml/s adjusted** mean change from baseline \pmsd at 24 months</p> <p>Prostate volume change from baseline at 24 months, mean %</p>	<p>Group 1: 11.9\pm6.8, SE 0.17 Group 2: 11.4\pm6.4, SE 0.16 Group 3: 10.1\pm6.4, SE 0.16</p> <p><u>Compared to baseline value</u> Group 1: -4.3 \pm6.0, SE 0.15 Group 2: - 4.9\pm6.0, SE 0.15 Group 3: - 6.2\pm6.0, SE 0.15 P value: < 0.001 for Grp 3 vs Grp1 and Grp 2, $P=0.0113$ for Grp 1 vs Grp 2</p> <p>Group 3 vs Group 1: -1.8 Group 3 vs Group 2: -1.3</p> <p><u>Compared to baseline value</u> Group 1: -1.1 Group 2: -1.1 Group 3: -1.4 P value: < 0.001 for Grp 3 vs Grp1 and Grp 2</p> <p>Group 1: 62 Group 2: 65 Group 3: 72 P value: < 0.001 for Grp 3 vs Grp1 and Grp 2</p> <p>Group 1: 0.9 \pm 4.8, SE 0.12 Group 2: 1.9 \pm 4.8, SE 0.12 Group 3: 2.4 \pm 4.8, SE 0.12 P value: ≤ 0.003 for Grp 3 vs Grp 1 and Grp 2, $P<0.001$ for Grp 1 vs Grp 2</p> <p>Group 1: 0.0% \pm 33.4 SE 0.84% Group 2: -28.0% \pm 24.3 SE 0.61% Group 3: -26.9% \pm 24.6 SE 0.62% P value < 0.001 for Grp 3 vs</p>	<p>Funding: GSK</p> <p>Limitations:</p> <ul style="list-style-type: none"> Only interim results available. Final 4-year results will be published at a later date (Autumn2009) <p>Additional outcomes: % of responders defined as</p> <ul style="list-style-type: none"> 25% or greater, 2points of more improvement in IPSS 30% or greater improvement in Qmax <p>Qmax improved significantly greater from baseline for combination vs. monotherapies from month-6.</p> <p>IPSS score improvement from baseline of combination vs. dutasteride was significant from month 3, vs. tamsulosin was significant from month</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																												
	<p>All patients N: 4,844 Dropouts: Age, mean \pmsd,(yr): 66.1 \pm 7.01 No. white ethnicity (%): 4,259 (88) IPSS mean \pm sd: 16.4 \pm 6.16 Duration since first LUTS mean\pmsd, (yr): 5.4 \pm 4.84 Prostate vol (cc): Mean \pm SD total: 55.0 \pm 23.58 Median total: 48.9 Mean \pm SD transition zone* 29.5 \pm 21.97 PSA serum, mean \pm sd:(ng/ml): 4.0 \pm 2.08 Qmax mean\pmsd (ml/sec): 10.7 \pm 3.62 Post-void residual vol, mean\pmsd, (ml): 67.7 \pm 64.87 No. sexually active (%): 3,529 (73) No. previous α-blocker use (%): 2,444 (50) No. previous 5-ARI use (%): 531 (11)</p> <p>Group 1(Tamsulosin) N: 1,611 Dropouts: Age, mean \pmsd,(yr): 66.2 \pm 7.00 No. white ethnicity (%): 1,405 (87) IPSS, mean \pm sd: 16.4 \pm 6.10 Duration since first LUTS mean \pm sd, (yr): 5.4 \pm 4.76 Prostate vol (cc): Mean \pm SD total: 55.8 \pm 24.18 Median total: 49.6</p>			<p>Grp 1</p> <p>PSA change from baseline at 24 months , mean % Group 1: +12.1% Group 2: -55.0% Group 3: -56.0%</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> </tr> </thead> <tbody> <tr> <td>Any Serious Drug related \dagger Leading to study withdrawal</td> <td>N=1611 N=1610 1039(64) 1015(63) 195(12) 209(13)</td> <td>N= 1623 1047(65) 193(12)</td> <td></td> </tr> <tr> <td>Drug related, leading to study withdrawal</td> <td>292(18) 258(16) 130(8) 145(9) 49(3) 48(3)</td> <td>386(24) 161(10) 81(5)</td> <td></td> </tr> </tbody> </table> <p>P value: P<0.001 for combination vs single treatments for any drug related event \dagger</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> </tr> </thead> <tbody> <tr> <td>Adverse events occurring in >1% patients</td> <td>N=1611</td> <td>N= 1623</td> <td>N=1610</td> </tr> <tr> <td>Erectile dysfunction</td> <td>61(3.8)</td> <td>97(6.0)</td> <td>119(7.4)</td> </tr> <tr> <td>Retrograde ejaculation</td> <td>18(1.1)</td> <td>10(0.6)</td> <td>68(4.2)</td> </tr> <tr> <td>Ejaculation failure</td> <td>13(0.8)</td> <td>8(0.5)</td> <td>39(2.4)</td> </tr> <tr> <td>Loss of libido</td> <td>14(0.9)</td> <td>21(1.3)</td> <td>27(1.7)</td> </tr> <tr> <td>Semen volume decreased</td> <td>13(0.8)</td> <td>5(0.3)</td> <td>29(1.8)</td> </tr> <tr> <td>Altered (decreased) libido</td> <td>27(1.7)</td> <td>45(2.8)</td> <td>55(3.4)</td> </tr> <tr> <td>Dizziness</td> <td>27(1.7)</td> <td>11(0.7)</td> <td>26(1.6)</td> </tr> <tr> <td>Breast enlargement</td> <td>13(0.8)</td> <td>29(1.8)</td> <td>23(1.4)</td> </tr> <tr> <td>Nipple pain</td> <td>5(0.3)</td> <td>10(0.6)</td> <td>19(1.2)</td> </tr> <tr> <td>Breast tenderness</td> <td>5(0.3)</td> <td>16(1.0)</td> <td>16(1.0)</td> </tr> </tbody> </table>		Grp 1	Grp 2	Grp 3	Any Serious Drug related \dagger Leading to study withdrawal	N=1611 N=1610 1039(64) 1015(63) 195(12) 209(13)	N= 1623 1047(65) 193(12)		Drug related, leading to study withdrawal	292(18) 258(16) 130(8) 145(9) 49(3) 48(3)	386(24) 161(10) 81(5)			Grp 1	Grp 2	Grp 3	Adverse events occurring in >1% patients	N=1611	N= 1623	N=1610	Erectile dysfunction	61(3.8)	97(6.0)	119(7.4)	Retrograde ejaculation	18(1.1)	10(0.6)	68(4.2)	Ejaculation failure	13(0.8)	8(0.5)	39(2.4)	Loss of libido	14(0.9)	21(1.3)	27(1.7)	Semen volume decreased	13(0.8)	5(0.3)	29(1.8)	Altered (decreased) libido	27(1.7)	45(2.8)	55(3.4)	Dizziness	27(1.7)	11(0.7)	26(1.6)	Breast enlargement	13(0.8)	29(1.8)	23(1.4)	Nipple pain	5(0.3)	10(0.6)	19(1.2)	Breast tenderness	5(0.3)	16(1.0)	16(1.0)	<p>9.</p> <p>IPSS-QOL improvement was significant from months 3 and 12 respectively.</p> <p>Notes: “investigator blinding to the treatment was maintained by an independent, unblinded reviewer who doubled the PSA values in subjects receiving dutasteride or combination therapy with the value randomly stated as the doubled value, or 0.1 units higher or lower.</p> <p>Methods published in Siami et al ²⁷⁹ The study recruitment was completed in 2005.</p> <p>The standard deviation values in the results were calculated by the NCCAC team from the SE values reported.</p>
	Grp 1	Grp 2	Grp 3																																																														
Any Serious Drug related \dagger Leading to study withdrawal	N=1611 N=1610 1039(64) 1015(63) 195(12) 209(13)	N= 1623 1047(65) 193(12)																																																															
Drug related, leading to study withdrawal	292(18) 258(16) 130(8) 145(9) 49(3) 48(3)	386(24) 161(10) 81(5)																																																															
	Grp 1	Grp 2	Grp 3																																																														
Adverse events occurring in >1% patients	N=1611	N= 1623	N=1610																																																														
Erectile dysfunction	61(3.8)	97(6.0)	119(7.4)																																																														
Retrograde ejaculation	18(1.1)	10(0.6)	68(4.2)																																																														
Ejaculation failure	13(0.8)	8(0.5)	39(2.4)																																																														
Loss of libido	14(0.9)	21(1.3)	27(1.7)																																																														
Semen volume decreased	13(0.8)	5(0.3)	29(1.8)																																																														
Altered (decreased) libido	27(1.7)	45(2.8)	55(3.4)																																																														
Dizziness	27(1.7)	11(0.7)	26(1.6)																																																														
Breast enlargement	13(0.8)	29(1.8)	23(1.4)																																																														
Nipple pain	5(0.3)	10(0.6)	19(1.2)																																																														
Breast tenderness	5(0.3)	16(1.0)	16(1.0)																																																														

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean \pm SD transition zone*: 30.5 \pm 24.47 PSA serum, mean \pm sd:(ng/ml): 4.0 \pm 2.08 Qmax mean \pm sd (ml/sec): 10.7 \pm 3.66 Post-void residual vol, mean \pm sd, (ml): 67.7 \pm 65.14 No. sexually active (%): 1,164 (72) No. previous α-blocker use (%): 819 (51) No. previous 5-ARI use (%): 172 (11)</p> <p>Group 2(Finasteride) N: 1,623 Dropouts: Age, mean \pmsd,(yr): 66.0 \pm 6.99 No. white ethnicity (%): 1,433 (88) IPSS, mean \pm sd: 16.4 \pm 6.03 Duration since first LUTS mean \pm sd, (yr): 5.3 \pm 4.69 Prostate vol (cc): Mean \pm SD total: 54.6 \pm 23.02 Median total: 48.4 Mean \pm SD transition zone*: 30.3 \pm 21.02 PSA serum, mean \pm sd:(ng/ml): 3.9 \pm 2.06 Qmax mean \pm sd (ml/sec): 10.6 \pm 3.57 Post-void residual vol, mean \pm sd, (ml): 67.4 \pm 63.49 No. sexually active (%): 1,189 (73) No. previous α-blocker use (%): 820 (51) No. previous 5-ARI use (%): 188 (12)</p> <p>Group 3: Tamsulosin + finasteride N: 1,610 Dropouts: Age, mean \pmsd,(yr): 66.0 \pm 7.05 No. white ethnicity (%): 1,421 (88) IPSS, mean \pm sd: 16.6 \pm 6.35</p>		<p>Other adverse events Breast neoplasm Floppy iris syndrome</p>	<p>0(0) 0(0) 0(0) 0(0) 0(0) 0(0)</p>	<p>* In a subset of 656 men. The baseline values were taken 4 weeks after screening, when all men received placebo treatment</p> <p>** General linear model adjusted for treatment, investigative site cluster, and baseline IPSS</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Duration since first LUTS mean \pm sd, (yr): 5.4 \pm 5.07</p> <p>Prostate vol (cc): Mean \pm SD total: 54.7 \pm 23.51 Median total: 48.9 Mean \pm SD transition zone*: 27.7 \pm 20.20</p> <p>PSA serum, mean \pm sd:(ng/ml): 4.0 \pm 2.05</p> <p>Qmax mean \pm sd (ml/sec): 10.9 \pm 3.62</p> <p>Post-void residual vol, mean \pm sd, (ml): 68.1 \pm 66.01</p> <p>No. sexually active (%): 1,176 (73)</p> <p>No. previous α-blocker use (%): 805 (50)</p> <p>No. previous 5-ARI use (%): 171 (11)</p>				

Evidence Table 11: Alpha-blockers vs. anticholinergics

See Evidence Table 9: Alpha-blockers vs. placebo
for Kaplan et al., 2006¹²³

Evidence Table 12: Alpha-blockers vs. phosphodiesterase-5 inhibitors

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kaplan et al., 2007¹³²</p> <p>Study design: RCT open label</p> <p>Setting: single-centre, Department of Urology, Weill Cornell Medical College, NY, USA</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: consecutive men with moderate to severe untreated LUTS and erectile dysfunction</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Moderate to severe untreated LUTS and self reported erectile dysfunction (not specific cut off points) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Contraindications to the study drugs <p>All patients N: 62</p> <p>Mean age: 63.4 ± 7.6</p> <p>Drop outs: 7 (11%) due to adverse events</p> <p>Group 1 (Sildenafil) N: 21</p> <p>Mean (± SD) Age: 64 ± 5.9</p> <p>Duration of LUTS, mths: 14.3 ± 2.4</p> <p>Duration of ED, mths: 25.6 ± 5.4</p> <p>Frequency: 9.3 ± 2.6</p> <p>Nocturia: 2.9 ± 0.6</p> <p>IPSS, mean ± SD: 17.3 ± 4.3</p> <p>IPSS moderate (8-19): 43%</p> <p>IPSS severe (>20): 57%</p> <p>IIEF-EF domain, mean ± SD: 14.3 ± 5.2</p> <p>IIEF Q3, mean ± SD: 2.1 ± 1.1</p> <p>IIEF Q5, mean ± SD: 2.3 ± 1.3</p> <p>Qmax, mean ± SD, mL/s: 9.7 ± 3.7</p> <p>PVR, mean ± SD, mL: 46 ± 14.3</p> <p>Dropouts: 2 (10%)</p> <p>Group 2 (Alfuzosin) N: 20</p> <p>Mean (± SD) Age: 62.6 ± 8.2</p> <p>Duration of LUTS, mths, mean ± SD: 12.4 ± 2.3</p> <p>Duration of ED, mths, mean ± SD: 22.5</p>	<p>Group 1: Sildenafil citrate 25 mg one daily at night</p> <p>Group 2: Alfuzosin 10mg once daily after the same meal</p> <p>Group 3: Sildenafil citrate 25 mg/day + Alfuzosin 10 mg/day</p> <p>Examination methods: Patients assessed at baseline and 12 weeks. IPSS taken and frequency and nocturia quantified with bladder diary. Qmax and PVR also assessed.</p> <p>Q3 frequency of penetration and Q4 frequency of maintained erection were analysed separately.</p>	<p>IPSS ± SD at 12 weeks P value calculated by NCGC as <i>t</i>-test with equal variances</p>	<p>Grp 1: 14.9 ± 4.2 Grp 2: 14.6 ± 3.7 Grp 3: 13.5 ± 4.2 P value grp 1 v grp 2 = 0.81</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> This was an open label study with no randomisation allocation and concealment methods reported. The outcomes are mainly subjective outcomes, and this makes it particularly at risk of biases. <p>Additional outcomes: % change from baseline for Qmax, PVR, frequency and nocturia IIEF Q3 % change from baseline and IIEF Q5 % change from baseline</p> <p>Notes: **Erectile Dysfunction assessed using the Erectile Function domain score of the 15-question IIEF, ie , ie Q1-5 and Q15 (Maximum score 30).</p> <p>This is different from IIEF-5, which consists</p>
			<p>IPSS change (%) from baseline at 12 weeks (<i>p</i> change from baseline <i>t</i>-test) Change (mean ±sd) calculated by NCGC from the difference in baseline and follow up values. % values as reported</p>	<p>Grp 1: -2.40 ±4.25 (11.8%) <i>p</i>=0.03 Grp 2: -2.30 ±3.91(15.6%) <i>p</i>=0.01 Grp 3: -2.70 ±3.96 (24.1%) <i>p</i>=0.002</p>	
			<p>Qmax mean ± SD P value calculated by NCGC as <i>t</i>-test with equal variances</p>	<p><u>at 12 weeks</u> Grp 1: 10.3 ± 2.4 Grp 2: 10.5 ± 2.3 Grp 3: 11.5 ± 2.9</p> <p><u>Change from baseline</u> Grp 1: 0.3±3.1 Grp 2: 1.1±2.3 Grp 3: 2.0±2.6</p>	
			<p>Frequency ± SD at 12 weeks P value calculated by NCGC as <i>t</i>-test with equal variances</p>	<p>Grp 1: 7.8 ± 1.7 Grp 2: 6.4 ± 2.1 Grp 3: 6.1 ± 2.2 P value grp 1 v grp 2 = 0.02</p>	
			<p>Nocturia ± SD at 12 weeks P value calculated by NCGC as <i>t</i>-test with equal variances</p>	<p><u>at 12 weeks</u> Grp 1: 2.1 ± 0.9 Grp 2: 1.8 ± 0.9 Grp 3: 1.8 ± 1.1</p> <p><u>Change from baseline</u> Grp 1:-0.8±0.8 Grp 2:-1.3±1.0 Grp 3:-1.1±1.0</p>	
<p>IIEF erectile function domain** ± SD at 12 weeks P value calculated by NCGC</p>	<p>Grp 1: 21.4 ± 5.7 Grp 2: 20.3 ± 5.2 Grp 3: 25.7 ± 4.9</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																												
	<p>± 4.9 Frequency, mean ± SD: 8.9 ± 2.5 Nocturia, mean ± SD: 3.1 ± 1.1 IPSS, mean ± SD: 16.9 ± 4.1 IPSS moderate (8-19): 45% IPSS severe (>20): 55% IIEF-EF , mean ± SD: 17.4 ± 4.9 IIEF Q3 , mean ± SD: 2.3± 1.3 IIEF Q5 , mean ± SD: 2.4 ± 1.2 Qmax , mean ± SD, mL/s: 9.4 ± 2.2 PVR, mean ± SD, mL: 54 ± 17.8 Dropouts: 2 (10%)</p> <p>Group 3 (Sildenafil + Alfuzosin) N: 21 Mean (± SD) Age: 63 ± 6.9 Duration of LUTS, mths mean±SD: 13.9±2.7 Duration of ED, mths, mean±SD: 26.9±5.4 Frequency, mean ± SD: 9.1 ± 2.2 Nocturia, mean ± SD: 2.89 ± 0.9 IPSS , mean ± SD: 16.2 ± 3.7 IPSS moderate (8-19): 48% IPSS severe (>20): 52% IIEF-EF mean± SD: 16.2 ± 3.7 IIEF Q3 , mean ± SD: 2.1 ± 1.1 IIEF Q5 , mean ± SD: 2.3 ± 1.3 Qmax , mean ± SD, mL/s: 9.5 ± 2.3 PVR , mean ± SD, mL: 53 ± 19.8 Dropouts: 3 (14%)</p>		<p>as t-test with equal variances</p> <p>IIEF erectile function domain** % change from baseline at 12 weeks (p change from baseline t-test)</p> <p>Adverse Events</p> <table border="1" data-bbox="1099 395 1440 635"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th>Grp 3</th> </tr> </thead> <tbody> <tr> <td>N</td> <td>21</td> <td>20</td> <td>21</td> </tr> <tr> <td>Withdrawals due to adverse events</td> <td>2</td> <td>2</td> <td>3</td> </tr> <tr> <td>Dizziness</td> <td>0</td> <td>2</td> <td>1</td> </tr> <tr> <td>Flushing</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Dyspepsia</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Gastric upset</td> <td>0</td> <td>0</td> <td>2</td> </tr> </tbody> </table>		Grp 1	Grp 2	Grp 3	N	21	20	21	Withdrawals due to adverse events	2	2	3	Dizziness	0	2	1	Flushing	1	0	0	Dyspepsia	1	0	0	Gastric upset	0	0	2	<p>P value grp 1 v grp 2 = 0.52</p> <p>Grp 1: 49.79%, p=0.01 Grp 2: 16.7%, p=0.11 Grp 3: 58.6%, p=0.002</p>	<p>of question Q2, Q4, Q5, Q7 and Q15 of the IIEF (maximum score 25).</p> <p>*Q3 - frequency of penetration and Q4 - frequency of maintained erection from the IIEF were analysed separately.</p> <p>% of IIEF change from baseline had been updated to correct publication error in original article.</p>
	Grp 1	Grp 2	Grp 3																														
N	21	20	21																														
Withdrawals due to adverse events	2	2	3																														
Dizziness	0	2	1																														
Flushing	1	0	0																														
Dyspepsia	1	0	0																														
Gastric upset	0	0	2																														

Evidence Table 13: 5-alpha reductase inhibitors vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Abrams et al., 1999⁷</p> <p>Setting: multi-centre, world wide</p> <p>Study design: RCT double blinded</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Men meeting objective evidence of obstruction after pressure flow studies</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> > 55 years Ambulatory Enlarged prostate by DRE Presence of LUTS <p>Exclusion criteria:</p> <ul style="list-style-type: none"> PSA > 10 ng/mL Need for immediate surgery PVR ≥300 mL Urethral strictures Chronic Bacterial prostatitis Neurogenic bladder Previous prostate or testicular surgery Prostate cancer or suspect Neurogenic bladder Acute UTI Use of drugs with anti-androgenic properties or alpha-blockers or plant extracts History of drug or alcohol abuse Evidence of renal or hepatic impairment History of recurrent renal or prostatic calculi <p>All patients N: 121 (out of 201 screened) Mean age: Drop outs: 15/121 (12.4%)</p> <p>Group 1 (Finasteride 5mg/dayl) N: 81</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Uroflowmetry performed at 4, 8, 12 months with voided volume of ≥ 150 mL. Prostate volume measured at baseline and month 12. IPSS assessed at 4, 8, 12 months</p>	<p>Mean change in IPSS ± SD from baseline at 1 year</p>	<p>Grp 1: -4.8 ± 6.4* (n=69) Grp 2: -3.3 ± 6.4* (n=37) P value: NS</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation & allocation concealment method not reported. Unclear whether examiners or investigators are masked. Primary outcomes are not changed in symptom score or adverse events <p>Additional outcomes: Detrusor pressure Free maximum flow rate</p> <p>Notes: Study was designed to detect differences in urodynamic parameters rather than symptom score.</p> <p>Randomisation was on a 2:1 basis</p> <p>* Standard deviation for change from baseline calculated using reported mean difference and confidence intervals for the between group comparison following methods from Cochrane Handbook</p> <p>Study reports that analysis of variance was used to</p>
			<p>Mean change in Qmax ± SD from baseline at 1 year</p>	<p>Grp 1: 1.1 ± 2.5 (n=69) Grp 2: -0.1 ± 1.5 (n=37) P value: 0.02</p>	
			<p>Withdrawals due to adverse events</p>	<p>Grp 1 Grp 2 3 3</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> Mean (\pm SD) Age: 68.1 \pm 6.1 IPSS \pm SD: 19.4 \pm 6.3 Qmax \pm SD, mL/s: 6.7 \pm 2.4 Prostate volume \pm SD, mL: 45.4 \pm 21.9 Number obstructed: 61 Number equivocal: 19 Dropouts: 12/81 (14.8%) </p> <p> Group 2 (Placebo 1/day) N: 40 Mean (\pm SD) Age: 67.4 \pm 7.2 IPSS \pm SD: 17.4 \pm 6.8 Qmax \pm SD, mL/s: 7.0 \pm 2.0 Prostate volume \pm SD, mL: 44.8 \pm 20.2 Number obstructed: 33 Number equivocal: 7 Dropouts: 3/40 (7.5%) </p>				<p>compare baseline to follow up with treatment centre and treatment group as variables.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments															
Andersen et al., 1995 ¹⁵	<p>Patient group: Men moderate symptoms of BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ≤ 80 years Ambulatory and good physical and mental health Qmax ≥5 ≤ 15 mL/s (at screening or start of placebo run-in) Enlarged prostate by DRE At least 2 symptoms indicting moderate BPH (increased frequency of urination or difficulty in urination) but not more than 2 severe symptoms <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Haematuria associated with UTI, prostatitis or bladder carcinoma Serum creatinine > 150 mmol/L or liver function tests ≥50% above normal Urethral strictures Chronic Bacterial prostatitis Previous prostate or testicular surgery Prostate cancer Neurogenic bladder ≥2 catheterisations for AUR in previous 2 years Significant abnormalities detected in screening examination Untreated UTI Use of drugs with anti-androgenic properties <p>All patients N: 707</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Physical examination including DRE was performed at baseline and months 12 and 24. Symptoms measured at baseline and months 1, 4, 8, 12, 16, 20 and 24 using modified Boyarsky scale (9 questions max score is 54) and obstructive symptoms totalled for Q1-5 as impairment in size and force of urinary stream, hesitancy or delay in starting urination, dribbling, interruption of stream, feeling of incomplete emptying (max score is 30)</p> <p>Flow rates measured using Dantec Urodyn 1000, PVR measured using portable ultrasound device at baseline and 12 & 24 months. Serum PSA at baseline and months 12 & 24.</p> <p>Subset of 416 patients had prostate volume measured by TRUS.</p>	<p>Mean change in total symptom score from baseline at 24 months (Boyarsky scale)</p> <p>Mean change in obstructive symptom score from baseline at 24 months (Boyarsky scale)</p> <p>Mean change in Qmax from baseline at 12 months estimated from graph with confidence intervals</p> <p>Mean change in Qmax from baseline at 24 months</p> <p>Mean change in Prostate volume from baseline at 24 months</p> <p>Median % change in PSA from baseline at 24 months</p> <p>Reason for withdrawal §</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>N</td> <td>353</td> <td>354</td> </tr> <tr> <td>Adverse Events</td> <td>39</td> <td>30</td> </tr> <tr> <td>Insufficient response</td> <td>13</td> <td>22</td> </tr> <tr> <td>Other (lost to follow up, protocol deviation, uncooperative)</td> <td>14</td> <td>12</td> </tr> </tbody> </table> <p>Adverse events – sexual dysfunction</p>		Grp 1	Grp 2	N	353	354	Adverse Events	39	30	Insufficient response	13	22	Other (lost to follow up, protocol deviation, uncooperative)	14	12	<p>Grp 1: -2.0 ± 6.2 *(n=347) Grp 2: 0.2 ± 7.6 * (n=346) P value: <0.01</p> <p>Grp 1: -1.5 ± 4.3 * (n=348) Grp 2: -0.2 ± 4.7 * (n=344) P value: <0.01</p> <p>Grp 1: 1.2 ± 3.1* (n=308) Grp 2: -0.3 ± 3.6* (n=309) P value: <0.01</p> <p>Grp 1: 1.5 ± 3.6* (n=308) Grp 2: -0.3 ± 3.1* (n=309) P value: <0.01</p> <p>Grp 1: -19.2 ± 23.1* (n=197) Grp 2: 11.5 ± 47.3 * (n=197) P value: <0.01</p> <p>Grp 1: -52% Grp 2: 6% P value: < 0.0001</p>	<p>Funding: Merck & Co, Inc.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation & allocation concealment method not reported. Unclear whether examiners or investigators are masked. Median changes from baseline reported. <p>Additional outcomes: Change in total symptom score at 12 months</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in to reduce placebo effect then randomised.</p> <p>Patients who withdrew were included in analysis using Last observation</p>
	Grp 1	Grp 2																		
N	353	354																		
Adverse Events	39	30																		
Insufficient response	13	22																		
Other (lost to follow up, protocol deviation, uncooperative)	14	12																		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean age: 65.5 (range 46-80) Drop outs: 130 (18.4%)</p> <p>Group 1 (Finasteride 5mg/day) N: 353 Mean (range) Age: NR Total symptom score: 13.4 ± NR (n=347) Total obstructive score: 8.8 ± NR (n=348) Qmax ± SD, mL/s: 10.2 ± NR (n=308) Prostate volume ± SD, mL: 40.6 ± NR (n=197) Dropouts: 66 (18.7%) see withdrawals§</p> <p>Group 2 (Placebo 1/day) N: 354 Mean (range) Age: NR Total symptom score: 13.1 ± NR (n=346) Total obstructive score: 8.6 ± NR (n=344) Qmax ± SD, mL/s: 10.5 ± NR (n=309) Prostate volume ± SD, mL: 41.7 ± NR (n=197) Dropouts: 64 (18.1%) see withdrawals§</p>				<p>Carried Forward.</p> <p>Study reports that analysis of variance used to compare outcomes but it unclear what variables were used in the model.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																																														
Beisland et al., 1992 ²⁸	<p>Patient group: men with symptomatic urinary obstruction</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> 40-80 years in good physical and mental health with symptoms of urinary obstructions and Qmax <15 ml/s documented by two measurements at screening. Enlarged prostate by DRE <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Clinical or laboratory abnormalities <p>All patients N: 182 Mean age: NR Drop outs: 14/182 (7.65)</p> <p>Group 1 (Finasteride 5mg/day) N: 94 Mean (range) Age: 66.6 (46-80) Total symptom score, mean ± SD: 8.8 ± 6.1 Total obstructive score, mean ± SD: 2.2 ± 4.0 Troublesome score, mean ± SD: Qmax ± SD, mL/s: 8.0 ± 3.0 Prostate volume ± SD, cm³: 44.2 ± 22.4 Drop outs: 7/94 (7.4%) see withdrawals§</p> <p>Group 2 (Placebo 1/day) N: 88 Drop outs: 3/88 (3.4%) Mean (range) Age: 68.0 (54-79)</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Symptoms were assessed using a modified Boyarksy scale modified which comprises 9 questions (max score is 36). Patients were treated as mild if the score was <6, moderate (6-13) and severe if scores were >13.</p> <p>Obstructive symptoms totalled for the following questions:</p> <ul style="list-style-type: none"> impairment of size and force of urinary stream hesitancy or delay in starting the flow of urine dribbling after urination feeling of incomplete emptying of the bladder interruption of urinary stream 	<p>Mean change in Qmax(ml/s) from baseline at 24 weeks</p> <p>Median % change in PSA from baseline at 12 weeks months</p> <p>Median % change in PSA from baseline at 24 weeks months</p> <p>Mediun % decrease iun prostate volume from baseline at 24 weeks</p> <p>§ Reason for withdrawal** (see notes)</p> <table border="1"> <thead> <tr> <th></th> <th>N</th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Adverse Events</td> <td>7</td> <td>7</td> <td>7</td> </tr> <tr> <td>No response</td> <td>6</td> <td>1</td> <td>3</td> </tr> <tr> <td>Other</td> <td>0</td> <td>3</td> <td>3</td> </tr> </tbody> </table> <p>Withdrawal due to sexual adverse events</p> <table border="1"> <thead> <tr> <th></th> <th>N</th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Adverse events</td> <td>93</td> <td>48</td> <td>48</td> </tr> <tr> <td>Insomnia and depression</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Deep vein thrombosis</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Urinary retention</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Decreased libido</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Impotence</td> <td>4</td> <td>4</td> <td>4</td> </tr> </tbody> </table>		N	Grp 1	Grp 2	Adverse Events	7	7	7	No response	6	1	3	Other	0	3	3		N	Grp 1	Grp 2	Adverse events	93	48	48	Insomnia and depression	1	0	0	Deep vein thrombosis	1	0	0	Urinary retention	1	0	0	Decreased libido	1	0	0	Impotence	4	4	4	<p>Grp 1: 1.6 ± 1.4* (n=87) Grp 2: 1.1 ± 1.4* (n=81) P value: 0.022(as reported)</p> <p>Grp 1: -22.4 Grp 2: No change P value < 0.001</p> <p>Grp 1: -32.4 Grp 2: No change P value < 0.001</p> <p>Grp 1: 22.5 Grp 2: 1.0 P value < 0.001</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Withdrawal due to sexual adverse events</td> <td>1</td> <td>1</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th></th> <th>N</th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Adverse events</td> <td>93</td> <td>48</td> <td>48</td> </tr> <tr> <td>Insomnia and depression</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Deep vein thrombosis</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Urinary retention</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Decreased libido</td> <td>1</td> <td>0</td> <td>0</td> </tr> <tr> <td>Impotence</td> <td>4</td> <td>4</td> <td>4</td> </tr> </tbody> </table>		Grp 1	Grp 2	Withdrawal due to sexual adverse events	1	1		N	Grp 1	Grp 2	Adverse events	93	48	48	Insomnia and depression	1	0	0	Deep vein thrombosis	1	0	0	Urinary retention	1	0	0	Decreased libido	1	0	0	Impotence	4	4	4	<p>Funding: Not stated. Most likely Merck Laboratories, as 4/12 authors were from Merck. .</p> <p>Limitations:</p> <ul style="list-style-type: none"> Method of randomisation and concealment not reported A modified Boyarksy scale was used <p>Additional outcomes: Change of total symptom score (Boyarksy scale) from baseline at 12 weeks for finasteride (-2.1) vs. placebo (-0.8) was significant (p=0.0046) for 12 weeks. Change for obstructive symptoms scores were -2.0 vs. -0.7 for 24 weeks (p=0.05) using analysis of covariance</p> <p>DHT level changes from baseline were also reported</p> <p>Notes: *Standard deviations for changes from baseline calculated from reported p values between groups using Cochrane methodology</p> <p>Analysis of covariance used to compare baseline parameters and % change from baseline.</p>
	N	Grp 1	Grp 2																																																																																
Adverse Events	7	7	7																																																																																
No response	6	1	3																																																																																
Other	0	3	3																																																																																
	N	Grp 1	Grp 2																																																																																
Adverse events	93	48	48																																																																																
Insomnia and depression	1	0	0																																																																																
Deep vein thrombosis	1	0	0																																																																																
Urinary retention	1	0	0																																																																																
Decreased libido	1	0	0																																																																																
Impotence	4	4	4																																																																																
	Grp 1	Grp 2																																																																																	
Withdrawal due to sexual adverse events	1	1																																																																																	
	N	Grp 1	Grp 2																																																																																
Adverse events	93	48	48																																																																																
Insomnia and depression	1	0	0																																																																																
Deep vein thrombosis	1	0	0																																																																																
Urinary retention	1	0	0																																																																																
Decreased libido	1	0	0																																																																																
Impotence	4	4	4																																																																																

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Total symptom score, mean \pm SD: 7.8 \pm 4.9</p> <p>Total obstructive score, mean \pm SD: 1.1 \pm 3.3</p> <p>Troublesome score, mean \pm SD: 6.8 \pm 3.9</p> <p>Qmax \pm SD, mL/s: 7.6 \pm 3.1</p> <p>Prostate volume \pm SD, cm³ 43.8 \pm 24.1</p>				<p>**6 year follow up reported by Ekman et al., 1998⁷⁸. The number of drop outs reported in this report was 14. Adverse events reported in more detail in BEISLAND 1992.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																		
Byrnes et al, 1995 ⁴⁰	<p>Patient group: Men attending community-based clinics for treatment of BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Clinical diagnosis of BPH based on moderate to severe symptoms with prostate gland enlargement on DRE PSA ≤ 10 ng/mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Urethral strictures Previous prostate surgery Pelvic radiotherapy Chronic Bacterial prostatitis Neurogenic bladder Recurrent UTI Use of drugs with anti-androgenic properties Use of hormonal therapy affecting prostate Prostate cancer or suspected <p>All patients N: 2417 included in safety analysis, 2342 in efficacy analysis Mean age: 65 Drop outs: 465 (19.2%)</p> <p>Group 1 (Finasteride 5mg/day) N: 1821 randomised 1759 efficacy Mean (range) Age: 65 (42-91) White/other: 1226 Black: 285 Hispanic: 248 AUA symptom score mild (<8): 33 AUA symptom score moderate (8-19): 1001</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Physical examination including DRE was performed at baseline and 12 mths. Serum dihydrotestosterone measured at baseline and mths 6 & 12 AUA-7 Symptom score, BPH Impact Index (BII) used for HRQoL, Patient satisfaction with urinary condition as extra question (0-6) and additional questions from modified BSIA instrument to measure interference with activities and extra question about adjustment of activities to cope with urinary symptoms were</p>	<p>Mean change in AUA-7 symptom score from baseline at 3 months Estimated from graph with confidence intervals. Numbers at follow up not clear so total for efficacy analysis used.</p> <p>Mean change in AUA-7 symptom score from baseline at 6 months estimated from graph with confidence intervals</p> <p>Mean change in AUA-7 symptom score from baseline at 12 months estimated from graph with confidence intervals</p> <p>Mean change in BPII at 12 months</p> <p>Mean change in patient global assessment at 12 months</p> <p>% Patients rating themselves “better” at 12 mths</p> <p>% Investigators rating patients “better” at 12 mths</p> <p>Reason for withdrawal §</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Total withdrawals</td> <td>343</td> <td>122</td> </tr> <tr> <td>Adverse Events</td> <td>100</td> <td>28</td> </tr> <tr> <td>Lost to follow up</td> <td>81</td> <td>30</td> </tr> <tr> <td>Treatment failure</td> <td>62</td> <td>24</td> </tr> <tr> <td>Protocol violation or other</td> <td>100</td> <td>40</td> </tr> </tbody> </table> <p>no significant differences</p>		Grp 1	Grp 2	Total withdrawals	343	122	Adverse Events	100	28	Lost to follow up	81	30	Treatment failure	62	24	Protocol violation or other	100	40	<p>Grp 1: -3.3 ± 7.7* (n=1759) Grp 2: -2.6 ± 7.8* (n=583) P value: <0.05</p> <p>Grp 1: -4.1 ± 7.7* (n=1759) Grp 2: -3.3 ± 7.8* (n=583) P value: <0.05</p> <p>Grp 1: -4.6 ± 9.6* (n=1759) Grp 2: -3.3 ± 8.6* (n=583) P value: <0.05</p> <p>Grp 1: -1.2 ± 4.2* (n=1711) Grp 2: -0.9 ± 3.7* (n=575) P value: <0.04 (ANOVA)</p> <p>Grp 1: 4.9 ± 2.1.2* (n=1714) Grp 2: 4.7 ± 1.2* (n=575) P value: 0.0001 (ANOVA)</p> <p>Grp 1: 56.2 % Grp 2: 44.2 % P value: <0.001</p> <p>Grp 1: 55.3 % Grp 2: 45.8 % P value: <0.001</p>	<p>Funding: Merck & Co, Inc.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation & allocation concealment method not reported.. Unclear whether examiners or investigators are masked. Numbers of patients remaining at each time point not clear for AUA score. <p>Additional outcomes: BPII + patient satisfaction question at 12 mths, activities of living score at 12 mths, general adjustment question at 12 mths, investigator global assessment at 12 mths</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in. Men with moderate to severe symptoms after run-in with good compliance were randomised in 3:1 ratio.</p> <p>*Standard deviations for changes from baseline calculated using</p>
	Grp 1	Grp 2																					
Total withdrawals	343	122																					
Adverse Events	100	28																					
Lost to follow up	81	30																					
Treatment failure	62	24																					
Protocol violation or other	100	40																					

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																												
	<p>AUA symptom score severe (20-35): 724 AUA symptom score unknown: 1 BII: 5.1 CI95% 4.9-5.2 BII + patient satisfaction: 8.8 CI95% 8.6-9.0 Activities of living score: 13.3 CI95% 12.8-13.8 Adjustment question: 1.4 CI95% 1.3-1.5 Dropouts: 343 (19.4%) for reasons see§</p> <p>Group 2 (Placebo 1/day) N: 596 randomised 583 efficacy Mean (range) Age: 65.1 (45-91) White/other: 397 Black: 95 Hispanic: 91 AUA symptom score mild (<8): 13 AUA symptom score moderate (8-19): 335 AUA symptom score severe (20-35): 235 AUA symptom score unknown: 0 BII: 5.0 CI95% 4.8-5.3 BII + patient satisfaction: 8.6 CI95% 8.3-9.0 Activities of living score: 12.8 CI95% 11.9-13.7 Adjustment question: 1.3 CI95% 1.2-1.4 Dropouts: 122 (20.4%) for reasons see§</p>	<p>taken at baseline and 3 mth intervals. Patient and investigator global assessment of change in urologic status also rated from 1 (much worse) to 7 (much better) every 3 mths. Patients with visual impairment had questionnaires read to them and Spanish versions provided.</p>	<p>Adverse events</p> <table> <thead> <tr> <th>N randomised</th> <th>Grp 1</th> <th>Grp 2</th> <th></th> </tr> <tr> <th></th> <th>1821</th> <th>596</th> <th></th> </tr> </thead> <tbody> <tr> <td>Impotence**</td> <td>102</td> <td>13</td> <td>p < 0.0001</td> </tr> <tr> <td>Libido decrease**</td> <td>53</td> <td>6</td> <td>p = 0.008</td> </tr> <tr> <td>Ejaculation disorder**</td> <td>38</td> <td>3</td> <td>p = 0.009</td> </tr> <tr> <td>Withdrawal due to sexual adverse events</td> <td>27</td> <td>3</td> <td>p = 0.06</td> </tr> <tr> <td>Acute urinary retention</td> <td>11</td> <td>4</td> <td>p = 0.77</td> </tr> </tbody> </table> <p>** Possibly, probably or definitely drug related</p>	N randomised	Grp 1	Grp 2			1821	596		Impotence**	102	13	p < 0.0001	Libido decrease**	53	6	p = 0.008	Ejaculation disorder**	38	3	p = 0.009	Withdrawal due to sexual adverse events	27	3	p = 0.06	Acute urinary retention	11	4	p = 0.77	<p>between groups</p>	<p>confidence intervals and Cochrane methodology</p> <p>Study reports that analysis of variance was used to compare baseline to follow up with race and treatment-by-race as variables. It is unclear whether the results presented have been adjusted for these variables.</p>
N randomised	Grp 1	Grp 2																															
	1821	596																															
Impotence**	102	13	p < 0.0001																														
Libido decrease**	53	6	p = 0.008																														
Ejaculation disorder**	38	3	p = 0.009																														
Withdrawal due to sexual adverse events	27	3	p = 0.06																														
Acute urinary retention	11	4	p = 0.77																														

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																				
<p>Finasteride Study Group, 1993⁹²</p> <p>Setting: multicentre worldwide</p> <p>Study design: RCT double blinded</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Men with BPH and symptoms of BOO</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> 40-80 years Good physical and mental health Qmax < 15 mL/s (from 2 measurements) Prostate volume ≥ 30 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Bacterial prostatitis Previous prostate or testicular surgery Prostate cancer PSA ≥ 40 ng/mL PVR > 350 mL Neurogenic bladder Repeated catheterisations Use of drugs with anti-androgenic properties <p>All patients N: 750 (all treatment arms) Mean age: NR Drop outs: NR</p> <p>Group 1 (Finasteride 5mg/day) N: 249 Mean (range) Age: 66 (46-83) Total obstructive score (max 20): 11.2 ± 3.8 Total symptom score (max 36): 18.6 ± 6.0 Qmax ± SD, mL/s: 9.2 ± 4.0 Prostate volume ± SD, mL: 47.0 ± 20.8 PSA ± SD, ng/mL: 5.8 ± 6.7</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Group 3: Finasteride 1 mg 1/day</p> <p>Results and baseline characteristics reported for normal dose finasteride arm 5mg/day only</p> <p>Examination methods: At baseline and months 3, 6 & 12 prostate volume measured by TRUS and Qmax measured at by Dantec Urolynx 1000 uroflowmeter, Boyarsky symptom questionnaire taken (9 questions). Testosterone, dihydrotestosterone, luteinising hormone measured at baseline and weeks 2, 8, 16, 24 and 9 and 12 months. Thyroxine and thyroid stimulating hormone measured at baseline and months 3 & 6. PSA measured at -2, 12, 24 weeks and 9 & 12 months</p>	<p>Median change in total symptom score (Boyarsky scale) from baseline at 12 months Estimated from graph</p> <p>Median change in Qmax from baseline at 12 months Estimated from graph</p> <p>% patients achieving ≥ 3 mL/s flow increase</p> <p>Median % change in prostate volume from baseline at 12 months</p> <p>Median % change in PSA from baseline at 12 months</p> <p>Adverse Events</p> <table border="1"> <thead> <tr> <th></th> <th>N</th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Withdrawals due to adverse events</td> <td></td> <td>249</td> <td>255</td> </tr> <tr> <td>Impotence</td> <td></td> <td>1</td> <td>0</td> </tr> <tr> <td>Acute urinary retention</td> <td></td> <td>12</td> <td>1</td> </tr> <tr> <td></td> <td></td> <td>3</td> <td>3</td> </tr> </tbody> </table> <p>p < 0.001</p>		N	Grp 1	Grp 2	Withdrawals due to adverse events		249	255	Impotence		1	0	Acute urinary retention		12	1			3	3	<p>Grp 1: 3.3 Grp 2: 2.0 P value = signif (value NR)</p> <p>Grp 1: 1.38 Grp 2: 0.42 P value = 0.025</p> <p>Grp 1: 31.0 % Grp 2: 21.0 %</p> <p>Grp 1: 22.4 % Grp 2: 5.0 % P value < 0.001</p> <p>Grp 1: 46.0 % Grp 2: 0 (no change) % P value < 0.001</p>	<p>Funding: Merck</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation & allocation concealment method not reported. Unclear whether examiners or investigators are masked. Median changes from baseline reported. Dropouts not clearly reported <p>Additional outcomes: % change from baseline for plasma dihydrotestosterone</p> <p>Notes: Eligible patients entered a 2 week month single blind placebo run-in to reduce placebo effect then randomised.</p>
	N	Grp 1	Grp 2																						
Withdrawals due to adverse events		249	255																						
Impotence		1	0																						
Acute urinary retention		12	1																						
		3	3																						

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Dropouts: Not clear. 1 patients withdrew due to impotence but others not mentioned</p> <p>Group 2 (Placebo 1/day) N: 255 Mean (range) Age: 66 (46-81) Total obstructive score (max 20): 11.1 ± 3.7 Total symptom score (max 36): 18.2 ± 5.9 Qmax ± SD, mL/s: 8.6 ± 3.4 Prostate volume ± SD, mL: 46.3 ± 23.4 PSA ± SD, ng/mL: 5.7 ± 7.2 Dropouts: NR</p>				<p>Analysis of variance used to compare outcomes with treatment centre and treatment group and treatment-centre interaction as model parameters</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																						
Gormley et al., 1992 ¹⁰⁵	Patient group: Men with BPH and symptoms of urinary obstruction	Group 1: Finasteride 5 mg 1/day	Mean symptom score(Boyarsky) at 12 months	Grp 1: 7.5 ± 5.2 (n=257) Grp 2: 8.8 ± 6.1 (n=263) P value: <0.05	Funding: Merck & Co, Inc. Limitations: <ul style="list-style-type: none"> • Randomisation & allocation concealment method not reported. • Unclear whether key examiners or investigators are masked. Additional outcomes: Median PSA at follow up, Median change in prostatic volume % at follow up. Mean Qmax + SE at follow up as graph. Notes: Eligible patients entered 2 week single blind placebo run-in. ITT analysis with missing data from last observation carried forward. Analysis of variance used to																																						
Finasteride study group	Inclusion criteria: <ul style="list-style-type: none"> • 40-83 years • Enlarged prostate gland enlargement on DRE • Qmax < 15 mL/s with voided volume of ≥ 150 mL • Men with very low urinary flow rates unless at risk for total obstruction Exclusion criteria: <ul style="list-style-type: none"> • Prostate cancer or suspected • PVR > 350 mL • Serum PSA ≥ 40 µg/L • UTI • Chronic prostatitis • Neurogenic bladder 	Group 2: Placebo 1/day	Mean obstruction score(Boyarsky) at 12 months	Grp 1: 5.1 ± 3.6 (n=257) Grp 2: 5.9 ± 3.8 (n=263) P value: <0.001																																							
Setting: multi-centre, 25 centres in USA and 5 in Canada		Group 3: Finasteride 1 mg 1/day	Mean Qmax at 12 months	Grp 1: 11.2 ± 4.7 (n=257) Grp 2: 9.8 ± 3.7 (n=263) P value: <0.001																																							
Study design: RCT double blinded		Results and baseline characteristics reported for normal dose finasteride arm 5mg/day only	Mean Prostate volume at 12 months	Grp 1: 47.5 ± 23.6 (n=257) Grp 2: 59.8 ± 39.4 (n=263) P value: <0.001																																							
Evidence level: 1+		Examination methods: Men were examined monthly by the same investigator for symptoms (Boyarsky – 9 questions max score 36), obstructive symptoms (Boyarsky – first 5 questions max score 20), side effects and compliance.	Reason for withdrawal *	<table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Total Adverse Events</td> <td>40</td> <td>37</td> </tr> <tr> <td>Lost to follow up</td> <td>16</td> <td>18</td> </tr> <tr> <td>Treatment failure</td> <td>3</td> <td>4</td> </tr> <tr> <td>Other</td> <td>12</td> <td>9</td> </tr> <tr> <td></td> <td>9</td> <td>6</td> </tr> </tbody> </table>			Grp 1	Grp 2	Total Adverse Events	40	37	Lost to follow up	16	18	Treatment failure	3	4	Other	12	9		9	6																				
	Grp 1	Grp 2																																									
Total Adverse Events	40	37																																									
Lost to follow up	16	18																																									
Treatment failure	3	4																																									
Other	12	9																																									
	9	6																																									
Duration of follow-up: 12 months	All patients N: 895 (all study arms) Mean age: 64 Drop outs: 105/895 (11.7%) Group 1 (Finasteride 5mg/dayl) N: 297 Mean (range) Age: 64 (40-80) White: 286 Black: 6 Other: 5 Total Symptom score ± SD: 10.2 ± 5.5 Obstructive symptom score ± SD: 7.0 ± 3.6 Qmax ± SD, mL/s: 9.6 ± 3.7 Prostate volume, mL: 58.6 ± 30.5 Serum PSA ± SD, µg/L: 3.6 ± 4.2	Flow rate measured using Urodyn 1000, PVR using TRUS. Prostate volume measured using MRI at baseline, 3, 6 & 12 mths,, ophthalmic examination at 12 mths; serum amino-transferases, urea nitrogen, creatinine, Na, K, Ca and glucose	Adverse events **	<table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>N randomised</td> <td>297</td> <td>300</td> </tr> <tr> <td>Impotence</td> <td>10</td> <td>5</td> </tr> <tr> <td>Libido decrease</td> <td>14</td> <td>4</td> </tr> <tr> <td>Ejaculation disorder</td> <td>13</td> <td>5</td> </tr> <tr> <td>Breast pain</td> <td>1</td> <td>0</td> </tr> <tr> <td>Digestive system</td> <td>8</td> <td>6</td> </tr> <tr> <td>Dizziness</td> <td>0</td> <td>2</td> </tr> <tr> <td>Headache</td> <td>2</td> <td>2</td> </tr> <tr> <td>Asthenia</td> <td>3</td> <td>3</td> </tr> <tr> <td>lens opacity</td> <td>0</td> <td>2</td> </tr> <tr> <td>lens change</td> <td>2</td> <td>0</td> </tr> <tr> <td>Withdrawal due to sexual dysfunction</td> <td>4</td> <td>1</td> </tr> </tbody> </table>		Grp 1	Grp 2	N randomised	297	300	Impotence	10	5	Libido decrease	14	4	Ejaculation disorder	13	5	Breast pain	1	0	Digestive system	8	6	Dizziness	0	2	Headache	2	2	Asthenia	3	3	lens opacity	0	2	lens change	2	0	Withdrawal due to sexual dysfunction	4	1
	Grp 1	Grp 2																																									
N randomised	297	300																																									
Impotence	10	5																																									
Libido decrease	14	4																																									
Ejaculation disorder	13	5																																									
Breast pain	1	0																																									
Digestive system	8	6																																									
Dizziness	0	2																																									
Headache	2	2																																									
Asthenia	3	3																																									
lens opacity	0	2																																									
lens change	2	0																																									
Withdrawal due to sexual dysfunction	4	1																																									
			** Possibly, probably or definitely drug related																																								

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>PVR ± SD, mL: 73 ± 89 Dropouts: 40 (13%) for reasons see*</p> <p>Group 2 (Placebo 1/day) N: 300 Mean (range) Age: 64 (45-82) White: 288 Black: 8 Other: 4 Total Symptom score ± SD: 9.8 ± 5.3 Obstructive symptom score ± SD: 6.7 ± 3.5 Qmax ± SD, mL/s: 9.6 ± 3.5 Prostate volume, mL: 61.0 ± 36.5 Serum PSA ± SD, µg/L: 4.1 ± 4.8 PVR ± SD, mL: 73 ± 91 Dropouts: 37 (12%) for reasons see*</p>	<p>measured every 3 mths. Compliance determined by counting number of tablets remaining and serum dihydrotestosterone measurements</p>			<p>compare outcomes with treatment centre and treatment group as model parameters..</p>

See Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors

for Lepor et al., 1996¹⁶³ .

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																												
<p>Marberger et al., 1998¹⁸¹</p> <p>PROWESS study group</p> <p>Setting: multi-centre, 285 worldwide</p> <p>Study design: RCT double blinded (patients and investigators)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Men moderate symptoms of BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 50 - 75 years • Good general health • Enlarged prostate gland enlargement on DRE • Qmax 5 - 15 mL/s with a voided volume \geq 150mL (2 measurements) • No more than 2 severe symptoms on modified Boyarsky scale • PSA < 10 ng/mL • PVR < 150 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Dysuria, haematuria • Previous prostate or bladder surgery • Concurrent use of alpha-blockers or anti-androgens • Recurrent UTI • Chronic prostatitis • Bladder cancer • Abnormalities on clinical examination • Liver function tests >50% above upper limit of normal • Allergies • History of drug or alcohol abuse • Prostate cancer or suspected • Neurogenic bladder • Urinary catheterisation for AUR twice during previous 2 years • Poor compliance during placebo run in. 	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Total and obstructive symptom score on modified Boyarsky scale measured at baseline and every 4 months. Prostate volume measured at baseline and 1 and 2 years by TRUS.</p>	<p>Mean change \pm SD in total symptom score at 1 year (Boyarsky scale)</p>	<p>Grp 1: -2.9 \pm NR</p> <p>Grp 2: -1.9 \pm NR</p> <p>P value: \leq0.001 (ANOVA)</p>	<p>Funding: Merck & Co, Inc. manufacturers of finasteride</p> <p>Limitations: Standard deviations for Qmax were not reported.</p> <p>Additional outcomes: Change in obstructive symptom score at 1 and 2 years % change in prostate volume</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in prior to computer generated randomisation.</p> <p>Sample size of 3000 to detect change in symptom score of 1.4 \pm 7 from baseline and change of 1.1 \pm 5 mL/s in Qmax and 11% \pm 40 change in</p>																												
			<p>Mean change \pm SD in total symptom score at 2 years(Boyarsky scale)</p>	<p>Grp 1: -3.2 \pm NR</p> <p>Grp 2: -1.5 \pm NR</p> <p>P value: \leq0.001 (ANOVA)</p>																													
			<p>Mean change in Qmax \pm SD at 1 year</p>	<p>Grp 1: 1.2 \pm NR</p> <p>Grp 2: 0.6 \pm NR</p> <p>P value: 0.01 (ANOVA)</p>																													
			<p>Mean change in Qmax \pm SD at 2 year</p>	<p>Grp 1: 1.5 \pm NR</p> <p>Grp 2: 0.7 \pm NR</p> <p>P value: 0.002 (ANOVA)</p>																													
			<p>Mean % change in prostate volume from baseline at 1 year</p>	<p>Grp 1: -13 \pm NR</p> <p>Grp 2: +5 \pm NR</p> <p>P value: \leq0.01 (ANOVA)</p>																													
			<p>Mean % change in prostate volume from baseline at year</p>	<p>Grp 1: -15 \pm NR</p> <p>Grp 2: +9 \pm NR</p> <p>P value: \leq0.001 (ANOVA)</p>																													
			<p>Reason for withdrawal *</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Total discontinuations</td> <td>331</td> <td>360</td> </tr> <tr> <td>Adverse Events</td> <td>111</td> <td>144</td> </tr> <tr> <td>Lack of improvement</td> <td>50</td> <td>64</td> </tr> <tr> <td>Protocol deviation</td> <td>25</td> <td>14</td> </tr> <tr> <td>Patient compliance</td> <td>40</td> <td>40</td> </tr> <tr> <td>Loss to follow up</td> <td>70</td> <td>55</td> </tr> <tr> <td>Other</td> <td>36</td> <td>47</td> </tr> </tbody> </table>			Grp 1	Grp 2	Total discontinuations	331	360	Adverse Events	111	144	Lack of improvement	50	64	Protocol deviation	25	14	Patient compliance	40	40	Loss to follow up	70	55	Other	36	47					
				Grp 1		Grp 2																											
			Total discontinuations	331		360																											
			Adverse Events	111		144																											
Lack of improvement	50	64																															
Protocol deviation	25	14																															
Patient compliance	40	40																															
Loss to follow up	70	55																															
Other	36	47																															
<p>Drug related adverse events (>1%)</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th></th> </tr> </thead> <tbody> <tr> <td>Total in safety analysis</td> <td>1577</td> <td>1591</td> <td></td> </tr> <tr> <td>Decreased libido</td> <td>63</td> <td>44</td> <td></td> </tr> <tr> <td>Impotence</td> <td>104</td> <td>74</td> <td>p <0.05</td> </tr> <tr> <td>Ejaculation disorder</td> <td>33</td> <td>9</td> <td>p <0.05</td> </tr> <tr> <td>Urinary retention</td> <td>17</td> <td>35</td> <td>p <0.05</td> </tr> <tr> <td>Asthenia/fatigue</td> <td>11</td> <td>24</td> <td>p <0.05</td> </tr> <tr> <td>Rash</td> <td>17</td> <td>21</td> <td></td> </tr> </tbody> </table>		Grp 1	Grp 2		Total in safety analysis	1577	1591		Decreased libido	63	44		Impotence	104	74	p <0.05	Ejaculation disorder	33	9	p <0.05	Urinary retention	17	35	p <0.05	Asthenia/fatigue	11	24	p <0.05	Rash	17	21		
	Grp 1	Grp 2																															
Total in safety analysis	1577	1591																															
Decreased libido	63	44																															
Impotence	104	74	p <0.05																														
Ejaculation disorder	33	9	p <0.05																														
Urinary retention	17	35	p <0.05																														
Asthenia/fatigue	11	24	p <0.05																														
Rash	17	21																															

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<ul style="list-style-type: none"> Planned fatherhood <p>All patients N: 2902 in efficacy analysis (368 excluded from some centres for poor clinical practice) and 3168 included in safety analysis Mean age: Drop outs:</p> <p>Group 1 (Finasteride 5mg/day) N: 1450 Mean (± SD) Age: 63.0 ± 6.3 Total Symptom score (Boyarksy) ± SD: 14.5 ± 7.3 Obstructive score ± SD: 9.3 ± 4.6 Qmax ± SD, mL/s: 11.2 ± 5.9 Prostate volume, mL: 38.7 ± 20.1 Dropouts: 331/1450 (23%) see*</p> <p>Group 2 (Placebo 1/day) N: 1452 Mean (± SD) Age: 63.4 ± 6.1 Total Symptom score (Boyarksy) ± SD: 14.3 ± 7.2 Obstructive score ± SD: 9.1 ± 4.5 Qmax ± SD, mL/s: 10.9 ± 3.6 Prostate volume, mL: 39.2 ± 20.2 Dropouts: 360/1452 (23%) see*</p>		<p>Headache 33</p> <p>Withdrawal due to sexual problem 22</p> <p>UTI 28</p> <p>Hypertension 48</p> <p>Myocardial infarction or angina 44</p> <p>Abdominal Pain 38</p> <p>Gastric problems (pain, gastritis, diarrhoea) 72</p> <p>Respiratory (infection or bronchitis) 55</p> <p>Influenza or pharyngitis 57</p> <p>Back pain 27</p> <p>Dysuria 16</p> <p>Haematuria 10</p> <p>BPH worsening 35</p>	<p>36</p> <p>16</p> <p>40</p> <p>58</p> <p>29</p> <p>36</p> <p>64</p> <p>61</p> <p>55</p> <p>46</p> <p>13</p> <p>24</p> <p>64</p> <p>p <0.05</p> <p>p <0.05</p>	<p>prostate volume of power=99% and α 0.05.</p> <p>Data collected for those patients that discontinued</p> <p>** Mean change and SD from baseline were estimated from graphs for mean change and standard error.</p> <p>Analysis of variance used to compare outcomes but it's not clear what variables have been included in the model</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>McConnell et al., 1998¹⁹⁰</p> <p>Study also reported in Roehrborn et al., 2000²⁵⁸</p> <p>PLESS study group</p> <p>Setting: multi-centre, 95 centres in USA</p> <p>Study design: RCT double blinded</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 years</p>	<p>Patient group: Men moderate to severe symptoms of BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Enlarged prostate gland enlargement on DRE Qmax < 15 mL/s PVR < 300 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous prostate or bladder surgery Concurrent use of alpha-blockers or anti-androgens Recurrent UTI Chronic prostatitis PSA >10 ng/mL (those with PSA > 4 ng/mL had a TRUS biopsy to rule out prostate cancer) <p>All patients N: 3040 randomised but 1 centre closed (n=24) so data available for 3016 patients Mean age: Drop outs: 1157/3040 (38%)</p> <p>Group 1 (Finasteride 5mg/day) N: 1524 Mean (± SD) Age: 64.0 ± 6.3 White: 94.9 % Black: 3% Other: 2.1% Quasi AUA Symptom score ± SD: 15.2 ± 5.6 Qmax ± SD, mL/s: 10.9 ± 3.9 Prostate volume, mL: 54 ± 25 Serum PSA ± SD, µg/L: 2.8 ± 2.1 Dropouts: 524/1524 (34%) see*</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Patients were evaluated every 4 months for symptom score, flow rate (>150mL) and side effects. PSA was measured every 4 months for 1 year and every 8 months thereafter. Blood components and DRE performed every year and biopsy if clinically indicated.</p> <p>Prostate volume was measured in a subset of 10% of patients at 13 sites using MRI.</p> <p>At the beginning of the study symptom score was assessed using a symptom score validated by Bolognese et al., 1992 comprising the same components as the AUA but with a slightly different score.</p> <p>The AUA symptom score was then adopted and the data from both scores combined as a Quasi AUA 0-34 points (1-5 for 6 questions and</p>	<p>Mean change ± SD in Quasi-AUA score at 1 year**</p> <p>Mean change ± SD in Quasi-AUA score at 2 year**</p> <p>Mean change ± SD in Quasi-AUA score at 3 year**</p> <p>Mean change ± SD in Quasi-AUA score at 4 year**</p> <p>Mean change in Qmax ± SD at 1 year**</p> <p>Mean change in Qmax ± SD at 2 year**</p> <p>Mean change in Qmax ± SD at 3 year**</p> <p>Mean change in Qmax ± SD at 4 year**</p> <p>Mean change (%) in prostate volume at 1 year</p> <p>Mean change (%) in prostate volume at 2 year</p>	<p>Grp 1: -2.4 ± 4.5 (n=1314) Grp 2: -1.6 ± 4.5 (n=1296) P value: NR</p> <p>Grp 1: -2.9 ± 6.4 (n=1153) Grp 2: -1.3 ± 6.2 (n=1101) P value: NR</p> <p>Grp 1: -3.1 ± 6.1 (n=1047) Grp 2: -1.3 ± 5.8 (n=961) P value: NR</p> <p>Grp 1: -3.3 ± 5.8 (n=965) Grp 2: -1.1 ± 5.5 (n=853) P value: NR</p> <p>Grp 1: 1.3 ± 3.1 (n=928) Grp 2: 0.2 ± 3.0 (n=899) P value: NR</p> <p>Grp 1: 1.8 ± 5.6 (n=786) Grp 2: 0.4 ± 5.4 (n=720) P value: NR</p> <p>Grp 1: 1.8 ± 5.3 (n=691) Grp 2: 0.0 ± 4.9 (n=608) P value: NR</p> <p>Grp 1: 2.0 ± 4.9 (n=588) Grp 2: 0.2 ± 4.9 (n=496) P value: NR</p> <p>Grp 1: -1.6 (n=144) Grp 2: +5 (n=136) P value: NR</p> <p>Grp 1: -1.8 (n=130) Grp 2: +9 (n=119) P value: NR</p>	<p>Funding: Merck & Co, Inc. manufacturers of finasteride</p> <p>Limitations:</p> <ul style="list-style-type: none"> High discontinuation rate at >30% for both arms though efforts were made to retrieve data (see notes) Unclear whether key examiners or investigators are masked. <p>Additional outcomes: % change in prostate volume</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in prior to computer</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Group 2 (Placebo 1/day) N: 1516 Mean (± SD) Age: 63.9 ± 6.6 White: 995.5.9 % Black: 3% Other: 1.5% Quasi AUA Symptom score ± SD: 15.2 ± 5.8 Qmax ± SD, mL/s: 11.1 ± 4.8 Prostate volume, mL: 55 ± 26 Serum PSA ± SD, µg/L: 2.8 ± 2.1 Dropouts: 633/1516 (42%) see *	1-4 for 1 question)	Mean change (%) in prostate volume at 3 year	Grp 1: -17 (n=116) Grp 2: +11 (n=98) P value: NR	generated randomisation stratified according to centre Those discontinuing study were also contacted at 6 months after discontinuing study and at the 4 year end point. Complete outcome data was collected for 92% in both treatment groups including discontinuations. ** Mean change and SD from baseline were estimated from graphs for mean change and standard error.
Mean change (%) in prostate volume at 4 year			Grp 1: -17 (n=102) Grp 2: +14 (n=85) P value: NR		
Reason for withdrawal *			Grp 1 Grp 2		
Total discontinuations			524 633		
Adverse Events			176 166		
Lack of improvement	99 104				
Worsening of disease	23 56				
Need for surgery or medical therapy	80 172				
Loss to follow up	52 36				
Other	94 99				
Spontaneous or precipitated AUR Acute urinary retention defined as spontaneous (no precipitating factors) or precipitated (stroke, UTI, pre surgery etc)	Grp 1: 42/1503 Grp 2: 99/1513 P value: NR				
Drug related adverse events (>1%) in year 1	Grp 1 Grp 2				
Decreased libido	1503 1513				
Impotence	96 51	p =0.002			
Ejaculation disorder	122 56	p <0.001			
Breast tenderness	12 2	p =0.003			
Breast enlargement	6 2	NR			
Rash	8 2	p=0.04			
	8 3	NR			

See Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?

for McConnell et al., 2003¹⁷⁰.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments		
<p>Nickel et al., 1996²²⁰</p> <p>Setting: multi-centre, 28 sites in Canada</p> <p>PROSPECT study</p> <p>Study design: RCT double blinded. Patients and investigators.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Men moderate symptoms of BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ≤ 80 years Ambulatory and in good health Qmax 5 - 15 mL/s (at screening or start of placebo run-in) Enlarged prostate by DRE At least 2 symptoms indicating moderate BPH (increased frequency of urination or difficulty in urination) but not more than 2 severe symptoms Serum PSA ≤ 10 ng/mL PVR ≤ 150 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or suspect Neurogenic bladder ≥2 catheterisations for AUR in previous 2 years Previous prostate or testicular surgery Urethral strictures Chronic Bacterial prostatitis Serum creatinine > 150 mmol/L or liver function tests ≥50% above normal Use of drugs with anti-androgenic properties Haematuria associated with UTI, prostatitis or bladder carcinoma Any condition that might jeopardise the patient's ability to complete the study 	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: At baseline and 12 and 24 months patients received a physical examination including DRE, urodynamics, serum PSA, liver function tests, and urinalysis.</p> <p>Primary outcomes for symptom score and flow rates measured every 4 months. Symptoms assessed using the Boyarksy scale modified by Bolognese et al. which comprises 9 questions (max score is 54) and obstructive symptoms totalled for Q1 -5 as impairment in size and force of urinary stream, hesitancy or delay in starting urination, dribbling, interruption of stream, feeling of incomplete emptying (max score is 30)</p> <p>A quasi IPSS score was also developed using the seven items that corresponded from the Boyarsky scale and condensing the 2 highest values on the 6 point scale to 1.</p>	<p>Mean change in Quasi-IPSS ± SD from baseline at 4 months</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: -1.0 ± 4.9*</p> <p>Grp 2: -1.0 ± 5.3*</p> <p>P value: NS</p>	<p>Funding: Merck Frost Canada, inc.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Quasi IPSS score Data estimated from graph. Unclear how many patients remaining at each time interval. <p>Additional outcomes: Mean change in total symptom score and obstructive score from baseline and % change in prostate volume from baseline.</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in to reduce placebo effect then randomised by computer generated</p>		
			<p>Mean change in Quasi-IPSS ± SD from baseline at 1 year</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: -1.5 ± 5.4*</p> <p>Grp 2: -1.0 ± 5.3*</p> <p>P value: <0.05</p>			
			<p>Mean change in Quasi-IPSS ± SD from baseline at 2 year</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: -1.7 ± 6.7*</p> <p>Grp 2: -0.5 ± 6.3*</p> <p>P value: <0.01</p>			
			<p>Mean change in Qmax ± SD from baseline at 4 months</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: 0.7 ± 3.8*</p> <p>Grp 2: 0.65 ± 6.2*</p> <p>P value: NS</p>			
			<p>Mean change in Qmax ± SD from baseline at 1 year</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: 0.95 ± 6.0*</p> <p>Grp 2: 0.3 ± 4.2*</p> <p>P value: <0.05</p>			
			<p>Mean change in Qmax ± SD from baseline at 2 years</p> <p>Number of patients remaining is unclear so use ITT figures</p>	<p>Grp 1: 1.25 ± 4.3*</p> <p>Grp 2: 0.25 ± 4.9*</p> <p>P value: <0.01</p>			
			<p>Mean change in % prostate volume from baseline at 1 year</p>	<p>Grp 1: -19</p> <p>Grp 2: +7</p> <p>P value: ≤0.01</p>			
			<p>Mean change in % prostate volume from baseline at 2 year</p>	<p>Grp 1: -21</p> <p>Grp 2: +9</p> <p>P value: ≤0.01</p>			
			<p>Median % change in PSA from baseline at 24 months</p>	<p>Grp 1: -52%</p> <p>Grp 2: 6%</p> <p>P value: < 0.0001</p>			
			<p>Reason for withdrawal §</p> <p style="text-align: right;">N</p> <p style="text-align: right;">Adverse Events</p>	<table border="1"> <thead> <tr> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>64</td> <td>77</td> </tr> <tr> <td>28</td> <td>40</td> </tr> </tbody> </table>		Grp 1	Grp 2
Grp 1	Grp 2						
64	77						
28	40						

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients N: 613 Mean age: NR Drop outs: 141 (23%)</p> <p>Group 1 (Finasteride 5mg/day) N: 310 Mean (range) Age: 63 (46-79) Total symptom score: 15.8 ± 7.6 Total obstructive score: 10.2 ± 4.8 Qmax ± SD, mL/s: 11.1 ± 3.7 Prostate volume ± SD, mL: 44.1 ± 23.5 Dropouts: 64/310 (20.6%) see withdrawals§</p> <p>Group 2 (Placebo 1/day) N: 303 Mean (range) Age: 63.5 (47-80) Total symptom score: 16.6 ± 7.2 Total obstructive score: 10.7 ± 4.5 Qmax ± SD, mL/s: 10.9 ± 3.5 Prostate volume ± SD, mL: 45.8 ± 22.4 Dropouts: 77/303 (25.4%) see withdrawals§</p>		<p>Insufficient response 16 Lost to follow up 5 Protocol violation 6 Other 9</p> <p>Other adverse events Urinary retention or surgery 19 Non-drug related mortality 5</p> <p>Adverse events related to sexual function N 31 Decreased libido 49 Impotence 24 Ejaculation disorder</p>	<p>16 19 5 9 6 3 9 6</p> <p>Grp 1 Grp 2 19 31 p=0.08 5 3</p> <p>Grp 1 Grp 2 104 43 31 19 49 19 p <0.01 24 5 p <0.01</p>	<p>sequence. Allocation preserved using sealed opaque envelopes. Analysis was ITT</p> <p>*Standard deviations for changes from baseline calculated using confidence intervals and Cochrane methodology</p> <p>Analysis of variance used to compare outcomes with treatment centre and treatment group as model parameters.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Polat et al., 1997 ²³⁹ Setting: single centre, Turkey Study design: RCT Evidence level: 1+ Duration of follow-up: 12 months	<p>Patient group: men with BPH Inclusion criteria:</p> <ul style="list-style-type: none"> • 50-80 years • In good health • Prostate volume >30 ml • Qmax <15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Prostate cancer or suspect <p>All patients N: 123 Mean age: NR</p> <p>Group 1 (Finasteride 5mg/day) N: 62 Mean (range) Age: 61 (45-80) AUA symptom score: 15.1 ± NR Qmax ± SD, mL/s: 9.9 ± NR Prostate volume ± SD, mL: 39.1 ± NR PVR ± SD, mL: 96.2 ± NR Serum PSA ± SD, ng/mL: 2.2 ± NR Dropouts: 23/62 (37%)</p> <p>Group 2 (Placebo 1/day) N: 61 Mean (range) Age: 59 (44-80) AUA symptom score: 15.3 ± NR Qmax ± SD, mL/s: 10.1 ± NR Prostate volume ± SD, mL: 38.2 ± NR PVR ± SD, mL: 100.0 ± NR Serum PSA ± SD, ng/mL: 2.32 ± NR Dropouts: 0</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Prostate volume (TRUS), AUA symptom score, Qmax, serum PSA, PVR and adverse events were recorded at 3, 6, 9 and 12 months</p>	<p>Mean AUA score ± SD at 3 months</p> <p>Mean AUA score ± SD at 6 months</p> <p>Mean AUA score ± SD at 12 months</p> <p>Mean Qmax ± SD at 3 months</p> <p>Mean Qmax ± SD at 6 months</p> <p>Mean Qmax ± SD at 12 months</p> <p>Mean PSA (ng/dl) at 3 months</p> <p>Mean PSA (ng/dl) at 6 months</p> <p>Mean PSA (ng/dl) at 12 months</p> <p>Prostate volume (cm³) at 3 months</p> <p>Prostate volume (cm³) at 6 months</p>	<p>Grp 1: 11.6 ± 5.3* Grp 2: 14.1 ± 5.3* P value: <0.01</p> <p>Grp 1: 10.9 ± 6.4* Grp 2: 13.9 ± 6.4* P value: <0.01</p> <p>Grp 1: 10.5 ± 9.0* Grp 2: 13.7 ± 9.0* P value: <0.05</p> <p>Grp 1: 10.5 ± NR Grp 2: 10.3 ± NR P value: NS</p> <p>Grp 1: 10.6 ± NR Grp 2: 10.4 ± NR P value: NS</p> <p>Grp 1: 13.2 ± 4.6* Grp 2: 10.4 ± 4.6* P value: <0.001</p> <p>Grp 1: 1.6 ± NR Grp 2: 2.3 ± NR P value: ≤0.01</p> <p>Grp 1: 1.4 ± NR Grp 2: 2.3 ± NR P value: ≤0.001</p> <p>Grp 1: 1.2 ± NR Grp 2: 2.3 ± NR P value: ≤0.001</p> <p>Grp 1: 32.4 ± NR Grp 2: 38.1 ± NR P value: ≤0.01</p> <p>Grp 1: 31.1 ± NR Grp 2: 38.0 ± NR P value: ≤0.01</p>	<p>Funding: Merck Frost Canada, inc.</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method, allocation concealment and blinding not reported. • High dropout rate in Finasteride arm • Reasons for withdrawal not explained. <p>Additional outcomes: % reduction in PSA</p> <p>Notes: * Standard deviations for changes from baseline calculated using p values for intergroup comparison following the Cochrane methodology</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments				
			Prostate volume (cm ³) at 12 months	Grp 1: 30.0 ± NR Grp 2: 38.0 ± NR P value: ≤0.01					
			Adverse events Impotence	<table border="0"> <tr> <td>Grp 1</td> <td>Grp 2</td> </tr> <tr> <td>1/62</td> <td>0/61</td> </tr> </table>	Grp 1	Grp 2	1/62	0/61	
Grp 1	Grp 2								
1/62	0/61								

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																								
<p>Roehrborn et al., 2002²⁵⁷</p> <p>A priori design for pooled analysis of parallel studies ARIA 3001, 3002, 3003 with identical inclusion/exclusion criteria.</p> <p>Study also reported in O'Leary et al., 2003²²⁹ and O'Leary et al., 2008²³⁰</p> <p>Setting: multi-centre, 400 sites in 19 countries</p> <p>Study design: RCT double blind. Patients and investigators masked.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Men with a clinical diagnosis of BPH (according to medical history, DRE and physical examination)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ≥ 50 years Prostate volume (TRUS) ≥ 30 mL AUA-7 ≥ 12 Qmax ≤ 15 mL/s on 2 consecutive voids of ≥125 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> PVR > 250 mL History of prostate cancer Previous prostate or bladder surgery Previous AUR within 3 months of screening Serum PSA <1.5 ng/mL or >10 ng/mL Concurrent use of alpha-blockers or anti-androgens <p>All patients N: 4325 Mean age: NR Drop outs: 1374/4325 (32%)</p> <p>Group 1 (Dutasteride 0.5mg/day) N: 2167 White: 91% Mean (± SD) Age: 66.5 ± 7.6 AUA Symptom score ± SD: 17.0 ± 6.0</p>	<p>Group 1: Dutasteride 0.5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: AUA score and Qmax were evaluated at baseline and months 1, 3, 6 and every 6 months thereafter. Total prostate volume by TRUS was measured at baseline and months 6, 12, 24 and additionally in month 1 for ARIA 3001 and in month 3 for ARIA 3002. PSA analysis was completed at baseline and months 1, 3, 6, 12, 18 and 24.</p> <p>O'Leary et al., 2008²³⁰ reports quality of life measures.</p> <p>Symptom Problem Index SPI - 7 questions about frequency and urgency with a scale of 0-28 where 0= no problem and 4=big problem. SPI is similar to AUA.</p>	<p>Mean change ± SD in AUA score from baseline at 2 years (ITT analysis)</p> <p>Mean change in Qmax ± SD from baseline at 2 years (ITT analysis)</p> <p>Mean change in total prostate volume ± SD from baseline at 2 years (ITT analysis)</p> <p>Mean change in Serum PSA ± SD from baseline at 2 years (ITT analysis)</p> <p>Mean change SPI ± SD from baseline at 2 years (ITT analysis)</p> <p>Mean change BSIA ± SD from baseline at 2 years (ITT analysis)</p> <p>Mean change BPWB ± SD from baseline at 2 years (ITT analysis)</p> <p>Reason for withdrawal *</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Total discontinuations</td> <td>657</td> <td>717</td> </tr> <tr> <td>Adverse Events</td> <td>193</td> <td>192</td> </tr> <tr> <td>Lack of improvement</td> <td>134</td> <td>212</td> </tr> <tr> <td>Protocol violation</td> <td>43</td> <td>50</td> </tr> <tr> <td>Consent withdrawn</td> <td>129</td> <td>135</td> </tr> <tr> <td>Loss to follow up</td> <td>67</td> <td>52</td> </tr> <tr> <td>Other/missing</td> <td>91</td> <td>76</td> </tr> </tbody> </table> <p>Spontaneous or precipitated AUR Acute urinary retention defined as spontaneous (no precipitating factors) or precipitated (stroke, UTI, pre surgery etc)</p>		Grp 1	Grp 2	Total discontinuations	657	717	Adverse Events	193	192	Lack of improvement	134	212	Protocol violation	43	50	Consent withdrawn	129	135	Loss to follow up	67	52	Other/missing	91	76	<p>Grp 1: -4.5 ± 6.6 (n=2167) Grp 2: -2.3 ± 6.8 (n=2158) P value: <0.001</p> <p>Grp 1: 2.2 ± 5.2 (n=2167) Grp 2: 0.6 ± 4.7 (n=2158) P value: <0.001</p> <p>Grp 1: -14.6 ± 13.5 (n=2167) Grp 2: 0.8 ± 14.3 (n=2158) P value: <0.001</p> <p>Grp 1: -3.1 ± 2.0 (n=2167) Grp 2: 0.5 ± 2.1 (n=2158) P value: <0.001</p> <p>Grp 1: -2.2 ± 5.8 (n=2167) Grp 2: -0.8 ± 5.8 (n=2158) P value: <0.001</p> <p>Grp 1: -1.7 ± 5.5 (n=2167) Grp 2: -1.5 ± 6.0 (n=2158) P value: <0.001</p> <p>Grp 1: -1.5 ± 3.9 (n=2167) Grp 2: -0.6 ± 4.0 (n=2158) P value: <0.001</p> <p>Grp 1: 42/1503 Grp 2: 99/1513 P value: NR</p>	<p>Funding: GSK of dutasteride</p> <p>Limitations:</p> <p>Additional outcomes: Serum DHT and transition zone volume. BSLA – BPH Specific lifestyle adaptations. (19 questions)</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in prior to randomisation by computer generated block sequence. Author confirms allocation concealment was preserved.</p> <p>Paper reports that a linear model was used to compare baseline and follow up data</p>
	Grp 1	Grp 2																											
Total discontinuations	657	717																											
Adverse Events	193	192																											
Lack of improvement	134	212																											
Protocol violation	43	50																											
Consent withdrawn	129	135																											
Loss to follow up	67	52																											
Other/missing	91	76																											

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																		
	<p>Qmax ± SD, mL/s: 10.1 ± 3.5 Prostate volume, mL: 54.9 ± 23.9 Serum PSA ± SD, ng/L: 4.0 ± 2.1 SPI (QoL): 11.7 ± 6.1 BSIA (QoL): 8.7 ± 6.2 BPWB (QoL): 11.0 ± 4.2 Dropouts: 657/2167 (30%) see*</p> <p>Group 2 (Placebo 1/day) N: 2158 White: 92% Mean (± SD) Age: 66.1 ± 7.4 AUA Symptom score ± SD: 17.1 ± 6.1 Qmax ± SD, mL/s: 10.4 ± 3.6 Prostate volume, mL: 54.0 ± 21.9 Serum PSA ± SD, ng/L: 4.0 ± 2.1 SPI (QoL): 11.8 ± 6.1 BSIA (QoL): 8.9 ± 6.2 BPWB (QoL): 11.0 ± 4.3 Dropouts: 717/2158 (33%) see *</p>	<p>BPH-specific interference with activities BSIA – 7 questions about how often urinary problems interfered with everyday activities with a scale of 0-28 where 0= none of the time and 4=all of the time. BPH-Specific Psychological Well Being (BPWB) – 6 questions about how often urinary condition has affected mental health with a scale of 5-25 where 1=not at all and 5=almost always</p>	<p>Drug related adverse events over 2 years</p> <p>N</p> <p>Decreased libido</p> <p>Impotence</p> <p>Ejaculation disorder</p> <p>Gynaecomastia</p>	<table border="1"> <thead> <tr> <th>Grp 1</th> <th>Grp 2</th> <th></th> </tr> </thead> <tbody> <tr> <td>2167</td> <td>2158</td> <td></td> </tr> <tr> <td>91</td> <td>46</td> <td>p <0.001</td> </tr> <tr> <td>158</td> <td>86</td> <td>p <0.001</td> </tr> <tr> <td>48</td> <td>17</td> <td>p <0.001</td> </tr> <tr> <td>50</td> <td>16</td> <td>p <0.001</td> </tr> </tbody> </table>	Grp 1	Grp 2		2167	2158		91	46	p <0.001	158	86	p <0.001	48	17	p <0.001	50	16	p <0.001	<p>for continuous variables with baseline values, treatment, protocol and investigator cluster as model parameters.</p>
Grp 1	Grp 2																						
2167	2158																						
91	46	p <0.001																					
158	86	p <0.001																					
48	17	p <0.001																					
50	16	p <0.001																					

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																										
<p>Tenover et al., 1997²⁹³</p> <p>Setting: multi-centre, 97 centres in the USA recruitment from April 1993 to October 1994.</p> <p>Study design: RCT double blind. Patients and investigators masked.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: men seeking treatment for symptomatic BPH from a primary care physician.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ≥ 45 years Moderate to severe AUA Enlarged prostate gland on DRE PSA ≤ 10 ng/mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Urethral stricture History of repeated catheterisations Previous pelvic radiotherapy Recurrent urinary retention Previous prostate or bladder surgery Chronic prostatitis Neurogenic bladder Recurrent UTI Concurrent use of alpha-blockers or anti-androgens Prostate cancer suspects unless biopsy ruled out cancer <p>All patients N: 2315 (2112 in efficacy analysis and baseline characteristics) Mean age: NR Drop outs:</p> <p>Group 1 (Finasteride 5mg/day) N: 1589 Mean (± SD) Age: 63.6 ± 8.7</p>	<p>Group 1: Finasteride 5 mg 1/day</p> <p>Group 2: Placebo 1/day</p> <p>Examination methods: Physical examination including DRE was performed at baseline and 12 mths. Serum dihydrotestosterone measured at baseline and mths 6 & 12</p> <p>AUA-7 Symptom score, BPH Impact Index (BII) used for HRQoL, Patient satisfaction with urinary condition as extra question (0-6) and additional questions from modified BSIA instrument to measure interference with activities and extra question about adjustment of activities to cope with urinary symptoms were taken at baseline and 3 mth intervals.</p> <p>Patient and investigator global assessment of change in urologic status also rated from 1 (much worse) to 7 (much better) every 3 mths.</p> <p>Patients with visual impairment had questionnaires read to them and Spanish versions</p>	<p>Adjusted mean change in AUA score* from baseline at 12 months</p> <p>Adjusted mean change in BII score** from baseline at 12 months</p> <p>Adjusted mean change in general adjustment question** from baseline at 12 months</p> <p>Adjusted mean change in BSIA score** from baseline at 12 months</p> <p>Reason for withdrawal \$</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Total discontinuations</td> <td>288</td> <td>95</td> </tr> <tr> <td>Adverse Events (all)</td> <td>118</td> <td>36</td> </tr> <tr> <td>Lack of improvement</td> <td>43</td> <td>14</td> </tr> <tr> <td>Protocol violation or patient request</td> <td>54</td> <td>20</td> </tr> <tr> <td>Loss to follow up</td> <td>73</td> <td>25</td> </tr> </tbody> </table> <p>Acute urinary retention</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Grp 1:</td> <td>34/1736</td> <td></td> </tr> <tr> <td>Grp 2:</td> <td>23/579</td> <td></td> </tr> <tr> <td>P value:</td> <td colspan="2">0.644</td> </tr> </tbody> </table> <p>Drug related adverse events (possibly, probably or definitely drug related)</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> <th></th> </tr> </thead> <tbody> <tr> <td>N Randomised</td> <td>54</td> <td>13</td> <td>p =0.243</td> </tr> <tr> <td>Withdrawals due to drug related AE</td> <td>85</td> <td>17</td> <td>p =0.038</td> </tr> <tr> <td>Decreased libido</td> <td>128</td> <td>19</td> <td>p <0.001</td> </tr> <tr> <td>Impotence</td> <td>57</td> <td>5</td> <td>p =0.001</td> </tr> <tr> <td>Ejaculation disorder</td> <td>38</td> <td>8</td> <td>p =0.213</td> </tr> <tr> <td>Withdrawal due to sexual AE</td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Grp 1	Grp 2	Total discontinuations	288	95	Adverse Events (all)	118	36	Lack of improvement	43	14	Protocol violation or patient request	54	20	Loss to follow up	73	25		Grp 1	Grp 2	Grp 1:	34/1736		Grp 2:	23/579		P value:	0.644			Grp 1	Grp 2		N Randomised	54	13	p =0.243	Withdrawals due to drug related AE	85	17	p =0.038	Decreased libido	128	19	p <0.001	Impotence	57	5	p =0.001	Ejaculation disorder	38	8	p =0.213	Withdrawal due to sexual AE				<p>Grp 1: -4.96 ± NR Grp 2: -3.71 ± NR P value: <0.01</p> <p>Grp 1: -1.12 CI95% -1.32 to -0.92 Grp 2: -0.70 CI95% -1.00 to -0.40 P value: 0.007</p> <p>Grp 1: -0.26 CI95% -0.35 to -0.17 Grp 2: -0.10 CI95% -0.23 to 0.03 P value: 0.019</p> <p>Grp 1: -2.65 CI95% -3.25 to -2.06 Grp 2: -2.21 CI95% -3.09 to -1.32 P value: 0.343</p>	<p>Funding: Merck & Co., Inc</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment was not clear <p>Additional outcomes: Changes in lipid profiles from baseline</p> <p>Notes: Eligible patients entered 1 month single blind placebo run-in prior to randomisation in a 3:1 ratio</p> <p>* Mean AUA symptom score was adjusted for treatment, centre and baseline age.</p> <p>** Mean BII score, general adjustment question, BSIA, Patient global assessment and investigator global assessment were</p>
	Grp 1	Grp 2																																																													
Total discontinuations	288	95																																																													
Adverse Events (all)	118	36																																																													
Lack of improvement	43	14																																																													
Protocol violation or patient request	54	20																																																													
Loss to follow up	73	25																																																													
	Grp 1	Grp 2																																																													
Grp 1:	34/1736																																																														
Grp 2:	23/579																																																														
P value:	0.644																																																														
	Grp 1	Grp 2																																																													
N Randomised	54	13	p =0.243																																																												
Withdrawals due to drug related AE	85	17	p =0.038																																																												
Decreased libido	128	19	p <0.001																																																												
Impotence	57	5	p =0.001																																																												
Ejaculation disorder	38	8	p =0.213																																																												
Withdrawal due to sexual AE																																																															

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>White/other: 1473 Black: 76 Hispanic: 40 AUA symptom score* ± SD: 19.03 ± NR BII**: 4.76 CI95% 4.61-4.9 General adjustment question**: 1.29 CI95% 1.21-1.36 BSIA**: 12.7 CI95% 12.16-13.24 Dropouts: 288/1736 (16.65) for reasons see§</p> <p>Group 2 (Placebo 1/day) N: 523 Mean (± SD) Age: 62.7 ± 8.9 White/other: 482 Black: 28 Hispanic: 13 AUA symptom score* ± SD: 18.35 ± NR BII**: 4.67 CI95% 4.45-4.9 General adjustment question**: 1.21 CI95% 1.09-1.33 BSIA**: 12.75 CI95% 11.93-13.57 Dropouts: 95/579 (16.4%) for reasons see§</p>	provided.			<p>adjusted for treatment, centre, baseline AUA and age covariates.</p> <p>A graph was presented in the study with adjusted AUA score at follow up but it was not clear if the mean was with a standard deviation or CI95%</p>

Evidence Table 14: Anticholinergics vs. placebo

See Evidence Table 9: Alpha-blockers vs. placebo
for Kaplan et al., 2006¹²³.

Evidence Table 15: Phosphodiesterase-5 inhibitors vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>McVary et al., 2007b¹⁹⁷</p> <p>Study design: Randomised controlled trial</p> <p>Setting: US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: Men 45 years and older with a history of LUTS secondary to BPH of 6 months or longer were recruited from 21 centres in US from November 2004 to July 2005. Patients agreed not to use other BPH medications during this study.</p> <p>Inclusion criteria: IPSS of 13 or greater and a Qmax of 4-15ml/s on a voided volume of 125ml or greater was required.</p> <p>Exclusion criteria: patients without treatment compliance during run in phase (<70%) were excluded. Men with PSA >10ng/ml, recent finasteride or dutasteride treatment, history of radical prostatectomy or other pelvic surgery; neurological condition affecting bladder function; recent lower urinary tract instrumentation, urinary retention or bladder stones; history of urethral obstruction due to strictures, valves, sclerosis or tumour; detrusor-sphincter dyssynergia; urinary tract inflammation or infection; intravesical obstruction secondary to the prostate median lobe; prostate cancer; PVR 200ml or greater; certain cardiovascular diseases, clinically significant renal or hepatic insufficiency; recent history of stroke or spinal cord injury; current treatment with nitrates, cancer chemotherapy,</p>	<p>Run-in period: Eligible patients entered 4 week single blind run in period with placebo dosed once daily.</p> <p>Group 1: PHOSPHODIESTERASE 5 INHIBITORS Tadalafil 5mg once daily for six weeks, followed by dose escalation to 20mg for remaining 6 weeks. Medication ingested at same time every day.</p> <p>Group 2: PLACEBO</p>	<p>Mean (SE) IPSS at 6 weeks</p>	<p>Baseline Group1 (n=138): 17.4 Group 2 (n=143): 18.5</p> <p>6 weeks Group1 (n=135): 14.5 Group 2 (n=136): 17.0</p> <p>Change from baseline: Group 1: -2.8 (0.5) Group 2: -1.2 (0.5); p=0.003</p> <p>Difference between change from baseline: 1.7 (95% CI: 0.5-2.9); p=0.003</p>	<p>Funding: NR</p> <p>Limitations: Randomisation method and allocation concealment unclear.</p> <p>Additional outcomes: Comparisons from before placebo run-in to endpoint were reported. BII reported and IPSS results for obstructive and irritative domains reported separately. Voided volume and average urinary flow were also reported.</p> <p>Notes: * All reports of erection increased were from 1 study site, reported in response to specific questioning by the investigator and described as secondary to sexual stimulation.</p> <p>Least square means calculations used for analysis. NCGC calculated SD for meta-analysis from</p>
			<p>Mean (SE) IPSS at 12 weeks</p>	<p>Baseline Group1 (n=138): 17.5 Group 2 (n=143): 18.3</p> <p>12 weeks Group1 (n=136): 13.3 Group 2 (n=138): 16.1</p> <p>Change: Group 1: -3.8 (0.5) Group 2: -1.7 (0.5); p<0.001</p> <p>Difference between change from baseline: 2.1 (95% CI: 0.9-3.3); p<0.001</p>	
			<p>Responders (defined as patients with an IPSS change from baseline or 3 points or greater)</p>	<p>6 weeks: Group 1: 49.3% Group 2: 36.4%; p=0.03</p> <p>12 weeks: Group 1: 60.9% Group 2: 42.7%; p<0.01</p>	
			<p>Mean (SE) IPSS quality of life question at 6 weeks</p>	<p>Baseline Group1 (n=138): 3.6 Group 2 (n=143): 3.8</p> <p>6 weeks Group1 (n=136): 3.1 Group 2 (n=138): 3.5</p> <p>Change from baseline: Group1: -0.5 (0.1) Group 2: -0.2 (0.1); p=0.017</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>antiandrogens or a potent cytochrome P450 3A4 inhibitor; or uncontrolled diabetes.</p> <p>All patients N: 281</p> <p>Group 1 N: 138 Ethnicity/race: Black 10.9%, white 79%, Hispanic 6.5%, other 3.6% Mean (range) Age: 62 (45.1-82.4) Dropouts: 13 (adverse events=5, lost to follow up=1, patient decision=2, other =5)</p> <p>Group 2 N: 143 Mean (range) Age: 61 (45.0-82.3) Ethnicity/race: Black 8.4%, white 83.2%, Hispanic 7%, other 1.4% Dropouts: 17 (adverse events=2, lack of efficacy=1, lost to follow up=5, patient decision=6, other=3)</p>		<p>Mean (SE) IPSS quality of life question at 12 weeks</p>	<p>Baseline Group1 (n=136): 3.6 Group 2 (n=138) : 3.8 12 weeks Group1 (n=136): 2.8 Group 2 (n=138): 3.3 Change from baseline: Group1: -0.7 (0.1) Group 2: -0.3 (0.1); p=0.004</p>	Cochrane calculations.
		<p>% of yes responses to question: Has the treatment you have been taking since your last visit improved your urinary symptoms?</p>	<p>6 weeks Group 1 (n=136): 55.9 Group 2 (n=138): 32.6; p<0.001 12 weeks Group 1 (n=136): 57.4 Group 2 (n=138): 37.7; p<0.001</p>		
		<p>Mean (SE) Qmax, ml/sec at 6 weeks</p>	<p>Baseline Group1 (n=110): 11.7 Group 2 (n=111) : 11.2 12 weeks Group1 (n=110): 12.2 Group 2 (n=111): 11.8 Change from baseline: Group1: 1.1 (0.6) Group 2: 1.0 (0.6); p=0.46</p>		
			<p>Mean (SE) Qmax, ml/sec at 12 weeks</p>	<p>Baseline Group1 (n=116): 11.8 Group 2 (n=121) : 11.1 12 weeks Group1 (n=116): 12.3 Group 2 (n=121): 12.1 Change from baseline: Group1: 0.5 (0.5) Group 2: 0.9 (0.5); p=0.72</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean (SE) PVR, ml at 6 weeks	Baseline Group1 (n=132): 58.0 Group 2 (n=135) : 58.5 12 weeks Group1 (n=132): 57.2 Group 2 (n=136): 53.8 Change from baseline: Group1: 3.6 (7.0) Group 2: 0.1 (6.7); p=0.66	
			Mean (SE) PVR, ml at 12 weeks	Baseline Group1 (n=132): 58.0 Group 2 (n=135) : 58.2 12 weeks Group1 (n=132): 57.9 Group 2 (n=136): 54.2 Change from baseline: Group1: 1.4 (6.5) Group 2: -2.6 (6.2); p=0.69	
			Mean (SE) IPSS change from baseline in men that were sexually active	6 weeks Group 1 (n=80): -3.2±0.7 Group 2 (n=76): -0.7±0.7; p=0.001 12 weeks Group 1 (n=80): -4.4± 0.7 Group 2 (n=76): -1.8± 0.7; p=0.001	
			Mean (SE) IIEF EF domain change from baseline in men that were sexually active	6 weeks Group 1(n=80): 6.0±0.9 Group 2(n=76): 0.6±0.9; p<0.001 12 weeks Group 1(n=80): 7.7± 0.9 Group 2 (n=76): 1.4± 1.0; p<0.001	
			Discontinuation due to treatment emergent adverse events	Group 1: 3.6% Group 2: 1.4%	
			Treatment emergent adverse events with a frequency of 2% or greater at 12 weeks	Erection increased* Group 1: 7 (5.1%) Group 2: 2 (1.4%) Dyspepsia	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 6 (4.3%) Group 2: 0 Back pain Group 1: 5 (3.6%) Group 2: 2 (1.4%) Headache Group 1: 4 (2.9%) Group 2: 1 (0.7%) Nasopharyngitis Group 1: 3 (2.2%) Group 2: 0 Upper respiratory tract infection Group 1: 3 (2.2%) Group 2: 1 (0.7%) Serious adverse events: Group 1: 0 Group 2: 1 (0.7%) AUR: Group 1: 0 Group 2: 0	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>McVary et al., 2007^{c196}</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: USA</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: men with erectile dysfunction and LUTS/BPH from 41 urology clinics and clinical research centres.</p> <p>Inclusion criteria: Men ≥45 years, had a clinical diagnosis of ED (score ≤25 on the erectile function domain of the International Index of Erectile Function) and IPSS ≥12.</p> <p>Exclusion criteria: Men with confirmed or suspected prostate malignancy, serum prostate-specific antigen >10ng/ml, previous invasive intervention for BPH, or previous prostate or bladder/pelvic rations or surgery. Those with PSA between 4-10ng/ml required two additional forms of documentation to confirm the absence of clinically evident malignancy. Men with acute urinary tract disease or cystoscopy with in 4 weeks of the trial, calculi in the urinary tract or acute urinary retention within 6 months of the trial, recurrent urinary tract infections or catheterisation for outflow obstruction in the year before the trial, or other known or suspected causes of urinary symptoms other than BPH, hypotension, hypertension orthostatic hypotension or significant cardiovascular disease. Men were excluded if used nitrates, had hepatic or renal dysfunction, poorly controlled</p>	<p>Group 1: Phosphodiesterase 5 inhibitors Sildenafil citrate: 50mg once daily with each night at bedtime or 30 minutes to 1hr before sexual activity. After 2 weeks the dose increased to 100mg but could be decreased to 50mg if the higher dose was not tolerated.</p> <p>Group 2: Placebo</p>	<p>Mean (SD) IIEF – erectile function domain (1-30; higher scores indicate better treatment outcome)</p> <p>Least mean change in IPSS score</p> <p>Least mean change in Qmax, ml</p> <p>Least mean change in IPSS quality of life score</p> <p>LS mean (SE) EDITS score (end of treatment satisfaction score; 0-100)</p> <p>Number (%) of patients reporting adverse events</p> <p>Number (%) of treatment related adverse events</p> <p>Headache</p> <p>Flushing</p> <p>Dyspepsia</p> <p>Rhinitis</p>	<p>Baseline Group 1: 13.4 Group 2: 13.2</p> <p>Change from baseline Group 1: 9.2 (1.0) Group 2: 1.9 (1.0) Mean change: 9.17, 95% CI: 7.25-11.09 vs. 1.86, 95% CI: -0.03, 3.74; p<0.0001</p> <p>Group 1 (n=182): -6.3 (-8.1, -4.6) Group 2 (n=178): -1.9 (-3.7, -0.2) P<0.001</p> <p>Group 1: 0.31 (-1.6, 2.2) Group 2: 0.16 (-1.7, 2.1) P=0.8</p> <p>Group 1: -0.97 (-1.32, -0.62) Group 2: -0.29 (-0.64, 0.05) P<0.001</p> <p>Group 1: 71.2±3.2 Group 2: 41.7±3.2; p<0.0001</p> <p>Group 1: 100/189 (53%) Group 2: 78/180 (43%)</p> <p>Group 1: 86/189 (%) Group 2: 25/180 (%)</p> <p>Group 1: 21/189 (11%) Group 2: 6/180 (3%)</p> <p>Group 1: 9/189 (5%) Group 2: 1/180 (1%)</p> <p>Group 1: 12/189 (6%) Group 2: 2/180 (1%)</p> <p>Group 1: 8/189 (4%) Group 2: 3/180 (2%)</p>	<p>Funding: Supported by Pfizer, Inc.</p> <p>Limitations: Actual figures and SD not provided for IPSS, Qmax and IPSS QoL question.</p> <p>Additional outcomes: BPHII score, SEAR questionnaire (self-esteem and relationship questionnaire)</p> <p>Notes: 8 week open label extension study after this 12 week study.</p> <p>Least square means calculations used for analysis. NCGC calculated SD for meta-analysis from Cochrane calculations.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>diabetes or a history of retinitis pigmentosa. Use of antimuscarinics, 5-alpha-reductase inhibitors within 6 months or alpha blockers within 4 weeks during study. PDE5 inhibitor or any other treatment for ED must have terminated therapy 4 weeks or more before the study.</p> <p>All patients N: 370 Mean age: 60 (9) Drop outs: 1 not treated/withdrew</p> <p>Group 1 N: 187 Mean (±SD) ED: 5.7 (4.6) years Ethnicity/race: White: 84%; Black: 10% Discontinuations:21</p> <p>Group 2 N: 179 Mean (±SD) ED: 5.6 (5.1) years Ethnicity/race: white: 80%; black: 13% Discontinuations: 25</p>		<p>Discontinuations due to adverse events</p> <p>Serious adverse events</p> <p>Discontinuations due to serious adverse events</p>	<p>Group 1: 9/189 (5%) Group 2: 2/180 (1%)</p> <p>Group 1: 2/189 (1%) Group 2: 3/180 (2%)</p> <p>Group 1: 1/189 (1%) Group 2: 0</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Roehrborn et al., 2008b ²⁶¹	Patient group: Men with a history of LUTS secondary to BPH of 6 months longer.	Group 1: PDE5I Tadalafil 2.5mg once daily	Least squares mean (SE) IPSS change from baseline	Group1 (n=208): -3.88 (0.50) Group 2 (n=212): -4.87 (0.49) Group 3 (n=216): -5.17 (0.49) Group 4 (n=208): -5.21 (0.50) Group 5 (n=210): -2.27 (0.49) P<0.001 (tad v placebo)	Funding: Eli Lilly and Co. Limitations: method of randomisation and allocation concealment unclear. Additional outcomes: BPH-II score Notes: None.
Study design: RCT	Inclusion criteria:	Group2: PDE5I Tadalafil 5 mg once daily			
Setting: 92 centres in 10 countries	<ul style="list-style-type: none"> At least 45 years old IPSS of 13 or greater Qmax of 4-15ml/s from pre-void bladder volume between 150-550ml with a voided volume of 125ml or greater. 	Group 3: PDE5I Tadalafil 10 mg once daily	Least squares mean (SE) IPSS quality of life change from baseline	Group1 (n=208): -0.74 (0.11) Group 2 (n=212): -0.86 (0.11) Group 3 (n=216): -0.92 (0.10) Group 4 (n=208): -0.88 (0.11) Group 5 (n=210): -0.49 (0.11) P<0.01 (tad v placebo)	
Evidence level: 1+	Exclusion criteria:	Group 4: PDE5I Tadalafil 20 mg once daily	Least squares mean (SE) Qmax change from baseline	Group1 (n=208): 1.41 (0.39) Group 2 (n=212): 1.64 (0.39) Group 3 (n=216): 1.58 (0.38) Group 4 (n=208): 1.96 (0.39) Group 5 (n=210): 1.24 (0.40) P=Not sig. (tad v placebo)	
Duration of follow-up: 12 weeks	<ul style="list-style-type: none"> PSA > 10ng/ml PVR volume was 300ml or greater at screening visit 1 Patients reporting use of other BPH or ED treatments underwent a 4 week treatment free screening/ washout period. Penile or pelvic surgery, radiotherapy, lower urinary tract malignancy, trauma or recent instrumentation, urinary retention or bladder stones, History of urethral obstruction Neurological condition Detrusor sphincter dyssynergia, intravesical obstruction secondary to the prostate median lobe, Urinary tract inflammation or infection Prostate cancer. Renal or hepatic insufficiency, Cardiovascular conditions, history of stroke or spinal cord 	Group 5: Placebo once daily	% Yes LUTS GAQ end point (GAC question: Has the treatment you have been taking since your last visit improved your urinary symptoms)	Group1 (n=208): 61.9 Group 2 (n=212): 69.2 Group 3 (n=216): 73.0 Group 4 (n=208): 74.2 Group 5 (n=210): 54.8 P<0.05 (tad v placebo)	
			Least squares mean (SE) sexually active ED IIEF-EF change from baseline (55% of patients)	Group1 (n=208): 5.59 (1.01) Group 2 (n=212): 6.97 (1.01) Group 3 (n=216): 7.98 (1.0) Group 4 (n=208): 8.34 (1.01) Group 5 (n=210): 2.20 (1.03) P<0.001 (tad v placebo)	
			Treatment emergent adverse events	Headache Group1: 5/209 Group 2: 6/212 Group 3: 11/216 Group 4: 7/209 Group 5: 6/211	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>injury, cancer chemotherapy, uncontrolled diabetes</p> <p>All patients N: 1058</p> <p>Group 1 N: 209 Mean Age: 62.03 Ethnicity/race: White 88.46%, Hispanic 9.62%, black 1.44%, other 0.48% Mean % ED history: 64.9% Dropouts: 27</p> <p>Group 2 N: 212 Mean Age: 61.95 Ethnicity/race: White 84.43%, Hispanic 11.79%, black 3.30%, other 0.47% Mean % ED history: 67.92% Dropouts: 30</p> <p>Group 3 N: 216 Mean Age: 62.22 Ethnicity/race: White 86.11%, Hispanic 11.11%, black 2.31%, other 0.46% Mean % ED history: 69.44% Dropouts: 41</p> <p>Group 4 N: 209 Mean Age: 62.55 Ethnicity/race: White 84.21%, Hispanic 11.96%, black 2.39%, other 1.44% Mean % ED history: 69.38% Dropouts: 47</p>			<p>Dyspepsia Group 1: 2/209 Group 2: 10/212 Group 3: 6/216 Group 4: 10/209 Group 5: 0/211</p> <p>Back Pain Group 1: 3/209 Group 2: 2/212 Group 3: 10/216 Group 4: 12/209 Group 5: 1/211</p> <p>Myalgia Group 1: 3/209 Group 2: 3/212 Group 3: 6/216 Group 4: 6/209 Group 5: 0/211</p> <p>Nasopharyngitis Group 1: 7/209 Group 2: 4/212 Group 3: 2/216 Group 4: 5/209 Group 5: 2/211</p> <p>Diarrhoea Group 1: 2/209 Group 2: 6/212 Group 3: 1/216 Group 4: 5/209 Group 5: 3/211</p> <p>Gastroesophageal reflux disease Group 1: 2/209 Group 2: 2/212 Group 3: 6/216 Group 4: 3/209 Group 5: 0/211</p> <p>Extremity pain Group 1: 3/209 Group 2: 5/212 Group 3: 2/216 Group 4: 3/209</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 5 N: 212 Mean Age: 61.75 Ethnicity/race: White 84.83%, Hispanic 13.74%, black 1.42%, other 0% Mean % ED history: 67.30% Dropouts: 27</p>		<p>Discontinuation due to adverse events</p>	<p>Group 5: 0/211 Influenza Group1: 4/209 Group 2: 4/212 Group 3: 1/216 Group 4: 2/209 Group 5: 1/211 Bronchitis Group1: 3/209 Group 2: 1/212 Group 3: 5/216 Group 4: 0/209 Group 5: 1/211 Muscle spasms Group1: 2/209 Group 2: 0/212 Group 3: 2/216 Group 4: 5/209 Group 5: 0/211 Urinary retention Group1: 0/209 Group 2: 0/212 Group 3: 0/216 Group 4: 0/209 Group 5: 1/211</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Stief et al.,2008²⁸⁷</p> <p>Study design: Randomised control trial.</p> <p>Setting: multi-centre, Germany</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 8 weeks.</p>	<p>Patient group: Men with BPH/LUTS from 16 centres in Germany from October 2005-June 2006.</p> <p>Inclusion criteria: Men aged 45-64 years with a history of BPH/LUTS for at least 6 months before commencing the study and an IPSS\geq12 at screening. Patients completed a 4 week run-in phase during which no study medications was administered.</p> <p>Exclusion criteria: contraindications to vardenafil, spinal cord injury, prostatitis, history of prostate or bladder cancer, bladder or urethra stricture, urinary retention (PVR\geq100ml), pelvic trauma or surgery, history of any malignancies, and life expectancy of less than 3 yr. concomitant use of nitrates or NO donors, androgens or anti-androgens, anticoagulants, cytochrome P-50 3A4 inhibitors, any treatment for ED or alpha1-adrenocoetpro antagonists were prohibited. Alpha blockers – if withdrawn at screening, subjects would fail o be eligible for study drug treatment, previous or current use of 5-alpha reductase inhibitors.</p> <p>All patients: N: 222</p> <p>Group 1 N: 109 Mean (\pmSD) Age: 56.5 (5.4) years Ethnicity: White 100% Dropouts: 4 (1=not received medication, 3=did not provide efficacy data) Premature discontinuation=13</p>	<p>Group 1: Phosphodiesterase 5 (PDE5) inhibitors 10mg Vardenafil twice daily</p> <p>Group 2: Placebo Matched placebo tablet twice daily (12-h dosing interval).</p>	<p>Mean IPSS symptom score*</p>	<p>Baseline Group1: 16.8 Group 2: 16.8 8 weeks Group1 (n=105): 11.0 Group 2 (n=110): 13.2 Between group difference in change from baseline: 2.3 (0.90-3.64), p=0.0013</p>	<p>Funding: This study was sponsored by Bayer Healthcare AG, Leverkusen, Germany. Bayer healthcare AG involved in the design and conduct of the study; management, analysis and interpretation of the data; and preparation, review and approval of the manuscript.</p> <p>Limitations: No SD values provided for further analysis. [NCC emailed author for this information]</p> <p>Additional outcomes: IPSS also reported by irritative and obstructive sub score.</p> <p>Notes: Serious adverse events reported included myocardial infarction, chest pain, and cardiac rehabilitation therapy (one patient) and hypertensive crisis in the intervention group. The placebo group comprised of haematochezia, a meniscus injury and knee surgery. None were considered related to study medication.</p>
			<p>Mean Qmax, ml/s*</p>	<p>Baseline Group1: 15.9 Group 2: 15.9 8 weeks Group1 (n=105): 17.5 Group 2 (n=110): 16.9 Between group difference in change from baseline: -0.6 (-2.62–1.43), p=0.5614</p>	
			<p>Mean PVR volume</p>	<p>Baseline Group1: 28.0 Group 2: 26.9 8 weeks Group1 (n=105): 27.0 Group 2 (n=110): 28.8 Between group difference in change from baseline: 1.8 (-7.39 to 10.99); p=0.6994</p>	
			<p>International Index of Erectile Function – Erectile function (IIEF-EF) score</p>	<p>Baseline Group1: 15.9 Group 2: 15.9 8 weeks Group1 (n=105): 23.4 Group 2 (n=110): 17.4 Between group difference in change from baseline: -6.0 (-7.77 to 4.16), p=0.0001</p>	
			<p>Total Urolife Qulatiy of life-9 score</p>	<p>-9.3 (95% CI: -12.79, -5.71) P<0.0001</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	ITT population=105 Group 2 N: 113 Mean (±SD) Age: 55.4 (5.7) years Ethnicity: White 98.2%; Black 0.9%; Asian 0.9%. Dropouts: 3 (3=did not provide efficacy data) Premature discontinuation=14 ITT population=110		Number (%) of adverse events (treatment-emergent adverse events affecting at least 2% of patients)	Any event: Group 1 (n=108): 32 (29.6%) Group 2 (n=113): 18 (15.9%) Headache: Group 1: 14 (13.0%) Group 2: 2 (1.8%) Dyspepsia: Group 1: 8 (7.4%) Group 2: 0 Flushing: Group 1: 7 (6.5%) Group 2: 1 (0.9%) Diarrhoea: Group 1: 5 (4.6%) Group 2: 1 (0.9%) Gastrointestinal reflux disease: Group 1: 3 (2.8%) Group 2: 0 Back pain: Group 1: 3 (2.8%) Group 2: 0 Serious adverse events Group 1: 2 Group 2: 3	* Least square means analysis reported for outcomes. NCGC calculated estimated SD for mean change in IPSS/Qmax from Cochrane handbook formula.

Evidence Table 16: Diuretics vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Reynard et al., 1998a ²⁴⁹ Study design: Randomised controlled trial Setting: Hospital, UK Evidence level: 1+ Duration of follow-up: 4 weeks.	Patient group: elderly men presenting with lower urinary tract symptoms and completed 7day frequency volume chart. Inclusion criteria: aged over 50 years, with nocturnal polyuria (defined as night time diuresis defined as the production of >33% of the 24-h urine volume between midnight and 8am). Exclusion criteria: serum creatinine >150umol.L, previous lower urinary tract surgery, symptomatic heart failure, taking medication active on the lower urinary tract including those taking any diuretic, concomitant neurological disease which could potentially affect lower urinary tract function, and clinical evidence of prostate cancer or diabetes mellitus. All patients N: 49 Number obstructed: 19/41 Drop outs: 6 (withdrew) Group 1 N: 21 Mean (±SD) Age: 70 Dropouts: 3 (evening frequency). Group 2 N: 22 Mean (±SD) Age: 69 Dropouts: 3=(lack of efficacy or evening frequency)	Two week placebo period. In second week a frequency volume chart was completed with the IPSS symptom score. Group 1: Diuretic Frusemide 40mg Afternoon dose taken 6 hours before their usual bedtime. Group 2: Placebo	Reduction in night time frequency	Group 1: -0.5 Group 2: 0 P=0.014	Funding: NR. Limitations: Method of randomisation, allocation concealment not reported. Actual figures not reported. Additional outcomes: No significant correlation between the % night time voided volume and changes in night time frequency, night time voided volume or % voided volume. Figures not reported. Notes: Day time defined as 08.00 and 23.59h and night time as between 00.00 and 07.59h.
			Increase in daytime frequency	Group 1: +1.9 Group 2: -0.1 P<0.001	
			Correlation for % night time voided volume at entry to the study against change in night-time voiding frequency	Spearman's correlation coefficient: 0.25 P=0.3	
			Increase in daytime voided volume, mL	Group 1: +365 Group 2: -31 P=0.002	
			Night time voided volume, mL	Group 1: -120 Group 2: +9 P=0.065	
			Reduction in night-time voiding frequency of one or more	Group 1: 7/19 Group 2: 1/20 P=0.02	
			Night time voiding frequency was reduced 2 or more	4/19 0/20	
			Correlation between % night time voided volume at entry and reduction in night time voided volume	Spearman's correlation coefficient: 0.03 P=0.9	
			Total urine output (24h), mL	Group 1: 1663 Group 2: 1780 P=0.2	
			% change of night time voided volume	Group 1: -18% Group 2: 0% P=0.001	
Correlation between % night time voided volume and change in % night time voided volume	Spearman's correlation coefficient = 0.43, p=0.08				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Change in IPSS	Group 1: +1 Group 2: 0 P=0.9	
			Patients reported that intervention 'helped'	Group 1: 14/21 Group 2: 5/22 P<0.001	

Evidence Table 17: Desmopressin vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Cannon et al., 1999⁴¹ Study design: RCT-cross over trial</p> <p>Setting: UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Two-2 week periods</p>	<p>Patient group: Men with nocturia</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Men >50 years Nocturnal polyuria confirmed after 48 hours of inpatient monitoring or a 1-week FV chart, which showed in excess of a third of their 24-hour urine volume being produced overnight <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Nocturnal enuresis or incontinence Significant cardiovascular, renal or hepatic disease, diabetes, UTI or concomitant medication active on the lower urinary tract <p>All patients N: 20 Mean age, mean (range): 70.5(52-80) years Drop outs: 2</p>	<p>Group 1: Desmopressin 20 microgram nasal spray, administered just before going to bed each evening</p> <p>Group 2: Placebo nasal spray, administered just before going to bed each evening</p>	<p>24-h volume, (ml) mean, se: (measured using FV-chart*)</p>	<p>Baseline: 1646.6 se107.6 Group 1: 1567.4 se 96.7 Group 2: 1713.5 se 119.4 P value (paired t-test): Not sig</p>	<p>Funding: Ferring Pharmaceuticals</p> <p>Limitations:</p> <ul style="list-style-type: none"> Cross over study Small sample size Method of randomisation allocation and concealment was not described. <p>Additional outcomes: Adverse events: For 20 microgram of desmopressin: dry throat plus cough (1), increased sputum (1), and fluid retention plus hyponatraemia (1). For placebo: headache (1), flu like illness (1). Another 2 patients had fluid retention symptoms while receiving the 40microgram dose.</p> <p>Notes: This is a cross over study. Patient had 1 week run in with placebo, and then allocated to desmopressin 20 microgram or placebo for 2 weeks, before crossing over for another 2 weeks. *FV chart resulted were collected at the second week. ** The 24 hour urine collection was done on the last day of the treatment period.</p>
			<p>Nocturnal frequency mean, se: (measured using FV-chart*)</p>	<p>Baseline: 3.0 se 0.3 Group 1: 2.7 se 0.33 Group 2: 3.1 se 0.3 P value (paired t-test): Not sig</p>	
			<p>Nocturnal volume (ml)mean, se: (measured using FV-chart*)</p>	<p>Baseline: 749.6 se 67.5 Group 1: 633.9 se 60.8 Group 2: 809.1 se 78.7 P value (paired t-test): <0.01</p>	
			<p>Nocturnal percentage (%) (measured using FV-chart*)</p>	<p>Baseline: 45.7 se 3.1 Group 1: 40.5 se 3.1 Group 2: 46.9 se 3.3 P value (paired t-test): <0.05</p>	
			<p>24-h volume, (ml) mean, se: (24 hour urine collection**)</p>	<p>Baseline: 1487.2 se110.5 Group 1: 1419 se 121.20 Group 2: 1400.6 se 88.5 P value (paired t-test):</p>	
			<p>Nocturnal volume (ml)mean, se: (24 hour urine collection**)</p>	<p>Baseline: 718.3 se 79.1 Group 1: 562.0 se 73.5 Group 2: 726.7 se74 P value (paired t-test): <0.01</p>	
			<p>Nocturnal percentage (%) (24 hour urine collection**)</p>	<p>Baseline: 47.3 se 3.5 Group 1: 39.2 se 3.5 Group 2: 50.6 se 3.5 P value (paired t-test): <0.001</p>	
			<p>Hyponatremia and hyposmolaemia (withdrawn early from study, sodium 127mmol/L, hypoosmolaemia 263mosmol/kg)</p>	<p>Group 1: 1/20 Group 2: 0/20</p>	

Evidence Table 18: Non steroidal anti-inflammatory drugs (NSAIDs) vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments												
<p>Falahatkar et al., 2008⁸⁸</p> <p>Study design: RCT, double blinded</p> <p>Setting: Iran, Jan to May 2007</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 month</p>	<p>Patient group: BPH patients with refractory nocturia</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> BPH with ≥ 2 voids per night Mean night time voided volume of $< 30\%$ of the 24 hour volume IPSS ≥ 8 Prostate volume $> 20\text{cm}^3$ Prescribed alpha-blockers or alpha blockers or finasteride (if prostate volume $> 30\text{cm}^3$) for 2-3 months but incidence of nocturia remained ≥ 2 times per night Negative urine culture findings Normal renal function <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous prostate surgery or other invasive procedures for testing of BPH Prostate cancer, or PSA $> 10\text{ng/ml}$. Men with PSA 4.1 to 10ng/ml were required to provide ultrasound guided biopsy <p>All patients N: 80 Mean age: range 49 to 80years Drop outs: 0</p> <p>Group 1 - Celecoxib N: 40 Mean (\pmSD) Age: 64.3 ± 7.7 (49-80) Dropouts: 0 IPSS, mean \pmsd: 18.2 ± 3.4 Qmax, ml/s, mean \pmsd: 12.5 ± 2.1</p>	<p>Group 1: COX II selective NSAID (celecoxib) 100mg capsule at 9PM</p> <p>Group 2: Placebo</p>	<p>IPSS</p>	<p>At 1 month Group 1: 15.5 ± 4.2 Group 2: 18.0 ± 3.9 P values:</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation allocation and concealment not reported Small sample size Short length of follow up <p>Additional outcomes: Authors reported that not baseline parameters did not influence level of response</p> <p>Notes: None</p>												
			<p>Qmax, ml/s, mean \pmsd</p>	<p>At 1 month Group 1: 12.9 ± 2.7 Group 2: 12.3 ± 2.5 P value:</p>													
			<p>Nocturia frequency</p>	<p>At 1 month Group 1: 2.5 ± 1.9 Group 2: 5.1 ± 1.9 P value:</p>													
			<p>Nocturia frequency, classified as excellent if decreased ≥ 2 voids/night or disappeared, improved if decreased by 1 void/night and no change.</p>	<p>At 1 month</p> <table border="1"> <thead> <tr> <th>change</th> <th>Excellent</th> <th>improved</th> <th>no</th> </tr> </thead> <tbody> <tr> <td>Group 1: 28(70) 7(17.5)</td> <td></td> <td>5(12.5)</td> <td></td> </tr> <tr> <td>Group 2: 3(7.5) 31(77.5)</td> <td></td> <td>6(15)</td> <td></td> </tr> </tbody> </table> <p>Values in brackets are percentages</p>		change	Excellent	improved	no	Group 1: 28(70) 7(17.5)		5(12.5)		Group 2: 3(7.5) 31(77.5)		6(15)	
			change	Excellent		improved	no										
Group 1: 28(70) 7(17.5)		5(12.5)															
Group 2: 3(7.5) 31(77.5)		6(15)															
<p>Adverse events – mild gastric discomfort</p>	<p>At 1 month Group 1: 4/40 Group 2: 0/40 P value: 0.11 [calculated by NCGC using Fisher's exact test]</p>																

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Nocturia frequency, mean±sd: 5.17±2.1</p> <p>Prostate volume, ml, mean±sd:18.25 ±4.5</p> <p>PSA level, ng/ml, mean±sd:2.62±1.16</p> <p>Group 2 - Placebo N: 40 Mean (±SD) Age: 64.9±7.05 (50-80) Dropouts:0</p> <p>IPSS, mean ±sd: 18.4±3.1</p> <p>Qmax, ml/s, mean±sd:12.1±2.1</p> <p>Nocturia frequency, mean±sd:5.30±2.4</p> <p>Prostate volume, ml, mean±sd:50.11 ±5.6</p> <p>PSA level, ng/ml, mean±sd: 2.68±1.18</p>				

Evidence Table 19: Combination therapy: 5-Alpha reductase inhibitor and alpha-blocker

See Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors
for Debruyne et al., 1998⁶⁹.

See Evidence Table 9: Alpha-blockers vs. placebo
for Kirby et al., 2003¹⁴⁷.

See Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors
for Lepor et al., 1996¹⁶⁴.

See Evidence Table 2: How does PSA predict symptom progression (in terms of symptom score)?
for McConnell et al., 2003¹⁷⁰.

See Evidence Table 10: Alpha blocker vs. 5-alpha reductase inhibitors
for Roehborn et al., 2008²⁶³

Evidence Table 20: Combination therapy: Anticholinergic added to alpha-blocker

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Macdiarmid et al., 2008¹⁷⁹</p> <p>Study design: RCT, double blinded, multicentre March 2004 to June 2005</p> <p>Setting: Double blinded RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks post randomisation. All patients received 4 weeks of tamsulosin between screening and randomisation</p>	<p>Patient group: Men with LUTS who remained symptomatic despite 4 weeks of alpha blocker therapy</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Age ≥ 45 years Diagnosed with LUTS, had urgency and frequency, with or without urge incontinence Qmax of 4ml/s with voided volumes of 125mL and post void residual volume of ≤ 150mL on at least 2 occasions <p>After receiving ≥4 weeks of 0.4mg tamsulosin, they should still have:</p> <ul style="list-style-type: none"> IPSS ≥13 and IPSS storage component (Question 2, 4 and 7) ≥8. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> History of urinary retention, bladder or prostate cancer PSA ≥4 ng/ml Angle closure glaucoma Surgical or procedural treatment of the prostate <p>Amendments in protocol in July 2004</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> Qmax of 8 ml/s with voided volumes of 125mL and post void residual volume of ≤ 150mL on at least 2 occasions <p>Discontinuation criteria:</p> <ul style="list-style-type: none"> Qmax decreased to 5mL/s or less Post void residual volume >300mL 	<p>Group 1: Oxybutynin ER + 0.4 mg tamsulosin Oxybutynin ER dose was 10mg/day, the recommended starting dose</p> <p>Group 2: 0.4mg Tamsulosin + placebo</p> <p>Note: All patients received 4 weeks of 0.4mg tamsulosin before randomisation</p>	<p>IPSS, mean±sd at various time points and change from baseline</p> <p>P values provided in paper based on ANCOVA using baseline values as the covariates</p>	<p><u>At week 4</u> <u>Change</u></p> <p>Group 1: 15.9±6.7 -4.4±5.6 Group 2: 16.6±5.8 -3.8±5.5 P value: 0.24</p> <p><u>At Week 8</u> <u>Change</u></p> <p>Group 1: 14.5±7.3 -5.7±6.3 Group 2: 16.0±6.7 -4.4±6.0 P value: 0.03</p> <p><u>At week 12</u> <u>Change</u></p> <p>Group 1: 13.3±7.4 -6.9±6.5 Group 2: 15.2±6.9 -5.2±6.2 P value: 0.006</p>	<p>Funding: Ortho Urology, US (oxybutynin manufacturer)</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation allocation and concealment not described The criteria for excluding about ½ of the screened population from randomisation not provided Characteristics at screening visit not provided This study only randomised patients who remained symptomatic despite ≥4 weeks of treatment with alpha blocker and should only be generalised to this group of patients (this is likely to augment the difference seen between the two intervention groups) <p>Additional outcomes: SPI (symptom problem index) values were also reported</p>
			<p>IPSS-QoL (maximum 6 points) at various at various time points and change from baseline</p> <p>P values provided in paper based on ANCOVA using baseline values as the covariates</p>	<p><u>Week 4</u> <u>Change</u></p> <p>Group 1: 3.2±1.3 -0.9±1.4 Group 2: 3.5±1.3 -0.5±1.3 P value: 0.006</p> <p><u>Week 8</u> <u>Change</u></p> <p>Group 1: 3.0±1.5 -1.2±1.5 Group 2: 3.4±1.4 -0.6±1.3 P value: <.001</p> <p><u>Week 12</u> <u>Change</u></p> <p>Group 1: 2.8±1.5 -1.3±1.5 Group 2: 3.2±1.5 -0.8±1.4 P value: 0.001</p>	
			<p>IPSS-Storage (maximum 15 points), mean ± sd at various time points and change from baseline</p> <p>P values provided in paper based on ANCOVA using baseline values as the covariates</p>	<p><u>At week 4</u> <u>Change</u></p> <p>Group 1: 7.7±2.9 -2.6±2.7 Group 2: 8.2±2.6 -1.9±2.6 P value: 0.008</p> <p><u>At Week 8</u> <u>Change</u></p> <p>Group 1: 7.0±3.2 -3.3±3.0 Group 2: 7.9±3.0 -2.1±2.8 P value: <.001</p> <p><u>At week 12</u> <u>Change</u></p> <p>Group 1: 6.5±3.2 -3.7±3.0 Group 2: 7.6±3.1 -2.4±2.9 P value : <.001</p>	
			<p>Qmax (ml/s), mean±sd</p> <p>P value and change values</p>	<p><u>At 12 weeks</u> <u>Change</u></p> <p>Group 1: 15.5±8.4 -0.2±7.8 Group 2: 14.7±8.4 0.1±7.6</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																																																									
	<p>All patients N: 420 randomised out of 818 screened Mean age: 62.9±9.1 Drop outs: 2 (took <1 dose of medications)</p> <p>Group 1- Oxybutynin ER + 0.4 mg tamsulosin N: 209 Age, mean ±sd: 62.6±9.0 Dropouts: Years since LUTS diagnosis, years, mean±sd:5.0±5.7 IPSS, mean±sd:20.2±5.0 IPSS-QoL, mean±sd:4.1±1.1 Qmax, ml/s, mean±sd:15.7±7.1 Post void residual volume, ml, mean±sd: 50.7±42.9</p> <p>Group 2 N: 209 Age, mean ±sd: 63.3±9.2 Dropouts: Years since LUTS diagnosis, years, mean±sd:5.0±4.7 IPSS, mean±sd:20.5±4.9 IPSS-QoL, mean±sd:4.0±1.0 Qmax, ml/s, mean±sd:14.6±6.6 Post void residual volume, ml, mean±sd: 45.8±41.4</p>		<p>calculated by NCGC</p> <p>Post void residual volume (ml), mean±sd</p> <p>P value and change values calculated by NCGC</p>	<p>P value: NS</p> <table border="1"> <thead> <tr> <th colspan="2">At 12 weeks</th> <th>Change</th> </tr> </thead> <tbody> <tr> <td>Group 1:</td> <td>69.7±75.3</td> <td>18.2±77.3</td> </tr> <tr> <td>Group 2:</td> <td>53.7±52.9</td> <td>7.8±47.5</td> </tr> </tbody> </table> <p>P value: NS</p> <table border="1"> <thead> <tr> <th></th> <th>Group 1</th> <th>Group 2</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Any adverse events</td> <td>89(42.6)</td> <td>89(42.6)</td> <td>NS</td> </tr> <tr> <td>Serious adverse events</td> <td>5(2.4)</td> <td>6(2.9)</td> <td>NS</td> </tr> <tr> <td>AEs leading to withdrawal</td> <td>21(10)</td> <td>20(9.6)</td> <td>NS</td> </tr> <tr> <td>Dry mouth</td> <td>32(15.3)</td> <td>10(4.8)</td> <td><.001</td> </tr> <tr> <td>Infections and infestations</td> <td>18(8.6)</td> <td>22(10.5)</td> <td>NS</td> </tr> <tr> <td>Renal and urinary AEs</td> <td>10(4.8)</td> <td>10(4.8)</td> <td>NS</td> </tr> <tr> <td>AUR (with or without Foley catheter)</td> <td>0(0)</td> <td>0(0)</td> <td>NS</td> </tr> <tr> <td>Nervous system disorders</td> <td>8(3.8)</td> <td>9(4.3)</td> <td>NS</td> </tr> <tr> <td>Constipation</td> <td>1(0.5)</td> <td>4(1.9)</td> <td>NS</td> </tr> </tbody> </table> <table border="1"> <thead> <tr> <th></th> <th>Group 1</th> <th>Group 2</th> <th>P value</th> </tr> </thead> <tbody> <tr> <td>Reasons for study discontinuation</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Adverse events</td> <td>21/209</td> <td>20/209</td> <td>NS</td> </tr> <tr> <td>Lack of efficacy</td> <td>4/209</td> <td>6/209</td> <td>NS</td> </tr> <tr> <td>Patient choice</td> <td>5/209</td> <td>0/209</td> <td>NS</td> </tr> <tr> <td>Others (include PVR> 300ml and Qmax <5ml/s)</td> <td>14/209</td> <td>8/209</td> <td>NS</td> </tr> </tbody> </table>	At 12 weeks		Change	Group 1:	69.7±75.3	18.2±77.3	Group 2:	53.7±52.9	7.8±47.5		Group 1	Group 2	P value	Any adverse events	89(42.6)	89(42.6)	NS	Serious adverse events	5(2.4)	6(2.9)	NS	AEs leading to withdrawal	21(10)	20(9.6)	NS	Dry mouth	32(15.3)	10(4.8)	<.001	Infections and infestations	18(8.6)	22(10.5)	NS	Renal and urinary AEs	10(4.8)	10(4.8)	NS	AUR (with or without Foley catheter)	0(0)	0(0)	NS	Nervous system disorders	8(3.8)	9(4.3)	NS	Constipation	1(0.5)	4(1.9)	NS		Group 1	Group 2	P value	Reasons for study discontinuation				Adverse events	21/209	20/209	NS	Lack of efficacy	4/209	6/209	NS	Patient choice	5/209	0/209	NS	Others (include PVR> 300ml and Qmax <5ml/s)	14/209	8/209	NS	<p>Notes: There were 6/209 vs. 1/209 patients with PVR >300ml (all withdrawn from study) in group 1 vs. group 2 respectively. There were 14/209 vs. 13/209 patients with Qmax<5 ml/s (8/209 vs. 12/209 at endpoint) respectively. The number patients discontinued as per protocol did not tally with the number of patients who had PVR>300ml</p>
At 12 weeks		Change																																																																												
Group 1:	69.7±75.3	18.2±77.3																																																																												
Group 2:	53.7±52.9	7.8±47.5																																																																												
	Group 1	Group 2	P value																																																																											
Any adverse events	89(42.6)	89(42.6)	NS																																																																											
Serious adverse events	5(2.4)	6(2.9)	NS																																																																											
AEs leading to withdrawal	21(10)	20(9.6)	NS																																																																											
Dry mouth	32(15.3)	10(4.8)	<.001																																																																											
Infections and infestations	18(8.6)	22(10.5)	NS																																																																											
Renal and urinary AEs	10(4.8)	10(4.8)	NS																																																																											
AUR (with or without Foley catheter)	0(0)	0(0)	NS																																																																											
Nervous system disorders	8(3.8)	9(4.3)	NS																																																																											
Constipation	1(0.5)	4(1.9)	NS																																																																											
	Group 1	Group 2	P value																																																																											
Reasons for study discontinuation																																																																														
Adverse events	21/209	20/209	NS																																																																											
Lack of efficacy	4/209	6/209	NS																																																																											
Patient choice	5/209	0/209	NS																																																																											
Others (include PVR> 300ml and Qmax <5ml/s)	14/209	8/209	NS																																																																											

Evidence Table 21: Combination therapy: phosphodiesterase-5-inhibitor added to alpha-blocker

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																	
<p>Bechara et al., 2008²⁷</p> <p>Study design: double blinded, cross over study</p> <p>Setting: single-centre in Argentina</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Week 12</p>	<p>Patient group: LUTS and erectile dysfunction</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> > 50 years Clinical diagnosis of LUTS by medical history and physical examination At least 6 months of LUTS; IPSS\geq12, Total PSA \leq4.0ng/ml Qmax > 5ml/s with minimum voided volume of >125ml <p>Exclusion criteria:</p> <ul style="list-style-type: none"> History or evidence of prostate cancer Previous prostate surgery or other invasive procedure to treat BPH Post void residual volume >250ml History of AUR \leq3 months of screening visit Use of alpha reductase inhibitors or phytotherapy \leq 6 months; alpha blockers or PDE5-I \leq2 weeks Cardiovascular comorbidities and uncontrolled diabetes Comorbidities which may interfere with urinary flow or symptoms. <p>All patients N: 30 out of 40 patients screened Drop outs: 3 (2 adverse events, 1 personal reasons) Age, mean (range): 63.7(51-78) Sexually active: 28/30 (93.3%) IPSS, mean (range): 19.4 (12-34) IPSS-QoL, mean (range): 4.1 (0-6) Qmax, ml/s, mean (range): 9.6 (4 to 14) **IIEF-EF mean(range):17(1-29)</p>	<p>Group 1: Tamsulosin 0.4mg/day+ tadalafil 20mg/day For 6 weeks, at about the same time each day</p> <p>Group 2: Tamsulosin 0.4mg/day +placebo For 6 weeks, at about the same time each day</p> <p>The capsules were identical and prepared by a third party (pharmacist) in numbered containers</p> <p>Cross over design: The patients were randomised to treatment Group 1 or Group 3 at Visit 1 (week 0). At week 6, end point measures were collected and patients switched over to the other treatment group. At week 12, end points were measured again.</p>	<p>IPSS change from baseline at end of 6 week treatment, mean \pmSD</p> <p>IPSS-QoL at end of 6 week treatment, mean \pmSD</p> <p>Qmax, ml/s, mean \pm SD</p> <p>IIEF-EF mean \pm SD</p> <p>Adverse Events</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Headache</td> <td>12</td> <td>0</td> </tr> <tr> <td>Hypotension</td> <td>2</td> <td>1</td> </tr> <tr> <td>Dizziness</td> <td>0</td> <td>1</td> </tr> <tr> <td>Dyspepsia</td> <td>3</td> <td>1</td> </tr> <tr> <td>Diarrhoea</td> <td>0</td> <td>1</td> </tr> <tr> <td>Ejaculation disorder</td> <td>0</td> <td>1</td> </tr> <tr> <td>Altered vision</td> <td>0</td> <td>1</td> </tr> </tbody> </table> <p>Withdrawals due to adverse events</p> <table border="1"> <thead> <tr> <th></th> <th>Grp 1</th> <th>Grp 2</th> </tr> </thead> <tbody> <tr> <td>Headache</td> <td>1/30</td> <td>0/30</td> </tr> <tr> <td>Rashes</td> <td>0/30</td> <td>1/30</td> </tr> </tbody> </table>		Grp 1	Grp 2	Headache	12	0	Hypotension	2	1	Dizziness	0	1	Dyspepsia	3	1	Diarrhoea	0	1	Ejaculation disorder	0	1	Altered vision	0	1		Grp 1	Grp 2	Headache	1/30	0/30	Rashes	0/30	1/30	<p>Grp 1: -9.2\pm5.08 Grp 2: -6.7\pm3.87 *P value: <0.05</p> <p>Grp 1: 1.6, no SD Grp 2: 2.3, no SD *P value: <0.05</p> <p>Grp 1: 12.6, no sd Grp 2: 11.7, no sd *P value: >0.05</p> <p>Grp 1: 23.2, no sd Grp 2: 16.9, no sd *P value:<0.001</p> <p>Grp 1 Grp 2</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> This is a cross-over RCT. There was no washout period to provide verification that patients had returned to their baseline level. The sample size is small <p>Additional outcomes: IIEF-EF, GAQ (Global Assessment Quality) and a visual analogue scale (no mention of validations)</p> <p>Notes: *P values were as reported in paper. Authors reported using Tukey Cramer test with multiple comparisons **IIEF-EF>25 points was reported as 28/30(93.3%) at baseline in Table1. These numbers did not tally with mean IIEF (sexual function domain) of 15 points at baseline (Table3) and number of men with ED who completed study (19/27).</p> <p>Erectile Function domain of the 15-question IIEF (Q1-5 and Q15, maximum score 30) was used. This is different from IIEF-5, which consists of Q2, Q4, Q5, Q7 and Q15 of the IIEF (maximum score 25)</p>
	Grp 1	Grp 2																																				
Headache	12	0																																				
Hypotension	2	1																																				
Dizziness	0	1																																				
Dyspepsia	3	1																																				
Diarrhoea	0	1																																				
Ejaculation disorder	0	1																																				
Altered vision	0	1																																				
	Grp 1	Grp 2																																				
Headache	1/30	0/30																																				
Rashes	0/30	1/30																																				

See Evidence Table 12: Alpha-blockers vs. phosphodiesterase-5 inhibitors
for Kaplan et al., 2007¹³²

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Liguori et al., 2009¹⁶⁷</p> <p>Study design: RCT open label,</p> <p>Setting: Multicentre (5) in Italy from Feb to Dec2007</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: Men with LUTS and previously untreated erectile dysfunction</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Men aged 50 to 75 years with previously untreated ED and a history of LUTS secondary to BPH for 6 months or longer IPSS>8 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Contraindications to the study drugs Using medications to control bladder symptoms or had ever taken alpha blockers, PDE5-I, or 5 alpha reductase inhibitors. Bladder tumours, urethral strictures, neurogenic bladder dysfunction History of prostatitis, prostate cancer; prostate surgery, radiotherapy PSA level>20 ng/ml Acute urinary retention or indwelling catheter Infection on urinalysis <p>All patients N: 66 Mean age: 61 years (range 50 to 75) Drop outs: 8/66 (Baseline data excluded patients who dropped out of study)</p> <p>Group 1 (Tadalafil) N: 21 Dropouts: 2 /21 Mean (± SD) Age: 60.8±8 IPSS mean± SD: 13.8±5.6 IIEF-EF, mean ±sd: 14.1 IIEF Q15 mean± SD: 2.5 Qmax mean± SD, mL/s: 13.1</p>	<p>Group 1: Tadalafil 20 mg every other day</p> <p>Group 2: Alfuzosin 10 mg/day</p> <p>Group 3: tadalafil 20 mg every other day + alfuzosin 10 mg/day</p>	<p>IPSS</p> <p>Note: The change from baseline values were calculated by NCGC</p>	<p><u>Baseline:</u> Grp 1: 13.8±5.6 Grp 2: 15.7±4.8 Grp 3: 15.3±4.5 <u>At 12 weeks</u> Grp 1: 12.5±5.6 Grp 2: 10.6±3.6 Grp 3: 9.0±4.0 <u>Change from baseline</u> Grp 1: -1.3±5.6 Grp 2: -5.2±4.2 Grp 3: -6.3±4.3</p>	<p>Funding: Reported no conflicts of interest</p> <p>Limitations:</p> <ul style="list-style-type: none"> This was an open label study with no randomisation allocation and concealment methods reported. The outcomes are mainly subjective outcomes, and this makes it particularly at risk of biases. <p>Additional outcomes: Changes in IPSS (obstructive), IPSS (irritative) IIEF-EF, and IIEF Q15 were also reported</p> <p>Notes: **Erectile Dysfunction assessed using the Erectile Function domain score of the 15-question IIEF, ie, ie Q1-5 and Q15 (Maximum score 30).</p> <p>This is different from IIEF-5, which consists of question Q2, Q4, Q5, Q7 and Q15 of the IIEF (maximum score 25).</p>
			<p>IPSS % change from baseline at 12 weeks The P values reported were for 12 weeks compared to baseline</p>	<p>Grp 1: -8.4, p=NS Grp 2: -27.2, p=0.003 Grp 3: -41.6, p<0.001</p>	
			<p>IPSS-QoL</p>	<p><u>Baseline:</u> Grp 1: 3.5±1.1 Grp 2: 3.4±0.9 Grp 3: 3.2±1 <u>At 12 weeks</u> Grp 1: 2.5±1.2 Grp 2: 2.1±0.9 Grp 3: 1.6±0.8 <u>Change from baseline</u> Grp 1: 1±1.2 Grp 2: 1.3±0.9 Grp 3: 1.6±0.9</p>	
			<p>Qmax, ml/s mean ±sd</p>	<p><u>Baseline:</u> Grp 1: 13.1±4.3 Grp 2: 12.3±5.4 Grp 3: 11.9±2.7 <u>At 12 weeks</u> Grp 1: 14.3±5.2 Grp 2: 14.0±3.7 Grp 3: 15.0±4.0</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments						
	<p>Group 2 (Alfuzosin) N: 22 Dropouts: 4/22 Mean (± SD) Age: 61.3±6.8 IPSS mean± SD:15.7±4.8 IIEF-EF, mean ±sd:14.2 IIEF Q15 mean± SD: 2.8 Qmax mean± SD, mL/s:12.3</p> <p>Group 3 (Tadalafil + Alfuzosin) N: 23 Dropouts: 2/23 Mean (± SD) Age: 63 ± 6.9 IPSS mean± SD:15.3±4.5 IIEF-EF, mean ±SD: 14.6 IIEF Q15 mean± SD: 2.4 Qmax mean± SD, mL/s:11.9</p>		<p>Nocturia (as recorded in voiding diary)</p> <p>Withdrawals due to AE</p> <p>The reason for withdrawals were</p>	<p><u>Change from baseline</u> Grp 1: 1.2±4.8 Grp 2: 1.7±4.6 Grp 3: 3.1±3.4</p> <p><u>Baseline:</u> Grp 1: 1.7±1 Grp 2: 1.9±0.9 Grp 3: 1.9±0.9 <u>At 12 weeks</u> Grp 1: 1.1±1.1 Grp 2: 1.0±0.7 Grp 3: 1.1±0.9 <u>Change from baseline</u> Grp 1: -0.6±1.1 Grp 2: -0.9±0.8 Grp 3: -0.8±0.9</p> <table border="0"> <tr> <td>Grp 1</td> <td>Grp 2</td> <td>Grp 3</td> </tr> <tr> <td>1/21</td> <td>3/22</td> <td>2/23</td> </tr> </table> <p>Group 1: back pain, head aches Group 2 :dizziness, constipations Group 3: myalgia, dizziness, sensation of heaviness</p>	Grp 1	Grp 2	Grp 3	1/21	3/22	2/23	
Grp 1	Grp 2	Grp 3									
1/21	3/22	2/23									

Evidence Table 22: Holmium laser enucleation (or resection) of the prostate HoLEP (HoLRP) vs. transurethral resection of the prostate

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ahyai et al., 2007¹¹</p> <p>Study design: RCT</p> <p>Setting: Urology department, Berlin</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 36 months</p>	<p>Patient group: Patients with lower urinary tract symptoms due to BPH.</p> <p>Inclusion criteria: AUA of 12 or more, Qmax of 12ml/s or less, PVR volume > 50ml, Schafer grade of II or more in pressure flow studies, and a total prostate volume <100cc in transrectal ultrasound.</p> <p>Exclusion criteria: previous prostate or uerthral surgery and voiding disorders not related to benign prostatic hyperplasia. Prostate carcinoma excluded by biopsy.</p> <p>All patients N: 200</p> <p>Group 1 N: 100 Mean Age: 68.0 Dropouts: 25 (prostate cancer=3, stricture=4, refused follow-up=6, bladder neck contracture=3, moved away=3, polymorbidity=2, death=3, BPH recurrence=1)</p> <p>Group 2 N: 100 Mean Age: 68.7 Dropouts: 31 (prostate cancer=10, stricture=3, refused follow-up=4, bladder neck contracture=3, moved away=1, polymorbidity=5, death=3, transition cell carcinoma=2)</p>	<p>Group 1: HoLEP 40-50 Hz, 80-100W used. Saline used as irrigation fluid and electrolyte-free solution for electrocautery loop tissue fragmentation. Postoperative bladder irrigation used as necessary until haematuria had settled sufficiently to remove catheter. Median postoperative catheterisation=1 day Median Hospital stay=2 days</p> <p>Group 2: TURP standard tungsten wire loop with a cutting current of 160 W and coagulating current of 80 W. Postoperative bladder irrigation used as necessary until haematuria had settled sufficiently to remove catheter. Median postoperative catheterisation=2 day Median Hospital stay=3 days</p>	<p>Mean (SD) AUA</p>	<p>Baseline: Group 1 (n=100): 22.1 (3.8) Group 2 (n=100): 21.4 (5.2); p=0.56</p> <p>6 months: Group 1 (n=94): 2.2 (1.6) Group 2 (n=89): 3.7 (3.4); p=0.006</p> <p>12 months: Group 1 (n=89): 1.7 (1.8) Group 2 (n=86): 3.9 (3.9); p<0.001</p> <p>18 months: Group 1 (n=82): 1.3 (1.5) Group 2 (n=78): 4.0 (3.8); p<0.0001</p> <p>24 months: Group 1 (n=80): 1.7 (1.7) Group 2 (n=75): 3.9 (3.7); p<0.0001</p> <p>36 months: Group 1 (n=75): 2.7 (3.2) Group 2 (n=69): 3.3 (3.0); p=0.17</p>	<p>Funding: Financial interest and/or other relationship with Lumenis, Inc and Karl Storz, Inc.</p> <p>Limitations: Allocation concealment and blinding unclear.</p> <p>Notes: Linked to Kuntz 2004¹⁵¹ – follow up for 24 months.</p>
			<p>Mean (SD) Qmax, ml/s</p>	<p>Baseline: Group 1: 4.9 (3.8) Group 2: 5.9 (3.9); p=0.08</p> <p>6 months: Group 1: 25.1 (6.9) Group 2: 25.1 (9.4); p=0.72</p> <p>12 months: Group 1: 27.9 (9.9) Group 2: 27.7 (12.2); p=0.76</p> <p>18 months: Group 1: 27.5 (9.2) Group 2: 28.2 (11.2); p=0.89</p> <p>24 months: Group 1: 28.0 (9.0) Group 2: 29.1 (10.9); p=0.82</p> <p>36 months: Group 1: 29.0 (11.0) Group 2: 27.5 (9.9); p=0.41</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean (SD) PVR, ml	Baseline: Group 1: 237 (163) Group 2: 216 (177); p=0.08 6 months: Group 1: 4.8 (12.5) Group 2: 16.7 (16.9); p=0.03 12 months: Group 1: 5.3 (15.3) Group 2: 26.6 (60.4); p<0.001 18 months: Group 1: 1.6 (11.5) Group 2: 16.3 (28.4); p<0.0001 24 months: Group 1: 5.6 (19.9) Group 2: 19.9 (29.6); p<0.0001 36 months: Group 1: 8.4 (16.0) Group 2: 20.2 (33.0); p<0.012	
			Peri-operative complications	Blood transfusion Group 1: 0 Group 2: 2 (2%) Recatheterisation Group 1: 0 Group 2: 5 (5%) Mortality Group 1: 0 Group 2: 0	
			Complications at 36 months	Urethral stricture Group 1: 4 (4.1%) Group 2: 3 (3.3%) Bladder neck contracture Group 1: 3 (3.1%) Group 2: 3 (3.3%) BPH recurrence: Group 1: 1 (1.0%) Group 2: 0 Reoperation: Group 1: 7.2% Group 2: 6.6%	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Urinary incontinence at 12 months	Preoperatively: Group 1: 27/89 Group 2: 33/86 Post operatively: Group 1: 5/89 Group 2: 5/86	
			Stress incontinence developed after surgery	Group 1: 1 Group 2: 1	
			Potency following preoperative erectile dysfunction (insufficient for sexual intercourse)	Group 1: 2/43 Group 2: 0/41	
			Resolved erectile dysfunction postoperatively	Group 1: 1 Group 2: 1	
			Decreased potency at 12 months compared to preoperative level	Group 1: 10/89 (11.2%) Group 2: 9/86 (10.5%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gupta et al., 2006¹⁰⁸</p> <p>Study design: RCT</p> <p>Setting: India</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: Patients with BPH who were candidates for TURP were selected from July 2002 to December 2003.</p> <p>Inclusion criteria: glands of >40g</p> <p>Exclusion criteria: patients with a previous history of prostatic and urethral surgery, neurovesical dysfunction and carcinoma of the prostate were excluded from the study.</p> <p>All patients N: 150</p> <p>Group 1 N: 50 Mean (\pmSD) Age: 65.88 (10.1) Dropouts: NR</p> <p>Group 2 N: 50 Mean (\pmSD) Age: 65.67 (7.5) Dropouts: NR</p> <p>Group 3 N: 50 Mean (\pmSD) Age: 67.68 (9.8) Dropouts: NR</p>	<p>Group 1: HoLEP Power settings were 80-100W. Operative duration: 75.4 minutes</p> <p>Group 2: TURP 80W cutting and 50W coagulation used. Operative duration: 64.1 minutes</p> <p>Group 3: TUVRP 180W cutting and 80W coagulation used. Operative duration: 55.9 minutes</p>	<p>Mean (SD) IPSS:</p>	<p>Baseline: Group 1: 23.4 (4.5) Group 2: 23.3 (3.9) Group 3: 24.9 (3.9) 6 months: Group 1: 5.2 (0.31) Group 2: 6.1 (0.42) Group 3: 5.9(0.25) 12 months: Group 1: 5.2 (0.17) Group 2: 5.6 (0.32) Group 3: 5.4 (0.28)</p>	<p>Funding: NR</p> <p>Limitations: No mention of drop outs in the study.</p> <p>Additional outcomes: Irrigation, haemoglobin decrease, serum sodium decrease.</p> <p>Notes: None.</p>
			<p>Mean (SD) Qmax</p>	<p>Baseline: Group 1: 5.15 (4.4) Group 2: 4.5(3.9) Group 3: 4.65 (3.6) 6 months: Group 1: 23.1(1.2) Group 2:20.7 (1.32) Group 3: 22.5 (0.95) 12 months: Group 1: 25.1 (1.06) Group 2: 23.7 (1.58) Group 3: 23.6(0.96)</p>	
			<p>Mean (SD) PVR, mL</p>	<p>Baseline: Group 1: 112.0(155.9) Group 2: 84.0(129.7) Group 3: 103 (174.1) 6 months: Group 1: <20 Group 2: <20 Group 3: <20 12 months: Group 1: <20 Group 2: <20 Group 3: <20</p>	
			<p>Mean (SD) blood loss, mL</p>	<p>Group 1: 40.6 (37.3) Group 2: 140.5 (60.7) Group 3: 68.6 (42.7)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean (SD) catheter duration, hours	Group 1: 28.6 (20.5) Group 2: 45.7 (12.7) Group 3: 36.2 (8.3)	
			Mean (SD) nursing contact time, minutes	Group 1: 28.1 (8.4) Group 2: 48.3 (9.2) Group 3: 37.2 (6.7)	
			Number (%) complications	Re-catheterisation: Group 1: 2 (4) Group 2: 3 (6) Group 3: 3 (6) Fever: Group 1: 1 (2) Group 2: 1 (2) Group 3: 2 (4) Hyponatraemia: Group 1: 0 Group 2: 1 (2) Group 3: 1 (2) Blood transfusion: Group 1: 0 Group 2: 1 (2) Group 3: 0 Capsular perforation: Group 1: 1 (2) Group 2: 0 Group 3: 0 Bladder mucosal injury: Group 1: 2 (4) Group 2: 0 Group 3: 0 Death (pneumonia): Group 1: 0 Group 2: 0 Group 3: 1 (2) Transient dysuria: Group 1: 5 (10) Group 2: 1 (2) Group 3: 9 (18) Stricture: Group 1: 1 (2)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 2 (4) Group 3: 1 (2) Incontinence: Group 1: 1 (2) Group 2: 1 (2) Group 3: 0	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Mavuduru RM 2009 ¹⁸⁷ Study design: RCT Evidence level: 1+ Setting: Chandigarh, India Duration of follow-up: 9 months	Patient group: Patients who underwent surgery for BPH. Inclusion criteria: Exclusion criteria: Patients with a history of previous prostatic or urethral surgery, and documented cases of prostate carcinoma. All patients N: 30 Group 1: TURP N: 15 Age (mean): 66.46±5.79 Drop outs: 0 Group 2: HoLEP N: 15 Age (mean): 69.86±9.6 Drop outs: 0	Group 1: Transurethral resection of the prostate (TURP). TURP was performed by standard technique using a 26-Fr continuous flow resectoscope (Karl Storz) with a cutting current of 100-120 D and coagulating current of 50-60 W. The intraoperative irrigation fluid used was 1.5% glycine, the TURP chips were removed by Ellick's evacuator. Group 2: Holmium laser enucleation of the prostate (HoLEP) Instrumentation included 550nm end-firing flexible quartz, and a continuous flow resectoscope consisting of a 27-Fr outer sheath, an inner rotating sheath with a self-designed working element. HoLEP was performed by standard technique as described by Gillig et al. The machine used was Versapulse Holmium Laser, with a frequency of 35-40 Hz and a power setting of 2 joules. The irrigant used was normal saline.	Mean ±SD symptom score- IPSS	Baseline: Group 1: 21.4±3.7 Group 2: 22.53±4.79 3 months: Group 1: 2.86±1.72 Group 2: 2.26±1.57 p value: 0.329 9 months: Group 1: 3.57±1.03 Group 2: 4.32±1.25 p value: 0.37	Funding: NR Limitations: Small study size and duration of follow up is less than 1 year. Additional outcomes: Intraoperative data including weight of gland resected and volume of irrigation fluid.
			Mean ± SD PVR volume (ml)	Baseline: Group 1: 103 ±27 Group 2: 91±30 3 months: Group 1: 13.66±14.0 Group 2: 13±8.61 p value: 0.87 9 months: Group 1: 35.66±15.0 Group 2: 43±10.61 p value: 0.97	
			Mean ± SD Uroflowmetry	Baseline: Group 1: 6.9 ±2.5 Group 2: 5.79±2.7 3 months: Group 1: 27.8±6.5 Group 2: 28.6±6.2 p value: 0.721 9 months: Group 1: 27.8±6.5 Group 2: 28.6±6.2 p value: 0.64	
			Operative time (minutes)	Group 1: 43±9.36 Group 2: 53±9.84 p value: <0.01	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Duration of catheterization (hours)	Group 1: 78.20±17.84 Group 2: 46.42±14.25 p value: <0.001	
			Adverse events	Transient dysuria Group 1: 3/15 (40%) Group 2: 1/15 (6.66%) Recatheterization Group 1: 1/15 (6.66%) Group 2: 1/15 (6.66%) Bleeding Group 1: 2/14 (13.33%) Group 2: nil Incontinence Group 1: nil Group 2: 2/15 (13.33%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Montorsi et al., 2004²⁰²</p> <p>Study design: RCT</p> <p>Setting: 2 centre study (Milan and Bergamo)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: consecutive patients with symptomatic obstructive BPH from January to October 2002.</p> <p>Inclusion criteria: Age <75 years, peak urinary flow rate <15ml/s, post void residual urine <100cc, medical therapy failure, transrectal ultrasound adenoma volume <100gm and urodynamic obstruction.</p> <p>Exclusion criteria: Neurogenic bladder, diagnosis of prostate cancer and any previous prostatic, bladder neck or urethral surgery.</p> <p>All patients N: 100</p> <p>Group 1 N: 52 Mean Age: 65.14 Mean TRUS volume (gm): 70.3 Dropouts: NR</p> <p>Group 2 N: 48 Mean Age: 64.5 Mean TRUS volume (gm): 56.2 Dropouts: NR</p>	<p>Group 1: HoLEP Tissue morcellation of the prostatic lobes into fragments that were retrieved from the bladder cavity. Energy delivered by a 360u fibre. Enucleation performed at 2.0J and 35Hz.</p> <p>Total operative time: 74±19.5 minutes. Catheterisation time 31±13 hours Hospital stay 59±19.9 hours</p> <p>Group 2: TURP Using a standard tungsten wire loop with a cutting current of 80W and a coagulation g current of 160W. Following procedure catheter inserted into bladder and irrigation started.</p> <p>Total operative time: 57±15 minutes. Catheterisation time 57.78±17.5 hours Hospital stay 85.8±18.9 hours</p>	<p>Mean (SD) IPSS</p>	<p>Baseline: Group 1: 21.6±6.7 Group 2: 21.9±7.2 6 months: Group 1: 3.9±2.9 Group 2: 2.9±2.6 12 months: Group 1: 4.1±2.3 Group 2: 3.9±3.6 p=0.58</p>	<p>Funding: NR</p> <p>Limitations: Number of drop outs not reported. Prostate size significantly different at baseline.</p> <p>Additional outcomes: Average flow reported. Orgasmic function, sexual desire, intercourse satisfaction.</p> <p>Notes: Linked with Rigatti 2006²⁵¹</p>
			<p>Mean (SD) QoL question</p>	<p>Baseline: Group 1: 4.6±1.11 Group 2: 4.7±1.0 6 months: Group 1: 1±0.8 Group 2: 0.6±0.2 12 months: Group 1: 1.4±0.9 Group 2: 0.8±1.28 p=0.31</p>	
			<p>Mean (SD) maximum flow (ml/s)</p>	<p>Baseline: Group 1: 8.2±3.2 Group 2: 7.8±3.6 6 months: Group 1: 23.1±8.6 Group 2: 26.5±15.5 12 months: Group 1: 25.1±7.2 Group 2: 24.7±10 p=0.25</p>	
			<p>Mean detrusor pressure at max flow (cmH2O)</p>	<p>Baseline: Group 1: 77.3 Group 2: 81.8 12 months Group 1: 36.2 Group 2: 38.5 p=0.85</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean Schafer grade	Baseline: Group 1: 3.4 Group 2: 3.5 12 months Group 1: 0.9 Group 2: 1.2; p=0.55	
			Mean (SD) Erectile function (International Index of Erectile Function IIEF-15)	Preoperatively: Group 1: 22.3±3.6 Group 2: 21.4±3.1 6 months: Group 1: 23.5±3.6 Group 2: 23.4±3.5 12 months: Group 1: 23.8±3.9 Group 2: 24.1±3.7	
			Number (%) of early Adverse events	Bladder mucosal injury Group 1: 10 (18.2%) Group 2: 0 Re-intervention for bleeding Group 1: 1 (1.7%) Group 2: 1 (2.2%) Transurethral resection syndrome Group 1: 0 Group 2: 1(2.2%) Early acute urinary retention Group 1: 3 (5.3%) Group 2: 1 (2.2%) Dysuria (burning) Group 1: 33 (58.9%) Group 2: 13 (29.5%) Transitory urge incontinence Group 1: 25 (44%) Group 2: 17 (38.6%)	
			Adverse events at 6 & 12 month follow up (%)	Urethral stricture: Group 1: 1 (1.7%) Group 2: 4 (7.4%) Stress incontinence: Group 1: 1 (1.7%) Group 2: 1 (2.2%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Westenberg et al., 2004³¹⁸</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Setting: Tauranga Hospital, New Zealand.</p> <p>Duration of follow-up: 48 months</p>	<p>Patient group: Candidates for surgery for LUTS and obstruction due to BPH at Tauranga Hospital from April 1996 to August 1997.</p> <p>Inclusion criteria: Age 80 years or younger, AUA score ≥ 8, peak urinary flow rate ≤ 15ml/s, transrectal ultrasound volume of the prostate < 100ml, post void residual volume < 400ml and Schafer grade ≥ 2.</p> <p>Exclusion criteria: Catheterised patients and those who had undergone previous urethral or prostatic surgery. All patients had a digital rectal examination and SPA before enrolment to excluded men with carcinoma of the prostate.</p> <p>All patients N: 120</p> <p>Group 1 N: 61 Mean (\pmSD) Age: 66.9 ± 6.5 Dropouts at 48m: 18 (2 died cardiovascular disease, 5 required reoperation, 6 intercurrent illness, 5 lost to follow up).</p> <p>Group 2 N: 59 Mean (\pmSD) Age: 66.8 ± 7.4 Dropouts at 48m: 29 (7 died – cardiovascular or malignant disease, 8 required reoperation, 4 intercurrent diseases, 10 lost to follow up).</p>	<p>Group 1 Holmium laser resection (HoLRP). Maximum average power of 80W was used. General or spinal anaesthesia required in all cases. Postoperative bladder irrigation was only used if deemed necessary by the surgeon. Catheter removed the morning after surgery. Mean catheter time: 26.2 ± 11.71.</p> <p>Group 2 TURP using a cutting current of 160W and a coagulating current of 80W. General or spinal anaesthesia was used. Bladder irrigation was used and catheter removed before patient discharged from hospital. Mean catheter time: 47.5 ± 17.37.</p>	<p>AUA score</p> <p>Quality of Life score:</p>	<p>Baseline: Group 1 (n=61): 21.9 ± 6.2 Group 2 (n=59): 23.0 ± 5.9</p> <p>3 months: Group 1 (n=61): 5.6 ± 5.1 Group 2 (n=59): 5.7 ± 5.2</p> <p>6 months: Group 1 (n=61): 3.8 ± 3.8 Group 2 (n=59): 5.0 ± 4.5</p> <p>12 months: Group 1 (n=53): 4.2 ± 6.0 Group 2 (n=49): 4.3 ± 4.1</p> <p>18 months: Group 1: 2.9 ± 5.3 Group 2: 4.5 ± 5.3</p> <p>24 months: Group 1 (n=45): 3.4 ± 4.9 Group 2 (n=41): 3.7 ± 4.9</p> <p>48 months: Group 1 n=43: 5.2 ± 5.9 Group 2 (n=30): 6.6 ± 5.0; P=0.32</p> <p>Baseline: Group 1 (n=61): 4.5 ± 1.1 Group 2 (n=59): 4.7 ± 1.1</p> <p>3 months: Group 1 (n=61): 1.4 ± 1.5 Group 2 (n=59): 1.6 ± 1.4</p> <p>6 months: Group 1 (n=61): 1.1 ± 1.3 Group 2 (n=59): 1.5 ± 1.4</p> <p>12 months: Group 1 (n=53): 0.88 ± 1.4 Group 2 (n=49): 1.6 ± 1.5</p> <p>18 months: Group 1 (n=61): 0.72 ± 1.1 Group 2 (n=59): 1.3 ± 1.1</p> <p>24 months: Group 1 (n=45): 0.98 ± 1.3 Group 2 (n=41): 1.0 ± 1.3</p>	<p>Funding: Financial interest and/or other relationship with Lumenis, Inc.</p> <p>Limitations: Allocation concealment and blinding unclear.</p> <p>Additional outcomes: Detrusor pressure at 6 months.</p> <p>Notes: Linked to Gilling 1999¹⁰⁴, Gilling 2000¹⁰³ and Fraundorfer 2001⁹⁷</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				48 months: Group 1 (n=43): 1.1±1.1 Group 2 (n=30): 1.4±1.4; P=0.37	
			Qmax (ml/s)	Baseline: Group 1 (n=61): 8.9±3.0 Group 2 (n=59): 9.1±3.2 3 months: Group 1 (n=61): 22.8±10.0 Group 2 (n=59): 20.2±9.5 6 months: Group 1 (n=61): 23.9±8.7 Group 2 (n=59): 22.4±9.0 12 months: Group 1 (n=53): 25.2±11.9 Group 2 (n=49): 20.4±8.5 18 months: Group 1: 25.1±9.3 Group 2: 19.2±9.3 24 months: Group 1 (n=45): 25.0±11.1 Group 2 (n=41): 20.9±11.1 48 months: Group 1 (n=43): 22.3±14.2 Group 2 (n=30): 18.5±8.2; P=0.23	
			TRUS volume (cc)	Baseline: Group 1: 44.3±19.0 (11-92) Group 2: 44.6±20.7 (11.5-95) 6 months: Group 1: 29.3 (11-61) Group 2: 27.3 (10-75)	
			Post void residual (ml)	Baseline: Group 1: 87.8±88.4 (0-346) Group 2: 84.7±81.7 (0-373) 6 months: Group 1: 26.7 (0-245) Group 2: 34.3 (0-295)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Adverse events:		
			Perioperative blood transfusions:	Group 1: 0/61 Group 2: 4/59	
			Recatheterised	Group 1: 5/61 Group 2: 8/59	
			Reoperations	Group 1: 5/61 Group 2: 8/59	
			Urinary tract infections	Group 1: 3/61 Group 2: 5/59	
			Strictures	Group 1: 6/61 Group 2: 6/59	
			Deep vein thrombosis	Group 1: 0/61 Group 2: 1/59	
			Incontinence	Group 1: 1/61 Group 2: 2/59	
			Deaths (due to cardiovascular or malignant disease)	12 months: Group 1: 1/61 Group 2: 1/59 48 months: Group 1: 2/61 Group 2: 7/59	
			% UI (preoperatively/48 months follow up)	Group 1: 50%/20% Group 2: 47%/17%	
			Patients with decreased erection quality at 48m	Group 1: 8% Group 2: 17%	
			% of men potent	Baseline: Group 1: 50% Group 2: 70% 48 months: Group 1: 53% Group 2: 60%	
			Retrograde ejaculation	Group 1: 24/25 (96.0%) Group 2: 32/37 (86.5%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Wilson et al., 2006 ³²³ Study design: RCT Setting: New Zealand Evidence level: 1+ Duration of follow-up: 24 months	<p>Patient group: Men at urology service at Hospital between June 1997 and December 2000 and considered for surgical treatment for bladder outlet obstruction secondary to BPH.</p> <p>Inclusion criteria: TRUS volume of 40-200g, Qmax of 15ml/s or less, AUA symptom score of 8 or greater, PVR of less than 400ml and urodynamic Schaffer grade 2 or greater.</p> <p>Exclusion criteria: prostatic carcinoma, catheterised patients and those with a history of previous urethral or prostatic surgery.</p> <p>All patients N: 61</p> <p>Group 1 N: 31 Mean (\pmSD) Age: 71.7 (1.1) Dropouts: 9 (one died preoperatively)</p> <p>Group 2 N: 30 Mean (\pmSD) Age: 70.3 (1.0) Dropouts: 4</p>	<p>Group 1: HoLEP Maximum power 100W and a Versacut morcellator was used. Post operative Foley catheter irrigation was performed if deemed necessary; most patients were treated with a Foley catheter, which was normally removed the day after surgery. Mean catheter time: 17.7 hrs Mean hospital time: 27.6 hrs</p> <p>Group 2: TURP Tungsten cutting wire at 160W cutting and 80 W coagulating current. Irrigating Foley catheter inserted and bladder irrigation was used as necessary until haematuria had settled sufficiently to remove the catheter. Mean catheter time: 44.9 hrs Mean hospital time: 49.9h hrs</p>	<p>Mean (SD) AUA symptom score</p> <p>Mean (SD) QoL</p> <p>Mean (SE) Qmax, ml/s</p>	<p>Baseline (n=60) Group1: 26\pm6.02 Group 2: 23.7\pm6.57 3 months (n=56) Group1 (n=28) 4.8\pm4.23 Group 2 (n=29): 3.4\pm4.85 6 months (n=54) Group1 (n=26): 6.0\pm5.10 Group 2 (n=29): 4.8\pm3.77 12 months (n=52) Group1 (n=25): 4.3\pm3.5 Group 2 (n=27): 5.0\pm4.68 24 months (n=48) Group1 (n=22): 6.1\pm4.69 Group 2 (n=26): 5.2\pm4.08</p> <p>Baseline: Group1: 4.8\pm1.1 Group 2: 4.7\pm1.1 3 months: Group1: 1.8\pm2.12 Group 2: 1.9\pm3.23 6 months Group1: 1.6\pm1.53 Group 2: 1.5\pm1.08 12 months Group1: 1.5\pm2.5 Group 2: 1.4\pm1.56 24 months Group1: 1.25\pm0.94 Group 2: 1.25\pm1.02</p> <p>Baseline: Group1: 8.4\pm0.5 Group 2: 8.3\pm0.4 3 months: Group1: 24.2\pm1.7 Group 2: 18.9\pm1.9 6 months Group1: 26.4\pm1.8 Group 2: 20.8\pm2.3</p>	<p>Funding: Supported by Pub Charity, Inc. Financial interest and/or other relationship with Lumenis, Inc, Tel Aviv, Israel.</p> <p>Limitations: Reported Tan 2003 results but these differ to some of the figures quoted in Wilson 2006. Used same results as HTA report.</p> <p>Additional outcomes: PSA before and after in selected patients. PVR at 6 months.</p> <p>Notes: Linked to Tan 2003²⁹² Calculated SD from SE figures given in study.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				12 months Group1: 21.8±2.1 Group 2: 18.4±2.8 24 months Group1: 21.0±2.0 Group 2: 19.3±2.2	
			PdetQmax (cmH2O)	Preoperative Group1: 73.2±4.4 Group 2: 85.8±5.4 6 months Group1: 20.8±2.8 Group 2: 40.7±2.7 P<0.001	
			Schaffer grade	Preoperative Group1: 3.5±0.2 Group 2: 3.7±0.2 6 months Group1: 0.2±0.09 Group 2: 1.2±0.2 P<0.001	
			TRUS volume (cc)	Preoperative Group1: 77.8±5.6 Group 2: 70.0±5.0 6 months Group1: 28.4±1.8 Group 2: 46.6±4.4 P<0.001	
			Onset of erectile dysfunction at 24 months	Group 1: 2 Group 2: 2	
			Retrograde ejaculation	Group 1: 12/16 Group 2: 8/13	
			Preoperative incontinence	Group1: 15/31 (48%) Group 2: 11/30 (38%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Patients that regained continence post operatively	Group1: 6/15 Group 2: 8/11	
			Adverse events at 24 months	Blood transfusion Group1: 0 Group 2: 1 Re-catheterisation Group1: 5 Group 2: 4 Re-operation Group1: 0 Group 2: 2 Urinary tract infections Group1: 0 Group 2: 2 Strictures Group1: 1 Group 2: 3 Deaths (cardiovascular causes) Group1: 0 Group 2: 1	

Evidence Table 23: Thulium laser resection vs. transurethral resection of the prostate

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Xia et al., 2008³³⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Setting: China</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: consecutive BPH patients from November 2004 to December 2005.</p> <p>Inclusion criteria: age < 85yr, maximum urinary flow rate <15ml/s, post void residual urine volume <150ml, medical therapy failure, transrectal ultrasound adenoma volume <100g and urodynamic obstruction.</p> <p>Exclusion criteria: neurogenic bladder, diagnosis of prostate cancer and any previous prostatic, bladder-neck or urethral surgery, and the presence of an indwelling catheter.</p> <p>All patients N: 100</p> <p>Group 1 N: 52 Age (mean): 68.9±7.7 TRUS volume (ml): 59.2±17.7 Drop outs: 0</p> <p>Group 2 N: 48 Age (mean): 69.3±7.3 TRUS volume (ml): 55.1±16.3 Drop outs: 0</p>	<p>Group 1: Thulium laser resection of prostate – tangerine technique. Epidural anaesthesia was achieved. An average power of 50-W thulium lasers operated in continuous wave mode was used. Energy delivered via 550um end-firing fibres. Saline irrigation used. Procedure similar to peeling a tangerine.</p> <p>Group 2: TURP Standard tungsten wire loop with a cutting power of 160W and a coagulating current of 80W. Irrigation started until haematuria had sufficiently decreased.</p> <p>Postoperative care for all patients: Following both procedures, triple lumen catheter inserted into the bladder. Patients kept in hospital 3 days following catheter removal. 500mg levofloxacin used 1 hour before operation and in the postoperative days (once a day).</p>	<p>Mean ±SD symptom score- IPSS</p>	<p>Baseline: Group 1: 21.9±6.7 Group 2: 20.8±5.8</p> <p>6 months: Group 1: 4.0±2.4 Group 2: 3.8±2.8</p> <p>12 months: Group 1: 3.5±2.9 Group 2: 3.9±2.7</p>	<p>Funding: NR</p> <p>Limitations: Allocation concealment and method of randomisation unclear.</p> <p>Additional outcomes: Haemoglobin, serum sodium decrease, resected weight.</p> <p>Notes: None.</p>
			<p>Mean ± SD quality of life</p>	<p>Baseline: Group 1: 4.7±0.9 Group 2: 4.5±1.1</p> <p>6 months: Group 1: 1.1±1.1 Group 2: 0.9±1.0</p> <p>12 months: Group 1: 1.0±0.9 Group 2: 0.9±0.8</p>	
			<p>Mean ± SD Qmax (ml/s)</p>	<p>Baseline: Group 1: 8.0±2.8 Group 2: 8.3±3.0</p> <p>6 months: Group 1: 24.5±9.2 Group 2: 23.3±10.5</p> <p>12 months: Group 1: 23.7±6.0 Group 2: 24.1±6.4</p>	
			<p>Mean ± SD PVR volume (ml)</p>	<p>Baseline: Group 1: 93.1 ±32.1 Group 2: 85.0±36.7</p> <p>6 months: Group 1: 7.1±6.6 Group 2: 6.7±6.3</p> <p>12 months: Group 1: 5.2±4.8 Group 2: 6.1±5.6</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Catheterisation time (hours)	Group 1: 45.7±25.8 Group 2: 87.4±33.8 p value: <0.0001	
			Hospital stay (hours)	Group 1: 115.1±25.5 Group 2: 161.1±33.8 p value: <0.0001	
			Operative time (minutes)	Group 1: 46.3±16.2 Group 2: 50.4±20.7 P=0.28	
			Adverse events	Blood transfusion Group 1: 0 Group 2: 2 (4.2%) TUR Group 1: 0 Group 2: 1 (2.1%) Urinary tract infection Group 1: 2 (3.9%) Group 2: 4 (8.3%) Recatheterisation Group 1: 0 Group 2: 0 Transitory urge incontinence Group 1: 12 (23.1%) Group 2: 15 (31.3%) Retrograde ejaculation Group 1: 18/33 (55%) Group 2: 20/31 (65%) Urethral stricture Group 1: 1 (1.9%) Group 2: 3 (6.3%) Stress incontinence Group 1: 0 Group 2: 1 (2.1%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			IIEF-5 scores	Preoperative: Group 1: 19.3±6.1 Group 2: 20.0±5.2 6 months Group 1: 20.4±6.0 Group 2: 21.7±4.8 12 months: Group 1: 21.0±5.8 Group 2: 21.4±5.3 P=0.67	
			Mean ± SD PdetQmax(cmH2O)	Preoperative: Group 1: 85.9±29.3 Group 2: 83.4±33.3 12 months: Group 1: 38.1±17.5 Group 2: 38.9±17.3 P=0.80	
			Schafer grade	Preoperative: Group 1: 3.8±1.1 Group 2: 3.6±1.2 12 months: Group 1: 0.71±0.67 Group 2: 0.79±0.77 P=0.58	

Evidence Table 24: Holmium laser enucleation of the prostate (HoLEP) vs. transurethral incision of the prostate (HoBNI)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Aho et al., 2005¹⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Men with bladder outflow obstruction (BOO) and small prostate (<40g)</p> <p>Setting: Urology department, New Zealand, between July 1998 to May 2001</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> ▪ Qmax less than 15 ml/s ▪ AUA symptom score ≤8 ▪ Prostate volume (measured by TRUS) ≤40cc ▪ PVR<400ml ▪ Schafer grade ≥2 <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> ▪ Known prostate cancer, or suspected prostate cancer (increased PSA and/or suspicious of DRE underwent TRUS biopsy) ▪ Catheterised patients ▪ History of urethral surgery ▪ On anticoagulants or had coagulation defects <p>All patients N: 40</p> <p>Drop outs:</p> <p>Group 1 - HoLEP N: 20</p>	<p>Group 1: HoLEP Performed under general anaesthesia by 1 of 2 surgeons. (technique described in another paper)</p> <p><u>Energy used (kJ), mean (range):</u> 74.2 (56-104)*</p> <p><u>Operative time, mins, mean, SD (range):</u> 29.7±6.1(18-43) *</p> <p><u>As outpatient procedure:</u> 15/19 (the above values are for 19 patients- 1 died preoperatively)</p> <p>Group 2: Ho BNI Performed under general anaesthesia by 1 in 3 surgeons. Incisions made at the 5 and 7 o' clock positions from just distal to each urethral orifice to either side of the verumontanum down to the depth of the surgical capsule. No tissue was excised.</p> <p><u>Energy used (kJ), mean (range):</u> 13.3 (5-26)*</p> <p><u>Operative time, mins, mean, SD (range):</u> 7.0±3.3(2-17) *</p> <p><u>As outpatient procedure:</u> 14/20</p>	<p>IPSS symptom score, mean ±SD, (range)</p>	<p><u>At 1 months</u> Group 1: 8.7±5.8 (0-21) Group 2: 6.2±6.8 (0-30) Relative risk: 95% CI: <u>At 3 months</u> Group 1: 6.8±5.5 (1-21) Group 2: 6.2±6.7 (0-22) Relative risk: 95% CI: <u>At 6 months</u> Group 1: 7.9±6.6 (0-26) Group 2: 9.1±8.4 (1-28) Relative risk: 95% CI: <u>At 12 months</u> Group 1: 8.9±8.5 (1-31) Group 2: 6.1±5.6 (1-16) Relative risk: 95% CI: p value: NS at anytime point</p>	<p>Funding: Supported by Pub Charity, Inc</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Number of patients with urinary incontinence was significantly different pre-operatively. ▪ Reporting of adverse event – definitions and follow-up period ▪ There was imbalance in the number of incontinence cases at baseline.: 2/20 vs. 11/20 ▪ Retrograde ejaculation outcome was based on the number of patients who were able to comment (sexually active?). The number of patients who were able to comment was not reported. <p>Additional outcomes: Death – 1 in HoLEP (pre-operative), 1 in BNI at 6th month (cardiac)</p> <p>Notes: Sample size calculation was provided. As sample size of 40 would be required to detect HoLEP is superior (Qmax change of 12ml/s</p>
			<p>IPSS QoL score mean ±SD, (range)</p>	<p><u>At 1 months</u> Group 1: 2.2±1.6 (0-6) Group 2: 1.4±1.6 (0-6) Relative risk: 95% CI: <u>At 3 months</u> Group 1: 1.8±1.4 (0-6) Group 2: 1.8±1.5 (0-6) Relative risk: 95% CI: <u>At 6 months</u> Group 1: 2.0±1.4 (0-5) Group 2: 2.1±1.5 (0-5) Relative risk: 95% CI: <u>At 12 months</u> Group 1: 1.7±0.9 (0-5) Group 2: 1.5±0.9 (0-3)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean): 65.1±11.5 (range not provided)</p> <p>Drop outs at 0/1/3/6/12 months: 0/1/2/3/4, 1 patient died pre-operatively</p> <p>IPSS symptom: 25.2±5.9(15-34)</p> <p>IPSS QoL: 5.2±0.8 (4-6)</p> <p>Qmax: 8.3±3.0(4-14)</p> <p>PdetQmax H₂O: 72.0±29.1(45-145)</p> <p>Schafer Grade: 3.2±1.3(2-6)</p> <p>Prostate Volume, PV: 30.3±6.6(14-39)</p> <p>Urinary incontinence: 2/20[#]</p> <p>Erectile dysfunction: 10/20</p> <p>Group 2 - HoBNI</p> <p>N: 20</p> <p>Age (mean): 64.9±10.1 (44-79)</p> <p>Drop outs at 0/1/3/6/12 months: 0/0/2/3/8</p> <p>IPSS symptom: 24.2±5.1(14-35)</p> <p>IPSS QoL: 5.0 ±1.0 (3-6)</p> <p>Qmax: 9.7±1.3(8-12)</p> <p>PdetQmax H₂O: 71.0±30.2(40-128)</p> <p>Schafer Grade: 3.2±1.3(2-6)</p> <p>Prostate Volume, PV: 30.5±5.9(18-39)</p> <p>Urinary incontinence: 11/20[#]</p> <p>Erectile dysfunction: 9/20</p> <p>[#]P value =0.006, calculated by NCGC team using Fisher's exact test</p>	<p>Both groups</p> <ul style="list-style-type: none"> ▪ <u>Maximal lasing power:</u> 100 W (2J at 50 Hz) ▪ Versacut™ morcellator ▪ <u>Catheters:</u> Two way catheters unless post-operative bladder irrigation was necessary. Catheters removed at the hospital or in the community the morning following surgery. ▪ <u>Discharged from hospital:</u> the afternoon or evening following surgery <p>*P value<0.001</p>		<p>Relative risk:</p> <p>95% CI:</p> <p>p value: NS at anytime point</p>	<p>compared to 8ml/s in BNI), at a power of 80% and p of 0.05</p>
		Qmax , mean ±SD, (range)	<p><u>At 1 months</u></p> <p>Group 1: 19.9±6.9(9-40)</p> <p>Group 2: 18.7±8.0(9-40)</p> <p>Relative risk:</p> <p>95% CI:</p> <p><u>At 3 months</u></p> <p>Group 1: 20.7±7.6 (7-36)</p> <p>Group 2: 18.5 ±9.2 (10-36)</p> <p>Relative risk:</p> <p>95% CI:</p> <p><u>At 6 months</u></p> <p>Group 1: 20.2±8.0 (5-33)</p> <p>Group 2: 17.4±7.3 (3-31)</p> <p>Relative risk:</p> <p>95% CI:</p> <p><u>At 12 months</u></p> <p>Group 1: 21.6±7.7 (10-38)</p> <p>Group 2: 17.4±4.6 (12-24)</p> <p>Relative risk:</p> <p>95% CI:</p> <p>p value: NS at anytime point</p>		
		PdetQmax (cm H ₂ O), mean ±SD, (range)	<p><u>At 6 months</u></p> <p>Group1: 29.1±11.1 (15-50)</p> <p>Group 2: 43.2±25.4 (2-100)</p> <p>Relative risk:</p> <p>95% CI:</p> <p>p value:<0.01</p>		
		Urodynamic obstruction, Schafer grade, mean ±SD, (range)	<p><u>At 6 months</u></p> <p>Group1: 0.5 ±0.7(0-5)</p> <p>Group 2: 1.6±1.4 0-5</p> <p>Relative risk:</p> <p>p value:<0.01</p>		
		Urodynamically obstructed No definition. 4 patients in HoBNI group subsequently had	<p><u>At 6 months</u></p> <p>Group1: 0/19</p> <p>Group 2: 5/20 (25%)</p>		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			HoLEP. See "Reoperation"	Relative risk: 95% CI: p value: NR	
			Prostate Volume , (g) mean \pm SD, (range). Measured using TRUS	<u>At 6 months</u> Group 1: 22.2 \pm 7.1(11-35) Group 2: 31.5 \pm 8.0(21-49) Relative risk: p value: <0.05	
			Catheter duration , mean \pm SD (range), hours	Group 1: 22.9 \pm 6.9(12-48) Group 2: 23.2 \pm 1.9(17-25) Relative risk: 95% CI: p value: NS	
			Post-op complications (early): Recatheterisation	Group 1: 0/19 Group 2: 2/20 Relative risk: p value: NR	
			Post-op complication: Reoperation: Patients had HoLEP between 6-16 months because of persistent LUTS	Group 1: 0/19 (within 1 year) Group 2: 4/20 Relative risk: p value:	
			Post-op complications: Submeatal Strictures	Group 1: 1 (dilated) Group 2: 1 (meatotomy) Relative risk: p value: NS	
			Incontinence % with incontinence <u>Note:</u> Patients in Group 2 (BNI) who had reoperation was not assessed.	<u>At 12 months</u> Group 1: 4/16 (44%) - Group 2: 0/13 (0%) Relative risk: p value: <0.01 None of the patients required pads	
			Erectile function: (No change /Worsened/ Improved)	<u>At 12 months</u> Group 1: 11/2/3 Group 2: 10/1/2 Relative risk: p value: NS	
			Post-op complications:	Group 1: 100%	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Retrograde ejaculation in sexually, % (in patients who are able to “comment” on it, number of patients not stated)	Group 2: 80% Relative risk: p value: reported as <0.01	
			Hospital time: mean \pm SD (range), hours	Group1: 12.3 \pm 7.0 (7-28) Group 2: 13.7 \pm 8.5 (7-28) Relative risk: 95% CI: p value: NS	

Evidence Table 25: Holmium laser enucleation of the prostate (HoLEP) vs. open prostatectomy (OP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kuntz et al., 2008¹⁵²</p> <p>Study design: RCT</p> <p>Setting: Department of Urology- Germany</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 5 years</p>	<p>Patient group: Candidates for surgical therapy of lower urinary symptoms and obstruction due to a prostate larger than 100 gm.</p> <p>Inclusion criteria: AUA\geq8, (Q_{max}) of \leq12 ml/s, post void residual urine volume \geq 50 ml, Schafer grade \geq 2.</p> <p>Exclusion criteria: Previous prostate or urethral surgery and non-BPH-related voiding disorders. Preoperatively, prostate carcinoma was screened for and excluded by prostate biopsy if indicated. There was no upper limit for prostate size.</p> <p>All patients N: 120 Drop outs: 46</p> <p>Group 1: N: 60 Mean \pmSD (range) Age: 69.2 \pm 8.4 (56-89) Schaffer grade: 4.3 \pm 1.12 (3-6) Postvoid residual volume (ml): 280 \pm 273 (50-1,000) Peak urinary flow rate (ml/s): 3.8 \pm 3.6 (0-10) Dropouts: 18 (died=3, intercurrent illness=3, moving=6, prostate cancer=3, reoperations=3)</p>	<p>Group 1: HoLEP HoLEP was carried out at 80 or 100 W with a high-powered Ho:YAG laser (2.0 J; 40-50 Hz). It involved retrograde enucleation of the median and lateral lobes from the apex toward the bladder. When the trial started, a mechanical tissue morcellator was not yet commercially available. Therefore in the first 50 of the 60 HoLEP patients, fragmentation of the lobes was performed by traditional electrocautery loop resection whilst the devascularised lobes were still connected to the surgical capsule by a narrow pedicle. In the last 10 of the 60 HoLEP patients, the lobes were enucleated in their entirety, pushed into the bladder, and fragmented with the use of a mechanical tissue morcellator.</p> <p>Group 2: Open prostatectomy (OP) Open prostatectomy was performed by a</p>	<p>Mean \pm SD AUA symptom score:</p>	<p>Preoperatively: Group 1: 22.1 \pm 3.3 (n=60) Group 2: 21.0 \pm 3.6 (n=60);</p> <p>3 months Group 1: 3.3 \pm 2.7 (n=54) Group 2: 3.6 \pm 2.7 (n=50)</p> <p>6months Group 1: 2.4 \pm 1.9 (n=54) Group 2: 2.8 \pm 3.9 (n=50)</p> <p>1-year: Group 1: 2.3 \pm 2.0 (n=56) Group 2: 2.3 \pm 1.7 (n=49); P value: 0.94</p> <p>2-year: Group 1: 2.3 \pm 2.2 (n=53) Group 2: 2.4 \pm 1.6 (n=46); P value: 0.89</p> <p>3 year. Group 1: 3.0 \pm 3.1 (n=48) Group 2: 2.8 \pm 1.6 (n=40); P value: 0.82</p> <p>4-year: Group 1: 3.0 \pm 3.1(n=45) Group 2: 2.8 \pm 1.9 (n=36); P value: 0.68</p> <p>5-year: Group 1: 3.0 \pm 3.2 (n=42) Group 2: 3.0 \pm 1.7 (n=32); P value: 0.98</p>	<p>Funding: Prof. Kuntz is a consultant for the companies Lumenis and Karl Storz.</p> <p>Limitations: Allocation concealment and blinding unclear.</p> <p>Notes: Linked with Kuntz 2002¹⁵⁰ and Kuntz2004¹⁵¹</p>
			<p>Mean \pm SD peak flow (ml/s)</p>	<p>Preoperatively: Group 1: 3.8 \pm 3.6 (n=60) Group 2: 3.6 \pm 3.8 (n=60); P value: 0.60</p> <p>3 months: Group 1: 27.6 \pm 7.0 (n=54) Group 2: 27.3 \pm 6.2 (n=50); P value: 0.66</p> <p>1-year: Group 1: 27.4 \pm 9.7 (n=56) Group 2: 28.3 \pm 7.5 (n=49); P value: 0.86</p> <p>2-year: Group 1: 26.7 \pm 8.3 (n=53) Group 2: 27.4 \pm 6.8 (n=46); P value: 0.65</p> <p>3-year: Group 1: 27.0 \pm 9.8 (n=48) Group 2: 25.3 \pm 6.9 (n=40); P value: 0.32</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2: N: 60 Mean \pmSD (range) Age: 71.2 \pm 8.3 (54-89) Schaffer grade: 4.3 \pm 0.79 (3-6) Postvoid residual volume (ml): 292 \pm 191 (50-1,000) Peak urinary flow rate (ml/s): 3.6 \pm 3.8 (0-12) Dropouts: 28 (died=8, intercurrent illness=3, moving=7, prostate cancer=6, reoperation=4)</p>	suprapubic transvesical approach via midline incision. The bladder catheter was routinely removed on the seventh postoperative day.		<p>4-year: Group 1: 27.7 \pm 9.6 (n=45) Group 2: 25.0 \pm 8.3 (n=36); P value: 0.20 5-year: Group 1: 24.3 \pm 10.1 (n=42) Group 2: 24.4 \pm 7.4 (n=32); P value: 0.97</p>	
			Mean \pm SD Residual volume (ml)	<p>Preoperatively: Group 1: 280 \pm 273 (n=60) Group 2: 292 \pm 191 (n=60); P value: 0.43 1-year: Group 1: 5.8 \pm 16.7 (n=56) Group 2: 6.4 \pm 12.3 (n=49); P value: 0.83 2-year: Group 1: 1.7 \pm 6.5 (n=53) Group 2: 2.4 \pm 6.8; P value: 0.61 3-year: Group 1: 6.1 \pm 12.1 (n=48) Group 2: 4.4 \pm 10.5 (n=40); P value: 0.50 4-year: Group 1: 8.6 \pm 13.5 (n=45) Group 2: 6.5 \pm 12.1 (n=36); P value: 0.48 5-year: Group 1: 10.6 \pm 24.4 Group 2: 5.3 \pm 11.2 (n=32); P value: 0.25</p>	
			Mortality (follow up 60 months)	Group 1: n=3 Group 2: n= 8	
			Mortality (3 months postoperatively)	Group 1: n=0 Group 2: n= 2	
			Complications (6 months postoperatively):	<p><u>Blood transfusion</u> Group 1: 0 Group 2: 8 (13.3%); P value: 0.003 <u>Reoperation for secondary coagulation of bleeding arteries (18)</u> Group 1: 3 Group 2: 3; P value: NR <u>Reoperation for secondary apical resections</u> Group 1: 2 Group 2: 0; P value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Re-interventions (60months)	Bladder neck contracture- holium laser incision: Group 1: 1 (1.7%) Group 2: 3 (5.0); P value: 0.60 Visual urethrotomy (from stricture): Group 1: 2 (3.3%) Group 2: 1 (1.7); P value: 0.61	
			Mean +/- SD Post-op stay (hrs.)	Group 1: 69.6 +/- 36.4 (24-192) Group 2: 251.0 +/- 45.5 (216-552) P value: <0.0001	
			Recatheterisation	Group 1: 3 (5%) Group 2: 3 (5%)	
			Incontinence	Group 1: 5/60 Group 2: 6/60	
			Erectile dysfunction	Group 1: 5/54 Group 2: 5/50	
			Retrograde ejaculation (in sexually active patients; 58%)	Group 1: 70% Group 2: 79%	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Naspro et al., 2006²¹⁰</p> <p>Study design: RCT</p> <p>Setting: Italy</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24-months</p>	<p>Patient group: Consecutive patients from March 2003 to December 2004 who suffered from BPH-related obstructed voiding symptoms with prostate volume >70 g, as determined by transrectal ultrasound and who had not responded to pharmacologic therapy.</p> <p>Inclusion criteria: Postvoiding residue <150 ml, peak urinary flow rate <15 ml/s, and urodynamic obstruction (Schafer grade >2).</p> <p>Exclusion criteria: Neurogenic bladder, history of adenocarcinoma of the prostate, or any previous prostatic, bladder-neck, or urethral surgery.</p> <p>All patients N: 80 Drop outs: 15</p> <p>Group 1: N: 41 Mean (\pmSD) Age: 66.26 (\pm 6.55) Total serum PSA ng/ml mean (\pmSD): 6.33 \pm 3.45 Incidental adenocarcinoma: 2 (4.8%) Dropouts: 6</p> <p>Group 2: N: 39 Mean (\pmSD) Age: 67.27 (\pm 6.72) Total serum PSA ng/ml mean (\pmSD):</p>	<p>Group 1: HoLEP The surgical technique included enucleation of the prostatic lobes with subsequent tissue morcellation into the fragments, which were retrieved from the bladder cavity.</p> <p>Total mean operative time: 72.09 \pm 21.22</p> <p>Group 2: OP Standard transvesicle approach.</p> <p>Total mean operative time: 58.31 \pm 11.95</p>	<p>Mean (SD) IPSS</p>	<p>Baseline: Group 1: 20.11 \pm 5.84 Group 2: 21.60 \pm 3.24; p value: 0.27</p> <p>1-month: Group 1: 6.9 \pm 4.2 Group 2: 4.7 \pm 2.1; p value: 0.20</p> <p>3-month: Group 1: 3.9 \pm 2.9 Group 2: 2.9 \pm 2.6; p value: 0.46</p> <p>12-month: Group 1: 8.45 \pm 5.87 Group 2: 8.40 \pm 6.0; p value: 0.98</p> <p>24-month: Group 1 (n=35): 7.9 \pm 6.2 Group 2: (n= 30): 8.1 \pm 7.1; p value: 0.44</p>	<p>Funding: NR</p> <p>Limitations: Allocation concealment and blinding unclear.</p> <p>Notes: None.</p>
			<p>Q_{max}</p>	<p>Baseline: Group 1: 7.83 \pm 3.42 Group 2: 8.32 \pm 2.37; p value: 0.64</p> <p>1-month: Group 1: 26.6 \pm 8.7 Group 2: 24.3 \pm 6.8; p value: 0.53</p> <p>3-month: Group 1: 22.2 \pm 8.6 Group 2: 25.5 \pm 10.5; p value: 0.57</p> <p>12-month: Group 1: 22.32 \pm 3.8 Group 2: 24.21 \pm 6.49; p value: 0.27</p> <p>24-month: Group 1 (n=35): 19.19 \pm 6.3 Group 2: (n= 30): 20.11 \pm 8.8; p value: 0.91</p>	
			<p>QOL question</p>	<p>Baseline: Group 1: 4.07 \pm 0.93 Group 2: 4.44 \pm 0.96; p value: 0.17</p> <p>1-month: Group 1: 1.4 \pm 1.4 Group 2: 1.3 \pm 0.7; p value: 0.76</p> <p>3-month:</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	6.99 +/- 4.28 Incidental adenocarcinoma: 3 (7.6%) Dropouts: 9			Group 1: 1 +/- 0.8 Group 2: 0.6 +/- 0.2; p value: 0.18 12-month: Group 1: 1.7 +/- 0.94 Group 2: 1.77 +/- 0.83; p value: 0.85 24-month: Group 1 (n=35): 1.5 +/- 0.87 Group 2 (n= 30): 1.66 +/- 0.76; p value: 0.76	
			Mean detrusor pressure at maximum flow rate (P_{detqmax})cm H₂O	Baseline: Group 1: 80.6 (44-130) Group 2:: 83.1 (41-147); p value: 0.94 12-month: Group 1: 30.6 (22-80) Group 2:: 34.8 (18-88); p value: 0.66	
			Schafer grade (LinPURR):	Baseline: Group 1: 3.8 (2-6) Group 2:: 3.1 (2-6); p value: 0.33; 12-month: Group 1: 0.7 (0-4) Group 2:: 0.8 (0-4); p value: 0.18	
			Perioperative morbidity (surgery to 3months)	Bladder mucosal injury: Group 1: 3 (7.3%) Group 2:: 0 (2-6); p value: < 0.001 Transitory urge incontinence: Group 1: 14 (34.1%) Group 2:: 17 (38.6%); p value: 0.2 Dysuria (burning): Group 1: 28 (68.2%) Group 2:: 16 (41.0%); p value: <0.001 Stress incontinence: Group 1: 1 (2.4%) Group 2: 1 (2.5%); p value: 0.9 Reintervention for bleeding: Group 1: 1(2.4%) Group 2:: 0; p value: 0.9 Early acute urinary retention: Group 1: 5 (12.1%) Group 2:: 2 (5.1%); p value: 0.11	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			<p>Complications 12-month follow-up:</p>	<p>Urge incontinence: Group 1: 2 (5.4%) Group 2: 3 (8.5%); p value: 0.03</p> <p>Dysuria (burning): Group 1: 4 (10.8%) Group 2: 3 (8.5%); p value: 0.02</p> <p>Bladder-neck/urethral strictures: Group 1: 2 (5.4%) Group 2: 2 (5.7%); p value: 0.3</p> <p>Overall reintervention: Group 1: 2 (5.4%) Group 2: 2 (5.7%); p value: 0.55</p> <p>Prostate cancer: Group 1: 4 (10.8%) Group 2: 4 (11.4%); p value: 0.4</p> <p>24-month follow-up:</p> <p>Prostate cancer: Group 1: 0 Group 2: 0; p value:</p> <p>Dysuria (burning): Group 1: 1 (2.8%) Group 2: 1 (3.3%); p value: 0.02</p> <p>Bladder-neck/urethral strictures: Group 1: 1 (2.8%) Group 2: 1 (3.3%); p value: 0.3</p>	
			<p>Mean +/- SD IIEF domains</p>	<p>baseline: Group 1: 20.3 +/- 6.6 Group 2: 21.1 +/- 5.3; p value: 0.5</p> <p>3 months: Group 1: 21.4 +/- 2.6 Group 2: 20.6 +/- 5.5; p value: 0.67</p> <p>6 months: Group 1: 22.8 +/- 2.1 Group 2: 24.6 +/- 4.0; p value: 0.55</p> <p>12 months: Group 1: 25.2 +/- 4.2 Group 2: 23.5 +/- 1.8; p value: 0.31</p> <p>24 months: Group 1: 22.3 +/- 4.0 Group 2: 21.9 +/- 5.6; p value: 0.21</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>Autologous blood transfusion: Group 1: 2 (4%) Group 2: 5 (12.8%) p value: < 0.001</p> <p>Homologous blood transfusion: Group 1: 0 Group 2: 2 (5.1%) p value: < 0.007</p> <p>Catheterisation time: Group 1: 1.5 +/- 1.07 Group 2: 4.1 +/- 0.5 p value: < 0.0001</p> <p>Hospital stay, d: Group 1: 2.7 +/- 1.1 Group 2: 5.43 +/- 1.05 p value: < 0.0001</p>	

Evidence Table 26: Laser coagulation vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Anson1995¹⁸</p> <p>McAllister2000¹⁸⁸</p> <p>Study design: RCT, open label, (multi-centre)</p> <p>Setting: United Kingdom</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 5 years</p>	<p>Patient group: Patients with BPH</p> <p>Setting: From March 1992, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Age > 50 years old American Society of Anaesthesiologist (ASA) Grade 1 to 3 Prostatic urethral length > 24mm Urinary flow rates consistent with outlet obstruction <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ASA Grade > 3 Known history or suspicion of prostate cancer Renal impairment Life expectancy < 6 months On medication such as anticoagulants <p>All patients N: 151, out of 166 candidates Age, mean, (range) (years): 68.1 (52-84)</p> <p>Drop outs</p> <ul style="list-style-type: none"> 1-year review: 137/151 5-year review: 42/151 (109 patients were traced from 151 at the 5-year review) <p>Group 1-Laser coagulation N: 76 Drop outs:</p> <ul style="list-style-type: none"> At 1-year review: 9/76 (11.8%) 	<p>Group 1- Laser coagulation (ELAP) Procedure: Nd:YAG, using Urolase fibre. Energy was applied at 60W for 6S at the 2, 5, 7, and 10 o'clock positions, modified according to prostate length and presence of median lobe. Room temperature sterile water was used for irrigation</p> <p>Power: 60W</p> <p>Group 2 –TURP Procedure: Standard electroresection, by experienced urologists</p>	<p>All cause mortality</p> <p>AUA-6 symptom score, mean (95% CI):</p> <p>Qmax, mean (95% CI):</p>	<p><u>“immediate post-operative period”</u> Group 1: 0/76 Group 2: 0/75 p value: NS <u>Week 52 (1 year)</u> Group 1: 1/76 Group 2: 1/75 p value: NS</p> <p><u>Week 4</u> Group 1: 13.5 (95%CI: 12.0 to 15.0) Group 2: 8.7 (95%CI: 7.6 to 9.8) p value: NS <u>Week 12</u> Group 1: 8.7 (95%CI: 7.3 to 10.1) Group 2: 6.4 (95%CI: 5.2 to 7.6) p value: NS <u>Week 26</u> Group 1: 7.9 (95%CI: 6.4 to 9.4) Group 2: 5.9 (95%CI: 4.6 to 7.2) p value: NS <u>Week 52</u> Group 1: 7.7 (95%CI: 6.3 to 9.1) Group 2: 5.1 (95%CI: 3.8 to 6.4) p value: < 0.05 <u>5 years</u> Group 1: 6.3, n=28 Group 2: 6.5, n=39 p value: NS</p> <p><u>Week 12</u> Group 1: 15.9 (95%CI: 13.6 to 18.2) Group 2: 21.3 (95%CI: 19.0 to 23.6) p value: < 0.05 <u>Week 26</u> Group 1: 15.6 (95%CI: 13.7 to 17.5) Group 2: 19.9 (95%CI: 17.4 to 22.4) p value: NS <u>Week 52</u> Group 1: 15.4 (95%CI: 13.6 to 17.2) Group 2: 21.8 (95%CI: 18.5 to 25.1)</p>	<p>Funding: Bard Europe Division</p> <p>Limitations:</p> <ul style="list-style-type: none"> Open label study Randomisation concealment method not described Only 44% of patients available at 5-year follow up, and no sd was provided. <p>Additional outcomes:</p> <ul style="list-style-type: none"> Pulmonary embolism – 1 patient in TURP group had PE after operation Deep vein thrombosis: 1 patient in laser group vs. 2 patients in TURP group had DVT <p>Notes: 5 year data not used in meta-analysis due to small number of available data compared to original sample size</p> <p>McAllister2000 reported the 5 year follow up period</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<ul style="list-style-type: none"> ▪ At 5-year review: 19/76 (25%) Age: mean (95% CI): 67.9 (66.3-69.5) Drop outs: Not stated AUA-6 symptom score, mean (95% CI): 18.1(17.1-19.1) Qmax, mean (95% CI): 9.6(8.8-10.4) Post void residual volume: mean (95% CI): 113(91-135) Sexually active: 27/76 (36%) <p>Group 2 - TURP N: 75 Drop outs:</p> <ul style="list-style-type: none"> ▪ At 1-year review: 5/75(6.7%) ▪ At 5-year review: 24/75(32%) <p>Age: mean (95% CI): 68.3(66.5-70.1)</p> <p>AUA-6 symptom score, mean (95% CI): 18.2(17.1-19.3)</p> <p>Qmax, mean (95% CI): 10.0 (9.1-10.9)</p> <p>Post void residual volume: mean (95% CI): 121(93-148)</p> <p>Sexually active:24/75 (32%)</p>		<p>Post void residual volume: mean (95% CI):</p> <p>Post-operative complications: Blood transfusion: (Mean of 2.7 units blood)</p> <p>Post-operative complications: Retrograde ejaculation (among patients who were sexually active preoperatively)</p> <p>Post-operative complications: Clot retention</p> <p>Post-operative</p>	<p>p value: NS <u>5 years</u> Group 1: 17.8, n=24 Group 2: 20.0, n=36 p value: NS</p> <p><u>Week 12</u> Group 1: 70.3 (95%CI: 51.1 to 89.3) Group 2: 21.3 (95%CI: 43.9 to 80.3) p value: NS</p> <p><u>Week 26</u> Group 1: 90.1 (95%CI: 61.6 to 118.0) Group 2: 19.9 (95%CI: 17.4 to 22.4) p value: <0.05</p> <p><u>Week 52</u> Group 1: 69.2 (95%CI:48.1 to 90.3) Group 2: 45.9 (95%CI:30.5 to 61.3) p value: <0.05</p> <p><u>5 years</u> Group 1: 76, n=24 Group 2: 55, n=35 p value: NS</p> <p>Group 1: 0/76 Group 2: 3/75 p value: NS</p> <p><u>Up to week 52 (1 year)</u> Group 1: 9/27 (33%) Group 2: 15/24 (63%) p value: NS</p> <p><u>Up to week 52 (1 year)</u> Group 1: 1/76 Group 2: 5/75 p value: NS</p> <p><u>Up to week 4</u></p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			complications: urinary tract infection (positive culture). 22/28 of patients in the ELAP group received prophylaxis	Group 1: 18/76 Group 2: 5/75 RR: 3.55 (95% CI: 1.47 to 8.97) p value: <0.01 <u>Up to week 52 (1 year)</u> Group 1: 28/76 Group 2: 7/75 RR: 3.95 (95% CI: 1.92 to 8.48) p value: <0.01	
			Post-operative complications: Dysuria	<u>Up to week 52 (1 year)</u> Group 1: 25/76 Group 2: 6/75 RR: 4.11 (95% CI: 1.88 to 9.42) p value: <0.01	
			Post-operative complications: epididymorchitis	<u>Up to week 52 (1 year)</u> Group 1: 2/76 Group 2: 1/75 p value: NS	
			Post-operative complications: Reoperation- by week 52, 2 had bladder neck incision, 3 had TURP	<u>Up to week 52 (1 year)</u> Group 1: 5/76 Group 2: 0/75 p value: <u>5 years</u> Group 1: 18/47 (38%) Group 2: 8/51 (16%) p value: <0.006	
			Hospitalisation days, mean (95% CI)	Group 1: 2.7(95%CI: 2.2 to 3.2) Group 2: 4.3 (95%CI: 3.3 to 5.3) p value: NS	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Chacko et al., 2001 ⁴⁸ CLASP study- acute urinary retention Study design: RCT, multicentre, open label Setting: UK Evidence level: 1+ Duration of follow-up: 7.5 months	Patient group: men with acute painful, urinary retention Setting: 3 centres in UK Inclusion criteria: Acute painful, urinary retention. All patients without strong history of LUTS underwent at least one trial without catheter Exclusion criteria: <ul style="list-style-type: none"> Prostate cancer or previous prostatic surgery; prostate size > 120ml; Life expectancy < 6 months; Urinary retention associated with recent operation, constipation or drugs which could cause acute urinary dysfunction, Neurogenic bladder dysfunction; Serum creatinine >250 µmol/L. All patients Number of eligible patients: 155 N randomised: 148 Mean age: Drop outs: Group 1-Laser coagulation N: 74 Dropouts:	Group 1- Laser coagulation Procedure: Nd:YAG/ Non-contact VLAP, side-firing fibre (Bard Urolase), using standard fixed spot technique Power: 60W ND: YAG for 60s, depends on prostate size. For prostate size with urethral length of >25 mm, additional set of laser was used. If median lobe was present, 60W for 30s was applied for each side of lobe. Energy: 33.93kJ (mean total delivered) Catheter protocol: Suprapubic catheter, voiding trial 1-2 wks after discharge. Other: All patients received antibiotic prophylaxis and anti-inflammatory suppository. Group 2 –TURP Procedure: Standard electroresection Catheter protocol: suprapubic; duration depends on success voiding after urine is clear.	All cause mortality Not treatment related	Group 1: 2/74 Group 2: 4/74 p value: NS	Funding: Laser machines provided by Bard Diagnostics, Redmond, Washington. Limitations: <ul style="list-style-type: none"> Open label study, with main outcomes using patient reported measures. The actual values of data and standard deviations were not reported for many outcomes – only reported p values or whether it was statistically significant – not suitable for meta-analysis Additional outcomes: <ul style="list-style-type: none"> Myocardial infarction during hospital stay Composite outcomes categories, and categorical outcomes for IPSS and Qmax Notes: <ul style="list-style-type: none"> Sample size calculation was performed. In the laser group, 7/74 patients were converted to the
			IPSS, mean change from baseline (±SD): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: -10.1 (95%CI: -12.8, -7.3), n=54 Group 2: -13.5 (95%CI -15.8, -11.2), n=48 p value: 0.26 Both groups stats sig compared to baseline	
			IPSS-QoL, mean(±SD): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: -3.10 (95%CI -3.65, -2.55), n=49 Group 2: -3.42 (95%CI -3.89, -2.95), n=45 Adjusted difference: : 0.26 (0.81-0.30)- page 169 P value: 0.37 Both groups stats sig compared to baseline	
			Post-op complications: Transurethral resection syndrome	Group 1: 0/74 Group 2: 2/74 P value: NS	
			Post-op complications: Blood transfusion (units and criteria not stated)	Group 1: 0/74 Group 2: 4/74 P value: NS	
			Post-op complications: Heavy bleeding (criteria not stated)	Group 1: 2/74 Group 2: 3/74 P value: NS	
			Post-op complications: Septicaemia	Group 1: 3/74 Group 2: 4/74 P value: NS	
			Post-op complications: Incontinence	Group 1: 0/74 Group 2: 3/74 P value: NS	
			Post-op complications: Reoperation (surgery	Group 1: 7/74 Group 2: 1/74	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Received as allocated: 57/74 Age, mean (\pmSD): 74.2 \pm 7.9 IPSS, mean (\pmSD): 20.3 \pm 9.3 IPSS-QoL, median(IQR): 5 (4-6) Ethnicity (% white): 97.3</p> <p>Group 2 - TURP N: 74 Dropouts: Received as allocated: 68/74 Age, mean (\pmSD): 72.7 \pm 7.3 IPSS, mean (\pmSD): 19.4 \pm 7.6 IPSS-QoL, median(IQR): 5 (4-6) Ethnicity (% white): 97.3</p>	<p>Other: All patients received antibiotic prophylaxis and anti-inflammatory suppository.</p>	<p>due to “unacceptable symptoms” or retention after 8 weeks)</p> <p>Post-op complications: Urinary retention (>8 weeks)</p> <p>LOS, geometric mean, days</p>	<p>P value: NS</p> <p>Group 1: 1/74 Group 2: 0/74 P value: NS</p> <p>Group 1: 3.4 (95% CI 2.8 to 4.0) Group 2: 5.8 (95% CI 5.2 to 6.5)</p> <p>Relative risk: 1.73 95% CI: 1.40-2.14 P value: <0.0001</p>	<p>standard surgery in theatre, and 3 refused treatment.</p> <ul style="list-style-type: none"> ▪ In the TURP group, 5 refused or deferred treatment. ▪ A total of 1073 patients were considered for inclusion of the 3 linked CLASP trial, and 570 were entered. 318 (29.5%) were not eligible because of \geq1 exclusion criteria. The rest did not enter for various reasons. There were 240 patients in the uncomplicated LUTS trial, 148 in the acute urinary retention trial and 82 in the chronic retention trial.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Cowles et al., 1995⁵⁶</p> <p>Study design: RCT, open label, multicentre</p> <p>Setting: United states</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Bladder outlet obstruction due to BPH</p> <p>Setting: Multicentre, United States in August 1991 to June 1992</p> <p>Inclusion criteria: Bladder outlet obstruction due to BPH, not in urinary retention</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ Physical status exceeding category III of the American Society of Anaesthesiologists ▪ Adenocarcinoma of the prostate ▪ Bladder neck to verumontanum length less than 2.4cm ▪ Life expectancy of < 6 months ▪ < 50 years ▪ Clinically significant illness ▪ Medication (hormonal therapy, alpha blockers, finasteride) that would have precluded participation in the study ▪ Medical condition (such as recent myocardial infarction, coagulopathy, recent stroke, sepsis) that investigators deemed unsuitable for one or more procedures <p>(the protocol had subsequently</p>	<p>Group 1- Laser coagulation</p> <p>Procedure: Nd; YAG laser, using Urolase fibre to the lateral lobes of the prostate at 3 and 9 o'clock positions for 60s each, and at 6 & 12 o'clock for 30s each, respectively.</p> <p>For patients with length of verumontanum and bladder neck >4 cm, treatment was repeated in 2 transverse planes, one just distal to the bladder and one just proximal to the verumontanum</p> <p>Average number of laser applications: 5.5±2.1 Cumulative duration of laser application: 4.2±1.5 minutes</p> <p>Power: 40W</p> <p>Energy: 5760-11520 J per patient, depending on prostate size.</p>	<p>AUA-6 symptom score</p>	<p>At 12 months, compared to baseline</p> <p>Group 1: -9.0 ±8.9, range -27 to 8</p> <p>Group 2: -13.3 ±7.5, range -29 to 7</p> <p>p value: <0.04</p>	<p>Funding: partially funded by CS Bard</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ The baseline AUA-6 was significantly lower for laser coagulation group. Statistical adjustment with ANCOVA reported ▪ Not stated which QoL instrument was used ▪ Impotence outcome- not certain if these are newly acquired cases ▪ Time point/period of complication measurement not stated <p>Additional outcomes:</p> <ul style="list-style-type: none"> ▪ Number of patients "non-serious" complications such as pain, hesitancy etc ▪ % of quality of life improved, at 12 months compared to baseline for Laser vs. TURP: 43/55 (78.2%) vs. 53/57 (93.0%) ▪ Post-op complications: (Bleeding (drop > 2.2g/dl of Hb in 24 hours post-procedure): 1/46 (2.2%) vs. 18/45 (40%). RR= 0.05 (95%
			<p>Post void residual volume, ml</p>	<p>At 12 months, compared to baseline</p> <p>Group 1: -55.4±124.3, range -425 to 220</p> <p>Group 2: 138.8±162.3 range -728 to 130</p> <p>p value: <0.01</p>	
			<p>Qmax, ml/s</p>	<p>At 12 months, compared to baseline</p> <p>Group 1: 5.3±6.9</p> <p>Group 2: 7.0±9.5</p> <p>p value: 0.27</p>	
			<p>Reoperation with VLAP or TURP (by 12months): 2 patients had VLAP: 1 patient had residual bladder neck tissue and later diagnosed with cancer. The other had residual apical lobe. 4 others had TURP.</p>	<p>Group 1: 2/56</p> <p>Group 2: 0/59</p> <p>p value: NS</p>	
			<p>Post-op complications: Blood transfusions</p>	<p>Group 1: 0/56 (0%)</p> <p>Group 2: 2/59(3.4%)</p> <p>p value: NS</p>	
			<p>Urinary retention</p>	<p>Group 1: 17/56 (30.4%)</p> <p>Group 2: 5/59 (8.5 %)</p> <p>Relative risk: 3.58(95% CI: 1.50, 9.00)</p> <p>p value: <0.005</p>	
<p>Urinary tract infection</p>	<p>Group 1: 3/56 (5.4%)</p> <p>Group 2: 1/59 (1.7%)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>changed to report patients with urinary retention, but these patients were not part of the cohort reported in this study)</p> <p>All patients N: 115</p> <p>Group 1-Laser coagulation N: 56 Dropouts: Age, mean (\pmSD): 65.8\pm6.7 **AUA – 6 symptom score, mean (\pmSD): 18.7\pm6.0 Prostate volume, ml:42.2\pm19.0 Qmax, ml/s: 8.9\pm3.6 Post void residual volume, ml: 162.7\pm126.6 Previous BPH therapy: 9/56(9.1%)</p> <p>Group 2 - TURP N: 59 Dropouts: Age, mean (\pmSD): 67.0\pm7.8 **AUA– 6 symptom score, mean (\pmSD): 20.8\pm4.8 Prostate volume, ml: 38.6\pm20.2 Qmax, ml/s: 9.5\pm5.2 Post void residual volume, ml: 206.7\pm181.9 Previous BPH therapy: 17/59(28.8%)</p>	<p>Anaesthesia: Spinal: 36/56 (64.2%) General: 20/56 (35.7%) Intravenous sedation only: 2(3.6%)</p> <p>Group 2 –TURP Procedure: Standard prostate resection using wire loop electrocautery under direct vision</p> <p>Anaesthesia: Spinal: 54/59(93.1%) General: 5/59(8.6%) Intravenous sedation only: 0/59(0%)</p> <p>For BOTH groups: Discharged when deemed medically fit, minimum of 24 hours hospitalisation post surgery for observation</p>	<p>Strictures (urethral and meatal stenosis): 6 patients in TURP group had urethral strictures. 1 patient in laser and 3 in TURP group had meatal stenosis</p> <p>Bladder neck contracture</p> <p>Incontinence</p> <p>Impotence (not stated how many were sexually active or whether these are newly acquired cases)</p> <p>Deep vein thrombosis</p> <p>Post TURP syndrome</p> <p>Clot retention</p> <p>Hospitalisation duration, days</p> <p>Duration of procedure, min</p>	<p>p value: NS</p> <p>Group 1: 1/56 (0%) Group 2: 9/59 (10.2%) RR: 0.12 (95% CI: 0.02, 0.67) p value: 0.02**</p> <p>Group 1: 0/56 (0%) Group 2: 3/59 (5.1%) p value: NS</p> <p>Group 1: 0/56 (0%) Group 2: 2/59 (3.4%) p value: NS</p> <p>Group 1: 3/56 (5.4%) Group 2: 2/59 (3.4%) p value: NS</p> <p>Group 1: 0/56 (0%) Group 2: 1/59 (1.7%) p value: NS</p> <p>Group 1: 0/56 (0%) Group 2: 2/59 (3.4%) p value: NS</p> <p>Group 1: 0/56 (0%) Group 2: 3/59 (5.1%) p value: NS</p> <p>Group 1: 1.8\pm1.1 Group 2: 3.1\pm0.9 p value: <0.01 **</p> <p>Group 1: 23.4\pm11.1 Group 2: 45.2\pm21.5 p value: <0.01 **</p>	<p>CI: 0.01-0.28), p value: <0.01 for Laser vs. TURP</p> <ul style="list-style-type: none"> Total number of patients with \geq1 serious complication, (impotence, UTI, meatal stenosis, urethral stricture, clot retention, bladder neck contracture, blood transfusions, TUR syndrome, incontinence, deep vein thrombosis, extravasation of irrigation fluid, prostatitis) was 6/56 in laser vs. 21/59 in TURP, RR = 0.30 (95% CI: 0.13, 0.66), p<0.01. <p>Notes: ** AUA-6 score was significantly lower in VLAP group. This required adjustment in data analysis using ANCOVA (analysis of covariance)</p> <p>**calculated by NCGC team using Fisher's exact test</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Donovan et al., 2000⁷⁴</p> <p>CLASP study-acute urinary retention</p> <p>Study design: RCT, multicentre, open label</p> <p>Setting: UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 7.5 months</p>	<p>Patient group: men with uncomplicated LUTS symptoms</p> <p>Setting: 3 centres in UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS score of ≥ 8, with physician and patient agreement that the symptoms require intervention Qmax <15ml/s when voided volume >200ml, <13ml/s when voided volume between 150-200ml and <10ml/s when voided volume between 100 to 149ml measured on two occasions, with the higher value between these two used for analysis >300ml post void volume urine on ultrasound <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or previous prostatic surgery; prostate size > 120ml; Life expectancy < 6 months; Urinary retention associated with recent operation, constipation or drugs which could cause acute urinary dysfunction, Neurogenic bladder dysfunction; Serum creatinine >250 $\mu\text{mol/L}$. <p>All patients N: 340</p> <p>Drop outs:</p>	<p>Group 1- Laser coagulation Procedure: Nd:YAG/ Non-contact VLAP, side-firing fibre (Bard Urolase), using standard fixed spot technique Power: 60W ND: YAG for 60s, depends on prostate size. For prostate size with urethral length of >25 mm, additional set of laser was used. If median lobe was present, 60W for 30s was applied for each side of lobe. Energy: 28684J Catheter protocol: Suprapubic catheter, removed when clinically appropriate. Other: All patients received antibiotic prophylaxis and anti-inflammatory suppository.</p> <p>Group 2 –TURP Procedure: Standard electroresection Catheter protocol: Suprapubic catheter.</p> <p>Group 3 – Conservative management Procedure: Men were given general advice and bladder training as deemed clinically</p>	<p>All cause mortality Not treatment related</p>	<p>Group 1: 5/117 Group 2: 0/117 Group 3: 1/106 p value: NS for all groups</p>	<p>Funding: Laser machines provided by Bard Diagnostics, Redmond, Washington.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Open label study, with main outcomes using patient reported measures. However, this paper specified that clinicians measuring outcomes were different from surgeons conducting the surgery <p>Additional outcomes:</p> <ul style="list-style-type: none"> Composite outcomes categories, and categorical outcomes for IPSS and Qmax <p>Notes: Sample size calculation performed Please see Chacko2001 for the acute urinary retention population of CLASP trial and Gujral 2000 for the chronic urinary retention population.</p>
			<p>IPSS, mean change from baseline (95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -10.8 (95% CI: -12.5,-9.0), n=96 Group 2: -12.3 (95% CI: -13.8,-10.7), n=89 Group 3: -1.3 (95% CI: -2.8,0.2), n=85 Adjusted difference: Group 1 vs. Group 2: -1.7 (95% CI: -3.6,0.1) p value: NS</p> <p>Statistically significant for surgical procedures vs. conservative</p>	
			<p>IPSS-QoL, mean (95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -1.9 (95% CI: -2.3, -1.6), n=93 Group 2: -2.2 (95% CI: -2.5, -1.8), n=85 Group 3: -0.4 (95% CI: -0.7, -0.1), n=85 Adjusted difference: Group 1 vs. Group 2: -0.2 (95% CI: -0.6,0.2) p value: NS</p>	
			<p>Qmax, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: 5.8 (95% CI: 4.5, 7.2), n=102 Group 2: 9.7 (95% CI: 7.7, 11.6), n=98 Group 3: 0.2 (95% CI: -0.4, 0.8), n=92 Adjusted difference: Group 1 vs. Group 2: 3.9 (95% CI:1.9, 5.8) p value: <0.05</p>	
		<p>Post void residual volume, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -73.4(95% CI:-91.3, -55.5), n=100 Group 2: -74.0 (95% CI:-89.2, -58.8), n=98 Group 3: 2.19 (95% CI:-23.1, -27.5, n=90 Adjusted difference:</p>		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 1-Laser coagulation N: 117 Dropouts: 1/117 Age, mean (\pmSD): 67.4\pm8.1 IPSS, mean (\pmSD): 19.1\pm6.6 IPSS-QoL, median(range): 4(2-6) Qmax, mean, (\pmSD): 10.4\pm2.9 Post void residual urine, mean, (\pmSD): 123.7\pm91.8 Prostate volume, mean, (\pmSD): 40.7\pm21.4 No obstructed (%): 90/117 (78.3) No equivocal and/or unobstructed (%): 25/117 (21.7)</p> <p>Group 2 - TURP N: 117 Dropouts: 2/117 Age, mean (\pmSD): 66.4\pm7.9 IPSS, mean (\pmSD): 19.2\pm6.7 IPSS-QoL, median(range): 4(0-6) Qmax, mean, (\pmSD): 10.3\pm2.7 Post void residual urine, mean, (\pmSD): 104.2\pm69.5 Prostate volume, mean, (\pmSD): 38.1\pm19.1 No obstructed (%): 91/117(78.4) No equivocal and/or unobstructed (%): 25/117(21.6)</p> <p>Group 3 – Conservative management N: 106 Dropouts: 5/106 Age, mean (\pmSD): 67.2\pm7.8 IPSS, mean (\pmSD): 18.8\pm6.5 IPSS-QoL, median(range): 4(1-6) Qmax, mean, (\pmSD): 9.9\pm2.7 Post void residual urine, mean, (\pmSD): 119.1\pm90.4</p>	appropriate		<p>Group 1 vs. Group 2: -13.4 (95% CI: -32.9, -6.1) p value: NS</p> <p>Post-op complications: Blood transfusion (units and criteria not stated) Group 1: 1/117 Group 2: 1/117 p value: NS</p> <p>Post-op complications: Perforation Group 1: 0/117 Group 2: 2/117 p value: NS</p> <p>Post-op complications: Septicaemia Group 1: 0/117 Group 2: 2/117 p value: NS</p> <p>Post-op complications: Urinary tract infection (symptomatic) Group 1: 3/117 Group 2: 2/117 p value: NS</p> <p>Time to catheter removal geometric mean, days Group 1: 2.2(95%CI 1.9 to 2.4) Group 2: 3.9(95%CI 3.7 to 4.2) Relative risk: 1.83 95% CI: 1.58 to 2.11 P value: <0.0001</p> <p>LOS, geometric mean (95% CI) days Group 1: 11.8(95%CI: 10.2 to 13.7) Group 2: 2.4 (95%CI: 2.1 to 2.9) Relative risk: 4.79 95% CI: 3.88 to 5.91 p value: <0.0001</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Prostate volume , mean, (\pm SD): 36.8 \pm 17.2 No obstructed (%): 82/106(77.4) No equivocal and/or unobstructed (%): 24/106(22.6)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gujral et al., 2000¹⁰⁷</p> <p>CLASP study-chronic urinary retention</p> <p>Study design: RCT, multicentre, open label</p> <p>Setting: UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 7.5 months</p>	<p>Patient group: men with chronic urinary retention</p> <p>Setting: 3 centres in UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS score ≥ 8, suggesting moderate to severe symptoms Low Qmax; < 15 ml/s when voided volume > 200 ml, < 13 ml/s when voided volume between 150-200 ml and < 10 ml/s when voided volume between 100 to 149 ml measured on two occasions, with the higher value between these two used for analysis > 300 ml post void volume urine on ultrasound <p>Exclusion criteria: CLASP criteria</p> <ul style="list-style-type: none"> Prostate cancer or previous prostatic surgery; prostate size > 120 ml; Life expectancy < 6 months; dysfunction; Neurogenic bladder Serum creatinine > 250 μmol/L. <p>Criteria specific to Chronic urinary retention group</p> <ul style="list-style-type: none"> Long term medication active on the lower urinary tract <p>All patients N: 82 Drop outs: 2</p>	<p>Group 1 - Laser coagulation Procedure: Nd:YAG/ Non-contact VLAP, side-firing fibre (Bard Urolase), using standard fixed spot technique Power: 60W ND: YAG for 60s, depends on prostate size. For prostate size with urethral length of > 25 mm, additional set of laser was used. If median lobe was present, 60W for 30s was applied for each side of lobe. Energy: 33.8kJ or 0.94kJ/ml of prostate tissue</p> <p>Catheter protocol: Suprapubic catheter, removed when clinically appropriate. Other: All patients received antibiotic prophylaxis and anti-inflammatory suppository.</p> <p>Group 2 –TURP Procedure: Standard electroresection</p>	<p>All cause mortality Not treatment related</p>	<p>Group 1: 0/38 Group 2: 1/44 p value: NS</p>	<p>Funding: Laser machines provided by Bard Diagnostics, Redmond, Washington.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Open label study, with main outcomes using patient reported measures. However, this paper specified that clinicians measuring outcomes were different from surgeons conducting the surgery <p>Additional outcomes:</p> <ul style="list-style-type: none"> Composite outcomes categories, and categorical outcomes for IPSS and Qmax <p>Notes: Sample size calculation performed, to detect 30% differences in binary outcomes and SD of 0.63 for continuous outcomes at a power of 80%</p>
			<p>IPSS, mean change from baseline (95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -12.2 (95%CI: -15.7, -8.7), n=29 Group 2: -14.2, (95% CI: 17.2,-11.2), n=33 Adjusted difference: -3.6 (95%CI-7.2 to -0.1) p value: 0.048</p>	
			<p>IPSS-QoL, mean (95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -2.8(95%CI: -3.4, -2.1), n=30 Group 2: -3.2(95%CI: -3.9, -2.6) ,n=33 Adjusted difference: -0.6(95% CI:-1.3 to 0.1) p value: NS</p>	
			<p>Qmax, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: 5.7 (95%CI: 2.6, 8.8), n=33 Group 2: 9.4 (95%CI: 6.5, 12.2) ,n=40 Adjusted difference: 1.1 (95%CI: -3.0 to 5.3) p value: NS</p>	
			<p>Post void residual volume, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA</p>	<p>Group 1: -329 (95%CI: -377, -281), n=33 Group 2: -464(95%CI: -553, -374) ,n=40 Adjusted difference: -27.5 (95%CI: -68.1 to 13.0) p value: NS</p>	
			<p>Post-op complications: Confusion (TUR syndrome)</p>	<p>Group 1: 0/38 Group 2: 1/44 p value: NS</p>	
			<p>Post-op complications: Blood transfusion (units and criteria not stated)</p>	<p>Group 1: 0/38 Group 2: 3/44 p value: NS</p>	
			<p>Post-op complications: Heavy bleeding (4 no</p>	<p>Group 1: 0/38 Group 2: 6/44 p value: NS</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 1-Laser coagulation N: 38 Dropouts: 2/38 Received as allocated: 30 Age, mean (\pmSD): 70.2\pm6.8 IPSS, mean (\pmSD): 20.9\pm6.4 IPSS-QoL, , mean, (\pmSD): 5.0\pm2.6 Prostate volume, mean, (\pmSD): 40.7\pm19.9 Qmax, mean, (\pmSD): 11.2\pm5.3 Post void residual urine, mean, (\pmSD): 438\pm151</p> <p>Group 2 - TURP N: 44 Dropouts: 0 Received as allocated: 44 Age, mean (\pmSD): 70.6\pm5.8 IPSS, mean (\pmSD): 19.5\pm7.2 IPSS-QoL, mean, (\pmSD): 4.5\pm2.6 Prostate volume, mean, (\pmSD): 49.7\pm21.8 Qmax, mean, (\pmSD): 8.5\pm3.6 Post void residual urine, mean, (\pmSD): 545\pm275</p>		termination, 2 cases termination		Please see Chacko2001 for the acute urinary retention population of CLASP trial and Donovan2000 for the uncomplicated LUTS symptom population.
			Post-op complications: Perforation	Group 1: 0/38 Group 2: 1/44 p value: NS	
			Post-op complications: Septicaemia	Group 1: 1/38 Group 2: 3/44 p value: NS	
			Post-op complications: Urinary tract infection (symptomatic)	Group 1: 1/38 Group 2: 2/44 p value: NS	
			Post-op complications: Reoperation (performed resection after laser therapy due to "unacceptable levels of symptoms")	Group 1: 3/38 Group 2: 0/44 p value: NS	
			Time to catheter removal geometric mean, days	Group 1: 25.5(95%CI 20.2 to 28.3) Group 2: 3.0 (95%CI 2.3 to 3.9) Relative risk: 8.62 95% CI: 6.04, 12.29 p value: <0.0001	
			LOS, geometric mean (95% CI) days	Group 1: 2.2(95%CI 1.7 to 2.8) Group 2: 4.4(95%CI 3.9 to 4.9) Relative risk: 2.01 95% CI: 1.54 to 2.61 P value: <0.0001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kursh et al., 2003¹⁵⁶</p> <p>Study design: RCT, open label</p> <p>Setting: US, tertiary care hospitals</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: Bladder outflow obstruction secondary to BPH</p> <p>Setting: six US tertiary care hospitals between Nov 1997 and Feb 1999</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> AUASI ≥ 13 Qmax < 1.5 ml/s for 2 s with an adequately filled bladder PVR between 30 and 300 ml Prostatic length ≥ 1.5 cm Prostatic volume ≤ 75 cm³ <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Any condition or history of illness or surgery which may pose additional risk to the patient such as unstable angina, significant renal impairment (creatinine > 1.8 mg/dL), or poorly controlled diabetes mellitus. History of prostate cancer; suspected prostate cancer (based on digital rectal examination or PSA level > 4 ng/mL) – must be ruled out with biopsy Acute urinary retention Acute or chronic prostatitis, cystolithiasis, neurogenic bladder, bladder neck contracture, or active urinary tract infection. Taking terazosin, doxazosin or tamsulosin within 14 days of 	<p>Group 1- Laser coagulation Performed with the Indigo 830e (830nm) laser system.</p> <p>Procedure: Slightly flexible laser fibre was inserted through the urethra and into the prostate using a standard cystoscope. A 1-cm long diffuser tip radiates heat in all directions at a low power (20W). The heat produces an olive-shaped area of coagulation necrosis about 2 x 2.5 cm or a volume of approximately 4 cm³.</p> <p>Power: 20W</p> <p>Energy: NR</p> <p>Catheter protocol: patients discharged with catheter in place, which was usually removed in 1 week.</p> <p>Other:</p>	<p>AUASI score, median:</p>	<p><u>At 6 months</u> Group 1: 7.0 Group 2: 6.0 Difference: 1.0 (95% CI: -3.0 to 3.0) p value: Not sig</p> <p><u>At 24 months</u> Group 1: 9.0 Group 2: 7.0 Difference: 2.0 (95% CI: -3.0 to 4.0) p value: Not sig</p>	<p>Funding: Indigo Medical Inc (the laser system manufacturer). First author a paid consultant of the parent company (Ethicon Endo-Surgery)</p> <p>Limitations:</p> <ul style="list-style-type: none"> Patient reported outcomes methods were not clearly reported. It was unclear which questionnaires were used to evaluate QoL and sexual function. Only point estimates (median) were reported for continuous variables. Only 61% (73/120) of targeted sample size was recruited. Enrolment stopped early because of low patient participation. <p>Additional outcomes:</p> <ul style="list-style-type: none"> Median prostate volume and PSA level post surgery were reported.
			<p>Qmax (ml/s), median</p>	<p><u>At 6 months</u> Group 1: 14.3 Group 2: 16.6 Difference: -2.3 (95% CI: -0.4 to -6.5) p value: < 0.05</p> <p><u>At 24 months</u> Group 1: 13.9 Group 2: 16.5 Difference: -2.6 (95% CI: -7.6 to 0.4) p value: Not sig</p>	
			<p>Post-void residual volume (ml), mean \pm SD (note that the baseline value was significantly different)</p>	<p><u>At 6 months</u> Group 1: 42.4 Group 2: 46.0 Difference: -3.6 (95% CI: -12.6 to 27.3) p value: NS</p> <p><u>At 24 months</u> Group 1: 57.7 Group 2: 44.0 Difference: 13.7(95% CI: -15.2 to 40.3) p value: NS</p>	
			<p>Post-op complications: Blood transfusion</p>	<p>Group 1: 0/37 Group 2: 0/35 p value: NS</p>	
<p>Post-op complications: Development of</p>	<p>Group 1: 0/37 Group 2: 2/35</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>enrolment; finasteride or phytotherapy and anticholinergic within one month of enrolment.</p> <p>All patients N: Age, range, years: 50-81 Drop outs: 1 patient withdrew consent before treatment group assignment</p> <p>Group 1-Laser coagulation N: 37 Dropouts: Age, mean (years): 67.6 Ethnicity, white (%): 30/37 (81%) AUASI, median: 24.0 Qmax, median (ml/s): 9.2 PVR, median (ml): 81 PSA, median (ng/ml): 2.3 Prostate volume, median (cm³):41.5</p> <p>Group 2 - TURP N: 35 Dropouts: Age, mean: 69.3 Ethnicity, white (%): 29/35(83%) AUASI, median: 23.0 Qmax, median (ml/s): 9.1 PVR, median (ml): 87.5 PSA, median (ng/ml): 2.3 Prostate volume, median (cm³): 40</p>	<p>Usually performed as an outpatient procedure.</p> <p>Anaesthesia: general/spinal/topical: 17/15/5</p> <p>Group 2 –TURP Procedure: Standard radiofrequency monopolar loop procedure</p> <p>Catheter protocol: Generally removed one day post-operatively, before discharge</p> <p>Others: Anaesthesia: general/spinal/topical: 11/24/0</p> <p>Both groups: received antibiotics – choice at discretion of individual investigators</p>	<p>anaemia (hematocrite less than 30%)</p> <p>Post-op complications: reoperation (2 patients retreated within 6 months, 1 with ILC and 1 with TURP. 4 additional patients receive TURP within 1 year)</p> <p>Post-op complications: Incontinence (1 case of urge incontinence and another case of stress incontinence requiring pads)</p> <p>LOS, median (range), (days)</p> <p>Sexual function score (Name of questionnaire not provided. Stated that the range was 0-30, higher scores better)</p>	<p>p value: NS</p> <p><u>At 6 months</u> Group 1: 2/37 Group 2: 0/35 Relative risk: NE p value:: NS</p> <p><u>At 12 and 24 months</u> Group 1: 6/37 Group 2: 0/35 Relative risk: NE p value: 0.02</p> <p>Group 1: 0/37 Group 2: 2/35 Relative risk: 0 (0-1.77) p value:: NS</p> <p>Group 1: 7.0 (3 to 145) Group 2: 33.5 (10 to 120) p value: NR</p> <p><u>At 6 months</u> Group 1: 19.0 Group 2: 5.0 Difference: 14.0 (95% CI: 3.0 to 14.0) p value: <0.05</p> <p><u>At 24 months</u> Group 1: 19.5 Group 2: 10.0 Difference: 9.5 (95% CI: -1.0 to 12.0) p value: Not sig</p>	<p>▪ “Problems from Symptom Index” score and “American Urological Association QoL Assessment” score were reported. However, it what unclear which questionnaire were used from the paper. There was no significant difference between treatment arms in these outcomes.</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Liedberg et al., 2003¹⁶⁶</p> <p>Study design: RCT, open label</p> <p>Setting: Hospital, Sweden</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 1 year</p>	<p>Patient group: moderate to severe BPH</p> <p>Setting: Department of urology, hospital in Sweden, Dec 1997 to Feb 2000</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 12 Qmax \leq 15ml/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Indwelling urinary catheter Prostatic carcinoma Clinical suspicion of neurogenic bladder disturbance <p>All patients N: 38</p> <p>Drop outs: 7/38 (3 due to prostate cancer), one was randomised to ILC but received TURP; 1 did not wish to undergo surgery and 2 could not undergo surgery due to undercurrent illness.</p> <p>Group 1-Laser coagulation N: 20</p> <p>Drop outs: Not stated</p> <p>IPSS, median (IQR): 19(16-24)</p> <p>Qmax, median (IQR): 8(7-10) [n=19]</p> <p>Prostate volume, median (IQR): 49(41-75)</p> <p>Post void residual volume: median (IQR): 96(64-190)</p> <p>Group 2 - TURP N: 11</p> <p>Dropouts: Not stated</p>	<p>Group 1- Laser coagulation Procedure: Performed with the Indigo 830e (830nm) laser system.</p> <p>Each puncture site was treated for 3 min with a target temperature of 85C. The prostate was punctured under visual control and the target was one puncture for every 4ml of prostate.</p> <p>Power setting not stated.</p> <p>Catheter protocol: suprapubic catheter, removed when PVR <150ml</p> <p>Others: Norfloxacin 400mg twice daily while catheter was in place</p> <p>Group 2 –TURP Procedure: Standard electroresection.</p>	<p>IPSS, median (IQR):</p> <p>Qmax (ml/s), median (IQR):</p> <p>Post void residual volume (ml), median (IQR):</p> <p>Post-op complications: Clot retention (requiring transurethral clot evacuation under general anaesthesia)</p> <p>Peri-operative complications: Bleeding (blood loss, median (IQR), (ml))</p> <p>Post-op complications: Catheterisation</p> <p>Post-op complications: urinary tract infections</p>	<p>At 3 months Group 1: 10(4-15), n=20 Group 2: 4(2-7), n=11 p value: NS</p> <p>At 12 months Group 1: 11(6-14), n=19 Group 2: 6(3-10), n=9 p value: NS</p> <p>At 3 months Group 1: 11(8-15), n=19 Group 2: 12(9-18), n=10 p value: NS</p> <p>At 12 months Group 1: 11(6-12), n=18 Group 2: 14(10-19), n=9 p value: NS</p> <p>At 3 months Group 1: 74(38-140), n=19 Group 2: 0(0-53), n=10 p value: NS</p> <p>At 12 months Group 1: 126(25-190), n=19 Group 2: 22(3-62), n=8 p value: NS</p> <p>Group 1: 1/20 Group 2: 0/11 p value: NS</p> <p>Group 1: 0(0-50) Group 2: 350(200-514) p value: <0.001</p> <p>Group 1: 24(14-34) Group 2: 2(1-2) p value: <0.001</p> <p>Group 1: 13/20 Group 2: 1/11 p value: <0.007</p>	<p>Funding: Partly finance by FroU-Kronoberg</p> <p>Limitations:</p> <ul style="list-style-type: none"> Open label study with subjective patient reported outcomes. Study stopped early (targeted N=50) due to prolonged rate of catheterisation and high rate of UTI Large number of exclusions from TURP group resulted in imbalance of sample <p>Additional outcomes: Prostate volume post operation</p> <p>Notes: Age of subjects not reported</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>IPSS, median (IQR): 17(17-24) Qmax, median (IQR): 8(6-9) [n=10] Prostate volume, median (IQR):47(37-61) Post void residual volume: median (IQR): 117(67-200)</p>		<p>Post-op complications: urethral stricture</p> <p>Post-op complications: bladder neck stenosis</p> <p>Post-op complications: Retrograde ejaculation</p> <p>Hospitalisation, median (IQR), (days):</p>	<p>Group 1: 0/20 Group 2: 0/11 p value: NS</p> <p>Group 1: 0/20 Group 2: 0/11 p value: NS</p> <p>Group 1: 1/20 Group 2: 3/11 p value: NS (0.084)</p> <p>Group 1: 2.5 (0.25 to 3.8) Group 2: 3 (3 to 4) p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Martenson et al., 1999¹⁸³</p> <p>Study design: RCT, open label</p> <p>Setting: Netherlands</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: BPH patients</p> <p>Setting: Department of Urology, University Hospital Nijmegen, Netherlands Oct 1994 to April 1996</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ▪ Prostate volume >25 cm³ ▪ age >45 years ▪ Duration of symptoms > 3 months ▪ IPSS12 ▪ Peak uroflow <15ml/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ Prostate carcinoma ▪ Bacterial prostatitis ▪ Urethral stricture ▪ Neurogenic bladder dysfunction ▪ Urinary tract infection ▪ Use of drugs influencing bladder function ▪ History of TURP ▪ Diabetes mellitus ▪ Bladder residual urine >350ml <p>All patients N: 44 Mean age: NR Drop outs: NR</p> <p>Group 1-Laser coagulation N: 30 IPSS, mean \pmsd: 21.7\pm6.1 IPSS-QoL, mean \pmsd: 4.1\pm1.4 Qmax, mean\pmsd, (ml/s):7.3\pm3.8 PVR, mean\pmsd, (ml):116\pm146 Normal erectile function: 28/30</p>	<p>Group 1- Laser coagulation Procedure: Performed with the Indigo 830 (830nm) laser system.</p> <p>Each individual fibre placement received 1420 J in a standard for 4 min treatment cycle</p> <p>Power: 10 W, decreased to 5 W</p> <p>Catheter protocol: Suprapubic catheters were removed when adequate voiding was demonstrated at scheduled follow up (1, 2 or 4 weeks)</p> <p>Group 2 –TURP Procedure: Standard procedure. 24Fr resectoscope used in combination with glycine irrigation fluid.</p> <p>Catheter protocol: Removed according to individual needs</p>	IPSS, mean \pm sd	<p><u>At 3 months (12 weeks)</u> Group 1: 11.8\pm6.9 Group 2: 4.7\pm4.0 p value: NS</p> <p><u>At 6 months (26 weeks)</u> Group 1: 10.3\pm5.4 Group 2: 3.8\pm2.4 p value: NS</p> <p><u>At 12 months (52 weeks)</u> Group 1: 12.4\pm7.7 Group 2: 3.5\pm2.9 p value: NS</p> <p><u>At 24 months (104 weeks)</u> Group 1: 12.0\pm4.9 Group 2: 5.0\pm4.4 p value: NS</p>	<p>Funding: Indigo- the laser manufacturer</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Small sample size, with no power calculation provided ▪ Patient age not reported ▪ T-tests were used <p>Additional outcomes: The paper also reported the results of another non-randomised phase II study which temperature-sensing laser system</p> <p>Notes: The patients were randomised 2:1 in this study.</p>
			IPSS-QoL, mean \pm sd	<p><u>At 3 months (12 weeks)</u> Group 1: 2.3\pm1.4 Group 2: 0.9\pm1.3 p value: NS</p> <p><u>At 6 months (26 weeks)</u> Group 1: 2.2\pm1.4 Group 2: 0.5\pm0.7 p value: NS</p> <p><u>At 12 months (52 weeks)</u> Group 1: 2.2\pm1.5 Group 2: 0.6\pm0.8 p value: NS</p> <p><u>At 24 months (104 weeks)</u> Group 1: 2.2\pm1.5 Group 2: 0.7\pm0.9 p value: NS</p>	
			Qmax, mean \pm sd, (ml/s):	<p><u>At 3 months (12 weeks)</u> Group 1: 12.5\pm5.4 Group 2: 25.8\pm9.7 p value: NS</p> <p><u>At 6 months (26 weeks)</u></p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 - TURP N: 14 IPSS, mean \pmsd: 21.6\pm7.7 IPSS-QoL, mean \pmsd: 4.0\pm1.3 Qmax, mean\pmsd, (ml/s):9.3\pm3.2 PVR, mean\pmsd, (ml):88\pm126 12/14</p>		<p>PVR, mean\pmsd, (ml):</p>	<p>Group 1: 11.1\pm4.5 Group 2: 18.2\pm6.6 p value: NS <u>At 12 months (52 weeks)</u> Group 1: 11.9\pm5.5 Group 2: 25.7\pm11.1 p value: NS <u>At 24 months (104 weeks)</u> Group 1: 10.3\pm4.4 Group 2: 20.1\pm13.7 p value: NS</p> <p><u>At 3 months (12 weeks)</u> Group 1: 58\pm103 Group 2: 12\pm19 p value: NS <u>At 6 months (26 weeks)</u> Group 1: 60\pm56 Group 2: 14\pm27 p value: NS <u>At 12 months (52 weeks)</u> Group 1: 59\pm77 Group 2: 14\pm21 p value: NS <u>At 24 months (104 weeks)</u> Group 1: 94\pm128 Group 2: 63\pm100 p value: NS</p> <p>Post-op complications: Blood transfusion Group 1: 0/30 Group 2: 0/14 p value: NS</p> <p>Post-op complications: Clot retention Group 1: 0/30 Group 2: 0/14 p value: NS</p> <p>Post-op complications: In continence (up to 24 months), definition of incontinence not provided Group 1: 0/30 Group 2: 0/14 p value: NS</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Post-op complications: urinary tract infections	Group 1: 10/30 Group 2: 4/14 RR: 4.67(95% CI : 0.94 to 27.8) p value: NS	
			Post-op complications: Reoperation (up to 24 months)	Group 1: 6/30 Group 2: 1/14 RR: 2.8(95%CI: 0.51 to 17.5) p value: NS	
			Post-op complications: Retrograde ejaculation	Group 1: 0/30 Group 2: 3/14 p value: NS (0.084)	
			Length of catheterisation, mean \pm sd (days)	Group 1: 27 \pm 23 Group 2: 3 \pm 1	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Rodrigo Aliaga et al., 1998 ²⁵³ (data extracted from HTA report) Study design: Setting: Spain Evidence level: 1+ Duration of follow-up: 6 months	Patient group: <ul style="list-style-type: none"> patients with BPH Inclusion criteria: <ul style="list-style-type: none"> prostate size 20–60 g; symptom score; IPSS score \geq 15 Exclusion criteria: <ul style="list-style-type: none"> age < 50 years All patients N: 41 Drop outs: Group 1 –TUIP N: 20 Age, years, mean\pmsd (range): NR Residual volume, mean \pm SD (ml): 89 \pm 92 Group 2 -TURP N: 21 Age, years, mean\pmsd (range): NR Residual volume, mean \pm SD (ml): 146 \pm 133	Group 1- TUIP/BNI Group 2 - TURP All Patients left hospital 24–72 hours postoperatively if no complications	IPSS score, mean \pm SD	Baseline Group 1: 24.2 \pm 7.7 Group 2: 24.4 \pm 10.3 3 months Group 1: 4.3 \pm 4.5 Group 2: 4.8 \pm 4.8 6 months Group 1: 5.7 \pm 6.2 Group 2: 3.7 \pm 3.8	Funding: NR Limitations: <ul style="list-style-type: none"> No information of randomisation allocation and concealment methods Baseline prognostic factors were reported as not equal in quality assessment (uncertain which factor this referred to) Additional outcomes: Irritative symptoms Quality of life score (WHO) Length of hospital stay Catheter duration Residual volume Notes: None.
			Qmax, ml/s, mean \pmsd (range)	Baseline Group 1: 8.7 \pm 5.5 Group 2: 8.3 \pm 4.5 3 months Group 1: 22 \pm 12.2 Group 2: 18.6 \pm 8.5 6 months Group 1: 20.6 \pm 8.7 Group 2: 20.6 \pm 10.1	
			Blood transfusion	Group 1: 0/20 Group 2: 1/21 P value: Not sig	
			Reoperation	Group 1: 1/20 Group 2: 1/21 P value: Not sig	
			Retrograde ejaculation	Group 1: 14/20 Group 2: 15/21	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Sengor et al., 1996²⁷¹</p> <p>Study design: RCT, open label</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Symptomatic bladder outlet obstruction due to BPH referred to urology clinic</p> <p>Setting: urology clinic, single-centre, Istanbul, Turkey</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Significant voiding symptoms to request therapy Qmax ≤15 ml/s and Qave ≤ 10 ml/s from uroflowmetric volume of ≥ 150 ml Age >50 years <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Prostate cancer- Induration or nodularity of prostate on DRE or PSA > 4.0 mg/ml further examined for cancer. Infections (treated with suitable antibiotics preoperatively) <p>All patients N: 60 Age: 50-85 Drop outs: NR</p> <p>Group 1 - Laser N: 30 Mean age (yrs): 66 (range 50-85) Drop outs: Erectile dysfunction: 7/30 AUA, mean ± SD: 21.8 ± 7.6 Prostate volume (TRUS) ml: 55 (30-80)</p>	<p>Group 1 Under spinal or general anaesthesia Ultraline side firing Nd:YAG laser fibre 600µm using SMA-905 adapter and standard Nd:YAG laser generator at 60W through 21F cystoscope. Bladder was continuously irrigated with saline. No indwelling catheter was used but supra public tubes were clamped 4-5 days after treatment and removed after successful urination.</p> <p>Group 2 TURP in standard manner under spinal anaesthesia using Storz 26F resectoscope with mannitol solution for irrigation. A 3-way Foley catheter was inserted and bladder irrigated with normal saline for 24-48 h.</p> <p>Examination methods: Patients followed at 3 and 6 months using AUA symptom score, Qmax</p>	<p>AUA score, mean ± SD:</p>	<p><u>At 3 months</u> Group 1: 8.5±4.2 Group 2: 9.8±3.1 p value: NS (P=0.17), calculated by NCGC team using t-tests. Reported as 0.034</p> <p><u>At 6 months</u> Group 1: 7.8±2.6 Group 2: 9.3±4.2 p value: NS (P=0.1), calculated by NCGC team using t-tests</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Outcome assessment was not masked. Randomisation and allocation method not reported. Statistical methods and sample size calculation not reported Baseline values of post void residual volume significantly different between groups. <p>Additional outcomes: % of mean change was reported for AUA score, Qmax and residual volume but standard deviations were not provided</p> <p>Notes: None.</p>
			<p>Qmax (ml/s), mean ± SD:</p>	<p><u>At 3 months</u> Group 1: 18.9±3.1 Group 2: 20.7±2.6 p value: 0.01, calculated by NCGC team using t-tests. Reported as 0.025</p> <p><u>At 6 months</u> Group 1: 18.2±2.1 Group 2: 19.8±2.5 p value: <0.01, calculated by NCGC team using t-tests, reported as NS</p>	
			<p>Post void residual volume (ml), mean ± SD (note that the baseline value was significantly different)</p>	<p><u>At 3 months</u> Group 1: 50.4±30 Group 2: 70±27 p value: NS</p> <p><u>At 6 months</u> Group 1: 47±19 Group 2: 68±22 p value: NS</p>	
			<p>Post-op complications: Transurethral resection syndrome</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NS</p>	
<p>Post-op complications: Blood transfusion (units and criteria not stated)</p>	<p>Group 1: 0/30 Group 2: 2/30 p value: NS</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>*PVR mean \pm SD: 110 \pm 68 Qmax mean \pm SD (ml/s): 8.7 \pm 2.3</p> <p>Group 2 - TURP N: 30 Mean age (yrs): 61 (55-70) Drop outs: Erectile dysfunction: 3/30 AUA, mean \pm SD: 22.1 \pm 2.6 Prostate volume (TRUS) ml: 47 (30-50) *PVR, mean \pm SD: 155 \pm 40 Qmax, mean \pm SD (ml/s): 8.4 \pm 2.8</p> <p>*P =0.003,calculated by t-test by NCGC team</p>	and PVR measurements	<p>Post-op complications: urethral strictures (6 months follow up)</p> <p>Post-op complications: Retrograde ejaculation (6 months follow up)</p> <p>Operation time, mean (range), (min):</p> <p>LOS, mean (range), days</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NS</p> <p>Group 1: 1/23 (3%) Group 2: 24/27 (80%) Relative risk:: 0.05 (95% CI: 0.01-0.19) p value: <0.001</p> <p>Group 1: 43 (15-70) Group 2: 56 (45-90) P value : NR</p> <p>Group 1: 1.6 (1-3) Group 2: 5.9 (4-7) P value : NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Suvakovic et al., 1996 ²⁹⁰ Study design: RCT, open label Evidence level: 1+ Duration of follow-up: 1 year	Patient group: Consecutive patients with prostatic symptoms Setting: Urology department, South Cleveland University, UK Inclusion Criteria: <ul style="list-style-type: none"> • Qmax ≤15mL/s for a voided volume of ≥150 mL • Age • Significant voiding symptoms (AUA score >15) • PSA level <2.5 ng/mL • Prostate volume <40g (assessed by TRUS, DRE and cystoscopy) • Length of the prostatic urethra >4 cm Exclusion Criteria: <ul style="list-style-type: none"> • Malignancy All patients N: 40 Group 1 - VLAP – side fire free beam alone N: 10 Age (mean): 67.5(8.7) IPSS: 15.7(5.1) Qmax ml/s: 10.5 (3.7) Residual Vol mL: 47.4(48.1)	Group 1: VLAP – side fire free beam alone 4 spot thermocoagulation at the 10, 2, 4 and 8 o'clock positions. Laser delivered at 60W for 60s. Group 2 : CLAP- contact laser alone Nd: YAG laser applied at 40W for vaporising and coagulating the prostate with a minimum depth of penetration. A 16 F two – way catheter was inserted into the bladder and removed after 24 h. Group 3 : Hybrid – side fire free beam and debridement As in VLAP, plus debridement of coagulated tissue using a 26F continuous irrigating resectoscope. At the end of the procedure, a 16 F two –way catheter was inserted into the bladder and removed after 24 h Group 4 : TURP	IPSS symptom score, mean±sd. Values for 12 months follow up reported in paper, but n was not reported	At 3 months Group 1: 16.8±15.0, n=10 Group 2: 9.7±2.6, n=10 Group 3: 8.1±5.4, n=8 Group 4: 12.8±5.9, n=10 P value: NS# P value for Group 1 vs. Group 3 was reported to be <0.01 in paper, but this could not be repeated. At 6 months Group 1: 16.2±4.2, n=9 Group 2: 18.7±7.5, n=9 Group 3: 19.4±3.4, n=4 Group 4: 19.0±0.8, n=10 P value: NS#	Funding: NR Limitations: <ul style="list-style-type: none"> • Small sample size, n of 10 in each arm • Unclear which statistical test was used for data – discrepancies in the stat sig reported for AUA score for 3 months and calculated by NCGC team. • Number of participants followed up at 12 months not reported. Additional outcomes: Operation duration for each procedure Notes: # values calculated by NCGC team based on mean and sd reported. It was not possible to calculate using Kruskal Wallis test without the
			Qmax ml/s, mean±sd Values for 12 months follow up reported in paper, but n was not reported	At 3 months Group 1: 14.8±5.4, n=10 Group 2: 15.6±13.5, n=10 Group 3: 15.1±7.3, n=8 Group 4: 17.8±3.8, n=10 P value: NS At 6 months Group 1: 16.2±4.2, n=9 Group 2: 18.7±7.5, n=9 Group 3: 19.4±3.4, n=4 Group 4: 19.0±0.8, n=10 P value: NS#	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Prostate size (by TRUS), g: 23.6(6.4) PSA (ng/mL): 2.3(0.8)</p> <p>Group 2 - CLAP- contact laser alone N: 10 Age (mean): 62.6(5.8) IPSS: 18 (6.0) Qmax ml/s: 12.2 (3.8) Residual Vol mL: 139.6(103) Prostate size (by TRUS), g: 24(5.8)</p> <p>Group 3 - Hybrid – side fire free beam and debridement N: 10 Age (mean): 64.1(6.9) IPSS: 17(6.0) Qmax ml/s: 11.8(4.1) Residual Vol mL: 68.3(64) Prostate size (by TRUS), g: 27(12.3)</p> <p>Group 4 - CLAP- TURP Standard resection N: 10 Age (mean): 66.1(5.1) IPSS: 18.8 (4.5) Qmax ml/s: 11.1(6.4) Residual Vol mL: 161.8(104) Prostate size (by TRUS), g: 22(5)</p>	<p>Standard resection using a 26 F continuous irrigating resectoscope. A 22 F three-way urethral catheter was inserted into the bladder and irrigation was continued up to 24 h. The catheter was removed after 48 h and the patients discharged home 3-4 days after the procedure.</p>	<p>Catheter duration, mean, hours (range or standard deviations not reported)</p> <p>Length of hospitalisation, (hours)</p>	<p>Group 1: 24, n=10 Group 2: 24, n=10 Group 3: 20, n=10 Group 4: 48, n=10 p value: reported as <0.05 between group 4 and “lasers”</p> <p>Group 1: 30, n=10 Group 2: 30, n=10 Group 3: 24, n=10 Group 4: 84, n=10 p value: reported as <0.05 between group 4 and “lasers”</p>	<p>raw data.</p> <p>All patients received preoperative oral antibiotics and controlled for more than 5 days post-operatively.</p>

Evidence Table 27: Laser vaporisation vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bouchier-Hayes et al., 2006³²</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 weeks</p>	<p>Patient group: Patients referred with LUTS to urology outpatient department</p> <p>Setting: single centre, Melbourne, Australia</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Age >50 years Referral by GP Flow rate ≤ 15 mL/s IPSS ≥ 12 Gland 15-85 cm³ on TRUS Obstructed Abrams-Griffiths (A-G) nomogram Able to complete QoL, Bother Score & Baseline Sexual Function Questionnaire (BSFQ) questionnaires Able to give informed consent <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Known or suspected prostate cancer Chronic retention Taking α-blocker or herbal remedy On anticoagulants On finasteride or dutasteride <p>All patients N: 95 Drop outs: 19 (25%)*</p> <p>Group 1 - Laser N: 38 Mean age (yrs): 65.2 range (51-81) Drop outs: NR* IPSS: NR</p>	<p>Group 1 Photoselective vaporisation was performed using 80W KTP using Greenlight laser system and StarPulse quasi-continuous wave laser (Laserscope) emitting green light at 532 nm. A 600 μm laser fibre with 70° lateral deflecting quartz element used through continuous flow cystoscope with saline irrigation. Catheters left situ at the discretion of the surgeon.</p> <p>Group 2 TURP in standard manner through 25F resectoscope sheath using ValleyLab diathermy machine with 3-way 22F Foley catheter on continuous saline irrigation</p> <p>Intervention performed by registrars in training or fellows in the department, all of whom had performed <5 laser prostatectomies each</p>	Change IPSS symptom score from baseline at 6 weeks**	Group1: 14.0 ± 9.8 (n=38) Group 2: 12.9 ± 10.6 (n=38) p value: Not Signif. (NCGC calculated p=0.63)	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Baseline values for Qmax and IPSS, QoL, bother and BSFQ not reported **Follow up period not clear for main outcome data or complications. Might be 6 weeks as number of patients with data at 6 weeks is 76 Outcome assessment was not masked. Randomisation method not reported. Allocation concealment not reported <p>Notes: 12 months data in publication at October 2008</p>
			Change in flow rate (Qmax) from baseline at 6 weeks**	Group1: 11.96 ± 8.23 (n=38) Group 2: 8.56 ± 9.08 (n=38) p value: Not Signif. (NCGC calculated p=0.09)	
			Change in QoL score from baseline at 6 weeks**	Group1: 2.65 ± 2.1 (n=38) Group 2: 2.91 ± 2.04 (n=38) p value: Not Signif.	
			Change in bother score from baseline at 6 weeks**	Group1: 2.65 ± 2.1 (n=38) Group 2: 1.61 ± 1.22 (n=38) p value: Not Signif.	
			Change in prostate volume from baseline at 6 weeks**	Group1: 125 ± 198 (n=38) Group 2: 86 ± 124.38 (n=38) p value: Not Signif.	
			Post-op complications Failure to void: (follow up period 6 weeks**)	Group1: 4/38 Group 2: 3/38 p value: NR	
			Post-op complications Stricture: (follow up period 6 weeks**)	Group1: 0/38 Group 2: 5/38 p value: NR	
			Post-op complications urine retention: (follow up period 6 weeks**)	Group1: 3/38 Group 2: 1/38 p value: NR	
			Post-op complications number of patients with blood transfusion (follow up period 6 weeks**)	Group1: 0/38 Group 2: 1/38 p value: NR	
			Post-op complications	Group1: 2/38	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Erectile dysfunction: NR Prostate volume (TRUS) ml: 42.4 range (16.5-82.6) Qmax: NR Operation time: 30.2 mins range (9-70) Mean catheterisation time (days): 0.5 ± 0.4 Mean length of stay (days): 1.1 ± 0.3</p> <p>Group 2 - TURP N: 38 Mean age (yrs): 66.2 range (55-80) Drop outs: NR* IPSS: NR Erectile dysfunction: NR PVR (TRUS) ml: 33.2 range (15.4-67.5) Qmax: NR Operation time: 31.3 mins range (5-70) Mean catheterisation time (days): 1.9 ± 1.3 Mean length of stay (days): 3.4 ± 1.2</p> <p>*3 patients dropped out after randomisation but groups not defined. Only 76 patients has data at 6 weeks postoperatively</p>	<p>and between 35 & 325 TURPs</p> <p>Examination methods: Patients followed at 6 weeks, 3, 6, 12 months by same investigator</p> <p>During follow up Qmax, IPSS, QoL, bother and BSFQ all completed and TRUS, urodynamics and serum PSA measured at 6 months</p>	<p>number of patients Peri-operative urinary tract infections (follow up period 6 weeks**)</p> <p>Post-op complications number of patients TUR syndrome (follow up period 6 weeks**)</p> <p>Post-op complication: Haemorrhage necessitating readmission: (follow up period 6 weeks**)</p>	<p>Group 2: 3/38 p value: NR</p> <p>Group 1: 0/38 Group 2: 1/38 p value: NR</p> <p>Group 1: 1/38 Group 2: 3/38 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Carter et al., 1999^{44,45}</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Patients from urology outpatient department with BPE severe enough to warrant operation</p> <p>Setting: single centre, UK</p> <p>Inclusion Criteria: (based on British Laser Urological Evaluation Society (BLUES))</p> <ul style="list-style-type: none"> • Qmax ≤ 15 ml/s • Voided volume > 150 ml • PVR < 300 ml • IPSS ≥ 12 <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • History of acute retention • Histological diagnosis of prostate adenocarcinoma • Prostate volume > 100 ml (TRUS) • Neurogenic bladder <p>All patients N: 204</p> <p>Drop outs: 13 (9 violated entry criteria, 2 with calculi, 2 with urethral strictures)</p> <p>Group 1 - Laser N: 95</p> <p>Mean age ± SD (yrs): 67.9 ± 7.8</p> <p>Drop outs: NR</p> <p>IPSS: 20.3 ± NR</p> <p>Erectile dysfunction: NR</p> <p>Mean Prostate volume (TRUS) ml ± SD: 41.6 ± 17.3</p> <p>Mean PSA ng/ml ± SD: 3.8 ± 2.7</p> <p>Mean Creatinine mmol/l ± SD: 95.3 ± 15.7</p>	<p>Group 1 Hybrid laser performed using Laserscope 40W KTP/60W Nd:YAG generator system abd AddStat laser delivery fibres producing forward or side beams through a 21 F laser cystoscope (Storz). 30W KTP treatment to create bladder neck incisions and vaporisation then Nd:YAG 60W used to coagulate. Catheter protocol: Urethral catheter removed either 1 or 2 days or 1-2 weeks</p> <p>Group 2 TURP in standard manner through 24 or 26 Fr resectoscope. Catheters removed postoperatively when clinically indicated</p> <p>All patients: Antibiotics: single dose Gentamicin at operation and catheter removal.</p> <p>Intervention</p>	<p>Mean (SD) IPSS symptom score at 6 months</p>	<p>Group 1 (n=90): 6.7 (4.0) Group 2 (n=89): 6.4 (4.0)</p>	<p>Funding: Partially funded by Somerset Health Authority</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Baseline values for were not reported with standard deviations • Follow up outcomes Qmax and IPSS, QoL scores not reported with standard deviations. Only as graphs. • Outcome assessment was not masked. • Allocation concealment not clear if opaque sequential envelopes were used • *Unclear which follow up complications refer to and how many patients remained. ITT analysis used for late complications <p>Notes: Mean and standard deviations for IPSS and Qmax data estimated from graphs.</p>
			<p>Mean (SD) IPSS symptom score at 12 months</p>	<p>Group 1 (n=86): 6.6 (3.6) Group 2 (n=84): 5.9 (4.7)</p>	
			<p>Mean (SD) Qmax at 6 months</p>	<p>Group 1 (n=90): 19.1 (5.1) Group 2 (n=89): 19.6 (5.1)</p>	
			<p>Mean (SD) Qmax at 12 months</p>	<p>Group 1 (n=86): 19.8 (5.8) Group 2 (n=84): 20.9 (6.5)</p>	
			<p>Early post-op complications: Failure to void as inpatient following catheter removal (follow up period up to 6 months)</p>	<p>Group 1: 26/81 Group 2: 5/96 p value: <0.00001 (calculated by NCGC Fishers exact test)</p>	
			<p>Late post-op complications: urinary tract infection (follow up period > 6 weeks to 1 year)*</p>	<p>Group 1: 2/95 Group 2: 6/96 p value: Not signif. (calculated by NCGC Fishers exact test)</p>	
			<p>Late post-op complications: urethral stricture (follow up period > 6 weeks to 1 year)*</p>	<p>Group 1: 2/95 Group 2: 9/96 p value: 0.06 (calculated by NCGC Fishers exact test)</p>	
			<p>Late post-op complications: acute retention (follow up period > 6 weeks to 1 year)*</p>	<p>Group 1: 2/95 Group 2: 0/96 p value: Not signif. (calculated by NCGC Fishers exact test)</p>	
			<p>Late post-op complications: incontinence (follow up period > 6 weeks to 1 year)*</p>	<p>Group 1: 1/95 Group 2: 0/96 p value: Not signif. (calculated by NCGC Fishers exact test)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> Qmax: 9.0 ± NR PVR: 109 ± NR Operation time: 37.4 ± 12.1 mins 3.4 ± 1.2 Median catheterisation time (days): NR Median length of stay (days): 2 (0-9) </p> <p> Group 2 - TURP N: 96 Mean age ± SD (yrs): 67.0 ± 7.5 Drop outs: NR IPSS: 19.8 ± NR Erectile dysfunction: NR Mean Prostate volume (TRUS) ml ± SD: 41.7 ± 19.4 Mean PSA ng/ml ± SD: 3.2 ± 2.4 Mean Creatinine mmol/l ± SD: 99.7 ± 27 Qmax: 9.5 ± NR PVR: 135 ± NR Operation time: 35.7 ± 10.8 mins Median catheterisation time (days): NR Median length of stay (days): 2 (2-14) </p>	<p> performed by: 1 of 3 consultants, 2 Snr registrars, 1 clinical research fellow or 1 staff-grade urologist. </p> <p> Examination methods: Patients followed at 6 weeks, 6, 12 months </p> <p> During follow up IPSS, Symptom problem index (SPI), BPH impact Index (BPHII), Short Form 36 (HRQoL) questionnaires completed and uroflometry (Dantec Uroflow 1200), TRUS to find PVR. </p>	<p> Late post-op complications: Re-operation (follow up period > 6 weeks to 1 year)* </p>	<p> Group1: 2/95 Group 2: 1/96 p value: Not signif. (calculated by NCGC Fishers exact test) </p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Horasanli et al., 2008¹²²</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients referred to urology clinic with symptoms of BOO due to BPH</p> <p>Setting: single centre, dept urology, Memorial Hospital, Istanbul, Turkey</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Prostate volume 70-100 mL (TRUS) or PVR >150 mL with IPSS score > 7 <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Urethral strictures PVR > 400mL Previous prostatic, bladder or urethral surgery Prostate malignancy Indwelling catheters Refusal of consent <p>All patients N: 76 Drop outs: NR*</p> <p>Group 1 - Laser N: 39 Mean age ± SD (yrs): 69.2 ± 7.1 (range 59-78) IPSS Score: 18.9 ± 5.1 IIEF-5: 19.9 ± 5.1</p>	<p>Group 1 Photoselective vaporisation performed using KTP/532 emitting green light at 80W via a 6F side-firing fibre through 24F continuous flow cystoscope. A 20F 3-way Foley catheter was left in place and bladder irrigated with saline for 24 hours.</p> <p>Group 2 TURP in standard manner under general anaesthesia using Storz 26F continuous flow resectoscope. A 20F 3-way Foley catheter was left in place and bladder irrigated with saline for 24-48 hours.</p> <p>All patients: Antibiotics before and after..</p> <p>Intervention performed by: 5 surgeons</p>	<p>IPSS symptom score at 3 months</p>	<p>Group 1: 11.2 ± 7.6 Group 2: 6.1 ± 5.4 p value: 0.01 (calculated by NCGC as t test with unequal variances using ITT analysis)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method not reported Allocation concealment not reported Masking of outcome assessment not reported Drop out numbers not clear so ITT analysis used <p>Notes: * Drop out numbers not clear so ITT analysis used.</p>
			<p>Change in IPSS symptom score from baseline at 3 months</p>	<p>Group 1: 7.7 ± NR Group 2: 14.1 ± NR p value: NR</p>	
			<p>IIEF-5 at 3 months</p>	<p>Group 1: 19.0 ± 3.8 Group 2: 20.0 ± 4.7 p value: Not signif. (calculated by NCGC as t test with equal variances using ITT analysis)</p>	
			<p>Change in IIEF-5 from baseline at 3 months</p>	<p>Group 1: 0.9 ± NR Group 2: 0.1 ± NR p value: NR</p>	
			<p>flow rate (Qmax) at 3 months</p>	<p>Group 1: 14.1 ± 8.7 Group 2: 21.3 ± 12.8 p value: 0.006 (calculated by NCGC as t test with unequal variances using ITT analysis)</p>	
			<p>Change in flow rate (Qmax) from baseline at 3 months</p>	<p>Group 1: 5.5 ± NR Group 2: 12.1 ± NR p value: NR</p>	
			<p>IPSS symptom score at 6 months</p>	<p>Group 1: 13.1 ± 5.8 Group 2: 6.4 ± 7.9 p value: 0.0001 (calculated by NCGC as t test with equal variances using ITT analysis)</p>	
			<p>Change in IPSS symptom score from baseline at 6 months</p>	<p>Group 1: 5.8 ± NR Group 2: 13.8 ± NR p value: NR</p>	
			<p>IIEF-5 at 6 months</p>	<p>Group 1: 19.0 ± 5.2 Group 2: 21.0 ± 6.8 p value: Not signif. (calculated by NCGC as t test with equal variances using ITT analysis)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean Prostate vol (TRUS) ml ± SD: 86.1 ± 8.8</p> <p>Mean PSA ng/ml ± SD: 5.2 ± 4.5</p> <p>Qmax ml/s ± SD : 8.6 ± 5.2</p> <p>PVR ml ± SD: 183.0 ± 50.1</p> <p>Operating time (min ± SD): 87 ± 18.3</p> <p>Mean catheterisation time (days): 1.7 ± 0.8</p> <p>Mean length of stay (days): 2.0 ± 0.7</p> <p>Drop outs: NR</p> <p>Group 2 - TURP</p> <p>N: 37</p> <p>Mean age ± SD (yrs): 68.3 ± 6.7 (range 58-76)</p> <p>IPSS Score: 20.2 ± 6.8</p> <p>IIEF-5: 20.1 ± 5.5</p> <p>Mean Prostate vol (TRUS) ml ± SD: 88.0 ± 9.2</p> <p>Mean PSA ng/ml ± SD: 4.7 ± 3.8</p> <p>Qmax ml/s ± SD : 9.2 ± 5.6</p> <p>PVR ml ± SD: 176.9 ± 45.3</p> <p>Operating time (min ± SD): 51 ± 17.2</p> <p>Mean catheterisation time (days): 3.9 ± 1.2</p> <p>Mean length of stay (days): 4.8 ± 1.2</p> <p>Drop outs: NR</p>	<p>(consultant or experienced SpR)</p> <p>Examination methods:</p> <p>Patients followed at 3 and 6 months. All patients were assessed preoperatively and at follow ups for IPSS score, International Index of Erectile Dysfunction (IIEF-5), PSA, Qmax, PVR. In addition, postoperatively, data on length of stay, operating time, catheter removal time, and complications were collected.</p>	<p>Change in IIEF-5 from baseline at 6 months</p> <p>flow rate (Qmax) at 6 months</p> <p>Change in flow rate (Qmax) from baseline at 3 months</p> <p>Early post-op complications: patients requiring transfusion (follow up period up to 6 months)</p> <p>Early post-op complication: urinary retention (follow up period up to 6 months)</p> <p>Early post-op complications: urinary tract infection (follow up period up to 6 months)</p> <p>Early post-op complications: urethral stricture (follow up period up to 6 months)</p> <p>Early post-op complications: incontinence (follow up period up to 6 months)</p> <p>Reoperation rate (follow up period up to 6 months)</p>	<p>Group1: 0.9 ± NR</p> <p>Group 2: -0.9 ± NR (IIEF-5 increased)</p> <p>p value: NR</p> <p>Group1: 14.1 ± 8.7</p> <p>Group 2: 21.3 ± 12.8</p> <p>p value: 0.002 (calculated by NCGC as t test with unequal variances using ITT analysis)</p> <p>Group1: 4.7 ± NR</p> <p>Group 2: 11.5 ± NR</p> <p>p value: NR</p> <p>Group1: 0/39 *</p> <p>Group 2: 3/37 *</p> <p>p value: Not signif (calculated by NCGC Fishers exact test)</p> <p>Group1: 6/39 *</p> <p>Group 2: 1/37 *</p> <p>p value: Not signif (calculated by NCGC Fishers exact test)</p> <p>Group1: 6/39 *</p> <p>Group 2: 5/37 *</p> <p>p value: Not signif (calculated by NCGC Fishers exact test)</p> <p>Group1: 2/39 *</p> <p>Group 2: 3/37 *</p> <p>p value: Not signif (calculated by NCGC Fishers exact test)</p> <p>Group1: 0/72 **</p> <p>Group 2: 1/76 **</p> <p>p value: Not signif. (calculated by NCGC Fishers exact test)</p> <p>Group1: 7/39 *</p> <p>Group 2: 0/37 *</p> <p>p value: 0.01 (calculated by NCGC Fishers exact test)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Keoghane et al., 2000^{142,144} & Keoghane et al., 1996^{140,141,143}</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 5 years</p>	<p>Patient group: Patients referred to hospital requiring surgery for BPE</p> <p>Setting: single centre, UK</p> <p>Inclusion Criteria: NR</p> <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Previous surgery or instrumentation for BPE • Prostate malignancy • Insufficient knowledge of English to answer questionnaire • Refusal of consent <p>All patients N: 148</p> <p>Drop outs: *at 5 years 63/148 (43%); 17 (7 laser and 10 TURP) had died., 8 unable to respond to questionnaires through disease and 38 lost to follow up.</p> <p>Group 1 - Laser N: 72</p> <p>Mean age ± SD (yrs): 69 ± 8 (range 51-95)</p> <p>Drop outs: *</p> <p>AUA 7 Score: 19.9 ± 7.7 (n=54)</p> <p>Bother score: 5.8 ± 3.0 (n=59)</p> <p>Mean SF36 (physical) ±SD: 43.69 ±12.58 (n=51)</p> <p>Mean SF36 (mental) ±SD: 47.07 ±11.2 (n=51)</p>	<p>Group 1 Vaporisation using MD60 Nd:YAG (Selected Laser Technologies) with 600 µm fibre incorporating sapphire-tipped probe. Irrigation using saline.</p> <p>Group 2 TURP in standard manner using Storz equipment and irrigation with glycine</p> <p>All patients: Oral ciprofloxacin prophylaxis before surgery.</p> <p>After treatment 22F 3-way catheter inserted and continuous irrigation commenced. Catheter removed when clinically indicated</p> <p>Intervention performed by: 5 surgeons (consultant or experienced SpR)</p> <p>Examination methods: Patients followed at 4 weeks, 3, 12, 24, 36 months to 5 years</p> <p>Patients received cystourethroscopy after randomisation to assess</p>	AUA 7 symptom score from baseline at 3 months	Group1: 9.6 ± 7.5 (n=55) Group 2: 6.5 ± 5.1 (n=62) p value: 0.03	<p>Funding: Oxford Regional Health Authority</p> <p>Limitations: **Patient numbers for primary and secondary outcomes and complications were unclear so ITT analysis used.</p> <p>Notes: Randomisation by random number tables and allocation concealment through sealed envelopes although opacity was not reported. Patients and investigators were masked to treatment allocation</p> <p>Change from baseline at 5 years were reported for AUA score but SDs were not reported.</p>
			Change in AUA 7 symptom score from baseline at 3 months	Group1: 10.1 ± 9.7 (n=47) Group 2: 13.6 ± 6.9 (n=54) p value: NS	
			AUA 7 symptom score from baseline at 12 months	Group1: 8.7 ± 6.5 (n=53) Group 2: 5.8 ± 5.4 (n=60) p value: 0.006	
			Change in AUA 7 symptom score from baseline at 12 months	Group1: 10.9 ± 8.4 (n=44) Group 2: 13.3 ± 7.8 (n=53) p value: not signif. (NCGC t-test)	
			AUA 7 symptom score from baseline at 2 years	Group1: 7.8 ± 6.6 (n=45) Group 2: 5.7 ± 6.0 (n=52) p value: 0.018	
			Change in AUA 7 symptom score from baseline at 2 years	Group1: 11.7 ± 9.7 (n=35) Group 2: 13.7 ± 7.7 (n=47) p value: not signif. (NCGC t-test)	
			AUA 7 symptom score from baseline at 3 years	Group1: 8.9 ± 6.6 (n=37) Group 2: 6.5 ± 6.5 (n=41) p value: 0.001	
			Change in AUA 7 symptom score from baseline at 3 years	Group1: 11.0 ± 9.7 (n=37) Group 2: 12.9 ± 7.9 (n=41) p value: not signif. (NCGC t-test)	
			Change in flow rate (Qmax) from baseline at 12 months	Group1: 6.2 ± 15.0 (n=32) Group 2: 9.4 ± 12.5 (n=37) p value: not signif. (NCGC t-test)	
			Change in flow rate (Qmax) from baseline at 24 months	Group1: 5.2 ± 7.0 (n=18) Group 2: 4.9 ± 7.5 (n=26) p value: not signif. (NCGC t-test)	
Change in flow rate (Qmax) from baseline at 24 months	Group1: 1.8 ± 6.2 (n=24) Group 2: 2.1 ± 6.9 (n=24) p value: not signif. (NCGC t-test)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Erectile dysfunction (difficulty maintaining erection): 9/38 (24%)</p> <p>Mean Prostate volume ml ± SD: 54.2 ± 26.3 (n=44)</p> <p>Qmax: 11.8 ± 4.5 (n=48)</p> <p>PVR: NR</p> <p>Median catheterisation time (days): 1 (0-9)</p> <p>Median length of stay (days): 3 (1-10)</p> <p>Group 2 - TURP</p> <p>N: 76</p> <p>Mean age ± SD (yrs): 70 ± 8 (range 47-84)</p> <p>Drop outs: *</p> <p>AUA 7 Score: 19.4 ± 6.5 (n=63)</p> <p>Bother score: 5.9 ± 2.3 (n=68)</p> <p>Mean SF36 (physical) ±SD: 44.66 ±12.12 (n=57)</p> <p>Mean SF36 (mental) ±SD: 47.75 ±10.47 (n=57)</p> <p>Erectile dysfunction (difficulty maintaining erection): 20/50 (40%)</p> <p>Mean Prostate volume ml ± SD: 51.9 ± 24.1 (n=48)</p> <p>Qmax: 11.4 ± 5.0 (n=54)</p> <p>PVR: NR</p> <p>Median catheterisation time (days): 2 (1-20)</p> <p>Median length of stay (days): 4 (1-8)</p>	<p>length of urethra and bladder pathology and residual volume.</p> <p>AUA score assessed preoperatively and at 4 weeks.</p> <p>Qmax was a secondary outcome measurement methods not reported.</p>	Erectile Dysfunction (difficulty maintaining erection) at 3 months	<p>Group1: 7/38</p> <p>Group 2: 12/50</p> <p>p value: Not signif. (calculated by NCGC Chi squared test)</p>	
			Bother score at 3 months	<p>Group1: 2.9 ± 3.0 (n=54)</p> <p>Group 2: 2.4 ± 3.0 (n=64)</p> <p>p value: Not Signif.</p>	
			Early post-op complications: Failure to void as inpatient following catheter removal (follow up period first 3 months)	<p>Group1: 17/72 **</p> <p>Group 2: 8/76 **</p> <p>p value: Not signif. (calculated by NCGC Chi squared test)</p>	
			Early post-op complications: patients requiring transfusion (follow up period first 3 months)	<p>Group1: 0/72 **</p> <p>Group 2: 13/76 **</p> <p>p value: 0.0001 (calculated by NCGC Fishers exact test)</p>	
			Late post-op complications: urinary tract infection (follow up period first 3 months)	<p>Group1: 1/72 **</p> <p>Group 2: 3/76 **</p> <p>p value: Not signif. (calculated by NCGC Fishers exact test)</p>	
			Late post-op complications: urethral stricture ((follow up period first 3 months)	<p>Group1: 0/72 **</p> <p>Group 2: 3/76 **</p> <p>p value: Not signif. (calculated by NCGC Fishers exact test)</p>	
			Late post-op complications: incontinence (follow up period first 3 months)	<p>Group1: 0/72 **</p> <p>Group 2: 1/76 **</p> <p>p value: Not signif. (calculated by NCGC Fishers exact test)</p>	
			Reoperation rate at 5 years	<p>Group1: 13/72</p> <p>Group 2: 11/76</p> <p>p value: Not signif. (calculated by NCGC Fishers exact test)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Mottet et al., 1999²⁰⁶</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Patients in urology clinics</p> <p>Setting: multi-centre, Nimes & Paris, France</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • Qmax <12ml/s • age >45 years • PVR <250ml • AUA > 13 • PSA < 10ng/ml • informed consent <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • history of prostatic or urethral surgery • prostate >60g • diabetes • bladder or neurogenic disease <p>All patients N: 36 Age: 66 (range 50-77) Drop outs: 17 (at 12 mths)</p> <p>Group 1 - Laser N: 23 Mean age (yrs): 67 Drop outs: 11 without outcome data at 12 months IPSS: 20 Madsen score: 15 Erectile dysfunction: NR Prostate volume (TRUS) ml: 39 Qmax ml/s: 8 Operation time mins: 75 Mean catheterisation time (days): 1.6</p>	<p>Group 1 Dual length VersaPulse Select Laser at 60W-80W of holmium:YAG energy in pulsed mode through 550µm fibre or side-firing fibre in 24F cystoscope. 6 patients also received additional Nd:YAG vaporisation. 20 or 24F Foley placed without irrigation and removed the next day.</p> <p>Group 2 TURP in standard manner under spinal anaesthesia with glycine irrigation followed by postoperative saline irrigation until urine was clear. Catheter was then removed.</p> <p>Intervention performed by same 2 experienced surgeons</p> <p>Examination methods: Patients followed at 1, 3, 6, 12 months</p> <p>During preoperative assessment and follow up DRE, Qmax, IPSS and Madsen score, PSA and TRUS all completed. Patients were also questioned about potency</p>	<p>Mean IPSS at 3 months</p>	<p>Group 1: 7.7 ± NR (n=22) Group 2: 7.5 ± NR (n=12) p value = NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Outcomes were reported without standard deviations • Outcome assessment was not masked. • Randomisation method not reported. • Allocation concealment not reported <p>Additional outcomes: Madsen score at follow up</p> <p>Notes: Randomisation on 2:1 model</p>
			<p>Mean IPSS at 6 months</p>	<p>Group 1: 6.2 ± NR (n=20) Group 2: 7.7 ± NR (n=11) p value = NR</p>	
			<p>Mean IPSS at 12 months</p>	<p>Group 1: 5.9 ± NR (n=12) Group 2: 7.5 ± NR (n=7) p value = NR</p>	
			<p>Mean Qmax at 3 months</p>	<p>Group 1: 22.8 ± NR (n=22) Group 2: 18.3 ± NR (n=12) p value = NR</p>	
			<p>Mean Qmax at 6 months</p>	<p>Group 1: 17.5 ± NR (n=20) Group 2: 16.6 ± NR (n=11) p value = NR</p>	
			<p>Mean Qmax at 12 months</p>	<p>Group 1: 19.3 ± NR (n=12) Group 2: 17.6 ± NR (n=7) p value = NR</p>	
			<p>Early Post-op complications number of patients with blood transfusion</p>	<p>Group 1: 0/23 Group 2: 0/13</p>	
			<p>Post-op complications number of patients incontinence at 6 months</p>	<p>Group 1: 1/23 Group 2: 0/13</p>	
			<p>Reoperation rate</p>	<p>Group 1: 1/23 Group 2: 2/13</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>± NR Mean length of stay (days): 2.2 ± NR</p> <p>Group 2 - TURP N: 13 Mean age (yrs): 64 Drop outs: 6 without outcome data at 12 months IPSS: 24 Madsen score: 17 Erectile dysfunction: NR Prostate volume (TRUS) ml: 34 Qmax ml/s: 8 Operation time mins: 40 Mean catheterisation time (days): 3.1 ± NR Mean length of stay (days): 2.1 ± NR</p>	<p>and ejaculation status.</p> <p>Length of stay, catheterisation time, reoperation rate also recorded</p>			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Shingleton et al., 2002²⁷⁵ & Shingleton et al., 1999²⁷⁷ &</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 years</p>	<p>Patient group: Patients with failed α-blockers therapy for voiding symptoms</p> <p>Setting: single-centre, Istanbul, Turkey</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • peak urine flow rate <15ml/s • age >45 years • failure of • medical therapy (α-blockers) • able to undergo regional/general anaesthesia • medical therapy discontinued 1 month before surgery <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Prostate cancer <p>All patients N: 100 Age: 66 (range 50-77) Drop outs:</p> <p>Group 1 - Laser N: 50 Mean age \pm SD (yrs): 68.2 \pm 7.9 Ethnicity: 38/50 (76%) white. Mean AUA score \pm SD: 22.5 \pm 6.0 Erectile dysfunction (full): 22/50 (44%) Prostate volume (TRUS) ml: 32.2 \pm 21.4 Mean PSA ng/ml \pm SD: 2.7 \pm 2.3 Mean Qmax \pm SD (ml/s): 8.2 \pm 3.2 Operation time mins: 43 (15-70)</p>	<p>Group 1 Laserscope KTP/Nd:YAG with Laserscope ADD or ADD/stat fibre. 36W was used first for vaporisation then 60W for further vaporisation and coagulation. A catheter was placed for between 1-5 days depending on size of prostate and energy used</p> <p>Group 2 TURP in standard manner using Circon/ACMI continuous flow resectoscope with mannitol solution.</p> <p>Laser intervention performed by one surgeon and TURPs by senior residents under same surgeon.</p> <p>Examination methods: All patients had AUA symptom score, serum PSA, TRUS, pressure flow urodynamics preoperatively and were followed up with AUA score, PSA and uroflowmetry measurements at 1, 3,</p>	AUA symptom score at 3 months	Group 1: 7.0 \pm NR (n=48) Group 2: 4.0 \pm NR (n=48) p value = 0.01	<p>Funding: In part by Laserscope</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Reasons for drop out were not reported and there were more patients at 3 years than 2 years • Outcome assessment was not masked. • Allocation concealment not reported • Changes from baseline were not reported <p>Additional outcomes: Prostate volume at follow up, serum PSA at follow up Other complications including retrograde ejaculation.</p> <p>Notes: Computer generated randomisation. *ITT analysis used for statistical analysis</p>
			AUA symptom score at 6 months	Group 1: 7.0 \pm NR (n=46) Group 2: 4.0 \pm NR (n=48) p value = 0.01	
			AUA symptom score at 12 months	Group 1: 6.0 \pm 6.0 (n=40) Group 2: 3.8 \pm 4.1 (n=33) p value = 0.03 (calculated by NCGC using t test with equal variances *)	
			AUA symptom score at 18 - 24 months	Group 1: 5.9 \pm 5.7 (n=23) Group 2: 4.6 \pm 4.2 (n=19) p value = 0.19 (calculated by NCGC using t test with equal variances *)	
			AUA symptom score at 36 months	Group 1: 9.9 \pm 6.7 (n=29) Group 2: 7.7 \pm 5.6 (n=33) p value = 0.07 (calculated by NCGC using t test with equal variances *)	
			Qmax at 3 months	Group 1: 15.0 \pm 5.7 (n=48) Group 2: 16.0 \pm 8.0 (n=48) p value = 0.60	
			Qmax at 6 months	Group 1: 15.8 \pm 6.9 (n=46) Group 2: 16.3 \pm 6.4 (n=48) p value = 0.77	
			Qmax at 12 months	Group 1: 14.6 \pm 5.9 (n=40) Group 2: 16.2 \pm 7.2 (n=33) p value = 0.23 (calculated by NCGC using t test with equal variances *)	
			Qmax at 18-24 months	Group 1: 14.9 \pm 5.4 (n=23) Group 2: 14.3 \pm 6.3 (n=19) p value = 0.6 (calculated by NCGC using t test with equal variances*)	
Qmax at 36 months	Group 1: 12.3 \pm 5.3. (n=29) Group 2: 12.8 \pm 5.6 (n=33) p value = 0.64 (calculated by NCGC using t test with equal variances *)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Drop outs: Mean catheterisation time (days): NR Mean length of stay (days): NR</p> <p>Group 2 - TURP N: 50 Mean age ± SD (yrs): 67.4 ± 7.3 Ethnicity: 34/50 (68%) white. Mean AUA score ± SD: 21.2 ± 6.1 Erectile dysfunction (full): 21/50 (42%) Prostate volume (TRUS) ml: 29.6 ± 15.4 Mean PSA ng/ml ± SD: 3.2 ± 2.2 Mean Qmax ± SD (ml/s): 7.3 ± 3.7 Operation time mins: 56 (45-90) Drop outs: Mean catheterisation time (days): NR Mean length of stay (days): NR</p>	6, 12, 18, 24, 36, 48, 60 and 72 months	<p>Post-op complications number of patients with urethral stricture (follow up period 12 months)*</p> <p>Post-op complications number of patients incontinence (follow up period 12 months)*</p> <p>Post-op complications number of patients with urinary retention (follow up period 12 months)</p>	<p>Group 1: 1/50 Group 2: 1/50 p value: NR</p> <p>Group 1: 1/50 Group 2: 1/50 p value: NR</p> <p>Group 1: 3/50 Group 2: 1/50 p value: NR</p>	

Study Details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Suvakovic & Hindmarsh, 1996²⁹⁰</p> <p>Study design: RCT, open label</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Consecutive patients with prostatic symptoms</p> <p>Setting: Urology department, South Cleveland University, UK</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • Qmax ≤15mL/s for a voided volume of ≥150 mL • Age • Significant voiding symptoms (AUA score >15) • PSA level <2.5 ng/mL • Prostate volume <40g (assessed by TRUS, DRE and cystoscopy) • Length of the prostatic urethra >4 cm <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • Malignancy <p>All patients N: 40</p> <p>Group 1 - CLAP- contact laser alone N: 10 Age (mean): 62.6(5.8) IPSS: 18 (6.0) Qmax ml/s: 12.2 (3.8) Residual Vol mL: 139.6(103) Prostate size (by TRUS), g: 24(5.8) Mean catheterisation time (days): 1 ± NR</p>	<p>Group 1 : CLAP- contact laser alone Nd: YAG laser applied at 40W for vaporising and coagulating the prostate with a minimum depth of penetration. a 16 F two-way catheter was inserted into the bladder and removed after 24 h.</p> <p>Group 2 : TURP Standard resection using a 26 F continuous irrigating resectoscope. A 22 F three-way urethral catheter was inserted into the bladder and irrigation was continued up to 24 h. The catheter was removed after 48 h and the patients discharged home 3-4 days after the procedure.</p> <p>All patients received preoperative oral antibiotics and controlled for more than 5 days post-operatively</p> <p>Examination methods: At 3, 6 12 months AUA</p>	<p>IPSS symptom score, mean ± SD at 3 months</p>	<p>Group 1: 9.7 ± 2.6, n=10 Group 2: 12.8 ± 5.9, n=10 p value: 0.15 (calculated by NCGC using t test with unequal variances using ITT analysis)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Small sample size, n of 10 in each arm • Unclear which statistical test was used for data – discrepancies in the stat sig reported for AUA score for 3 months and calculated by NCGC team. • Randomisation method and allocation concealment not reported. • Masking of outcome assessment not reported. • Number of participants followed up at 12 months not reported. • Complications were poorly reported <p>Notes: None.</p>
			<p>IPSS symptom score, mean ± SD at 6 months</p>	<p>Group 1: 8.7 ± 5.4, n=9 Group 2: 8.5 ± 3.0, n=10 p value: 0.91 (calculated by NCGC using t test with unequal variances using ITT analysis)</p>	
			<p>IPSS symptom score, mean ± SD at 12 months *Values for 12 months follow up reported in paper, but n was not reported</p>	<p>Group 1: 8.7 ± 4.9, * Group 2: 7.2 ± 6.1, * p value: 0.55 (calculated by NCGC using t test with equal variances using ITT analysis)</p>	
			<p>Qmax mean ± SD at 3 months</p>	<p>Group 1: 15.6 ± 13.5, n=10 Group 2: 17.8 ± 3.8, n=10 p value: NR</p>	
			<p>Qmax mean ± SD at 6 months</p>	<p>Group 1: 18.7 ± 7.5, n=9 Group 2: 19.0 ± 0.8, n=10 p value: NR</p>	
			<p>Qmax mean ± SD at 12 months *Values for 12 months follow up reported in paper, but n was not reported</p>	<p>Group 1: 23.5 ± 5.9, * Group 2: 15.2 ± 2.7, * p value: NR</p>	
			<p>Post-op complications: Catheter duration, mean, hours (range or standard deviations NR)</p>	<p>Group 1: 24, n=10 Group 2: 48, n=10 p value: NR</p>	
			<p>Post-op complications Length of hospitalisation, (hours)</p>	<p>Group 1: 30, n=10 Group 2: 84, n=10</p>	

Study Details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean length of stay (days): 1.3 ± NR</p> <p>Group 2 - TURP Standard resection N: 10 Age (mean): 66.1(5.1) IPSS: 18.8 (4.5) Qmax ml/s: 11.1(6.4) Residual Vol mL: 161.8(104) Prostate size (by TRUS), g: 22(5) Mean catheterisation time (days): 2 ± NR Mean length of stay (days): 3.5 ± NR</p>	<p>score, PSA, flow rate, PVR measured and TRUS performed</p>			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Tuhkanen et al., 2001³⁰⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24 months</p>	<p>Patient group: Patients with BPH and BOO that were referred to the outpatient clinic at Kuopio university hospital from January 1995 to November 1997.</p> <p>Setting: Urology department, Finland</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Obstructed if min. voiding pressure > 40cm water prostate volume 40-100ml (TRUS) <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> prostate cancer or surgery urinary retention <p>All patients N: 46 Drop outs: 9 (20%)</p> <p>Group 1 N: 21 Age (mean): 67 (55-78) Mean (range) symptom score (DanPSS-1): 18.6 (5-40) Prostate volume: 55 (42-83) Qmax ml/s (range): 8.5 (2.3-17.2) PVR ml (range): 125 (0-350) Drop outs: 4 (1=died cardiac infarct 5 months post-operatively; 3=underwent TURP - Mean prostate size: 55 (42-83)ml Mean catheterisation time (days): NR Mean length of stay (days): 4.0 (2-9)</p> <p>Group 2 - N: 25 Age (mean): 67 (46-77) Mean (range) symptom score</p>	<p>Group 1: laser (hybrid) Initial noncontact Nd:YAG coagulation 40W power asset for 90 sec burn times. Followed by a contact Nd:YAG vaporisation to open prostatic urethra. Vaporised at 40W. Urethral catheter was inserted for one day. Postoperatively the suprapubic catheter removed when the patient could urinate and residual urine was less than 150ml. Spinal anaesthesia.</p> <p>Group 2: TURP 28 F Storz resectoscope without application of the suprapubic catheter. Spinal anaesthesia.</p> <p>Examination methods: Patients reviewed at 3, 6, 12, 24 mths DanPSS-1, urinalysis, serum creatinine, serum PSA, Qmax, PVR, DRE were recorded at each visit. TRUS was performed for suspicious cancer cases</p>	<p>Mean (range) symptom score (DanPSS-1)</p>	<p><u>At 3 months</u> Group 1 (n=21): 10.0 (0-49) Group 2 (n=22): 5.6 (0-27)</p> <p><u>At 6 months</u> Group 1 (n=19): 5.5 (0-21) Group 2 (n=21): 4.7 (0-22)</p> <p><u>At 24 months</u> Group 1 (n=17): 7.2 (0-25) Group 2 (n=20): 3.4 (0-21)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method, allocation concealment and masking of outcome assessment were not reported uses DanPSS-1 score standard deviations not reported <p>Additional outcomes: Average urinary flow rate reported.</p> <p>Notes: Linked to Tuhkanen 1999a³⁰¹</p>
			<p>Qmax mL/sec (range)</p>	<p><u>At 3 months</u> Group 1: 13.7 (4.9-27.5) Group 2: 21.0 (3.2-41.9)</p> <p><u>At 6 months</u> Group 1: 14.4 (7.9-20.7) Group 2: 19.6 (4.1-43.2)</p> <p><u>At 24 months</u> Group 1: Group 2: 20.6 (9.5-38.9)</p>	
			<p>Residual urinary volume, ml</p>	<p><u>At 3 months</u> Group 1: 77 (0-162) Group 2: 54 (0-210)</p> <p><u>At 6 months</u> Group 1: 69 (0-160) Group 2: 45 (0-177)</p> <p><u>At 24 months</u> Group 1: 114 (28-202) Group 2: 58 (0-166)</p>	
			<p>Reoperation rate (24 months follow-up):</p>	<p>Group 1: 3/21 Group 2: 2/25</p>	
			<p>Retrograde ejaculation at 3 months</p>	<p>Group 1: 3/16 Group 2: 12/14</p>	
			<p>Complications</p>	<p><u>Transfusion:</u> Group 1: 1/21 Group 2: 2/25</p> <p><u>Mortality</u> Group 1: 1(myocardial infarction at 5 m)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>(DanPSS-1): 22.8 (5-69) Prostate volume: 55 (40-95) Qmax ml/s (range): 7.2 (3.7-14.8) PVR ml (range): 138 (0-450) Drop outs: 5 (2=prostatic adenocarcinoma at initial operation, 1=internal urethrotomy for distal urethral stricture at 5 months; 1=died unknown causes at 13 months; 1=re-TURP due to overflow incontinence) Mean prostate size: 55 (40-94)ml Mean catheterisation time (days): NR Mean length of stay (days): 3.5 (1-8)</p>			<p>Group 2: 1 (unknown at 13 m) <u>Stricture (internal urethrotomy treatment)</u> Group 1: 0/21 Group 2: 1/25 <u>Incontinence (overflow at 13m)</u> Group 1: 0/21 Group 2: 1/24 <u>Urinary retention</u> (at 17 months and underwent TURP) Group 1: 2/21 Group 2: 0/25</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Tuhkanen et al., 2003²⁹⁹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 4 years</p>	<p>Patient group: LUTS with confirmed BOO recruited from September 1994 – January 1998. Prostate volume less than 40ml.</p> <p>Setting: Finland</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> • minimum volume of ≥ 120ml • minimum voiding detrusor pressure > 40 cm water <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> • prostate cancer, prostate surgery or history of TUIP or TURP • prostate size > 40ml • urethral structure • neurogenic bladder dysfunction • residual volume > 350ml <p>All patients N: 52 Drop outs: 10</p> <p>Group 1 N: 26 Age (mean): 68 (56-82) Median (range) DanPSS-1 symptom score: 18 (5-54) Qmax (mean \pm SD) ml/s: 9.0 ± 3.8 Mean prostate volume (range) ml: 30 (15-37) Median PVR ml (range): 87 (0-331) Mean catheterisation time (days): NR Mean length of stay (days): 3.4 (2-7) Drop outs: 4 (3 died of BPH-unrelated</p>	<p>Group 1: Contact laser vaporisation Porsatic urethra vaporised with an Nd:YAG laser at a power setting 40W. Urethral catheter inserted for one day. Spinal anaesthesia. Ciproflaving eve and morning of operation.</p> <p>Group 2: TURP Ciproflaving eve and morning of operation. Spinal anaesthesia.</p> <p>Examination methods: Patients reviewed at 3, 6, 12, 24 and 48 mths DanPSS-1, urinalysis, serum creatinine, serum PSA, Qmax, PVR, DRE were recorded at each visit. Urodynamics and TRUS were performed at 6 months and 4 years</p>	<p>Median (range) DanPSS-1 symptom score</p>	<p><u>At 3 months: mean</u> Group 1 (n=25): 6 (7) Group 2 (n=25): 5 (6) <u>At 6 months: mean</u> Group 1: 6 (9) Group 2: 5 (7) <u>At 48 months</u> Group 1: (n=22): 5 (0-34) Group 2: (n=20): 4 (0-18)</p>	<p>Funding: Financially supported by University of Kuopio.</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method, allocation concealment and masking of outcome assessment were not reported • uses DanPSS-1 score • Patient numbers not clear at 6 months • 2 patients in TURP group refused follow-up due to good subjective outcomes. <p>Notes: Median values reported at baseline and 48 months in Tuhkanen 2003. Earlier study (Tuhkanen 1999) reports mean (SD) for baseline, 3 months and 6 months.</p>
			<p>Mean (SD) Qmax, mL/s</p>	<p><u>At 3 months</u> Group 1: 15.0 (5.2) Group 2: 19.0 (9) <u>At 6 months</u> Group 1: 17.9 (7.1) Group 2: 21.1 (9.7) <u>At 48 months – median (range)</u> Group 1: 14.3 (10.1-33.6) Group 2: 16.1 (7.7-39.6)</p>	
			<p>PVR, ml</p>	<p><u>At 3 months – mean (SD)</u> Group 1: 44 (39) Group 2: 36 (39) <u>At 6 months - mean (SD)</u> Group 1: 50 (64) Group 2: 32 (37) <u>At 48 months – median (range)</u> Group 1: 60 (0-380) Group 2: 10 (0-90) P<0.05</p>	
			<p>UTI (epididymitis) ejaculation at 6 mths</p>	<p>Group 1: 0/26 Group 2: 1/26</p>	
			<p>Retrograde ejaculation at 6 mths</p>	<p>Group 1: 1/16 (6%) Group 2: 13/16 (81%)</p>	
			<p>Mortality at 4 years</p>	<p>Group 1: 3/26 Group 2: 1/26</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>causes and one underwent TURP at 2 years postoperatively due to gross haematuria, residual adenoma tissue and bladder stones)</p> <p>Group 2 - N: 26 Age (mean): 67 (55-77) Median (range) DanPSS-1 symptom score: 18 (4-46) Qmax (mean ± SD) ml/s: 8.2 ± 3.2 Mean prostate volume (range) ml: 28 (15-38) Median PVR ml (range): 83 (8-350) Mean catheterisation time (days): NR Mean length of stay (days): 2.9 (2-5) Drop outs: 6 (1 died of BPH-unrelated causes, 2 diagnosed with prostatic carcinoma, one patient with bladder neck stenosis and underwent a re-TURP, 2 refused reviews due to good subjective outcomes).</p>		<p>Reoperation rate at 4 years</p>	<p>Group 1: 1/26 Group 2: 1/26</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Van Melick et al., 2003 ³⁰⁹	Patient group: men over 45 years with LTUS associated with BPH that were recruited from their clinic from 1996 to 2001	Group 1: Laser vaporisation Transurethral catheter post-operation SLT Nd:Yag (MTRL sapphire tip) through Morgenstern scope irrigated with isotonic salt solution. Pre-procedural antibiotics and transurethral catheter postoperatively.	Mean (± SD) symptom score (IPSS) at 6 months	Group 1 (n=33): 5.9 ± 5.5 Group 2 (n=37): 3.2 ± 2.7	Funding: NR. Limitations: <ul style="list-style-type: none"> Randomisation method was not described and masking of outcome assessment was not reported. High attrition rate at 1-7 years and 4-7 years Additional outcomes: Frequency during day, frequency during night, symptom problem index and BPH impact index. Uroflowmetry also reported. Notes: Links with Van Melick 2002 (up to 6 months), Van Melick 2003 Follow up time varied individually as all patients were analysed within a 2 month period. Depending on the individual follow-up time, patient divided into two groups: those with a follow-up time between 1 and 4 years and those with follow up time between 4 and 7 years.
Study design: RCT	Setting: Netherlands		Mean (± SD) symptom score (IPSS) at 12 months	Group 1 (n=37): 3.6 ± 3.4 Group 2 (n=41): 4.1 ± 4.8	
Evidence level: 1+	Inclusion Criteria: patient with lower urinary tract symptoms suggestive of BPH; met ISC criteria for BPH, Schafer obstruction score ≥ 2, prostate size between 20-65ml. Exclusion Criteria: age ≤ 45 yrs		Mean (± SD) symptom score (IPSS) at 1-4 years	Group 1 (n=10): 9.3 ± 5.2 Group 2 (n=15): 5.8 ± 7.5	
Duration of follow-up: Up to 7 years	All patients N: 95		Mean (± SD) symptom score (IPSS) at 4-7 years	Group 1 (n=17): 8.3 ± 6.4 Group 2 (n=15): 7.3 ± 7.1	
	Group 1 N: 45 Age (mean) ± SD: 67 ± 9 IPSS (mean) ± SD: 18.9 ± 6.8 Mean prostate size, ml: 37 ± 11 Mean (SD) Global quality of life score: 3.7 ± 1.6 Mean Qmax ± SD ml/s: 12 ± 4 Follow-up 1 to 4 years = 15 Follow-up 4 to 7 years = 15 ± 0.4 Mean length of stay (days): 3.8 ± 1.3 Mean catheterisation time (days): 2.1 ± 0.9 Drop outs: 8 at one year post-operatively (procedure during surgery changed for medical reasons=3, equipment failure resulting in TURP)=2, reoperation –TURP=1, reoperation – due to stricture =2)	Group 2: TURP Stabdard 24FR resectoscope using glycine for irrigation. Suprapubic catheter if required peri-operatively. Pre-procedural antibiotics and transurethral catheter postoperatively.	Mean (SD) Global quality of life score at 6 months	Group 1: 0.8 ± 1.0 Group 2: 0.5 ± 0.5	
			Mean (SD) Global quality of life score at 12 months	Group 1: 0.6 ± 0.9 Group 2: 0.6 ± 0.8	
			Mean (SD) Global quality of life score at 1-4 years	Group 1: 2.0 ± 1.0 Group 2: 1.1 ± 1.2	
			Mean (SD) Global quality of life score at 4-7 years	Group 1: 1.4 ± 1.2 Group 2: 1.3 ± 1.3	
			Qmax mean ± SD at 6 months	Group 1: 25 ± 9 Group 2: 26 ± 6	
			Qmax mean ± SD at 12 months	Group 1: 27 ± 12 Group 2: 23 ± 10	
			Qmax mean ± SD at 1-4 years	Group 1: 19 ± 6 Group 2: 20 ± 5	
		Examination methods: Urodynamic studies (cystometry and pressure flow) at baseline and 1-6 weeks, 3, 6, 12 months after treatment	Qmax mean ± SD at 4-7 years	Group 1: 19 ± 9 Group 2: 17 ± 8	
			Post-op complications: urethral stricture (within 12 mths)	Group 1: 2/45 Group 2: 2/50	
			Post-op complications: mortality (within 12 mths)	Group 1: 0/45 Group 2: 2/50	
			Post-op complications: transfusion required (within 12 mths)	Group 1: 0/45 Group 2: 1/50	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean prostate size, ml ± SD: 37 ± 11</p> <p>Mean ± SD Global quality of life score: 3.8 ± 1.5</p> <p>Mean Qmax ± SD ml/s: 11 ± 4</p> <p>Follow-up 1 to 4 years = 10</p> <p>Follow-up 4 to 7 years = 17</p> <p>Mean length of stay (days): 3.9 ± 0.9</p> <p>Mean catheterisation time (days): 2.8 ± 3.1</p> <p>Drop outs: 9 at one year post-operatively (surgery cancelled=1, mortality=2, morbidity=2, emigrated=1, reoperation (TURP) =2, reoperation (stricture)=1)</p>		<p>Post-op complications: urinary retention (within 12 mths)</p> <p>Reoperation rate (TURP) within 12 mths</p>	<p>Group 1: 5/45</p> <p>Group 2: 0/50</p> <p>Group 1: 1/45</p> <p>Group 2: 2/50</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Zorn et al., 1999³³²</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: military beneficiaries with symptomatic BPH – recruited from June 1995 to June 1996</p> <p>Setting: Walter Reed Army Medical Centre and Madigan Army Medical Centre, US</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> symptomatic BPH Qmax<15ml/s Age > 50 AUA score 13 or more PVR>125ml Prostate volume <45g <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> previous surgical therapy for BPH known prostate, bladder, urethral or neurological conditions that could affect the bladder <p>All patients N: 33</p> <p>Group 1 N: 21 Age (mean): 70.6 Drop outs: 3 IPSS: 24.0 Prostate size: 29.9 Qmax (mean) ml: 8.7 AUA symptom score (mean): 24.0 Mean length of stay (days): 1.2 ± NR Mean catheterisation time (days): 1.1 ± NR</p> <p>Group 2 - N: 12</p>	<p>Group 1: Laser vaporisation contact laser vaporisation of the prostate (CLVP)</p> <p>Nd:YAG laser. Power (w): CLVP 50-60. Performed under general or regional anaesthesia</p> <p>Group 2 : TURP Performed under general or regional anaesthesia.</p>	AUA symptom score	<p><u>At 1 month</u> Group1: 9.6 (n=20) Group 2: 11.0 (n=12)</p> <p><u>At 6 months</u> Group1: 9.1 (n=19) Group 2: 8.2 (n=10)</p> <p><u>At 12 months</u> Group1: 8.4 (n=18) Group 2: 4.7 (n=7)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method, allocation concealment and masking of outcome assessment were not reported Standard deviations were not reported. <p>Additional outcomes: Results for 5 patients that had CHRP (see notes).</p> <p>Notes: There was another group of patients (n=5) with prostate volumes >45 mL that underwent coagulation and haemostatic resection of the prostate (CHRP).</p> <p>2:1 randomisation method</p>
			Qmax	<p><u>At 1 month</u> Group1: 19.3 (n=20) Group 2: 21.4 (n=12)</p> <p><u>At 6 months</u> Group1: 20.0 (n=18) Group 2: 23.1 (n=10)</p> <p><u>At 12 months</u> Group1: 20.0 (n=18) Group 2: 26.9 (n=6)</p>	
			Transfusions	Group 1: 0/21 Group 2: 0/12	
			Re-catheterisation	Group 1: 3/21 (14.0%) Group 2: 3/12 (25.0%)	
			Urethral strictures	Group 1: 0/21 Group 2: 0/12	
			Reoperations:	Group 1: 0/21 Group 2: 0/12	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age (mean): 69.0 Drop outs: 5 (1 diagnosed with prostate cancer and had radical prostatectomy so not included in baseline data) IPSS: 24.7 Prostate size: 33.9 Qmax (mean) ml: 9.0 AUA symptom score (mean): 24.7 Mean length of stay (days): 2.5 ± NR Mean catheterisation time (days): 1.7 ± NR</p>				

Evidence Table 28: Laser vs. open prostatectomy

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Skolarikos et al., 2008 ²⁸¹ Study design: Randomised controlled trial Setting: Greece Evidence level: 1+ Duration of follow-up: 18 months	Patient group: Men recruited from March 2005 to April 2006. Inclusion criteria: Age > 50 years, LUTS due to BPH, prostate volume on TRUS >80cc, IPSS>12, medical therapy failure, no alpha blockers during the last month, no 5AR over the last 3 months, post void residual<150ml, peak urinary flow arte<12ml/sec. Exclusion criteria: neurogenic bladder, history of adenocarcinoma of the prostate, urethral stricture, previous prostatic, bladder neck or urethral surgery, no urethral catheter at baseline, history of bladder cancer, indwelling urethral catheter. All patients N: 125 Drop outs: NR Group 1 N: 65 Median (25-75 centile) Age: 74 (67-80) Group 2 N: 60 Median (25-75centile) Age:67.5 (65-74)	Group 1: Laser Photoselective vaporisation PVP) using high power potassium titanyl phosphate laser (KTP) PVP performed with an 80 watt KTP side-firing laser system. A flexible green light PV ADDStat fiber was used through a modified 23F continuous irrigation 12* Storz cystoscope. Isotonic saline used for irrigation. At end of procedure a 20F triple lumen catheter was inserted into the bladder for irrigation to start. Group 2: Open prostatectomy (OP) Transvesical approach used. At end of the procedure a 22F triple lumen catheter inserted into the bladder and irrigation was initiated. A suprapubic catheter was inserted whenever the surgeon thought extra irrigation needed.	Median (25-75 centile) Symptom score, IPSS	Baseline Group 1: 20 (15-22.5) Group 2: 21 (16.2-23.7); p=0.399 1 month Group 1: 12 (12-13.5) Group 2: 12 (10-16); p=0.019 3 months Group 1: 10 (8-12) Group 2: 10 (7-12); p=0.743 6 months Group 1: 9 (7-12) Group 2: 9 (7-12); p=0.224 12 months Group 1: 9 (7-12) Group 2: 8 (7-12); p=0.128 18 months Group 1: 10 (7-12) Group 2: 8.5 (7-12); p=0.063	Funding: NR Limitations: Patients significantly older at baseline in the laser group. Allocation concealment method unclear. Additional outcomes: 1, 3, 6, 12 month outcomes for prostate size, PSA, post void residual and IIEF scores. Notes: 5 laser patients the resectoscope was used at some point of the operation to achieve hemostasis. When optimal view restored, the KTP laser reused to finish operation.
			Median (25-75 centile) IPSS quality of life question	Baseline Group 1: 3 (2-4) Group 2: 3 (2.25-4) p=0.520 1 month Group 1: 2 (1-2) Group 2: 2 (1-2) p=0.283 3 months Group 1: 1 (1-2) Group 2: 2 (1-2) p=0.995 6 months Group 1: 1 (1-2) Group 2: 1 (0.25-1) p=0.024 12 months Group 1: 1 (1-2) Group 2: 1 (1-1) p=0.035 18 months Group 1: 1 (1-2) Group 2: 1 (1-1) p=0.001	
			Median (25-75 centile) Qmax, ml/s	Baseline Group 1: 8.6 (6.7-10.5) Group 2: 8 (5.8-10.2) p=0.283	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>1 month Group 1: 13.4 (10.7-15) Group 2: 12.5 (10.7-15) p=0.552</p> <p>3 months Group 1: 16 (14-18) Group 2: 15.1 (12.6-17) p=0.255</p> <p>6 months Group 1: 16 (13.9-18.8) Group 2: 15.6 (12.8-17.1) p=0.220</p> <p>12 months Group 1: 16 (13.7-19) Group 2: 15.1 (13-17.5) p=0.186</p> <p>18 months Group 1: 16 (13.5-18.9) Group 2: 15 (13-17.4) p=0.271</p>	
			Median (25-75 centile) PVR, ml	<p>Baseline Group 1: 97 (6-124) Group 2: 89 (50-120)</p> <p>18 months Group 1: 15 (0-33.5) Group 2: 12 (0-25); p=0.281</p>	
			Median (25-75 centile) IIEF-5	<p>Baseline Group 1: 12 (8-16) Group 2: 12 (7-16)</p> <p>18 months Group 1: 12 (7-17) Group 2: 12 (9-17); p=0.987</p>	
			Median (25-75 centile) P-size, ml	<p>Baseline Group 1: 93 (85-100) Group 2: 96 (86.2-100)</p> <p>18 months Group 1: 55 (45-65) Group 2: 10 (5.5-15); p<0.001</p>	
			Median (25-75 centile) PSA, ng/dl	<p>Baseline Group 1: 6.2 (3.1-8.44) Group 2: 6.3 (2.9-8.6)</p> <p>18 months Group 1: 2.4 (1.8-3.6) Group 2: 2 (1.4-2.6); p=0.025</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Median (25th-75th centile) Catheter removal (hours)	Group 1: 24 (20-36) Group 2: 120 (96-144); p< 0.001	
			Median (25th-75th centile) Hospital stay (hours)	Group 1: 48 (24-48) Group 2: 144 (120-144); p< 0.001	
			Median (25th-75th centile) Operation time (minutes)	Group 1: 80 (70-90) Group 2: 50 (45-60); p< 0.001	
			Number (%) Adverse events	Stress/urge incontinence Group 1: 0 Group 2: 0 Intra-operative TURP-hemotaxis Group 1: 5 (7.69) Group 2: 0 Peri-operative blood transfusion Group 1: 0 Group 2: 8 (13.3) Transurethral resection syndrome Group 1: 0 Group 2: NR Urethrogragia Group 1: 1 (1.54) Group 2: 0 Pulmonary infection Group 1: 0 Group 2: 1 (1.67) Prolonged dysuria Group 1: 5 (7.6) Group 2: 7 (11.6) Culture confirmed UTIs Group 1: 14 (21.5) Group 2: 16 (27) Re-catheterisation Group 1: 7 (10.7) Group 2: 10 (16.67) Re-operation Group 1: 3 (4.62); urethral strictures (2), persistent bladder outlet flow obstruction symptoms (1)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 3 (5); urethral stricture (1), bladder neck contracture (2) Mortality Group 1: 1 (liver cancer) Group 2: 0	

Evidence Table 29: Laser vs. transurethral microwave thermotherapy (TUMT)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Norby et al., 2002a²²³</p> <p>Study design: Randomised controlled trial (RCT)</p> <p>Evidence level: 1+</p> <p>Setting: Denmark (two centres)</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Men ≥ 50 years between May 1996 and November 1999.</p> <p>Inclusion criteria: IPSS ≥ 7, QoL ≥ 3, obstructed according to ICS nomogram or Qmax < 12 mL/s; able to understand project information and have written consent.</p> <p>Exclusion criteria: suspicion of prostate cancer; PVR > 350 mL or urinary catheter; prostatic urethra < 25 mm long, neurological disease or diabetes with abnormal cystometry; previous prostate operation; ongoing UTI; previous diagnosis of rectal cancer, intake of medication known to influence voiding; severe peripheral arterial insufficiency; previous pelvic radiation therapy; general health condition contraindicating surgery.</p> <p>All patients N: 118 Mean age: 66 Drop outs: 8 (6.7%)</p> <p>Group 1 N: 48 Mean age (SD): 65 (8) Median catheter duration: 3 days Median prostate volume, ml = 44 Dropouts: 4 (diagnosis changed for 3 and 2 declined surgery, of which one reported IPSS at 6m and included in</p>	<p>Group 1: LASER Interstitial laser coagulation. NdYag: 7-20W. Median length of stay was 3 days. Median catheter duration was 3 days</p> <p>Group 2: TUMT Transurethral microwave thermotherapy (TUMT). Prostatron 2.0 (n=8) or 2.5 (n=37). Performed as an outpatient procedure (four stayed overnight and 1 patient for 2 nights). Median catheter duration was 7-14 days</p> <p>Control: TUIP (n=3) or TURP (n=18). Median catheterisation was 2 days and hospital stay 5 days.</p>	<p>Mean (SD) IPSS:</p>	<p>Baseline: Group 1: 21.4 (5.8), n=44 Group 2: 20.5 (5.7), n=46 Group 3: 21.3 (6.6), n=22</p> <p>6 Months: Group 1: 9.5 (6.6), n=44 Group 2: 9.5 (7.1), n=44 Mean difference: 0.00 [-2.86, 2.86] Group 3: 6.8 (5.7), n=22</p>	<p>Funding: Supported by a grant from Vejle County, Denmark.</p> <p>Limitations: Had to stop early due to financial restrictions and did not reach target enrolment population.</p> <p>Additional outcomes: - Effect on prostatic volume. - Results also compared to control group that had either TURP or TUIP. - Overall satisfaction scores reported in comparison to control group. Figures not provided. - Subgroup analysis comparing results from TUMT 2.0 v TUMT 2.5.</p> <p>Notes: Reported in Cochrane Systematic Review by Hoffman 2000.</p> <p>UTI defined as 'symptomatic UTI requiring antibiotic treatment (infections treated both in the outpatient clinical and in primary health care were included)'. * Erectile dysfunction and retrograde ejaculation was only estimated amongst those who had answered</p>
			<p>Median (IQR) IPSS Quality of life:</p>	<p>Baseline: Group 1: 4 (4-4), n=44 Group 2: 4 (4-4), n=46 Group 3: 4 (4-5), n=22</p> <p>6 Months: Group 1: 1 (1-2), n=44 Group 2: 2 (1-3), n=44 Group 3: 1 (1-2), n=22</p>	
			<p>Mean (SD) peak urinary flow (Qmax mL/s):</p>	<p>Baseline: Group 1: 10.2 (4.0), n=44 Group 2: 9.1 (4.2), n=46 Group 3: 9.6 (3.2), n=22</p> <p>6 Months: Group 1: 16.2 (8.5), n=43 Group 2: 13.2 (6.9), n=44 Group 3: 20.6 (12.8), n=22</p>	
			<p>Median (IQR) post void residual, mL</p>	<p>Baseline: Group 1: 117 (50-180), n=44 Group 2: 110 (50-210), n=46 Group 3: 75 (17-193), n=22</p> <p>6 Months: Group 1: 58 (14-118), n=43 Group 2: 48 (24-129), n=44 Group 3: 23 (3-48), n=22</p>	
<p>Urinary retention:</p>	<p>Group 1: 4/44 (9%) Group 2: 3/46 (7%) Group 3: 1/22 (5%)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>results).</p> <p>Group 2 N: 46 Mean age (SD): 66 (7) Median catheter duration: 7-14 days; with longer catheterisation required after higher energy procedures. Median prostate volume, ml = 43 Drop outs: 2 (one had TURP, other had apoplexy at 4m and only had 3m follow-up)</p> <p>Group 3 N: 24 Mean age (SD): 68 (7) Median prostate volume, ml = 44 Drop outs: 2 (prostate cancer)</p>		<p>Urinary tract infection:</p> <p>Transurethral resection syndrome (TUR)</p> <p>Transfusion:</p> <p>Stricture:</p> <p>Urinary incontinence:</p> <p>Development of erectile dysfunction:*</p> <p>Development of retrograde ejaculation:</p> <p>Reoperation for BPO</p> <p>Mortality</p>	<p>Group 1: 27/44 (61%) Group 2: 14/46 (30%) Group 3: 3/22 (14%)</p> <p>Group 1: 0/44 (0%) Group 2: 0/46 (0%) Group 3: 1/22 (5%)</p> <p>Group 1: 0/44 (0%) Group 2: 0/46 (0%) Group 3: 2/22 (9%)</p> <p>Group 1: 1/44 (2%) Group 2: 0/46 (0%) Group 3: 1/22 (5%)</p> <p>Group 1: 0/44 (0%) Group 2: 0/46 (0%) Group 3: 1/22 (5%)</p> <p>Group 1: 4/18 (29%) Group 2: 2/22 (9%) Group 3: 1/7 (14%)</p> <p>Group 1: 9/26 (35%) Group 2: 6/27 (22%) Group 3: 7/14 (50%)</p> <p>Group 1: 0/44 (0%) Group 2: 1/46 (2%) Group 3: 0/22 (0%)</p> <p>Group 1: 0/44 (0%) Group 2: 0/46 (0%) Group 3: 0/22 (0%)</p>	<p>the relevant questions both at baseline and at the 6 month follow-up. Each question was scored from 0 to 3. For evaluation of ejaculation, patients scoring 0, 1 and 2 (i.e. normal amount, slightly reduced and greatly reduced amount of semen) were classified as having antegrade ejaculation. Patients scoring 3 (i.e. no ejaculation) were classified as having retrograde ejaculation.</p>

Evidence Table 30: Laser vs. transurethral vaporisation of the prostate (TUVP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Abdelkhalek et al., 2003⁴</p> <p>Study design: RCT, open label</p> <p>Setting: Egypt</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 4 years</p>	<p>Patient group: Symptomatic bladder outlet obstruction due to BPH</p> <p>Setting: Urology and Nephrology Centre, Mansoura University, Egypt. (March 1995 to March 1997)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ▪ Qmax ≤10ml/s ▪ Serum PSA level of < 4 ng/mL ▪ IPSS of ≥15 ▪ Prostate volume of 20-80mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ Urethral stricture ▪ Contracted bladder ▪ Large vesicle diverticulum ▪ Neuropathic bladder <p>All patients N: 180 Age, mean ±SD Drop outs: 40/180</p> <p>Group 1-Laser prostatectomy N: 90 Dropouts: 28/90</p> <p>Age, mean (years): 63.3±6.5 IPSS, mean (±SD): 27.9±5.3 IPSS-QoL, mean (±SD): 5±0.8 Qmax, mean, (±SD): 6.9±2.8 Post void residual urine, mean,</p>	<p>Group 1- Laser prostatectomy: combination of coagulation and vaporisation methods:</p> <p>i) Side firing coagulation of two lateral lobes using fibres with a lateral beam angle of 90° at 40W for 90s at each coagulation spot in the 2, 4, 8, 10 and 12 o'clock positions.</p> <p>ii) Vaporisation of the median lobe using contact (sapphire) tips at 60W in a retrograde fashion.</p> <p>Power: 40W Nd: YAG for 60s at each lateral lobe at 9 and 3 o'clock positions, and 30s each at 6 and 12 o'clock positions.</p> <p>Group 2 –TUVP Procedure: TUVP delivered using Vaportrode™ under the 250 to 300 W of pure cutting current in an antegrade fashion. The median lobe was vaporised first, and continued down the</p>	<p>All cause mortality (due to cardiopulmonary disease)</p> <p>IPSS, mean± SD:</p> <p>IPSS-QoL mean ± SD:</p>	<p>Group 1: 1/90 Group 2: 2/90 P value: NS</p> <p><u>At 1 year</u> Group 1: 13.3±6 Group 2: 5.6±3.5 p value: 0.003</p> <p><u>At 2 year</u> Group 1: 12.2±5.6 Group 2: 5.2±3.3 p value: 0.006</p> <p><u>At 3 year</u> Group 1: 13.1±5.7 Group 2: 4.8±2.6 p value: 0.002</p> <p><u>At 4 year</u> Group 1: 11.9±6.1 Group 2: 3.7±1.3 p value: <0.001</p> <p><u>At 1 year</u> Group 1: 3.4±0.4 Group 2: 1.4±0.5 p value: 0.008</p> <p><u>At 2 year</u> Group 1: 3.2±0.5 Group 2: 1.4±0.4 p value: 0.009</p> <p><u>At 3 year</u> Group 1: 3.3±0.6 Group 2: 1.4±0.5 p value: 0.009</p> <p><u>At 4 year</u> Group 1: 3.1±1.0 Group 2: 1.3±0.5 p value: <0.001</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Open label study with subjective patient reported outcomes. ▪ Randomisation and concealment methods not reported <p>Additional outcomes:</p> <ul style="list-style-type: none"> ▪ Prostate and adenoma volume at 1 and 4 years ▪ An additional 6 and 2 reoperations were completed for the laser and TUVP groups respectively after the 4-year follow up. <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	(\pm SD): 120 \pm 97.5 Prostate volume , mean (\pm SD):43.8 \pm 13.4 Group 2 - TUVP N: 90 Dropouts: 12/90 Age , mean (years): 62.9 \pm 5.9 IPSS , mean (\pm SD): 26.0 \pm 5.8 IPSS-QoL, mean (\pm SD): 4.8 \pm 0.9 Qmax , mean, (\pm SD): 6.4 \pm 2.5 Post void residual urine , mean, (\pm SD): 125 \pm 97.5 Prostate volume , mean; 47.4 \pm 16.1	surgical capsule until a wide prostatic cavity was created, followed by careful coagulation.	Qmax (ml/s), mean \pm SD: Post void residual volume (ml), mean \pm SD Post-op complications: Bleeding at surgery (definition not provided) Post-op complications: Haematuria Post-op complications: urinary retention Post-op complications: Urethral Stricture (urethral stricture, apparent after 6 months) Post-op complications: Bladder neck stenosis	At baseline Group 1: 6.9 \pm 2.8 Group 2: 6.4 \pm 2.5 p value: 0.256 At 1 year Group 1: 15.1 \pm 6.0 Group 2: 20.8 \pm 7.4 p value: 0.029 At 4 year Group 1: 13.6 \pm 3.6 Group 2: 21.4 \pm 4.1 p value: <0.001 At 1 year Group 1: 61.3 \pm 49.2 Group 2: 22.1 \pm 22 p value: <0.001 At 4 years Group 1: 64.6 \pm 29.8 Group 2: 25.1 \pm 12.8 p value: <0.001 Group 1: 0/90 Group 2: 1/90 p value: NS Group 1: 0/90 Group 2: 2/90 p value: NS Group 1: 9/90 Group 2: 2/90 p value: NS Up to 1 year Group 1: 0/90 Group 2: 2/90 p value: NS Up to 1 year Group 1: 2/90 Group 2: 2/90 p value: NS	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Post-op complications: Retrograde ejaculation	<u>At 1 year</u> Group 1: 16/90 Group 2: 57/90 p value: <0.001	
			Post-op complications: Impotence (among patients who were potent at baseline)	<u>At 1 year</u> Group 1: 0/49 Group 2: 4/53 p value: 0.04	
			Post -op complications: Reoperation (cumulative) Details of type reoperation provided.	<u>At 1 year</u> Group 1: 10/89 Group 2: 3/889 p value: 0.04 <u>At 2 year</u> Group 1: 18/90 Group 2: 5/90 p value: <0.05 <u>At 3 year</u> Group 1: 27/90 Group 2: 8/90 p value: <0.05 <u>At 4 year</u> Group 1: 35/90 Group 2: 11/90 p value: <0.001	
			Operation time, mean (range), (min):	Group 1: 37.5±15 Group 2: 36.6±16.4 p value: NS	
			Catheter period (days)mean ±SD	Group 1: 6.8 (0.9) Group 2: 2.3 (0.5) p value: <0.001	
			Length of hospital stay, (days) mean ±SD	Group 1: 1.1±0.5 Group 2: 2.2±0.8 p value: NS	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Shingleton et al., 1998 ²⁷⁶ Study design: RCT Setting: USA Evidence level: 1+ Duration of follow-up: 6 months	Patient group: consecutive patients with benign prostatic hyperplasia Inclusion criteria: Consecutive patients (no further information) Exclusion criteria: Not stated All patients N: 31 Randomised (ratio 2:1) Group 1 N: 11 Mean (range) Age: 67.5 (60-82) Mean prostate volume (cc): 34.6 (9.2 to 87.7) Erectile function: Full: 3/11 (27%) Partial: 5/11 (45%) None: 3/11 (27%) Dropouts: Not stated Group 2 N: 20 Mean (range) Age: 66.7 (48-77) Mean prostate volume (cc): 34.6 (13.7 to 66.4) Dropouts: Not stated Erectile function: Full: 4/20 (25%) Partial: 7/20 (35%) None: 9/20 (47%)	Group 1: VLAP + KTP (contact laser – vaporisation) KTP laser set at 40 watts for initial vaporisation of all median and lateral lobe tissue. Nd:YAG beam used at 60 watts for 60 sec to create a series of craters in lateral lobes of the prostate. Catheter protocol: Catheter put in place without accompanying bladder irrigation. Group 2: Transurethral Electro vaporisation (TVP) High energy electrical current to vaporise tissue and create a zone of coagulation surrounding vaporised tissue cavity. Catheter protocol Set at initial 275 watts, but increased to 300 watts in all patients. The coagulation setting was 40watts for all patients. Catheter protocol: After procedure a 22F three way catheter was put in place and standard irrigation with normal saline begun.	AUA symptom score, mean (range)	Baseline: Group 1: 19 (13-27) Group 2: 22.1(8-31) 3 months: Group 1: 5.9 (1-12) Group 2: 5.2 (2-24) 6 months: Group 1: 5.0 (0-10) Group 2: 5.2 (1-19) P value: NS between arms, stat sig compared to baseline	Funding: NR Limitations: <ul style="list-style-type: none"> Randomisation allocation and concealment not reported No specific inclusion or exclusion criteria were stated in this paper. No statistical methods provided. Additional outcomes: <ul style="list-style-type: none"> 1 month outcomes % of patients who had improved more than 50 % compared to baseline at 6th month follow up Notes: QoL was reported to be collected in method section but was not reported. Shingleton1998A – reported on the urodynamics outcome of a subset of the patients in this cohort (10 patients in each arm). However, the basis of selecting this subset of patients was not
			Qmax, mean (range)	Baseline: Group 1: 10.7 (0-11.8) Group 2: 7.7 (3.4-13.2) 3 months: Group 1: 17.6 (6.2-22) Group 2: 17.5 (7.6-24.9) 6 months: Group 1: 16.5 (7.1-24.9) Group 2: 14.3 (7.8-27.1) P value: NS for all P value: NS between arms, stat sig compared to baseline	
			Post-op complications: Clot retention	Group 1: 0/11 Group 2: 2/20 p value: NS	
			Post-op complications: haematuria (2 patient in laser group had clot retention)	Group 1: 2/11 Group 2: 6/20 p value: NS	
			Post-op complications: Post operative urinary retention	Group 1: 3/11 Group 2: 1/20 p value: NS	
Stricture (urethral stricture)	Group 1: 1/11 Group 2: 0/20 p value: NS				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Post-op complications: Development of erectile dysfunction	Group 1: 1/11 Group 2: 2/20 p value: NS	provided. Inclusion/exclusion criteria from Shingleton1998A Inclusion: >45 years, Qmax <15ml, no history of carcinoma and ability to undergo general anaesthesia.
Operation time, mean, (min):	Group 1: 27.5 Group 2: 46 p value: <0.05				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Van Melick et al., 2003³⁰⁸</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 7 years:</p>	<p>Patient group: men over 45 years with LTUS associated with BPH that were recruited from their clinic from 1996 to 2001</p> <p>Setting: Netherlands</p> <p>Inclusion Criteria: patient with lower urinary tract symptoms suggestive of BPH; met ISC criteria for BPH, Schafer obstruction score ≥ 2, prostate size between 20-65ml.</p> <p>Exclusion Criteria: age ≤ 45 yrs</p> <p>All patients N: 141</p> <p>Group 1 N: 45 Age (mean): 67\pm9 Drop outs: 8 at one year post-operatively (procedure during surgery changed for medical reasons=3, equipment failure resulting in TURP)=2, reoperation – TURP=1, reoperation – due to stricture =2) Mean prostate size, ml: 37\pm11 Follow-up 1 to 4 years = 15 Follow-up 4 to 7 years=15</p> <p>Group 2 N: 50 Age (mean): 66\pm8 Drop outs: 9 at one year post-operatively (surgery cancelled=1, mortality=2, morbidity=2, emigrated=1, reoperation (TURP) =2, reoperation (stricture)=1) Mean prostate size, ml: 38\pm9 Follow-up 1 to 4 years = 10</p>	<p>Group 1: Laser vaporisation Transurethral catheter post-operation SLT Nd:Yag Pre-procedural antibiotics and transurethral catheter postoperatively.</p> <p>Group 2: TURP Suprapubic catheter if required peri-operatively. Pre-procedural antibiotics and transurethral catheter postoperatively.</p> <p>Group 3: Electro vaporisation Performed with a Vaportrode element using glycine for irrigation. Pre-procedural antibiotics and transurethral catheter postoperatively.</p>	<p>Mean (SD) symptom score (IPSS)</p> <p>Mean (SD) Global quality of life score:</p>	<p><u>At baseline:</u> Group 1: 18.3\pm8.2 Group 2: 16.6\pm5.6 Group 3: 20.3\pm6.8</p> <p><u>At 6 months</u> Group 1 (n=33): 5.9\pm5.5 Group 2 (n=37): 3.2\pm2.7 Group 3: 3.8\pm2.7</p> <p><u>At 1 year</u> Group 1 (n=37): 3.6\pm3.4 Group 2 (n=41): 4.1\pm4.8 Group 3: 4.8\pm4.9</p> <p><u>At 1-4 years</u> Group 1 (n=10): 9.3\pm5.2 Group 2 (n=15): 5.8\pm7.5 Group 3: 8.4\pm8.7</p> <p><u>At 4-7 years</u> Group 1 (n=17): 8.3\pm6.4 Group 2 (n=15): 7.3\pm7.1 Group 3: 7.0\pm5.6</p> <p><u>At baseline:</u> Group 1: 3.6\pm1.6 Group 2: 3.9\pm1.6 Group 3: 4.3\pm1.3</p> <p><u>At 6 months</u> Group 1: 0.8\pm1.0 Group 2: 0.5\pm0.5 Group 3: 1.0\pm0.8</p> <p><u>At 1 year</u> Group 1: 0.6\pm0.9 Group 2: 0.6\pm0.8 Group 3: 1.0\pm0.9</p> <p><u>At 1-4 years</u> Group 1: 2.0\pm1.0 Group 2: 1.1\pm1.2 Group 3: 1.0\pm1.2</p> <p><u>At 4-7 years</u> Group 1: 1.4\pm1.2 Group 2: 1.3\pm1.3 Group 3: 1.4\pm0.8</p>	<p>Funding: NR.</p> <p>Limitations: Open label study</p> <p>Additional outcomes: Frequency during day, frequency during night, symptom problem index and BPH impact index. Uroflowmetry also reported.</p> <p>Notes: Links with Van Melick 2002³⁰⁷, Van Melick 2003³⁰⁸.</p> <p>Follow up time varied individually as all patients were analysed within a 2 month period. Depending on the individual follow-up time, patient divided into two groups: those with a follow-up time between 1 and 4 years and those with follow up time between 4 and 7 years.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments	
	Follow-up 4 to 7 years=17 Group 3 N: 46 Age (mean): 64±10 Drop outs: 22 Mean prostate size, ml: 35±12 Follow-up 1 to 4 years = 12 Follow-up 4 to 7 years=12		Mean (SD) maximal flow (mL/s)	At baseline: Group 1: 9±3 Group 2: 13±4 Group 3: 9±3 At 6 months Group 1: 25±9 Group 2: 26±6 Group 3: 24±11 At 1 year Group 1: 27±12 Group 2: 23±10 Group 3: 28±6 At 1-4 years Group 1: 19±6 Group 2: 20±5 Group 3: 23±6 At 4-7 years Group 1: 19±9 Group 2: 17±8 Group 3: 16±11		
			Stricture	Group 1: 2/45 Group 2: 2/50 Group 3: 1/46		
			Incontinence Reported in HTA (ncc study)	Group 1: 14/45 (8%) Group 2: 4/50 (39%) Group 3: 15%		
			Reoperation by TURP	Group 1: 1/45 Group 2: 2/50 Group 3: 2/46		
			Blood transfusion	Group 1: 0/45 Group 2: 1/50 Group 3: 0/46		
			Urinary retention	Group 1: 5/45 Group 2: 0/50 Group 3: 0/46		
			Urinary tract infection (after one week)	Group 1: 4/45 (9%) Group 2: 5/50 (10%) Group 3: 5%		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean (SD) operative time, minutes:	Group 1: 58 (11) Group 2: 58 (26) Group 3: 50 (16)	
			Mean (SD) postoperative hospital days	Group 1: 3.8 (1.3) Group 2: 3.9 (0.9) Group 3: 3.4 (0.9)	
			Mortality: *cardiac failure, hepatic failure (HTA reports 3 v 4)	Group 1: 0/45 Group 2: 2/50* Group 3: 0/46	

Evidence Table 31: Laser coagulation vs. laser vaporisation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Bryan et al, 2000 ³⁷ Study design: RCT, single centre – open study Evidence level: 1+ Duration of follow-up: 2 years	Patient group: Bladder outlet obstruction, BOO due to benign prostatic hyperplasia, BPH. Setting: Urology department, UK hospital Inclusion Criteria: Ambulant male patients with BOO due to BPH, confirmed with pressure/flow urodynamics. Exclusion Criteria: <ul style="list-style-type: none"> ▪ Neurological disorders affecting the urinary tract ▪ Previous prostatic or urethral surgery ▪ Clinical evidence of prostatic or vesicle malignancy ▪ Acute urinary tract infection ▪ Prostate gland volume of <20mm³ On medication known to influence voiding function. All patients N: 38 Drop outs: 0 Group 1 - CLAP N: 21 Age (mean): 72.25, SE1.68 Drop outs: 0	Laser prostatectomy was carried out using a SLT (Surgical Laser Technologies, Oaks, Pa, USA) neodymium:YAG laser system with semi-rigid endoscopic fibre (SREF15) set a 40W Group 1-CLAP A chiselled probe (MD6) with a distal end incorporating a 6 mm sapphire tipped round probe was used. The probe was brought back to the verumontanum and then pushed forward to produce furrows. Mean operating time:37.7min SEM1.6 Group 2 - VLAP Laser energy applied using a side firing free beam probe (SFB 1.0), to the lateral lobes 1 cm distal	IPSS symptom score The data was shown in a graph, and values only reported for 6 th and 24 th month.	<u>At 1, 3, 12th months</u> Group1 : No reported Group 2 : NR P value: NS <u>At 6 months</u> Group1: 8.3 ± 6.4*** Group 2: 12.5** ± 6.4*** p value: 0.05 <u>At 24 months</u> Group1: 13.5 ± 8.26* Group 2: 13.3 ± 7.36* p value: NS <u>Compared to baseline</u> Group 1: P value= 0.006 Group 2: P value= 0.002	Funding: Not stated Limitations: <ul style="list-style-type: none"> ▪ No sample size calculation provided- small sample size ▪ 38% in CLAP and 24% in VLAP group did not perform urodynamics at 6 months to determine obstruction Additional outcomes: <ul style="list-style-type: none"> ▪ Mean operating time ▪ Increased irritative symptoms which returned to normal after 1 month (5 in VLAP, 4 in CLAP) Notes: *SD estimated following the Cochrane handbook method using p values reported for change from baseline. ** estimated from graph shown. Likely error in the value from text (21.3) ***SD estimated from standard error bars
			Qmax	<u>At 12 months</u> Group1: 16.6 ± 7.37* Group 2: 17.5 ± 6.50* P value: NS <u>Compared to baseline</u> Group 1: P value= 0.006 Group 2: P value= 0.002 <u>At 24 months</u> Group1: 15.5 ± 7.35* Group 2: 15.9 ± 10.15* P value: NS <u>Compared to baseline</u> Group 1: P value= 0.02 Group 2: P value= 0.1	
			PdetQmax (cm H₂O)	<u>At 6 months</u> Group1: 54.6 Group 2: 56.4 p value: 0.4 Both Sig different compared to baseline p<0.005	
			Post-op complications (early): Catheter duration, mean (range), days	Group1: 4.5(1-31) Group 2: 13.2 (7-70) p value: NR#	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>IPSS: 20.9, SE1.6 Erectile dysfunction: 10, SE 21 (47.6%) Qmax:10.0, SE 0.68 PdetQmax H₂O: 79.4, SE 9.4 Unequivocal obstruction, proven urodynamically: 19/21</p> <p>Group 2 - VLAP N: 17 Age (mean): 71.88, SE 1.59 Drop outs: 0 IPSS: 21.8, SE 1.5 Qmax:10.0, SE 0.8 PdetQmax H₂O: 91.9, SE 9.8 Erectile dysfunction: 8/17 (47.1%) Unequivocal obstruction, proven urodynamically: 16/17</p>	<p>to the bladder neck at 40W for 90s each of 4 quadrants: 2, 4, 8, and 10 o' clock positions.</p> <p>Mean operating time: 24.5min</p>	<p>Post-op complications (early): Required Catheter > 7 days</p> <p>Post-op complications (early): Bladder irrigation</p> <p>Post-op complications (early): Blood transfusion</p> <p>Post-op complications (early): Peri-operative urinary tract infections</p> <p>Post-op complications: Developed erectile dysfunction</p> <p>Post-op complication: Reoperation:</p> <p>Unequivocal obstruction, proven urodynamically, at 6 months \$</p>	<p>Group1: 2/21 Group 2: 7/17 Relative risk: NS</p> <p>Group1: 5/21 Group 2: 0/17 Relative risk: 9.00 95% CI: 0.53-152.1 p value: NS</p> <p>Group1: 1/21 Group 2: 0/17 Relative risk: 2.45 95% CI: 0.11-56.7 p value: NS</p> <p>Group1: 1/21 Group 2: 2/17 Relative risk: 0.40 95% CI: 0.04-4.09 p value: NS</p> <p>Group1: 1/21 Group 2: 1/17 Relative risk: 0.81 95% CI: 0.05-12.01 p value: NS</p> <p>Group1: 1/21 Group 2: 2/17 Relative risk: 0.40 95% CI: 0.04-4.09 p value: NS</p> <p>Group1: 3/13 Group 2: 6/13 Relative risk: 0.50 95% CI: 0.16-1.58 p value: NS</p>	<p>from graph because p value for change from baseline was not reported in the results #No SD provided \$ 9 in the CLAP and 4 in the VLAP group were infirm or refused to do urodynamics at 6 months post-op</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Narayan et al., 1995²⁰⁹</p> <p>Study design: RCT, multi-centre, open study</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Moderate to severe obstruction, including 8 patients in chronic retention and had indwelling Foley catheter*</p> <p>Setting: US, in two Veteran Affairs medical centres</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Consecutive patients with moderate to severe obstructive symptoms as defined by AUA symptom score ≥ 13 (midway of the scale between mild and moderate obstructive symptoms) $Q_{max} < 15 \text{ ml/s}$, with or without significant post void residual volume <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Prostate cancer <p>All patients N: 64 Drop outs:</p>	<p>Group 1 CLAP- Evaporation Standard cystourethroscopy was performed before laser ablation.</p> <p>Laser applied initially at the 5 and 7 o'clock position at 60W until circular fibres of the bladder neck visible. Next, the median lobe was treated with laser at 45 degrees angle from the lobe from the right to left sides and vice versa. The ablation was completed by laser application at the 6 o'clock position deep enough to visualise the bladder neck muscle fibres and a smooth, bladder neck between 5 and 7 o'clock positions.</p> <p>Prostate evaporation was then performed. Fibre held <u>in contact</u> with area treated and dragged at rate of 1 cm/20 to 30s. At the beginning each furrow dragging was commenced when bubbling was noted signifying evaporation of</p>	<p>IPSS symptom score, only mean value reported, no standard deviation provided</p> <p>Q_{max} (ml/s), only mean value reported, no standard deviation provided</p>	<p><u>At 1 months</u> Group 1: 9.9 Group 2: 9.8</p> <p><u>At 3 months</u> Group 1: $7.0 \pm 14.81^*$ Group 2: $8.4 \pm 13.18^*$</p> <p><u>At 6 months (N=52)</u> Group 1: $5.0 \pm 16.73^*$ Group 2: $5.1 \pm 16.35^*$</p> <p><u>At 12 months (N=15)</u> Group 1: $5.3 \pm 16.45^*$ Group 2: $5.2 \pm 16.25^*$</p> <p>P value: NR, not sig between arms at all time points (All $P < 0.001$ compared to baseline)</p> <p><u>At 1 months</u> Group 1: 17 Group 2: 12.0</p> <p><u>At 3 months</u> Group 1: $19.7 \pm 12.79^*$ Group 2: $16.3 \pm 14.00^*$</p> <p><u>At 6 months (N=52)</u> Group 1: $20.0 \pm 13.08^*$ Group 2: $16.4 \pm 9.04^*$</p> <p><u>At 12 months (N=15)</u> Group 1: $19.9 \pm 12.98^*$ Group 2: $16.9 \pm 11.46^*$</p> <p>P value: < 0.05 for all time points. (All $P < 0.05$ compared to baseline)</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> No mention of blinding of outcomes assessors. Relatively small sample size- not sample size calculation provided. There was a trend (not statistically significant) of older patients, with larger prostate size, higher number in retention, lower Q_{max} and higher post void residual volume in the evaporation group. Most continuous variable outcomes only reported mean values- not standard

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 1 – CLAP-evaporation N: 32 Age (mean, range): 66.0(49-78) Prostate volume (mean, range); 51.7(16-120) N patient in retention: 6/32 Median lobe: 5/32 <u>Data excluding patients with chronic urinary retention (n=26):</u> AUA symptom score: 22.4(14-35) Qmax: 6.4(0-15) Post void residual volume: 276.6(20-960)</p> <p>Group 2 – VLAP-Coagulation N: 32 Age (mean, range): 64.1(48-92) Prostate volume (mean, range); 41.4 (20-62) N patient in retention: 3/32 Median lobe: 4/32 <u>Data excluding patients with chronic urinary retention (n=29):</u> AUA symptom score: 22.1(15-30) Qmax: 70(0-14) Post void residual volume: 210(0-250)</p> <p>* Patients who were in chronic retention were assigned “0” Qmax and not assigned any AUA score. These results were analysed separately.</p>	<p>tissue. Dragging the fibre at this rate resulted in furrow 5-7 mm deep and with a 3-4mm rim of coagulated tissue.</p> <p>Group 2 VLAP-Coagulation (modified visual laser ablation technique) Laser application at 60W for 60s to 11-19 spots (depending on prostate size). Spots included 5 and 7 o’clock positions at the bladder neck, the 6’ o clock position for the median lobe and the 5, 7, 11, and 1 o’clock position for each cm length of the prostate. Each spot covered a 1 cm area.</p> <p>Fibre held 2-4 mm away from tissue to ensure coagulation and not evaporation.</p> <p><u>Antibiotic prophylaxis:</u> All patients received cefazolin 1g/ml perioperatively and trimethoprim-sufamethoxazole double strength twice daily; one hospital provide 24-48 hours of prophylaxis whereas another provided 10 days</p>	<p>Post void residual volume (ml), only mean value reported, no standard deviation provided</p> <p>Catheter duration, Median (range), days</p> <p>Post-op complications (early): Blood transfusion</p> <p>Post-op complications (early): Epididymitis</p> <p>Peri-operative urinary tract infections (patients operated in 2 hospitals, all perioperative UTIs in hospital which only provide 24-48 of prophylaxis.</p> <p>Post-op complications: Developed erectile dysfunction</p> <p>Post-op complications: Incontinence</p>	<p><u>At 1 months</u> Group 1: 49 Group 2: 46 <u>At 3 months</u> Group 1: 31 Group 2: 20 <u>At 6 months(N=52)</u> Group 1: 29 Group 2: 24 <u>At 12 months (N=15)</u> Group 1: 26 Group 2: 28 P value: NR, not sig between arms at all time points (All P<0.05 compared to baseline)</p> <p>Group 1: 1.9 (1-10) Group 2: 2.1 (1-21) p value: NS</p> <p>Group 1: 0/32 Group 2: 0/32 p value: NS</p> <p>Group 1: 0/32 Group 2: 0/32 p value: NS</p> <p>Group 1: 2/32 Group 2: 1/32 Relative risk: 2.00 95% CI: (0.19-20.97) p value: NS #</p> <p>Group 1: 0/32 Group 2: 0/32 p value: NS</p> <p>Group 1: 0/32 Group 2: 0/32 p value: NS</p>	<p>deviation.</p> <p>Additional outcomes: Qmax, AUA symptom score and post void residual volume for 8 patients in chronic retention analysed and reported separately. There was no significant difference in terms of improvement in AUA symptom score or Qmax.</p> <p>Notes: # Calculated by NCGC team using Mantel Haenszel test in Rev Man version 5. Values reported in paper were based on chi-square test (Pearson)</p> <p>*SDs estimated following Cochrane methods using p values for change from baseline</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Post-op complication: Reoperation:	Group 1: 0/32 Group 1: 5/32 Relative risk: 0.09 95% CI: 0.01-1.58 p value: NS	
			Post-op complication: Post operative retention (Longer than 7 days after catheter removal)	Group 1: 2/32 Group 2: 8/32 Relative risk: 0.25 95% CI: 0.06-0.94 p value: <0.05#	
			“Bothersome irritative symptoms” > 14 days	Group 1: 10/32 Group 2: 11/32 Relative risk: 0.87 95% CI: 0.31-2.47 P value: NS	

See Evidence Table 26: Laser coagulation vs. transurethral resection of the prostate (TURP)

Evidence Table 32: Holmium laser resection of the prostate (HoLRP) vs. laser coagulation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gilling et al., 1998¹⁰²</p> <p>Study design: RCT, open study</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Men with symptomatic benign prostatic hyperplasia</p> <p>Setting: Urology department, New Zealand</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> Qmax ≤15ml/s AUA symptom score >8 Urodynamically proven bladder outlet obstruction – defined as Schaefer grade of ≥2 and at detrusor pressure at peak flow (PdetQmax) value in the obstructed or equivocal region of Abrams-Griffiths nomogram <p>Exclusion Criteria:</p> <ul style="list-style-type: none"> Age ≥85 years Prostate volume (measured by TRUS), >100ml <p>All patients N: 44 Drop outs: 0</p> <p>Group 1 -HoLRP N: 22 Drop outs: <i>All values provided as mean (range)</i> Age : 64 (44-81) IPSS: 24(14-33) Qmax, ml/s: 8(3-15) PVR (TRUS volume), mL: 42(20-72) PdetQmax H₂O: 72(37-117) Shaffer Grade: 4 (2-5) Residual volume: 179 (30-40)</p>	<p>Group 1- HoLRP Retrograde approach to the incision of the first and median lobe and then each lateral lobe in turn. This was performed using a 550micrometer bare quartz fibre passed down a continuous-flow resectoscope. Power setting was 60W.</p> <p><u>Energy (kJ), mean (range):</u> 67 (32-165) <u>Mean lasing time, mean (range)*:</u> 27.2min (13-75) <u>Resection weight, g, mean (range):</u> <u>Estimated:</u> 21(10-60) <u>Actual :</u> 5 (2-13)</p> <p>Catheter removed at 6 the following morning and discharged once voided successfully.</p> <p>Group 2 - VLAP Standard 4-quarant Nd:YAG lasing technique. A total of at least 1kJ/g of measures tissue was delivered using a 60W for 60s at each treatment site.</p> <p><u>Energy (kJ), mean (range):</u> 53 (25-102) <u>Mean lasing time, mean (range)*:</u> 27.2min (13-75) <u>Resection weight, g, mean</u></p>	<p>IPSS symptom score, mean (range). All not sig between treatment arms.</p>	<p><u>At 1 month</u> Group1: 8(0-16) Group 2: 11(2-26) p value: Not Sig</p> <p><u>At 3 months</u> Group1: 4(0-12) Group 2: 8(0-26) p value: Not Sig</p> <p><u>At 6 months</u> Group1: 5(1-16) Group 2: 7(0-22) p value: Not Sig</p> <p><u>At 12 months</u> Group1: 4(0-9) Group 2: 5(1-18) p value: Not Sig</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> No details of randomisation method and concealment was provided Small sample size- sample size calculation not provided Open study <p>Additional outcomes:</p> <ul style="list-style-type: none"> % of men requiring analgesia for dysuria symptoms (64% VLAP, 41% for HoLRP) Mean duration of surgery – stats sig <p>Notes: None.</p>
			<p>Dysuria score , mean, (no SD given) Measured using a visual analogue scale (VAS), ranging from 0 (no voiding symptom), 10 (severe dysuria)</p>	<p>First 10 post-operative days Group1: 2 Group 2: 4 p value: <0.05 <u>First 5 days after catheter removal</u> Group1: 2.1 (Day 1- 5) Group 2: 3.7 (Day 6-10) p value: <0.05</p>	
			<p>Qmax, mL/s, mean (range)</p>	<p><u>At 1 months</u> Group1: 21(10-56) Group 2: 13(4-27) p value: <0.01</p> <p><u>At 3 months</u> Group1: 20(12-30) Group 2: 15(5-27) p value: <0.05</p> <p><u>At 6 months</u> Group1: 21(12-32) Group 2: 15(5-24) p value: <0.01</p> <p><u>At 12 months</u> Group1: 22(8-41)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Prostate length, cm: 3(2-5) Group 2 - VLAP N: 22 Drop outs: 0 <i>All values provided as mean (range)</i> Age: 68(45-80) IPSS: 23(13-35) Qmax, ml/s: 8(3-15) PVR (TRUS volume), mL: 49(24-80) PdetQmax H₂O: 77(42-113) Shaffer Grade: 4 (2-5) Residual volume: 131 (40-227) Prostate length, cm: 3(2-6)	(range): <u>Estimated:</u> 24(5-60) <u>Actual</u> : not stated All patients discharged the morning after surgery. Catheters removed routinely on the 5 th post-operative day. * Stats sig between groups		Group 2: 18(10-33) p value: NS	
			Residual volume, mL, mean (range)	At 3 months Group 1: 40 (5-163) Group 2: 73(20-211) p value: NS	
			PdetQmax (cm H₂O)	At 3 months Group 1: 39 (21-63) Group 2: 51 (37-85) p value: <0.05	
			Urodynamic obstruction, at 3 months, Schafer grade Abrams- Griffiths nomogram, % still obstructed, N not provided	Group 1: 1.9 (0-4) Group 2: 1.0 (0-3) 95% CI: NR p value: <0.05 Group 1: 6% Group 2: 21% 95% CI: NR p value: NR	
			Catheter duration, mean (range), days	Group 1: 1.4 (1-8) Group 2: 11.6(3-8) 95% CI: NR p value: <0.0001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Post-op complications (early): Recatheterisation	Group 1: 2/22 (9%) Group 2: 8/22 (36%) Relative risk: 0.25 95% CI: 0.06-1.05 p value: NR	
			Post-op complications (early): Blood transfusion	Group 1: 0/22 Group 2: 0/22 p value: NS	
			Post-op complications (early): Catheter irrigation (for hematuria)	Group 1: 0/22 Group 2: 0/22 p value: NS	
			Post-op complications (early): Peri-operative urinary tract infections	Group 1: 0/22 Group 2: 3/22 (13.6%) Relative risk: 0.14 95% CI: 0.01-2.61 p value: NS	
			Post-op complications: Retrograde ejaculation in sexually active patients (Number sexually active not stated)	Group 1: 0/NR Group 2: 0/NR p value: NS	
			Post-op complication: Reoperation: 3 in VLAP group had to be reoperated because of persistent urinary retention. 1 in the HoLRP group – urethral dilatation for submeatal stricture	Group 1: 1/22 Group 2: 3/22 Relative risk: 0.33 95% CI: 0.04-2.96 p value: NS	

Evidence Table 33: Holmium laser enucleation of the prostate (HoLEP) vs. laser vaporisation

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Elzayat 2009 ⁸⁰ Study design: RCT Evidence level: 1+ Setting: Canada Duration of follow-up: 12 months	<p>Patient group: Between March 2005 and April 2007 men with LUTS secondary to BPH were recruited at McGill University Health centre, Canada.</p> <p>Inclusion criteria: prostate size 60cc or smaller, IPSS of 9 or greater, Qmax < 15ml/s.</p> <p>Exclusion criteria: previously diagnosed with prostate cancer, urethral stricture or neurogenic bladder or previous prostate surgery.</p> <p>All patients N: 109</p> <p>Group 1 N: 57 Mean age ± SD: 72.7±10.3 Drop outs: 13</p> <p>Group 2 N: 52 Mean age ± SD: 71.6 ±10.3 Drop outs: 10</p>	<p>Group 1: holmium laser ablation of the prostate (HoLAP) Performed using an 80 to 100 watt holmium laser generator and 550um side firing laser fibre. Laser setting ranged from 2.0J and 50Hz to 3.2J and 30Hz.</p> <p>Group 2: photoselective vaporisation (PVP) Performed using the green light laser system with 80 Watt output and side firing laser fibre with a 600 um core diameter.</p> <p>Both procedures: Patient under general or regional anaesthesia and normal saline was used as an irrigant. Continuous flow 26Fr resectoscope with laser fibre stabilising bridge at the tip of the inner sheath was used. After each laser procedure a standard 22Fr 2-way catheter was inserted.</p> <p>Catheter routinely removed the next morning after surgery and when patient is able to void adequately he is discharged from the</p>	<p>Mean (SD) symptom score (IPSS)</p> <p>Mean (SD) quality of life from IPSS score</p> <p>Mean (SD) Qmax</p>	<p>Baseline: Group 1 (n=57): 20 (6.8) Group 2 (n=52): 18.4 (6.6)</p> <p>1 month: Group 1(n=54): 8.7 (6.5) Group 2(n=48): 8.9 (5.4)</p> <p>3 months: Group 1(n=44): 8.4 (7) Group 2(n=39):5.8 (4.4)</p> <p>6 months: Group 1(n=40):7.8 (5.7) Group 2(n=39):7.7 (6.9)</p> <p>12 months: Group 1(n=44):6.2 (3.9) Group 2(n=42):8.2 (6.2); p=0.22</p> <p>Baseline: Group 1 (n=57): 3.8 (1.5) Group 2 (n=52): 3.6 (1.4)</p> <p>1 month: Group 1(n=54): 1.8 (1.6) Group 2(n=48): 1.9 (1.6)</p> <p>3 months: Group 1(n=44): 1.5 (1.4) Group 2(n=39): 1.2(1.1)</p> <p>6 months: Group 1(n=40):1.6 (1.3) Group 2(n=39):1.2 (1.1)</p> <p>12 months: Group 1(n=44):1.6 (1.2) Group 2(n=42):1.5 (1.4); p=0.81</p> <p>Baseline: Group 1 (n=57): 6.7 (3.9) Group 2 (n=52): 6.4 (3.9)</p> <p>1 month: Group 1(n=54): 17.1 (7.5) Group 2(n=48): 18.8 (8.5)</p> <p>3 months: Group 1(n=44): 18.4 (6.4) Group 2(n=39): 18.7 (9.9)</p>	<p>Funding: Author Elhilali has financial interest and/or other relationship with Lumenis and Laserscope.</p> <p>Limitations: Reasons for drop out not reported. Allocation concealment not reported.</p> <p>Additional outcomes: IIEF erectile function domain score was reported. Level of haemoglobin and serum Na. PSA was reported.</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
		hospital.		6 months: Group 1 (n=40): 17.4 (5.9) Group 2 (n=39): 19.4 (8.5) 12 months: Group 1 (n=44): 17.2 (8.4) Group 2 (n=42): 18.4 (8.4); p=0.66	
			Mean (SD) PVR	Baseline: Group 1 (n=57): 205 (197) Group 2 (n=52): 215 (208) 1 month: Group 1 (n=54): 47.4 (93) Group 2 (n=48): 56.2 (79.5) 3 months: Group 1 (n=44): 57.2 (104) Group 2 (n=39): 73.7 (96) 6 months: Group 1 (n=40): 55 (100) Group 2 (n=39): 67.5 (90) 12 months: Group 1 (n=44): 68.9 (90) Group 2 (n=42): 66 (101); p=0.92	
			Mean (SD) laser time, minutes	Group 1: 69.8 (31.6) Group 2: 55.5 (21) P=0.008	
			Mean (SD) catheterisation, days	Group 1: 2.1 (2.7) Group 2: 1.65 (1.6) P=0.29	
			Mean (SD) hospital stay, days	Group 1: 0.87 (0.3) Group 2: 0.96 (0.27) P=0.15	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Number (%) complications	Intraoperative bleeding Group 1: 0 Group 2: 3 (5.7) Blood transfusions Group 1: 0 Group 2: 0 Hematuria Group 1: 1 (1.7) Group 2: 1 (1.9) Irritative symptoms Group 1: 13 (22.8) Group 2: 10 (19.2) Re-catheterisation Group 1: 7 (12.2) Group 2: 6 (11.5) Clot retention Group 1: 1 (1.7) Group 2: 1 (1.9) Stress incontinence Group 1: 1 (1.7) Group 2: 2 (3.8) Urge incontinence Group 1: 4 (7) Group 2: 3 (5.7) Urinary tract infection Group 1: 3 (5.3) Group 2: 2 (3.8)	
			Number (%) late postoperative complications	Urethral stricture Group 1: 1 (1.7) Group 2: 3 (5.7) BNC Group 1: 2 (3.5) Group 2: 4 (7.7) Reoperation Group 1: 2 (3.5) Group 2: 1 (1.9)	
			Mean prostate volume (cc) at 6 months	Group 1: 19.8 Group 2: 24.4; p=NS	

Evidence Table 34: Transurethral microwave thermotherapy (TUMT) vs. no treatment

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Abbou et al., 1995³</p> <p>Study design: Randomised controlled trial</p> <p>Setting: France</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Men recruited from 7 urological departments with symptomatic prostatism that had voiding disorders for at least 3 months.</p> <p>Inclusion criteria: Men >50 years, peak flow rate <15mL/s for a voided volume of ≥150mL; and residual urine <300mL/s. No suspicion of prostate cancer, prostate weight between 30 and 80g; PSA level <10ng/mL for a prostatic weight <60g or a PSA level <15ng/mL for a prostatic weight ≥60g; serum creatinine level <160mol/L; no infection.</p> <p>Exclusion criteria: undergone previous surgery on the prostate or bladder; mental incapacity; any chronic disease potentially hindering follow-up; diabetes; participation in any clinical protocol within at least 3 months; any other urological disease; any medical treatment of voiding disorders within 15 days of inclusion; taken diuretics in the previous 3 months; anticoagulant therapy; allergy to lidocaine or colorectal disease.</p> <p>All patients N: 200 (includes transrectal arms)</p> <p>Group 1 N: 66 Mean (±SD) Age: 65 (8) Mean (±SD) prostate weight: 45g (15) Dropouts: 17% (complementary medical or surgical treatment for worsening obstructive symptoms; one lost to follow-up and 1 withdrew during treatment)</p> <p>Group 2 N: 31 Mean (±SD) Age: 66 (7) Mean (±SD) prostate weight: 44g (11) Dropouts: 38% (complementary medical or surgical treatment for worsening obstructive</p>	<p>Group 1: Transurethral hyperthermia (TUMT) Three devices used for transurethral treatment (Thermex II, Technorex, Israel; Prostatecare, Brucker Spectrospin, France; BSD-50, BSD medical Corp, USA).</p> <p>Prostate temperature was monitored by an integrated microwave generator and controlled each device through a fibre-optic temperature monitor.</p> <p>One session given that lasted between 1-3 hours depending on the device used. Deliver a temperature compatible with hyperthermia treatment (45°C).</p> <p>Group 2: SHAM Single session with the temperature maintained at 37°C.</p>	<p>Number (%) of complications during treatment</p> <p>Number (%) of early post-treatment complications</p> <p>% Objective response rates (PFR)*</p> <p>% Subjective response (Madsen score)*</p>	<p>Urethral bleeding: Group 1: 2 (3) Group 2: 0</p> <p>Urethral pain Group 1: 1 (1.5) Group 2: 0</p> <p>Acute retention: Group 1: 1 (1.5) Group 2: 0</p> <p>Urethral bleeding: Group 1: 18 (27) Group 2: 9 (29)</p> <p>Cystitis Group 1: 12 (18) Group 2: 6 (19)</p> <p>Acute retention: Group 1: 0 Group 2: 0</p> <p>Urinary tract infection: Group 1: 0 Group 2: 1 (3)</p> <p>Prostatitis Group 1: 1 (1.5) Group 2: 1 (3)</p> <p>Other: Group 1: 4 (6) Group 2: 0</p> <p>Group 1 (n=66) : 14 Group 2 (n=29): 17</p> <p>Group 1 (n=66): 50 Group 2 (n=29): 17 P<0.05</p>	<p>Funding: Grant from Comité d'Évaluation et de Diffusion des Innovations Technologiques (CEDIT). Assistance Publique – Hôpitaux de Paris. Devices were lent by the following companies: Biodan, Brucker, BSD, Direc and Tecnomatrix.</p> <p>Limitations: Unclear if allocation concealment used. All withdrawals included in the analysis as non-responders, except for two patients who excluded for reasons unrelated to treatment.</p> <p>Additional outcomes: Study randomised patients to transrectal hyperthermia and transrectal sham arm but results not reported.</p> <p>Notes: * responder defined as patients showing excellent, good or moderate responses according to each of the criteria analysed separately (Madsen decrease >30%; a PFR>10mL/s with a PFR increase>30%) Non responders were patients who withdrew during treatment (because of complications complementary treatment or refusal to continue) and</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	symptoms; one lost to follow-up)				patients who had a Madsen score decrease <30%, PFR<10mL/s or a PFR>10mL/s but with an increase <30%.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Albala et al., 2002¹²</p> <p>Study design: Randomised controlled study</p> <p>Setting: US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Male patients between 50-80 years old with a diagnosis of symptomatic BPH to a sufficient degree that treatment was warranted.</p> <p>Inclusion criteria: AUA symptom index > 13 and a bother score >11. Peak flow rates were <12mL/s and the post voiding residual volume was <125mL. Prostate volume between 30-100cc without a significant intravesical middle lobe.</p> <p>All patients N: 200</p> <p>Group 1 N: 125 Mean (\pmSD) Age: 65.2 (7.3) Mean (\pmSD) volume: 50.5 (18.6) cc Dropouts: NR Number reporting AUA scores indicates that was 6 drop outs at 12 months. Complications reported for 121 out of 125 randomised patients.</p> <p>Group 2 N: 65 Mean (\pmSD) Age: 64.6 (7.1) Mean (\pmSD) volume: 47.1 (17.9) cc Dropouts: NR Complications reported for 62 of 65 patients.</p>	<p>Performed in urology offices or clinics.</p> <p>Group 1: TUMT TherMatrix TMx-2000 that directly heats the transition zone to greater than 50 degrees C. 60-90W. Toradol, narcotic analgesic and lorazepam were given orally 45 minutes before treatment. Prior to catheter insertion lidocaine jelly injected into the urethra and allowed to remain in place for 15 minutes. Treatment temperature delivered to peak tissue temperature of 50 to 55°C. After temperature had increased to 50 degrees the treatment was continued for 40 minutes under computer control. Foley catheter inserted into bladder following treatment and left in place from 2 to 4 days.</p> <p>Group 2: SHAM Placement of the microwave catheter for the treatment period without energy delivery and received the same post treatment care as the active treatment patients.</p>	AUA symptom index (SI)	<p>Baseline: Group1 (n=125): 22.5 Group 2 (n=65): 22.8</p> <p>3 months: Group1 (n=124): 12.4 Group 2 (n=NR): 17</p> <p>6 months: Group1 (n=115): 12.1 Group 2 : NR</p> <p>12 months: Group1 (n=119): 11.9 Group 2: NR</p>	<p>Funding: NR</p> <p>Limitations: Symptom scores only reported fro TUMT arm for 6 and 12 months.</p> <p>Additional outcomes: Bother and quality of life scores reported but only for the treatment arm.</p> <p>Notes: Patients were unblinded at 3 months and sham treated patients offered options of having active treatment. Results for treatment arm only includes patients randomised to active treatment and not those that crossed over at 3 months (intention to treat analysis used).</p>
			AUASI Change (12 months)	Group1: -10.6 (-47.1%) Group 2: NR	
			PFR change , mL/sec (12months)	Group1: +5.0 (58.1%) Group 2: NR	
			Number of complications	<p>Recatheterisation Group 1: 20/121 (16.8%) Group 2: 0/62 (0%)</p> <p>Dysuria Group 1: 8/121 (6.6%) Group 2: 3/62 (4.8%)</p> <p>Urgency Group 1: 0/121 (0%) Group 2: 0/62 (0%)</p> <p>Gross haematuria Group 1: 11/121 (9.1%) Group 2: 0/62 (0%)</p> <p>Bladder spasm Group 1: 5/121 (4.1%) Group 2: 0/62 (0%)</p> <p>Urethral stricture Group 1: 0/121 (0%) Group 2: 0/62 (0%)</p> <p>Ejaculatory dysfunction pain Group 1: 0/121 (0%) Group 2: 0/62 (0%)</p> <p>Rectal damage fistula Group 1: 0/121 (0%) Group 2: 0/62 (0%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bdesha et al., 1994²⁶</p> <p>Study design: RCT</p> <p>Setting: UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: patients with significant symptoms of prostatism and unequivocally benign glands recruited.</p> <p>Inclusion criteria: symptoms of prostatism at least 6 months in duration, symptom score >10, peak flow rate of <15ml/s and/or residual urine volume of greater than 50ml.</p> <p>Exclusion criteria: upper tract dilatation, impaired renal function, acute urinary retention, residual urine volume >200ml, prostatic malignancy, significant middle lobe hypertrophy, large gland, coexisting urinary tract pathological condition or previous prostatic surgery.</p> <p>All patients N: 42</p> <p>Group 1 N: 22 Mean Age: 63.7 years Drop outs: 0</p> <p>Group 2 N: 18 Mean Age: 62.6 years Drop outs: 2 lost to follow-up</p>	<p>Dedicated day care unit. Anaesthetised with topical lidocaine gel and a catheter passed to empty the bladder. Balloon inflated and the catheter pulled back to position the microwave antenna accurately within the prostatic urethra. Rectal temperature monitoring probe was placed the microwave catheter was connected to the microwave device. LEO Microthermer used and delivers a maximum power output of 20W at 915MHz and automatic power cut-off when rectal temperature increases to greater than 42.5°C. Heated pad placed across lower abdomen of all patients to minimise speculation of which treatment arm patients were in.</p> <p>Group 1: TUMT Single active 90 minute treatment</p> <p>Group 2: SHAM Sham treatment for the same time when no power was delivered.</p>	<p>Mean [SD] (95% CI) AUA symptom scores at 3 months</p>	<p>Baseline: Group 1: 19.2 (16.3-22.1) Group 2: 18.8 (16.0-21.7) 3 months: Group 1: 7.1 [5.00] (5.0-9.2) Group 2: 16.2 [7.35] (12.8-19.6)</p>	<p>Funding: NR</p> <p>Limitations: Randomisation method unclear.</p> <p>Additional outcomes: Reported results of sham patients that went onto have active treatment. Scores for force of stream, hesitancy, intermittent voiding and incomplete voiding.</p> <p>Notes: SD reported from HTA report.</p> <p>Patients in the sham arm that showed no improvement after 3 months were offered the active treatment. One patient had sham treatment for 3 months and then retreated with active treatment and subsequently had urinary retention followed by reoperation of transurethral prostatectomy.</p>
			<p>Mean (95% CI) peak flow rate (ml/s)</p>	<p>Baseline: Group 1: 12.3 (10.7-13.9) Group 2: 10.8 (9.2-12.4) 3 months: Group 1: 14.6 [5.98] (12.1-17.1) Group 2: 9.8 [2.81] (8.5-11.1)</p>	
			<p>Mean (95% CI) Residual volume, ml</p>	<p>Baseline: Group 1: 104 (85-125) Group 2: 80 (57-103) 3 months: Group 1: 52 (34-70) Group 2: 94 (71-117)</p>	
			<p>Mean (95% CI) number of daytime voids (frequency)</p>	<p>Baseline: Group 1: 9.4 (7.3-11.4) Group 2: 7.4 (5.4-9.4) 3 months: Group 1: 5.5 (4.4-6.5) Group 2: 7.4 (5.9-8.9)</p>	
			<p>Mean (95% CI) number of voids (nocturia)</p>	<p>Baseline: Group 1: 3.5 (2.5-4.4) Group 2: 3.5 (2.5-4.6) 3 months: Group 1: 1.6 (0.9-2.3) Group 2: 3.3 (2.9-3.7)</p>	
<p>Mean (95% CI) urgency</p>	<p>Baseline: Group 1: 3.5 (2.8-4.2) Group 2: 2.8 (1.6-3.1) 3 months:</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 1.1 (0.5-1.8) Group 2: 1.6 (0.9-2.5)	
			Retrograde ejaculation (new cases) * number with antegrade ejaculation preoperatively not reported	Group 1: 0/NR Group 2: 0/NR	
			% correctly guesses which treatment arm they were in	Group 1: 86% Group 2: 50%	
			Successful outcomes (defined as a decrease in symptom scores with greater than a 50% decrease) at 3 months	Group 1: 18/22 Group 2: 2/20	
			Reoperation (at 3 months patients in sham arm offered active treatment)	Group 1: 0/22 Group 2: 16/20	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments	
Blute et al., 1996 ³¹ Study design: RCT Setting: US Evidence level: 1+ Duration of follow-up: 12 months	Patient group: patients with symptomatic BPH. Inclusion criteria: peak urine flow rate <10ml/s; residual volume 100-200ml; Madsen score >8; prostate length 35-50 mm from TRUS. Exclusion criteria: Prostate cancer; transurethral or rectal surgery; urinary retention; any medications that affect prostate symptoms; antiandrogen therapy; upper UT pathology shown by ultrasound; metallic implants; symptoms suggesting neuropathological bladder; serum creatinine >2mg/dl; bladder stones; uncontrolled dysrhythmias or cardiac pacemaker; asymmetric median lobe enlargement; patients at high risk from prostatic disease. All patients N: 115 Drop outs: NR Group 1 N: 78 Mean (±SD) Age: 66.9 (7.8) Group 2 N: 37 Mean (±SD) Age: 66.9 (7.1)	Outpatient procedure. Antibodies and nonsteroidal anti-inflammatory agent given before therapy. Group 1: TUMT – Prostatron (Prostasoft) Rectal thermometry probe inserted and treatment catheter with Foley balloon located by transabdominal ultrasound and TURS; anaesthesia: 89% had only local anaesthetic (lidocaine), 11% had midazolam-fentanyl intravenously; blood pressure, pulse and temperature monitored every 15 minutes during treatment; observation for 2 hours. Group 2: SHAM No sedation; urethral coolant circulated; NSAIDs given before therapy. Treatment ran for 60 minutes.	Mean (SD) AUA scores Baseline Group 1 (n=64): 19.7 (7.2) Group 2 (n=31): 21.9 (6.3) 6 weeks: Group 1 (n=59): 12.8 (6.6) Group 2 (n=28): 17.1 (6.9) 3 months: Group 1 (n=64): 11.3 (6.3) Group 2 (n=31): 16.3 (7.6)	Baseline Group 1 (n=74): 7.2 (1.6) Group 2 (n=34): 7.4 (1.6) 6 weeks: Group 1 (n=72): 10.7 (4.1) Group 2 (n=32): 8.5 (3.7) 3 months: Group 1 (n=74): 11.5 (4.0) Group 2 (n=34): 9.4 (3.7)	Funding: NR Limitations: Drop outs and reasons not reported. Additional outcomes: PSA levels at baseline and at 6 months. Madsen symptom scores reported. Notes: Sham group offered active treatment at 3 months. Reported that no sexual dysfunction following procedure but no indication of patients that previously had dysfunction.	
			Mean (SD) peak flow rates (mL/s)			Baseline Group 1 (n=71): 140.9 (35.9) Group 2 (n=33): 142.1 (35.5) 3 months: Group 1 (n=71): 145.5 (126.1) Group 2 (n=33): 147.2 (107.7)
			Mean (SD) residual urine by catheter, mL			Any positive change Group 1: 60/75 (80%) Group 2: 11/37 (29.7%) No change Group 1: 12/75 (16.0%) Group 2: 23/37 (62.2%) Uncertain Group 1: 3/75 (4.0%) Group 2: 3/37 (8.1%)
			Number (%) of improved symptoms assessed by the patient at 3 months			Any positive change Group 1: 63/75 (84%) Group 2: 13/37 (35.1%) No change Group 1: 8/75 (10.7%)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 23/37 (62.2%) Uncertain Group 1: 4/75 (5.3%) Group 2: 1/37 (2.7%)	
			Number (%) complications at 3 months	Haematuria: Group 1: 54/78 (69.2%) Group 2: 19/37 (51.3%) Urethral bleeding Group 1: 16/78 (20.5%) Group 2: 5/37 (13.5%) Urethral discharge Group 1: 2/78 (2.6%) Group 2: 0 Urinary retention Group 1: 20/78 (25.6%) Group 2: 0 Other urinary tract Group 1: 11/78 (14.1%) Group 2: 4/37 (10.8%) Reproductive (including genital dermatology) Group 1: 8/78 (10.3%) Group 2: 0 Rectal (including proctoscopy findings) Group 1: 4/78 (5.1%) Group 2: 4/37 (10.8%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Brehmer et al, 1999³⁴</p> <p>Study design: RCT</p> <p>Setting: Sweden</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Men with LUTS dominated by hesitancy, slow urination and an enlarged prostate.</p> <p>Inclusion criteria: maximum flow rate of <12mL/s</p> <p>Exclusion criteria: indwelling catheter, median prostatic lobe, a prostate gland estimated as >50g, suspected prostatic malignancy, neurological disease and previous surgery for prostatic disease.</p> <p>All patients N: 44 Mean age (range): 70.4 (53-83) Drop outs: 2</p> <p>Group 1 N: 14 Dropouts: 1 (withdrew as had repeated transient ischaemic attacks and developed early dementia)</p> <p>Group 2 N: 16 Dropouts: 0</p> <p>Group 3 N: 14 Dropouts: 1 (prostatic carcinoma)</p>	<p>An ECP system (Comair, Sweden) equipped with a 22F catheter with a microwave antenna (915MHz), a fibre-optic system for measuring the temperature in the urethra and, by a rectal probe in the rectum. The two-way urethral catheter has a circulation cooling system that reduces the heat delivered to the urethral wall. Maximum heating is achieved within 30s and the temperature limit is 46 degrees in the urethral and 43 in the rectum. If unable to void a urethral catheter inserted and left in place for 3 days. All patients received antibiotics for 5 days.</p> <p>Group 1: TUMT for 30 minutes</p> <p>Group 2: TUMT for 60 minutes</p> <p>Group 3: SHAM Only water at 20° was circulated in the treatment catheter and a computer monitor, visible to the patient, showed a simulated heat treatment curve, similar to that produced during TUMT.</p>	Qmax, mL/s	<p>Baseline: Group 1: 8.7 Group 2: 7.0 Group 3: 7.9</p> <p>4 months: Group 1: 12.3 Group 2: 9.9 Group 3: 8.3</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, allocation concealment unclear. Baseline urodynamic scores similar between groups but A scores were significantly higher in the 30 minute TUMT group (Group 1). Complications reported as whole rather than by group.</p> <p>Additional outcomes: Frequency and timed void before and after treatment. % improved in different variables reported (but actual figures reported in full).</p> <p>Notes: ICS score defined as a Questionnaire with 32 questions (A questions about symptoms and B question about the bother related to the symptom. Maximum A and B scores are 124 and 92 respectively. High score indicates worse symptoms.</p>
			Treatment failure	<p>Group 1 & 2: 5/30 (17%) Group 3: 7/14</p>	
			Reoperation	<p>Group 1: 0/14 Group 2: 3/16 Group 3: 7/14</p>	
			ICS A score (with % decrease) * See notes for definition of score	<p>Before Group 1: 58 Group 2: 49 Group 3: 46</p> <p>4 months: Group 1: 44 (25) Group 2: 41 (16) Group 3: 44 (4)</p>	
			ICS B score (with % decrease) * See notes for definition of score	<p>Before Group 1: 40 Group 2: 36 Group 3: 36</p> <p>4 months: Group 1: 30 (34) Group 2: 30 (17) Group 3: 31 (14)</p>	
			% improvement using quality of life score (from ICS questionnaire last question - with 7 points indicating worst situation possible)	<p>Group 1: 25% Group 2: 4% Group 3: 0%</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Dewiltd et al., 1996⁶⁷</p> <p>Links with Delarosette 1994⁶⁴ and Francisca 1997⁹⁵</p> <p>Study design: Randomised controlled trial</p> <p>Setting: 2 centres – London and Nijmegen, Netherlands</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: From June 1991 to December 1992 patients recruited.</p> <p>Inclusion criteria: >45 years; complaining of symptoms of bladder outlet obstruction for >3 months, have a Madsen symptom score of >8 and urinary free-flow rate estimates of <15 mL/s during two voids of >150mL.</p> <p>Exclusion criteria: prostate cancer, prostatitis, urethral stricture, intravesical pathology, neurogenic bladder dysfunction UTI, isolated enlargement of the middle lobe, a residual urine volume of ≥300mL, use of drugs influencing bladder or prostate function, previous transurethral resection of the prostate or transurethral incision, a metallic pelvic implant, disorders of blood flow or coagulation, diabetes, mental incapacity or inability to give informed consent.</p> <p>All patients N: 93</p> <p>Group 1 N: 47</p> <p>Mean (±SD) Age: 66.3 (8.1)</p> <p>Dropouts: 2 (had TURP)</p> <p>At 12 months: 14 (TURP=4, Lost to follow-up=5, second TUMT=4, death (not related to treatment)=1)</p> <p>Group 2 N: 46</p> <p>Mean (±SD) Age: 63.9 (6.0)</p> <p>Drop outs: 3 (lost to follow up=2, technical failure=1)</p> <p>At 12 months: 33 (5 lost to follow up, technical failure=1 and 27 had TUMT at 3 months)</p>	<p>Group 1: TUMT Transurethral thermotherapy Prostatron, Prostatsoft 2.0</p> <p>Group 2: SHAM Procedure simulated but without applying microwave energy. Real time treatment profile displayed on the computer screen as done in active treatment and explained to the patient. Sequence of temperature, calibration and checks were identical in both groups.</p>	<p>Mean (95% CI) of Madsen symptom score</p> <p>Baseline Group 1: 13.7 (12.7-14.7) Group 2: 12.9 (11.9-13.9)</p> <p>3 months Group 1: 4.7 (3.6-5.9) Group 2: 10.4 (8.9-11.8)</p> <p>12 months Group 1: 4.2 (3.0-5.3) Group 2: 8.2 (5.5-11.0)</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation and use of allocation concealment are unclear. Some significant baseline differences between the two centre. London centre had significantly older patients, more obstructive symptoms and greater residual volume.</p> <p>Additional outcomes: Reports results for SHAM group when they have had an active treatment as 3 months following no improvement. Voided fraction reported.</p> <p>Notes: When patients had no improvement after 3 months, whether he had received sham or active treatment, a second genuine TUMT was performed on request.</p>	
			<p>Mean (95% CI) of peak flow rate, mL./s</p> <p>Baseline Group 1: 9.2 (8.4-9.9) Group 2: 9.6 (8.8-10.4)</p> <p>3 months Group 1: 13.4 [6.16] (11.7-15.3) Group 2: 9.7 [3.30] (11.7-15.3)</p> <p>12 months Group 1: 13.4 [5.13] (11.6-15.1) Group 2: 10.5 [4.79] (7.9-13.1)</p>		
			<p>Mean (95% CI) of post void residual urine, mL</p> <p>Baseline Group 1: 93.9 (71.8-116.0) Group 2: 84.7 (64-105.1)</p> <p>3 months Group 1: 34.2 (19.4-46.8) Group 2: 104.1 (74.7-133.4)</p> <p>12 months Group 1: 49.72 (33-66.3) Group 2: 56.3 (16.9-95.7)</p>		
			<p>Mortality</p> <p>Group 1: 1/47 Group 2: 0/46</p>		
			<p>Retention</p> <p>Group 1: 10/47 Group 2: 1/46</p>		
			<p>Reoperation</p> <p>Group 1: 8/47 Group 2: 27/46</p>		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Larson et al., 1998¹⁵⁹</p> <p>Study design: RCT</p> <p>Setting: 5 centres in US.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months.</p>	<p>Patient group: symptomatic BPH patients enrolled between September 1994 and June 1996</p> <p>Inclusion criteria: Qmax ≤12mL/s with voided volume ≥12mL/s with voided volume ≥25mL, AUA symptom score ≥9, 3-5cm preprostatic urethral length as determined by cystoscopy or TURS, No disproportionately enlarged or prominent prostatic median lobe on cystoscopy, life expectancy ≥1 year.</p> <p>Exclusion criteria: UTI within 1 week of study enrolment, gross hematuria, acute urinary retention, prostate weight >100g, concomitant medications, use of alpha antagonists or antiandrogens, coexisting disease that could mimic obstructive bladder neck syndrome, coexisting illness or specific obstructive symptoms caused by neurogenic bladder; bladder stones, renal failure, cardiac failure, prostate cancer, urethral stricture, severe bladder neck contracture, bladder cancer, urinary sphincter abnormalities, prostatitis or hepatic failure. Continuous or intermittent urinary catheterisation within 2 weeks or study, previous prostate surgery or non medical treatment for BPH, penile implant or artificial</p>	<p>Group 1: TUMT Urologix Targis system used. Microwave energy for one hour. Outpatient setting without anaesthesiologist or anaesthetist. The catheter provides urethral cooling via circumferential cooking compartments and monitors temperatures. The thermoablation system automatically interrupts microwave power if urethral temperatures reach 44.5°C or higher or rectal temperatures over 42.5. Topical lidocaine anaesthesia used for catheterisation. Microwave power applied in increments to achieve target temperature of 40 degrees. Treatment administered for one hour. Given 3 day prescription of prophylactic oral antibiotics and catheterisation for 36 to 60 hours.</p> <p>Group 2: SHAM Underwent procedures identical to those in active arm but the microwave energy not applied. Coolant temperature was increased in increments from 8 to 20° over the same time period as microwave power was increased in active group. Given 3 day prescription of prophylactic oral antibiotics and catheterisation for 36 to 60</p>	<p>Mean (SD) / [range] symptom score (AUA)</p>	<p>Baseline: Group 1 (n=124): 20.8 [19.8-21.9] Group 2 (n=42): 21.3 [19.3-23.3]</p> <p>3 months: Group 1 (n=123): 9.60 (5.94) Group 2 (n=40): 14.50 (6.77)</p> <p>6 months: Group 1 (n=120): 10.50 (7.26) Group 2 (n=35): 14.30 (6.34)</p>	<p>Funding: Supported by a grant from Urologix, Inc.</p> <p>Limitations: Method of randomisation and whether allocation concealment used were not reported.</p> <p>One enrollee who had been assigned to the sham group was inadvertently made aware of his group assignment and consequently this patient's schedule study treatment was cancelled. Prostate volume 17% greater in sham group at baseline.</p> <p>Additional outcomes: PSA levels before and after treatment. 6 week results for symptom score and Qmax. Prostate volume reported but only for active group.</p> <p>Notes: SD for Qmax and symptom scores was</p>
			<p>Mean (SD) / [range] Qmax</p>	<p>Baseline: Group 1 (n=106): 7.8 [7.4-8.2] Group 2 (n=39): 7.8 [7.00-8.6]</p> <p>3 months: Group 1 (n=102): 11.70 (5.41) Group 2 (n=37): 9.20 (3.72)</p> <p>6 months: Group 1 (n=101): 11.80 (5.89) Group 2 (n=31): 9.80 (4.00)</p>	
			<p>Mean [range] post void residual, mL</p>	<p>Baseline: Group 1 (n=105): 99.1 [82.0-116.1] Group 2 (n=39): 103.6 [79.4-127.8]</p> <p>3 months: Group 1 (n=103): 68.4 [52.9-83.8] Group 2 (n=37): 93.0 [57.6-128.4]</p> <p>6 months: Group 1 (n=101): 84.5 [67.8-101.2] Group 2 (n=31): 84.4 [58.3-110.6]</p>	
			<p>Quality of life score (SD) evaluated by patient responses to the question of how they would feel if their current urinary symptoms were to continue indefinitely</p>	<p>Baseline: Group 1 (n=120): 4.2 (95% CI: 4.0-4.4) Group 2 (n=35): 4.0 (95% CI: 3.6-4.3)</p> <p>6 months: Group 1 (n=120): 2.20 (1.40) Group 2 (n=35): 2.90 (1.20)</p>	
<p>Complications</p>	<p>Blood transfusions Group 1: 0/125 Group 2: 0/44</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>urinary sphincter, previous pelvic or rectal surgery, metallic implants in the pelvic area, cardiac pacemaker, desire for future offspring, likely non compliance.</p> <p>All patients N: 169 Mean age: 45-85 years Drop outs:</p> <p>Group 1 N: 125 Mean (range) Age: 66.0 (64.7-67.4) Dropouts: 5 (prostate cancer=2, need for further treatment for BPH=2, died of unrelated causes=1)</p> <p>Group 2 N: 44 Mean (range) Age: 65.9 (63.4-68.3) Dropouts: 9 (study procedure cancelled=1, missed prostatitis at screening=1, need for further treatment for BPH=7)</p>	hours.	<p>Urinary retention Group 1: 10/125 Group 2: 1/44</p> <p>Urinary tract infection Group 1: 11/125 Group 2: 2/44</p> <p>Stricture Group 1: 3/125 Group 2: 0/44</p> <p>Urinary incontinence Group 1: 5/125 Group 2: 0/44</p> <p>Reoperation Group 1: 2/125 Group 2: 27/44</p> <p>Ejaculatory disorders: Group 1: 5/125 Group 2: 0/44</p> <p>Mortality: Group 1: 1/125 Group 2: 0/44</p> <p>Number (%) that correctly identified intervention received</p> <p>Number of patients experiencing discomfort during the procedure</p>	<p>Group 1: 100/112 (90%) Group 2: 21/37 (50%)</p> <p>None or mild: Group 1: 65/125 (52.0%) Group 2: 37/42 (88.1%)</p> <p>Moderate: Group 1: 57/125 (45.6%) Group 2: 5/42 (11.9%)</p> <p>Severe Group 1: 3/125 (2.4%) Group 2: 0/42 (0%)</p>	<p>calculated in HTA report.</p> <p>After 6 months follow up continued on unblinded basis, with follow up to one year by mail in questionnaire only. After 6 months evaluation sham group patients could elect to undergo microwave or other treatment for BPH.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nawrocki et al., 1997²¹⁸</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with symptoms associated with bladder outlet obstruction and BPH.</p> <p>Inclusion criteria: symptoms of lower urinary tract dysfunction thought to be due to benign enlargement of the prostate meriting surgical treatment, Qmax<15mL/s and voided volume 150mL or more, Pdet max of 70cmH2O or more.</p> <p>Exclusion criteria: Complications of bladder outlet obstruction (retention, residual urine volume >350mL, renal failure, recurrent urinary tract infection, bladder calculus, bladder diverticulum); suspicion of malignancy, short prostate, presence of a prominent middle lobe projecting asymmetrically into the bladder, presence of a urethral stricture, previous prostate or pelvic surgery or radiotherapy, presence of metal within the lower trunk or upper legs, uncontrolled cardiac dysrhythmias or presence of a cardiac pacemaker, neurological disorders, inability to understand treatment procedure, presence of other treatment which may affect LUT function.</p> <p>All patients N: 120 Median age: 70 (56-80) years Drop outs: NR (only that urodynamic data incomplete in 4 patients).</p> <p>Group 1 N: 38 Group 2 N: 40 Group 3 N: 42</p>	<p>Group 1: TUMT Prostasoft v 2.0. 1 hour treatment with microwaves performed with the patient under local anaesthesia and as an out-patient.</p> <p>Group 2: SHAM Simulated TUMT with identical procedure as active treatment but treatment device emitted no microwaves during the procedure. The machine noise, treatment duration and graphical computer display were all simulated by placebo software on disk. Heat simulated using a heat pad.</p> <p>Group 3: No treatment</p>	<p>Median (range) AUA symptom score:</p>	<p>Baseline: Group 1: 19 (7-31) Group 2: 17.5 (7-28) Group 3: 18 (10-29) 6 months: Group 1: 9.5 (1-27) Group 2: 9.5 (0-30) Group 3: 17 (4-28)</p>	<p>Funding: Research was in part supported by a LORS grant from the South East Thames Regional Research Committee. This work in part contributed to the award of an MS thesis from University of London.</p> <p>Limitations: Allocation concealment use was unclear and drop outs not reported.</p> <p>Additional outcomes: Minimum urethral opening pressure, maximum detrusor pressure, voided volume, detrusor instability, functional bladder capacity.</p> <p>Notes: Active and sham arms included in the meta-analysis.</p> <p>37% judged that they knew which treatment that they had. Of which 59% were correct. Operators judged correctly 68% of time.</p>
			<p>Mean (SD) Qmax, mL/s</p>	<p>Baseline: Group 1: 8.83 (2.32) Group 2: 9.44 (2.78) Group 3: 8.79(2.66) 6 months: Group 1: 9.94 (3.08) Group 2: 9.49 (2.88) Group 3: 8.47 (1.92)</p>	
			<p>Mean (SD) residual urine volume, mL</p>	<p>Baseline: Group 1: 85.7 (56.6) Group 2: 96.5 (56.3) Group 3: 86.0 (62.7) 6 months: Group 1: 85.8 (51.2) Group 2: 106.3 (84.5) Group 3: 82.7 (52.7)</p>	
			<p>Mean (SD) prostate volume, mL</p>	<p>Baseline: Group 1: 41.2 (14.6) Group 2: 46.7 (16.8) Group 3: 46.4 (19.9) 6 months: Group 1: 45.6 (17.6) Group 2: 48.9 (19.7) Group 3: 45.2 (17.9)</p>	
			<p>Urinary retention</p>	<p>Group 1: 4/38 (10.5%) Group 2: 0/40</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ogden et al., 1993²³² (abstract only but data extracted in HTA systematic review)</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: UK</p> <p>Evidence level: Abstract only</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: Recruitment dates from September 1991.</p> <p>Inclusion criteria: peak urine flow rate <15ml/s on two occasions; residual volume ≤350ml. Madsen score>8 for 6 months, prostate urethral length 35-50mm.</p> <p>Exclusion criteria: prostate cancer from DRE; heat to prostate or pelvic surgery/radiotherapy; urinary retention; alpha blockers within 4 weeks; antiandrogens within 1 year; anything affecting prostate of bladder; prostatitis or UTI; renal dysfunction; peripheral arterial disease; diabetic neuropathy; UT disease; bladder disease; mental incapacity; dementia, inability to give informed consent; neurological disorders affecting bladder function; disorders of blood flow or coagulation; history or uncontrolled cardiac arrhythmias or cardiac pacemaker; metallic pelvic implant; prominent isolated median lobe; intravesical pathology; renal impairment due to chronic retention; urethral stricture inhibiting catheterisation.</p> <p>All patients N: 43</p> <p>Group 1 N: 22 Mean (±SD) Age: 68.3 (64.1-72.5)</p> <p>Group 2 N: 21 Mean (±SD) Age: 67.1 (63.7-70.3)</p>	<p>Group 1: TUMT Catheter protocol – inserted for retention for one week.</p> <p>Group 2: SHAM Catheter protocol – inserted for retention for one week.</p>	<p>Mean (95% CI) Madsen score</p>	<p>Group 1: 14.5 (12.9-16.1) Group 2: 14.2 (12.7-15.7)</p>	<p>Funding: Unknown</p> <p>Limitations: HTA appraisal of study reports unclear method of randomisation and no allocation concealment. Patients blinded but assessors were not.</p> <p>Additional outcomes: Voided volume and residual volume reported in the HTA report.</p> <p>Notes: If patient saw no improvement in 3 months after sham or TUMT a second TUMT was performed on request.</p>
			<p>Mean (95% CI) Qmax, ml/s</p>	<p>Baseline: Group 1: 8.5 (7.5-9.5) Group 2: 8.6 (7.6-9.6) 3 months: Group 1: (n=21) 13.0 (5.84) Group 2: (n=19) 9.2 (4.45)</p>	
			<p>Mean (95% CI) Quality of life score</p>	<p>Group 1: 13.4 (10.7-16.1) Group 2: 13.3 (9.2-17.4)</p>	
			<p>Urinary tract infection</p>	<p>Group 1: 5/22 Group 2: 1/21</p>	
			<p>Urinary retention</p>	<p>Group 1: 5/22 Group 2: 0/21</p>	
			<p>Reoperation</p>	<p>Group 1: 1/22 Group 2: 1/21</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Trachtenberg et al., 1998^{29,6}</p> <p>Linked to Tan 2005</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: multicentre, US and Canada</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Men over 55 years</p> <p>Inclusion criteria: AUA >13; peak urinary flow rate <12 ml/s and voided volume >125ml. serum PSA <10ng/ml; prostate volume between 25-100ml; bladder neck to verumontanum distance <30mm.</p> <p>All patients N: 220</p> <p>Group 1 N: 147 Mean (rang)) Age: 66.2 (54.4-82.7) Dropouts: between 2-5</p> <p>Group 2 N: 73 Mean (range) Age: 66.0 (55.1-78.1) Dropouts: 3</p>	<p>Group 1: TUMT Dornier Urowave used which operates at 915MHz. Generator capable of delivering up to 90W of power. Safety threshold set at 50°C in the urethra and 42.5°C in the rectum. Outpatient procedure without general anaesthesia. Peri-treatment antibiotic prophylaxis at the investigators choice. Following treatment a Foley catheter was inserted and left indwelling for 2-5 days.</p> <p>Group 2: SHAM 60minute pre-programmed treatment cycle without the application of power.</p>	<p>Mean (range) AUA symptom score</p> <p>Mean (range) AUA bother score</p> <p>Mean peak flow, ml/s</p> <p>Complications</p>	<p>Baseline: Group 1: 23.6 [5.6] (12-35) Group 2: 23.9 [5.6] (13-35)</p> <p>3 months: Group 1: 11.6 Group 2: 16.4</p> <p>6 months: Group 1: 12.6 Group 2: 17.9</p> <p>Baseline: Group 1: 18.5 (0-28) Group 2: 18.6 (0-28)</p> <p>6 months: Group 1: 8.7 Group 2: 12.6</p> <p>Baseline: Group 1: 7.7 (3.5-11.5) Group 2: 8.1 (4.0-11.9)</p> <p>3 months: Group 1: 11.0 Group 2: 9.7</p> <p>6 months: Group 1: 10.6 Group 2: 9.6</p> <p>Pain Group 1: 80% Group 2: 56%</p> <p>Occurrences ejaculatory dysfunction Group 1: 30/147 Group 2: 1/73</p> <p>Irritative voiding: Group 1: 21/147 Group 2: 4/73</p> <p>haematuria Group 1: 19/147 Group 2: 1/73</p> <p>UTI Group 1: 11/147 Group 2: 2/73</p>	<p>Funding: NR</p> <p>Limitations: Randomisation method unclear and reason for dropouts not reported. Results report one stricture in the active treatment compared to none in the sham arm. Conversely, the conclusion reports no strictures in the study so have excluded this outcome.</p> <p>Additional outcomes: Prostate volume and PSA baseline scores. Quality of life question (0-6) but only reported figures for baseline scores.</p> <p>Notes: At 6 months follow-up patients on sham treatments were offered active treatment.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>Urinary retention: Group 1: 8/147 Group 2: 0/73</p> <p>Scrotal abscess Group 1: 6/147 Group 2: 1/73</p> <p>Rectal disorder: Group 1: 8/147 Group 2: 2/73</p> <p>Pelvic pain: Group 1: 5/147 Group 2: 1/73</p> <p>Penile disorder: Group 1: 5/147 Group 2: 0/73</p> <p>Urinary incontinence Group 1: 0/147 Group 2: 0/73</p> <p>Bladder spasm: Group 1: 1/147 Group 2: 1/73</p> <p>Split urinary stream: Group 1: 0/147 Group 2: 1/73</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments	
Zerbib et al, 1994 ³³¹ Study design: Randomised controlled study Setting: France Evidence level: 1+ Duration of follow-up: 3 months	Patient group: symptomatic BPH patients. Inclusion criteria: candidates for prostatectomy. All had failed one conservative treatment (e.g. alpha-blockers) and the symptoms were of sufficient severity such that prostatectomy was indicated. Exclusion criteria: anterior rectal wall thickness >10mm or <2mm; anterior to posterior thickness of prostate >55mm. All patients N: 68 Mean age: 69.5±10.44 (53-88) Drop outs: NR Group 1 N: 38 Group 2 N: 30	Group 1: TUMT Prostatic hyperthermia treatments were performed using Prostathermer. Intraprostatic temperature maintained at 43±0.5°C. 1 hour session per week for 5 consecutive weeks. Outpatient without anaesthesia. Group 2: SHAM Intraprostatic temperature maintained at 37±0.5°C by radiofrequency power. One hour session per week for 5 consecutive weeks.	Mean (SD) peak flow, ml/s Baseline Group 1: 7.6 (3.8) Group 2: 10.6 (5.8) 3 Months: Group 1: 9.60 (5.80) Group 2: 10.8 (5.4)	Baseline Group 1: 7.6 (3.8) Group 2: 10.6 (5.8) 3 Months: Group 1: 9.60 (5.80) Group 2: 10.8 (5.4)	Funding: NR. Limitations: Randomisation method and allocation concealment unclear. Baseline peak flow significantly different between arms. Inclusion and exclusion criteria not defined. No complications reported. Additional outcomes: Siroky S.D. and adjusted flow scores. Response rate (objective criteria) reported. Notes: 3 month result for peak flow for TUMT group not reported in study – result obtained from HTA report.	
			Mean (SD) voided volume, ml Baseline Group 1: 151 (92.0) Group 2: 145 (86.3) 3 Months: Group 1: 154 (90) Group 2: 166 (91.3)			Baseline Group 1: 151 (92.0) Group 2: 145 (86.3) 3 Months: Group 1: 154 (90) Group 2: 166 (91.3)
			Mean (SD) Residual volume, ml Baseline Group 1: 110 (88.8) Group 2: 84.2 (76.6) 3 Months: Group 1: 67 (101.6) Group 2: 81.2 (66.8)			Baseline Group 1: 110 (88.8) Group 2: 84.2 (76.6) 3 Months: Group 1: 67 (101.6) Group 2: 81.2 (66.8)
			Objective score (simplified version of the Siroky nomogram, lower scores indicates a higher degree of urinary obstruction)			Baseline Group 1: 17.8 (8.5) Group 2: 24.8 (10.3) 3 Months: Group 1: 25.8 (12.0) Group 2: 24.3 (11.8)
Subjective score, ranging from 6 (sever disturbance) to 38 (no disturbance)	Baseline Group 1: 16.7 (7.8) Group 2: 19.4 (8.2) 3 Months: Group 1: 23.0 (10.8) Group 2: 23.6 (7.0)	Baseline Group 1: 16.7 (7.8) Group 2: 19.4 (8.2) 3 Months: Group 1: 23.0 (10.8) Group 2: 23.6 (7.0)				

Evidence Table 35: Transurethral microwave thermotherapy (TUMT) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ahmed et al., 1997⁹</p> <p>Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Single centre, UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Patients presenting with symptomatic, uncomplicated BPH.</p> <p>Inclusion criteria: residual urine volume \leq300 ml; AUA score \geq 12; urine flow rate $<$ 15ml/s, prostate volume 25-100ml by TRUS; symptomatic uncomplicated BPH $>$ 1 year; pdet max $>$70cm H₂O; informed consent; obstructed on Abrams-Griffith nomogram; suitable for either treatment.</p> <p>Exclusion criteria: $<$55years; prostate cancer; previous prostatic surgery; acute or chronic retention; mental incapacity; severe cardiovascular disease; rectal surgery or disease; pelvic mass surgery; cardiac pace marker; metallic implants; uncontrolled coagulation disorder; meatal stricture; upper tract dilation; obstructive uropathy; bladder calculi; bladder diverticuli; recurrent prostatic haematuria; active drugs; previous medication for BPH; prostatic abscess; active UTI; recurrent UTI; prominent middle lobe.</p> <p>Group 1 N: 30 Mean (range) age: 69.36 (56-88) Mean AUA score (95% CI): 18.5 (17.1-20.1) Dropouts: 0</p> <p>Group 2 N: 30 Mean (range) age: 69.45 (58-82) Mean AUA score (95% CI): 18.4 (16.7-20.1) Dropouts: 0</p>	<p>Group 1: TUMT With urethral cooling in a high energy protocol (Prostratron version 2.5). Temperature 43.5 degrees, power at 70W.</p> <p>60 minute session under topical anaesthesia with instillagel. 3 required parenteral pethidine. Antibiotics: gentamycin (80mg) before treatment and oral trimethoprim, 200mg, 2 times day for 5 days.</p> <p>Group 2: TURP No post operative irrigation was used. Urethral catheter was removed 3 or 4 days after surgery.</p>	<p>Mean (range) [SD] AUA symptom scores:</p>	<p>Baseline: Group 1: 18.5 (17.1-20.1) Group 2: 18.4 (16.7-20.1) 6 months: Group 1: 5.3 (3.9-6.4) [3.5] Group 2: 5.2 (3.9-6.5) [3.6]</p>	<p>Funding: NR</p> <p>Limitations: 3 drop outs after randomisation were substituted. One emigrated to Australia; one developed severe UTI requiring hospital admission and one patient could not be catheterised with the treatment catheter.</p> <p>Method of randomisation and use of blinding unclear.</p> <p>Additional outcomes: None</p> <p>Notes: Urodynamic outcomes improved in TURP group but not after TUMT.</p>
			<p>AUA symptom score decreased $>$ 50%</p>	<p>Group 1: 18/30 (60%) Group 2: 30/30 (100%)</p>	
			<p>Qmax (mL/s):</p>	<p>Baseline: Group 1: 10.1 (9.2-10.9) Group 2: 9.5 (8.9-10.1) 6 months: Group 1: 9.1 (8.0-10.2) Group 2: 14.6 (13.4-15.8)</p>	
			<p>Pdet max (cmH₂O):</p>	<p>Baseline: Group 1: 98.5 (70.1-116.9) Group 2: 96.7 (85.5-103.9) 6 months: Group 1: 105.6 (73.7-117.5) Group 2: 48.8 (44.3-52.7)</p>	
			<p>PVR (mL):</p>	<p>Baseline: Group 1: 94.4 (70.0-112.8) Group 2: 109.1 (88.2-130.0) 6 months: Group 1: 104.9 (78.9-130.9) Group 2: 32.5 (22.5-40.5)</p>	
			<p>Prostate volume (mL):</p>	<p>Baseline: Group 1: 36.6 (31.8-41.4) Group 2: 46.1 (38.1-54.1) 6 months: Group 1: 34.5 (29.7-39.3) Group 2: 25.4 (19.4-31.4)</p>	
			<p>Blood transfusion:</p>	<p>Group 1: 0/30</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 4/30	
			Urinary tract infection:	Group 1: 1/30 Group 2: 3/30	
			Strictures:	Group 1: 0/30 Group 2: 1/30	
			Retrograde ejaculation (sexually active men only):	Group 1: 4/18 Group 2: 12/19	
			Hematuria:	Group 1: 1/30 Group 2: 0/30	
			Erectile dysfunction:	Group 1: 0/18 Group 2: 4/19	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Delarosette et al., 2003⁶⁵ Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Netherlands</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Median 33 months.</p>	<p>Patient group: From January 1996 to March 1997 patients with LUTS suggestive of BPH were recruited.</p> <p>Inclusion criteria: age \geq 45 years; duration of LUTS \geq 3 months, prostate volume \geq 30 mL; urethral length \geq 25mm; peak urine flow rate \leq 15ml/s; Residual urine volume \leq 350 ml; and severe co morbidity.</p> <p>Exclusion criteria: acute prostatitis or urinary tract infection; prostate carcinoma; previous prostatic surgery; heart pacemaker; neurological disorders affecting lower urinary tract function; isolate prostate middle lobe protruding in bladder; urethral stricture.</p> <p>All patients N: 155 Group 1: 82 Group 2: 73 Drop outs: 11 (10 refused and 1 died) – 4 from Group 1 and 7 in Group 2. Not included in the ITT analysis as no follow-up data.</p> <p>Group 1 N: 78 Mean (\pmSD) Age: 67(\pm8.3)</p>	<p>Group 1: TUMT Prostatron device and Prostatsoft 2.5 software. Administered under local anaesthesia. Outpatient procedure.</p> <p>Group 2: TURP Under spinal anaesthesia. Mean in-hospital stay of 5.3 days.</p>	<p>Mean (SD) symptom score IPSS</p>	<p>Baseline: Group 1 (n=78): 20 (6.7) Group 2 (n=66): 20 (6.2) 3 months: Group 1: (n=57): 10.5 (7.9) Group 2 (n=55): 5.3 (5.2) 1 year: Group 1 (n=58): 8.1 (6.0) Group 2 (n=48): 3.2 (3.0) 2 years: Group 1 (n=46): 9.3 (7.3) Group 2 (n=38): 3.7 (4.9) 3 years: Group 1 (n=35): 11.5 (6.4) Group 2 (n=33): 2.6 (2.2)</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, allocation concealment and blinding unclear.</p> <p>Additional outcomes: Cost analysis was performed.</p> <p>Notes: Links with Francisca 1999, Francisca 2000, Floratos 2001.</p>
			<p>Mean (SD) IPSS Quality of life question</p>	<p>Baseline: Group 1 (n=78): 4 (0.9) Group 2 (n=66): 4(1.1) 1 year: Group 1 (n=58): 1.9 (1.3) Group 2 (n=48): 0.6 (0.7) 2 years: Group 1 (n=46): 1.9 (1.0) Group 2 (n=38): 0.9 (1.1) 3 years: Group 1 (n=35): 2.3 (1.2) Group 2 (n=33): 0.6 (0.8)</p>	
			<p>Mean (SD) Maximum urinary flow (Q_{max}, mL/s)</p>	<p>Baseline: Group 1: 9.2 (3.1) Group 2: 7.8 (2.8) 3 months: Group 1 (n=54): 15.5 (12.1) Group 2 (n=47): 25.0 (7.5) 1 year: Group 1: 14.9 (7.2) Group 2: 23.8 (10.4) 2 years: Group 1: 13.7 (6.4) Group 2: 22.5 (11.4)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean (\pmSD) IPSS: 20 (\pm6.7) Dropouts: 23 (5 lost to follow up and 2 died unrelated causes, 16 re-treated by TURP=8, laser prostatectomy=1, cystolithotripsy=2, internal optical urethrotomy=1, TUMT=1, alpha blockers=3).</p> <p>Group 2 N: 66 Mean (\pmSD) Age: 66 (\pm8.2) Mean (\pmSD) IPSS: 20 (\pm6.3) Dropouts: 21 (11 lost to follow up and 2 died of unrelated causes, 8 retreated by bladder neck incisions=3, internal optical urethrotomy=2, physiotherapy=1, medication=2).</p>			<p>3 years: Group 1: 11.7 (5.8) Group 2: 22.8 (11.6)</p>	
			Mean (SD) post void residual (PVR, mL)	<p>Baseline: Group 1: 68 (85) Group 2: 97 (99) 1 year: Group 1: 55 (69) Group 2: 20 (49) 2 years: Group 1: 91 (116) Group 2: 29 (39) 3 years: Group 1: 94 (114) Group 2: 35 (56)</p>	
			Patients with re-treatment:	<p>Group 1: 16/78 22.9% (12.5-33.2) Group 2: 8/66 13.2 (4.5-21.9), P=0.215</p>	
			Kaplan-Meier risk of retreatment (36 months)	<p>Group 1: 22.9 (12.5-33.2)% Group 2: 13.2 (4.5-21.9)%, P=0.215</p>	
			Urinary retention:	<p>Group 1: 2/78 (3%) Group 2: 0/66 (0%)</p>	
			Urinary incontinence:	<p>Group 1: 0/78 (0%) Group 2: 1/66 (2%)</p>	
			Stricture:	<p>Group 1: 1/78 (1%) Group 2: 2/66 (3%)</p>	
			Mortality (unrelated causes)	<p>Group 1: 2/78 (3%) Group 2: 2/66 (3%)</p>	
			Retrograde ejaculation (reported in HTA 2008)	<p>Group 1: 24/36 (67%) Group 2: 5/42 (12%)</p>	
			Erectile dysfunction	<p>Group 1: 7/35 (20%) Group 2: 9/53 (17%)</p>	
			Reoperation	<p>Group 1: 13/78 (17%) Group 2: 5/66 (8%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Mattiasson et al., 2007¹⁸⁶ and Wagrell et al., 2002³¹²</p> <p>Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Sweden, Denmark and USA</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 60 months</p>	<p>Patient group: Patients from ten centres in Scandinavia and the United States recruited between October 1998 and November 1999.</p> <p>Inclusion criteria: symptomatic BPH, peak urine flow rate \leq 13ml/s; ml; IPSS score \geq 13; prostate volume 30-100ml.</p> <p>All patients N: 154 eligible Drop outs: 8 withdrawn before treatment</p> <p>Group 1 N: 100 Mean (\pmSD) Age: 67 (8) Mean (\pmSD) IPSS: 21 (5.4) Dropouts before intervention: 3 (screening failures and not treated) Withdrawn at 12m: 9 Withdrawn at 60m: 38 (adverse events=5, treatment failure=10, patient request=22, other =1)</p> <p>Group 2 N: 46 Mean (\pmSD) Age: 69 (8) Mean (\pmSD) IPSS: 20.4 (5.9) Dropouts before intervention: 5 (screening failures and not treated) Withdrawn: 4 Withdrawn at 60m: 12 (reasons: adverse events=4, treatment failure=2, patient request=5 and other=1)</p>	<p>Group 1: TUMT PLFT technique. Given as outpatient procedure requiring sedo-analgesic with or without local anaesthetic. Diazepam, ketorolac, or ketobemidone or combinations of these. Mean duration of treatment 57 (27-80) minutes. Catheter after treatment: 14\pm8 days before removal.</p> <p>Group 2: TURP Urethral catheter usually removed after 3\pm4 days.</p>	<p>Mean (SD) IPSS</p>	<p>Baseline: Group1 (n=99): 21.0 (5.4) Group 2 (=46): 20.4 (5.9) 3 months: Group1 (n=85): 8.4 (5.5) Group 2 (n=41): 6.7 (4.3) 6 months: Group1 (n=95): 7.4 (6.2) Group 2 (n=43): 5.9 (5.0) 12 months: Group1 (n=93): 7.2 (6.2) Group 2 (n=43): 7.1 (6.6) P=0.603 24 months: Group1 (n=77): 7.2 (5.9) Group 2 (n=38): 4.6 (4.4) 36 months: Group 1 (n=68): 8.2 (6.9) Group 2 (n=35): 5.0 (3.9) 48 months: Group 1: (n=56): 7.1 (5.4) Group 2: (n=30):6.4 (6.6) 60 months: Group 1 (n=63): 7.4 (4.8) Group 2 (n=34): 6.0 (5.8)</p>	<p>Funding: ProstaLund. Authors (Wagrell, Schelin, Larson, Mattiasson) are paid consultants to the sponsor of this study.</p> <p>Limitations: Method of randomisation, allocation concealment and blinding not reported.</p> <p>Additional outcomes: Detrusor pressure Qmax at 3 and 6 months.</p> <p>Notes: % of responders at 12 months defined as those with an IPSS of 7 or less or > 50% gain compared with baseline and/or a Qmax of 15mL/s or greater and/or > 50% gain.</p> <p>Links with Wagrell 2004³¹³</p>
			<p>Mean (SD) IPSS Quality of life:</p>	<p>Baseline: Group1 (n=99): 4.3 (1.0) Group 2 (n=46): 4.2 (1.1) 3 months: Group1 (n=84): 1.5 (1.4) Group 2 (n=41): 1.1 (1.6) 6 months: Group1 (n=93): 1.3 (1.4) Group 2 (n=42): 1.0 (1.5) 12 months: Group1 (n=93): 1.4 (1.3) Group 2 (n=43): 1.5 (1.7) 24 months: Group 1 (n=77): 1.3 (1.2)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2 (n=38): 0.9 (1.3) 36 months: Group 1 (n=68): 1.3 (1.2) Group 2 (n=35): 1.0 (1.4) 48 months: Group 1: (n=56): 1.2 (1.0) Group 2: (n=30): 1.0 (1.3) 60 months: Group 1 (n=63): 1.1 (0.9) Group 2 (n=34): 1.1 (1.2)	
			Urinary flow rate (Qmax mL/s):	Baseline: Group1 (n=79): 7.6 ± 2.7 Group 2 (n=35): 7.9 ± 2.7 3 months: Group1 (n=81): 12.8 ± 6.1 Group 2 (n=41): 14.6 ± 9.0 6 months: Group1 (n=91): 13.5 ± 6.1 Group 2 (n=43): 13.8 ± 6.8 12 months: Group1 (n=73): 13.3 ± 6.0 Group 2 n=31): 15.2 ± 7.8 24 months: Group 1 (n=77): 12.4 ± 5.3 Group 2 (n=37): 15.6 ± 9.6 36 months: Group 1 (n=66): 11.9 ± 4.9 Group 2 (n=34): 13.5 ± 7.4 48 months: Group1 (n=49): 12.3 ± 5.7 Group 2 (n=30): 14.7 ± 7.57 60 months: Group 1 (n=61): 11.4 (4.9) Group 2 (n=32): 13.6 (7.8)	
			Mean (SD) residual urine in mL	Baseline: Group1 (n=99): 106 ± 77 Group 2 (n=45): 94 ± 82 12 months:	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1 (n=86): 49 ± 70 Group 2 (n=38): 54 ± 77 24 months: Group 1 (n=75): 56 (63) Group 2 (n=38): 40 (48) 36 months: Group 1 (n=68): 47 (62) Group 2 (n=34): 54 (118) 48 months: Group 1 (n=55): 60 (59) Group 2 (n=29): 55 (53) 60 months: Group 1 (n=63): 70 (90) Group 2 (n=32): 51 (45)	
			Reduction in prostate volume (after 12 months):	Group 1 (n=16): 30% Group 2 (n=13): 51%	
			Additional BPH treatment (5 year follow-up)	Group 1: 10/100 (10%) Group 2: 2/46 (4.3%)	
			Mortality (27 days after treatment)	Group 1: 0/100 Group 2: 1/46	
			Complications	Micturition urgency at 12months: Group 1: 37/100 (37%) Group 2: 6/46 (13%) Urinary retention: 0-12 months: Group 1: 19/100 (19%) Group 2: 6/46 (13%) 12-60 months Group 1: 2/80 (2.5%) Group 2: 0/39 Urinary tract infection: 12 months: Group 1: 18/100 (18%) Group 2: 9/46 (20%) 12-60 months: Group 1: 0/80 Group 2: 1/39 (2.6%)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>Haematuria: 12 months Group 1: 13/100 (13%) Group 2: 18/46 (39%) 12-60 months Group 1: 5/80 (6.3%) Group 2: 0</p> <p>Erectile dysfunction: 12 months: Group 1: 6/100 (6%) Group 2: 5/46 (11%) 12-60 months: Group 1: 6/80 (7.5%) Group 2: 6/39 (15.4%)</p> <p>Transient incontinence 12 months: Group 1: 3/100 (3%) Group 2: 6/46 (13%) 12-60 months: Group 1: 1/80 (1.3%) Group 2: 2/39 (5.1%)</p> <p>TUR syndrome: Group 1: 0/100 Group 2: 1/46</p> <p>Reoperation (up to 60 months): Group 1: 8/100 Group 2: 1/46</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Dahlstrand et al., 1993⁶² Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Sweden</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Inclusion criteria: residual urine volume \leq 350ml; Madsen score \geq 8; prostate length 35-50mm from TRUS. Qmax $<$ 15m/s (twice); BPH; anaesthetic risk group 1-3; obstructive symptoms $>$ 3 months.</p> <p>Exclusion criteria: $<$45 years; suspicion or known prostate cancer or bladder cancer; previous surgery for cancer of prostate or radiotherapy; rectal surgery; prior surgery or heat treatment of BPH; large median lobe; neurogenic bladder disorder; mental incapacity, dementia or inability to give informed consent; neurological disorders that may affect bladder function; peripheral arterial disease; disorder of haemostasis or serum creatinine $>$2mg/dl; uncontrolled cardiac dysrhythmias, or cardiac pacemaker; total hip replacement or other metallic implants; indwelling or condom catheter; post void residual urine $>$350ml; urethral stricture; bladder stones; adrenergic blockers antiandrogen medication or other medication that might affect prostate or bladder; bacterial prostatitis or UTI at time of treatment ; prostatic urethral length of $>$50mm or $<$35mm by transrectal US; anaesthesia risk category 4 or 5.</p> <p>All patients N: 79</p>	<p>Group 1: TUMT Prostatron, Power: 60W; Temperature: urethral: 44.5 degrees and rectal 42.5 degrees.</p> <p>If no voiding use indwelling catheter for 3-5 days. No general anaesthesia but intraurethral topical lidocaine HCl jelly 2% and NSAID. Postoperative oral norfloxacin 400mg twice per day for 5 days. Treatment time 60 minutes.</p> <p>Group 2: TURP performed by urologists were senior registrar or above. Mean operative time: 60.9 minutes. Hospital stay: 5 ± 1.9 days</p>	<p>Mean (SD) Madsen symptom score</p>	<p>Baseline: Group 1 (n=39): 11.2\pm3.1 Group 2(n=39): 13.3\pm4.2 3 months: Group 1(n=37): 2.3\pm2.7 Group 2(n=39): 1.6\pm2.5 6 months: Group 1(n=28): 3.1\pm3.0 Group 2(n=23): 0.9\pm1.6 12 months: Group 1(n=25): 2.7\pm2.9 Group 2(n=22): 0.9\pm2.2</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, allocation concealment and blinding not reported.</p> <p>Additional outcomes: Maximum capacity change. Additional follow-up 6-8 weeks after surgery.</p> <p>Notes: * Catheterisation required but removed within 3-5 days.</p>
			<p>Mean (SD) residual urine volume (ml)</p>	<p>Baseline: Group 1 (n=39): 105\pm88 Group 2 (n=40): 116\pm97 3 months: Group 1(n=37): 55\pm51 Group 2(n=39): 31\pm25 6 months: Group 1(n=28): 68\pm69 Group 2(n=24): 17\pm10 12 months: Group 1 (n=24): 47\pm51 Group 2 (n=22): 22\pm16</p>	
			<p>Mean (SD) maximum flow rate (ml/s)</p>	<p>Baseline: Group 1 (n=39): 8.0\pm2.8 Group 2 (n=40): 7.9\pm3.2 3 months: Group 1 (n=35): 12.2\pm4.9 Group 2 (n=37): 18.7\pm6.0 6 months: Group 1 (n=32):12.0\pm4.5 Group 2 (n=24):18.8\pm5.9 12 months: Group 1 (n=24): 12.3\pm4.7 Group 2 (n=22): 17.7\pm6.5</p>	
			<p>Reoperation:</p>	<p>Group 1: 4/39 (10.2%) Group 2: 0/40</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Drop outs: 4</p> <p>Group 1 N: 39 Mean Age: 68 Prostate volume: 33ml Mean Madsen \pmSD: 11.2\pm 3.1 Dropouts: 0</p> <p>Group 2 N: 40 Mean Age:70 Prostate volume: 37ml Mean Madsen \pm SD: 13.3\pm 4.2 Dropouts: 4 (sever hepatitis=1, cancer discovered=2, refusal for TURP=1).</p>		<p>Re-catheterisation due to unable to void:</p> <p>Transient urgency after surgery</p> <p>Transient urinary leakage</p> <p>Bleeding and rehospitalisation</p> <p>Internal urethrotomy due to stricture</p> <p>Urinary tract infections</p> <p>Men with retrograde ejaculation following surgery (previously with antegrade ejaculations)</p> <p>% Reduction in prostate size (6m)</p> <p>Unstable detrusor contractions</p> <p>Sexually active men</p>	<p>Group 1: 8/39* Group 2: 2/40</p> <p>Group 1: 7/39 Group 2: 4/40</p> <p>Group 1: 0/39 Group 2: 1/40 (2.5%)</p> <p>Group 1: 0/39 Group 2: 3/40</p> <p>Group 1: 0/39 Group 2: 3/40</p> <p>Group 1: 3/39 Group 2: 0/40</p> <p>Group 1: 0 Group 2: 4/16</p> <p>Group 1: 0 Group 2: 47</p> <p>Baseline Group 1: 6/21 Group 2: 5/13 After surgery: Group 1: 8/21 Group 2: 2/13</p> <p>All men who were sexually active before treatment remained so after.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Dahlstrand et al., 1995⁶³ Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Sweden</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Inclusion criteria: residual urine volume \leq 350ml; Madsen score \geq 8; prostate length 35-50mm from TRUS.</p> <p>Exclusion criteria: prostate cancer or bladder cancer; previous surgery for cancer of prostate; prior treatment for BPH; indwelling catheter, urethral stricture; large median lobe; neurogenic bladder disorder, metallic hip implant.</p> <p>All patients N: 72 eligible – 69 randomised Drop outs: 10</p> <p>Group 1 N: 37 Mean Age: 67.9\pm9 Mean Madsen \pm SD: 12.1\pm 3 Dropouts: 2 (died=1, hernia operation=1)</p> <p>Group 2 N: 32 Mean Age:70\pm6 Mean Madsen \pm SD: 13.6\pm 3.9 Dropouts: 8 (TURP=2, abroad=1, refused=1, severe pancreatitis=1, neurological disease=1, reoperation with TUMT and then TURP=2)</p>	<p>Group 1: TUMT Prostatron (Prostasoft 2.0 software) – 60W. Treatment in single session as outpatient. Intra-urethraly applied lidocaine hydrochloride jelly used. Before treatment patients given indomethacin 50mg and norfloxacin 400mg was given; after treatment indomethacin given twice for one day and norfloxacin 400mg twice daily for 5 days.</p> <p>Group 2: TURP by senior registrar grade or above. Mean operation time=48\pm17 minutes. Mean hospital stay=3.9\pm1.3 days.</p>	<p>Madsen symptom score</p>	<p>Baseline: Group 1 (n=37): 12.1\pm3.0 Group 2 (n=32): 13.6\pm3.9 3 months: Group 1 (n=36): 2.9\pm3.0 Group 2 (n=32): 1.7\pm2.6 6 months: Group 1 (n=37): 2.6\pm2.6 Group 2 (n=32): 1.1\pm1.8 12 months: Group 1 (n=33): 2.2\pm2.4 Group 2 (n=31): 0.6\pm1.4 24 months: Group 1 (n=31): 2.3\pm3.0 Group 2 (n=30): 1.2\pm1.9</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, use of allocation concealment and blinding were not reported.</p> <p>Unsure if same study as Dahlstrand 1993 – HTA attempted to contact authors.</p> <p>Additional outcomes: Volume at first sensation to void after 6 months. Detrusor contractions and urethral resistance factor.</p> <p>Notes: Reoperation: TUMT group=4: 2 retreated by TURP, 2 by TUMT; the TUMT reoperations had TURP at 1 year due to unsatisfactory improvement. TURP group: reoperation from early complication=3 due to bleeding or to remove clots; 1 retreatment after 1 year due to bladder neck sclerosis.</p>
			<p>Reduction in symptom score > 50%</p>	<p>Group 1: 26/31 Group 2: 29/30</p>	
			<p>Maximum flow rate (mL/s)</p>	<p>Baseline: Group 1 (n=37): 8.6\pm2.5 Group 2 (n=32): 8.6\pm3.0 3 months: Group 1 (n=36): 11.6\pm4.2 Group 2 (n=32): 18.1\pm7.1 6 months: Group 1 (n=37): 11.8\pm3.9 Group 2 (n=31): 18.6\pm5.2 12 months: Group 1 (n=33): 12.6\pm3.9 Group 2 (n=31): 18.9\pm6.0 24 months: Group 1 (n=30): 12.3\pm4.4 Group 2 (n=29): 17.6\pm5.9</p>	
			<p>Residual urine volume (mL)</p>	<p>Baseline: Group 1 (n=37): 194\pm78 Group 2 (n=32): 1104\pm95 3 months: Group 1 (n=36): 147\pm45 Group 2 (n=32): 134\pm32 6 months:</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1 (n=37): 166±64 Group 2 (n=32): 134±30 12 months: Group 1 (n=33):152±64 Group 2 (n=31): 123±18 24 months: Group 1 (n=31):148±44 Group 2 (n=30):127±2	
			Prostate volume	Baseline: Group 1: 33.9±11.9 Group 2: 36.8 ±16 2 years: Group 1: 30.3 ±9.6 Group 2: 22.5±10.9	
			Reoperation:	Group 1: 4/37 Group 2: 1/32	
			Catheterisation due to failure to void	Group 1: 5/37 Group 2: 0/32	
			Transient rectal pain in perineum	Group 1: 1/37 Group 2: 0/32	
			Urethral stricture	Group 1: 0/37 Group 2: 2/32	
			Meatal stenosis	Group 1: 0/37 Group 2: 2/32	
			Urinary tract infection	Group 1: 5/37 Group 2: 4/32	
			Mortality (brain tumour)	Group 1: 0/37 Group 2: 1/32	
			Erectile dysfunction	Group 1: 0/37 Group 2: 0/32	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>D'Ancona et al., 1998⁶¹ Reported in systematic review HTA 2008</p> <p>Study design: RCT</p> <p>Setting: Netherlands</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2.5 years</p>	<p>Patient group: Between January 1994 and August 1995 patients recruited.</p> <p>Inclusion criteria: unequivocal BPH candidates for TURP. Qmax 15ml/s; residual volume <350ml; Madsen score ≥ 8; prostate length 25-50mm, Prostate Volume 30-100ml; 45 years plus.</p> <p>Exclusion criteria: prostate cancer; prior prostate surgery; urinary retention requiring catheterisation; medications prescribed for prostate/bladder treatment; neurogenic disorders affecting bladder function; diabetic neuropathy; possible microwave sensitive implants (pacemaker, hip prosthesis); renal impairment or obstructed bladder neck due to enlarged median lobe of prostate</p> <p>All patients N: 52</p> <p>Group 1 N: 31 Mean Age ± SD: 69.6 ± 8.5 Mean IPSS ± SD: 18.3 ± 6.3 Dropouts: 14 (6 TURP, 1 died, 5 refused or lost to follow up, 2 medication)</p> <p>Group 2 N: 21 Mean Age ± SD: 69.3 ± 5.9 Mean IPSS ± SD: 16.7 ± 5.6</p>	<p>Group 1: TUMT – Prostatron software version 2.5. Total mean energy applied 151.8kJ. 100mg suppository of diclofenac administered and 2mg of medazolam injected. No additional anaesthesia during treatment. Out patient. Prolonged catheterisation: 12.7 days.</p> <p>Group 2: TURP by 2 urologists and resection performed under spinal anaesthesia. Mean length of hospital stay 4.1. Mean catheterisation 4.1 days.</p>	<p>Mean (SD) IPSS score:</p>	<p>Baseline: Group 1 (n=31): 18.3 (6.3) Group 2 (n=21): 16.7 (5.6) 3 months: Group 1 (n=31): 15.1 (8.2) Group 2 (n=21): 5.1 (3.1) 6 months: Group 1 (n=28): 6.7 (5.5) Group 2 (n=20): 4.0 (2.1) 12 months: Group 1 (n=27): 5.0 (2.7) Group 2 (n=17): 3.4 (2.2) 30 months: Group 1 (n=17): 7.9 (6.3) Group 2 (n=12): 6.3 (4.8)</p>	<p>Funding: NR</p> <p>Limitations: Method of randomisation, allocation concealment and blinding unclear.</p> <p>Additional outcomes: Madsen score, voided volumes, URA and LPURR.</p> <p>Notes: Links with D'Ancona 1997⁶⁰</p>
			<p>Qmax (mL/s)</p>	<p>Baseline: Group 1 (n=31): 9.3 (3.9) Group 2 (n=21): 9.3 (3.4) 3 months: Group 1 (n=31): 15.5 (8.0) Group 2 (n=21): 19.6 (11.2) 6 months: Group 1 (n=38): 17.0 (7.5) Group 2 (n=20): 15.3 (5.9) 12 months: Group 1 (n=27): 17.1 (7.8) Group 2 (n=17): 19.3 (29.8) 30 months: Group 1 (n=17): 15.1 (9.6) Group 2 (n=12): 19.1 (8.2)</p>	
			<p>PVR (mL)</p>	<p>Baseline: Group 1 (n=31): 49.5 (69.9) Group 2 (n=21): 91.1 (104.7) 3 months: Group 1 (n=31): 25.5 (58.1) Group 2 (n=21): 10.5 (24.5) 6 months: Group 1 (n=28): 30.6 (41.0)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Drop outs: 9 (4 refused or lost to follow up, 1 bladder neck incision, 1 bladder carcinoma, 1 at own request, 2 dementia)			Group 2 (n=20): 52.7 (70.7) 12 months: Group 1 (n=27): 70.4 (81.3) Group 2 (n=17): 23.6 (29.8) 30 months: Group 1 (n=17): 27.4 (49.1) Group 2 (n=12): 9.3 (14.6)	
			Pdet Qmax (cmH2O)	Baseline Group 1: 77.7 (40.0) Group 2: 65.4 (24.9) 6 months: Group 1: 54.0 (15.9) Group 2: 38.5 (24.5)	
			Prostate volume (mL)	Baseline Group 1: 43.4 (11.8) Group 2: 44.9 (15.3) 3 months: Group 1: 36.6 (10.0) Group 2: 23.0 (8.8)	
			Reoperation:	Group 1: 2/31 (6.4%) Group 2: 1/21 (4.8%)	
			Blood transfusions	Group 1: 0/31 Group 2: 0/21	
			UTI	Group 1: 5/31 (16%) Group 2: 1/21 (4%)	
			Irritative voiding symptom	Group 1: 9 (29%) Group 2: 4 (19%)	
			Hematuria	Group 1: 0 Group 2: 3 (14%)	
			Mortality	Group 1: 1 Group 2: 0	

Evidence Table 36: Transurethral vaporisation of the prostate (TUVP) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Çetinkaya et al.,1996⁴⁷</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months after surgery</p>	<p>Patient group: moderate or severe symptoms of prostatism</p> <p>Setting: single centre, urology clinic, Ankara Nummune Hospital, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Peak urine flow rate <15 AUA moderate to severe <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Patients who had previously undergone a prostate operation or who had any abnormality of kidney and liver function, urethral strictures, neurogenic deficits, bladder stones Those with confirmed or suspected prostate cancer. <p>All patients N: 46 Drop outs: NR</p> <p>Group 1: N: 23 Age (mean ± SD): 68.4 ± 8.3 Mean prostate size ± SD: 48.4 ± 9.7 ml (TRUS) Operative duration ± SD: 41.6 ± 22.1 min Solution volume used ± SD: 16.0 ± 10.2 ml Catheterisation time (days): 1.4 ± 0.8 days Length of stay (days): NR Drop outs: NR</p> <p>Group 2: N: 23 Age (mean ± SD): 62.5 ± 10.1</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Storz Spike 5mm 2-system electrode. cutting mode: 240-300 W & coagulation mode: 40-70 W TUVP continued until capsule was visible</p> <p>Group 2: Transurethral resection of the prostate (TURP) Conventional electroresection</p> <p>All patients: Glycine was used as irrigant. Indwelling catheter placed after surgery and removed when urine was clear.</p> <p>Examination Methods Preoperative: Baseline prostate volume (TRUS), digital rectal examination, uroflowmetry, haemocrit & Na⁺ levels, AUA symptom score, post void residual (PVR) Postoperative PVR, symptom score and uroflowmetry taken 3 months after catheter removed. Haemocrit & Na⁺ levels taken 24 h after surgery</p>	<p>Mean change in AUA symptom score from baseline at 3 months</p> <p>Mean change in Qmax from baseline at 3 months</p> <p>Mean change in PVR from baseline at 3 months</p> <p>Complications: transfusion</p> <p>Complications: re-catheterisation required (retention)</p> <p>Complications: urethral or meatal stricture:</p>	<p>Group 1: -20.89 Group 2: -21.31 p value: NR</p> <p>Group 1: 16.37 Group 2: 17.49 p value: NR</p> <p>Group 1: -211.52 Group 2: -199.05 p value: NR</p> <p>Group 1: 0/23 Group 2: 2/23</p> <p>Group 1: 4/23 Group 2: 0/23</p> <p>Group 1: 1/23 Group 2: 0/23</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masking of outcome assessment not reported Symptom score and Qmax were not reported at 3 months or at baseline Standard deviations not reported for changes from baseline Not clear whether ITT analysis performed Drop outs not reported <p>Additional outcomes: Irritative symptoms after catheter removal more in TUVP group.</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean prostate size \pm SD: 48.8 \pm 15.4 ml (TRUS)</p> <p>Operative duration \pm SD: 52.4 \pm 20 min</p> <p>Solution volume used \pm SD: 19.8 \pm 8.6 ml</p> <p>Catheterisation time (days): 1.9 \pm 0.8 days</p> <p>Length of stay (days): NR</p> <p>Drop outs: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Ekengren et al., 2000 ⁷⁷	<p>Patient group: men scheduled for surgery for obstruction</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Roller-ball 27050 electrode (Stortz) Cutting mode: 240 W</p> <p>Group 2: Transurethral resection of the prostate (TURP) Conventional electroresection</p> <p>All patients: Operations performed using 26F resectoscope. Ringer's solution with heparin used to replace blood lost measured using a photometer. Irrigating fluid of mannitol & ethanol and fluid absorption using ethanol method.</p> <p>Preoperative: Baseline prostate volume & PVR (TRUS), IPSS, uroflowmetry (Flo-LabII), serum PSA, Quality of Life Score (QoL) score,</p> <p>Postoperative prostate volume & PVR (TRUS), IPSS, uroflowmetry (Flo-LabII), serum PSA, Quality of Life Score (QoL) score</p>	<p>Median IPSS score (range) at 12 months</p> <p>Mean \pm SD IPSS at 12 months*</p> <p>Median Qmax mL/s (range) at 12 months</p> <p>Mean Qmax \pm SD mL/s at 12 months*</p> <p>Median QoL score (range) at 12 months</p> <p>Mean \pm SD QoL at 12 months*</p> <p>Complications: mortality</p> <p>Complications: transfusion</p> <p>Complications: urethral stricture</p> <p>Complications: urinary retention</p> <p>Complications: reoperation rate</p>	<p>Group 1: 4.5 (0-24) Group 2: 4.0 (0-100) p value: Not</p> <p>Group 1: 7.0 \pm 6.5 ** Group 2: 9.3 \pm 19.8 ** p value: NR</p> <p>Group 1: 10 (4-19) Group 2: 11 (0-19) p value: Not sig.</p> <p>Group 1: 10.7 \pm 4.1 (n=23) Group 2: 11.1 \pm 4.4 (n=28) p value: NR</p> <p>Group 1: 1.5 (0-6) Group 2: 1.0 (0-6) p value: Not sig.</p> <p>Group 1: 1.8 \pm 1.6 (n=23) Group 2: 1.8 \pm 2.0 (n=28) p value: NR</p> <p>Group 1: 2/26 Group 2: 0/28</p> <p>Group 1: 0/26 Group 2: 0/28</p> <p>Group 1: 2/26 Group 2: 0/28</p> <p>Group 1: 0/26 Group 2: 1/28</p> <p>Group 1: 2/26 Group 2: 1/28</p>	<p>Funding: Supported by the Board of Research and Education of Stockholm County Council</p> <p>Limitations:</p> <ul style="list-style-type: none"> Patients and investigators were unmasked to treatment allocation Not clear whether ITT analysis performed **Values for mean IPSS given by author were very different to the median reported in the study values at baseline were >35 <p>Additional outcomes: Significantly higher blood loss during the operation for TURP. Unable to check p value.</p> <p>Notes: *Requested Mean IPSS, Qmax, QoL and follow up data from author. Author reports that data were skewed hence presented as median and range. Author reported</p>
<p>Study design: RCT Unmasked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months after surgery</p>	<p>Setting: single centre, department of surgery and urology, Söder Hospital, Stockholm, Sweden</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria: NR</p> <p>All patients N: 54 Drop outs: 3 died (TUVP)</p> <p>Group 1: N: 26 Median age (range): 71 (49-82) Median IPSS (range): 22 (1-100) Median QoL score (range): 4.5 (2-6) Mean QoL score \pm SD: 4.6 \pm 1.2* Median PSA (range): 4 (2-23) ng/mL Median PVR (range): 55 (0-3000) mL Median Qmax (range): 4 (0-8) mL/s Mean Qmax \pm SD: 3.7 \pm 2.4 mL/s* Median prostate vol. (range): 50 (25-90) mL (TRUS) Median operative duration (range): 30 (15-80) min Median blood loss (range): 75 (8-400) mL Drop outs: 3 (1 died from myocardial infarction, 1 died (catheter) and 1 with urethral stricture lost to follow up)</p> <p>Group 2: N: 28 Median age (range): 70 (48-83)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Median IPSS (range): 25 (13-100) Median QoL score (range): 5.5 (3-6) Mean QoL score \pm SD: 5.2 \pm 1.0* Median PSA (range): 6 (1-82) ng/mL Median PVR (range): 100 (0-3000) mL Median Qmax (range): 2 (0-10) mL/s Mean Qmax \pm SD: 2.8 \pm 3.0 mL/s* Median prostate vol. (range): 39 (20-80) mL (TRUS) Median operative duration (range): 33 (10-90) min Median blood loss (range): 150 (10-726) mL Drop outs: 0</p>				<p>randomisation performed by drawing of sealed envelopes from a box prior to surgery</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Erdagi et al., 1999⁸³</p> <p>Study design: RCT Unmasked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months after surgery</p>	<p>Patient group: men with symptomatic BPH</p> <p>Setting: single centre, Turkish High Specialisation Hospital, Ankara, Turkey</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria: NR</p> <p>All patients N: 40 Drop outs: NR</p> <p>Group 1: N: 20 Mean age (range): 64.2 (56-82) Mean IPSS (range): 20.6 (12-27) (n=15*) Mean Qmax ml/s (range): 5.1 (0-11.27) (n=15*) Mean PVR ml (range): 68 (20-150) Mean prostate weight. (range): 32.5 (20-48) (TRUS) Mean operative duration (range): 61.5 min Mean operative blood loss ml: 117.6 Catheterisation time (days): 1.1 Drop outs: NR</p> <p>Group 2: N: 20 Mean age (range): 66.1 (58-75) Mean IPSS (range): 21.5 (11-30) (n=15*) Mean Qmax ml/s (range): 4.6 (0-9.6) (n=15*)</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) VaporTrode® rollerball electrode (Storz) at 240W for cutting and 40W for coagulation.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard 0.012 inch loop</p> <p>All patients: Operations performed using 26F resectoscope under continuous 1.5% mannitol solution.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, PSA, uroflowmetry using Synectics Urodynamics Polygraph System, PVR by ultrasonography and prostate volume by TRUS. Assessed at 1, 3 & 6 months postoperatively</p>	<p>Mean IPSS score (range) at 3 months</p>	<p>Group 1: 0.9 ± NR (0-4) Group 2: 5.3 ± NR (1-12) p value: Not sig.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Mean and standard deviations not reported for outcomes at baseline or end point. Randomisation method and allocation concealment not reported Masking of patients or outcome assessment not reported Dropouts not reported Small sample size <p>Notes: Mann Whitney test was used for statistical analysis</p>
			<p>Mean IPSS score (range) at 6 months</p>	<p>Group 1: 0.6 ± NR (0-3) Group 2: 3.9 ± NR (1-9) p value: 0.92 (Mann Whitney-U)</p>	
			<p>Mean Qmax mL/s (range) at 3 months</p>	<p>Group 1: 21.0 ± NR Group 2: 17.0 ± NR p value: NR</p>	
			<p>Mean Qmax mL/s (range) at 6 months</p>	<p>Group 1: 21.4 ± NR Group 2: 17.7 ± NR p value: 0.04 (Mann Whitney-U)</p>	
			<p>Catheterisation time (days)</p>	<p>Group 1: 1.1 ± NR Group 2: 3.4 ± NR p value: <0.001</p>	
			<p>Complications: transfusion</p>	<p>Group 1: 0/20 Group 2: 9/20 p value: NR NCC_AC calculate p=0.01 Fishers exact test</p>	
			<p>Complications: retrograde ejaculation</p>	<p>Group 1: 2/20 Group 2: 12/20</p>	
			<p>Complications: UTI</p>	<p>Group 1: 1/20 Group 2: 5/20 p value: NR NCC_AC calculate p=0.18 Fishers exact test</p>	
			<p>Complications: Urethral Stricture</p>	<p>Group 1: 0/20 Group 2: 1/20 p value: NR NCC_AC calculate p=1.00 Fishers exact test</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean PVR ml (range): 123 (0-600) Mean prostate weight. (range): 37 (15-60) (TRUS) Mean operative duration (range): 67.7 min Mean operative blood loss ml: 491 Catheterisation time (days): 3.4 Drop outs: NR</p> <p>*10 patients with chronic retention with indwelling catheter also included did not have baseline IPSS or Qmax data</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Fowler et al, 2005⁹⁴</p> <p>Study design: RCT Single masked (though patients on regional anaesthetic may have known which operation they had)</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient group: men considering surgery for BPH</p> <p>Setting: multi-centre, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Must have completed pre-treatment evaluation with current criteria for prostate surgery. Able to give written informed consent to randomisation and treatment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous bladder outlet surgery clinical evidence of prostate cancer Physical status >ASA 3 Medications that (in investigators opinion) would preclude entry into trial Clinically significant acute illness Known disease of central or peripheral nervous system. Prostate cancer. <p>All patients N: 235 45/235 patients in acute retention</p> <p>Drop outs: Number of patients completing study NR</p> <p>Group 1: N: 115 Mean age (± SD): 70.2 ± NR Mean IPSS (± SD): 20.7 ± 7.2 (n=107) Mean EuroQoL score: 0.78 ± 0.23 (n=112) Mean IPSS QoL: 4.6 ± 1.7 (n=109) Mean PSA (± SD): 4.7 ± NR ng/mL (n=101) Mean PVR (± SD): 181 ± NR mL (n=91) Mean Qmax (SD): 10.1 ± 4.35 mL/s (n=94)</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Circon-ACMI 24.5 Fr continuous flow rectoscope with new Circon- ACMI Fluted VaporTrode® electrode for each patient. 180W for cut and 55W for coagulation</p> <p>Group 2: Transurethral resection of the prostate (TURP) Circon-ACMI 24.5 Fr continuous flow rectoscope with new wire loop for each patient. Cutting mode: 120-140 W. Coagulation mode: 50-60 W</p> <p>All patients: Irrigating fluids varied between glycine and glycine & ethanol depending on the centre.. 3-way catheters were removed when degree of haematuria was permitted.</p> <p>Preoperative: Baseline blood tests (FBC, urea, PSA), Uroflow using Dantec Urodyn 1000 (2 flow rates >150mL if possible), PVR using TRUS</p>	<p>Mean change in IPSS Score from baseline ± SD at 2 mths</p>	<p>Group 1: 9.8 ± 7.2 (n=105) Group 2: 11.8 ± 7.7 (n=110) p value NR</p>	<p>Funding: Supported the INAHTA Health Technology Assessment programme</p> <p>Limitations:</p> <ul style="list-style-type: none"> Baseline data was not available for all outcomes Drop outs reported for primary outcome rather than those completing study Investigators were not masked to treatment allocation <p>Additional outcomes: Change in General Health related EuroQoL score from baseline Erectile dysfunction, failed ejaculation, change in ejaculatory function, change in PVR and prostate volume. Additional procedures</p> <p>Notes: Randomisation method was computer generated by study organisers and allocation concealment by sequentially numbered opaque envelopes.</p>
			<p>Mean change in IPSS Score from baseline ± SD at 6 mths</p>	<p>Group 1: 8.5 ± 7.4 (n=106) Group 2: 6.9 ± 5.5 (n=108) p value NR</p>	
			<p>Mean change in IPSS Score from baseline ± SD at 2 years</p>	<p>Group 1: 8.6 ± 7.2 (n=90) Group 2: 7.5 ± 5.8 (n=77) p value NR</p>	
			<p>Mean change in IPSS QoL Score from baseline ± SD at 2 mths</p>	<p>Group 1: 2.6 ± 1.82 (n=105) Group 2: 2.3 ± 1.73 (n=109) p value NR</p>	
			<p>Mean change in IPSS QoL Score from baseline ± SD at 6 mths</p>	<p>Group 1: 2.0 ± 1.63 (n=107) Group 2: 1.6 ± 1.34 (n=108) p value NR</p>	
			<p>Mean change in IPSS QoL Score from baseline ± SD at 2 years</p>	<p>Group 1: 1.9 ± 1.62 (n=89) Group 2: 1.8 ± 1.34 (n=80) p value NR</p>	
			<p>Mean change in Qmax from baseline ± SD at 2 mths</p>	<p>Group 1: 19.12 ± 11.76 (n=108) Group 2: 21.23 ± 10.20 (n=111) p value NR</p>	
			<p>Mean change in Qmax from baseline ± SD at 6 mths</p>	<p>Group 1: 19.60 ± 11.04 (n=109) Group 2: 22.29 ± 10.25 (n=109) p value NR</p>	
			<p>Duration of catheterisation (days)</p>	<p>Group 1: 4.9 ± 11.6* (CI95% 2.7-7.1) n=107 Group 2: 3.1 ± 4.4* (CI95% 2.3-3.9) n=116 p value: 0.93</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 4.4 ± 3.6* (CI95% 3.8-5.1) n=115 Group 2: 4.6 ± 4.2* (CI95% 3.9-5.4) n=120 p value: 0.47</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean prostate vol. (SD): 54.3 ± NR mL (TRUS) (n=100) Serum creatinine (mmol/L): 105 ± NR (n=100) Number of patients with ED: 34/109 Drop outs: 6/115 violated protocol. Number of patients completing study NR</p> <p>Group 2: N: 120 Mean age (± SD): 69.7 ± NR Mean IPSS (± SD): 20.7 ± 6.9 (n=114) Mean EuroQoL score: 0.74 ± 0.25 (n=116) Mean IPSS QoL: 4.9 ± 0.98 (n=114) Mean PSA (± SD): 4.6 ± NR ng/mL (n=99) Mean PVR (± SD): 171 ± NR mL (n=94) Mean Qmax (SD): 10.52 ± 5.04 mL/s (n=97) Mean prostate vol. (SD): 51.1 ± NR mL (TRUS) (n=103) Serum creatinine (mmol/L): 104 ± NR (n=106) Number of patients with ED: 48/110 Drop outs: 6/120 violated protocol Number of patients completing study NR</p>	<p>7.5 MHz, Cystometry and questionnaires: IPSS, EuroQoL, Sexual Function from ICS-BPH questionnaire.</p> <p>Postoperative Assessment at 2 months, 6 months: Blood tests (FBC & urea only) Uroflow using Dantec Urodyn 1000 (2 flow rates >150mL if possible), PVR using TRUS 7.5 MHz, cystometry and questionnaires: IPSS, EuroQoL, Sexual Function from ICS-BPH questionnaire. IPSS Score, ICS-BPH & EuroQoL repeated 2 years as well.</p>	<p>Complications: transfusion</p> <p>Complications: reoperation rate (TUIP)</p> <p>Complications: urethral or meatal stricture. Reported as number of meatotomies, otis urethrotomies and urethral dilatations</p>	<p>Group 1: 2/115 Group 2: 9/120 P value: 0.04 (Chi-squared)</p> <p>Group 1: 5/115 Group 2: 17/120 P value: NR</p> <p>Group 1: 64/115 Group 2: 66/120</p>	<p>*SD calculated from confidence intervals and sample size according to section 7.7.3.2 of the Cochrane Handbook Number of patients in each group was not reported for length of stay data but states that data collected for all but 3 patients. Use numbers randomised for calculation.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gallucci et al, 1998⁹⁹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: men symptomatic men with BPH who were urodynamically obstructed</p> <p>Setting: multi-centre, 9 centres, Italy</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Complete urinary retention • Bladder calculi • Neurogenic bladder • Prostate weight >70g • Bladder cancer • Mental illness • Prostate cancer or suspect <p>All patients N: 150 Drop outs: 0</p> <p>Group 1: N: 70 Mean age (range): NR Mean IPSS \pm SD: 18.84 \pm 5.69 Mean Qmax ml/s \pm SD: 7.26 \pm 3.1 Mean PVR ml \pm SD: 84.7 \pm 95.3 Mean prostate weight \pm SD (g): 36.61 \pm 12.72 Drop outs: 0</p> <p>Group 2: N: 80 Mean age (range): NR Mean IPSS \pm SD: 18.19 \pm 5.90 Mean Qmax ml/s \pm SD: 8.78 \pm 10.38 Mean PVR ml \pm SD: 64.61 \pm 77.37</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) VaporTrode® rollerball electrode (Circon ACMI) at 200-250W for cutting.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard diathermic loop</p> <p>All patients: Operations performed using 22.5F resectoscope under continuous 5% mannitol solution. 3-way catheter inserted. Prophylactic antibiotics were used.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, PSA, Blood, TRUS, uroflowmetry (opening pressure, detrusor pressure, Qmax and PVR using <6f catheters). Flow rate at months 1 & 6 and pressure flow at 3 months. IPSS assessed at 1, 3, 6 & 12 months postoperatively</p>	<p>Mean IPSS score \pm SD at 3 months</p>	<p>Group 1: 5.50 \pm 4.77 Group 2: 5.52 \pm 4.11 p value: Not sig.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method and allocation concealment not reported • Masking of outcome assessment not reported <p>Additional outcomes: Detrusor and opening pressure at 3 months. Transient stress incontinence.</p> <p>Notes: No patients were lost to follow up</p> <p>SD calculated from standard error and and sample size according to section 7.7.3.2 of the Cochrane Handbook numbers randomised for calculation.</p>
			<p>Mean IPSS score \pm SD at 6 months</p>	<p>Group 1: 4.94 \pm 4.69 Group 2: 3.77 \pm 3.31 p value: Not sig.</p>	
			<p>Mean IPSS score \pm SD at 12 months</p>	<p>Group 1: 4.04 \pm 4.27 Group 2: 3.52 \pm 3.04 p value: Not sig.</p>	
			<p>Mean Qmax mL/s \pm SD at 3 months</p>	<p>Group 1: 18.18 \pm 7.7 Group 2: 19.21 \pm 8.14 p value: Not sig.</p>	
			<p>Mean Qmax mL/s \pm SD at 6 months</p>	<p>Group 1: 20.13 \pm 9.62 Group 2: 20.77 \pm 8.5 p value: Not sig.</p>	
			<p>Mean Qmax mL/s \pm SD at 12 months</p>	<p>Group 1: 20.31 \pm 6.02 Group 2: 20.30 \pm 6.35 p value: Not sig.</p>	
			<p>Catheterisation time (days)</p>	<p>Group 1: 1.96 \pm 1.09 Group 2: 2.71 \pm 1.07 p value: <0.0001</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 3.9 \pm 2.01 Group 2: 4.69 \pm 1.97 p value: <0.0001</p>	
			<p>Complications: incontinence (at 12 mths)</p>	<p>Group 1: 4/70 Group 2: 3/80 p value: NR</p>	
			<p>Complications: Urethral Stricture</p>	<p>Group 1: 3/70 Group 2: 3/80 p value: NR</p>	
<p>Complications: transfusion</p>	<p>Group 1: 0/70 Group 2: 0/80 p value: NR</p>				
<p>Complications: transient urinary retention</p>	<p>Group 1: 12/70 Group 2: 3/80 p value: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean prostate weight \pm SD (g): 36.59 \pm 12.25 Drop outs: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Hammadeh et al., 2003¹¹⁰ linked to Hammadeh et al., 2000¹¹¹ & Hammadeh et al., 1998¹⁰⁹</p> <p>Study design: RCT Investigator masked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 5 years</p>	<p>Patient group: men with bladder outflow obstruction due to BPH considering surgery</p> <p>Setting: single-centre, Whipps Cross Hospital, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 13 QoL index \geq 3 Qmax \leq 15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Complete urinary retention Neurogenic bladder Previous prostatic or urethral surgery Bladder calculi Prostate cancer or suspect Receiving anticoagulant therapy <p>All patients N: 104 (109 randomised but 5 excluded for medical problems or social circumstances) Drop outs: *51 at 5 years: 6 TURP and 3 TUVP died from cardiopulmonary disease, 12 TURP and 16 TUVP lost to follow up. Remaining 14 patients unaccounted for.</p> <p>Group 1: N: 52 Mean age (\pm SD): 67.5 \pm 6.7 (52-82) Mean IPSS \pm SD: 26.5 \pm 4.5</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Circon VaporTrode® roller-ball at 240W for cutting & 60W coagulation.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop with 145W cutting & 60W coagulation</p> <p>All patients: Operations performed using 27F resectoscope using continuous glycine. 3-way catheter inserted. TURP patients were irrigated postoperatively until bleeding stopped.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up visits at 6 weeks, 3, 6 & 12 months, 2, 3 5 years postoperatively</p>	<p>Mean IPSS score \pm SD at 1 year</p>	<p>Group 1: 4.4 \pm 3.8 (n=51) Group 2: 5.9 \pm 5.2 (n=51) p value: 0.3</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Dropouts were only partially reported. <p>Additional outcomes:</p> <p>Notes: Patients allocated by nurse drawing a sealed opaque envelope prior to surgery.</p>
			<p>Mean IPSS score \pm SD at 2 years</p>	<p>Group 1: 4.3 \pm 3.5 (n=47) Group 2: 6.3 \pm 4.6 (n=47) p value: 0.02</p>	
			<p>Mean IPSS score \pm SD at 3 years</p>	<p>Group 1: 4.1 \pm 3.3 (n=40) Group 2: 7.1 \pm 6.2 (n=40) p value: 0.01</p>	
			<p>Mean IPSS score \pm SD at 5 years</p>	<p>Group 1: 5.9 \pm 6.3 (n=26) Group 2: 8.6 \pm 7.1 (n=27) p value: 0.16</p>	
			<p>Mean Qmax mL/s \pm SD at 1 year</p>	<p>Group 1: 22.5 \pm 9.0 (n=51) Group 2: 20.8 \pm 7.7 (n=51) p value: 0.4</p>	
			<p>Mean Qmax mL/s \pm SD at 2 years</p>	<p>Group 1: 22.4 \pm 7.7 (n=47) Group 2: 21.2 \pm 8.5 (n=47) p value: 0.5.</p>	
			<p>Mean Qmax mL/s \pm SD at 3 years</p>	<p>Group 1: 22.2 \pm 8.5 (n=40) Group 2: 18.0 \pm 7.1 (n=40) p value: 0.02</p>	
			<p>Mean Qmax mL/s \pm SD at 5 years</p>	<p>Group 1: 21.0 \pm 9 (n=26) Group 2: 17.9 \pm 13.1 (n=27) p value: 0.17</p>	
			<p>Mean IPSS QoL \pm SD at 1 year</p>	<p>Group 1: 1.2 \pm 1.0 (n=51) Group 2: 1.5 \pm 1.0 (n=51) p value: 0.3</p>	
			<p>Mean IPSS QoL \pm SD at 2 years</p>	<p>Group 1: 1.1 \pm 1.0 (n=47) Group 2: 1.7 \pm 1.1 (n=47) p value: 0.004</p>	
			<p>Mean IPSS QoL \pm SD at 3 years</p>	<p>Group 1: 1.0 \pm 0.9 (n=40) Group 2: 1.6 \pm 1.4 (n=40) p value: 0.04</p>	
			<p>Mean IPSS QoL \pm SD at 5 years</p>	<p>Group 1: 1.1 \pm 1.2 (n=26) Group 2: 1.7 \pm 1.4 (n=27) p value: 0.09</p>	
			<p>Catheterisation time (days) hours reported converted to days</p>	<p>Group 1: 0.87 \pm 0.29 Group 2: 1.94 \pm 0.52 p value: <0.001</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 2.2 \pm 0.59 Group 2: 3.19 \pm 0.76 p value: <0.001</p>	
			<p>Complications: transfusion (early)</p>	<p>Group 1: 0/52 Group 2: 1/52 p value: 0.3</p>	
<p>Complications: urinary retention (early)</p>	<p>Group 1: 12/52 Group 2: 4/52 p value: 0.04</p>				
<p>Complications: UTI</p>	<p>Group 1: 3/52</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>IPSS QoL \pm SD: 4.9 ± 0.9 Mean Qmax ml/s \pm SD: 8.9 ± 3.2 Mean PVR ml \pm SD: 131.0 ± 78.5 Mean prostate weight \pm SD (g): 32.0 ± 9.1 Drop outs: *</p> <p>Group 2: N: 52 Mean age (\pm SD): 70.2 ± 7.2 (52-87) Mean IPSS \pm SD: 26.6 ± 4.8 IPSS QoL \pm SD: 5.0 ± 0.7 Mean Qmax ml/s \pm SD: 8.6 ± 3.2 Mean PVR ml \pm SD: 101.0 ± 87.93 Mean prostate weight \pm SD (g): 27.0 ± 12.2 Drop outs: *</p>		<p>(early)</p> <p>Complications: TUR (early)</p> <p>Complications: urethral stricture (long term)</p> <p>Complications: incontinence (long term)</p> <p>Complications: Retrograde ejaculation</p> <p>Reoperation rate</p> <p>Mortality at 5 years (cardiopulmonary)</p>	<p>Group 2: 2/52 p value: 0.7</p> <p>Group 1: 0/52 Group 2: 0/52 p value: 0.7</p> <p>Group 1: 2/52 Group 2: 2/52 p value: NR</p> <p>Group 1: 0/52 Group 2: 0/52 p value: NR</p> <p>Group 1: 21/52 Group 2: 28/52 p value: NR</p> <p>Group 1: 2/52 Group 2: 2/52 p value: NR</p> <p>Group 1: 3/52 Group 2: 6/52 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Kaplan et al., 1998 ¹³³ Study design: RCT Examiner masked Evidence level: 1+ Duration of follow-up: 12 months	Patient group: men with moderate to severe LUTS Setting: single-centre, department of urology, Columbia University, New York, USA Inclusion criteria: <ul style="list-style-type: none"> AUA symptom score ≥ 10 Qmax ≤ 15 mL/s Prostate volume 15-60g (TRUS) Exclusion criteria: <ul style="list-style-type: none"> < 50 years old Neurogenic bladder Previous prostatic or urethral surgery On medications known to affect voiding function Prostate or bladder cancer All patients N: 64 Drop outs: 3 at 1 year Group 1: N: 32 Mean age (\pm SD): 68.9 \pm 8.7 Mean AUA \pm SD: 19.4 \pm 3.5 Mean Qmax ml/s \pm SD: 7.2 \pm 2.8 Mean PVR ml \pm SD: 77.8 \pm 20.3 Mean prostate volume \pm SD: 47.8 \pm 22.3 Operative time \pm SD: 47.6 \pm 17.6 mins Drop outs: 2 Group 2:	Group 1: Transurethral vaporisation of the prostate (TUVP) Fluted roller-ball electrode at 240-270W for cutting Group 2: Transurethral resection of the prostate (TURP) Standard loop All patients: Operations performed using 27F continuous flow resectoscope. Examination methods Preoperative: Baseline AUA symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry (Dantec Urodyn). Follow up visits at 1, 3, 6 and 12 months postoperatively	Mean AUA score \pm SD at 3 months	Group 1: 9.2 \pm 2.7 (n=32) Group 2: 8.6 \pm 2.5 (n=32) p value: Not sig.	Funding: Partial funding: Grant RR-0045 from National Institutes of Health Limitations: <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masked outcome assessment was not reported Additional outcomes: PVR at follow up Notes: Statistical analysis was performed by third party who was masked to treatment allocation
			Mean AUA score \pm SD at 6 months	Group 1: 7.4 \pm 2.9 (n=32) Group 2: 7.9 \pm 3.1 (n=32) p value: Not sig.	
			Mean AUA score \pm SD at 12 months	Group 1: 6.6 \pm 2.4 (n=30) Group 2: 6.1 \pm 1.9 (n=31) p value: Not sig.	
			Mean Qmax mL/s \pm SD at 3 months	Group 1: 14.8 \pm 3.9 (n=32) Group 2: 16.8 \pm 3.6 (n=32) p value: 0.03 (NCGC calculate as t-test with equal variance)..	
			Mean Qmax mL/s \pm SD at 6 months	Group 1: 15.6 \pm 3.2 (n=32) Group 2: 18.1 \pm 4.2 (n=32) p value: 0.01 (NCGC calculate as t-test with equal variance)..	
			Mean Qmax mL/s \pm SD at 12 months	Group 1: 16.9 \pm 4.1 (n=30) Group 2: 19.6 \pm 4.9 (n=31) p value: 0.02 (NCGC calculate as t-test with equal variance).	
			Catheterisation time (days) hours reported converted to days	Group 1: 0.54 \pm 0.19 Group 2: 2.81 \pm 0.57 p value: <0.01	
			Length of hospital stay (days)	Group 1: 1.3 \pm 0.5 Group 2: 2.6 \pm 0.9 p value: <0.03	
			Complications: transfusion	Group 1: 0/32 Group 2: 1/32 p value: NR	
			Complications: UTI	Group 1: 5/32 Group 2: 4/32 p value: NR	
Complications: TUR	Group 1: 0/32 Group 2: 1/32 p value: NR				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 32 Mean age (\pm SD): 72.8 \pm 6.9 Mean AUA \pm SD: 18.3 \pm 4.7 Mean Qmax ml/s \pm SD: 8.3 \pm 3.6 Mean PVR ml \pm SD: 66.9 \pm 15.7 Mean prostate volume \pm SD: 41.5 \pm 19.7 Operative time \pm SD: 34.6 \pm 11.2 mins Drop outs: 1</p>		<p>Complications: urethral stricture</p> <p>Complications: incontinence</p> <p>Retrograde ejaculation</p>	<p>Group 1: 1/32 Group 2: 1/32 p value: NR</p> <p>Group 1: 0/32 Group 2: 0/32 p value: NR</p> <p>Group 1: 17/32 Group 2: 13/32 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kupeli et al., 1998¹⁵³ KUPELI A 1998 (forest plot)</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: men with symptomatic BPH</p> <p>Setting: single-centre, department of urology, Ankara Hospital, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> AUA symptom score ≥ 7 Qmax ≤ 15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate volume ≥ 60g (TRUS) < 50 years old Neurogenic bladder Previous prostatic or urethral surgery On medications known to affect voiding function Prostate or bladder cancer <p>All patients N: 66 Drop outs: 6 at 6 months and 10 at 1 year.</p> <p>Group 1: N: 30 Mean age (range): 65.7 (52-72) Mean AUA (range): 13.7 (7-29) Mean Qmax ml/s (range): 8.3 (2.7 -11.8) Mean prostate volume \pm SD: 43.57 \pm 12.01 Operative time \pm SD: 38.61 \pm 7.32 mins Drop outs: 3 at 6 months and 4 at 1 year.</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Storz spike electrode: cutting 180-250W (mean 220W) and coagulation 40-70W (mean 60W)</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop</p> <p>All patients: Operations performed using 24F continuous flow resectoscope with 1.5% glycine as an irrigant</p> <p>Examination methods Preoperative: Baseline AUA symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up visits to collect AUA symptom score and Qmax collected at 6 and 12 months postoperatively</p>	<p>Mean AUA score (range) at 6 months</p>	<p>Group 1: 7.9 \pm NR (0-12) (n=27) Group 2: 7.3 \pm NR (1-12) (n=33) p value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Allocation concealment not reported Masked outcome assessment was not reported Standard deviations were missing from primary outcome measures (AUA symptom score and Qmax) and p values not reported <p>Notes: Randomisation by flipping a coin</p>
			<p>Mean AUA score (range) at 12 months</p>	<p>Group 1: 6.1 \pm NR (0-11) (n=26) Group 2: 7.0 \pm NR (1-14) (n=30) p value: NR</p>	
			<p>Mean Qmax (range) at 6 months</p>	<p>Group 1: 13.8 \pm NR (8.2-16.4) (n=27) Group 2: 14.3 \pm NR (7.2-17.5) (n=33) p value: NR</p>	
			<p>Mean Qmax (range) at 12 months</p>	<p>Group 1: 17.3 \pm NR (11.5-23.8) (n=26) Group 2: 19.6 \pm NR (9.4-24.5) (n=30) p value: NR</p>	
			<p>Catheterisation time (days)</p>	<p>Group 1: 1.61 \pm 0.8 Group 2: 3.83 \pm 1.39 p value: <0.0001</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 1.92 \pm 0.89 Group 2: 4.16 \pm 1.46 p value: <0.0001</p>	
			<p>Complications: transfusion</p>	<p>Group 1: 0/30 Group 2: 2/36 p value: NR</p>	
			<p>Complications: UTI</p>	<p>Group 1: 4/30 Group 2: 3/36 p value: NR</p>	
			<p>Complications: urinary retention</p>	<p>Group 1: 1/30 Group 2: 0/36 p value: NR</p>	
			<p>Complications: reoperation rate</p>	<p>Group 1: 1/30 Group 2: 0/36 p value: NR</p>	
<p>Complications: urethral stricture</p>	<p>Group 1: 0/30 Group 2: 0/36 p value: NR</p>				
<p>Complications: incontinence</p>	<p>Group 1: 1/30 Group 2: 1/36 p value: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	1 year Group 2: N: 36 Mean age (range): 62.4 (56-70) Mean AUA (range): 14.6 (8-32) Mean Qmax ml/s (range): 8.8 (3.0 -12.4) Mean prostate volume ± SD: 41.46 ± 10.7 Operative time ± SD: 41.40 ± 7.95 mins Drop outs: 3 at 6 months and 6 at 1 year				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kupeli et al., 1998¹⁵⁴ KUPELI B 1998 (forest plot)</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months (mean 4.2 months)</p>	<p>Patient group: men with moderate to severe symptoms of BPH Setting: single-centre, department of urology, Ankara Hospital, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • IPSS symptom score ≥ 8 • Qmax < 15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurogenic bladder • Previous prostatic surgery • Prostate cancer <p>All patients N: 60 Drop outs: 0</p> <p>Group 1: N: 30 Mean age (\pm SD): 62.4 \pm 3.2 Mean IPSS score: 19.4 \pm NR Mean Qmax ml/s (\pm SD): 7.9 \pm 2.1 Mean prostate size (g) \pm SD: 48.9 \pm 8.7 Operative time \pm SD: 47.3 \pm NR mins Drop outs: 0</p> <p>Group 2: N: 30 Mean age (\pm SD): 59.8 \pm 2.6 Mean IPSS score: 21.6 \pm NR Mean Qmax ml/s (\pm SD): 9.2 \pm 2.6 Mean prostate size (g) \pm SD: 51.7 \pm 9.1 Operative time \pm SD: 41.6 \pm NR mins Drop outs: 0</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Storz spike electrode: cutting mean 250-300W</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop (80-120W)</p> <p>All patients: Operations performed using 24F continuous flow resectoscope</p> <p>Examination methods Preoperative: Baseline AUA symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up visits to collect AUA symptom score and Qmax collected at 6 and 12 months postoperatively</p>	<p>Mean IPSS score at 3 months</p>	<p>Group 1: 4.1 \pm 22.25* Group 2: 5.2 \pm 23.85* p value: Not sig.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method and allocation concealment not reported • Masked outcome assessment was not reported • Standard deviations were missing from primary outcome measure IPSS symptom score • Dropouts were not mentioned. Assume all patients completed study at 3 months <p>Notes: *SD for change from baseline estimated using Cochrane methods with $p \approx 0.01$</p>
			<p>Mean Qmax (\pm SD) at 3 months</p>	<p>Group 1: 17.7 \pm 4.1 Group 2: 19.7 \pm 3.2 p value: 0.05 (NCGC calculated using <i>t</i> test with equal variances)</p>	
			<p>Catheterisation time (days) hours reported converted to days</p>	<p>Group 1: 2 \pm NR Group 2: 4 \pm NR p value: <0.05</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 2.5 \pm NR Group 2: 4.5 \pm NR p value: <0.05</p>	
			<p>Complications: transfusion</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NR</p>	
			<p>Complications: TUR</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NR</p>	
			<p>Complications: UTI</p>	<p>Group 1: 4/30 Group 2: 3/36 p value: NR</p>	
			<p>Complications: urinary retention</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NR</p>	
			<p>Complications: urethral stricture</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NR</p>	
			<p>Complications: retrograde ejaculation</p>	<p>Group 1: 23/30 Group 2: 13/30 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nathan & Wickham 1996²¹¹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: men requiring TURP</p> <p>Setting: single-centre, department of minimally invasive therapy, Guy's Hospital, UK</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Patients with indwelling catheters • Patients on anticoagulant therapy • Neurogenic bladder • Previous prostatic surgery <p>All patients N: 40 Drop outs: NR</p> <p>Group 1: N: 20 Mean age (range): 65.4 (57-77) Mean IPSS score: 21.9 ± 4.2 Mean IPSS QoL ± SD: 4.9 ± 0.7 Mean Qmax ml/s (± SD): 10.2 ± 4.4 PVR mL (range): 130 (0-300) Mean prostate size (g) ± SD: 53.5 ± 28 Operative time ± SD: 39.2 ± NR mins Drop outs: 0</p> <p>Group 2: N: 30 Mean age (range): 69.2 (57-81)</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) VaporTrode® electrode: cutting 200W and 40W</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop: cutting 120W and coagulation 60W</p> <p>All patients: Operations performed using 24Ch continuous flow resectoscope. A 3-way catheter was inserted.</p> <p>Examination methods Preoperative: Baseline IPSS symptom score and IPSS QoL, , TRUS, uroflowmetry. Follow up visits at 4, 8, 12 weeks for IPSS and uroflowmetry</p>	<p>Mean IPSS score at 3 months (follow up interval not clear)</p>	<p>Group 1: 2.86 ± 2.8 Group 2: 3.1 ± 2.3 p value: NR.</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method and allocation concealment not reported • Masked outcome assessment was not reported • Follow up interval for postoperative measurements not clear • There were significant baseline differences in IPSS score and Qmax. • Dropouts were not mentioned. Assume all patients completed study at 3 months <p>Notes: None.</p>
			<p>Mean IPSS QoL score at 3 months (follow up interval not clear)</p>	<p>Group 1: 0.5 ± 7 Group 2: 0.9 ± 0.9 p value: NR</p>	
			<p>Mean Qmax ± SD mL/s at 3 months (follow up interval not clear)</p>	<p>Group 1: 21.3 ± 5.9 Group 2: 20.6 ± 2.6 p value: NR</p>	
			<p>Catheterisation time (days) hours reported converted to days</p>	<p>Group 1: 0.58 Group 2: 1.9 p value: NR</p>	
			<p>Length of hospital stay (days)</p>	<p>Group 1: 1.85 Group 2: 3.45 p value: <0.0001</p>	
			<p>Complications: transfusion</p>	<p>Group 1: 0/20 Group 2: 2/20 p value: NR</p>	
			<p>Complications: UTI at 3 months</p>	<p>Group 1: 0/20 Group 2: 0/20 p value: NR</p>	
			<p>Complications: TUR</p>	<p>Group 1: 0/20 Group 2: 0/20 p value: NR</p>	
			<p>Complications: incontinence (urgency & frequency) at 3 months</p>	<p>Group 1: 0/30 Group 2: 0/30 p value: NR</p>	
			<p>Complications: reoperation rate</p>	<p>Group 1: 1/20 Group 2: 3/20 p value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean IPSS score: 17.0 ± 4.3 Mean IPSS QoL \pm SD: 4.9 ± 0.7 Mean Qmax ml/s (\pm SD): 7.2 ± 3.5 PVR mL (range): 120 (0-380) Mean prostate size (g) \pm SD: 53.4 ± 21 Operative time \pm SD: $37.4 \pm$ NR mins Drop outs: 0</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nuhoglu et al., 2005²²⁷</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 5 years</p>	<p>Patient group: men with LUTS association with BPH</p> <p>Setting: single-centre, Ankara, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS >15 Qmax < 10 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Suspected prostate cancer Neurogenic bladder Previous prostatic or urethral surgery <p>All patients N: 77</p> <p>Drop outs: 33 at 5 years (5 died, 5 dropped out and 19 could not be contacted. 4 patients are unaccounted for in the study report)</p> <p>Group 1: N: 37 Mean age (\pm SD): 64.5 \pm 8.7 Mean IPSS score: 17.3 \pm 6.8 Mean Qmax ml/s (\pm SD): 6.3 \pm 2.1 PVR mL (range): 88 \pm 20 Mean prostate volume mL \pm SD: 39 \pm 8.1 Operative time \pm SD: 45 \pm 13.2 mins Drop outs: 16 at 5 years. Mean follow up time yrs: 5.7 \pm 0.6</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Storz spike loop: cutting 250W and 100W coagulation</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop:</p> <p>All patients: Operations performed using 24F continuous flow resectoscope using glycine as irrigant. A 3-way catheter was inserted. Antibiotic prophylaxis applied to surgeon's discretion</p> <p>Examination methods Preoperative: Baseline DRE, IPSS symptom score, urinalysis, PSA, TRUS, uroflowmetry. Follow up visits at 1 & 3 months and >5 years thereafter</p>	Mean IPSS score \pm SD at 3 months	Group 1: 4.7 \pm 3.1 (n=35) Group 2: 4.8 \pm 4.2 (n=38) P value: Not sig.	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masked outcome assessment was not reported Dropouts were not reported completely <p>Additional outcomes: PVR and average flow at 3 months and \geq 5 years. Serum electrolytes</p> <p>Notes: None.</p>
			Mean IPSS score \pm SD at \geq 5 years	Group 1: 6.5 \pm 3.2 (n=21) Group 2: 6.1 \pm 3.5 (n=23) P value: Not sig.	
			Mean Qmax \pm SD mL/s at 3 months	Group 1: 17.7 \pm 2.3 Group 2: 17.5 \pm 3.3 P value: Not sig.	
			Mean Qmax \pm SD mL/s at \geq 5 years	Group 1: 12.9 \pm 3.1 Group 2: 13.8 \pm 2.9 P value: Not sig.	
			Catheterisation time (days) hours reported converted to days	Group 1: 0.92 \pm 0.24 Group 2: 3.15 \pm 0.52 p value: <0.001	
			Complications: transfusion	Group 1: 0/37 Group 2: 2/40 p value: NR	
			Complications: urinary retention	Group 1: 1/37 Group 2: 0/40 p value: NR	
			Complications: retrograde ejaculation	Group 1: 5/37 Group 2: 4/40 p value: NR	
			Complications: reoperation rate	Group 1: 1/37 Group 2: 0/40 p value: NR	
			Complications: urethral stricture	Group 1: 1/37 Group 2: 0/40 p value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2: N: 40 Mean age (\pm SD): 65.1 \pm 9.4 Mean IPSS score: 17.6 \pm 7.2 Mean Qmax ml/s (\pm SD): 5.9 \pm 2.6 PVR mL (range): 95 \pm 26 Mean prostate volume mL \pm SD: 39 \pm 7.7 Operative time \pm SD: 42 \pm 9.5 mins Drop outs: 17 at 5 years Mean follow up time yrs: 5.7 \pm 0.9</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Patel et al, 1997²³⁵</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: men with symptomatic BOO</p> <p>Setting: single-centre, department of urology, UCLA, USA</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS moderate or severe (n=6) Qmax < 15 mL/s Acute urinary retention (n=6) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> UTI Neurogenic bladder <p>All patients N: 12</p> <p>Drop outs:</p> <p>Group 1: N: 6 Mean age (range): 67 (60-85) Mean IPSS score (range): 29.6 (28-31)* Mean Qmax ml/s (range): 10 (7.3-13.1) Mean prostate volume mL (range): 54 (25-90) TRUS Operative time (range): 64.3 (40-120) mins Median energy used: 1657.5 (1286-2010) kJ Drop outs: NR</p> <p>Group 2: N: 6 Mean age (range): 65.8 (59-71) Mean IPSS score (range): 23.3</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) VaporTrode® grooved bar electrode (Circon ACMI) cutting 130-190W and 40W coagulation</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop resection. cutting 120-170W and 40W coagulation</p> <p>All patients: Operations performed using 25F continuous flow resectoscope using water as irrigant.</p> <p>Examination methods Preoperative: Baseline IPSS symptom score, urinalysis, TRUS, uroflowmetry. Follow up visits at 3 months</p>	<p>Mean IPSS score (range) at 3 months*</p> <p>Mean Qmax (range) mL/s at 3 months</p> <p>Catheterisation time (days)</p> <p>Length of hospital stay (days)</p>	<p>Group 1: 3.5 (2-4) Group 2: 3.2 (1-5) P value: NR</p> <p>Group 1: 21.4 (17.2-25.3) Group 2: 22.6 (19.3-25.2) P value: NR</p> <p>Group 1: 2 (1-3) Group 2: 2.6 (1-5) p value: NR</p> <p>Group 1: 1.8 (1-2) Group 2: 2.6 (2-4) p value: NR</p>	<p>Funding: Equipment loaned from Circon ACMI</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masked outcome assessment was not reported Dropouts were not reported Small sample size pilot study Adverse events poorly reported <p>Additional outcomes: PVR and average flow at 3 months and ≥ 5 years. Serum electrolytes</p> <p>Notes: Randomised after stratification for prostate volume (TRUS) *IPSS score for patients without retention for baseline but unclear whether IPSS postoperative results were for all patients</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>(17-29)* Mean Qmax ml/s (range): 7.5 (5.1-11) Mean prostate volume mL (range): 64.6 (31.5-119) TRUS Operative time (range): 66 (27-95) mins Median energy used: 753 (555-977) kJ Drop outs: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Shokeir et al., 1997 ²⁷⁸	Patient group: men symptomatic LUTS	Group 1: Transurethral vaporisation of the prostate (TUVP) Storz grooved roller electrode: cutting mean 240W (200-300) and mean 70W (50-80W) coagulation	Mean AUA-7 score ± SD at 3 months	Group 1: 4.5 ± 1.9 Group 2: 4.8 ± 2.2 P value: Not sig.	Funding: NR Limitations: <ul style="list-style-type: none"> • Randomisation method and allocation concealment not reported • Masked outcome assessment was not reported • Dropouts were not reported Additional outcomes: PVR at each follow up and serum electrolytes Notes: None.
Study design: RCT	Setting: multi-centre, department of urology, New Jeddah and King Hafid Madina Hospitals, Saudi Arabia		Mean AUA-7 score ± SD at 6 months	Group 1: 4.6 ± 1.2 Group 2: 4.5 ± 1.3 P value: Not sig.	
Evidence level: 1+	Inclusion criteria: <ul style="list-style-type: none"> • AUA-7 Symptom score >15 • Qmax < 12 mL/s • Prostate size < 60g measured by TRUS 	Group 2: Transurethral resection of the prostate (TURP) Standard loop:	Mean AUA-7 score ± SD at 12 months	Group 1: 5.2 ± 1.4 Group 2: 4.7 ± 1.5 P value: Not sig.	
Duration of follow-up: 12 months	Exclusion criteria: <ul style="list-style-type: none"> • Neurogenic bladder • Prostate cancer • Bladder stone • Previous prostatic surgery • Prostate size > 60g measured by TRUS • Patients with acute urinary retention • Patients with indwelling catheter 	All patients: Operations performed using 26F continuous flow resectoscope using glycine as irrigant. A 3-way catheter was inserted.	Mean Qmax ± SD mL/s at 3 months	Group 1: 19.4 ± 2.2 Group 2: 19.4 ± 2.1 P value: Not sig.	
Mean 14.4 months (12-17)		Examination methods Preoperative: Baseline serum electrolytes, AUA-7 symptom score, urinalysis, PSA, TRUS, uroflowmetry (Qmax from 3 voids >150mL, Urodyn Dantec). Follow up visits at 1, 3, 6 and 12 months	Mean Qmax ± SD mL/s at 6 months	Group 1: 19.2 ± 2.0 Group 2: 19.3 ± 2.0 P value: Not sig.	
			Mean Qmax ± SD mL/s at 12 months	Group 1: 20.1 ± 3.2 Group 2: 18.2 ± 3.0 P value: Not sig.	
			Catheterisation time (days)	Group 1: 1.1 ± 0.4 Group 2: 2.0 ± 0.8 p value: <0.001	
			Length of hospital stay (days)	Group 1: 1.5 ± 0.7 Group 2: 2.5 ± 1.0 p value: <0.001	
			Complications: transfusion	Group 1: 0/35 Group 2: 0/35 p value: NR	
			Complications: TUR	Group 1: 0/35 Group 2: 0/35 p value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>mins Mean follow up time mths: 14.3 ± 2.1 Drop outs: NR</p> <p>Group 2: N: 35 Mean age (± SD): 68.4 ± 9.6 Mean AUA-7 score: 25.1 ± 5.5 Mean Qmax ml/s (± SD): 6.9 ± 1.7 PVR mL (range): 77.1 ± 20.3 Mean prostate volume mL ± SD: 39 ± 7.7 Operative time ± SD: 39.7 ± 8.8 mins Mean follow up time mths: 14.5 ± 1.8 Drop outs: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Van Melick et al., 2003³⁰⁹ Links with Van Melick et al., 2002³⁰⁷ (up to 6 months) and Van Melick et al., 2003³⁰⁸ (up to 12 months)</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 7 years</p>	<p>Patient group: men over 45 years with LUTS associated with BPH that were recruited from their clinic from 1996 to 2001</p> <p>Setting: single-centre, University Medical Centre Utrecht, Netherlands</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> met ISC criteria for BPH Schafer obstruction score ≥ 2 prostate size between 20-65ml. <p>Exclusion Criteria: age ≤ 45 yrs</p> <p>All patients N: 96</p> <p>Group 1 N: 46</p> <p>Age (mean) \pm SD: 64 \pm 10 IPSS (mean) \pm SD: 20.2 \pm 6.6 Mean prostate size, ml: 35 \pm 11 Mean (SD) Global quality of life score: 4.1 \pm 1.4 Mean Qmax \pm SD ml/s: 11 \pm 4 Follow-up 1 to 4 years = 12 Follow-up 4 to 7 years = 12</p> <p>Drop outs: 12 at one year post-operatively (procedure during surgery changed for medical reasons=2, surgery cancelled=1, equipment failure resulting in TURP)=1, surgery incorrectly performed=4, morbidity=1, reoperation –TURP=2, reoperation – due to stricture =1)</p>	<p>Group 1: Laser vaporisation VaporTrode® (Circon ACMI) power settings were not reported</p> <p>Group 2: TURP Standard resection. Suprapubic catheter if required perioperatively.</p> <p>All patients: Standard 24FR resectoscope using glycine for irrigation. Pre-procedural antibiotics and transurethral 20F catheter postoperatively.</p> <p>Examination methods: Urodynamic studies (cystometry and pressure flow) at baseline and 1-6 weeks, 3, 6, 12 months after treatment</p>	<p>Mean (\pm SD) symptom score (IPSS) at 6 months</p>	<p>Group 1: 7.2 \pm 6.7 (n=33) Group 2: 5.3 \pm 5.1 (n=37)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method was not described and masking of outcome assessment was not reported. Significant baseline difference in IPSS score Not all patients were evaluated with urodynamics during the follow up period Numbers of patients completing IPSS score not clear at 6 & 12 mths <p>Additional outcomes: Frequency during day, frequency during night, symptom problem index and BPH impact index. Uroflowmetry also reported.</p> <p>Notes: Follow up time varied individually as all patients were analysed within a 2 month period. Depending on the individual follow-up time,</p>
			<p>Mean (\pm SD) symptom score (IPSS) at 12 months</p>	<p>Group 1: 6.7 \pm 6.4 (n=34) Group 2: 4.6 \pm 4.8 (n=41)</p>	
			<p>Mean (\pm SD) symptom score (IPSS) at 1-4 years*</p>	<p>Group 1: 8.4 \pm 8.7 (n=12) Group 2: 5.8 \pm 7.5 (n=15)</p>	
			<p>Mean (\pm SD) symptom score (IPSS) at 4-7 years*</p>	<p>Group 1: 7.0 \pm 5.6 (n=12) Group 2: 7.3 \pm 7.1 (n=15)</p>	
			<p>Mean (SD) Global quality of life score at 6 months</p>	<p>Group 1: 1.6 \pm 1.6 Group 2: 0.9 \pm 1.2</p>	
			<p>Mean (SD) Global quality of life score at 12 months</p>	<p>Group 1: 1.4 \pm 1.4 Group 2: 0.9 \pm 1.2</p>	
			<p>Mean (SD) Global quality of life score at 1-4 years*</p>	<p>Group 1: 1.0 \pm 1.2 Group 2: 1.1 \pm 1.2</p>	
			<p>Mean (SD) Global quality of life score at 4-7 years*</p>	<p>Group 1: 1.4 \pm 0.8 Group 2: 1.3 \pm 1.3</p>	
			<p>Qmax mean \pm SD at 3 months</p>	<p>Group 1: 20 \pm 10 (n=19) Group 2: 25 \pm 11 (n=15)</p>	
			<p>Qmax mean \pm SD at 6 months</p>	<p>Group 1: 23 \pm 10 (n=33) Group 2: 24 \pm 7 (n=37)</p>	
			<p>Qmax mean \pm SD at 12 months</p>	<p>Group 1: 28 \pm 6 (n=34) Group 2: 23 \pm 10 (n=41)</p>	
			<p>Qmax mean \pm SD at 1-4* years</p>	<p>Group 1: 23 \pm 6 Group 2: 20 \pm 5</p>	
			<p>Qmax mean \pm SD at 4-7* years</p>	<p>Group 1: 16 \pm 11 Group 2: 17 \pm 8</p>	
			<p>Catheterisation time (days)</p>	<p>Group 1: 1.9 \pm 0.6 Group 2: 2.1 \pm 0.7 p value: NR</p>	
<p>Length of hospital stay (days)</p>	<p>Group 1: 3.4 \pm 0.9 Group 2: 3.9 \pm 0.9 p value: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 N: 50 Age (mean) ± SD: 66 ± 8 IPSS (mean) ± SD: 16.8 ± 6.0 Mean prostate size, ml ± SD: 37 ± 11 Mean ± SD Global quality of life score: 3.8 ± 1.5 Mean Qmax ± SD ml/s: 11 ± 4 Follow-up 1 to 4 years = 10 Follow-up 4 to 7 years = 17 Drop outs: 9 at one year post-operatively (surgery cancelled=1, mortality=2, morbidity=2, emigrated=1, reoperation (TURP) =2, reoperation (stricture)=1)</p>		<p>Post-op complications: urethral stricture (within 12 mths)</p> <p>Post-op complications: mortality (within 12 mths)</p> <p>Post-op complications: transfusion required (within 12 mths)</p> <p>Post-op complications: urinary retention (within 12 mths)</p> <p>Reoperation rate (TURP) within 12 mths</p>	<p>Group 1: 1/46 Group 2: 2/50</p> <p>Group 1: 0/46 Group 2: 2/50</p> <p>Group 1: 0/46 Group 2: 1/50</p> <p>Group 1: 0/46 Group 2: 0/50</p> <p>Group 1: 2/46 Group 2: 2/50</p>	<p>patient divided into two groups: those with a follow-up time between 1 and 4 years and those with follow up time between 4 and 7 years. * follow up = 2.8 yrs for TUVP 1-4 yrs and 5.4 yrs for category 4-7 years. For TURP mean follow up = 2.7 yrs for category 1-4 yrs and 5.7 yrs for category 4-7 yrs.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Wang et al., 2002 ³¹⁵ Study design: RCT Evidence level: 1+ Duration of follow-up: 24 months	<p>Patient group: NR Setting: China Inclusion criteria: NR Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or suspect Neurogenic bladder Urethral stricture <p>All patients N: 206 Drop outs:</p> <p>Group 1: N: 97 Mean age (range): 72 (62-85) Mean IPSS score (range): 20 (8-30) Mean Qmax ml/s (range): 7 (2-13) Mean PVR ml (range): 120 (60-400) Mean prostate volume mL (range): NR Operation time (range) mins: 35 (25–70) Drop outs: 1 (death due to cardiovascular event)</p> <p>Group 2: N: 109 Mean age (range): 71 (61-84) Mean IPSS score (range): 20 (9-31) Mean Qmax ml/s (range): 7 (3-12) Mean PVR ml (range): 131 (60–380) Operation time (range) mins: 35 (25–70) Mean prostate volume mL (range): NR Drop outs: NR</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Electrode not specified. Power 240-260W</p> <p>Group 2: Transurethral resection of the prostate (TURP) Power 100-140W</p> <p>Examination methods Preoperative: Not reported in HTA report</p>	<p>Mean IPSS score (range) at 12 months</p> <p>Mean IPSS score (range) at 24 months</p> <p>Complications: TUR syndrome</p> <p>Complications: mortality</p> <p>Complications: incontinence</p> <p>Complications: strictures</p>	<p>Group 1: 4 (4–20) n=109 Group 2: 3 (1–17) n=96 P value: NR</p> <p>Group 1: 5 (4-23) n=38 Group 2: 4 (2-21) n=43 P value: Not sig.</p> <p>Group 1: 3/97 Group 2: 5/109</p> <p>Group 1: 1/97 Group 2: 0/109</p> <p>Group 1: 5/97 Group 2: 1/109</p> <p>Group 1: 5/97 Group 2: 2/109</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masked outcome assessment was not reported Unable to obtain copy of reference to check figures <p>Notes: Data taken from HTA report.</p>

Evidence Table 37: Bipolar transurethral vaporisation of the prostate (TUVP) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Dunsmuir et al., 2003⁶⁷⁶</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months (mean 9 months)</p>	<p>Patient group: men with LUTS secondary to BPH being considered for surgery</p> <p>Setting: single-centre: Department of Urology, Monash Medical Centre, Melbourne, Australia.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> <80 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Acute urinary retention Anticoagulant therapy Prostate volume >80mL Prostate cancer or suspect Previous prostate surgery <p>All patients N: 51 Drop outs: 0</p> <p>Group 1: N: 30 Mean age \pm SD: 63 \pm 7.1 Mean AUA \pm SD: 24.0 \pm 6.9 Mean Qmax \pm SD, mL/s: 9.6 \pm 3.0 Mean PVR \pm SD, mL: 112 \pm 13.3 Mean prostate volume \pm SD, mL: 36 \pm 19 QoL \pm SD: 12 \pm 3.4 Operative time \pm SD, min: 33 \pm NR Drop outs: 0</p> <p>Group 2: N: 35 Mean age \pm SD: 60 \pm 6.5 Mean AUA \pm SD: 17.0 \pm 6.2 Mean Qmax \pm SD, mL/s: 10.4 \pm 3.1 Mean PVR \pm SD, mL: 96 \pm 11.4 Mean prostate volume \pm SD, mL: 42 \pm 21 QoL \pm SD: 11 \pm 3.2</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop</p> <p>All patients:</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, QoL, Qmax, PVR assessed and follow up of IPSS, QoL, PVR and Qmax at 3, 6 12 months</p>	<p>Mean \pm SD IPSS at 3 months</p> <p>Mean \pm SD IPSS at 6 months</p> <p>Mean \pm SD IPSS at 12 months</p> <p>Mean \pm SD Qmax at 3 months</p> <p>Mean \pm SD Qmax at 6 months</p> <p>Mean \pm SD Qmax at 12 months</p> <p>Catheterisation time (days) converted into days</p> <p>Length of stay (days) reported as time to discharge</p> <p>Complications: urinary retention (re-catheterisation)</p>	<p>Group 1: 5.7 \pm NR (n=30) Group 2: 8.2 \pm NR (n=21) P value: NR</p> <p>Group 1: 7.1 \pm NR (n=24) Group 2: 5.7 \pm NR (n=20) P value: NR</p> <p>Group 1: 5.0 \pm NR (n=20) Group 2: 6.4 \pm NR (n=20) P value: NR</p> <p>Group 1: 18.0 \pm NR (n=30) Group 2: 20.0 \pm NR (n=21) P value: NR</p> <p>Group 1: 18.5 \pm NR (n=24) Group 2: 17.0 \pm NR (n=20) P value: NR</p> <p>Group 1: 17.0 \pm NR (n=20) Group 2: 15.0 \pm NR (n=20) P value: NR</p> <p>Group 1: 0.8 \pm NR Group 2: 0.7 \pm NR P value: 0.92</p> <p>Group 1: 1.45 \pm NR Group 2: 1.55 \pm NR P value: 0.88</p> <p>Group 1: 10/30 Group 2: 1/21 P value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Masking of outcome assessment was not reported Mean \pm SD were not reported for IPSS and Qmax. Data were estimated from graph. Intermediate report, not all patients randomised have received surgery or been followed up for 12 mths. <p>Notes: Randomisation by drawing tickets from previously sealed box containing equal numbers of tickets for each type of surgery.</p> <p>QoL score was based on AUA symptom scoring section C with a maximum score of 19</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Operative time \pm SD, min: 26 \pm NR Drop outs: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Hon et al., 2006¹²¹</p> <p>Study design: RCT Observer masked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Mean 9 months</p>	<p>Patient Group: Men with BOO undergoing surgery</p> <p>Setting: single centre: Shrewsbury & Telford Hospital, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> NR <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous myocardial infarction Prostate cancer or suspect Previous history of prostatic surgery Serum creatinine >200 mmol/L Prostate volume > 80 mL Neurogenic bladder Urethral stricture <p>All patients N: 160 Dropouts: NR</p> <p>Group 1 N: 81 Mean age \pm SD: 66.1 \pm 8.5 Mean IPSS \pm SD: 21.3 \pm 6.2 Mean Qmax \pm SD, mL/s: 12.0 \pm 6.4 Mean PVR \pm SD, mL: 147 \pm 156 Mean prostate volume \pm SD, mL: 38.0 \pm 17.5 IPSS QoL \pm SD: 4.2 \pm 1.1 History of urinary retention: 17/81 Catheter in situ: 8/81 9.9% Operative time \pm SD, min: 32.6 \pm 13.4 Drop outs: 0</p> <p>Group 2 N: 79</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system with Plasma V™ bar (320-450kHz) at 160W cutting and 80W coagulation. Isotonic saline as irrigant</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop and irrigation with mannitol/sorbitol.</p> <p>All patients Underwent Otis urethrotomy before prostatectomy and received continuous irrigation with saline.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up for IPSS, QoL, PVR and Qmax</p>	<p>Mean \pm SD IPSS at 9 months</p>	<p>Group 1: 7.7 \pm 6.8 (n=73) Group 2: 6.9 \pm 5.8 (n=76) P value: 0.44</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Reasons for missing data at follow up were not reported Data presented for mean overall follow up <p>Additional outcomes: Irrigation volumes.</p> <p>Notes: Randomisation using sequentially numbered opaque envelopes containing computer generated numbers.</p>
			<p>Mean \pm SD Qmax at 9 months</p>	<p>Group 1: 25.6 \pm 15.6 (n=73) Group 2: 23.5 \pm 15.2 (n=76) P value: 0.41</p>	
			<p>Mean \pm SD QoL at 9 months</p>	<p>Group 1: 1.7 \pm 1.5 (n=73) Group 2: 1.5 \pm 1.5 (n=76) P value: 0.64</p>	
			<p>Length of Stay \pm SD, days reported as mean postoperative stay</p>	<p>Group 1: 3.0 \pm 0.9 (n=81) Group 2: 3.4 \pm 1.1 (n=79) P value: 0.04</p>	
			<p>Complications: Transfusion</p>	<p>Group 1: 0/81 Group 2: 4/79 P value: 0.02</p>	
			<p>Complications: urinary retention (re-hospitalisation)</p>	<p>Group 1: 1/81 Group 2: 2/79 P value: NR</p>	
			<p>Complications: urethral stricture</p>	<p>Group 1: 0/81 Group 2: 1/79 P value: NR</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> Mean age \pm SD: 68.1 \pm 7.5 Mean IPSS \pm SD: 20.6 \pm 7.0 Mean Qmax \pm SD, mL/s: 11.9 \pm 6.0 Mean PVR \pm SD, mL: 182 \pm 180 Mean prostate volume \pm SD, mL: 40.0 \pm 17.1 IPSS QoL \pm SD: 4.3 \pm 1.3 History of urinary retention: 18/79 Catheter in situ: 13/79 16% Operative time \pm SD, min: 28.5 \pm 15.2 Drop outs: 0 </p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Karaman et al., 2005¹³⁷ and Kaya et al., 2007¹³⁹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: men with BOO secondary to BPH</p> <p>Setting: single centre: Department of Urology, Haydarparsa Numune Training & Research Hospital, Istanbul, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Severe LUTS on IPSS score requiring treatment Qmax < 15 mL/s or obstructive pressure flow study Prostatic volume <60 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or suspect after biopsy for DRE or PSA >4 ng/mL Untreated UTI Previous history of prostatic surgery Neurogenic bladder Urethral stricture <p>All patients N: 75 Dropouts: NR</p> <p>Group 1 N: 38 Median Age (range), yrs: 66 (49-80) IPSS ± SD: 21.0 ± 3.8 Mean ± SD Qmax, mL/s: 6.0 ± 2.1 Mean prostate volume ± SD, mL: 50.0 ± 2.0 Operation time ± SD, min: 40.3 ± 15 Dropouts: NR</p> <p>Group 2</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ tissue management system (160Ω, 320-450kHz, 254-350V) using saline irrigant/ 80-100 V coagulation</p> <p>Group 2: TURP Standard loop through 26F continuous flow resectoscope with glycine irrigant.</p> <p>All patients 3-way catheter inserted and irrigation continued until urine was clear. Catheter was before the patient was discharged</p> <p>All operations performed by the same surgeons</p> <p>Examination methods Preoperative: Baseline IPSS, Qmax and PVR, PSA, blood, urinalysis, TRUS</p> <p>Postoperative: IPSS and Qmax repeated at follow up of 3, 6 & 12 mths</p>	Mean ± SD IPSS at 3 months	Group 1: 5.0 ± 3.4 (n=38) Group 2: 9.0 ± 2.9 (n=37) P value: <0.001	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method, allocation concealment and masking of outcome assessment were not reported Dropouts NR. Unclear whether all patients completed follow up <p>Notes: Long term follow up for 2 and 3 years was available for 25 Group 1 patients and 15 group 2 patients reported in Kaya et al., 2007¹³⁹</p>
			Mean ± SD IPSS at 6 months	Group 1: 6.0 ± 2.7 (n=38) Group 2: 10.0 ± 2.6 (n=37) P value: <0.001	
			Mean ± SD IPSS at 12 months	Group 1: 7.0 ± 8.7 (n=38) Group 2: 12.0 ± 2.6 (n=37) P value: <0.001	
			Mean ± SD IPSS at 2 years	Group 1: 7.1 ± 1.5 (n=25) Group 2: 5.2 ± 1.1 (n=15) P value: <0.05	
			Mean ± SD IPSS at 3 years	Group 1: 7.6 ± 1.4 (n=25) Group 2: 5.7 ± 1.2 (n=15) P value: <0.05	
			Mean ± SD Qmax at 3 months	Group 1: 17.0 ± 2.3 (n=38) Group 2: 18.0 ± 2.0 (n=37) P value: NS	
			Mean ± SD Qmax at 6 months	Group 1: 17.0 ± 1.3 (n=38) Group 2: 17.0 ± 3.3 (n=37) P value: NS	
			Mean ± SD Qmax at 12 months	Group 1: 16.0 ± 1.3 (n=38) Group 2: 15.0 ± 0.7 (n=37) P value: NS	
			Mean ± SD Qmax at 2 years	Group 1: 12.5 ± 2.1 (n=25) Group 2: 20.8 ± 2.4 (n=15) P value: <0.05	
			Mean ± SD Qmax at 3 years	Group 1: 14.4 ± 2.6 (n=25) Group 2: 21.8 ± 3.1 (n=15) P value: <0.05	
			Catheterisation time (days) converted into days	Group 1: 1.5 ± 0.4 Group 2: 2.8 ± 1.1 P value: <0.001	
			Length of stay (days)	Group 1: 1.5 ± 0.4	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 37 Median Age (range), yrs: 65 (54-78) IPSS \pm SD: 22.0 \pm 4.6 Mean \pm SD Qmax, mL/s: 6.0 \pm 3.1 Mean prostate volume \pm SD, mL: 51.1 \pm 1.0 Operation time \pm SD, min: 55.0 \pm 11.0 Dropouts: NR</p>		<p>equal to catheterisation time</p> <p>Complications: Transfusion</p> <p>Complications: TUR</p> <p>Complications: urethral stricture</p> <p>Complications: retrograde ejaculation</p> <p>Complications: erectile dysfunction</p>	<p>Group 2: 2.8 \pm 1.1 P value: <0.001</p> <p>Group 1: 0/38 Group 2: 2/37 P value: NR</p> <p>Group 1: 0/38 Group 2: 0/37 P value: NR</p> <p>Group 1: 2/38 Group 2: 2/37 P value: NR</p> <p>Group 1: 31/38 (82%) Group 2: 32/37 (86%) P value: NR</p> <p>Group 1: 13% Group 2: 12% P value: NR</p>	

Evidence Table 38: Transurethral needle ablation (TUNA) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Cimentepe et al., 2003⁵⁴</p> <p>Study design: RCT</p> <p>Setting: May 1999 to 2000, Turkey</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 18 months</p>	<p>Patient group:</p> <ul style="list-style-type: none"> Patients with lower urinary tract symptoms attributable to BPH. <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Lower urinary tract symptoms due to BPH Age > 40 Q_{max} < 15 mL/sec IPSS > 13 Prostate weight 20-70 g No suspicion of prostate malignancy (according to DRE and PSA) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Urethral stricture Bladder neck contracture Previous prostate surgery Bladder stones or tumours Neurogenic bladder Prominent median lobe <p>All patients N: 59 patients enrolled Drop outs: 0</p> <p>Group 1-TUNA N: 26 Dropouts: 0 Age, years, mean (±SD): 60.1 ± 7.3 IPSS, mean (±SD): 22.9 ± 3.8 IPSS-QoL, mean (±SD): 4.8 ± 0.75 Q_{max}, ml/s, mean (±SD): 9.8 ± 3.6 Prostate size, g, mean (±SD): 46.1 ± 11.2</p>	<p>Group 1: TUNA TEAP system (Vidamed Inc.) Radiofrequency (RF)-powered generator that delivers a dual 465-kHz RF signal.</p> <p>The TEAP procedure was performed with the patient in the lithotomy position under spinal or epidural anaesthesia.</p> <p>The number of treatments for each lateral lobe was determined according to the length of the prostatic urethra. The procedure was performed at 1-cm intervals starting 1 cm from the bladder neck to 1 cm proximal to the verumontanum.</p> <p>The RF energy was delivered continuously and slowly increased to achieve a minimum of 50°C on the shields after 4 minutes of treatment. At the same time, it has been shown that the temperature at the tips of the needles is increased to approx. 100°C. This temperature should be maintained for 1.5 minutes to create lesions. Therefore the device tip was kept</p>	<p>IPSS, mean ± SD</p>	<p><u>Baseline:</u> Group 1: 22.9 ± 3.8 Group 2: 24.1 ± 3.8 p value: 0.41</p> <p><u>3 months:</u> Group 1: 9.7 ± 2.8 Group 2: 8.3 ± 2.9 p value: 0.25</p> <p><u>18 months:</u> Group 1: 8.5 ± 3.2 Group 2: 8.6 ± 1.8 p value: 0.90</p>	<p>Funding: Not reported. Authors from Department of Urology Faith University, School of Medicine, Ankara, Turkey.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Method of randomisation, allocation concealment, ITT and sample size calculation was not reported It was unclear how patients were recruited and screened, and how many of those screened were enrolled Unequal number of patients in both arms, 27% more patients in the TURP arm <p>Additional outcomes:</p> <ul style="list-style-type: none"> 1 patient in TUNA group had acute urinary retention requiring recatheterisation,
			<p>IPSS-QOL, mean ± SD</p>	<p><u>Baseline:</u> Group 1: 4.8 ± 0.75 Group 2: 5.2 ± 0.65 p value: 0.11</p> <p><u>3 months:</u> Group 1: 2.1 ± 0.5 Group 2: 1.9 ± 0.5 p value: 0.30</p> <p><u>18 months:</u> Group 1: 1.8 ± 1.3 Group 2: 1.7 ± 0.5 p value: 0.35</p>	
			<p>Q_{max}, mean ± SD (ml/s)</p>	<p><u>Baseline:</u> Group 1: 9.8 ± 3.6 Group 2: 9.2 ± 3.4 p value: 0.66</p> <p><u>3 months:</u> Group 1: 16.7 ± 4.5 Group 2: 23.1 ± 5.3 p value: 0.002</p> <p><u>18 months:</u> Group 1: 17.7 ± 4.2 Group 2: 23.3 ± 4.9 p value: 0.004</p>	
			<p>Complications: Blood transfusion, (2 patients in TEAP and all patients in TURP)</p>	<p>Group 1: 0/26 (7.7%) Group 2: 0/33 (100%) P value: Not stat sig</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>PVR, ml, mean(\pmSD):67.4\pm29.4</p> <p>Group 2-TURP N: 33 Dropouts: 0 Age, years, mean (\pmSD): 63.3 \pm5.9 IPSS, mean (\pmSD): 24.1 \pm3.8 IPSS-QoL, mean (\pmSD): 5.2\pm0.65 Qmax, ml/s, mean(\pmSD):9.2\pm3.4 Prostate size, g, mean(\pmSD):49.1\pm17.7 PVR, ml, mean(\pmSD):76.1\pm50.1</p> <p>(all parameters not stat sig between two groups)</p>	<p>firmly pressed against the prostate, and the RF power was applied for 5.5 minutes for each lesion.</p> <p>Catheter protocol: catheter was left indwelling for 12-24 hours. Discharge: discharged home on the same day.</p> <p>Group 2: TURP Performed under spinal or epidural anaesthesia. Catheter protocol: catheter was left indwelling for 48- 72 hours. Discharge: hospitalised for a minimum of 48 hours.</p> <p>All patients received analgesics and antibiotics</p>	<p>group had transient bleeding-haematuria after operation)</p> <p>Complications: Retrograde ejaculation (all patients were sexually active pre-operatively)</p> <p>Complications: Urethral stricture</p> <p>Complications: Reoperation, 18 months follow-up) n/N (%)</p> <p>Complications: Slight stress incontinence: (definition not provided)</p> <p>Complications: Erectile impairment (deterioration in achieving and maintaining erection)</p> <p>Duration of operation, minutes, mean\pmSD</p>	<p>18 months follow-up Group 1: 0/26 (0) Group 2: 16/33 (48.5) RR: 0.0 (95% CI: 0.0 to 0.25) P value: <0.01</p> <p>18 months follow-up Group 1: 0/26 (0) Group 2: 2/33 (6.0) P value: Not stat sig</p> <p>18 months follow-up Group 1: 2/26 (7) Group 2: 0/33 (0) P value: Not stat sig</p> <p>18 months follow-up Group 1: 0/26 (0) Group 2: 1/33 (0.3) P value: Not stat sig</p> <p>18 months follow-up Group 1: 0/26 (0) Group 2: 4/33 (12) P value: Not stat sig</p> <p>Group 1: 44.3\pm7.8 Group 2: 55.9\pm12.4 P value: 0.06</p>	<p>unclear how many in the TURP group</p> <ul style="list-style-type: none"> Prostate size at <u>18 months:</u> g), mean \pm SD: TEAP: 41.9 \pm 10.9, TURP: 34.3 \pm 10.4, p value: 0.08 Post void residual volume (mL), mean \pm SD <u>3 months:</u> Group 1: 45.3 \pm 16.7 Group 2: 32.4\pm 17.4 p value: 0.07 <u>18 months:</u> Group 1: 46.4 \pm 17.5 Group 2: 30.3 \pm 18.7 p value: 0.03 <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Hill2004¹¹⁶</p> <p>Study design: RCT</p> <p>Setting: 7 medical centres across the US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 5-years</p> <p>Links with: BRUSKEWITZ 1998³⁶ – 1 year study</p> <p>ROEHRBORN 1999B²⁵⁹ – 6 months data</p>	<p>Patient group: Men with LUTS secondary to BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ▪ Men 50 years or older who have LUTS secondary to BPH a minimum of three months in duration. ▪ I-PSS of greater than 13, a PFR of 12 ml per second or less with a minimum voided volume of at least 125 ml and a prostate size of between 20 and 75 gm, as determined by TRUS. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ Active urinary tract infection ▪ urinary retention or PVR greater than 350 cc ▪ abnormal renal function, ▪ PSA greater than 10 ng/ml (If serum PSA between 4 to 10 ng/ml, TRUS guided prostate 	<p>Group 1: TUNA TEAP device consisted of a hand piece similar to a rigid 18 Fr cystoscope with a 0-degree optical lens, light source and irrigation system, an RF generator that operated a frequency of 460 kHz and 2, 18 gauge needle electrodes to deliver RF energy to the prostate. Temperatures at the centre of the lesion reached 90C to 110C with a gradient decreased of 5C to 15C for 2 to 3 mm such that peripheral temperatures attained 50C to 54C.</p>	<p>IPSS, mean ±SEM</p>	<p><u>Baseline</u> Group 1: 24.0 ± 0.8 (n=65) Group 2: 24.1 ± 0.8 (n=55) P value: NR</p> <p><u>1 year follow up</u> Group 1: 11.7 ± 1.0 (n=56) Group 2: 7.8 ± 0.9 (n=44) P value: 0.0049</p> <p><u>2 year follow up</u> Group 1: 15.0 ± 1.3 (n=43) Group 2: 9.5 ± 1.1 (n=35) P value: 0.0028</p> <p><u>3 year follow up</u> Group 1: 15.2 ± 1.3 (n=38) Group 2: 10.1 ± 1.4 (n=31) P value: 0.0079</p> <p><u>4 year follow up</u> Group 1: 13.2 ± 1.5 (n=24) Group 2: 7.6 ± 1.6 (n=21) P value: 0.0137</p> <p><u>5 year follow up</u> Group 1: 10.7 ± 1.4 (n=18) Group 2: 10.8 ± 1.6 (n=22) P value: 0.9813</p>	<p>Funding: Authors report financial interest and/or other relationship with Glaxo, Merek, Medtronic and Celsion. Funding for trial not reported.</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Randomisation well described but concealment of allocation is not described. ▪ Number of withdrawals and drop-outs is described for 1-year follow up but not for the 5-year period. ▪ Sample size calculation was mentioned, but assumptions used were not described ▪ There were

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>biopsies were performed to exclude prostate cancer), biopsy proven prostate cancer</p> <ul style="list-style-type: none"> an enlarged median lobe neurogenic bladder and/or sphincter abnormalities previous non-pharmacological prostate treatment Prostate gland size < 34 or greater than 64 mm in transverse diameter, Current therapy affecting prostate physiology or other medical conditions that would pose an unacceptable patient risk. <p>All patients N: 121 patients Drop outs: 15 lost to follow-up at 1 year</p> <p>Group 1-TUNA N: 65 Age, years, mean (\pmSE): 66 ± 1.0 IPSS, mean (\pmSD): 24 ± 0.8 Dropouts: 6 lost to follow up at 1 year PVR, ml, mean \pmSEM : 91.8 ± 10.0 (n=65)</p> <p>Group 2- TURP N: 56 Age, years, mean (\pmSE): 66 ± 1.0 IPSS, mean \pmSD: 24.1 ± 0.8 Dropouts: 9 lost to follow up at 1 year PVR, ml, mean \pmSEM : 81.9 ± 9.3 (n=56)</p>	<p>Group 2: TURP Each TURP was done at one of the reporting centres. The patient received general or spinal anaesthesia. Resection was performed using standard techniques and a urethral catheter was left indwelling for 24 to 48 hours postoperatively.</p>	<p>Qmax (ml/s), mean\pmSEM</p>	<p><u>Baseline</u> Group 1: 8.8 ± 0.3 (n=65) Group 2: 8.8 ± 0.3 (n=56) P value: NR</p> <p><u>1 year follow up</u> Group 1: 14.6 ± 1.0 (n=53) Group 2: 21.1 ± 1.3 (n=43) P value: <0.0001</p> <p><u>2 year follow up</u> Group 1: 12.5 ± 0.7 (n=40) Group 2: 21.3 ± 1.4 (n=33) P value: 0.0001</p> <p><u>3 year follow up</u> Group 1: 13.0 ± 1.3 (n=33) Group 2: 19.1 ± 2.0 (n=26) P value: 0.0106</p> <p><u>4 year follow up</u> Group 1: 11.7 ± 1.4 (n=18) Group 2: 18.9 ± 2.5 (n=17) P value: 0.0142</p> <p><u>5 year follow up</u> Group 1: 11.4 ± 1.2 (n=13) Group 2: 18.6 ± 2.3 (n=15) P value: 0.0143</p>	<p>discrepancies in the baseline and follow up values of 3 papers reporting the study.</p> <ul style="list-style-type: none"> Quality of life scale – it was unclear how this was calculated in Bruskewitz1998 and Hill2004. The mean score was more the maximum of IPSS-QoL Scale. Only Roehborn1999B reported used of IPSS-QOL. <p>Additional outcomes: Percent improvement over baseline for AUA, QOL, PFR and PVR (table 3)</p> <p>Procedure related mortality: 0 in both arms</p>
			<p>QoL score, mean \pmSEM (Unclear what scales were used)</p>	<p><u>Baseline</u> Group 1: 11.8 ± 0.5 (n=64) Group 2: 12.6 ± 0.5 (n=56) P value: NR</p> <p><u>1 year follow up</u> Group 1: 4.3 ± 0.5 (n=55) Group 2: 3.7 ± 0.7 (n=45) P value: 0.4814</p> <p><u>2 year follow up</u> Group 1: 6.0 ± 0.7 (n=43) Group 2: 3.7 ± 0.7 (n=33) P value: 0.0309</p> <p><u>3 year follow up</u> Group 1: 5.4 ± 0.7 (n=40) Group 2: 4.7 ± 1.0 (n=32) P value: 0.5275</p>	<p>PVR, ml, mean \pmSEM: <u>1 year follow up</u> Group 1: 80.3 ± 11.0 (n=52) Group 2: 47.1 ± 7.0 (n=43) P value: 0.0173</p> <p><u>2 year follow up</u> Group 1: 74.1 ± 12.6 (n=40) Group 2: 34.6 ± 5.6 (n=31)</p> <p><u>3 year follow up</u> Group 1: 78.2 ± 13.7 (n=32) Group 2: 50.7 ± 10.4 (n=26) P value: 0.1285</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p><u>4 year follow up</u> Group 1: 5.2 ± 0.9 (n=22) Group 2: 3.7 ± 1.0 (n=21) P value: 0.2316</p> <p><u>5 year follow up</u> Group 1: 3.8 ± 0.7 (n=18) Group 2: 4.0 ± 0.8 (n=22) P value: 0.719</p>	<p><u>4 year follow up</u> Group 1: 138.2 ± 45.7 (n=19) Group 2: 39.5 ± 13.1 (n=17) P value: 0.0564</p> <p><u>5 year follow up</u> Group 1: 60.4 ± 21.8 (n=13) Group 2: 27.4 ± 7.9 (n=17) P value: 0.128</p>
			<p>QoL- IPSS Scale, mean ±SD (only reported in Roehborn1999B)</p>	<p><u>Baseline</u> Group 1: 4.6±1.1 Group 2: 4.8±1.1</p> <p><u>6 months follow up:</u> TEAP: 2.0 (sd not provided) Group 2: 1.5 P<0.001</p>	<p>Notes: Where there were discrepancies, values from Hill2004 were used.</p>
			<p>Stricture formation/scar tissue</p>	<p><u>Five-year follow up</u> Group 1: 1/65(1.5) Group 2: 4/56(7.1)</p>	
			<p>Retrograde ejaculation:</p>	<p><u>Five-year follow up</u> Group 1: 0/65 Group 2: 23/56 (41.1)</p>	
			<p>Urinary incontinence:</p>	<p><u>Five-year follow up</u> Group 1: 2/65(3.1) Group 2: 12/56 (21.4)</p>	
			<p>Reoperation: (The 9 men in TEAP group received TURP, the TURP patient received TUIP). One additional patient received radical prostatectomy for prostate cancer.</p>	<p><u>Five-year follow up</u> Group 1: 9/65(13.8) Group 2: 1/56(1.8)</p>	
			<p>Erectile dysfunction:</p>	<p><u>Five-year follow up</u> Group 1: 2/65(3.1) Group 2: 12/56(21.4)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hindley2001 ¹¹⁸ Study design: RCT Setting: UK Evidence level: 1+ Duration of follow-up: 2-year Links with MOSTAFID1997 ²⁰⁵	Inclusion criteria: <ul style="list-style-type: none"> Men > 50 years referred to an integrated prostate-assessment unit for cystometry. Urodynamically confirmed bladder outlet obstruction (BOO) due to BPH, defined as Pdet Q_{max} value within the obstructed area of the Abrams Griffith pressure/flow nomogram. Bothersome LUTS, defined as an IPSS≥13 and an IPSS QOLscore ≥3 Written informed consent. Exclusion criteria: <ul style="list-style-type: none"> History of any illness or surgery that might confound the results of the study, and that produce symptoms which might be confused with those produced by BPH, or that pose additional risk to the patient. Confirmed or suspected malignancy of the prostate by DRE or biopsy. PSA level >4 ng/mL unless T1 carcinoma of the prostate excluded by TRUS-guided biopsy. Previous prostatic surgery or thermotherapy Pharmacological treatment of symptomatic BPH within the last 6 months. Confirmed or suspected bladder cancer. Previous rectal surgery other than haemorrhoidectomy. Previous pelvic irradiation. History of cystolithiasis, haematuria or bladder pathology, urethral strictures, bladder neck contracture, active urinary tract infection or prostatitis. Previous history of neurogenic disorder including Parkinson's disease, multiple sclerosis, stroke and diabetic neuropathy. 	Group 1: TUNA A simple disposable 7 F RF needle-electrode was inserted into the lateral lobes of the prostate and, where appropriate, the median lobe of the prostate, using a catheterising endoscope. A standard surgical diathermy generator was used to produce the 10 W of coagulation for 3 min. After treatment, patients were catheterised and allowed home on first-operative day. The catheter was removed and a trial of voiding carried out 7 days after treatment. Group 2: TURP Patients undergoing TURP were operated on by an experienced surgeon according to the normal principles of prostatic resection.	Mortality There were no deaths during the 2-year follow-up.	Funding: NR	
			IPSS, median (interquartile range)	Effect size <u>Baseline</u> Group 1: 20 (15-23) (n=25) Group 2: 22 (18-15) (n=25) <u>6-months:</u> Group 1: 9 (6-23) (n=20) Group 2: 3 (2-6) (n=22) <u>1 year:</u> Group 1: 6 (4-10) (n=19) Group 2: 3 (2-6) (n=19) <u>2 years:</u> Group 1: 8 (5-13) (n=19) Group 2: 3 (1-5) (n=19) P value: NR for all time points	Limitations: <ul style="list-style-type: none"> Small sample size Drop outs accounted for but intention to treat analyses not conducted. Patients (2 in TEAP 1 in TURP) who refused cystometry at 6 months were also excluded
			QoL score, median (inter-quartile range)	Effect size <u>Baseline</u> Group 1: 4 (3-5) (n=25) Group 2: 5 (4-5) (n=25) <u>6-months:</u> Group 1: 2 (1-3) (n=20) Group 2: 1 (0-2) (n=22) <u>1 year:</u> Group 1: 1 (1-3) (n=19) Group 2: 1 (0-2) (n=19) <u>2 years:</u> Group 1: 2 (1-3) (n=19) Group 2: 1 (0-2) (n=19) P value: NR for all time points	Additional outcomes: <ul style="list-style-type: none"> Post void residual volume (mL), mean ±SD: <u>6-months:</u> Group 1: 50 (44) (n=20) Group 2: 87 (74)(n=22) <u>1 year:</u> Group 1: 104 (109) (n=19) Group 2: 21 936) (n=19) <u>2 years:</u> Group 1: 89 (81) (n=19) Group 2: 32 (42) (n=19) P value: NR
Q_{max} (mL/s), mean ±SD	Effect size <u>Baseline</u> Group 1: 8.5 (3.7) (n=25) Group 2: 9.0 (3.6) (n=25) <u>6-months:</u> Group 1: 9.8 (4.0) (n=20) Group 2: 18.4 (7.7) (n=22) <u>1 year:</u> Group 1: 9.7 (5.0) (n=19) Group 2: 22 (10.3) (n=19)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<ul style="list-style-type: none"> ▪ Patients wishing to maintain potential fertility. ▪ PVR >250 mL (measured by ultrasonography) ▪ Compromised renal function with a serum creatinine >180 mg/L or radiological evidence of upper tract dilatation. ▪ Unable to provide at least one voided volume of >150 mL. ▪ Unable to give informed consent. <p>All patients N: 50 Drop outs: 12</p> <p>Group 1-TUNA N: 25 Dropouts: 5 Age, years, mean (range): 66 (56-82) IPSS, mean (IQ range): 20 (15-23) Post void residual volume (mL), mean ±SD: 55 (44) PdetQ_{max}(cmH₂O), mean ±SD: 92 (12)</p> <p>Group 2-TURP N: 25 Dropouts: 3 Age, years, mean (range): 71 (56-88) IPSS, mean (IQ range): 22 (18-25) Post void residual volume (mL): 74 (53) PdetQ_{max}(cmH₂O), mean ±SD: 99 (10)</p>	<p>At the end of the procedure a 22 F three-way urethral catheter was inserted to allow bladder irrigation; after a successful trial of voiding the patient was allowed home.</p> <p>Prophylactic antibiotic cover with 120 mg IV gentamicin was given preoperatively in both groups.</p>	<p>Blood transfusion: (2 units each)</p> <p>Incontinence (all were urge incontinence, with detrusor instability)</p> <p>Urinary retention (post-op) (Failed trial of voiding)</p> <p>Clot retention:</p> <p>Urinary tract infection:</p> <p>Persistent dysuria:</p> <p>Treatment failure: Defined as patient dissatisfaction with treatment or the development of complications from persisting BOO, including evidence of detrusor dysfunction, incomplete bladder emptying, urinary retention, infection or upper tract obstruction.</p>	<p><u>2 years:</u> Group 1: 8.6 (3.5) (n=19) Group 2: 18.1 (7.1) (n=19) P value: NR for all time points</p> <p>Group 1: 0/20 Group 2: 3/22</p> <p>Group 1: 2/20 Group 2: 2/22</p> <p>Group 1: 1/20 Group 2: 0/22</p> <p>Group 1: 0/20 Group 2: 1/22</p> <p>Group 1: 4/20 Group 2: 4/22</p> <p>Group 1: 4/20 Group 2: 0/22</p> <p><u>2-year follow-up:</u> Group 1: 2/25 Group 2: 0/25 One patient was dissatisfied with the outcome at 8 months. Another patient was dissatisfied at 2 years. Both patients were found to have persistent BOO at urodynamic assessment and underwent TURP.</p>	<ul style="list-style-type: none"> ▪ PdetQ_{max}(cmH₂O), mean ±SD <p>6-months: Group 1: 70 (12) (n=20) Group 2: 44 (11) (n=22) P value: NR</p> <p>2 years: Group 1: 71 (36) (n=12) Group 2: 36 (8) (n=9) P value: NR</p> <p>Notes: The methodology stated in MOSTAFID1997²⁰⁵.</p> <p>The PdetQ_{max} was the primary outcomes variable in the study design</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kim et al., 2006¹⁴⁶ (data obtained from HTA report)</p> <p>Study design: RCT</p> <p>Setting: Korea, recruitment from January 1998–December 2002</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient group: Patients with symptomatic BPE</p> <p>Inclusion criteria: NR</p> <p>Exclusion criteria: NR</p> <p>All patients N: 94/110/89/110 204 randomised, from 223 eligible for TEAP vs. TURP 199 randomised from 212 eligible for Laser coagulation vs. TURP 220 randomised out of 235 eligible for TUNA vs. TURP Drop outs: overall drop out not reported</p> <p>Group 1-TEAP N: 94 Dropouts: Unknown Age, years, mean or median (range) : 66.2 (49–88) QoL score, mean: 4.4 Qmax (ml/s), mean or median: 7.2 Residual volume, (ml), mean or median: 126.1 Prostate size, (ml), mean or median: 36.4</p> <p>Group 2- TUNA N: 110 Dropouts: Unknown Age, years, mean or median(range): 66.4 (48–80) IPSS QoL score, mean: 4.3 Qmax (ml/s), mean or median: 7.0 Residual volume, (ml), mean or median: 257 Prostate size, (ml), mean or median: 40.6</p>	<p>Group 1-TEAP Prostajec device (American Medical Systems, Minnetonka, MN, USA)</p> <p>Group 2 - TUNA VidaMed TUNA system (VidaMed Inc.)⁴</p> <p>Group 3 - Laser Coagulation: Other: procedure: Indigo 830e laser optic system (Ethicon Endosurgery)</p> <p>Group 4 - TURP</p>	IPSS, mean:	<u>Baseline</u> TEAP: 19.5 TUNA: 20.8 Coag; 21.1 TURP: 24.0 <u>3 months</u> TEAP: 9.6 TUNA: 10.8 TURP: 10.6 <u>12 months</u> TEAP: 7.5 TUNA: 11.6 TURP: 8.8	<p>Funding: Unknown</p> <p>Limitations:</p> <ul style="list-style-type: none"> Uncertain whether the data reported was mean or median Randomisation allocation, concealment and blinding had been rated as “unclear” Baseline severity of TEAP vs. TURP patient may differ: <ol style="list-style-type: none"> “medium sized” prostates in TEAP vs. large prostate sizes in TURP Mean IPSS at baseline level was numerically higher in TURP compared to TEAP. Uncertain length of follow up for complications <p>Additional outcomes: (values not reported in HTA reported) Duration of operation, Recatheterisation, Retrograde ejaculation, Erectile dysfunction Reoperation, IPSS-QoL, Length of hospital stay Qmax, Residual volume , Prostate size</p>
			Blood transfusion	TEAP: 0/94 TUNA: 0/100 TURP: 19/101 <u>TEAP vs. TURP</u> RR (95% CI): 0.03(0.00 to 0.45) P value: 0.01 <u>TUNA vs. TURP:</u> RR (95% CI): 0.03(0.00 to 0.42) P value: Sig	
			Urinary retention	TEAP: 2/94 TUNA: 4/100 TURP: 4/101 <u>TEAP vs. TURP</u> RR (95% CI): 0.54 (0.10 to 2.87) P value: 0.47 <u>TUNA vs. TURP:</u> RR (95% CI): 1.01 (0.26 to 3.93) P value: Not sig	
			Urinary tract infection	TEAP: 5/94 TUNA: 10/100 TURP: 7/101	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 3 - Laser Coagulation N: 89 Dropouts: Unknown Age, years, mean or median(range): 68.7 (50–89) IPSS QoL score, mean: 4.7 Qmax (ml/s), mean or median: 8.6 Residual volume, (ml), mean or median: 219 Prostate size, (ml), mean or median: 42.7</p> <p>Group 4 -TURP N: 110 Dropouts: Unknown, 9/110? Age, years, mean or median(range): 7.4 (60–87) QoL score, mean: 4.7 Qmax (ml/s), mean or median:11.9 Residual volume, (ml), mean or median: 187 Prostate size, (ml), mean or median: 44.2</p>		<p>Stricture (in the TURP arm, this was recorded as 7 in TEAP vs. TURP and 5 in TUNA vs. TURP- 5 urethral + 2 bladder neck)</p> <p>Retrograde ejaculation</p> <p>Urinary incontinence</p> <p>Reoperation</p>	<p><u>TEAP vs. TURP</u> RR (95% CI): 0.77(0.25 to 2.34) P value: 0.64 <u>TUNA vs. TURP:</u> RR (95% CI): 1.44(0.57 to 3.64) P value: Not sig</p> <p>TEAP: 0/94 TUNA: 0/100 TURP: 7/101</p> <p><u>TEAP vs. TURP</u> RR (95% CI): 0.07(0.00 to 1.24) P value: 0.07 <u>TUNA vs. TURP:</u> RR (95% CI): P value:</p> <p>TEAP: NR TUNA:5/100 TURP: 39/101</p> <p><u>TUNA vs. TURP:</u> RR (95% CI):0.13(0.05 to 0.32) P value: Not sig</p> <p><u>TEAP vs. TURP</u> RR (95% CI): 0.12(0.01 to 2.19) P value: 0.15 <u>TUNA vs. TURP:</u> RR (95% CI): 1.01 (0.26 to 3.93) P value: Not sig</p> <p>TEAP: NR TUNA: 0/100 TURP: 0/101 <u>TUNA vs. TURP:</u> RR (95% CI): P value:</p>	<p>Notes: Evidence Table produced with data from Evidence Table of the HTA report.</p> <p>Values for complications obtained from Figure 11 of HTA report (page 49).</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Duration of operation, minutes, mean (range)	TEAP: NR TUNA: 37(25-60) TURP: 51(20-85)	
			Length of hospitalisation, days, mean (range)	TEAP: NR TUNA: 1.3(1-3) TURP: 6.5(6-8)	

Evidence Table 39: Transurethral incision of the prostate (TUIP) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Dorflinger et al., 1992⁷⁵</p> <p>Study design: RCT</p> <p>Setting: Denmark</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> bladder neck to seminal crest < 2 cm <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostatic cancer previous prostatic or major pelvic surgery; high operative risk or overt neurological or psychiatric disease; patients with urethral stricture; prostate size > 20 g <p>All patients N: 60 Sexually/not sexually active: 44/8 Drop outs:</p> <p>Group 1-TUIP N: 29 Age, years, median: 69 Symptom score, Madsen Iversen (median) : 15 Qmax (ml/s), median:10 Urinary retention:9/29 (31%);</p> <p>Group 2 -TURP N: 31 Age, years, median: 71 Symptom score, Madsen Iversen (median) : 15 Qmax (ml/s), median:8 Urinary retention:5/31 (16%)</p>	<p>Group 1-TUIP 24Fr resectoscope and Collings knife used. An incision to the depth of the surgical capsule was made at the 7 o clock position</p> <p>Catheter protocol: A balloon catheter was inserted into the bladder and left in until urine was clear</p> <p>Group 2-TURP 24Fr resectoscope used and prostatic tissue resected in a standard fashion</p>	<p>Symptom score, Madsen Iversen (range of 1-27) , median.</p> <p>Only included data from “successfully treated patients”</p>	<p><u>At baseline</u> Group 1: 14.5, n=22 Group 2: 16, n=29 p value: Not sig</p> <p><u>At 3 month follow up</u> Group 1: 2.5, n=22 Group 2: 1, n=29 p value: Not sig</p> <p><u>At 12 months follow up</u> Group 1: 2, n=21 Group 2: 2, n=26 p value: Not sig</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported Only median values were reported for most outcomes <p>Additional outcomes:</p> <ul style="list-style-type: none"> Median values for Obstructive and Irritative components of Madsen Iversen score at baseline, 3 months and 6 months follow up. Total voided volume 1/44 patient was made sexually inactive by the operations No bladder neck contracture
			<p>Qmax, ml/s, mean ± SD:</p>	<p><u>At baseline</u> Group 1: 10.0, n=22 Group 2: 8.0, n=29 p value: Not sig</p> <p><u>At 3 month follow up</u> Group 1: 15.2, n=22 Group 2: 18.8, n=29 p value: Not sig</p> <p><u>At 12 months follow up</u> Group 1: 14.5, n=21 Group 2: 20.2, n=26 p value: 0.025 (Mann Whitney signed rank test)</p>	
			<p>Blood transfusion</p>	<p>Group 1: 0/29 Group 2: 4/31 p value: 0.11</p>	
			<p>Retrograde ejaculation (among patients who were sexually active before and after the operations)</p>	<p>Group 1: 1/19 Group 2: 12/24 Relative risk: 0.11 (95% CI: 0.02 to 0.51) p value: 0.002 [RR calculated by NCGC team]</p>	
			<p>Erectile dysfunction</p>	<p>Group 1: 1/19 Group 2: 4/24</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				p value: Not sig	Notes: Appropriate statistical tests were used Preliminary results reported in Dorflinger 1987
			Urethral stricture	Group 1: 0/29 Group 2: 1/31 p value: Not sig	
			Reoperation (data from study abstract)	<u>At 12 months follow up</u> Group 1: 8/29 Group 2: 4/31 P value: Not sig	
			Length of hospitalisation, days, median	Group 1: 3 Group 2: 3 p value: Not sig	
			Length of indwelling catheterisation, min, median	Group 1: 2 Group 2: 2 p value: Not sig	
			Length of operation, min, median	Group 1: 15 Group 2: 30 p value: <0.001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hellstrom1986 ^{11,4} Study design: RCT Setting: Finland Evidence level: 1+ Duration of follow-up: 6 months	Patient group: Patients with symptomatic bladder outlet obstruction Inclusion criteria: <ul style="list-style-type: none"> ▪ Prostate size < 30 g ▪ Symptoms of infravesicle obstruction, including hesitancy, weakened stream, urgency and a feeling of inadequate emptying. All patients N: 24 Group 1-TUIP N: 11 Age (mean): 63 (54-77) Drop outs: Not reported Group 2 -TURP N: 13 Age (mean): 59 (54-63) Drop outs: Not reported	Group 1 TUIP – bladder neck incision using a vertical knife electrode to make a deep diathermy incision from the right ureteral orifice to the verumontanum through the bladder neck and prostatitis tissue. Group 2 TURP Using 26F continuous flow resectoscope by cutting from the bladder neck to the verumontanum and step by step to the prostatic capsule. After both operations a 22F 3-way indwelling catheter was left in position for 3 days and prophylactic sulphatrimethoprim medication used for about 2 weeks.	All cause mortality (myocardial infarction in TURP and colon cancer in TUIP)	Group 1: 1/24 Group 2: 1/25 p value: Not sig	Funding: Not reported Limitations: <ul style="list-style-type: none"> ▪ No symptom scores were collected ▪ Randomisation method reported but concealment method unclear Additional outcomes: None Notes: None
			Mean (SD) Qmax	Group 1: 12.9 (6) Group 2: 16.5 (6)	
			Transfusion	Group 1: 0/11 Group 2: 0/13	
			Acute urinary retention	Group 1: 0/11 Group 2: 0/13	
			UTI	Group 1: 0/11 Group 2: 0/13	
			Stricture	Group 1: 1/11 Group 2: 0/13	
Retrograde ejaculation	Group 1: 0/11 Group 2: 8/12				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Johnson et al., 1998¹²⁹</p> <p>Study design: RCT, open</p> <p>Setting: Sweden. Feb to Sept 1991</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 60 months</p>	<p>Patient group: small to medium BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Admitted from the waiting list for surgical treatment of BPH No previous treatment for BPH Estimated prostate weight at DRE 20-40g, or 20-40mL by TRUS Distance from verumontanum to bladder neck < 4.0cm1 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Bladder stone or cancer Cystitis Clinical prostatic cancer Prominent median lobe of the prostate Adequate follow up difficult for geographical, psychological or social reasons <p>All patients</p> <p>N:</p> <p>Age, years, mean (±SD):</p> <p>Drop outs:</p> <p>Group 1</p> <p>N: 43</p> <p>Drop outs: 2 (reoperated after failing to void post catheter removal)</p> <p>Age, years, mean (range): 70.2 (52-87)</p> <p>Madsen Iversen, mean (95% CI): 15.4 (6-27)</p>	<p>Group 1-TUIP</p> <p>Catheter protocol: overnight</p> <p>Others: Perioperative heparin :13 Antibiotics:17</p> <p>Group 2-TURP</p> <p>Resected in a standard manner from bladder neck to verumontanum out to the prostate capsule</p> <p>Catheter protocol: overnight</p> <p>Others: Perioperative heparin:17 Antibiotics: 14 Resection weight, g, mean (range): 18.8 (8-45)</p> <p>For both groups: Anti provided to those who had indwelling catheter preoperatively, diabetes mellitus or with positive urine culture</p>	<p>All cause mortality (due to cerebrovascular lesion at 8 weeks)</p> <p>Symptom score (Madsen Iversen, total score), mean (95% CI)</p> <p>Qmax, ml/s, mean (95% CI) estimated from graph for follow ups:</p>	<p>Group 1: 0/43 Group 2: 1/42 p value: Not sig</p> <p><u>At baseline</u> Group 1: 15.4 (6-27), n=43 Group 2: 15.8 (5-28), n=42</p> <p><u>At 3 months:</u> Group 1: 3.5(0-21), n=41 Group 2: 3.8(0-16), n=39</p> <p><u>At 6 months:</u> Group 1: 4.3(0-21),n=36 Group 2: 3.5(0-18),n=34</p> <p><u>At 12 months:</u> Group 1: 3.6(0-15),n=31 Group 2: 2.8(0-11),n=32</p> <p><u>At 24 months:</u> Group 1: 4.5(0-14),n=33 Group 2: 4.7(0-17),n=31</p> <p><u>At 60 months:</u> Group 1: 4.5(0-14),n=22 Group 2: 4.7(0-17),n=24 p value: Not sig between groups; Sig compared to baseline</p> <p><u>At baseline</u> Group 1: 9 (7.5-11) ,n=34 Group 2: 8.5 (7.5-9.5), n=36</p> <p><u>At 3 months:</u> Group 1: 20, n=41 Group 2: 15, n=39</p> <p><u>At 60 months:</u> Group 1: 15, n=22 Group 2: 12, n=24 p value: Reported sig difference between groups at 3, 6, 12 and 24 months. Not sig diff between groups at 60 months. All sig better than baseline except at 60 months</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported Patients who were reoperated not included in analysis <p>Additional outcomes:</p> <ul style="list-style-type: none"> Cystoscopy at 24 and 60 months to investigate healing and incision Post void residual volume, blood loss in volume, number of preoperative positive cultures. 3 patients in TURP group was detected with cancer <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Prostate size, ml, mean (range): 26.2(20.0-37.6) Residual volume, ml, mean (range): 139 (0–650) Indwelling catheter: 7/43</p> <p>Group 2 N: 42 Drop outs: 2 (1 lost to follow up at 8 weeks, 1 died) Age, years, mean (\pmSD): 70.8 (56–85) Madsen Iversen, mean (95% CI): 15.8 (5–28) Prostate size, ml, mean (range): 25.4(20.0-39.8) Residual volume ml, mean (range): 109 (0–400) Indwelling catheter: 8/42</p>		Blood transfusion	<p>Group 1: 0/43 Group 2: 1/42 p value: Not sig</p>	
			Urinary retention , 2 cases from TUIP group failed to void after catheter removal. 1 from TURP group had urinary retention 3 weeks post surgery and a bladder neck stricture was incised 3 weeks later	<p>Group 1: 2/43 Group 2: 1/42 p value: Not sig</p>	
			Reoperation rate (repeated when it was impossible to remove the indwelling catheter or symptoms scores deteriorated, combined with a maximum urinary flow rate of \geq 150ml)	<p>Group 1: 10/43 (within 1-38 months) Group 2: 3/42 (within 2-25 months) Relative risk: 3.26 (95% CI: 1.06 to 10.65) p value: 0.04</p>	
			Catheter duration , days, mean (range)	<p>Group 1: 2.8 (1-15) Group 2: 1.4(1-5) P value: Sig</p>	
			Duration of operation , min, mean (range)	<p>Group 1: 15 (5-40) Group 2: 32 (15-60) P value: Sig</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Larsen et al., 1987¹⁵⁸</p> <p>Study design: RCT, open</p> <p>Setting: US, Veteran Affairs</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group:</p> <ul style="list-style-type: none"> Men with symptoms of prostatism due to BPH <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Estimated prostate weight at cystoscopy to be ≤ 20g <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Severe neurologic and or psychiatric disease Previous TURP Urethral stricture Urinary retention Clinical suspicion of cancer of the prostate Previous major intrapelvic surgical procedures <p>All patients N: 40 Drop outs: 3 (2 lost to follow up- 1 had operation cancelled)</p> <p>Group 1 –TURP N: 19 Age, years, median (range): 63(51-73) Estimated prostate weight, g, median(range): 20(10-20) Duration of symptoms, months, median(range): 24(6-240)</p> <p>Group 2 –TURP N: 18 Age, years, median (range): 61(43-74) Estimated prostate weight, g, median(range): 20(15-20)</p>	<p>Group 1- TUIP Performed using Colling's knife at the 6 pm position extending form the internal urethral orifice to the verumontanum down through the prostate and the capsule.</p> <p>A 3-way Foley catheter with continuous irrigation was used for bladder drainage.</p> <p>Group 2 – TURP performed using method described by Blandy JP 1978.</p> <p>All patients received antibiotic prophylaxis</p>	<p>Symptom score (Madsen Iversen, Total score), median (range)</p>	<p><u>Baseline</u> Group 1: 17(9-23), n=19 Group 2: 17(9-23), n=18 <u>At 3-month follow up</u> Group 1: 2(0-19), n=19 Group 2: 2(0-12), n=18 <u>At 12-month follow up</u> Group 1: 2(0-19), n=12 Group 2: 2(0-7), n=11 p value: Not sig between groups; <0.05, compared to baseline values using Mann Whitney signed rank test</p>	<p>Funding: US Veterans Administration and Danish Medical Research Council grant</p> <p>Limitations:</p> <ul style="list-style-type: none"> Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported Relevance of study – published in 1987 <p>Additional outcomes: Voided volume, post void residual volume</p> <p>Notes: None.</p>
			<p>Symptom score (Madsen Iversen, Irritative score), median (range)</p>	<p><u>Baseline</u> Group 1: 13(5-16), n=19 Group 2: 12(4-16)18 <u>At 3-month follow up</u> Group 1: 0(0-15), n=19 Group 2: 1(0-7), n=18 <u>At 12-month follow up</u> Group 1: 0(0-8), n=12 Group 2: 0(0-5), n=11 p value: Not sig between groups; <0.05, compared to baseline values using Mann Whitney signed rank test</p>	
			<p>Symptom score (Madsen Iversen, Obstructive score), median (range)</p>	<p><u>Baseline</u> Group 1: 5(2-8), n=19 Group 2: 5(2-8), n=18 <u>At 3-month follow up</u> Group 1: 1(0-5), n=19 Group 2: 1(0-6), n=18 <u>At 12-month follow up</u> Group 1: 1(0-3), n=12 Group 2: 1(0-6), n=11 p value: <0.05, compared to baseline values using Mann Whitney signed rank test</p>	
			<p>Qmax, ml/s, median (range)</p>	<p><u>Baseline</u> Group 1: 7.4(2.7-27.3), n=15</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Duration of symptoms, months, median(range): 24(0.5-72)			<p>Group 2: 8.6(1.7-15.5), n=16 At 3-month follow up</p> <p>Group 1: 14.4(2.6-34.6), n=15</p> <p>Group 2: 18.5(5.3-45.3), n=16 At 12-month follow up</p> <p>Group 1: 16.3(6.4-34.7), n=11</p> <p>Group 2: 20.6(9.0-41.3), n=11</p> <p>p value: Not sig between groups; <0.05, compared to baseline values using Mann Whitney signed rank test</p>	
			Urinary tract infections (within 1 month of surgery)	<p>Group 1: 2/19</p> <p>Group 2: 3/18</p> <p>P value: Not sig</p>	
			Post operative bleeding (definition not provided)	<p>Group 1: 1/19</p> <p>Group 2: 2/18</p> <p>P value: Not sig</p>	
			Recatheterisation (2 cases due to bleeding and clot retention in TURP, and 1 case due to haematuria on 10 th day for TUIP)	<p>Group 1: 1/19</p> <p>Group 2: 2/18</p> <p>P value: Not sig</p>	
			Retrograde ejaculation (based on number of patients who were potent and had antegrade ejaculation preoperatively)	<p>Group 1: 2/10</p> <p>Group 2: 8/10</p> <p>Relative risk: 0.25 (95% CI: 0.09 to 0.71)</p> <p>p value: 0.02</p> <p>[calculated by NCGC using Fisher's exact test]</p>	
			Catheterisation, hours median (range)	<p>Group 1: 1(1-2)</p> <p>Group 2: 2(2-7)</p> <p>p value: Not sig between groups; <0.01 (Mann Whitney signed rank test)</p>	
			Hospital stay, days, median (range)	<p>Group 1: 2.5(1-4)</p> <p>Group 2: 4.5(3-10)</p> <p>p value: Not sig between groups; <0.01 (Mann Whitney signed rank test)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Li et al., 1987¹⁶⁵</p> <p>Study design: RCT, open</p> <p>Setting: Hong Kong</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 3 months</p>	<p>Patient group: Patient with prostatism presented with acute urinary retention</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Acute urinary retention Ambulatory Diagnosis confirmed with urethroscopy with use of local anaesthesia before operation <p>Exclusion criteria:</p> <ul style="list-style-type: none"> medical diseases such as ischaemic heart disease, stroke, diabetes mellitus. <p>All patients N: 59</p> <p>Group 1 –TUIP N: 29 Dropouts: 0 Age, years, mean (±SD): 65±1.4 Prostate size, g, mean(±SD): NR</p> <p>Group 2 –TURP N: 30 Dropouts: 0 Age, years, mean (±SD): 70±1.7 Prostate size, g, mean(±SD): NR</p>	<p>Group 1-TUIP Bladder neck resection was performed with diathermy loops. A 24 or 26F continuous irrigation Wolf resectoscope was used. The prostate was resected at the 4 and 8 o'clock positions until the capsule was reached. Homeostasis was secured before the capsule of the prostate was incised. Incisions were made with the same diathermy loop until extracapsular fat was reached. The incision extended from the verumontanum to the level below the trigone. The prostatic chips, which weighted approximately 5 g were sent for pathological examination</p> <p>Group 2-TURP The usual complete resection of the prostatic adenoma to the capsule was performed. A 22F 3-way Foley catheter was used with traction on a 40 to 50 ml balloon and irrigation with normal saline in both situations.</p>	<p>Mortality (at operation)</p>	<p>Group 1: 0/29 Group 2: 0/30 p value: Not sig</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Baseline parameters, except age, not reported (patients were in acute urinary retention). Method of concealment not reported. No symptom scores were collected <p>Additional outcomes: Bleeding or extravasation requiring further operation=0</p> <p>Notes: All the surgeries were only performed by 2 “experienced urologists”</p>
			<p>Qmax (ml/s), mean ±se [baseline values not reported]</p>	<p>At 3 months Group 1: 22.8±2.9 Group 2: 18.5±2.7 p value: Not sig</p>	
			<p>Perioperative complications: Blood transfusions determined by anaesthetist based on blood pressure, pulse rate, and general condition or observation on the return of irrigation fluid</p>	<p>Group 1: 2/29 Group 2: 13/30 Relative risk: 95% CI: p value: 0.004</p>	
			<p>Perioperative complications: UTI</p>	<p>Group 1: 5/29 Group 2: 13/30 Relative risk: 95% CI: p value: 0.05</p>	
			<p>Perioperative complications: TUR syndrome</p>	<p>Group 1: 0/29 Group 2: 0/30 p value: Not sig</p>	
			<p>Post operative complications: Acute urinary retention</p>	<p>Group 1: 0/29 Group 2: 0/30 p value: Not sig</p>	
			<p>Recatheterisation (due to secondary haemorrhage)</p>	<p>Group 1: 0/29 Group 2: 2/30 p value: Not sig</p>	
			<p>Urinary incontinence (transient, 2 weeks for the TURP group)</p>	<p>Group 1: 1/29 Group 2: 2/30 p value: Not sig</p>	
			<p>Urethral stricture (at bulbous urethra asymptomatic, detected using cystoscopy)</p>	<p>At 3 months Group 1: 0/29 Group 2: 1/30 p value: Not sig</p>	
			<p>Bladder neck stenosis (asymptomatic, detected using cystoscopy)</p>	<p>At 3 months Group 1: 0/29 Group 2: 1/30</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				p value: Not sig	
			Length of operation, min, mean±se	Group 1: 19±2.9 Group 2: 36±3.6 p value: 0.0002	
			Length of hospitalisation, days, mean ± se	Group 1: 5.6±0.6 Group 2: 8.0±1.3 p value: Not sig	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nielsen1988²²¹</p> <p>Study design: RCT</p> <p>Setting: Odense University Hospital, Denmark</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Up to 1 year</p>	<p>Patient group: Consecutive patients with symptomatic benign BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> ▪ patients with symptomatic bladder outlet obstruction cause by prostate hypertrophy ▪ Age >60 <p>All patients N: 49</p> <p>Drop outs: 4 at 12 months (2 deaths, 2 refused to attend follow up)</p> <p>Group 1-TUIP N: 24 Age, years, median: 69(60-85) Qmax (ml/s), median; 5(5-10) Prostate weight, g, estimated: <30: 3 30-50:14 >50: 7</p> <p>Group 2 -TURP N: 25 Age, years, median: 73(61-83) Qmax (ml/s), median; 5(5-13) Prostate weight, g, estimated: <30: 7 30-50:14 >50: 4</p>	<p>Group 1-TUIP After cystoscopy, a resectoscope was inserted and a cut was made along the sulcus, using the Stortz diathermy knife, either at 5 or 7 o'clock from the left or right ureteric orifice to the level of the verumontanum, and deepened along its whole length until reaching the fat layer.</p> <p>Group 2-TURP The whole of the prostatic gland resected using a cutting loop.</p> <p>For both groups: Haemostasis was achieved using electrocoagulation. Prophylactic antibiotics not used</p> <p>Anaesthesia: spinal or general</p> <p>Catheter protocol: A catheter (18 to 22 F) was inserted and withdrawn as soon as urine became clear.</p>	<p>All cause mortality (myocardial infarction in TURP and colon cancer in TUIP)</p>	<p>Group 1: 1/24 Group 2: 1/25 p value: Not sig</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ No symptom scores were collected ▪ Randomisation method reported but concealment method unclear <p>Additional outcomes:</p> <p>Notes: Sample size calculation provided for this study – assumption that TURP was 30% better (not stated which outcome) that TUIP, at the 90% power and Type I error or 0.05.</p> <p>Authors reported statistical significance based on fisher's exact test or Mann Whitney test (appropriate)</p> <p>Sexual function, eg retrograde ejaculation not reported</p>
			<p>Qmax, ml/s, mean</p>	<p><u>At baseline</u> Group 1: 5(5-10), n=24 Group 2: 5(5-13), n=25 p value: Not sig <u>At 2 month follow up</u> Group 1: 10(7-18), n=24 Group 2: 17(6-32) n=25 p value: <0.02 <u>At 12 months follow up</u> Group 1: 9(5-25), n=22 Group 2: 12(5-28), n=23 p value: Not sig</p>	
			<p>Perioperative complication; Blood transfusion</p>	<p>Group 1: 1/24 Group 2: 20/25 Relative risk: p value: <0.02</p>	
			<p>Septicaemia</p>	<p>Group 1: 1/24 Group 2: 2/25 p value: >0.1</p>	
			<p>Acute urinary retention (required reoperation, TURP)</p>	<p>Group 1: 3/24 Group 2: 0/25 p value: Not sig</p>	
			<p>Clot retention (reoperation required)</p>	<p>Group 1: 1/24 Group 2: 1/25 p value: Not sig</p>	
			<p>Incontinence</p>	<p>Group 1: 0/24 Group 2: 1/25 p value: Not sig</p>	
			<p>Successful (incontinence or increased frequency of micturation was not considered not successful results)</p>	<p><u>At 2 month follow up</u> Group 1: 24/24, n=24 Group 2: 20/25 n=25 p value: Not sig <u>At 12 months follow up</u> Group 1: 21/22, n=22</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 18/23, n=23 p value: Not sig	
			Reoperation rate (At 2 months, 3 patients in the TUIP group had urinary retention group had required TURP. 1 patient from each group had clot retention and had to be operated again)	<u>At 2 month follow up</u> Group 1: 4/24 Group 2: 1/25 <u>At 12 month follow up</u> This was not clearly reported	
			Stricture (4 patients in TURP group had stricture, 2 had internal urethrotomy and 2 by dilatation)	<u>At 2 month follow up</u> Group 1: 0/24 Group 2: 4/25	
			Length of catheterisation days, median (range)	Group 1: 1(1-2) Group 2: 1(1-4) p value : >0.1	
			Length of operation , minutes, median (range)	Group 1: 18 (10-35) Group 2: 45(20-80) p value: <0.01	
			Length of hospitalisation , days, median, (range)	Group 1: 3(2-13) Group 2: 3(2-18) p value: >0.1	

See Evidence Table 26: Laser coagulation vs. transurethral resection of the prostate (TURP)

for Rodrigo et al., 1998²⁵³

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Riehmann et al, 1995²⁵⁰</p> <p>Study design: RCT</p> <p>Setting: Jan 1985 to Aug 1990, Madison, Wisconsin, US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Mean 34 months (range 7 to 82 months)</p>	<p>Inclusion criteria: patients with bladder outlet obstruction symptoms</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ prostatic urethra > 3 cm or median lobe > 2g ▪ previous prostatic or major pelvic surgery ▪ high operative risk or overt neurological or psychiatric disease <p>All patients Number of eligible patients: 120 Number of patients randomised: 117 Drop outs: 5 (1 received radical prostatectomy after TURP specimen revealed cancer of the prostate, 1 had bladder perforation during the surgery and 1 patient who had TUIP initially had a TURP before the one month follow up) Mean age:</p> <p>Group 1-TUIP N: 61 Drop outs: Age, years, mean (range):65(51–77) Madsen Iversen score, mean: 15.5 Qmax , ml/s mean: 9 (n = 52)</p>	<p>Group 1-TUIP Performed using a Coling's knife at the 6 o'clock position from the bladder neck distally to the verumontanum. The incision extended through the posterior prostatic capsule</p> <p>Group 2-TURP The prostate was resected completely and circumferentially to the anatomic capsule from the bladder neck to the verumontanum.</p> <p>Mean weight of tissue resected : 15 g (range from 1 to 37 g)</p> <p>For both groups Procedures were performed by staff members or residents supervised for staff</p>	<p>All cause mortality (one death in the TURP group was due to saddle pulmonary embolism, classified as operative death)</p> <p>Madsen Iversen, (range of 1-27), mean±se [Values estimated from graph]</p>	<p>Group 1: 14/61 Group 2: 8/56 p value: Not sig</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> ▪ Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported ▪ Results reported graphically-actual values not stated ▪ Qmax significantly higher in TURP group preoperatively <p>Additional outcomes:</p> <ul style="list-style-type: none"> ▪ Madsen Iversen symptom score – results reported in graph, no statistical difference between two groups' pre and post operatively. The scores were significantly lower compared to baseline for both procedures. ▪ Overall subjective assessment of surgical outcomes ▪ Perforation during surgery- 1 case (did not state which arm) <p>Notes: Christensen1990⁵² reported the preliminary results</p>
			<p>Qmax, ml/s, mean± SD:</p>	<p><u>At baseline</u> Group 1: 15.5, n=61 Group 2: 15.5, n=56 p value: Not sig <u>At 3 month follow up</u> Group 1: 6 SE1 n=51 Group 2: 6, SE1 n=52 p value: Not sig <u>At 12 months follow up</u> Group 1: 6 SE 0.5, n=50 Group 2: 5.5 SE 0.5, n=46 p value: Not sig <u>At 24 months follow up</u> Group 1: 7 SE 1, n=41 Group 2: 5 SE 1.5, n=40 p value: Not sig <u>At 36 months follow up</u> Group 1: 8 SE 1, n=22 Group 2: 6.5 SE 1.5, n=19 p value: Not sig <u>At 48 months follow up</u> Group 1: 10.5 SE 1, n=17 Group 2: 9.5 SE 1.5, n=17 p value: Not sig <u>At 60 months follow up</u> Group 1: 9.5 SE 1, n=8 Group 2: 9.5 SE 1.5, n=15 p value: Not sig <u>At 72 months follow up</u> Group 1: 10 SE 1, n=6 Group 2: 9.5 SE 1.5, n=11 p value: Not sig All stat sig compared to baseline</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2-TURP N: 56 Drop outs: Age, years, mean (range):64 (42–78) Madsen Iversen score, mean: 15 Qmax , ml/s mean:11 (n = 50)</p>		<p>[Values estimated from graph]</p>	<p>Group 1: 9, n=52 Group 2: 11, n=50 p value: Stat sig, p<0.015 <u>At 3 month follow up</u> Group 1: 15 SE2 n=42 Group 2: 20, SE2 n=44 p value: Stat sig, p<0.015 <u>At 12 months follow up</u> Group 1: 16 SE 2 , n=42 Group 2: 19 SE 2, n=37 p value: Not sig <u>A 24 months follow up</u> Group 1: 12.5 SE 1, n=32 Group 2: 17 SE 2, n=31 p value: Stat sig, p<0.015 <u>At 72 months follow up</u> Group 1: 13 SE 4, n=4 Group 2: 19 SE 5, n=8 p value: Not sig Not sig compared to baseline for 72 month follow up</p>	
			<p>Reoperation (TURP group – 8 TUIP or resection of bladder neck contracture, 1 further TURP, TUIP group- 12 received TURP, 1 received another TUIP)</p>	<p>Group 1: 13/61 Group 2: 9/56 p value: Not sig</p>	
			<p>Retrograde ejaculation (among patients who were sexually active before an after surgery)</p>	<p>Group 1: 8/23 Group 2: 15/22 Relative risk: 95% CI: p value: 0.02</p>	
			<p>Duration of operation time, mean, (range)</p>	<p>Group 1: 23 (7 to 95) Group 2: 55 (5 to 135) P value: 0.001</p>	
			<p>Catheter duration, day, mean,(range)</p>	<p>Group 1: 1.4 (1-3) Group 2: 2.5(1-12) P value: 0.001</p>	
			<p>Length of hospital stay day, mean,(range)</p>	<p>Group 1: 3.0 (1-8) Group 2: 4.3 (2-14)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				P value: 0.001	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Saporta et al, 1996 ²⁶⁸ Study design: RCT Setting: Not stated (Israel/Turkey) Evidence level: 1+ Duration of follow-up: 72 months	Inclusion criteria: <ul style="list-style-type: none"> patients with obstructive BPH symptoms prostate weight at DRE \leq 40g Exclusion criteria: <ul style="list-style-type: none"> chronic urinary retention urethral stricture, bladder cancer, prostatitis; clinical and suspicion of prostatic cancer; prominent median lobe of prostate neurogenic bladder All patients N: 40 Age, years, mean (\pmSD): 66.85 \pm 2.28 Drop outs: 4 Group 1 N: 20 Drop outs: 3 Age, years, mean (\pmSE): 66.85 \pm 2.28 Prostate size, g, mean(\pmSE): 29.55 \pm 0.94(20-37) Sexually active with antegrade ejaculation: 16/20 [†] Group 2 N: 20 Drop outs: 1 at 3 rd year Age, years, mean (\pmSE): 71.45 \pm 1.15 Prostate size, g, mean(\pmSE): 30.0 \pm 1.51(19-40) Sexually active with antegrade ejaculation: 10/20 [†]	Group 1-TUIP Incision with Collings knife from interureteric ridge from 6 o'clock to verumontanum as deep as fat layer Catheter protocol: 20Fr Foley for 18–24 hours Group 2-TURP Low pressure continuous flow with trocar cystostomy Catheter protocol: 14Fr Foley through trocar cystostomy channel and 20Fr Foley through urethra; irrigated for 18–24 hours; 14Fr Foley removed next day, 20Fr 48 hours after procedure For both groups: spinal, epidural or general were used	Symptom score, Madsen Iversen (range of 1-27), mean \pm se (range)	<u>At baseline</u> Group 1: 14.7 \pm 0.96 (7-21) Group 2: 14.3 \pm 0.93 (6-22) p value: Not sig <u>At 1st year</u> Group 1: 5.29 \pm 0.62 (2-13), n=17 Group 2: 4.95 \pm 0.74 (1-14), n=20 p value: Not sig <u>At 3rd year</u> Group 1: 7.0 \pm 0.64 (3-14), n=17 Group 2: 5.79 \pm 0.85 (1-18), n=19 p value: Not sig	Funding: Not stated Limitations: <ul style="list-style-type: none"> Baseline slightly different Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported Patients who were reoperated not included in analysis Additional outcomes: There was a third arm of balloon dilatation. Notes: Appropriate non-parametric tests used for this study [†] Unequal number of patients with retrograde ejaculation at baseline
			Global assessment of symptoms (marked/moderate or slight improvement/no improvement or worse, %) Patients who required additional treatment were recorded as no improvement	<u>At 1st year</u> Group 1: 80/5/15 Group 2: 85/10/5 p value: Not sig <u>At 3rd year</u> Group 1: 50/30/20 Group 2: 60/35/5 p value: Not sig	
			Qmax, ml/s, mean \pm se(range)	<u>At baseline</u> Group 1: 7.35 \pm 0.56 (3.7-12) Group 2: 6.5 \pm 0.43(3.2-11.9) p value: Not sig <u>At 1st year</u> Group 1: 14.58 \pm 1.05(5.3-5.7), n=17 Group 2: 17.29 \pm 1.16(8.2 -7.1), n=20 p value: Not sig <u>At 3rd year</u> Group 1: 12.65 \pm 1.04(4.1-23.3), n=17 Group 2: 14.36 \pm 1.14(5.5-25.5), n=19 p value: Not sig	
			Retrograde ejaculation [†]	<u>At 3rd year</u> Group 1: 3/16	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 2: 9/10 RR: 0.21 (0.14-0.49) P value: 0.001 [calculated by NCGC team using Fisher's exact test]	
			Reoperation rate For TURP patient- 1 internal urethrotomy in 3 rd year. For TUIP patients, 2 had TURP and 1 had another TUIP at 1 year	<u>At 1st year</u> Group 1: 3/20 Group 2: 0/20 P value: NR <u>At 3rd year</u> Group 1: 3/20 Group 2: 1/20 P value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Soonawalla and Pardanani 1992²⁸⁵</p> <p>Study design: RCT</p> <p>Setting: India</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24 months</p>	<p>Inclusion criteria:</p> <ul style="list-style-type: none"> patients with prostate hypertrophy <p>Exclusion criteria:</p> <ul style="list-style-type: none"> prostatic cancer or suspicion of malignancy prostate size >30g <p>All patients N: 220 Age: 45-87 years</p> <p>Group 1-TUIP N: 110 Age, years, mean: 62.2 Qmax (ml/s), mean; 7.91 Prostate weight, g, mean: 14.8 Sexually active: 60/110</p> <p>Group 2 -TURP N: 110 Age, years, mean: 65.0 Qmax (ml/s), mean; 8.04 Prostate weight, g, mean: 15.6 Sexually active: 49/110</p>	<p>Group 1-TUIP A single incision at the 5 or 7 o'clock position extending from below the ureteric orifice up to the verumontanum was made the Coling's knife and deepened up to the perivesicle and periprostatic fat along its entire length Anaesthesia: general Anaesthesia (69) and spinal (24), local (17 cases) Catheter protocol: 24Fr Foley; 24–48hours</p> <p>Group 2-TURP Catheter protocol: 24Fr Foley; ≤ 48hours</p> <p>For both groups: Anaesthesia: general Anaesthesia (88) and spinal (20) and epidural (2 cases)</p>	<p>All cause mortality (myocardial infarction- 1 each in TUIP and TURP, 1 septicemia in TURP)</p>	<p>Group 1: 1/110 Group 2: 2/110 p value: Not sig[#]</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Methods of randomisation and concealment and whether subjects were blinded to treatment received were not reported No symptom scores were collected <p>Additional outcomes:</p> <ul style="list-style-type: none"> 4/7 of the patients with retention after TUIP had repeat TUIP, and 3 had resection. All 4 TURP patients with urinary retention had reoperation. % of patients satisfied (excellent/fair) vs. not satisfied (no change/worse)-determined "subjectively", methods not reported <p>Notes: # Relative risk (RR) and/or P value calculated by NCGC team using Fisher's exact test</p>
			<p>Qmax, ml/s, mean</p>	<p>At baseline Group 1: 7.91, n=110 Group 2: 8.04, n=110 At 3 month follow up Group 1: 19.38, n=110 Group 2: 20.69 n=110 At 12 months follow up Group 1: 19.45, n=70 Group 2: 20.10, n=67 At 24 months follow up Group 1: 18.91, n=70 Group 2: 19.86, n=67 p value: Not sig for all time points</p>	
			<p>Perioperative complication; Blood transfusion (mean number of units transfused per patient was 0.44)</p>	<p>Group 1: 0/110 Group 2: 38/110 Relative risk: 0.0(95% CI: 0.00 to 1.00)[#] p value: <0.001[#]</p>	
			<p>TUR Syndrome</p>	<p>Group 1: 0/110 Group 2: 7/110 RR: 0.00 (95%CI: 0.00 to 0.53)[#] p value: 0.01[#] [RR and P value calculated by NCGC team]</p>	
			<p>Haemorrhage, 3 intraoperative, requiring open surgery, 2 post-operative haemorrhage</p>	<p>Group 1: 0/110 Group 2: 5/110 p value: Not sig[#]</p>	
			<p>Perforation requiring open surgery</p>	<p>Group 1: 2/110 Group 2: 3/110 p value: Not sig[#]</p>	
			<p>Acute urinary retention (failure to void upon catheter removal)</p>	<p>Group 1: 7/110 Group 2: 4/110 p value: Not sig[#]</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Acute renal failure	Group 1: 0/110 Group 2: 1/110 p value: Not sig #	
			Retrograde ejaculation (among sexually active patients before and after the operations)	Group 1: 14/60 Group 2: 13/49 p value: Not sig #	
			Erectile dysfunction	Group 1: 0/60 Group 2: 0/49 p value: Not sig#	
			Epididymo-orchitis	Group 1: 5/110 Group 2: 2/110 p value: Not sig#	
			Urethral stricture	Group 1: 5/110 Group 2: 3/110 p value: Not sig#	
			Incontinence	Group 1: 2/110 Group 2: 4/110 p value: Not sig#	
			Length of hospitalisation, days, mean	Group 1: 6.03 Group 2: 7.16 p value: NR	
			Length of indwelling catheterisation, min, mean	Group 1: 2.62 Group 2: 3.01 p value: NR	
			Length of operation, min, mean	Group 1: 20.4(10-40) Group 2: 59.2(30-95) p value: NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Tkocz and Prajsner 2002²⁹⁵</p> <p>Study design: RCT</p> <p>Setting: Poland</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24 months</p>	<p>Patient group: Men with moderate symptoms of BPH caused by a small prostate</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> prostate size < 30g <p>Exclusion criteria:</p> <ul style="list-style-type: none"> presence of median lobe <p>All patients N: 100 Mean age: 68±6.7(51 to 78) years Drop outs: 0 (no drop outs reported)</p> <p>Group 1 N: 50 Dropouts: 0 Age, years, mean (±SD): Not reported separately for each group IPSS, mean (±SD): 17.1±2.2 IPSS-QoL, mean (±SD): 4.6±0.5 Prostate size (incised adenoma), g, mean(±SD): 27±2 Residual volume, mean ± SD (ml): 75 ± 22 Pdetmax, cmH₂O, mean ± SD: 84 ± 10</p> <p>Group 2 N: 50 Dropouts: 0 Age, years, mean (±SD): Not reported separately for each group IPSS, mean (±SD): 17.1±1.9 IPSS-QoL, mean (±SD): 4.4±0.3 Prostate size (resected adenoma),</p>	<p>Group 1-TUIP Incisions with a Collins blade, from the urethral orifice to the level of the urethral colliculus, deeply reaching the perivesicle fat. All incisions were performed bilaterally, thus resulting in the full opening of the neck and prostatic urethra.</p> <p>Catheter protocol: Foley 18-French catheter left in the urethra for 24 hours</p> <p>Group 2-TURP Performed using the resectoscope, calibre 24-French.</p> <p>All: subarachnoid anaesthesia with hyperbaric lidocaine</p>	<p>Symptom score, IPSS (range of 1-35), mean±sd</p> <p>IPSS-QoL(range of 1-6) mean±sd</p> <p>Qmax, ml/s, mean± SD:</p> <p>Blood transfusion</p> <p>Retrograde ejaculation</p> <p>Detrusor instability</p>	<p><u>At baseline</u> Group 1: 17.1±2.2 Group 2: 17.1±1.9 P value: Not sig</p> <p><u>At 24 months:</u> Group 1: 4.1±1.8 Group 2: 5.1±1.9 p value: Not sig between groups; <0.01 compared to baseline</p> <p><u>At baseline</u> Group 1: 4.6±0.5 Group 2: 4.4±0.3 <u>At 24 months:</u> Group 1: 2.1±0.3 Group 2: 1.9±0.6 p value: Not sig between groups; <0.01 compared to baseline</p> <p><u>At baseline</u> Group 1: 7.6±1.8 Group 2: 6.9 ±1.5 <u>At 24 months:</u> Group 1: 16.9±1.9 Group 2: 17.6±1.7 p value: Not sig between groups; <0.01 compared to baseline</p> <p>Group 1: 0/50 Group 2: 1/50 p value: Not sig</p> <p>Group 1: 6/50 Group 2: 16/50 Relative risk: 0.38(95% CI: 0.16 to 0.84) P value: 0.03</p> <p><u>Baseline:</u> Group 1: 31/50 Group 2: 30/50 <u>At 24 months</u></p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Methods of randomisation and concealment not reported Patient diary- no mention of content, validation and duration of method of data collection and analysis <p>Additional outcomes: Urodynamic parameters such as Pdetop, PdetQmax, CysCapF etc</p> <p>Notes: No patient reported to have dropped out from study</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>g, mean(\pmSD); 28.2\pm2</p> <p>Residual volume, ml, mean \pmSD : 68 \pm21</p> <p>Pdef_{max}, cmH₂O, mean, \pmSD : 85 \pm8</p>		<p></p> <p>Weakening of detrusor post operation (“lazy” and incomplete voiding, returned to normal by 24 months)</p> <p>Urinary frequency, diurnal (recorded through diary. Diary kept for 7 days after preliminary examination (baseline. No mention of how many days data were collected for follow up)</p> <p>Urinary frequency, nocturnal (recorded through diary. Diary kept for 7 days after preliminary examination (baseline. No mention of how many days data were collected for follow up)</p>	<p>Group 1: 15/50 Group 2: 11/50 P value: Not sig</p> <p><u>Post-op (time not provided)</u> Group 1: 4/50 Group 2: 11/50 P value: Not sig <u>At 24 months</u> Group 1: 0/50 Group 2: 0/50</p> <p><u>Baseline:</u> Group 1: 7.8\pm0.9 Group 2: 7.2\pm1.2 <u>At 24 months</u> Group 1: 4.9\pm1.1 Group 2: 5.2\pm1.0 P value: Not sig between groups; <0.001 compared to baseline</p> <p><u>Baseline:</u> Group 1: 2.8\pm0.9 Group 2: 2.4\pm0.8 <u>At 24 months</u> Group 1: 1.1\pm0.5 Group 2: 0.9\pm0.5 P value: Not sig between groups; <0.001 compared to baseline</p>	

Evidence Table 40: Botulinum toxin vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Maria et al., 2003¹⁸²</p> <p>Study design: RCT, double blinded</p> <p>Setting: Jan to Dec 2000</p> <p>Department of Surgery, University Hospital of Agostino Gemelli, Rome</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 months for blinded study, 12 months for open label on the active arm</p>	<p>Patient group: Men with symptomatic BPH</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Age 50 to 80 years with symptomatic BPH Moderate to severe symptoms of urinary obstruction as determined by the AUA score Qmax ≤ 15 ml/s with a voided volume of ≥150mL An enlarged prostate gland on digital rectal examination <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic voiding disorders Prostate or bladder cancer or a serum PSA level of 10 ng/ml or more Previously had surgery or treated with botulinum toxin <p>All patients N: 30 (out of 42 assessed for eligibility, 8 did not meet inclusion criteria, 4 refused) Drop outs: 0</p> <p>Group 1 N: 15 Age, years, mean (±SD): 69.4±4.9 Prostate vol ml, mean ± (SD): 52.6±10.6 Residual vol, ml, mean±(SD): 126.3±38.3</p> <p>Group 2 N: 15 Age, years, mean (±SD): 68.2±3.9 Prostate volume ml, mean ± (SD): 52.3±10.0 Residual volume, ml, mean±(SD): 118.0±39.7</p>	<p>Group 1 Botulinum toxin Received 200U of botulinum toxin</p> <p>Group 2 – Placebo Received saline solution</p> <p>For both groups: 4 ml of solution injected in to the prostate, divided into 2 injections of equal volume (2 mL) into each lobe of the gland.</p> <p>With patient lying on the left side, a 22-gauge spinal needle (0.7 X 90-mm Yale spinal needle, Becton Dickinson, Spain) was inserted in the perineum in the anterior midline approximately 1.5 to 2.0 cm from the anus. The injection sites were visualised using transrectal ultrasonography.</p> <p>No sedation or anaesthesia was used during the procedure</p>	<p>AUA symptom score, mean±sd:</p> <p>(No data reported for group 2 after 2nd month)</p> <p>Qmax, ml/s, mean±sd</p> <p>(No data reported for group 2 after 2nd month)</p> <p>Urinary incontinence (at 1 and 2 months)</p>	<p><u>Baseline</u> Group 1: 23.2±4.1 Group 2: 23.3±3.9</p> <p><u>1 month</u> Group 1: 10.6±1.7 Group 2: 23.4±3.5</p> <p><u>2 month</u> Group 1: 8.0±1.6 Group 2: 23.3±3.3</p> <p><u>6 month (open label)</u> Group 1: 9.1±3 <u>12 month (open label)</u> Group 1: 8.9±3.2 P values: Sig *</p> <p><u>Baseline</u> Group 1: 8.1±2.2 Group 2: 8.8±2.5</p> <p><u>1 month</u> Group 1: 14.9±2.1 Group 2: 8.8±2.3</p> <p><u>2 month</u> Group 1: 15.4±1.7 Group 2: 8.7±2.3</p> <p><u>6 month (open label)</u> Group 1: 14.6±4.1 <u>12 month (open label)</u> Group 1: 15±2.9 P values: Sig *</p> <p>Group 1: 0/15 Group 2: 0/15</p>	<p>Funding: Not stated</p> <p>Limitations:</p> <ul style="list-style-type: none"> Small sample size – no calculation provided Uncertain whether all outcomes/side effects relevant to the patient had been reported (eg pain) <p>Additional outcomes: Prostate volume, serum PSA, and residual volume at 1 and 2-months follow up. Also reported the 6 and 12 months follow up results for the botulinum toxin group</p> <p>Prostate size reduction at 1 and 2 months were significant for the botulinum toxin arm</p> <p>Notes: * P values <0.001 for Group 1 compared to baseline, and between Group 1 and 2 at 1 and 2 months</p>

Evidence Table 41: Transurethral vaporesction of the prostate (TUVRP) vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gotoh et al., 1999¹⁰⁶</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: men with moderate to severe LUTS</p> <p>Setting: multi-centre, Department of Urology, Nagoya University School of Medicine, Japan</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS ≥ 10 Qmax < 15 mL/s Prostate volume ≥ 30 ml or higher than normal PSA <p>Exclusion criteria: NR</p> <p>All patients N: 53 Drop outs: 2</p> <p>Group 1: N: 25 Mean age (\pm SD): 69.7 \pm 6.3 Mean IPSS \pm SD: 19.6 \pm 7.5 Mean Qmax ml/s \pm SD: 7.3 \pm 2.8 Mean PVR ml \pm SD: 56.7 \pm 51.4 Mean prostate volume \pm SD (mL): 47.8 \pm 16.4 Operative time \pm SD mins: 60 \pm 28 Resected weight (g): 29.4 \pm 15.1 Drop outs: 2 excluded because cancer found</p> <p>Group 2: N: 28 Mean age (\pm SD): 66.5 \pm 15.7 Mean IPSS \pm SD: 18.9 \pm 7.3 Mean Qmax ml/s \pm SD: 9.4 \pm 2.8 Mean PVR ml \pm SD: 41.9 \pm 25.5</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVRP) Bandloop cutting 230–250W</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop cutting 120W</p> <p>All patients: Same surgeon performed all procedures at each different hospital</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, PSA, Blood, TRUS, uroflowmetry. Flow rate at months 1 & 6 and pressure flow at 3 months. IPSS assessed at 3 months postoperatively</p>	<p>Mean IPSS score \pm SD at 3 months</p> <p>Mean Qmax mL/s \pm SD at 3 months</p> <p>Catheterisation time (days)</p> <p>Complications: transfusion</p> <p>Complications: TUR</p> <p>Complications: Urethral Stricture</p> <p>Complications: UTI</p> <p>Complications: incontinence</p>	<p>Group 1: 3.7 \pm 2.4 (n=23) Group 2: 3.8 \pm 2.3 (n=28) p value: Not sig.</p> <p>Group 1: 23.6 \pm 13.9 Group 2: 21.2 \pm 9.4 p value: Not sig.</p> <p>Group 1: 3.4 \pm 1.3 Group 2: 3.3 \pm 1.3 p value: Not sig.</p> <p>Group 1: 0/25 Group 2: 0/28 p value: NR</p> <p>Group 1: 0/25 Group 2: 0/28 p value: NR</p> <p>Group 1: 0/25 Group 2: 0/28 p value: NR</p> <p>Group 1: 0/25 Group 2: 0/28 p value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Author confirmed no masking of outcome assessment and no allocation concealment Significant differences at baseline for Qmax <p>Additional outcomes: Urinalysis</p> <p>Notes: Author reports randomisation by drawing envelopes</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean prostate volume ± SD (mL): 44.7 ± 15.2 Operative time ± SD mins: 61.1 ± 29 Resected weight (g): 36.5 ± 17.6 Drop outs: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Gupta et al., 2006¹⁰⁸</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: Patients with BPH who were candidates for TURP were selected from July 2002 to December 2003.</p> <p>Setting: single centre: All India Institute of Medical Sciences, New Delhi, India</p> <p>Inclusion criteria: glands of >40g</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Previous history of prostatic and urethral surgery • Neurovesical dysfunction • Carcinoma of the prostate <p>All patients N: 100 Dropouts: NR</p> <p>Group 1 N: 50 Mean ± SD Age: 67.68 ± 9.8 IPSS ± SD: 24.9 ± 3.9 Mean SD Qmax: 4.65 ± 3.6 Mean SD PVR, mL: 103 ± 174.1 Mean prostate size ± SD, g: 62.6 ± 14.8 Resectate ± SD g: 24.8 ± 12.7 Operation duration ±SD min: 55.9 ± 18.1 Patients with catheter: 19/50 Dropouts: NR</p> <p>Group 2 N: 50 Mean ±SD Age: 65.67 ± 7.5 IPSS ± SD: 23.3 ± 3.9 Mean SD Qmax: 4.5 ± 3.9 Mean SD PVR, mL: 84.0 ± 129.7 Resectate ±SD g: 18.9 ± 12.9</p>	<p>Group 1: TUVRP Wing (Wolf) loop: 180W cutting and 80W coagulation</p> <p>Group 2: TURP Standard tungsten wire loop 80W cutting and 50W coagulation</p> <p>All patients 27F continuous-flow resectoscope. 22 F Foley catheter inserted and irrigation with saline. Catheter removed when urine clear.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 1, 3, 6, 12 months for complications and IPSS, PVR, Qmax reassessed at 6 & 12 months</p>	<p>Mean (SD) IPSS at 6 months</p>	<p>Group 1: 5.9 ± 0.25 Group 2: 6.1 ± 0.42 P value: NS</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method and allocation concealment were not reported. • Outcome assessment was not masked • Drop outs NR so patient numbers at follow up unclear <p>Additional outcomes: Irrigation, haemoglobin decrease, serum sodium decrease.</p> <p>Notes: HOLEP arm of study not reported. *ANOVA analysis used to compare 3 groups</p>
			<p>Mean (SD) IPSS at 12 months</p>	<p>Group 1: 5.4 ± 0.28 Group 2: 5.6 ± 0.32 P value: NS</p>	
			<p>Mean (SD) Qmax at 6 months</p>	<p>Group 1: 22.5 ± 0.95 Group 2: 20.7 ± 1.32 P value: NS</p>	
			<p>Mean (SD) Qmax at 12 months</p>	<p>Group 1: 23.6 ± 0.96 Group 2: 23.7 ± 1.58 P value: NS</p>	
			<p>Mean (SD) catheter duration, days (converted from hours)</p>	<p>Group 1: 1.51 ± 0.35 Group 2: 1.90 ± 0.53 P value: Significant*</p>	
			<p>Complications: urinary retention (re-catheterisation)</p>	<p>Group 1: 3/50 Group 2: 3/50</p>	
			<p>Complications: TUR Syndrome</p>	<p>Group 1: 1/50 Group 2: 1/50</p>	
			<p>Complications: Transfusion</p>	<p>Group 1: 0/50 Group 2: 1/50</p>	
			<p>Complications: Mortality (pneumonia)</p>	<p>Group 1: 1/50 Group 2: 0/50</p>	
			<p>Complications: urethral stricture</p>	<p>Group 1: 1/50 Group 2: 2/50</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean prostate size \pm SD, g: 59.8 ± 16.5 Operation duration \pmSD min: 64.1 ± 13.1 Patients with catheter: 16/50 Dropouts: NR				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Helke et al., 2001¹¹³</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: Patients moderate or severe voiding dysfunction and BPE.</p> <p>Setting: single centre: University Hospital Carl Gustav Carus, Dresden, Germany</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Enlarged prostate on DRE At least moderate LUTS IPSS > 10 and/or PVR >60 mL Patients with recent urinary retention and indwelling catheters < 6 weeks duration <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous prostatic surgery Neurogenic bladder disorders Known urethral strictures Prostate cancer Indwelling catheter > 6 weeks duration Severe neurological disease Psychiatric abnormalities Reduced patient compliance <p>All patients N: 185 Dropouts: 37</p> <p>Group 1 N: 93 Mean ± SD Age: 67.3 ± 7.73 (47-85) IPSS ± SD: 17.29 ± 6.06 Mean SD Qmax: 10.8 ± 4.76 Mean SD PVR, mL: 76.0 ± 60.5 Mean prostate volume ± SD, mL: 48.8 ± 21.21 Resectate ± SD g: 21.98 ± 13.47</p>	<p>Group 1: TUVRP Vaporising loop 1mm: 250W cutting</p> <p>Group 2: TURP Standard loop 0.3 mm: 150W cutting</p> <p>All patients 26F intermittent flow resectoscope. Irrigation with Purisole 0.96% alcohol. Antibiotic prophylaxis was given and catheter removed 2-3 days after surgery.</p> <p>TUVRP performed by 5 urologists with experience of at least 5 TUVRP patients each</p> <p>Examination methods Preoperative: Baseline ASA, New York Heart Association scores, IPSS Symptom score, AUA bother score, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 3, 6, 12 months for PVR and flow rates at 12 months. Symptom score follow up by postal questionnaire</p>	<p>Mean (SD) IPSS at 12 months</p> <p>Mean (SD) Qmax at 12 months</p> <p>Complications: incontinence</p> <p>Complications: Transfusion</p> <p>Complications: urethral stricture</p> <p>Complications: reoperation</p>	<p>Group 1: 4.66 ± 4.3 (n=79) Group 2: 5.21 ± 5.1 (n=69) P value: NS</p> <p>Group 1: 22.19 ± 12.3 Group 2: 22.12 ± 10.6 P value: NS</p> <p>Group 1: 0/93 Group 2: 0/92</p> <p>Group 1: 6/93 Group 2: 9/92</p> <p>Group 1: 5/93 Group 2: 7/92</p> <p>Group 1: 9/93 Group 2: 5/92</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment were not reported. Outcome assessment was not masked Significant difference reported between baseline Qmax p = 0.02 Significant difference found between baseline PVR p = 0.02 which was not reported as significant. <p>Additional outcomes: IPSS & Bother score were reported graphically at 3, 6 and 12mths</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Operation duration \pmSD min: 71.02 \pm 27.5 Indwelling catheter: 28/93 Dropouts: 14 (2 patients underwent radical prostatectomy and were excluded, 11 lost to follow up and incomplete outcome data for 1)</p> <p>Group 2 N: 92 Mean \pmSD Age: 68.7 \pm 8.38 (53-89) IPSS \pm SD: 18.29 \pm 7.49 Mean SD Qmax: 8.5 \pm 5.19 Mean SD PVR, mL: 101.8 \pm 84.1 Resectate \pmSD g: 18.9 \pm 12.9 Mean prostate volume \pm SD, mL: 49.9 \pm 22.1 Operation duration \pmSD min: 65.68 \pm 25.8 Indwelling catheter: 32/93 Dropouts: 23 (4 patients underwent radical prostatectomy and were excluded, 14 lost to follow up and incomplete outcome data for 5)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kupeli et al., 2001¹⁵⁵</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient Group: Moderate to severe symptoms of prostatism</p> <p>Setting: single centre: Ankara University, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 8 Qmax < 15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Carcinoma of the prostate History of prostate surgery <p>All patients N: 100 Dropouts: NR</p> <p>Group 1 N: 50 Mean \pm SD Age: 61.4 \pm 3.2 IPSS \pm SD: 19.4 \pm NR Mean SD Qmax: 7.9 \pm 2.1 Mean prostate size \pm SD, g: 57.8 \pm 4.1 Resectate \pm SD g: NR Operation duration \pmSD min: 48.2 \pm NR Previous medical treatment: 32/50 Preoperative retrograde ejaculation: 50/50 Preoperative erectile dysfunction: 14/50 Dropouts: NR</p> <p>Group 2 N: 50 Mean \pmSD Age: 58.9 \pm 3.6 IPSS \pm SD: 21.6 \pm NR Mean SD Qmax: 9.2 \pm 2.6</p>	<p>Group 1: TUVRP Wing (Wolf) loop: 205-300W cutting</p> <p>Group 2: TURP Storz 24F loop: 80-120W cutting</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 6 months</p>	<p>Mean (SD) IPSS at 6 months</p>	<p>Group 1: 4.0 \pm NR Group 2: 5.0 \pm NR* P value: NS</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment were not reported. Outcome assessment was not masked No mention of drop outs in the study Standard deviations for IPSS NR Significance difference in baseline Qmax p=0.007 Almost all patients had retrograde ejaculation prior to surgery <p>Additional outcomes: Haemocrit and sodium</p> <p>Notes: None.</p>
			<p>Mean (SD) Qmax at 6 months</p>	<p>Group 1: 26.7 \pm 3.7 Group 2: 24.6 \pm 3.4 P value: NR</p>	
			<p>Mean (SD) catheter duration, days (converted from hours)</p>	<p>Group 1: 2 \pm NR Group 2: 4 \pm NR P value: <0.05</p>	
			<p>Mean (SD) length of stay, days</p>	<p>Group 1: 2.5 \pm NR Group 2: 4.5 \pm NR P value: <0.05</p>	
			<p>Complications: urinary retention (re-catheterisation)</p>	<p>Group 1: 0/50 Group 2: 0/50</p>	
			<p>Complications: TUR Syndrome</p>	<p>Group 1: 0/50 Group 2: 0/50</p>	
			<p>Complications: Transfusion</p>	<p>Group 1: 0/50 Group 2: 0/50</p>	
			<p>Complications: Incontinence</p>	<p>Group 1: 0/50 Group 2: 0/50</p>	
			<p>Complications: Retrograde ejaculation</p>	<p>Group 1: 26/50 Group 2: 27/50</p>	
			<p>Complications: urethral stricture</p>	<p>Group 1: 0/50 Group 2: 0/50</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean prostate size \pm SD, g: 56.7 \pm 6.3 Resectate \pm SD g: NR Operation duration \pmSD min: 42.7 \pm NR Previous medical treatment: 31/50 Preoperative retrograde ejaculation: 44/50 Preoperative erectile dysfunction: 19/50 Dropouts: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Liu et al., 2006¹⁶⁸</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 years</p>	<p>Patient Group: Patients with BOO due to BPH on waiting list for surgery</p> <p>Setting: single centre: Taipei City Hospital, Taiwan</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 15 IPSS QoL \geq 3 Qmax \leq 12 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> PSA \geq 4 ng/mL Neurogenic bladder Carcinoma of the prostate History of prostate or urethral surgery Bladder stones Patients on anticoagulant therapy <p>All patients N: 76 Dropouts: NR</p> <p>Group 1 N: 44 Mean \pm SD Age: 66.0 \pm 6.6 IPSS \pm SD: 26.8 \pm 4.7 IPSS QoL \pm SD: 4.1 \pm 0.6 Mean SD Qmax: 6.9 \pm 2.1 Mean SD PVR, mL: 142 \pm 48 Mean prostate volume \pm SD, mL: 60.5 \pm 10.9 Resectate \pm SD g: 32.2 \pm 7.1 Operation duration \pm SD min: 49.4 \pm 8.0</p>	<p>Group 1: TUVRP Wedge resection loop: 200W cutting and 60W coagulation</p> <p>Group 2: TURP Standard wire loop 110W cutting and 60W coagulation.</p> <p>All patients 27F continuous-flow resectoscope. 22 F Foley catheters inserted.</p> <p>TUVRP performed by 3 urologists with experience of at least 10 TUVRP patients each</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 3, 6, 12 months and 2 years</p> <p>Sexual function was assessed by face to face or telephone questionnaire</p>	Mean (SD) IPSS at 3 months	Group 1: 8.2 \pm 2.2 (n=42) Group 2: 7.9 \pm 1.8 (n=30) P value: 0.53	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Unbalanced baseline numbers Allocation concealment unclear Outcome assessment was not masked Number of patients remaining at 2 years was unclear and reasons for incomplete outcome data not given. <p>Notes: Randomisation by drawing envelopes</p>
			Mean (SD) IPSS at 2 years	Group 1: 9.0 \pm 3.1 Group 2: 8.4 \pm 2.6 P value: 0.45	
			Mean (SD) IPSS QoL at 3 months	Group 1: 1.7 \pm 0.5 (n=36) Group 2: 1.5 \pm 0.7 (n=26) P value: 0.57	
			Mean (SD) IPSS QoL at 2 years	Group 1: 1.6 \pm 0.6 Group 2: 1.4 \pm 0.7 P value: 0.48	
			Mean (SD) Qmax at 3 months	Group 1: 20.7 \pm 2.8 (n=29) Group 2: 21.6 \pm 2.0 (n=21) P value: 0.2	
			Mean (SD) Qmax at 2 years	Group 1: 19.6 \pm 3.7 Group 2: 21.2 \pm 2.7 P value: 0.12	
			Mean (SD) catheter duration, days (converted from hours)	Group 1: 1.06 \pm 0.18 Group 2: 1.66 \pm 0.38 P value: <0.0001	
			Mean (SD) length of stay, days	Group 1: 1.65 \pm 0.2 Group 2: 2.06 \pm 0.35 P value: <0.0001	
			Complications: urinary retention (re-catheterisation)	Group 1: 3/44 Group 2: 4/32	
			Complications: TUR Syndrome	Group 1: 0/44 Group 2: 2/32	
Complications: Transfusion	Group 1: 1/44 Group 2: 2/32				
Complications: Incontinence	Group 1: 2/44 Group 2: 1/32				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Dropouts: NR</p> <p>Group 2 N: 32 Mean ±SD Age: 64.7 ± 6.3 IPSS ± SD: 25.6 ± 3.5 IPSS QoL ± SD: 4.0 ± 0.7 Mean SD Qmax: 6.9 ± 1.9 Mean SD PVR, mL: 131 ± 41 Resectate ±SD g: 35.5 ± 4.3 Mean prostate volume ± SD, mL: 58.4 ± 8.4 Operation duration ± SD min: 52.9 ± 6.0 Dropouts: NR</p>		<p>Complications: Reoperation rate</p> <p>Complications: urethral stricture</p> <p>Complications: retrograde ejaculation * answered by those men who were sexually active preoperatively in each group</p>	<p>Group 1: 2/44 Group 2: 3/32</p> <p>Group 1: 3/44 Group 2: 2/32</p> <p>Group 1: 10/17 Group 2: 7/13</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Netto et al., 1999 ²¹⁹	<p>Patient group: moderate to severe symptomatic BPH</p> <p>Setting: single-centre, division of urology, Unicamp & Hospital Beneficencia Portuguesa, São Paulo, Brazil</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Patients with >1 symptomatic and uncomplicated BPH IPSS >12 Qmax < 15 mL/s Voided volume ≥150mL PVR <250 mL Prostate volume 25-90 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Exposure to α-antagonists, anticholinergics, cholinergics, diuretics, estrogens, androgens, antihypertensive medications or other agents within the previous 2 weeks Prostate cancer Urethral stricture Urinary tract stone disease Neurogenic bladder Hydronephrosis UTI within 3 months prior to surgery Pelvic irradiation Previous prostatic surgery <p>All patients N: 78 Drop outs: NR</p> <p>Group 1: N: 40</p>	<p>Group 1: Transurethral vaporisation of the prostate (TUVP) Standard loop: cutting 250-300 without haemostasis</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop: cutting 50-80 and haemostasis mode 50W</p> <p>All patients: Operations performed using 24F continuous flow resectoscope using a 3% mannitol as irrigant. A 22F Foley catheter was inserted. Oral antibiotics for 1 week.</p> <p>Examination methods Preoperative: Baseline IPSS symptom score, urinalysis, PSA, TRUS, uroflowmetry. Follow up visits at 3, 6, 12 months and annually thereafter</p>	<p>Mean IPSS score at mean follow up 17 months (follow up interval not clear for each group)</p> <p>Mean Qmax ± SD mL/s at mean follow up 17 months (follow up interval not clear for each group)</p> <p>Catheterisation time (days) hours reported converted to days</p> <p>Length of hospital stay (days)</p> <p>Complications: retrograde ejaculation</p> <p>Complications: TUR</p> <p>Complications: urethral stricture</p>	<p>Group 1: 3.83 ± 4.62 Group 2: 8.68 ± 2.30 p value: <0.00001 (calculated by NCGC as <i>t</i> test with unequal variances) conflicts with study finding p=0.88</p> <p>Group 1: 15.43 ± 3.4 Group 2: 16.16 ± 2.48 p value: 0.28 (calculated by NCGC as <i>t</i> test with equal variances) conflicts with study finding p=0.02</p> <p>Group 1: 0.77 ± 0.29 Group 2: 1.68 ± 0.36 p value: <0.00001 (calculated by NCGC as <i>t</i> test with equal variances)</p> <p>Group 1: 1.55 ± 0.75 Group 2: 2.63 ± 0.63 p value: <0.0001</p> <p>Group 1: 26/40 (65%) Group 2: 12/38 (32%) p value: NR</p> <p>Group 1: 0/40 Group 2: 0/38 p value: NR</p> <p>Group 1: 0/40 Group 2: 0/38 p value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Masked outcome assessment was not reported Follow up interval for each group not clear only overall mean follow up reported. There were significant baseline differences in IPSS score Dropouts were not reported. P values reported conflicted with outcome measures. <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> Mean age (range): 66.8 (52-80) Mean IPSS score: 19.65 ± 6.14 Mean Qmax ml/s (± SD): 7.88 ± 2.51 PVR mL (range): 73.0 ± 5.81 Mean prostate volume mL ± SD: 46.88 ± 17.1 Operative time ± SD: 29.78 ± 11.78 mins Resectate ± SD, g: 21.6 ± NR Drop outs: NR </p> <p> Group 2: N: 38 Mean age (range): 65 (51-82) Mean IPSS score: 24.29 ± 6.48 Mean Qmax ml/s (± SD): 6.77 ± 3.08 PVR mL (range): 88.64 ± 8.43 Mean prostate volume mL ± SD: 53.4 ± 21 Operative time ± SD: 56.32 ± 8.36 mins Resectate ± SD, g: 22.3 ± NR Drop outs: NR </p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Tallic et al., 2000²⁹¹</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months (Mean follow up 9.2 mths for TUVRP and 8.8 mths for TURP)</p>	<p>Patient Group: Patients with BOO due to BPH on waiting list for surgery</p> <p>Setting: single centre: King Khalid University Hospital, Saudi Arabia</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Men with urinary retention IPSS > 15 Qmax < 15 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Carcinoma of the prostate History of prostate or urethral surgery <p>All patients N: 68 Dropouts: NR</p> <p>Group 1 N: 34 Mean \pm SD Age: 70.9 \pm 9.3 IPSS \pm SD: 24.9 \pm 6 Mean SD Qmax: 7.5 \pm 3.5 Mean prostate size \pm SD, g: 52.4 \pm 18.7 Resectate \pm SD g: 22.4 \pm 10.5 Men with urinary retention: 15/34 Operation duration \pm SD min: 42.4 \pm 15 Urinary retention: 15/34 Dropouts: NR</p> <p>Group 2 N: 34 Mean \pmSD Age: 70.4 \pm 8.8 IPSS \pm SD: 20.1 \pm 6.8 Mean SD Qmax: 9.1 \pm 6.3 Resectate \pmSD g: 20.2 \pm 9.5</p>	<p>Group 1: TUVRP Wing resection loop: 250W cutting and 80W coagulation</p> <p>Group 2: TURP Standard wire loop 150W cutting and 50W coagulation.</p> <p>All patients 27F continuous-flow resectoscope. Foley catheters inserted with saline irrigation</p> <p>TUVRP performed by 3 urologists with experience of at least 10 TUVRP patients each</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, blood, uroflowmetry. Follow up every 3 months</p> <p>Sexual history questionnaire was answered by those men that were sexually active</p>	<p>Mean (SD) IPSS at 6 months</p>	<p>Group 1: 4.0 \pm 3.4 Group 2: 5.6 \pm 3.1 P value: 0.03</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment not reported Outcome assessment was not masked Significant baseline differences in Qmax p=0.02 & IPSS p<0.0001 Dropouts were not reported <p>Additional outcomes: Haematocrit, haemoglobin, serum sodium</p> <p>Notes: None.</p>
			<p>Mean (SD) Qmax at 6 months</p>	<p>Group 1: 19.0 \pm 6.5 Group 2: 15.2 \pm 10.0 P value: 0.01</p>	
			<p>Mean (SD) catheter duration, days (converted from hours)</p>	<p>Group 1: 0.96 \pm 0.43 Group 2: 1.5 \pm 0.72 P value: <0.0001</p>	
			<p>Complications: TUR Syndrome</p>	<p>Group 1: 0/34 Group 2: 0/34</p>	
			<p>Complications: Transfusion</p>	<p>Group 1: 0/34 Group 2: 0/34</p>	
			<p>Complications: urethral stricture</p>	<p>Group 1: 3/34 Group 2: 4/34</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Men with urinary retention: 18/34 Mean prostate size \pm SD, g: 57.2 \pm 22.5 Operation duration \pm SD min: 35.9 \pm 12.8 Urinary retention: 18/34 Dropouts: NR</p>				

Evidence Table 42: Bipolar TUVRP vs. transurethral resection of the prostate (TURP)

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Fung et al., 2005⁹⁸</p> <p>Study design: RCT Observer and patient masked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient group: men on waiting list for surgery for BPH with acute or chronic retention, failure to remove catheter and</p> <p>Setting: single-centre: Division of Urology, Pamela Youde Nethersole Eastern Hospital, Hong Kong, China</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • IPSS >20 • Qmax <10 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Neurogenic bladder • Urethral stricture • Anticoagulant therapy • Bladder stone • Prostate cancer or suspect • Previous prostate surgery <p>All patients N: 60 Drop outs: 9</p> <p>Group 1: N: 29 (n=21) Mean age (range): 72.5 (59-91) Mean IPSS ± SD: 15.82 ± NR Mean IPSS QoL ± SD: 3.55 ± NR Mean PVR ± SD, mL: NR Mean prostate volume ± SD, mL: NR Resection time (range), min: 36.6 (12-76) Resected weight (range), g: 18.6 (1-57) Patients with urinary retention: 17 Drop outs: 8 for machine failure</p> <p>Group 2: N: 31 (n=30)</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system through 27F resectoscope at 240W for vaporisation and 60W for coagulation.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop through 27F continuous flow resectoscope. Cutting 120W and coagulation 60W</p> <p>All patients: Surgery performed by a consultant, senior medical officer or senior registrar with experience of performing TURP.</p> <p>A 22F 3-way catheter was inserted with saline irrigant until effluent was clear. Catheter removed the following morning</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, QoL, assessed and follow up of IPSS, QoL and Qmax at 3 months</p>	<p>Mean ± SD IPSS change from baseline at 3 months</p>	<p>Group 1: 8.81 ± NR (n=21) Group 2: 9.63 ± NR (n=30) P value: 0.86</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • 8 dropouts in Group 1 due to machine failure • Allocation concealment was not reported <p>Additional outcomes: reduction in serum sodium and haemoglobin</p> <p>Notes: Randomisation using computer generated numbers</p>
			<p>Mean ± SD change in Qmax from baseline at 3 months</p>	<p>Group 1: 16.57 ± NR (n=21) Group 2: 14.71 ± NR (n=30) P value: 0.96</p>	
			<p>Mean ± SD IPSS QoL change from baseline at 3 months</p>	<p>Group 1: 0.55 ± NR (n=21) Group 2: 1.54 ± NR (n=30) P value: 0.17</p>	
			<p>Mean ± SD Qmax at 12 months</p>	<p>Group 1: 17.0 ± NR (n=20) Group 2: 15.0 ± NR (n=20) P value: NR</p>	
			<p>Catheterisation time (days)</p>	<p>Group 1: 1.14 ± NR Group 2: 1.21 ± NR P value: 0.59</p>	
			<p>Complications: urinary retention (re-catheterisation)</p>	<p>Group 1: 4/21 Group 2: 3/30 P value: NR</p>	
			<p>Complications: urinary retention UTI</p>	<p>Group 1: 4/21 Group 2: 4/30 P value: NR</p>	
<p>Complications: TUR</p>	<p>Group 1: 0/21 Group 2: 0/30 P value: NR</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> Mean age (range): 73 (59-88) Mean IPSS \pm SD: 19.36 \pm NR Mean IPSS QoL \pm SD: 3.64 \pm NR Mean PVR \pm SD, mL: NR Mean prostate volume \pm SD, mL: NR Resection time (range), min: 32.9 (12-105) Resected weight (range), g: 25.1 (4-100) Patients with urinary retention: 25 Drop outs: 1 (patient contracted sepsis) </p>				

Evidence Table 43: Transurethral ethanol ablation of the prostate (TEAP) vs. transurethral resection of the prostate (TURP)

See Evidence Table 9: Alpha-blockers vs. placebo
for Kim et al. 2006 ¹⁴⁶

Evidence Table 44: Transurethral resection of the prostate (TURP) vs. watchful waiting

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Wasson et al., 1995³¹⁶ & Anon1993¹</p> <p>Study design: RCT</p> <p>Setting: US, July 1986 to 1989.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 years (average of 2,8 years)</p>	<p>Patient group: Consecutive male veterans referred to urology clinics because of BPH symptoms</p> <p>Inclusion criteria: Score of 10-20 on the Madsen Iversen symptom score (moderate or somewhat severe)</p> <p>Exclusion criteria:</p> <ul style="list-style-type: none"> ▪ <55 years old ▪ History of prostate surgery or radiation treatment ▪ Unable to walk ▪ Had active urinary tract infection not responding to treatment ▪ Received diagnosis of prostate or bladder cancer ▪ Residual volume > 350 ml ▪ Low total score on a scale that rates BPH on a the basis of cystoscopy, the symptom interview and bladder ultrasonography ▪ Serious medical conditions that would have made surgery inappropriate for follow-up unlikely (e.g: uncontrolled diabetes, neurogenic bladder, cirrhosis, active alcoholism, bleeding diathesis, psychosis, and late stage cardiac or respiratory disease) ▪ Serum creatinine concentration >3.0 mg /dl or had doubled in the previous year 	<p>Group 1: TURP Performed by the chief surgical resident or staff surgeon. No description of the procedure was provided</p> <p>Group 2: Watchful waiting No specific description</p> <p>All patients: All participants were told to avoid ingesting coffee, alcohol, and other liquids after dinner and were informed about medications that might make their symptoms worse. Physicians were asked to avoid prescribing medications such as alpha-adrenergic antagonists that might confound the results of the trial. A referral to a urologist was considered if there was an indication of treatment failure or a patient requested such referral.</p> <p>All participants were followed in general medical clinic six to eight weeks after randomisation and</p>	<p>All cause mortality (no deaths associated with surgery)</p>	<p><u>At 3 year follow up</u> Group 1: 13/280 Group 2: 10/276 Relative risk: 1.28 (95% CI: 0.57 to 2.87) P value: Not sig</p>	<p>Funding: Cooperative Studies Programme of the Department of Veteran Affairs Medical Research Service</p> <p>Limitations: Randomisation allocation and concealment</p> <p>Additional outcomes:</p> <ul style="list-style-type: none"> ▪ Residual volume ▪ Perioperative complications: 5 perforation of capsule, 1 thrombophlebitis. 10 men found to have prostate cancer ▪ Factors predicting improvement, and influence of patient reported bother from urinary symptoms on outcomes of surgery and watchful waiting (see outcomes measure) <p>Notes: Related publication: Anon1993 published</p>
			<p>Symptom scores, mean (±SD) : Range: 0 to 27, (Madsen Iversen questionnaire) higher values more severe</p>	<p><u>At baseline</u> Group 1: 146±3.0 Group 2: 14.6±2.8 p value: Not sig <u>At 3 year follow up</u> Group 1: 4.9±4.0 Group 2: 9.1±4.7 p value: <u>Change from baseline</u> Group 1: -9.6±5.0 Group 2: -5.5±5.2 p value: <0.001</p>	
			<p>Qmax, mean (±SD) :</p>	<p><u>At baseline</u> Group 1: 11.6±6.4 Group 2: 12.5±7.5 p value: Not sig <u>At 3 year follow up</u> Group 1: 17.8±9.1 Group 2: 12.7±7.6 p value: <0.001 <u>Change from baseline</u> Group 1: 6.3±9.7 Group 2: 0.4±9.2 p value: <0.001</p>	
			<p>Perioperative complications: Recatheterisation</p>	<p>Group 1: 9/280 Group 2: 0/276 p value: <0.05*</p>	
			<p>Perioperative complications: transfusion</p>	<p>Group 1: 3/280 Group 2: 0/276 p value: Not sig</p>	
<p>Perioperative complications: Urinary</p>	<p>Group 1: 2/280</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>All patients A total of 800 patients screened 591 eligible for randomisations 30 did not provide informed consent, and 5 were found to be ineligible. N: 556 Drop outs: 71/556 [41/556 withdrew consent, 30 lost to follow up] Age, years, mean (±SD): 66±5</p> <p>Group 1 N: 280 Dropouts: 38/280, [24/280 withdrew consent, 14/280 lost to follow up] Age, years, mean (±SD): 65.6±5.2 White race, %: 91.4 **QoL scores, mean (±SD) :</p> <ul style="list-style-type: none"> ▪ Bother from urinary difficulties: 43.8±29.3 ▪ Sexual performance: 43.3±32.7 ▪ Activities of daily living: 66.5±27.2 ▪ General well being: 72.8±27.9 ▪ Social activities: 75.6±23.5 <p>Problems with dripping urine or wetting of plans: 46.0 Erective dysfunction: 60.7</p> <p>Group 2 N=276 Dropouts: 33/276 [17/276 withdrew consent, 16/276 lost to follow up] Age, years, mean (±SD):66.2±5.3 White race, %: 93.1 **QoL scores, mean (±SD) :</p> <ul style="list-style-type: none"> ▪ Bother from urinary difficulties: 	followed-up twice a year	<p>tract infection</p> <p>Incontinence (new persistent urinary incontinence requiring use of pads, clamps or condom)</p> <p>Treatment failure (Any of these events: death, repeated or intractable UTI, a residual volume of >350ml, development of bladder calculus, new urinary incontinence; a symptom score of ≥24 at one visit of a symptom score of ≥21 at 2 consecutive visits, doubling of baseline serum creatinine concentration)</p> <p>Reoperation/received surgery (in the watchful waiting arm) Reason: 9 bladder neck contracture, 9 urethral strictures, 8 received second TURP (4 due to adenoma). In the watchful waiting group: 20 treatment failure (11 high volume residual urine, 8 urinary symptoms, 5 intractable urinary retention)</p> <p>QoL scores - Bother from urinary difficulties, mean (±SD) :</p>	<p>Group 2: 0/276 p value: Not sig</p> <p><u>At 3 year follow up</u> Group 1: 4/280 Group 2: 4/276 Relative risk: 0.99(95% CI: 0.25- 3.90) P value: Not sig</p> <p><u>At 3 year follow up</u> Group 1: 23/280 Group 2: 47/276 Relative risk: 0.47 (95% CI: 0.29 to 0.72) p value: <0.05</p> <p><u>At 3 year follow up</u> Group 1: 26/280 Group 2: 65/276 Relative risk: 0.39 (95% CI: 0.26 to 0.60) p value: <0.05</p> <p><u>At baseline</u> Group 1: 43.8±29.3 Group 2: 46.3±29.3 p value: Not sig <u>At 3 year follow up</u> Group 1: 75.7±23.9 Group 2: 57.6±28.3 p value: <u>Change from baseline</u> Group 1: 29.6±29.4 Group 2: 9.6±29.7 p value: <0.001</p>	<p>the patient reported outcomes aspects</p> <p>Intention to treat analyses used. Data for all men, including those who had dropped out were analysed based on the group assigned.</p> <p>*Calculated by NCGC team using Fisher's exact test ** Score on a scale ranging from 0 (greatest impairment) to 100 (least impairment)</p> <p>Average period of follow up; 2.8 years</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>46.3±29.3</p> <ul style="list-style-type: none"> ▪ Sexual performance: 42.5±30.3 ▪ Activities of daily living: 69.0±26.6 ▪ General well being: 71.2±28.8 ▪ Social activities: 74.2±23.1 <p>Problems with dripping urine or wetting of plans: 44.4 Erective dysfunction: 63.7</p>		<p>QoL scores - Sexual performance: mean (±SD) :</p> <p>QoL scores - Activities of daily living: mean (±SD) :</p> <p>QoL scores - General well being: mean (±SD) :</p> <p>QoL scores - Social activities: mean (±SD) :</p>	<p><u>At baseline</u> Group 1: 43.3±32.7 Group 2: 42.5±30.3 <u>At 3 year follow up</u> Group 1: 36.0±26.0 Group 2: 35.6±25.6 <u>Change from baseline</u> Group 1: -3.0±27.9 Group 2: -3.2±26.6 p values: Not sig</p> <p><u>At baseline</u> Group 1: 66.5±27.2 Group 2: 69.0±26.6 p value: Not sig <u>At 3 year follow up</u> Group 1: 86.4±20.1 Group 2: 75.6±27.1 p value: <u>Change from baseline</u> Group 1: 19.6±26.5 Group 2: 6.4±30.3 p value: <0.001</p> <p><u>At baseline</u> Group 1: 72.8±27.9 Group 2: 71.2±28.8 <u>At 3 year follow up</u> Group 1: 76.2±27.8 Group 2: 71.4±31.0 <u>Change from baseline</u> Group 1: 3.0±25.5 Group 2: 0.1±28.3 p values: Not sig</p> <p><u>At baseline</u> Group 1: 75.6±23.5 Group 2: 74.2±23.1 <u>At 3 year follow up</u> Group 1: 75.5±25.3 Group 2: 73.1±25.5 <u>Change from baseline</u></p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				<p>Group 1: -1.6±24.3 Group 2: -1.7±23.5 p values: Not sig</p>	
			<p>Factors predicting improvement from bother from urinary difficulties at follow up (logistic regression, “improvement” not defined. Factors in model were baseline variables of bother from urinary difficulties, treatment assignment, age, symptom score, residual urinary volume, urinary volume after voiding, bladder trabeculation, Qmax)</p>	<p>2 factors were significant: Treatment assigned: odds ratio 5.7 (95% CI: 1.9 to 17.3) High bother score (>55) at baseline (for surgery group only, odds ration of 6.6(95% CI: 3.0 to 14.3) for surgery group, odds ratio of 1.4 (95% CI: 0.8 to 2.5) for watchful waiting group.</p> <p><u>In the TURP group, % improved</u> High bother: 134/148 (91%) Less bother: 45/73 (62%)</p> <p><u>In the watchful waiting group, % receiving surgery</u> High bother: 48/155 (31%) Low bother: 16/97(16%)</p>	
			<p>Association of symptom severity with QoL aspects (Perception of urinary difficulty(UD), sexual function (SF), Activities of daily living (ADL), general well being (GWB), Social activities(SA))</p>	<p>Nocturia: UD, ADL, GWB, Dribbling: UD Urgency: Sig for all Hesistancy: SF Frequency: UD, ADL, GWB, SA</p>	

Evidence Table 45: Bipolar transurethral resection of the prostate (TURP) vs. TURP

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Autorino et al., 2009²⁰</p> <p>De Sio et al., 2006⁶⁶ reported 12 month outcomes.</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 48 months</p>	<p>Patient group: men with LUTS including those with urinary retention from failed medical therapy</p> <p>Setting: Seconda Università di Napoli, Università magna Graecia, Catanzaro & Università Federico, Naples, Italy.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> >50 years AUR if catheter failed after medical therapy and CUR after unresponsiveness to medical therapy IPSS >18 Qmax < 15mL/s Prostate volume > 30 ml or higher than normal PSA <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or suspect Neurogenic bladder Bladder stone and/or diverticula Urethral stricture Maximum bladder capacity >500mL Previous prostate surgery Warfarin therapy <p>All patients N: 70</p> <p>Drop outs: 7 (refused follow-up=3; moved away=2; death, other causes=2)</p> <p>Group 1: N: 35 Mean age ± SD: 59.0 ± 5.9 Mean IPSS ± SD: 24.8 ± 4.0 Mean Qmax ± SD, mL/s: 7.1 ± 2.0 Mean PVR ± SD, mL: 80.0 ± 22.5</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop</p> <p>All patients: 26F resectoscope. Insertion of 22F 3-way Dufour catheter and irrigation with saline until urine was clear</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, QoL, Qmax, PVR, PSA assessed and follow up of IPSS, QoL, PVR and Qmax at 3, 6 12 months</p>	Mean ± SD IPSS at 3 months	Group 1: 8.0 ± NR (n=35) Group 2: 8.0 ± NR (n=35) P value: NR	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Allocation concealment not reported. Masking of IPSS and Qmax were not reported but catheterisation time was masked as primary outcome. 3 and 6 month outcomes estimated from graphs <p>Additional outcomes: Bladder irrigation time PVR at longer follow up periods.</p> <p>Notes: Randomisation sequence was computer generated</p> <p>NCC calculated average SD per arm from P values and means reported [from Cochrane handbook].</p>
			Mean ± SD IPSS at 6 months	Group 1: 5.0 ± NR (n=35) Group 2: 5.5 ± NR (n=35) P value: NR	
			Mean ± SD IPSS at 12 months	Group 1: 3.9 ± 3.32 (n=35) Group 2: 3.8 ± 3.32 (n=35) P value: 0.9	
			Mean ± SD IPSS at 24 months	Group 1: 4.5 ± 3.84 (n=33) Group 2: 4.8 ± 3.84 (n=34) P value: 0.75	
			Mean ± SD IPSS at 36 months	Group 1: 6.8 ± 5.19 (n=33) Group 2: 6.2 ± 5.19 (n=33) P value: 0.64	
			Mean ± SD IPSS at 48 months	Group 1: 6.9 ± 3.57 (n=32) Group 2: 6.4 ± 3.57 (n=31) P value: 0.58	
			Mean ± SD IPSS QoL at 3 months	Group 1: 2.1 ± NR (n=35) Group 2: 1.4 ± NR (n=35) P value: NR	
			Mean ± SD IPSS QoL at 6 months	Group 1: 1.1 ± NR (n=35) Group 2: 1.0 ± NR (n=35) P value: NR	
			Mean ± SD IPSS QoL at 12 months	Group 1: 1.0 ± 2.16 (n=35) Group 2: 0.8 ± 2.16 (n=35) P value: 0.7	
			Mean ± SD IPSS QoL at 24 months	Group 1: 1.1 ± 2.49 (n=33) Group 2: 1.2 ± 2.49 (n=34) P value: 0.87	
Mean ± SD IPSS QoL at 36 months	Group 1: 1.2 ± 1.27 (n=33) Group 2: 1.3 ± 1.27 (n=33) P value: 0.75				
Mean ± SD IPSS QoL at 48 months	Group 1: 1.3 ± 1.74 (n=32)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Mean prostate volume \pm SD, mL: 51.6 \pm 3.9 IPSS QoL \pm SD: 4.2 \pm 1.0 Operative time \pm SD, min: 49 \pm NR Resection time \pm SD, min: 33 \pm NR Resected weight (g): 20 \pm NR Drop outs: 3</p> <p>Group 2: N: 35 Mean age \pm SD: 61.0 \pm 5.9 Mean IPSS \pm SD: 24.38 \pm 5.0 Mean Qmax \pm SD, mL/s: 6.3 \pm 3.0 Mean PVR \pm SD, mL: 75.5 \pm 35.5 Mean prostate volume \pm SD, mL: 47.5 \pm 5.1 IPSS QoL \pm SD: 3.9 \pm 1.0 Operative time \pm SD, min: 53 \pm NR Resection time \pm SD, min: 39 \pm NR Resected weight (g): 24 \pm NR Drop outs: 4</p>			<p>Group 2: 1.4 \pm 1.74 (n=31) P value: 0.82</p>	
			Mean \pm SD Qmax at 3 months	<p>Group 1: 21.5 \pm NR (n=35) Group 2: 20.5 \pm NR (n=35) P value: NR</p>	
			Mean \pm SD Qmax at 6 months	<p>Group 1: 20.5 \pm NR (n=35) Group 2: 20.0 \pm NR (n=35) P value: NR</p>	
			Mean \pm SD Qmax at 12 months	<p>Group 1: 20.8 \pm 7.73 (n=35) Group 2: 22.3 \pm 7.73 (n=35) P value: 0.42</p>	
			Mean \pm SD Qmax at 24 months	<p>Group 1: 20.2 \pm 14.37 (n=33) Group 2: 22.0 \pm 14.37 (n=34) P value: 0.61</p>	
			Mean \pm SD Qmax at 36 months	<p>Group 1: 20.5 \pm 7.3 (n=33) Group 2: 21.5 \pm 7.3 (n=33) P value: 0.58</p>	
			Mean \pm SD Qmax at 48 months	<p>Group 1: 19.8 \pm 7.15 (n=32) Group 2: 21.2 \pm 7.15 (n=31) P value: 0.44</p>	
			Catheterisation time (days) converted into days	<p>Group 1: 3.0 \pm NR Group 2: 4.2 \pm NR P value: <0.05</p>	
			Length of stay (days) converted into days reported at time to discharge	<p>Group 1: 3.3 \pm NR Group 2: 4.5 \pm NR P value: <0.05.</p>	
			Complications: transfusion	<p>Group 1: 1/35 Group 2: 0/35 P value: NS</p>	
			Complications: TUR	<p>Group 1: 0/35 Group 2: 0/35 P value: NS</p>	
			Complications: urinary retention	<p>Group 1: 0/35 Group 2: 0/35 P value: NS</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Late complications at 48 months	Stricture Group 1: 1/32 Group 2: 2/31; p=0.6 Bladder neck contracture Group 1: 1/32 Group 2: 1/31; p=0.8 BPH recurrence Group 1: 1/32 Group 2: 1/31; p=0.8 Reoperation Group 1: 2/32 Group 2: 3/31; p=0.15	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bhansali et al., 2009³⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Men with BPH related LUTS that necessitated surgical intervention between May 2004 and December 2005.</p> <p>Setting: Institute of Urology in Pune, India</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> >45 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> AUA <18 Qmax >12 Gland size < 60g Neurologic illness Renal insufficiency, bladder stone Urethral stricture, prostate carcinoma Receiving 5AR inhibitors <p>All patients N: 70 Drop outs: 3</p> <p>Group 1: N: 35 Preop Qmax: 4.367 Gland size: 82.38 Mean age ± SD: NR</p> <p>Group 2: N: 35 Preop Qmax: 4.194 Gland size: 82.61 Mean age ± SD: NR</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) PK superpulse using 26F Gyrus Superpulse PK resectoscope and physiologic saline with 1% ethanol as irrigation fluid. Generator settings were 160 and 80 for cutting and coagulation, respectively.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop 26F resectoscope and an electro-surgical generator with glycine as irrigation fluid. Generator settings were 110 for cutting and 70 for coagulation.</p> <p>All patients: 500mg ciprofloxacin and 80mg gentamicin 1 hour preoperatively. All patients catheterised with 20F triple lumen Foley catheter at end of surgery, and irrigation started.</p>	<p>Mean (SD) Qmax at 3 months</p>	<p>Group 1 (n=34): 19.85 (3.939) Group 2 (n=33): 19.23 (5.176) P=0.582</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Dropouts not explained Allocation concealment method unclear <p>Notes: None.</p>
			<p>Mean (SD) Qmax at 9 months</p>	<p>Group 1 (n=34): 17.41 (2.840) Group 2 (n=33): 17.76 (3.269) P=0.645</p>	
			<p>Mean (SD) Qmax at 12 months</p>	<p>Group 1 (n=34): 16.6 (2.640) Group 2 (n=33): 15.9 (3.126) P=0.715</p>	
			<p>Mean (SD) Blood loss</p>	<p>Group 1 (n=34): 195.97 (50.079) Group 2 (n=33): 361.52 (97.599) P=0.000</p>	
			<p>Mean (SD) Time catheterised</p>	<p>Group 1 (n=34): 19.05 (3.920) Group 2 (n=33): 39.25 (10.223) P=0.000</p>	
			<p>Mean (SD) Hospital stay</p>	<p>Group 1 (n=34): 79.21 (14.251) Group 2 (n=33): 81.09 (15.438) P=0.605</p>	
			<p>Average tissue resected, g</p>	<p>Group 1: 42.8 Group 2: 45.0</p>	
			<p>Mean AUASS at baseline</p>	<p>Group 1: 26.3 Group 2: 24.6</p>	
			<p>Mean AUASS at 3 months</p>	<p>Group 1: 6.5 Group 2: 6.8</p>	
			<p>Mean AUASS at 9 months</p>	<p>Group 1: 8.2 Group 2: 8.0</p>	
			<p>Mean AUASS at 12 months</p>	<p>Group 1: 8.8 Group 2: 9.1</p>	
			<p>TUR</p>	<p>Group 1: 0% Group 2: 12.2%</p>	
			<p>Strictures</p>	<p>Group 1: 5 Group 2: 4</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Bladder neck contracture	Group 1: 1 Group 2: 0	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Erturhan et al., 2007⁸⁴</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: Patients with BPH and moderate to severe LUTS</p> <p>Setting: single centre: Sahinbey Medical Center, University of Gaziantep, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 18 or PVR > 50 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer or suspect Previous history of prostatic surgery Neurogenic bladder Urethral stricture <p>All patients N: 240 Dropouts: NR</p> <p>Group 1 N: 120 Mean age (range): 68.5 (52-90) Mean IPSS \pm SD: 25.0 \pm 5.0 Mean Qmax \pm SD, mL/s: 10.9 \pm 1.2 Mean PVR \pm SD, mL: 114 \pm 19 Mean prostate volume \pm SD, mL: 43 \pm 9 IPSS QoL \pm SD: 2.0 \pm 1.0 Operative time \pm SD, min: 36 \pm 19 Drop outs: 0</p> <p>Group 2 N: 120 Mean age (range): 67.4 (68-74) Mean IPSS \pm SD: 24.0 \pm 6.0 Mean Qmax \pm SD, mL/s: 9.29 \pm 1.7 Mean PVR \pm SD, mL: 135 \pm 25 Mean prostate volume \pm SD, mL: 42 \pm 11 IPSS QoL \pm SD: 3.0 \pm 1.0</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system with Plasma Sect electrode (200W, 160Ω, 320-450kHz, 254-350V) 27F continuous flow resectoscope with isotonic saline as irrigant</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop: 120W cutting and 80W coagulation. 26F continuous flow resectoscope with glycine 5% irrigant</p> <p>All patients 22 F Foley catheter inserted and irrigation with saline. Catheter removed when urine clear.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 1 and 12 months for IPSS, QoL, PVR and Qmax</p>	Mean \pm SD IPSS at 1 months	Group 1: 5.0 \pm 2.0 (n=120) Group 2: 5.0 \pm 2.0 (n=120) P value: NS	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment were not reported. Outcome assessment was not masked <p>Additional outcomes: Irrigation volumes.</p> <p>Notes: None.</p>
			Mean \pm SD IPSS at 12 months	Group 1: 4.0 \pm 2.0 (n=120) Group 2: 4.0 \pm 2.0 (n=120) P value: NS	
			Mean \pm SD Qmax at 1 months	Group 1: 17.4 \pm 2.5 (n=120) Group 2: 16.4 \pm 3.5 (n=120) P value: <0.001 P=0.01 calculated by NCGC using t-test with unequal variances	
			Mean \pm SD Qmax at 12 months	Group 1: 19.5 \pm 3.5 (n=120) Group 2: 18.5 \pm 3.0 (n=120) P value: <0.001 P=0.02 calculated by NCGC using t-test with unequal variances	
			Mean \pm SD QoL at 1 months	Group 1: 2.1 \pm 1.0 (n=120) Group 2: 2.1 \pm 1.0 (n=120) P value: NS	
			Mean \pm SD QoL at 12 months	Group 1: 2.1 \pm 1.0 (n=120) Group 2: 2.1 \pm 1.0 (n=120) P value: NS	
			Mean \pm SD catheter duration, days	Group 1: 3.0 \pm 1.1 (n=120) Group 2: 4.5 \pm 1.1 (n=120) P value: <0.001	
			Length of Stay \pm SD, days reported as time to discharge	Group 1: 3.0 \pm 1.2 (n=120) Group 2: 5.0 \pm 1.2 (n=120) P value: <0.001	
			Complications: Transfusion	Group 1: 1/120 Group 2: 7/120 P value: <0.0001	
			Complications: urinary retention (re-catheterisation)	Group 1: 2/120 Group 2: 5/120 P value: 0.083	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Operative time \pm SD, min: 57 \pm 24 Drop outs: 0		Complications: TUR Syndrome Complications: Reoperation rate Complications: Incontinence Complications: Mortality Complications: urethral & meatal stricture	Group 1: 0/120 Group 2: 2/120 P value: 0.15 Group 1: 0/120 Group 2: 5/120 P value: 0.025 Group 1: 0/120 Group 2: 0/120 Group 1: 0/120 Group 2: 0/120 Group 1: 5/120 Group 2: 4/120	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ho et al., 2007¹²⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months.</p>	<p>Patient Group: Patients awaiting TURP for failed medical therapy (alpha-blockers or 5-alpha reductase inhibitors), UTI or haematuria</p> <p>Setting: single centre: Department of Urology, Singapore General Hospital, Singapore</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> >50 years Fit for anaesthesia IPSS > 18 Qmax < 15 mL/s Patients with acute urinary retention and failed trial of voiding without catheter also included <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Previous prostatic surgery Neurogenic bladder disorders Bladder stones Renal impairment Hydronephrosis Prostate cancer or suspect Urethral strictures <p>All patients N: 100 Dropouts: 0</p> <p>Group 1 N: 48 Mean ± SD Age, yrs: 66.6 ± 6.8 IPSS ± SD: 22.6 ± 5.5</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Olympus TURIS system with 180W cutting and 100W coagulation</p> <p>Group 2: TURP Standard loop: 100W cutting and 50W coagulation with glycine 5% as irrigant.</p> <p>All patients 26F Olympus continuous flow resectoscope. 20F Foley 3-way catheter inserted for bladder irrigation and removed after 1 or 2 days.</p> <p>All operations performed by 2 senior consultants</p> <p>Examination methods Preoperative: Baseline IPSS, QoL, Qmax and PVR, PSA, Na⁺, creatinine and Hb. Postoperative: Na⁺, Hb repeated after 6 hours and IPSS and Qmax assessed at 1, 3, 6, 12 months follow up visits</p>	Mean ± SD IPSS at 3 months	Group 1: 9.0 ± NR (n=48) Group 2: 7.5 ± NR (n=52) P value: NS	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Allocation concealment not reported Outcome assessment was not masked Mean values are estimated from graph for IPSS and Qmax. P values were not provided for change from baseline so SDs could not be estimated <p>Additional outcomes: Decline in post op serum Na⁺ and Hb</p> <p>Notes: Computer randomisation</p>
			Mean ± SD IPSS at 6 months	Group 1: 7.0 ± NR (n=48) Group 2: 7.0 ± NR (n=52) P value: NS	
			Mean ± SD IPSS at 12 months	Group 1: 6.0 ± NR (n=48) Group 2: 6.0 ± NR (n=52) P value: NS	
			Mean ± SD Qmax at 3 months	Group 1: 19.5 ± NR (n=48) Group 2: 16.5 ± NR (n=52) P value: NS	
			Mean ± SD Qmax at 6 months	Group 1: 17.5 ± NR (n=48) Group 2: 18.0 ± NR (n=52) P value: NS	
			Mean ± SD Qmax at 12 months	Group 1: 17.0 ± NR (n=48) Group 2: 17.5 ± NR (n=52) P value: NS	
			Complications: Transfusion	Group 1: 1/48 Group 2: 1/52 P value: NS	
			Complications: TUR	Group 1: 0/48 Group 2: 2/52 P value: <0.05	
			Complications: urethral stricture	Group 1: 3/48 Group 2: 1/52 P value: NS	
			Complications: urinary retention (re-catheterisation)	Group 1: 5/48 Group 2: 4/52 P value: NS	
Complications: UTI	Group 1: 2/48 Group 2: 2/52 P value: NS				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p> PSA \pm SD, ng/mL: 2.8 \pm 1.0 Mean \pm SD Qmax, mL/s: 6.8 \pm 4.8 Mean prostate volume \pm SD, mL: 56.5 \pm 17.9 Resectate \pm SD, g: 29.8 \pm 11.2 Resection time \pm SD, min: 59 \pm 18 Number with AUR: 24/48 Number with failed medical therapy: 20/48 Number with UTI/Haematuria: 4/48 Dropouts: 0 </p> <p> Group 2 N: 52 Mean \pm SD Age, yrs: 66.5 \pm 7.2 IPSS \pm SD: 24.6 \pm 6.0 PSA \pm SD, ng/mL: 2.2 \pm 0.5 Mean \pm SD Qmax, mL/s: 6.5 \pm 3.2 Mean prostate volume \pm SD, mL: 54.8 \pm 19.2 Resectate \pm SD, g: 30.6 \pm 9.8 Resection time \pm SD, min: 58 \pm 16 Number with AUR: 21/52 Number with failed medical therapy: 25/52 Number with UTI/Haematuria: 6/52 Dropouts: 0 </p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
lori et al., 2008 ¹²⁴	<p>Patient Group: Patients scheduled for surgery for obstruction</p> <p>Setting: single centre: Department of Urology, University of Rome, Italy</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Obstruction class 2-5 on Schaefer nomogram <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Bladder stones Urethral stricture Renal insufficiency Current finasteride medical therapy <p>All patients N: 51 Dropouts: 0</p> <p>Group 1 N: 25 Mean age (range): 65.0 ± 5.0 Mean IPSS ± SD: 21.0 ± 2.0 Mean Qmax ± SD, mL/s: 7.0 ± 1.0 Mean PVR ± SD, mL: 99 ± 58 Mean prostate volume ± SD, mL: 49 ± 11 IPSS QoL ± SD: 3.0 ± 1.0 Resection time ± SD, min: 39 ± 19 Drop outs: 0</p> <p>Group 2 N: 26 Mean age (range): 63.0 ± 5.0 Mean IPSS ± SD: 20.0 ± 4.0 Mean Qmax ± SD, mL/s: 8.7 ± 2.0 Mean PVR ± SD, mL: 96 ± 97</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system with Plasma Sect electrode (200W, 160Ω, 320-450kHz, 254-350V) 27F continuous flow resectoscope with isotonic 0.9% saline as irrigant</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop. 26F continuous flow resectoscope with mannitol as irrigant</p> <p>All patients 22 F Foley catheter inserted and irrigation with saline. Catheter removed when urine clear and patient had passed a stool.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, QoL DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 12 months for IPSS, QoL, PVR and Qmax</p>	<p>Mean ± SD IPSS at 12 months</p> <p>Mean ± SD Qmax at 12 months</p> <p>Mean ± SD QoL at 12 months</p> <p>Mean ± SD catheter duration, days (converted from hours)</p> <p>Length of Stay ± SD, days (converted from hours)</p> <p>Complications: Transfusion</p> <p>Complications: urinary retention (re-catheterisation)</p> <p>Complications: TUR Syndrome</p>	<p>Group 1: 7.0 ± 1.7 (n=25) Group 2: 6.7 ± 4.0 (n=26) P value: NR</p> <p>Group 1: 24.2 ± 5.0 (n=25) Group 2: 23.2 ± 9.0 (n=26) P value: NR</p> <p>Group 1: 1.1 ± 1.0 (n=25) Group 2: 1.1 ± 1.0 (n=26) P value: NR</p> <p>Group 1: 0.96 ± 0.2 (n=25) Group 2: 1.33 ± 0.2 (n=26) P value: <0.0001</p> <p>Group 1: 2.0 ± 0.04 (n=25) Group 2: 2.1 ± 0.13 (n=26) P value: 0.9</p> <p>Group 1: 0/25 Group 2: 0/26</p> <p>Group 1: 1/25 Group 2: 0/26</p> <p>Group 1: 0/25 Group 2: 0/26</p>	<p>Funding: NR</p> <p>Limitations: None.</p> <p>Additional outcomes: Irrigation time, postoperative Schaefer obstruction class</p> <p>Notes: Randomisation by drawing opaque sealed envelopes.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean prostate volume \pm SD, mL: 48 \pm 91 IPSS QoL \pm SD: 3.6 \pm 1.0 Resection time \pm SD, min: 39 \pm 19 Drop outs: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Michielsen et al., 2007²⁰⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 month</p>	<p>Patient Group: Men with obstruction due to BPH</p> <p>Setting: single centre: Department of Urology, Virrije Universiteit, Brussels, Belgium</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS \geq 13 Qmax < 15 mL/s QoL \geq 3 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Carcinoma of the prostate History of prostate or urethral surgery Bladder stones Patients on anticoagulant therapy <p>All patients N: 238 Dropouts: 0</p> <p>Group 1 N: 118 Mean \pm SD Age: 73.8 \pm 8.1 (53-92) IPSS \pm SD: NR Mean SD Qmax: NR Mean prostate size \pm SD, g: NR Resectate \pm SD g: 21.0 \pm NR Operation duration \pmSD min: 56 \pm 25 Dropouts: 0</p> <p>Group 2 N: 50 Mean \pm SD Age: 73.1 \pm 8.6 (52-92) IPSS \pm SD: NR Mean SD Qmax: NR</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Olympus TURIS system with 270W cutting and 75W coagulation</p> <p>Group 2: TURP Standard loop with 26F resectoscope: 175W cutting and 75W coagulation</p> <p>All patients 22 F Foley catheter inserted and irrigation with saline until bleeding ended.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry.</p> <p>Postoperative: Full blood count was performed</p>	<p>Mean \pm SD catheter duration, days</p> <p>Mean \pm SD length of stay, days</p> <p>Complications: urinary retention (re-catheterisation)</p> <p>Complications: TUR Syndrome</p> <p>Complications: Transfusion</p> <p>Complications: reoperation (transurethral revision)</p>	<p>Group 1: 4.0 \pm 3.0 Group 2: 4.5 \pm 3.5 P value: 0.2</p> <p>Group 1: 4.9 \pm NR Group 2: 5.1 \pm NR P value: 0.6</p> <p>Group 1: 3/118 Group 2: 5/120</p> <p>Group 1: 0/118 Group 2: 1/120</p> <p>Group 1: 4/118 Group 2: 1/120</p> <p>Group 1: 0/118 Group 2: 2/120</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Unclear whether sealed envelopes were opaque. Primary outcome in study is not IPSS or Qmax Follow up very short to capture early complications only <p>Additional outcomes: Haemoglobin, sodium, potassium, chloride. Differences in operative times for staff v trainees</p> <p>Notes: None.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean prostate size \pm SD, g: NR Resectate \pm SD g: 21.3 \pm NR Operation duration \pm SD min: 44 \pm 20 Dropouts: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nuhoglu et al., 2006²²⁸</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 months</p>	<p>Patient Group: Patients with LUTS</p> <p>Setting: single centre: Ministry of Health Ankara Training & Teaching Hospital, Turkey</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS > 15 Qmax < 10 mL/s <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Carcinoma of the prostate History of prostate or urethral surgery Bladder stones Patients on anticoagulant therapy <p>All patients N: 57 Dropouts: 7 (5 patients could not be contacted, 1 died and 1 left study)</p> <p>Group 1 N: 27 Mean ± SD Age, years: 64.6 ± 8.8 IPSS ± SD: 17.6 ± 6.1 Mean SD Qmax: 6.9 ± 2.8 Mean SD PVR, mL: 96 ± 27 Mean prostate volume ± SD, mL: 47 ± 7.7 Operation duration ± SD min: 55 ± 9.7 Number of patients on alpha-blockers: 18/27 Dropouts: 3</p> <p>Group 2 N: 30 Mean ± SD Age, years: 65.2 ± 9.3</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system with Plasma Sect electrode (200W, 160Ω, 320-450kHz, 254-350V) Resection performed on PK3 mode with 340V. Saline irrigant</p> <p>Group 2: TURP 25F Storz resectoscope with glycine as irrigant.</p> <p>All patients All patients received antibiotic prophylaxis. 22 F Foley catheters inserted and continuous irrigation with saline for 1 postoperative day. Catheters removed when urine clear and discharge after free micturation.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, DRE, urinalysis, PSA, Blood, TRUS, uroflowmetry. Follow up at 1 and 12 months for IPSS, Qmax, PVR, and prostate volume. Complications were assessed at end of the first year.</p>	<p>Mean ± SD IPSS at 1 months</p>	<p>Group 1: 4.8 ± 3.4 (n=27) Group 2: 4.7 ± 3.1 (n=30) P value: NS</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method and allocation concealment were not reported Outcome assessment was not masked <p>Additional outcomes: Sodium, Haemocrit, Haemoglobin</p> <p>Notes: None.</p>
			<p>Mean ± SD IPSS at 12 months</p>	<p>Group 1: 5.4 ± 3.7 (n=24) Group 2: 5.2 ± 3.2 (n=26) P value: NS</p>	
			<p>Mean ± SD Qmax at 1 months</p>	<p>Group 1: 17.6 ± 4.3 (n=27) Group 2: 17.7 ± 2.3 (n=30) P value: NS</p>	
			<p>Mean ± SD Qmax at 12 months</p>	<p>Group 1: 17.1 ± 2.7 (n=24) Group 2: 17.9 ± 3.1 (n=26) P value: NS</p>	
			<p>Mean ± SD catheter duration, days (converted from hours)</p>	<p>Group 1: 1.96 ± 0.23 (n=27) Group 2: 3.15 ± 0.52 (n=30) P value: 0.009</p>	
			<p>Complications: Transfusion</p>	<p>Group 1: 1/27 Group 2: 2/30</p>	
			<p>Complications: urinary retention (re-catheterisation)</p>	<p>Group 1: 1/27 Group 2: 0/30</p>	
			<p>Complications: TUR Syndrome</p>	<p>Group 1: 0/27 Group 2: 0/30</p>	
			<p>Complications: Incontinence</p>	<p>Group 1: 0/27 Group 2: 0/30</p>	
			<p>Complications: Reoperation rate</p>	<p>Group 1: 0/27 Group 2: 0/30</p>	
<p>Complications: urethral stricture</p>	<p>Group 1: 1/27 Group 2: 0/30</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>IPSS \pm SD: 17.3 \pm 5.8 Mean SD Qmax: 7.3 \pm 2.1 Mean SD PVR, mL: 88 \pm 20 Mean prostate volume \pm SD, mL: 49 \pm 8.1 Operation duration \pm SD min: 52 \pm 13.2 Number of patients on alpha-blockers: 21/30 Dropouts: 4</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Patankar et al., 2006 ²³⁴	<p>Patient group: men with LUTS associated with BPH</p> <p>Setting: single-centre. Institute of urology & BJ Medical College, Pune, India</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> >45 years AUA score ≥ 18 Qmax < 10 mL/s Prostate volume 35-70 mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer Previous prostatic surgery <p>All patients N: 104 Drop outs: 1</p> <p>Group 1: N: 53 Mean age: 64 Mean AUA score \pm SD: 23.3 \pm 4.85 Mean Qmax \pm SD, mL/s: 5.9 \pm 1.98 Mean PVR \pm SD, mL: NR Mean prostate volume \pm SD, mL: 51.3 \pm 12.44 Operative time \pm SD, mins: 49.99 \pm 12.35 Resectate \pm SD, g: NR Drop outs: 1</p> <p>Group 2: N: 51 Mean age: 62 Mean AUA score \pm SD: 23.73 \pm 4.6 Mean Qmax \pm SD, mL/s: 6.4 \pm 1.77</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PK Superpulse system: Cutting 150V and 120V coagulation with saline irrigant.</p> <p>Group 2: Transurethral resection of the prostate (TURP) Standard loop through 24F resectoscope with glycine as irrigant</p> <p>All patients: Preoperative antibiotics. One consultant performed all the operations. A 20 3-way catheter was inserted and irrigation continued until returning fluid was clear for a minimum of 6 hours. Post irrigation catheter was removed if urine remained clear.</p> <p>Examination methods Preoperative: Baseline AUA score, urinalysis, PSA, TRUS, uroflowmetry. Uroflowmetry and AUA score repeated 21 days</p>	<p>Mean AUA score at 3 weeks</p> <p>Mean Qmax \pm SD mL/s at 3 weeks</p> <p>Catheterisation time (days) hours reported converted to days</p> <p>Complications: transfusion</p> <p>Complications: UTI</p>	<p>Group 1: 6.11 \pm 1.02 Group 2: 7.7 \pm 1.86 P value: NS</p> <p>Group 1: 19.16 \pm 1.9 Group 2: 20.67 \pm 1.63 P value: NS</p> <p>Group 1: 0.77 \pm 0.11 Group 2: 1.77 \pm 0.63 P value: <0.05</p> <p>Group 1: 0/53 Group 2: 1/51 p value: 0.5</p> <p>Group 1: 6/53 Group 2: 7/51 p value: 0.74</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Short follow up interval <p>Notes: Randomisation via drawing opaque envelopes</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean PVR ± SD, mL: NR Mean prostate volume ± SD, mL: 52.26 ± 10.71 Operative time ± SD, mins: 49.99 ± 12.35 Resectate ± SD, g: NR Drop outs: 0				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Seckiner et al., 2006 ²⁷⁰ Study design: RCT Observer masked Evidence level: 1+ Duration of follow-up: 12 months	Patient Group: Not specified Setting: single centre: Department of Urology, Zonguldak Karaelmas University School of Medicine, Turkey Inclusion criteria: <ul style="list-style-type: none"> IPSS \geq 8 Qmax < 15 mL/s Prostate volume 30-70g on TRUS Exclusion criteria: <ul style="list-style-type: none"> < 50 years Neurogenic bladder Carcinoma of the prostate or bladder History of prostate or urethral surgery On current medication known to affect voiding function All patients N: 48 Dropouts: 4 Group 1 N: 24 Mean \pm SD Age: 61.2 \pm 9.3 IPSS \pm SD: 24.1 \pm 5.2 IPSS QoL \pm SD: 4.4 \pm 0.6 Mean \pm SD Qmax, mL/s: 8.5 \pm 2.9 Mean PVR \pm SD, mL: 88 \pm 74 Mean prostate size \pm SD, mL: 49.4 \pm 18.9 Resectate \pm SD, g: 36.6 \pm 14.4 Operation duration \pm SD, min: 52.9 \pm 12.8	Group 1: Bipolar transurethral resection of the prostate (B-TURP) Gyrus PlasmaKinetic™ system with Plasma Sect electrode (200W, 160 Ω , 320-450kHz, 254-350V) set to 160W cutting and 80W. Resection performed through 27F resectoscope with saline as irrigant. Group 2: TURP Standard wire loop 120W cutting and 80W coagulation through 26F resectoscope with glycine 5% as irrigant All patients All operations were performed by the same surgeon. Bladder irrigation carried out for not more than 12 hours Examination methods Preoperative: Baseline IPSS Symptom score, QoL, DRE, urinalysis, blood, TRUS, uroflowmetry. IPSS and Qmax were recorded at 1, 3, 6 & 12 months, PVR at 3, 6 & 12 months and TRUS at 6 months.	Mean \pm SD IPSS at 3 months	Group 1: 9.3 \pm 3.9 (n=24) Group 2: 10.6 \pm 6.3 (n=24) P value: NS	Funding: NR Limitations: <ul style="list-style-type: none"> Allocation concealment with opaque sealed envelopes was not used Additional outcomes: Bleeding score, serum haemoglobin and sodium Notes: Randomisation using random number tables
			Mean \pm SD IPSS at 6 months	Group 1: 7.4 \pm 2.2 (n=24) Group 2: 6.0 \pm 6.7 (n=23) P value: NS	
			Mean \pm SD IPSS at 12 months	Group 1: 8.7 \pm 4.1 (n=23) Group 2: 8.3 \pm 2.9 (n=21) P value: NS	
			Mean \pm SD Qmax at 3 months	Group 1: 17.7 \pm 9.1 (n=24) Group 2: 18.6 \pm 9.1 (n=24) P value: NS	
			Mean \pm SD Qmax at 6 months	Group 1: 23.4 \pm 10.6 (n=24) Group 2: 16.2 \pm 12.0 (n=23) P value: NS	
			Mean \pm SD Qmax at 12 months	Group 1: 18.8 \pm 6.9 (n=23) Group 2: 15.7 \pm 6.3 (n=21) P value: NS	
			Mean \pm SD IPSS QoL at 3 months	Group 1: 1.8 \pm 1.0 (n=24) Group 2: 2.1 \pm 1.2 (n=24) P value: NS	
			Mean \pm SD IPSS QoL at 6 months	Group 1: 1.6 \pm 0.7 (n=24) Group 2: 1.6 \pm 1.3 (n=23) P value: NS	
			Mean \pm SD IPSS QoL at 12 months	Group 1: 1.8 \pm 0.8 (n=23) Group 2: 2.0 \pm 0.8 (n=21) P value: NS	
			Mean \pm SD catheter duration, days	Group 1: 3.1 \pm 0.6 Group 2: 3.1 \pm 1.4 P value: 0.98	
Complications: urethral stricture	Group 1: 2/24 Group 2: 1/24				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Dropouts: 1 patient where measurements were not obtained</p> <p>Group 2 N: 24 Mean \pm SD Age: 63.9 \pm 10.9 IPSS \pm SD: 23.2 \pm 4.9 IPSS QoL \pm SD: 4.7 \pm 0.9 Mean \pm SD Qmax, mL/s: 8.3 \pm 3.1 Mean PVR \pm SD, mL: 138 \pm 115 Mean prostate size \pm SD, mL: 41.4 \pm 14.5 Resectate \pm SD, g: 31.9 \pm 13.2 Operation duration \pm SD, min: 52.9 \pm 16.3 Dropouts: 3 patients where measurements were not obtained</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Singh et al., 2005²⁸⁰</p> <p>Study design: RCT Observer masked</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3 months</p>	<p>Patient Group: Patients with symptomatic BPH requiring surgical intervention</p> <p>Setting: single centre: Department of Urology, Muljibhai Patel Urological Hospital, Gujarat, India</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> >50 IPSS > 7 Qmax < 12 mL/s PCAR (from TRUS) >0.75 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Neurogenic bladder Renal insufficiency Bladder stone Urethral stricture Current finasteride therapy <p>All patients N: 60 Dropouts: NR</p> <p>Group 1 N: 30 Mean ± SD Age: 68.9 ± 7.6 IPSS ± SD: 20.5 ± 4.8 IPSS QoL ± SD: 4.6 ± 0.9 Mean ± SD Qmax, mL/s: 5.8 ± 3.0 Mean PVR ± SD, mL: 124 ± 58 Resectate ± SD, g: 24.0 ± 18.2 Operation duration ± SD, min: 39.3 ± 17.8 Number of patients with retention: 10/30 Dropouts: NR</p>	<p>Group 1: Bipolar transurethral resection of the prostate (B-TURP) ACMI Vista CTR Controlled Tissue Resection system through 25.6F resectoscope and cauterly setting of 6-8 for cutting and 7 for coagulation with saline as irrigant.</p> <p>Group 2: TURP Standard wire loop through 25.5F resectoscope.</p> <p>All patients All operations were performed by the same surgeon. A 20F 3-way catheter was placed and saline irrigation continued as required.</p> <p>Examination methods Preoperative: Baseline IPSS Symptom score, QoL, PCAR (TRUS), PSA, Blood, uroflowmetry.</p> <p>IPSS, QoL, Qmax at 1 and 3 months. Patients were given a questionnaire on postoperative complications on haematuria, dysuria, urgency, incontinence and pain weekly after surgery up to 4 weeks.</p>	<p>Mean ± SD IPSS at 3 months</p> <p>Mean ± SD Qmax at 3 months</p> <p>Mean ± SD IPSS QoL at 3 months</p> <p>Mean ± SD catheter duration, days</p> <p>Mean ± SD length of stay, days</p> <p>Complications: TUR</p> <p>Complications: UTI</p> <p>Complications: urethral stricture</p>	<p>Group 1: 5.3 ± NR Group 2: 6.2 ± NR P value: NR</p> <p>Group 1: 19.0 ± NR Group 2: 17.8 ± NR P value: NR</p> <p>Group 1: 1.1 ± NR Group 2: 1.0 ± NR P value: NR</p> <p>Group 1: 2.52 ± 0.5 Group 2: 3.41 ± 0.53 P value: 0.02</p> <p>Group 1: 3.02 ± 0.55 Group 2: 3.88 ± 0.58 P value: 0.02</p> <p>Group 1: 0/30 Group 2: 0/30</p> <p>Group 1: 3/30 Group 2: 4/30</p> <p>Group 1: 2/30 Group 2: 1/30</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Allocation concealment with opaque envelopes not clear. Unclear if all the patients completed study Standard deviations not reported for IPSS, Qmax or QoL and could not be estimated because there were p values for change from baseline <p>Additional outcomes: Haematuria, dysuria, urgency, incontinence and pain results from questionnaire.</p> <p>Notes: Randomised by drawing envelopes</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 N: 30 Mean \pm SD Age: 67.9 \pm 9.8 IPSS \pm SD: 21.6 \pm 6.3 IPSS QoL \pm SD: 4.47 \pm 1.0 Mean \pm SD Qmax, mL/s: 5.1 \pm 2.0 Mean PVR \pm SD, mL: 136 \pm 52 Resectate \pm SD, g: 27.6 \pm 13.4 Operation duration \pm SD, min: 36.9 \pm 14.6 Number of patients with retention: 11/30 Dropouts: NR</p>				

Evidence Table 46: Conservative vs. surgery

Bladder training vs. TURP

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Donovan et al., 2000 ⁷⁴	Patient group: men with uncomplicated LUTS symptoms	Group 1- Laser coagulation Procedure: Nd:YAG/ Non-contact VLAP, side-firing fibre (Bard Urolase), using standard fixed spot technique Power: 60W ND: YAG for 60s, depends on prostate size. For prostate size with urethral length of >25 mm, additional set of laser was used. If median lobe was present, 60W for 30s was applied for each side of lobe. Energy: 28684J Catheter protocol: Suprapubic catheter, removed when clinically appropriate. Other: All patients received antibiotic prophylaxis and anti-inflammatory suppository.	IPSS, mean change from baseline (95%CI): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: -10.8 ± 8.64* (95% CI: -12.5,-9.0), n=96 Group 2: -12.3 ± 7.36* (95% CI: -13.8,-10.7), n=89 Group 3: -1.3 ± 5.29* (95% CI: -2.8,0.2), n=85 p value: Group 2 v Group 3 - NR	Funding: Laser machines provided by Bard Diagnostics, Redmond, Washington. Limitations: • Open label study, with main outcomes using patient reported measures • The clinician following up patients was different to the surgeon although it was not stated whether the clinician was masked to treatment allocation
ClasP study	Setting: 3 centres in UK			Group 1: -1.9 ± 1.7* (95% CI: -2.3, -1.6), n=93 Group 2: -2.2 ± 1.62* (95% CI: -2.5, -1.8), n=85 Group 3: -0.4 ± 1.39* (95% CI: -0.7, -0.1), n=85 p value: Group 2 v Group 3 - NR	
Study design: RCT, multicentre, open label	Inclusion criteria: • IPSS score of ≥8, with physician and patient agreement that the symptoms require intervention • Qmax <15ml/s when voided volume >200ml, <13ml/s when voided volume between 150-200ml and <10ml/s when voided volume between 100 to 149ml measured on two occasions, with the higher value between these two used for analysis • >300ml post void volume urine on ultrasound		IPSS-QoL, mean (95%CI): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: 5.8 ± 6.87* (95% CI: 4.5, 7.2), n=102 Group 2: 9.7 ± 9.73* (95% CI: 7.7, 11.6), n=98 Group 3: 0.2 ± 2.9* (95% CI: -0.4, 0.8), n=92 p value: Group 2 v Group 3 - NR	Additional outcomes: Composite outcomes categories, and categorical outcomes for IPSS and Qmax
Evidence level: 1+	Exclusion criteria: • Prostate cancer or previous prostatic surgery; • prostate size > 120ml; • Life expectancy < 6 months; • Urinary retention associated with recent operation, constipation or drugs which could cause acute urinary dysfunction, • Neurogenic bladder	Group 2 –TURP Procedure: Standard electroresection Catheter protocol: Suprapubic catheter.	Qmax, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: -73.4(95% CI:-91.3, -55.5), n=100 Group 2: -74.0 (95% CI:-89.2, -58.8), n=98 Group 3: 2.19 (95% CI:-23.1, -27.5), n=90 p value: Group 2 v Group 3 - NR	Notes: Randomisation using computer generated numbers in blocks of 6 Allocation concealed using consecutive
Duration of follow-up: 7.5 months			Post void residual volume, mean(95%CI): Adjusted for centre and baseline symptom score, ANCOVA	Group 1: 5/117 Group 2: 0/117	
			All cause mortality Not treatment related		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>dysfunction;</p> <ul style="list-style-type: none"> Serum creatinine >250 µmol/L. <p>All patients N: 340 Drop outs:</p> <p>Group 1-Laser coagulation N: 117 Dropouts:1/117 Age, mean ± SD: 67.4 ± 8.1 IPSS, mean ± SD: 19.1 ± 6.6 IPSS-QoL, median(range): 4(2-6) Qmax, mean, ± SD: 10.4 ± 2.9 Post void residual urine, mean, ± SD: 123.7 ± 91.8 Prostate volume, mean, ± SD: 40.7 ± 21.4 No obstructed (%): 90/117 (78.3) No equivocal and/or unobstructed (%): 25/117 (21.7)</p> <p>Group 2 - TURP N: 117 Dropouts:2/117 Age, mean ± SD: 66.4 ± 7.9 IPSS, mean ± SD: 19.2 ± 6.7 IPSS-QoL, median(range): 4(0-6) Qmax, mean, ± SD: 10.3 ± 2.7 Post void residual urine, mean, ± SD: 104.2 ± 69.5 Prostate volume, mean, ± SD: 38.1 ± 19.1 No obstructed (%): 91/117(78.4) No equivocal and/or unobstructed (%): 25/117(21.6)</p> <p>Group 3 – Conservative management N: 106</p>	<p>Group 3 – Conservative management Procedure: Men were given general advice and bladder training as deemed clinically appropriate</p>	<p>Post-op complications: Blood transfusion (units and criteria not stated)</p> <p>Post-op complications: Perforation</p> <p>Post-op complications: Septicaemia</p> <p>Post-op complications: Urinary tract infection (symptomatic)</p> <p>Time to catheter removal geometric mean, days</p> <p>LOS, geometric mean (95% CI) days</p>	<p>Group 3: 1/106 p value: NS for all groups</p> <p>Group 1: 1/117 Group 2: 1/117 p value: NS</p> <p>Group 1: 0/117 Group 2: 2/117 p value: NS</p> <p>Group 1: 0/117 Group 2: 2/117 p value: NS</p> <p>Group 1: 3/117 Group 2: 2/117 p value: NS</p> <p>Group 1: 2.2 (95%CI 1.9 to 2.4) Group 2: 3.9 (95%CI 3.7 to 4.2) Relative risk: 1.83 95% CI: 1.58 to 2.11 P value: <0.0001</p> <p>Group 1: 11.8(95%CI: 10.2 to 13.7) Group 2: 2.4 (95%CI: 2.1 to 2.9) Relative risk: 4.79 95% CI: 3.88 to 5.91 p value: <0.0001</p>	<p>opaque sealed envelopes.</p> <p>Sample size calculation performed Please see Chacko et al., 2001⁴⁸ for the acute urinary retention population of CLASP trial and Gujral et al., 2000¹⁰⁷ for the chronic urinary retention population.</p> <p>* SD estimated using methods detailed in the Cochrane handbook for change from baseline with confidence intervals</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Dropouts: 5/106</p> <p>Age, mean \pm SD: 67.2 \pm 7.8</p> <p>IPSS, mean \pm SD: 18.8 \pm 6.5</p> <p>IPSS-QoL, median(range): 4(1-6)</p> <p>Qmax, mean, \pm SD: 9.9 \pm 2.7</p> <p>Post void residual urine, mean, \pm SD: 119.1 \pm 90.4</p> <p>Prostate volume, mean, \pm SD: 36.8\pm17.2</p> <p>No obstructed (%): 82/106(77.4)</p> <p>No equivocal and/or unobstructed (%): 24/106(22.6)</p>				

Catheters vs. TURP

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Ghalayini et al., 2005¹⁰⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with chronic urinary retention (CUR)</p> <p>Setting: 2 centres in Jordan and UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> IPSS >7 CUR defined by PVR > 300mL measured by ultrasonography on 2 occasions <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Prostate cancer Previous prostatic surgery Uncontrolled renal impairment Life expectancy <6 months Neurogenic bladder dysfunction Inability to perform clean intermittent self catheterisation. <p>All patients N: 51 Drop outs: 10</p> <p>Group 1 – CISC N: 29 (baseline variables for only 24 patients who completed the study) Age, mean (± SD): 69 ± 7.3 IPSS, mean (± SD): 23.2 ± 6.1 IPSS-QoL, mean (± SD): 4.2 ± 1.1 Qmax, mean (± SD), mL/s: 5.5 ± 4.2 PVR, mean (± SD), mL: 963 ± 503 Dropouts: 5 (3 withdrawn and 2 lost to follow up)</p>	<p>Group 2 – Clean intermittent self catheterisation (CISC) Patients were taught how to use a 12 or 14 F catheter every 6 hours.</p> <p>Group 1 – TURP Procedure: Standard electroresection</p> <p>Examination methods: Prior to start men had cystometry and PFS. Men were reviewed at 3 and 6 months after TURP or start of CISC for IPSS, serum creatinine, urine culture and PFS at 6 months. Men in the CISC group with urodynamic evidence of BOO at 6 months were advised to have TURP at the end of the study.</p>	<p>IPSS, mean change from baseline at 6 months (95%CI):</p> <p>IPSS QoL, mean change from baseline at 6 months (95%CI):</p>	<p>Group 1: -12.25 ± 7.77* (95% CI: -15.53,-8.97), n=24 Group 2: -20.29 ± 8.86* (95% CI: -24.85,-15.74), n=17 p value: NR</p> <p>Group 1: -2.54 ± 1.35* (95% CI: -3.11,-1.97), n=24 Group 2: -3.00 ± 1.46* (95% CI: -3.75,-2.25), n=17 p value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Randomisation method, allocation concealment and masking of outcome assessment were not reported. Complications were listed but not by group <p>Additional outcomes: At 6 months, PVR, voiding, end-filling and end-void pressures</p> <p>Notes: * SD estimated using methods detailed in the Cochrane handbook for change from baseline with confidence intervals</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2 - TURP N: 22 (baseline variables for only 17 patients who completed the study) Age, mean (\pm SD): 67 ± 8 IPSS, mean (\pm SD): 25.8 ± 4.2 IPSS-QoL, mean (\pm SD): 4.4 ± 0.9 Qmax, mean (\pm SD), mL/s: 5.2 ± 3.4 PVR, mean (\pm SD), mL: 954 ± 531 Dropouts: 5 lost to follow up</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Kadow et al., 1988¹³⁰</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with prostatism and proven BOO</p> <p>Setting: single-centre, UK</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Men with prostatism <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Haematuria Prostate cancer Normal peak flow rate and pattern after urodynamics <p>All patients N: 38 Drop outs: 0</p> <p>Group 1 – Conservative N: 17 Age, mean (\pm SD): 64.5 \pm NR Qmax, mean (\pm SD), mL/s: 9.8 \pm 2.1 PVR, mean (\pm SD), mL: 115 \pm 305 Day-time frequency, mean \pm SD: 8.25 \pm 11.34 Nocturia, voids \pm SD: 1.7 \pm 4.6 Dropouts: 0</p> <p>Group 2 - TURP N: 21 Age, mean (\pm SD): 66.5 \pm NR Qmax, mean (\pm SD), mL/s: 8.5 \pm 9.53 PVR, mean (\pm SD), mL: 86.2 \pm 369 Day-time frequency, mean \pm SD: 7.76 \pm 16.59 Nocturia, voids \pm SD: 2.6 \pm 5.6 Dropouts: 0</p>	<p>Group 2 – Conservative treatment Instruction on bladder training for 1 month consisting of weekly visits of encouragement to increase interval between day-time voids and reduce fluid intake < 1 litre/day. Advice on timing was given to those with nocturia. Frequency/volume charts were analysed at each visit. Those with bladder instability after a cystometrogram at the end of training were given Pro-Banthine for urgency symptoms (10 patients). All patients were encouraged to continue bladder training throughout 6 month period</p> <p>Group 1 – TURP Procedure: Standard electroresection with histological conformation of BPH</p> <p>Examination methods: Prior to start men completed a frequency/volume chart for 7 days then voiding water cystometry. Reassessment after 6 months</p>	<p>Q max \pm SD at 6 months</p>	<p>Group 1: 11.2 \pm 3.42, n=17 Group 2: 19.0 \pm 4.08, n=21 p value: NR</p>	<p>Funding: NR</p> <p>Limitations:</p> <p>Additional outcomes: Voiding patterns, day time frequency, nocturia, Max voided volume, average voided volume, maximum intervals between voids, P det max, PVR after treatment.</p> <p>Notes: Marked cards in identical envelopes were used for randomisation</p>

Evidence Table 47: What is the effectiveness of alpha-blockers in treating men after acute urinary retention?

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Lucas et al., 2005¹⁷⁶</p> <p>Study design: Randomised controlled study</p> <p>Setting: 8 hospitals and one in Ireland.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 3-8 days depending on normal practice of hospital.</p>	<p>Patient group: Men with acute urinary retention (AUR) secondary to benign prostatic hyperplasia recruited from March 1997 to December 2000 from an Accident and Emergency department. .</p> <p>Inclusion criteria: Men with acute urinary retention, who had been catheterised in the previous 72 hours.</p> <p>Exclusion criteria: Men with initial catheterisation volumes of >1500mL or <500mL ; evidence of renal or hepatic dysfunction; previous surgery on the urinary tract; other diseases of the bladder; any malignancy; retention-enhancing medications; allergies; and sever cardiac disease.</p> <p>All patients N: 149 Mean age: 69.4 (range: 51-91) years Drop outs: 8 not evaluable and not included in ITT analysis.</p> <p>Group 1 N: 71 Mean (\pmSD) Age: NR Dropouts: NR</p> <p>Group 2 N: 70 Mean (\pmSD) Age: NR Dropouts: NR</p>	<p>Group 1: Alpha-blocker Tamsulosin hydrochloride 0.4mg in a modified-release capsule once daily. Medication given after breakfast or lunch on the first dose, then after each day's breakfast. Duration of treatment was decided by each site to be either three or 8 doses, according to their normal practice.</p> <p>Group 2: placebo</p>	<p>Successful trial without catheter (defined as a flow rate of >5mL/s, >100mL voided volume, and a residual volume of \leq200mL)</p>	<p>Group 1: 24/71 (34%) Group 2: 17/70 (24%) p value: 0.193</p>	<p>Funding: Sponsored by a grant from Yamanouchi Pharma Ltd.</p> <p>Limitations: None</p> <p>Notes: Definition of success in treatment of AUR has yet to be universally agreed. The initial definition was not significant but the authors conducted secondary analysis using revised criteria of success. This was completed before breaking randomisation code.</p> <p>Some patients were catheterised for 3 day and others for 8; to allow for variations in practice across the sites. Differences in outcome between the two were not statistically significant.</p>
			<p>Secondary analysis: (success defined as any of two free-flow criteria described above)</p>	<p>Group 1: 41/71 (58%) Group 2: 28/70 (40%) p value: 0.02</p>	
			<p>Secondary analysis: Success defined as flow rate >5mL/s, voided volume >100mL</p>	<p>Group 1: 37/71 (52%) Group 2: 24/70 (34%) p value: 0.019</p>	
			<p>Secondary analysis: (defined as a flow rate of >5mL/s, >100mL voided volume, and a residual volume of \leq250mL)</p>	<p>Group 1: 43/71 (61%) Group 2: 29/70 (41%) p value: 0.013</p>	
			<p>Patients not re-catheterised</p>	<p>Group 1: 34/71 (48%) Group 2: 18/70 (26%) p value: 0.011 OR: 2.47, 95% CI: 1.23-4.97</p>	
			<p>Patients re-catheterised</p>	<p>Group 1: 37/71 (52%) Group 2: 52/70 (74%)</p>	
			<p>Adverse events</p>	<p>Dizziness Group 1: 7/71 (10%) Group 2: 2/70 (3%)</p> <p>Somnolence Group 1: 4/71 (6%) Group 2: 2/70 (3%)</p> <p>Mortality (carcinomatosis; not due to intervention) Group 1: 1/71 (1%) Group 2: 0/70 (0%)</p>	
<p>Patients withdrew due to adverse events</p>	<p>Group 1: 7 (9%) Group 2: 1 (1%)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
McNeill et al., 1999 ¹⁹³ Study design: Randomised controlled trial Setting: Scotland (4 centres) Evidence level: 1+ Duration of follow-up: Treatment for 48 hours. Follow-up of successful patients for mean 7.2 months	Patient group: patients with a first episode of acute urinary retention related to benign prostatic obstruction were recruited between September 1996 and March 1998 from 4 centres in Scotland. Inclusion criteria: 55 years or over; residual volume of 0.5-1.5L on catheterisation. Exclusion criteria: patients unwilling or unable to give informed consent; significant renal and/or hepatic disease; depressive illness on medication; extra-pyramidal disorders; neurological disease; confirmed or suspected urethral stricture; dipstick detected UTI, acute or chronic prostatitis. History of unstable angina pectoris, myocardial infarction, transient ischaemic attacks, cerebrovascular accident of congestive cardiac failure during the previous 6 months, current or previous orthostatic hypotension. Patient taking monoamine oxidase inhibitors, cholinergic or anticholinergic drugs, calcium-channel blockers, or alpha blocking drugs. Other antihypertensive drugs were not altered whilst the patient was receiving the trial medication. Phytotherapy or finasteride use did not exclude patients from study but their use was recorded. Known hypersensitivity to afluzosin or alpha blockers. Patients requiring	Group 1: alpha-blocker Sustained-release alfuzosin, an alpha1-selective blocker, (5mg twice daily, with no dose titration) for 48 hours. Catheter removed after 24 hour of treatment and final dose was given on the afternoon after catheter removal. Group 2: placebo Identical procedure as intervention but with placebo (twice daily for 48 hours).	Number (%) of patients successful: (defined as able to void successfully after removal of catheter and not re-catheterised within 24h) Number (%) of patient successful using per-protocol analysis (excluding patient that withdrew and failed to complete medication) Mean (SD) age for all patients: Mean (SD) age by success in each group: Logistic regression analysis of treatment versus outcome adjusted for age Logistic regression using per-protocol analysis: All reported adverse events	Group 1: 22/40 (55%) Group 2: 12/41 (29%), P=0.034 Odds Ratio (OR): 2.95 (95% CI 1.08-8.21) Group 1: 22/39 (56%) Group 2: 12/41 (29%), P=0.026 Odds Ratio (OR): 3.13 (95% CI 1.13-8.76) Successful: 68.4 (7.8) Unsuccessful: 72.9 (8.1) P=0.02 Group 1: Successful: 69.1 (8.7) Unsuccessful: 69.6 (7.3), p=0.81 Group 2: Successful: 67.2 (6.1) Unsuccessful: 75.0 (8.1), p=0.005 P=0.052 OR: 2.55, 95% CI 0.99-6.58 P=0.039 OR: 2.72, 95% CI 1.05-7.08 Faint: Group 1: 1/40 Group 2: 0/41 Dizziness: Group 1: 1/40 Group 2: 0/41 Headache: Group 1: 1/40 Group 2: 0/41	Funding: Financial support for the study was received from Lorex Synthelabo UK & Ireland; authors received financial support from Lorex Synthelabo to attend and present their work at scientific meetings. Limitations: The mean age was 5 years lower in the intervention group (significant difference). Following power calculation the authors planned to recruit 100 per arm to detect a 20% difference in outcome with 95% power. Unable to reach this number before the trial medication expired. The difference in outcome between the groups was >20% and power of the study is reflected in statistical significance of the results. Additional outcomes: Comparison of variables between successful and unsuccessful patients. Non significant results

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>suprapubic catheterisation where urethral catheterisation was unsuccessful; patients who had a suprapubic catheter as a primary procedure were not excluded. Postoperative retention after major abdominal/pelvic surgery. Large residual volume, clot retention secondary to haematuria of any cause.</p> <p>All patients N: 81</p> <p>Group 1 N: 40 Mean (\pmSD) Age: 67.7 (13.6) Dropouts: 1 (withdrew following a faint after the first dose of the trial medication)</p> <p>Group 2 N: 41 Mean (\pmSD) Age: 72.7 (8.33) Dropouts: 0</p>			<p>Atrial fibrillation* Group 1: 1/40 Group 2: 0/41</p>	<p>for mean residual volume on catheterisation, mean duration of catheterisation and prostate size.</p> <p>Additional follow-up of 11/34 (32%) successful patients experiencing a further episode of AUR and/or requiring a prostatectomy (mean follow-up of 7.2 months).</p> <p>Notes: Atrial fibrillation 8 hours after last dose, which was later resolved. A subsequent 24-h ECG revealed previously undiagnosed asymptomatic paroxysmal atrial tachycardia, which was treated with sotalol.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>McNeill et al., 2004¹⁹⁴</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: 71 centres across Europe and South Africa.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: Treatment for 3 days.</p>	<p>Patient group: patients presenting with a first episode of spontaneous AUR related to BPH between January 2000 and March 2002.</p> <p>Inclusion criteria: Minimum age of 51 yrs; urine retention volume 500-1500ml at catheterisation</p> <p>Exclusion criteria: Patients with mental disorders, in a trial within last 3 months, patients with neurogenic bladder dysfunction, isolated bladder neck disease, prostatitis, carcinoma of prostate, history of prostatic and urethral surgery, urethral stricture, bladder stones, clot retention secondary to hematuria; residual volume <500ml or >1500ml, AUR not related to BPH; Parkinson's disease, insulin dependent diabetes, multiple sclerosis, stroke or myocardial infarction within last 6 months, hepatic abnormalities, unstable or severe heart failure, history of postural hypotension or syncope, hypersensitivity to α-blockers, evolutive neoplastic disease; patients who received sympathomimetics within the previous week, received 5α-reductase inhibitors within previous 3 months or α-blocker in previous month, received tricyclic antidepressants, anticholinergics, sympathomimetics or first generation antihistamines within previous months, patients receiving disopyramide.</p> <p>All patients: N: 363</p> <p>Drop outs: 3 (results missing)</p> <p>Group 1: N: 238 Mean (\pmSD) Age: 69.3 (8.5) Dropouts: 4 (postural hypotension=2, catheter related infection=1 and treatment unrelated haemorrhoids=1)</p>	<p>All patients: urethral bladder catheterisation was performed. Catheter removed after minimum of two doses of study drug and each patient received one additional tablet the day after catheter removal.</p> <p>Group 1: Alpha-blocker 10mg alfuzosin once daily for three days</p> <p>Group 2: Placebo Once daily for three days.</p>	<p>Success (defined as patient returned to satisfactory voiding within the first 24 hours following removal of the urethral catheter without re-catheterisation)</p> <p>Number of patients experiencing at least one adverse event</p>	<p>Group 1: 146/236 (61.9%) Group 2: 58/121 (47.9%) p value: 0.012</p> <p>Group 1: 20/238 (8.4%) Group 2: 16/122 (13.1%)</p>	<p>Funding: NR.</p> <p>Limitations: Breakdown of adverse events not listed.</p> <p>Additional outcomes: Logistic regression analysis of successful trial without catheter. Age 65 years plus and drained volume 1000ml or greater adversely influenced the successful voiding rate.</p> <p>Backward multiple logistic regression.</p> <p>Notes: Randomisation in a 2:1 ratio for intervention: placebo.</p> <p>Extension study carried out following patients that had a successful trial without catheter.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2: N: 122 Mean (\pmSD) Age: 69.4 (8.0) Dropouts: 1 (catheter related infection)</p>				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Shah et al., 2002²⁷²</p> <p>Study design: Randomised controlled trial</p> <p>Setting: St Lukes Hospital and Bradford Royal infirmary, UK</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 2 weeks for primary study and follow up of successful patients at 2 years.</p>	<p>Patient group: patients presenting with acute urinary retention at the hospital between March 1998 and December 1999.</p> <p>Exclusion criteria: patients with cardiac disease contra-indicating the use of alpha blockers, receiving medical therapy for bladder outflow obstruction, patients with bladder calculi, prostate cancer, renal impairment, urethral stricture, urinary infection, neurogenic bladder dysfunction, bladder tumour and clot retention.</p> <p>All patients N: 81 Mean age: 68.6 (46-88) years Drop outs: 19 (urethral stricture=1, patient request for removal=9, adverse events=1, other reasons including suprapubic catheter, aortic aneurysm and other severe co-morbidity=8)</p> <p>Group 1 N: 34 Mean (\pmSD) Age: 69.5 (56-88) Dropouts: 0</p> <p>Group 2 N: 28 Mean (\pmSD) Age: 67.7 (46-84) Dropouts: 0</p>	<p>Group 1: Alpha-Blocker Alfuzosin SR 5mg twice a day. Catheter removed after a minimum of three doses or 36 hours of admission.</p> <p>Group 2: Placebo Catheter removed after a minimum of three doses or 36 hours of admission.</p> <p>All patients: if trial without catheter was unsuccessful a second trial was given 2 weeks later. During this period patients continued their trial medication. If unsuccessful again patients were offered alternative treatment options.</p>	<p>Successful voiding (defined as being able to void with a residual volume of < 200ml)</p>	<p>Group 1: 17/34 (50%) Group 2: 16/28 (57%) OR: 0.86 (95% CI: 0.38, 1.98; p=0.72)</p>	<p>Funding: Lorex Synthelabo Pharma</p> <p>Limitations: Method of randomisation and allocation concealment not reported. Baseline characteristics not addressed except for age.</p> <p>Additional outcomes: Additional outcomes for patients that had an unsuccessful trial without catheter and were given alfuzosin.</p> <p>Notes: The mean age and range at baseline was lower in the placebo group.</p>
			<p>Unsuccessful voiding and re-catheterised</p>	<p>Group 1: 17/34 (50%) Group 2: 12/28 (43%)</p>	
			<p>TURP following successful trial without catheter (open labelled study where all patients on alfuzosin)</p>	<p>Year 1: 13/30 (43%) Year 2: 6/15 (40%)</p>	

Evidence Table 48: Phytotherapy vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Bent et al., 2006²⁹</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Northern California, US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: Men who had moderate to severe symptoms of benign prostatic hyperplasia. Recruited from San Francisco Veterans Affairs Medical Center and the surrounding area by direct mailings, letters to primary care providers, posters and newspapers and local radio adverts between July 2001 and May 2004.</p> <p>Inclusion criteria: Over 49 years, AUA of 8 or more, peak urinary flow rate <15ml/s. Eligible if had stopped taking alpha-blocker at least one month before randomisation or discontinued taking saw palmetto or a 5 alpha-reductase inhibitor 6 months before randomisation.</p> <p>Exclusion criteria: high risk for urinary retention; history of prostate cancer; surgery for BPH; urethral stricture or neurogenic bladder; had a creatinine level >2.0mg per decilitre; PSA >4ng; using medications known to affect urination; severe concomitant disease.</p> <p>All patients N: 225</p> <p>Group 1 N: 112 Mean (±SD) Age: 62.9 (8.0) Dropouts: 5 Discontinued medication: 5 (outcomes assessments completed)</p> <p>Group 2 N: 113</p>	<p>All patients: One month placebo run in period – excluded if rate of adherence was <75%.</p> <p>Group 1: Saw palmetto extract (160mg twice a day with meals) Carbon dioxide extract in a soft gelatine capsule – manufactured in one batch for product consistency.</p> <p>Group 2: Placebo Similar appearing placebo in soft brown gelatine capsules. Twice a day with meals.</p>	<p>Mean (SE) change in AUA symptom index score</p>	<p>Group 1 (n=112): -0.68 (0.35) [95% CI: -0.37 to 0.01] Group 2 (n=113): -0.72 (0.35) [95% CI: -1.40 to -0.04] Difference=0.04 [-0.93 to 1.01]</p>	<p>Funding: Grant from the national institute of diabetes and digestive and kidney diseases and by a grant from the National Centre for Complementary and Alternative medicine.</p> <p>Limitations: BPH impact score significantly different at baseline.</p> <p>Additional outcomes: Prostate transitional zone volume, BPH impact index score reported. Subgroup analyses of AUASI outcome when stratified by varying baseline levels.</p> <p>Notes: Most commonly reported nonserious adverse events also reported – no significance difference between the groups.</p>
			<p>Mean (SE) difference maximum urinary flow rate, ml/min</p>	<p>Group 1: 0.42 (0.34) Group 2: -0.01 (0.34) Difference=-1.22 [-3.90 to 1.47]</p>	
			<p>Mean (SE) Prostate volume (ml)</p>	<p>Group 1: 3.76 (0.98) Group 2: 4.98 (0.96) Difference=0.43 [-0.52 to 1.38]</p>	
			<p>Mean (SE) residual volume, ml</p>	<p>Group 1: 14.10 (7.24) Group 2: 18.62 (7.14) Difference=-4.51 [-24.44 to 15.42]</p>	
			<p>SF-36 score (scores range from 0-100; higher scores indicate better quality of life)</p>	<p>Mental subscale: Group 1: -0.72 (0.72) Group 2: 0.47 (0.71) Difference=-1.18 [-3.16 to 0.79]</p> <p>Physical subscale: Group 1: 0.10 (0.67) Group 2: -0.51 (0.66) Difference=0.61 [-1.24 to 2.45]</p>	
			<p>Sexual function (O'Leary scale) range from 0-4; with higher scores indicating better function</p>	<p>Group 1: -0.06 (0.10) Group 2: 0.07 (0.10) Difference=-0.13 [-0.40 to 0.14]</p>	
			<p>Serious adverse events</p>	<p>cardiovascular Group 1: 2 Group 2: 7</p> <p>Elective orthopaedic surgery Group 1: 3 Group 2: 3</p> <p>Gastrointestinal bleeding Group 1: 2</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Mean (\pm SD) Age: 63.0 (7.4) Dropouts: 4 Discontinued medication: 5 (outcomes assessment completed)			Group 2: 1 Bladder cancer Group 1: 0 Group 2: 1 Colon cancer: Group 1: 0 Group 2: 1 Elective hernia repair Group 1: 0 Group 2: 1 Hematoma Group 1: 0 Group 2: 1 Melanoma Group 1: 1 Group 2: 0 Prostate cancer Group 1: 0 Group 2: 1 Shortness of breath Group 1: 0 Group 2: 1 Rhabdomyolysis Group 1: 0 Group 2: 1 Total Group 1: 8/112 (n=6) Group 2: 18/113 (n=11)	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Safarinejad et al., 2005²⁶⁵</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Iran</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with LUTS due to BPH, 1-3 years in duration presenting to the outpatient urology clinic.</p> <p>Inclusion criteria: no cancer laboratory findings were normal; and patient had no lower urinary tract problem other than BPH.</p> <p>Exclusion criteria: loss to follow-up, surgical intervention for BPH, discontinuation of study medication; alpha blocker, 5-alpha reductase inhibitor or other drug therapy during trial and follow-up, any combination of Urtica dioica with other phototherapeutic agent and insufficient follow-up.</p> <p>All patients: N: 620</p> <p>Group 1 N: 305 Completed by: 287 Mean (range) Age: 64 (57-71) Dropouts: 36; follow-up=25, surgical intervention =5, medication discontinued=2, other pharmacological treatment=4</p> <p>Group 2 N: 315 Completed by: 271 Mean (range) Age: 62 (53-73) Dropouts: follow-up=36, surgical intervention =14, medication discontinued=10, other pharmacological treatment=9</p>	<p>Group 1: Urtica dioica 120mg three times daily Herbal blend contained a standard preparation of 100mg of urtica dioica root extract in 1ml. Ingested three times daily with meals.</p> <p>Group 2: placebo three times daily placebo.</p>	<p>Mean (SD) IPSS</p>	<p>Baseline Group 1: 19.8 (4.9) Group 2: 19.2 (4.6) 6 months Group 1: 11.8 (4) Group 2: 17.7 (3.1)</p>	<p>Funding: NR</p> <p>Limitations: Number completed trial was used for analysis. Reasons for drop-outs gives different total number of dropouts but this may have included the extension study.</p> <p>Additional outcomes: Serum PSA and serum testosterone also reported.</p> <p>Notes: After the 6 month randomised trial placebo patients were switched to the active treatment until 18 months.</p>
			<p>Mean (SD) Qmax (mL/s)</p>	<p>Baseline Group 1: 10.7 (2.4) Group 2: 10.8 (2.8) 6 months Group 1: 18.9 (4.7) Group 2: 14.2 (3.7)</p>	
			<p>Mean (SD) PVR, mL</p>	<p>Baseline Group 1: 73 (32.6) Group 2: 74 (29.6) 6 months Group 1: 36 (25.5) Group 2: 71 (24.4)</p>	
			<p>Mean (SD) Prostate volume, cc</p>	<p>Baseline Group 1: 40.1 (6.8) Group 2: 40.8 (6.2) 6 months Group 1: 36.3 (4.2) Group 2: 40.6 (5.1)</p>	
			<p>Patients reporting improved LUTS</p>	<p>Group 1: 232/287 (86%) Group 2: 43/271 (16%) P<0.001</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Shi et al., 2008 ²⁷⁴ Study design: Randomised controlled trial Setting: China Evidence level: 1+ Duration of follow-up: 12 weeks	Patient group: men between 49-75 years old with newly diagnosed LTS associated with BPH based on urological symptoms, including nocturia, incomplete emptying, urinary frequency, intermittence, weak urine stream, straining and urgency. Inclusion criteria: digital rectal examination showing an enlarged prostate but no signs of prostate cancer, serum creatinine >160umol/l, bacterial count less than 1000,000/ml, serum PSA 4ng/ml or less, IPSS greater than 12, uroflowmetry with MFR no more than 15ml per second and voiding volume greater than 150ml. Urinalysis by dipstick and microscopic examination of the spun urine specimen were performed to rule out urinary tract infection or hematuria. All patients had refused conventional therapy or elected watchful waiting. Exclusion criteria: history of prostate cancer and the use of any drugs, herbs or other non-prescription preparations for LUTS associated with BPH within 4 weeks of screening, including finasteride, alpha or beta blockers, diuretics, calcium channel blockers and anticholinergic drugs. Abnormal lab parameters, including PSA>4, serum	Group 1: 2 Prostataplex soft gels daily Group 2: 2 placebo soft gels daily	Mean (SD) IPSS Number of patients with an IPSS improvement (defined as decrease of 3 points or greater) Mean (SD) Qmax, ml/s Mean (SD) Relative urinary resistance Mean (95%CI) Blood urea nitrogen at 12 weeks mg/dl Mean (95% CI) Prostate size, cm3 Mean (95% CI) PSA, ng/ml	Baseline Group 1: 16.85 (6.48) Group 2: 14.46 (4.32) 12 weeks: Group 1: 14.83 (6.42) Group 2: 14.13 (4.25) Group 1: 18/46 (39.1%) Group 2: 1/46 (2.2%) P<0.001 Baseline Group 1: 12.40; 95%CI:11.90-12.89 Group 2: 12.89; 95% CI: 2.22-13.56 12 weeks: Group 1: 14.07 (2.56) Group 2: 11.74 (1.23) P<0.001 Baseline Group 1: 2.97; 95% CI: 2.60-3.35 Group 2: 2.88; 95%CI: 2.57-3.19 12 weeks: Group 1: 2.35 (0.83) Group 2: 3.02 (1.18) P=0.002 Group 1: 3.872 (3.426-4.318) Group 2: 3.809 (3.414-4.203) P=0.832 Group 1: 45.62 (43.85-47.39) Group 2: 45.90 (44.04-47.76) P=0.826 Group 1: 1.845 (1.617-2.073) Group 2: 1.694 (1.505-1.882)	Funding: NR. Limitations: Significant baseline difference in IPSS scores (lower in placebo group) Baseline IPSS for control was reported differently in the text as 14.46 and 14.27. Additional outcomes: Compliance rates reported as > 95% for both groups at each time point. Notes: Prostataplex, contains mainly saw palmetto.

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>creatinine >160umol/l, urine bacterial count>100,000/ml, BUN more than 8mg/dl, MFR >15ml/s and voiding volume <150ml, previous bladder or prostate surgery, micturition problems associated with identified bladder pathology, urethral stricture, recurrent urinary tract infections, known renal or hepatic or cardiac insufficiency, diabetes mellitus, recent myocardial infarction, known alcohol abuse, known sensitivity to the ingredients in the product, significant depression or other psychiatric disease, any other cancer in the last 5 years except skin cancer and being on anticoagulation therapy.</p> <p><u>All patients</u> N: 94 Mean age: 49-75 Drop outs: 2</p> <p><u>Group 1</u> N: 46 Dropouts: 0</p> <p><u>Group 2</u> N: 48 Dropouts: 2 lost to follow-up</p>		<p>Mean (95% CI) Creatinine, mg/dl</p>	<p>Group 1: 1.107.80 (100.24-115.36) Group 2: 115.43 (109.13-121.73)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Willets et al., 2003³²²</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Australia</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 12 weeks</p>	<p>Patient group: men with symptoms of benign prostatic hyperplasia screened between January 1999 and March 2000.</p> <p>Inclusion criteria: Men with at least three symptoms of prostatism, (increased frequency of urination, nocturia, hesitancy, dribbling and poor stream); Under 80 years, with a maximum urinary flow rate of 5-15mL/s for a voiding volume of 150mL and a normal PSA level (<4ng/mL) within previous 3 months.</p> <p>Exclusion criteria: insulin-dependent diabetes, severe cardiopulmonary disease or significant CNS disease. Men who had used androgens, 5alpha reductase inhibitors, alpha blocker or herbal preparations in the last 4 weeks. Men with a history of prostate cancer, adenomas, urethral bladder, uretric or renal abnormalities, urogenital surgery ,renal stones, strictures or scarring , acute urinary retention or allergy to study treatment.</p> <p>All patients N: 100</p> <p>Group 1 N: 50 Mean (SEM) Age: 62.1 (1.2) Dropouts: 4 (discontinued due to acute bladder retention, abdominal pain, high PSA, arthralgia)</p> <p>Group 2 N: 50 Mean (SEM) Age: 63.9 (1.3) Dropouts: 3 (atrial fibrillation, dysuria, urinary incontinence)</p>	<p>Group 1: Serenoa repens 320mg (2X160mg of CO2 extract)</p> <p>Group 2: Placebo Paraffin oil (2 capsules a day)</p>	<p>Mean IPSS</p>	<p>Group 1: 12 Group 2: 13 1.74 (-0.54 to 4.03; p=0.131)</p>	<p>Funding: Blackmores Ltd.</p> <p>Limitations: At baseline the men in the placebo arm had significantly higher IPSS scores and more had symptoms of incontinence than in the intervention arm.</p> <p>Qmax reported for 62 men who attended initial and final visits and who voided >150mL but number in each group not provided. Therefore, further analysis can not be conducted.</p> <p>Additional outcomes: Multivariate regression analysis.</p> <p>Notes: Mean IPSS scores estimated from a graph as exact figures not given.</p>
			<p>Mean (95% CI) [SD] Quality of life score (IPSS question)</p>	<p>Baseline: Group 1: 3.66 (3.35-3.97) Group 2: 4.0 (3.58-4.42) 12 weeks: Group 1: 3.17 (2.76-3.58) [1.38] Group 2: 3.31 (2.85-3.77) [1.57] Treatment effect: 0.18 (-0.16 to 0.53); p=0.292</p>	
			<p>Mean Qmax, mL/s</p>	<p>Baseline (n=62): Group 1: 11.1 (10.3-11.8) Group 2: 11.2 (10.5-11.9) 12 Weeks (n=62): Group 1: 12.6 (11.0-14.2) Group 2: 15.6 (13.2-18.1)</p>	
			<p>IIEF scores (reported for 74 sexually active men)</p>	<p>Baseline Group 1: 51.5 (43.9-59.1) Group 2: 49.4 (43.3-55.4) 12 weeks: Group 1: 55.11 (48.4-61.8) Group 2: 48.7 (41.9-55.4)</p>	
			<p>Serious adverse events leading to withdrawal</p>	<p>Acute urinary retention Group 1: 1 Group 2: 0 Atrial fibrillation Group 1: 0 Group 2: 1 Abdominal pain Group 1: 1 Group 2: 0</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Wilt et al., 1999³²⁸</p> <p>Reports on four studies.</p> <p>Study design: Systematic review – Cochrane review</p> <p>Setting: Germany (3 studies) and UK (one study)</p> <p>Evidence level: 1++</p> <p>Duration of follow-up: 4-26 weeks</p>	<p>Patient group: Men with mild to moderate symptomatic benign prostatic hyperplasia.</p> <p>Inclusion criteria: Treatment duration of at least 30 days.</p> <p>Exclusion criteria: None reported</p> <p>All patients N: 519 Mean age: 65.4 (34-85) yrs Mean IPSS score=15.2 points (n=377) Mean peak urine flow=10.2mL/s (n=519) Mean prostate size=49.1 cc (n=262) Drop outs: 41 (7.9%)</p> <p>Group 1 Dropouts: 7.8%</p> <p>Group 2 Dropouts: 8.0%</p>	<p>Group 1: Phytotherapy Beta-sitosterols derived from South African star grass, Hypoxis rooperi or from species of Pinus and Picea.</p> <p>Three studies contained non-glucosidic B-sitosterol, but dosages ranged from 60mg/day to 195mg/day. Two studies utilised a preparation that contains at least 70% non-glucosidic B-sitosterol and one utilised a preparation with a non-glucosidic B-sitosterol concentration of 50%. One study utilised a preparation that contained 100% B-sitosterol-B-D-glucoside. The other 3 trials had a quantity of the b-sitosterol derivative, B-sitosterol-b-D-glucoside was less than 5% of the daily B-sitosterol.</p> <p>Group 2: placebo</p>	<p>Mean difference Symptom score (IPSS)</p>	-4.91 (95% CI: -6.29 to -3.53); 2 studies (n=342)	<p>Funding: Internal support from: Department of Veterans Affairs Health Services Research and Development Program, USA and Minneapolis/VISN-13 Center for chronic Diseases Outcomes Research, USA.</p> <p>Limitations: Allocation concealment and method of randomisation was unclear in 2 of the 4 studies. Different studies used varying doses and preparations of B-sitosterols.</p> <p>Additional outcomes:</p> <ul style="list-style-type: none"> - Boyarsky quality of life score in one study. - Physician overall evaluation of efficacy. - Sensitivity analysis of peak and residual volume without study Kadow 1986. Increases significance for intervention. <p>Notes: IPSS symptom scores from 0 to 35.</p>
			<p>Mean difference Nocturia; times per evening</p>	-1.00 (95% CI: -1.75 to -0.25); one study (n=80)	
			<p>Mean difference Peak urine flow, mL/s</p>	3.91 (95% CI: 0.91 to 6.90); 4 studies (n=474)	
			<p>Mean difference urine flow</p>	2.60 (95% CI: 1.30 to 3.90)	
			<p>Mean difference Residual volume, mL ; 4 studies</p>	-28.62 (95% CI: -41.42 to -15.83); 4 studies (n=475)	
			<p>Mean difference in reduction in prostate size</p>	-6.19 (95% CI: -15.29 to 2.91); 2 studies (n=216)	
			<p>% of patients with adverse events</p>	<p>Gastrointestinal: Group 1: 1.6 Group 2: 0</p> <p>Impotence: Group 1: 0.5 Group 2: 0</p>	
			<p>Mean difference of Boyarsky quality of life scale</p>	-4.50 [-6.05, -2.95]; one study (n=200)	
<p>Patient overall evaluation of efficacy (rated very good or good)</p>	8.25 [3.22, 21.13]; one study (n=80)				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Wilt et al., 2002a³²⁶</p> <p>Study design: Cochrane systematic review 21 RCTS included but 17 included that were compared to placebo.</p> <p>Setting: Europe and USA</p> <p>Evidence level: 1++</p> <p>Duration of follow-up: Mean study duration 13 weeks (4 -48 weeks range).</p>	<p>Patient group: Men with lower urinary tract symptoms consistent with benign prostatic hyperplasia.</p> <p>Inclusion criteria: Treatment duration of at least 30 days</p> <p>All patients N: 3139 (1408 in this comparison) Mean age: 65 years (40-88) Drop outs: 319 (10%) [0-18% range]</p>	<p>Group 1: Serenoa repens (SR) - alone or in combination)</p> <p>Group 2: placebo Also compares against other interventions.</p>	Mean difference symptom score (0-19)	-1.41 [-2.52, -0.30]; one study (n=205) P=0.013	<p>Funding: Internal sources of support:</p> <ul style="list-style-type: none"> • Management decision and research center- department of veterans affairs, USA • Minneapolis/VISN-13 Center for Chronic Diseases Outcomes Research, USA. <p>Limitations: Studies utilised different doses of serenoa repens but most frequently reported dose was 160mg twice per day.</p> <p>Additional outcomes: Also reported:</p> <ul style="list-style-type: none"> • SR/urtica vs. finasteride. • SR vs. pygeum africanum • SR vs. gestonorone <p>Notes: Results did not substantially change when restricted analysis to studies that had adequate allocation concealment or were double blinded. Meta-analysis used randoms effect model for all comparisons.</p>
			Mean change in IPSS score (score from 0-35)	-2.20 [-4.70, 0.30]; one study (n=79) P=0.084	
			Patient reported self rating from improved symptoms (men rating very good to good)	RR=1.76 [1.21, 2.56]; 6 studies (n=659) P=0.0029	
			Physician assessed improvement of symptoms	RR=1.72 [1.11, 2.66]; 3 studies (n=524) P=0.015	
			Mean difference Nocturia (times/evening)	-0.76 [-1.21, -0.31]; 10 studies (n=634) P=0.00084	
			Weighted mean difference Qmax, mL/s	1.86 [0.60, 3.12]; 9 studies (n=723) P=0.0038	
			Mean urine flow, ml/s	2.23 [1.18, 3.27]; 4 studies (n=382) P=0.000028	
			Residual volume, mL	-22.95 [-42.33, -3.56]; 6 studies (n=450) P=0.020	
			Prostate size	-2.14 [-10.93, 6.65]; 2 studies (n=243) P=0.63	
			Study withdrawals	0.72 [0.39, 1.32]; 7 studies (n=595) P=0.29	
			IPSS total score, mean change (serenoa repens/sabal urtica)	-3.50 [-6.75, -0.25]; one study (n=40) P=0.035	
Qmax (serenoa repens/sabal urtica)	1.60 [-1.67, 4.87]; one study (n=40) P=0.34				

Evidence Table 49: Phytotherapy combinations vs. placebo

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Lopatkin et al., 2005¹⁷¹</p> <p>Study design: Randomised controlled trial</p> <p>Setting: Multi centre,</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 24 weeks</p>	<p>Patient group: Male outpatients ≥ 50 years suffering from LUTS caused by BPH.</p> <p>Inclusion criteria: maximum urinary flow rate < 15 ml/s; change in maximum urinary flow between screening and end of run-in period 3 ml/s or less; urinary output > 100 ml at baseline; IPSS total score 14 or greater; IPSS quality of life 4 or greater. Written informed consent.</p> <p>Exclusion criteria: Inability to give informed consent or to complete self-ratings; previous or scheduled surgery involving pelvis or urinary tract; urethral stricture disease or a history of pelvic radiation therapy; PSA > 10 ng/ml; large residual urine > 350 ml; symptomatic urinary tract infection; chronic bacterial prostatitis; patients with diabetes mellitus, diabetic neuropathy or prostate carcinoma; serious general and specific risks; concomitant medication affecting the micturition pattern.</p> <p>All patients: N: 257</p> <p>Group 1 N: 129 Mean (±SD) Age: 68 (7) Dropouts: 4 (informed consent revoked=1; adverse events=3)</p> <p>Group 2 N: 128 Mean (±SD) Age: 67 (7) Dropouts: 3 (lost to follow-up=1, non-compliance=1; informed consent revoked=1)</p>	<p>Group 1: Phytotherapy combination of sabal/urtica 2 X 1 capsule daily of 160mg sabal fruit extract WS1473 and 120mg urtica root extract WS 1031 per capsule (PRO 160/120).</p> <p>Group 2: Placebo 2X1 capsule day (capsule identical in appearance to intervention).</p> <p>All patients: Placebo run in phase 2 weeks.</p>	<p>Mean (SD) total changes IPSS</p>	<p>Baseline Group 1 (n=127): 18 (4) Group 2 (n=126): 18 (3) Week 16 Group 1 (n=127): -4 (4) Group 2 (n=126): -3 (5) Week 24 Group 1 (n=127): -6 (4) Group 2 (n=126): -5 (5) P=0.03</p>	<p>Funding: NR</p> <p>Limitations: Baseline assessments: Initial diagnosis of BPH was systematically longer in patients randomised to intervention.</p> <p>Additional outcomes: Per protocol analysis also completed to assess robustness of results. Sub-analysis of IPSS score by irritative and obstructive components and by individual question. Sub-analysis of moderate and severe baseline IPSS scores and number in mild, moderate and severe IPSS category after 24 weeks.</p> <p>Notes: This trial was followed by an open label extension period where all patients received the intervention.</p> <p>2 patients from each group terminated trial early without any data for the primary outcome measure, and were excluded from the analysis.</p>
			<p>Mean (SD) changes in Qmax, ml/s</p>	<p>Baseline Group 1: 10.4 (2.4) Group 2: 10.5 (2.6) Week 24 Group 1: +1.8 (4.6) Group 2: +1.9 (4.5) P=0.59</p>	
			<p>Adverse events</p>	<p>Group 1: 23/129 (17.8%) Group 2: 24/128 (18.8%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Melo et al., 2002¹⁹⁹</p> <p>Study design: Randomised controlled trial.</p> <p>Setting: NR</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: Men with urinary symptoms.</p> <p>Inclusion criteria: ≥50 years, urinary symptoms assessed by IPSS with minimal score of 12, quality of life index of at least 3 points, rectal examination consistent with BPH and Maximum urinary flow rate between 5 and 15mL/s.</p> <p>Exclusion criteria: NR</p> <p>All patients N: 49 Drop outs: NR</p> <p>Group 1 N: 27 Mean (range) Age: 65.3 (52-86) Dropouts: NR</p> <p>Group 2 N: 22 Mean (range) Age: 65 (50-79) Dropouts: NR</p>	<p>Group 1: PHYTOTHERAPY COMBINATION 25mg Pygeum africanum and 300mg stinging nettle (1 PO bid).</p> <p>Group 2: PLACEBO Placebo (bid)</p>	<p>Mean (SD) IPSS score</p>	<p>Baseline Group 1: 19.3 (5.2) Group 2: 20.0 (5.9) 6 months Group 1: 14.6 (7.3) Group 2: 15.6 (7.9); P=0.658</p>	<p>Funding: NR.</p> <p>Limitations: No dropouts were reported in the study and method of randomisation was unclear.</p> <p>Additional outcomes: Comparison of ≥30% and 50% drop in IPSS, QoL and increase in Qmax.</p> <p>Notes: Baseline Qmax was better in the intervention group but Not sig.ly different.</p>
			<p>Mean (SD) quality of life index</p>	<p>Baseline Group 1: 3.81 (0.83) Group 2: 3.95 (1.09) 6 months Group 1: 3.33 (1.27) Group 2: 3.73 (1.52)</p>	
			<p>Mean (SD) Qmax</p>	<p>Baseline Group 1: 11.4 (3.1) Group 2: 10.2 (2.4); P=0.066 6 months Group 1: 12.5 (6.1) Group 2: 11.4 (3.8); P=0.770</p>	
			<p>Adverse events</p>	<p>Headache Group 1: 1/27 (3.7%) Group 2: 1/22 (4.5%) Chest pain Group 1: 0/27 Group 2: 1/22 (4.5%) Epigastric pain Group 1: 4/27 (14.8%) Group 2: 0/22 Drowsiness Group 1: 1/27 (3.7%) Group 2: 1/22 (4.5%) Vertigo Group 1: 0/27 Group 2: 1/22 (4.5%)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Preuss et al., 2001²⁴²</p> <p>Study design: Randomised controlled trial</p> <p>Setting: 3 sites, US</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 90 days</p>	<p>Patient group: Men with diagnosis of BPH.</p> <p>Inclusion criteria: no evidence of cancer by digital rectal and/or PSA examinations; maximal urinary flow rates were to be between 5-15ml/s for a voided volume in excess of 100ml. Read, speaks and understand English and written informed consent obtained.</p> <p>Exclusion criteria: Age over 80 years, presence of any tumour, malformation, or infection of the genitourinary tract; sever concomitant medical condition, severe laboratory abnormalities at baseline; finasteride within the last 4 weeks; patients being treated with antibiotics for genitourinary tract infections.</p> <p>All patients: N: 144 Drop outs: 17</p> <p>Group 1 N: 75 Mean (\pmSD) Age: Dropouts:5 (withdrew consent=1, lost to follow-up=1)</p> <p>Group 2 N: 69 Mean (\pmSD) Age: Dropouts:12 (adverse events=3, withdrew=5, lost to follow-up=3; protocol violation=1)</p>	<p>Group 1: phytotherapy 2 pills of combined natural products Cernitin 378mg, saw palmetto complex and phytosterol (saw palmetto fruit standardised to 40-50% free fatty acids and B-sitosterol standardised to 43%) 286g, and Vitamin E 100 IU.</p> <p>Group 2: Control 2 pills of placebo</p>	<p>Mean AUA scores</p> <p>Mean (SEM) [SD] change in AUA symptom index</p> <p>Mean (SEM) [SD] maximum flow rate, ml/min</p> <p>Mean (SEM) Average flow rate, ml/min</p> <p>Mean (SEM) Bladder volume, ml</p> <p>Adverse events</p>	<p>Baseline Group 1 (n=70): 18.9 Group 2 (n=57): 17.7</p> <p>Day 45 Group 1 (n=70): 14.6 Group 2 (n=57): 15.0</p> <p>Day 90 Group 1 (n=70): 12.7 Group 2 (n=57): 14.5 ANOVA p=0.014</p> <p>Group 1 (n=70): -6.171 (0.766) [6.41] Group 2 (n=57): -3.241 (0.774) [5.84] P=0.009</p> <p>Baseline Group 1 (n=70): 11.2 (0.8) Group 2 (n=57): 12.1 (0.9)</p> <p>Day 90 Group 1 (n=70): 11.8 (0.7) [5.86] Group 2 (n=57): 13.1 (1.0) [7.55]</p> <p>Baseline Group 1 (n=70): 6.0 (0.4) Group 2 (n=57): 6.1 (0.5)</p> <p>Day 90 Group 1 (n=70): 6.0 (0.5) Group 2 (n=57): 6.8 (0.5)</p> <p>Baseline Group 1 (n=70): 58.9 (11.4) Group 2 (n=57): 59.6 (12.8)</p> <p>Day 90 Group 1 (n=70): 57.5 (12.8) Group 2 (n=57): 40.7 (10.4)</p> <p>Flatulence: Group 1: 3 Group 2: 0</p> <p>Lower abdominal rash: Group 1: 0 Group 2: 1</p> <p>Dizziness</p>	<p>Funding: Rexall/Sundown, Inc, Boca Raton, FL through the National Research Council for Health, Washington DC and Meridian ID.</p> <p>Limitations: Baseline levels not reported.</p> <p>Additional outcomes: AUA scores for each of 7 questions reported. Comparison of PSA changes.</p> <p>Notes: SD calculated by NCC.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				Group 1: 0 Group 2: 1 Headache Group 1: 1 Group 2: 1 Nausea/GI distress Group 1: 0 Group 2: 2 Urinary tract infection: Group 1: 1 Group 2: 0 Ear infection: Group 1: 0 Group 2: 1 Lumbar spine surgery Group 1: 0 Group 2: 1 Herpes Zoster Group 1: 1 Group 2: 0 Elevated BP: Group 1: 0 Group 2: 1 Chest pain: Group 1: 0 Group 2: 1 Right arm laceration Group 1: 1 Group 2: 0	

Evidence Table 50: Phytotherapy vs. Alpha-blockers

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																				
Debruyne et al., 2002 ⁶⁸ Study design: RCT Patients masked to treatment Evidence level: 1+ Duration of follow-up: 12 months	Patient group: men with BPH Setting: multicentre, 98 centres across 9 European countries. Inclusion criteria: <ul style="list-style-type: none"> IPSS >10 Qmax between 5-15 mL/sec with a urine volume of \geq 150 mL and PVR <150mL Prostate volume \geq25 mL Serum PSA <4ng/mL Men with serum PSA 4-10 ng/mL required to have free/total PSA ratio of \geq15% to be enrolled 50 - 85 years 90% compliance after a 4 week placebo run in. Exclusion criteria: <ul style="list-style-type: none"> Prostate cancer Known history of bladder disease (cancer, bladder neck surgery, neurogenic) Urethral strictures Pelvic radiotherapy Lower urinary tract infection Chronic bacterial prostatitis Any disease affecting micturation Patients with clinically significant cardiovascular disease, haematuria, type II diabetes, history of hepatic failure or abnormal liver function tests. Patients on concomitant medication likely to interfere with study medication. 	Group 1: Serenoa repens (saw palmetto), Permixon® 320 mg/day Group 2 Tamsulosin 0.4 mg/day Examination methods: Each patient evaluated at baseline then at 6, 13, 26, 39 and 52 weeks for IPSS and uroflowmetry. At weeks 26 and 52 TRUS was performed and blood and serum PSA taken at week 52. **Patient completed the validated male sexual function (MSF-4) questionnaire of 4 questions (0-5 points each): <ul style="list-style-type: none"> interest in sex quality of erection achieving orgasm achieving ejaculation 	IPSS \pm SD at 12 mths	Group 1: 10.8 \pm 5.5, n=269 Group 2: 11.0 \pm 6.0, n=273 p value: 0.99	Funding: Grant from Pierre Fabre Médicament, Castres, France, manufacturer of Permixon®. Authors have served as consultants or speakers for, or have received research grants from Pierre Fabre Médicament. Limitations: <ul style="list-style-type: none"> Randomisation method was not clear Allocation concealment was not clear Masking of outcome assessment was not clear. Only the per protocol data was available at follow up. Additional outcomes: Notes: Masking of treatments to patients was achieved by providing tamsulosin in a green coloured size 0 capsule similar to Permixon® Serious advent events defined as fatal, life																																				
			Qmax \pm SD at 12 mths	Group 1: 12.7 \pm 5.2, n=267 Group 2: 13.0 \pm 4.9, n=265 p value: 0.79																																					
			MSF-4 \pm SD at 12 mths	Group 1: 8.8 \pm 5.4, n=267 Group 2: 8.2 \pm 5.0, n=266 p value: 0.69																																					
			Serum PSA \pm SD at 12 mths	Group 1: 2.8 \pm 2.3, n=266 Group 2: 2.9 \pm 2.5, n=268 p value: 0.50																																					
			Prostate Volume \pm SD at 6 mths	Group 1: 47.0 \pm 20.9, n=269 Group 2: 48.2 \pm 22.7, n=270 p value: 0.27																																					
			Incidence of Adverse Events	<table border="1"> <thead> <tr> <th></th> <th>Group 1: (%)</th> <th>Group 2: (%)</th> </tr> </thead> <tbody> <tr> <td>349</td> <td>354</td> <td></td> </tr> <tr> <td>N</td> <td>1 (0.3)</td> <td>4 (1.1)</td> </tr> <tr> <td>Decreased Libido</td> <td>2 (0.6)</td> <td>15 (4.2)</td> </tr> <tr> <td>Ejaculation Disorders</td> <td>4 (1.1)</td> <td>5 (1.4)</td> </tr> <tr> <td>Asthenia</td> <td>6 (1.7)</td> <td>5 (1.4)</td> </tr> <tr> <td>Fatigue</td> <td>10 (2.9)</td> <td>6 (1.7)</td> </tr> <tr> <td>Dizziness</td> <td>30 (8.6)</td> <td>43 (12.1)</td> </tr> <tr> <td>Rhinitis</td> <td>4 (1.1)</td> <td>3 (0.8)</td> </tr> <tr> <td>Hypotension postural</td> <td>28 (8.0)</td> <td>37 (10.5)</td> </tr> <tr> <td>Headache</td> <td>3 (0.9)</td> <td>2 (0.64)</td> </tr> <tr> <td>Dry Mouth</td> <td></td> <td></td> </tr> </tbody> </table>			Group 1: (%)	Group 2: (%)	349	354		N	1 (0.3)	4 (1.1)	Decreased Libido	2 (0.6)	15 (4.2)	Ejaculation Disorders	4 (1.1)	5 (1.4)	Asthenia	6 (1.7)	5 (1.4)	Fatigue	10 (2.9)	6 (1.7)	Dizziness	30 (8.6)	43 (12.1)	Rhinitis	4 (1.1)	3 (0.8)	Hypotension postural	28 (8.0)	37 (10.5)	Headache	3 (0.9)	2 (0.64)	Dry Mouth		
				Group 1: (%)		Group 2: (%)																																			
349	354																																								
N	1 (0.3)	4 (1.1)																																							
Decreased Libido	2 (0.6)	15 (4.2)																																							
Ejaculation Disorders	4 (1.1)	5 (1.4)																																							
Asthenia	6 (1.7)	5 (1.4)																																							
Fatigue	10 (2.9)	6 (1.7)																																							
Dizziness	30 (8.6)	43 (12.1)																																							
Rhinitis	4 (1.1)	3 (0.8)																																							
Hypotension postural	28 (8.0)	37 (10.5)																																							
Headache	3 (0.9)	2 (0.64)																																							
Dry Mouth																																									
Reasons for withdrawal*	<table border="1"> <thead> <tr> <th></th> <th>Group 1: n=54</th> <th>Group 2: n=56</th> </tr> </thead> <tbody> <tr> <td>Serious Adverse Events</td> <td></td> <td></td> </tr> <tr> <td>Non-serious adverse events</td> <td>3</td> <td>8</td> </tr> <tr> <td>Acute urinary retention</td> <td>10</td> <td>13</td> </tr> <tr> <td>Lack of efficacy</td> <td>4</td> <td>3</td> </tr> <tr> <td>Sexual dysfunction</td> <td>15</td> <td>8</td> </tr> <tr> <td>Other events</td> <td>1</td> <td>2</td> </tr> <tr> <td>Other events</td> <td>2</td> <td>1</td> </tr> <tr> <td>Patient decision</td> <td>14</td> <td>15</td> </tr> <tr> <td>Lost to follow up</td> <td>2</td> <td>2</td> </tr> </tbody> </table>		Group 1: n=54	Group 2: n=56	Serious Adverse Events			Non-serious adverse events	3	8	Acute urinary retention	10	13	Lack of efficacy	4	3	Sexual dysfunction	15	8	Other events	1	2	Other events	2	1	Patient decision	14	15	Lost to follow up	2	2										
	Group 1: n=54	Group 2: n=56																																							
Serious Adverse Events																																									
Non-serious adverse events	3	8																																							
Acute urinary retention	10	13																																							
Lack of efficacy	4	3																																							
Sexual dysfunction	15	8																																							
Other events	1	2																																							
Other events	2	1																																							
Patient decision	14	15																																							
Lost to follow up	2	2																																							

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<ul style="list-style-type: none"> • Hypersensitivity to study drugs • Participation in another trial within previous 3 mths <p>All patients N: 704 randomised but only 685 included in ITT analysis Mean age: 65.2 yrs Drop outs: 110 (16.1%)*</p> <p>Group 1 N: 340 Mean (± SD) Age: 65.6 ± 7.4 BMI (± SD): 26.7 ± 3.6 IPSS (± SD): 15.5 ± 4.8 MSF-4 (± SD): 8.3 ± 5.3** Qmax (± SD), mL/s: 10.9 ± 3.9 Prostate volume (± SD), mL: 48.0 ± 18.2 Serum PSA (± SD), ng/mL: 2.8 ± 2.0 Dropouts: 54*</p> <p>Group 2 N: 345 Mean (± SD) Age: 64.9 ± 7.6 BMI (± SD): 26.7 ± 3.7 IPSS (± SD): 15.2 ± 5.2 MSF-4 (± SD): 7.7 ± 5.0** Qmax (± SD), mL/s: 11.3 ± 4.3 Prostate volume (± SD), mL: 47.7 ± 18.6 Serum PSA (± SD), ng/mL: 2.8 ± 2.2 Dropouts: 56*</p>		Other	3 4	threatening, disabling resulting in hospitalisation or associated with cancer

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Engelmann U et al., 2006⁸²</p> <p>Study design: RCT</p> <p>Setting: 23 private urological practices in Germany.</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 60 weeks</p>	<p>Patient group: Outpatients suffering from BPH that did not require surgery.</p> <p>Inclusion criteria: A maximum urinary flow rate ≤ 12 ml/s at a urinary volume ≥ 150 ml was required. Aged 50 years old and above. Initial IPSS score of ≥ 13 points and an IPSS QoL assessment score ≥ 3.</p> <p>Exclusion criteria: Patients whose peak urinary flow rate changed by more than 3 ml/s during a 2-week placebo run-in phase were excluded. Patients with a residual urinary volume > 150 ml, congested urinary tract passages, an indication for BPH surgery, urinary tract infection, prostate carcinoma, diabetes, neurogenic or bladder dysfunction as well as patients previously treated with 5α-reductase inhibitors.</p> <p>All patients N: 140 Drop outs: 9/140</p> <p>Group 1 N: 71 Age \pm SD, years: 65 \pm 8 Time since diagnosis of BPH (years): 3.1 \pm 4 Dropouts: 11</p> <p>Group 2 N: 69 Age \pm SD, years: 65 \pm 8 Time since diagnosis of BPH (years): 3.61 \pm 4.5 Dropouts: 8</p> <p>Exclusions after randomization</p>	<p>Group 1: PRO 160/120 160mg Sabal fruit extract and 120mg Urtica root per capsule.</p> <p>Group 2: Tamsulosin Slow-release capsules containing 0.4mg active ingredient.</p> <p>For both drugs placebo capsules were available which were indistinguishable from their pharmacologically active counterparts in all aspects of their outer appearance.</p> <p>(After screening patients entered a single blind placebo run in phase of two weeks.)</p> <p>Examination methods: Visits scheduled after 8, 16, 24, 36, 48 and 60 weeks of</p>	<p>Median IPSS total score</p>	<p>Baseline Group 1: 20 Group 2: 20</p> <p>24 weeks Group 1: 13 Group 2: 12</p> <p>60 weeks Group 1: 10 Group 2: 9</p>	<p>Funding: NR</p> <p>Limitations: Median scores reported. Details of adverse events not reported.</p> <p>Additional outcomes: Subgroup analysis of patients with IPSS baseline score of ≤ 19 and IPSS baseline score ≥ 20</p> <p>Erectile function score – median score change for both groups = 0.</p> <p>Notes: Randomization was performed in balanced blocks, by means of a validated EDP random number generator program.</p>
			<p>Median improvement from baseline in LUTS-associated QoL (single item, range 0 [very good] -6 [very bad]).</p>	<p>Group 1: 2 Group 2: 1</p>	
			<p>Adverse events (details not reported)</p>	<p>Group 1: 15 patients (21.1%) reported 18 events Group 2: 19 patients (27.5%) reported 23 events.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Revoked informed consent: 2 Adverse event during placebo run-in: 2 Not meeting selection criteria: 5	double blind treatment.			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																								
<p>Hizli & Uygur, 2007¹¹⁹</p> <p>Study design: RCT open label</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with symptomatic BPH</p> <p>Setting: Department of Urology, Oncology, Education and research, Ankara Hospital, Turkey.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • IPSS ≥ 10 • Qmax 5-15 mL/s • PVR ≤ 150 mL • Prostate volume ≥ 25 mL • PSA ≤ 4 ng/mL <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • History of bladder disease affecting micturation • Urethral stenosis • Pelvic radiotherapy • Prostate cancer • Infections of urinary tract or chronic bacterial prostatitis • Clinically significant cardiovascular disease • Haematuria • Type II diabetes • Severe hepatic failure or abnormal liver function tests • Known hypersensitivity to study drugs • Participation in another trial within previous 3 months <p>All patients N: 60 Age (range): 43-73 years Drop outs:</p> <p>Group 1 N: 20</p>	<p>Group 1: Serenoa repens (Prostagood®) 320 mg/day</p> <p>Group 2 Tamsulosin 0.4 mg/day</p> <p>Group 3 Serenoa repens (Prostagood®) 320 mg/day + Tamsulosin 0.4 mg/day</p> <p>Examination methods: IPSS, QoL, Qmax by uroflowmetry recorded at baseline and months 2, 4, 6</p>	<p>IPSS \pm SD reduction from baseline at 6 mths</p>	<p>Group 1: -6.1 ± 2.7 Group 2: -4.6 ± 3.3 Group 3: -4.9 ± 2.3 p value: 0.16 (Kruskal-Wallis)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Randomisation method not reported • Allocation concealment not reported • Masking of outcome assessment not reported • Open label • Small study <p>Additional outcomes: No patients withdrew from the study due to adverse events.</p> <p>Notes: Notes</p>																																								
			<p>IPSS QoL \pm SD reduction from baseline at 6 mths</p>	<p>Group 1: -2.6 ± 0.9 Group 2: -2.1 ± 0.8 Group 3: -2.2 ± 1.0 p value: 0.14 (Kruskal-Wallis)</p>																																									
			<p>Qmax \pm SD increase from baseline at 6 mths</p>	<p>Group 1: 3.2 ± 2.2 Group 2: 3.7 ± 2.6 Group 3: 4.2 ± 2.5 p value: 0.38 (Kruskal-Wallis)</p>																																									
			<p>Prostate volume \pm SD decrease from baseline at 6 mths</p>	<p>Group 1: -0.7 ± 2.2 Group 2: -1.0 ± 2.2 Group 3: -0.8 ± 2.0 p value: 0.61 (Kruskal-Wallis)</p>																																									
			<p>PSA \pm SD decrease from baseline at 6 mths</p>	<p>Group 1: -2.0 ± 0.3 Group 2: -0.1 ± 0.2 Group 3: -3.5 ± 0.2 p value: 0.07 (Kruskal-Wallis)</p>																																									
			<p>Incidence of Adverse Events</p>	<table border="1"> <thead> <tr> <th></th> <th>Group 1: (%)</th> <th>Group 2: (%)</th> <th>Group 3: (%)</th> </tr> </thead> <tbody> <tr> <td>N</td> <td>20</td> <td>20</td> <td>20</td> </tr> <tr> <td>Decreased Libido</td> <td>-</td> <td>4 (20)</td> <td>1 (5)</td> </tr> <tr> <td>Ejaculation Disorders</td> <td>-</td> <td>7 (35)</td> <td>3 (15)</td> </tr> <tr> <td>Asthenia</td> <td>-</td> <td>-</td> <td>1 (5)</td> </tr> <tr> <td>Fatigue</td> <td>-</td> <td>2 (10)</td> <td>-</td> </tr> <tr> <td>Dizziness</td> <td>-</td> <td>2 (10)</td> <td>-</td> </tr> <tr> <td>Rhinitis</td> <td>-</td> <td>2 (10)</td> <td>-</td> </tr> <tr> <td>Hypotension postural</td> <td>-</td> <td>3 (15)</td> <td>-</td> </tr> <tr> <td>Dry Mouth</td> <td>-</td> <td>5 (25)</td> <td>1 (5)</td> </tr> </tbody> </table>			Group 1: (%)	Group 2: (%)	Group 3: (%)	N	20	20	20	Decreased Libido	-	4 (20)	1 (5)	Ejaculation Disorders	-	7 (35)	3 (15)	Asthenia	-	-	1 (5)	Fatigue	-	2 (10)	-	Dizziness	-	2 (10)	-	Rhinitis	-	2 (10)	-	Hypotension postural	-	3 (15)	-	Dry Mouth	-	5 (25)	1 (5)
				Group 1: (%)		Group 2: (%)	Group 3: (%)																																						
N	20	20	20																																										
Decreased Libido	-	4 (20)	1 (5)																																										
Ejaculation Disorders	-	7 (35)	3 (15)																																										
Asthenia	-	-	1 (5)																																										
Fatigue	-	2 (10)	-																																										
Dizziness	-	2 (10)	-																																										
Rhinitis	-	2 (10)	-																																										
Hypotension postural	-	3 (15)	-																																										
Dry Mouth	-	5 (25)	1 (5)																																										

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Age \pm SD, years: 56.8 \pm 7.8 IPSS \pm SD: 18.0 \pm 4.9 IPSS QoL \pm SD: 4.2 \pm 1.1 Qmax \pm SD, mL/s: 9.4 \pm 2.9 Prostate volume \pm SD, mL: 35.2 \pm 10.3 PVR \pm SD, mL: 67.4 \pm 27.7 PSA \pm SD, ng/mL: 1.9 \pm 0.9 BMI \pm SD, kg/m²: 26.7 \pm 2.5 Dropouts: 0</p> <p>Group 2 N: 20 Age \pm SD, years: 58.9 \pm 5.7 IPSS \pm SD: 16.2 \pm 4.7 IPSS QoL \pm SD: 3.5 \pm 1.1 Qmax \pm SD, mL/s: 10.5 \pm 2.8 Prostate volume \pm SD, mL: 38.6 \pm 11.6 PVR \pm SD, mL: 65.5 \pm 33.3 PSA \pm SD, ng/mL: 2.1 \pm 0.9 BMI \pm SD, kg/m²: 28.0 \pm 3.4 Dropouts: 0</p> <p>Group 3 N: 20 Age \pm SD, years: 60.2 \pm 6.3 IPSS \pm SD: 15.6 \pm 3.2 IPSS QoL \pm SD: 3.5 \pm 1.1 Qmax \pm SD, mL/s: 9.9 \pm 2.4 Prostate volume \pm SD, mL: 31.2 \pm 4.2 PVR \pm SD, mL: 63.7 \pm 23.7 PSA \pm SD, ng/mL: 1.7 \pm 0.7 BMI \pm SD, kg/m²: 27.8 \pm 2.3 Dropouts: 0</p>				

Evidence Table 51: Phytotherapy vs. 5-Alpha Reductase inhibitors

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																																				
<p>Carraro et al., 1996⁴³</p> <p>Study design: RCT Placebo controlled</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 6 months</p>	<p>Patient group: men with BPH and symptoms of BOO</p> <p>Setting: multicentre, 87 centres across 9 European countries.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • BPH diagnosed by DRE • IPSS >6 • Qmax between 4-15 mL/sec with a urine volume of ≥ 150 mL and PVR <200mL • Prostate volume >25 mL • Serum PSA <10 ng/mL for prostates ≤ 60mL • Serum PSA < 15 ng/mL for prostates > 60mL (measured before or 3 days after DRE & TRUS) • > 50 years • 2 week washout period after previous alpha-blockers or Pygeum • Good physical and mental condition <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Prostate cancer • Known history of bladder disease (cancer, bladder neck surgery, neurogenic) • Lower urinary tract infection • Any disease affecting micturation • Abnormal liver function (twice upper normal limit of serum aminotransferases and/or bilirubin, creatinine >160 μmol/L • Diuretics or drugs with antiandrogen or alpha receptor properties administered over previous 3 months for hypertension, 	<p>Group 1: Serenoa repens (saw palmetto), Permixon® 160 mg + placebo 2/day morning and evening for 26 weeks.</p> <p>Group 2 Finasteride (Proscar®) 5mg + placebo 1/day in the morning then 2 x placebo in the evening</p> <p>Examination methods: Each patient was examined prior to baseline and at 6, 13 and 26 weeks by the same investigator. At each visit Qmax (at 200 mL voided volume), IPSS, IPSS QoL and sexual function score (0-20 points) were determined. At weeks 13 & 26 TRUS and PSA were performed.</p>	<p>IPSS \pm SD at 6 mths</p>	<p>Group 1: 9.9 \pm 5.4, n=467 Group 2: 9.5 \pm 5.5, n=484 p value: 0.17 (CI 95%: -0.17, 0.96)</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> • Masking of outcome assessment was not clear. • Allocation concealment by packaging of drugs was not clear. <p>Additional outcomes: % patients with Qmax <10 mL/s or Qmax ≥ 10 mL/s at baseline and at 6 mths against % patients with IPSS <18 or IPSS ≥ 18 at baseline and at 6 mths.</p> <p>Notes: Computer generated randomisation sequence **Sexual function comprised 4 questions in the male sexual function questionnaire MSF-4 (0-5 points each) on interest in sex, quality of erection, achieving orgasm & ejaculation</p>																																				
			<p>IPSS QoL score \pm SD at 6 mths</p>	<p>Group 1: 2.25 \pm 1.29, n=467 Group 2: 2.15 \pm 1.26, n=484 p value: 0.14 (CI 95%: -0.04, 0.24)</p>																																					
			<p>Sexual Function Score \pm SD at 6 mths</p>	<p>Group 1: 7.9 \pm 5.4, n=467 Group 2: 9.3 \pm 5.7, n=484 p value: <0.0001 (CI 95%: -1.52, 0.96)</p>																																					
			<p>Qmax \pm SD at 6 mths</p>	<p>Group 1: 13.3 \pm 6.7, n=467 Group 2: 14.0 \pm 7.4, n=484 p value: 0.035 (CI 95%: -1.46, -0.054)</p>																																					
			<p>Prostate Volume \pm SD at 6 mths</p>	<p>Group 1: 41.5 \pm 20.5 n=467 Group 2: 36.7 \pm 17.2 n=484 p value: <0.001 (CI 95%: 1.11, 1.18)</p>																																					
			<p>Serum PSA at 6 mths</p>	<p>Group 1: 3.22 \pm 4.00, n=467 Group 2: 1.99 \pm 1.98, n=484 p value: <0.001 (CI 95%: 1.33, 1.45)</p>																																					
			<p>Inter current clinical events</p>	<table> <thead> <tr> <th></th> <th>Group 1: (%)</th> <th>Group 2: (%)</th> </tr> </thead> <tbody> <tr> <td>Hypertension</td> <td>17 (3.1)</td> <td>12 (2.2)</td> </tr> <tr> <td>Decreased Libido</td> <td>12 (2.2)</td> <td>16 (3.0)</td> </tr> <tr> <td>Abdominal pain</td> <td>10 (1.8)</td> <td>15 (2.8)</td> </tr> <tr> <td>Impotence</td> <td>8 (1.5)</td> <td>15 (2.8)</td> </tr> <tr> <td>Back pain</td> <td>9 (1.6)</td> <td>3 (0.6)</td> </tr> <tr> <td>Diarrhoea</td> <td>5 (0.9)</td> <td>6 (1.1)</td> </tr> <tr> <td>Influenza-type symptoms</td> <td>5 (0.9)</td> <td>6 (1.1)</td> </tr> <tr> <td>Urinary retention</td> <td>7 (1.3)</td> <td>3 (0.6)</td> </tr> <tr> <td>Headache</td> <td>7 (1.3)</td> <td>3 (0.4)</td> </tr> <tr> <td>Nausea</td> <td>3 (0.5)</td> <td>6 (1.1)</td> </tr> <tr> <td>Constipation</td> <td>2 (0.4)</td> <td>6 (1.1)</td> </tr> </tbody> </table>			Group 1: (%)	Group 2: (%)	Hypertension	17 (3.1)	12 (2.2)	Decreased Libido	12 (2.2)	16 (3.0)	Abdominal pain	10 (1.8)	15 (2.8)	Impotence	8 (1.5)	15 (2.8)	Back pain	9 (1.6)	3 (0.6)	Diarrhoea	5 (0.9)	6 (1.1)	Influenza-type symptoms	5 (0.9)	6 (1.1)	Urinary retention	7 (1.3)	3 (0.6)	Headache	7 (1.3)	3 (0.4)	Nausea	3 (0.5)	6 (1.1)	Constipation	2 (0.4)	6 (1.1)
				Group 1: (%)		Group 2: (%)																																			
Hypertension	17 (3.1)	12 (2.2)																																							
Decreased Libido	12 (2.2)	16 (3.0)																																							
Abdominal pain	10 (1.8)	15 (2.8)																																							
Impotence	8 (1.5)	15 (2.8)																																							
Back pain	9 (1.6)	3 (0.6)																																							
Diarrhoea	5 (0.9)	6 (1.1)																																							
Influenza-type symptoms	5 (0.9)	6 (1.1)																																							
Urinary retention	7 (1.3)	3 (0.6)																																							
Headache	7 (1.3)	3 (0.4)																																							
Nausea	3 (0.5)	6 (1.1)																																							
Constipation	2 (0.4)	6 (1.1)																																							

Study details	Patients	Interventions	Outcome measures	Effect size	Comments																					
	<p>cerebrovascular insufficiency.</p> <ul style="list-style-type: none"> • Prior treatment with Permixon® or Finasteride <p>All patients N: 1098 Mean age: 64.5 yrs Drop outs: 147 (13.4%)</p> <p>Group 1 N: 553 Mean (range) Age: 64.3 (49-87) BMI (range): 26 (17-38) IPSS (± SD): 15.7 ± 5.8 IPSS QoL (± SD): 3.63 ± 1.28 MSF-4 (± SD): 8.4 ± 5.5** Qmax (± SD), mL/s: 10.6 ± 2.8 PVR (± SD), mL: 52 ± 44 Prostate volume (± SD), mL: 43.0 ± 19.6 Serum PSA (± SD), ng/mL: 3.26 ± 3.41 Dropouts: 86*</p> <p>Group 2 N: 545 Mean (range) Age: 64.7 (49-88) BMI (range): 25.9 (18-36) IPSS (± SD): 15.7 ± 5.7 IPSS QoL (± SD): 3.66 ± 1.17 MSF-4 (± SD): 8.5 ± 5.5** Qmax (± SD), mL/s: 10.8 ± 3.1 PVR (± SD), mL: 52 ± 44 Prostate volume (± SD), mL: 44.0 ± 20.6 Serum PSA (± SD), ng/mL: 3.23 ± 3.34 Dropouts: 61*</p>		<p>Dysuria</p> <p>Reasons for withdrawal*</p> <p>Side effects Lack of efficacy Patient decision Lost to follow up Mortality (non drug related) Other</p>	<table border="1"> <thead> <tr> <th></th> <th>Group 1: n=86</th> <th>Group 2: n=61</th> </tr> </thead> <tbody> <tr> <td>Lack of efficacy</td> <td>28</td> <td>14</td> </tr> <tr> <td>Patient decision</td> <td>0</td> <td>2</td> </tr> <tr> <td>Lost to follow up</td> <td>28</td> <td>20</td> </tr> <tr> <td>Mortality (non drug related)</td> <td>5</td> <td>7</td> </tr> <tr> <td>Other</td> <td>1 (heart attack)</td> <td>1 (fatal MI)</td> </tr> <tr> <td></td> <td>24</td> <td>17</td> </tr> </tbody> </table>		Group 1: n=86	Group 2: n=61	Lack of efficacy	28	14	Patient decision	0	2	Lost to follow up	28	20	Mortality (non drug related)	5	7	Other	1 (heart attack)	1 (fatal MI)		24	17	
	Group 1: n=86	Group 2: n=61																								
Lack of efficacy	28	14																								
Patient decision	0	2																								
Lost to follow up	28	20																								
Mortality (non drug related)	5	7																								
Other	1 (heart attack)	1 (fatal MI)																								
	24	17																								

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Sökeland, 2000²⁸²</p> <p>Study design: RCT Placebo controlled</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 1 year</p>	<p>Patient group: men with BPH (Aiken stages I to II)</p> <p>Setting: multicentre, University of Münster, Germany.</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> NR <p>Exclusion criteria:</p> <ul style="list-style-type: none"> < 50 years BPH III or above (Aiken) PSA > 10 ng/mL Prostate cancer Use of other prostate medications Infections Severe concomitant disease requiring therapy <p>All patients N: 516 Age (range): 50 - 88 Drop outs: 27 (5%) 489 available for efficacy analysis</p> <p>Group 1 N: 261 IPSS (\pm SD): 11.3 \pm 6.5 (n=258) Qmax (\pm SD), mL/s: 12.4 \pm 4.5 (n=245) Prostate volume (\pm SD), mL: 42.7 \pm 27.8 (n=215) Dropouts: 16</p> <p>Group 2 N: 255 Dropouts: 11 IPSS (\pm SD): 11.8 \pm 6.6 (n=255)</p>	<p>Group 1: Combination phytotherapy PRO 160/120 (serenoa repens (saw palmetto) extract 160 mg and Urtica (nettle) extract 120 mg) 2/day + 1 placebo 1/day</p> <p>Group 2 Finasteride (Proscar®) 5mg 1/day + 1 placebo 2/day</p> <p>Examination methods: Qmax, average flow and IPSS measured.</p>	<p>IPSS \pm SD at 6 mths</p> <p>IPSS \pm SD at 12 mths</p> <p>Qmax \pm SD at 3 mths</p> <p>Qmax \pm SD at 6 mths</p> <p>Qmax \pm SD at 12 mths</p> <p>Prostate volume \pm SD at 12 mths</p> <p>Number of adverse events (details not reported in Cochrane review or Sökeland, 2000) but the</p>	<p>Group 1: 8.2 \pm 5.8, n=233 Group 2: 8.0 \pm 5.7, n=230 p value: 0.66</p> <p>Group 1: 6.5 \pm 5.8, n=230 Group 2: 6.2 \pm 5.2, n=223 p value: 0.54</p> <p>Group 1: 14.2 \pm 6.0, n=240 Group 2: 14.6 \pm 6.6, n=242 p value: 0.46</p> <p>Group 1: 14.6 \pm 6.2, n=245 Group 2: 15.1 \pm 7.1, n=244 p value: 0.34</p> <p>Group 1: 14.6 \pm 6.4, n=233 Group 2: 15.4 \pm 6.8, n=232 p value: 0.19</p> <p>Group 1: 42.4 \pm NR Group 2: 37.2 \pm NR p value: NR</p> <p>Group 1: 74 in 52 patients Group 2: 96 in 54 patients</p> <p>Note: the abstract for Sökeland & Albrecht, 1997²⁸³ states that there were less cases of diminished ejaculation volume, erectile dysfunction and headache for those patients on PRO160/120</p>	<p>Funding: NR</p> <p>Limitations:</p> <ul style="list-style-type: none"> Safety information was not reported in the 2000 study and not available from the Wilt et al., 2002³²⁶ Cochrane Review. Neither standard deviations or p values <p>Notes: Additional methods information is available from first publication, Sökeland & Albrecht, 1997²⁸³, translated from German in the Wilt et al., 2002³²⁶ Cochrane Review.</p> <p>Randomisation was computer generated and allocation concealment was reported as being adequate in the Cochrane Review</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	Qmax (\pm SD), mL/s: 12.8 ± 4.0 (n=241) Prostate volume (\pm SD), mL: 44.0 ± 26.6 (n=216)				

Evidence Table 52 Provision of information

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Barry et al., 1997 ²³ Study design: RCT Evidence level: 1+ Duration of follow-up: 1 Year	Patient group: Men with clinical diagnosis of BPH. Setting: Urologic practices of Group Health Cooperative of Puget Sound (staff model health maintenance organisation) in Washington; 2 practices were located in Seattle and Tacoma. Exclusion criteria: Evidence of prostate cancer, obstructive nephropathy, post void residual >350mL, recurrent or refractory urinary infection, acute retention, previous prostate surgery, repeated gross hematuria, clot retention, bladder stones, comorbid conditions, inability to understand English. All patients N: 227 Group 1 N: 104 Age (mean): 66.4 (SD: 8.6) AUA score (mean): 16.6 (SD: 6.7) Drop outs: 1 Group 2 N: 123 Age (mean): 66.2 (SD: 8.2) AUA score (mean): 15.9 (SD: 7.0) Drop outs: 7	Group 1: Computer and interactive video-based shared decision-making program (SDP) to educate men about their condition and its treatments. - short questionnaire before viewing; so a subset of items entered into computer to tailor programme to viewer. - 30 minute segment explaining importance of participation in the treatment decision and outlines the choices of watchful waiting, medical or surgical treatment. Estimates of outcome probabilities given. - then there is an interactive segment that allows for review of old material and inspection of 30 minutes of new material in optional modules on acute retention, sexual dysfunction, incontinence, new treatments, BPH and prostate cancer, blood transfusion, symptom response to surgery. Group 2: Brochure to provide basic information about the prostate gland and disease that can affect it, including BPH. No quantitative information about treatment outcomes provided.	Treatment selection at 3 months:	Prostatectomy: Group 1: 5/104 (4.8%) Group 2: 8/123 (6.5%) Medication: Group 1: 14/104 (13.5%) Group 2: 14/123 (11.4%) Watchful waiting: Group 1: 85/104 (81.7%) Group 2: 101/123 (82.1%) P=0.8	Funding: Grant Nos. HS 06540 and 08397 from the Agency for Health Care Policy and Research. The development of the first edition of the SDP for BPH was funded by a grant from the John A. Hartford Foundation. Limitations: 2 phases of recruitment (pre-consent randomisation phase and post consent randomisation phase). Additional outcomes: Mean change in autonomy preference scores. Notes: * p values from a repeated measures analysis of covariance over all assessment points, controlling for age, practice site, marital status, education, income and race.
			Men undergone prostatectomy at 1 year:	Group 1: 8/104 (7.7%) Group 2: 16/123 (13.0%) p value: 0.28 Absolute diff: 5.3% (CI: -2.5%, +13.0%)	
			Mean BPH knowledge score: at 2 weeks	Group 1: 11.5 (SEM 0.5) Group 2: 6.7 (SEM 0.4) p value: <0.001	
			Mean (SE) satisfaction scores for decision process: 12 months	Group 1: 74.77 (1.72) Group 2: 69.26 (1.89) p value*: 0.03	
			Mean (SE) satisfaction scores for decision made: 12 months	Group 1: 75.16 (1.80) Group 2: 71.74 (1.75) p value: 0.21	
			Mean (SE) changes of AUA symptom score: 12 months	Group 1: -0.88 (0.74) Group 2: -1.45 (0.58) p value: 0.58	
			Mean (SE) change in BPH impact score: 12 months	Group 1: -1.05 (0.25) Group 2: -0.59 (0.25) p value: 0.12	
			Mean (SE) changes in general health score at 12 months:	Group 1: 0.61 (1.58) Group 2: -4.99 (1.44) p value: 0.02	
Mean (SE) changes in physical functioning score at 12 months:	Group 1: 0.15 (1.40) Group 2: -3.74 (1.18) p value: 0.02				

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
			Mean (SE) changes in social functioning score at 12 months:	Group 1: -1.46 (1.85) Group 2: -3.52 (1.71) p value: 0.17	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Brown et al., 2007 ³⁵ Study design: RCT Evidence level: 1+ Duration of follow-up: 12 months	<p>Patient group: men over 40 with uncomplicated lower urinary tract symptoms who were referred for the first time by their GP (from January 2003 and April 2004).</p> <p>Setting: Outpatient departments of 2 urological centres in London, a teaching hospital and a district general hospital.</p> <p>Exclusion criteria: medical treatment in the previous three months, recent surgery, complications potentially related to their symptoms or severe comorbidity.</p> <p>All patients N: 140 Drop outs: 25</p> <p>Group 1: N: 73 Age (mean): 63.3 (11.1) Drop outs: 14 at 12M Mean (SD) duration of symptoms (years): 3.9 (4.0) Mean (SD) IPSS: 16.9 (5.1) Mean (SD) AUA-QoL score: 4.0 (1.0)</p>	<p>Group 1: Self management and standard care group Small group sessions (5-8 men), each lasting between 1.5 and 2 hours, which were scheduled one, two and six weeks after randomisation. The aim of these sessions was to bring about modification of lifestyle (fluid management, avoidance of caffeine, and use of alcohol) and specific changes in behaviour (bladder training, double voiding, and urethral milking). Facilitated by urology nurses trained to enhance self management skills and provided support by brainstorming and group discussion. This intervention group also received standard care (as described below).</p> <p>Group 2: Standard care Standard care began with watchful waiting. Escalation to medical treatment and surgery was left to the discretion of the clinician and patient.</p> <p>All patients, irrespective of treatment allocation, received standard written information about lower urinary tract symptoms.</p>	<p>Number (%) of men with treatment failure:</p> <p>Failure defined as a rise of 3 points or more on the international prostate symptom score, use of drugs to control lower urinary tract symptoms, acute urinary retention, or surgical intervention) during follow-up.</p>	<p>3-month outcome: Group 1: 7/71 (10%) Group 2: 27/65 (42%) Difference (95% CI): 32 (18 to 46) p value: <0.001</p> <p>6-month outcome: Group 1: 13/69 (19%) Group 2: 39/64 (61%) Difference (95% CI): 42 (27 to 57) p value: <0.001</p> <p>12-month outcome: Group 1: 18/59 (31%) Group 2: 44/56 (79%) Difference (95% CI): 48 (32 to 64) p value: <0.001</p>	<p>Funding: BUPA Foundation Project Grant. Author CTB received a research fellowship from the Royal College of Surgeons of England, funded by Cazenove & Co. Author JvdM is funded by a national public health career scientist award from the Department of Health and NHS R&D Programme.</p> <p>Limitations: The study was underpowered as according to their calculations 84 men in each group were necessary to have a 90% chance to detect a 3 point reduction in mean international prostate symptom score at 5% level of significance with SD of 6.</p> <p>Additional outcomes: Reasons for treatment failure at 3, 6 and 12 months. BPH index score.</p> <p>Notes:</p>
			<p>Mean (SD) International Prostate Symptom Score (IPSS) (Score: 0-35; the higher the score the worse the symptoms)</p>	<p>3-month outcome: Group 1:(n= 71): 10.7 (5.9) Group 2: (n=64): 16.4 (5.8) Difference (95% CI): 5.7 (3.7 to 7.7), p value: <0.001</p> <p>6-month outcome: Group 1(n= 67): 10.4 (6.1) Group 2(n=61): 16.9 (6.4) Difference (95% CI): 6.5 (4.3 to 8.7), p value: <0.001</p> <p>12-month outcome: Group 1: (n=53): 10.2 (6.1) Group 2:(n=51): 15.4 (6.6) Difference (95% CI): 5.1 (2.7 to 7.6), p value: <0.001</p>	
			<p>Mean (SD) AUA-QoL score: (lower score the</p>	<p>3-month outcome: Group 1:(n= 71): 2.8 (1.2)</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>Group 2: N: 67 Age (mean): 63.4 (10.4) Drop outs: 11 at 12M Mean (SD) duration of symptoms (years): 4.3 (6.7) Mean (SD) IPSS: 15.9 (6.5) Mean (SD) AUA-QoL score: 3.3 (1.1)</p>		<p>better quality of life)</p>	<p>Group 2:(n=64): 3.4 (1.1) Difference (95% CI): 0.6 (0.2 to 1.0), p value: < 0.001</p> <p>6-month outcome: Group 1:(n= 67): 2.6 (1.3) Group 2:(n=61): 3.3 (1.4) Difference (95% CI): 0.7 (0.2 to 1.2), p value: 0.008</p> <p>12-month outcome: Group 1: (n=54): 2.6 (1.3) Group 2: (n=52): 3.1 (1.2) Difference (95% CI): 0.5 (0 to 1.0) p value: 0.03</p>	<p>Compliance with self management programme was high; 68 (93%) patients attended all three sessions. The five patients who did not attend were included in the self management group for analysis.</p> <p>Self management group included more men with university degree and fewer men with no qualification.</p>

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Murray et al., 2001²⁰⁷</p> <p>Study design: RCT</p> <p>Evidence level: 1+</p> <p>Duration of follow-up: 9 months</p>	<p>Patient group: Men with benign prostatic hypertrophy</p> <p>Setting: Primary care</p> <p>Inclusion criteria: Men with benign prostatic hypertrophy. No more details provided.</p> <p>Exclusion: Men with any clinical suggestion of carcinoma of the prostate or if they had chronic retention of the urine, recent urinary tract infection, a history of acute urinary retention or prostate surgery, severe visual or hearing impairment, or severe learning difficulties or mental illness.</p> <p>All patients N: 112 Drop outs: 10</p> <p>Intervention group N: 57 Age (mean +/- SD): 63.7 +/- 8.4 Drop outs: 3 Mean (SD) American Urological Association score: 15.64 (6.57) Up to secondary education; n (%): 25 (44) Beyond secondary education; n (%): 32 (56) Mean (SD) Spielberg state trait anxiety inventory: 33.93 (13.09)</p> <p>Control group</p>	<p>Group 1: Interactive multimedia programme with booklet and printed summary. Treatment options discussed were surgery, balloon dilatation of the prostate, drugs, and watchful waiting. Information comprised probabilities of the risks and benefits of each treatment, calculated on the basis of information on age, severity of symptoms, and general health entered by the patient at the beginning of the session. All patients saw the core interactive video disc, lasting about 45 minutes; viewing optional sections for further information took up to 60 min. more. A research nurse started the programme, taught the patient how to use it, and then withdrew.</p> <p>Group 2: Normal care from GP practitioner.</p>	<p>Mean (SD) decisional conflict score at three months: Higher scores indicated increased uncertainty.</p>	<p>Group 1: 2.3 (0.4) Group 2: 2.6 (0.5) Mean difference (95% CI): -0.3 (-0.5 to -0.1), p <0.01</p>	<p>Funding: NHS national research and development programme, the BUPA Foundation, and the King's Fund.</p> <p>Limitations: The initial aim of the study was to detect a difference in anxiety, however, recruitment rate was low and it was not possible to recruit the 210 patients needed from the sample size calculation.</p> <p>Additional outcomes: Cost per patient for a number of item. Only total costs are reported in this table.</p> <p>Authors found no difference between the two groups in the trends over time in the EQ-5D responses nor in the SF-36 scores. Data not provided.</p> <p>Anxiety scores: the Spielberg scores were similar at the final assessment in the two groups (Mann-Whitney U test). No data provided.</p> <p>Resource volumes per patient over nine months of trial.</p> <p>Notes: Decisional conflict score contains three subscales that</p>
			<p>Mean (SD) decisional conflict score at nine months:</p>	<p>Group 1: 2.23 (0.38) Group 2: 2.55 (0.50) Mean difference (95% CI): -0.33 (-0.51 to -0.14)</p>	
			<p>GPs perceptions of decision making at three months. Values are numbers and (%). Question: Who do you think made the treatment decision?</p>	<p>Mainly or only GP: Group 1 (n=48): 1 (2) Group 2 (n=49): 5 (10) % difference (95% CI): -8 (-17.5 to 1.3) GP and patient together: Group 1: 25 (52) Group 2: 32 (65) % difference (95% CI): -13 (-32.6 to 6.2) Mainly or only patient: Group 1: 22 (46) Group 2: 12 (25) % difference (95% CI): 21 (2.8 to 39.9) X²= 6.458, df=2; p=0.04</p>	
			<p>Patients' perceptions of decision making at three months. Question: Who do you think made the treatment decision?</p>	<p>Mainly or only GP: Group 1 (n=57): 5(9) Group 2 (n=48): 4 (8) % difference (95% CI): 1 (-10.3 to 11.2) GP and patient together: Group 1: 34 (60) Group 2: 42 (88) % difference (95% CI): -28 (-43.7 to 12.0) Mainly or only patient: Group 1: 18 (32) Group 2: 2 (4) % difference (95% CI):</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
	<p>N: 55 Age (mean +/- SD): 63.9 +/- 8.4 Drop outs: 7 Mean (SD) American Urological Association score: 14.85 (7.10) Up to secondary education; n (%): 28 (51) Beyond secondary education; n (%): 27(49) Mean (SD) Spielberg state trait anxiety inventory: 32.01 (10.49)</p>		<p>American Urological Association scores</p> <p>Total costs in pounds sterling (at 1999 prices) per patient: Mean (SD)</p>	<p>28 (14.1 to 40.7) $X^2= 13.078, df=2; p=0.001$</p> <p>Scores improved in both groups over the study period. Median change in score: Group 1: -1 Group 2: -2 Mann-Whitney U test, $p=0.8$</p> <p>Excluding intervention: Group 1 (n=57): 310.3 (602.0) Group 2 (n=48): 188.8 (300.4) Mean difference (95% CI): 121.5 (-58.9 to 302.0)</p> <p>including intervention: Group 1: 594.1 (602.0) Group 2: 188.8 (300.4) Mean difference (95% CI): 405.4 (224.9 to 585.8) $P<0.001$</p>	<p>elicit uncertainty about choosing between alternatives, awareness of modifiable factors contributing to the uncertainty, and perceived effectiveness of decision making process. Higher scores indicated increased uncertainty in each subscale. Subscales combined to give a total decisional conflict score.</p>

Evidence Table 53 Economic evidence

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Annemans 2005¹⁷ UK</p> <p>Economic analysis: cost-effectiveness analysis</p> <p>Study design Decision analysis*</p> <p>Time horizon: 6 months</p> <p>Discount rates: Costs: NA Effects: NA</p>	<p>Patient group: patients hospitalised for acute urinary retention</p>	<p>Intervention 1: Alfuzosin 10mg once daily used for 3 days during the initial hospitalization followed by TWOC (mean duration 55hours). If TWOC is successful treatment with Alfuzosin for 6 months.</p> <p>Intervention 2: Immediate inpatient prostatectomy</p> <p>Intervention 3: Placebo followed by TWOC (mean duration 55hours) and placebo if TWOC is successful.</p>	Successful TWOC*	<p>Int 1: 62%</p> <p>Int 2: NA</p> <p>Int 3: 48%</p> <p>p value: 0.012</p>	<p>Funding: Sanofi-Aventis</p> <p>Limitations: Short follow-up.</p> <p>Additional outcomes: After successful TWOC, 17% of patients treated with Alfuzosin for 6 months require prostatectomy compared to 24% of patients treated with placebo.</p> <p>Notes: * based on the ALFAUR Study¹⁹⁵ **based on 2002 Reference Costs inflated to 2003 (inflator 1.035)</p>
			<p>Mean cost per patient over 6 months** 2002 GBP cost of hospitalisation, prostatectomy and TURP, drugs, unsuccessful TWOC (prostatectomy), tests.</p>	<p>Int 1: 2,029</p> <p>Int 2: 2,378</p> <p>Int 3: 2,921</p> <p>p value: NR</p>	
			<p>Incremental costs over 6 months (based on 1,000 Monte Carlo simulations)</p>	<p>Int 3 vs. Int 1: 349 (95% CI 64-624)</p> <p>Int 2 vs. Int 1: 892 (95% CI 644-1121)</p> <p>Int 2 vs. Int 3 : 543 (95% CI 228 - 776)</p> <p>p value : Sig</p>	
			<p>Cost-effectiveness cost per successful TWOC</p>	<p>Int 1 dominates Int 2 and 3</p>	
			<p>Sensitivity analysis Monte Carlo simulation</p>	<p>If the proportion of patients having an immediate prostatectomy after a failed TWOC is higher, Alfuzosin is more cost-saving.</p> <p>If surgery after successful TWOC is done in an elective setting, Alfuzosin is more cost saving.</p>	

Study details	Patients	Interventions*	Outcome measures	Effect size	Comments
<p>DiSantostefano 2006⁷¹ USA</p> <p>Economic analysis: Cost-utility analysis</p> <p>Study design Decision analysis</p> <p>Time horizon: 20 years</p> <p>Discount rates: Costs: 3% Effects: 3%</p>	<p>Patient group: men aged 65 years with moderate to severe LUTS and uncomplicated BPH, with no contraindications to any of the drugs.</p> <p>Group A: moderate symptoms (IPSS 8-19)</p> <p>Group B: severe symptoms (IPSS 20-35)</p>	<p>Intervention 1: Watchful waiting (WW)</p> <p>Intervention 2: Alpha-blockers (AB)</p> <p>Intervention 3: 5-Alpha reductase inhibitors (5-ARI)</p> <p>Intervention 4: High-energy transurethral microwave thermotherapy (TUMT)</p> <p>Intervention 5: Transurethral resection of the prostate (TURP)</p>	QALYs – Group A	<p>Intervention 1: 10.68 Intervention 2: 10.76 Intervention 3: 10.71 Intervention 4: 10.69 Intervention 5: 10.63 p value: NR</p>	<p>Funding: National Research Service Award Institutional Training Grant from the Institute of Aging; grant from the Agency for Healthcare Research and Quality. Conflict of Interest: the author is an employee of GlaxoSmithKline.</p> <p>Limitations: Partial applicability. The lack of long-term studies and differences between patient populations might have biased the results in favour of pharmaceuticals.</p> <p>Notes: * Combination of AB and 5-ARI was an additional intervention compared in the study but it was excluded because its effectiveness was based only on experts opinion. ** GBP calculated by using the 2008 PPP</p>
			QALYs – Group B	<p>Intervention 1: 9.79 Intervention 2: 9.88 Intervention 3: 9.83 Intervention 4: 10.30 Intervention 5: 10.47 p value: NR</p>	
			Mean cost per patient** – Group A 2004 USD, cost of GP visits, tests, drugs, surgery, complications (strictures, incontinence)	<p>Intervention 1: \$ 4,419 (£ 2,793) Intervention 2: \$ 6,666 (£ 4,213) Intervention 3: \$ 8,891 (£ 5,619) Intervention 4: \$ 7,982 (£ 5,045) Intervention 5: \$ 8,599 (£ 5,435) p value: NR</p>	
			Mean cost per patient** – Group B 2004 USD, cost of GP visits, tests, drugs, surgery, complications (strictures, incontinence)	<p>Intervention 1: \$ 4,403 (£ 2,783) Intervention 2: \$ 6,664 (£ 4,212) Intervention 3: \$ 8,888 (£ 5,617) Intervention 4: \$ 7,983 (£ 5,045) Intervention 5: \$ 8,558 (£ 5,409) p value: NR</p>	
			Cost-effectiveness** – incremental cost per QALY	<p>Group A Int 2 vs. Int 1: \$ 28,088 (£17,752) Int 3, 4 and 5 are dominated by Int 2. Int 6 is dominated by Int 5.</p> <p>Group B Int 2 vs. Int 1: \$ 25,122 (£ 15,877) Int 3 is dominated by Int 2. Int 4 vs. Int 2: \$ 3,140 (£ 1,984) Int 5 vs. Int 2: \$ 3,210 (£ 2,029) Int 5 vs. Int 1: \$ 6,110 (£ 3,861) Int 5 vs. Int 4: \$ 3,382 (£ 2,137)</p>	
			Sensitivity analysis One-way sensitivity analysis	If switching between treatments was not permitted, TURP would cost \$30,204 (£ 19,090) more than AB for each QALY	

Study details	Patients	Interventions*	Outcome measures	Effect size	Comments
			<p>Probabilistic sensitivity analysis</p>	<p>gained for moderate symptoms patients.</p> <p>The overall results did not change with the age of the patient.</p> <p>If effectiveness of TUMT is set equal to TURP, TUMT dominates TURP.</p> <p>For a willingness to pay equal to \$50,000 alpha-blockers have about a 70% probability of being cost-effective for patients with moderate symptoms. For the same willingness to pay, TURP had almost a 90% probability of being cost-effective for patients with severe symptoms.</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Fader 2008⁸⁶ UK</p> <p>Economic analysis: Cost-effectiveness analysis</p> <p>Study design RCT (cross-over)*</p> <p>Duration of follow-up: One month</p> <p>Discount rates: Costs: NA Effects: NA</p>	<p>Patient group: moderate/heavily incontinent adults (urinary or urinary/faecal) living in the community</p> <p>All patients N: 85 IPSS: NR Age (mean): 52.8 M/F: 49/36 Drop outs: 0</p>	<p>Intervention 1: Insert</p>	<p>Proportion of patients willing to buy a product used during the day if they had to bear the cost</p>	<p>Int 1: 39% Int 2: 50% Int 3: 43% Int 4: 39% Int 5: 38% p value: NR</p>	<p>Funding: commissioned by the Health Technology Assessment Programme. Some of the authors have received research grant money and travel grant money from SCA AB (absorbent pad manufacturing company)</p> <p>Limitations: The study included women and faecal incontinence as well. Not a full economic evaluation. Effectiveness was not measured in terms of any of the clinical outcomes included in our Guideline.</p> <p>Notes: *crossover design in which each participant tested all products within their group in random order. Only trial 2a is included and reported. ** scale from 0 – 100 to assess patients' preference for a product. *** Visual Analogue Scale score is not a clinical outcome of interest and an incremental cost-effectiveness analysis based on this outcome would not be useful.</p>
		<p>Intervention 2: Diaper</p>	<p>Proportion of patients willing to buy a product used during the night if they had to bear the cost</p>	<p>Int 1: 33% Int 2: 52% Int 3: 39% Int 4: 33% Int 5: 53% p value: NR</p>	
		<p>Intervention 3: Pull-up</p>	<p>Mean Visual Analogue Scale score** (day use – night use)</p>	<p>Int 1: 48 – 53 Int 2: 66 – 64 Int 3: 73 – 62 Int 4: 60 – 54 Int 5: 34 – 43 p value: NR</p>	
		<p>Intervention 4: T-shaped</p>	<p>Mean monthly cost per patient (day – night) 2005 GBP, cost of supplying the product, assuming three products per day and one per night are used. Cost of laundering washable products is not included.</p>	<p>Int 1: £44 - £23 Int 2: £47 - £15 Int 3: £79 - £25 Int 4: £75 - £25 Int 5: £9 - £6 p value: NR</p>	
		<p>Intervention 5: Washables</p>	<p>Cost-effectiveness</p>	<p>NA***</p>	
		<p>Sensitivity analysis</p>	<p>Different types of products within the same category have different costs and performance. The results are very sensitive to these variations.</p>		

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Fehrling2007 ⁹⁰ Sweden Economic analysis: Cost consequences analysis Study design Within group comparison Duration of follow-up: 3 months Discount rates: Costs: NA Effects: NA	Patient group: patients with an overactive bladder with or without incontinence All patients N: 60 IPSS: Age: the majority was 70 or older M/F: 31/29 Drop outs: 0	Treatment: 10 session (twice weekly for 5 weeks) of Maximal Functional Electrical Stimulation (MFES) at the highest tolerable amplitude	Number of patients with: up to 8 voids per day > 8voids per day - NR	Before treatment: 11 – 44 – 5 After treatment: 11 – 30 – 19 p value: NR	Funding: Swedish Research Council, Sahlgrenska university Hospital, and the Martha and Gustaf Agrens research Foundation. Limitations: Within group study. The outcomes are not clear-cut. Only the cost of the intervention is considered. Mixed male and female population. Notes: * the total sum is 61 while N=60 **Cost of treatment for each successfully treated patient is reported (€17,000) but success is not defined. *** calculated by using the 2008 PPP for Germany
			Number of patients with the following degree of leakage: No leakage - Minor - Moderate - Severe- NR	Before treatment*: 17 – 11 – 16 – 13 – 4 After treatment: 21 – 12 – 10 – 11 – 6 p value: NR	
			Mean cost per patient 2007 Euro, cost of 10 sessions.	Before treatment: NR After treatment: €3,500 (£2,640***) p value:	
			Cost-effectiveness	NR**	
			Sensitivity analysis)	NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Fraundorfer2001 ⁹⁷ New Zealand Economic analysis: Cost consequences Study design RCT* ¹⁰⁴ Duration of follow-up: 1 year Discount rates: Costs: NA Effects: NA	Patient group: men with urodynamically proved outflow obstruction due to BPH, AUA score of 8 or greater, independent peak urinary flow rate (Qmax) of 15 mL/s or less, and bladder outflow obstruction confirmed by pressure flow urodynamic studies (Schafer grade 2 or more). <u>All patients</u> N: 120 <u>Group 1</u> N: 61 Mean (±SD) Age: 66.9±6.5 <u>Group 2</u> N: 59 Mean (±SD) Age: 66.8±7.4	Group 1 Holmium laser resection (HoLRP) Group 2 TURP	Qmax (mL/s) ± SD	Group 1: 25.2 ± 11.9 Group 2: 20.4 ± 8.5 p value: <0.05	Funding: partially funded by Coherent Medical Group. Clinical study authors have financial interest and/or other relationship with Lumenis, Inc. Limitations: Not a full economic evaluation. Partially applicable. In real practice HoLEP might be less successful as it requires high level of skills and experience. Additional outcomes: Group 1 had a shorter LOS and lower complication rate. Notes: * The two year follow-up study ³¹⁸ was reviewed for clinical effectiveness **calculated by using the 2008 PPP
			AUA score	Group 1: 4.2 ± 6.0 Group 2: 4.3 ± 4.1 p value: Not Sig	
			Mean cost per patient 2001 NZD cost of consumables, hospital facility use, operations, clinic visits, capital equipment, and unplanned events.	Group 1: 2,012 (£857**) Group 2: 2,663 (£1,134**) p value: NR	
			Cost-effectiveness	NA	
			Sensitivity analysis	NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Hillman 1996 ¹¹⁷ USA Economic analysis: Cost consequences and cost-effectiveness Study design Multicentre RCT ²⁶² Duration of follow-up: 12 months Discount rates: Costs: NA Effects: NA	Patient group: men 55 years or older with a clinical diagnosis of BPH and at least moderate IPSS. All patients N: 2084 IPSS: 20.1 Age (mean and range): 65.7 (46 – 94) Drop outs*: 867 Group 1 N: 1053 (1010 in economic analysis) IPSS: 20.1 Age (mean): 65.7 Drop outs*: 396 Group 2 N: 1031 (983 in economic analysis) IPSS: 20.1 Age (mean): 65.7 Drop outs*: 471	Group 1: Alpha-blockers (Terazosin), 1mg daily for 3 days followed by 2mg daily for the remainder of the first 4 weeks. The medication dose was titrated upward at the investigator's discretion until a satisfactory response was achieved (improvement of 35% or more of IPSS). Group 2: Placebo	Mean change in IPSS ± SE	Group 1: -7.6 ± 0.2 Group 2: -3.7 ± 0.2 p value: <0.001	Funding: Abbott Laboratories, Abbott Park, Illinois. Limitations: Partial applicability. Placebo was used instead of watchful waiting. Short follow up. Notes: *Patients withdrawn because of adverse events and lack of efficacy were respectively 168 and 93 in group 1, and 114 and 220 in group 2 (p<0.001). **Calculated by using the 2008 PPP *** calculated by NCGC
			Mean change in IPSS – Quality of Life ± SE	Group 1: -3.6 ± 0.1 Group 2: -1.8 ± 0.1 p value: <0.001	
			Mean cost per patient 1992 USD, cost of visits (home, GP and urologist), inpatient care, medication.	Group 1: \$2,932 (£1,865**) Group 2: \$3,404 (£2,165**) p value: NR	
			Cost-effectiveness*** incremental cost per IPSS point change	Group 1 dominates Group 2	
			Sensitivity analysis one-way SA	Overall results were not sensitive to outlier costs, costs assigned by patient-reported events, regional vs. satellite patients, costs of patients completing a full year of therapy, costs of improperly randomised patients.	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Johansen 2007^{1,27} Norway</p> <p>Economic analysis: cost analysis</p> <p>Study design decision analysis*</p> <p>Time horizon: 4 years</p> <p>Discount rates: Costs: 5% Effects: NA</p>	<p>Patient group: men with BPH</p>	<p>Intervention 1: Alpha-blockers (Tamsulosin)</p> <p>Intervention 2: 5-Alpha-reductase inhibitors (Dutasteride and Finasteride)</p> <p>Intervention 3: TURP</p>	<p>Mean cost per patient over 4 years 2006 NOK, cost of drugs, tests, visits to GP, pre-TURP visits to urologist, TURP, surgical follow-up, prostate cancer evaluation following TURP, post-TURP antibiotics, cost of AUR.</p>	<p>Int 1: 16,933 (£1,219**) Int 2***: 13,946 (£ 1,004**) Int 3: 46,109 (£ 3,320**) p value: NR</p>	<p>Funding: NR. One of the authors was an employee of GlaxoSmithKline.</p> <p>Limitations: Risk of AUR and TURP for Tamsulosin was assumed to be equal to the placebo arm of the trials.</p> <p>Notes: *improvement rates, risk of AUR and TURP were taken from Phase-III trials¹ for Dutasteride, assumed to be equal for Finasteride. Risk of AUR and TURP of Tamsulosin was assumed to be equal to the placebo arm of those trials. Improvement rate of Tamsulosin was obtained from Phase-III trials and improvement rate of TURP was based on clinical opinion. ** Calculated by using the 2008 PPP ***cost of Dutasteride. Finasteride was more costly than Dutasteride but less costly than Tamsulosin.</p>
			<p>Cost-effectiveness</p>	<p>NA</p>	
			<p>Sensitivity analysis One-way and multi-way SA</p>	<p>The overall results were not sensitive to the following changes in one-way, two-way and multi-way SA:</p> <ul style="list-style-type: none"> - Time-horizon increased to lifetime. - Decrease or increase costs of TURP and AUR by 10%. - Inclusion of indirect costs. - Probability of AUR decreased by 10% after TURP/any intervention. - Probability of TURP after AUR reduced by 25%. - Decrease symptoms improvement by 10%. - Change in discount rate (0-8%). 	

¹ <http://www.gsk-clinicalstudyregister.com/files/pdf/883.pdf>, <http://www.gsk-clinicalstudyregister.com/files/pdf/895.pdf>, <http://www.gsk-clinicalstudyregister.com/files/pdf/3241.pdf>

Study details	Patients	Interventions*	Outcome measures	Effect size	Comments
<p>Johnson 1999¹²⁸ UK</p> <p>Economic analysis: cost-consequences analysis</p> <p>Study design decision analysis</p> <p>Time horizon: 5 years</p> <p>Discount rates: Costs: 6% Effects: 6%</p>	<p>Patient group: 60 years old patients with uncomplicated moderate to severe benign prostatic hyperplasia</p>	<p>Intervention 1: Watchful waiting. If ineffective it will be followed by second line (Doxazosin or Finasteride) and if necessary surgery.</p> <p>Intervention 2: Alpha-blockers (Doxazosin). If ineffective or have side effects it will be followed by second line (Finasteride or watchful waiting) and if necessary surgery.</p> <p>Intervention 3: 5-alpha-reductase inhibitors (Finasteride). If ineffective or have side effects it will be followed by second line (Doxazosin or watchful waiting) and if necessary surgery.</p>	<p>Patients discontinuing treatment over 5 years</p>	<p>Int 1: 46.0% Int 2: 39.1% Int 3: 42.0% p value: NR</p>	<p>Funding: Pfizer International</p> <p>Limitations: It was not clear how the response-years gained were calculated.</p> <p>Notes: * Surgery was excluded from the interventions compared as this was a mix of TURP and open prostatectomy. ** Obtained from the meta-analysis described by the American Agency for Health Care Policy and Research</p>
			<p>Patients with improved symptoms**</p>	<p>Int 1: 42% Int 2: 74% Int 3: 67% p value: NR</p>	
			<p>Improvement in symptom score from baseline**</p>	<p>Int 1: 32% Int 2: 48% Int 3: 31% p value: NR</p>	
			<p>Response-years gained</p>	<p>Int 1: 0.57 Int 2: 0.81 Int 3: 0.60 p value: NR</p>	
			<p>Mean cost per patient over 5 years 1999 GBP; cost of GP and urologist consultations, laboratory procedures, examinations, medications, surgical procedures, complications.</p>	<p>Int 1: £791 Int 2: £1427 Int 3: £1720 p value: NR</p>	
			<p>Cost-effectiveness</p>	<p>NR</p>	
			<p>Sensitivity analysis One-way SA</p>	<p>Results not sensitive to cost of surgery, response rates, discontinuation rates, response degree, and time horizon</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Keoghane2000 ¹⁴² UK Economic analysis: cost-effectiveness analysis Study design RCT Duration of follow-up: 36 months (costs only 24 months) Discount rates: Costs: NR Effects: NR	Patient group: all patients presenting for TURP who had not undergone previous surgery. <u>All patients</u> N: 152 (100 for cost analysis) Drop outs: NR <u>Group 1</u> N: 47 for cost analysis AUA score (SD): 19.9 (7.7) <u>Group 2</u> N: 53 for cost analysis AUA score (SD): 19.4 (6.5)	Group 1 Vaporisation using MD60 Nd:YAG (Selected Laser Technologies) with 600 µm fibre incorporating sapphire-tipped probe. Irrigation using saline. Group 2 TURP in standard manner using Storz equipment and irrigation with glycine	Mean change in AUA 7 symptom score from baseline at 12 months (±SD)	Group1: 10.9 ± 8.4 (n=44) Group 2: 13.3 ± 7.8 (n=53) p value: not Sig (NCGC-ACC t-test)	Funding: Oxford Regional Health Authority Limitations: Surgeons had limited experience with the laser technique which may have caused the high failure rate with this treatment. Additional outcomes: Duration of catheterisation and complications favour Contact Laser. Reoperation rate was 18% in Group 1 and 9% in Group 2. Inpatient stay was 3.5 days in Group 1 and 3.9 days in Group 2. Notes: * In the study prices were up-rated using the NHS hospital and community price index.
			Mean change in AUA 7 symptom score from baseline at 24 months (±SD)	Group1: 11.7 ± 9.7 (n=35) Group 2: 13.7 ± 7.7 (n=47) p value: not Sig (NCGC-ACC t-test)	
			Mean change in AUA 7 symptom score from baseline at 36 months (±SD)	Group1: 11.0 ± 9.7 (n=37) Group 2: 12.9 ± 7.9 (n=41) p value: not Sig (NCGC-ACC t-test)	
			Change in flow rate (Qmax) from baseline at 3 years	Group1: 1.8 ± 6.2 (n=24) Group 2: 2.1 ± 6.9 (n=24) p value: Not Sig (NCGC-ACC t-test)	
			Mean cost per patient at 2 years 1997 GBP*, cost of operation, hospitalisation, outpatient visits, GP and nurse visits, re- operation, capital costs and overheads.	Group 1: £1,252 Group 2: £971 p value: Sig	
			Cost-effectiveness cost per change in AUA score	TURP is dominant	
			Sensitivity analysis One way	If inpatient stay in Group 1 is reduced to 1.5 days laser becomes less costly by £50.	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Lourenco 2008 ¹⁷⁴ UK Economic analysis: Cost-utility analysis Study design Decision analysis Time horizon: 10 years Discount rates: Costs: 3.5% Effects: 3.5%	Patient group: Men at the age of 70 years with BPE, presence of LUTS with a measure of IPSS>7, no complications and TURP indicated (medical treatment either contraindicated or failed). Mean start age 70 years.	Intervention 1: TUVP	QALYs*	Int 1: 0.3668 Int 2: 0.3625 Int 3: 0.3679 Int 4: 0.3673 Int 5: 0.3631 Int 6: 0.3684 Int 7: 0.3684 Int 8: 0.3684 p value: NR	Funding: NHS R&D Health Technology Assessment Programme Limitations: Cost of equipment was included only for some strategies. Duration and cost of operations were equal in all the strategies. Training costs not included. Some interventions (TURP) are used to identify prostate cancer. Additional diagnostic tests would be necessary of another strategy is adopted. Additional outcomes: Other sequences of treatments starting with TURP or TUMT were dominated. When compared to TURP alone, only TUVP, KTP and all the strategies involving a second operation starting with TUMT are not cost-effective. Expected value of partial perfect information was £4,187,062 for TUVP
		Intervention 2: TUMT	Mean cost per patient* 2006 GBP, cost of procedure, short-term complications (acute urinary retention, bladder neck contracture or urethral stricture, blood transfusion, transurethral syndrome, urinary tract infections), long-term complications (incontinence: 95% oxybutinin, 5% artificial sphincter), equipment for KTP, HoLEP and TUMT only.	Int 1: £152 Int 2: £155 Int 3: £160 Int 4: £174 Int 5: £223 Int 6: £166 Int 7: £167 Int 8: £167 p value: NR	
		Intervention 3: HoLEP	Cost-effectiveness incremental cost per QALY	Int 3 vs. Int 1: £7,273 Int 6 vs. Int 3: £12,000 Int 2 dominated by Int 1. Int 3 vs. Int 2: £833. Int 4 dominated by Int 3, 6, 7, 8. Int 5 dominated by any interventions. Int 7 and 8 dominated by Int 6**.	
		Intervention 4: TURP	Sensitivity analysis Probabilistic sensitivity analysis	At the threshold of £20,000/QALY, Int 6 has a probability of being cost-effective of about 80%.	
		Intervention 5: KTP	One way sensitivity analysis	If LOS TURP is 2 days instead of 3 days, Int 8 is cost-effective. Results not sensitive to start age, utility of 'incontinence no remission' state = utility of 'incontinence remission' state, utility of IPSS<8 is 0.97 instead of 1, risk data from	
		Intervention 6: TUVP followed by HoLEP if it fails			
		Intervention 7: TUVP followed by TURP if it fails			
		Intervention 8: TUVP followed by repeated TURP if it fails			

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
				all studies instead of UK studies only, test for obstruction after TUVp.	epidemiology and £1,652,886 for HoLEP epidemiology. Notes: * results per patient of Monte Carlo simulation with 10,000 samples where 25,000 new individuals enter the model each year. ** Int 8 vs. 6 ICER=£90,576/QALY when results are calculated per population

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
McDonald 2004 ¹⁹² Canada Economic analysis: Cost-utility analysis Study design Decision analysis* Time horizon: 15 years Discount rates: Costs: 5% Effects: 5%	Patient group: men 65 years old with moderate to severe symptoms of BPH and an enlarged prostate as determined by digital rectal examination who choose not to undergo immediate surgical treatment.	Intervention 1: Watchful waiting (WW) Intervention 2: Alpha-blockers (Doxazosin) Intervention 3: 5-alpha-reductase inhibitors (Finasteride) Intervention 4: Combination therapy with Doxazosin and Finasteride.	QALYs gained Mean cost per patient** 2003 CAD, cost of drugs (including 10% pharmacy mark-up charge and dispensing fee), visits (one full and one partial per year plus two partial for Group 1), hospitalisation, surgery, surgical complications, tests. Cost-effectiveness ** incremental cost per QALY gained Sensitivity analysis One way SA.	Int 1: 8.608 Int 2: 8.787 Int 3: 8.709 Int 4: 8.930 p value: NR Int 1: \$2,254 (£ 1,181) Int 2: \$4,615 (£ 2,418) Int 3: \$6,167 (£ 3,231) Int 4: \$9,477 (£ 4,966) p value: NR Int 2 vs. Int 1***: \$13,190 (£ 6,912) Int 3 dominated by Int 2. Int 4 vs. Int 2: \$34,000 (£ 17,816) Considering only patients with PSA > 1.3 ng/ml or PSA > 3.2 ng/ml the results were similar. Results were not sensitive to discounting, probability of TURP following AUR, cost of TURP, cost of AUR. Combination is no longer cost-effective when AUR rates are obtained from MTOPS instead of PLESS, treatment effect is decreased by 50%, or QALY weights from Baladi ¹⁹⁶ are used. Finasteride is more cost-effective than Doxazosin if it improves IPSS past year 4 by 2 points.	Funding: Merck Frosst Canada Ltd. Limitations: Partially applicable. Additional outcomes: Incremental cost per AUR averted and incremental cost per TURP averted. Notes: * based mainly on the PLESS ²⁵⁶ and MTOPS studies ¹⁹¹ ** GBP calculated by using the 2008 PPP *** calculated by NCGC

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Medicare Services Advisory Committee¹⁹⁸ Australia</p> <p>Economic analysis: cost-utility analysis</p> <p>Study design Decision analysis</p> <p>Time horizon: 20 years</p> <p>Discount rates: Costs: 5% Effects: 5%</p>	<p>Patient group: Patients with symptomatic benign prostatic hyperplasia.</p>	<p>Intervention 1: TUNA</p> <p>Intervention 2: TURP</p>	QALY	<p>Int 1: 12.2869 Int 2: 12.3082 p value: NR</p>	<p>Funding: Report prepared from the National Health and Medical Research Council Clinical Trials Centre, University of Sydney for the Medical Services Advisory Committee.</p> <p>Limitations: Utilities were obtained from expert opinion and not elicited with recognised methods.</p> <p>Notes: * Calculated by using the 2008 PPP</p>
			<p>Mean cost per patient 1999 AUD, cost of procedures, cost of side effects, cost of treatment failure (GP visits, surgery, hospitalisation, medical treatment).</p>	<p>Int 1: \$8,296 (£4,165*) Int 2: \$6,910 (£3,469*) p value: NR</p>	
			<p>Cost-effectiveness cost per QALY gained</p>	TURP dominates TUNA	
			<p>Sensitivity analysis One-way SA</p>	<p>TUNA is cost-effective when either: probability that TURP fails within 6 months $\geq 20\%$; time horizon = 5 years; annual failure rate of TUNA $\leq 2.4\%$; probability of having TURP after TUNA fails = 100%</p>	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Murray2001 ²⁰⁷ UK	Patient group: Men with benign prostatic hypertrophy in 33 general practices in the UK.	Group 1: Interactive multimedia programme with booklet and printed summary. Treatment options discussed were surgery, balloon dilatation of the prostate, drugs, and watchful waiting. Information comprised probabilities of the risks and benefits of each treatment, calculated on the basis of information on age, severity of symptoms, and general health entered by the patient at the beginning of the session. All patients saw the core interactive video disc, lasting about 45 minutes; viewing optional sections for further information took up to 60 min. more. A research nurse started the programme, taught the patient how to use it, and then withdrew.	Mean (SD) decisional conflict score at nine months	Group 1: 2.23 (0.38) Group 2: 2.55 (0.50) p value: sig	Funding: NHS national research and development programme, the BUPA Foundation, and the King's Fund.
Economic analysis: cost consequences analysis	All patients N: 112 Drop outs: 10		Median change in American Urological Association scores	Group 1: -1 Group 2: -2 p value: 0.8	Limitations: Results on EQ-5D scores were not reported. The intervention might be different to the clinical practice with a consequent overestimation of costs.
Study design RCT	Group 1 N: 57 Age (mean +/- SD): 63.7 +/- 8.4 Drop outs: 3 Mean (SD) American Urological Association score: 15.64 (6.57)		Mean cost per patient 1999 GBP, Cost of equipment and staff time, consultations with GPs, referrals to urologists, other referrals, drugs, tests, diagnostic and surgical procedures.	Group 1: 594 Group 2: 188 p value: <0.001	Additional outcomes: No difference in health utility scores (EQ-5D) and anxiety scores (data not provided). Mean decisional conflict score at 3 months (-0.3). GPs and patients' perception of decision making at 3months was significantly different between the two groups with higher proportion of GPs and patients perceiving that the treatment decision had been mainly or only by the patients in Group 1.
Duration of follow-up: 9 months	Group 2 N: 55* Age (mean +/- SD): 63.9 +/- 8.4 Drop outs: 7 Mean (SD) American Urological Association score: 14.85 (7.10)	Group 2: Normal care from GP practitioner.	Cost-effectiveness	NR	
Discount rates: Costs: NA Effects: NA			Sensitivity analysis	NR	Notes: *Only 48 included in the economic analysis

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Nathan 1996²¹¹ UK</p> <p>Economic analysis: cost consequence</p> <p>Study design (e.g. RCT, Decision analysis, etc)</p> <p>Duration of follow-up: 3 months</p> <p>Discount rates: Costs: Effects:</p>	<p>Patient group: men requiring TURP</p> <p>All patients N: 40 Drop outs: 0</p> <p>Group 1 N: 20 Mean age (range): 65.4 (57-77) Mean IPSS score: 21.9 ± 4.2 Mean IPSS QoL ± SD: 4.9 ± 0.7 Mean Qmax ml/s (± SD): 10.2 ± 4.4 Drop outs: 0</p> <p>Group 2: N: 30 Mean age (range): 69.2 (57-81) Mean IPSS score: 17.0 ± 4.3 Mean IPSS QoL ± SD: 4.9 ± 0.7 Mean Qmax ml/s (± SD): 7.2 ± 3.5 Drop outs: 0</p>	<p>Group 1: Transurethral electrovaporisation of the prostate (TVP)</p> <p>Group 2: TURP</p>	Mean IPSS score at 3 months (follow up interval not clear)	Group 1: 2.86 ± 2.8 Group 2: 3.1 ± 2.3 p value: NR	<p>Funding: NR</p> <p>Limitations: Cost components included in the analysis were only those that significantly differed between interventions.</p> <p>Additional outcomes: There were more complications in the TURP group. There was no statistically significant or appreciable difference in the success rates among the two groups.</p>
			Mean IPSS QoL score at 3 months (follow up interval not clear)	Group 1: 0.5 ± 7 Group 2: 0.9 ± 0.9 p value: NR	
			Mean Qmax ± SD mL/s at 3 months (follow up interval not clear)	Group 1: 21.3 ± 5.9 Group 2: 20.6 ± 2.6 p value: NR	
			Mean cost per patient 1996 GBP, cost of fibres and consumables, transfusions, and hospital stay.	Group 1: £1,730 Group 2: £2,373 p value: NR	
			Cost-effectiveness	NR	
			Sensitivity analysis	NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Noble 2002²²² UK</p> <p>Economic analysis: Cost-consequences analysis</p> <p>Study design RCT⁷⁴</p> <p>Duration of follow-up: 7.5 months</p> <p>Discount rates: Costs: NA Effects: NA</p>	<p>Patient group: men with uncomplicated lower urinary tract symptoms (no acute or chronic urinary retention)</p> <p>All patients N: 340 Drop outs:</p> <p>Group 1 N: 117 Dropouts: 1/117 Age, mean (\pmSD): 67.4\pm8.1 IPSS, mean (\pmSD): 19.1\pm6.6 IPSS-QoL, median(range): 4(2-6)</p> <p>Group 2 N: 117 Dropouts: 2/117 Age, mean (\pmSD): 66.4\pm7.9 IPSS, mean (\pmSD): 19.2\pm6.7 IPSS-QoL, median(range): 4(0-6)</p> <p>Group 3 N: 106 Dropouts: 5/106 Age, mean (\pmSD): 67.2\pm7.8 IPSS, mean (\pmSD): 18.8\pm6.5 IPSS-QoL, median(range): 4(1-6)</p>	<p>Group 1: Laser therapy with a noncontact side firing neodymium:YAG probe</p> <p>Group 2: Standard transurethral prostate resection</p> <p>Group 3: conservative management</p>	Mean difference in IPSS from baseline	<p>Group 1: -10.8 Group 2: -12.3 Group 3: -1.3 p value: NR</p>	<p>Funding: Bard UK provided the laser fibres. South West and Northern Regional National Health Service Research and Development Directorates.</p> <p>Limitations: Resource use data were available only for 30% of the patients population. The conclusions of the study were incorrect.</p> <p>Additional outcomes: Patient costs were higher for noncontact laser.</p> <p>Notes: * calculated by NCGC using mean cost and mean change in health-related quality of life utility</p>
			Mean difference in IPSS quality of life from baseline	<p>Group 1: -1.9 Group 2: -2.2 Group 3: -1.3 p value: NR</p>	
			Mean change in QALY from baseline	<p>Group 1: 0.044 Group 2: 0.016 Group 3: -0.001 p value: NR</p>	
			Mean cost per patient 1998 GBP, cost of resources used in investigations, staff time, equipment, medication, hospital stay, rehospitalisation for catheter-free trial, other rehospitalisation, outpatient visits, GP and nursing visits, consumables (catheter bags, pads and other aids)	<p>Group 1: £1,223 Group 2: £928 Group 3: £45 p value: NR</p>	
			Cost-effectiveness* cost per QALY gained	<p>Group 1 vs. Group 2: £10,536 Group 1 vs. Group 3: £26,178</p>	
			Sensitivity analysis one-way	Cost of probes, their multiple use, and machinery lifetime were varied with no considerable difference in results.	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Norby2002 ²²³ Denmark Economic analysis: CEA Study design RCT ²²⁴ Duration of follow-up: 6 months Discount rates: Costs: NA Effects: NA	Patient group: Men \geq 50 years between May 1996 and November 1999. <u>All patients*</u> N: 113 <u>Group 1</u> N: 45 IPSS (\pmSD): 21.4 \pm 5.8 <u>Group 2</u> N: 46 IPSS (\pmSD): 20.5 \pm 5.7	Group 1: Interstitial laser coagulation (ILC). Group 2: Transurethral microwave thermotherapy (TUMT).	Mean difference in IPSS at 6 months from baseline (\pm SD)	Group 1: 12.0 \pm 7.5 Group 2: 11.2 \pm 9.2 p value: Not sig	Funding: Vejle County, Denmark. Limitations: Small sample size for economic analysis. Short follow-up. Limited applicability. Notes: * 22 patients were randomised to a mix of TUIP and TURP and therefore excluded. In the results this group dominates Group 1. **ITT analysis was used for clinical outcomes but not for costs **Data collected in 20 patients only. *** Calculated by using the 2008 PPP ****Incremental analysis done by NCGC
			Mean cost per patient** 1999 DKK, cost of hospitalisation, medications, examinations, follow-up visits, GP visits, nurse visits, and re-operations.	Group 1: 14,398 (£1,152 ^{***}) Group 2: 10,508 (£841 ^{***}) p value: NR	
			Cost-effectiveness**** Group 1 vs. Group 2: DKK cost per 1-point of reduction in IPSS	Group 1 vs. Group 2: DKK 4,862 (£ 388 ^{***}) per point	
			Sensitivity analysis One way	If TUMT catheters were reused once, Group 1 vs. Group 2 ICER = DKK 7,981 (£ 638 ^{***}) If ITT analysis is applied, Group 1 vs. Group 2 ICER = DKK 4,161 (£ 332 ^{***})	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
Salonia 2006 ²⁶⁷ Italy Economic analysis: cost analysis Study design RCT Duration of follow-up: NR Discount rates: Costs: NR Effects: NR	Patient group: consecutive patients with symptomatic benign prostatic hyperplasia in a large prostate (70 to 220 g) and documented bladder outlet obstruction. <u>All patients</u> N: 63 Group 1 N: 29 IPSS: 21.6 Age (mean): 68.0 Drop outs: Group 2 N: 34 IPSS: 19.6 Age (mean): 67.4 Drop outs:	Group 1: Open prostatectomy Group 2: HoLEP	Operative time (minutes)	Group 1: 57.5 Group 2: 73.4 p value: 0.002	Funding: Scientific Institute San Raffaele Hospital, Milan Limitations: Partial applicability. Additional outcomes: The amount of unplanned events was not significantly different. Notes: *calculated by using the 2008 PPP
			Catheterisation time (hours)	Group 1: 106.3 Group 2: 35.3 p value: 0.0001	
			Hospital stay (hours)	Group 1: 131.0 Group 2: 64.6 p value: <0.0001	
			Mean cost per patient 2004 Euro, costs associated with the procedures (operating room time, disposables, blood transfusion) and hospital stay. Medical salaries were not included. Capital cost for HoLEP was 85% of actual capital cost. Holmium fibres were used at least 10 times.	Group 1: 2,869 (£2,079*) Group 2: 2,356 (£1,708*) p value: NR	
			Cost-effectiveness	NR	
			Sensitivity analysis	NR	

Study details	Patients	Interventions	Outcome measures	Effect size	Comments
<p>Stovsky2006²⁸⁸ USA</p> <p>Economic analysis: Cost consequences analysis</p> <p>Study design Decision analysis</p> <p>Time horizon: 2 years</p> <p>Discount rates: Costs: NR Effects: NR</p>	<p>Patient group: patients with lower urinary tract symptoms indicative of BOH requiring procedural management with of the interventions indicated.</p>	<p>Intervention 1: Photoselective vaporisation</p> <p>Intervention 2: TURP</p> <p>Intervention 3: TUNA</p> <p>Intervention 4: TUMT Targis</p> <p>Intervention 5: TUMT Prostatron 2.5</p>	% change from baseline IPSS at 2 years	<p>Int 1: 76 Int 2: 66 Int 3: 44 Int 4: 46 Int 5: 39 p value: NR</p>	<p>Funding: All the authors had financial interest and/or relationship with Laserscope</p> <p>Limitations: Discount rate NR. Partially applicable: cost of inpatient stay in the USA is higher than in the UK, which favours laser.</p> <p>Additional outcomes: Qmax and QoL were also reported. The cost-effectiveness results did not change if those outcomes were used.</p> <p>Notes: * based on the assumption that PVP was performed in a hospital outpatient setting, TUNA and TUMT at a physician office site of service, TURP in a hospital inpatient setting, ILC at a physician office site of service (86%), ambulatory surgery centre (9%) and hospital outpatient setting (5%) ** converted into GBP by using the 2008 PPP ***incontinence, UTI, impotence, dysuria/irritative voiding, bladder neck stenosis/stricture, urinary retention, hematuria **** calculated by NCGC-ACC</p>
			% change from baseline Quality of Life score at 2 years	<p>Int 1: 83 Int 2: 73 Int 3: 61 Int 4: 52 Int 5: 24 p value: NR</p>	
			% Qmax at 2 years from baseline	<p>Int 1: 221 Int 2: 117 Int 3: 28 Int 4: 45 Int 5: 45 p value: NR</p>	
			Mean cost per patient* 2005 USD**, cost of intervention, follow-up care, adverse events***, re-treatment. Cost of pharmacological therapy not included.	<p>Int 1: \$ 3,589 (£ 2,315) Int 2: \$ 4,927 (£ 3,178) Int 3: \$ 6,179 (£ 3,985) Int 4: \$ 5,699 (£ 3,676) Int 5: \$ 5,488 (£ 3,562) p value: NR</p>	
			Cost-effectiveness**** cost per 1-point of %reduction in IPSS	<p>Intervention 2 dominates Interventions 3, 4 and 5. Intervention 1 dominates all the other interventions, including 2.</p>	
			Sensitivity analysis One way Threshold SA	<p>If ILC performed in a less costly setting, it is still dominated by PVP. When retreatment rate of PVP = 17%, PVP and TURP are cost equivalent.</p>	

Appendix E – Forest Plots

1.1	FREE UROFLOWMETRY (PEAK URINARY FLOW)	521
	<i>Figure E-1: Sensitivity and specificity of free uroflowmetry (Qmax) in the diagnosis of bladder outlet obstruction</i>	521
	<i>Figure E-2: Summary receiver operating characteristic (SROC) curve for uroflowmetry Qmax in the diagnosis of bladder outlet obstructions</i>	522
2.1	PELVIC FLOOR MUSCLE TRAINING (PFMT)	523
2.1.1	PFMT VS. CONTROL	523
	<i>Figure E-3: PFMT vs. Control: Number of post-prostatectomy men who were incontinent</i>	523
	<i>Figure E-4: PFMT vs. Control: Mean urine lost (g) per 24 hours (pad test) in post-prostatectomy men</i>	524
	<i>Figure E-5: PFMT vs. Control: Number of post-TURP men who were incontinent</i>	524
2.2	BIOFEEDBACK	525
2.2.1	BIOFEEDBACK + PFMT VS. CONTROL.....	525
	<i>Figure E-6: PFMT + Biofeedback vs. no intervention: Number of men who were incontinent at follow up</i>	525
2.3	ELECTRICAL STIMULATION (ES)	525
2.3.1	ES + PFMT VS. CONTROL.....	525
	<i>Figure E-7: ES + PFMT vs. no intervention: Number of men who were incontinent at follow up</i>	525
3.1	ALPHA-BLOCKERS	526
3.1.1	ALPHA-BLOCKERS VS. PLACEBO	526
	<i>Figure E-8: Alpha-blockers vs. Placebo: Symptom score (random effects analysis)</i>	526
	<i>Figure E-9: Alpha-blockers vs. Placebo: Qmax (ml/s) (random effects analysis)</i>	526
	<i>Figure E-10: Alpha-blockers vs. Placebo: Quality of life – IPSS question (random effects analysis)</i>	527
	<i>Figure E-11: Alpha-blockers vs. Placebo: Adverse events (cardiovascular and neurological) - asthenia (fatigue) and headache</i>	528
	<i>Figure E-12: Alpha-blockers vs. Placebo: Adverse events (cardiovascular and neurological) - postural hypotension and rhinitis</i>	529
	<i>Figure E-13: Alpha-blockers vs. Placebo: Adverse events - erectile dysfunction /impotence</i>	529
	<i>Figure E-14: Alpha-blockers vs. Placebo: Adverse events - dizziness and retrograde ejaculation (random effects analysis)</i>	530
	<i>Figure E-15: Alpha-blockers vs. Placebo: Withdrawal from study due to adverse events</i>	531
3.1.2	ALPHA-BLOCKERS VS. 5-ALPHA REDUCTASE INHIBITORS (5-ARI).....	532
	<i>Figure E-16: Alpha-blockers vs. 5-ARI: Symptom score</i>	532
	<i>Figure E-17: Alpha-blockers vs. 5-ARI: Quality of life (IPSS-question)</i>	532
	<i>Figure E-18: Alpha-blockers vs. 5-ARI: Qmax (ml/s)</i>	533
	<i>Figure E-19: Alpha-blockers vs. 5-ARI: Prostate volume (ml)</i>	533
	<i>Figure E-20: Alpha-blockers vs. 5-ARI: PSA (ng/ml)</i>	534
	<i>Figure E-21: Alpha-blockers vs. 5-ARI: Adverse events (cardiovascular or neurological)</i>	535
	<i>Figure E-22: Alpha-blockers vs. 5-ARI: Adverse events (sexual or urological)</i>	536
	<i>Figure E-23: Alpha-blockers vs. 5-ARI: Adverse events - postural hypotension and ejaculatory abnormality (random effects analysis)</i>	537
	<i>Figure E-24: Alpha-blockers vs. 5-ARI: Ejaculatory abnormality – subgroup analysis of tamsulosin and other alpha-blockers</i>	537
	<i>Figure E-25: Alpha-blockers vs. 5-ARI: Withdrawal from study due to adverse events (random effects analysis)</i>	538
3.1.3	ALPHA-BLOCKERS VS. ANTICHOLINERGICS	538
3.1.4	ALPHA-BLOCKERS VS. PHOSPHODIESTERASE 5-INHIBITORS (PDE5-I).....	538
3.2	5-ALPHA REDUCTASE INHIBITORS (5-ARI)	539
3.2.1	5-ARI VS. PLACEBO.....	539
	<i>Figure E-26: 5-ARI vs. Placebo: Symptom score at 3 months, 6 months 2 years and 4 years or longer (random effects analysis)</i>	539
	<i>Figure E-27: 5-ARI vs. Placebo: Symptom score at 2 years- subgroup analysis</i>	539
	<i>Figure E-28: 5-ARI vs. Placebo: Symptom score at 12 months and 3 years</i>	540

Figure E-29: 5-ARI vs. Placebo: Qmax (ml/s) at 3 months, 6 months, 2 years, 3 years and 4 years or longer	540
Figure E-30: 5-ARI vs. Placebo: Qmax (ml/s) at 12 months (random effects analysis)	541
Figure E-31: 5-ARI vs. Placebo: Prostate volume(ml) at 1 year follow up	541
Figure E-32: 5-ARI vs. Placebo: Prostate volume (ml) at 2 years follow up (random effects analysis)	541
Figure E-33: 5-ARI vs. Placebo: PSA (ng/ml) level at 2 year follow up	541
Figure E-34: 5-ARI vs. Placebo: Adverse events (cardiovascular and neurological)	542
Figure E-35: 5-ARI vs. Placebo: Adverse events (sexual and urological)	543
Figure E-36: 5-ARI vs. Placebo: Withdrawal from study due to adverse events	544
3.2.2 5-ALPHA REDUCTASE INHIBITORS (5-ARI) VS. ALPHA-BLOCKERS	544
3.3 ANTICHOLINERGICS	545
3.3.1 ANTICHOLINERGICS VS. PLACEBO	545
Figure E-37: Anticholinergics vs. Placebo: Adverse events	545
Figure E-38: Anticholinergics vs. Placebo: Withdrawal from study due to adverse events	545
3.3.2 ANTICHOLINERGICS VS. ALPHA-BLOCKERS	546
Figure E-39: Anticholinergics vs. Alpha-blockers: Adverse events	546
3.4 PHOSPHODIESTERASE-5-INHIBITORS (PDE5-I)	547
3.4.1 PDE5-I VS. PLACEBO	547
Figure E-40: PDE5-I vs. Placebo: Symptom score	547
Figure E-41: PDE5-I vs. Placebo: Quality of life (IPSS question)	547
Figure E-42: PDE5-I vs. Placebo: Qmax(ml/s)	547
Figure E-43: PDE5-I vs. Placebo: Adverse events	548
3.4.2 PDE5-I VS. ALPHA-BLOCKERS	549
Figure E-44: PDE5-I vs. Alpha-blockers: Symptom score	549
Figure E-45: PDE5-I vs. Alpha-blockers: Qmax (ml/s)	549
Figure E-46: PDE5-I vs. Alpha-blockers: Voiding frequency	549
Figure E-47: PDE5-I vs. Alpha-blockers: Nocturia	549
Figure E-48: PDE5-I vs. Alpha-blockers: Adverse events	550
3.5 DIURETICS	550
3.5.1 DIURETICS VS. PLACEBO	550
3.6 DESMOPRESSIN	550
3.6.1 DESMOPRESSIN VS. PLACEBO	550
Figure E-49: Desmopressin vs. Placebo: Adverse events	550
3.7 NSAIDS	551
3.7.1 NSAIDS VS. PLACEBO	551
Figure E-50: NSAIDs vs. Placebo: Symptom score at 1 month	551
Figure E-51: NSAIDs vs. Placebo: Qmax (ml/s) at 1 month	551
Figure E-52: NSAIDs vs. Placebo: Nocturia frequency at 1 month	551
Figure E-53: NSAIDs vs. Placebo: Adverse events (1 month follow up)	551
3.8 COMBINATION THERAPY: ALPHA-BLOCKERS PLUS 5-ALPHA REDUCTASE INHIBITORS(5-ARI)	552
3.8.1 COMBINATION (ALPHA-BLOCKERS + 5-ARI) VS. ALPHA-BLOCKERS	552
Figure E-54: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Symptom score	552
Figure E-55: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Qmax (ml/s)	552
Figure E-56: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Prostate volume(ml)	553
Figure E-57: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: PSA (ng/ml)	553
Figure E-58: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (cardiovascular or neurological)	554
Continued Figure E-58: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (cardiovascular or neurological)	555
Figure E-59: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (sexual or urological)	556
Figure E-60: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Withdrawal from study due to adverse events	557
3.8.2 COMBINATION (ALPHA-BLOCKERS + 5-ARI) VS. 5-ARI	557
Figure E-61: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Symptom score	557
Figure E-62: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Qmax(ml/s)	558

Figure E-63: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Prostate volume (ml)	558
Figure E-64: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: PSA (ng/ml)	559
Figure E-65: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (Cardiovascular or neurological)	560
Continued Figure E-65: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (cardiovascular or neurological)	561
Figure E-66: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (sexual or urological)	562
Figure E-67: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Ejaculatory abnormality (random effects analysis)	563
Figure E-68: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Ejaculatory abnormality subgroup analysis ..	563
Figure E-69: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Withdrawal from study due to adverse events (random effects analysis)	563
3.8.3 COMBINATION (ALPHA-BLOCKERS + 5-ARI) VS. PLACEBO	564
Figure E-70: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Symptom score	564
Figure E-71: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Qmax (ml/s)	564
Figure E-72: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Prostate volume (ml)	564
Figure E-73: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Change in PSA (ng/ml)	565
Figure E-74: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events (cardiovascular and neurological)	566
Figure E-75: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events – postural hypotension (random effects analysis)	567
Figure E-76: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events (sexual or urological)	567
Figure E-77: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Withdrawal from study due to adverse events (random effects analysis)	567
3.9 COMBINATION THERAPY : ANTI-CHOLINERGIC PLUS ALPHA-BLOCKERS	568
3.9.1 COMBINATION (ANTI-CHOLINERGIC + ALPHA-BLOCKERS) VS. ALPHA-BLOCKERS	568
Figure E-78: Combination (Anti-cholinergic + Alpha-blockers) vs. Alpha-blockers: Adverse events	568
3.9.2 ANTI-CHOLINERGIC ADDED ON TO ALPHA-BLOCKERS VS. ALPHA-BLOCKERS	568
Figure E-79: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Symptom score at 3 months	568
Figure E-80: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Quality of life (IPSS question) at 3 months	569
Figure E-81: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Qmax (ml/s) at 3 months	569
Figure E-82: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Adverse events (3-months follow up) ..	569
3.9.3 COMBINATION (ANTI-CHOLINERGIC + ALPHA-BLOCKERS) VS. ANTICHOLINERGICS	570
Figure E-83: Combination (Anti-cholinergic + Alpha-blockers) vs. Anticholinergics: Adverse events	570
3.9.4 COMBINATION (ANTI-CHOLINERGIC + ALPHA-BLOCKERS) VS. PLACEBO	571
Figure E-84: Combination (Anti-cholinergic + Alpha-blockers) vs. Placebo: adverse events	571
3.10 COMBINATION (PDE5-I + ALPHA-BLOCKERS)	571
3.10.1 COMBINATION (PDE5-I + ALPHA-BLOCKERS) VS. ALPHA-BLOCKERS	571
Figure E-85: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Symptom score	571
Figure E-86: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Quality of life (IPSS question) ..	572
Figure E-87: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Qmax (ml/s)	572
Figure E-88: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Frequency at 3-month	572
Figure E-89: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Nocturia at 3 months	572
Figure E-90: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events	573
Continued Figure E-90: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events	574
Figure E-91: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events resulting in withdrawal at 3-month	574
3.10.2 COMBINATION (PDE5-I + ALPHA-BLOCKERS) VS. PDE5-I	575
Figure E-92: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: symptom score (random effects analysis) ..	575
Figure E-93: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Quality of life (IPSS-QoL) up to 3-month ..	575
Figure E-94: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Qmax (ml/s) at 3-month	575
Figure E-95: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Frequency at 3-month	575
Figure E-96: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Nocturia at 3-month	576
Figure E-97: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Adverse events (only those resulting in withdrawals reported)	576
Figure E-98: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Withdrawal from study due to adverse events ..	576
4.1 HOLMIUM LASER ENUCLEATION OF THE PROSTATE (HOLEP)	577
4.1.1 HOLEP VS. TRANSURETHRAL RESECTION OF THE PROSTATE (TURP)	577
Figure E-99: HoLEP vs. TURP: Symptom score at 3 months, 36 months and 48 months	577

Figure E-100: HoLEP vs. TURP: Symptom score at 6, 12 and 24 months (random effects analysis).....	578
Figure E-101: HoLEP vs. TURP: Quality of life (IPSS question) – 3, 24 and 48 months.....	578
Figure E-102: HoLEP vs. TURP: Quality of life (IPSS question) – 6 to 12 months (random effects analysis) ..	579
Figure E-103: HoLEP vs. TURP: Qmax(ml/s) at 3 months and longest available follow up.....	579
Figure E-104: HoLEP vs. TURP: All cause mortality and complications.....	580
Continued Figure E-104: HoLEP vs. TURP: All cause mortality and complications	581
4.1.2 THULIUM LASER RESECTION VS. TURP.....	582
Figure E-105: Thulium laser resection vs. TURP: Symptom score – 6 months postoperatively.....	582
Figure E-106: Thulium laser resection vs. TURP: Symptom score – 12 months postoperatively	582
Figure E-107: Thulium laser resection vs. TURP: Qmax(ml/s) – 12 months postoperatively.....	582
Figure E-108: Thulium laser resection vs. TURP: Quality of life (IPSS question) – 6 and 12 months	582
Figure E-109: Thulium laser resection vs. TURP: Complications	583
4.1.3 HOLEP VS. TRANSURETHRAL INCISION OF THE PROSTATE (TUIP)	584
Figure E-110: HoLEP vs. TUIP: Symptom score.....	584
Figure E-111: HoLEP vs. TUIP: quality of life (IPSS question)	584
Figure E-112: HoLEP vs. TUIP: Qmax(ml/s)	585
Figure E-113: HoLEP vs. TUIP: All cause mortality and complications.....	585
4.1.4 HOLEP VS. OPEN PROSTATECTOMY (OP)	586
Figure E-114: 1 HoLEP vs. OP: Symptom score.....	586
Figure E-115: 1 HoLEP vs. OP: quality of life (IPSS question)	587
Figure E-116: 1 HoLEP vs. OP: Qmax(ml/s) at 3 months (random effects analysis) and longest available follow up (fixed effects analysis).....	587
Figure E-117: 1 HoLEP vs. OP: All cause mortality and complications	588
4.2 LASER TREATMENTS	589
4.2.1 LASER COAGULATION TECHNIQUES VS. TURP.....	589
Figure E-118: 1 Laser Coagulation Techniques vs. TURP: Symptom score at 3 and 6 months (random effects analysis), 12 months and 24 months (change and endpoints).....	589
Figure E-119: Laser Coagulation Techniques vs. TURP: Quality of life (IPSS question), change and endpoints.....	590
Figure E-120: Laser Coagulation Techniques vs. TURP: Qmax (ml/s).....	590
Figure E-121: Laser Coagulation Techniques vs. TURP: All cause mortality and complications	591
Continued Figure E-121 Laser Coagulation Techniques vs. TURP: All cause mortality and complications	592
Figure E-122: Laser Coagulation Techniques vs. TURP: Complications – retrograde ejaculation (random effects analysis).....	592
4.2.2 LASER COAGULATION TECHNIQUES VS. TURP IN AUR PATIENTS	593
Figure E-123: Laser Coagulation Techniques vs. TURP in AUR patients: Symptom score change.....	593
Figure E-124: Laser Coagulation Techniques vs. TURP in AUR patients: Quality of life (IPSS question), change.....	593
Figure E-125: Laser Coagulation Techniques vs. TURP in AUR patients: Complications	593
4.2.3 LASER VAPORISATION TECHNIQUES VS. TURP.....	594
Figure E-126: Laser Vaporisation Techniques vs. TURP: Symptom score at 3 months and 6 months (random effects analysis).....	594
Figure E-127: Laser Vaporisation Techniques vs. TURP: Symptom score at 1, 2, 3 and 5 years (fixed effects analysis)	594
Figure E-128: Laser Vaporisation Techniques vs. TURP: quality of life (IPSS question)	595
Figure E-129: Laser Vaporisation Techniques vs. TURP: Qmax(ml/s) – 3 months(fixed effect analysis) and longest available follow up(random effects analysis)	595
Figure E-130: Laser Vaporisation Techniques vs. TURP: All cause mortality and complications.....	597
Continued Figure E-130: Laser Vaporisation Techniques vs. TURP : Complications	598
Figure E-131: Laser Vaporisation Techniques vs. TURP: Complications – retrograde ejaculation (random effects analysis).....	598
4.2.4 LASER (PHOTOSELECTIVE VAPORISATION) VS. OPEN PROSTATECTOMY(OP).....	599
Figure E-132: Laser (photoselective vaporisation) vs. OP: Complications.....	599
4.2.5 LASER COAGULATION VS. TUMT (TRANSURETHRAL MICROWAVE THERMOTHERAPY)	599
Figure E-133: Laser coagulation vs. TUMT –Symptom score at 6 months	599
Figure E-134: Laser coagulation vs. TUMT – Qmax(ml/s) at 6 months.....	599
Figure E-135: Laser coagulation vs. TUMT: Complications.....	600
4.2.6 LASER VS. TUVV (TRANSURETHRAL VAPORISATION OF THE PROSTATE)	601
Figure E-136: Laser vs. TUVV: Symptom score (random effects analysis)	601
Figure E-137: Laser vs. TUVV – Quality of life (IPSS question)	602
Figure E-138: Laser vs. TUVV – Qmax(ml/s) at 6 month, 12 month(fixed effect analysis) and longest available follow up (random effects analysis).....	602

Figure E-139: Laser vs. TUVp – All cause mortality and complications	604
4.2.7 LASER VS. LASER.....	605
Figure E-140: Laser Vaporization Techniques vs. Laser Coagulation Techniques: Symptom score at 3 months (random effects analysis)	605
Figure E-140b: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Symptom score at 6, 12 and 24 months (fixed effect analysis)	605
Figure E-141: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Qmax (ml/s) at 3 months and longest available follow up.....	606
Figure E-142: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Complications.....	606
Figure E-143: HoLRP vs. Laser coagulation: Complications.....	607
Figure E-144: HoLAP vs. Laser vaporisation: Symptom score.....	607
Figure E-145: HoLAP vs. Laser vaporisation: quality of life (IPSS question)	608
Figure E-146: HoLAP vs. laser vaporisation: Qmax(ml/s) at 3 and longest available follow up(12 months)	608
Figure E-147: HoLAP vs. laser vaporisation: All cause mortality and complications.....	609
4.3 TRANSURETHRAL MICROWAVE THERMOTHERAPY (TUMT).....	609
4.3.1 TUMT vs. SHAM PROCEDURE	609
Figure E-148: TUMT vs. SHAM: Symptom score at 3 and 6 months.....	609
Figure E-149: TUMT vs. SHAM: Qmax(ml/s) and 3 months and at long term follow up	610
Figure E-150: TUMT vs. SHAM: All cause mortality and complications.....	611
Figure E-151: TUMT vs. SHAM: Complications – reoperatoions (random effects analysis)	612
4.3.2 TUMT vs. TURP	612
Figure E-152: TUMT vs. TURP: Symptom score at 3, 12 and 36 months (random effects analysis)	612
Figure E-153: TUMT vs. TURP: Symptom score at 6, 24, 48 and 60 months postoperatively	613
Figure E-154: TUMT vs. TURP: Qmax(ml/s) at 3 months and longest available follow up (random effects analysis)	614
Figure E-155: TUMT vs. TURP: Quality of life (IPSS question) at 3 and 6 months postoperatively	614
Figure E-156: TUMT vs. TURP: quality of life (IPSS question) at 12 months postoperatively (random effects analysis)	615
Figure E-157: TUMT vs. TURP: quality of life (IPSS question) at 48 and 60 months postoperatively	615
Figure E-158: TUMT vs. TURP: All cause mortality and complications.....	616
Continued Figure E-158: TUMT vs. TURP: Complications.....	617
Figure E-159: TUMT vs. TURP: Complications - Incontinence and retrograde ejaculation (random effects analysis)	617
4.3.3 TUMT vs. LASER	618
4.4 TUVp.....	618
4.4.1 TUVp vs. TURP.....	618
Figure E-160: TUVp vs. TURP: Symptom score at 3, 6 and 12 months and 5 years or more postoperatively (fixed effects model)	618
Figure E-161: TUVp vs. TURP: Symptom score at 2 and 3 years postoperatively (random effects analysis) ..	619
Figure E-162: TUVp vs. TURP: Quality of life (IPSS question)	619
Figure E-163: TUVp vs. TURP: Quality of life (IPSS question) – 1 year and 2 year postoperatively (random effects analysis)	620
Figure E-164: TUVp vs. TURP: Qmax(ml/s) at 3 months (fixed effect analysis) and longest available follow up(random effects analysis)	620
Figure E-165: TUVp vs. TURP: All cause mortality and complications.....	621
Continued Figure E-165: TUVp vs. TURP: All cause mortality and complications	622
Figure E-166: TUVp vs. TURP: Complications – retrograde ejaculation (random effects analysis)	623
4.4.2 BIPOLAR TUVp vs. TURP	623
Figure E-167: Bipolar TUVp vs. TURP: Symptom score.....	623
Figure E-168: Bipolar TUVp vs. TURP: Qmax(ml/s) at 3 months and longest available follow up.....	624
Figure E-169: Bipolar TUVp vs. TURP: All cause mortality and complications.....	624
4.4.3 TUVp vs. LASER.....	625
4.5 TRANSURETHRAL NEEDLE ABLATION OF THE PROSTATE (TUNA)	626
4.5.1 TUNA vs. TURP	626
Figure E-170: TUNA vs. TURP: Symptom score	626
Figure E-171: TUNA vs. TURP: Quality of life (IPSS question).....	627
Figure E-172: TUNA vs. TURP: Qmax(ml/s)	627
Figure E-173: TUNA vs. TURP: All cause mortality and complications	628
4.6 TRANSURETHRAL INCISION OF THE PROSTATE (TUIP)	629

4.6.1	TUIP vs. TURP	629
	Figure E-174: TUIP vs. TURP: Symptom score.....	629
	Figure E-175: TUIP vs. TURP: Quality of life (IPSS question)	629
	Figure E-176: TUIP vs. TURP: Qmax (ml/s).....	630
	Figure E-177: TUIP vs. TURP: All cause mortality and complications.....	631
	Figure E-178: TUIP vs. TURP: Complications – retrograde ejaculation (random effects analysis)	632
4.6.2	TUIP vs. TURP IN AUR PATIENTS.....	632
	Figure E-179: TUIP vs. TURP in AUR patients: All cause mortality and complications	632
4.6.3	TUIP vs. HOLEP	632
4.7	BOTULINUM TOXIN IN THE PROSTATE	633
4.7.1	BOTULINUM TOXIN VS. PLACEBO	633
	Figure E-180: Botulinum toxin vs. placebo: Symptom score at 1- and 2-month follow up.....	633
	Figure E-181: Botulinum toxin vs. placebo: Qmax (ml/s) at-2 month follow up	633
	Figure E-182: Botulinum toxin vs. placebo: Complications (urinary incontinence) – 2 month follow up	633
4.8	TRANSURETHRAL VAPOURESECTION OF THE PROSTATE (TUVRP)	634
4.8.1	TUVRP vs. TURP	634
	Figure E-183: TUVRP vs. TURP: Symptom score at 3 months, 1 year and 2 years follow up	634
	Figure E-184: TUVRP vs. TURP: Symptom score at 6 months follow up (random effects analysis)	634
	Figure E-185: TUVRP vs. TURP: Quality of life (IPSS question)	635
	Figure E-186: TUVRP vs. TURP: Qmax (ml/s)	635
	Figure E-187: TUVRP vs. TURP: All cause mortality and complications	636
	Figure E-188: TUVRP vs. TURP: Complications – retrograde ejaculation (random analysis)	637
4.8.2	BIPOLAR TUVRP vs. TURP	637
	Figure E-189: Bipolar TUVRP vs. TURP: Symptom score at 3-month follow up.....	637
	Figure E-190: Bipolar TUVRP vs. TURP: Quality of life (IPSS question) at 3-month follow up	637
	Figure E-191: Bipolar TUVRP vs. TURP: Qmax(ml/s) at 3-month follow up	637
	Figure E-192: Bipolar TUVRP vs. TURP: Complications.....	638
4.9	TRANSURETHRAL ETHANOL ABLATION OF THE PROSTATE (TEAP).....	638
4.9.1	TEAP vs. TURP	638
	Figure E-193: TEAP vs. TURP: Complications	638
4.10	OPEN PROSTATECTOMY (OP)	639
4.10.1	OPEN PROSTATECTOMY vs. HOLEP	639
4.10.2	OPEN PROSTATECTOMY vs. LASER VAPORISATION.....	639
4.11	TRANSURETHRAL RESECTION OF THE PROSTATE TURP	639
4.11.1	TURP vs. WATCHFUL WAITING	639
	Figure E-194: TURP vs. Watchful waiting: Qmax (ml/s)	639
	Figure E-195: TURP vs. Watchful waiting: All cause mortality and complications	640
4.11.2	BIPOLAR TURP vs. TURP	641
	Figure E-196: Bipolar TURP vs. TURP: Symptom score	641
	Figure E-197: Bipolar TURP vs. TURP: Quality of life (IPSS question)	642
	Figure E-198: Bipolar TURP vs. TURP: Qmax (ml/s) at 3 months or longest available follow up	643
	Figure E-199: Bipolar TURP vs. TURP: All cause mortality and complications	644
	Continued Figure E-199b: Bipolar TURP vs. TURP: All cause mortality and complications	645
4.11.3	TURP vs. TUVP	645
4.11.4	TURP vs. TUNA	645
4.11.5	TURP vs. LASER.....	646
4.11.6	TURP vs. TUMT.....	646
4.11.7	TURP vs. TUIP.....	646
4.11.8	TURP vs. HOLEP	646
4.11.9	TURP vs. TUVP	646
4.11.10	TURP vs. BIPOLAR TUVP.....	646
4.11.11	TURP vs. TUVRP.....	646
4.11.12	TURP vs. BIPOLAR TUVRP	646
4.11.13	TURP vs. TEAP.....	646
7.1.1	BLADDER TRAINING vs. TURP	648

Figure E-200: Bladder training vs. TURP: Symptom score change at 6 months follow up	648
Figure E-201: Bladder training vs. TURP: Symptom score change at 6 months follow up	648
Figure E-202: Bladder training vs. TURP: Qmax (ml/s) change at 6 months follow up	648
7.1.2 SELF-CATHETERISATION VS. TURP	648
Figure E-203: Self catheterisation vs. TURP in men with chronic urinary retention: Symptom score change at 6 months follow up.....	648
Figure E-204: Self catheterisation vs. TURP in men with chronic urinary retention: quality of life (IPSS question) change at 6 months follow up	648
8.1.1 ACUTE URINARY RETENTION	649
Figure E-205: Alpha-blockers vs. placebo in men with acute urinary retention: Able to void	649
Figure E-206: Alpha-blockers vs. placebo in men with acute urinary retention: Re-catheterisation.....	649
8.2 CHRONIC RETENTION	649
9.1 PHYTOTHERAPY VS. PLACEBO	650
9.1.1 BETA-SITOSTEROL	650
Figure E-207: Beta-sitosterol vs. placebo: Symptom score.....	650
Figure E-208: Beta-sitosterol vs. placebo: Qmax (ml/s)	650
9.1.2 SERENOA REPENS.....	650
Figure E-209: Serenoa repens vs. placebo: Symptom score	650
Figure E-210: Serenoa repens vs. placebo: Qmax (ml/s)	651
Figure E-211: Serenoa repens vs. placebo: Quality of life (IPSS question)	651
9.1.3 URTICA DIOCIA	651
Figure E-212: Urtica dioica vs. placebo: Symptom score.....	651
Figure E-213: Urtica dioica vs. placebo: Qmax (ml/s).....	651
9.1.4 PYGEUM	652
Figure E-214: Urtica dioica vs. placebo: Qmax(ml/s)	652
9.1.5 CERNILTON	652
Figure E-215: Cernilton vs. placebo: Qmax (ml/s).....	652
9.1.6 PHYTOTHERAPY COMBINATIONS.....	652
Figure E-216: Combination of serenoa repens and urtica dioica vs. placebo: Symptom score	652
Figure E-217: Combination of serenoa repens and urtica dioica vs. placebo: Qmax (ml/s)	652
Figure E-218: Combination of pygeum and urtica dioica vs. placebo: Symptom score	653
Figure E-219: Combination of pygeum and urtica dioica vs. placebo: Qmax (ml/s)	653
Figure E-220: Combination of pygeum and urtica dioica vs. placebo: Quality of life (IPSS question)	653
Figure E-221: Combination of cernitin, serona repens, phytosterol and Vitamin E vs. placebo: Symptom score.....	653
Figure E-222: Combination of cernitin, serona repens, phytosterol and Vitamin E vs. placebo: Qmax (ml/s)	653
9.2 PHYTOTHERAPY VS. ALPHA-BLOCKERS.....	654
9.2.1 SERENOA REPENS VS. ALPHA-BLOCKERS.....	654
Figure E-223: Phytotherapy vs. Alpha-blockers: Symptom score	654
Figure E-224: Phytotherapy vs. Alpha-blockers: Quality of life (IPSS question)	654
Figure E-225: Phytotherapy vs. Alpha-blockers: Qmax (ml/s)	654
Figure E-226: Phytotherapy vs. Alpha-blockers: Urinary retention	655
9.3 PHYTOTHERAPY VS. 5-ARI	655
9.3.1 SERENOA REPENS VS. 5-ARI	655
Figure E-227: Serenoa repens vs. 5-alpha-reductase inhibitors: Symptom score	655
Figure E-228: Serenoa repens vs. 5-alpha-reductase inhibitors: quality of life (IPSS question)	655
Figure E-229: Serenoa repens vs. 5-alpha-reductase inhibitors: Qmax (ml/s) at longest available follow up.....	655
Figure E-230: Serenoa repens vs. 5-alpha-reductase inhibitors: Urinary retention	656
9.3.2 SERENOA REPENS AND URTICA DIOCIA VS. 5-ARI	656
Figure E-231: Serenoa repens and urtica dioica vs. 5-alpha-reductase inhibitors: Symptom score	656
Figure E-232: Serenoa repens and urtica dioica vs. 5-alpha-reductase inhibitors: Qmax (ml/s) at 3 months and 12 months.....	656
10.1 EDUCATIONAL INTERVENTION VS. NO INTERVENTION	657
Figure E-233: Interactive video vs. no intervention: Decisional conflict score	657
10.2 SELF MANAGEMENT VS. STANDARD CARE	657
Figure E-234: Self management vs. standard care: symptom score.....	657
Figure E-235: Self management vs. standard care: Treatment failure.....	657


1. Diagnostic Tests

1.1 Free Uroflowmetry (Peak Urinary Flow)

Figure E-1: Sensitivity and specificity of free uroflowmetry (Qmax) in the diagnosis of bladder outlet obstruction


Free Uroflowmetry Qmax < 10 mL/s

Study	TP	FP	FN	TN	Sensitivity	Specificity
Oelke 2007	51	23	24	62	0.68 [0.56, 0.78]	0.73 [0.62, 0.82]
Poulsen 1994	68	23	31	31	0.69 [0.59, 0.78]	0.57 [0.43, 0.71]
Reynard 1996	47	8	47	54	0.50 [0.40, 0.60]	0.87 [0.76, 0.94]
Reynard 1998	252	107	288	250	0.47 [0.42, 0.51]	0.70 [0.65, 0.75]



Free Uroflowmetry Qmax < 12 mL/s

Study	TP	FP	FN	TN	Sensitivity	Specificity
Reynard 1996	62	16	33	46	0.65 [0.55, 0.75]	0.74 [0.62, 0.84]



Free Uroflowmetry Qmax < 15 mL/s

Study	TP	FP	FN	TN	Sensitivity	Specificity
Oelke 2007	74	52	1	33	0.99 [0.93, 1.00]	0.39 [0.28, 0.50]
Poulsen 1994	89	37	10	17	0.90 [0.82, 0.95]	0.31 [0.20, 0.46]
Reynard 1996	81	29	14	33	0.85 [0.77, 0.92]	0.53 [0.40, 0.66]
Reynard 1998	440	221	100	136	0.81 [0.78, 0.85]	0.38 [0.33, 0.43]





Figure E-2: Summary receiver operating characteristic (SROC) curve for uroflowmetry Qmax in the diagnosis of bladder outlet obstructions



2. Conservative Interventions

2.1 Pelvic Floor Muscle Training (PFMT)

2.1.1 PFMT vs. Control

Figure E-3: PFMT vs. Control: Number of post-prostatectomy men who were incontinent




Figure E-4: PFMT vs. Control: Mean urine lost (g) per 24 hours (pad test) in post-prostatectomy men






Figure E-5: PFMT vs. Control: Number of post-TURP men who were incontinent



2.2 Biofeedback

2.2.1 Biofeedback + PFMT vs. Control


Figure E-6: PFMT + Biofeedback vs. no intervention: Number of men who were incontinent at follow up



2.3 Electrical Stimulation (ES)

2.3.1 ES + PFMT vs. Control

Figure E-7: ES + PFMT vs. no intervention: Number of men who were incontinent at follow up



3. Pharmacological Interventions

3.1 Alpha-blockers

3.1.1 Alpha-blockers vs. placebo

Figure E-8: Alpha-blockers vs. Placebo: Symptom score (random effects analysis)




Figure E-9: Alpha-blockers vs. Placebo: Qmax (ml/s) (random effects analysis)




Figure E-10: Alpha-blockers vs. Placebo: Quality of life – IPSS question (random effects analysis)




Figure E-11: Alpha-blockers vs. Placebo: Adverse events (cardiovascular and neurological) - asthenia (fatigue) and headache




Figure E-12: Alpha-blockers vs. Placebo: Adverse events (cardiovascular and neurological) - postural hypotension and rhinitis




Figure E-13: Alpha-blockers vs. Placebo: Adverse events - erectile dysfunction /impotence




Figure E-14: Alpha-blockers vs. Placebo: Adverse events - dizziness and retrograde ejaculation (random effects analysis)





Figure E-15: Alpha-blockers vs. Placebo: Withdrawal from study due to adverse events



3.1.2 Alpha-blockers vs. 5-Alpha reductase inhibitors (5-ARI)

Figure E-16: Alpha-blockers vs. 5-ARI: Symptom score




Figure E-17: Alpha-blockers vs. 5-ARI: Quality of life (IPSS-question)




Figure E-18: Alpha-blockers vs. 5-ARI: Qmax (ml/s)




Figure E-19: Alpha-blockers vs. 5-ARI: Prostate volume (ml)




Figure E-20 Alpha-blockers vs. 5-ARI: PSA (ng/ml)






Figure E-21: Alpha-blockers vs. 5-ARI: Adverse events (cardiovascular or neurological)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998 and Rigatti2003, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Figure E-22: Alpha-blockers vs. 5-ARI: Adverse events (sexual or urological)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998 and Rigatti2003, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Figure E-23: Alpha-blockers vs. 5-ARI: Adverse events - postural hypotension and ejaculatory abnormality (random effects analysis)




Figure E-24: Alpha-blockers vs. 5-ARI: Ejaculatory abnormality – subgroup analysis of tamsulosin and other alpha-blockers





Figure E-25: Alpha-blockers vs. 5-ARI: Withdrawal from study due to adverse events (random effects analysis)

The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998 and Rigatti2003, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

3.1.3 Alpha-blockers vs. Anticholinergics

See section 3.3.2 Anticholinergics vs. Alpha-blockers

3.1.4 Alpha-blockers vs. Phosphodiesterase 5-inhibitors (PDE5-I)

See section 3.4.2 PDE5-I vs. Alpha-blockers

3.2 5-alpha reductase inhibitors (5-ARI)

3.2.1 5-ARI vs. placebo

Figure E-26: 5-ARI vs. Placebo: Symptom score at 3 months, 6 months 2 years and 4 years or longer (random effects analysis)




Figure E-27: 5-ARI vs. Placebo: Symptom score at 2 years- subgroup analysis




Figure E-28: 5-ARI vs. Placebo: Symptom score at 12 months and 3 years




Figure E-29: 5-ARI vs. Placebo: Qmax (ml/s) at 3 months, 6 months, 2 years, 3 years and 4 years or longer




Figure E-30: 5-ARI vs. Placebo: Qmax (ml/s) at 12 months (random effects analysis)




Figure E-31: 5-ARI vs. Placebo: Prostate volume(ml) at 1 year follow up




Figure E-32: 5-ARI vs. Placebo: Prostate volume (ml) at 2 years follow up (random effects analysis)




Figure E-33: 5-ARI vs. Placebo: PSA (ng/ml) level at 2 year follow up




Figure E-34: 5-ARI vs. Placebo: Adverse events (cardiovascular and neurological)




Figure E-35: 5-ARI vs. Placebo: Adverse events (sexual and urological)





Figure E-36: 5-ARI vs. Placebo: Withdrawal from study due to adverse events



3.2.2 5-Alpha reductase inhibitors (5-ARI) vs. Alpha-blockers

See section 3.1.2: Alpha-blockers vs. 5-Alpha reductase inhibitors (5-ARI)

3.3 Anticholinergics

3.3.1 Anticholinergics vs. placebo

Figure E-37: Anticholinergics vs. Placebo: Adverse events






Figure E-38: Anticholinergics vs. Placebo: Withdrawal from study due to adverse events



3.3.2 Anticholinergics vs. Alpha-blockers

Figure E-39: Anticholinergics vs. Alpha-blockers: Adverse events



3.4 Phosphodiesterase-5-inhibitors (PDE5-I)

3.4.1 PDE5-I vs. placebo

Figure E-40: PDE5-I vs. Placebo: Symptom score




Figure E-41: PDE5-I vs. Placebo: Quality of life (IPSS question)




Figure E-42: PDE5-I vs. Placebo: Qmax(ml/s)





Figure E-43: PDE5-I vs. Placebo: Adverse events



3.4.2 PDE5-I vs. Alpha-blockers

Figure E-44: PDE5-I vs. Alpha-blockers: Symptom score




Figure E-45: PDE5-I vs. Alpha-blockers: Qmax (ml/s)




Figure E-46: PDE5-I vs. Alpha-blockers: Voiding frequency




Figure E-47: PDE5-I vs. Alpha-blockers: Nocturia





Figure E-48: PDE5-I vs. Alpha-blockers: Adverse events



3.5 Diuretics

3.5.1 Diuretics vs. placebo


Forest plots were not prepared for this comparison. Please see Evidence Table 16 in Appendix D for details.

3.6 Desmopressin

3.6.1 Desmopressin vs. placebo

Forest plots were not prepared for the efficacy outcomes of this cross over trial. Please see Evidence Table 17 in Appendix D for details.

Figure E-49: Desmopressin vs. Placebo: Adverse events



This is a cross over trial and a paired test would be more appropriate. Forest plots prepared for illustration purpose.

3.7 NSAIDS

3.7.1 NSAIDS vs. placebo

Figure E-50: NSAIDs vs. Placebo: Symptom score at 1 month




Figure E-51: NSAIDs vs. Placebo: Qmax (ml/s) at 1 month




Figure E-52: NSAIDs vs. Placebo: Nocturia frequency at 1 month





Figure E-53: NSAIDs vs. Placebo: Adverse events (1 month follow up)



one type of adverse event was reported.

Only

3.8 Combination therapy: Alpha-blockers plus 5-alpha reductase inhibitors(5-ARI)

3.8.1 Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers

Figure E-54: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Symptom score




Figure E-55: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Qmax (ml/s)




Figure E-56: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Prostate volume(ml)




Figure E-57: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: PSA (ng/ml)






Figure E-58: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (cardiovascular or neurological)




The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Continued Figure E-58: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (cardiovascular or neurological)




The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Figure E-59: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Adverse events (sexual or urological)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)


Figure E-60: Combination (Alpha-blockers + 5-ARI) vs. Alpha-blockers: Withdrawal from study due to adverse events



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

3.8.2 Combination (Alpha-blockers + 5-ARI) vs. 5-ARI

Figure E-61: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Symptom score



Test for subgroup differences: Chi² = 15.94, df = 3 (P = 0.001), I² = 81.2%

Figure E-62: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Qmax(ml/s)




Figure E-63: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Prostate volume (ml)




Figure E-64: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: PSA (ng/ml)






Figure E-65: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (Cardiovascular or neurological)




The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Continued Figure E-65: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (cardiovascular or neurological)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Figure E-66: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Adverse events (sexual or urological)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003, 2 years for Roehrborn2008 and 4 years for McConnell2003)

Figure E-67: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Ejaculatory abnormality (random effects analysis)




Figure E-68: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Ejaculatory abnormality subgroup analysis





Figure E-69: Combination (Alpha-blockers + 5-ARI) vs. 5-ARI: Withdrawal from study due to adverse events (random effects analysis)



The studies were arranged in the forest plots based on duration of follow up (6 months for Debruyne1998, 1 year for Lepor1996 and Kirby2003 and 2 years for Roehrborn2008)

3.8.3 Combination (Alpha-blockers + 5-ARI) vs. placebo

Figure E-70: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Symptom score




Figure E-71: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Qmax (ml/s)




Figure E-72: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Prostate volume (ml)




Figure E-73: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Change in PSA (ng/ml)





Figure E-74: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events (cardiovascular and neurological)



The studies were arranged in the forest plots based on duration of follow up (1 year for Lepor1996 and Kirby2003 and 4 years for McConnell2003)

Figure E-75: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events – postural hypotension (random effects analysis)




Figure E-76: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Adverse events (sexual or urological)





Figure E-77: Combination (Alpha-blockers + 5-ARI) vs. Placebo: Withdrawal from study due to adverse events (random effects analysis)




The studies were arranged in the forest plots based on duration of follow up (1 year for Lepor1996 and Kirby2003 and 4 years for McConnell2003)

3.9 Combination Therapy : Anti-cholinergic plus Alpha-blockers

3.9.1 Combination (Anti-cholinergic + Alpha-blockers) vs. Alpha-blockers

Figure E-78: Combination (Anti-cholinergic + Alpha-blockers) vs. Alpha-blockers: Adverse events



3.9.2 Anti-cholinergic added on to Alpha-blockers vs. Alpha-blockers

Figure E-79: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Symptom score at 3 months




Figure E-80: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Quality of life (IPSS question) at 3 months




Figure E-81: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Qmax (ml/s) at 3 months






Figure E-82: Anti-Ch added on to Alpha-blockers vs. Alpha-blockers: Adverse events (3-months follow up)




3.9.3 Combination (Anti-cholinergic + Alpha-blockers) vs. Anticholinergics

Figure E-83: Combination (Anti-cholinergic + Alpha-blockers) vs. Anticholinergics: Adverse events



3.9.4 Combination (Anti-cholinergic + Alpha-blockers) vs. Placebo

Figure E-84: Combination (Anti-cholinergic + Alpha-blockers) vs. Placebo: adverse events



3.10 Combination (PDE5-I + Alpha-blockers)

3.10.1 Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers

Figure E-85: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Symptom score




Figure E-86: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Quality of life (IPSS question)




Figure E-87: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Qmax(ml/s)




Figure E-88: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Frequency at 3-month




Figure E-89: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Nocturia at 3 months





Figure E-90: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events



Continued Figure E-90: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events





Figure E-91: Combination (PDE5-I + Alpha-blockers) vs. Alpha-blockers: Adverse events resulting in withdrawal at 3-month



3.10.2 Combination (PDE5-I + Alpha-blockers) vs. PDE5-I

Figure E-92: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: symptom score (random effects analysis)




Figure E-93: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Quality of life (IPSS-QoL) up to 3-month




Figure E-94: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Qmax (ml/s) at 3-month




Figure E-95: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Frequency at 3-month




Figure E-96: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Nocturia at 3-month




Figure E-97: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Adverse events (only those resulting in withdrawals reported)





Figure E-98: Combination (PDE5-I + Alpha-blockers) vs. PDE5-I: Withdrawal from study due to adverse events



4. Surgery

4.1 Holmium Laser Enucleation of the Prostate (HoLEP)

4.1.1 HoLEP vs. Transurethral resection of the prostate (TURP)

Figure E-99: HoLEP vs. TURP: Symptom score at 3 months, 36 months and 48 months




Figure E-100: HoLEP vs. TURP: Symptom score at 6, 12 and 24 months (random effects analysis)





Figure E-101: HoLEP vs. TURP: Quality of life (IPSS question) – 3, 24 and 48 months



Test for subgroup differences: Chi² = 0.72, df = 2 (P = 0.70), I² = 0%

Figure E-102: HoLEP vs. TURP: Quality of life (IPSS question) – 6 to 12 months (random effects analysis)







Figure E-103: HoLEP vs. TURP: Qmax(ml/s) at 3 months and longest available follow up



Test for subgroup differences: Chi² = 1.11, df = 1 (P = 0.29), I² = 9.5%

Figure E-104: HoLEP vs. TURP: All cause mortality and complications

Continued Figure E-104: HoLEP vs. TURP: All cause mortality and complications



4.1.2 Thulium laser resection vs. TURP

Figure E-105: Thulium laser resection vs. TURP: Symptom score – 6 months postoperatively




Figure E-106: Thulium laser resection vs. TURP: Symptom score – 12 months postoperatively




Figure E-107: Thulium laser resection vs. TURP: Qmax(ml/s) – 12 months postoperatively




Figure E-108: Thulium laser resection vs. TURP: Quality of life (IPSS question) – 6 and 12 months






Figure E-109: Thulium laser resection vs. TURP: Complications



4.1.3 HoLEP vs. Transurethral Incision of the Prostate (TUIP)


Figure E-110: HoLEP vs. TUIP: Symptom score



Test for subgroup differences: Chi² = 1.16, df = 2 (P = 0.56), I² = 0%

* Only one study using holmium laser for bladder neck incision (HoBNI) was found.

Figure E-111: HoLEP vs. TUIP: quality of life (IPSS question)



Test for subgroup differences: Chi² = 0.28, df = 2 (P = 0.87), I² = 0%

Figure E-112: HoLEP vs. TUIP: Qmax(ml/s)





Figure E-113: HoLEP vs. TUIP: All cause mortality and complications



4.1.4 HOLEP vs. Open prostatectomy (OP)

Figure E-114: 1 HoLEP vs. OP: Symptom score




Figure E-115: 1 HoLEP vs. OP: quality of life (IPSS question




Figure E-116: 1 HoLEP vs. OP: Qmax(ml/s) at 3 months (random effects analysis) and longest available follow up (fixed effects analysis)





Figure E-117: 1 HoLEP vs. OP: All cause mortality and complications

4.2 Laser treatments

4.2.1 Laser Coagulation Techniques vs. TURP

Figure E-118: 1 Laser Coagulation Techniques vs. TURP: Symptom score at 3 and 6 months (random effects analysis), 12 months and 24 months (change and endpoints)




Figure E-119: Laser Coagulation Techniques vs. TURP: Quality of life (IPSS question), change and endpoints.




Figure E-120: Laser Coagulation Techniques vs. TURP: Qmax (ml/s)





Figure 121: Laser Coagulation Techniques vs. TURP: All cause mortality and complications



Continued Figure 121 Laser Coagulation Techniques vs. TURP: All cause mortality and complications





Figure E-122: Laser Coagulation Techniques vs. TURP: Complications – retrograde ejaculation (random effects analysis)



4.2.2 Laser Coagulation Techniques vs. TURP in AUR patients

Figure E-123: Laser Coagulation Techniques vs. TURP in AUR patients: Symptom score change




Figure E-124: Laser Coagulation Techniques vs. TURP in AUR patients: Quality of life (IPSS question), change





Figure E-125: Laser Coagulation Techniques vs. TURP in AUR patients: Complications



4.2.3 Laser Vaporisation Techniques vs. TURP

Figure E-126: Laser Vaporisation Techniques vs. TURP: Symptom score at 3 months and 6 months (random effects analysis)




Figure E-127: Laser Vaporisation Techniques vs. TURP: Symptom score at 1, 2, 3 and 5 years (fixed effects analysis)




Figure E-128: Laser Vaporisation Techniques vs. TURP: quality of life (IPSS question)





Figure E-129: Laser Vaporisation Techniques vs. TURP: Qmax(ml/s) – 3 months(fixed effect analysis) and longest available follow up(random effects analysis)







Figure E-130: Laser Vaporisation Techniques vs. TURP: All cause mortality and complications



Continued Figure E-130: Laser Vaporisation Techniques vs. TURP : Complications






Figure E-131: Laser Vaporisation Techniques vs. TURP: Complications – retrograde ejaculation (random effects analysis)



4.2.4 Laser (photoselective vaporisation) vs. Open prostatectomy(OP)

Figure E-132: Laser (photoselective vaporisation) vs. OP: Complications



4.2.5 Laser coagulation vs. TUMT (Transurethral Microwave Thermotherapy)

Figure E-133: Laser coagulation vs. TUMT –Symptom score at 6 months




Figure E-134: Laser coagulation vs. TUMT – Qmax(ml/s) at 6 months





Figure E-135: Laser coagulation vs. TUMT: Complications



4.2.6 Laser vs. TUVP (Transurethral Vaporisation of the Prostate)

Figure E-136: Laser vs. TUVP: Symptom score (random effects analysis)




Figure E-137: Laser vs. TUVP – Quality of life (IPSS question)






Figure E-138: Laser vs. TUVP – Qmax(ml/s) at 6 month, 12 month(fixed effect analysis) and longest available follow up (random effects analysis)




Test for subgroup differences: Chi² = 5.55, df = 1 (P = 0.02), I² = 82.0%



No 3 month data was available for this comparison.

Figure E-139: Laser vs. TUVP – All cause mortality and complications



0.001 0.1 1 10 1000
Laser TUVP

4.2.7 Laser vs. laser

4.2.7.1 Laser Vaporisation Techniques vs. Laser Coagulation Techniques

Figure E-140: Laser Vaporization Techniques vs. Laser Coagulation Techniques: Symptom score at 3 months (random effects analysis)




Figure E-140b: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Symptom score at 6, 12 and 24 months (fixed effect analysis)




Figure E-141: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Qmax (ml/s) at 3 months and longest available follow up






Figure E-142: Laser Vaporisation Techniques vs. Laser Coagulation Techniques: Complications




4.2.7.2 Holmium laser resection of the prostate(HoLRP) vs. Laser coagulation

Figure E-143: HoLRP vs. Laser coagulation: Complications



4.2.7.3 Holmium Laser Ablation of the Prostate(HoLAP) vs. Laser Vaporisation

Figure E-144: HoLAP vs. Laser vaporisation: Symptom score



Test for subgroup differences: Chi² = 7.37, df = 2 (P = 0.03), I² = 72.9%
 Only one study was using photoselective laser vaporisation (PVP) method was found

Figure E-145: HoLAP vs. Laser vaporisation: quality of life (IPSS question)




Figure E-146: HoLAP vs. laser vaporisation: Qmax(ml/s) at 3 and longest available follow up(12 months)






Figure E-147: HoLAP vs. laser vaporisation: All cause mortality and complications



4.3 Transurethral Microwave Thermotherapy (TUMT)

4.3.1 TUMT vs. Sham procedure

Figure E-148: TUMT vs. SHAM: Symptom score at 3 and 6 months



Test for subgroup differences: Chi² = 1.53, df = 1 (P = 0.22), I² = 34.4%

Figure E-149: TUMT vs. SHAM: Qmax(ml/s) and 3 months and at long term follow up




Figure E-150: TUMT vs. SHAM: All cause mortality and complications





Figure E-151: TUMT vs. SHAM: Complications – reoperatoions (random effects analysis)



4.3.2 TUMT vs. TURP

Figure E-152: TUMT vs. TURP: Symptom score at 3, 12 and 36 months (random effects analysis)






Figure E-153: TUMT vs. TURP: Symptom score at 6, 24, 48 and 60 months postoperatively






Figure E-154: TUMT vs. TURP: Qmax(ml/s) at 3 months and longest available follow up (random effects analysis)





Figure E-155: TUMT vs. TURP: Quality of life (IPSS question) at 3 and 6 months postoperatively



Test for subgroup differences: Chi² = 0.06, df = 1 (P = 0.80), I² = 0%

Figure E-156: TUMT vs. TURP: quality of life (IPSS question) at 12 months postoperatively (random effects analysis)








Figure E-157: TUMT vs. TURP: quality of life (IPSS question) at 48 and 60 months postoperatively



Test for subgroup differences: Chi² = 0.31, df = 1 (P = 0.58), I² = 0%

Figure E-158: TUMT vs. TURP: All cause mortality and complications



Continued Figure E-158: TUMT vs. TURP: Complications





Figure E-159: TUMT vs. TURP: Complications - Incontinence and retrograde ejaculation (random effects analysis)



4.3.3 TUMT vs. Laser

See section 4.2.5 Laser coagulation vs. TUMT (Transurethral Microwave Thermotherapy)

4.4 TUVP

4.4.1 TUVP vs. TURP

Figure E-160: TUVP vs. TURP: Symptom score at 3, 6 and 12 months and 5 years or more postoperatively (fixed effects model)




Figure E-161: TUVP vs. TURP: Symptom score at 2 and 3 years postoperatively (random effects analysis)





Figure E-162: TUVP vs. TURP: Quality of life (IPSS question)



Test for subgroup differences: Chi² = 13.06, df = 3 (P = 0.005), I² = 77.0%

Figure E-163: TUVP vs. TURP: Quality of life (IPSS question) – 1 year and 2 year postoperatively (random effects analysis)




Figure E-164: TUVP vs. TURP: Qmax(ml/s) at 3 months (fixed effect analysis) and longest available follow up (random effects analysis)





Figure E-165: TUVP vs. TURP: All cause mortality and complications



Continued Figure E-165: TUVP vs. TURP: All cause mortality and complications





Figure E-166: TUVP vs. TURP: Complications – retrograde ejaculation (random effects analysis)



4.4.2 Bipolar TUVP vs. TURP

Figure E-167: Bipolar TUVP vs. TURP: Symptom score




Figure E-168: Bipolar TUVP vs. TURP: Qmax(ml/s) at 3 months and longest available follow up






Figure E-169: Bipolar TUVP vs. TURP: All cause mortality and complications





4.4.3 TUVP vs. Laser

See section 4.2.6 Laser vs. TUVP (Transurethral Vaporisation of the Prostate)

4.5 Transurethral Needle Ablation of the Prostate (TUNA)

4.5.1 TUNA vs. TURP

Figure E-170: TUNA vs. TURP: Symptom score




Figure E-171: TUNA vs. TURP: Quality of life (IPSS question)




Figure E-172: TUNA vs. TURP: Qmax(ml/s)





Figure E-173: TUNA vs. TURP: All cause mortality and complications



4.6 Transurethral Incision of the Prostate (TUIP)

4.6.1 TUIP vs. TURP

Figure E-174: TUIP vs. TURP: Symptom score




Figure E-175: TUIP vs. TURP: Quality of life (IPSS question)




Figure E-176: TUIP vs. TURP: Qmax (ml/s)




Figure E-177: TUIP vs. TURP: All cause mortality and complications






Figure E-178: TUIP vs. TURP: Complications – retrograde ejaculation (random effects analysis)



4.6.2 TUIP vs. TURP in AUR patients

Figure E-179: TUIP vs. TURP in AUR patients: All cause mortality and complications



4.6.3 TUIP vs. HOLEP

See 4.1.3HoLEP vs. Transurethral Incision of the Prostate (TUIP)

4.7 Botulinum toxin in the prostate

4.7.1 Botulinum toxin vs. placebo

Figure E-180: Botulinum toxin vs. placebo: Symptom score at 1- and 2-month follow up




Figure E-181: Botulinum toxin vs. placebo: Qmax (ml/s) at-2 month follow up





Figure E-182: Botulinum toxin vs. placebo: Complications (urinary incontinence) – 2 month follow up



4.8 Transurethral Vapouresection of the Prostate (TUVRP)

4.8.1 TUVRP vs. TURP

Figure E-183: TUVRP vs. TURP: Symptom score at 3 months, 1 year and 2 years follow up




Figure E-184: TUVRP vs. TURP: Symptom score at 6 months follow up (random effects analysis)




Figure E-185: TUVRP vs. TURP: Quality of life (IPSS question)




Figure E-186: TUVRP vs. TURP: Qmax (ml/s)




Figure E-187: TUVRP vs. TURP: All cause mortality and complications





Figure E-188: TUVRP vs. TURP: Complications – retrograde ejaculation (random analysis)



4.8.2 Bipolar TUVRP vs. TURP

Figure E-189: Bipolar TUVRP vs. TURP: Symptom score at 3-month follow up




Figure E-190: Bipolar TUVRP vs. TURP: Quality of life (IPSS question) at 3-month follow up




Figure E-191: Bipolar TUVRP vs. TURP: Qmax(ml/s) at 3-month follow up






Figure E-192: Bipolar TUVRP vs. TURP: Complications



4.9 Transurethral Ethanol Ablation of the Prostate (TEAP)

4.9.1 TEAP vs. TURP

Figure E-193: TEAP vs. TURP: Complications



4.10 Open Prostatectomy (OP)

4.10.1 Open prostatectomy vs. HOLEP

See section 4.1.4 on HOLEP vs. Open prostatectomy (OP)

4.10.2 Open prostatectomy vs. laser vaporisation

See section 4.2.4 on Laser (photoselective vaporisation) vs. Open prostatectomy(OP)

4.11 Transurethral Resection of the Prostate TURP

4.11.1 TURP vs. Watchful Waiting

Figure E-194: TURP vs. Watchful waiting: Qmax (ml/s)





Figure E-195: TURP vs. Watchful waiting: All cause mortality and complications



4.11.2 Bipolar TURP vs. TURP

Figure E-196: Bipolar TURP vs. TURP: Symptom score




Figure E-197: Bipolar TURP vs. TURP: Quality of life (IPSS question)




Figure E-198: Bipolar TURP vs. TURP: Qmax (ml/s) at 3 months or longest available follow up






Figure E-199: Bipolar TURP vs. TURP: All cause mortality and complications



Continued Figure E-199b: Bipolar TURP vs. TURP: All cause mortality and complications



4.11.3 TURP vs. TUVP

See section 4.4.1 TUVP vs. TURP

4.11.4 TURP vs. TUNA

See section 4.5.1 TUNA vs. TURP

4.11.5 TURP vs. Laser

See sections 4.2.1 Laser Coagulation Techniques vs. TURP, 4.2.2 Laser Coagulation Techniques vs. TURP in AUR patients, 4.2.3 Laser Vaporisation Techniques vs. TURP

4.11.6 TURP vs. TUMT

See section 4.3.2 TUMT vs. TURP

4.11.7 TURP vs. TUIP

See section 4.6.1 TUIP vs. TURP

4.11.8 TURP vs. HoLEP

See section 4.1.1 HoLEP vs. TURP

4.11.9 TURP vs. TUVP

See section 4.4.1 TUVP vs. TURP

4.11.10 TURP vs. Bipolar TUVP

See section 4.4.2 Bipolar TUVP vs. TURP

4.11.11 TURP vs. TUVRP

See section 4.8.1 TUVRP vs. TURP

4.11.12 TURP vs. Bipolar TUVRP

See section 4.8.2 Bipolar TUVRP vs. TURP

4.11.13 TURP vs. TEAP

See section 4.9.1 TEAP vs. TURP

5. Surgical vs. Medical Interventions

There are no forest plots for this section

6. Medical vs. Conservative Interventions

No results found — no forest plots

7. Surgical vs. Conservative Interventions

7.1.1 Bladder training vs. TURP

Figure E-200: Bladder training vs. TURP: Symptom score change at 6 months follow up




Figure E-201: Bladder training vs. TURP: Symptom score change at 6 months follow up





Figure E-202: Bladder training vs. TURP: Qmax (ml/s) change at 6 months follow up



7.1.2 Self-catheterisation vs. TURP

Figure E-203: Self catheterisation vs. TURP in men with chronic urinary retention: Symptom score change at 6 months follow up





Figure E-204: Self catheterisation vs. TURP in men with chronic urinary retention: quality of life (IPSS question) change at 6 months follow up



8. Urinary retention

8.1.1 Acute urinary retention

Figure E-205: Alpha-blockers vs. placebo in men with acute urinary retention: Able to void





Figure E-206: Alpha-blockers vs. placebo in men with acute urinary retention: Re-catheterisation



8.2 Chronic retention

See forest plots in section surgery vs. conservative and conservative

9. Alternative and complementary therapies

9.1 Phytotherapy vs. placebo

9.1.1 Beta-sitosterol

Figure E-207: Beta-sitosterol vs. placebo: Symptom score





Figure E-208: Beta-sitosterol vs. placebo: Qmax (ml/s)



9.1.2 Serenoa repens

Figure E-209: Serenoa repens vs. placebo: Symptom score




Figure E-210: Serenoa repens vs. placebo: Qmax (ml/s)





Figure E-211: Serenoa repens vs. placebo: Quality of life (IPSS question)



9.1.3 Urtica dioica

Figure E-212: Urtica dioica vs. placebo: Symptom score






Figure E-213: Urtica dioica vs. placebo: Qmax (ml/s)




9.1.4 Pygeum

Figure E-214: *Urtica dioica* vs. placebo: Qmax(ml/s)



9.1.5 Cernilton

Figure E-215: Cernilton vs. placebo: Qmax (ml/s)



9.1.6 Phytotherapy combinations

Figure E-216: Combination of serenoa repens and urtica dioica vs. placebo: Symptom score




Figure E-217: Combination of serenoa repens and urtica dioica vs. placebo: Qmax (ml/s)




Figure E-218: Combination of pygeum and urtica dioica vs. placebo: Symptom score




Figure E-219: Combination of pygeum and urtica dioica vs. placebo: Qmax (ml/s)




Figure E-220: Combination of pygeum and urtica dioica vs. placebo: Quality of life (IPSS question)




Figure E-221: Combination of cernitin, serona repens, phytosterol and Vitamin E vs. placebo: Symptom score





Figure E-222: Combination of cernitin, serona repens, phytosterol and Vitamin E vs. placebo: Qmax (ml/s)



9.2 Phytotherapy vs. Alpha-blockers

9.2.1 *Serenoa repens* vs. Alpha-blockers

Figure E-223: Phytotherapy vs. Alpha-blockers: Symptom score




Figure E-224: Phytotherapy vs. Alpha-blockers: Quality of life (IPSS question)




Figure E-225: Phytotherapy vs. Alpha-blockers: Qmax (ml/s)






Figure E-226: Phytotherapy vs. Alpha-blockers: Urinary retention



9.3 Phytotherapy vs. 5-ARI


9.3.1 Serenoa repens vs. 5-ARI

Figure E-227: Serenoa repens vs. 5-alpha-reductase inhibitors: Symptom score




Test for subgroup differences: Not applicable

Figure E-228: Serenoa repens vs. 5-alpha-reductase inhibitors: quality of life (IPSS question)




Test for subgroup differences: Not applicable

Figure E-229: Serenoa repens vs. 5-alpha-reductase inhibitors: Qmax (ml/s) at longest available follow up



Test for subgroup differences: Not applicable

Figure E-230: Serenoa repens vs. 5-alpha-reductase inhibitors: Urinary retention



9.3.2 Serenoa repens and urtica dioica vs. 5-ARI

Figure E-231: Serenoa repens and urtica dioica vs. 5-alpha-reductase inhibitors: Symptom score






Figure E-232: Serenoa repens and urtica dioica vs. 5-alpha-reductase inhibitors: Qmax (ml/s) at 3 months and 12 months



10. Provision of information

10.1 Educational intervention vs. no intervention

Figure E-233: Interactive video vs. no intervention: Decisional conflict score



10.2 Self management vs. standard care

Figure E-234: Self management vs. standard care: symptom score





Figure E-235: Self management vs. standard care: Treatment failure



Appendix F - Cost-effectiveness analysis

1.1 Introduction

Two original cost-effectiveness analyses were carried out to answer the clinical questions on transurethral resection of the prostate (TURP) vs. laser (Chapter 8), and the clinical question on Alpha-blockers (AB) alone or in combination with 5-Alpha Reductase-Inhibitors (5-ARI) (Chapter 6). Throughout the guideline we refer to these two analyses respectively as 'NCGC Surgery Model' and 'NCGC Combination model'.

1.2 Methods

A review of the literature was conducted followed by economic modelling of the cost-effectiveness of the listed interventions in England and Wales. The literature search and review methods can be found in Chapter 2.

Our aim in constructing the models was to determine the most cost-effective strategy in men considering respectively surgery and medical treatment. Those would be mainly men with moderate to severe lower urinary tract symptoms (LUTS).

We found a number of economic evaluations in the published literature (Chapters 6 and 8), among which a Health Technology Assessment (HTA) model of good quality¹⁷². However the Guideline Development Group (GDG) felt that they needed an original model with slightly different assumptions and data in order to make a recommendation with confidence.

The following general principles were adhered to:

- The GDG was consulted during the construction and interpretation of the model.
- When published data was not available we used expert opinion to populate the model.
- Model assumptions were reported fully and transparently.
- The results were subject to sensitivity analysis and limitations were discussed.
- We followed the methods of the NICE reference case²¹⁵. Therefore costs were calculated from a health services perspective. Health gain was

measured in terms of quality-adjusted life-years (QALYs) gained. Both future costs and QALYs were discounted at 3.5%.

- The model employed a cost-effectiveness threshold of £20,000 per QALY gained.
- The model was peer-reviewed by another health economist at the NCGC.

1.2.1 Software

The cost-effectiveness analyses were conducted using TreeAge Pro 2008.

1.3 NCGC Surgery model

1.3.1 General method

We based the model on two of the main outcomes considered in our systematic review of the clinical evidence (Chapter 2.4): mean IPSS change from baseline and adverse events. We chose IPSS change because it better expresses the change in quality of life as felt by the patient compared to other clinical measures such as Qmax. Consequently, it was easier to find data linking utility values to levels of symptoms.

Since LUTS are a lifelong condition, we built a Markov model with a life time horizon and we changed this in a sensitivity analysis. The cycle length is three months, as this was deemed the minimum clinically meaningful time interval to detect differences in patients undergoing surgery.

All the probabilities, costs and health utilities were converted in order to reflect the three-month values.

The treatments compared in our analysis are TURP and Holmium Laser Enucleation of Prostate (HoLEP). TURP is the current standard practice and HoLEP was one of the alternative treatments that were significantly effective as compared to TURP. Transurethral electrovaporisation of prostate (TUVAP) was another effective treatment as compared to TURP but the available economic evidence was considered sufficient to prove it cost-effective.

Patients in the studies included in our clinical review had a moderate-to-severe level of symptoms. Therefore patients in our model were defined as men with moderate-to-severe LUTS who are suitable for either TURP or HoLEP.

Both arms of the model have the same structure (Figure 236): after the intervention, the patient can either have a significant remission of symptoms (success) or no remission/minor remission (failure).

Short-term complications identified in the clinical review (see Appendix E) were assumed to be resolved within 3 months (the cycle length) and could occur with a probability independent from the success. Incontinence is the only long-term adverse event and in some cases it requires an artificial urinary sphincter (AUS). If the man still has storage LUTS together with incontinence, he will not undergo further de-obstructive surgery, therefore he will remain in this health state throughout the model.

Men who initially had a successful outcome can have deterioration in symptoms and end up with residual LUTS state. Some of them will undergo further de-obstructive surgery if incontinence is not present, and some will be medically treated. The second surgery is always TURP, even in the HoLEP arm, as the experts in the GDG believe that HoLEP is unlikely to be performed twice. We varied the structure between the two arms in a structural sensitivity analysis where we assumed TURP was not possible after HoLEP either.

The list of the health states that are part of the model is reported in Table 1.

Table 1 - Health states

HEALTH STATES
(Moderate-to-Severe) LUTS
Remission
LUTS + Incontinence
LUTS + Incontinence AUS
Incontinence
Incontinence AUS

The experts of the GDG members have defined a significant remission of symptoms after surgery as a change in IPSS greater than five. This was agreed after considering that the minimally important difference is estimated as 3 points²⁴ but a more consistent improvement is expected after an invasive intervention. It was agreed that a change by 5 points would constitute a treatment success.




Figure 236 - Model structure. The health states are represented by the six blue circles on the top right corner. The arrows represent the possible transitions from a state to another or to the same state.

For each strategy the expected healthcare costs and expected QALYs were calculated by estimating the costs and QALYs for each state and then multiplying them by the proportion of patients who would be in that state as determined by the strategy taken.

We performed a probabilistic sensitivity analysis (SA) to test the robustness of the results against the imprecision of these estimates and the other model parameters, and to obtain more accurate estimates of expected costs and QALYs.

We identified sensitive parameters with a threshold analysis and then conducted multi-way sensitivity analyses on those parameters at decision point.

1.3.2 Key assumptions

The experts in the GDG were consulted in order to make the following assumptions:

- a) After a relapse in symptoms, only 5% of patients will undergo a second TURP. The remaining 95% are treated medically.
- b) The probability of success of the same intervention when performed a second time is 75% the probability of success when performed for the first time.

- c) The proportion of men with incontinence after surgery/laser requiring an AUS is 5%. The remaining 95% are treated medically or with incontinence products (catheters, pads, etc).

1.3.3 Probability of success - TURP

We searched for an RCT which reported the probability of success of either TURP or HoLEP as defined in our model (change in IPSS \geq 5). We found only one large multicentre RCT⁹⁴ where 120 of the randomised patients received TURP while the other 115 received TUVP. Data from this study⁹⁴ that were used in the model are reported in Table 2.

Table 2 - Data on TURP used in the model (a)

	Data used in the model
IPSS at baseline (IPSS pre)	20.7 (SD 6.9)
IPSS at 6 months (IPSS post)	6.9 (SD 5.5)
Probability of success of TURP at 6 months	85.4%
Probability of success of TURP at 24 months	84.0%

(a) From Fowler et al. (2005)⁹⁴

1.3.4 Probability of success - HoLEP

We could not find similar data for HoLEP so we adopted an alternative approach, linking the probability of success of the two interventions using the IPSS change data from our clinical review.

Table 3 - Effectiveness from meta-analysis

	HoLEP vs. TURP
Weighted Mean Difference (WMD) from baseline IPSS at 6 months	- 0.52
WMD from baseline IPSS at 24 months	- 0.80

1.3.4.1 Setting up the precondition

IPSS_{post} is the mean IPSS after the intervention and it is equal to:

$$I \quad IPSS_{post} = P_{success} * IPSS_{success} + (1 - P_{success}) * IPSS_{fail}$$

Where IPSS_{fail} and IPSS_{success} are respectively the mean IPSS in the group of patients whose treatment has failed and the mean IPSS in the group of patients whose treatment was successful.

By assuming that IPSS_{fail} is the same for both TURP and HoLEP and also that IPSS_{success} is the same for both, we can estimate the success rate for HoLEP.

1.3.4.2 Deriving IPSS after a TURP failure

$$\text{II } IPSS_{\text{fail}} = IPSS_{\text{pre}} - \Delta IPSS_{\text{fail}}$$

Where $\Delta IPSS_{\text{fail}}$ is the change in IPSS in patients for whom the intervention has failed. By definition this must be ≤ 4 . Assuming in some patients the symptoms might have deteriorated, we can consider the range -1 to 4, and use the central value 1.5, which is then varied in a sensitivity analysis. Substituting this value in II and using the data from TURP we get $IPSS_{\text{fail}} = 20.7 - 1.5 = 19.2$

1.3.4.3 Deriving IPSS after a successful TURP

We can rearrange equation I as

$$\text{III } IPSS_{\text{success}} = (IPSS_{\text{post}} - (1 - P_{\text{success}}) \times IPSS_{\text{fail}}) / P_{\text{success}}$$

Using data from Table 2 and our result for $IPSS_{\text{fail}}$ from 1.3.4.2 we get:

$$\text{IV } IPSS_{\text{success}} = (6.9 - 14.6\% \times 19.2) / 85.4\% = 4.8$$

1.3.4.4 Deriving IPSS after HoLEP

The mean difference in change in IPSS from baseline to 6 months was -0.52 compared with TURP (Chapter 8.3.1). The IPSS 6 months after HoLEP is simply the IPSS at 6 months for TURP plus this difference:

$$\text{V } IPSS_{\text{post}} = 6.9 - 0.52 = 6.4$$

1.3.4.5 Calculating the probability of HoLEP success at 6 months

We rearranged equation I to give us:

$$\text{VI } P_{\text{success}} = (IPSS_{\text{post}} - IPSS_{\text{fail}}) / (IPSS_{\text{success}} - IPSS_{\text{fail}})$$

Substituting the values derived above (1.3.4.2, 1.3.4.3, 1.3.4.4) we get:

$$\text{VII } P_{\text{success}} = (6.4 - 19.2) / (4.8 - 19.2) = 88.9\%$$

1.3.5 Probability of relapse

According to the data reported in Fowler et al (2005)⁹⁴, TURP was more effective after 6 months than after 24 months, as only 84% of patients had an improvement in symptoms by at least 5 points at 24 months compared to 85.4% of patients at 6 months Table 2. To mimic what happens in real practice, where a relapse in symptoms sometimes follows an initial improvement, it was necessary to incorporate a time-dependant probability of relapse after an initial success.

The probability of relapse between these two intervals (6 months and 24 months) is calculated as follows:

$$\text{VIII } (P_{\text{success } 6 \text{ months}} - P_{\text{success } 24 \text{ months}}) / P_{\text{success } 6 \text{ months}}$$

Which in case of TURP is equal to $(85.4\% - 84\%) / 85.4\% = 1.6\%$

We converted the probability of relapse of TURP over 18 months into a 3-month rate, which is the cycle length of the model, by using the formula:

$$\text{IX } 1 - \exp((\ln(1 - \text{relapse18months}))/6)$$

We used the same probability of relapse for HoLEP (a conservative assumption).

1.3.6 Probability of complications

Several complications of HoLEP and TURP were identified in the systematic review (Appendix E). In our economic model we only included those that would require additional treatment and generate additional costs.

To calculate the probability of complications following TURP (Table 4), we aggregated data from the TURP arm in every study included in our review, excluding the duplicates. We then compared the incidences of adverse events after TURP with those reported in the AUA¹⁴ and we found no considerable difference.

The incidence of complications following HoLEP (Table 4) was estimated by multiplying their probability after TURP by the risk ratio (RR) of HoLEP compared to TURP.

Table 4 - Probability of complications

	TURP	HoLEP	
	Probability	RR vs. TURP	Probability
Incontinence	4.0%	1.19	4.8%
Blood transfusion	6.2%	0.27	1.8%
Acute urinary retention (AUR)	3.9%	0.71	2.8%
Urinary tract infections	6.9%	0.45	3.1%
Transurethral syndrome	2.0%	0.31	0.6%
Strictures	7.2%	0.69	5.0%

All the adverse events were assumed to occur within three months after the intervention, and so within the same cycle in the model. All of them have associated one-off costs (see 1.3.11) and no detriment in quality of life with the exception of incontinence which has a lifetime cost and disutility (1.3.8).

1.3.7 Life expectancy

The mean age of the men when entering the model was 71 as this was the mean age of men in the diagnosis-related group 'Hyperplasia of prostate' in the Hospital Episode Statistics 2006/07.

Life expectancy in patients with LUTS was assumed to be the same as the general population in England and Wales. The remaining life expectancy for men aged 71 is 12.99 years, as reported in the Life Tables for the general population of England and Wales in the year 2005-2007 from the Government Actuary Department

(<http://www.gad.gov.uk/Documents/Demography/EOL/ILT%202005-07/wltewm0507.xls>).

1.3.8 Quality of life

The utility scores in Table 5 are a measure of the quality of life associated with LUTS and incontinence. A systematic search for quality of life in men with LUTS and with incontinence was performed (Appendix C). Studies were included if they reported utility values for the states of LUTS or incontinence.

Studies reporting utilities specific to non-compared interventions were excluded.

Two studies^{21,198} were excluded because the values were obtained from consensus rather than from patients or general public.

Kok et al (2002)¹⁴⁹ reported utility values according to the obstructive and irritative dimension of IPSS. However, using this study to estimate an average utility score for LUTS would have required further assumptions on the nature of the symptoms.

Ackerman et al (2000)⁸ assessed the preference of 13 patients to health states with the standard gamble technique. We excluded this study due to the small sample size but we used it as an alternative source of data in the sensitivity analysis.

Trueman et al (1999)²⁹⁷ designed a survey to collect EQ-5D scores by symptoms severity in 1115 men in the UK. The results of this study²⁹⁷ were used in our model and are reported in Table 5. Although the population in the model is made of men with moderate-to-severe LUTS we used the utility value for severe LUTS as 20.7 was the average IPSS of this population.

We found a UK study⁵⁸ reporting the deterioration in quality of life caused by incontinence. A multivariate analysis of EQ-5D scores, found that after controlling for age, gender and body mass index, incontinence was associated with a reduction in the EQ-5D score by 0.11 (SE 0.026). This value was subtracted from the remission and LUTS utility scores for the health states respectively characterised by symptoms remission and Incontinence and LUTS and Incontinence. The values thus obtained are reported in Table 5.

Among patients with incontinence, 5% require an artificial urinary sphincter while the remaining 95% are treated pharmacologically or with incontinence products. The utility score does not differ for these two subgroups.

Other adverse events were assumed to be negligible in terms of quality of life because they could be promptly treated.

Table 5 - Utility values

	Utility score
Remission (a)	0.91
LUTS (a)	0.71
Remission + Incontinence (a, b)	0.80
LUTS + Incontinence (a, b)	0.60

(a) Source: Trueman et al (1999)²⁹⁷

(b) Source: Currie et al (2006)⁵⁸

1.3.9 Calculating QALYs gained

For each strategy, the expected QALYs in each cycle are calculated as follows:

$$X \quad \text{Expected QALYs} = \sum (U_i \times P_i)$$

where

U_i = the utility score for health state i

P_i = the proportion of patients in health state i

and where health state i could be any of the health states reported in Table 1.

The proportion of patients in each health state depends on the effectiveness of the treatment, in terms of symptoms improvement and incontinence, and on the proportion of patients still alive, which falls as the number of cycles and therefore age increases.

The overall *lifetime expected QALYs* are given by the sum of QALYs calculated for each cycle. The *incremental QALYs gained* associated with a treatment strategy are calculated as the difference between the expected QALYs with that strategy and the expected QALYs with the comparator.

1.3.10 Cost of interventions

We adopted a bottom-up approach to calculate the intervention cost as differentiating the total costs for the two intervention was not possible by using national sources (NHS Reference Costs or Tariffs) or published evidence. In fact, no UK study could be found which reported the cost of HoLEP as this is performed only in a few UK centres only while TURP is a widespread technique. For this reason we decided to include only the capital cost of the HoLEP equipment as the TURP equipment is already present in every Urology centre. Only disposables used in TURP were included in the calculation.

We contacted the UK supplier of HoLEP equipment (SIGMACON) to obtain precise data on the cost of the machine and the cost and number of uses of disposables. We assumed the life span of the machine is 10 years. As we want to estimate the cost of the machine per patient, the GDG had to estimate the number of patients per centre undergoing surgery for LUTS in a year.

We found the cost of TURP disposables in a study⁹⁴ and the GDG estimated the number of uses. The data thus collected are reported in Table 6.

In addition to the cost of equipment, other factors influencing the total costs are the operating theatre cost, the length of stay after the intervention, and the complications. The costs of operating theatre and hospital stay are reported in Table 6 while the costs of complications are described in 1.3.11.

Table 6 – Resources used and costs

	HoLEP	Source
Cost of HoLEP machine	£150,000	UK supplier (SIGMACON)
Lifespan of HoLEP	10 years	Assumption
Number of patients per year per HoLEP machine	280	Expert opinion
Cost of morcellator blades (HoLEP)	£595 each	UK supplier (SIGMACON)
Number of uses per blade	10	UK supplier (SIGMACON)
Cost of fibres (HoLEP)	£550 each	UK supplier (SIGMACON)
Number of uses per fibre	20	UK supplier (SIGMACON)
Cost of loops (TURP)	£47	Expert opinion
Number of uses per loop	10	Expert opinion
Operating time TURP	60 minutes	Systematic review (Appendix E) (a)
Operating time HoLEP	75 minutes	Systematic review (Appendix E) (a)
Cost of urology operating theatre	£9 per minute	Local cost estimate
Median length of hospital stay after TURP (b)	3 days	Hospital Episode Statistics 2006/07
Median length of hospital stay after HoLEP (b)	2 days	Hospital Episode Statistics 2006/07
Mean cost per bed day	£204	National Schedule of Reference Costs 2006-07 for NHS Trust & PCT Combined – HRG LB25C

(a) Mean number of times reported in Gupta et al (2006)¹⁰⁸ and Montorsi et al (2004)²⁰².

(b) The median was used as an estimate of the mean to exclude outliers probably due to complications.

The annual cost of the HoLEP machine is a function of the capital cost of the machine, its life span and the discount rate according to the formula:

$$XI \quad E = K * r / [1 - (1 + r)^{-n}]$$

where E = annual cost of the machine

K = capital outlay (cost of purchasing the machine)

r = discount rate / interest rate = 3.5%

n = lifespan

The total cost of a single intervention can be represented by the formula:

$$\text{XII } TC_i = E/np + cDisp_i + opT_i * cTheatre + cComp * pComp_{A-i}$$

Where TC_i = total cost of the intervention i

E = annual cost of machine (only HoLEP)

np = number of patients using the machine per year

$cDisp_i$ = cost of disposables of intervention i

opT_i = operating time of intervention i

$cTheatre$ = cost of theatre per minute

$cComp_A$ = cost of treating complication A (Table 7)

$pComp_{A-i}$ = probability of complication A after intervention i (Table 4)

where i is either TURP or HoLEP and A is any complication described in Table 7.

1.3.11 Cost of complications

The complications included in the model and their probabilities are reported in 1.3.6. The GDG estimated the resources used to treat each complication as shown in Table 7 with the exception of acute urinary retention for which we used a UK economic study¹⁷. When a procedure could be performed as a daycase or inpatient, we checked this proportion in the Hospital Episode Statistics 2006/07².

Table 7 - Cost of complications

	COST	SOURCE
Blood transfusion	£635 (a)	Varney et al (2003) ³¹⁰
Stricture	£706 (b)	National Schedule of Reference Costs 2006-07 – HRG code LB30B
Acute urinary retention	£2,029 (c)	Annemans et al (2005) ¹⁷
Trans-urethral syndrome	£1,710 (d)	National Schedule of Reference Costs 2006-07: 1) High Dependency Unit – 0 organs supported XC07ZHDU; plus 2) Excess bed day - HRG LB25C
Urinary tract infections	£742 (e)	National Schedule of Reference Costs 2006-07 – HRG code LA04C

(a) cost of a transfusion of red blood cells

(b) weighted cost - £509 x 54% (daycase) + £938 x 46% (inpatient)

(c) cost of the most cost-effective intervention to treat AUR in the study

(d) cost of two days in HDU and two days in normal ward

$$(e) \text{ weighted cost} - £376 \times 10\%(\text{daycase}) + £783 \times 90\%(\text{inpatient})$$

Incontinence is a complication but it is also a health state in the model so its cost is calculated separately in 1.3.12.

1.3.12 Cost of health states

The possible health states in which a patient could be in the model are listed in Table 1. By collecting information on the resources used while in these states from the GDG experts, we calculated the costs reported in Table 8.

When the patient has a remission of symptoms, we assumed no further treatment would be necessary and this state has no cost associated.

If after the intervention a patient still has LUTS, he would undergo urodynamic studies to investigate the cause of the intervention failure. He would then be treated with either anticholinergics or alpha-blockers and be recalled for a visit every six months. We assumed that 50% would be treated with anticholinergics and 50% with alpha-blockers. The details of the cost calculations are reported in Table 8.

Table 8 - Cost of residual LUTS state

Resources used	Proportion of patients using the resource	Unit cost of resource	Total cost per month per patient
Alpha-blockers	50%	£0.35 (a)	£5.32
5mg Oxybutynin twice daily	25%	£0.39 (b)	£5.93
Other Anticholinergics	25%	£1.05 (c)	£15.97
One visit every 6 months	100%	£75 (d)	12.50
TOTAL			£39.72
Urodynamic studies (one-off)	100%	£165 (e)	-

(a) Average cost per day of Alfuzosin, Tamsulosin, Doxazosin, and Prazosin (BNF 57)

(b) Cost of treatment per day (BNF 57)

(c) Average cost per day of Darifenacin, Solifenacin, Tolterodine, Trospium, Propiverine and Fesoterodine (BNF 57)

(d) From National Schedule of Reference Costs 2006-07 – Consultant led follow-up attendance – outpatient face-to-face – Urology

(e) From National Schedule of Reference Costs 2006-07 - Outpatient procedure LB42Z

To estimate the cost of incontinence in men treated with drugs or products we searched for UK cost-of-illness studies excluding those studies conducted in women. We did not find any so we estimated the resources and their costs with the help of experts from the GDG (

Table 9).

Table 9 - Cost of incontinence in men treated with products or drugs

Resources used	Proportion of patients using the resource	Unit cost of resource	Total cost per month per patient (f)
3 ISC catheters per day	25%	£1.30	£29.66
1 indwelling catheter every 6 weeks	25%	£6.00	£1.08
5mg Oxybutynin twice daily	50%	£0.39 (a)	£5.93
Other anticholinergics	50%	£1.05 (b)	£15.97
1 pad a day	25%	£0.34	£2.58
1 leg bag per week	25%	£2.50	£2.71
1 overnight bag per night	25%	£0.10	£0.76
1 bag support, leg sleeve and Stalock Bard per week	25%	£6.00	£6.50
Sheath appliances	25%	£40.00 (c)	£10.00
1 district nurse visit per week	100%	£21.00 (d)	£91.00
1 specialist nurse visit every 6 months	100%	£66.00 (e)	£11.00
TOTAL			£177.19

(a) Cost of treatment per day (BNF 57)

(b) Average cost per day of Darifenacin, Solifenacin, Tolterodine, Trospium, Propiverine and Fesoterodine (BNF 57)

(c) Estimate on cost per month rather than number of items.

(d) From Curtis (2008)⁵⁹ – cost of district nurse per home visit including travel, excluding qualification

(e) From Curtis (2008)⁵⁹ – cost of specialist nurse per hour of client contact, excluding qualification

(f) These figures account for the proportion of patients who use that resource

In the model, 5% of the men with incontinence have an AUS implanted. The costs associated with this intervention are the one-off cost of urodynamic studies, the cost of implanting the AUS and the recurrent visits. The AUS needs to be re-implanted on average every ten years and this is taken into account in the model with a recurrent cost of the operation (Table 10).

Table 10 - Cost of artificial urinary sphincter (AUS)

Resources used	Frequency	Unit cost of resource	Source of cost
AUS implant	10 years	£4,137	National Schedule of Reference Costs 2006-07– HRG code LB21Z
Urology visit	6 months	£75	National Schedule of Reference Costs 2006-07– Consultant led follow-up attendance – outpatient face-to-face – Urology
Urodynamic studies	One-off	£165	National Schedule of Reference Costs 2006-07 - Outpatient procedure LB42Z

The costs associated with the ‘LUTS + Incontinence’ state are similar to the costs of the Incontinence state, while the ‘LUTS + Incontinence AUS’ state generates the same costs as the ‘LUTS+Incontinence AUS’ state with the addition of the anticholinergics (in 50% of the men) and alpha-blockers (in the other 50%).

For each strategy, the expected cost per cohort of patients is calculated as follows:

$$\text{XIII Expected cost} = C_s + \sum_{j=1}^{40} \sum_{i=1}^6 C_i P_{ij}$$

where

C_s = cost of the initial strategy (TURP or HoLEP)

C_i = cost of health state i

P_{ij} = proportion of patients in health state i in cycle j

and where health state i could be any stage in Table 1.

The proportion of patients in a health state depends on the magnitude of the improvement in symptoms specific to each treatment, its probability of causing incontinence, and on the proportion of patients still alive according to the mortality rate for the general population of England and Wales.

The overall lifetime expected costs are given by the sum of costs calculated for each cycle. The incremental cost associated with a treatment strategy is calculated as the difference between the expected cost with that strategy and the expected cost with the comparator.

1.3.13 Probabilistic sensitivity analysis

A probabilistic sensitivity analysis was performed to assess the robustness of the model results to plausible variations in the model parameters.

Probability distributions were assigned to each model parameter, where there was some measure of parameter variability (Table 11). We then re-calculated the main results 10000 times, and each time all the model parameters were set simultaneously, selecting from the respective parameter distribution at random.

Table 11 - Parameters and distributions used in the probabilistic sensitivity analysis

Description of variable	Mean value	Probability distribution	Parameters	Source
IPSS post treatment with TURP after 6 months	6.9	Normal	SD = 0.5102	Fowler et al (2005) ⁹⁴
IPSS post treatment with TURP after 2 years	7.5	Normal	SD = 0.6633	Fowler et al (2005) ⁹⁴
Initial IPSS	20.7	Normal	SD=0.6633	Fowler et al (2005) ⁹⁴

IPSS change when treatment fails	1.5	Triangular	Min=0 Likeliest=1.5 Max=3	Assumption
Weighted mean difference of IPSS at 6 months	0.52	Normal	SD=0.4235	Systematic review of clinical effectiveness
Weighted mean difference of IPSS at 2 years	0.8	Normal	SD=0.9847	Systematic review of clinical effectiveness
Capital cost of HoLEP	£150,000	None		UK Supplier SIGMACON
Lifespan of HoLEP machine (years)	10	Gamma (a)	$\alpha = 61.46$ $\lambda = 6.146$	Assumption
Number of patients per year	280	Gamma (a)	$\alpha = 61.46$ $\lambda = 0.2195$	Assumption
Cost of each blade	£595	None		UK Supplier SIGMACON
Cost of each fibre	£550	None		UK Supplier SIGMACON
Cost of each loop	£47	None		Experts opinion
Number of uses of a blade	10	Triangular (b)	Min=5 Likeliest=10 Max=15	UK Supplier SIGMACON
Number of uses of a fibre	20	Triangular (b)	Min=15 Likeliest=20 Max=25	UK Supplier SIGMACON
Number of uses of a loop	10	Triangular	Min=5 Likeliest=10 Max=15	Experts opinion
Cost of operating theatre per minute	£9	Gamma (a)	$\alpha = 61.46$ $\lambda = 6.829$	Local cost estimate
Operating time – HoLEP (minutes)	75	Triangular	Min=55 Likeliest=75 Max=95	Gupta at al (2006) ¹⁰⁸ and Montorsi at el (2004) ²⁰²
Operating time – TURP (minutes)	60	Triangular	Min=45 Likeliest=60 Max=75	Gupta at al (2006) ¹⁰⁸ and Montorsi at el (2004) ²⁰²
Cost bed day	£204	Gamma (c)	$\alpha = 4.925$ $\lambda = 0.0241$	National Schedule of Reference Costs 2006-07 Excess Bed Day HRG code LB25C
Hospital stay after HoLEP (days)	2	Triangular (d)	Min=1 Likeliest=2 Max=3	Hospital Episode Statistics 2006/07

Hospital stay after TURP (days)	3	Triangular (d)	Min=2 Likeliest=3 Max=4	Hospital Episode Statistics 2006/07
Cost of residual LUTS state	see 1.3.12	None		NCGC calculations
Cost of incontinence per three months (see 1.3.12)	£510	Gamma (a)	$\alpha = 61.46$ $\lambda = 0.1205$	NCGC calculation of cost of health states
Cost of AUS	£4,137	Gamma (c)	$\alpha = 7.089$ $\lambda = 0.0017$	National Schedule of Reference Costs 2006-07 HRG code L25 – LB21Z
Cost of treating AUR	£2,029	Gamma (a)	$\alpha = 61.46$ $\lambda = 0.0303$	Annemans2005 ¹⁷
Cost of treating TUR	See Table 7			
Cost of HDU per day	£651	Gamma (c)	$\alpha = 5.096$ $\lambda = 0.0078$	National Schedule of Reference Costs 2006-07 HDU – 0 organs supported XC07ZH DU
Cost of multichannel cystometry	£165	Gamma (c)	$\alpha = 4.094$ $\lambda = 0.0248$	National Schedule of Reference Costs 2006-07 Outpatient procedure LB42Z
Cost of treating strictures – daycase	£509	Gamma (c)	$\alpha = 4.055$ $\lambda = 0.008$	National Schedule of Reference Costs 2006-07 non elective LB30B
Cost of treating strictures – inpatient	£938	Gamma (c)	$\alpha = 3.344$ $\lambda = 0.0036$	National Schedule of Reference Costs 2006-07 non elective LB30B
Cost of blood transfusion	£635	Gamma (a)	$\alpha = 61.46$ $\lambda = 0.0968$	Varney et al (2003) ³¹⁰
Cost of treating UTI – daycase	£376	Gamma (c)	$\alpha = 3.926$ $\lambda = 0.0104$	National Schedule of Reference Costs 2006-07 LA04C
Cost of treating UTI - inpatient	£783	Gamma (c)	$\alpha = 3.079$ $\lambda = 0.0039$	National Schedule of Reference Costs 2006-07 LA04C
Cost of urology visit	£75	Gamma (c)	$\alpha = 7.898$ $\lambda = 0.1053$	National Schedule of Reference Costs 2006-07 Consultant led follow-up attendance, face-to-face - Urology
Number of visits every 3 months	0.5	Triangular	Min=0.25 Likeliest=0.5 Max=1	Experts opinion
Probability of AUR after TURP (see 1.3.6)	3.9%	Beta	$\alpha = 88$ $\beta = 2184$	Systematic review of clinical effectiveness

Proportion of patients with incontinence requiring an AUS	5%	Triangular	Min=2.5% Likeliest=5% Max=7.5%	Experts opinion
Probability of incontinence after TURP (see 1.3.6)	4.0%	Beta	$\alpha = 84$ $\beta = 2036$	Systematic review of clinical effectiveness
Probability of strictures after TURP (see 1.3.6)	7.2%	Beta	$\alpha = 180$ $\beta = 2316$	Systematic review of clinical effectiveness
Proportion of treating strictures - inpatient: daycase	0.46 : 0.54	None		Hospital Episodes Statistics 2006-07
Probability of success at 6 months after TURP	85%	Beta	$\alpha = 88$ $\beta = 15$	Fowler et al (2005) ⁹⁴
Probability of success at 2 years after TURP	84%	Beta	$\alpha = 63$ $\beta = 12$	Fowler et al (2005) ⁹⁴
Probability of blood transfusion after TURP (see 1.3.6)	6.2%	Beta	$\alpha = 197$ $\beta = 2977$	Systematic review of clinical effectiveness
Probability of TUR after TURP (see 1.3.6)	2.0%	Beta	$\alpha = 29$ $\beta = 1454$	Systematic review of clinical effectiveness
Probability of UTI after TURP (see 1.3.6)	6.9%	Beta	$\alpha = 111$ $\beta = 1488$	Systematic review of clinical effectiveness
Proportion of treating UTI - inpatient: daycase	0.9 : 0.1	None		Hospital Episodes Statistics 2006-07
Proportion of patients being re-operated after a first failure	5%	Triangular	Min=0% Likeliest=5% Max=10%	Experts opinion
Relative Risk of AUR – HoLEP vs. TURP	0.72	Log-normal	SD=0.313	Systematic review of clinical effectiveness
Relative Risk of incontinence – HoLEP vs. TURP	1.26	Log-normal	SD=0.213	Systematic review of clinical effectiveness
Relative Risk of strictures – HoLEP vs. TURP	0.69	Log-normal	SD=0.356	Systematic review of clinical effectiveness
Relative Risk of blood transfusion – HoLEP vs. TURP	0.27	Log-normal	SD=0.615	Systematic review of clinical effectiveness
Relative Risk of TUR – HoLEP vs. TURP	0.31	Log-normal	SD=1.685	Systematic review of clinical effectiveness
Relative Risk of UTI – HoLEP vs. TURP	0.45	Log-normal	SD=0.636	Systematic review of clinical effectiveness
Utility of severe LUTS	0.71	Beta	$\alpha = 80.23$ $\beta = 32.77$	Trueman et al (1999) ²⁹⁷

Utility of Remission	0.91	Beta	$\alpha = 33.67$ $\beta = 3.33$	Trueman et al (1999) ²⁹⁷
Disutility from incontinence	0.11	Normal	SD = 0.026	Currie et al (2006) ⁵⁸
Effectiveness when procedure is performed the second time compared to first time	75%	Triangular	Min=50% Likeliest=75% Max=100%	Experts opinion
Discount rate (cost and QALYs)	3.5%	None		

(a) We approximated the standard error (SE) of the mean by assuming the width of the 95% CI was 50% of the mean using the following equation: $SE=0.25 \times \text{mean} / Z_{0.0975}$

(b) Based on experts opinion

(c) We used the interquartile range (IQR) to approximately estimate the SE of the mean using the following equation: $SE=0.5 \times \text{IQR} / Z_{0.75}$

(d) Based on the range from HES 2006/07

1.3.14 Results of the cost-effectiveness analysis

We analysed the data deterministically (Table 12) and probabilistically (Table 13). We found that the results of the model were sensitive to various parameters and this is reflected in the extreme confidence intervals obtained with the probabilistic SA.

In the base case analysis HoLEP is more cost-effective than TURP but this result is overturned by minimal changes in variables (Table 12).

Table 12 - HoLEP vs. TURP - Results of base case analysis

	Mean cost (£)	QALYs	Incremental cost per QALY gained (HoLEP vs. TURP)	Sensitivity analysis
TURP	2,938	8.5761	-	TURP is cost-effective if: - probability that TURP fails <12% - probability that HoLEP fails >13.5% - RR of Incontinence (HoLEP vs TURP) >1.51 - WMD in IPSS change <0.17 - TURP is not possible after HoLEP - probability of incontinence after TURP and RR incontinence (HoLEP/TURP) are varied together (Figure 237).
HoLEP	2,920	8.6019	HoLEP dominates (a)	

(a) HoLEP dominates means that HoLEP is both more effective and less costly. Hence the ICER cannot be calculated.




Figure 237 - Two-way sensitivity analysis of probability of incontinence after TURP and RR of Incontinence HoLEP vs TURP. The green and blue areas of the graph represent respectively the combinations of the two parameters where TURP or HoLEP is cost-effective.

The instability of this conclusion is even more evident from the results of the probabilistic SA (Table 13).

Table 13 - Probabilistic SA results - HoLEP vs. TURP

Mean incremental cost/mean QALYs gained	95% CI – lower limit (£/QALY)	95% CI – upper limit (£/QALY)	Probability of being cost-effective at £20,000/QALY
TURP dominates (a)	HoLEP dominates	TURP dominates	HoLEP 48% TURP 52%

(a) TURP dominates means that TURP is both more effective and less costly. Hence the ICER cannot be calculated.

The probability of HoLEP being cost-effective (48%) is very close to the probability of TURP being cost-effective (52%) at a willingness to pay of £20,000/QALY (the NICE threshold). The probabilities are very similar for other willingness to pay thresholds (Figure 238).




Figure 238 - Acceptability curve of HoLEP and TURP

The uncertainty can also be graphically represented by plotting the results of the incremental analysis for all the 10,000 simulations into a cost-effectiveness plane (Figure 239). Each point represents the ICER of TURP vs. HoLEP for each simulation. The dotted line represents the £20,000/QALY threshold while the ellipse delimits the 95% confidence interval.




Figure 239 - Incremental cost-effectiveness scatterplot - HoLEP vs TURP

1.3.15 Discussion

HoLEP and TURP could be equally cost-effective.

TURP is the current standard of care in the UK while HoLEP is a relatively new technique practiced in a small number of UK centres. Although our analysis shows that HoLEP is at least as cost-effective as TURP, careful considerations should be given to recommending its widespread use.

The cost-effectiveness of HoLEP seems to be associated with the skills of the surgeons as the probabilities of complications depend on the expertise of the surgeon performing the operation. The probabilities as reported in the studies included in our clinical review, where HoLEP was performed by specialised surgeons, might be largely different from the actual events following an operation performed by a trainee surgeon. Therefore we might have overestimated the effectiveness of HoLEP.

Another overestimation might be due to the blood transfusion rate after TURP as estimated from our review of clinical studies. Some of the included studies¹⁴⁵ reported a blood transfusion rate after TURP higher than the average.

The major limitation of our model is the arbitrary definition of success (IPSS change of at least 5 points). Although other authors⁹⁴ have adopted this definition, it is still debatable whether a change of 5 points could be considered a remission in symptoms. Other authors¹⁷² have used an improvement by 10% in IPSS as a proxy for success but this was judged to be even more optimistic by our experts, as this would equate to 2 points of improvement when the baseline score is 20.

The results of our study are based on trial data for men with moderate-to-severe symptoms with a mean baseline IPSS of 20.7. For men with less severe symptoms, TURP might be more cost-effective as it is less costly, while for men with more severe symptoms HoLEP might be more cost-effective as it is more effective than TURP at improving symptoms.

We compared the results of our study with the economic analysis from the HTA¹⁷² included in our review and we found similar results and conclusions. In this study¹⁷², HoLEP was more effective and less costly than TURP but the results were highly sensitive to several parameters. Unlike this study¹⁷² our model takes into account the capital cost of HoLEP which might explain the higher cost of HoLEP compared to TURP in the mean results of the probabilistic analysis.

From an NHS perspective, the results of our study would suggest training new surgeons in HoLEP could improve outcomes and save costs if performed correctly. However, a shift from TURP to HoLEP would have to be gradual for it to be cost-effective since purchasing the new equipment might not warrant the improved outcomes which were marginal. It is important to note that there is still inadequate long-term data for HoLEP. However, if a centre has to replace old equipment and surgeons trained in HoLEP are available, HoLEP could be an efficient option.

In conclusion, given the learning curve associated with the new technique and the cost of purchasing the new equipment, the GDG felt it was reasonable to recommend HoLEP only in centres specialised in the technique.

1.3.16 Conclusions

- HoLEP and TURP are similarly cost-effective
- In settings where HoLEP is not currently performed, TURP is more cost-effective because of the capital cost and the learning curve

1.4 NCGC Combination model

An economic model comparing Alpha-Blockers (AB) with a combination of AB and 5-Alpha-Reductase Inhibitors (Comb) was developed further to the exclusion of any economic evidence focusing on this comparison. The main outcomes considered were the change in IPSS from baseline and the treatment adverse events which were expressed in quality of life measures. Patients in this model are men who have moderate lower urinary tract symptoms and are selected for medical treatment. Studies specifically conducted on patients with a prostate size larger than the average^{191,263} were not used to estimate IPSS change as it was the GDG opinion that this would have favoured the Combination intervention.

We built a Markov model with a lifetime horizon (Figure 240) and we chose a cycle length of six months as it was the shortest follow up period in our clinical review of effectiveness (Chapter 6.10.1). All the probabilities, costs and health utilities were converted in order to reflect the six-month values. The time horizon was shortened to 5 years in a sensitivity analysis.

After a treatment period of six months, men can have either a meaningful improvement in IPSS (treatment success) or a negligible/no improvement (treatment failure). During this period they can also experience various adverse events which are independent from the treatment success. However, a proportion of those men experiencing adverse events will discontinue treatment, going back to the LUTS state. Men who had a treatment failure to start with will go to the LUTS state (with or without adverse events) but they can still have an improvement in the following six month cycle. Some men in the LUTS state will undergo TURP and they will feed into the TURP model (1.3).




Figure 240 - Structure of the combination model. The squared boxes represent the chance nodes in the model while the circles are the possible health states.

The list of the health states that are part of the combination model is reported in Table 14.

Table 14 - Health states of combination model

HEALTH STATES
(Moderate) LUTS
Remission
LUTS +adverse events
Remission + adverse events
TURP

While in the Surgery model a significant remission of symptoms was a change in IPSS greater than five, in the Combination model we used the 3 point estimate by Barry et al (1995)²⁴.

For each strategy the expected healthcare costs and expected QALYs were calculated by estimating the costs and QALYs for each state and then multiplying them by the proportion of patients who would be in that state as determined by the strategy taken.

We performed a probabilistic sensitivity analysis (PSA) to test the robustness of the results against the imprecision of these estimates and the other model parameters, and to obtain more accurate estimates of expected costs and QALYs.

1.4.1 Key assumptions

The experts in the GDG were consulted in order to make the following assumptions:

- a) Patients are kept on treatment for all their life if the treatment is effective and there are no adverse events.
- b) If the treatment does not work (i.e. IPSS improves by less than 3 points) the treatment is kept for one year then it is discontinued.
- c) 50% of the patients who discontinue the treatment after one year undergo TURP.
- d) If adverse events have not occurred during the first two years, they will never occur.

The following assumption was based on the conclusions of our clinical review:

- a) After the first year the treatment effectiveness is stable (no improvement or deterioration in IPSS are possible).

1.4.2 Probability of success

We could not find any studies reporting the proportion of successful treatment where success was defined as an improvement of at least 3 points of IPSS. We assumed that the IPSS change was normally distributed and we used the standard deviation (SD) from the mean to obtain the proportion of cases within the 3-point cut-off (Table 15). This was calculated as:

$$\text{Success rate} = 1 - \Phi_{\mu\sigma^2}(\text{IPSS}) \text{ where IPSS}=3,$$

where μ =mean IPSS, σ^2 =IPSS variance= IPSS SD squared (Table 15), 3 is the IPSS cut-off for success and where $\Phi_{\mu\sigma^2}(\text{IPSS})$ gives the cumulative distribution function for a normal distribution with mean μ and variance σ^2 .

Table 15 - Probability of treatment success when the cut-off is 3 points

	Mean IPSS change (a)	SD of IPSS change (a)	Proportion of treatment success
AB – 6 months	6.3	5.8	72%
Comb – 6 months	6.1	7.4	66%
AB – 12 months	7.1	5.7	76%
Comb – 12 months	7.3	5.8	77%

a) Source: clinical review.

As the figures in Table 15 suggest, treatment success is more likely achieved at 12 months than 6 months. Therefore men in the model for whom treatment has failed in the first six months can still experience a remission in the following 6 months. The probability of remission is simply the difference between the probability of success at 12 months and the probability of success at 6 months (Table 16).

Table 16 - Probability of symptoms remission at 12 months

	P success 6 months	P success 12 months	P remission between 6 and 12 months (a)
AB	72%	76%	14.3%
Comb	66%	77%	16.6%

a) $(P \text{ success } 12 \text{ months} - P \text{ success } 6 \text{ months}) / (1 - P \text{ success } 6 \text{ months})$

We changed the definition of success in sensitivity analyses where we defined success as an improvement by at least 5 or at least 8 points.

1.4.3 Probability of adverse events and withdrawals

We looked for RCT data on adverse events and withdrawals due to adverse events. We realised it was not feasible to estimate the incidence of specific adverse events and their specific probability of causing withdrawals from treatment. Consequently we adopted a three-step approach:

1. estimate the overall probability of a man experiencing a drug-related adverse event with AB and with combinations
2. estimate the probability of an adverse event leading to treatment discontinuation with AB and with combination
3. once an adverse event occurs, estimate the probability of specific adverse events

We found a large RCT²⁶³ reporting both drug related adverse events and drug-related adverse events leading to study withdrawals. With these data (Table 17) we were able to perform step 1 and 2 (Table 17).

Table 17 - Probability of discontinuation in patients with adverse events*

	Number of drug-related adverse events x	Number of drug-related adverse events leading to withdrawal y	Probability of drug-related adverse events	Probability of discontinuation in patients with adverse events $z=y/x$
AB	258	48	16%	18.6%
Comb	386	80	24%	20.7%

* From Roehrborn et al (2008)²⁶³

Figure 241 and Figure 242 illustrate how these values were used in the model.




Figure 241 - Adverse events in the AB arm of the model




Figure 242 - Adverse events in the combination arm of the model

For step 3 we used the evidence from the review of clinical effectiveness (Chapter 6.10.1). Various adverse events were reported in the included studies and in order to avoid double-counting we grouped those adverse events that could be similar in symptoms. The most common adverse event was used to represent the group (Table 18). Therefore whilst in the clinical review postural hypotension, headache, syncope and dizziness are all reported, it is likely to be an overlap of those symptoms and just dizziness (the most frequent one) is reported as part of that group. Similarly decreased libido was grouped together with impotence or erectile dysfunction.

In our model we did not use the incidences reported in the included studies (Chapter 6.10.1) but these were used to calculate the probability of each type being the adverse event occurring (Table 18).

Table 18 - Incidence and proportion of adverse events

	Incidence		Proportion of adverse events	
	AB X_i	Comb Y_i	AB $X_i/\sum X_i$	Comb $Y_i/\sum Y_i$
Dizziness	4.8%	4.3%	22%	16%
Fatigue	3.6%	4.2%	17%	16%
Rhinitis	6.6%	7.8%	31%	29%
Ejaculatory abnormality	0.6%	3.0%	3%	11%
Impotence/erectile dysfunction	3.0%	5.9%	14%	22%
Breast enlargement	1.8%	1.4%	8%	5%
Acute urinary retention (AUR)	1.0%	0.4%	5%	1%
TOTAL	21.4%	27.0%	100%	100%

The probability of each adverse event group was used in the model to estimate the detriment in quality of life and additional costs due to adverse events (see 1.4.5 and 1.4.7).

1.4.4 Life expectancy

Men in the Combination Model were assumed to be on average 60 years old.

Life expectancy in patients with LUTS was assumed to be the same as the general population in England and Wales. The remaining life expectancy for men aged 60 is 21.22 years, as reported in the Life Tables for the general population of England and Wales in the year 2005-2007 from the Government Actuary Department (<http://www.gad.gov.uk/Documents/Demography/EOL/ILT%202005-07/wltewm0507.xls>).

1.4.5 Quality of life

The same sources used in the Surgery Model for quality of life estimates of the residual LUTS and remission states (1.3.8) were used in the Combination Model. However, while men in the Surgery Model had on average severe symptoms, in the Combination Model men have moderate symptoms.

The health states ‘Remission + Adverse events’ and ‘LUTS + Adverse events’ are made of the Remission or LUTS utility value and the disutility (decrease in utility) due to adverse events.

Being the spectrum of adverse events in the AB arm different from that in the combination arm (1.4.3), the adverse events health states will also have different utility values in the different arms.

The utility value of the LUTS + adverse events state for intervention y will be calculated as:

$$\text{XIV } u_{\text{LUTS-AE}y} = u_{\text{LUTS}} + \sum(\text{disutility}_{\text{AE}i} * p_{\text{AE}iy})$$

where u_{LUTS} is the utility values of Moderate LUTS reported in Table 19,

$\text{disutility}_{\text{AE}i}$ is the disutility of the adverse event i where i is any of the adverse events reported in Table 18,

and $p_{\text{AE}iy}$ is the proportion of the adverse event i for the intervention y , where y could be either AB or combination.

From equation **XIV** it can be deduced that the utility of these health states depend on the intervention being the proportion of adverse events the variable parameter.

We conducted a search in the CEA Registry (<https://research.tufts-nemc.org/cear/default.aspx>) to find quality of life values associated with the adverse events reported in Table 18.

Two studies^{289,311} were found which reported the one-day disutilities deriving from dizziness, fatigue and rhinitis. We assumed that those symptoms were experienced half the time; therefore the original value was halved in our analysis (Table 19) but this assumption was varied in sensitivity analyses.

One study²³⁸ reported the disutility due to breast enlargement.

In a study by Dedhia et al (2008)⁷⁰ patients with LUTS were interviewed and their time-trade off scores for various adverse events collected. The utility values reported in this study were 0.71 for ejaculatory abnormality and 0.73 for erectile dysfunction in men with LUTS. If we assume that the utility decrements are additive, we can calculate the disutility due to these adverse events as the difference of the utility of LUTS and the utility of adverse event in presence of LUTS:

$$\text{XV } \text{disutility}_{\text{AE}} = u_{\text{LUTS}} - u_{\text{LUTS+AE}}$$

By substituting the values from the study⁷⁰ in formula **XV** we obtain the disutilities reported in Table 19.

Table 19 - Utility values used in the Combination Model

	Utility score	Source
Remission	0.91	Trueman et al (1999) ²⁹⁷
Moderate LUTS	0.78	Trueman et al (1999) ²⁹⁷
Disutility breast enlargement	- 0.05	Penson et al (2005) ²³⁸
Disutility dizziness (a)	- 0.11	Vera-Llonch et al (2008) ³¹¹
Disutility ejaculatory abnormality	-0.07	Dedhia et al (2008) ⁷⁰
Disutility fatigue (a)	-0.125	Vera-Llonch et al (2008) ³¹¹
Disutility impotence	-0.05	Dedhia et al (2008) ⁷⁰
Disutility rhinitis (a)	-0.095	Sullivan et al (2004) ²⁸⁹
Disutility AB adverse events	- 0.088	Weighted average of above disutilities
Disutility Comb adverse events	- 0.086	Weighted average of above disutilities

(a) Assuming symptoms are experienced half the time.

The disutility due to Acute Urinary Retention (AUR) was not included in the model as this complication was assumed to be treated and resolved within six months. The cost associated with this adverse event is already explained in the Surgery Model (see 1.3.11).

1.4.6 Calculating QALYs gained

See 1.3.9.

1.4.7 Cost of interventions and health states

The cost components of the health states in the model are made of the continuous cost of drug therapy and the cost of visits (Table 20). During the first six-month cycle men are treated with either AB or Combination and have a follow-up visit. The cost of the initial treatment is kept for at least another cycle unless there is a discontinuation due to adverse events. If the treatment is discontinued only the cost of a visit is included in the cost of a cycle.

Table 20 - Resources used in the health states of the model

HEALTH STATE	RESOURCES USED
Moderate LUTS - initial	Drugs (AB or Comb) + 1 follow-up visit
Moderate LUTS - residual	1 follow-up visit
Remission	Drugs (AB or Comb)
LUTS +adverse events	1 follow-up visit
Remission + adverse events	Drugs (AB or Comb)

The cost details of the resources used in the health states are reported in Table 21.

Table 21 - Cost of resources used

Resource	Total cost per patient over six months	Source
Alpha-blockers	£65	BNF 57 (a)
Combination (5-ARI+AB)	£186	BNF 57 (b)
Follow-up visit	£75	National Schedule of Reference Costs 2006-07 – Consultant led follow-up attendance – outpatient face-to-face – Urology

a) Based on the average cost per day of Alfuzosin, Tamsulosin, Doxazosin, and Prazosin = £ 0.35

b) Based on the cost of AB and on the average cost per day of Dutasteride and Finasteride = £0.66

In addition, some costs are associated with particular events in the model: the cost of treating AUR when adverse events occur (adjusted by the proportion of AUR in the adverse events) and the cost of TURP if the therapy fails and the man considers surgery. In this event the model feeds directly into the Surgery Model described in 1.3 where the cost components are the same ones described in 1.3.10 and 1.3.11 for the TURP strategy.

1.4.8 Probabilistic sensitivity analysis

A probabilistic sensitivity analysis was performed to assess the robustness of the model results to plausible variations in the model parameters.

The same method described for the Surgery Model (1.3.13) was used for the Combination Model. The same parameters used in the TURP arm of the Surgery Model were used in the Combination Model when men undergo TURP after a treatment failure. All the other parameters and their distributions are listed in Table 22.

Table 22 - Parameters and distributions used in the probabilistic sensitivity analysis

Description of variable	Mean value	Probability distribution	Parameters	Source
Mean IPSS change at 6 months – AB	6.3	Normal	SD= 5.8	Systematic review of clinical effectiveness
Mean IPSS change at 6 months – Comb	6.1	Normal	SD=5.6	Systematic review of clinical effectiveness
Mean IPSS change at 12 months – AB	7.1	Normal	SD=5.7	Systematic review of clinical effectiveness
Mean IPSS change at 12 months – Comb	7.3	Normal	SD=5.8	Systematic review of clinical effectiveness

Probability of success at 6 months – AB	72%	None (function of IPSS change)		See 1.4.2
Probability of success at 6 months - Comb	66%	None (function of IPSS change)		See 1.4.2
Probability of success at 12 months – AB	76%	None (function of IPSS change)		See 1.4.2
Probability of success at 12 months - Comb	77%	None (function of IPSS change)		See 1.4.2
Probability of remission at 12 months – AB	14.3%	None (function of probability of success)		See 1.4.2
Probability of remission at 12 months - Comb	16.6%	None (function of probability of success)		See 1.4.2
Cost of Alpha-blockers treatment over 6 months	£65	None		BNF 57
Cost of combination treatment over 6 months	£186	None		BNF 57
Cost of urology visit	£75	Gamma (a)	$\alpha = 7.898$ $\lambda = 0.1053$	National Schedule of Reference Costs 2006-07 Consultant led follow-up attendance, face-to-face - Urology
Cost of treating AUR	£2,029	Gamma (b)	$\alpha = 61.46$ $\lambda = 0.0303$	Annemans et al (2005) ¹⁷
Probability of adverse events - AB	16%	Beta	$\alpha = 258$ $\beta = 1353$	Roehrborn et al (2008) ²⁶³
Probability of adverse events - Comb	24%	Beta	$\alpha = 386$ $\beta = 1224$	Roehrborn et al (2008) ²⁶³
Probability of discontinuing in men with adverse events - AB	18.6%	Beta	$\alpha = 48$ $\beta = 210$	Roehrborn et al (2008) ²⁶³
Probability of discontinuing in men with adverse events - Comb	20.7%	Beta	$\alpha = 80$ $\beta = 306$	Roehrborn et al (2008) ²⁶³
Proportion of breast enlargement/adverse events AB	8%	Dirichlet	0.08,	Systematic review of clinical effectiveness

Proportion of dizziness/adverse events AB	22%	Dirichlet	0.22,	Systematic review of clinical effectiveness
Proportion of fatigue/adverse events AB	17%	Dirichlet	0.17,	Systematic review of clinical effectiveness
Proportion of ejaculatory abnormality/adverse events AB	3%	Dirichlet	0.03,	Systematic review of clinical effectiveness
Proportion of impotence/adverse events AB	14%	Dirichlet	0.14,	Systematic review of clinical effectiveness
Proportion of rhinitis/adverse events AB	31%	Dirichlet	0.31,	Systematic review of clinical effectiveness
Proportion of AUR/adverse events AB	5%	Dirichlet	0.05	Systematic review of clinical effectiveness
Proportion of breast enlargement/adverse events - Comb	5%	Dirichlet	where each parameter refers to proportion of each type of adverse event	Systematic review of clinical effectiveness
Proportion of dizziness/adverse events - Comb	16%	Dirichlet	0.16,	Systematic review of clinical effectiveness
Proportion of fatigue/adverse events – Comb	16%	Dirichlet	0.11,	Systematic review of clinical effectiveness
Proportion of ejaculatory abnormality/adverse events AB	11%	Dirichlet	0.22,	Systematic review of clinical effectiveness
Proportion of impotence/adverse events – Comb	22%	Dirichlet	0.29,	Systematic review of clinical effectiveness
Proportion of rhinitis/adverse events – Comb	29%	Dirichlet	0.01	Systematic review of clinical effectiveness
Proportion of AUR/adverse events – Comb	1%	Dirichlet	where each parameter refers to proportion of each type of adverse event	Systematic review of clinical effectiveness
Proportion of men undergoing TURP after treatment failure	50%	Triangular	Min=0% Likeliest=50% Max=100%	Experts opinion
Utility of Moderate LUTS	0.78	Beta	$\alpha = 80.23$ $\beta = 32.77$	Trueman et al (1999) ²⁹⁷
Utility of Remission	0.91	Beta	$\alpha = 33.67$ $\beta = 3.33$	Trueman et al (1999) ²⁹⁷
Disutility from breast enlargement	0.05	Beta	$\alpha = 23.7$ $\beta = 450.3$	Penson et al (2005) ²³⁸

Disutility from dizziness	0.11	Beta	$\alpha = 6.22$ $\beta = 50.32$	Vera-Llonch et al (2008) ³¹¹
Disutility from fatigue	0.125	Beta	$\alpha = 6.097$ $\beta = 42.681$	Vera-Llonch et al (2008) ³¹¹
Disutility from ejaculatory abnormality	0.07	Beta	$\alpha = 14.81$ $\beta = 196.76$	Dedhia et al (2008) ⁷⁰
Disutility from impotence/erectile dysfunction	0.05	Beta	$\alpha = 6.706$ $\beta = 127.406$	Dedhia et al (2008) ⁷⁰
Disutility from rhinitis	0.19	Beta	$\alpha = 20.604$ $\beta = 87.836$	Dedhia et al (2008) ⁷⁰
Discount rate (cost and QALYs)	3.5%	None		NICE Reference Case

(a) We used the interquartile range (IQR) to approximately estimate the standard error (SE) of the mean using the following equation: $se=0.5 \times IQR / Z_{0.75}$

(b) We approximated the SE of the mean by assuming the width of the 95% CI was 50% of the mean using the following equation: $se=0.25 \times \text{mean} / Z_{0.975}$

1.4.9 Results

Alpha-blockers generate less cost and more QALYs compared to combinations (Table 23).

Table 23 - Results of base case analysis - Combination vs. Alpha-blockers

	Mean cost (£)	QALYs	Incremental cost (£) per QALY gained	Sensitivity analysis
Alpha-blockers	3,824	12.4347	-	One-way SA: Combination is cost-effective if probability of adverse events with AB > 29% (16% in base case). Results were not sensitive to other changes in parameters or structure.
Combination	6,411	12.4276	Dominated	

In a set of one-way sensitivity analyses, where the low and high values were respectively half or double the base case value, we identified the parameters that might have changed the results. The only variable to which the model was sensitive was the probability of adverse events with AB. We explored this uncertainty further through a two-way SA where the probability of adverse events with AB was co-varied with the probability of adverse events with combination (Figure 243).




Figure 243 - Two-way SA on probability of adverse events with AB (x axis) and comb (y axis). The area in green is where AB is cost-effective, while the area in blue is where combination is cost-effective. The black dot represents the base case values.

If we consider a 95% confidence interval the base case results did not reach statistical significance (Table 24).

Table 24 - Results of probabilistic SA - Comb vs. AB

Mean ICER (£/QALY)	95% CI – lower limit (£/QALY)	95% CI – upper limit (£/QALY)	Probability of being cost-effective at £20,000/QALY
Comb dominated	3,850	Comb dominated	AB 90% Comb 10%

However, at a willingness to pay of £20,000/QALY alpha-blockers have a 90% probability of being cost-effective (Figure 244).




Figure 244 - Acceptability curve of AB and Comb

1.4.10 Discussion

5-ARI and AB have a different mechanism of action and the combination of the two could enhance the effectiveness on men with LUTS. Our review of clinical evidence (Chapter 6.10.1) has shown that the long-term (one year) improvement in IPSS is higher with combinations than with AB. However there are extra costs associated with the improvement and more side effects. The results of our model show that after weighting the advantages (improvement in IPSS) and disadvantages (costs and side effects) combinations are not cost-effective in a general population of men with LUTS.

We based our model on studies where men had a normal prostate size. We have deliberately excluded those studies conducted on men with large prostates as 5-ARI are believed to be more effective in this group of men. A specific model for that population could be built once good data are available.

We encountered some challenges when building our model: defining success of treatment according to an IPSS improvement by 3 points might have been arbitrary even if based on a previous study²⁴; however, when we changed this definition to up to 10 points the overall results did not change.

Other assumptions were made while building the model but those did not have an impact on the conclusions.

Adverse events were a core component of the model and their incidence was the only parameter to which the results were sensitive. When we changed the probability of adverse events with AB and combinations simultaneously we noted that if the probability was lower with combination than with AB the former would have been more cost-effective than the latter. Nevertheless, as AB are part of the combination it would be very unlikely that their adverse events while used in combination would be less frequent than when they are used alone.

This is the only model which compares AB and combination using randomized data. A cost-utility analysis by McDonald et al (2004)¹⁹² concluded that combinations were more cost-effective than Doxazosin but the clinical data were obtained from men with large prostate for one arm and men with normal prostate for the other arm. This explains the higher value-for-money of combination in this study compared to ours. Conversely the cost-utility analysis by DiSantostefano et al (2006)⁷¹ reached our same conclusions, yet the effectiveness data on combinations were not based on RCTs but on assumptions.

1.4.11 Conclusions

- Combination of alpha-blockers with 5-ARI was not cost-effective in a general population of men with LUTS.
- Clinical data on men with large prostate might be useful to assess the cost-effectiveness in this group where combinations are presumed to be more effective.

Appendix G - Recommendations for research

1.1 Multichannel cystometry

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome).</p>	<p>Question: What is the clinical and cost effectiveness of multichannel cystometry in improving patient related outcomes in men being considered for bladder outlet surgery? Patients: Bothering LUTS not responding to conservative therapy (catheterised patients excluded). Intervention: Pressure flow studies. Comparison: Two groups, awaiting bladder outlet surgery, randomised either to pre-operative pressure flow studies, or not Outcome: Primary outcome-patient-related outcome (IPSS, EQ5D), secondary outcomes-adverse events, flow rate, residual urine, pdetQmax.</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>This research would clarify whether this test could improve the outcome of surgery. If the result is positive, this could improve the chance of a good outcome from surgery.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>As above, it would add to knowledge about the utility of pressure flow studies and allow them to be recommended or not recommended in future revisions of guidance.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>It would allow the NHS to know whether resources should be committed to the test or not.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>NSF for older people, Integrated Continence Services.</p>
<p><u>Current evidence base</u> What is the current evidence base? What</p>	<p>There are currently no randomised controlled trials comparing multichannel</p>

<p>are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>cystometry to no intervention in men before surgery.</p>
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>No specific consideration.</p>
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p><i>Design:</i> A randomised comparative trial of men awaiting bladder outlet surgery, to be randomised to either a pressure flow study or not, before their surgery. The results of the pressure flow study would be used in subsequent counselling of patients in a protocol-driven way, before the proposed surgery, and <i>might</i> result in surgery not being done.</p> <p><i>Outcome:</i> As above.</p>
<p><u>Feasibility</u> Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The research would be ethically and technically feasible.</p>
<p><u>Other comments</u> Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>The National Institute for Health Research (NIHR) would be an appropriate funding source. The normal service delivery cost to participants would be taken over by the research during the trial, thus relieving the service delivery budget. Since the NIHR is an NHS funded body the costs of care would simply be shifted from one NHS budget to another. Additional costs would be those associated with conducting the research itself.</p>
<p><u>Importance</u> How important is the question to the</p>	<p>High. The research is essential to inform future updates of key recommendations in</p>

overall guideline? The research recommendation should be categorised into one of the following categories of importance:

- High: the research is essential to inform future updates of key recommendations in the guideline
- Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates
- Low: the research is of interest and will fill existing evidence gaps.

the guideline.

1.2 Catheterisation

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>What are the clinical and cost effectiveness and associated adverse events of intermittent catheterisation compared to indwelling suprapubic or urethral catheterisation for men with voiding difficulty and chronic retention of urine?</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>The number of men judged unfit to undergo de-obstructing surgery is steadily increasing given the increasing proportion of older men in the population. Current practice varies widely across the UK with no established standard for long term management and no systematic review of practice. The research could establish the best approach to management in these men in the longer term and so bring more effective treatment, better focused on each patient's need, and consequent cost-efficiency gains.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>NICE currently cannot give clear guidance on this topic because of an inadequate evidence base.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>Catheters are currently used variably across the UK with no systematic approach to management except for men with spinal cord injury. The aim of catheterisation, to drain the bladder so as to protect the upper renal tracts and maintain continence may not be achieved acceptably. Evidence-based guidance on the selection of the most suitable mode of catheterisation will benefit the quality of life of patients, ensure the efficient use of skilled staff and may reduce the costs of waste of unsuitable or sub-optimal product use.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>None currently relevant.</p>
<p><u>Current evidence base</u> What is the current evidence base? What</p>	<p>There is no currently no evidence for these interventions.</p>

<p>are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>This treatment predominantly affects older people.</p>
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>A randomised controlled study of the interventions:</p> <ul style="list-style-type: none"> a) intermittent catheterisation b) indwelling suprapubic catheterisation c) indwelling urethral catheterisation <p>Outcomes of interest: quality of life, healthcare resource utilisation, adverse events (including leakage, skin breakdown, infection, erosion and death).</p>
<p><u>Feasibility</u> Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The major issues with this trial would be the identification of cases and the studying of them in a primary care environment.</p> <p>An adequate population of men with this problem already exists precisely because of the absence of any consensus strategy for this group.</p>
<p><u>Other comments</u> Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>None.</p>

<p><u>Importance</u> How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none">• High: the research is essential to inform future updates of key recommendations in the guideline• Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates• Low: the research is of interest and will fill existing evidence gaps.	<p>High. Surgery is indicated as therapy for retention – but may not be appropriate in the presence of impaired bladder function (underactive) or where comorbidity precludes it.</p>
---	---

1.3 Products for men with urinary incontinence

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>What is the clinical and cost effectiveness and associated adverse events of absorbent pads compared to sheath collectors for men with urinary incontinence?</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>The number of patients in this group is steadily increasing with more radical prostatectomies and an ageing demographic. Current practise varies widely across the UK with no established standards of good practice. The research could establish the best approach to continence management in these men and so bring more effective treatment, better focussed on each patient's needs, and consequently cost-efficiency gains.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>NICE currently cannot give clear guidance on this topic because of an inadequate evidence base.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>Containment products are currently used variably across the UK. It is rare that any element of bladder training or recognition and treatment of bladder dysfunction is recognised as part of the continence management problem. The aim, so often, is simply to keep the patient socially dry; and even that is not always achieved acceptably. Evidence-based guidance on the selection of the most suitable containment product and its subsequent management will benefit the quality of life of patients, use skilled nurse/career resources more efficiently and reduce the costs of waste of unsuitable or sub-optimal product use.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>There is currently no national service framework for men with LUTS and incontinence or difficulty with bladder emptying.</p>
<p><u>Current evidence base</u> What is the current evidence base? What are the problems with the current</p>	<p>There is no currently no level 1 evidence for pads and sheaths.</p>

<p>evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>There are no equality issues.</p>
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>A randomised controlled trial to compare these interventions. Outcomes of interest would be symptom severity, quality of life, changes in measured leakage, and occurrence of adverse events.</p>
<p><u>Feasibility</u> Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The major issues with this trial would be the identification of cases and the studying of them in a primary care environment.</p> <p>An adequate population of men with this problem already exists precisely because of the absence of any consensus strategy for this group.</p>
<p><u>Other comments</u> Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>In general, manufacturers have been reluctant to fund randomised controlled trials. Currently the D4D project is addressing unmet needs. Work with specialist and patient advocacy groups and manufacturers will be essential.</p>
<p><u>Importance</u> How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline • Medium: the research is relevant to the recommendations in the guideline, but the 	<p>High. This is a population of men who have been rendered incontinent by surgery. The impact on their quality of life is profound and there is currently only one realistic treatment option for more major incontinence namely surgery which many men find unacceptable. It is important that solutions are found for this growing number of men.</p>

<p>research recommendations are not key to future updates</p> <ul style="list-style-type: none">• Low: the research is of interest and will fill existing evidence gaps.	
--	--

1.4 Laser vapourisation techniques

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>What is the clinical and cost effectiveness and associated adverse events of laser vaporisation techniques compared to TURP in men with moderate to severe bothersome LUTS considering surgery for bladder outlet obstruction? Assessed by symptom severity, quality of life, and adverse events.</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>The potential advantages of reduced blood loss, shorter hospital stay and earlier return to normal activities make laser vaporisation techniques attractive to patients and healthcare providers although there is uncertainty around degree of symptom improvement and improvement in quality of life in the short and longer term.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>NICE cannot give clear guidance on this intervention because the evidence base is inadequate. The proposed research will add new knowledge.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>Green Light laser use in the NHS is increasing at a rapid rate with approximately 70 units in the UK using it (~ 60% NHS and ~ 40% private sector) from personal communication with representatives of American Medical Systems Inc and clinical units. This is despite a lack of clinical and cost-effectiveness data to support this practice.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document should be specified.</p>	<p>None</p>
<p><u>Current evidence base</u> What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>A recent NCCHTA commissioned systematic review suggests that TURP should remain the standard of care and specifically that green Light Laser was unlikely to be cost-effective in the economic model and thereby arguing against its unrestricted use in the NHS until further evidence of effectiveness and cost-reduction is obtained ^{19,172-174}.</p>
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus</p>	<p>Not applicable</p>

<p>on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>Primary research (RCT). Comparator is TURP. Careful consideration must be given to treatment strategies within the trial design such as incorporating early versus delayed intervention.</p>
<p><u>Feasibility</u> Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>Proposed research can be carried out in a realistic timescale and at an acceptable cost. There are no ethical issues. A potential risk is that KTP laser vaporisation use may diminish without adequate assessment.</p>
<p><u>Other comments</u> Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>NCCHTA would be the obvious funder</p>
<p><u>Importance</u> How important is the question to the overall guideline? The research recommendation should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none"> • High: the research is essential to inform future updates of key recommendations in the guideline • Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates • Low: the research is of interest and will fill existing evidence gaps. 	<p>High</p>

1.5 Male slings

<p><u>PICO question</u> Each research recommendation should be formulated as an answerable question or a set of closely related questions. This should use the <u>PICO framework</u> (patient, intervention, comparison and outcome)</p>	<p>In men with mild to moderate post prostatectomy urinary incontinence (P), what is the clinical or cost effectiveness of a male sling or an implanted adjustable compression device (IC), when assessed by symptom severity, quality of life, changes in measured leakage, and occurrence of adverse events (O).</p> <p>Possible interventions include: Retrobulbar ‘non-compressive’ male sling, adjustable compression sling and implanted adjustable compression devices.</p> <p>Paraurethral injections have been used but are not recommended by the recent WHO International Consultation on Incontinence.</p>
<p><u>Importance to patients or the population.</u> What would be the impact of any new or altered guidance on the population? (for example, acceptability to patients, quality of life, morbidity or disease prevalence, severity of disease or mortality).</p>	<p>This increasingly prevalent group of men have, until recently, had no acceptable treatment option other than insertion of an artificial urinary sphincter but many men consider this treatment to be too invasive and too prone to complication or failure. A number of new interventions have been devised but there is no clarity on which of these offers the best outcomes. This research could lead to clear recommendations and effective treatment for the majority of these men.</p>
<p><u>Relevance to NICE guidance</u> How would the answer to this question change future NICE guidance (that is, generate new knowledge and/or evidence)?</p>	<p>NICE currently cannot give clear guidance on this topic because of an inadequate evidence base.</p>
<p><u>Relevance to the NHS</u> What would be the impact on the NHS and (where relevant) the public sector of any new or altered guidance (for example, financial advantage, effect on staff, impact on strategic planning or service delivery)?</p>	<p>This group of men currently depend on containment alone for control of their incontinence – there are likely to be cost savings from effective incontinence treatment. Insertion of an artificial urinary sphincter, whilst of recognised efficacy, carries a significant cost. Guidance is needed on the most suitable surgical options for this group of men.</p>
<p><u>National priorities</u> Is the question relevant to a national priority area (such as a national service framework or white paper)? The relevant document</p>	<p>There is currently no national service framework for men with LUTS or incontinence.</p>

should be specified.	
<p><u>Current evidence base</u> What is the current evidence base? What are the problems with the current evidence base? (that is, why is further research required?) Reference should be made to the section of the full guideline that describes the current evidence base, including details of trials and systematic reviews. The date on which the final literature search was undertaken should be specified.</p>	<p>There is currently no level 1 evidence for these surgical interventions because they are relatively new and have not been subjected to randomised controlled trials.</p> <p>NICE Interventional Procedures Committee has reported on Male slings (mostly “Invance”) and non circumferential extraurethral compression devices.</p>
<p><u>Equality</u> Does the research recommendation address equality issues? For example, does it focus on groups that need special consideration, or focus on an intervention that is not available for use by people with certain disabilities?</p>	<p>There are no equality issues.</p>
<p><u>Study design</u> It should also specify the most appropriate study design to address the proposed question(s). Primary research or secondary research (for example, systematic reviews) can be recommended.</p>	<p>A randomised controlled trial comparing up to three current interventions; retrobulbar “non compressive” male sling, adjustable compression sling, and implanted adjustable compression device is recommended. However other new devices are being introduced rapidly into the market place with little or no clinical data to underpin marketing.</p>
<p><u>Feasibility</u> Can the proposed research be carried out in a realistic timescale and at an acceptable cost? As part of cost-effectiveness analysis, formal value-of-information methods may also sometimes be used to estimate the value for money of additional research. Are there any ethical or technical issues?</p>	<p>The major issues with this trial would be the centralisation of cases into centres able to offer the surgery and the training of participating surgeons since the procedures proposed are still relatively new. An adequate population of men with this problem already exists precisely because of the absence of any really effective treatment for this group.</p>
<p><u>Other comments</u> Any other important issues should be mentioned, such as potential funders or outcomes of previous attempts to address this issue or methodological problems. However, this is not a research protocol.</p>	<p>In general, manufacturers have been reluctant to fund randomised controlled trials and prefer to sponsor the establishment of surgical registries. Whilst these facilitate the involvement of a greater number of surgeons and cases, the risk of bias is very high. It may be that independent registries are a better way to establish the associated risks of surgery because of the feasibility of including all patients, not just those eligible for inclusion in an RCT.</p>
<p><u>Importance</u> How important is the question to the overall guideline? The research recommendation</p>	<p>High. This is a population of men who have been rendered incontinent by surgery which may or may not cure their cancer. The</p>

<p>should be categorised into one of the following categories of importance:</p> <ul style="list-style-type: none">• High: the research is essential to inform future updates of key recommendations in the guideline• Medium: the research is relevant to the recommendations in the guideline, but the research recommendations are not key to future updates• Low: the research is of interest and will fill existing evidence gaps.	<p>impact on their quality of life is profound and there is currently only one realistic treatment option which many men find unacceptable. It is important that solutions are found for this growing number of men.</p>
---	--

Appendix H – International Prostate Symptom Score (IPSS)

INTERNATIONAL-PROSTATE SYMPTOM SCORE (I-PSS)							
	Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always	
1. Over the past 4 weeks, how often have you had a sensation of not emptying your bladder completely after you finished urinating?	0	1	2	3	4	5	
2. Over the past 4 weeks, how often have you had to urinate again less than two hours after you finished urinating?	0	1	2	3	4	5	
3. Over the past 4 weeks, how often have you found you stopped and started again several times when you urinated?	0	1	2	3	4	5	
4. Over the past 4 weeks, how often have you found it difficult to postpone urination?	0	1	2	3	4	5	
5. Over the past 4 weeks, how often has your urinary stream been weaker than usual?	0	1	2	3	4	5	
6. Over the past 4 weeks, how often have you had to push or strain to begin urination?	0	1	2	3	4	5	
	None	1 time	2 times	3 times	4 times	5 or more times	
7. Over the past 4 weeks, how many times, in general, did you get up to urinate from the time you went to bed at night until the time you got up in the morning?	0	1	2	3	4	5	
Total I-PSS Score S =							
QUALITY OF LIFE DUE TO URINARY SYMPTOMS							
	Delighted	Pleased	Mostly satisfied	Mixed - neither satisfied nor dissatisfied	Mostly dissatisfied	Unhappy	Terrible
1. If you were to spend the rest of your life with your urinary condition just the way it is now, how would you feel about that?	0	1	2	3	4	5	6
Quality of life assessment index L =							

International – Prostate Symptom Score © (I-PSS©) Michael J Barry, 1992. All rights reserved.

I-PSS contact information and permission to use:
 MAPI Research Trust, Lyon, France.
 E-mail: contact@mapl-trust.org
 Website: www.mapl-trust.org

I-PSS 1_ Standard_UK English_Mapi Institute_ID2831

International Prostate Symptom Score – Acute version

INTERNATIONAL-PROSTATE SYMPTOM SCORE							
Question	Not at all	Less than 1 time in 5	Less than half the time	About half the time	More than half the time	Almost always	
1. Over the past week, how often have you had a sensation of not emptying your bladder completely after you finished urinating?	0	1	2	3	4	5	
2. Over the past week, how often have you had to urinate again less than two hours after you finished urinating?	0	1	2	3	4	5	
3. Over the past week, how often have you found you stopped and started again several times when you urinated?	0	1	2	3	4	5	
4. Over the past week, how often have you found it difficult to postpone urination?	0	1	2	3	4	5	
5. Over the past week, how often has your urinary stream been weaker than usual?	0	1	2	3	4	5	
6. Over the past week, how often have you had to push or strain to begin urination?	0	1	2	3	4	5	
	None	1 time	2 times	3 times	4 times	5 or more times	
7. Over the past week, how many times, in general, did you get up to urinate from the time you went to bed at night until the time you got up in the morning?	0	1	2	3	4	5	
QUALITY OF LIFE DUE TO URINARY SYMPTOMS							
Question	Delighted	Pleased	Mostly satisfied	Mixed - neither satisfied nor dissatisfied	Mostly dissatisfied	Unhappy	Terrible
1. If you were to spend the rest of your life with your urinary condition just the way it is now, how would you feel about that?	0	1	2	3	4	5	6

International – Prostate Symptom Score © (I-PSS©) Michael J Barry, 1992. All rights reserved.

I-PSS contact information and permission to use:
 MAPI Research Trust, Lyon, France.
 E-mail: contact@mapr-trust.org
 Website: www.mapr-trust.org

I-PSS1 Acute- UK English –ID3546 15/02/99

Bibliography

1. A comparison of quality of life with patient reported symptoms and objective findings in men with benign prostatic hyperplasia. The Department of Veterans Affairs Cooperative Study of transurethral resection for benign prostatic hyperplasia. *Journal of Urology* 1993, **150**(5 Pt 2):1696-700. (Guideline Ref ID: ANON1993)
2. Hospital Episode Statistics 2006-07 www.hesonline.nhs.uk (Guideline Ref ID: ANON2007)
3. Abbou CC, Payan C, Viens-Bitker C, Richard F, Boccon-Gibod L, Jardin A *et al.* Transrectal and transurethral hyperthermia versus sham treatment in benign prostatic hyperplasia: a double-blind randomized multicentre clinical trial. The French BPH Hyperthermia. *British Journal of Urology* 1995, **76**(5):619-24. (Guideline Ref ID: ABBOU1995)
4. Abdel-Khalek M, El Hammady S, Ibrahim E-H. A 4-year follow-up of a randomized prospective study comparing transurethral electrovaporization of the prostate with neodymium: YAG laser therapy for treating benign prostatic hyperplasia. *BJU International* 2003, **91**(9):801-5. (Guideline Ref ID: ABDELKHALEK2003)
5. Abrams P. Urodynamic effects of doxazosin in men with lower urinary tract symptoms and benign prostatic obstruction. Results from three double-blind placebo-controlled studies. *European Urology* 1997, **32**(1):39-46. (Guideline Ref ID: ABRAMS1997A)
6. Abrams P, Donovan JL, de la Rosette JJ, Schafer W. International Continence Society "Benign Prostatic Hyperplasia" Study: background, aims, and methodology. *Neurourology and Urodynamics* 1997, **16**(2):79-91. (Guideline Ref ID: ABRAMS1997)
7. Abrams P, Schafer W, Tammela TL, Barrett DM, Hedlund H, Rollema HJ *et al.* Improvement of pressure flow parameters with finasteride is greater in men with large prostates. Finasteride Urodynamics Study Group. *Journal of Urology* 1999, **161**(5):1513-7. (Guideline Ref ID: ABRAMS1999)
8. Ackerman SJ, Rein AL, Blute ML, Beusterian K, Sullivan EM, Tanio CP *et al.* Cost effectiveness of microwave thermotherapy in patients with benign prostatic hyperplasia. Part I: methods. *Urology* 2000, **56**(6):972-80. (Guideline Ref ID: ACKERMAN2000)
9. Ahmed M, Bell T, Lawrence WT, Ward JP, Watson GM. Transurethral microwave thermotherapy (Prostatron version 2.5) compared with transurethral resection of the prostate for the treatment of benign prostatic hyperplasia: a randomized, controlled, parallel study. *British Journal of Urology* 1997, **79**(2):181-5. (Guideline Ref ID: AHMED1997)
10. Aho TF, Gilling PJ, Kennett KM, Westenberg AM, Fraundorfer MR, Frampton CM. Holmium laser bladder neck incision versus holmium enucleation of the prostate as outpatient procedures for prostates less than 40 grams: a randomized trial. *Journal of Urology* 2005, **174**(1):210-4. (Guideline Ref ID: AHO2005)

11. Ahyai SA, Lehrich K, Kuntz RM. Holmium laser enucleation versus transurethral resection of the prostate: 3-year follow-up results of a randomized clinical trial. *European Urology* 2007, **52**(5):1456-63. (Guideline Ref ID: AHYAI2007)
12. Albala DM, Fulmer BR, Turk TM, Koleski F, Andriole G, Davis BE et al. Office-based transurethral microwave thermotherapy using the TherMatrix TMx-2000. *Journal of Endourology* 2002, **16**(1):57-61. (Guideline Ref ID: ALBALA2002)
13. Albertsen PC, Pellissier JM, Lowe FC, Girman CJ, Roehrborn CG. Economic analysis of finasteride: a model-based approach using data from the Proscar Long-Term Efficacy and Safety Study. *Clinical Therapeutics* 1999, **21**(6):1006-24. (Guideline Ref ID: ALBERTSEN1999)
14. American Urological Association. (2006) Guideline on the management of benign prostatic hyperplasia (BPH). American Urological Association. (Guideline Ref ID: AUA2006)
15. Andersen JT, Ekman P, Wolf H, Beisland HO, Johansson JE, Kontturi M et al. Can finasteride reverse the progress of benign prostatic hyperplasia? A two-year placebo-controlled study. The Scandinavian BPH Study Group. *Urology* 1995, **46**(5):631-7. (Guideline Ref ID: ANDERSEN1995)
16. Andersen M, Dahlstrand C, Høye K. Double-blind trial of the efficacy and tolerability of doxazosin in the gastrointestinal therapeutic system, doxazosin standard, and placebo in patients with benign prostatic hyperplasia. *European Urology* 2000, **38**(4):400-9. (Guideline Ref ID: ANDERSEN2000)
17. Annemans L, Cleemput I, Lamotte M, McNeill A, Hargreave T. The economic impact of using alfuzosin 10 mg once daily in the management of acute urinary retention in the UK: a 6-month analysis. *BJU International* 2005, **96**(4):566-71. (Guideline Ref ID: ANNEMANS2005)
18. Anson K, Nawrocki J, Buckley J, Fowler C, Kirby R, Lawrence W et al. A multicenter, randomized, prospective study of endoscopic laser ablation versus transurethral resection of the prostate. *Urology* 1995, **46**(3):305-10. (Guideline Ref ID: ANSON1995)
19. Armstrong N, Vale L, Deverill M, Nabi G, McClinton S, N'Dow J et al. Surgical treatments for men with benign prostatic enlargement: cost effectiveness study. *BMJ* 2009, **338**:b1288. (Guideline Ref ID: ARMSTRONG2009)
20. Autorino R, Damiano R, Di LG, Quarto G, Perdon S, D'Armiento M et al. Four-year outcome of a prospective randomised trial comparing bipolar plasmakinetic and monopolar transurethral resection of the prostate. *European Urology* 2009, **55**(4):922-31. (Guideline Ref ID: AUTORINO2009)
21. Baladi JF, Menon D, Otten N. An economic evaluation of finasteride for treatment of benign prostatic hyperplasia. *Pharmacoeconomics* 1996, **9**(5):443-54. (Guideline Ref ID: BALADI1996)
22. Bales GT, Gerber GS, Minor TX, Mhoon DA, McFarland JM, Kim HL et al. Effect of preoperative biofeedback/pelvic floor training on continence in men undergoing radical prostatectomy. *Urology* 2000, **56**(4):627-30. (Guideline Ref ID: BALES2000)
23. Barry MJ, Cherkin DC, Chang Y, Fowler FJ, Jr., Skates S. A randomized trial of a multimedia shared decision-making program for men facing a treatment decision for

- benign prostatic hyperplasia. *Disease Management and Clinical Outcomes* 1997, **1**(1):5-14. (Guideline Ref ID: BARRY1997)
24. Barry MJ, Williford WO, Chang Y, Machi M, Jones KM, Walker-Corkery E *et al.* Benign prostatic hyperplasia specific health status measures in clinical research: how much change in the American Urological Association symptom index and the benign prostatic hyperplasia impact index is perceptible to patients? *Journal of Urology* 1995, **154**(5):1770-4. (Guideline Ref ID: BARRY1995B)
 25. Bautista OM, Kusek JW, Nyberg LM, McConnell JD, Bain RP, Miller G *et al.* Study design of the Medical Therapy of Prostatic Symptoms (MTOPS) trial. *Controlled Clinical Trials* 2003, **24**(2):224-43. (Guideline Ref ID: BAUTISTA2003)
 26. Bdesha AS, Bunce CJ, Snell ME, Witherow RO. A sham controlled trial of transurethral microwave therapy with subsequent treatment of the control group. *Journal of Urology* 1994, **152**(2 Pt 1):453-8. (Guideline Ref ID: BDESHA1994)
 27. Bechara A, Romano S, Casabe A, Haime S, Dedola P, Hernandez C *et al.* Comparative efficacy assessment of tamsulosin vs. tamsulosin plus tadalafil in the treatment of LUTS/BPH. Pilot study. *Journal of Sexual Medicine* 2008, **5**(9):2170-8. (Guideline Ref ID: BECHARA2008)
 28. Beisland HO, Binkowitz B, Brekkan E, Ekman P, Kontturi M, Lehtonen T *et al.* Scandinavian clinical study of finasteride in the treatment of benign prostatic hyperplasia. *European Urology* 1992, **22**(4):271-7. (Guideline Ref ID: BEISLAND1992)
 29. Bent S, Kane C, Shinohara K, Neuhaus J, Hudes ES, Goldberg H *et al.* Saw palmetto for benign prostatic hyperplasia. *New England Journal of Medicine* 2006, **354**(6):557-66. (Guideline Ref ID: BENT2006)
 30. Bhansali M, Patankar S, Dobhada S, Khaladkar S. Management of large (>60 g) prostate gland: PlasmaKinetic Superpulse (bipolar) versus conventional (monopolar) transurethral resection of the prostate. *Journal of Endourology* 2009, **23**(1):141-5. (Guideline Ref ID: BHANSALI2009)
 31. Blute ML, Patterson DE, Segura JW, Tomera KM, Hellerstein DK. Transurethral microwave thermotherapy v sham treatment: double-blind randomized study. *Journal of Endourology* 1996, **10**(6):565-73. (Guideline Ref ID: BLUTE1996)
 32. Bouchier-Hayes DM, Anderson P, Van Appledorn S, Bugeja P, Costello AJ. KTP laser versus transurethral resection: early results of a randomized trial. *Journal of Endourology* 2006, **20**(8):580-5. (Guideline Ref ID: BOUCHIERHAYES2006)
 33. Brawer MK, Adams G, Epstein H. Terazosin in the treatment of benign prostatic hyperplasia. Terazosin Benign Prostatic Hyperplasia Study Group. *Archives of Family Medicine* 1993, **2**(9):929-35. (Guideline Ref ID: BRAWER1993)
 34. Brehmer M, Wiksell H, Kinn A. Sham treatment compared with 30 or 60 min of thermotherapy for benign prostatic hyperplasia: a randomized study. *BJU International* 1999, **84**(3):292-6. (Guideline Ref ID: BREHMER1999)
 35. Brown CT, Yap T, Cromwell DA, Rixon L, Steed L, Mulligan K *et al.* Self management for men with lower urinary tract symptoms: randomised controlled trial. *British Medical Journal* 2007, **334**(7583):25-8. (Guideline Ref ID: BROWN2007)

36. Bruskewitz R, Issa MM, Roehrborn CG, Naslund MJ, Perez-Marrero R, Shumaker BP *et al.* A prospective, randomized 1-year clinical trial comparing transurethral needle ablation to transurethral resection of the prostate for the treatment of symptomatic benign prostatic hyperplasia. *Journal of Urology* 1998, **159**(5):1588-93. (Guideline Ref ID: BRUSKEWITZ1998)
37. Bryan NP, Hastie KJ, Chapple CR. Randomised prospective trial of contact laser prostatectomy (CLAP) versus visual laser coagulation of the prostate (VLAP) for the treatment of benign prostatic hyperplasia. 2-year follow-up. *European Urology* 2000, **38**(3):265-71. (Guideline Ref ID: BRYAN2000)
38. Burgio KL, Goode PS, Urban DA, Umlauf MG, Locher JL, Bueschen A *et al.* Preoperative biofeedback assisted behavioral training to decrease post-prostatectomy incontinence: a randomized, controlled trial. *Journal of Urology* 2006, **175**(1):196-201. (Guideline Ref ID: BURGIO2006)
39. Burgio KL, Stutzman RE, Engel BT. Behavioral training for post-prostatectomy urinary incontinence. *Journal of Urology* 1989, **141**(2):303-6. (Guideline Ref ID: BURGIO1989)
40. Byrnes CA, Morton AS, Liss CL, Lippert MC, Gillenwater JY. Efficacy, tolerability, and effect on health-related quality of life of finasteride versus placebo in men with symptomatic benign prostatic hyperplasia: a community based study. CUSP Investigators. Community based study of Proscar. *Clinical Therapeutics* 1995, **17**(5):956-69. (Guideline Ref ID: BYRNES1995)
41. Cannon A, Carter PG, McConnell AA, Abrams P. Desmopressin in the treatment of nocturnal polyuria in the male. *BJU International* 1999, **84**(1):20-4. (Guideline Ref ID: CANNON1999)
42. Carbin BE, Bauer P, Friskand M, Moyse D. Efficacy of alfuzosine (an alpha 1-adrenoreceptor blocking drug) in benign hyperplasia of the prostate. *Scandinavian Journal of Urology and Nephrology Supplementum* 1991, **138**:73-5. (Guideline Ref ID: CARBIN1991)
43. Carraro JC, Raynaud JP, Koch G, Chisholm GD, Di Silverio F, Teillac P *et al.* Comparison of phytotherapy (Permixon) with finasteride in the treatment of benign prostate hyperplasia: a randomized international study of 1,098 patients. *Prostate* 1996, **29**(4):231-40. (Guideline Ref ID: CARRARO1996)
44. Carter A, Sells H, Speakman M, Ewings P, MacDonagh R, O'Boyle P. A prospective randomized controlled trial of hybrid laser treatment or transurethral resection of the prostate, with a 1-year follow-up. *BJU International* 1999, **83**(3):254-9. (Guideline Ref ID: CARTER1999)
45. Carter A, Sells H, Speakman M, Ewings P, O'Boyle P, MacDonagh R. Quality of life changes following KTP/Nd:YAG laser treatment of the prostate and TURP. *European Urology* 1999, **36**(2):92-8. (Guideline Ref ID: CARTER1999A)
46. Carter HB, Landis P, Wright EJ, Parsons JK, Metter EJ. Can a baseline prostate specific antigen level identify men who will have lower urinary tract symptoms later in life? *Journal of Urology* 2005, **173**(6):2040-3. (Guideline Ref ID: CARTER2005)
47. Cetinkaya M, Ulusoy E, Adsan O, Saglam H, Ozturk B, Basay S. Comparative early results of transurethral electroresection and transurethral electrovaporization in benign prostatic

- hyperplasia. *British Journal of Urology* 1996, **78**(6):901-3. (Guideline Ref ID: CETINKAYA1996)
48. Chacko KN, Donovan JL, Abrams P, Peters TJ, Brookes ST, Thorpe AC *et al.* Transurethral prostatic resection or laser therapy for men with acute urinary retention: the Clasp randomized trial. *Journal of Urology* 2001, **166**(1):166-70. (Guideline Ref ID: CHACKO2001)
 49. Chapple CR, Al Shukri SH, Gattegno B, Holmes S, Martinez-Sagarra JM, Scarpa RM *et al.* Tamsulosin oral controlled absorption system (OCAS) in patients with lower urinary tract symptoms suggestive of benign prostatic hyperplasia (LUTS/BPH): Efficacy and tolerability in a placebo and active comparator controlled phase 3a study. *European Urology, Supplements* 2005, **4**(2):33-44. (Guideline Ref ID: CHAPPLE2005)
 50. Chapple CR, Carter P, Christmas TJ, Kirby RS, Bryan J, Milroy EJ *et al.* A three month double-blind study of doxazosin as treatment for benign prostatic bladder outlet obstruction. *British Journal of Urology* 1994, **74**(1):50-6. (Guideline Ref ID: CHAPPLE1994)
 51. Chapple CR, Wyndaele JJ, Nordling J, Boeminghaus F, Ypma AF, Abrams P. Tamsulosin, the first prostate-selective alpha 1A-adrenoceptor antagonist. A meta-analysis of two randomized, placebo-controlled, multicentre studies in patients with benign prostatic obstruction (symptomatic BPH). European Tamsulosin Study Group. *European Urology* 1996, **29**(2):155-67. (Guideline Ref ID: CHAPPLE1996)
 52. Christensen MM, Aagaard J, Madsen PO. Transurethral resection versus transurethral incision of the prostate. A prospective randomized study. *Urologic Clinics of North America* 1990, **17**(3):621-30. (Guideline Ref ID: CHRISTENSEN1990)
 53. Christensen MM, Bendix HJ, Rasmussen PC, Jacobsen F, Nielsen J, Norgaard JP *et al.* Doxazosin treatment in patients with prostatic obstruction. A double-blind placebo-controlled study. *Scandinavian Journal of Urology and Nephrology* 1993, **27**(1):39-44. (Guideline Ref ID: CHRISTENSEN1993)
 54. Cimentepe E, Unsal A, Saglam R. Randomized clinical trial comparing transurethral needle ablation with transurethral resection of the prostate for the treatment of benign prostatic hyperplasia: results at 18 months. *Journal of Endourology* 2003, **17**(2):103-7. (Guideline Ref ID: CIMENTEPE2003)
 55. Cockrum PC, Finder SF, Ries AJ, Potyk RP. A pharmacoeconomic analysis of patients with symptoms of benign prostatic hyperplasia. *Pharmacoeconomics* 1997, **11**(6):550-65. (Guideline Ref ID: COCKRUM1997)
 56. Cowles RS, III, Kabalin JN, Childs S, Lepor H, Dixon C, Stein B *et al.* A prospective randomized comparison of transurethral resection to visual laser ablation of the prostate for the treatment of benign prostatic hyperplasia. *Urology* 1995, **46**(2):155-60. (Guideline Ref ID: COWLES1995)
 57. Crawford ED, Wilson SS, McConnell JD, Slawin KM, Lieber MC, Smith JA *et al.* Baseline factors as predictors of clinical progression of benign prostatic hyperplasia in men treated with placebo. *Journal of Urology* 2006, **175**(4):1422-6. (Guideline Ref ID: CRAWFORD2006)

58. Currie CJ, McEwan P, Poole CD, Odeyemi IA, Datta SN, Morgan CL. The impact of the overactive bladder on health-related utility and quality of life. *BJU International* 2006, **97**(6):1267-72. (Guideline Ref ID: CURRIE2006)
59. Curtis L. (2008) Unit costs of health and social care 2008. Personal Social Services Research Unit. (Guideline Ref ID: CURTIS2008)
60. d'Ancona FC, Francisca EA, Witjes WP, Welling L, Debruyne FM, de la Rosette JJ. High energy thermotherapy versus transurethral resection in the treatment of benign prostatic hyperplasia: results of a prospective randomized study with 1 year of followup. *Journal of Urology* 1997, **158**(1):120-5. (Guideline Ref ID: DANCONA1997)
61. d'Ancona FC, Francisca EA, Witjes WP, Welling L, Debruyne FM, de la Rosette JJ. Transurethral resection of the prostate vs high-energy thermotherapy of the prostate in patients with benign prostatic hyperplasia: long-term results. *British Journal of Urology* 1998, **81**(2):259-64. (Guideline Ref ID: DANCONA1998)
62. Dahlstrand C, Geirsson G, Fall M, Pettersson S. Transurethral microwave thermotherapy versus transurethral resection for benign prostatic hyperplasia: preliminary results of a randomized study. *European Urology* 1993, **23**(2):292-8. (Guideline Ref ID: DAHLSTRAND1993)
63. Dahlstrand C, Walden M, Geirsson G, Pettersson S. Transurethral microwave thermotherapy versus transurethral resection for symptomatic benign prostatic obstruction: a prospective randomized study with a 2-year follow-up. *British Journal of Urology* 1995, **76**(5):614-8. (Guideline Ref ID: DAHLSTRAND1995)
64. de la Rosette JJ, De Wildt MJ, Alivizatos G, Froeling FM, Debruyne FM. Transurethral microwave thermotherapy (TUMT) in benign prostatic hyperplasia: placebo versus TUMT. *Urology* 1994, **44**(1):58-63. (Guideline Ref ID: DELAROSSETTE1994)
65. de la Rosette JJ, Floratos DL, Severens JL, Kiemeny LA, Debruyne FM, Pilar LM. Transurethral resection vs microwave thermotherapy of the prostate: a cost-consequences analysis. *BJU International* 2003, **92**(7):713-8. (Guideline Ref ID: DELAROSSETTE2003B)
66. de Sio M, Autorino R, Quarto G, Damiano R, Perdona S, di Lorenzo G et al. Gyrus bipolar versus standard monopolar transurethral resection of the prostate: a randomized prospective trial. *Urology* 2006, **67**(1):69-72. (Guideline Ref ID: DESIO2006)
67. De Wildt MJ, Hubregtse M, Ogden C, Carter SS, Debruyne FM, de la Rosette JJ. A 12-month study of the placebo effect in transurethral microwave thermotherapy. *British Journal of Urology* 1996, **77**(2):221-7. (Guideline Ref ID: DEWILDT1996)
68. Debruyne F, Koch G, Boyle P, Da Silva FC, Gillenwater JG, Hamdy FC et al. Comparison of a phytotherapeutic agent (Permixon) with an alpha-blocker (Tamsulosin) in the treatment of benign prostatic hyperplasia: a 1-year randomized international study. *European Urology* 2002, **41**(5):497-506. (Guideline Ref ID: DEBRUYNE2002)
69. Debruyne FM, Jardin A, Colloi D, Resel L, Witjes WP, Delauche-Cavallier MC et al. Sustained-release alfuzosin, finasteride and the combination of both in the treatment of benign prostatic hyperplasia. European ALFIN Study Group. *European Urology* 1998, **34**(3):169-75. (Guideline Ref ID: DEBRUYNE1998)

70. Dedhia RC, Calhoun E, McVary KT. Impact of phytotherapy on utility scores for 5 benign prostatic hyperplasia/lower urinary tract symptoms health states. *Journal of Urology* 2008, **179**(1):220-5. (Guideline Ref ID: DEDHIA2008)
71. Disantostefano RL, Biddle AK, Lavelle JP. An evaluation of the economic costs and patient-related consequences of treatments for benign prostatic hyperplasia. *BJU International* 2006, **97**(5):1007-16. (Guideline Ref ID: DISANTOSTEFANO2006)
72. Djavan B, Fong YK, Chaudry A, Reissigl A, Anagnostou T, Bagheri F et al. Progression delay in men with mild symptoms of bladder outlet obstruction: a comparative study of phytotherapy and watchful waiting. *World Journal of Urology* 2005, **23**(4):253-6. (Guideline Ref ID: DJAVAN2005)
73. Djavan B, Milani S, Davies J, Bolodeoku J. The impact of tamsulosin oral controlled absorption system (OCAS) on nocturia and the quality of sleep: Preliminary results of a pilot study. *European Urology, Supplements* 2005, **4**(2):61-8. (Guideline Ref ID: DJAVAN2005D)
74. Donovan JL, Peters TJ, Neal DE, Brookes ST, Gujral S, Chacko KN et al. A randomized trial comparing transurethral resection of the prostate, laser therapy and conservative treatment of men with symptoms associated with benign prostatic enlargement: The CLasP study. *Journal of Urology* 2000, **164**(1):65-70. (Guideline Ref ID: DONOVAN2000)
75. Dorflinger T, Jensen FS, Krarup T, Walter S. Transurethral prostatectomy compared with incision of the prostate in the treatment of prostatism caused by small benign prostate glands. *Scandinavian Journal of Urology and Nephrology* 1992, **26**(4):333-8. (Guideline Ref ID: DORFLINGER1992)
76. Dunsmuir WD, McFarlane JP, Tan A, Dowling C, Downie J, Kourambas J et al. Gyrus bipolar electrovaporization vs transurethral resection of the prostate: a randomized prospective single-blind trial with 1 y follow-up. *Prostate Cancer & Prostatic Diseases* 2003, **6**(2):182-6. (Guideline Ref ID: DUNSMUIR2003)
77. Ekengren J, Haendler L, Hahn RG. Clinical outcome 1 year after transurethral vaporization and resection of the prostate. *Urology* 2000, **55**(2):231-5. (Guideline Ref ID: EKENGREN2000)
78. Ekman P. Maximum efficacy of finasteride is obtained within 6 months and maintained over 6 years. Follow-up of the Scandinavian Open-Extension Study. The Scandinavian Finasteride Study Group. *European Urology* 1998, **33**(3):312-7. (Guideline Ref ID: EKMAN1998)
79. Elhilali MM, Ramsey EW, Barkin J, Casey RW, Boake RC, Beland G et al. A multicenter, randomized, double-blind, placebo-controlled study to evaluate the safety and efficacy of terazosin in the treatment of benign prostatic hyperplasia. *Urology* 1996, **47**(3):335-42. (Guideline Ref ID: ELHILALI1996)
80. Elzayat EA, Al-Mandil MS, Khalaf I, Elhilali MM. Holmium laser ablation of the prostate versus photoselective vaporization of prostate 60 cc or less: short-term results of a prospective randomized trial. *Journal of Urology* 2009, **182**(1):133-8. (Guideline Ref ID: ELZAYAT2009)

81. Emberton M, Neal DE, Black N, Harrison M, Fordham M, McBrien MP *et al.* The National Prostatectomy Audit: the clinical management of patients during hospital admission. *British Journal of Urology* 1995, **75**(3):301-16. (Guideline Ref ID: EMBERTON1995)
82. Engelmann U, Walther C, Bondarenko B, Funk P, Schlafke S. Efficacy and safety of a combination of sabal and urtica extract in lower urinary tract symptoms. A randomized, double-blind study versus tamsulosin. *Arzneimittel-Forschung* 2006, **56**(3):222-9. (Guideline Ref ID: ENGELMANN2006)
83. Erdagi U, Akman RY, Sargin SY, Yazicioglu A. Transurethral electrovaporization of the prostate versus transurethral resection of the prostate: a prospective randomized study. *Archivio Italiano di Urologia, Andrologia* 1999, **71**(3):125-30. (Guideline Ref ID: ERDAGI1999)
84. Erturhan S, Erbagci A, Seckiner I, Yagci F, Ustun A. Plasmakinetic resection of the prostate versus standard transurethral resection of the prostate: a prospective randomized trial with 1-year follow-up. *Prostate Cancer & Prostatic Diseases* 2007, **10**(1):97-100. (Guideline Ref ID: ERTURHAN2007)
85. Ezz el Din K, Koch WF, De Wildt MJ, Debruyne FM, de la Rosette JJ. The predictive value of microscopic haematuria in patients with lower urinary tract symptoms and benign prostatic hyperplasia. *European Urology* 1996, **30**(4):409-13. (Guideline Ref ID: EZZ1996)
86. Fader M, Cottenden A, Getliffe K, Gage H, Clarke-O'Neill S, Jamieson K *et al.* Absorbent products for urinary/faecal incontinence: a comparative evaluation of key product designs. *Health Technology Assessment* 2008, **12**(29):1-208. (Guideline Ref ID: FADER2008)
87. Fader M, Macaulay M, Pettersson L, Brooks R, Cottenden A. A multi-centre evaluation of absorbent products for men with light urinary incontinence. *Neurourology and Urodynamics* 2006, **25**(7):689-95. (Guideline Ref ID: FADER2006)
88. Falahatkar S, Mokhtari G, Pourreza F, Asgari SA, Kamran AN. Celecoxib for treatment of nocturia caused by benign prostatic hyperplasia: a prospective, randomized, double-blind, placebo-controlled study. *Urology* 2008, **72**(4):813-6. (Guideline Ref ID: FALAHATKAR2008)
89. Fawzy A, Braun K, Lewis GP, Gaffney M, Ice K, Dias N. Doxazosin in the treatment of benign prostatic hyperplasia in normotensive patients: a multicenter study. *Journal of Urology* 1995, **154**(1):105-9. (Guideline Ref ID: FAWZY1995)
90. Fehrling M, Fall M, Peeker R. Maximal functional electrical stimulation as a single treatment: is it cost-effective? *Scandinavian Journal of Urology and Nephrology* 2007, **41**(2):132-7. (Guideline Ref ID: FEHRLING2007)
91. Filocamo MT, Li M, V, Del Popolo G, Cecconi F, Marzocco M, Tosto A *et al.* Effectiveness of early pelvic floor rehabilitation treatment for post-prostatectomy incontinence. *European Urology* 2005, **48**(5):734-8. (Guideline Ref ID: FILOCAMO2005)
92. Finasteride Study Group. Finasteride (MK-906) in the treatment of benign prostatic hyperplasia. The Finasteride Study Group. *Prostate* 1993, **22**(4):291-9. (Guideline Ref ID: ANON1993A)
93. Floratos DL, Sonke GS, Rapidou CA, Alivizatos GJ, Deliveliotis C, Constantinides CA *et al.* Biofeedback vs verbal feedback as learning tools for pelvic muscle exercises in the early

- management of urinary incontinence after radical prostatectomy. *BJU International* 2002, **89**(7):714-9. (Guideline Ref ID: FLORATOS2002)
94. Fowler C, McAllister W, Plail R, Karim O, Yang Q. Randomised evaluation of alternative electrosurgical modalities to treat bladder outflow obstruction in men with benign prostatic hyperplasia. *Health Technology Assessment* 2005, **9**(4):iii-30. (Guideline Ref ID: FOWLER2005)
95. Francisca EA, d'Ancona FC, Hendriks JC, Kiemeny LA, Debruyne FM, de la Rosette JJ. Quality of life assessment in patients treated with lower energy thermotherapy (Prostasoft 2.0): results of a randomized transurethral microwave thermotherapy versus sham study. *Journal of Urology* 1997, **158**(5):1839-44. (Guideline Ref ID: FRANCISCA1997)
96. Franke JJ, Gilbert WB, Grier J, Koch MO, Shyr Y, Smith JA. Early post-prostatectomy pelvic floor biofeedback. *Journal of Urology* 2000, **163**(1):191-3. (Guideline Ref ID: FRANKE2000)
97. Fraundorfer MR, Gilling PJ, Kennett KM, Dunton NG. Holmium laser resection of the prostate is more cost effective than transurethral resection of the prostate: results of a randomized prospective study. *Urology* 2001, **57**(3):454-8. (Guideline Ref ID: FRAUNDORFER2001)
98. Fung BT, Li SK, Yu CF, Lau BE, Hou SS. Prospective randomized controlled trial comparing plasmakinetic vaporessection and conventional transurethral resection of the prostate. *Asian Journal of Surgery* 2005, **28**(1):24-8. (Guideline Ref ID: FUNG2005)
99. Gallucci M, Puppo P, Perachino M, Fortunato P, Muto G, Breda G et al. Transurethral electrovaporization of the prostate vs. transurethral resection. Results of a multicentric, randomized clinical study on 150 patients. *European Urology* 1998, **33**(4):359-64. (Guideline Ref ID: GALLUCCI1998)
100. Ghalayini IF, Al Ghazo MA, Pickard RS. A prospective randomized trial comparing transurethral prostatic resection and clean intermittent self-catheterization in men with chronic urinary retention. *BJU International* 2005, **96**(1):93-7. (Guideline Ref ID: GHALAYINI2005)
101. Gillenwater JY, Conn RL, Chrysant SG, Roy J, Gaffney M, Ice K et al. Doxazosin for the treatment of benign prostatic hyperplasia in patients with mild to moderate essential hypertension: a double-blind, placebo-controlled, dose-response multicenter study. *Journal of Urology* 1995, **154**(1):110-5. (Guideline Ref ID: GILLENWATER1995)
102. Gilling PJ, Cass CB, Malcolm A, Cresswell M, Fraundorfer MR, Kabalin JN. Holmium laser resection of the prostate versus neodymium:yttrium-aluminum-garnet visual laser ablation of the prostate: a randomized prospective comparison of two techniques for laser prostatectomy. *Urology* 1998, **51**(4):573-7. (Guideline Ref ID: GILLING1998)
103. Gilling PJ, Kennett KM, Fraundorfer MR. Holmium laser resection v transurethral resection of the prostate: results of a randomized trial with 2 years of follow-up. *Journal of Endourology* 2000, **14**(9):757-60. (Guideline Ref ID: GILLING2000)
104. Gilling PJ, Mackey M, Cresswell M, Kennett K, Kabalin JN, Fraundorfer MR. Holmium laser versus transurethral resection of the prostate: a randomized prospective trial with 1-year followup. *Journal of Urology* 1999, **162**(5):1640-4. (Guideline Ref ID: GILLING1999)

105. Gormley GJ, Stoner E, Bruskewitz RC, Imperato-McGinley J, Walsh PC, McConnell JD *et al.* The effect of finasteride in men with benign prostatic hyperplasia. The Finasteride Study Group. *New England Journal of Medicine* 1992, **327**(17):1185-91. (Guideline Ref ID: GORMLEY1992)
106. Gotoh M, Okamura K, Hattori R, Nishiyama N, Kobayashi H, Tanaka K *et al.* A randomized comparative study of the Bandloop versus the standard loop for transurethral resection of the prostate. *Journal of Urology* 1999, **162**(5):1645-7. (Guideline Ref ID: GOTOH1999)
107. Gujral S, Abrams P, Donovan JL, Neal DE, Brookes ST, Chacko KN *et al.* A prospective randomized trial comparing transurethral resection of the prostate and laser therapy in men with chronic urinary retention: The CLasP study. *Journal of Urology* 2000, **164**(1):59-64. (Guideline Ref ID: GUJRAL2000)
108. Gupta N, Sivaramakrishna, Kumar R, Dogra PN, Seth A. Comparison of standard transurethral resection, transurethral vapour resection and holmium laser enucleation of the prostate for managing benign prostatic hyperplasia of >40 g. *BJU International* 2006, **97**(1):85-9. (Guideline Ref ID: GUPTA2006)
109. Hammadeh MY, Fowles GA, Singh M, Philp T. Transurethral electrovaporization of the prostate--a possible alternative to transurethral resection: a one-year follow-up of a prospective randomized trial. *British Journal of Urology* 1998, **81**(5):721-5. (Guideline Ref ID: HAMMADEH1998B)
110. Hammadeh MY, Madaan S, Hines J, Philp T. 5-year outcome of a prospective randomized trial to compare transurethral electrovaporization of the prostate and standard transurethral resection. *Urology* 2003, **61**(6):1166-71. (Guideline Ref ID: HAMMADEH2003)
111. Hammadeh MY, Madaan S, Singh M, Philp T. A 3-year follow-up of a prospective randomized trial comparing transurethral electrovaporization of the prostate with standard transurethral prostatectomy. *BJU International* 2000, **86**(6):648-51. (Guideline Ref ID: HAMMADEH2000)
112. Hansen BJ, Nordling J, Mensink HJ, Walter S, Meyhoff HH. Alfuzosin in the treatment of benign prostatic hyperplasia: effects on symptom scores, urinary flow rates and residual volume. A multicentre, double-blind, placebo-controlled trial. ALFECH Study Group. *Scandinavian Journal of Urology and Nephrology Supplementum* 1994, **157**:169-76. (Guideline Ref ID: HANSEN1994)
113. Helke C, Manseck A, Hakenberg OW, Wirth MP. Is transurethral vaporesction of the prostate better than standard transurethral resection? *European Urology* 2001, **39**(5):551-7. (Guideline Ref ID: HELKE2001)
114. Hellström P, Lukkarinen O, Kontturi M. Bladder neck incision or transurethral electroresection for the treatment of urinary obstruction caused by a small benign prostate? A randomized urodynamic study. *Scandinavian Journal of Urology and Nephrology* 1986, **20**(3):187-92. (Guideline Ref ID: HELLSTRÖM1986)
115. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.0.0 [updated February 2008] www.cochrane-handbook.org (Guideline Ref ID: HIGGINS2008)

116. Hill B, Belville W, Bruskewitz R, Issa M, Perez-Marrero R, Roehrborn C *et al.* Transurethral needle ablation versus transurethral resection of the prostate for the treatment of symptomatic benign prostatic hyperplasia: 5-year results of a prospective, randomized, multicenter clinical trial. *Journal of Urology* 2004, **171**(6 Pt 1):2336-40. (Guideline Ref ID: HILL2004)
117. Hillman AL, Schwartz JS, Willian MK, Peskin E, Roehrborn CG, Oesterling JE *et al.* The cost-effectiveness of terazosin and placebo in the treatment of moderate to severe benign prostatic hyperplasia. *Urology* 1996, **47**(2):169-78. (Guideline Ref ID: HILLMAN1996)
118. Hindley RG, Mostafid AH, Brierly RD, Harrison NW, Thomas PJ, Fletcher MS. The 2-year symptomatic and urodynamic results of a prospective randomized trial of interstitial radiofrequency therapy vs transurethral resection of the prostate. *BJU International* 2001, **88**(3):217-20. (Guideline Ref ID: HINDLEY2001)
119. Hizli F, Uygur MC. A prospective study of the efficacy of *Serenoa repens*, tamsulosin, and *Serenoa repens* plus tamsulosin treatment for patients with benign prostate hyperplasia. *International Urology and Nephrology* 2007, **39**(3):879-86. (Guideline Ref ID: HIZLI2007)
120. Ho HS, Yip SK, Lim KB, Fook S, Foo KT, Cheng CW. A prospective randomized study comparing monopolar and bipolar transurethral resection of prostate using transurethral resection in saline (TURIS) system. *European Urology* 2007, **52**(2):517-22. (Guideline Ref ID: HO2007)
121. Hon NH, Brathwaite D, Hussain Z, Ghiblawi S, Brace H, Hayne D *et al.* A prospective, randomized trial comparing conventional transurethral prostate resection with PlasmaKinetic vaporization of the prostate: physiological changes, early complications and long-term followup. *Journal of Urology* 2006, **176**(1):205-9. (Guideline Ref ID: HON2006)
122. Horasanli K, Silay MS, Altay B, Tanriverdi O, Sarica K, Miroglu C. Photoselective potassium titanyl phosphate (KTP) laser vaporization versus transurethral resection of the prostate for prostates larger than 70 mL: a short-term prospective randomized trial. *Urology* 2008, **71**(2):247-51. (Guideline Ref ID: HORASANLI2008)
123. Hunter KF, Moore KN, Cody DJ, Glazener CM. Conservative management for postprostatectomy urinary incontinence. *Cochrane Database of Systematic Reviews* 2007, **Issue 2**:CD001843. (Guideline Ref ID: HUNTER2007)
124. Iori F, Franco G, Leonardo C, Laurenti C, Tubaro A, Amico F *et al.* Bipolar transurethral resection of prostate: clinical and urodynamic evaluation. *Urology* 2008, **71**(2):252-5. (Guideline Ref ID: IORI2008)
125. Jahnsen S, Dalen M, Gustavsson G, Pedersen J. Transurethral incision versus resection of the prostate for small to medium benign prostatic hyperplasia. *British Journal of Urology* 1998, **81**(2):276-81. (Guideline Ref ID: JAHNSON1998)
126. Jakobsson L. Indwelling catheter treatment and health-related quality of life in men with prostate cancer in comparison with men with benign prostatic hyperplasia. *Scandinavian Journal of Caring Sciences* 2002, **16**(3):264-71. (Guideline Ref ID: JAKOBSSON2002)
127. Johansen TE, Istad JA. Long-term cost analysis of treatment options for benign prostatic hyperplasia in Norway. *Scandinavian Journal of Urology and Nephrology* 2007, **41**(2):124-31. (Guideline Ref ID: JOHANSEN2007)

128. Johnson N, Kirby R. Treatments for benign prostatic hyperplasia: an analysis of their clinical and economic impact in the United Kingdom and Italy. *Journal of drug assessment* 1999, **2**(3):371-86. (Guideline Ref ID: JOHNSON1999)
129. Johnson TMJ, Busby-Whitehead J, Ashford-Works C, Clarke MK, Fowler L, Williams ME. Promoting help-seeking behavior for urinary incontinence. *Journal of Applied Gerontology* 1998, **17**(4):419-41. (Guideline Ref ID: JOHNSON1998)
130. Kadow C, Feneley RC, Abrams PH. Prostatectomy or conservative management in the treatment of benign prostatic hypertrophy? *British Journal of Urology* 1988, **61**(5):432-4. (Guideline Ref ID: KADOW1988)
131. Kalsi V, Popat RB, Apostolidis A, Kavia R, Odeyemi IA, Dakin HA et al. Cost-consequence analysis evaluating the use of botulinum neurotoxin-A in patients with detrusor overactivity based on clinical outcomes observed at a single UK centre. *European Urology* 2006, **49**(3):519-27. (Guideline Ref ID: KALS12006)
132. Kaplan SA, Gonzalez RR, Te AE. Combination of alfuzosin and sildenafil is superior to monotherapy in treating lower urinary tract symptoms and erectile dysfunction. *European Urology* 2007, **51**(6):1717-23. (Guideline Ref ID: KAPLAN2007)
133. Kaplan SA, Laor E, Fatal M, Te AE. Transurethral resection of the prostate versus transurethral electrovaporization of the prostate: a blinded, prospective comparative study with 1-year followup. *Journal of Urology* 1998, **159**(2):454-8. (Guideline Ref ID: KAPLAN1998)
134. Kaplan SA, Roehrborn CG, Chancellor M, Carlsson M, Bavendam T, Guan Z. Extended-release tolterodine with or without tamsulosin in men with lower urinary tract symptoms and overactive bladder: effects on urinary symptoms assessed by the International Prostate Symptom Score. *BJU International* 2008, **102**(9):1133-9. (Guideline Ref ID: KAPLAN2008)
135. Kaplan SA, Roehrborn CG, McConnell JD, Meehan AG, Suryanawansi S, Lee JY et al. Long-term treatment with finasteride results in a clinically significant reduction in total prostate volume compared to placebo over the full range of baseline prostate sizes in men enrolled in the MTOPS trial. *Journal of Urology* 2008, **180**(3):1030-2. (Guideline Ref ID: KAPLAN2008B)
136. Kaplan SA, Roehrborn CG, Rovner ES, Carlsson M, Bavendam T, Guan Z. Tolterodine and tamsulosin for treatment of men with lower urinary tract symptoms and overactive bladder: a randomized controlled trial. *JAMA* 2006, **296**(19):2319-28. (Guideline Ref ID: KAPLAN2006)
137. Karaman MI, Kaya C, Ozturk M, Gurdal M, Kirecci S, Pirincci N. Comparison of transurethral vaporization using PlasmaKinetic energy and transurethral resection of prostate: 1-year follow-up. *Journal of Endourology* 2005, **19**(6):734-7. (Guideline Ref ID: KARAMAN2005)
138. Kawabe K, Ueno A, Takimoto Y, Aso Y, Kato H. Use of an alpha 1-blocker, YM617, in the treatment of benign prostatic hypertrophy. YM617 Clinical Study Group. *Journal of Urology* 1990, **144**(4):908-11. (Guideline Ref ID: KAWABE1990)
139. Kaya C, Ilktac A, Gokmen E, Ozturk M, Karaman IM. The long-term results of transurethral vaporization of the prostate using plasmakinetic energy. *BJU International* 2007, **99**(4):845-8. (Guideline Ref ID: KAYA2007)

140. Keoghane SR, Cranston DW, Lawrence KC, Doll HA, Fellows GJ, Smith JC. The Oxford Laser Prostate Trial: a double-blind randomized controlled trial of contact vaporization of the prostate against transurethral resection; preliminary results. *British Journal of Urology* 1996, **77**(3):382-5. (Guideline Ref ID: KEOGHANE1996A)
141. Keoghane SR, Doll HA, Lawrence KC, Jenkinson CP, Cranston DW. The Oxford Laser Prostate Trial: sexual function data from a randomized controlled clinical trial of contact laser prostatectomy. *European Urology* 1996, **30**(4):424-8. (Guideline Ref ID: KEOGHANE1996)
142. Keoghane SR, Lawrence KC, Gray AM, Doll HA, Hancock AM, Turner K *et al.* A double-blind randomized controlled trial and economic evaluation of transurethral resection vs contact laser vaporization for benign prostatic enlargement: a 3-year follow-up. *BJU International* 2000, **85**(1):74-8. (Guideline Ref ID: KEOGHANE2000)
143. Keoghane SR, Lawrence KC, Jenkinson CP, Doll HA, Chappel DB, Cranston DW. The Oxford Laser Prostate Trial: sensitivity to change of three measures of outcome. *Urology* 1996, **47**(1):43-7. (Guideline Ref ID: KEOGHANE1996B)
144. Keoghane SR, Sullivan ME, Doll HA, Kourambas J, Cranston DW. Five-year data from the Oxford Laser Prostatectomy Trial. *BJU International* 2000, **86**(3):227-8. (Guideline Ref ID: KEOGHANE2000A)
145. Kim JY, Moon KH, Yoon CJ, Park TC. Bipolar transurethral resection of the prostate: a comparative study with monopolar transurethral resection. *Korean Journal of Urology* 2006, **47**(5):493-7. (Guideline Ref ID: KIM2006A)
146. Kim TS, Choi S, Rhew HY, Ahn JH, Jang JH, Cho MH. Comparative study on the treatment outcome and safety of TURP, ILC, TUNA and TEAP for patients with benign prostatic hyperplasia. *Korean Journal of Urology* 2006, **47**(1):13-9. (Guideline Ref ID: KIM2006)
147. Kirby RS, Roehrborn C, Boyle P, Bartsch G, Jardin A, Cary MM *et al.* Efficacy and tolerability of doxazosin and finasteride, alone or in combination, in treatment of symptomatic benign prostatic hyperplasia: the Prospective European Doxazosin and Combination Therapy (PREDICT) trial. *Urology* 2003, **61**(1):19-26. (Guideline Ref ID: KIRBY2003)
148. Kobelt G, Spangberg A, Mattiasson A. The cost of feedback microwave thermotherapy compared with transurethral resection of the prostate for treating benign prostatic hyperplasia. *BJU International* 2004, **93**(4):543-8. (Guideline Ref ID: KOBELT2004)
149. Kok ET, McDonnell J, Stolk EA, Stoevelaar HJ, Busschbach JJ. The valuation of the International Prostate Symptom Score (IPSS) for use in economic evaluations. *European Urology* 2002, **42**(5):491-7. (Guideline Ref ID: KOK2002)
150. Kuntz RM, Lehrich K. Transurethral holmium laser enucleation versus transvesical open enucleation for prostate adenoma greater than 100 gm.: a randomized prospective trial of 120 patients. *Journal of Urology* 2002, **168**(4 Pt 1):1465-9. (Guideline Ref ID: KUNTZ2002)
151. Kuntz RM, Lehrich K, Ahyai S. Transurethral holmium laser enucleation of the prostate compared with transvesical open prostatectomy: 18-month follow-up of a randomized trial. *Journal of Endourology* 2004, **18**(2):189-91. (Guideline Ref ID: KUNTZ2004A)

152. Kuntz RM, Lehrich K, Ahyai SA. Holmium laser enucleation of the prostate versus open prostatectomy for prostates greater than 100 grams: 5-year follow-up results of a randomised clinical trial. *European Urology* 2008, **53**(1):160-6. (Guideline Ref ID: KUNTZ2008)
153. Kupeli B, Yalcinkaya F, Topaloglu H, Karabacak O, Gunlusoy B, Unal S. Efficacy of transurethral electrovaporization of the prostate with respect to standard transurethral resection. *Journal of Endourology* 1998, **12**(6):591-4. (Guideline Ref ID: KUPELI1998A)
154. Kupeli S, Baltaci S, Soygur T, Aytac S, Yilmaz E, Budak M. A prospective randomized study of transurethral resection of the prostate and transurethral vaporization of the prostate as a therapeutic alternative in the management of men with BPH. *European Urology* 1998, **34**(1):15-8. (Guideline Ref ID: KUPELI1998)
155. Kupeli S, Yilmaz E, Soygur T, Budak M. Randomized study of transurethral resection of the prostate and combined transurethral resection and vaporization of the prostate as a therapeutic alternative in men with benign prostatic hyperplasia. *Journal of Endourology* 2001, **15**(3):317-21. (Guideline Ref ID: KUPELI2001)
156. Kursh ED, Concepcion R, Chan S, Hudson P, Ratner M, Eyre R. Interstitial laser coagulation versus transurethral prostate resection for treating benign prostatic obstruction: a randomized trial with 2-year follow-up. *Urology* 2003, **61**(3):573-8. (Guideline Ref ID: KURSH2003)
157. Laguna MP, Kiemeny LA, Debruyne FM, de la Rosette JJ. Baseline prostatic specific antigen does not predict the outcome of high energy transurethral microwave thermotherapy. *Journal of Urology* 2002, **167**(4):1727-30. (Guideline Ref ID: LAGUNA2002)
158. Larsen EH, Dørflinger T, Gasser TC, Gravensen PH, Bruskewitz RC. Transurethral incision versus transurethral resection of the prostate for the treatment of benign prostatic hypertrophy. A preliminary report. *Scandinavian Journal of Urology and Nephrology Supplementum* 1987, **104**:83-6. (Guideline Ref ID: LARSEN1987)
159. Larson TR, Blute ML, Bruskewitz RC, Mayer RD, Ugarte RR, Utz WJ. A high-efficiency microwave thermoablation system for the treatment of benign prostatic hyperplasia: results of a randomized, sham-controlled, prospective, double-blind, multicenter clinical trial. *Urology* 1998, **51**(5):731-42. (Guideline Ref ID: LARSON1998)
160. Lepor H. Phase III multicenter placebo-controlled study of tamsulosin in benign prostatic hyperplasia. Tamsulosin Investigator Group. *Urology* 1998, **51**(6):892-900. (Guideline Ref ID: LEPOR1998A)
161. Lepor H, Auerbach S, Puras-Baez A, Narayan P, Soloway M, Lowe F *et al.* A randomized, placebo-controlled multicenter study of the efficacy and safety of terazosin in the treatment of benign prostatic hyperplasia. *Journal of Urology* 1992, **148**(5):1467-74. (Guideline Ref ID: LEPOR1992)
162. Lepor H, Jones K, Williford W. The mechanism of adverse events associated with terazosin: an analysis of the Veterans Affairs cooperative study. *Journal of Urology* 2000, **163**(4):1134-7. (Guideline Ref ID: LEPOR2000)
163. Lepor H, Williford WO, Barry MJ, Brawer MK, Dixon CM, Gormley G *et al.* The efficacy of terazosin, finasteride, or both in benign prostatic hyperplasia. Veterans Affairs

- Cooperative Studies Benign Prostatic Hyperplasia Study Group. *New England Journal of Medicine* 1996, **335**(8):533-9. (Guideline Ref ID: LEPOR1996)
164. Lepor H, Williford WO, Barry MJ, Haakenson C, Jones K. The impact of medical therapy on bother due to symptoms, quality of life and global outcome, and factors predicting response. Veterans Affairs Cooperative Studies Benign Prostatic Hyperplasia Study Group. *Journal of Urology* 1998, **160**(4):1358-67. (Guideline Ref ID: LEPOR1998)
165. Li MK, Ng AS. Bladder neck resection and transurethral resection of the prostate: a randomized prospective trial. *Journal of Urology* 1987, **138**(4):807-9. (Guideline Ref ID: LI1987)
166. Liedberg F, Adell L, Hagberg G, Palmqvist IB. Interstitial laser coagulation versus transurethral resection of the prostate for benign prostatic enlargement--a prospective randomized study. *Scandinavian Journal of Urology and Nephrology* 2003, **37**(6):494-7. (Guideline Ref ID: LIEDBERG2003)
167. Liguori G, Trombetta C, De GG, Pomara G, Maio G, Vecchio D *et al.* Efficacy and safety of combined oral therapy with tadalafil and alfuzosin: An integrated approach to the management of patients with lower urinary tract symptoms and erectile dysfunction. Preliminary report. *Journal of Sexual Medicine* 2009, **6**(2):544-52. (Guideline Ref ID: LIGUORI2009)
168. Liu CK, Lee WK, Ko MC, Chiang HS, Wan KS. Transurethral electrovapor resection versus standard transurethral resection treatment for a large prostate: a 2-year follow-up study conducted in Taiwan. *Urology international* 2006, **76**(2):144-9. (Guideline Ref ID: LIU2006)
169. Lloyd SN, Buckley JF, Chilton CP, Ibrahim I, Kaisary AV, Kirk D. Terazosin in the treatment of benign prostatic hyperplasia: a multicentre, placebo-controlled trial. *British Journal of Urology* 1992, **70**(Suppl 1):17-21. (Guideline Ref ID: LLOYD1992)
170. Logan K, Shaw C, Webber I, Samuel S, Broome L. Patients' experiences of learning clean intermittent self-catheterization: a qualitative study. *Journal of Advanced Nursing* 2008, **62**(1):32-41. (Guideline Ref ID: LOGAN2008)
171. Lopatkin N, Sivkov A, Walther C, Schlafke S, Medvedev A, Avdeichuk J *et al.* Long-term efficacy and safety of a combination of sabal and urtica extract for lower urinary tract symptoms--a placebo-controlled, double-blind, multicenter trial. *World Journal of Urology* 2005, **23**(2):139-46. (Guideline Ref ID: LOPATKIN2005)
172. Lourenco T, Armstrong N, N'Dow J, Nabi G, Deverill M, Pickard R *et al.* Systematic review and economic modelling of effectiveness and cost utility of surgical treatments for men with benign prostatic enlargement. *Health Technology Assessment* 2008, **12**(35):iii-169. (Guideline Ref ID: LOURENCO2008B)
173. Lourenco T, Pickard R, Vale L, Grant A, Fraser C, MacLennan G *et al.* Minimally invasive treatments for benign prostatic enlargement: systematic review of randomised controlled trials. *British Medical Journal* 2008, **337**:a1662. (Guideline Ref ID: LOURENCO2008A)
174. Lourenco T, Pickard R, Vale L, Grant A, Fraser C, MacLennan G *et al.* Alternative approaches to endoscopic ablation for benign enlargement of the prostate: systematic review of randomised controlled trials. *British Medical Journal* 2008, **337**:a449. (Guideline Ref ID: LOURENCO2008)

175. Lowe FC, McDaniel RL, Chmiel JJ, Hillman AL. Economic modelling to assess the costs of treatment with finasteride, terazosin, and transurethral resection of the prostate for men with moderate to severe symptoms of benign prostatic hyperplasia. *Urology* 1995, **46**(4):477-83. (Guideline Ref ID: LOWE1995)
176. Lucas MG, Stephenson TP, Nargund V. Tamsulosin in the management of patients in acute urinary retention from benign prostatic hyperplasia. *BJU International* 2005, **95**(3):354-7. (Guideline Ref ID: LUCAS2005)
177. Macaulay M, Clarke-O'Neill S, Fader M, Pettersson L, Cottenden A. A pilot study to evaluate reusable absorbent body-worn products for adults with moderate/heavy urinary incontinence. *Journal of Wound, Ostomy and Continence Nursing* 2004, **31**(6):357-66. (Guideline Ref ID: MACAULAY2004A)
178. Macaulay M, Clarke-O'Neill S, Fader M, Pettersson L, Cottenden A. Are washable absorbents effective at containing urinary incontinence? *Nursing Times* 2004, **100**(12):58-62. (Guideline Ref ID: MACAULAY2004)
179. MacDiarmid SA, Peters KM, Chen A, Armstrong RB, Orman C, Aquilina JW et al. Efficacy and safety of extended-release oxybutynin in combination with tamsulosin for treatment of lower urinary tract symptoms in men: randomized, double-blind, placebo-controlled study. *Mayo Clinic Proceedings* 2008, **83**(9):1002-10. (Guideline Ref ID: MACDIARMID2008)
180. Manassero F, Traversi C, Ales V, Pistolesi D, Panicucci E, Valent F et al. Contribution of early intensive prolonged pelvic floor exercises on urinary continence recovery after bladder neck-sparing radical prostatectomy: Results of a prospective controlled randomized trial. *Neurourology and Urodynamics* 2007, **26**(7):985-9. (Guideline Ref ID: MANASSERO2007)
181. Marberger MJ. Long-term effects of finasteride in patients with benign prostatic hyperplasia: a double-blind, placebo-controlled, multicenter study. PROWESS Study Group. *Urology* 1998, **51**(5):677-86. (Guideline Ref ID: MARBERGER1998)
182. Maria G, Brisinda G, Civello IM, Bentivoglio AR, Sganga G, Albanese A. Relief by botulinum toxin of voiding dysfunction due to benign prostatic hyperplasia: results of a randomized, placebo-controlled study. *Urology* 2003, **62**(2):259-64. (Guideline Ref ID: MARIA2003)
183. Martenson AC, de la Rosette JJ. Interstitial laser coagulation in the treatment of benign prostatic hyperplasia using a diode laser system: results of an evolving technology. *Prostate Cancer & Prostatic Diseases* 1999, **2**(3):148-54. (Guideline Ref ID: MARTENSON1999)
184. Martorana G, Giberti C, Di Silverio F, Von Heland M, Rigatti P, Colombo R et al. Effects of short-term treatment with the alpha 1-blocker alfuzosin on urodynamic pressure/flow parameters in patients with benign prostatic hyperplasia. *European Urology* 1997, **32**(1):47-53. (Guideline Ref ID: MARTORANA1997)
185. Mathewson-Chapman M. Pelvic muscle exercise/biofeedback for urinary incontinence after prostatectomy: an education program. *Journal of Cancer Education* 1997, **12**(4):218-23. (Guideline Ref ID: MATHEWSONCHAPMAN1997)
186. Mattiasson A, Wagrell L, Schelin S, Nordling J, Richthoff J, Magnusson B et al. Five-year follow-up of feedback microwave thermotherapy versus TURP for clinical BPH: a prospective randomized multicenter study. *Urology* 2007, **69**(1):91-6. (Guideline Ref ID: MATTIASSON2007)

187. Mavuduru RM. Comparison of HoLEP and TURP in terms of efficacy in the early postoperative period and perioperative morbidity. *Urologia Internationalis* 2009, **82**(2):130-5. (Guideline Ref ID: MAVUDURU2009)
188. McAllister WJ, Absalom MJ, Mir K, Shivde S, Anson K, Kirby RS *et al.* Does endoscopic laser ablation of the prostate stand the test of time? Five-year results from a multicentre randomized controlled trial of endoscopic laser ablation against transurethral resection of the prostate. *BJU International* 2000, **85**(4):437-9. (Guideline Ref ID: MCALLISTER2000)
189. McConnell JD, Barry MJ, Bruskewitz RC. (1994) Benign prostatic hyperplasia: diagnosis and treatment. Rockville, MD: Agency for Health Care Policy and Research, Public Health Service, US Department of Health and Human Services. (Guideline Ref ID: MCCONNELL1994)
190. McConnell JD, Bruskewitz R, Walsh P, Andriole G, Lieber M, Holtgrewe HL *et al.* The effect of finasteride on the risk of acute urinary retention and the need for surgical treatment among men with benign prostatic hyperplasia. Finasteride Long-Term Efficacy and Safety Study Group. *New England Journal of Medicine* 1998, **338**(9):557-63. (Guideline Ref ID: MCCONNELL1998)
191. McConnell JD, Roehrborn CG, Bautista OM, Andriole GL, Jr., Dixon CM, Kusek JW *et al.* The long-term effect of doxazosin, finasteride, and combination therapy on the clinical progression of benign prostatic hyperplasia. *New England Journal of Medicine* 2003, **349**(25):2387-98. (Guideline Ref ID: MCCONNELL2003)
192. McDonald H, Hux M, Brisson M, Bernard L, Nickel JC. An economic evaluation of doxazosin, finasteride and combination therapy in the treatment of benign prostatic hyperplasia. *Canadian Journal of Urology* 2004, **11**(4):2327-40. (Guideline Ref ID: MCDONALD2004)
193. McNeill SA, Daruwala PD, Mitchell ID, Shearer MG, Hargreave TB. Sustained-release alfuzosin and trial without catheter after acute urinary retention: a prospective, placebo-controlled. *BJU International* 1999, **84**(6):622-7. (Guideline Ref ID: MCNEILL1999)
194. McNeill SA, Hargreave TB, Members of the Alfaur Study Group. Alfuzosin once daily facilitates return to voiding in patients in acute urinary retention. *Journal of Urology* 2004, **171**(6 Pt 1):2316-20. (Guideline Ref ID: MCNEILL2004A)
195. McNeill SA, Hargreave TB, Roehrborn CG, Alfaur study group. Alfuzosin 10 mg once daily in the management of acute urinary retention: results of a double-blind placebo-controlled study. *Urology* 2005, **65**(1):83-9. (Guideline Ref ID: MCNEILL2005)
196. McVary KT, Monnig W, Camps JL, Jr., Young JM, Tseng LJ, van den Ende G. Sildenafil citrate improves erectile function and urinary symptoms in men with erectile dysfunction and lower urinary tract symptoms associated with benign prostatic hyperplasia: a randomized, double-blind trial. *Journal of Urology* 2007, **177**(3):1071-7. (Guideline Ref ID: MCVARY2007C)
197. McVary KT, Roehrborn CG, Kaminetsky JC, Auerbach SM, Wachs B, Young JM *et al.* Tadalafil relieves lower urinary tract symptoms secondary to benign prostatic hyperplasia. *Journal of Urology* 2007, **177**(4):1401-7. (Guideline Ref ID: MCVARY2007B)
198. Medicare Services Advisory Committee. Transurethral needle ablation (TUNA) for the treatment of benign prostatic hyperplasia Cochrane Database of Systematic Reviews (Guideline Ref ID: MSAC2002)

199. Melo EA, Bertero EB, Rios LAS, Mattos J. Evaluating the efficiency of a combination of *Pygeum africanum* and stinging nettle (*Urtica dioica*) extracts in treating benign prostatic hyperplasia (BPH): Double-blind, randomized, placebo controlled trial. *International Brazilian Journal of Urology* 2002, **28**(5):418-25. (Guideline Ref ID: MELO2002)
200. Michielsen DP, Debacker T, De Boe V, Van Lersberghe C, Kaufman L, Braeckman JG et al. Bipolar transurethral resection in saline--an alternative surgical treatment for bladder outlet obstruction? *Journal of Urology* 2007, **178**(5):2035-9. (Guideline Ref ID: MICHIELSEN2007)
201. Mohanty NK, Nayak RL, Malhotra V, Arora RP. A double-blind placebo controlled study of tamsulosin in the management of benign prostatic hyperplasia in an Indian population. *Annals of the College of Surgeons of Hong Kong* 2003, **7**(3):88-93. (Guideline Ref ID: MOHANTY2003)
202. Montorsi F, Naspro R, Salonia A, Suardi N, Briganti A, Zanoni M et al. Holmium laser enucleation versus transurethral resection of the prostate: results from a 2-center, prospective, randomized trial in patients with obstructive benign prostatic hyperplasia. *Journal of Urology* 2004, **172**(5 Pt 1):1926-9. (Guideline Ref ID: MONTORSI2004)
203. Moore KN, Griffiths D, Hughton A. Urinary incontinence after radical prostatectomy: a randomized controlled trial comparing pelvic muscle exercises with or without electrical stimulation. *BJU International* 1999, **83**(1):57-65. (Guideline Ref ID: MOORE1999A)
204. Moore KN, Schieman S, Ackerman T, Dzus HY, Metcalfe JB, Voaklander DC. Assessing comfort, safety, and patient satisfaction with three commonly used penile compression devices. *Urology* 2004, **63**(1):150-4. (Guideline Ref ID: MOORE2004)
205. Mostafid AH, Harrison NW, Thomas PJ, Fletcher MS. A prospective randomized trial of interstitial radiofrequency therapy versus transurethral resection for the treatment of benign prostatic hyperplasia. *British Journal of Urology* 1997, **80**(1):116-22. (Guideline Ref ID: MOSTAFID1997)
206. Mottet N, Anidjar M, Bourdon O, Louis JF, Teillac P, Costa P et al. Randomized comparison of transurethral electroresection and holmium: YAG laser vaporization for symptomatic benign prostatic hyperplasia. *Journal of Endourology* 1999, **13**(2):127-30. (Guideline Ref ID: MOTTET1999)
207. Murray E, Davis H, Tai SS, Coulter A, Gray A, Haines A. Randomised controlled trial of an interactive multimedia decision aid on benign prostatic hypertrophy in primary care. *British Medical Journal* 2001, **323**(7311):493-6. (Guideline Ref ID: MURRAY2001)
208. Narayan P, Tewari A. A second phase III multicenter placebo controlled study of 2 dosages of modified release tamsulosin in patients with symptoms of benign prostatic hyperplasia. *Journal of Urology* 1998, **160**(5):1701-6. (Guideline Ref ID: NARAYAN1998)
209. Narayan P, Tewari A, Aboseif S, Evans C. A randomized study comparing visual laser ablation and transurethral evaporation of prostate in the management of benign prostatic hyperplasia. *Journal of Urology* 1995, **154**(6):2083-8. (Guideline Ref ID: NARAYAN1995)
210. Naspro R, Suardi N, Salonia A, Scattoni V, Guazzoni G, Colombo R et al. Holmium laser enucleation of the prostate versus open prostatectomy for prostates >70 g: 24-month follow-up. *European Urology* 2006, **50**(3):563-8. (Guideline Ref ID: NASPRO2006)

211. Nathan MS, Wickham JEA. TVP: a cheaper and effective alternative to TURP. *Minimally invasive therapy and allied technologies* 1996, **5**(3):292-6. (Guideline Ref ID: NATHAN1996)
212. National Collaborating Centre for Primary Care. Medicines concordance and adherence: involving adults and carers in decisions about prescribed medicines: draft for consultation. <http://www.nice.org.uk/nicemedia/pdf/MedicinesConcordanceDraftFullGuidelineForConsultation.doc> [accessed 15-9-2008]. (Guideline Ref ID: NCCPC2008)
213. National Institute for Clinical Excellence. Sacral nerve stimulation for urge incontinence and urgency-frequency <http://www.nice.org.uk/IPG064> [accessed 13-1-2009]. (Guideline Ref ID: IPG0642004)
214. National Institute for Health and Clinical Excellence. The guidelines manual 2007 www.nice.org.uk (Guideline Ref ID: NICE2007)
215. National Institute for Health and Clinical Excellence. Guide to the methods of technology appraisals <http://www.nice.org.uk/media/B52/A7/TAMethodsGuideUpdatedJune2008.pdf> [accessed 19-12-2008]. (Guideline Ref ID: NATIONALINSTITU2008)
216. National Institute for Health and Clinical Excellence. Suburethral synthetic sling insertion for stress urinary incontinence in men <http://www.nice.org.uk/IPG256> [accessed 13-1-2009]. (Guideline Ref ID: IPG2562008)
217. National Institute for Health and Clinical Excellence. The guidelines manual 2009 <http://www.nice.org.uk> [accessed 13-1-2009]. (Guideline Ref ID: NICE2009)
218. Nawrocki JD, Bell TJ, Lawrence WT, Ward JP. A randomized controlled trial of transurethral microwave thermotherapy. *British Journal of Urology* 1997, **79**(3):389-93. (Guideline Ref ID: NAWROCKI1997)
219. Netto NR, Jr., De Lima ML, Lucena R, Lavoura NS, Cortado PL, Netto MR. Is transurethral vaporization a remake of transurethral resection of the prostate? *Journal of Endourology* 1999, **13**(8):591-4. (Guideline Ref ID: NETTO1999)
220. Nickel JC, Fradet Y, Boake RC, Pommerville PJ, Perreault JP, Afridi SK *et al.* Efficacy and safety of finasteride therapy for benign prostatic hyperplasia: results of a 2-year randomized controlled trial (the PROSPECT study). PROscar Safety Plus Efficacy Canadian Two year Study. *Canadian Medical Association Journal* 1996, **155**(9):1251-9. (Guideline Ref ID: NICKEL1996)
221. Nielsen HO. Transurethral prostatotomy versus transurethral prostatectomy in benign prostatic hypertrophy. A prospective randomised study. *British Journal of Urology* 1988, **61**(5):435-8. (Guideline Ref ID: NIELSEN1988)
222. Noble SM, Coast J, Brookes S, Neal DE, Abrams P, Peters TJ *et al.* Transurethral prostate resection, noncontact laser therapy or conservative management in men with symptoms of benign prostatic enlargement? An economic evaluation. *Journal of Urology* 2002, **168**:2476-82. (Guideline Ref ID: NOBLE2002)
223. Norby B, Nielsen HV, Frimodt-Moller PC. Cost-effectiveness of new treatments for benign prostatic hyperplasia: results of a randomized trial comparing the short-term cost-effectiveness of transurethral interstitial laser coagulation of the prostate, transurethral

- microwave thermotherapy and standard transurethral resection or incision of the prostate. *Scandinavian Journal of Urology and Nephrology* 2002, **36**(4):286-95. (Guideline Ref ID: NORBY2002)
224. Nørby B, Nielsen HV, Frimodt-Møller PC. Transurethral interstitial laser coagulation of the prostate and transurethral microwave thermotherapy vs transurethral resection or incision of the prostate: results of a randomized, controlled study in patients with symptomatic benign prostatic hyperplasia. *BJU International* 2002, **90**(9):853-62. (Guideline Ref ID: NORBY2002A)
225. Nordling J. Efficacy and safety of two doses (10 and 15 mg) of alfuzosin or tamsulosin (0.4 mg) once daily for treating symptomatic benign prostatic hyperplasia. *BJU International* 2005, **95**(7):1006-12. (Guideline Ref ID: NORDLING2005A)
226. Nordling J, Wagrell L, Larson T, Duelund J, Kroyer K, Mattiasson A. ProstaLund feedback treatment (PLFT) versus TURP: a prospective randomized multicenter study with 36 month follow up. *BJU International* 2005, **95**(Suppl 5):75. (Guideline Ref ID: NORDLING2005)
227. Nuhoglu B, Ayyildiz A, Fidan V, Ersoy E, Huri E, Germiyanoglu C. Transurethral electrovaporization of the prostate: is it any better than standard transurethral prostatectomy? 5-year follow-up. *Journal of Endourology* 2005, **19**(1):79-82. (Guideline Ref ID: NUHOGLU2005)
228. Nuhoglu B, Ayyildiz A, Karaguzel E, Cebeci O, Germiyanoglu C. Plasmakinetic prostate resection in the treatment of benign prostate hyperplasia: results of 1-year follow up. *International Journal of Urology* 2006, **13**(1):21-4. (Guideline Ref ID: NUHOGLU2006)
229. O'Leary MP, Roehrborn C, Andriole G, Nickel C, Boyle P, Hofner K. Improvements in benign prostatic hyperplasia-specific quality of life with dutasteride, the novel dual 5alpha-reductase inhibitor. *BJU International* 2003, **92**(3):262-6. (Guideline Ref ID: OLEARY2003)
230. O'Leary MP, Roehrborn CG, Black L. Dutasteride significantly improves quality of life measures in patients with enlarged prostate. *Prostate Cancer & Prostatic Diseases* 2008, **11**(2):129-33. (Guideline Ref ID: OLEARY2008)
231. Oelke M, Hofner K, Jonas U, de la Rosette JJ, Ubbink DT, Wijkstra H. Diagnostic accuracy of noninvasive tests to evaluate bladder outlet obstruction in men: detrusor wall thickness, uroflowmetry, postvoid residual urine, and prostate volume. *European Urology* 2007, **52**(3):827-34. (Guideline Ref ID: OELKE2007)
232. Ogden C, Reddy P, Johnson H, Carter S. Sham vs TUMT: a randomized study with cross over. *Journal of Urology* 1993, **149**(4 Supp):250A. (Guideline Ref ID: OGDEN1993)
233. Parekh AR, Feng MI, Kirages D, Bremner H, Kaswick J, Aboseif S. The role of pelvic floor exercises on post-prostatectomy incontinence. *Journal of Urology* 2003, **170**(1):130-3. (Guideline Ref ID: PAREKH2003)
234. Patankar S, Jamkar A, Dobhada S, Gorde V. PlasmaKinetic Superpulse transurethral resection versus conventional transurethral resection of prostate. *Journal of Endourology* 2006, **20**(3):215-9. (Guideline Ref ID: PATANKAR2006)
235. Patel A, Fuchs GJ, Gutierrez-Aceves J, Ryan TP. Prostate heating patterns comparing electrosurgical transurethral resection and vaporization: a prospective randomized study. *Journal of Urology* 1997, **157**(1):169-72. (Guideline Ref ID: PATEL1997)

236. Paterson J, Dunn S, Kowanko I, van Loon A, Stein I, Pretty L. Selection of continence products: Perspectives of people who have incontinence and their carers. *Disability and Rehabilitation: An International, Multidisciplinary Journal* 2003, **25**(17):955-63. (Guideline Ref ID: PATERSON2003)
237. Paterson J, Pinnock CB, Marshall VR. Pelvic floor exercises as a treatment for post-micturition dribble. *British Journal of Urology* 1997, **79**(6):892-7. (Guideline Ref ID: PATERSON1997)
238. Penson DF, Ramsey S, Veenstra D, Clarke L, Gandhi S, Hirsch M. The cost-effectiveness of combined androgen blockade with bicalutamide and luteinizing hormone releasing hormone agonist in men with metastatic prostate cancer. *Journal of Urology* 2005, **174**(2):547-52. (Guideline Ref ID: PENSON2005)
239. Polat O, Ozbey I, Gul O, Demirel A, Bayraktar Y. Pharmacotherapy of benign prostatic hyperplasia: inhibitor of 5 alpha-reductase. *International Urology and Nephrology* 1997, **29**(3):323-30. (Guideline Ref ID: POLAT1997)
240. Porru D, Campus G, Caria A, Madeddu G, Cucchi A, Rovereto B *et al.* Impact of early pelvic floor rehabilitation after transurethral resection of the prostate. *Neurourology and Urodynamics* 2001, **20**(1):53-9. (Guideline Ref ID: PORRU2001)
241. Poulsen AL, Schou J, Puggaard L, Torp-Pedersen S, Nordling J. Prostatic enlargement, symptomatology and pressure/flow evaluation: Interrelations in patients with symptomatic BPH. *Scandinavian Journal of Urology and Nephrology Supplementum* 1994, **157**:67-73. (Guideline Ref ID: POULSEN1994)
242. Preuss HG, Marcusen C, Regan J, Klimberg IW, Welebir TA, Jones WA. Randomized trial of a combination of natural products (cernitin, saw palmetto, B-sitosterol, vitamin E) on symptoms of benign prostatic hyperplasia (BPH). *International Urology and Nephrology* 2001, **33**(2):217-25. (Guideline Ref ID: PREUSS2001)
243. Ragnarson TG, Hjelmgren J, Malmberg L. Under what conditions is feedback microwave thermotherapy (ProstaLund Feedback Treatment) cost-effective in comparison with alpha-blockade in the treatment of benign prostatic hyperplasia and lower urinary tract symptoms? *Scandinavian Journal of Urology and Nephrology* 2006, **40**(6):495-505. (Guideline Ref ID: RAGNARSON2006)
244. Razzaghi MR, Habibi G, Djavid GE, Gholamrezaee H. Laser prostatectomy versus transurethral resection of prostate in the treatment of benign prostatic hyperplasia. *Saudi medical journal* 2007, **28**(1):68-72. (Guideline Ref ID: RAZZAGHI2007)
245. Resnick MI, Roehrborn CG. Rapid onset of action with alfuzosin 10 mg once daily in men with benign prostatic hyperplasia: a randomized, placebo-controlled trial. *Prostate Cancer & Prostatic Diseases* 2007, **10**(2):155-9. (Guideline Ref ID: RESNICK2007)
246. Revicki D, Hays RD, Cella D, Sloan J. Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. *Journal of Clinical Epidemiology* 2008, **61**(2):102-9. (Guideline Ref ID: REVICKI2008)
247. Reynard JM, Cannon A, Yang Q, Abrams P. A novel therapy for nocturnal polyuria: a double-blind randomized trial of frusemide against placebo. *British Journal of Urology* 1998, **81**(2):215-8. (Guideline Ref ID: REYNARD1998A)

248. Reynard JM, Peters TJ, Lim C, Abrams P. The value of multiple free-flow studies in men with lower urinary tract symptoms. *British Journal of Urology* 1996, **77**(6):813-8. (Guideline Ref ID: REYNARD1996)
249. Reynard JM, Yang Q, Donovan JL, Peters TJ, Schafer W, De la Rosette JJMC et al. The ICS-'BPH' Study: Uroflowmetry, lower urinary tract symptoms and bladder outlet obstruction. *British Journal of Urology* 1998, **82**(5):619-23. (Guideline Ref ID: REYNARD1998)
250. Riehmman M, Knes JM, Heisey D, Madsen PO, Bruskewitz RC. Transurethral resection versus incision of the prostate: a randomized, prospective study. *Urology* 1995, **45**(5):768-75. (Guideline Ref ID: RIEHMANN1995)
251. Rigatti L, Naspro R, Salonia A, Centemero A, Ghezzi M, Guazzoni G et al. Urodynamics after TURP and HoLEP in urodynamically obstructed patients: are there any differences at 1 year of follow-up? *Urology* 2006, **67**(6):1193-8. (Guideline Ref ID: RIGATTI2006)
252. Rigatti P, Brausi M, Scarpa RM, Porru D, Schumacher H, Rizzi CA et al. A comparison of the efficacy and tolerability of tamsulosin and finasteride in patients with lower urinary tract symptoms suggestive of benign prostatic hyperplasia. *Prostate Cancer & Prostatic Diseases* 2003, **6**(4):315-23. (Guideline Ref ID: RIGATTI2003)
253. Rodrigo Aliaga M, Valls Blasco F, Jimenez Cruz JF. Lasers as an alternative to the endoscopic surgery in BPH. *Actas Urologicas Espanolas* 1998, **22**(1):17-22. (Guideline Ref ID: RODRIGO1998)
254. Roehrborn CG. Efficacy and safety of once-daily alfuzosin in the treatment of lower urinary tract symptoms and clinical benign prostatic hyperplasia: a randomized, placebo-controlled trial. *Urology* 2001, **58**(6):953-9. (Guideline Ref ID: ROEHRBORN2001A)
255. Roehrborn CG. Alfuzosin 10 mg once daily prevents overall clinical progression of benign prostatic hyperplasia but not acute urinary retention: results of a 2-year placebo-controlled study. *BJU International* 2006, **97**(4):734-41. (Guideline Ref ID: ROEHRBORN2006)
256. Roehrborn CG, Boyle P, Bergner D, Gray T, Gittelman M, Shown T et al. Serum prostate-specific antigen and prostate volume predict long-term changes in symptoms and flow rate: results of a four-year, randomized trial comparing finasteride versus placebo. PLESS Study Group. *Urology* 1999, **54**(4):662-9. (Guideline Ref ID: ROEHRBORN1999)
257. Roehrborn CG, Boyle P, Nickel JC, Hoefner K, Andriole G, A.R.I.A. Efficacy and safety of a dual inhibitor of 5-alpha-reductase types 1 and 2 (dutasteride) in men with benign prostatic hyperplasia. *Urology* 2002, **60**(3):434-41. (Guideline Ref ID: ROEHRBORN2002A)
258. Roehrborn CG, Bruskewitz R, Nickel GC, Glickman S, Cox C, Anderson R et al. Urinary retention in patients with BPH treated with finasteride or placebo over 4 years. Characterization of patients and ultimate outcomes. The PLESS Study Group. *European Urology* 2000, **37**(5):528-36. (Guideline Ref ID: ROEHRBORN2000)
259. Roehrborn CG, Burkhard FC, Bruskewitz RC, Issa MM, Perez-Marrero R, Naslund MJ et al. The effects of transurethral needle ablation and resection of the prostate on pressure flow urodynamic parameters: analysis of the United States randomized study. *Journal of Urology* 1999, **162**(1):92-7. (Guideline Ref ID: ROEHRBORN1999B)

260. Roehrborn CG, McConnell JD, Saltzman B, Bergner D, Gray T, Narayan P *et al.* Storage (irritative) and voiding (obstructive) symptoms as predictors of benign prostatic hyperplasia progression and related outcomes. *European Urology* 2002, **42**(1):1-6. (Guideline Ref ID: ROEHRBORN2002)
261. Roehrborn CG, McVary KT, Elion-Mboussa A, Viktrup L. Tadalafil administered once daily for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a dose finding study. *Journal of Urology* 2008, **180**(4):1228-34. (Guideline Ref ID: ROEHRBORN2008B)
262. Roehrborn CG, Oesterling JE, Auerbach S, Kaplan SA, Lloyd LK, Milam DE *et al.* The Hytrin Community Assessment Trial study: a one-year study of terazosin versus placebo in the treatment of men with symptomatic benign prostatic hyperplasia. HYCAT Investigator Group. *Urology* 1996, **47**(2):159-68. (Guideline Ref ID: ROEHRBORN1996A)
263. Roehrborn CG, Siami P, Barkin J, Damiao R, Major-Walker K, Morrill B *et al.* The effects of dutasteride, tamsulosin and combination therapy on lower urinary tract symptoms in men with benign prostatic hyperplasia and prostatic enlargement: 2-year results from the CombAT study. *Journal of Urology* 2008, **179**(2):616-21. (Guideline Ref ID: ROEHRBORN2008)
264. Rovner ES, Kreder K, Sussman DO, Kaplan SA, Carlsson M, Bavendam T *et al.* Effect of tolterodine extended release with or without tamsulosin on measures of urgency and patient reported outcomes in men with lower urinary tract symptoms. *Journal of Urology* 2008, **180**(3):1034-41. (Guideline Ref ID: ROVNER2008A)
265. Safarinejad MR. *Urtica dioica* for treatment of benign prostatic hyperplasia: a prospective, randomized, double-blind, placebo-controlled, crossover study. *Journal of Herbal Pharmacotherapy* 2005, **5**(4):1-11. (Guideline Ref ID: SAFARINEJAD2005)
266. Saint S, Lipsky BA, Baker PD, McDonald LL, Ossenkop K. Urinary catheters: what type do men and their nurses prefer? *Journal of the American Geriatrics Society* 1999, **47**(12):1453-8. (Guideline Ref ID: SAINT1999)
267. Salonia A, Suardi N, Naspro R, Mazzoccoli B, Zanni G, Gallina A *et al.* Holmium laser enucleation versus open prostatectomy for benign prostatic hyperplasia: an inpatient cost analysis. *Urology* 2006, **68**(2):302-6. (Guideline Ref ID: SALONIA2006)
268. Saporta L, Aridogan IA, Erlich N, Yachia D. Objective and subjective comparison of transurethral resection, transurethral incision and balloon dilatation of the prostate. A prospective study. *European Urology* 1996, **29**(4):439-45. (Guideline Ref ID: SAPORTA1996)
269. Schulman CC, De Sy W, Vandendris M, Tomas M, Santoni JP. Belgian multicenter clinical study of alfuzosin, a selective alpha 1-blocker, in the treatment of benign prostatic hyperplasia. The Alfuzosin Belgian Group. *Acta Urologica Belgica* 1994, **62**(4):15-21. (Guideline Ref ID: SCHULMAN1994)
270. Seckiner I., Yesilli C, Akduman B, Mungan NA. A prospective randomized study for comparing bipolar plasmakinetic resection of the prostate with standard TURP. *Urology international* 2006, **76**(2):139-43. (Guideline Ref ID: SECKINER2006)
271. Sengor F, Kose O, Yucebas E, Beysel M, Erdogan K, Narter F. A comparative study of laser ablation and transurethral electroresection for benign prostatic hyperplasia: results of a 6-

- month follow-up. *British Journal of Urology* 1996, **78**(3):398-400. (Guideline Ref ID: SENGOR1996)
272. Shah T, Palit V, Biyani S, Elmasry Y, Puri R, Flannigan GM. Randomised, placebo controlled, double blind study of alfuzosin SR in patients undergoing trial without catheter following acute urinary retention. *European Urology* 2002, **42**(4):329-32. (Guideline Ref ID: SHAH2002)
273. Shaw C, Logan K, Webber I, Broome L, Samuel S. Effect of clean intermittent self-catheterization on quality of life: a qualitative study. *Journal of Advanced Nursing* 2008, **61**(6):641-50. (Guideline Ref ID: SHAW2008)
274. Shi R, Xie Q, Gang X, Lun J, Cheng L, Pantuck A *et al.* Effect of saw palmetto soft gel capsule on lower urinary tract symptoms associated with benign prostatic hyperplasia: a randomized trial in Shanghai, China. *Journal of Urology* 2008, **179**(2):610-5. (Guideline Ref ID: SHI2008)
275. Shingleton WB, Farabaugh P, May W. Three-year follow-up of laser prostatectomy versus transurethral resection of the prostate in men with benign prostatic hyperplasia. *Urology* 2002, **60**(2):305-8. (Guideline Ref ID: SHINGLETON2002)
276. Shingleton WB, Renfro LD, Kolski JM, Fowler JE. A randomized prospective study of transurethral electrovaporization vs laser ablation of the prostate in men with benign prostatic hypertrophy. *Scandinavian Journal of Urology and Nephrology* 1998, **32**(4):266-9. (Guideline Ref ID: SHINGLETON1998)
277. Shingleton WB, Terrell F, Renfro DL, Kolski JM, Fowler JE, Jr. A randomized prospective study of laser ablation of the prostate versus transurethral resection of the prostate in men with benign prostatic hyperplasia. *Urology* 1999, **54**(6):1017-21. (Guideline Ref ID: SHINGLETON1999)
278. Shokeir AA, al Sisi H, Farage YM, el Maaboud MA, Saeed M, Mutabagani H. Transurethral prostatectomy: a prospective randomized study of conventional resection and electrovaporization in benign prostatic hyperplasia. *British Journal of Urology* 1997, **80**(4):570-4. (Guideline Ref ID: SHOKEIR1997)
279. Siami P, Roehrborn CG, Barkin J, Damiao R, Wyczolkowski M, Duggan A *et al.* Combination therapy with dutasteride and tamsulosin in men with moderate-to-severe benign prostatic hyperplasia and prostate enlargement: the CombAT (Combination of Avodart and Tamsulosin) trial rationale and study design. *Contemporary Clinical Trials* 2007, **28**(6):770-9. (Guideline Ref ID: SIAMI2007)
280. Singh H, Desai MR, Shrivastav P, Vani K. Bipolar versus monopolar transurethral resection of prostate: randomized controlled study. *Journal of Endourology* 2005, **19**(3):333-8. (Guideline Ref ID: SINGH2005)
281. Skolarikos A, Papachristou C, Athanasiadis G, Chalikopoulos D, Deliveliotis C, Alivizatos G. Eighteen-month results of a randomized prospective study comparing transurethral photoselective vaporization with transvesical open enucleation for prostatic adenomas greater than 80 cc. *Journal of Endourology* 2008, **22**(10):2333-40. (Guideline Ref ID: SKOLARIKOS2008)

282. Sokeland J. Combined sabal and urtica extract compared with finasteride in men with benign prostatic hyperplasia: analysis of prostate volume and therapeutic outcome. *BJU International* 2000, **86**(4):439-42. (Guideline Ref ID: SOKELAND2000)
283. Sokeland J, Albrecht J. [Combination of Sabal and Urtica extract vs. finasteride in benign prostatic hyperplasia (Aiken stages I to II). Comparison of therapeutic effectiveness in a one year double-blind study]. *Urologe A* 1997, **36**(4):327-33. (Guideline Ref ID: SOKELAND1997)
284. Soloway M, Snyder J, Stone N, Laddu A. Terazosin for the treatment of benign prostatic hyperplasia in the elderly: a 6-month double-blind study. *Journal of the American Geriatrics Society* 1992, **40**:SA 11. (Guideline Ref ID: SOLOWAY1992)
285. Soonawalla PF, Pardanani DS. Transurethral incision versus transurethral resection of the prostate. A subjective and objective analysis. *British Journal of Urology* 1992, **70**(2):174-7. (Guideline Ref ID: SOONAWALLA1992)
286. Stepanov VN, Siniakova LA, Sarrazin B, Raynaud JP. Efficacy and tolerability of the lipidosterolic extract of *Serenoa repens* (Permixon) in benign prostatic hyperplasia: a double-blind comparison of two dosage regimens. *Advances in Therapy* 1999, **16**(5):231-41. (Guideline Ref ID: STEPANOV1999)
287. Stief CG, Porst H, Neuser D, Beneke M, Ulbrich E. A randomised, placebo-controlled study to assess the efficacy of twice-daily vardenafil in the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia. *European Urology* 2008, **53**(6):1236-44. (Guideline Ref ID: STIEF2008)
288. Stovsky MD, Griffiths R, I, Duff SB. A clinical outcomes and cost analysis comparing photoselective vaporization of the prostate to alternative minimally invasive therapies and transurethral prostate resection for the treatment of benign prostatic hyperplasia. *Journal of Urology* 2006, **176**(4):1500-6. (Guideline Ref ID: STOVSKY2006)
289. Sullivan PW, Nichol MB. The economic impact of payer policies after the Rx-to-OTC switch of second-generation antihistamines. *Value in health* 2004, **7**(4):402-12. (Guideline Ref ID: SULLIVAN2004)
290. Suvakovic N, Hindmarsh JR. A step towards day case prostatectomy. *British Journal of Urology* 1996, **77**(2):212-4. (Guideline Ref ID: SUVAKOVIC1996)
291. Talic RF, El Tiraifi A, El Faqih SR, Hassan SH, Attassi RA, Abdel-Halim RE. Prospective randomized study of transurethral vaporization resection of the prostate using the thick loop and standard transurethral prostatectomy. *Urology* 2000, **55**(6):886-90. (Guideline Ref ID: TALIC2000)
292. Tan AH, Gilling PJ, Kennett KM, Frampton C, Westenberg AM, Fraundorfer MR. A randomized trial comparing holmium laser enucleation of the prostate with transurethral resection of the prostate for the treatment of bladder outlet obstruction secondary to benign prostatic hyperplasia in large glands (40 to 200 grams). *Journal of Urology* 2003, **170**(4 Pt 1):1270-4. (Guideline Ref ID: TAN2003)
293. Tenover JL, Pagano GA, Morton AS, Liss CL, Byrnes CA. Efficacy and tolerability of finasteride in symptomatic benign prostatic hyperplasia: a primary care study. Primary Care Investigator Study Group. *Clinical Therapeutics* 1997, **19**(2):243-58. (Guideline Ref ID: TENOVER1997)

294. Tibaek S, Klarskov P, Hansen BL, Thomsen H, Andresen H, Jensen CS *et al.* Pelvic floor muscle training before transurethral resection of the prostate: A randomized, controlled, blinded study. *Scandinavian Journal of Urology and Nephrology* 2007, **41**(4):329-34. (Guideline Ref ID: TIBAEK2007)
295. Tkocz M, Prajsner A. Comparison of long-term results of transurethral incision of the prostate with transurethral resection of the prostate, in patients with benign prostatic hypertrophy. *Neurourology and Urodynamics* 2002, **21**(2):112-6. (Guideline Ref ID: TKOCZ2002)
296. Trachtenberg J, Roehrborn CG. Updated results of a randomized, double-blind, multicenter sham-controlled trial of microwave thermotherapy with the Dornier Urowave in patients with symptomatic benign prostatic hyperplasia. Urowave Investigators Group. *World Journal of Urology* 1998, **16**(2):102-8. (Guideline Ref ID: TRACHTENBERG1998)
297. Trueman P, Hood SC, Nayak US, Mrazek MF. Prevalence of lower urinary tract symptoms and self-reported diagnosed 'benign prostatic hyperplasia', and their effect on quality of life in a community-based survey of men in the UK. *BJU International* 1999, **83**(4):410-5. (Guideline Ref ID: TRUEMAN1999)
298. Tubaro A, La Vecchia C. The relation of lower urinary tract symptoms with life-style factors and objective measures of benign prostatic enlargement and obstruction: An Italian survey. *European Urology* 2004, **45**(6):767-72. (Guideline Ref ID: TUBARO2004)
299. Tuhkanen K, Heino A, Aaltomaa S, Ala-Opas M. Long-term results of contact laser versus transurethral resection of the prostate in the treatment of benign prostatic hyperplasia with small or moderately enlarged prostates. *Scandinavian Journal of Urology and Nephrology* 2003, **37**(6):487-93. (Guideline Ref ID: TUHKANEN2003)
300. Tuhkanen K, Heino A, Ala-Opas M. Two-year follow-up results of a prospective randomized trial comparing hybrid laser prostatectomy with TURP in the treatment of big benign prostates. *Scandinavian Journal of Urology and Nephrology* 2001, **35**(3):200-4. (Guideline Ref ID: TUHKANEN2001)
301. Tuhkanen K, Heino A, Alaopas M. Hybrid laser treatment compared with transurethral resection of the prostate for symptomatic bladder outlet obstruction caused by a large benign prostate: a prospective, randomized trial with a 6-month follow-up. *BJU International* 1999, **84**(7):805-9. (Guideline Ref ID: TUHKANEN1999A)
302. U.S.Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Biologics Evaluation and Research (CBER), Center for Devices and Radiological Health (CDRH). Guidance for industry patient-reported outcome measures: use in medical product development to support labelling claims - draft guidance <http://www.ispor.org/workpaper/FDA%20PRO%20Guidance.pdf> [accessed 7-12-2009]. (Guideline Ref ID: ANON2006B)
303. Vale JA, Bdesha AS, Witherow RO. An analysis of the costs of alternative treatments for benign prostatic hypertrophy. *Journal of the Royal Society of Medicine* 2007, **88**(11):644P-8P. (Guideline Ref ID: VALE1995)
304. Van Kampen M, De Weerd W, Van Poppel H, De Ridder D, Feys H, Baert L. Effect of pelvic-floor re-education on duration and degree of incontinence after radical prostatectomy: a randomised controlled trial. *Lancet* 2000, **355**(9198):98-102. (Guideline Ref ID: VANKAMPEN2000)

305. van Kerrebroeck P, Jardin A, Laval KU, Van Cangh P. Efficacy and safety of a new prolonged release formulation of alfuzosin 10 mg once daily versus alfuzosin 2.5 mg thrice daily and placebo in patients with symptomatic benign prostatic hyperplasia. ALFORTI Study Group. *European Urology* 2000, **37**(3):306-13. (Guideline Ref ID: VANKERREBROECK2000)
306. van Melick HH, Van Venrooij GE, Boon TA. Long-term follow-up after transurethral resection of the prostate, contact laser prostatectomy, and electrovaporization. *Urology* 2003, **62**(6):1029-34. (Guideline Ref ID: VANMELICK2003A)
307. van Melick HH, Van Venrooij GE, Eckhardt MD, Boon TA. A randomized controlled trial comparing transurethral resection of the prostate, contact laser prostatectomy and electrovaporization in men with benign prostatic hyperplasia: urodynamic effects. *Journal of Urology* 2002, **168**(3):1058-62. (Guideline Ref ID: VANMELICK2002)
308. van Melick HH, Van Venrooij GE, Eckhardt MD, Boon TA. A randomized controlled trial comparing transurethral resection of the prostate, contact laser prostatectomy and electrovaporization in men with benign prostatic hyperplasia: analysis of subjective changes, morbidity and mortality. *Journal of Urology* 2003, **169**(4):1411-6. (Guideline Ref ID: VANMELICK2003)
309. Van Melick HHE, Van Venrooij GEP, Boon TA. Laser prostatectomy in patients on anticoagulant therapy or with bleeding disorders. *Journal of Urology* 2003, **170**(5):1851-5. (Guideline Ref ID: VANMELICK2003B)
310. Varney SJ, Guest JF. The annual cost of blood transfusions in the UK. *Transfusion Medicine* 2003, **13**(4):205-18. (Guideline Ref ID: VARNEY2003)
311. Vera-Llonch M, Brandenburg NA, Oster G. Cost-effectiveness of Add-on Therapy with Pregabalin in Patients with Refractory Partial Epilepsy. *Epilepsia* 2008, **49**(3):431-7. (Guideline Ref ID: VERALLONCH2008)
312. Wagrell L, Schelin S, Nordling J, Richthoff J, Magnusson B, Schain M et al. Feedback microwave thermotherapy versus TURP for clinical BPH--a randomized controlled multicenter study. *Urology* 2002, **60**(2):292-9. (Guideline Ref ID: WAGRELL2002)
313. Wagrell L, Schelin S, Nordling J, Richthoff J, Magnusson B, Schain M et al. Three-year follow-up of feedback microwave thermotherapy versus TURP for clinical BPH: a prospective randomized multicenter study. *Urology* 2004, **64**(4):698-702. (Guideline Ref ID: WAGRELL2004)
314. Walden M, Acosta S, Carlsson P, Pettersson S, Dahlstrand C. A cost-effectiveness analysis of transurethral resection of the prostate and transurethral microwave thermotherapy for treatment of benign prostatic hyperplasia: two-year follow-up. *Scandinavian Journal of Urology and Nephrology* 1998, **32**(3):204-10. (Guideline Ref ID: WALDEN1998)
315. Wang ZL, Wang XF, Li B, Ji JT, Hou SC, Shao SX. Comparative study of transurethral electrovaporisation of prostate versus transurethral resection of prostate on benign prostatic hyperplasia. *Zhong hua nan ke xue* 2002, **8**(6):428-30. (Guideline Ref ID: WANG2002)
316. Wasson JH, Reda DJ, Bruskewitz RC, Elinson J, Keller AM, Henderson WG. A comparison of transurethral surgery with watchful waiting for moderate symptoms of benign prostatic hyperplasia. The Veterans Affairs Cooperative Study Group on Transurethral Resection of

- the Prostate. *New England Journal of Medicine* 1995, **332**(2):75-9. (Guideline Ref ID: WASSON1995)
317. Watson V, Ryan M, Brown CT, Barnett G, Ellis BW, Emberton M. Eliciting preferences for drug treatment of lower urinary tract symptoms associated with benign prostatic hyperplasia. *Journal of Urology* 2004, **172**(6 Pt 1):2321-5. (Guideline Ref ID: WATSON2004)
318. Westenberg A, Gilling P, Kennett K, Frampton C, Fraundorfer M. Holmium laser resection of the prostate versus transurethral resection of the prostate: results of a randomized trial with 4-year minimum long-term followup. *Journal of Urology* 2004, **172**(2):616-9. (Guideline Ref ID: WESTENBERG2004)
319. Wild D, Eremenco S, Mear I, Martin M, Houchin C, Gawlicki M *et al.* Multinational trials-recommendations on the translations required, approaches to using the same language in different countries, and the approaches to support pooling the data: the ISPOR Patient-Reported Outcomes Translation and Linguistic Validation Good Research Practices Task Force report. *Value Health* 2009, **12**(4):430-40. (Guideline Ref ID: WILD2009)
320. Wild D, Grove A, Martin M, Eremenco S, McElroy S, Verjee-Lorenz A *et al.* Principles of good practice for the translation and cultural adaptation process for Patient-Reported Outcomes (PRO) Measures: report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value Health* 2005, **8**(2):94-104. (Guideline Ref ID: WILD2005)
321. Wille S, Sobottka A, Heidenreich A, Hofmann R. Pelvic floor exercises, electrical stimulation and biofeedback after radical prostatectomy: results of a prospective randomized trial. *Journal of Urology* 2003, **170**(2 Pt 1):490-3. (Guideline Ref ID: WILLE2003)
322. Willetts KE, Clements MS, Champion S, Ehsman S, Eden JA. Serenoa repens extract for benign prostate hyperplasia: a randomized controlled trial. *BJU International* 2003, **92**(3):267-70. (Guideline Ref ID: WILLETTS2003)
323. Wilson LC, Gilling PJ, Williams A, Kennett KM, Frampton CM, Westenberg AM *et al.* A randomised trial comparing holmium laser enucleation versus transurethral resection in the treatment of prostates larger than 40 grams: results at 2 years. *European Urology* 2006, **50**(3):569-73. (Guideline Ref ID: WILSON2006)
324. Wilt TJ. Tamsulosin for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 2002, **Issue 4**:CD002081. (Guideline Ref ID: WILT2002)
325. Wilt TJ, Howe RW, Rukts I, MacDonald R. Terazosin for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 2000, **Issue 1**:CD003581. (Guideline Ref ID: WILT2000A)
326. Wilt TJ, Ishani A, MacDonald R. Serenoa repens for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 2002, **Issue 3**:CD001423. (Guideline Ref ID: WILT2002A)
327. Wilt TJ, Ishani A, MacDonald R, Rukts I, Stark G. Pygeum africanum for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 1998, **Issue 1**:CD001044. (Guideline Ref ID: WILT1998A)
328. Wilt TJ, Ishani A, MacDonald R, Stark G, Mulrow C, Lau J. Beta-sitosterols for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 1999, **Issue 3**:CD001043. (Guideline Ref ID: WILT1999)

329. Wilt TJ, MacDonald R, Ishani A, Ruktts I, Stark G. Cernilton for benign prostatic hyperplasia. *Cochrane Database of Systematic Reviews* 1998, **Issue 3**:CD001042. (Guideline Ref ID: WILT1998)
330. Xia SJ, Zhuo J, Sun XW, Han BM, Shao Y, Zhang YN. Thulium laser versus standard transurethral resection of the prostate: a randomized prospective trial. *European Urology* 2008, **53**(2):382-9. (Guideline Ref ID: XIA2008)
331. Zerbib M, Steg A, Conquy S, Debre B. Hyperthermia: a randomized prospective study applying hyperthermia or a sham procedure in obstructive benign hyperplasia of the prostate. *Progress in Clinical and Biological Research* 1994, **386**:439-48. (Guideline Ref ID: ZERBIB1994)
332. Zorn BH, Bauer JJ, Ruiz HE, Thrasher JB. Randomized trial of safety and efficacy of transurethral resection of the prostate using contact laser versus electrocautery. *Techniques in Urology* 1999, **5**(4):198-201. (Guideline Ref ID: ZORN1999)