

NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE

INTERVENTIONAL PROCEDURES PROGRAMME

Interventional procedure overview of radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

Interdigital (Morton's) neuroma affects one of the nerves leading to the toes. It can cause a burning pain in the ball of the foot and pain, tingling and numbness in the toes.

In this procedure, a thin probe is inserted at the base of one of the toes and into the affected nerve. An electric current passed through the probe destroys the nerve with radiofrequency heat energy.

Introduction

The National Institute for Health and Care Excellence (NICE) has prepared this interventional procedure (IP) overview to help members of the Interventional Procedures Advisory Committee (IPAC) make recommendations about the safety and efficacy of an interventional procedure. It is based on a rapid review of the medical literature and specialist opinion. It should not be regarded as a definitive assessment of the procedure.

Date prepared

This IP overview was prepared in 30 March 2015.

Procedure name

- Radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

Specialist societies

- British Orthopaedic Foot and Ankle Society – foot surgery
- The Society of Chiropractors and Podiatrists – surgical podiatry
- British Society of Skeletal Radiologists.

Description

Indications and current treatment

Symptomatic interdigital (Morton's) neuroma is caused by perineural fibrosis which creates scar tissue, resulting in compression of an interdigital nerve. It usually occurs between the metatarsal heads of the third and fourth toes but can sometimes occur between the second and third toes. Symptoms include severe intermittent pain, a burning sensation, and paraesthesia in the front part of the sole of the foot extending into the toes.

Initial management of symptomatic Morton's neuroma includes rest, anti-inflammatory medications, using an orthosis in the shoe and wearing a different type of shoe. Injection of steroids and local anaesthetic may be used. Persistent symptoms may be treated by cryoablation or surgical removal of the nerve (neurectomy).

What the procedure involves

Radiofrequency ablation (RFA) for symptomatic Morton's neuroma is a percutaneous treatment which is usually done as an outpatient procedure. A local anaesthetic block of the posterior tibial nerve is done by injecting it posterior to the medial malleolus under ultrasound guidance. Further local anaesthetic is injected around the neuroma. Using imaging guidance (X-ray fluoroscopy, ultrasound or MRI), an RFA probe attached to a generator is inserted into the web space between the toes and into the centre of the nerve. Controlled pulses of radiofrequency energy are delivered, which cause thermal ablation of the nerve. After the procedure, a steroid injection is usually given to reduce pain and inflammation. Patients are discharged as soon as comfortable and advised to limit their walking for 1 or 2 days. Any pain is managed with analgesics. The procedure can be repeated if necessary after a few weeks.

Literature review

Rapid review of literature

The medical literature was searched to identify studies and reviews relevant to radiofrequency ablation for symptomatic interdigital (Morton's) neuroma. The following databases were searched, covering the period from their start to 30-03-15: MEDLINE, PREMEDLINE, EMBASE, Cochrane Library and other databases. Trial registries and the Internet were also searched. No language restriction was applied to the searches (see appendix C for details of search strategy). Relevant published studies identified during consultation or resolution that are published after this date may also be considered for inclusion.

The following selection criteria (table 1) were applied to the abstracts identified by the literature search. Where selection criteria could not be determined from the abstracts the full paper was retrieved.

Table 1 Inclusion criteria for identification of relevant studies

Characteristic	Criteria
Publication type	Clinical studies were included. Emphasis was placed on identifying good quality studies. Abstracts were excluded where no clinical outcomes were reported, or where the paper was a review, editorial, or a laboratory or animal study. Conference abstracts were also excluded because of the difficulty of appraising study methodology, unless they reported specific adverse events that were not available in the published literature.
Patient	Patients with symptomatic interdigital (Morton's) neuroma.
Intervention/test	Radiofrequency ablation.
Outcome	Articles were retrieved if the abstract contained information relevant to the safety and/or efficacy.
Language	Non-English-language articles were excluded unless they were thought to add substantively to the English-language evidence base.

List of studies included in the IP overview

This IP overview is based on 162 patients from 4 case series.

Other studies that were considered to be relevant to the procedure but were not included in the main extraction table (table 2) have been listed in appendix A.

Table 2 Summary of key efficacy and safety findings on radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

Study 1 Finney W (1989)

Details

Study type	Retrospective case series
Country	USA (single centre)
Recruitment period	1977-86
Study population and number	n=71 patients (79 nerve lesions) 35% (28/79) presented in the 2 nd web space, 59% (47/79) in the 3 rd web space, 5% (4/79) in 4 th web space. Mean duration of symptoms: not reported.
Age and sex	mean 54 years; (67/71) female
Patient selection criteria	Patients with symptomatic Morton's neuroma treated previously (12 by surgical excision, 68 by a variety of pads, shoe alterations, orthotic devices, and cortisone injections).
Technique	Radiofrequency ablation (using Radionics nerve lesion generator) was performed. The foot is prepared and local anaesthesia injected proximally between the involved metatarsal necks and distally between the involved toes. An insulated or Teflon coated active electrode is passed dorsal to plantar, between the metatarsal necks and just inferior to them. An inactive electrode (used as grounding needle) is passed beneath the metatarsal ligament. The location is conformed through fluoroscopy or radiograph. The active electrode tip temperature is adjusted to 85°C and it is moved gently within the sagittal plane, dorsal to plantar and proximal to distal, until a pain response similar to neuroma is obtained. The temperature is maintained for 90 seconds. The symptoms resolve within 30–45 seconds until neurolysis occurs. 80% (57/71) patients had 1 procedure, 27% (19/71) had 2 and 3% (3/71) had 3 procedures.
Follow-up	Not reported.
Conflict of interest/source of funding	Not reported

Analysis

Follow-up issues: not clear.

Study design issues: small sample size, retrospective review of patients, patients responded to a mailed questionnaire. Pain responses were assessed by 3 different questions: pain assessment done with a 0–5 numerical pain intensity scale (0 being absence of pain and 5 being severe pain).

No ultrasound guidance was used during the procedure.

Study population issues: 41 were left feet and 38 were right feet. 8 patients had bilateral neuromas.

Key efficacy and safety findings

Efficacy	Safety								
<p>Number of patients analysed: 71 patients (79 nerve lesions)</p> <p>Pain score before and after treatment (on a pain scale 0-5)</p> <table border="1" data-bbox="94 306 941 457"> <thead> <tr> <th></th> <th>Mean pain rating</th> </tr> </thead> <tbody> <tr> <td>Preoperatively</td> <td>4.53</td> </tr> <tr> <td>Postoperatively</td> <td>2.02</td> </tr> <tr> <td>Difference*</td> <td>2.51 (or an average of 56% improvement in symptoms)</td> </tr> </tbody> </table> <p>*no statistical significance reported</p> <p>Symptom relief after treatment (rated as worse, same, improved or gone)</p> <p>71% (56/79) of the patients reported that their symptoms were improved or completely gone. 5% (4/79) patients stated their symptoms were worse and 24% (19/79) reported they were same.</p> <p>Satisfaction with treatment</p> <p>68% (54/71) patients said they were pleased with the treatment and 32% (25/71) were not pleased with the treatment. The patients with prior excisional surgery (n=12) responded similarly to RFA as those with no prior surgery (n=68).</p> <p>Absence from work: average 1.5 days (range 0-7 days).</p> <p>Postoperative pain medications were taken by 20% of the patients.</p>		Mean pain rating	Preoperatively	4.53	Postoperatively	2.02	Difference*	2.51 (or an average of 56% improvement in symptoms)	<p>Complications</p> <p>Burns: 2 patients incurred burns at the inactive (grounding) electrode site (as a result of the electrode being placed too superficially). These patients were absent from work for a week each.</p>
	Mean pain rating								
Preoperatively	4.53								
Postoperatively	2.02								
Difference*	2.51 (or an average of 56% improvement in symptoms)								
Abbreviations used: NPS, numerical pain score; RFA, radiofrequency ablation.									

Study 2 Genon M (2010)

Details

Study type	Retrospective case series (survey)
Country	Australia (single centre)
Recruitment period	2006-2008
Study population and number	n=37 patients (38 neuromas) 53% (20/38) presented in the 2 nd web space, 47% (18/38) in the 3 rd web space. Mean duration of symptoms: at least 12 months (in 28 patients)
Age and sex	Not reported
Patient selection criteria	Patients with symptomatic Morton's neuroma in whom conservative management failed
Technique	Radiofrequency ablation (using Neurotherm NT 500) was performed as a day procedure. Using an antiseptic technique, a guide needle with the insulating cannula was used to locate the neuroma. The needle was introduced from the dorsum of the appropriate inter-metatarsal space, 2-2.5cm proximal to the skin web fold. The needle was inserted deep to the transverse metatarsal ligament and adjusted to the point of maximal tenderness, assumed to be in the main body of the neuroma. Leaving the cannula <i>in situ</i> , the guide needle was removed and then an RFA electrode was inserted. Sedation was then administered. Current intensity increased until the temperature reached 90°C. The electrode was left in for 3 minutes to ablate the neuroma. After the procedure, local anaesthetic was injected. The patient was allowed to bear weight as tolerated and was discharged the same day.
Follow-up	average 10.6 months (range 3-21 months)
Conflict of interest/source of funding	None

Analysis

Follow-up issues: limited follow-up.

Study design issues: small sample size, retrospective review of patients, survey was undertaken. Pain assessment done with a 0–10 numerical pain intensity scale.

No ultrasound guidance was used during the procedure.

Study population issues: 22 were left feet and 16 were right feet.

Key efficacy and safety findings

Efficacy	Safety																															
<p>Number of patients analysed: 37 patients (38 neuromas)</p> <p>Resolution of symptoms % (n)</p> <table border="1" data-bbox="97 306 755 422"> <tr> <td>Complete relief</td> <td>18.4 (7/38)</td> </tr> <tr> <td>Partial relief</td> <td>55.3 (21/38)</td> </tr> <tr> <td>No relief</td> <td>26.3 (10/38)</td> </tr> </table> <p>Resolution of symptoms according to location of neuromas</p> <table border="1" data-bbox="97 489 755 632"> <thead> <tr> <th>Location of neuromas</th> <th>complete/partial relief % (n)</th> </tr> </thead> <tbody> <tr> <td>Second web space (n=20)</td> <td>50%</td> </tr> <tr> <td>Third web space (n=18)</td> <td>100%</td> </tr> </tbody> </table> <p>Patients who reported no relief had neuromas in 2nd web space. 84% of patients with 3rd web space neuromas were satisfied compared with 45% of those with neuromas in 2nd web space.</p> <p>Numerical pain scores (NPS)</p> <table border="1" data-bbox="97 800 902 1066"> <thead> <tr> <th></th> <th>pre-RFA</th> <th>post-RFA</th> </tr> </thead> <tbody> <tr> <td>Median NPS</td> <td>9.0 (8.0-9.0)</td> <td>5.0 (3.0-8.0)</td> </tr> <tr> <td>Median NPS in patients with complete relief (n=7)</td> <td>9.0</td> <td>2.0</td> </tr> <tr> <td>Median NPS in patients with partial relief (n=21)</td> <td>9.0</td> <td>5.0</td> </tr> <tr> <td>Median NPS in patients with no relief (n=10)</td> <td>9.0</td> <td>9.0</td> </tr> </tbody> </table> <p>Patient satisfaction % (n)</p> <table border="1" data-bbox="97 1136 755 1209"> <tr> <td>Satisfied</td> <td>86.8 (32/37)</td> </tr> <tr> <td>Not satisfied</td> <td>13.2 (5/37)</td> </tr> </table> <p>83.7% patients said they would have the procedure again.</p> <p>Repeat RFA treatment</p> <p>2 patients with no relief had repeat treatment but were not satisfied with the outcome. No further treatment was provided to these patients.</p> <p>Progressing to surgical excision</p> <p>Overall, 29% (11/38) patients (3 with partial relief and 8 with no relief) underwent surgical excision of the neuromas. The time to surgery was not reported. 6 had complete relief, 3 had partial relief, 1 had no change in symptoms and 1 got worse. The average NPS decreased from 6.9 to 2.7.</p>	Complete relief	18.4 (7/38)	Partial relief	55.3 (21/38)	No relief	26.3 (10/38)	Location of neuromas	complete/partial relief % (n)	Second web space (n=20)	50%	Third web space (n=18)	100%		pre-RFA	post-RFA	Median NPS	9.0 (8.0-9.0)	5.0 (3.0-8.0)	Median NPS in patients with complete relief (n=7)	9.0	2.0	Median NPS in patients with partial relief (n=21)	9.0	5.0	Median NPS in patients with no relief (n=10)	9.0	9.0	Satisfied	86.8 (32/37)	Not satisfied	13.2 (5/37)	<p>Complications</p> <p>None related to RFA.</p>
Complete relief	18.4 (7/38)																															
Partial relief	55.3 (21/38)																															
No relief	26.3 (10/38)																															
Location of neuromas	complete/partial relief % (n)																															
Second web space (n=20)	50%																															
Third web space (n=18)	100%																															
	pre-RFA	post-RFA																														
Median NPS	9.0 (8.0-9.0)	5.0 (3.0-8.0)																														
Median NPS in patients with complete relief (n=7)	9.0	2.0																														
Median NPS in patients with partial relief (n=21)	9.0	5.0																														
Median NPS in patients with no relief (n=10)	9.0	9.0																														
Satisfied	86.8 (32/37)																															
Not satisfied	13.2 (5/37)																															
Abbreviations used: NPS, numerical pain score; RFA, radiofrequency ablation.																																

Study 3 Moore JL (2012)

Details

Study type	Retrospective case series
Country	USA (single centre)
Recruitment period	2007-10
Study population and number	n=29 patients (32 feet) All neuromas located within the 2 nd or 3 rd intermetatarsal space.
Age and sex	age range 23-73 years; 76% (22/29) female
Patient selection criteria	Patients with symptomatic neuroma pain of the foot not relieved by routine conservative treatments (injections for 4-8 weeks), diagnosed by clinical presentation and physical examination. There were no inclusion or exclusion criteria in the study.
Technique	Maximum point of tenderness was marked and anaesthesia was first administered, then a sharp tip probe was inserted from dorsal to plantar at the area of maximum pain under fluoroscopy. This was replaced with an RFA electrode, and the radiofrequency thermoneurolysis therapy process initiated at a temperature of 85°C for 90 seconds. The probe was removed and 4 mg dexamethasone was injected into the surgical site and a bandage applied. Patients were advised to bear weight as tolerated, use anti-inflammatory drugs if needed and followed up at regular intervals.
Follow-up	Average 13 months (range 6 months to 3 years)
Conflict of interest/source of funding	None

Analysis

Follow-up issues: 2 patients were lost to follow-up after 1 month.

Study design issues: Small sample size, retrospective study. Final pain assessment was done at 6 months; none of the patients were using anti-inflammatory drugs.

Study population issues: 3 patients had neuroma pain bilaterally. 1 patient had type 2 diabetes. 4 patients were unable to accurately pinpoint the area of pain and a second RFA procedure was done distally to assure pain relief.

Key efficacy and safety findings

Efficacy	Safety
<p>Number of patients analysed: 29 patients (20 feet)</p> <p>Symptom relief (rated as some relief, complete relief or no relief) 83% (24/29) patients reported complete relief of symptoms (return to activity with no pain or disability) 1 month after RFA treatment. 17% (5/29) patients had minimal to no relief of symptoms (1 underwent open surgery, 2 were lost to follow-up and 2 were satisfied with minimal relief).</p> <p>Recurrence 1 patient reported recurrence at 9 months, which was successfully treated with a combination of anaesthetic and steroid injection (bupivacaine/triamcinolone).</p> <p>Return to normal activities All patients returned to normal shoes and activities within 2 days.</p> <p>Abbreviations used: RFA, radiofrequency ablation.</p>	<p>Complications Superficial cellulitis (after 5 days, treated with a week's course of antibiotics) in 1 patient.</p>

Study 4 Chuter GSJ (2013)

Details

Study type	Retrospective case series
Country	Australia (single centre)
Recruitment period	January –September 2011
Study population and number	n= 25 patients (30 feet) 40% (12/30) presented in the 2 nd web space, 60% (18/30) in the 3 rd web space. Mean duration of symptoms: 3.8 years (range 6 months to 15 years) Size of neuromas: average 10.7 mm (range 4-21mm)
Age and sex	Mean 55 years; 84% (21/25) female
Patient selection criteria	Patients with Morton's neuroma (confirmed by MRI, ultrasound, or clinical history and examination), persistent or recurrent symptoms, presented for neurectomy but referred for radiofrequency ablation, tried previous methods of conservative management (80% orthotics, 60% cortisone injections). Patients with previous excision or recurrence (e.g. stump neuromas) and 1 patient with neurofibromatosis were excluded.
Technique	Ultrasound-guided radiofrequency ablation (using Neurotherm NT 500) of Morton's neuroma performed by a single radiologist under local anaesthetic in an outpatient clinic. 5 cycles of 2 minutes each were used; with the probe tip maintain a temperature of 81°C. Steroid injection was used at the time of procedure. Treatment was repeated after 4 weeks if needed. Average number of treatment sessions was 1.6 (range 1-3). Patients were reviewed 4 weeks later with a follow-up ultrasound scan and RFA was repeated if still symptomatic. If unsatisfied after 3 treatment sessions, they were referred back to clinic for further management (surgical excision).
Follow-up	6 months
Conflict of interest/source of funding	None

Analysis

Follow-up issues: 6 months follow-up.

Study design issues: Small sample size.

Study population issues: only patients with primary Morton's neuroma were included. There were 19 left feet and 11 right feet. All patients had neuromas confirmed at the time of treatment ('focal well circumscribed anechoic lesion in the interdigital web space'). 8 feet had additional neuromas in the adjacent web space (2nd or 3rd) treated with radiofrequency ablation. 3 patients had asymptomatic bilateral bunions (hallux vagus) of which 2 had bilateral symptomatic neuromas.

Key efficacy and safety findings

Efficacy	Safety								
<p>Number of patients analysed: 25 patients (30 feet)</p> <p>Change in visual analogue pain scores (measured on a VAS scale 0-10)*</p> <table border="1" data-bbox="94 338 837 501"> <thead> <tr> <th></th> <th>Pre-treatment</th> <th>6 months post-treatment</th> <th>p value</th> </tr> </thead> <tbody> <tr> <td>VAS pain score on activity mean (range)</td> <td>6.0 (3-9)</td> <td>1.7 (0-8)</td> <td><0.001</td> </tr> </tbody> </table> <p>Only 4 patients at pain at rest prior to RFA and all had pain on activity. *higher score representing worst ever pain.</p> <p>There was no statistically significant difference in the mean reduction in VAS between 2nd (9 feet) and 3rd web space (13 feet) neuromas (4.3 versus 3.3, p=0.299).</p> <p>Symptom improvement</p> <p>The average overall symptom improvement (defined as 'as described by the patient', no further explanation given) was 76% (range 0-100%).</p> <p>Satisfactory outcome:</p> <p>At 6 months 87% (26/30) of feet had a satisfactory outcome (not defined in the paper) without the need for further surgery.</p> <p>No statistically significant correlation was found between pain or outcome, and any of the following: age, gender, site, size, number, symptom duration, activity level or previous treatment.</p> <p>32% (8/25) patients described the experience as unpleasant but would have the procedure again as first line treatment. Those who had surgery preferred RFA as first-line treatment.</p> <p>Progression to surgical excision: 10% (3/30). 1 patient had no improvement, 1 had 40% improvement but wanted surgery, and 1 had 100% improvement initially but had recurrence within 6 months. Subsequent surgery resolved symptoms in all.</p>		Pre-treatment	6 months post-treatment	p value	VAS pain score on activity mean (range)	6.0 (3-9)	1.7 (0-8)	<0.001	<p>Complications</p> <p>Temporary posterior tibial nerve irritation (for 3 weeks, resolved): 1</p> <p>Ongoing unchanged pain (no obvious cause): 1</p>
	Pre-treatment	6 months post-treatment	p value						
VAS pain score on activity mean (range)	6.0 (3-9)	1.7 (0-8)	<0.001						
Abbreviations used: VAS, visual analogue score; RFA, radiofrequency ablation.									

Efficacy

Pain scores

A case series of 25 patients (30 feet) with symptomatic interdigital (Morton's) neuroma reported a statistically significant reduction in pain scores on activity after ultrasound guided radiofrequency ablation (RFA) treatment. Pain scores were measured on a visual analogue scale (assessed on a scale of 0–10, with lower scores indicating less pain) and were an average 6.0 at baseline compared with 1.7 at 6-month follow-up ($p < 0.001$)⁴.

A case series of 37 patients (38 neuromas) for whom conservative management failed and who had RFA reported median numerical pain scores (assessed on a scale of 0–10, with lower scores indicating less pain). Pain scores decreased from 9.0 at baseline to 5.0 at an average follow-up of 10.6 months. In those who reported complete resolution of symptoms ($n=7$) and partial relief ($n=21$) the median numerical pain score reduced from 9.0 to 5.0. In those who had no relief ($n=10$) it did not change².

Symptom improvement

The case series of 25 patients reported that the average overall symptom improvement (as described by patients, not otherwise defined) was 76%⁴.

The case series of 37 patients (38 neuromas) reported that 74% of neuromas had complete or partial resolution of symptoms and 26% had no benefit from RFA treatment at an average follow-up of 10.6 months. All patients with neuromas in the third web space ($n=18$) reported complete or partial relief of symptoms compared with only 50% of those with second web space neuromas ($n=20$)².

A case series of 29 patients (32 feet) reported that 83% (24/29) patients had complete relief of symptoms at 1 month after RFA treatment and 17% (5/29) had minimal to no relief of symptoms³.

Repeat RFA treatment

The case series of 37 patients reported that 2 patients with no relief had repeat RFA treatment but were not satisfied with the outcome at an average follow-up of 10.6 months. No further treatment was offered².

Patient satisfaction

The case series of 37 patients reported that 87% (32/37) of patients were satisfied with RFA treatment, at an average follow-up of 10.6 months. Most patients (84%) said that they would have the procedure again².

The case series of 25 patients reported that at 6 months 87% (26/30) of feet had a satisfactory outcome (not defined) without the need for further surgery. It was reported that 32% (8/25) of patients described the experience as unpleasant but would have the procedure again as first-line treatment⁴.

The case series of 71 patients reported that 68% (54/71) of the patients were satisfied with RFA treatment and 32% (25/71) were not satisfied with RFA treatment¹.

Recurrence

The case series of 29 patients (32 feet) reported symptom recurrence in 1 patient at 9 months follow-up. This was successfully treated with an injection of steroid and local anaesthetic³.

Progression to surgical removal

Progression to surgical removal of the neuromas was reported in 29% (11/38) of patients (3 with partial relief and 8 with no relief) in the case series of 37 patients (38 neuromas) at an average follow-up of 10.6 months. Of the patients who had surgical removal, 6 patients had complete relief of symptoms, 3 had partial relief, 1 had no change in symptoms and 1 got worse. The average numerical pain score decreased from 6.9 to 2.7².

Surgical removal was reported in 10% (3/30) neuromas after the procedure in a case series of 25 patients. One patient had no improvement after RFA, 1 had 40% improvement but wanted surgery, 1 had 100% improvement initially but symptoms recurred within 6 months. Subsequent surgery resolved symptoms in all⁴.

Safety

Pain

Ongoing, unchanged pain after RFA treatment was reported in 1 patient in the case series of 25 patients. Further investigations found no obvious cause for his symptoms⁴.

Superficial cellulitis

Superficial cellulitis 5 days after RFA treatment was reported in 1 patient in a case series of 29 patients. This was treated with a course of antibiotics⁴.

Burns

Burns at the site of the inactive (grounding) electrode (explained by the authors as a result of the electrode being placed too superficially) were reported in

2 patients in a case series of 71 patients. These patients were each off work for a week⁴.

Temporary nerve irritation

Irritation of the posterior tibial nerve for 3 weeks after the procedure was reported in 1 patient in a case series of 25 patients. This resolved completely⁴.

Validity and generalisability of the studies

- There are no randomised controlled trials. Only 4 small retrospective case series have been published on this procedure.
- Most of the patients were women.
- One study used ultrasound and steroid injection in the treatment protocol⁴.
- There was a lack of long-term follow-up in the included studies.

Existing assessments of this procedure

There were no published assessments from other organisations identified at the time of the literature search.

Related NICE guidance

There is currently no NICE guidance related to this procedure.

Specialist advisers' opinions

Specialist advice was sought from consultants who have been nominated or ratified by their Specialist Society or Royal College. The advice received is their individual opinion and is not intended to represent the view of the society. The advice provided by Specialist Advisers, in the form of the completed questionnaires, is normally published in full on the NICE website during public consultation, except in circumstances but not limited to where comments are considered voluminous, or publication would be unlawful or inappropriate. Eleven Specialist Advisor Questionnaires for radiofrequency ablation for symptomatic interdigital (Morton's) neuroma were submitted and can be found on the [NICE website \[INSERT HYPER LINK TO MAIN IP PAGE\]](#).

Patient commentators' opinions

NICE's Public Involvement Programme sent **xxx** questionnaires to **xxx** NHS trusts for distribution to patients who had the procedure (or their carers). NICE received **xxx** completed questionnaires.

Section to be inserted if there is no patient commentary

NICE's Public Involvement Programme was unable to gather patient commentary for this procedure.

Section to be inserted if patient commentators raised no new issues

The patient commentators' views on the procedure were consistent with the published evidence and the opinions of the specialist advisers.

Section to be inserted if patient commentators raised new issues

The patient commentators raised the following issues about the safety/efficacy of the procedure which did not feature in the published evidence or the opinions of specialist advisers, and which the Committee considered to be particularly relevant:

- [insert additional efficacy and safety issues raised by patient commentators and highlighted by IPAC, add extra rows as necessary].
- [Last item in list].

Issues for consideration by IPAC

Studies on the use of radiofrequency ablation for symptomatic interdigital (Morton's) neuroma are limited.

References

1. Finney W et al (1989). Treatment of Morton's neuroma using percutaneous electrocoagulation. *Journal of the American Podiatric Medical Association*. 79: 615-8.
2. Genon MP, Chin TY, Bedi HS and Blackney MC (2010). Radio-frequency ablation for the treatment of Morton's neuroma. [Review]. *ANZ Journal of Surgery* 80 (9) 583-585.
3. Moore JL, Rosen R, Cohen J, and Rosen B (2012). Radiofrequency thermoneurolysis for the treatment of Morton's neuroma. *Journal of Foot & Ankle Surgery* 51: 20-22.
4. Chuter GS, Chua YP, Connell DA, and Blackney MC (2013). Ultrasound-guided radiofrequency ablation in the management of interdigital (Morton's) neuroma. *Skeletal Radiology* 42 (1) 107-111.

Appendix A: Additional papers on radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

The following table outlines the studies that are considered potentially relevant to the IP overview but were not included in the main data extraction table (table 2). It is by no means an exhaustive list of potentially relevant studies.

Article	Number of patients/follow-up	Direction of conclusions	Reasons for non-inclusion in table 2
Bregman PJ (2012). Radiofrequency thermoneurolysis for the treatment of Morton's neuroma. <i>Journal of Foot & Ankle Surgery</i> 51 (3) 405.			Letter in response to article.
Gobbi, F., Mazzetti, R., et al (2014). Ultrasound guided radiofrequency ablation of Morton's neuroma: Preliminary results. <i>Regional Anesthesia and Pain Medicine</i> . Conference: 33rd Annual European Society of Regional Anaesthesia and Pain Therapy, ESRA Congress 2014 Seville Spain. Conference Start: 20140903 Conference End: 20140906. Conference Publication: (var.pagings).39 (5 S (var.pagings) e302-e303.	Prospective cohort study 2013 n=16 (20 feet) patients Ultra-sound guided radiofrequency ablation Follow-up: 6 months	No post-procedure complications. At the end of the procedure (about 15 minutes) patients could immediately walk and go back to their daily activities. In 3 cases it was necessary to repeat the RF treatment. Dysesthesias reported by 75% of patients in the area of the treated nerve regressed on average after 4 weeks. Pre-treatment pain was present in 100% of cases. Based on preliminary data currently available, 75% of patients reported a virtually full reduction of pain after the RF treatment, 16% a very limited reduction, 9% no reduction. Conclusions: The preliminary study of ultra-sound guided radiofrequency ablation of MN shows a significant regression to disappearance of the pain symptoms. Available data on the duration of the treatment efficacy show a good short-medium term outcome (1-3 months) and are encouraging on the medium-long term (6 months).	Conference abstract.

Appendix B: Related NICE guidance for radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

There is currently no NICE guidance related to this procedure.

Appendix C: Literature search for radiofrequency ablation for symptomatic interdigital (Morton's) neuroma

Databases	Date searched	Version/files
Cochrane Database of Systematic Reviews – CDSR (Cochrane)	30/03/2015	Issue 3 of 12, March 2015
HTA database (Cochrane)	30/03/2015	Issue 1 of 4, January 2015
Cochrane Central Database of Controlled Trials – CENTRAL (Cochrane)	30/03/2015	Issue 2 of 12, February 2015
MEDLINE (Ovid)	30/03/2015	1946 to March week 4 2015
MEDLINE In-Process (Ovid)	30/03/2015	March 27, 2015
EMBASE (Ovid)	30/03/2015	1974 to 2015 March 27
PubMed	30/03/2015	n/a
BLIC (Dialog DataStar)	30/03/2015	n/a

Trial sources searched on

- Clinicaltrials.gov
- ISRCTN
- WHO International Clinical Trials Registry

Websites searched on

- National Institute for Health and Care Excellence (NICE)
- NHS England
- Food and Drug Administration (FDA) - MAUDE database
- Australian Safety and Efficacy Register of New Interventional Procedures – Surgical (ASERNIP – S)
- Australia and New Zealand Horizon Scanning Network (ANZHSN)
- EuroScan
- General internet search

The following search strategy was used to identify papers in MEDLINE. A similar strategy was used to identify papers in other databases.

- 1 ablation techniques/ (1014)
- 2 denervation/ (13141)
- 3 catheter ablation/ (22380)
- 4 radio waves/ (5740)
- 5 ((radiofrequen* or radio-frequen* or "radio frequen*") adj4 (denervat* or ablat*)).ti,ab. (12251)
- 6 (RFA or (rf adj4 ablat*)).ti,ab. (5394)

- 7 ((catheter* or prob* or needle* or electro* or heat*) adj4 (denervat* or ablat*)).ti,ab. (8971)
- 8 ((ultrasound* or ultra-sound*) adj4 guid*).ti,ab. (15543)
- 9 or/1-8 (61013)
- 10 Foot/ (20503)
- 11 nerve compression syndromes/ (9190)
- 12 Metatarsal Bones/ (2665)
- 13 Neuroma/ (2399)
- 14 Peripheral Nervous System Neoplasms/ (4012)
- 15 ((morton* or plantar* or interdigital* or inter-digital* or inter-metatarsal* or intermetatarsal*) adj4 (neurectom* or syndrome* or neuroma* or entrap* or neuralgi* or neuritis* or metatarsalgi*)).ti,ab. (620)
- 16 (metatarsophalangeal* adj4 (joint* or articulat*) adj4 (nerve* or compress* or pain* or numb* or burn* or sore* or swell* or discomfort* or ache* or dysesthesia* or tender* or tingl* or cramp*)).ti,ab. (109)
- 17 ((toe* or foot* or feet* or meta-tarsal* or metatarsal* or tarsal* or (plantar* adj4 digital*)) adj4 (nerve* or compress* or pain* or numb* or burn* or sore* or swell* or discomfort* or ache* or dysesthesia* or tender* or tingl* or cramp*)).ti,ab. (5495)
- 18 or/10-17 (42380)
- 19 9 and 18 (269)
- 20 animal/ not human/ (3909440)
- 21 19 not 20 (173)