

## Falls in older adults

### NICE guideline: methods

*NICE guideline <number>*

*Methods*

*October 2024*

*Draft for Consultation*

*Developed by the NICE Guideline  
Development Team*



## **Disclaimer**

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

## **Copyright**

© NICE, 2024. All rights reserved. Subject to Notice of rights.

# Contents

<b>1</b>	<b>Development of the guideline</b>	<b>5</b>
1.1	Remit	5
<b>2</b>	<b>Methods</b>	<b>6</b>
2.1	Developing the review questions and outcomes	6
2.1.1	Stratification	9
2.2	Searching for evidence	9
2.2.1	Clinical and health economics literature searches	9
2.3	Reviewing evidence	10
2.3.1	Types of studies and inclusion and exclusion criteria	11
2.4	Methods of combining evidence	12
2.4.1	Data synthesis for intervention reviews	12
2.4.2	Data synthesis for qualitative reviews	15
2.5	Appraising the quality of evidence by outcomes	15
2.5.1	Intervention reviews	15
2.5.2	Prediction rules/models	21
2.5.3	Qualitative reviews	22
2.5.4	Publication bias	24
2.6	Assessing clinical importance	24
2.7	Identifying and analysing evidence of cost effectiveness	25
2.7.1	Literature review	25
2.7.2	Undertaking new health economic analysis	27
2.7.3	Cost-effectiveness criteria	28
2.7.4	In the absence of health economic evidence	28
2.8	Developing recommendations	29
2.8.1	Research recommendations	30
2.8.2	Validation process	30
2.8.3	Updating the guideline	30
2.9	General terms	30
2.10	Clinical terms used in the guideline	42
	References	43

# 1 Development of the guideline

## 1.1 Remit

NICE received the remit for this guideline from NHS England.

The remit for this guideline is to fully update the guideline on: Falls in older people: assessing risk and prevention.

To see what this guideline covers and what this guideline does not cover, please see the guideline scope for the Falls in older people guideline.

## 2 Methods

This guideline was developed using the methods described in the 2014 NICE guidelines manual, updated 2020<sup>9</sup>

Declarations of interest were recorded according to the NICE conflicts of interest policy.

Sections 2.1 to 2.3 describe the process used to identify and review evidence. Sections 2.1.1 and 2.7 describe the process used to identify and review the health economic evidence.

### 2.1 Developing the review questions and outcomes

The review questions developed for this guideline were based on the key areas and draft review questions identified in the guideline scope. They were drafted by the technical team, refined and validated by the committee and signed off by NICE. A total of 9 review questions were developed in this guideline and outlined in Table 1.

The review questions were based on the following frameworks:

- population, intervention, comparator and outcome (PICO) for reviews of interventions (including test and treat)
- population, tests, and target condition for reviews of risk prediction test accuracy
- population, setting and context for qualitative reviews.

This use of a framework informed a more detailed protocol that guided the literature searching process, critical appraisal and synthesis of evidence, and facilitated the development of recommendations by the guideline committee. Full literature searches, critical appraisals and evidence reviews were completed for all the specified review questions.

**Table 1: Review questions**

Evidence report	Type of review	Review questions	Outcomes
A	Qualitative 1.1	What are the education and information needs (regarding prevention) of people after being identified and assessed to be at risk of falls, or had a fall?	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <p>Themes will be derived from the evidence identified for this review and not pre-specified.</p> <p>For information to guide the technical team, relevant themes may include:</p> <ul style="list-style-type: none"> <li>• Risk factors for falls and fall-related injuries.</li> <li>• Information on prevention interventions</li> <li>• The impact of fear of falling / concerns about falling</li> <li>• Getting up from the floor after a fall</li> </ul>

Evidence report	Type of review	Review questions	Outcomes
			<p>Themes that will not be covered by the evidence review but which can be found in other NICE guidance:</p> <ul style="list-style-type: none"> <li>• Accessing information/signposting to services</li> <li>• Self-management (including when to ask for help, condition-specific advice)</li> <li>• Social prescribing</li> <li>• Patient choice</li> </ul>
B	Prognostic 2.1	How accurate are assessments by clinicians, such as questioning, observation and examination, in identifying people at risk of falls?	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <p>Accuracy of estimation of risk of falls:</p> <p>Statistical outputs may include:</p> <ul style="list-style-type: none"> <li>• Discrimination (sensitivity, specificity, predictive values)</li> <li>• Area under the ROC curve (c-statistic, c-index)</li> <li>• Predicted risk versus observed risk (calibration)</li> <li>• Reclassification</li> </ul> <p>Other statistical measures: for example, D statistic, R2 statistic and Brier points</p>
C	Prognostic 2.2	How accurate are screening tools which quantify or categorise the degree of risk of falling in identifying people at risk of falls?	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <p>Accuracy of estimation of risk of falls:</p> <p>Statistical outputs may include:</p> <ul style="list-style-type: none"> <li>• Discrimination (sensitivity, specificity, predictive values)</li> <li>• Area under the ROC curve (c-statistic)</li> <li>• Predicted risk versus observed risk (calibration)</li> <li>• Reclassification</li> </ul> <p>Other statistical measures: for example, D statistic, R2 statistic and Brier points</p>
D	Prognostic 2.3	How accurate are electronic patient records for identifying people at risk of falls?	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <p>Accuracy of estimation of risk of falls:</p> <p>Statistical outputs may include:</p>

Evidence report	Type of review	Review questions	Outcomes
			<ul style="list-style-type: none"> <li>• Discrimination (sensitivity, specificity, predictive values)</li> <li>• Area under the ROC curve (c-statistic, c-index)</li> <li>• Predicted risk versus observed risk (calibration)</li> <li>• Reclassification</li> </ul> <p>Other statistical measures: for example, D statistic, R2 statistic and Brier points</p>
E	Prognostic 3.1	What methods of assessment are most accurate for identifying individual risk factors for risk of falls?	<p>For balance and gait assessment tools and wearable technology: All outcomes are considered equally important for decision making and therefore have all been rated as critical: Accuracy of estimation of risk of falls: Statistical outputs may include:</p> <ul style="list-style-type: none"> <li>• Discrimination (sensitivity, specificity, predictive values)</li> <li>• Area under the ROC curve (c-statistic)</li> <li>• Predicted risk versus observed risk (calibration)</li> <li>• Reclassification</li> </ul> <p>Other statistical measures: for example, D statistic, R2 statistic and Brier points.</p>
F	Intervention 4.1a	What are the most clinically effective and cost-effective methods for falls prevention in older people in: Community Hospital care settings.	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <ul style="list-style-type: none"> <li>• Rate of falls</li> <li>• Number of people sustaining one or more falls</li> <li>• Number of participants sustaining fall-related fractures</li> <li>• Adverse effects of the interventions (composite of all)</li> <li>• Validated health-related quality of life scores e.g. EQ-5D or similar.</li> </ul>
G	Intervention 4.1b	What are the most clinically effective and cost-effective methods for falls prevention in older people in: Hospital care settings.	<p>All outcomes are considered equally important for decision making and therefore have all been rated as critical:</p> <ul style="list-style-type: none"> <li>• Rate of falls</li> <li>• Number of people sustaining one or more falls</li> <li>• Number of participants sustaining fall-related fractures</li> <li>• Adverse effects of the interventions (composite of all)</li> <li>• Validated health-related quality of life scores e.g. EQ-5D or similar.</li> </ul>



Evidence report	Type of review	Review questions	Outcomes
H	Intervention 4.1c	What are the most clinically effective and cost-effective methods for falls prevention in older people in Residential care settings?	All outcomes are considered equally important for decision making and therefore have all been rated as critical: <ul style="list-style-type: none"> <li>• Rate of falls</li> <li>• Number of people sustaining one or more falls</li> <li>• Number of participants sustaining fall-related fractures</li> <li>• Adverse events of the interventions (composite of all)</li> <li>• Validated health-related quality of life scores.</li> </ul>
I	Intervention 4.3	What are the best methods for maximising participation, adherence, and continuation of falls prevention interventions?	All outcomes are considered equally important for decision making and therefore have all been rated as critical: <ul style="list-style-type: none"> <li>• Reduction in falls</li> <li>• Increased participation</li> <li>• Increased adherence</li> <li>• Continuation through follow-up period</li> <li>• Refusal/ non-response/ drop-out rate</li> </ul>

### 1      2.1.1 Stratification

2      In this guideline all reviews were stratified according to setting, of hospital, residential  
3      care and community, depending on where the person was situated. Only people  
4      aged 65 and over were included unless the population was aged 50 or over and had  
5      conditions that made them at high risk of falling, throughout the guideline these two  
6      populations were stratified. Within the intervention reviews where people were not  
7      aged 65 or over, we included studies where the mean age was 65 years or over. The  
8      Cochrane reviews reported in the guideline included those 60 years or over minus  
9      the standard deviation were over 60 years. We have included all the included  
10     Cochrane studies as they all had a mean age of 65 years or over.

## 11     2.2 Searching for evidence

### 12     2.2.1 Clinical and health economics literature searches

13     The full strategy including population terms, intervention terms, study types applied,  
14     the databases searched, and the years covered can be found in Appendix B of the  
15     evidence review.

16     Systematic literature searches were undertaken to identify published clinical and  
17     health economic evidence relevant to the review questions. These were run  
18     according to the parameters as stipulated within the NICE guideline's manual,  
19     <https://www.nice.org.uk/process/pmg20/chapter/identifying-the-evidence-literature-searching-and-evidence-submission>.  
20    

21     Databases were searched using relevant medical subject headings, free-text terms  
22     and where appropriate study-type filters. Studies published in languages other than

English were not reviewed, and where possible, searches were restricted to English language. Papers published or added to databases after this date were not considered. Where new evidence was identified, for example in consultation comments received from stakeholders, the impact on the guideline was considered, and the action agreed between the technical team and NICE staff with a quality assurance role. A Cochrane review (Dahota 2024), relevant to the Interventions for prevention of falls in the community review (psychological and educational interventions), was noted in the text because it will be published prior to this guideline.

Searches were quality assured using different approaches prior to being run. Medline search strategies were peer reviewed by a second information specialist using a QA process based on the PRESS checklist<sup>8</sup>. Key (seed) papers if provided, were checked if retrieved by the search.

Searching for unpublished literature was not undertaken. NICE do not have access to drug manufacturers' unpublished clinical trial results, so the clinical evidence considered by the committee for pharmaceutical interventions may be different from that considered by the MHRA and European Medicines Agency for the purposes of licensing and safety regulation.

Additional studies were added to the evidence base these consisted of references included in relevant systematic reviews, and those highlighted by committee members.

During the scoping stage, a search was conducted for guidelines and reports on the websites including:

- Guidelines International Network database ([www.g-i-n.net](http://www.g-i-n.net))
- National Guideline Clearing House ([www.guideline.gov](http://www.guideline.gov))
- National Institute for Health and Care Excellence (NICE) ([www.nice.org.uk](http://www.nice.org.uk))
- National Institutes of Health Consensus Development Program ([consensus.nih.gov](http://consensus.nih.gov))

## 2.3 Reviewing evidence

The evidence for each review question was reviewed using the following process:

- Potentially relevant studies were identified from the search results by reviewing titles and abstracts. The full papers were then obtained.
- Full papers were evaluated against the pre-specified inclusion and exclusion criteria set out in the protocol to identify studies that addressed the review question. The review protocols are included in an appendix to each of the evidence reports.
- Relevant studies were critically appraised using the preferred study design checklist as specified in the NICE guidelines manual.<sup>9</sup> The checklist used is included in the individual review protocols in each of the evidence reports.
- Key information was extracted about interventional study methods and results into EPPI reviewer version 5. Summary evidence tables were produced from data entered into EPPI Reviewer, including critical appraisal ratings. Key information about qualitative study methods and results were manually extracted into standard Word evidence tables (evidence tables are included in an appendix to each of the evidence reports).

- 1 • Summaries of the evidence were generated by outcome. Outcome data were  
2 combined, analysed and reported according to study design:
    - 3 ○ Randomised data were meta-analysed where appropriate and reported in  
4 GRADE evidence profiles.
    - 5 ○ Data from non-randomised studies were meta-analysed where appropriate and  
6 reported in GRADE evidence profiles.
    - 7 ○ Risk prediction tool data were meta-analysed where appropriate or presented  
8 as a range of values in GRADE evidence profiles.
    - 9 ○ Qualitative data were synthesised across studies using thematic analysis and  
10 presented as summary statements in GRADE CERQual tables.
  - 11 • A minimum of 10% of the abstracts were reviewed by two reviewers, with any  
12 disagreements resolved by discussion or, if necessary, a third independent  
13 reviewer.
  - 14 • All of the evidence reviews were quality assured by a senior systematic reviewer.  
15 This included checking:
    - 16 ○ papers were included or excluded appropriately
    - 17 ○ a sample of the data extractions
    - 18 ○ a sample of the risk of bias assessments
    - 19 ○ correct methods were used to synthesise data.
- 20 Discrepancies will be identified and resolved through discussion (with a third  
21 reviewer where necessary).

### 22 **2.3.1 Types of studies and inclusion and exclusion criteria**

23 The inclusion and exclusion of studies was based on the criteria defined in the review  
24 protocols, which can be found in an appendix to each of the evidence reports.  
25 Excluded studies (with the reasons for their exclusion) are listed in an appendix to  
26 each of the evidence reports. The committee was consulted about any uncertainty  
27 regarding inclusion or exclusion.

28 Conference abstracts were not generally considered for inclusion. If abstracts were  
29 included the authors were contacted for further information. Literature reviews,  
30 posters, letters, editorials, comment articles, unpublished studies and studies not in  
31 published in English language were excluded.

#### 32 **2.3.1.1 Type of studies**

33 Randomised controlled trials, non-randomised intervention studies, and other  
34 observational studies (including risk prediction tool studies) were included in the  
35 evidence reviews as appropriate.

36 For intervention reviews, randomised controlled trials (RCTs) were included where  
37 identified as because they are considered the most robust type of study design that  
38 can produce an unbiased estimate of the intervention effects. Non-randomised  
39 intervention studies were not considered appropriate for inclusion for intervention  
40 reviews because there was sufficient randomised evidence for the committee to  
41 make a decision. Refer to the review protocols in each evidence report for full details  
42 on the study design of studies that were appropriate for each review question.

43 For risk prediction review questions, prospective and retrospective cohort studies  
44 were included. Case-control studies were not included. For the risk prediction tool

1 reviews where a lot of evidence was expected, we limited to prospective cohort  
2 studies which were externally validated with a minimum of 100 participants included  
3 to ensure that the highest quality studies were included.

4 Systematic reviews and meta-analyses conducted to the same methodological  
5 standards as the NICE reviews were included within the evidence reviews in  
6 preference to primary studies, where they were available and applicable to the review  
7 questions and updated or added to where appropriate to the guideline review  
8 question. Individual patient data (IPD) meta-analyses were preferentially included if  
9 meeting the protocol and methodological criteria.

10 Five Cochrane reviews were identified that meet the topic and were included as the  
11 basis for many of the prevention intervention reviews. The Cameron 2018 Cochrane  
12 review<sup>1</sup> included fall prevention interventions older people in residential care and in  
13 hospitals. It used the Prevention of Falls Network Europe (ProFaNE) classification  
14 system to group interventions based on their subtype, which we followed in order to  
15 be consistent with the Cochrane review. Gillespie 2012 (updated 2018)<sup>4</sup> Cochrane  
16 review was used as the basis for the fall prevention intervention review for older  
17 people in the community. More recent Cochrane reviews had updated this,  
18 depending on the subtype of falls prevention. Sherrington (2019)<sup>15</sup> updated exercise  
19 interventions and Hopewell (2018)<sup>6</sup> updated the multifactorial/multi component  
20 interventions. We updated these Cochrane reviews to include all recent papers  
21 identified in our search. Clemson 2023<sup>2</sup> was a new review at the time of writing the  
22 guideline, where no further RCTs were identified, therefore we included this in its  
23 entirety. More recent Cochranes were identified that updated other parts, such as  
24 psychological and educational interventions, so these were also noted within the  
25 reviews, but the overall categorisation remained as in the older Cochrane reviews. It  
26 should be noted that the protocol for our intervention reviews did not match the  
27 Cochrane reviews entirely, but we tried to ensure the reviews aligned as much as  
28 possible.

### 29 **2.3.1.1.1 Qualitative studies**

30 In the qualitative reviews, studies using focus groups, or structured or semi-  
31 structured interviews were considered for inclusion. Survey data or other types of  
32 questionnaires were only included if they provided analysis from open-ended  
33 questions, but not if they reported descriptive quantitative data only.

## 34 **2.4 Methods of combining evidence**

### 35 **2.4.1 Data synthesis for intervention reviews**

36 Meta-analyses were conducted using Cochrane Review Manager (RevMan5)<sup>14</sup>  
37 software

#### 38 **2.4.1.1 Analysis of different types of data**

##### 39 ***Dichotomous outcomes***

40 Fixed-effects (Mantel–Haenszel) techniques were used to calculate risk ratios  
41 (relative risk, RR) for the binary outcomes. The absolute risk difference was also  
42 calculated using GRADEpro<sup>5</sup> software, using the median event rate in the control arm  
43 of the pooled results.

1 We followed the Cochrane review analysis of dichotomous outcomes by inputting as  
2 Generic Inverse Variance variables so that adjusted and unadjusted outcomes could  
3 be meta-analysed together. This resulted in absolute risk difference not being  
4 calculated for the outcome of number of fallers.

### 5 **Rate ratio**

6 Similar to risk ratios for rate ratios, we followed the Cochrane review and had a  
7 hierarchy of data:

- 8 1. Incidence rate ratio (IRR) which was available in the studies
- 9 2. Hazard ratios reported for rate of falls
- 10 3. We calculated the rate ratio using a rate data calculator

11 For binary variables where there were zero events in either arm or a less than 1%  
12 event rate, Peto odds ratios, rather than risk ratios, were calculated as they are more  
13 appropriate for data with a low number of events. Where there are zero events in  
14 both arms, the risk difference was calculated and reported instead.

### 15 **Continuous outcomes**

16 Continuous outcomes were analysed using an inverse variance method for pooling  
17 weighted mean differences.

18 Where the studies within a single meta-analysis had different scales of measurement  
19 for the same outcomes, standardised mean differences were used (providing all  
20 studies reported either change from baseline or final values rather than a mixture of  
21 both); each different measure in each study was 'normalised' to the standard  
22 deviation value pooled between the intervention and comparator groups in that same  
23 study.

24 The means and standard deviations of continuous outcomes are required for meta-  
25 analysis. However, in cases where standard deviations were not reported, the  
26 standard error was calculated if the p values or 95% confidence intervals (95% CI)  
27 were reported, and meta-analysis was undertaken with the mean and standard error  
28 using the generic inverse variance method in RevMan5<sup>14</sup>.

### 29 **Generic inverse variance**

30 If a study reported only the summary statistic and 95% CI the generic-inverse  
31 variance method was used to enter data into RevMan5.<sup>14</sup> If the control event rate was  
32 reported this was used to generate the absolute risk difference in GRADEpro.<sup>5</sup> If  
33 multivariate analysis was used to derive the summary statistic but no adjusted control  
34 event rate was reported no absolute risk difference was calculated.

### 35 **Complex analysis**

36 Where studies had used a crossover design, paired continuous data were extracted  
37 where possible, and forest plots were generated in RevMan5<sup>14</sup> with the generic  
38 inverse variance function. When a crossover study had categorical data and the  
39 number of subjects with an event in both interventions was known, the standard error  
40 (of the log of the risk ratio) was calculated using the simplified Mantel–Haenszel  
41 method for paired outcomes. Forest plots were also generated in RevMan5<sup>14</sup> with the  
42 generic inverse variance function. If paired continuous or categorical data were not  
43 available from the crossover studies, the separate group data were analysed in the

1 same way as data from parallel groups, on the basis that this approach would  
2 overestimate the confidence intervals and thus artificially reduce study weighting  
3 resulting in a conservative effect. Where a meta-analysis included a mixture of  
4 studies using both paired and parallel group approaches, all data were entered into  
5 RevMan5<sup>14</sup> using the generic inverse variance function.

#### 6 **2.4.1.2 Network meta-analysis**

7 Network meta-analysis was considered for the comparison of interventional  
8 treatments but was not pursued because of insufficient data available for the relevant  
9 outcomes.

#### 10 **2.4.1.3 Data synthesis for prediction rules/models**

11 Evidence for risk prediction rules or risk prediction tools were presented separately  
12 for discrimination and calibration. The discrimination data were analysed according to  
13 the principles of data synthesis for diagnostic accuracy studies as outlined in section  
14 **Error! Reference source not found..**

15 Coupled forest plots of the agreed primary paired outcome measures for decision  
16 making (sensitivity and specificity) with their 95% CIs across studies (at various  
17 thresholds) were produced for each test, using RevMan5.<sup>14</sup> In order to do this, 2 by 2  
18 tables (the number of true positives, false positives, true negatives and false  
19 negatives) were directly taken from the study if given, or else were derived from raw  
20 data or calculated from the set of test accuracy statistics.

21 Meta-analysis was conducted where appropriate, that is, when 3 or more studies  
22 were available per threshold. Predictive accuracy for the studies was pooled using  
23 the bivariate method for the direct estimation of summary sensitivity and specificity  
24 using a random-effects approach in WinBUGS software.<sup>16</sup> The advantage of this  
25 approach is that it produces summary estimates of sensitivity and specificity that  
26 account for the correlation between the 2 statistics. The bivariate method uses  
27 logistic regression on the true positives, true negatives, false positives and false  
28 negatives reported in the studies. Overall sensitivity and specificity and confidence  
29 regions were plotted (using methods outlined by Novielli 2010.<sup>12</sup>) Pooled median  
30 sensitivity and specificity and their 95% CIs were reported in the clinical evidence  
31 summary tables. For analyses with fewer than 3 studies included the results of the  
32 study with the lower sensitivity value was reported when there were 2 studies with  
33 the corresponding specificity for that study with the range stated for both studies, or  
34 reported individually for a single study.

35 Heterogeneity or inconsistency amongst studies was visually inspected in the forest  
36 plots.

37 If available, area under the ROC curve (AUC) data for each study were also plotted  
38 on a graph, for each risk tool. The AUC describes the overall predictive accuracy  
39 across the full range of thresholds. The following criteria were used for evaluating  
40 AUCs:

- 41 •  $\leq 0.50$ : worse than chance
- 42 • 0.50–0.60: very poor
- 43 • 0.61–0.70: poor

- 1 • 0.71–0.80: moderate
- 2 • 0.81–0.90: good
- 3 • 0.91–1.00: excellent or perfect test.

4 Heterogeneity or inconsistency amongst studies was visually inspected.

5 Calibration data such as r-squared ( $R^2$ ), if reported, were presented separately to the  
6 discrimination data. The results were presented for each study separately along with  
7 the quality rating for the study. Inconsistency and imprecision were not assessed for  
8 calibration data.

## 9 **2.4.2 Data synthesis for qualitative reviews**

10 The main findings for each included paper were identified and thematic analysis  
11 methods were used to synthesise this information into broad overarching themes  
12 which were summarised into the main review findings. The evidence was presented  
13 in the form of a narrative summary detailing the evidence from the relevant papers  
14 and how this informed the overall review finding plus a statement on the level of  
15 confidence for that review finding. Considerable limitations and issues around  
16 relevance were listed. A summary evidence table with the succinct summary  
17 statements for each review finding was produced including the associated quality  
18 assessment.

## 19 **2.5 Appraising the quality of evidence by outcomes**

### 20 **2.5.1 Intervention reviews**

21 The evidence for outcomes from the included RCTs and, where appropriate, non-  
22 randomised intervention studies, were evaluated and presented using the ‘Grading of  
23 Recommendations Assessment, Development and Evaluation (GRADE) toolbox’  
24 developed by the international GRADE working group  
25 (<http://www.gradeworkinggroup.org/>). The software (GRADEpro<sup>5</sup>) developed by the  
26 GRADE working group was used to assess the quality of each outcome, taking into  
27 account individual study quality and the meta-analysis results.

28 Each outcome was first examined for each of the quality elements listed and defined  
29 in Table 2.

30 **Table 2: Description of quality elements in GRADE for intervention studies**

Quality element	Description
Risk of bias	Limitations in the study design and implementation may bias the estimates of the treatment effect. Major limitations in studies decrease the confidence in the estimate of the effect. Examples of such limitations are selection bias (often due to poor allocation concealment), performance and detection bias (often due to a lack of blinding of the patient, healthcare professional or assessor) and attrition bias (due to missing data causing systematic bias in the analysis).
Indirectness	Indirectness refers to differences in study population, intervention, comparator and outcomes between the available evidence and the review question.
Inconsistency	Inconsistency refers to an unexplained heterogeneity of effect estimates between studies in the same meta-analysis.

Quality element	Description
Imprecision	Results are imprecise when studies include relatively few patients and few events (or highly variable measures) and thus have wide confidence intervals around the estimate of the effect relative to clinically important thresholds. 95% confidence intervals denote the possible range of locations of the true population effect at a 95% probability, and so wide confidence intervals may denote a result that is consistent with conflicting interpretations (for example a result may be consistent with both clinical benefit AND clinical harm) and thus be imprecise.
Publication bias	Publication bias is a systematic underestimate or overestimate of the underlying beneficial or harmful effect due to the selective publication of studies. A closely related phenomenon is where some papers fail to report an outcome that is inconclusive, thus leading to an overestimate of the effectiveness of that outcome.
Other issues	Sometimes randomisation may not adequately lead to group equivalence of confounders, and if so this may lead to bias, which should be taken into account. Potential conflicts of interest, often caused by excessive pharmaceutical company involvement in the publication of a study, should also be noted.

1 Details of how the 4 main quality elements (risk of bias, indirectness, inconsistency  
2 and imprecision) were appraised for each outcome are given below. Publication bias  
3 was considered with the committee. If there was reason to suspect it was present, it  
4 was explored with funnel plots.

#### 5 **2.5.1.1 Risk of bias**

6 The main domains of bias for RCTs are listed in Table 3. Each outcome had its risk  
7 of bias assessed within each study first using the appropriate checklist for the study  
8 design (Cochrane RoB 2 for RCTs, or ROBINS-I for non-randomised studies or  
9 ROBIS for systematic reviews). For each study, if there was no risk of bias in any  
10 domain, the risk of bias was given a rating of 'low risk of bias'. An overall judgment of  
11 'some concerns' was made if some concerns were present in at least one domain  
12 and the domain was judged to be at high risk of bias. An overall judgment of 'high  
13 risk of bias' was made if high risk domains in a way that substantially lowers  
14 confidence in the result. An overall rating of; not serious, serious or very serious, is  
15 applied in GRADEpro across all studies combined in a meta-analysis by taking into  
16 account the weighting of studies according to study precision.

17 **Table 3: Principal domains of bias in randomised controlled trials**

Limitation	Explanation
Selection bias (sequence generation and allocation concealment)	If those enrolling participants are aware of the group to which the next enrolled patient will be allocated, either because of a non-random sequence that is predictable, or because a truly random sequence was not concealed from the researcher, this may translate into systematic selection bias. This may occur if the researcher chooses not to recruit a participant into that specific group because of: <ul style="list-style-type: none"> <li>• knowledge of that participant's likely prognostic characteristics, and</li> <li>• a desire for one group to do better than the other.</li> </ul>
Performance and detection bias (lack of blinding)	Patients, caregivers, those adjudicating or recording outcomes, and data analysts should not be aware of the arm to which the participants are allocated. Knowledge of the group can influence: <ul style="list-style-type: none"> <li>• the experience of the placebo effect</li> <li>• performance in outcome measures</li> </ul>



Limitation	Explanation
	<ul style="list-style-type: none"> <li>the level of care and attention received, and</li> <li>the methods of measurement or analysis</li> </ul> all of which can contribute to systematic bias.
Attrition bias	Attrition bias results from an unaccounted-for loss of data beyond a certain level (a differential of at least 10% between groups). Loss of data can occur when participants are compulsorily withdrawn from a group by the researchers (for example, when a per-protocol approach is used) or when participants do not attend assessment sessions. If the missing data are likely to be different from the data of those remaining in the groups, and there is a differential rate of such missing data from groups, systematic attrition bias may result.
Selective outcome reporting	Reporting of some outcomes and not others on the basis of the results can also lead to bias, as this may distort the overall impression of efficacy.
Other limitations	For example: <ul style="list-style-type: none"> <li>Stopping early for benefit observed in randomised trials, in particular in the absence of adequate stopping rules.</li> <li>Use of unvalidated patient-reported outcome measures.</li> <li>Lack of washout periods to avoid carry-over effects in crossover trials.</li> <li>Recruitment bias in cluster-randomised trials.</li> </ul>

### 1      **2.5.1.2      Indirectness**

2      Indirectness refers to the extent to which the populations, interventions, comparisons  
 3      and outcome measures are dissimilar to those defined in the inclusion criteria for the  
 4      reviews. Indirectness is important when these differences are expected to contribute  
 5      to a difference in effect size or may affect the balance of harms and benefits  
 6      considered for an intervention. As for the risk of bias, each outcome had its  
 7      indirectness assessed within each study first. For each study, if there were no  
 8      sources of indirectness, indirectness was given a rating of 'directly applicable'. If  
 9      there was indirectness in just 1 source (for example in terms of population),  
 10     indirectness was given a rating of 'partially applicable', but if there was indirectness in  
 11     2 or more sources (for example, in terms of population and treatment) the  
 12     indirectness was given an 'indirectly applicable' rating. An overall rating of; not  
 13     serious, serious, or very serious, was applied GRADEpro across all studies by taking  
 14     into account the weighting of studies according to study precision.

### 15     **2.5.1.3      Inconsistency**

16     Inconsistency refers to an unexplained heterogeneity of results for an outcome  
 17     across different studies. When estimates of the treatment effect across studies differ  
 18     widely, this suggests true differences in the underlying treatment effect, which may  
 19     be due to differences in populations, settings or doses. Statistical heterogeneity was  
 20     assessed for each meta-analysis estimate by an I-squared ( $I^2$ ) inconsistency statistic.

21     Heterogeneity or inconsistency amongst studies was also visually inspected. Where  
 22     statistical heterogeneity as defined above was present or there was clear visual  
 23     heterogeneity not captured in the  $I^2$  value predefined subgrouping of studies was  
 24     carried out according to the protocol. See the review protocols for the subgrouping  
 25     strategy.

26     When heterogeneity existed within an outcome ( $I^2 > 50\%$ ), but no plausible  
 27     explanation could be found, the quality of evidence for that outcome was

1 downgraded. Inconsistency for that outcome was given a 'serious' rating if the  $I^2$  was  
2 50–74%, and a 'very serious' rating if the  $I^2$  was 75% or more.

3 If inconsistency could be explained based on pre-specified subgroup analysis (that is,  
4 each subgroup had an  $I^2 < 50\%$ ) then each of the derived subgroups were presented  
5 separately for that forest plot and GRADE profile (providing at least 2 studies  
6 remained in each subgroup). The committee took this into account and considered  
7 whether to make separate recommendations based on the variation in effect across  
8 subgroups within the same outcome. In such a situation the quality of evidence was  
9 not downgraded.

10 If all predefined strategies of subgrouping were unable to explain statistical  
11 heterogeneity, then a random effects (DerSimonian and Laird) model was employed  
12 to the entire group of studies in the meta-analysis. A random-effects model assumes  
13 a distribution of populations, rather than a single population. This leads to a widening  
14 of the confidence interval around the overall estimate. If, however, the committee  
15 considered the heterogeneity was so large that meta-analysis was inappropriate,  
16 then the results were not pooled and were described narratively.

#### 17 **2.5.1.4 Imprecision**

18 The criteria applied for imprecision were based on the 95% CIs for the pooled  
19 estimate of effect, and the minimal important differences (MID) for the outcome. The  
20 MIDs are the threshold for appreciable benefits and harms, separated by a zone  
21 either side of the line of no effect where there is assumed to be no clinically important  
22 effect. If either end of the 95% CI of the overall estimate of effect crossed 1 of the  
23 MID lines, imprecision was regarded as serious in the GRADEpro rating. This was  
24 because the overall result, as represented by the span of the confidence interval,  
25 was consistent with 2 interpretations as defined by the MID (for example, both no  
26 clinically important effect and clinical benefit were possible interpretations). If both  
27 MID lines were crossed by either or both ends of the 95% CI then imprecision was  
28 regarded as very serious. This was because the overall result was consistent with all  
29 3 interpretations defined by the MID (no clinically important effect, clinical benefit and  
30 clinical harm). This is illustrated in Figure 1.

31 The value / position of the MID lines is ideally determined by values reported in the  
32 literature. 'Anchor-based' methods aim to establish clinically meaningful changes in a  
33 continuous outcome variable by relating or 'anchoring' them to patient-centred  
34 measures of clinical effectiveness that could be regarded as gold standards with a  
35 high level of face validity. For example, a MID for an outcome could be defined by the  
36 minimum amount of change in that outcome necessary to make patients feel their  
37 quality of life had 'significantly improved'. MIDs in the literature may also be based on  
38 expert clinician or consensus opinion concerning the minimum amount of change in a  
39 variable deemed to affect quality of life or health.

40 In the absence of values identified in the literature, the alternative approach to  
41 deciding on MID levels is to use the modified GRADE 'default' values, as follows:

- 42 • For dichotomous outcomes the MIDs were taken to be RRs of 0.8\* and 1.25. For  
43 'positive' outcomes such as 'patient satisfaction', the RR of 0.8 is taken as the line  
44 denoting the boundary between no clinically important effect and a clinically  
45 important harm, whilst the RR of 1.25 is taken as the line denoting the boundary  
46 between no clinically important effect and a clinically important benefit. For  
47 'negative' outcomes such as 'bleeding', the opposite occurs, so the RR of 0.8 is  
48 taken as the line denoting the boundary between no clinically important effect and

1 a clinically important benefit, whilst the RR of 1.25 is taken as the line denoting the  
2 boundary between no clinically important effect and a clinically important harm.  
3 There aren't established default values for ORs and the same values (0.8 and  
4 1.25) are applied here but are acknowledged as arbitrary thresholds agreed by the  
5 committee.

- 6 ○ In cases where there are zero events in one arm of a single study, or some or  
7 all of the studies in one arm of a meta-analysis, the same process is followed  
8 as for dichotomous outcomes. However, if there are no events in either arm in  
9 a meta-analysis (or in a single un-pooled study) the sample size is used to  
10 determine imprecision using the following rule of thumb:
  - 11 – No imprecision: sample size  $\geq 350$
  - 12 – Serious imprecision: sample size  $\geq 70$  but  $< 350$
  - 13 – Very serious imprecision: sample size  $< 70$ .
- 14 ○ When there was more than one study in an analysis and zero events occurred  
15 in both groups for some but not all of the studies across both arms, the  
16 optimum information size was used to determine imprecision using the  
17 following guide:
  - 18 – No imprecision:  $> 90\%$  power
  - 19 – Serious imprecision: 80-90% power
  - 20 – Very serious imprecision:  $< 80\%$  power.
- 21 ● Time to event data, there aren't established default values for HRs, so the same  
22 values as dichotomous outcomes are applied here (0.8 and 1.25) but are  
23 acknowledged as arbitrary thresholds agreed by the committee.
- 24 ● For continuous outcome variables the MID was taken as half the median baseline  
25 standard deviation of that variable, across all studies in the meta-analysis. Hence  
26 the MID denoting the minimum clinically important benefit was positive for a  
27 'positive' outcome (for example, a quality-of-life measure where a higher score  
28 denotes better health), and negative for a 'negative' outcome (for example, a  
29 visual analogue scale [VAS] pain score). Clinically important harms will be the  
30 converse of these. If baseline values are unavailable, then half the median  
31 comparator group standard deviation of that variable will be taken as the MID. As  
32 these vary for each outcome per review, details of the values used are reported in  
33 the footnotes of the relevant GRADE summary table.
- 34 ● If standardised mean differences have been used, where the GC are able to  
35 specify a priority measure, the results are back converted to a mean difference on  
36 that scale for the assessment of imprecision and clinical importance. If it is not  
37 deemed appropriate to back-convert to a single scale, then the MID was set at the  
38 absolute value of +0.5. This follows because standardised mean differences are  
39 mean differences normalised to the pooled standard deviation of the 2 groups and  
40 are thus effectively expressed in units of 'numbers of standard deviations'. The 0.5  
41 MID value in this context therefore indicates half a standard deviation, the same  
42 definition of MID as used for non-standardised mean differences.

43 \*NB GRADE report the default values as 0.75 and 1.25. These are consensus  
44 values. This guideline follows NICE process to use modified values of 0.8 and 1.25  
45 as they are symmetrical on a relative risk scale.

46 For this guideline, the following MIDs for continuous or dichotomous outcomes were  
47 found in the literature and adopted for use:


1

**Table 4: Published or pre-agreed MIDs**

Outcome measure	MID	Source
EQ-5D	0.03	Consensus pragmatic MID used in some previous NICE guidelines
SF36	Physical component summary: 2 Mental component summary: 3 Physical functioning: 3 Role-physical: 3 Bodily pain: 3 General health: 2 Vitality: 2 Social functioning: 3 Role-emotional: 4 Mental health: 3	User's manual for the SF-36v2 Health Survey, Third Edition <sup>7</sup>

2

**Figure 1:** Illustration of precise and imprecise outcomes based on the 95% CI of dichotomous outcomes in a forest plot (Note that all 3 results would be pooled estimates, and would not, in practice, be placed on the same forest plot)



3

### 2.5.1.5 Overall grading of the quality of clinical evidence

4 Once an outcome had been appraised for the main quality elements, as above, an  
 5 overall quality grade was calculated for that outcome from the ratings from each of  
 6 the main quality elements were summed to give a score that could be anything from  
 7 high to very low. The evidence for each outcome started at High, and the overall  
 8 quality (or confidence in the evidence) remained High if there were no reasons for  
 9 downgrading, or became Moderate, Low or Very Low according to the number of  
 10 independent reasons for downgrading. The significance of these overall ratings is  
 11 explained in Table 5. The reasons for downgrading in each case are specified in the  
 12 footnotes of the GRADE tables.

Falls in older people Update: methods DRAFT (October 2024)

**Table 5: Overall quality of outcome evidence in GRADE**

Level	Description
High	Further research is very unlikely to change our confidence in the estimate of effect
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate
Very low	Any estimate of effect is very uncertain

## 2.5.2 Prediction rules/models

### 2.5.2.1.1 Risk of bias

Risk of bias and applicability of evidence for prognostic risk data were evaluated by study using the Prediction study Risk of Bias Assessment Tool (PROBAST) checklist. Risk of bias and applicability in risk prediction studies in PROBAST consists of 4 domains:

- patient selection
- predictors
- outcome
- analysis.

If data were meta-analysed, the quality for pooled studies was presented. If the data were not pooled, then a quality rating was presented for each study.

### 2.5.2.1.2 Inconsistency

Inconsistency for discrimination and outcomes was assessed by inspection of the primary outcome measures (sensitivity and specificity) using the point estimates and 95% CIs of the individual studies on the forest plots. Particular attention was placed on values above or below 50% (prediction based on chance alone) and the threshold set by the committee (the threshold above which it would be acceptable to recommend a rule/model). The evidence was downgraded by 1 increment if the CI varied across 2 areas (50-70%) and by 2 increments if the individual studies varied across 3 areas, 0–50%, 50–70% and 70–100%). Where only a single study reports an outcome, inconsistency is rated as 'not detected'. Inconsistency was not assessed for calibration outcomes.

### 2.5.2.1.3 Imprecision

In meta-analysed reclassification outcomes, or for non-pooled outcomes, the position of the 95% CIs in relation to the null line determined the existence of imprecision. If the 95% CI did not cross the null line, then no serious imprecision was recorded. If the 95% CI crossed the null line, then serious imprecision was recorded. For discrimination outcomes, the judgement of precision was based on visual inspection of the confidence region around the primary paired outcome measures for decision-making (sensitivity and specificity) from the meta-analysis, if a meta-analysis was conducted. Where a meta-analysis was not conducted, imprecision was assessed according to the range of point estimates or, if only one study contributed to the evidence, the 95% CI around the single study. The decision thresholds set by the committee were used to determine whether imprecision is not serious, serious or

1 very serious depending on whether confidence intervals cross zero, one or two  
2 thresholds. Imprecision was not assessed for calibration outcomes.

### 3 **2.5.2.1.4 Overall grading**

4 Quality rating started at High for prospective studies, and each major limitation  
5 brought the rating down by 1 increment to a minimum grade of Very Low, as  
6 explained for interventional reviews. For prognostic reviews prospective cohort  
7 studies with a multivariate analysis are regarded as the gold standard because RCTs  
8 are usually inappropriate for these types of review for ethical or pragmatic reasons.  
9 Furthermore, if the study is looking at more than 1 risk factor of interest then  
10 randomisation would be inappropriate as it can only be applied to 1 of the risk  
11 factors. This was presented in a modified GRADE evidence profile.

## 12 **2.5.3 Qualitative reviews**

13 Review findings from the included qualitative studies were evaluated and presented  
14 using the 'Confidence in the Evidence from Reviews of Qualitative Research'  
15 (CERQual) Approach developed by the GRADE-CERQual Project Group, a subgroup  
16 of the GRADE Working Group.

17 The CERQual Approach assesses the extent to which a review finding is a  
18 reasonable representation of the phenomenon of interest (the focus of the review  
19 question). Each review finding was assessed for each of the 4 quality elements listed  
20 and defined below in Table 6.

21 **Table 6: Description of quality elements in GRADE-CERQual for qualitative**  
22 **studies**

Quality element	Description
Methodological limitations	The extent of problems in the design or conduct of the included studies that could decrease the confidence that the review finding is a reasonable representation of the phenomenon of interest. Assessed at the study level using the CASP checklist.
Coherence	The extent to how clear and cogent the fit is between the data from the primary studies and the review finding.
Relevance	The extent to which the body of evidence from the included studies is applicable to the context (study population, phenomenon of interest, setting) specified in the protocol.
Adequacy	The degree of the confidence that the review finding is being supported by sufficient data. This is an overall determination of the richness (depth of analysis) and quantity of the evidence supporting a review finding or theme.

23 Details of how the 4 quality elements (methodological limitations, coherence,  
24 relevance and adequacy) were appraised for each review finding are given below.

### 25 **2.5.3.1 Methodological limitations**

26 Each review finding had its methodological limitations assessed within each study  
27 first using the CASP checklist. Based on the degree of methodological limitations,  
28 studies were evaluated as having minor, moderate or severe limitations. A summary  
29 of the domains and questions covered is given below.

**Table 7: Description of limitations assessed in the CASP checklist for qualitative studies**

Domain	Aspects considered
Are the results valid?	<ul style="list-style-type: none"> <li>• Was there a clear statement of the aims of the research?</li> <li>• Is qualitative methodology appropriate?</li> <li>• Was the research design appropriate to address the aims of the research?</li> <li>• Was the recruitment strategy appropriate to the aims of the research?</li> <li>• Was the data collected in a way that addressed the research issue?</li> <li>• Has the relationship between researcher and participants been adequately considered?</li> </ul>
What are the results?	<ul style="list-style-type: none"> <li>• Have ethical issues been taken into consideration?</li> <li>• Was the data analysis sufficiently rigorous?</li> <li>• Is there a clear statement of findings?</li> </ul>
Will the results help locally?	<ul style="list-style-type: none"> <li>• How valuable is the research?</li> </ul>

The overall assessment of the methodological limitations of the evidence was based on the limitations of the primary studies contributing to the review finding. The relative contribution of each study to the overall review finding and of the type of methodological limitation(s) were taken into account when giving an overall rating of concerns for this component.

#### 2.5.3.2 Relevance

Relevance is the extent to which the body of evidence from the included studies is applicable to the context (study population, phenomenon of interest, setting) specified in the protocol. As such, relevance is dependent on the individual review and discussed with the guideline committee.

#### 2.5.3.3 Coherence

Coherence is the extent to which the reviewer is able to identify a clear pattern across the studies included in the review, and if there is variation present (contrasting or disconfirming data) whether this variation is explained by the contributing study authors. For example, if a review finding in 1 study does not support the main finding and there is no plausible explanation for this variation, or if there is ambiguity in the descriptions in the primary data, then the confidence that the main finding reasonably reflects the phenomenon of interest is decreased.

#### 2.5.3.4 Adequacy

The judgement of adequacy is based on the confidence of the finding being supported by sufficient data. This is an overall determination of the richness (and quantity of the evidence supporting a review finding or theme. Rich data provide sufficient detail to gain an understanding of the theme or review finding, whereas thin data do not provide enough detail for an adequate understanding. Quantity of data is the second pillar of the assessment of adequacy. For review findings that are only supported by 1 study or data from only a small number of participants, the confidence that the review finding reasonably represents the phenomenon of interest might be decreased because there is less confidence that studies undertaken in other settings or participants would have reported similar findings. As with richness of data, quantity

of data is review dependent. Based on the overall judgement of adequacy, a rating of no concerns, minor concerns, or substantial concerns about adequacy was given.

### 2.5.3.5 Overall judgement of the level of confidence for a review finding

GRADE-CERQual is used to assess the body of evidence as a whole through a confidence rating representing the extent to which a review finding is a reasonable representation of the phenomenon of interest. For each of the above components, level of concern is categorised as either:

- no or very minor concerns
- minor concerns
- moderate concerns, or
- serious concerns.

The concerns from the 4 components (methodological limitations, coherence, relevance and adequacy) are used in combination to form an overall judgement of confidence in the finding. GRADE-CERQual uses 4 levels of confidence: high, moderate, low and very low confidence. The significance of these overall ratings is explained in Table 8. Each review finding starts at a high level of confidence and is downgraded based on the concerns identified in any 1 or more of the 4 components. Quality assessment of qualitative reviews is a subjective judgement by the reviewer based on the concerns that have been noted. An explanation of how such a judgement had been made for each component is included in the footnotes of the summary of evidence tables.

**Table 8: Overall level of confidence for a review finding in GRADE-CERQual**

Level	Description
High confidence	It is highly likely that the review finding is a reasonable representation of the phenomenon of interest.
Moderate confidence	It is likely that the review finding is a reasonable representation of the phenomenon of interest.
Low confidence	It is possible that the review finding is a reasonable representation of the phenomenon of interest.
Very low confidence	It is not clear whether the review finding is a reasonable representation of the phenomenon of interest.

### 2.5.4 Publication bias

## 2.6 Assessing clinical importance

The committee assessed the evidence by outcome in order to determine if there was, or potentially was, a clinically important benefit, a clinically important harm or no clinically important difference between interventions. To facilitate this where possible, binary outcomes were converted into absolute risk differences (ARDs) using GRADEpro<sup>5</sup> software: the median control group risk across studies was used to calculate the ARD and its 95% CI from the pooled risk ratio. However, for the outcomes of rate ratio and number of fallers because Generic Inverse Variance was used to summarise the data, we did not have absolute risk differences available for these outcomes. In these instances, the point estimate of the meta-analysed values was assessed according to whether it crossed the default of 0.80 to 1.25 for Risk ratio and Rate ratio.



1 The assessment of clinical benefit, harm, or no benefit or harm was based on the  
2 point estimate of absolute effect for intervention studies, which was standardised  
3 across the reviews. The committee considered for most of the dichotomous  
4 outcomes in the intervention reviews that if at least 100 more participants per 1000  
5 (10%) achieved the outcome of interest in the intervention group compared to the  
6 comparison group for a positive outcome then this intervention was considered  
7 beneficial. The same point estimate but in the opposite direction applied for a  
8 negative outcome. For adverse events 50 events or more per 1000 (5%) represented  
9 clinical harm.

10 For continuous outcomes if the mean difference was greater than the minimally  
11 important difference (MID) then this represented a clinical benefit or harm.

12 The published values used for imprecision and clinical importance are provided in  
13 **Table 4**. For continuous outcomes where the GRADE default MID has been used,  
14 the values for each outcome are provided in the footnotes of the relevant GRADE  
15 tables.

## 16 **2.7 Identifying and analysing evidence of cost** 17 **effectiveness**

18 The committee is required to make decisions based on the best available evidence of  
19 both clinical effectiveness and cost effectiveness. Guideline recommendations should  
20 be based on the expected costs of the different options in relation to their expected  
21 health benefits (that is, their 'cost effectiveness') rather than the total implementation  
22 cost. However, the committee will also need to be increasingly confident in the cost  
23 effectiveness of a recommendation as the cost of implementation increases.  
24 Therefore, the committee may require more robust evidence on the effectiveness and  
25 cost effectiveness of any recommendations that are expected to have a substantial  
26 impact on resources; any uncertainties must be offset by a compelling argument in  
27 favour of the recommendation. The cost impact or savings potential of a  
28 recommendation should not be the sole reason for the committee's decision.<sup>9</sup>

29 Health economic evidence was sought relating to the key clinical issues being  
30 addressed in the guideline. Health economists:

- 31 • Undertook a systematic review of the published economic literature.
- 32 • Undertook new cost-effectiveness analysis in priority areas.

### 33 **2.7.1 Literature review**

34 The health economists:

- 35 • Identified potentially relevant studies for each review question from the health  
36 economic search results by reviewing titles and abstracts. Full papers were then  
37 obtained.
- 38 • Reviewed full papers against prespecified inclusion and exclusion criteria to  
39 identify relevant studies (see below for details).
- 40 • Critically appraised relevant studies using economic evaluations checklists as  
41 specified in the NICE guidelines manual.<sup>9</sup>
- 42 • Extracted key information about the studies' methods and results into health  
43 economic evidence tables (which can be found in appendices to the relevant  
44 evidence reports).

- Generated summaries of the evidence in NICE health economic evidence profile tables (included in the relevant evidence report for each review question) – see below for details.

#### 2.7.1.1 Inclusion and exclusion criteria

Full economic evaluations (studies comparing costs and health consequences of alternative courses of action: cost–utility, cost-effectiveness, cost–benefit and cost–consequences analyses) and comparative costing studies that addressed the review question in the relevant population were considered potentially includable as health economic evidence.

Studies that only reported cost per hospital (not per patient), or only reported average cost effectiveness without disaggregated costs and effects were excluded. Literature reviews, abstracts, posters, letters, editorials, comment articles, unpublished studies and studies not in English were excluded. Studies published before 2004 and studies from non-OECD countries or the USA were also excluded, on the basis that the applicability of such studies to the present UK NHS context is likely to be too low for them to be helpful for decision-making.

Remaining health economic studies were prioritised for inclusion based on their relative applicability to the development of this guideline and the study limitations. However, in this guideline, no economic studies were excluded on the basis that more applicable evidence was available.

For more details about the assessment of applicability and methodological quality see **Table 9** below and the economic evaluation checklist (appendix H of the NICE guidelines manual<sup>9</sup>) and the health economics review protocol, which can be found in each of the evidence reports.

When no relevant health economic studies were found from the economic literature review, relevant UK NHS unit costs related to the compared interventions were presented to the committee to inform the possible economic implications of the recommendations.

#### 2.7.1.2 NICE health economic evidence profiles

NICE health economic evidence profile tables were used to summarise cost and cost-effectiveness estimates for the included health economic studies in each evidence review report. The health economic evidence profile shows an assessment of applicability and methodological quality for each economic study, with footnotes indicating the reasons for the assessment. These assessments were made by the health economist using the economic evaluation checklist from the NICE guidelines manual.<sup>9</sup> It also shows the incremental costs, incremental effects (for example, quality-adjusted life years [QALYs]) and incremental cost-effectiveness ratio (ICER) for the base case analysis in the study, as well as information about the assessment of uncertainty in the analysis. See **Table 9** for more details.

When a non-UK study was included in the profile, the results were converted into pounds sterling using the appropriate purchasing power parity.<sup>13</sup>

1

**Table 9: Content of NICE health economic evidence profile**

Item	Description
Study	Surname of first author, date of study publication and country perspective with a reference to full information on the study.
Applicability	An assessment of applicability of the study to this guideline, the current NHS situation and NICE decision-making: <sup>(a)</sup> <ul style="list-style-type: none"> <li>• Directly applicable – the study meets all applicability criteria or fails to meet 1 or more applicability criteria but this is unlikely to change the conclusions about cost effectiveness.</li> <li>• Partially applicable – the study fails to meet 1 or more applicability criteria, and this could change the conclusions about cost effectiveness.</li> <li>• Not applicable – the study fails to meet 1 or more of the applicability criteria, and this is likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.</li> </ul>
Limitations	An assessment of methodological quality of the study: <sup>(a)</sup> <ul style="list-style-type: none"> <li>• Minor limitations – the study meets all quality criteria, or fails to meet 1 or more quality criteria, but this is unlikely to change the conclusions about cost effectiveness.</li> <li>• Potentially serious limitations – the study fails to meet 1 or more quality criteria, and this could change the conclusions about cost effectiveness.</li> <li>• Very serious limitations – the study fails to meet 1 or more quality criteria, and this is highly likely to change the conclusions about cost effectiveness. Such studies would usually be excluded from the review.</li> </ul>
Other comments	Information about the design of the study and particular issues that should be considered when interpreting it.
Incremental cost	The mean cost associated with one strategy minus the mean cost of a comparator strategy.
Incremental effects	The mean QALYs (or other selected measure of health outcome) associated with one strategy minus the mean QALYs of a comparator strategy.
Cost effectiveness	Incremental cost-effectiveness ratio (ICER): the incremental cost divided by the incremental effects (usually in £ per QALY gained).
Uncertainty	A summary of the extent of uncertainty about the ICER reflecting the results of deterministic or probabilistic sensitivity analyses, or stochastic analyses of trial data, as appropriate.

2  
3

(a) *Applicability and limitations were assessed using the economic evaluation checklist in appendix H of the NICE guidelines manual<sup>9</sup>*

4

## 2.7.2 Undertaking new health economic analysis

5  
6  
7  
8  
9

As well as reviewing the published health economic literature for each review question, as described above, new health economic analysis was undertaken by the health economist in selected areas. Priority areas for new analysis were agreed by the committee after formation of the review questions and consideration of the existing health economic evidence.

10  
11  
12  
13  
14  
15

The committee identified home hazard assessment by an Occupational Therapist versus standard care as the highest priority area for original health economic modelling. This was due to there being clear clinical evidence that showed using an Occupational Therapist for home hazard assessment was beneficial. However, the committee wanted evidence to see if using an Occupational Therapist is cost effective,

1 The following general principles were adhered to in developing the cost-effectiveness  
2 analysis:

- 3 • Methods were consistent with the NICE reference case for interventions with  
4 health outcomes in NHS settings.<sup>3,9</sup>
- 5 • The committee was involved in the design of the model, selection of inputs and  
6 interpretation of the results.
- 7 • Model inputs were based on the systematic review of the clinical literature  
8 supplemented with other published data sources where possible.
- 9 • When published data were not available committee expert opinion was used to  
10 populate the model.
- 11 • Model inputs and assumptions were reported fully and transparently.
- 12 • The results were subject to sensitivity analysis and limitations were discussed.
- 13 • The model was peer-reviewed by another health economist.

14 Full methods and results of the cost-effectiveness analysis for home hazard  
15 assessment by an Occupational Therapist versus standard care are described in a  
16 separate economic analysis report.

### 17 **2.7.3 Cost-effectiveness criteria**

18 NICE sets out the principles that committees should consider when judging whether  
19 an intervention offers good value for money.<sup>9, 11</sup> In general, an intervention was  
20 considered to be cost effective (given that the estimate was considered plausible) if  
21 either of the following criteria applied:

- 22 • the intervention dominated other relevant strategies (that is, it was both less costly  
23 in terms of resource use and more clinically effective compared with all the other  
24 relevant alternative strategies), or
- 25 • the intervention cost less than £20,000 per QALY gained compared with the next  
26 best strategy.

27 If the committee recommended an intervention that was estimated to cost more than  
28 £20,000 per QALY gained, or did not recommend one that was estimated to cost less  
29 than £20,000 per QALY gained, the reasons for this decision are discussed explicitly  
30 in 'The committee's discussion of the evidence' section of the relevant evidence  
31 report, with reference to issues regarding the plausibility of the estimate or to factors  
32 set out in NICE methods manuals.<sup>9</sup>

33 When QALYs or life years gained are not used in the analysis, results are difficult to  
34 interpret unless one strategy dominates the others with respect to every relevant  
35 health outcome and cost.

### 36 **2.7.4 In the absence of health economic evidence**

37 When no relevant published health economic studies were found, and a new analysis  
38 was not prioritised, the committee made a qualitative judgement about cost  
39 effectiveness by considering expected differences in resource use between options  
40 and relevant UK NHS unit costs, alongside the results of the review of clinical  
41 effectiveness evidence.

42 The UK NHS costs reported in the guideline are those that were presented to the  
43 committee and were correct at the time recommendations were drafted. They may

1 have changed subsequently before the time of publication. However, we have no  
2 reason to believe they have changed substantially.

## 3 **2.8 Developing recommendations**

4 Over the course of the guideline development process, the committee was presented  
5 with:

- 6 • Summaries of clinical and health economic evidence and quality (as presented in  
7 evidence reports A–I).
- 8 • Evidence tables of the clinical and health economic evidence reviewed from the  
9 literature. All evidence tables can be found in appendices to the relevant evidence  
10 reports.
- 11 • Forest plots (in appendices to the relevant evidence reports).
- 12 • A description of the methods and results of the cost-effectiveness analysis  
13 undertaken for the guideline (in a separate economic analysis report).

14 Decisions on whether a recommendation could be made, and if so in which direction,  
15 were made on the basis of the committee’s interpretation of the available evidence,  
16 taking into account the balance of benefits, harms and costs between different  
17 courses of action. This was either done formally in an economic model, or informally.  
18 The net clinical benefit over harm (clinical effectiveness) was considered, focusing on  
19 the magnitude of the effect (or clinical importance), quality of evidence (including the  
20 uncertainty) and amount of evidence available. When this was done informally, the  
21 committee took into account the clinical benefits and harms when one intervention  
22 was compared with another. The assessment of net clinical benefit was moderated  
23 by the importance placed on the outcomes (the committee’s values and preferences),  
24 and the confidence the committee had in the evidence (evidence quality). Secondly,  
25 the committee assessed whether the net clinical benefit justified any differences in  
26 costs between the alternative interventions. When the clinical harms were judged by  
27 the committee to outweigh any clinical benefits, they considered making a  
28 recommendation not to offer an intervention. This was dependant on whether the  
29 intervention had any reasonable prospect of providing cost-effective benefits to  
30 people using services and whether stopping the intervention was likely to cause harm  
31 for people already receiving it.

32 When clinical and health economic evidence was of poor quality, conflicting or  
33 absent, the committee decided on whether a recommendation could be made based  
34 on its expert opinion. The considerations for making consensus-based  
35 recommendations include the balance between potential harms and benefits, the  
36 economic costs compared to the economic benefits, current practices,  
37 recommendations made in other relevant guidelines, patient preferences and equality  
38 issues. The consensus recommendations were agreed through discussions in the  
39 committee. The committee also considered whether the uncertainty was sufficient to  
40 justify delaying making a recommendation to await further research, taking into  
41 account the potential harm of failing to make a clear recommendation (see  
42 section 2.8.1 below).

43 The committee considered the appropriate ‘strength’ of each recommendation. This  
44 takes into account the quality of the evidence but is conceptually different. Some  
45 recommendations are ‘strong’ in that the committee believes that the vast majority of  
46 healthcare and other professionals and patients would choose a particular  
47 intervention if they considered the evidence in the same way that the committee has.

1 This is generally the case if the benefits clearly outweigh the harms for most people  
2 and the intervention is likely to be cost effective. However, there is often a closer  
3 balance between benefits and harms, and some patients would not choose an  
4 intervention whereas others would. This may happen, for example, if some patients  
5 are particularly averse to some side effect and others are not. In these circumstances  
6 the recommendation is generally weaker, although it may be possible to make  
7 stronger recommendations about specific groups of patients.

8 The committee focused on the following factors in agreeing the wording of the  
9 recommendations:

- 10 • The actions health professionals need to take.
- 11 • The information readers need to know.
- 12 • The strength of the recommendation (for example the word 'offer' was used for  
13 strong recommendations and 'consider' for weaker recommendations).
- 14 • The involvement of patients (and their carers if needed) in decisions on treatment  
15 and care.
- 16 • Consistency with NICE's standard advice on recommendations about drugs,  
17 waiting times and ineffective interventions (see section 9.2 in the NICE guidelines  
18 manual<sup>10</sup>).

19 The main considerations specific to each recommendation are outlined in 'The  
20 committee's discussion of the evidence' section within each evidence report.

### 21 **2.8.1 Research recommendations**

22 When areas were identified for which, good evidence was lacking, the committee  
23 considered making recommendations for future research. Decisions about the  
24 inclusion of a research recommendation were based on factors such as:

- 25 • the importance to patients or the population
- 26 • national priorities
- 27 • potential impact on the NHS and future NICE guidance
- 28 • ethical and technical feasibility.

### 29 **2.8.2 Validation process**

30 This guidance is subject to a 6-week public consultation and feedback as part of the  
31 quality assurance and peer review of the document. All comments received from  
32 registered stakeholders are responded to in turn and posted on the NICE website.

### 33 **2.8.3 Updating the guideline**

34 Following publication, and in accordance with the NICE guidelines manual, NICE will  
35 undertake a review of whether the evidence base has progressed significantly to alter  
36 the guideline recommendations and warrant an update.

## 37 **2.9 General terms**

38

Term	Definition
Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.
Algorithm (in guidelines)	A flow chart of the clinical decision pathway described in the guideline, where decision points are represented with boxes, linked with arrows.
Allocation concealment	The process used to prevent advance knowledge of group assignment in an RCT. The allocation process should be impervious to any influence by the individual making the allocation, by being administered by someone who is not responsible for recruiting participants.
Applicability	How well the results of a study or NICE evidence review can answer a clinical question or be applied to the population being considered.
Arm (of a clinical study)	Subsection of individuals within a study who receive one particular intervention, for example placebo arm.
Association	Statistical relationship between 2 or more events, characteristics or other variables. The relationship may or may not be causal.
Base case analysis	In an economic evaluation, this is the main analysis based on the most plausible estimate of each input. In contrast, see Sensitivity analysis.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable), with which subsequent results are compared.
Bayesian analysis	A method of statistics, where a statistic is estimated by combining established information or belief (the 'prior') with new evidence (the 'likelihood') to give a revised estimate (the 'posterior').
Before-and-after study	A study that investigates the effects of an intervention by measuring particular characteristics of a population both before and after taking the intervention, and assessing any change that occurs.
Bias	Influences on a study that can make the results look better or worse than they really are. (Bias can even make it look as if a treatment works when it does not.) Bias can occur by chance, deliberately or as a result of systematic errors in the design and execution of a study. It can also occur at different stages in the research process, for example, during the collection, analysis, interpretation, publication or review of research data. For examples see selection bias, performance bias, information bias, confounding factor, and publication bias.
Blinding	A way to prevent researchers, doctors and patients in a clinical trial from knowing which study group each patient is in so they cannot influence the results. The best way to do this is by sorting patients into study groups randomly. The purpose of 'blinding' or 'masking' is to protect against bias.  A single-blinded study is one in which patients do not know which study group they are in (for example whether they are taking the experimental drug or a placebo). A double-blinded study is one in which neither patients nor the researchers and doctors know which study group the patients are in. A triple blind study is one in which neither the patients, clinicians nor the people carrying out the statistical analysis know which treatment patients received.
Carer (caregiver)	Someone who looks after family, partners or friends in need of help because they are ill, frail or have a disability.
Case-control study	A study to find out the cause(s) of a disease or condition. This is done by comparing a group of patients who have the disease or condition (cases) with a group of people who do not have it (controls) but who

Term	Definition
	<p>are otherwise as similar as possible (in characteristics thought to be unrelated to the causes of the disease or condition). This means the researcher can look for aspects of their lives that differ to see if they may cause the condition.</p> <p>For example, a group of people with lung cancer might be compared with a group of people the same age that do not have lung cancer. The researcher could compare how long both groups had been exposed to tobacco smoke. Such studies are retrospective because they look back in time from the outcome to the possible causes of a disease or condition.</p>
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinical effectiveness	<p>How well a specific test or treatment works when used in the 'real world' (for example, when used by a doctor with a patient at home), rather than in a carefully controlled clinical trial. Trials that assess clinical effectiveness are sometimes called management trials.</p> <p>Clinical effectiveness is not the same as efficacy.</p>
Clinician	A healthcare professional who provides patient care. For example, a doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence-based medicine databases including the Cochrane Database of Systematic Reviews (reviews of randomised controlled trials prepared by the Cochrane Collaboration).
Cohort study	A study with 2 or more groups of people – cohorts – with similar characteristics. One group receives a treatment, is exposed to a risk factor or has a particular symptom and the other group does not. The study follows their progress over time and records what happens. See also observational study.
Comorbidity	A disease or condition that someone has in addition to the health problem being studied or treated.
Comparability	Similarity of the groups in characteristics likely to affect the study results (such as health status or age).
Concordance	This is a recent term whose meaning has changed. It was initially applied to the consultation process in which doctor and patient agree therapeutic decisions that incorporate their respective views, but now includes patient support in medicine taking as well as prescribing communication. Concordance reflects social values but does not address medicine-taking and may not lead to improved adherence.
Confidence interval (CI)	A range of values for an unknown population parameter with a stated 'confidence' (conventionally 95%) that it contains the true value. The interval is calculated from sample data, and generally straddles the sample estimate. The 'confidence' value means that if the method used to calculate the interval is repeated many times, then that proportion of intervals will actually contain the true value.
Confounding factor	<p>Something that influences a study and can result in misleading findings if it is not understood or appropriately dealt with.</p> <p>For example, a study of heart disease may look at a group of people that exercises regularly and a group that does not exercise. If the ages of the people in the 2 groups are different, then any difference in heart disease rates between the 2 groups could be because of age rather than exercise. Therefore, age is a confounding factor.</p>



Term	Definition
Consensus methods	Techniques used to reach agreement on a particular issue. Consensus methods may be used to develop NICE guidance if there is not enough good quality research evidence to give a clear answer to a question. Formal consensus methods include Delphi and nominal group techniques.
Control group	A group of people in a study who do not receive the treatment or test being studied. Instead, they may receive the standard treatment (sometimes called 'usual care') or a dummy treatment (placebo). The results for the control group are compared with those for a group receiving the treatment being tested. The aim is to check for any differences. Ideally, the people in the control group should be as similar as possible to those in the treatment group, to make it as easy as possible to detect any effects due to the treatment.
Cost–benefit analysis (CBA)	Cost–benefit analysis is one of the tools used to carry out an economic evaluation. The costs and benefits are measured using the same monetary units (for example, pounds sterling) to see whether the benefits exceed the costs.
Cost–consequences analysis (CCA)	Cost–consequences analysis is one of the tools used to carry out an economic evaluation. This compares the costs (such as treatment and hospital care) and the consequences (such as health outcomes) of a test or treatment with a suitable alternative. Unlike cost–benefit analysis or cost-effectiveness analysis, it does not attempt to summarise outcomes in a single measure (like the quality-adjusted life year) or in financial terms. Instead, outcomes are shown in their natural units (some of which may be monetary) and it is left to decision-makers to determine whether, overall, the treatment is worth carrying out.
Cost-effectiveness analysis (CEA)	Cost-effectiveness analysis is one of the tools used to carry out an economic evaluation. The benefits are expressed in non-monetary terms related to health, such as symptom-free days, heart attacks avoided, deaths avoided or life years gained (that is, the number of years by which life is extended as a result of the intervention).
Cost-effectiveness model	An explicit mathematical framework, which is used to represent clinical decision problems and incorporate evidence from a variety of sources in order to estimate the costs and health outcomes.
Cost–utility analysis (CUA)	Cost–utility analysis is one of the tools used to carry out an economic evaluation. The benefits are assessed in terms of both quality and duration of life and expressed as quality-adjusted life years (QALYs). See also utility.
Credible interval (CrI)	The Bayesian equivalent of a confidence interval.
Decision analysis	An explicit quantitative approach to decision-making under uncertainty, based on evidence from research. This evidence is translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Deterministic analysis	In economic evaluation, this is an analysis that uses a point estimate for each input. In contrast, see Probabilistic analysis
Diagnostic odds ratio	The diagnostic odds ratio is a measure of the effectiveness of a diagnostic test. It is defined as the ratio of the odds of the test being positive if the subject has a disease relative to the odds of the test being positive if the subject does not have the disease.

Term	Definition
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Disutility	The loss of quality of life associated with having a disease or condition. See Utility
Dominance	A health economics term. When comparing tests or treatments, an option that is both less effective and costs more is said to be 'dominated' by the alternative.
Drop-out	A participant who withdraws from a trial before the end.
Economic evaluation	An economic evaluation is used to assess the cost effectiveness of healthcare interventions (that is, to compare the costs and benefits of a healthcare intervention to assess whether it is worth doing). The aim of an economic evaluation is to maximise the level of benefits – health effects – relative to the resources available. It should be used to inform and support the decision-making process; it is not supposed to replace the judgement of healthcare professionals. There are several types of economic evaluation: cost–benefit analysis, cost–consequences analysis, cost-effectiveness analysis, cost-minimisation analysis and cost–utility analysis. They use similar methods to define and evaluate costs but differ in the way they estimate the benefits of a particular drug, programme or intervention.
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	A measure that shows the magnitude of the outcome in one group compared with that in a control group. For example, if the absolute risk reduction is shown to be 5% and it is the outcome of interest, the effect size is 5%. The effect size is usually tested, using statistics, to find out how likely it is that the effect is a result of the treatment and has not just happened by chance (that is, to see if it is statistically significant).
Effectiveness	How beneficial a test or treatment is under usual or everyday conditions, compared with doing nothing or opting for another type of care.
Efficacy	How beneficial a test, treatment or public health intervention is under ideal conditions (for example, in a laboratory), compared with doing nothing or opting for another type of care.
Epidemiological study	The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (for example, infection, diet) and interventions.
EQ-5D (EuroQol 5 dimensions)	A standardised instrument used to measure health-related quality of life. It provides a single index value for health status.
Evidence	Information on which a decision or guidance is based. Evidence is obtained from a range of sources including randomised controlled trials, observational studies, expert opinion (of clinical professionals or patients).
Exclusion criteria (literature review)	Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.
Exclusion criteria (clinical study)	Criteria that define who is not eligible to participate in a clinical study.
Extended dominance	If Option A is both more clinically effective than Option B and has a lower cost per unit of effect, when both are compared with a do-

Term	Definition
	nothing alternative then Option A is said to have extended dominance over Option B. Option A is therefore cost effective and should be preferred, other things remaining equal.
Extrapolation	An assumption that the results of studies of a specific population will also hold true for another population with similar characteristics.
Follow-up	Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been assessed in order to observe changes in health status or health-related variables.
Generalisability	The extent to which the results of a study hold true for groups that did not participate in the research. See also external validity.
Gold standard	A method, procedure or measurement that is widely accepted as being the best available to test for or treat a disease.
GRADE, GRADE evidence profile	A system developed by the GRADE Working Group to address the shortcomings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE evidence profile.
Harms	Adverse effects of an intervention.
Hazard Ratio	The hazard or chance of an event occurring in the treatment arm of a study as a ratio of the chance of an event occurring in the control arm over time.
Health economics	Study or analysis of the cost of using and distributing healthcare resources.
Health-related quality of life (HRQoL)	A measure of the effects of an illness to see how it affects someone's day-to-day life.
Heterogeneity or Lack of homogeneity	The term is used in meta-analyses and systematic reviews to describe when the results of a test or treatment (or estimates of its effect) differ significantly in different studies. Such differences may occur as a result of differences in the populations studied, the outcome measures used or because of different definitions of the variables involved. It is the opposite of homogeneity.
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.
Inclusion criteria (literature review)	Explicit criteria used to decide which studies should be considered as potential sources of evidence.
Incremental analysis	The analysis of additional costs and additional clinical outcomes with different interventions.
Incremental cost	The extra cost linked to using one test or treatment rather than another. Or the additional cost of doing a test or providing a treatment more frequently.
Incremental cost-effectiveness ratio (ICER)	The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another.
Incremental net benefit (INB)	The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained, then the INB is calculated as: $(£20,000 \times \text{QALYs gained}) - \text{Incremental cost}$ .

Term	Definition
Indirectness	The available evidence is different to the review question being addressed, in terms of PICO (population, intervention, comparison and outcome).
Intention-to-treat analysis (ITT)	An assessment of the people taking part in a clinical trial, based on the group they were initially (and randomly) allocated to. This is regardless of whether or not they dropped out, fully complied with the treatment or switched to an alternative treatment. Intention-to-treat analyses are often used to assess clinical effectiveness because they mirror actual practice: that is, not everyone complies with treatment and the treatment people receive may be changed according to how they respond to it.
Intervention	In medical terms this could be a drug treatment, surgical procedure, diagnostic or psychological therapy. Examples of public health interventions could include action to help someone to be physically active or to eat a healthier diet.
Length of stay	The total number of days a participant stays in hospital.
Licence	See 'Product licence'.
Life years gained	Mean average years of life gained per person as a result of the intervention compared with an alternative intervention.
Likelihood ratio	The likelihood ratio combines information about the sensitivity and specificity. It tells you how much a positive or negative result changes the likelihood that a patient would have the disease. The likelihood ratio of a positive test result (LR+) is sensitivity divided by (1 minus specificity).
Long-term care	Residential care in a home that may include skilled nursing care and help with everyday activities. This includes nursing homes and residential homes.
Logistic regression or Logit model	In statistics, logistic regression is a type of analysis used for predicting the outcome of a binary dependent variable based on one or more predictor variables. It can be used to estimate the log of the odds (known as the 'logit').
Loss to follow-up	A patient, or the proportion of patients, actively participating in a clinical trial at the beginning, but whom the researchers were unable to trace or contact by the point of follow-up in the trial
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
Meta-analysis	A method often used in systematic reviews. Results from several studies of the same test or treatment are combined to estimate the overall effect of the treatment.
Multivariate model	A statistical model for analysis of the relationship between 2 or more predictor (independent) variables and the outcome (dependent) variable.
Negative predictive value (NPV)	In screening or diagnostic tests: A measure of the usefulness of a screening or diagnostic test. It is the proportion of those with a negative test result who do not have the disease and can be interpreted as the probability that a negative test result is correct. It is calculated as follows: $TN/(TN+FN)$
Net monetary benefit (NMB)	The value in monetary terms of an intervention net of its cost. The NMB can be calculated for a given cost-effectiveness threshold. If the threshold is £20,000 per QALY gained, then the NMB for an intervention is calculated as: $(£20,000 \times \text{mean QALYs}) - \text{mean cost}$ .

Term	Definition
	The most preferable option (that is, the most clinically effective option to have an ICER below the threshold selected) will be the treatment with the highest NMB.
Non-randomised intervention study	<p>A quantitative study investigating the effectiveness of an intervention that does not use randomisation to allocate patients (or units) to treatment groups. Non-randomised studies include observational studies, where allocation to groups occurs through usual treatment decisions or people's preferences. Non-randomised studies can also be experimental, where the investigator has some degree of control over the allocation of treatments.</p> <p>Non-randomised intervention studies can use a number of different study designs, and include cohort studies, case-control studies, controlled before-and-after studies, interrupted-time-series studies and quasi-randomised controlled trials.</p>
Number needed to treat (NNT)	<p>The average number of patients who need to be treated to get a positive outcome. For example, if the NNT is 4, then 4 patients would have to be treated to ensure 1 of them gets better. The closer the NNT is to 1, the better the treatment.</p> <p>For example, if you give a stroke prevention drug to 20 people before 1 stroke is prevented, the number needed to treat is 20. See also number needed to harm, absolute risk reduction.</p>
Observational study	<p>Individuals or groups are observed, or certain factors are measured. No attempt is made to affect the outcome. For example, an observational study of a disease or treatment would allow 'nature' or usual medical care to take its course. Changes or differences in one characteristic (for example, whether or not people received a specific treatment or intervention) are studied without intervening.</p> <p>There is a greater risk of selection bias than in experimental studies.</p>
Odds ratio	A measure of treatment effectiveness. The odds of an event happening in the treatment group, expressed as a proportion of the odds of it happening in the control group. The 'odds' is the ratio of events to non-events.
Opportunity cost	The loss of other healthcare programmes displaced by investment in or introduction of another intervention. This may be best measured by the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.
Outcome	<p>The impact that a test, treatment, policy, programme or other intervention has on a person, group or population. Outcomes from interventions to improve the public's health could include changes in knowledge and behaviour related to health, societal changes (for example, a reduction in crime rates) and a change in people's health and wellbeing or health status. In clinical terms, outcomes could include the number of patients who fully recover from an illness or the number of hospital admissions, and an improvement or deterioration in someone's health, functional ability, symptoms or situation.</p> <p>Researchers should decide what outcomes to measure before a study begins.</p>
P value	<p>The p value is a statistical measure that indicates whether or not an effect is statistically significant.</p> <p>For example, if a study comparing 2 treatments found that one seems more effective than the other, the p value is the probability of obtaining these, or more extreme results by chance. By convention, if the p value is below 0.05 (that is, there is less than a 5% probability that the</p>

Term	Definition
	<p>results occurred by chance) it is considered that there probably is a real difference between treatments. If the p value is 0.001 or less (less than a 1% probability that the results occurred by chance), the result is seen as highly significant.</p> <p>If the p value shows that there is likely to be a difference between treatments, the confidence interval describes how big the difference in effect might be.</p>
Perioperative	The period from admission through surgery until discharge, encompassing the preoperative and postoperative periods.
Placebo	A fake (or dummy) treatment given to participants in the control group of a clinical trial. It is indistinguishable from the actual treatment (which is given to participants in the experimental group). The aim is to determine what effect the experimental treatment has had – over and above any placebo effect caused because someone has received (or thinks they have received) care or attention.
Polypharmacy	The use or prescription of multiple medications.
Posterior distribution	In Bayesian statistics this is the probability distribution for a statistic based after combining established information or belief (the prior) with new evidence (the likelihood).
Positive predictive value (PPV)	In screening or diagnostic tests: A measure of the usefulness of a screening or diagnostic test. It is the proportion of those with a positive test result who have the disease and can be interpreted as the probability that a positive test result is correct. It is calculated as follows: $TP/(TP+FP)$
Postoperative	Pertaining to the period after patients leave the operating theatre, following surgery.
Post-test probability	In diagnostic tests: The proportion of patients with that particular test result who have the target disorder (post-test odds/[1 plus post-test odds]).
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Preoperative	The period before surgery commences.
Pre-test probability	In diagnostic tests: The proportion of people with the target disorder in the population at risk at a specific time point or time interval. Prevalence may depend on how a disorder is diagnosed.
Prevalence	See Pre-test probability.
Prior distribution	In Bayesian statistics this is the probability distribution for a statistic based on previous evidence or belief.
Primary care	Healthcare delivered outside hospitals. It includes a range of services provided by GPs, nurses, health visitors, midwives and other healthcare professionals and allied health professionals such as dentists, pharmacists and opticians.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Probabilistic analysis	In economic evaluation, this is an analysis that uses a probability distribution for each input. In contrast, see Deterministic analysis.
Product licence	An authorisation from the MHRA to market a medicinal product.
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good

Term	Definition
	prognosis is associated with low rate of undesirable outcomes; poor prognosis is associated with a high rate of undesirable outcomes.
Prospective study	A research study in which the health or other characteristic of participants is monitored (or 'followed up') for a period of time, with events recorded as they happen. This contrasts with retrospective studies.
Publication bias	Publication bias occurs when researchers publish the results of studies showing that a treatment works well and don't publish those showing it did not have any effect. If this happens, analysis of the published results will not give an accurate idea of how well the treatment works. This type of bias can be assessed by a funnel plot.
Quality of life	See 'Health-related quality of life'.
Quality-adjusted life year (QALY)	A measure of the state of health of a person or group in which the benefits, in terms of length of life, are adjusted to reflect the quality of life. One QALY is equal to 1 year of life in perfect health. QALYS are calculated by estimating the years of life remaining for a patient following a particular treatment or intervention and weighting each year with a quality-of-life score (on a scale of 0 to 1). It is often measured in terms of the person's ability to perform the activities of daily life, freedom from pain and mental disturbance.
Randomisation	Assigning participants in a research study to different groups without taking any similarities or differences between them into account. For example, it could involve using a random numbers table or a computer-generated random sequence. It means that each individual (or each group in the case of cluster randomisation) has the same chance of receiving each intervention.
Randomised controlled trial (RCT)	A study in which a number of similar people are randomly assigned to 2 (or more) groups to test a specific drug or treatment. One group (the experimental group) receives the treatment being tested, the other (the comparison or control group) receives an alternative treatment, a dummy treatment (placebo) or no treatment at all. The groups are followed up to see how effective the experimental treatment was. Outcomes are measured at specific times and any difference in response between the groups is assessed statistically. This method is also used to reduce bias.
RCT	See 'Randomised controlled trial'.
Receiver operated characteristic (ROC) curve	A graphical method of assessing the accuracy of a diagnostic test. Sensitivity is plotted against 1 minus specificity. A perfect test will have a positive, vertical linear slope starting at the origin. A good test will be somewhere close to this ideal.
Reference standard	The test that is considered to be the best available method to establish the presence or absence of the outcome – this may not be the one that is routinely used in practice.
Reporting bias	See 'Publication bias'.
Resource implication	The likely impact in terms of finance, workforce or other NHS resources.
Retrospective study	A research study that focuses on the past and present. The study examines past exposure to suspected risk factors for the disease or condition. Unlike prospective studies, it does not cover events that occur after the study group is selected.

Term	Definition
Review question	In guideline development, this term refers to the questions about treatment and care that are formulated to guide the development of evidence-based recommendations.
Risk ratio (RR)	<p>The ratio of the risk of disease or death among those exposed to certain conditions compared with the risk for those who are not exposed to the same conditions (for example, the risk of people who smoke getting lung cancer compared with the risk for people who do not smoke).</p> <p>If both groups face the same level of risk, the risk ratio is 1. If the first group had a risk ratio of 2, subjects in that group would be twice as likely to have the event happen. A risk ratio of less than 1 means the outcome is less likely in the first group. The risk ratio is sometimes referred to as relative risk.</p>
Secondary outcome	An outcome used to evaluate additional effects of the intervention deemed a priori as being less important than the primary outcomes.
Selection bias	<p>Selection bias occurs if:</p> <ul style="list-style-type: none"> <li>a) The characteristics of the people selected for a study differ from the wider population from which they have been drawn, or</li> <li>b) There are differences between groups of participants in a study in terms of how likely they are to get better.</li> </ul>
Sensitivity	<p>How well a test detects the thing it is testing for.</p> <p>If a diagnostic test for a disease has high sensitivity, it is likely to pick up all cases of the disease in people who have it (that is, give a 'true positive' result). But if a test is too sensitive it will sometimes also give a positive result in people who don't have the disease (that is, give a 'false positive').</p> <p>For example, if a test were developed to detect if a woman is 6 months pregnant, a very sensitive test would detect everyone who was 6 months pregnant but would probably also include those who are 5 and 7 months pregnant.</p> <p>If the same test were more specific (sometimes referred to as having higher specificity), it would detect only those who are 6 months pregnant, and someone who was 5 months pregnant would get a negative result (a 'true negative'). But it would probably also miss some people who were 6 months pregnant (that is, give a 'false negative').</p> <p>Breast screening is a 'real-life' example. The number of women who are recalled for a second breast screening test is relatively high because the test is very sensitive. If it were made more specific, people who don't have the disease would be less likely to be called back for a second test but more women who have the disease would be missed.</p>
Sensitivity analysis	<p>A means of representing uncertainty in the results of economic evaluations. Uncertainty may arise from missing data, imprecise estimates or methodological controversy. Sensitivity analysis also allows for exploring the generalisability of results to other settings. The analysis is repeated using different assumptions to examine the effect on the results.</p> <p>One-way simple sensitivity analysis (univariate analysis): each parameter is varied individually in order to isolate the consequences of each parameter on the results of the study.</p>



Term	Definition
	<p>Multi-way simple sensitivity analysis (scenario analysis): 2 or more parameters are varied at the same time and the overall effect on the results is evaluated.</p> <p>Threshold sensitivity analysis: the critical value of parameters above or below which the conclusions of the study will change are identified.</p> <p>Probabilistic sensitivity analysis: probability distributions are assigned to the uncertain parameters and are incorporated into evaluation models based on decision analytical techniques (for example, Monte Carlo simulation).</p>
Significance (statistical)	A result is deemed statistically significant if the probability of the result occurring by chance is less than 1 in 20 ( $p < 0.05$ ).
Specificity	<p>The proportion of true negatives that are correctly identified as such. For example, in diagnostic testing the specificity is the proportion of non-cases correctly diagnosed as non-cases.</p> <p>See related term 'Sensitivity'.</p> <p>In terms of literature searching a highly specific search is generally narrow and aimed at picking up the key papers in a field and avoiding a wide range of papers.</p>
Stakeholder	<p>An organisation with an interest in a topic that NICE is developing a guideline or piece of public health guidance on. Organisations that register as stakeholders can comment on the draft scope and the draft guidance. Stakeholders may be:</p> <ul style="list-style-type: none"> <li>• manufacturers of drugs or equipment</li> <li>• national patient and carer organisations</li> <li>• NHS organisations</li> <li>• organisations representing healthcare professionals.</li> </ul>
State transition model	See Markov model
Stratification	When a different estimate effect is thought to underlie two or more groups based on the PICO characteristics. The groups are therefore kept separate from the outset and are not combined in a meta-analysis, for example, children and adults. Specified a priori in the protocol.
Sub-groups	Planned statistical investigations if heterogeneity is found in the meta-analysis. Specified a priori in the protocol.
Systematic review	A review in which evidence from scientific studies has been identified, appraised and synthesised in a methodical way according to predetermined criteria. It may include a meta-analysis.
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Transition probability	In a state transition model (Markov model), this is the probability of moving from one health state to another over a specific period of time.
Treatment allocation	Assigning a participant to a particular arm of a trial.
Univariate	Analysis which separately explores each variable in a data set.
Utility	In health economics, a 'utility' is the measure of the preference or value that an individual or society places upon a particular health state. It is generally a number between 0 (representing death) and 1 (perfect health). The most widely used measure of benefit in cost–utility analysis is the quality-adjusted life year, but other measures include disability-adjusted life years (DALYs) and healthy year equivalents (HYEs).

1

## 2.10 Clinical terms used in the guideline.

Term	Definition
Acute setting	A setting with onsite availability of the full range of diagnostic and therapeutic capabilities needed to diagnose and treat acute physical illnesses.
Assessment	An in-depth, and possibly ongoing, process of identifying risk factors.
Carer	Where the term 'carer' is used, this refers to unpaid carers, not paid carers such as care workers.
Comprehensive assessment	An assessment that aims to identify a person's risk factors for falling.
Comprehensive falls management / comprehensive falls interventions	Interventions tailored to address the risk factors identified in a comprehensive assessment. Individual interventions may be directly carried out by one or more health professionals in a specialist falls team (i.e. medication review by the team pharmacist or home hazard modification by the team occupational therapist) or by referrals for further action (i.e. referral to ophthalmology for consideration of cataract surgery).
Dementia	The diagnostic and statistical manual of mental disorders, fourth edition (DSM-IV 1994), expresses the internationally prevailing view of the concept of dementia being a form of memory disturbance, with at least one of the following disturbances of aphasia, apraxia, agnosia and disturbance in executive functioning.
Extended care	A care setting such as a nursing home or supported accommodation.
Home hazard assessment	The assessment of an older person's home environment and the identification of any hazards that may contribute to that person being at risk of falling.
Injurious fall	A fall resulting in a fracture or soft tissue damage that needs treatment.
Multidisciplinary	More than one healthcare professional from different disciplines.
Multifactorial assessment or multifactorial falls risk assessment	An assessment with multiple components that aims to identify a person's risk factors for falling.
Multifactorial interventions (MFI)	In multifactorial interventions, two or more categories of intervention are given, and these are linked to each individual's risk profile. An initial assessment is usually carried out by one or more health professionals and an intervention is then provided or recommendations given, or referrals made for further action. This guideline uses the term 'comprehensive falls management'. The evidence included in the guideline uses 'multifactorial interventions'.
Multiple interventions	In multiple interventions, the same combination of single categories of intervention was delivered to all participants in the group.
Non-acute setting	A setting focused on recovery and rehabilitation, symptom control or palliative care.
Primary prevention	Interventions that aim to prevent the first fall in a person who is vulnerable to falling because of, for example, unsteady gait, but who has not yet fallen.
Rehabilitation	Interventions that are targeted at people who have suffered an injurious fall.
Risk prediction tool	A tool that aims to calculate a person's risk of falling, either in terms of 'at risk/not at risk', or in terms of 'low/medium/high risk', etc.

Term	Definition
Secondary intervention	Interventions that are targeted at a person who has a history of falls.
Self-efficacy	A person's perception of their capability. High self-efficacy relates to increased confidence. This term is referred to in relation to reducing an older person's fear of falling.
Tailored	Intervention packages or programmes that are planned to meet the needs of the particular person.
Targeted	Interventions that are aimed at modifying a particular risk factor or factors.

1

2

## References

- 3 1. Cameron ID, Dyer SM, Panagoda CE, Murray GR, Hill KD, Cumming RG et  
4 al. Interventions for preventing falls in older people in care facilities and  
5 hospitals. The Cochrane database of systematic reviews 2018, Issue DOI:  
6 10.1002/14651858.cd005465.pub4.
- 7 2. Clemson L, Stark S, Pighills AC, Fairhall NJ, Lamb SE, Ali J et al.  
8 Environmental interventions for preventing falls in older people living in the  
9 community. The Cochrane database of systematic reviews 2023, Issue DOI:  
10 10.1002/14651858.cd013258.pub2.
- 11 3. Downey PA, Perry SB, Anderson JM. Screening postmenopausal women for  
12 fall and fracture prevention. Journal of geriatric physical therapy (2001). 2013;  
13 36(3):138-145
- 14 4. Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson  
15 LM et al. Interventions for preventing falls in older people living in the  
16 community. The Cochrane database of systematic reviews 2012, Issue 9.  
17 DOI: 10.1002/14651858.cd007146.pub3.
- 18 5. GRADE Working Group. The Grading of Recommendations Assessment,  
19 Development and Evaluation (GRADE) Working Group website. 2011.  
20 Available from: <http://www.gradeworkinggroup.org/> Last accessed:
- 21 6. Hopewell S, Adedire O, Copsey BJ, Boniface GJ, Sherrington C, Clemson L  
22 et al. Multifactorial and multiple component interventions for preventing falls in  
23 older people living in the community. The Cochrane database of systematic  
24 reviews 2018, Issue DOI: 10.1002/14651858.cd012221.pub2.
- 25 7. Maruish ME. User's Manual for the SF-36v2 Health Survey. 3rd ed. Quality  
26 Metric Incorporated. 2011. Available from:  
27 <https://books.google.co.uk/books?id=a0vYnQEACAAJ>
- 28 8. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C.  
29 PRESS Peer Review of Electronic Search Strategies: 2015 Guideline  
30 Statement. Journal of Clinical Epidemiology. 2016; 75:40-46
- 31 9. National Institute for Health and Care Excellence. Developing NICE  
32 guidelines: the manual. London. National Institute for Health and Care  
33 Excellence, 2014. Available from:

1 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20o>  
2 [vreview](http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20o)

- 3 10. National Institute for Health and Care Excellence. Developing NICE  
4 guidelines: the manual [updated October 2018]. London. National Institute for  
5 Health and Care Excellence, 2014. Available from:  
6 <http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20o>  
7 [vreview](http://www.nice.org.uk/article/PMG20/chapter/1%20Introduction%20and%20o)
- 8 11. National Institute for Health and Clinical Excellence. Social value judgements:  
9 principles for the development of NICE guidance. London. National Institute  
10 for Health and Clinical Excellence, 2008. Available from:  
11 [https://www.nice.org.uk/media/default/about/what-we-do/research-and-](https://www.nice.org.uk/media/default/about/what-we-do/research-and-development/social-value-judgements-principles-for-the-development-of-nice-guidance.pdf)  
12 [development/social-value-judgements-principles-for-the-development-of-nice-](https://www.nice.org.uk/media/default/about/what-we-do/research-and-development/social-value-judgements-principles-for-the-development-of-nice-guidance.pdf)  
13 [guidance.pdf](https://www.nice.org.uk/media/default/about/what-we-do/research-and-development/social-value-judgements-principles-for-the-development-of-nice-guidance.pdf)
- 14 12. Novielli N, Cooper NJ, Abrams KR, Sutton AJ. How is evidence on test  
15 performance synthesized for economic decision models of diagnostic tests? A  
16 systematic appraisal of Health Technology Assessments in the UK since  
17 1997. *Value in Health*. 2010; 13(8):952-957
- 18 13. Organisation for Economic Co-operation and Development (OECD).  
19 Purchasing power parities (PPP). 2012. Available from:  
20 <http://www.oecd.org/std/ppp> Last accessed: 23 July 2024.
- 21 14. Review Manager (RevMan) [Computer program]. Version 5. Copenhagen.  
22 The Nordic Cochrane Centre, The Cochrane Collaboration, 2015. Available  
23 from: <http://tech.cochrane.org/Revman>
- 24 15. Sherrington C, Fairhall NJ, Wallbank GK, Tiedemann A, Michaleff ZA, Howard  
25 K et al. Exercise for preventing falls in older people living in the community.  
26 The Cochrane database of systematic reviews 2019, Issue DOI:  
27 10.1002/14651858.cd012424.pub2.
- 28 16. WinBUGS [Computer programme] version 1.4. Cambridge. MRC Biostatistics  
29 Unit University of Cambridge, 2015. Available from: [http://www.mrc-](http://www.mrc-bsu.cam.ac.uk/software/bugs/the-bugs-project-winbugs/)  
30 [bsu.cam.ac.uk/software/bugs/the-bugs-project-winbugs/](http://www.mrc-bsu.cam.ac.uk/software/bugs/the-bugs-project-winbugs/)

31