

**NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE**

**Multiple Technology Appraisal**

**Prophylactic removal of impacted third molars**

**Draft scope**

**Remit/appraisal objective**

To appraise the clinical and cost effectiveness of the prophylactic removal of impacted mandibular third molars (review of NICE technology appraisal 1).

**Background**

Permanent molar teeth normally erupt from the age of 6 onwards, with the third molars (wisdom teeth) being the last to erupt, usually between the ages of 18 and 24 years. Wisdom teeth may erupt normally into correct dental alignment and function, or conversely develop in non- or minimally functional positions. Impaction occurs when there is prevention of complete eruption due to lack of space, obstruction or development in an abnormal position. This may result in a tooth erupting partially or not at all. Impaction may be associated with pathological changes including pericoronitis (inflammation of the gums surrounding the crown of a tooth), an increased risk of decay and disease in adjacent teeth, and orthodontic problems in later life.

The number of patients having their third molar removed has been thought to have increased to approximately 77,000 patients per year more in 2010<sup>1</sup>.

Third molar procedures are generally suitable for day care management and it is recognised that treatment under local anaesthesia with or without sedation is associated with reduced complication rates. Some of the risks associated with this procedure include infection, delayed healing, nerve damage (which can cause pain or a tingling sensation and numbness in the tongue, lower lip, chin, teeth and gums) and 'dry socket', which is a dull, aching sensation in the gum or jaw.

NICE technology appraisal 1 recommends that impacted wisdom teeth that are free from disease (healthy) should not be operated on. People who have impacted wisdom teeth that are not causing problems should visit their dentist for their usual check-ups. Only people who have diseased wisdom teeth, or other problems with their mouth, should have their wisdom teeth removed. Examples include untreatable tooth decay, abscesses, cysts or tumours, disease of the tissues around the tooth or where the tooth is in the way of other surgery. The guidance states that the standard routine programme of dental care by dental practitioners and/or paraprofessional staff, need be no different, in general, for pathology-free impacted third molars (those requiring no additional investigations or procedures).

Some studies have suggested that third molars that are at an angle facing towards the front of the mouth (mesioangular), away from the tooth next to it

or impacted horizontally may increase the risk of decay in adjacent second molars. This review of NICE technology appraisal 1 will allow for consideration of this evidence.

**The technology**

Reasons for prophylactically removing asymptomatic or pathology-free impacted third molars could be to reduce the risk of infection, untreatable decay, cysts, tumours, and destruction of adjacent teeth and bone.

Conventional extraction of a fully erupted wisdom tooth involves using dental extraction forceps. Surgical removal of a tooth is dependent upon its status such as the degree or complexity of impaction. Generally it involves raising of soft tissue flaps for adequate exposure of bone and/or tooth (using water-cooled rotary instruments with or without a chisel) and removal with forceps.

<b>Intervention(s)</b>	Prophylactic removal of third molars
<b>Population(s)</b>	People with impacted mandibular third molars
<b>Comparators</b>	Standard care without prophylactic removal of third molars
<b>Outcomes</b>	<p>The outcome measures to be considered include:</p> <ul style="list-style-type: none"> <li>• pathology associated with retention of third molars</li> <li>• post-operative complications following extraction</li> <li>• adverse effects of treatment</li> <li>• health-related quality of life.</li> </ul>
<b>Economic analysis</b>	<p>The reference case stipulates that the cost effectiveness of treatments should be expressed in terms of incremental cost per quality-adjusted life year.</p> <p>The reference case stipulates that the time horizon for estimating clinical and cost effectiveness should be sufficiently long to reflect any differences in costs or outcomes between the technologies being compared.</p> <p>Costs will be considered from an NHS and Personal Social Services perspective.</p>
<b>Other considerations</b>	<p>If evidence allows, consideration may be given to the following subgroups:</p> <ul style="list-style-type: none"> <li>• People with mesioangular third molars</li> <li>• People with impacted third molars with or without evidence of pathology</li> </ul>

<p><b>Related NICE recommendations and NICE Pathways</b></p>	<p>Related Technology Appraisals:</p> <p>‘Guidance on the extraction of wisdom teeth’ (2000). NICE Technology Appraisal 1. Under review as part of this appraisal.</p> <p>HealOzone for the treatment of tooth decay (occlusal pit and fissure caries and root caries) (2005). NICE technology appraisal 92. Static list.</p> <p>Related Guidelines:</p> <p>‘Dental checks: intervals between oral health reviews’ (2004). NICE guideline 19. Static list.</p> <p>Related Public Health Guidance/Guidelines:</p> <p>Oral health improvement approaches for local authorities and their partners’ (2014). NICE public health guideline 55.</p> <p>Related Quality Standards:</p> <p><a href="http://www.nice.org.uk/guidance/qualitystandards/qualitystandards.jsp">http://www.nice.org.uk/guidance/qualitystandards/qualitystandards.jsp</a></p> <p>‘Surgical site infection’ (2013). NICE quality standard 49.</p> <p>Related NICE Pathways:</p> <p>Oral and dental health (2015) NICE pathway</p> <p><a href="http://pathways.nice.org.uk/pathways/oral-health-improvement-for-local-authorities-and-their-partners">http://pathways.nice.org.uk/pathways/oral-health-improvement-for-local-authorities-and-their-partners</a></p>
<p><b>Related National Policy</b></p>	<p>Chapter 107. Specialist dentistry services for children and young people</p> <p><a href="http://www.england.nhs.uk/wp-content/uploads/2014/01/pss-manual.pdf">http://www.england.nhs.uk/wp-content/uploads/2014/01/pss-manual.pdf</a></p> <p>Department of Health, NHS Outcomes Framework 2015-2016, Dec 2014. Domain 4a (iii).</p> <p><a href="https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/385749/NHS_Outcomes_Framework.pdf">https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/385749/NHS_Outcomes_Framework.pdf</a></p>

### Questions for consultation

Is the remit ‘to appraise the clinical and cost effectiveness of the prophylactic removal of impacted mandibular third molars’ appropriate?

- Based on the studies considered during the review process, is it more appropriate for the remit of this review to focus only on the prophylactic removal of disease-free (healthy) impacted mandibular third molars; that

is, a partial update of TA1 (specifically recommendations 1.1 and 1.2 of the original guidance)?

- Is there any new evidence to suggest that recommendation 1.3 in the original guidance for third molars with evidence of pathology needs updating?
- Should the review focus only on people with mesioangular third molars? That is, should the population in the scope be 'People with impacted mesioangular mandibular third molars'?

Have all relevant comparators for the prophylactic removal of third molars been included in the scope?

What is considered to be established clinical practice in the NHS for people requiring prophylactic removal of third molars? How should standard care be defined?

Are the outcomes listed appropriate? Are there any other outcomes that should be included?

Are there any other subgroups of people in whom prophylactic removal of third molars is expected to be more clinically effective and cost effective or other groups that should be examined separately?

NICE is committed to promoting equality of opportunity, eliminating unlawful discrimination and fostering good relations between people with particular protected characteristics and others. Please let us know if you think that the scope may need changing in order to meet these aims. In particular, please tell us if the scope:

- could exclude from full consideration any people protected by the equality legislation who fall within the patient population for whom prophylactic removal of third molars might be carried out;
- could lead to recommendations that have a different impact on people protected by the equality legislation than on the wider population, e.g. by making it more difficult in practice for a specific group to access the technology;
- could have any adverse impact on people with a particular disability or disabilities.

Please tell us what evidence should be obtained to enable the Committee to identify and consider such impacts.

Do you consider the prophylactic removal of third molars to be innovative in its potential to make a significant and substantial impact on health-related benefits and how it might improve the way that current need is met (is this a 'step-change' in the management of the condition)?

Do you consider that the prophylactic removal of third molars can result in any potential significant and substantial health-related benefits that are unlikely to be included in the QALY calculation?

Please identify the nature of the data which you understand to be available to enable the Appraisal Committee to take account of these benefits.

### References

1. McArdle LW and Renton T (2012). The effects of NICE guidelines on the management of third molar teeth. *British Dental Journal* 213, E8