



University of Exeter

Medical School



Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

External Assessment Group report

NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE

Early Value Assessment Programme

Produced by

Peninsula Technology Assessment Group (PenTAG)
University of Exeter Medical School
South Cloisters
St Luke's Campus
Heavitree Road
Exeter
EX1 2LU

Authors

Maxwell S. Barnish
Alan Lovell
Sophie Robinson
Elham Nikram

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Date: May 2023

1 of 100

Caroline Farmer
Edward C.F. Wilson
Peninsula Technology Assessment Group (PenTAG), University
of Exeter Medical School, Exeter

Correspondence to Dr Alan Lovell

[REDACTED]
[REDACTED]

Date completed 22/05/2023

Contains confidential information: Yes

Number of attached appendices: 4

Purpose of the assessment report

The purpose of this External Assessment Group (EAG) report is to review the evidence currently available for included technologies and advise what further evidence should be collected to help inform decisions on whether the technologies should be widely adopted in the NHS. The report may also include additional analysis of the submitted evidence or new clinical and/or economic evidence. NICE has commissioned this work and the report forms part of the papers considered by the Medical Technologies Advisory Committee when it is making decisions about the early value assessment.

Declared interests of the authors

Description of any declared interests with related companies, and the matter under consideration. See [NICE's Policy on managing interests for board members and employees](#).

None.

Acknowledgements

The EAG acknowledges the administrative support of Sue Whiffin and Jenny Lowe (both PenTAG) and Specialist Committee Member (SCM) input from Robert Dudley (Gateshead Early Intervention in Psychosis service, CNTW Foundation Trust), Rhema Immanuel (East London and North East London NHS Foundation Trust) and Elizabeth Murphy (Greater

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Date: May 2023

2 of 100

Manchester Mental Health NHS Foundation Trust (GMMH NHS)). We also thank Prof Sam Vine (University of Exeter) for input on VR-based technology.

Copyright: © 2023, PenTAG, University of Exeter

Responsibility for report

The views expressed in this report are those of the authors and not those of NICE. Any errors are the responsibility of the authors.

Contents

Abbreviations	7
1. Executive summary	10
2. Decision problem.....	12
3. Overview of the technology	14
3.1. Purpose of the medical technology.....	14
3.2. Product properties	15
3.3. Comparator	17
4. Clinical context	18
4.1. Care pathway	18
4.2. User issues and preferences	19
5. Special considerations, including issues related to equality.....	20
6. Potential implementation issues	21
7. Clinical evidence selection	22
7.1. Search strategy	22
7.2. Study selection.....	22
8. Clinical evidence review	25
8.1. Overview of methodologies of all included studies.....	30
8.2. Study design, intervention and comparator	30
8.3. Participants and setting	31
8.4. Outcomes.....	31
8.5. Quality of included studies.....	32

8.6.	Results from the evidence base	33
8.7.	Additional indirectly relevant evidence.....	37
9.	Adverse events	39
10.	Economic evidence	40
10.1.	Published economic evidence	40
10.2.	Conceptual modelling.....	44
10.3.	Model Inputs.....	47
10.4.	Approach to Analysis.....	54
10.5.	Results from the economic modelling	56
10.6.	Commentary.....	59
11.	Interpretation of the evidence	62
11.1.	Interpretation of the clinical and economic evidence.....	62
11.2.	Integration into the NHS	64
11.3.	Evidence gap analysis.....	64
11.4.	Ongoing studies	66
11.5.	Summary and conclusions of evidence gap analysis.....	68
11.6.	Key areas for evidence generation	69
12.	Conclusions.....	70
12.1.	Conclusions from the clinical evidence	70
12.2.	Conclusions from the economic evidence	70
12.3.	Summary of the combined clinical and economic sections	71
13.	References.....	72
14.	Appendices	80

Appendix A: Searches for clinical and cost effectiveness evidence	80
Appendix B: PRISMA flow diagram	85
Appendix C: List of excluded studies and studies awaiting assessment	86
Appendix D. Additional study results	91
Appendix E: Data from Sensitivity Analyses	94

Abbreviations

Term	Definition
A&E	Accident and emergency
Afc	Agenda for change
ASD	Autism spectrum disorder
BAI	Beck Anxiety Inventory
CBT	Cognitive behavioural therapy
CEA	Cost-effectiveness analysis
CE mark	<i>Conformité européenne</i> (European conformity) marking
CI	Confidence interval
CORE-OM	Clinical Outcomes in Routine Evaluation – Outcome Measure
CRD	Centre for Reviews and Dissemination
DP	Decision problem
DTAC	Digital Technology Assessment Criteria
EAG	External assessment group
EE	Economic evaluation
EQ-5D	EuroQoL-5 dimensions
EQ-5D-5L	EuroQoL-5 dimensions 5-level
EVA	Early value assessment
GAD-7	Generalised Anxiety Disorder Assessment 7
GP	General practitioner
HRQoL	Health-related quality of life
HRSD	Hamilton Rating Scale for Depression
HTA	Health technology assessment
IAPT	Improving Access to Psychological Therapies
ICD	International Classification of Diseases
ICER	Incremental cost effectiveness ratio
ICTRP	International Clinical Trials Registry Platform
INAHTA	International Network of Agencies for Health Technology Assessment
IQR	Interquartile range
ITT	Intention to treat
MANCOVA	Multivariate analysis of covariance
MANOVA	Multivariate analysis of variance
MAUDE	Manufacturer and User Facility Device Experience

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

MCID	Minimally clinically important difference
MID	Minimally important difference
MeSH	Medical subject headings
MHRA	Medicines & Healthcare products Regulatory Agency
MTEP	Medical Technologies Evaluation Programme
N/A	Not applicable
NG	NICE guideline
NHS	National Health Service
NICE	National Institute for Health and Care Excellence
NLM	National Library of Medicine
NR	Not reported
O-AS	Oxford Agoraphobic Avoidance Scale
O-BAT	Oxford Behavioural Avoidance Test
ONS	Office for National Statistics
OWSA	One-way sensitivity analysis
PenTAG	Peninsula Technology Assessment Group
PHQ-9	Patient Health Questionnaire-9
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PSS	Personal social services
PSSRU	Personal Social Services Research Unit
PW	People with
QALY	Quality-adjusted life year
RCT	Randomised controlled trial
ReQoL	Recovering Quality of Life quality
RWE	Real world evidence
SA	Sensitivity analysis
SCM	Specialist Committee Member
SD	Standard deviation
SE	Standard error
SIGN	Scottish Intercollegiate Guidelines Network
TAU	Treatment as usual
UK	United Kingdom
UKCA	United Kingdom Conformity Assessed marking
VAS	Visual analogue scale
VR	Virtual reality

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

1. EXECUTIVE SUMMARY

Quality and relevance of clinical evidence

The EAG considered that the clinical effectiveness for all technologies was uncertain, reflecting the relatively early stage of development of the interventions. No evidence was available for Invirto. For the other three technologies, Amelia Virtual Care, gameChangeVR and XR Therapeutics, clinical effectiveness evidence was limited. There was one key effectiveness study per technology. For XR Therapeutics, evidence was limited to one single arm study of eight people with autism spectrum disorder (ASD), which reached equivocal conclusions about effectiveness. For Amelia Virtual Care, the included single-arm study measured acceptance and adherence of the technology (called Psious at the time of the trial, but now called Amelia), but did not report any clinical outcomes. In the gameChangeVR RCT, there was some evidence of a benefit of gameChangeVR over treatment as usual in terms of agoraphobia symptoms. However, it should be noted that the magnitude and duration of benefit is uncertain, the treatment as usual comparator profile was not profiled precisely in published information, there was no evidence of benefit on wider secondary outcome measures, and the evidence comes only from one trial.

Adverse event data were only provided for one intervention, gameChangeVR, and these data came from an RCT, rather than real world safety observation.

Quality and relevance of economic evidence

The EAG identified one published economic evaluation of the cost-effectiveness of gameChangeVR+treatment as usual (TAU) vs TAU in people with psychosis and agoraphobia, conducted alongside the RCT. The EAG noted the results were highly sensitive to a small number (n=4) of participants in an inpatient setting. EAG's exploratory modelling of gameChangeVR suggests a high degree of uncertainty and whilst point estimates suggest it may not represent a cost-effective use of NHS resources when judged against the NICE reference case, the uncertainty is such that its being cost-effective cannot be ruled out at this time.

Evidence Gap Analysis

There was uncertainty surrounding the clinical effectiveness of all interventions. A number of evidence gaps were identified. These included 1) the differences in populations and

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

outcomes studied for each intervention, 2) the absence of any evidence for Invirto, economic evidence for XR Therapeutics, and clinical and economic evidence for Amelia 3) differences in comparators across trials, 4) published evidence not being available for all outcomes, 5) an absence of evidence on the durability of the effect of VR-based therapies, and 6) safety data only being available for gameChangeVR.

2. DECISION PROBLEM

Table 1 details the final scope issued by NICE for this EVA, defined per element of assessment. SCMs generally considered the scope to be well aligned to NHS practice.

Table 1: Summary scope of the assessment

Element of assessment	Final scope issued by NICE
Population	People aged 16 years and over with agoraphobia or agoraphobic avoidance
Subgroups	<p>If the evidence allows the following subgroups will be considered:</p> <ul style="list-style-type: none"> • people with psychosis who have agoraphobia or agoraphobic avoidance • agoraphobia or agoraphobic avoidance that occurs with other mental health problems including but not limited to severe mental illness • high or severe agoraphobic avoidance
Interventions (proposed technologies)	<p>Virtual reality (VR) for agoraphobia and agoraphobic avoidance, delivered with the support of a mental health worker or as part of face-to-face therapy or teletherapy. Namely:</p> <ul style="list-style-type: none"> • Amelia Virtual Care (Amelia Virtual Care) • gameChangeVR (Oxford VR) • Invirto (Invirto) • XR Therapeutics (XR Therapeutics) <p>VR interventions would be offered in addition to standard care for co-occurring mental health conditions.</p>
Comparator	<p>Standard care which may include any combination of:</p> <ul style="list-style-type: none"> • Guided self-help • Cognitive behavioural therapy (CBT) • Exposure therapy • Applied relaxation • Antidepressants licensed for the treatment of panic disorder • Oral antipsychotic medication • Simple contact and monitoring with services.
Healthcare setting	Outpatient clinics, inpatient settings or home-based care

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Element of assessment	Final scope issued by NICE
Outcomes	<p>Intermediate measures for consideration may include:</p> <ul style="list-style-type: none"> • Patient choice and preferences • Acceptability and satisfaction • Accessibility and digital access • Intervention adherence and completion • Intervention-related adverse events • Device-related adverse events
	<p>Clinical outcomes for consideration may include:</p> <ul style="list-style-type: none"> • Change in agoraphobia symptoms • Change in other psychological symptoms • Global functioning and work and social adjustment • Rates of recovery, time to recovery • Rates of relapse or deterioration, time to relapse or deterioration
	<p>Patient-reported outcomes for consideration may include:</p> <ul style="list-style-type: none"> • Health-related quality of life • Recovering quality of life • Patient experience • Social contact
	<p>Costs will be considered from an NHS and Personal Social Services perspective. Costs for consideration should include:</p> <ul style="list-style-type: none"> • Costs of the standalone VR headsets • Costs of the technologies including license fees • Healthcare professional grade and time • Cost of other resource use (e.g. associated with managing anxiety, adverse events or complications): <ul style="list-style-type: none"> ○ GP or mental health team appointments ○ Healthcare professional training
Time horizon	The time horizon for estimating the clinical and economic value should be sufficiently long to reflect any differences in costs or outcomes.

Abbreviations: CBT, Cognitive Behavioural Therapy; EVA, Early Value Assessment; GP, General Practitioner; NICE, National Institute for Health and Care Excellence; SCM, Specialist Committee Member; VR, virtual reality

3. OVERVIEW OF THE TECHNOLOGY

3.1. Purpose of the medical technology

Agoraphobia is an anxiety disorder characterised by marked and excessive fear of being in situations where escape may be difficult or help may not be available^{1,2}. This experience may be described in terms of feeling threatened or worried about leaving home or other place of safety. Agoraphobia involves fear and avoidance of places or situations that might cause panic and feelings of being trapped, helpless or embarrassed. It constitutes a form of anxious avoidance of everyday situations and may co-occur with other mental health conditions, such as panic disorder, depression, social anxiety and psychosis.

Given the high prevalence of mental health conditions and the importance of early intervention, improving and widening mental health services has been identified as a key priority for the NHS.¹ The most recent Adult Psychiatric Morbidity Survey reports that only one in three people with a common mental health condition accesses treatment.³ Barriers to accessing face-to-face treatments such as individual cognitive behavioural therapy (CBT) include a shortage of trained healthcare professionals and limited clinical resources, while agoraphobia may further impact a person's ability to access mental health services and support and may lead to discontinuation, for example through difficulty tolerating exposure therapy. Agoraphobia may also often be untreated or undertreated when it co-occurs with other mental health conditions.

Virtual reality (VR)-based interventions may increase access to care by offering another treatment channel for people with agoraphobia or agoraphobic avoidance. VR is a simulated 3-dimensional environment with scenes and objects that people can explore and interact with, most typically using a VR headset. Alternatively, images can be projected onto a large hemispherical screen. This can create an immersive experience that is thought to trigger emotional responses similar to those in real-world situations. VR may be used as a tool in therapy sessions or as a vehicle to deliver a digital intervention with the support of a mental health worker, in particular exposure therapy. It can allow people to immerse themselves in real-world situations while being in the safety of their home or clinic. Virtual environments can be adjusted based on a person's needs and individual treatment plan. This could allow more gradual exposure to stressful situations, which may increase comfort and confidence in completing interventions.

Using VR as a treatment modality would support remote treatment delivery. This would allow some people to receive treatment at home and may address barriers to accessing treatment for those who cannot or prefer not to attend face-to-face treatment. This could facilitate faster access to symptom management. The scalable nature of VR-based interventions could allow mental health professionals to treat more people in less time and therefore use time and resources more efficiently compared with standard care interventions for agoraphobia and agoraphobic avoidance.

3.2. Product properties

This scope focuses on VR technologies for treating agoraphobia and agoraphobic avoidance that meet the following criteria:

- Can be used as a platform to treat agoraphobia and agoraphobic avoidance either as a digital intervention with the support of a mental health worker or as a tool in face-to-face therapy or teletherapy.
- Meet the standards within the digital technology assessment criteria (DTAC), including the criteria to have a European conformity (CE) or United Kingdom Conformity Assessed (UKCA) mark where required. Products may also be considered if they are actively working towards required CE or UKCA mark and meet all other standards within the DTAC.
- Are available for use in the NHS.

In total, four VR technologies for treating agoraphobia and/or agoraphobic avoidance were included in the scope. For the included technologies, the EAG noted that version changes may limit the generalisability of evidence. Furthermore, it should be noted that evidence evaluating the interventions did not always use the present brand names.

3.2.1. Amelia Virtual Care (Amelia Virtual Care)

Amelia Virtual Care is a VR platform designed to be used by therapists to support the treatment of mental health disorders. It is delivered under the guidance of a therapist in clinical settings or remotely using Amelia's smartphone app. It also offers a homework feature with virtual mindfulness and relaxation sessions. Amelia helps therapists to facilitate the delivery of evidence-based treatment including gradual exposure, mindfulness-based cognitive therapy and desensitisation. Amelia has over 100 virtual environments that can be configured and personalised to a person's needs using a simple control panel.⁴ Amelia was

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

previously called Psious. The company has confirmed that this was largely a re-branding exercise with no changes to the intervention content, though some changes had been made to environments and avatar options.

3.2.2. gameChangeVR (Oxford VR)

gameChangeVR was designed to treat agoraphobia and agoraphobic avoidance in people with psychosis. It delivers cognitive therapy within a VR environment and is compatible with a range of VR equipment that uses six degrees of freedom tracking (i.e the range of motion for objects within here dimensional space). This includes the HTC Vive, Meta Quest and Pico Neo headsets. The treatment includes repeated behavioural experiments using the headset to simulate different real-life situations (including visiting a café, shop, pub, street, doctor's office and bus) to help people test their fear expectations. It is delivered in around six weekly 30-minute sessions. Treatment is facilitated by a virtual coach to support the use of techniques and assist people to overcome their difficulties. It should also be supported by a mental health worker either remotely or in the room during sessions to help people maximise their learning from gameChangeVR in the real world. It may be used with outpatients in clinics or at home. The company suggested that it may also benefit people in inpatient settings. The EAG noted that the name of this technology has changed since the CE mark was issued. gameChangeVR was initially called Social Avoidance. The company request for information states that this name has not yet been updated on the CE mark, but this will be done on the next iteration of the Declaration of Conformity and Instructions for Use. The company advised that the latest version of the technology functions 'untethered'; that is to say, removing the need for a high specification desktop or laptop computer and allowing it to run on the latest stand-alone headset devices.

3.2.3. Invirto (Invirto)

Invirto offers app-based cognitive behavioural therapy (CBT) content and exposure exercises in VR using a VR headset. The programme includes psychoeducation via the app, interoceptive exposure, situational exposure with VR, anxiety diary, monitoring and progress reports, and relaxation and mindfulness exercises. Its programme for agoraphobia includes over 15 situational exposure scenarios such as driving a car, using an elevator, public transport and shopping. These are prepared and followed up as behavioural experiments in the app. Invirto also has programmes for panic disorder and social phobia. No company reference list was provided, and the company did not respond to queries by NICE. No

evidence was identified by the EAG and therefore this technology was noted but not explored in detail.

3.2.4. XR Therapeutics (XR Therapeutics)

XR Therapeutics (XRT) offers a VR-based treatment platform to help reduce anxieties and treat phobias including agoraphobia. It was designed to be combined with face-to-face CBT and allows therapists to tailor digital scenes to a person's individual needs. Treatment can be adapted in real time allowing therapists to manage the rate of exposure and the intensity of situations. Digital scenes can also be personalised in line with a person's background and cultural preferences. XRT does not require the use of a VR headset; VR technology is used to project digital scenes onto a curved white screen to recreate situations such as being in a supermarket in a safe setting. The company said this is easy to install, operate and maintain. The company has advised that originally the technology used third-party hardware called the Blue Room, as referenced in Maskey et al.⁵ However, XR Therapeutics has now developed the Immersive Studio, in-house hardware and software. The company states that the *"treatment is still based on the initial research, but we have significantly advanced the technology being used"* (Personal Communication, XR Therapeutics, April 2023).

3.3. Comparator

The comparator is standard care which may include guided self-help, CBT with exposure therapy, applied relaxation, antidepressants and/or simple contact and monitoring with services.

4. CLINICAL CONTEXT

The target population for this assessment is described in section 3.1.

4.1. Care pathway

The NHS recommends a stepped care approach for treating agoraphobia and any underlying panic disorder.¹ The first step involves recognition and accurate diagnosis, including identification of any comorbidities. The second step involves guided self-help. The third step involves more intensive treatments such as CBT or medication.

Psychological treatments for agoraphobia include self-help programmes and CBT with exposure therapy and is usually delivered in primary care or in local settings such as NHS Talking Therapies. However, agoraphobia can be treated in secondary care and inpatient settings, particularly due to co-occurring complex or severe mental health problems.

Treatment for agoraphobia and agoraphobic avoidance may encourage self-help techniques and lifestyle changes such as exercise to help people relieve and manage their symptoms. People may be offered individual guided self-help which is based on CBT and delivered with the support of a therapist. If needed or preferred, more intensive treatments should be offered such as CBT or applied relaxation.

The NICE clinical guideline on generalised anxiety disorder and panic disorder in adults⁶ recommends that people with moderate to severe panic disorder with or without agoraphobia should be offered CBT or an antidepressant. For those diagnosed with psychosis, the NICE clinical guideline⁷ recommends oral antipsychotic medication along with psychological interventions including family intervention and individual CBT. SCM input advised that only a minority of people with psychosis receive the recommended treatment and may instead be offered antipsychotic medication and simple contact and monitoring with services.

One SCM noted that the care pathways outlined in the scope reflect routine practice well. Generally, it was noted that there is no established care pathway for agoraphobia within a psychosis setting and that people with agoraphobia and psychosis may not necessarily be diagnosed with agoraphobia due to symptom overlap and a focus on treating psychosis symptoms as the primary diagnosis.

4.1.1. Current use of VR technologies to treat agoraphobia

The companies advised that their respective VR technologies are in current use within the NHS for agoraphobia or psychosis: Amelia Virtual Care (six NHS Trusts), gameChangeVR (two NHS Trusts on an investigational basis) and XR Therapeutics (four NHS Trusts). SCMs were not aware of VR technology use in routine NHS practice for agoraphobia.

It is anticipated that VR-based interventions⁴ would be offered after clinical assessment and diagnosis and as an alternative or addition to standard care psychological interventions for agoraphobia and agoraphobic avoidance. VR could be delivered by a therapist as part of face-to-face therapy or teletherapy or used as a standalone intervention with the support of a mental health worker such as an assistant psychologist, peer support worker or therapist. The level of support provided may vary depending on the intervention and the person's need.

4.2. User issues and preferences

It is anticipated that VR technologies may increase treatment options and access to care. They may enable some people to receive treatment at home. This may be especially beneficial for people with agoraphobia or agoraphobic avoidance who have difficulty leaving their homes to access standard care. VR technologies may also help people to test their fear expectations in a setting where they feel safe. People may feel more comfortable completing behavioural experiments in VR and this could increase their confidence in performing these tasks in real-world settings. People may be more motivated to use and engage with VR if they have sufficient digital skills and prefer remote or digital interventions to face-to-face therapy.

However, it should be taken into consideration that some people may have a preference for face-to-face therapy and may choose not to use VR technology. There may be some concerns about the level of support provided and uncertainty around how treatment may be delivered. Lay SCM input noted that some may be disappointed if scenes in VR interventions are not photorealistic. The realism of a VR environment is important both in terms of how it influences the perception of the user⁸ and how it impacts the intended outcomes of the VR experience.⁹

5. SPECIAL CONSIDERATIONS, INCLUDING ISSUES RELATED TO EQUALITY

A number of potential equality issues have been identified.

People using VR-based technologies at home would be provided with the VR device through their mental health service. However, some VR technologies require Wi-Fi to use the intervention or upload content. This could lead to challenges for those in rural areas with poor internet services or those who do not have internet access in their homes.

Additional support may be required for those with poor digital literacy or poor internet connectivity. Furthermore, people with visual or cognitive impairment, problems with manual dexterity, people with a learning disability or who are unable to read or understand health-related information, and people who cannot read English may need additional support to use the technologies, although the EAG is aware that software packages can be (and have been) adapted to overcome some of these issues. Some people would benefit from VR in languages other than English. In general, views of mental health problems and interventions may vary according to their ethnicity, religious or cultural background.

VR may not be suitable for use by people with photosensitive epilepsy, significant visual, auditory, or balance impairment, organic mental disorder, primary diagnosis of alcohol or substance disorder or personality disorder, significant learning disability, or active suicidal plans.⁴ Some VR interventions may involve moving around the room or standing. This may be difficult for some people with physical disabilities or additional accessibility needs.

6. POTENTIAL IMPLEMENTATION ISSUES

The NICE adoption and implementation team consulted clinical experts and noted several potential implementation issues for VR technologies:

- Safety and comfort – including dizziness, the amount of space needed, wearing glasses.
- Patient selection – ensuring VR technologies are used for people with the target condition.
- Acceptability.

However, some software can be (and has been) adopted to overcome some of these difficulties. The EAG also notes that not all interventions require the use of VR headsets. Other issues relevant to this appraisal are outlined in the scope.

7. CLINICAL EVIDENCE SELECTION

7.1. Search strategy

Search strategies were based on those devised during the initial scoping searches by NICE Information Services with some amendments. The search strategies used relevant search terms, comprising a combination of indexed keywords (e.g., Medical Subject Headings, MeSH) and free-text terms appearing in the titles and/or abstracts of database records and were adapted according to the configuration of each database. No date, language or publication status (published, unpublished, in-press, and in-progress) limits were applied. Searches for clinical and cost-effectiveness were combined and carried out in one search strategy.

Following deduplication, a total of 318 records of potentially relevant evidence on clinical and/or cost effectiveness were retrieved. Databases searched were Medline (including Medline in Process), Embase, PsycInfo, Cochrane, INAHTA, CEA Registry and ScharrHUD. Additional trial registries searched were Clinicaltrials.gov (NLM) and ICTRP (WHO). The websites of the individual companies were searched; NICE and SIGN websites were searched for related guidelines; MAUDE and MHRA were searched for adverse events data; the company submission references were also scanned for additional references.

The search strategies are presented in Appendix A.

7.2. Study selection

The abstracts and titles of references retrieved by the searches were screened for relevance. Full paper copies of potentially relevant studies were obtained. The retrieved articles were assessed for inclusion against pre-specified inclusion/exclusion criteria. At each stage of screening, a minimum of 10% of records were independently screened by a second reviewer. Discrepancies were resolved by discussion, with involvement of a third reviewer, where necessary. All duplicate papers were excluded.

This assessment looked across a range of evidence types, including RCTs and real-world evidence, to inform clinical effectiveness.

The following study types were excluded:

- Animal models

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

- Pre-clinical and biological studies
- Narrative reviews, editorials, opinion pieces
- Meeting abstracts for studies where full-text papers were available. If studies were only available as meeting abstracts, inclusion depended on sufficient information being available to offer meaningful critique.
- Studies not available in the English language.

Eligible studies assessed a scoped intervention (Amelia Virtual Care, gameChangeVR, XR Therapeutics or Invirto) in a population of adults (16+) with agoraphobia or agoraphobic avoidance.

Studies were not excluded if the comparator did not match the scope or if the outcomes did not match the scope, provided the outcomes appeared reasonable and could offer useful information in the context of the appraisal. Studies conducted in other populations were not included, but a brief commentary on these indirectly relevant studies was provided. This did not include studies conducted exclusively in a paediatric population, as it was not considered that these would inform the present decision problem. In the event that studies investigated a closely related population, such as a study with a majority of adult participants but some paediatric participants, these could be included, depending on what other evidence was identified.

A PRISMA flow diagram is provided as Appendix B.

Data were extracted from included studies by one reviewer into a bespoke database and a sample of at least 10% was checked by another reviewer. Generalisability to NHS practice was considered in the interpretation of the findings.

Due to time and resource constraints associated with conducting an EVA, the EAG did not conduct formal risk of bias assessment of the included studies. Generally, RCTs could be considered more robust than other study types, but it depends on the details of how each study was conducted, and how well the trial setting reflects routine clinical practice in terms for example of eligible population and staff attention, which could affect generalisability. For example, adverse event data may be better collected via cohorts or other longitudinal study designs.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

The EAG identified the key studies for detailed assessment for each technology, based on a preference for studies conducted within a UK setting and studies near the top of the hierarchy of evidence¹⁰ (such as RCTs) where available and then assessed whatever outcome data were available within these key studies, supplementing this with additional data from other studies where it was considered appropriate.

8. CLINICAL EVIDENCE REVIEW

The EAG identified a total of nine publications, comprising four unique studies, that were relevant to the present decision problem. One included paper¹¹ was an economic analysis of an included study and is explored in more detail in section 10.1 of this report.

Table 2 presents a detailed overview of the study design and characteristics of each included study. No evidence was identified for Invirto.

Table 2: Included clinical effectiveness studies

References	Study name	Country	Method	Sample	Intervention	Comparator(s)	Outcomes
<i>Amelia Virtual Care</i>							
Gelabert and Giner, 2018 ¹²	NR	Spain (Catalonia) ^a	Single-arm study	51 adults over 18 with a diagnosis of agoraphobia or panic disorder with agoraphobia (only 42 were included in analysis)	Amelia Virtual Care ^a (conference abstract, so no further details about intervention delivery available)	None	Follow-up was 6 months <ul style="list-style-type: none"> • Number of intervention sessions needed • Therapeutic adherence • Patient satisfaction
<i>gameChangeVR</i>							
Altunkaya et al, 2022; ¹¹ Bond et al, 2023; ¹³ Freeman et al, 2022a,b,c ¹⁴⁻¹⁶	gameChangeVR trial.	UK	RCT, with embedded qualitative study and economic evaluation	346 people (aged 16 or older) with a clinical diagnosis of schizophrenia spectrum disorder or an affective diagnosis with psychotic symptoms, and who have self-reported difficulties in going outside due to anxiety	gameChangeVR (approximately 6 sessions – with some variation if needed – of 30 minutes each over 6 weeks)	Treatment as usual (no published details, but information provided in correspondence from Prof Freeman is discussed in text)	Follow-up in primary analysis paper 6 months Primary: <ul style="list-style-type: none"> • Agoraphobic avoidance (Oxford Agoraphobic Avoidance Scale; Oxford-Behavioural Avoidance Test). Secondary: <ul style="list-style-type: none"> • Agoraphobia (Agoraphobia Mobility Inventory-Avoidance-scale), • Suicidal ideation (Columbia Suicidal Severity Rating Scale)

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

References	Study name	Country	Method	Sample	Intervention	Comparator(s)	Outcomes
							<ul style="list-style-type: none"> Paranoia (Revised Green et al Paranoid Thoughts Scale) Paranoia worries (Paranoia Worries Questionnaire) Depression (PHQ-9) Activity levels (actigraphy over 7 days; time budget) Quality of life (EQ-5D) Recovery Quality of Life questionnaire; Questionnaire about the Process of Recovery) Safety (serious adverse events; adverse event profile) Participant satisfaction Participant experiences with gameChangeVR Economic outcomes (see Section 10.1)
Knight et al, 2021; ¹⁷ Lambe et al, 2020 ¹⁸	gameChangeVR project	UK	Person-centred design process	Clinical psychologists, programmers, animators, designers, product managers,	Development of gameChangeVR (this study profiles the intervention development process rather	None	<ul style="list-style-type: none"> Multistakeholder perspectives to inform the development of gameChangeVR User acceptability ratings

References	Study name	Country	Method	Sample	Intervention	Comparator(s)	Outcomes
				producers, writers, researchers, 3D artists, mental health advocates, and people with lived experience of psychosis	than assessing the effectiveness of its delivery)		
<i>XR Therapeutics</i>							
Maskey et al, 2019 ⁵	NR	UK	Single-arm study	8 adults (18-60) with autism who reported 'fears and phobias')	XR Therapeutics ^a (2 visits each comprising 2 sessions of 20-30 minute with a 15 minute break)	None	Follow-up was 6 months <ul style="list-style-type: none"> • Participation • Retention • Symptom change (Target Behaviors) • Anxiety (Beck Anxiety Inventory, Generalized Anxiety Disorder 7, self-reported ratings of confidence in managing target anxiety situation) • Depression (Patient Health Questionnaire – 9) • Quality of life (WHOQOL-BREF).

a = confirmed through correspondence with the company.

Abbreviations: BDI, Beck Depression Inventory; dCBT, Computerised Cognitive Behavioural Therapy; CES-D, Center for Epidemiologic Studies Depression Scale; CORE-OM, Clinical Outcomes in Routine Evaluation – Outcome Measure; GAD, Generalised Anxiety Disorder assessment; HRSD, Hamilton Rating Scale for Depression; IAPT, Improving

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Access to Psychological Therapies; MADRS, Montgomery-Asberg Depression Rating Scale; PHQ, Patient Health Questionnaire; QIDS-SR, Quick Inventory of Depressive Symptomatology-Self Report; RCT, Randomised controlled trial; UK, United Kingdom; USA, United States.

8.1. Overview of methodologies of all included studies

All studies described in Table 2 had some methodological limitations or misalignment with the NICE decision problem for this appraisal.

8.2. Study design, intervention and comparator

There was a total of four unique included studies (two for gameChangeVR and one for each of Amelia and XR Therapeutics), published in nine papers. The included studies for gameChangeVR and XR Therapeutics were conducted in a UK setting, while the included study for Amelia was conducted in Spain. There was one RCT –for gameChangeVR.¹⁵

The comparator in the gameChangeVR trial¹⁵ was treatment as usual, which was not described clearly in the papers or company submission, although resource use was measured in the companion economic evaluation.¹¹ Some information on treatment as usual was provided in the supplementary material but not signposted in the main paper. Treatment as usual differed between trusts and centres, so could not be considered consistent across participants. Correspondence from the lead author of the gameChange trial, Professor Daniel Freeman, clarified that in the treatment as usual arm 92% of participants were on antipsychotic medication, 58% were on an antidepressant, 85% were seeing a care co-ordinator (believed to mainly be monthly) and around two thirds (exact value not provided) were seeing a psychiatrist every few months. This study took place in a UK setting, but standard care as a comparator also has known limitations due to variability in routine practice between centres.

There was no comparative evidence for Amelia or XR Therapeutics.

Evidence gap: No clinical effectiveness evidence was available for Amelia, and no RCT evidence was available for XR Therapeutics. Pivotal evidence for gameChangeVR was as adjunct to treatment as usual against treatment as usual alone and no details were provided about what this comprised in the main trial paper. However, this can be partially inferred from the companion economic evaluation and the lead author communicated some further details. Any future research involving a treatment as usual (TAU) arm needs careful definition of TAU.

8.3. Participants and setting

The key studies for gameChangeVR and XR Therapeutics were conducted in a UK setting. The evidence for Amelia was from a Spanish setting, which the EAG considered to be relatively comparable, although there are likely to be some differences in terms of how care is structured and delivered. Studies were generally conducted in an outpatient setting, although four participants in the gameChange trial were psychiatric inpatients and incurred particularly high costs in the model.

All included studies described participants who are broadly relevant to the decision problem. The evidence base for gameChangeVR was specifically in the context of schizophrenia spectrum disorder or psychosis, which aligned to a subgroup of interest in the NICE scope. The single-arm study by Gelabert and Giner (2018),¹² for Amelia considered participants with a primary diagnosis of agoraphobia or panic disorder with agoraphobia. The Maskey et al⁵ study for XR therapeutics included only eight participants consisting of adults with autism who had various ‘fears and phobias’, only two of whom reported phobias related to agoraphobic symptoms (open spaces and crowded buses). This study therefore is only partially relevant for this decision problem.

Evidence gap: Evidence for Amelia Virtual Care was only available in a Spanish setting. There was no evidence for XR therapeutics in a population with agoraphobia (only 2/8, 25%) of participants had fears related to those experienced by people with agoraphobia)

Generalisability gap: Available evidence for gameChangeVR was exclusively in a psychosis setting. The available study for XR Therapeutics considered exclusively people with autistic spectrum disorders. However, this specific focus may also have a benefit, as the intervention specifically considers the needs of people with autism which might address health inequalities.

8.4. Outcomes

None of the included studies reported all outcomes included in the NICE scope, but all reported some outcomes of interest. The gameChangeVR project^{17,18} and Gelabert et al for Amelia¹² focused on user experience and process outcomes. The gameChangeVR trial¹⁵ and Maskey et al⁵ studies presented clinical effectiveness

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

outcomes. Instruments used differed, but both studies presented measures of mood and social functioning or quality of life. The gameChangeVR trial reported measures of agoraphobia symptoms, while Maskey et al⁵ assessed symptom change using target behaviours. Safety data were presented for the gameChangeVR trial, in terms of serious adverse events, but not for the other interventions.

The EAG noted that change with respect to validated thresholds for minimally clinically important difference (MCID) were not always reported in the trial publications, and those reported were not clearly cited or justified. The EAG was unable to identify published MCID thresholds in an agoraphobia population for the key outcome measures, and therefore the interpretation of clinical effectiveness outcomes is uncertain. The EAG asked SCMs for input on this matter. While the SCMs reported that a variety of measures were used in their clinical practice, no specific MCIDs were identified. However, one suggested a potential rule of thumb based on 0.5 standard deviations.

Evidence gap: Evidence was not available for all scoped outcomes for all scoped interventions. Validated MCIDs may not be available for key outcome measures, and therefore change with respect to the MCID was not comprehensively assessed. No clinical outcomes were reported for Amelia.

Heterogeneity issue: There was variation in which outcomes were assessed between studies and what instruments were used.

8.5. Quality of included studies

A RCT was available for just one of the three technologies (gameChangeVR). UK-based evidence was available for two technologies (gameChangeVR and XR Therapeutics). Evidence for Amelia was available from Spain, which may be relatively generalisable to a UK context, but there will be some differences to consider in terms of culture and health system operation. The company advised that most of the environments used in Amelia are language-free, meaning they contain ambient audio but do not have a specific script or audio guide. This may aid transferability between countries, although it is noted that the environments have not specifically been validated in different languages. It is important to note that the Amelia environments are not intended as treatment protocols, but rather as tools for therapist use based on clinical judgement.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

In the principal evaluation of gameChangeVR,¹⁵ it should be noted that there was a substantial pause in recruitment from 16 March 2020 to 14 September 2020. It is uncertain if this would have impacted upon sampling consistency. Moreover, due to face-to-face restrictions caused by COVID, a protocol amendment was necessitated to replace the original primary outcome measure the Oxford Behavioural Avoidance Test (O-BAT) with a previous secondary outcome measure the Oxford Agoraphobic Avoidance Scale (O-AS). In total, 19 participants had VR therapy curtailed due to Covid-19-related policy decisions and 8 participants in the VR group could not receive any therapy. Treatment was unaffected for 84% of participants. These circumstances may result in somewhat reduced generalisability compared to trials conducted at other times.

The Maskey et al⁵ study for XR Therapeutics included only eight patients and was specifically conducted with a population of people with autism who had ‘fears and phobias’. The generalisability of this evidence to a broader agoraphobia population is therefore restricted. Additionally, gameChangeVR is positioned for agoraphobia specifically in a psychosis context, which is a subgroup of the NICE scope, but may not generalise to the broader scoped population.

Methodological gap: Studies for gameChange VR and XR Therapeutics assessed populations that were relevant to the NICE scope, but often did not match precisely to the full breath of the scoped population.

8.6. Results from the evidence base

The EAG summarises the results from the evidence base in this section, arranged by intervention as per the NICE scope. Detailed results for all eligible studies are presented in Appendix D.

8.6.1. Amelia Virtual Care

Evidence was available from one single-arm study of Amelia, conducted in a Spanish setting.

In the single-arm study by Gelabert and Giner (2018),¹² 98% of the 42 participants completed their course of psychotherapy within the previously established course of eight sessions. Two participants required two additional sessions of Amelia Virtual Care therapy beyond the protocol. The entire treatment protocol was completed by

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

82.4% of participants, with the main cause (55.6%) for non-completion being a lack of presence within the virtual environment and consequent perception of its usefulness. Presence relates to the extent participants can really feel like they are in the scenario being situated. On the Client Satisfaction Questionnaire-8,¹⁹ 57% of participants indicated a high or very high presence, while 12% indicated a null or low sense of presence. Across categories, there was an average satisfaction rating of 68%.

Overall, while there is evidence that Amelia has fairly good adherence and patient satisfaction, there is no evidence as to whether Amelia produces clinical benefit over and above existing care.

8.6.2. gameChangeVR

Evidence was available from one person-centred design study of how gameChangeVR was developed (the gameChangeVR project)^{17,18} and one RCT of gameChangeVR added on top of treatment as usual against treatment as usual alone¹⁴⁻¹⁶ (not specifically defined, the gameChangeVR trial), with embedded qualitative¹³ and economic¹¹ analyses – both conducted in a UK setting.

In the gameChangeVR project,^{17,18} a series of six key scenarios were developed based on participant input and feasibility of designing a suitable VR environment. These were: café (request or order), waiting room (give personal information), pub (unexpected event/erratic behaviour), bus (trapped/shut in), food shop (find an item), and street (safe place to unknown). In user testing, the success criterion was pre-determined as 90% of users rating gameChangeVR as immersive, easy to use and engaging. This was achieved, with all six participants rating gameChangeVR accordingly.

In the gameChangeVR trial, qualitative analysis¹³ showed that anxious avoidance was having a significant impact on participants' lives before the VR intervention, leaving some of them housebound and isolated. Overall, participants reported that using the intervention created an anxiety response that was useful for learning and practicing a different response while still within their safe environment. There was a need to balance the intensity of the anxiety response to a middle ground, so that the intervention was not boring (anxiety response too low) or that the intervention was not overly draining (anxiety response too high). The authors reported that the support

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

provided within the intervention meant that finding the right balance was “usually” possible (Bond et al, p.8). Those people who supplemented the intervention with activities to reinforce learning (e.g. writing notes, active reflection, discussions with care teams) generally had a better response to the intervention. Motivation to engage with the intervention, including undergoing the anxiety response within, was reported to be important. Those who were coping well with their condition had less motivation for this. Those who were struggling the most with agoraphobic avoidance expressed the greatest appreciation for, and gains from, gameChangeVR therapy.

Freeman et al¹⁵ found statistically significant benefits of VR exposure therapy on agoraphobic avoidance ($p=0.026$) and distress ($p=0.014$) measured by O-AS and the Questionnaire about the Process of Recovery.²⁰ However, the EAG considered that the magnitude of effect on reported outcome measures was not particularly large; the authors reported that a difference of 1 point on the O-AS avoidance measure was considered clinically meaningful, though the difference between arms was <1 . No minimally important difference (MID) for the O-AS distress was reported and the EAG was unable to identify this. However, the difference between arms (-4.33 , 95%CI $-7.78, -0.87$) appeared small in comparison to the scale (0-80). There was no statistically significant difference between arms in participants’ quality of life or other secondary outcome measures. The EAG also noted that numerical differences between treatment arms reduced between week 6 and week 26, suggesting that any benefit of gameChangeVR as compared to usual care was short-lived.

Consistent with the qualitative evidence suggesting that those with more severe symptoms may benefit more from gameChangeVR, Table 4 of the Freeman et al¹⁵ publication shows that those with severe symptoms showed larger improvements on agoraphobic avoidance and distress symptoms as measured by the O-AS (SCM input noted that while there is no specific criteria for severe symptoms, it would be fairly straightforward to identify people with severe agoraphobia on a case by case basis). In this paper, average avoidance was a score of 0 on the O-AS, moderate avoidance 1-2, high avoidance 3-5 and severe avoidance 6-8. In the case of the O-AS avoidance, the benefit was >1 indicating that the difference was clinically meaningful according to this threshold. The benefit for distress was $>10\%$ of the scale and was considered to be more likely to be clinically meaningful for patients, pending a validated MID threshold. These differences were also reasonably stable between 6 and 26 weeks. In comparison, benefits for participants with average, External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

moderate or high severity symptoms were absent or more uncertain. Only O-AS subscales were reported separately according to baseline severity, and therefore it was unclear if this pattern of effects was present across other scoped outcomes. Freeman et al¹⁶ found that participants with severe agoraphobia showed the greatest benefits from gameChangeVR therapy, exhibiting significant post-treatment improvements in agoraphobic avoidance, agoraphobic distress, ideas of reference, persecutory ideation, paranoia worries, recovery quality of life, and perceived recovery, but no significant improvements were found in depression, suicidal ideation, or health-related quality of life. Further data can be found in Table 3 of the publication.

Considering patient satisfaction, Freeman et al¹⁴ found that 65.8% of patients were very satisfied with VR therapy, 30.8% were mostly satisfied, 2.5% were indifferent or mildly dissatisfied, and 0.8% of patients were quite dissatisfied. Difficulties concentrating in VR (see adverse events Section 9) were associated with slightly lower satisfaction. Safety data are presented in Section 9. An economic analysis was also conducted¹¹ and this is discussed in Chapter 13.

While gameChangeVR received high levels of patient satisfaction from participants, there was evidence that it was only beneficial for reducing symptoms of agoraphobia in participants with severe symptoms at baseline (as opposed to those with average, moderate or high symptom severity). However, more evidence is needed to determine whether this effect was consistent across other outcomes, such as quality of life. Furthermore, base case economic results for gameChange were highly influenced by four inpatient people with psychosis who incurred particularly high costs. This was not explored in the gameChange trial publications.

8.6.3. Invirto

No eligible evidence was identified for this technology.

8.6.4. XR Therapeutics

Evidence was available from one single-arm study⁵ assessing eight adults with autism and 'fears and phobias' – conducted in a UK setting. This was the most relevant evidence available for this technology. There were no other adult studies for

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

this technology identified in this appraisal. There were two paediatric studies, which are discussed in Section 8.7

As this was a single-arm study, results can only show whether participants' scores improved over time and cannot show whether XR Therapeutics was more effective than standard care or whether any benefit of this technology remains following adjustment for confounding effects.

Retention and participation were achieved for all sessions. Members of an expert panel rated five out of eight participants as showing an improvement in their symptoms related to their phobia, however these participants did not include the two participants with phobias relevant to agoraphobia (open spaces and crowded buses), whose symptoms were reported to be "equivocally improved". This was a smaller potential improvement that allowed by the scale, which also included options for "normalised", "markedly improved", and "definitely improved". Two of the non-responders attributed their non-response to personal circumstances and routine changes respectively, while the third was making progress at the 6-month follow-up while not yet meeting response criteria. The sample did not show any benefit for general symptoms of anxiety (GAD-7 and BAI), depression (PHQ-9) or quality of life. Participants' self-reported confidence for managing their phobia appeared to improve between baseline and the end of session 4, though as with other outcomes, these data are challenging to interpret without a control arm. Across the sample, there was no difference in WHOQOL-BREF subscales following treatment, with the exception of a small improvement in the social subscale.

The company's reporting focused on statistical significance and the company attributed the absence of statistically significant benefit to the absence of specifically validated outcome measures for a population of people with autism. However, the EAG was not convinced by this explanation, given that the measures used (Table 2) have been widely used across populations, including people with autism.

8.7. Additional indirectly relevant evidence

In addition to the included studies, the EAG noted some additional indirectly relevant studies and studies that are outside of scope. These are listed with reasons for exclusion in Table 17 and Table 18, in Appendix C.²¹⁻⁵¹

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

The EAG considered that the most potentially relevant among these additional studies are Freeman et al²⁷ for gameChangeVR and Maskey et al⁵⁰ and Maskey et al⁵¹ for XR Therapeutics. Freeman et al²⁷ randomised 30 people with persecutory delusions to receive either virtual reality exposure or virtual reality cognitive therapy. Compared to exposure, cognitive therapy led to reductions in delusion conviction (reduction 22.0%, $P = 0.024$, Cohen's $d = 1.3$) and real-world distress (reduction 19.6%, $P = 0.020$, Cohen's $d = 0.8$). Additional Maskey et al studies were conducted in a paediatric population. However, given the presence of only one adult study for this intervention, it was considered that this evidence may be of interest. These were both conducted in a population of young people (up to age 14) with autistic spectrum conditions who have fears and phobias. Maskey et al⁵² reported a single blind RCT on a total of 35 young people from the North East of England compared to delayed treatment and found no statistically significant benefit on questionnaire outcomes, although a statistically significant improvement in Target Behaviours was found for the intervention group compared to the control group, from baseline to two months ($U=67.5$, $p=0.021$) and from baseline to six months post treatment ($U=53.0$, $p=0.007$), with six out of 16 participants counted as responders in the treatment group after six months compared to no participants in the control group. Maskey et al⁵¹ reported a single-arm study of eight young people from the North East of England and found that four participants were classed as responders at 12 months, three were classed as non-responders and one participant was lost to follow up. An earlier paper by Maskey et al,⁵⁰ also in a paediatric population with autism, but delivering the intervention differently than the later work, found that among nine young people, four overcame their phobias, eight out of nine were classed as treatment responders, this was maintained at 12-16 months follow-up and there was no loss to follow up in this analysis.

9. ADVERSE EVENTS

Information on adverse events were only available for one intervention (gameChangeVR). MAUDE (U.S. Food and Drug Administration) and MHRA (UK Government alerts) searches did not retrieve any results.

Freeman et al¹⁵ found that there was no statistically significant difference in the occurrence of serious adverse events between the gameChangeVR group (12 events in 8 patients) and the usual care alone group (eight events in seven patients, $p=0.37$). However, serious adverse events per participant were numerically higher in the gameChangeVR group and the EAG considers caution is required in interpreting the lack of statistical significance due to small sample size. A secondary analysis paper from this trial¹⁴ presents a broader adverse event profile. The most common adverse events were thinking about what might be happening in the room (14.2%), lasting headache (8.3%), and the headset causing feelings of panic (7.4%). However, there was no assessment of VR-specific adverse events using a standard questionnaire, such as the Simulator Sickness Questionnaire⁵³ to assess cybersickness, as used by Pot-Kolder et al⁵⁴ in a study of a non-scoped VR technology. This secondary analysis profiled adverse events only in the VR arm and not in the treatment as usual arm.

Evidence gap: Published safety information was only available for one intervention (gameChangeVR) investigated specifically in a psychosis context, and VR-specific adverse events were not assessed using a standard questionnaire. Moreover, safety data came only from an RCT, rather than real-world safety assessment.

10. ECONOMIC EVIDENCE

10.1. Published economic evidence

The search for economic evidence was conducted alongside the search for clinical evidence, detailed in Section 7.1. After screening, the only directly relevant study identified was Altunkaya et al (2022),¹¹ a within-trial cost-effectiveness analysis conducted alongside Freeman et al. 2022.¹⁵ The objective of the evaluation was to estimate the maximum cost-effectiveness price for gameChangeVR given conventional willingness-to-pay thresholds for QALYs routinely used in the NHS.

A full critique of the clinical trial can be found in Section 8, above. Briefly, 346 participants with psychosis and symptoms of agoraphobia were randomised to gameChangeVR+TAU or TAU (174 intervention, 172 comparator), with six-month follow-up. The frequency of participants' HRQoL and use of health and social care was recorded at baseline, six weeks and six months.

Participants completed two types of health-related questionnaires, EQ-5D-5L and Recovering Quality of Life (ReQoL). From this, two sets of utility scores were generated and consequently two sets of QALYs also produced. QALYs were calculated from utilities using the area under the curve approach.

GameChangeVR+TAU was associated with an incremental gain of +0.008 (-0.010 to 0.026) EQ-5D-based-QALYs (Table 3) and +0.003 (-0.011 to 0.017) ReQoL-based QALYs (Altunkaya et al 2022¹¹) compared with TAU alone.

Costs were calculated from an NHS and personal social services perspective (as per the NICE reference case⁵⁵), as well as a societal perspective. Health care services comprised GP contacts, contacts with psychiatrists, therapists and community mental health teams, hospitalisations, emergency department visits, outpatient appointments and paid help from NHS or social care services. Societal costs comprised criminal justice costs, private health care and carer time, but did not measure lost productivity from time-off-work. Criminal justice costs included police contacts, nights spent in a police cell or prison, psychiatric assessments whilst in custody and criminal or civil court appearances. Unit cost for paid care at home was based on the home care worker cost per hour reported on PSSRU 2019;⁵⁶ while the unit cost for unpaid care received from family and friends was based on minimum hourly wage in 2019.⁵⁷

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Overall, gameChangeVR was associated with a -£105 (-£1,135 to £924) difference in NHS & PSS costs (Table 4). These incremental costs excluded the cost of delivering gameChangeVR so were used to estimate the maximum cost that could be charged to yield an ICER below commonly accepted thresholds used by NICE and the NHS.

The EAG notes that the cost of VR headsets, training and intervention delivery was not included in the analysis, thus the maximum cost-effective prices represent the maximum per patient cost of delivering the entire intervention (software, headset, training, and intervention delivery), not just the licence cost for the software.

Taking an NHS and PSS perspective and using QALYs calculated using EQ-5D, the gameChangeVR intervention could cost up to £262 or £341 per person based on a £20,000 or £30,000 threshold for it to be cost-effective. When considering the intervention's impact on wider societal costs beyond the NHS and PSS perspective (which is beyond the scope of costs usually considered within NICE appraisals), the maximum cost-effective price was greater (Table 5).

Subgroup analyses were conducted stratifying participants by Oxford Agoraphobic Avoidance Scale, defined as (1) high or severe avoidance, (2) high or severe distress, (3) high or severe avoidance or distress, and (4) high or severe avoidance and distress. In each case the base case was the most pessimistic (i.e. the lowest maximum price). Table 5 reports the most optimistic subgroup by perspective based on a £20,000 per QALY threshold. From an NHS+PSS perspective in those with high or severe avoidance and distress, the maximum price yielding an ICER below £20,000 is £684 (£844 at £30,000/QALY). Full subgroup analyses are reported in Altunkaya et al. 2022.¹¹

The EAG noted the lack of statistically significant differences in incremental costs and QALYs and that the base case results were disproportionately driven by four psychiatric inpatient participants (three randomised to gameChangeVR+TAU and one to TAU). Specifically, the point estimate difference in NHS+PSS costs between arms increased from -£105 to +£233, and incremental QALYs fell from 0.008 to 0.006, resulting in there being no positive price at which gameChangeVR was cost-effective in the general population with psychosis and agoraphobia from an NHS+PSS perspective. However, there were subgroups for which gameChangeVR had the potential to be cost-effective: subgroups 1 (high or severe avoidance) and 4 (high or severe avoidance and distress), and 3 (high or severe avoidance or distress)

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

at a £30,000 threshold only. The most optimistic scenario yielded a max price of £125 at a £20,000 threshold (or £324 at £30,000) (Table 5).

Table 3 Utilities and QALYs accrued (EQ-5D-based)

	gC+TAU (n=174), mean (SE)	TAU (n=172), mean (SE)	Adjusted mean difference (95% CI)
Baseline	0.538 (0.021)	0.545 (0.020)	N/A
6 weeks	0.608 (0.021)	0.588 (0.022)	0.026 (-0.023-0.075)
6 months	0.570 (0.023)	0.568 (0.022)	0.007 (-0.043 to 0.057)
QALYs @ 6m	0.293 (0.010)	0.288 (0.009)	0.008 (-0.010 to 0.026)

gC: gameChangeVR; TAU: treatment as usual; N/A: not applicable. Source: Tables 2&4, Altunkaya et al (2022) ¹¹

Table 4 Summary costs, gameChangeVR+TAU vs TAU

Category	gC+TAU (n=174), mean (SE)	TAU (n=172), mean (SE)	Adjusted mean difference (95% CI)
NHS+PSS	£2695 (£619)	£2194 (£515)	-£105 (-£1135 to £924)
Criminal Justice	£42 (£20)	£2 (£2)	£38 (-0 to 77)
Caregiving	£2839 (£400)	£4403 (£860)	-£1,576 (-£3432 to £280)
Other private	£58 (£15)	£135 (£30)	-£88 (-£149 to -£26)
Societal	£5634 (£763)	£6733 (£993)	-£1,731 (-£3886 to £424)

Source: Table 4, Altunkaya et al (2022) ¹¹

Table 5 Maximum cost-effective price for different WTP thresholds, base case and selected sub-group analyses (EQ-5D-based QALYs)

	Incremental QALYs (95% CI)	Incremental cost (95% CI)*	Max price @ £20,000 per QALY	Max price @ £30,000 per QALY
NHS & PSS perspective				
Base Case (n=174, 172)	0.008 (-0.010 to 0.026)	-£105 (-£1135 to £924)	£262	£341
High or severe avoidance and distress (n=72, 90)	0.016 (-0.012 to 0.044)	-£365 (-£2399 to £1670)	£684	£844
Base Case excluding 4 inpatient participants (n=171, 171)	0.006 (-0.012, 0.025)	£233 (-£417 to £883)	N/A*	N/A*
High or severe avoidance excluding 4 inpatient participants (n=88, 98)	0.020 (-0.005 to 0.045)	£274 (-£699 to £1248)	£125	£324

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Societal perspective				
Base Case (n=174, 172)	0.008 (-0.010 to 0.026)	-£1731 (-£3886 to £424)	£1888	£1967
High or severe avoidance (n = 90, 99)	0.021 (-0.004 to 0.046)	-£2431 (-£6005 to £1142)	£2859	£3073
Base Case excluding 4 inpatient participants (n=171, 171)	0.006 (-0.012 to 0.025)	-£1315 (-£3314 to £683)	£1435	£1495
High or severe avoidance excluding 4 inpatient participants (n=88, 98)	0.020 (-0.005 to 0.045)	-£1911 (-£5195 to £1374)	£2310	£2509

**Societal perspective includes NHS+PSS plus criminal justice services, private health care and carer time. So: Tables 6 & S13, Altunkaya et al. 2022¹¹*

In summary, the evidence suggests that from an NHS+PSS perspective, the maximum acceptable per-patient price for delivering the gameChangeVR intervention ranges from below zero (in the general population with psychosis and agoraphobia, excluding inpatient participants) to £844 (subgroup with high or severe avoidance and distress, including psychiatric inpatient participants). However, this range underestimates uncertainty as it is based on point estimates from scenario analyses, ignoring confidence intervals around each.

10.1.1. Indirect evidence for economic outcomes

The EAG screened for indirectly relevant studies, as described in Section 8.7. We found no studies that provided information indirectly relevant to cost-effectiveness of the interventions. The only study which had discussed the benefit and cost was Segal (2011).²⁴ However, the paper only considered the therapist's perception of benefit and costs using virtual reality delivered treatments. Hence, it does not provide information on actual economic value. The EAG therefore concluded that there is no indirectly relevant study in this area.

10.1.2. Learnings relating to model structure and key issues impacting cost-effectiveness

The Altunkaya et al paper¹¹ was the only economic study considered of direct relevance to the decision questions. The EAG therefore used this as a starting point to develop a decision model to explore several uncertainties, namely duration of effect (i.e. time to relapse) and the effect of subsequent rounds of treatment.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

10.2. Conceptual modelling

The primary purpose of this analysis was to assess whether there is a plausible *prima facie* case for any of the VR interventions to be cost-effective, and if so, to identify evidence gaps to guide future data collection. Due to the high-level exploratory nature of the modelling, point estimate ICERs should be considered broadly indicative and not conclusive.

As no evidence was received from the manufacturer of Invirto (and no published evidence was identified by the EAG) and no clinical effectiveness evidence was available for Amelia, the EAG excluded these two technologies from further analysis. The two other interventions (gameChangeVR and XRT) were both trialled in different populations: gameChangeVR was trialled in people with psychosis and agoraphobia as an adjunct to TAU, and XRT is at an early stage of development but was explored in people with ASD and fears/phobias.

These therefore represent two different decision problems (DP):

1. What is the cost-effectiveness of gameChangeVR+TAU vs TAU in people with psychosis and agoraphobia? (DP1)
2. What is the cost-effectiveness of XRT vs TAU in people with ASD and agoraphobia? (DP2)

In DP2, in people with ASD, the EAG noted that the evidence base for XRT was very weak in the target population for this EVA (reflecting the early stage of development of its product). The EAG therefore excluded XRT from formal modelling and provided commentary on the available evidence and how it could be used in a future model to estimate the likely costs and outcomes associated with XR Therapeutics. Decision problem 1 is described below.

An early value assessment is reported using a decision analytic model for DP1, in people with psychosis and agoraphobia. The model drew from the one economic evaluation of relevance identified in the EAG's searches (gameChangeVR, Altunkaya et al. 2022¹¹), complemented by evidence sourced from the literature and supplemented with expert opinion. The analysis should be regarded as exploratory, focusing on the uncertainties in the data, rather than as a definitive estimate of the cost-effectiveness of the interventions against relevant comparator(s).

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Evidence Gap: Clarity on whether VR interventions are generalisable to more than the target patient groups observed in the included studies, and thus whether more than one of the interventions would be available to the same patient (although SCM input noted that such a situation is unlikely at the moment, given the limited availability of the interventions). If so, their relative cost-effectiveness must be compared to ensure the most efficient intervention is offered first. Direct head-to-head trials (rather than indirect statistical comparisons) are considered the least biased evidence source of comparative effectiveness.


10.2.1. Population

The analysis is structured as a decision problem with a distinct population: people with psychosis and agoraphobia. There are no limitations by age or other subgroupings in the analysis.

10.2.2. Model structure

The model is a two-state state-transition model (Markov model). The states are psychosis+agoraphobia and psychosis alone.

Figure 1: Structure of state-transition model



Health states labelled according to decision problem 1. State names for decision problem 1 are people with psychosis+agoraphobia and psychosis+no agoraphobia.

The transition period for the model is six months and it is run for a period of five years. A six-month transition period was chosen to reflect the follow-up period of the most relevant source study (Altunkaya 2022¹¹) as this enabled relatively easy translation of the data to the model. Five years were chosen as a reasonable time horizon over which to explore uncertainties in recurrence rates, and effectiveness of subsequent courses of therapy. (Note also Appendix F, supplied as a separate document.)

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

The choice of a state-transition model was driven purely by the need to explore uncertainties in longer term costs and effects of the different interventions. The key source study was a within-trial economic evaluation of gameChangeVR (Altunkaya 2022¹¹). This presented costs accrued and health state utilities at six months (*inter alia*) by treatment arm. A state-transition model requires discrete health states to be defined, such as 'responder' and 'non-responder'. Data are not reported in such a manner in the source study.¹¹ Furthermore, doing so requires dichotomising patients according to either their absolute score on an outcome measure, or a minimum change in score. The cut-offs for these are essentially arbitrary, usually based on subjective clinical opinion as to what constitutes a meaningful change or difference in scores, and any categorisation results in loss of information (e.g. all patients with an outcome score above the cut-off are classed as responders no matter what their actual score).

To avoid such a dichotomy in this analysis, we assume all patients receiving VR therapy 'respond' at six months such that the mean per person utility changes according to the (adjusted) mean difference observed in the clinical trial for gameChangeVR (+0.007 SE approx. 0.013, Table 2, Altunkaya et al 2022¹¹). This is the mean change in utility amongst all those who received the intervention. Therefore, 100% of patients transition from the 'agoraphobia' health state to the 'non-agoraphobia' health state in cycle 1. Subsequent cycles allowed the EAG to explore uncertainties around recurrence and effectiveness of second or more courses of VR therapy up to a time horizon of five years.

The model base case assumed those who relapse were offered additional rounds of therapy. Alternative assumptions were explored in scenario analysis.

Costs and outcomes accrued after the first year of analysis were discounted at 3.5% as recommended in the NICE manual.⁵⁵

10.2.3. Interventions and Comparators

Definitions of intervention and comparators were driven by availability of source data. Participants were allocated to gameChangeVR+TAU compared with TAU. As described in section 8.2 above, TAU was not specifically defined and may have varied from centre to centre.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Evidence gap: What is TAU and how best should it be defined in any future studies?

Evidence gap: What is the cost-effectiveness of gameChangeVR in place of TAU rather than as adjunct?

10.3. Model Inputs

10.3.1. Clinical Parameters

Clinical parameters were drawn from the gameChangeVR RCT¹⁵ and expert opinion.

10.3.1.1. Response

As described in section 10.2.2, the health state utility assigned to the response health state is derived from clinical trial results. This is the mean of all participants in the intervention arm of the study, and so includes all levels of response, from zero to 'maximum' or 'full'. For the purpose of the model, this utility difference is the driver of effect rather than a probability of response. We therefore assume a 100% probability of 'response' for all interventions.

Evidence gap: treatment effect estimates for XR and Amelia.

Evidence gap: (Optional) definitions of 'response', or 'partial' and 'complete' response according to an appropriate disease specific scale.

Evidence gap: Is there any difference in effect with who delivers the intervention(s)?

10.3.1.2. Relapse

No evidence was identified from the literature on relapse rates. The EAG consulted the manufacturers and clinical experts on plausible relapse rates.

Population with psychosis

The manufacturer of gameChangeVR stated that the mean outcome scores did not change from end of treatment to six months, and that a figure of 10% relapse at 12 months may be expected.

General population

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Amelia does not recommend a specific protocol for treatment and the manufacturer stated that the relapse rate is likely to vary depending on the protocol used. Using Craske and Barlow's protocol (pp 24-29),⁵⁸ it might be expected that:

...between 80% and 100% of patients undergoing these treatments will be panic free at the end of treatment and maintain these gains for up to 2 years. These results reflect substantially more durability than medication treatments. Furthermore, between 50 and 80% of these patients reach a point of "high end state", meaning within normative realms of symptoms and functioning, and many of the remainder have only residual symptomatology.

Population with ASD

XR therapeutics reported that in their initial studies on nine children with autism, eight improved and all improvements were maintained for 12 months. A study on eight adults with autism⁵² classified five as responders, all maintained at six months. Finally, a feasibility crossover RCT of VR+CBT for specific phobias in young people with ASD found improvement to be maintained at six months (Maskey 2019⁵²). In summary, according to the company, for those that showed improvements, they appeared to be maintained at six months, and there is some evidence of continued effect at 12 months. The company acknowledged that collected data to date cannot show that 100% of improvements would be maintained long term but stressed the desire to collect this evidence as part of any conditional approval by NICE.

Based on these subjective opinions, the EAG assumed a base case relapse rate of 25%, varied according to a uniform distribution between 0% and 50%.

Evidence gap: relapse rates over the short-, medium- and long-term following treatment (over six to 24 months or longer).

10.3.1.3. Response from second and subsequent courses of therapy

There are no data on whether a participant will respond as well to a second course of therapy post relapse. We include the relative risk of response for second and subsequent rounds of treatment purely to explore scenarios, with the relative risk set at one in our base case.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

10.3.2. Resource use and cost

Resource use and costs are calculated from an NHS and personal social services perspective (as per the NICE reference case). In addition, public sector (NHS & PSS plus criminal justice) and societal (defined as public sector plus time spent caring) costs are also calculated and reported for additional information as these were reported in Altunkaya et al. 2022.¹¹ All costs are adjusted to a 2021 price year.

10.3.2.1. Intervention costs

Intervention costs comprise licensing costs, therapist time (including delivery and training), and apportionment of capital cost of the VR headset. In the following the EAG describes the resource requirements for gameChange, before summarising quantities and costs included in the model.

gameChangeVR

gameChangeVR is recommended in those aged 16 and above and a course comprises six sessions over a period of six weeks, involving 30 mins of VR. In the RCT of gameChangeVR¹⁵, a mental health worker (peer support worker, assistant psychologist or clinical psychologist) was in the room with the participant, but they were not required to have previous experience of cognitive therapy. The mental health workers had a half day training session in delivery of the VR therapy, and weekly supervision was provided to them during the clinical trial. In its submission, the company stated that training of the mental health worker comprises three 90-minute sessions (4.5hrs), which is provided by the company as part of the licence fee.

In a session with a patient, the mental health worker first explained the therapy concepts, assisted with donning the headset, and started the programme. They then set homework tasks for the participants to apply the lessons learned in VR to real life situations. These tasks were reviewed by the worker. Sessions were held in an NHS clinic or in the participant's own home.

In the EAG's base case, the intervention was assumed delivered by a Band 4 mental health worker, under the supervision of a Band 8c clinical psychologist (Table 7).

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

gameChangeVR states a cost of [REDACTED] per patient per course of therapy

Specialist Committee Member Comments

Additional information from SCMs suggested that a full-time clinician is expected to have 14-16 clinical contacts per week. An assistant psychologist would expect to see three people per day if home based, and more if the patients were able to come to the clinic. The estimate of home visits is based on a 45-60 minute session, time to assemble and pack-up, travel time and note writing. SCMs also confirmed that staff with a wide range of skill levels and pay bands could deliver the intervention (most commonly people from bands 3-5, but sometimes higher).

Summary

Based on the descriptions above, the EAG estimated a course of treatment with gameChangeVR cost [REDACTED] (Table 6 and Table 7).

Table 6: Unit Costs

Item	Unit	Unit Cost	Source
Mental health worker	Hour	£33.00	AfC Band 4, equivalent to clinical psychology assistant practitioner (ch 17, and hourly cost from ch 10.1 PSSRU 2021)
Clinical psychologist	Hour	£105.00	consultant clinical psychologist (Band 8c, Ch 9, PSSRU 2021)
VR headset cost	Headset	£300	Notional cost.
gameChangeVR Licence	per patient per course	[REDACTED]	gameChangeVR company Rfl.

Table 7: Intervention Resource Use

Item	Quantity	Total Cost
gameChangeVR		
<i>Per session</i>		
Mental health worker intervention delivery	1 hour	£33.00
Mental health worker weekly supervision	1hr MH worker+clinical psychologist, assuming 15 sessions pw per MH worker	£9.20

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Training	4.5 hours with clinical psychologist and 6 MH workers in attendance, assuming training lasts 2 years before refresher required (eg due to staff turnover)	£0.17
VR Headset	One per MH worker conducting 15 sessions pw for 44 weeks/year, lasting 2 years	£0.64
<i>Total</i>		<i>£43.01</i>
<i>Per course</i>		
Per session costs above	Six sessions per course	£258.05
Licence cost		■
<i>Total per course, gameChangeVR</i>		■■■■

10.3.2.2. Other health service use costs

Altunkaya et al.¹¹ reported a point estimate difference in routine health service use of -£105 (95%CI -£1135 to £924) per patient, equivalent to -£112.15 in GBP2021, with an approximate standard error of £280.50 (Table 8).

Evidence gap: Are there any differences in other health service contacts between patients receiving VR-based therapy vs TAU?

10.3.2.3. Criminal justice costs

Altunkaya et al¹¹ reported a (borderline statistically significant) increase in criminal justice costs associated with gameChangeVR vs TAU in the population with psychosis. The EAG noted the magnitude was small (mean, 95%CI: +£38, £0 to £77 per patient). Adjusted to 2021 prices this equates to £40.59, with an approximate standard error of £10.49 per patient (Table 8).

Evidence gap: Is the finding of increased criminal justice costs of note and in need of further investigation or is it spurious?

10.3.2.4. Other costs

Whilst outside the NICE reference case, the EAG considered that other cost elements including informal caregiving and lost productivity from time off work may be of note.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Altunkaya et al¹¹ reported an informal care cost difference of -£1576 (-£3432 to £280) per participant in patients receiving gameChangeVR compared with TAU. Adjusted to GBP2021 this equates to -£1683 with an approximate SE of £506 per patient (Table 8).

The EAG noted that there was no evidence on changes in employment or hours worked associated with gameChangeVR. This was excluded from the EAG's analysis but the evidence gap noted.

Evidence gap: Is there a change in productivity / hours worked associated with successful treatment for agoraphobia in a population with psychosis?

Table 8: Incremental health service, criminal justice and carer costs associated with gameChangeVR vs TAU

Cost item	mean	SE	Distribution	Source/Notes
Incremental cost of NHS+PSS in pw psychosis vs pw psychosis+agoraphobia (NICE reference case cost perspective)	-£112.50	£280.50	Normal	Altunkaya et al. 2022 ¹¹ adjusted to 2021 prices using NHSCII ⁵⁹
Incremental cost of criminal justice services in pw psychosis vs pw psychosis+agoraphobia (non-reference case)	£40.59	£10.49	Normal	Altunkaya et al. 2022 ¹¹ adjusted to 2021 prices using NHSCII ⁵⁹
Incremental cost of time spent caring in pw psychosis vs pw psychosis+agoraphobia (non-reference case)	-£1683.28	£505.70	Normal	Altunkaya et al. 2022 ¹¹ adjusted to 2021 prices using NHSCII ⁵⁹

10.3.3. Health State Utilities

Baseline health state utility was set to the weighted mean of baseline utility across both arms of the gameChangeVR study (Table 2, Altunkaya et al. 2022¹¹). The increase in utility associated with treatment (representing the health state of people with psychosis without agoraphobia) was equal to the adjusted mean difference at six months (+0.007, 95%CI -0.043 to 0.057).

Table 9: Health State Utilities

DP	Health state	Utility, mean (SE)	Distribution	Source / Notes
1	People with psychosis and agoraphobia	0.541 (0.021)	B(319.08, 270.19)	Weighted mean of baseline utility, Altunkaya et al. 2022, ¹¹ Table 2

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

1	Treatment effect (gameChangeVR)	+0.007 (0.013)	N(0.007,0.013)	Altunkaya et al. 2022, ¹¹ Table 2. SE estimated from 95%CI
1	People with psychosis	0.548 (0.024)	-	Inferred from BL+treatment effect (previous two rows).

Evidence gap: Is EQ-5D sensitive to meaningful differences in HRQoL in people with agoraphobia (with and without psychosis)? Are disease specific scales (such as Beck Anxiety Inventory) with suitable mapping algorithms to health state utility suitable?

10.3.4. Model assumptions

A summary of the key assumptions in the decision modelling are in Table 10.

Table 10 Assumptions in exploratory modelling

1	gameChangeVR is delivered as per Freeman et al ¹⁵ as an adjunct to TAU rather than in place of TAU
3	The driver of effect in the model is change in utility following treatment rather than probability of response. Thus the 'probability of response' is 100% but the utility gain includes the values from all source study participants, as reflected in the standard deviation around mean utility difference.
4	The transition period of the model is 6 months, with a five year time horizon
6	Base case relapse rate is 25% per six months. Participants who relapse are offered repeat VR therapy
7	Repeat sessions are available when required and are as effective as the first session

10.4. Approach to Analysis

The EAG conducted a cost utility analysis estimating the incremental cost per incremental QALY gained from gameChangeVR compared with TAU in people with psychosis and agoraphobia. Analyses are conducted from the perspective of the NHS+PSS (NICE reference case). Only costs that differ between arms are measured and valued. Results for from a public sector and societal perspective are presented for information.

The EAG reports mean and SE costs and QALYs gained per patient in each arm, incremental cost-effectiveness ratios and probability of cost-effectiveness at £20,000 and £30,000 per QALY thresholds. Means and uncertainty distributions are generated from probabilistic analysis of 10,000 simulations, sampling from the distributions of input parameters specified in Table 8 and Table 9 above.

Additional sensitivity and scenarios were conducted as follows:

10.4.1. SA1: One-way sensitivity analysis, incremental utility gain from gameChangeVR

After excluding four psychiatric inpatient participants, Altunkaya et al.¹¹ reported no positive price at which gameChangeVR was cost-effective from an NHS+PSS perspective in the overall population. However, the analysis suggested there was the potential for the intervention to be cost-effective in those with 'high or severe avoidance'.¹¹ Whilst incremental QALYs in this subpopulation are reported (+0.020, 95%CI -0.005 to 0.0045 over six months, Table S13, Altunkaya et al. 2022¹¹), incremental utility is not. Therefore, as a proxy for this subgroup, the EAG conducted a one-way sensitivity analysis on incremental utility to determine the minimum required for gameChangeVR to be cost-effective from an NHS+PSS perspective.

10.4.2. SA2: Cost of VR headset

The cost of VR headsets varies substantially. In its base case the EAG assumed a cost of £300, which is a representative cost of a self-contained VR headset in 2023, although high specification headsets retail for close to £1,000 each. The EAG conducted a one-way sensitivity analysis, varying the cost of a headset from £0 to £1,000.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

10.4.3. SA3: Licence fees

License fees are a key cost input. The EAG therefore varied the fee for gameChangeVR to explore the impact on cost-effectiveness. Results are presented as point estimate ICERs as well as decision uncertainty (showing the probability of cost-effectiveness at £20,000 per QALY and £30,000 per QALY thresholds). The fees associated with a 50% probability of cost-effectiveness are identified.

10.4.4. SA4: Relapse rate

The EAG's base case assumed a 25% relapse rate per six months. This was varied between 0% and 100% in one-way sensitivity analysis.

10.4.5. SA5: Subsequent therapy availability

The EAG's base case assumed patients would get repeat therapy straightaway on relapse (i.e. within the same six month model cycle). The EAG explored a scenario where no repeat therapy was provided.

10.4.6. SA6: Subsequent therapy effectiveness

The EAG's base case assumed second and subsequent cycles of VR-based therapy were as effective as the first. The EAG conducted a one-way sensitivity analysis varying the effect of second and subsequent rounds of therapy from 0% to 100% of the effect of the first.

10.4.7. SA7: Two-way sensitivity analysis of licence fees versus relapse rate

This analysis explored the maximum cost-effective licence fees for the interventions as a function of the relapse rate.

10.4.8. SA8: Two-way sensitivity analysis of licence fees versus incremental utility from gameChangeVR

As a proxy for exploring the cost-effectiveness of gameChangeVR in severe subgroups, the EAG conducted a one-way analysis on utility gain. However, this analysis shows the cost-effectiveness of combinations of different licence fees against utility gain, so shows the maximum cost-effective licence fee as a function of the incremental gain in utility.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

All sensitivity analyses were conducted probabilistically, holding the target parameter at a given value whilst running the PSA on all other values as described in Wilson 2021.⁶⁰

10.5. Results from the economic modelling

10.5.1. Base Case Results

Base case results are reported in Table 11 below. VR therapy is an adjunct to standard care (TAU), costs reported are only those that differ between arms. Thus, the reported cost for TAU is zero.

Table 11: DP1 base case results

Perspective	Costs			QALYs				P(CE)	
	gC+TAU	TAU	Inc.	gC+TAU	TAU	Inc.	ICER	£20k	£30k
NHS+PSS	■	£0.00	■	■	■	■	■	26.3%	31.2%
Additional perspectives									
Public	■	£0.00	■	■	■	■	■	22.9%	27.5%
Societal	■	£0.00	■	■	■	■	■	99.0%	99.0%

Note QALYs vary by perspective due to Monte Carlo error

Whilst the point estimate ICER suggests that on average gameChangeVR is not cost-effective from an NHS+PSS perspective, the EAG base case suggests there is substantial decision uncertainty with around 25% to 30% probability of gameChangeVR being cost-effective from an NHS+PSS perspective at conventional thresholds of willingness to pay.

10.5.2. Scenario & Sensitivity Analyses

As previously stated, the point estimates from the decision modelling should be considered indicative only. The modelling does, however, provide a useful platform from which to explore a number of uncertainties. The results of the sensitivity analyses are presented below. Data from which figures are drawn are presented in Appendix E.

10.5.2.1. SA1: Incremental Utility gain

The gameChangeVR trial¹⁵ observed a higher treatment effect in the more severe subgroup. As a proxy for conducting an economic analysis in patients with more

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

severe agoraphobia, the EAG explored the ICER as a function of the utility difference from relief of agoraphobia in people with psychosis. Under the base case assumptions of the decision model, a mean utility gain from relief of agoraphobia of at least [REDACTED] is required to achieve an ICER below £30,000 per QALY gained, and at least [REDACTED] to achieve an ICER below £20,000 per QALY (Figure 2). (Note also Appendix F, supplied as a separate document)

Figure 2 SA1: OWSA on utility gain from relief of agoraphobia in people with psychosis

Figure redacted.

10.5.2.2. SA2: Cost of VR headset

Whilst an important cost element, the per session cost of the VR headset is very low, therefore the ICER is mostly insensitive to changes in the purchase price of the VR headset within the range considered (Figure 3).

Figure 3 SA2: OWSA on cost of VR headset

Figure redacted.

10.5.2.3. SA3: Licence fees

The point estimate results are highly sensitive to the licence fee charged, varying between dominant to £140,000 per QALY. The magnitude of the sensitivity is driven by the licensing model:

[REDACTED]
[REDACTED]
[REDACTED] (Figure 4).

Figure 4 OWSA licence fee per user per course, gameChangeVR

Figure redacted.

10.5.2.4. SA4: Relapse rate

Relapse rate is entered in the model as a six-month probability. One-way sensitivity analysis suggests the ICER of gameChangeVR deteriorates as the relapse rate increases (Figure 5).

Figure 5 OWSA on relapse rate

Figure redacted.

10.5.2.5. SA5: Subsequent therapy availability

Under the assumed base case relapse rate (25% per six-months), where repeat sessions are not available, the ICER is correspondingly less favourable. This is due to incurring the cost of the initial therapy, but the benefit only being sustained for a relatively short period (Table 12).

Table 12 Scenario analysis on availability of repeat therapy sessions

Available	Inc £	Inc QALY	ICER
No	████	████	████
Yes	████	████	████

10.5.2.6. SA6: Subsequent therapy effectiveness

As the relative effectiveness of subsequent therapy falls, the ICER deteriorates. This is because the same cost is incurred, but the relative benefit diminishes (Figure 6).

Figure 6 Relative Effectiveness of Subsequent Therapy

Figure redacted.

10.5.2.7. SA7: Two-way sensitivity analysis of licence fee vs relapse rate

As the six-month probability of relapse increases from zero, the maximum cost-effective licence fee declines accordingly. Figure 7 illustrates combinations of licence fee for gameChangeVR and probability of relapse associated with an ICER below £20,000 per QALY, between £20,000 and £30,000 and over £30,000.

Figure 7 Heatmap of licence fee vs 6m probability of relapse, gameChangeVR

Figure redacted.

10.5.2.8. SA8: Two-way sensitivity analysis of licence fee vs incremental utility

As the incremental utility associated with gameChangeVR increases, the licence fee associated with an ICER below £20,000 / QALY and below £30,000 increases (Figure 8).

Figure 8 Two-way sensitivity analysis, licence fee vs incremental utility, gameChangeVR

Figure redacted.

10.6. Commentary

10.6.1. gameChangeVR

gameChangeVR was the only intervention for which there was a published economic evaluation. Drawing directly on this (Section 10.1), the EAG noted that gameChangeVR was priced above that which would normally be considered cost-effective from an NHS+PSS perspective. Only by including broader societal costs (in particular, time spent caring) did the price point chosen by gameChangeVR become cost-effective.

Within-trial economic evaluations suffer from limitations including drawing only on one source of evidence of effect (the RCT on which they are based) and are limited in time horizon to the follow-up period of the study (six months in this case). The EAG therefore undertook some exploratory decision modelling to explore a number of scenarios and assumptions around relapse rate, effectiveness of repeat sessions, as well as assumed utility gains from relief of agoraphobia *inter alia* (described in section 10.4).

Under a base case assumption of 25% relapse every six months and assuming sustained effectiveness of repeat sessions, the ICER was

████████████████████, and thus is even less likely to represent an efficient use of NHS resources (Table 11). However, probabilistic analyses and one- and two-way sensitivity analyses suggested this finding was associated with substantial decision uncertainty and was highly sensitive to assumptions made in the model (section 10.5.2). In particular, two-way sensitivity analyses suggested there were scenarios under which combinations of licence fees and relapse rates and/or utility gains from relief of agoraphobia yield cost-

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

effectiveness estimates within that generally considered a good value for money investment by the NHS (that is, an ICER below £20,000 or £30,000). The EAG base case assumed gameChangeVR was delivered by a Band 4 mental health worker. If this is delivered by a higher band worker, then the ICER would be expected to increase further.

Bringing together the published economic evaluation of gameChangeVR conducted alongside the RCT and the decision modelling, the EAG notes that the within-trial evidence on cost-effectiveness (Altunkaya et al¹¹) was largely driven by four participants who were inpatients, and that excluding these, in the population with psychosis and agoraphobia, gameChangeVR was unlikely to be cost-effective. However, subgroup analysis amongst those with high or severe avoidance suggested a licence cost of up to £324 would yield an ICER at or below £30,000 per QALY. [REDACTED]

However, the EAG's analysis projecting the six-month findings to five years suggested that uncertainty in relapse rates / sustainability of treatment effect, the availability or otherwise of repeat courses of therapy, and their relative effectiveness, did have the capacity to alter the ICER substantially and there were scenarios where the ICER of gameChangeVR could be within NICE's willingness to pay threshold.

10.6.2. Amelia

There are no data on the clinical effectiveness of Amelia. The EAG notes that the company licences their technology at a cost of

[REDACTED]
[REDACTED]
[REDACTED]

However, in the absence of any evidence it is impossible to draw conclusions on the plausibility of this, and more research is needed to confirm the appropriateness of the company's price.

10.6.3. XR Therapeutics

There are no data on the relative effectiveness of XR Therapeutics. The EAG notes that the company charges a price of

[REDACTED]
[REDACTED]

[REDACTED]

[REDACTED]. In the absence of any evidence, it is impossible to draw conclusions on the plausibility of this, and more research is needed to confirm the appropriateness of the company's price.

11. INTERPRETATION OF THE EVIDENCE

11.1. Interpretation of the clinical and economic evidence

Following systematic searches, four unique clinical studies were identified, of which two were on gameChangeVR, one was on Amelia Virtual Care, and one on XR Therapeutics. RCT evidence was available for gameChangeVR. UK evidence was available for gameChangeVR and XR Therapeutics. All evidence on Amelia came from a Spanish setting; the EAG considered this to be fairly generalisable to a UK setting, with some limitations related to differences in culture and health system organisation. Evidence on gameChangeVR was specific to a population of schizophrenia spectrum disorder or psychosis, which was a relevant subgroup in the NICE scope.

In order to clarify how the scoped psychosis sub-group related to the studied population of psychosis or schizophrenia spectrum disorder in the gameChange trial, the EAG sought SCM input on how psychosis related to schizophrenia spectrum disorder. Advice was generally consistent but with different emphases. One SCM advised that *“psychosis is an umbrella term that encompasses conditions like schizophrenia”*. Another SCM advised that *“schizophrenia spectrum disorder refers to schizophrenia which is a psychotic disorder... Schizophrenia is a chronic mental health condition with distortion in thought and perception. Psychosis is when there is loss of touch with reality. Psychosis refers to symptoms that affect thought and perception. Psychosis is not a mental health diagnosis as such. Patients with schizophrenia experience psychosis.”* One SCM emphasised that psychosis *“is a symptom-based description, usually referring to positive symptoms such as hearing voices, seeing visions or having unusual beliefs (commonly paranoia). People can be described as have symptoms of psychosis even if they do not have a formal diagnosis”, while “schizophrenia spectrum disorder diagnosis refers to having a psychosis diagnosis, of which there are many subcategories.”*

Available evidence for the psychosis sub-population equates solely and in full to the available evidence for gameChangeVR. Evidence on XR Therapeutics was specific to people with autism with ‘fears and phobias’. This was considered relevant to the present decision problem, although not precisely aligned to the scope. Evidence was

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

not available for all scoped outcomes and safety data were only reported for gameChangeVR.

The EAG considered evidence for XR Therapeutics in the target population to be very limited. There was only one single-arm study of eight people with autism who had specific fears and phobias.⁵² Evidence of a benefit was only found in just over half of participants (five out of eight), and neither participant with fears thematically related to agoraphobia (fear of open spaces and fear of crowded buses) showed any benefit. Similarly, the only evidence for Amelia was a single-arm study, in which adherence and satisfaction was reported. While Amelia has fairly good adherence and patient satisfaction, no clinical outcomes were available in the scoped population, and hence there is no evidence with regards to its potential clinical benefit.

The evidence for gameChangeVR was suggestive of a small clinical benefit, however there was uncertainty as to its duration and how meaningful this benefit would be to participants. A subgroup analysis suggested that any clinical benefit was limited to those with very severe symptoms only, though limited data exploring outcomes by severity was reported. Moreover, there was no evidence of a benefit on a wider range of outcome measures. gameChangeVR was studied solely in a psychosis population, which is a subgroup of the scope. Effectiveness evidence was only available from one RCT and no further studies of any design.

The economic evidence for the three VR-based interventions is extremely limited. The EAG notes that gameChangeVR is priced at a point above that which would be considered cost-effective according to the NICE reference case (NHS+PSS perspective), but within that which would be considered value for money if a wider societal perspective is adopted (defined in Altunkaya 2022 as NHS+PSS, criminal justice and carer time costs). The EAG's exploratory modelling suggests the uncertainty is such that there are scenarios under which gameChangeVR may be cost-effective, for example in certain populations or with more optimistic interpretations of clinical effect estimates, but more evidence is required to confirm or refute this. Amelia is [REDACTED] while XR Therapeutics

[REDACTED]
[REDACTED]
[REDACTED]

██████████ Again, in the complete absence of randomised controlled evidence in the target population conclusions cannot be drawn as to either Ameila's or XR Therapeutic's cost-effectiveness.

11.2. Integration into the NHS

Information available to the EAG suggests that there is already some use of the scoped VR technologies within certain NHS trusts. Wider use would involve upscaling across more trusts. Potential challenges include ensuring sufficient appropriate staff resource and training to deliver such interventions. Furthermore, there are some challenges relating to access to digital technologies and service providers would have to consider supply of relevant equipment or signposting to relevant community resources, such as libraries, where equipment can be accessed, noting potential concerns regarding opening hours, access for patients who work during the day and are unable to leave the house, and confidentiality. When considering VR as a treatment option, it needs to be taken into account that there should be an alternative treatment modality for people for whom VR is unsuitable.

11.3. Evidence gap analysis

A summary of evidence gaps, pertaining to the intermediate and final outcomes from the scope, and those pertaining to decision modelling are summarised in Table 13. Evidence was focused around certain key outcomes and therefore there is limited information about certain additional scoped outcomes. A narrative assessment of evidence gaps in other methodological areas besides outcomes is presented within the clinical section of the report.

Table 13: Evidence Gap Analysis

Outcomes	Amelia Virtual Care	gameChangeVR	XR Therapeutics
Intermediate outcome: Patient choice and preferences	No studies RED	No studies RED	No studies RED
Intermediate outcome: Acceptability and satisfaction	One study AMBER	One study AMBER	One study AMBER
Intermediate outcome: Accessibility and digital access	No studies RED	No studies RED	No studies RED
Intermediate outcome: Intervention adherence and completion	One study AMBER	One study AMBER	One study AMBER
Intermediate outcome: Intervention-related adverse events	No studies RED	One study AMBER	No studies RED
Intermediate outcome: Device-related adverse events	No studies RED	One study AMBER	No studies RED
Clinical outcome: Change in agoraphobia symptoms	No studies RED	One study AMBER	One study, mixed results AMBER
Clinical outcome: Change in other psychological symptoms	No studies RED	One study, negative results RED	One study, negative results RED
Clinical outcome: Global functioning and work and social adjustment	No studies RED	No studies RED	No studies RED
Clinical outcome: Rates of recovery, time to recovery	No studies RED	No studies RED	No studies RED
Clinical outcome: Rates of relapse or deterioration, time to relapse or deterioration	No studies RED	No studies RED	No studies RED
Patient reported outcomes: Health-related quality of life	No studies RED	One study, negative results RED	One study, negative results RED
Patient reported outcomes: Recovering quality of life	No studies RED	One study, negative results RED	No studies RED
Patient reported outcomes: Patient experience	No studies RED	One study AMBER	No studies RED
Patient reported outcomes: Social contact	No studies RED	One study, negative result RED	No studies RED

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Modelling and economic outcomes

Effectiveness evidence: Populations & comparative data	Each intervention has been trialled in very different populations (eg gameChangeVR in people with psychosis and agoraphobia, Amelia in people with agoraphobia with or without panic disorder, and XR Therapeutics in people with autism and 'fears and phobias'). It is unknown whether any of the interventions are interchangeable between different populations and thus require head to head comparison RED
Effectiveness evidence: Comparative data	There is no randomised evidence on the effectiveness of Amelia or XR Therapeutics in the target population. RED
Effectiveness evidence: Comparative data	There is no evidence on durability of treatment effect and/or relapse rates after more than six months follow-up. RED
Effectiveness evidence: Comparative data	There is no evidence on effect of second or subsequent courses of therapy. RED
Effectiveness evidence: Comparative data	Is there an impact on other health service use from VR-based therapies? AMBER
Effectiveness evidence: Generalisability	Is there any difference in effect between who delivers the interventions? AMBER
Costs: Criminal justice costs	Is the impact of gameChangeVR on criminal justice costs in people with psychosis of meaningful? AMBER
Costs: Lost productivity	Is there a case for including time off work within economic evaluations of agoraphobia (outside NICE reference case)? The evidence base contains no data on lost productivity. RED
HRQoL: Health state utilities	Evidence on health state utilities is currently very weak RED

11.4. Ongoing studies

Ongoing studies, identified either through company lists or EAG searches, are listed below in Table 14.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Amelia Virtual Care	gameChangeVR	Invirto
[REDACTED]	No ongoing trials	DRKS00027001. ⁶¹ Evaluation of "Invirto aftercare" for anxiety disorders: a pilot study.
[REDACTED]		DRKS00027585. ⁶² Evaluation of "Invirto Therapy" for people with panic disorder: a randomized-controlled trial
[REDACTED]		
[REDACTED]		
[REDACTED]		

Table 14. Ongoing studies

Amelia Virtual Care	gameChangeVR	Invirto	XR Therapeutics
[REDACTED]	No ongoing trials	DRKS00027001. ⁶¹ Evaluation of "Invirto aftercare" for anxiety disorders: a pilot study.	No ongoing trials – monitoring of real-world outcomes is being conducted
[REDACTED]		DRKS00027585. ⁶² Evaluation of "Invirto Therapy" for people with panic disorder: a randomized-controlled trial	
[REDACTED]			
[REDACTED]			
[REDACTED]			

11.5. Summary and conclusions of evidence gap analysis

There are a number of evidence gaps in respect of the clinical evidence base as it pertains to the decision problem. These in part drive key uncertainties within the economic analysis. Key gaps included:

Population gaps

- The populations studied differed for each intervention, precluding direct comparison. While the population for Amelia corresponded best to the breadth of the NICE scope for this appraisal, the population studied for gameChangeVR was restricted to schizophrenia spectrum disorder or psychosis (psychosis being a subgroup in the NICE scope), and the population studied for XR Therapeutics was restricted to people with autism.
- No UK evidence was available for Amelia Virtual Care. Differences in health system organisation and treatment pathways may limit generalisability of international evidence.

Intervention gaps

- There is limited evidence for all interventions.
- There is no evidence on Invirto and no clinical effectiveness evidence for Ameila in adults with agoraphobia.
- Evidence for the different interventions was not balanced across populations and outcomes.

Comparator gaps

- There is uncertainty about how closely comparators matched routine practice, especially with regard to treatment as usual.
- There is no evidence on the durability of the effect of VR-based therapies / relapse rates after more than six months follow-up.

Outcome gaps

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

- Published evidence was not available for some outcomes. There was also heterogeneity in how clinical measures are reported. No clinical outcomes were reported for Amelia.
- Safety data were only available for gameChangeVR.

Other considerations

- Whilst outside the NICE reference case, employment status / lost time at work may be a key element of importance in any economic analysis of treatments for agoraphobia. Lost productivity was not measured in either of the RCTs identified by the EAG.

Included studies mostly suffered from methodological limitations, and bias in effect estimates could not be ruled out as a result.

11.6. Key areas for evidence generation

The EAG presents several specific evidence generation recommendations in Table 15.

Table 15: Evidence generation recommendations

Research question	Recommended study design	Outcomes
1. Would clinical effectiveness of scoped interventions be shown with longer follow-up?	Cohort study in RWE setting with standard care as comparator	Key scoped outcomes
2. Are statistically significant differences in outcomes clinically meaningful?	Validation study of minimally clinically significant differences	Agoraphobia symptoms, and potentially other key measures
3. What is the durability of effect / relapse rate associated with VR-based therapies?	Longer term follow-up of RCTs	Disease-specific and generic HRQoL tools, repeat presentation for treatment
4. Is Amelia effective (particularly in a UK setting)?	An RCT or comparative RWE study in a UK setting	Key scoped outcomes

12. CONCLUSIONS

12.1. Conclusions from the clinical evidence

A total of four eligible unique studies were available for consideration. The EAG was aware of a small number of ongoing studies, though it is unknown whether any will enhance the evidence base within the present scope. RCT evidence was available only for gameChangeVR. UK evidence was available for gameChangeVR and XR Therapeutics.

There was only one key effectiveness study for XR Therapeutics and gameChangeVR, and none for Amelia. The EAG considered evidence for XR Therapeutics in the target population to be very limited – there was only one single-arm study of eight people with autism with specific fears and phobias.⁵ Evidence of a benefit was only found in just over half of participants (five out of eight), and no benefit was apparent in those with fears thematically related to agoraphobia. The EAG noted some evidence of a benefit for gameChangeVR on symptoms of agoraphobia, however it was unclear whether this benefit would be clinically meaningful or would be durable. There was evidence that the intervention was only beneficial for those with very severe symptoms at baseline, but as few outcomes explored the effect of baseline severity, further evidence is needed to establish this. Furthermore, gameChangeVR was studied solely in a psychosis population, which is a subgroup of the scope only.

12.2. Conclusions from the economic evidence

At the price point chosen for gameChangeVR and drawing solely on the published within-trial analysis (Altunkaya et al¹¹), it is unlikely to be considered cost-effective under NICE's reference case analysis. However, the Altunkaya et al¹¹ analysis was limited in duration to six months. The EAG's exploratory analysis suggests that coupled with a price reduction, there are scenarios where the ICER could be within NICE's conventional threshold, but there is a great deal of uncertainty, so it is not possible to declare gameChangeVR cost-effective or not.

There is too little evidence on Amelia or XR Therapeutics to draw any conclusions as to cost-effectiveness.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

[REDACTED]

The key uncertainties are in 1) the effects of each of the interventions and 2) the duration of effect in those that do respond.

12.3. Summary of the combined clinical and economic sections

There was only one intervention for which RCT evidence is available. The EAG considered clinical effectiveness to be uncertain for all three technologies for which eligible evidence was available. The EAG did not consider there to be robust evidence for the clinical effectiveness of any scoped intervention. gameChangeVR, within the population studied, did show some evidence of potential benefit on agoraphobia symptoms, however there were considerable uncertainties about the interpretation and reliability of these findings.

The economic evidence suggests gameChangeVR is at best borderline cost-effective, but it is not strong enough to either rule it out or in as representing value for money. There is no evidence on which to draw any conclusions as to the cost-effectiveness of Amelia or XR Therapeutics,

[REDACTED]

[REDACTED].

Key evidence requirements are: 1) in populations where more than one VR-based intervention may be indicated, 2) to determine the clinical effectiveness of Amelia 3) to determine the relative effectiveness of all interventions compared with each other, and relevant control in relevant populations, 4) to assess clinical effectiveness for all technologies using a longer follow-up period, and 5) to assess MCIDs for key outcome measures in an agoraphobia population.

13. REFERENCES

1. National Health Service. Overview - Agoraphobia. 2022. Available from: <https://www.nhs.uk/mental-health/conditions/agoraphobia/overview/> (accessed April 2023).
2. World Health Organisation (WHO). ICD-11 for Mortality and Morbidity Statistics. 6B02 Agoraphobia. 2023. Available from: <https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/530592394> (accessed April 2023).
3. NHS Digital. Adult Psychiatric Morbidity Survey (APMS) 2022 – Survey Consultation Findings. 2022. Available from: <https://digital.nhs.uk/data-and-information/areas-of-interest/public-health/national-study-of-health-and-wellbeing/adult-psychiatric-morbidity-survey-2022-survey-consultation-findings> (accessed April 2023).
4. National Institute for Health and Care Excellence. Virtual reality for treating agoraphobia and agoraphobic avoidance: early value assessment. In development [GID-HTE10016]. Final scope. March 2023. Available from: <https://www.nice.org.uk/guidance/gid-hte10016/documents/final-scope> (accessed April 2023).
5. Maskey M, Rodgers J, Ingham B, Freeston M, Evans G, Labus M, et al. Using Virtual Reality Environments to Augment Cognitive Behavioral Therapy for Fears and Phobias in Autistic Adults. *Autism Adulthood*. 2019;1(2):134-45.
6. National Institute for Health and Care Excellence. Generalised anxiety disorder and panic disorder in adults: management. Clinical guideline [CG113]. Published: 26 January 2011. Last updated: 15 June 2020. Available from: <https://www.nice.org.uk/guidance/cg113> (accessed March 2023).
7. National Institute for Health and Care Excellence. Psychosis and schizophrenia in adults: prevention and management, Clinical guideline [CG178]. Published: 12 February 2014; Last updated: 01 March 2014. Available from: <https://www.nice.org.uk/Guidance/CG178> (accessed April 2023).
8. Harris DJ, Buckingham G, Wilson MR, Vine SJ. Virtually the same? How impaired sensory information in virtual reality may disrupt vision for action. *Exp Brain Res*. 2019;237(11):2761-6.
9. Harris DJ, Bird JM, Smart PA, Wilson MR, Vine SJ. A Framework for the Testing and Validation of Simulated Environments in Experimentation and Training. *Front Psychol*. 2020;11:605.
10. Murad MH, Noor A, Mouaz A, Fares A. New evidence pyramid. *Evidence Based Medicine*. 2016;21(4):125.
11. Altunkaya J, Craven M, Lambe S, Beckley A, Rosebrock L, Dudley R, et al. Estimating the Economic Value of Automated Virtual Reality Cognitive Therapy for Treating Agoraphobic Avoidance in Patients With Psychosis: Findings From the gameChange Randomized Controlled Clinical Trial. *Journal of Medical Internet Research*. 2022;24(11):e39248.
12. Gelabert JMG, C. Intervenció psicològica del' agorafòbia mitjançant realitat virtual: avaluació de paràmetres d'eficiència. [Psychological intervention for agoraphobia using virtual reality: Evaluation of efficiency parameters.]. Poster session presented at Jornades R+D+I TIC Salut Social; Vic, Spain 2018.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

13. Bond J, Kenny A, Pinfold V, Couperthwaite L, Kabir T, Larkin M, et al. A Safe Place to Learn: Peer Research Qualitative Investigation of gameChange Virtual Reality Therapy. *JMIR Serious Games*. 2023;11:e38065.
14. Freeman D, Rosebrock L, Waite F, Loe BS, Kabir T, Petit A, et al. Virtual reality (VR) therapy for patients with psychosis: satisfaction and side effects. *Psychol Med*. 2022;1-12.
15. Freeman D, Lambe S, Kabir T, Petit A, Rosebrock L, Yu LM, et al. Automated virtual reality therapy to treat agoraphobic avoidance and distress in patients with psychosis (gameChange): a multicentre, parallel-group, single-blind, randomised, controlled trial in England with mediation and moderation analyses. *The Lancet Psychiatry*. 2022;9(5):375-88.
16. Freeman D, Lambe S, Galal U, Yu LM, Kabir T, Petit A, et al. Agoraphobic avoidance in patients with psychosis: Severity and response to automated VR therapy in a secondary analysis of a randomised controlled clinical trial. *Schizophrenia Research*. 2022;250:50-9.
17. Knight I, West J, Matthews E, Kabir T, Lambe S, Waite F, et al. Participatory design to create a VR therapy for psychosis. *Design for Health*. 2021;5(1):98-119.
18. Lambe S, Knight I, Kabir T, West J, Patel R, Lister R, et al. Developing an automated VR cognitive treatment for psychosis: gameChange VR therapy. *Journal of Behavioral and Cognitive Therapy*. 2020;30(1):33-40.
19. Attkisson CC, Zwick R. The client satisfaction questionnaire. Psychometric properties and correlations with service utilization and psychotherapy outcome. *Eval Program Plann*. 1982;5(3):233-7.
20. Neil ST, Kilbride M, Pitt L, Nothard S, Welford M, Sellwood W, et al. The questionnaire about the process of recovery (QPR): A measurement tool developed in collaboration with service users. *Psychosis*. 2009;1(2):145-55.
21. Modrego-Alarcón M, López-Del-Hoyo Y, García-Campayo J, Pérez-Aranda A, Navarro-Gil M, Beltrán-Ruiz M, et al. Efficacy of a mindfulness-based programme with and without virtual reality support to reduce stress in university students: A randomized controlled trial. *Behav Res Ther*. 2021;142:103866.
22. Garcia-Palacios A, Botella C, Hoffman H, Fabregat S. Comparing acceptance and refusal rates of virtual reality exposure vs. in vivo exposure by patients with specific phobias. *Cyberpsychol Behav*. 2007;10(5):722-4.
23. Kirkham R, Batten C. Public perceptions on using Virtual Reality and Mobile Apps in Anxiety treatment. *The Journal of Undergraduate Research*. 2020;13.
24. Segal R, Bhatia M, Drapeau M. Therapists' perception of benefits and costs of using virtual reality treatments. *Cyberpsychol Behav Soc Netw*. 2011;14(1-2):29-34.
25. Shah LB, Torres S, Kannusamy P, Chng CM, He HG, Klainin-Yobas P. Efficacy of the virtual reality-based stress management program on stress-related variables in people with mood disorders: the feasibility study. *Arch Psychiatr Nurs*. 2015;29(1):6-13.
26. Brown P, Waite F, Lambe S, Jones J, Jenner L, Diamond R, et al. Automated Virtual Reality Cognitive Therapy (gameChange) in Inpatient Psychiatric Wards: Qualitative Study of Staff and Patient Views Using an Implementation Framework. *JMIR Form Res*. 2022;6(4):e34225.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

27. Freeman D, Bradley J, Antley A, Bourke E, DeWeever N, Evans N, et al. Virtual reality in the treatment of persecutory delusions: randomised controlled experimental study testing how to reduce delusional conviction. *Br J Psychiatry*. 2016;209(1):62-7.
28. Alsem SC, van Dijk A, Verhulp EE, de Castro BO. Using virtual reality to treat aggressive behavior problems in children: A feasibility study. *Clinical Child Psychology and Psychiatry*. 2021;26:1062 - 75.
29. Servera M, Sáez B, Mir. JMG. Feasibility of a virtual reality program to treat children with fear of darkness with nonexpert therapists. *Revista de Psicología Clínica con Niños y Adolescentes*. 2020;7:16-21.
30. Dehghan F, Jalali R, Bashiri H. The effect of virtual reality technology on preoperative anxiety in children: a Solomon four-group randomized clinical trial. *Perioper Med (Lond)*. 2019;8:5.
31. Bioulac S, de Sevin E, Sagaspe P, Claret A, Philip P, Micoulaud-Franchi JA, et al. [What do virtual reality tools bring to child and adolescent psychiatry?]. *Encephale*. 2018;44(3):280-5.
32. Kelson JN, Ridout B, Steinbeck K, Campbell AJ. The Use of Virtual Reality for Managing Psychological Distress in Adolescents: Systematic Review. *Cyberpsychol Behav Soc Netw*. 2021;24(10):633-41.
33. Hua Y, Qiu R, Yao WY, Zhang Q, Chen XL. The Effect of Virtual Reality Distraction on Pain Relief During Dressing Changes in Children with Chronic Wounds on Lower Limbs. *Pain Manag Nurs*. 2015;16(5):685-91.
34. Sarver NW, Beidel DC, Spitalnick JS. The feasibility and acceptability of virtual environments in the treatment of childhood social anxiety disorder. *J Clin Child Adolesc Psychol*. 2014;43(1):63-73.
35. Tennant M, Youssef GJ, McGillivray J, Clark TJ, McMillan L, McCarthy MC. Exploring the use of Immersive Virtual Reality to enhance Psychological Well-Being in Pediatric Oncology: A pilot randomized controlled trial. *Eur J Oncol Nurs*. 2020;48:101804.
36. Adjorlu A, Høeg ER, Mangano L, Serafin S. Daily Living Skills Training in Virtual Reality to Help Children with Autism Spectrum Disorder in a Real Shopping Scenario. 2017 IEEE International Symposium on Mixed and Augmented Reality (ISMAR-Adjunct). 2017:294-302.
37. Snider L, Majnemer A, Darsaklis V. Virtual reality as a therapeutic modality for children with cerebral palsy. *Dev Neurorehabil*. 2010;13(2):120-8.
38. Wallach HS, Safir MP, Bar-Zvi M. Virtual reality cognitive behavior therapy for public speaking anxiety: a randomized clinical trial. *Behav Modif*. 2009;33(3):314-38.
39. Segawa T, Baudry T, Bourla A, Blanc JV, Peretti CS, Mouchabac S, et al. Virtual Reality (VR) in Assessment and Treatment of Addictive Disorders: A Systematic Review. *Front Neurosci*. 2019;13:1409.
40. Botella C, Serrano B, Baños RM, Garcia-Palacios A. Virtual reality exposure-based therapy for the treatment of post-traumatic stress disorder: A review of its efficacy, the adequacy of the treatment protocol, and its acceptability. *Neuropsychiatric Disease and Treatment*. 2015;11.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

41. Botella C, Fernández-Álvarez J, Guillén V, García-Palacios A, Baños R. Recent Progress in Virtual Reality Exposure Therapy for Phobias: A Systematic Review. *Curr Psychiatry Rep.* 2017;19(7):42.
42. Gerardi M, Cukor J, Difede J, Rizzo A, Rothbaum BO. Virtual reality exposure therapy for post-traumatic stress disorder and other anxiety disorders. *Curr Psychiatry Rep.* 2010;12(4):298-305.
43. Guillén V, Baños RM, Botella C. Users' Opinion About a Virtual Reality System as an Adjunct to Psychological Treatment for Stress-Related Disorders: A Quantitative and Qualitative Mixed-Methods Study. *Front Psychol.* 2018;9:1038.
44. Meyerbröker K, Emmelkamp PM. Virtual reality exposure therapy in anxiety disorders: a systematic review of process-and-outcome studies. *Depress Anxiety.* 2010;27(10):933-44.
45. McCann RA, Armstrong CM, Skopp NA, Edwards-Stewart A, Smolenski DJ, June JD, et al. Virtual reality exposure therapy for the treatment of anxiety disorders: An evaluation of research quality. *Journal of Anxiety Disorders.* 2014;28(6):625-31.
46. Morina N, Ijntema H, Meyerbröker K, Emmelkamp PMG. Can virtual reality exposure therapy gains be generalized to real-life? A meta-analysis of studies applying behavioral assessments. *Behaviour Research and Therapy.* 2015;74:18-24.
47. Opiş D, Pintea S, García-Palacios A, Botella C, Szamosközi Ş, David D. Virtual reality exposure therapy in anxiety disorders: a quantitative meta-analysis. *Depress Anxiety.* 2012;29(2):85-93.
48. Powers MB, Emmelkamp PM. Virtual reality exposure therapy for anxiety disorders: A meta-analysis. *J Anxiety Disord.* 2008;22(3):561-9.
49. Turner WA, Casey LM. Outcomes associated with virtual reality in psychological interventions: where are we now? *Clin Psychol Rev.* 2014;34(8):634-44.
50. Maskey M, Lowry J, Rodgers J, McConachie H, Parr JR. Reducing specific phobia/fear in young people with autism spectrum disorders (ASDs) through a virtual reality environment intervention. *PLoS One.* 2014;9(7):e100374.
51. Maskey M, McConachie H, Rodgers J, Grahame V, Maxwell J, Tavernor L, et al. An intervention for fears and phobias in young people with autism spectrum disorders using flat screen computer-delivered virtual reality and cognitive behaviour therapy. *Research in Autism Spectrum Disorders.* 2019;59:58-67.
52. Maskey M, Rodgers J, Grahame V, Glod M, Honey E, Kinnear J, et al. A Randomised Controlled Feasibility Trial of Immersive Virtual Reality Treatment with Cognitive Behaviour Therapy for Specific Phobias in Young People with Autism Spectrum Disorder. *J Autism Dev Disord.* 2019;49(5):1912-27.
53. Fornells-Ambrojo M, Barker C, Swapp D, Slater M, Antley A, Freeman D. Virtual reality and persecutory delusions: safety and feasibility. *Schizophr Res.* 2008;104(1-3):228-36.
54. Pot-Kolder R, Geraets CNW, Veling W, van Beilen M, Staring ABP, Gijssman HJ, et al. Virtual-reality-based cognitive behavioural therapy versus waiting list control for paranoid ideation and social avoidance in patients with psychotic disorders: a single-blind randomised controlled trial. *Lancet Psychiatry.* 2018;5(3):217-26.
55. National Institute for Health and Care Excellence. NICE health technology evaluations: the manual, Process and methods [PMG36]. 2022. Available from:

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

<https://www.nice.org.uk/process/pmg36/chapter/introduction-to-health-technology-evaluation> (accessed April 2023).

56. Personal Social Services Research Unit (PSSRU). Unit Costs of Health and Social Care 2019. 2019. Available from: <https://www.pssru.ac.uk/project-pages/unit-costs/unit-costs-2019/> (accessed May 2023).
57. UK Government. National Minimum Wage and National Living Wage rates. 2023. Available from: <https://www.gov.uk/national-minimum-wage-rates> (accessed May 2023).
58. Barlow DHe. Panic Disorder and Agoraphobia (p. 1-64) Clinical handbook of psychological disorders: A step-by-step treatment manual. 6th ed. New York: Guilford Press; 2021.
59. Personal Social Services Research Unit (PSSRU). Unit Costs of Health and Social Care 2021. 2021. Available from: <https://www.pssru.ac.uk/project-pages/unit-costs/unit-costs-of-health-and-social-care-2021/> (accessed May 2023).
60. Wilson ECF. Methodological Note: Reporting Deterministic versus Probabilistic Results of Markov, Partitioned Survival and Other Non-Linear Models. Appl Health Econ Health Policy. 2021;19(6):789-95.
61. DRKS00027001. Evaluation of "Invirtio aftercare" for anxiety disorders: a pilot study. 2021.
62. DRKS00027585. Evaluation of "Invirtio Therapy" for people with panic disorder: a randomized-controlled trial. 2022.
63. Brown P, Waite F, Lambe S, Rosebrock L, Freeman D. Virtual Reality Cognitive Therapy in Inpatient Psychiatric Wards: Protocol for a Qualitative Investigation of Staff and Patient Views Across Multiple National Health Service Sites. JMIR Res Protoc. 2020;9(8):e20300.
64. ACTRN12609000959279. Virtual Reality Exposure Therapy for Agoraphobia. 2009.
65. ACTRN12615000927527. A randomised controlled trial comparing internet-delivered Cognitive Behavioural Therapy (iCBT), internet-delivered Mindfulness-Enhanced Cognitive Behavioural Therapy, internet-delivered Mindfulness Skills Training versus usual care for adults with depression and/or anxiety. <https://trialssearchwho.int/Trial2.aspx?TrialID=ACTRN12615000927527>. 2015.
66. Andersen NJ, Schwartzman D, Martinez C, Cormier G, Drapeau M. Virtual reality interventions for the treatment of anxiety disorders: A scoping review. Journal of Behavior Therapy & Experimental Psychiatry. 2023;81:101851.
67. Bentz D, Wang N, Ibach MK, Schick Tanz NS, Zimmer A, Papassotiropoulos A, et al. Effectiveness of a stand-alone, smartphone-based virtual reality exposure app to reduce fear of heights in real-life: a randomized trial. NPJ digital medicine. 2021;4(1).
68. Botella C, Garcia-Palacios A, Villa H, Banos R, Quero S, Alcaniz M, et al. Virtual reality exposure in the treatment of panic disorder and agoraphobia: A controlled study. Clinical Psychology & Psychotherapy. 2007;14(3):164-75.
69. Botella C, Villa H, Garcia Palacios A, Quero S, Banos RM, Alcaniz M. The use of VR in the treatment of panic disorders and agoraphobia. Studies in Health Technology & Informatics. 2004;99:73-90.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

70. Canais F, MacOrano R, Ferreira HA, Charraz A, Martins Y, Matos Pires A, et al. Virtual/augmented reality and neurofeedback in the treatment of agoraphobia and social phobia - A comparative and randomized study. *European Psychiatry*. 2020;63(Supplement 1):S303-S4.
71. Carl E, Stein AT, Levihn-Coon A, Pogue JR, Rothbaum B, Emmelkamp P, et al. Virtual reality exposure therapy for anxiety and related disorders: A meta-analysis of randomized controlled trials. *Journal of Anxiety Disorders*. 2019;61:27-36.
72. Castro WP, Sanchez MJR, Gonzalez CTP, Bethencourt JM, de la Fuente Portero JA, Marco RG. Cognitive-behavioral treatment and antidepressants combined with virtual reality exposure for patients with chronic agoraphobia. *International Journal of Clinical and Health Psychology*. 2014;14(1):9-17.
73. Chen J, Xie Z, Or C. Effectiveness of immersive virtual reality-supported interventions for patients with disorders or impairments: a systematic review and meta-analysis. *Health and Technology*. 2021;11(4):811-33.
74. Cottraux J. A comparative controlled study of virtual reality therapy and cognitive behavior therapy in panic disorder with agoraphobia. *Clinicaltrials.gov* [www.clinicaltrials.gov]. 2006.
75. Emmelkamp PMG, Meyerbroeker K. Virtual Reality Therapy in Mental Health. *Annual Review of Clinical Psychology*. 2021;17:495-519.
76. Freeman D, Lister R, Waite F, Yu LM, Slater M, Dunn G, et al. Automated psychological therapy using virtual reality (VR) for patients with persecutory delusions: study protocol for a single-blind parallel-group randomised controlled trial (THRIVE). *Trials*. 2019;20(1):87.
77. Freitas JRS, Velosa VHS, Abreu LTN, Jardim RL, Santos JAV, Peres B, et al. Virtual Reality Exposure Treatment in Phobias: a Systematic Review. *Psychiatric Quarterly*. 2021;92(4):1685-710.
78. Gomez-Busto FJ, Ortiz MI. Virtual reality and psychedelics for the treatment of psychiatric disease: A systematic literature review. *Clinical Neuropsychiatry*. 2020;17(6):365-80.
79. ISRCTN10661970. Investigating the efficacy of self-guided digital treatment with virtual reality for panic disorder and agoraphobia. 2022.
80. ISRCTN12497310. Virtual reality treatment for building confidence around people. 2018.
81. ISRCTN12882676. Using virtual reality (VR) to reduce social avoidance. <https://trialssearchwho.int/Trial2.aspx?TrialID=ISRCTN12882676>. 2019.
82. ISRCTN17308399. The gameChange trial: automated virtual reality therapy to help patients with psychosis feel more confident in everyday situations. 2019.
83. Jang DP, Ku JH, Shin MB, Choi YH, Kim SI. Objective validation of the effectiveness of virtual reality psychotherapy. *CyberPsychology & Behavior*. 2000;3(3):369-74.
84. KCT0007996. The Effect of Virtual-Reality-Based Exposure Training for Panic Disorder and Agoraphobia: A Pilot Study. 2022.
85. Krzystanek M, Surma S, Stokrocka M, Romanczyk M, Przybylo J, Krzystanek N, et al. Tips for Effective Implementation of Virtual Reality Exposure Therapy in

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Phobias-A Systematic Review. *Frontiers in psychiatry* Frontiers Research Foundation. 2021;12:737351.

86. Ling Y, Nefs HT, Morina N, Heynderickx I, Brinkman WP. A meta-analysis on the relationship between self-reported presence and anxiety in virtual reality exposure therapy for anxiety disorders. *PLoS ONE*. 2014;9(5) (no pagination).

87. Lorenzo Gonzalez M, Castro WP, Pitti Gonzalez CT, Bethencourt Perez JM, de la Fuente Portero JA, Marco RG. Efficacy of virtual reality exposure therapy combined with two pharmacotherapies in the treatment of agoraphobia. *International Journal of Clinical and Health Psychology*. 2011;11(2):189-203.

88. Lundin J, Lundstrom A, Gulliksen J, Blendulf J, Ejeby K, Nyman H, et al. Using 360-degree videos for virtual reality exposure in CBT for panic disorder with agoraphobia: a feasibility study. *Behavioural & Cognitive Psychotherapy*. 2022;50(2):158-70.

89. Malbos E, Rapee RM, Kavakli M. Creation of interactive virtual environments for exposure therapy through game-level editors: Comparison and tests on presence and anxiety. *International Journal of Human-Computer Interaction*. 2013;29(12):827-37.

90. Malbos E, Rapee RM, Kavakli M. Isolating the effect of Virtual Reality Based Exposure Therapy for agoraphobia: a comparative trial. *Studies in Health Technology & Informatics*. 2011;167:45-50.

91. Manyande A, Cyna AM, Yip P, Chooi C, Middleton P. Non-pharmacological interventions for assisting the induction of anaesthesia in children. *Cochrane Database of Systematic Reviews*. 2015(7).

92. Martin HV, Botella C, Garcia-Palacios A, Osma J. Virtual reality exposure in the treatment of panic disorder with agoraphobia: A case study. *Cognitive and Behavioral Practice*. 2007;14(1):58-69.

93. Meyerbroeker K, Morina N, Kerkhof GA, Emmelkamp PM. Virtual reality exposure therapy does not provide any additional value in agoraphobic patients: a randomized controlled trial. *Psychotherapy & Psychosomatics*. 2013;82(3):170-6.

94. Meyerbroker K, Morina N, Kerkhof G, Emmelkamp PM. Virtual reality exposure treatment of agoraphobia: a comparison of computer automatic virtual environment and head-mounted display. *Studies in Health Technology & Informatics*. 2011;167:51-6.

95. NCT00129610. Study of Virtual Reality Therapy and Cognitive Behavior Therapy in Panic Disorder With Agoraphobia. 2005.

96. NCT00734370. Virtual Reality Exposure Therapy in Agoraphobic Participants. 2008.

97. NCT03101332. Virtual Reality for Panic Disorder With Agoraphobia. 2017.

98. NCT03845101. CBT Versus CBT With Virtual Reality Exposure for Social Anxiety Disorder and Agoraphobia. 2019.

99. NCT03973541. CBT With VR Based Exposure for Social Anxiety Disorder. <https://clinicaltrials.gov/show/NCT03973541>. 2019.

100. NCT04695249. Using 360° Videos in the Treatment of Anxiety Disorders.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

101. NCT05319509. Pilot Study of Virtual Reality Therapy for Students With Anxiety. 2022.
102. NCT05510804. Efficacy of an App-based CBT Featuring Virtual Reality for Anxiety Disorders. 2021.
103. Pelissolo A, Zaoui M, Aguayo G, Yao SN, Roche S, Ecochard R, et al. Virtual reality exposure therapy versus cognitive behavior therapy for panic disorder with agoraphobia: A randomized comparison study. *Journal of Cybertherapy and Rehabilitation*. 2012;5(1):35-43.
104. Pitti CT, Penate W, de la Fuente J, Bethencourt JM, Roca-Sanchez MJ, Acosta L, et al. The combined use of virtual reality exposure in the treatment of agoraphobia. *Actas Espanolas de Psiquiatria*. 2015;43(4):133-41.
105. Pompoli A, Furukawa TA, Imai H, Tajika A, Efthimiou O, Salanti G. Psychological therapies for panic disorder with or without agoraphobia in adults: a network meta-analysis. *Cochrane Database of Systematic Reviews*. 2016(4).
106. Quero S, Perez-Ara M, Breton-Lopez J, Garcia-Palacios A, Banos RM, Botella C. Acceptability of virtual reality interoceptive exposure for the treatment of panic disorder with agoraphobia. *British Journal of Guidance & Counselling*. 2014;42(2):123-37.
107. Vincelli F, Molinari E, Riva G. CBT virtual reality assisted for the treatment of panic disorders with agoraphobia: a controlled study. 35th international meeting of the society for psychotherapy research; 2004 june 16 - 19; rome. 2004:35.
108. Vincelli F, Anolli L, Bouchard S, Wiederhold BK, Zurloni V, Riva G. Experiential cognitive therapy in the treatment of panic disorders with agoraphobia: a controlled study. *Cyberpsychology & Behavior*. 2003;6(3):321-8.
109. Vincelli F, Riva G. Virtual reality: a new tool for panic disorder therapy. *Expert Review of Neurotherapeutics*. 2002;2(3):377-83.
110. Vincelli F, Choi YH, Molinari E, Wiederhold BK, Riva G. A VR-based multicomponent treatment for panic disorders with agoraphobia. *Studies in Health Technology & Informatics*. 2001;81:544-50.
111. Wechsler TF, Kumpers F, Muhlberger A. Inferiority or Even Superiority of Virtual Reality Exposure Therapy in Phobias?-A Systematic Review and Quantitative Meta-Analysis on Randomized Controlled Trials Specifically Comparing the Efficacy of Virtual Reality Exposure to Gold Standard in vivo Exposure in Agoraphobia, Specific Phobia, and Social Phobia. *Frontiers in Psychology*. 2019;10:1758.
112. Wiebe A, Kannen K, Selaskowski B, Mehren A, Thone AK, Pramme L, et al. Virtual reality in the diagnostic and therapy for mental disorders: A systematic review. *Clinical Psychology Review*. 2022;98:102213.

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

14. APPENDICES

Appendix A: Searches for clinical and cost effectiveness evidence

Table 16: Resources searched for clinical and cost effectiveness studies

Database/Resource	Host	Date Searched	Results
MEDLINE and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily	Ovid	28.3.23	54
Embase	Ovid	28.3.23	117
APA PsycINFO	Ovid	28.3.23	92
CDSR / CENTRAL	Cochrane Library: Wiley	29.3.23	122
INAHTA HTA database	https://database.inahta.org/	30.3.23	7
Company websites		29.3.23	13
Guidelines	SIGN/NICE	29.3.23	2
ClinicalTrials.gov	http://www.clinicaltrials.gov/	29.3.23	24
WHO ICTRP	https://trialsearch.who.int/	29.3.23	31
MHRA	https://www.gov.uk/drug-device-alerts	29.3.23	0
MAUDE	https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfmaude/search.cfm	29,3,23	0
ScharrHUD	https://www.scharrhud.org/	29.3.23	0
CEA Registry	https://cear.tuftsmedicalcenter.org/	29.3.23	2
Total records retrieved			464
Total records after deduplication			318

Ovid MEDLINE(R) ALL <1946 to March 27, 2023>

- 1 Agoraphobia/ 2670
- 2 agoraphobi*.tw. 3609
- 3 ((phobi* or anxi* or fear*) adj3 (crowd* or 'open spac*' or 'go* out' or 'leav* home' or 'leav* house')).tw. 89

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

- 4 1 or 2 or 3 4469
- 5 exp virtual reality/ 5404
- 6 Virtual Reality Exposure Therapy/ 862
- 7 Augmented Reality/ 1085
- 8 (VR or 'virtual realit*').tw. 20841
- 9 (haptic adj2 technolog*).tw. 136
- 10 (VRCBT or VR-CBT).tw. 19
- 11 ("automated therap*" or "VR therap*" or "VR cognitive therap*" or "virtual reality therap*" or "virtual reality exposure" or VRET or "virtual reality based exposure" or VRBET).tw. 617
- 12 ("extended realit*" or "augmented realit*" or "mixed realit*").tw. 4379
- 13 ('game change' or gameChange or 'oxford VR' or BehaVR or 'HTC Vive' or 'Meta Quest' or 'Pico Neo').af. 134
- 14 ('amelia vr' or 'amelia virtual care').af. 0
- 15 invirto.af. 0
- 16 ('xr therapeutics' or XRT).af. 1203
- 17 or/5-1626907
- 18 4 and 17 54

Embase <1974 to 2023 March 27>

- 1 exp Agoraphobia/ 6838
- 2 agoraphobi*.tw. 4781
- 3 ((phobi* or anxi* or fear*) adj3 (crowd* or 'open spac*' or 'go* out' or 'leav* home' or 'leav* house')).tw. 127
- 4 1 or 2 or 3 7795
- 5 exp virtual reality/ 25414
- 6 Virtual Reality Exposure Therapy/ 910
- 7 Augmented Reality/ 2082
- 8 (VR or 'virtual realit*').tw. 28766
- 9 (haptic adj2 technolog*).tw. 162

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

- 10 (VRCBT or VR-CBT).tw. 26
- 11 ("automated therap*" or "VR therap*" or "VR cognitive therap*" or "virtual reality therap*" or "virtual reality exposure" or VRET or "virtual reality based exposure" or VRBET).tw. 817
- 12 ("extended realit*" or "augmented realit*" or "mixed realit*").tw. 5237
- 13 ('game change' or gameChange or 'oxford VR' or BehaVR or 'HTC Vive' or 'Meta Quest' or 'Pico Neo').af. 251
- 14 ('amelia vr' or 'amelia virtual care').af. 0
- 15 invirto.af. 1
- 16 ('xr therapeutics' or XRT).af. 2916
- 17 or/5-1647116
- 18 4 and 17 117

APA PsycInfo <1806 to March Week 3 2023>

- 1 exp Agoraphobia/ 2961
- 2 agoraphobi*.tw. 5508
- 3 ((phobi* or anxi* or fear*) adj3 (crowd* or 'open spac*' or 'go* out' or 'leav* home' or 'leav* house')).tw. 108
- 4 1 or 2 or 3 5617
- 5 exp virtual reality/ 11366
- 6 Virtual Reality Exposure Therapy/ 249
- 7 Augmented Reality/ 909
- 8 (VR or 'virtual realit*').tw. 10254
- 9 (haptic adj2 technolog*).tw. 48
- 10 (VRCBT or VR-CBT).tw. 15
- 11 ("automated therap*" or "VR therap*" or "VR cognitive therap*" or "virtual reality therap*" or "virtual reality exposure" or VRET or "virtual reality based exposure" or VRBET).tw. 655
- 12 ("extended realit*" or "augmented realit*" or "mixed realit*").tw. 1612
- 13 ('game change' or gameChange or 'oxford VR' or BehaVR or 'HTC Vive' or 'Meta Quest' or 'Pico Neo').af. 532
- 14 ('amelia vr' or 'amelia virtual care').af. 0

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

15 invirto.af. 0
 16 ('xr therapeutics' or XRT).af. 19
 17 or/5-16 16048
 18 4 and 17 92

Cochrane Library

#1 MeSH descriptor: [Agoraphobia] explode all trees 472
 #2 agoraphobi* 1262
 #3 ((phobi* or anxi* or fear*) NEAR/2 (crowd* or 'open spac*' or 'go* out' or 'leav* home' or 'leav* house')) 448
 #4 #1 or #2 or #3 1692
 #5 MeSH descriptor: [Virtual Reality] explode all trees 784
 #6 MeSH descriptor: [Virtual Reality Exposure Therapy] explode all trees 267
 #7 MeSH descriptor: [Augmented Reality] explode all trees 57
 #8 (VR or 'virtual realit*') 8193
 #9 (haptic adj2 technolog*) 7
 #10 (VRCBT or VR-CBT) 27
 #11 ('automated therap*' or 'VR therap*' or 'VR cognitive therap*' or 'virtual reality therap*' or 'virtual reality exposure' or VRET or 'virtual reality based exposure' or VRBET) 8383
 #12 ('extended realit*' or 'augmented realit*' or 'mixed realit*') 1225
 #13 'extended realit*' 224
 #14 ('game change' or gameChange or 'oxford VR' or BehaVR or 'HTC Vive' or 'Meta Quest' or 'Pico Neo') 1598
 #15 ('amelia vr' or 'amelia virtual care') 1
 #16 invirto 2
 #17 ('xr therapeutics' or XRT) 355
 #18 #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 14741
 #19 #4 and #18 142

[55 in CDSR and 67 in CENTRAL]

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

INAHTA

((((((((((((agoraphobia)[mh]) OR (agoraphobi*)) OR (virtual reality)[mh])) OR ("virtual reality")) OR ("extended reality")) OR ("augmented reality")) OR ("mixed reality")))) OR ((gameChange OR amelia OR invitro or XR)))

ClinicalTrials.gov

Search string	Results
Agoraphobia/Virtual Reality/all studies	7
Agoraphobia/VR/all studies	3
GameChange/all studies	1
Oxford vr/all studies	8
Amelia virtual care/all studies	0
Invitro/all studies	1
Xr therapeutics/phobia/all studies	4

ICTRP – using basic search

Search string	Results
Agoraphobia and 'virtual reality'	9
Agoraphobia and VR	3
GameChange	4
'Oxford vr'	4
'Amelia virtual care'	0
Invitro	3
'xr therapeutics'	8

CEA Registry and ScharrHUD


agoraphobia or virtual reality or VR

NICE and SIGN

Agoraphobia or virtual reality or VR

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Appendix B: PRISMA flow diagram



From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71. For more information, visit: <http://www.prisma-statement.org/>

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Date: May 2023

86 of 100

Appendix C: List of excluded studies and studies awaiting assessment

Table 17. List of excluded full-text English-language publications studies from company lists, with reasons

Excluded study	Reason for exclusion
<i>Amelia Virtual Care</i>	
Adjorlu et al ³⁶	Population (paediatric)
Alsem et al ²⁸	Population (paediatric)
Bioulac et al ³¹	Population (paediatric)
Botella et al ⁴¹	Population (phobias)
Botella et al ⁴⁰	Population (post-traumatic stress disorder)
Dehghan et al ³⁰	Population (paediatric)
Garcia-Palacios et al ²²	Population (specific phobias)
Gerardi et al ⁴²	Population (post-traumatic stress disorder and other anxiety disorders)
Guillen et al ⁴³	Population (stress-related disorders)
Hua et al ³³	Population (paediatric)
Kelson et al ³²	Population (paediatric)
Kirkham & Batten ²³	Population (anxiety)
McCann et al ⁴⁵	Population (anxiety disorders)
Meyerbroeker & Emmelkamp ⁴⁴	Population (anxiety disorders)
Modrego-Alarcon et al ²¹	Population (students with stress)
Morina et al ⁴⁶	Population (specific phobias)
Opris et al ⁴⁷	Population (anxiety disorders)

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Excluded study	Reason for exclusion
Powers & Emmelkamp ⁴⁸	Population (anxiety disorders)
Segal et al ²⁴	Population (broad psychological)
Segawa et al ³⁹	Population (addictive disorders)
Shah et al ²⁵	Population (mood disorders)
Servera et al ²⁹	Population (paediatric)
Snider et al ³⁷	Population (paediatric)
Tennant et al ³⁵	Population (paediatric)
Turner & Casey ⁴⁹	Population (broad psychological)
Wallach et al ³⁸	Population (public speaking anxiety)
Wong Sarver et al ³⁴	Population (paediatric)
<i>GameChangeVR</i>	
Brown et al ⁶³	Population (general psychiatric)
Freeman et al ²⁷	Population (persecutory delusions)
<i>XR Therapeutics</i>	
Maskey et al ⁵²	Population (paediatric)
Maskey et al ⁵¹	Population (paediatric)

Table 18: List of excluded full-text publications from EAG search, with reasons

Excluded study	Reason for exclusion
ACTRN12609000959279 ⁶⁴	Intervention
ACTRN12615000927527 ⁶⁵	Intervention

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Excluded study	Reason for exclusion
Andersen et al ⁶⁶	Intervention
Bentz et al ⁶⁷	Intervention
Botella et al ⁶⁸	Intervention
Botella et al ⁶⁹	Intervention
Canais et al ⁷⁰	Intervention
Carl et al ⁷¹	Intervention
Castro et al ⁷²	Intervention
Chen et al ⁷³	Intervention
CN-00595152 ⁷⁴	Intervention
DRKS00027001 ⁶¹	No results
DRKS00027585 ⁶²	No results
Emmelkamp et al ⁷⁵	Intervention
Freeman et al ⁷⁶	Article type
Freitas et al ⁷⁷	Intervention
Gomez-Busto & Ortiz ⁷⁸	Intervention
ISRCTN10661970 ⁷⁹	Intervention
ISRCTN12497310 ⁸⁰	No results
ISRCTN12882676 ⁸¹	Intervention
ISRCTN17308399 ⁸²	No results
Jang et al ⁸³	Intervention
KCT0007996 ⁸⁴	Intervention
Krzystanek et al ⁸⁵	Intervention

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Excluded study	Reason for exclusion
Ling et al ⁸⁶	Intervention
Lorenzo Gonzalez et al ⁸⁷	Intervention
Lundin et al ⁸⁸	Intervention
Malbos et al ⁸⁹	Intervention
Malbos et al ⁹⁰	Intervention
Manyande et al ⁹¹	Population (paediatric)
Martin et al ⁹²	Intervention
Meyerbroeker et al ⁹³	Intervention
Meyerbroker et al ⁹⁴	Intervention
NCT00129610 ⁹⁵	Intervention
NCT00734370 ⁹⁶	Intervention
NCT03101332 ⁹⁷	Intervention
NCT03845101 ⁹⁸	Intervention
NCT03973541 ⁹⁹	Intervention
NCT04695249 ¹⁰⁰	Intervention
NCT05319509 ¹⁰¹	Population (students with anxiety)
NCT05510804 ¹⁰²	Intervention
Pelissolo et al ¹⁰³	Intervention
Pitti et al ¹⁰⁴	Intervention
Pompoli et al ¹⁰⁵	Intervention
Pot-Kolder et al ⁵⁴	Intervention
Quero et al ¹⁰⁶	Intervention

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Excluded study	Reason for exclusion
Vincelli et al ¹⁰⁷	Intervention
Vincelli et al ¹⁰⁸	Intervention
Vincelli et al ¹⁰⁹	Intervention
Vincelli et al ¹¹⁰	Intervention
Wechsler et al ¹¹¹	Intervention
Wiebe et al ¹¹²	Intervention

Appendix D. Additional study results

This table presents results for clinical effectiveness outcomes. Further details compared to the results presented in the main clinical section are provided where relevant. However, there has been a focus on making the results understandable rather than presenting all minutiae. Safety and economic outcomes are discussed in relevant report sections and not included in this table.

Table 19: Study results for clinical effectiveness

Papers	Study name	Results
<i>Amelia Virtual Care</i>		
Gelabert et al ¹²	NR	98% of the 42 participants completed their course of psychotherapy within the previously established course of eight sessions. Two participants required 2 additional sessions of Amelia Virtual Care therapy beyond the protocol. The entire treatment protocol was completed by 82.4% of participants, with the main cause (55.6%) for non-completion being a lack of presence within the virtual environment and consequent perception of its usefulness. On the Client Satisfaction Questionnaire-8, ¹⁹ 57% of participants indicated a high or very high presence, while 12% indicated a null or low sense of presence. Across categories, there was an average satisfaction rating of 68%.
<i>gameChangeVR</i>		
Knight et al, ¹⁷ Lambe et al ¹⁸	gameChangeVR project	A series of six key scenarios were developed based on participant input and feasibility of designing a suitable VR environment. These were: café (request or order), waiting room (give personal information), pub (unexpected event/erratic behaviour), bus (trapped/shut in), food shop (find an item), and street (safe place to unknown). In user testing, the success criterion was pre-determined as 90% of users rating gameChangeVR as immersive, easy to use and engaging. This was achieved, with all six participants rating gameChangeVR accordingly.
Altunkaya et al 2022, ¹¹ Bond et al 2023, ¹³ Freeman et al 2022a, ¹⁴ Freeman et al 2022b, ¹⁵ Freeman et al 2022c ¹⁶	gameChangeVR trial	Qualitative analysis showed that anxious avoidance was having a significant impact on participants' lives before the VR intervention, leaving some of them housebound and isolated. Those who were struggling the most with agoraphobic avoidance expressed the greatest appreciation for, and gains from, gameChangeVR therapy. Five key superordinate themes were identified: i) experience and cost of anxious avoidance without treatment, ii) reasons to try: curiosity and motivation for trying VR treatment, iii) a place to practice different or new

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Papers	Study name	Results
		<p>responses to anxiety, iv) the security of knowing VR scenarios are not real, despite experiencing an anxiety response, and can therefore be a safe place to learn and build confidence and v) taking learning from the VR intervention into the real world. There are no subordinate themes within themes 1 and 2. Subordinate themes within theme 3 are: a) an immersive experience, b) a chance to observe anxiety, and c) new ways of responding. Subordinate themes within theme 4 are: a) the sweet spot of safety and anxiety and b) calibrating for a personalised approach. Subordinate themes within theme 5 are: a) from training wheels to real-world practice and b) one thing to hold onto. Overall, participants reported that using the intervention created an anxiety response that was useful for learning and practicing a different response while still within their safe environment. There was a need to balance the intensity of the anxiety response to a middle ground, so that the intervention was not boring (anxiety response too low) or that the intervention was not overly draining (anxiety response too high). The authors reported that the support provided within the intervention meant that finding the right balance was “usually” possible (Bond et al, p.8). Those people who supplemented the intervention with activities to reinforce learning (e.g. writing notes, active reflection, discussions with care teams) generally had a better response to the intervention. Motivation to engage with the intervention, including undergoing the anxiety response within, was reported to be important. Those who were coping well with their condition had less motivation for this.</p> <p>In the primary quantitative analysis, Freeman et al¹⁵ found that compared to the usual care alone group, participants in the VR therapy group had greater reductions in agoraphobic avoidance (p=0.026) and distress (p=0.014) at follow-up, measured by O-AS. Between-group differences were greater using the O-BAT, where data were available. No between-group differences were found for secondary outcome measures (excluding O-BAT, which was initially a primary outcome), except for recovery as assessed by the Questionnaire about the Process of Recovery.²⁰ The difference in O-AS was statistically significant, but was small. A difference of -0.47 (scale 0-8) and 4.3 on a scale of 0-100. A difference of 0.47 is not going to change the severity classification of avoidance as assessed by the scale (average 0, moderate 1, high 3, severe 6).</p> <p>Half (55%) of participants found a VR intervention appealing and described feeling intrigued by what it would be like.¹³ In the secondary quantitative analyses, Freeman et al¹⁴ found that 65.8% of patients were very satisfied with VR therapy, 30.8% were mostly satisfied, 2.5% were indifferent or mildly dissatisfied, and 0.8% of patients were quite dissatisfied. Difficulties concentrating in VR (see adverse events Section 9) were associated with slightly lower</p>

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

Papers	Study name	Results
		satisfaction. Meanwhile, Freeman et al ¹⁶ found that participants with severe agoraphobia showed the greatest benefits from gameChangeVR VR therapy, exhibiting significant post-treatment improvements in agoraphobic avoidance, agoraphobic distress, ideas of reference, persecutory ideation, paranoia worries, recovery quality of life, and perceived recovery, but no significant improvements were found in depression, suicidal ideation, or health-related quality of life. Further data can be found in Table 3 of the publication.
<i>XR Therapeutics</i>		
Maskey et al ⁵	NR	Retention and participation were achieved for all sessions. Five out of eight participants achieved improvement to symptoms related to target behaviours. Two of the non-responders attributed this to personal circumstances and routine changes respectively, while the third was making progress at the 6-month follow-up while not yet meeting response criteria. No consistent pattern of reliable or observable changes was found on any of the standardized questionnaire measures, relating to anxiety, depression and quality of life. There is selective presentation of numerical results in the company publication. Therefore, reporting only narrative results here prevents undue focus on highlighted values.

Appendix E: Data from Sensitivity Analyses

Data informing the figures reported in section 10.5.2 are reported in tables below.

Table:20 SA1: Incremental utility gain from gameChangeVR

Incremental utility	Inc £	Inc QALY	ICER
█	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████
███	██████	███	██████████
████	██████	███	██████████

Table:21 SA2: Cost of VR Headset

	DP1 (gameChangeVR)		
£ of headset	Inc £	Inc QALY	ICER
0	██████	███	██████████
50	██████	███	██████████
100	██████	███	██████████

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

150	████	██	████
200	████	██	████
250	████	██	████
300	████	██	████
350	████	██	████
400	████	██	████
450	████	██	████
500	████	██	████
550	████	██	████
600	████	██	████
650	████	██	████
700	████	██	████
750	████	██	████
800	████	██	████
850	████	██	████
900	████	██	████
950	████	██	████
1000	████	██	████

Table:22 SA3: Licence fees

DP1 (gameChangeVR)			
£ per person per course	Inc £	Inc QALY	ICER
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████
█	████	██	████

External assessment group report: Virtual reality for treating agoraphobia and agoraphobic avoidance [GID-HTE10016]

■	■	■	■
■	■	■	■
■	■	■	■
■	■	■	■
■	■	■	■
■	■	■	■
■	■	■	■
■	■	■	■

Table:23 SA4 Relapse Rate

P(relapse/6m)	DP1 (gameChangeVR)		
	Inc £	Inc QALY	ICER
0	■	■	■
0.05	■	■	■
0.1	■	■	■
0.15	■	■	■
0.2	■	■	■
0.25	■	■	■
0.3	■	■	■
0.35	■	■	■
0.4	■	■	■
0.45	■	■	■
0.5	■	■	■
0.55	■	■	■
0.6	■	■	■
0.65	■	■	■
0.7	■	■	■
0.75	■	■	■
0.8	■	■	■
0.85	■	■	■
0.9	■	■	■
0.95	■	■	■
1	■	■	■

Table:24 SA6 Relative Effectiveness of Repeat Therapy

relative effectiveness	DP1 (gameChangeVR)		
	Inc £	Inc QALY	ICER
0	██████	██	██████
0.05	██████	██	██████
0.1	██████	██	██████
0.15	██████	██	██████
0.2	██████	██	██████
0.25	██████	██	██████
0.3	██████	██	██████
0.35	██████	██	██████
0.4	██████	██	██████
0.45	██████	██	██████
0.5	██████	██	██████
0.55	██████	██	██████
0.6	██████	██	██████
0.65	██████	██	██████
0.7	██████	██	██████
0.75	██████	██	██████
0.8	██████	██	██████
0.85	██████	██	██████
0.9	██████	██	██████
0.95	██████	██	██████
1	██████	██	██████

Table: 25SA7: Heatmap of licence cost of gameChange vs probability of relapse

Table redacted.

Table: 26SA8: Heatmap of licence cost of gameChangeVR vs incremental utility

Table redacted.