

Division of ankyloglossia (tongue-tie) for breastfeeding

Interventional procedures guidance

Published: 14 December 2005

www.nice.org.uk/guidance/ipg149

Your responsibility

This guidance represents the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, healthcare professionals are expected to take this guidance fully into account, and specifically any special arrangements relating to the introduction of new interventional procedures. The guidance does not override the individual responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

All problems (adverse events) related to a medicine or medical device used for treatment or in a procedure should be reported to the Medicines and Healthcare products Regulatory Agency using the [Yellow Card Scheme](#).

Commissioners and/or providers have a responsibility to implement the guidance, in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity, and foster good relations. Nothing in this guidance should be interpreted in a way that would be inconsistent with compliance with

those duties. Providers should ensure that governance structures are in place to review, authorise and monitor the introduction of new devices and procedures.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should assess and reduce the environmental impact of implementing NICE recommendations wherever possible.

1 Recommendations

- 1.1 Current evidence suggests that there are no major safety concerns about division of ankyloglossia (tongue-tie) and limited evidence suggests that this procedure can improve breastfeeding. This evidence is adequate to support the use of the procedure provided that normal arrangements are in place for consent, audit and clinical governance.
- 1.2 Division of ankyloglossia (tongue-tie) for breastfeeding should only be performed by registered healthcare professionals who are properly trained.
- 1.3 Publication of further controlled trials on the effect of the procedure on successful long-term breastfeeding will be useful.

2 The procedure

2.1 Indications

- 2.1.1 Ankyloglossia, also known as tongue-tie, is a congenital anomaly characterised by an abnormally short lingual frenulum, which may restrict mobility of the tongue. It varies from a mild form in which the tongue is bound only by a thin mucous membrane, to a severe form in which the tongue is completely fused to the floor of the mouth. Breastfeeding difficulties may arise, such as problems with latching (getting the mother and baby appropriately positioned to breastfeed successfully), sore nipples and poor infant weight gain.
- 2.1.2 Many tongue-ties are asymptomatic and cause no problems. Some babies with

tongue-tie have breastfeeding difficulties. Conservative management includes breastfeeding advice, and careful assessment is important to determine whether the frenulum is interfering with feeding and whether its division is appropriate. Some practitioners believe that if division is required, this should be undertaken as early as possible. This may enable the mother to continue to breastfeed, rather than having to feed artificially.

2.2 Outline of the procedure

- 2.2.1 In early infancy, division of the tongue-tie is usually performed without anaesthesia, although local anaesthetic is sometimes used. The baby's head is stabilised, and sharp, blunt-ended scissors are used to divide the lingual frenulum. There should be little or no blood loss and feeding may be resumed immediately. After the early months of life, general anaesthesia is usually required.

2.3 Efficacy

- 2.3.1 One randomised controlled trial compared division of tongue-tie with 48 hours of intensive support from a lactation consultant. Mothers reported that 95% of babies (19 of 20) had improved breastfeeding 48 hours after tongue-tie division, compared with 5% of babies (1 of 20) in the control group ($p < 0.001$).
- 2.3.2 In 1 case series of 215 babies, 80% of mothers (173 of 215) reported improved breastfeeding 24 hours after the procedure. In another case series of 123 babies, 100% of mothers (70 of 70) reported improved latch after the procedure, and the 53 mothers with nipple pain noted significant improvement immediately after the procedure. In a third case series, 100% of babies (36 of 36) were reported to have normal tongue motion at 3 months. For more details, refer to the [sources of evidence](#).
- 2.3.3 There were conflicting opinions among the specialist advisors and some stated that it is difficult to be certain whether any perceived improvement in breastfeeding is due to division of the tongue-tie.

2.4 Safety

- 2.4.1 Few adverse effects were reported. One case series reported that, after the procedure, 2% of babies (4 of 215) had an ulcer under the tongue for more than 48 hours. Two studies, including a total of 159 babies, stated that there were no complications.
- 2.4.2 Two studies reported that 8% (3 of 36) and 18% (39 of 215) of babies slept through the procedure. For more details, refer to the sources of evidence.
- 2.4.3 The specialist advisors stated that adverse effects were likely to be rare. Potential adverse events include bleeding, infection, ulceration, pain, damage to the tongue and submandibular ducts, and recurrence of the tongue-tie.

2.5 Other comments

- 2.5.1 It was recognised that breastfeeding is a complex interaction between mother and child, and that many factors can affect the ability to feed. Skilled breastfeeding support is an integral part of the management of breastfeeding difficulties.
- 2.5.2 Public consultation highlighted that this procedure may also be relevant for bottle feeding, but it was noted that this was not included in the scope or in the literature search for this guidance.

3 Further information

- 3.1 NICE has developed a [guideline on postnatal care](#). The Health Development Agency has also published a systematic review on breastfeeding.

Sources of evidence

The evidence considered by the interventional procedures advisory committee is described in the [overview](#).

Information for patients

NICE has produced [information on this procedure for patients and carers](#). It explains the nature of the procedure and the guidance issued by NICE, and has been written with patient consent in mind.

4 Update information

Minor changes since publication

January 2012: Minor maintenance.

ISBN: 978-1-4731-6442-0

Endorsing organisation

This guidance has been endorsed by [Healthcare Improvement Scotland](#).