

# IP1096- Faecal microbiota transplant for recurrent Clostridium difficile infection

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

#### **Procedure:**

#### OPCS-4 code(s):

FMT using a colonoscope:

H21.8 Other specified other therapeutic endoscopic operations on colon

Y37.8 Other specified introduction of other substance into organ NOC

FMT using a nasoduodenal tube:

G57.8 Other specified other operations on duodenum

Y37.8 Other specified introduction of other substance into organ NOC

FMT using a nasogastric tube:

G48.8 Other specified other operations on stomach

Y37.8 Other specified introduction of other substance into organ NOC

FMT via rectal enema:

H62.8 Other specified other operations on bowel

Y37.8 Other specified introduction of other substance into organ NOC

### Diagnosis or health condition:

ICD-10 code(s):

A04.7 Eneterocolitis due to Clostridium difficil



## Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see here.