

IPG56 - Complete cytoreduction for pseudomyxoma peritonei (Sugarbaker technique)

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

This procedure cannot be expressed in the OPCS-4 classification by a single code. The current guidance is to code each organ removed as per normal coding rules, and to combine this with the ICD-10 diagnosis code.

For the intraperitoneal chemotherapy, the following OPCS-4 code is assigned:

T48.2 Introduction of cytotoxic substance into peritoneal cavity

Note: The collection of codes classifying procurement is not required, therefore codes from categories **X70** and **X71** should not be assigned unless there is a local requirement to do so.

Diagnosis or health condition:

ICD-10 code(s):

C78.6 Secondary malignant neoplasm of retroperitoneum and peritoneum

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).