

NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE

INTERVENTIONAL PROCEDURES PROGRAMME

Interventional procedure overview of epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

The tough covering of a spinal disc can sometimes tear, allowing the soft centre to bulge through. This is called herniation, also known as 'slipped disc'. It may cause pain in the back and leg (sciatica), and numbness and weakness in the leg. In this procedure the bulging part of the disc is removed using an endoscope (a thin, flexible tube with a camera on the end) and other instruments inserted through a small cut between the buttocks and up the spinal canal to the mid-back. The aim is to relieve pain by removing parts of the disc that press against the spinal nerve.

Introduction

The National Institute for Health and Care Excellence (NICE) has prepared this interventional procedure (IP) overview to help members of the interventional procedures advisory committee (IPAC) make recommendations about the safety and efficacy of an interventional procedure. It is based on a rapid review of the medical literature and specialist opinion. It should not be regarded as a definitive assessment of the procedure.

Date prepared

This IP overview was prepared in January 2016 and updated in October 2016.

Procedure name

- Epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

Specialist societies

- British Association of Spinal Surgeons

Description

Indications and current treatment

Lumbar disc herniation occurs when the nucleus pulposus of an intervertebral disc protrudes through a weakening or tear in the surrounding annulus fibrosus. Symptoms include pain in the back or leg, and numbness or weakness in the leg. Serious neurological sequelae including painful foot drop, bladder dysfunction, or cauda equina syndrome, may sometimes occur.

Conservative treatments include analgesics, non-steroidal anti-inflammatory medication and manual therapy. Epidural corticosteroid injections can also be used to reduce nerve pain in the short term. Lumbar discectomy is considered if there is severe nerve compression or persistent symptoms that are unresponsive to conservative treatment. Surgical techniques include open discectomy or minimally invasive alternatives using percutaneous endoscopic approaches. The choice of technique may be guided by several factors, including the presenting symptoms and signs and the location and size of the disc involved.

What the procedure involves

Epiduroscopic lumbar discectomy through the sacral hiatus for sciatica is usually done with the patient under sedation and local anaesthesia.

Under fluoroscopic guidance, a needle is inserted through the sacral hiatus. Over a guidewire a dilator is used to create a working channel through which a flexible endoscope can be steered into the anterior epidural space. The endoscope can reach nerve roots as high as the mid-lumbar lumbar spine bilaterally. When the appropriate disc level is reached, a laser optic fibre is introduced through the working channel of the endoscope to ablate disc tissue. The aim is to relieve pain by removing parts of the disc that press against the spinal nerve.

Literature review

Rapid review of literature

The medical literature was searched to identify studies and reviews relevant to epiduroscopic lumbar discectomy through the sacral hiatus for sciatica. The following databases were searched, covering the period from their start to 6 June 2016: MEDLINE, PREMEDLINE, EMBASE, Cochrane Library and other databases. Trial registries and the Internet were also searched. No language restriction was applied to the searches (see appendix C for details of search strategy). Relevant published studies identified during consultation or resolution that are published after this date may also be considered for inclusion.

The following selection criteria (table 1) were applied to the abstracts identified by the literature search. Where selection criteria could not be determined from the abstracts the full paper was retrieved.

Table 1 Inclusion criteria for identification of relevant studies

Characteristic	Criteria
Publication type	Clinical studies were included. Emphasis was placed on identifying good quality studies. Abstracts were excluded where no clinical outcomes were reported, or where the paper was a review, editorial, or a laboratory or animal study. Conference abstracts were also excluded because of the difficulty of appraising study methodology, unless they reported specific adverse events that were not available in the published literature.
Patient	Patients with sciatica.
Intervention/test	Epiduroscopic lumbar discectomy through the sacral hiatus
Outcome	Articles were retrieved if the abstract contained information relevant to the safety and/or efficacy.
Language	Non-English-language articles were excluded unless they were thought to add substantively to the English-language evidence base.

List of studies included in the IP overview

This IP overview is based on approximately 636 patients from 2 non-randomised comparative studies and 3 case series¹⁻⁵.

Other studies that were considered to be relevant to the procedure but were not included in the main extraction table (table 2) have been listed in appendix A.

Table 2 Summary of key efficacy and safety findings on epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

Study 1 Kim JD (2011)

Details

Study type	Non-randomised comparative study
Country	Korea
Recruitment period	2008–11
Study population and number	Patients with lumbar spinal stenosis n=98 patients (78 adhesiolysis, foraminoplasty + discectomy versus 20 adhesiolysis + foraminoplasty without discectomy)
Age and sex	Mean age: adhesiolysis, foraminoplasty + discectomy group, 58.5 years; adhesiolysis + foraminoplasty without discectomy group, 65.6 years Sex: adhesiolysis, foraminoplasty + discectomy group, 51% (40/78) male; adhesiolysis + foraminoplasty without discectomy, 55% (11/20) male
Patient selection criteria	Inclusion criteria: patients with chronic low back pain and radiculopathy which failed to respond to at least 3 months of conservative treatments (epidural steroid injections or oral analgesia) were included. Exclusion criteria: patients with renal failure, chronic liver disease, calcification and adhesion inside the spinal canal due to chronic spinal stenosis, osseous lesions that resulted in degenerative conditions or patients who were pregnant were excluded.
Technique	All procedures were performed using local anaesthesia. Adhesiolysis, foraminoplasty + discectomy group: A flexible endoscope was inserted into the anterior and posterior epidural space through an incision in the sacral hiatus region. The endoscope was advanced to the site of stenosis, and then adhesiolysis and foraminoplasty were performed. Subsequently, a discectomy was performed using a Ho:YAG laser to cauterise and decompress the intervertebral disc nucleus. Adhesiolysis + foraminoplasty without discectomy group: no discectomy was performed following adhesiolysis and foraminoplasty.
Follow-up	Adhesiolysis, foraminoplasty + discectomy group, mean of 20.7 months Adhesiolysis + foraminoplasty without discectomy group, mean of 23 months
Conflict of interest/source of funding	Not reported

Analysis

Follow-up issues: No patients were lost to follow-up.

Study design issues: Study assessed the added benefit of performing endoscopic laser lumbar discectomy after adhesiolysis and foraminoplasty. All procedures were performed by 1 surgeon. Visual analogue scores for radicular pain ranged from 0–10 with lower scores indicating less pain. Roland Morris disability questionnaire scores ranged from 0–24 with lower scores indicating less disability. The Macnab classification system rates the results of spine surgery using a 4-point scale: excellent: no pain or restriction of activity; good: occasional back or leg pain of sufficient severity to interfere with the patients' ability to do normal work or capacity to enjoy themselves in leisure hours; fair: improved functional capacity, but handicapped by intermittent pain of sufficient severity to curtail or modify work or leisure activities; poor: no improvement or insufficient improvement to enable an increase in activities; further operative intervention needed.

Study population issues: Diagnoses in the adhesiolysis, foraminoplasty + discectomy group included lumbar spinal stenosis, 53.8% (42/78); lumbar disc extrusion, 32.1% (25/78); postoperative pain after lumbar disc surgery, 10.3% (8/78); and chronic refractory low back pain

IP overview: Epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

caused by a lesion that was not identified before surgery, 3.8% (3/78). Diagnoses in the adhesiolysis + foraminoplasty without discectomy group included lumbar spinal stenosis, 90% (18/20); lumbar disc extrusion, 5% (1/20); and postoperative pain after lumbar disc surgery, 5% (1/20).

Other issues: there is an overlap in authorship between this study and Lee GW (2014) and there may be some overlap in patient populations.

Key efficacy and safety findings

Efficacy		Safety		
Number of patients analysed: 98 (78 adhesiolysis, foraminoplasty + discectomy versus 20 adhesiolysis + foraminoplasty without discectomy)				
Changes in outcome measures				
		Mean score		
Outcome measure	Group	Baseline	1 month	Final follow-up
VAS score for pain	Discectomy	7.6	4.9	3.6
	Without discectomy	8.5	4.6	6.1
RMDQ score	Discectomy	18.8	11.4	10.6
	Without discectomy	11.3	9.6	11.4
<ul style="list-style-type: none"> No p values were reported for analyses of changes within groups or comparisons between groups. 				
Postoperative outcomes at final follow-up according to Macnab criteria				
	Proportion of patients: % (n/N)			
Group	Excellent	Good	Fair	poor
Discectomy	32 (25/78)	24 (19/78)	26 (20/78)	18 (14/78)
Without discectomy	15 (3/20)	10 (2/20)	20 (4/20)	55 (11/20)
Endoscopy-related adverse events (in all patients) <ul style="list-style-type: none"> Transient headaches were reported in 8% (8/98) of all patients included in the study. Focal infection was reported in 2% (2/98) of all patients included in the study. Transient hyperaesthesia was reported in 1 patient. Pain over the endoscope insertion site was reported in 1 patient Meningitis was reported in 1 patient. Symptoms resolved after bed rest and symptomatic treatment (undefined). 				
Laser-related adverse events (in the discectomy group) <ul style="list-style-type: none"> Foot drop was reported in 3% (2/78) of patients in the END + laser lumbar discectomy group. Symptoms resolved within 6 months. Paraesthesia was reported in 19% (15/78) of patients in the END + laser lumbar discectomy group. Symptoms resolved with 6 months. 				
Abbreviations used: RMDQ, Roland Morris disability questionnaire; VAS, visual analogue scale				

Study 2 Lee GW (2014)

Details

Study type	Non-randomised comparative study
Country	Korea
Recruitment period	2008–12
Study population and number	Patients with lumbar spinal stenosis n=57 patients (32 adhesiolysis, foraminoplasty + discectomy versus 25 adhesiolysis + foraminoplasty without discectomy)
Age and sex	Mean age: adhesiolysis, foraminoplasty + discectomy group, 65.2 years; adhesiolysis + foraminoplasty without discectomy group, 67.1 years Sex: adhesiolysis, foraminoplasty + discectomy group, 47% male; adhesiolysis + foraminoplasty without discectomy group, 44% male
Patient selection criteria	Inclusion criteria: patients with central canal stenosis concurrently with or without foraminal stenosis because of hypertrophied ligamentum flavum, facet hypertrophy, and bony compression (such as bony spur), were included. All patients had low back pain, with or without sciatica, which failed to respond to 3 months of conservative treatment (including oral medication, physical therapy and unspecified injections). Exclusion criteria: patients with failed back surgery syndrome, spondylolysis or lumbar disc herniation without stenotic lesions were excluded.
Technique	All procedures were done using local anaesthesia. Adhesiolysis, foraminoplasty + discectomy group: A flexible endoscope was inserted into the anterior and posterior epidural space through an incision in the sacral hiatus region. The endoscope was advanced to the site of stenosis, and then adhesiolysis and foraminoplasty were performed. Subsequently, a discectomy was performed using a Ho:YAG laser to cauterise and decompress the intervertebral disc nucleus. Adhesiolysis + foraminoplasty without discectomy group: no discectomy was done after adhesiolysis and foraminoplasty.
Follow-up	Minimum of 2 years
Conflict of interest/source of funding	Not reported

Analysis

Follow-up issues: Authors state that 32 patients were enrolled in the adhesiolysis, foraminoplasty + discectomy group and 25 patients were enrolled in the adhesiolysis + foraminoplasty without discectomy group. They then report results for 27 patients in the adhesiolysis, foraminoplasty + discectomy group and 20 patients in the adhesiolysis + foraminoplasty without discectomy group but do not explicitly state that the missing patients were lost to follow-up.

Study design issues: Study assessed the added benefit of performing endoscopic laser lumbar discectomy after adhesiolysis and foraminoplasty. All procedures were performed by 1 surgeon. Visual analogue scale (VAS) scores for back pain and VAS scores for radicular pain ranged from 0–10 with lower scores indicating less pain. Roland Morris disability questionnaire scores ranged from 0–24 with lower scores indicating less disability.

Study population issues: There were various causes of lumbar spinal stenosis, including thickened ligament flavum, hypertrophic facet joints and herniated lumbar disc; however, authors did not state proportions in each group.

Other issues: there may be some patient population overlap with Kim JD (2011).

Key efficacy and safety findings

Efficacy				Safety
Number of patients analysed: 47 (27 adhesiolysis, foraminoplasty + discectomy versus 20 adhesiolysis + foraminoplasty without discectomy)				Epiduroscopy-related adverse events (in all patients, n=57) <ul style="list-style-type: none"> • Transient headaches were reported in 5% (3/57) of all patients included in the study. • Focal infection of the entry site was reported in 4% (2/57) of all patients included in the study. • Meningitis was reported in 1 patient. Symptoms resolved after bed rest and treatment with medication. Laser-related adverse events (in the discectomy group) <ul style="list-style-type: none"> • Transient mild motor paralysis was reported in 1 patient. Symptoms resolved within 1 month after the procedure.
Clinical outcomes (mean±standard deviation)				
Follow-up	Discectomy n=27	Without discectomy n=20	p value	
VAS for low back pain				
Baseline	8.1±0.7	8.5±1.3	0.41	
1 month	3.1±0.3	3.8±0.4	0.11	
6 months	3.0±0.7	4.1±0.5	0.03	
12 months	3.5±0.6	4.9±0.3	0.02	
24 months	4.4±0.5	6.7±0.9	<0.01	
	p=0.01*	p=0.12*		
VAS for leg pain				
Baseline	6.2±0.9	6.7±0.8	0.27	
1 month	2.8±0.4	3.1±0.7	0.06	
6 months	2.9±0.3	3.2±0.4	0.07	
12 months	3.9±0.4	4.5±0.8	0.03	
24 months	4.7±0.6	5.2±0.6	0.05	
	p=0.07*	p=0.15*		
RMDQ scores				
Baseline	13.2±0.9	12.6±1.2	0.32	
1 month	7.2±0.5	8.6±0.6	0.04	
6 months	6.8±0.5	8.5±0.9	0.01	
12 months	7.1±0.6	9.7±1.1	<0.01	
24 months	8.5±0.3	10.4±0.5	<0.01	
	p=0.03*	p=0.09*		
* baseline versus 24-month follow-up				
Note: Where there are discrepancies in the numbers reported, data have been extracted from the tables of the article rather than the text.				
Abbreviations used: RMDQ, Roland Morris disability questionnaire; VAS, visual analogue scale				

Study 3 Richter (2011)

Details

Study type	Case series
Country	Multiple: United states, South Korea, Italy, Spain, Belgium and Mexico
Recruitment period	2009–11
Study population and number	Patients with low back pain with or without sciatica caused by 'discogenic pathology' (assumed to be disc herniation) n=154 patients
Age and sex	Not reported
Patient selection criteria	Inclusion criteria: patients with severe back pain with or without leg pain with radiculopathic features were included. All patients had symptoms that failed to respond to conservative treatment options (not specified).
Technique	All procedures were performed using local anaesthesia and with the patient under sedation. A flexible endoscope was inserted into the anterior epidural space through an incision in the sacral hiatus region. Following a laminotomy, the endoscope was advanced to the site of stenosis and a laser discectomy was performed.
Follow-up	Not reported
Conflict of interest/source of funding	Not reported

Analysis

Follow-up issues: Authors did not report whether there were any losses to follow-up.

Study design issues: Patients were recruited from 8 participating centres. The number of clinicians undertaking the procedures was not stated. Visual analogue scale (VAS) scores for pain ranged from 0–10 with lower scores indicating less pain. Roland Morris disability questionnaire scores ranged from 0–24 with lower scores indicating less disability.

Study population issues: Demographic characteristics of the study population were not reported. Furthermore, authors did not report or stratify the types of discogenic pathology in the study population.

Key efficacy and safety findings

Efficacy	Safety
<p>Number of patients analysed: 154</p> <ul style="list-style-type: none"> • VAS scores for pain scores decreased from 7.5±1.0 at baseline to 3.4±1.3 at follow-up (p<0.005). • Roland Morris disability questionnaire scores decreased from 18.1±4.0 at baseline to 10.3±3.8 at final follow-up (p<0.005). <p>82% of patients had a successful outcome at follow-up (based on the modified MacNab scale; 'excellent', 'very good' and 'good' categories were noted as 'successful', and 'poor' and 'terrible' were noted as 'failure').</p>	<ul style="list-style-type: none"> • Study did not actively monitor the occurrence of adverse events.
Abbreviations used: VAS, visual analogue scale	

Study 4 Jo DH (2013)

Details

Study type	Case series
Country	Korea
Recruitment period	2011–12
Study population and number	Patients with refractory low back or lower extremity pain n=77 patients
Age and sex	Mean 58 years; 39% (30/77) male
Patient selection criteria	Inclusion criteria: patients with refractory low back or lower extremity pain, which did not improve with non-invasive conservative treatment including fluoroscopically-guided epidural steroid injection, or when 50% or more of the pain returned within 1 week.
Technique	Epiduroscopic laser neural decompression. Technique is not described in detail.
Follow-up	1 month
Conflict of interest/source of funding	Not reported

Analysis

Follow-up issues: 1 patient was lost to follow-up at 2 weeks and 3 patients were lost to follow-up at 1 month.

Study design issues: Retrospective review of medical records. The degree of pain relief was assessed on a 5-point scale: 1=very bad, 2=bad, 3=no change, 4=good, 5=very good. The baseline symptom scores were not reported.

Study population issues: 4 patients had failed back surgery syndrome, 44 patients had lumbar disc herniation, 11 had lumbar spinal stenosis, 16 had lumbar disc herniation and lumbar spinal stenosis, and 2 patients had lumbar facet joint syndrome. Symptom duration was 5 years or more for 24 patients and between 1 and 5 years for 23 patients.

Key efficacy and safety findings

Efficacy			Safety
Number of patients analysed: 77			The sutured area opened after removal of stitches from the operated area 1 week after the procedure in 1 patient.
Symptom relief scores at follow-up % (n)			
	2-week follow-up (n=76)	1-month follow-up (n=74)	
5 (very good)	42.1 (32)	10.8 (8)	
4 (good)	46.1 (35)	74.3 (55)	
3 (no change)	9.2 (7)	10.8 (8)	
2 (bad)	2.6 (2)	1.4 (1)	
1 (very bad)	0	2.7 (2)	
81.8% (63/77) of patients had improved 1 month after the procedure.			

Study 5 Lee SH (2016)**Details**

Study type	Case series
Country	Korea
Recruitment period	2011–13
Study population and number	Patients with low back pain and radiculopathy n=250
Age and sex	Mean 46.5 years; 60% (150/250) male
Patient selection criteria	Low back pain and radicular pain (247 patients had herniated nucleus pulposus and 3 had a discal cyst, proven by MRI) refractory to medical and physical therapy for more than 2 weeks. Exclusion criteria: pain originating from infection, patients with a bleeding tendency, history of previous spine surgery when the symptoms and neurological findings were thought to be solely from scar tissue formation after an operation.
Technique	Epiduroscopic laser decompression. The procedures were done with the patient under local anaesthesia. Under direct vision with epiduroscopy, adhesiolysis and the size of the herniated disc were reduced by a Ho:YAG laser. The ruptured discs were decompressed until the epiduroscopic images confirmed nerve root decompression.
Follow-up	3 months
Conflict of interest/source of funding	None

Analysis

Follow-up issues: There were no losses to follow-up. Postoperative outcome scores were obtained in the outpatient clinic and by telephone interview. Postoperative MRIs were not available for all patients.

Study design issues: Prospective case series. Efficacy was assessed by a visual analogue scale score for both low back pain and radiculopathy, and the Oswestry Disability Index (ODI) questionnaire.

Study population issues: Lesion levels: 20 at L3/4, 130 at L4/5 and 100 at L5/S1. The mean duration of symptoms before the procedure was 5 months.

Key efficacy and safety findings

Efficacy				Safety
Number of patients analysed: 250				Complications <ul style="list-style-type: none"> • Headache, n=3 • Epidural pneumocephalus, n=1
<i>Changes in the visual analogue scale (VAS) score (leg and back) and Oswestry Disability Index (ODI) at 2 weeks and 3 months postoperatively,</i>				
Follow-up period	VAS (leg)	VAS (back)	ODI	
Preoperative	7.12±0.8	5.93±0.8	50.34±0.8	
2 weeks	3.58±0.7	4.10±0.9	19.16±0.9	
3 months	2.60±1.3	2.69±0.1	11.65±0.6	
p value	<0.01	<0.01	<0.01	
Recurrence rate=5.8% (7/250) (5 patients were subsequently treated by open laminectomy and discectomy).				

Efficacy

Pain relief

A non-randomised comparative study of 98 patients compared treatment by endoscopic adhesiolysis, foraminoplasty and discectomy (n=78) with endoscopic adhesiolysis and foraminoplasty without discectomy (n=20). Visual analogue scale (VAS) scores for radicular pain (ranging from 0–10, with lower scores indicating less pain) improved from 7.6 to 3.6 with discectomy and from 8.5 to 6.1 without discectomy at final follow-up (p values not reported; mean follow-up periods were 21 and 23 months respectively)¹. A non-randomised comparative study of 57 patients compared treatment by endoscopic adhesiolysis, foraminoplasty and discectomy (n=32) with endoscopic adhesiolysis and foraminoplasty without discectomy (n=25). The improvement in VAS score for low back pain (scores range from 0–10, with lower scores indicating less pain) was statistically significant with discectomy (from 8.1 to 4.4; p=0.01) but not without discectomy (from 8.5 to 6.7; p=0.12) at 24-month follow-up. The difference between the groups was statistically significant (p<0.01)². In the same study, improvements in VAS scores for leg pain were not statistically significant (from 6.2 to 4.7; p=0.07 and from 6.7 to 5.2; p=0.15 respectively) at 24-month follow-up. The difference between the groups was statistically significant (p=0.05). In a case series of 154 patients, there was a statistically significant decrease in VAS scores for pain from 7.5 at baseline to 3.4 at follow-up (p<0.005)³. In a case series of 77 patients, 81.8% (63/77) of patients had improved symptoms at 1-month follow-up⁴. In a case series of 250 patients, the mean VAS score for leg pain decreased from 7.1 at baseline to 2.6 (p<0.01) and the mean VAS score for back pain decreased from 5.9 at baseline to 2.7 (p<0.01) at 3-month follow-up⁵.

Functional capacity

In the non-randomised comparative study of 98 patients, Roland Morris disability questionnaire scores (ranging from 0–24, with lower scores indicating less disability) changed from 18.8 to 10.6 with discectomy and from 11.3 to 11.4 without discectomy at final follow-up (p values not reported; mean follow-up periods were 21 and 23 months respectively)². In the non-randomised comparative study of 57 patients, the change in Roland Morris disability questionnaire scores was statistically significant with discectomy (from 13.2 to 8.5; p=0.03) but not without discectomy (from 12.6 to 10.4; p=0.09) at 24-month follow-up. The difference between the groups was statistically significant (p<0.01)². In the case series of 154 patients, the change in Roland Morris disability questionnaire scores was statistically significant, from 18.1 at baseline to 10.3 at follow-up (p<0.005)³. In the case series of 250 patients, the Oswestry Disability Index score (ranging from 0–100) improved from 50 at baseline to 12 at 3-month follow-up (p<0.01)⁵.

Safety

Transient paralysis

Transient mild motor paralysis was reported in 1 patient from the discectomy group (n=32) in a non-randomised comparative study of 57 patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy or endoscopic adhesiolysis and foraminoplasty without discectomy. Symptoms resolved 1 month after the procedure². Foot drop was reported in 3% (2/78) of patients in the discectomy group in the non-randomised comparative study of 98 patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy (n=78) or endoscopic adhesiolysis and foraminoplasty without discectomy (n=20)¹. Symptoms resolved within 6 months.

Dysaesthesia

Transient hyperaesthesia was reported in 1 patient in the non-randomised comparative study of 98 patients. The authors did not state which group this patient was in¹. Paraesthesia was reported in 19% (15/78) of patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy in the same study; symptoms resolved within 6 months.

Transient headaches

Transient headaches were reported in 8% (8/98) and 5% (3/57) of patients in the 2 non-randomised comparative studies of patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy or endoscopic adhesiolysis and foraminoplasty without discectomy. The authors did not state which groups these patients were in^{1,2}. Headache was reported in 1% (3/250) of patients in a case series of 250 patients⁵.

Focal infection

Focal infection was reported in 2% (2/98) and 4% (2/57) of patients in the 2 non-randomised comparative studies of patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy or endoscopic adhesiolysis and foraminoplasty without discectomy. The authors did not state which groups these patients were in^{1,2}.

Meningitis

Meningitis was reported in 1 patient each in the 2 non-randomised comparative studies of patients treated by endoscopic adhesiolysis, foraminoplasty and discectomy or endoscopic adhesiolysis and foraminoplasty without discectomy. The authors did not state which treatment groups these patients were in. Symptoms resolved after bed rest and symptomatic treatment^{1,2}.

Epidural pneumocephalus

IP overview: Epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

Epidural pneumocephalus was reported in 1 patient in the case series of 250 patients (no further information given)⁵.

Validity and generalisability of the studies

- Most of the studies were done in Korea. The results from this population may not be generalisable to the UK population.
- There may be some patient overlap, particularly between the 2 non-randomised comparative studies^{1,2}.
- Two studies evaluated the efficacy of percutaneous endoscopic laser lumbar discectomy as an adjunct to adhesiolysis and foraminoplasty^{1,2}.
- One study specifically excluded patients with failed back surgery syndrome².
- Studies used a number of different subjective scales but few objective efficacy outcome measures were reported.

Existing assessments of this procedure

There were no published assessments from other organisations identified at the time of the literature search.

Related NICE guidance

Below is a list of NICE guidance related to this procedure. Appendix B gives details of the recommendations made in each piece of guidance listed.

Interventional procedures

- Percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain. NICE interventional procedure guidance 545 (2016). Available from <http://www.nice.org.uk/guidance/ipg545>
- Percutaneous intradiscal electrothermal treatment of the intervertebral disc annulus for low back pain and sciatica. NICE interventional procedure guidance 544 (2016). Available from <http://www.nice.org.uk/guidance/ipg544>
- Percutaneous coblation of the intervertebral disc for low back pain and sciatica. NICE interventional procedure guidance 543 (2016). Available from <http://www.nice.org.uk/guidance/ipg543>

- Insertion of an annular disc implant at lumbar discectomy. NICE interventional procedure guidance 506 (2014). Available from <http://www.nice.org.uk/guidance/ipg506>
- Percutaneous intradiscal laser ablation in the lumbar spine. NICE interventional procedure guidance 357 (2010). Available from <http://www.nice.org.uk/guidance/ipg357>
- Prosthetic intervertebral disc replacement in the lumbar spine. NICE interventional procedure guidance 306 (2009). Available from <http://www.nice.org.uk/guidance/ipg306>
- Automated percutaneous mechanical lumbar discectomy. NICE interventional procedure guidance 141 (2005). Available from <http://www.nice.org.uk/guidance/ipg141>
- Percutaneous endoscopic laser thoracic discectomy. NICE interventional procedure guidance 61 (2004) Available from <http://www.nice.org.uk/guidance/ipg61>
- Endoscopic laser foraminoplasty. NICE interventional procedure guidance 31 (2003). Available from <http://www.nice.org.uk/guidance/ipg31>

NICE guidelines

- Low back pain and sciatica in over 16s: assessment and management. NICE guideline 59 (2016). Available from: <http://www.nice.org.uk/guidance/ng59>

Specialist advisers' opinions

Specialist advice was sought from consultants who have been nominated or ratified by their Specialist Society or Royal College. The advice received is their individual opinion and is not intended to represent the view of the society. The advice provided by Specialist Advisers, in the form of the completed questionnaires, is normally published in full on the NICE website during public consultation, except in circumstances but not limited to, where comments are considered voluminous, or publication would be unlawful or inappropriate. Three Specialist Adviser questionnaires for epiduroscopic lumbar discectomy through the sacral hiatus for sciatica were submitted and can be found on the [NICE website](#).

Patient commentators' opinions

NICE's Public Involvement Programme was unable to gather patient commentary for this procedure.

Issues for consideration by IPAC

None other than those described above.

References

1. Kim JD, Jang JH, Jung GH et al. (2011) Epiduroscopic laser disc and neural decompression. *Journal of neurosurgical review supplement* 1: 14-19
2. Lee GW, Jang SJ, Kim JD (2014) The efficacy of epiduroscopic neural decompression with Ho:YAG laser ablation in lumbar spinal stenosis. *Eur J Orthop Surg Traumatol. Suppl* 1:S231-7. doi: 10.1007/s00590-013-1407-7.
3. Richter EO, Abramova MV, Cantu F et al. (2011) Anterior epiduroscopic neural decompression: Eight-center experience in 154 patients. *European Journal of Pain* 5: 401-407
4. Jo DH, Yang HJ (2013) The survey of the patient received the epiduroscopic laser neural decompression. *Korean Journal of Pain* 26: 27–31
5. Lee SH, Lee SH, Lim KT. (2016) Trans-sacral epiduroscopic laser decompression for symptomatic lumbar disc herniation: a preliminary case series. *Photomedicine and Laser Surgery* 34: 121–9

Appendix A: Additional papers on epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

No additional papers were identified.

Appendix B: Related NICE guidance for epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

Guidance	Recommendations
-----------------	------------------------

Interventional procedures	<p>Percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain. NICE interventional procedure guidance 545 (2016)</p> <p>1.1 Current evidence on percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain raises no major safety concerns. The evidence on its efficacy is limited in quantity and quality. Therefore, this procedure should only be used with special arrangements for clinical governance, consent and audit or research.</p> <p>1.2 Clinicians wishing to do percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain should:</p> <ul style="list-style-type: none"> • Inform the clinical governance leads in their NHS trusts. • Ensure that patients understand the uncertainty about the procedure's efficacy and provide them with clear written information. In particular, patients should be informed about other treatment options, about the possibility that the procedure may not relieve their symptoms, and about the risk of a flare-up of their pain after treatment. In addition, the use of NICE's information for the public is recommended. • Audit and review clinical outcomes of all patients having percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain (see section 7.2). <p>1.3 NICE encourages further research into percutaneous intradiscal radiofrequency treatment of the intervertebral disc nucleus for low back pain. Further research should include details of patient selection, the duration of patients' symptoms, and a precise account of the technique used for treatment. Outcome measures should include pain relief and quality of life. Long-term follow-up data should include details of any subsequent procedures.</p> <p>Percutaneous electrothermal treatment of the intervertebral disc annulus for low back pain and sciatica. NICE interventional procedure guidance 544 (2016)</p> <p>1.1 Current evidence on percutaneous electrothermal treatment of the intervertebral disc annulus for low back pain and sciatica raises no major safety concerns. The evidence on efficacy is inconsistent and of poor quality. Therefore, this procedure should only be used with special arrangements for clinical governance, consent and audit or research.</p> <p>1.2 Clinicians wishing to do percutaneous electrothermal treatment of the intervertebral disc annulus for low back pain and sciatica should:</p> <ul style="list-style-type: none"> • Inform the clinical governance leads in their NHS trusts. • Ensure that patients understand the uncertainty about the procedure's efficacy and provide them with clear written
---------------------------	---

	<p>information. In particular, patients should be informed about other treatment options, about the possibility that the procedure may not relieve their symptoms, and about the risk of a flare-up of their pain following treatment. In addition, the use of NICE's information for the public is recommended.</p> <ul style="list-style-type: none"> • Audit and review clinical outcomes of all patients having percutaneous intradiscal radiofrequency treatment of the intervertebral disc annulus (see section 7.2). <p>1.3 NICE encourages further research into percutaneous electrothermal treatment of the intervertebral disc annulus. Further research should document details of patient selection, including the duration of their symptoms. It should report precise details of the technique used for treatment. Outcome measures should include pain relief and quality of life. Long-term follow-up data should include details of any subsequent procedures.</p> <p>Percutaneous coblation of the intervertebral disc for low back pain and sciatica. NICE interventional procedure guidance 543 (2016)</p> <p>1.1 Current evidence on percutaneous coblation of the intervertebral disc for low back pain and sciatica raises no major safety concerns. The evidence on efficacy is adequate and includes large numbers of patients with appropriate follow-up periods. Therefore, this procedure may be used provided that normal arrangements are in place for clinical governance, consent and audit.</p> <p>1.2 As part of the consent process, patients should be informed that there is a range of treatment options available to them and also that further procedures may be needed.</p> <p>Insertion of an annular disc implant at lumbar discectomy. NICE interventional procedure guidance 506 (2014)</p> <p>1.1 Current evidence on the safety and efficacy of insertion of an annular disc implant at lumbar discectomy is limited in quantity and quality. Therefore, this procedure should only be used with special arrangements for clinical governance, consent and audit or research.</p> <p>1.2 Clinicians wishing to undertake insertion of an annular disc implant at lumbar discectomy should take the following actions:</p> <p>Inform the clinical governance leads in their NHS trusts.</p> <p>Ensure that patients and their carers understand the uncertainty about the procedure's safety and efficacy and provide them with clear written information. In addition, the use of NICE's information for the public is recommended.</p>
--	--

	<p>1.3 NICE encourages further research on insertion of an annular disc implant at lumbar discectomy, particularly comparative trials. All studies should report details of patient selection and recurrence rates.</p> <p>1.4 Clinicians should enter details about all patients undergoing insertion of an annular disc implant at lumbar discectomy onto the British Spine Registry and review clinical outcomes locally.</p> <p>Percutaneous intradiscal laser ablation in the lumbar spine. NICE interventional procedure guidance 357 (2010)</p> <p>1.1 Current evidence on the safety and efficacy of percutaneous intradiscal laser ablation in the lumbar spine is adequate to support the use of this procedure provided that normal arrangements are in place for clinical governance, consent and audit.</p> <p>1.2 Patients selected for the procedure should be limited to those with severe pain refractory to conservative treatment, in whom imaging studies show bulging of an intact disc, and who do not have neurological deficit requiring surgical decompression.</p> <p>Prosthetic intervertebral disc replacement in the lumbar spine. NICE interventional procedure guidance 306 (2009)</p> <p>1.1 Current evidence on the safety and efficacy of prosthetic intervertebral disc replacement in the lumbar spine is adequate to support the use of this procedure provided that normal arrangements are in place for clinical governance, consent and audit.</p> <p>1.2 A multidisciplinary team with specialist expertise in the treatment of degenerative spine disease should be involved in patient selection for prosthetic intervertebral disc replacement in the lumbar spine. The procedure should only be carried out in patients for whom conservative treatment options have failed or are contraindicated.</p> <p>1.3 The current evidence includes studies with a maximum follow-up of 13 years, but the majority of evidence is from studies with shorter durations of follow-up. NICE encourages clinicians to continue to collect and publish data on longer-term outcomes, which should include information about patient selection and the need for further surgery.</p> <p>Automated percutaneous mechanical lumbar discectomy. NICE interventional procedure guidance 141 (2005)</p> <p>1.1 Current evidence suggests that there are no major safety concerns associated with automated percutaneous mechanical lumbar discectomy. There is limited evidence of efficacy based on uncontrolled case series of heterogeneous groups of patients, but evidence from small randomised controlled trials shows conflicting</p>
--	---

	<p>results. In view of the uncertainties about the efficacy of the procedure, it should not be used without special arrangements for consent and for audit or research.</p> <p>1.2 Clinicians wishing to undertake automated percutaneous mechanical lumbar discectomy should take the following actions.</p> <ul style="list-style-type: none"> • Inform the clinical governance leads in their Trusts. • Ensure that patients understand the uncertainty about the procedure's efficacy and provide them with clear written information. In addition, use of the Institute's information for the public is recommended. • Audit and review clinical outcomes of all patients having automated mechanical percutaneous lumbar discectomy. The Institute may review the procedure upon publication of further evidence. <p>Percutaneous endoscopic laser thoracic discectomy. NICE interventional procedure guidance 61 (2004)</p> <p>1.1 Current evidence on the safety and efficacy of percutaneous endoscopic laser thoracic discectomy does not appear adequate for this procedure to be used without special arrangements for consent and for audit or research.</p> <p>1.2 Clinicians wishing to undertake percutaneous endoscopic laser thoracic discectomy should take the following action.</p> <ul style="list-style-type: none"> • Inform the clinical governance leads in their Trusts. • Ensure that patients understand the uncertainty about the procedure's safety and efficacy and provide them with clear written information. Use of the Institute's information for the public is recommended. • Audit and review clinical outcomes of all patients having percutaneous endoscopic laser thoracic discectomy. <p>1.3 Further research will be useful in reducing the current uncertainty and clinicians are encouraged to collect longer-term follow-up data. The Institute may review the procedure upon publication of further evidence.</p> <p>Endoscopic laser foraminoplasty. NICE interventional procedure guidance 31 (2003)</p> <p>1.1 Current evidence of the safety and efficacy of endoscopic laser foraminoplasty does not appear adequate to support the use of this procedure without special arrangements for consent and for audit or research. Clinicians wishing to undertake endoscopic laser foraminoplasty should inform the clinical governance leads in their Trusts. They should ensure that patients offered the procedure understand the uncertainty about its safety and efficacy and should provide them with clear written information. Use of the Institute's</p>
--	---

	<p>information for the public is recommended. Clinicians should ensure that appropriate arrangements are in place for audit or research. Further research into safety and efficacy outcomes will be useful in reducing the current uncertainty. NICE is not undertaking further investigation at present.</p>
--	---

NICE guidelines	<p>Low back pain and sciatica in over 16s: assessment and management. NICE guideline 59 (2016).</p> <p>1.1 Assessment of low back pain and sciatica</p> <p>Alternative diagnoses</p> <p>1.1.1 Think about alternative diagnoses when examining or reviewing people with low back pain, particularly if they develop new or changed symptoms. Exclude specific causes of low back pain, for example, cancer, infection, trauma or inflammatory disease such as spondyloarthritis. If serious underlying pathology is suspected, refer to relevant NICE guidance on:</p> <ul style="list-style-type: none"> • Metastatic spinal cord compression in adults • Spinal injury • Spondyloarthritis • Suspected cancer <p>Risk assessment and risk stratification tools</p> <p>1.1.2 Consider using risk stratification (for example, the STarT Back risk assessment tool) at first point of contact with a healthcare professional for each new episode of low back pain with or without sciatica to inform shared decision-making about stratified management.</p> <p>1.1.3 Based on risk stratification, consider:</p> <ul style="list-style-type: none"> • simpler and less intensive support for people with low back pain with or without sciatica likely to improve quickly and have a good outcome (for example, reassurance, advice to keep active and guidance on self-management) • more complex and intensive support for people with low back pain with or without sciatica at higher risk of a poor outcome (for example, exercise programmes with or without manual therapy or using a psychological approach). <p>Imaging</p> <p>1.1.4 Do not routinely offer imaging in a non-specialist setting for people with low back pain with or without sciatica.</p> <p>1.1.5 Explain to people with low back pain with or without sciatica that if they are being referred for specialist opinion, they may not need imaging.</p> <p>1.1.6 Consider imaging in specialist settings of care (for example, a musculoskeletal interface clinic or hospital) for people with low back pain with or without sciatica only if the result is likely to change management.</p> <p>1.2 Non-invasive treatments for low back pain and sciatica</p> <p>Non-pharmacological interventions</p> <p>Self-management</p>
-----------------	---

	<p>1.2.1 Provide people with advice and information, tailored to their needs and capabilities, to help them self-manage their low back pain with or without sciatica, at all steps of the treatment pathway. Include:</p> <ul style="list-style-type: none"> • information on the nature of low back pain and sciatica • encouragement to continue with normal activities. <p>Exercise</p> <p>1.2.2 Consider a group exercise programme (biomechanical, aerobic, mind–body or a combination of approaches) within the NHS for people with a specific episode or flare-up of low back pain with or without sciatica. Take people's specific needs, preferences and capabilities into account when choosing the type of exercise.</p> <p>Orthotics</p> <p>1.2.3 Do not offer belts or corsets for managing low back pain with or without sciatica.</p> <p>1.2.4 Do not offer foot orthotics for managing low back pain with or without sciatica.</p> <p>1.2.5 Do not offer rocker sole shoes for managing low back pain with or without sciatica.</p> <p>Manual therapies</p> <p>1.2.6 Do not offer traction for managing low back pain with or without sciatica.</p> <p>1.2.7 Consider manual therapy (spinal manipulation, mobilisation or soft tissue techniques such as massage) for managing low back pain with or without sciatica, but only as part of a treatment package including exercise, with or without psychological therapy.</p> <p>Acupuncture</p> <p>1.2.8 Do not offer acupuncture for managing low back pain with or without sciatica.</p> <p>Electrotherapies</p> <p>1.2.9 Do not offer ultrasound for managing low back pain with or without sciatica.</p> <p>1.2.10 Do not offer percutaneous electrical nerve stimulation (PENS) for managing low back pain with or without sciatica.</p> <p>1.2.11 Do not offer transcutaneous electrical nerve stimulation (TENS) for managing low back pain with or without sciatica.</p> <p>1.2.12 Do not offer interferential therapy for managing low back pain with or without sciatica.</p> <p>Psychological therapy</p> <p>1.2.13 Consider psychological therapies using a cognitive behavioural approach for managing low back pain with or without sciatica but only as part of a treatment package including exercise, with or without manual therapy (spinal manipulation, mobilisation or soft tissue techniques such as massage).</p> <p>Combined physical and psychological programmes</p>
--	--

	<p>1.2.14 Consider a combined physical and psychological programme, incorporating a cognitive behavioural approach (preferably in a group context that takes into account a person's specific needs and capabilities), for people with persistent low back pain or sciatica:</p> <ul style="list-style-type: none"> • when they have significant psychosocial obstacles to recovery (for example, avoiding normal activities based on inappropriate beliefs about their condition) or • when previous treatments have not been effective. <p>Return-to-work programmes</p> <p>1.2.15 Promote and facilitate return to work or normal activities of daily living for people with low back pain with or without sciatica.</p> <p>Pharmacological interventions</p> <p>1.2.16 For recommendations on pharmacological management of sciatica, see NICE's guideline on neuropathic pain in adults.</p> <p>1.2.17 Consider oral non-steroidal anti-inflammatory drugs (NSAIDs) for managing low back pain, taking into account potential differences in gastrointestinal, liver and cardio-renal toxicity, and the person's risk factors, including age.</p> <p>1.2.18 When prescribing oral NSAIDs for low back pain, think about appropriate clinical assessment, ongoing monitoring of risk factors, and the use of gastroprotective treatment.</p> <p>1.2.19 Prescribe oral NSAIDs for low back pain at the lowest effective dose for the shortest possible period of time.</p> <p>1.2.20 Consider weak opioids (with or without paracetamol) for managing acute low back pain only if an NSAID is contraindicated, not tolerated or has been ineffective.</p> <p>1.2.21 Do not offer paracetamol alone for managing low back pain.</p> <p>1.2.22 Do not routinely offer opioids for managing acute low back pain (see recommendation 1.2.20).</p> <p>1.2.23 Do not offer opioids for managing chronic low back pain.</p> <p>1.2.24 Do not offer selective serotonin reuptake inhibitors, serotonin–norepinephrine reuptake inhibitors or tricyclic antidepressants for managing low back pain.</p> <p>1.2.25 Do not offer anticonvulsants for managing low back pain.</p> <p>1.3 Invasive treatments for low back pain and sciatica</p> <p>Non-surgical interventions</p> <p>Spinal injections</p> <p>1.3.1 Do not offer spinal injections for managing low back pain.</p> <p>Radiofrequency denervation</p> <p>1.3.2 Consider referral for assessment for radiofrequency denervation for people with chronic low back pain when:</p> <ul style="list-style-type: none"> • non-surgical treatment has not worked for them and • the main source of pain is thought to come from structures supplied by the medial branch nerve and
--	--

	<ul style="list-style-type: none"> • they have moderate or severe levels of localised back pain (rated as 5 or more on a visual analogue scale, or equivalent) at the time of referral. <p>1.3.3 Only perform radiofrequency denervation in people with chronic low back pain after a positive response to a diagnostic medial branch block.</p> <p>1.3.4 Do not offer imaging for people with low back pain with specific facet joint pain as a prerequisite for radiofrequency denervation.</p> <p>Epidurals</p> <p>1.3.5 Consider epidural injections of local anaesthetic and steroid in people with acute and severe sciatica.</p> <p>1.3.6 Do not use epidural injections for neurogenic claudication in people who have central spinal canal stenosis.</p> <p>Surgical interventions</p> <p>Surgery and prognostic factors</p> <p>1.3.7 Do not allow a person's BMI, smoking status or psychological distress to influence the decision to refer them for a surgical opinion for sciatica.</p> <p>Spinal decompression</p> <p>1.3.8 Consider spinal decompression for people with sciatica when non-surgical treatment has not improved pain or function and their radiological findings are consistent with sciatic symptoms.</p> <p>Spinal fusion</p> <p>1.3.9 Do not offer spinal fusion for people with low back pain unless as part of a randomised controlled trial.</p> <p>Disc replacement</p> <p>1.3.10 Do not offer disc replacement in people with low back pain.</p>
--	--

Appendix C: Literature search for epiduroscopic lumbar discectomy through the sacral hiatus for sciatica

Databases	Date searched	Version/files
Cochrane Database of Systematic Reviews – CDSR (Cochrane)	06/06/2016	Issue 6 of 12, June 2016
HTA database (Cochrane)	06/06/2016	Issue 6 of 12, June 2016
Cochrane Central Database of Controlled Trials – CENTRAL (Cochrane)	06/06/2016	Issue 6 of 12, June 2016
MEDLINE (Ovid)	06/06/2016	1946 to May Week 4 2016
MEDLINE In-Process (Ovid)	06/06/2016	June 03, 2016
EMBASE (Ovid)	06/06/2016	1974 to 2016 Week 23
PubMed	06/06/2016	-
BLIC (British Library)	06/06/2016	-

The following search strategy was used to identify papers in MEDLINE. A similar strategy was used to identify papers in other databases.

1	epiduroscop*.tw. (63)
2	(epidural* adj4 (endoscop* or myeloscop*)).tw. (46)
3	(ELND or (epiduroscop* adj4 laser* adj4 neural* adj4 decompress*)).tw. (109)
4	(flex* adj4 endoscop*).tw. (2127)
5	(sacral* adj4 (hiatus* or vertebra* or canal*)).tw. (819)
6	((trans* adj4 sacral*) or trans-sacral*).tw. (406)
7	or/1-6 (3513)
8	Low Back Pain/ (15837)
9	(low* adj4 (backpain* or back pain* or backache* or back ache*)).tw. (18606)
10	LBP.tw. (4320)
11	lumbago*.tw. (1093)
12	Sciatica/ (4451)
13	sciatic*.tw. (22474)
14	(chronic* adj4 back pain*).tw. (5601)

15	((low* extremity or leg*) adj4 (weak* or pain* or ache* or discomfort*).tw. (8270)
16	((displac* or degenerat* or degrad* or deteriorat* or decay* or slip* or extru* or hernia* or prolaps* or protru* or compress* or an?ulus or bulg*) adj4 (disc* or disk*).tw. (22198)
17	((discogenic* or diskogenic* or radicular or non-dermatomal) adj4 pain*).tw. (2695)
18	Radiculopathy/ (3958)
19	(lumbar adj4 radiculopath*).tw. (584)
20	(nerv* adj4 root* adj4 imping*).tw. (98)
21	(cauda adj4 equina adj4 syndrome*).tw. (1085)
22	paraparesis/ (643)
23	paraparesis.tw. (4700)
24	(spinal adj4 fibrosis).tw. (80)
25	or/8-24 (83743)
26	7 and 25 (231)