

IPG681- Pressurised intraperitoneal aerosol chemotherapy for peritoneal carcinomatosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

T48.2 Introduction of cytotoxic substance into peritoneal cavity

Y75.2 Laparoscopic approach to abdominal cavity NEC

X72–X73 Delivery of chemotherapy for neoplasm

Note: An OPCS-4 code from the range **X72-X73** from the National Tariff Chemotherapy Regimens List 2017-18 is assigned, as per the instructions in National Clinical Coding Standards OPCS-4 reference book **PCSX27: Delivery of chemotherapy for neoplasm (X72-X73)**.

Note: Only delivery of chemotherapy codes from categories **X72** and **X73** are required for national reporting purposes. The collection of codes classifying procurement is not required.

Note: It is not possible to classify the ‘pressurised aerosolised’ aspect of this treatment within OPCS-4.

Diagnosis or health condition:

ICD-10 code(s):

C78.6 Secondary malignant neoplasm of retroperitoneum and peritoneum

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).