

IPG719 - Endoscopic balloon dilation for subglottic or tracheal stenosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

Balloon dilatation of tracheal stenosis with the use of a bronchoscope

E48.8 Other specified therapeutic fiberoptic endoscopic operations on lower respiratory tract

Y40.3 Balloon dilation of organ NOC

Z24.3 Trachea

Balloon dilatation of subglottic stenosis with the use of a laryngoscope

E35.8 Other specified other therapeutic endoscopic operations on larynx

Y40.3 Balloon dilation of organ NOC

Z88.5 Subglottis

Diagnosis or health condition:

ICD-10 code(s):

Tracheal stenosis

J39.8 Other specified diseases of upper respiratory tract

Or

Q32.1 Other congenital malformations of trachea

Subglottic stenosis

J38.6 Stenosis of larynx

Or

J95.5 Postprocedural subglottic stenosis

Or

Q31.1 Congenital subglottic stenosis

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).