

## IP923/2 - Percutaneous transluminal renal sympathetic denervation for resistant hypertension

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

### **Procedure:**

#### **OPCS-4 code(s):**

Renal sympathetic denervation using radiofrequency:

**L43.6 Percutaneous transluminal radiofrequency denervation of renal artery**

**Y53.4 Approach to organ under fluoroscopic control**

**Z94.- Laterality of operation**

Renal sympathetic denervation using ultrasound:

**L43.8 Other specified transluminal operations on renal artery**

**Y11.5 Ultrasonic destruction of organ NOC**

**Y53.4 Approach to organ under fluoroscopic control**

**Z94.- Laterality of operation**

### **Diagnosis or health condition:**

#### **ICD-10 code(s):**

**I10.X Essential (primary) hypertension**

**I15.- Secondary hypertension**

## **Clinical coding recommendations for NICE guidance**

For each published interventional procedure guidance, we work with NHS Digital to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS Digital Information Representation Services work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS Digital is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).