

IPG77 - Balloon dilatation of systemic to pulmonary arterial shunts in children

SNOMED CT provides clinical terms for entry into the patient record to record clinical information relevant to that encounter; the mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure and device:

SNOMED CT preferred term (concept ID)

Fluoroscopy guided balloon dilatation of systemic to pulmonary artery shunt (853551000000106)

OPCS-4 code(s):

L05.4 Percutaneous transluminal balloon dilation of interposition tube prosthesis between pulmonary artery and aorta

Y53.1 Approach to organ under radiological control

Diagnosis or health condition:

SNOMED CT preferred term (concept ID)

Partial obstruction of systemic to pulmonary artery shunt (448651008)

Complete obstruction of systemic to pulmonary artery shunt (448622009)

Clinical coding recommendations for NICE guidance

For each published interventional procedure and medical technologies guidance, we work with NHS Digital to provide relevant clinical coding information.

SNOMED CT provides clinical terms for entry into the patient record to store clinical information relevant to that encounter.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

The UK Edition of SNOMED CT is managed by the Clinical Terminology Service of NHS Digital. For further information including licensing, see [UK Terminology Centre — NHS Digital](#).

The Clinical Classifications Service of NHS Digital is the central definitive source for clinical coding guidance and determines the coding standards associated with the classifications (OPCS-4 and ICD-10) to be used across the NHS. The Clinical Classifications Service and NICE work collaboratively to ensure the most appropriate classification codes are provided. [Clinical Classifications Service — NHS Digital](#).