

IP881/2 Transfemoral carotid artery stent placement for asymptomatic extracranial carotid stenosis

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

Procedure:

OPCS-4 code(s):

L31.4 Percutaneous transluminal insertion of stent into carotid artery

Plus

L76.- Endovascular placement of stent

Or

L89.- Other endovascular placement of stent

Or

O20.- Endovascular placement of stent graft

Plus

Y79.3 Transluminal approach to organ through femoral artery

Plus

Y53.- Approach to organ under image control

Or

Y68.- Other approach to organ under image control

Note: Multiple codes from **Y53** and **Y68** may be assigned where different types of image control are used.

Diagnosis or health condition:

ICD-10 code(s):

I65.2 Occlusion and stenosis of carotid artery

Or

I65.3 Occlusion and stenosis of multiple and bilateral precerebral arteries

Clinical coding recommendations for NICE guidance

For each published interventional procedure guidance, we work with NHS England to provide relevant clinical coding information.

The mandated classifications (OPCS-4 or ICD-10) provide a method to collect and aggregate data to allow accurate and consistent data analysis.

NICE and NHS England Terminology and Classifications Delivery Service work collaboratively to ensure the most appropriate OPCS-4 and ICD-10 codes are provided.

NHS England is the national release centre for the UK edition of ICD-10 and OPCS-4. Further Classifications information, see [here](#).