

## Addendum to Intrapartum care: care for healthy women and babies

### Appendix G Evidence tables

#### G.1 Intermittent auscultation compared with cardiotocography on admission

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Mitchell,K., The effect of the labour electronic fetal monitoring admission test on operative delivery in low-risk women: a randomised controlled trial, Evidence Based Midwifery, 6, 18-26, 2008</p> <p><b>Ref Id</b></p> <p>66879</p> <p><b>Country/ies where the study was carried out</b></p> <p>England</p> <p><b>Study type</b></p> <p>Randomised controlled trial</p>	<p><b>Sample size</b></p> <p>See entry in systematic review by Devane 2012</p> <p><b>Characteristics</b></p> <p><b>Parity (n (%))</b></p> <p>- 0 Cardiotocograph (CTG): 203 (70) Auscultation: 199 (68)</p> <p>- 1 or more CTG: 95 (30) Auscultation: 85 (32)</p> <p><b>Inclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p>	<p><b>Interventions</b></p> <p>Admission CTG</p> <p>Intermittent auscultation</p>	<p><b>Details</b></p> <p><b>Care during labour</b></p> <p>Following the admission CTG, the decision to end tracing and start intermittent monitoring was left up to the midwives and clinicians caring for the woman. The CTG was stopped when it was considered normal (as defined by the 2001 NICE inherited guideline on the use of EFM). This meant that the length of CTG could vary between the 15 minute admission test and the whole labour period.</p>	<p><b>Results</b></p> <p>All priority outcomes of interest were reported by the authors of the systematic review</p>	<p><b>Limitations</b></p> <p>See entry in systematic review by Devane 2012</p> <p><b>Other information</b></p> <p>MOST STUDY DETAILS ARE REPORTED IN DEVANE 2012. THIS ENTRY ONLY REPORTS EXTRA DETAILS THAT WERE NOT REPORTED IN THE COCHRANE REVIEW, WHICH THE TECHNICAL TEAM FELT WERE IMPORTANT CONSIDERATIONS WHEN INTERPRETING THE RESULTS</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To test the relationship between the labour electronic fetal monitoring (EFM) admission test and obstetric intervention</p> <p><b>Study dates</b></p> <p>15th December 2002 to 30th June 2006</p> <p><b>Source of funding</b></p> <p>Initial grant from the Buckinghamshire Hospitals NHS Trust's Research Department and establishment of a research midwife role in the unit</p>	<p><b>Exclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p>		<p>Women allocated to auscultation were intermittently monitored during labour. However, regardless of allocation, if the woman was considered to have become higher risk, continuous EFM was offered and recommended as per unit policy.</p> <p>Analysis was by intention to treat</p>		
<p><b>Full citation</b></p> <p>Cheyne,H., Dunlop,A., Shields,N., Mathers,A.M., A randomised</p>	<p><b>Sample size</b></p> <p>See entry in systematic review by Devane 2012</p>	<p><b>Interventions</b></p> <p>Admission EFM</p> <p>Intermittent auscultation</p>	<p><b>Details</b></p> <p><u>Care during labour</u></p> <p>Following randomisation, women received</p>	<p><b>Results</b></p> <p>All priority outcomes of interest reported in trial are reported in the systematic</p>	<p><b>Limitations</b></p> <p>See Devane 2012 for risk of bias assessment</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>controlled trial of admission electronic fetal monitoring in normal labour, Midwifery, 19, 221-229, 2003</p> <p><b>Ref Id</b></p> <p>158779</p> <p><b>Country/ies where the study was carried out</b></p> <p>Scotland</p> <p><b>Study type</b></p> <p>Randomised controlled trial</p> <p><b>Aim of the study</b></p> <p>To test the hypothesis that admission electronic fetal monitoring (EFM) for healthy pregnant women in spontaneous labour would lead to an increase in continuous EFM when compared to</p>	<p><b>Characteristics</b></p> <p><b><u>Women having artificial rupture of membranes (n (%))</u></b></p> <p>Cardiotocgraph (CTG): 65 (44%) Auscultation: 60 (36%)</p> <p><b><u>Primiparous women (n (%))</u></b></p> <p>CTG: 65 (44%) Auscultation: 76 (46%)</p> <p><b>Inclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p> <p><b>Exclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p>	<p>with a hand-held Doppler device</p>	<p>either a routine 20 minute period of EFM at the time of admission to the Midwives Birth Unit, or auscultation immediately following a contraction for a minimum of 60 seconds.</p> <p>With the exception of the randomised intervention, women received the same admission assessment, i.e. history taking, blood pressure measurement, temperature recording, abdominal palpation, and vaginal examination.</p> <p>Subsequently, all women were monitored using intermittent auscultation, at 15 minute intervals in the first stage of labour and at 5</p>	<p>review (Devane 2012)</p>	<p>MOST STUDY DETAILS ARE REPORTED IN DEVANE 2012. THIS ENTRY ONLY REPORTS EXTRA DETAILS THAT WERE NOT REPORTED IN THE COCHRANE REVIEW, WHICH THE TECHNICAL TEAM FELT WERE IMPORTANT CONSIDERATIONS WHEN INTERPRETING THE RESULTS</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>women who have no admission EFM</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>North Glasgow University Hospitals NHS Trust</p>			<p>minute intervals, or after a contraction, during the second stage of labour. EFM was used, where required, in accordance with the guidelines for the unit. However, it should be noted that in addition to the women who received continuous EFM during labour (as reported in the systematic review), a further 125 (84%) of women in the CTG arm and 61 (37%) of the auscultation arm received additional EFM during labour.</p> <p>The reasons were (n (%)):</p> <ul style="list-style-type: none"> <li>- Admission EFM not discontinued CTG: 80 (64)</li> <li>Auscultation: 1 (2)</li> <li>- FHR abnormalities noted</li> </ul>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>CTG: 29 (23) Auscultation: 13 (21)</p> <p>- EFM commenced on transfer to labour ward CTG: 10 (8) Auscultation: 33 (54)</p> <p>- Meconium stained liquor CTG: 2 (2) Auscultation: 9 (15)</p> <p>- Other CTG: 4 (3) Auscultation: 5 (8)</p>		
<p><b>Full citation</b></p> <p>Devane,D., Lalor,J.G., Daly,S., McGuire,W., Smith,V., Cardiotocography versus intermittent auscultation of fetal heart on admission to labour ward for assessment of fetal wellbeing, Cochrane Database of</p>	<p><b>Sample size</b></p> <p>Trials: N = 4 Women: N = 13296</p> <p><b>Characteristics</b></p> <p><u>Cheyne (2003)</u> - Inclusion criteria: Healthy women with a normal pregnancy, presenting in</p>	<p><u>Admission CTG:</u> Defined as a commonly used screening test, comprising a short, usually 20 minute long, recording of the FHR and uterine activity</p>	<p><u>Searching for studies</u> The Trials Search Co-ordinator was contacted on 17 May 2011, and asked to search the Cochrane Pregnancy and Childbirth Group's Trials Register. In addition, CENTRAL,</p>	<p><b>Results</b></p> <p><u>Mode of birth (number/total)</u> a. Caesarean section CTG: 248/5657 Auscultation: 207/5681</p> <p>RR 1.20 (95% CI 1.00 to 1.44) Heterogeneity: I<sup>2</sup> = 0.0% Test for overall</p>	<p><b>Limitations</b></p> <p>The systematic review did not have any serious limitations.</p> <p>Impey (2003) included women with an early amniotomy, and only included women with clear amniotic fluid. The study also included some women (&lt; 5%) who had a previous caesarean section (CS) and who went into labour prior to 37 completed weeks' gestation. However, the authors of the review contacted the study authors, who provided data for women who went into labour at 37-42 weeks and without a previous CS, and the data for these women were used in the main analysis in the systematic review.</p> <p>Mires (2001) randomised women in the third trimester, and between randomisation and admission in labour, 37% of women developed a complication, so that only 2367 were judged to be low risk in labour. The</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Systematic Reviews, 2, CD005122-, 2012</p> <p><b>Ref Id</b></p> <p>157062</p> <p><b>Country/ies where the study was carried out</b></p> <p>Included trials were conducted in England, Scotland and Ireland</p> <p><b>Study type</b></p> <p>Systematic review of randomised controlled trials</p> <p><b>Aim of the study</b></p> <p>To compare the effects of admission cardiotocograph (CTG) with intermittent auscultation of the fetal heart rate (FHR) on maternal and infant outcomes for pregnant women without risk factors</p>	<p>spontaneous labour and who were eligible for admission to the Midwives Birth Unit</p> <p>- Exclusion criteria: Women with risk factors</p> <p>- N = 344 women randomised on admission in labour</p> <p>- Admission CTG: Routine 20 minute period at time of admission</p> <p>- Intermittent Auscultation: Fetal heart was auscultated during and immediately following a contraction for a minimum of 60 seconds</p> <p><u>Impey (2003)</u></p> <p>- Inclusion criteria: Admitted in labour, singleton pregnancy, less than 42 completed weeks' gestation, no suspicion or evidence of antenatal fetal compromise, no adverse obstetric history, clear</p>	<p><u>Intermittent auscultation:</u></p> <p>Intermittent surveillance of the FHR using a hand-held Doppler device or a Pinard stethoscope</p> <p>Both tests were performed upon the woman's admission to the labour ward.</p>	<p>MEDLINE, CINAHL and Dissertation Abstracts were searched. The reference list of identified studies was also searched, and any studies assessed for eligibility. No language restrictions were applied.</p> <p>No studies were excluded.</p> <p><u>Data collection and analysis</u></p> <p>Two review authors independently assessed studies for inclusion. They then extracted data into a predesigned form and resolved discrepancies through discussion. Data were entered into RevMan and checked for accuracy. If there was any unclear information, the</p>	<p>effect: Z = 2.00, p = 0.045</p> <p>[Note: the interpretation of this result by the authors of the systematic review is as follows. "Given that (i) the 95% CI just reaches 1.00 and (ii) the absence of measurable heterogeneity in this outcome analysis (T<sup>2</sup> = 0.00, I<sup>2</sup> = 0%), the probability is that admission CTG increases the caesarean section rate by approximately 20%."]</p> <p>[4 trials: Cheyne 2003, Impey 2003, Mires 2001, Mitchell 2008]</p> <p><u>b. Instrumental vaginal birth</u></p> <p>CTG: 782/5657</p> <p>Auscultation: 716/5681</p> <p>RR 1.10 (95% CI 0.95 to 1.27)</p> <p>Heterogeneity: I<sup>2</sup> =</p>	<p>low risk subgroup data were provided by the authors, and these were used in the analysis in the systematic review.</p> <p>The following represents the review author's risk of bias for the included studies. Overall, all studies were assessed as being at low risk of bias:</p> <p><u>Cheyne 2003</u></p> <ul style="list-style-type: none"> <li>- Random sequence generation: low risk of bias</li> <li>- Allocation concealment: low risk of bias</li> <li>- Blinding of outcome assessors: high risk of bias; they were not blinded</li> <li>- Incomplete outcome data: low risk of bias; the trial publication reported that 22 women (7%) were excluded from the analysis (21 not in labour, 1 missing randomisation card); however, the review authors contacted the trial authors and received data for 21/22 of them</li> <li>- Selective reporting: low risk of bias</li> </ul> <p><u>Impey 2003</u></p> <ul style="list-style-type: none"> <li>- Random sequence generation: low risk</li> <li>- Allocation concealment: low risk of bias</li> <li>- Blinding of outcome assessors: low risk of bias - data were entered and neonatal assessment was performed without knowledge of treatment allocation</li> <li>- Incomplete outcome data: low risk of bias; loss to follow-up was 0.5% in CTG arm and 0.6% in auscultation arm</li> <li>- Selective reporting: low risk of bias</li> </ul> <p><u>Mires 2001</u></p> <ul style="list-style-type: none"> <li>- Random sequence generation: low risk of bias</li> <li>- Allocation concealment: low risk of bias</li> <li>- Blinding of outcome assessors: low risk of bias; data analysts were blind to randomisation code</li> <li>- Incomplete outcome data: low risk of bias</li> <li>- Selective reporting: low risk of bias</li> <li>- Other bias: between randomisation (third trimester) and admission in labour, 1384 women (37%) developed a complication that warranted continuous FHR monitoring in labour; the authors provided data for the low-risk women separately and these were used for the analysis in the systematic review</li> </ul>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>for intrapartum hypoxia</p> <p><b>Study dates</b></p> <p>Content was assessed as up-to-date on 14 November 2011</p> <p><b>Source of funding</b></p> <p>Health Research Board, Ireland</p>	<p>amniotic fluid, maternal temperature of 37.5 degrees or less at admission</p> <p>- N = 8628 women randomised on admission in labour</p> <p>- Admission CTG: 20 minute admission CTG immediately after early amniotomy performed on diagnosis of labour in women presenting to delivery ward</p> <p>- Intermittent auscultation: Performed for 1 minute after a contraction every 15 minutes in the first stage of labour and every 5 minutes in the second stage. It was performed after early amniotomy on diagnosis of labour in women presenting to the delivery ward.</p> <p><u>Mires (2001)</u></p> <p>- Inclusion criteria: Booked for hospital birth, attended a</p>		<p>authors were contacted to provide details.</p> <p><u>Quality assessment</u></p> <p>Risk of bias was assessed independently by two authors using the The Cochrane Collaboration's tool for assessing risk of bias. The following criteria were considered:</p> <ul style="list-style-type: none"> <li>- Sequence generation</li> <li>- Allocation concealment</li> <li>- Blinding: due to the intervention, it would not be possible to blind participants or those providing care; however, the authors reported that they did consider whether outcome assessors were blinded</li> <li>- Incomplete outcome data: low risk was defined as 20% or less missing data, and high risk as more</li> </ul>	<p>38%</p> <p>Test for overall effect: <math>Z = 1.28</math>, <math>p = 0.20</math></p> <p>[4 trials: Cheyne 2003, Impey 2003, Mires 2001, Mitchell 2008]</p> <p><b>Fetal and neonatal deaths (number/total)</b></p> <p>CTG: 5/5658</p> <p>Auscultation: 5/5681</p> <p>RR 1.01 (95% CI 0.30 to 3.47 )</p> <p>Heterogeneity: <math>I^2 = 0.0\%</math></p> <p>Test for overall effect: <math>Z = 0.02</math>, <math>p = 0.98</math></p> <p>[4 trials: Cheyne 2003, Impey 2003, Mires 2001, Mitchell 2008]</p> <p><b>Major neonatal morbidity (number/total)</b></p> <p><u>a. Hypoxic ischaemic encephalopathy</u></p> <p>CTG: 6/1186</p> <p>Auscultation:</p>	<p><u>Mitchell 2008</u></p> <ul style="list-style-type: none"> <li>- Random sequence generation: low risk of bias</li> <li>- Allocation concealment: low risk of bias</li> <li>- Blinding of outcome assessors: unclear risk of bias - no details reported</li> <li>- Incomplete outcome data: low risk of bias</li> <li>- Selective reporting: low risk of bias</li> </ul> <p><b>Other information</b></p> <p>The systematic review is available online at: <a href="http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005122.pub4/full">http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005122.pub4/full</a></p> <p>The authors identified one trial which was ongoing - the ADCAR trial; it was unclear when this trial would be published.</p> <p><b>Monitoring during labour</b></p> <p>3 trials reported the number of women having continuous EFM in labour and in 2 of the trials, the difference was significant:</p> <p><b>Cheyne 2003:</b></p> <ul style="list-style-type: none"> <li>- CTG: 10/157 (6.4%)</li> <li>- Auscultation: 10/177 (5.6%) (NS)</li> </ul> <p>[Note: a further 125 women from the CTG arm and 61 women from the auscultation arm received additional EFM during labour]</p> <p><b>Impey 2003:</b></p> <ul style="list-style-type: none"> <li>- CTG: 2341/4017 (58.3%)</li> <li>- Auscultation: 1686/4039 (41.7%) (<math>p &lt; 0.00001</math>)</li> </ul> <p><b>Mires 2001:</b></p> <ul style="list-style-type: none"> <li>- CTG: 672/1185 (56.7%)</li> <li>- Auscultation: 551/1178 (46.8%) (<math>p &lt; 0.00001</math>)</li> </ul> <p><b>Total:</b></p> <ul style="list-style-type: none"> <li>- CTG: 3023/5359</li> </ul>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>hospital or community based consultant led clinic in the third trimester, and had no obstetric complications at that visit that would warrant continuous monitoring of FHR (pre-eclampsia or hypertension in previous or current pregnancy, essential hypertension, diabetes, suspected intrauterine growth restriction (IUGR), placental abruption or praevia or bleeding of unknown origin, multiple pregnancy, fetal malformation, previous caesarean section, breech presentation, or rhesus isoimmunisation)</p> <p>- N = 3752 women randomised during third trimester.</p> <p>- Admission CTG: 20 minute CTG on admission in spontaneous uncomplicated</p>		<p>than 20% missing data</p> <p>- Selective reporting bias: established by cross-checking the outcomes reported in the methods and results sections of the included publications</p> <p>- Other sources of bias</p> <p><u>Missing data</u> Levels of attrition were noted for the studies. Sensitivity analysis was performed to explore the effect of including studies with high attrition. All analyses were carried out on an intention-to-treat basis. Denominators were the number randomised, minus any women whose outcomes were known to be missing.</p> <p><u>Analysis</u></p>	<p>5/1181</p> <p>RR 1.19 (95% CI 0.37 to 3.90) Heterogeneity: NA Test for overall effect: Z = 0.29, p = 0.77</p> <p>[1 trial: Mires 2001]</p> <p><u>b. Neonatal seizures</u> CTG: 10/4017 Auscultation: 14/4039</p> <p>RR 0.72 (95% CI 0.32 to 1.61) Heterogeneity: I<sup>2</sup> = Test for overall effect: Z = , p =</p> <p>[1 trial: Impey 2003]</p> <p><u>Admission to NICU (number/total)</u> CTG: 219/5656 Auscultation: 213/5675</p> <p>RR 1.03 (95% CI 0.86 to 1.24) Heterogeneity: I<sup>2</sup> = 0.0% Test for overall</p>	<p>- Auscultation: 2247/5394 (RR 1.30 [95% CI 1.14 to 1.48])</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>labour - Intermittent auscultation: Auscultation of the fetal heart with hand-held Doppler device during and immediately after 1 contraction</p> <p><u>Mitchell (2008)</u></p> <p>- Inclusion criteria: Labouring women considered to be at 'low risk' of fetal or maternal complications on admission</p> <p>- Exclusion criteria: Any minor maternal medical complication (e.g. diabetes or essential hypertension), previous caesarean section, preterm labour (less than 37 completed weeks), multiple pregnancy, prolonged pregnancy (more than 42 weeks), prolonged membrane rupture (more than 24 hours), induction of labour, meconium-</p>		<p>Statistical analysis was performed in RevMan. A random effects model was used. This was because the authors felt that there was sufficient clinical heterogeneity to expect that the underlying treatment effect would differ. In Impey 2003, only women whose liquor was known to be clear were included. In the other trials, membrane rupture and clear liquor were not inclusion criteria</p>	<p>effect: Z = 0.32, p = 0.75</p> <p>[4 trials: Cheyne 2003, Impey 2003, Mires 2001, Mitchell 2008]</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>stained liquor, maternal pyrexia, rhesus sensitisation, polyhydramnios, oligohydramnios, pre-eclampsia or blood pressure over 140/90 mmHg, abnormal presentation or lie (e.g. breech, transverse), high head (5/5ths palpable per abdomen), antepartum or intrapartum haemorrhage, known or suspected IUGR, any known or suspected fetal medical complication, abnormal Doppler artery velocimetry, known fetal malformation, poor obstetric history (e.g. history of stillbirth), unbooked - N = 582 women randomised on admission in labour</p> <p>- Admission CTG: 15-minute CTG on admission in spontaneous uncomplicated</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>labour - Intermittent auscultation: Auscultation of the fetal heart for 1 continuous minute using a Pinard stethoscope or Doppler ultrasound device, after a contraction, at least every 15 minutes in the first stage of labour and every 5 minutes in the second stage</p> <p><b>Inclusion criteria</b></p> <p>Randomised and quasi-randomised trials comparing admission CTG with intermittent auscultation of the FHR</p> <p><b>Exclusion criteria</b></p> <p>None reported</p>				
<p><b>Full citation</b></p> <p>Impey,L., Reynolds,M.,</p>	<p><b>Sample size</b></p>	<p><b>Interventions</b></p> <p>Admission CTG</p>	<p><b>Details</b></p> <p><u>Care during labour</u></p>	<p><b>Results</b></p> <p>All priority outcomes were</p>	<p><b>Limitations</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>MacQuillan,K., Gates,S., Murphy,J., Sheil,O., Admission cardiocography: A randomised controlled trial, Lancet, 361, 465-470, 2003</p> <p><b>Ref Id</b> 60264</p> <p><b>Country/ies where the study was carried out</b>  Ireland</p> <p><b>Study type</b>  Randomised controlled trial</p> <p><b>Aim of the study</b>  To compare the effect on neonatal outcomes of admission CTG versus intermittent auscultation of the fetal heart rate</p> <p><b>Study dates</b></p>	<p>See entry in systematic review by Devane 2012</p> <p><b>Characteristics</b></p> <p>The following relate to the whole study population, not the low risk subgroup from the systematic review.</p> <p><b>Induction of labour (n/total (%))</b> Cardiotocograph (CTG): 765/4298 (18) Auscultation: 749/4282 (17)</p> <p><b>Major congenital anomaly (n/total (%))</b> CTG: 27/4298 (1) Auscultation: 18/4282 (&lt;1)</p> <p><b>Parity (n/total (%))</b> - 0 CTG: 2093/4298 (49) Auscultation: 2077/4282 (49) - 1 to 3 CTG: 2121/4298 (49) Auscultation:</p>	Intermittent auscultation	<p>In the intermittent auscultation group, auscultation was performed for 1 minute after a contraction, every 15 minutes in the first stage of labour and every 5 minutes in the second stage. EFM was used only if any of the following occurred: a deceleration in fetal heart rate or persistent tachycardia on auscultation; meconium in liquor or heavily blood stained liquor; maternal temperature of 38 degrees or higher; labour lasting longer than 8 hours.</p> <p>In the CTG group, the CTG was reviewed by the admitting midwife after 20 minutes. If the baseline FHR was 110-160 bpm, variability</p>	<p>reported in the systematic review (see Devane 2012)</p>	<p>There is indirectness of population due to the proportion of women who had induction of labour</p> <p><b>Other information</b></p> <p>All women appear to have had an early amniotomy.</p> <p>MOST STUDY DETAILS ARE REPORTED IN DEVANE 2012. THIS ENTRY ONLY REPORTS EXTRA DETAILS THAT WERE NOT REPORTED IN THE COCHRANE REVIEW, WHICH THE TECHNICAL TEAM FELT WERE IMPORTANT CONSIDERATIONS WHEN INTERPRETING THE RESULTS</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>August 1997 to April 2001</p> <p><b>Source of funding</b></p> <p>Research Committee of the National Maternity Hospital, Dublin</p>	<p>2115/4282 (49)</p> <p>- <math>\geq</math> 4</p> <p>CTG: 81/4298 (2)</p> <p>Auscultation: 90/4282 (2)</p> <p><b>Inclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p> <p><b>Exclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p>		<p>was visually assessed as more than 5 per minutes, decelerations were absent, and if there was more than one acceleration, it was classified as normal. Subsequent care was then the same as the intermittent auscultation group. If the criteria for normal were not met, CTG was continued until birth; 58% of the CTG arm and 42% of the auscultation arm had continuous EFM during labour (this is reported as an outcome in the systematic review)</p>		
<p><b>Full citation</b></p> <p>Mires,G., Williams,F.,</p>	<p><b>Sample size</b></p>	<p><b>Interventions</b></p> <p>Admission CTG</p>	<p><b>Details</b></p> <p>The reasons for which women</p>	<p><b>Results</b></p> <p><b><u>Metabolic acidosis at birth</u></b></p>	<p><b>Limitations</b></p> <p>For the outcome of metabolic acidosis, 1003/3751 (26.7%) of the whole study population, corresponding to 641/2367 (27.1%) of the low-risk</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Howie, P., Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk obstetric population, BMJ, 322, 1457-1460, 2001</p> <p><b>Ref Id</b> 97907</p> <p><b>Country/ies where the study was carried out</b>  Scotland</p> <p><b>Study type</b>  Randomised controlled trial</p> <p><b>Aim of the study</b>  To compare the effect of admission CTG and Doppler auscultation of the fetal heart on neonatal outcome and level of obstetric</p>	<p>See entry in systematic review by Devane 2012</p> <p><b>Characteristics</b></p> <p><b><u>Women having artificial rupture of membranes (n/total)</u></b></p> <p><u>a. All women</u> Cardiotocograph (CTG): 1065/1864 Auscultation: 1031/1879</p> <p><u>b. Low-risk women</u> CTG: 640/1185 Auscultation: 614/1175</p> <p>Proportion of nulliparous and multiparous women in the trial was not reported</p> <p><b>Inclusion criteria</b></p> <p>See entry in systematic review by Devane 2012</p> <p><b>Exclusion criteria</b></p>	<p>Intermittent auscultation with Doppler</p>	<p>were excluded from the 'low-risk' subgroup analysis are listed here. Some women could have had more than one reason (n (%)):</p> <ul style="list-style-type: none"> <li>- Antepartum haemorrhage: 159 (4.2)</li> <li>- Raised blood pressure: 271 (7.2)</li> <li>- Suspected small for gestational age: 56 (1.5)</li> <li>- Preterm labour: 48 (1.30)</li> <li>- Gestational diabetes: 2 (0.1)</li> <li>- Fetal anomaly: 2 (0.1)</li> <li>- Reduced fetal movements and suspected fetal compromise: 63 (1.7)</li> <li>- Meconium stained liquor: 99 (2.6)</li> <li>- Intrauterine death: 3 (0.1)</li> <li>- Persistent breech: 67 (1.8)</li> <li>- Membranes ruptured before labour: 164 (4.4)</li> <li>- Induction of</li> </ul>	<p><b><u>(defined as umbilical cord pH &lt; 7.20 with a base deficit of &gt; 8.0 mmol/l)</u></b></p> <p><u>a. All women</u> CTG: 252/1370 Auscultation: 262/1378</p> <p><u>b. Low-risk women</u> CTG: 159/876 Auscultation: 154/860</p>	<p>women, had no outcome data available.</p> <p>Power calculation and sample size estimate were changed as the trial went along, once after the interim analysis and once following an audit of the data available.</p> <p>A significantly higher proportion of women randomised to CTG had an abnormal FHR pattern at the start of labour, when compared to women randomised to auscultation.</p> <p>Part of the reason that the original trial needed to be accessed was to establish the trial protocol for monitoring in labour. No details were reported beyond those reported in the Cochrane review, therefore it cannot be established whether the admission CTG compared with intermittent auscultation on admission was the only way in which monitoring during labour differed. The following data for the number of women receiving continuous monitoring in labour were reported:</p> <p><b><u>Continuous fetal heart rate monitoring in labour (n/total (%))</u></b></p> <p><u>a. All women</u> CTG: 1246/1865 (66.8) Auscultation: 1128/1882 (59.9)</p> <p><u>b. Low-risk women</u> CTG: 672/1186 (56.7) Auscultation: 551/1178 (46.8)</p> <p><b>Other information</b></p> <p>MOST STUDY DETAILS ARE REPORTED IN DEVANE 2012. THIS ENTRY ONLY REPORTS EXTRA DETAILS THAT WERE NOT REPORTED IN THE COCHRANE REVIEW, WHICH THE TECHNICAL TEAM FELT WERE IMPORTANT CONSIDERATIONS WHEN INTERPRETING THE RESULTS</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>intervention in a low-risk obstetric population</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Chief Scientists Office of the Scottish Executive</p>	<p>See entry in systematic review by Devane 2012</p>		<p>labour: 833 (22.2)</p> <ul style="list-style-type: none"> <li>- Baby born before arrival at hospital: 19 (0.5)</li> <li>- Elective caesarean section: 61 (1.6)</li> <li>- Woman withdrew from trial: 31 (0.8)</li> <li>- Other: 44 (1.2)</li> </ul> <p>Total: 1384 (36.9)</p> <p>In the confirmed low-risk women, 21.5% of those randomised to CTG were considered to have an abnormal fetal heart trace at the onset of labour, compared with 3.6% in the Doppler group (p &lt; 0.0001)</p>		

## G.2 Intermittent auscultation compared with cardiotocography during labour

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Grant,A., O'Brien,N., Joy,M.T., Hennessy,E., MacDonald,D., Cerebral palsy among children born during the Dublin randomised trial of intrapartum monitoring, Lancet, 2, 1233-1236, 1989</p> <p><b>Ref Id</b></p> <p>164086</p> <p><b>Country/ies where the study was carried out</b></p> <p>Ireland</p> <p><b>Study type</b></p> <p>Randomised controlled trial</p> <p><b>Aim of the study</b></p> <p>To confirm that the absence of neonatal signs (such as seizures) suggestive of intrapartum asphyxia is strong evidence that asphyxia was not the cause of later cerebral palsy</p>	<p><b>Sample size</b></p> <p>N = 13079</p> <p>(number of live-born babies during the trial)</p> <p><b>Characteristics</b></p> <p>See entry of MacDonald 1985 for details</p> <p><b>Inclusion criteria</b></p> <p>See entry of MacDonald 1985 for details</p> <p><b>Exclusion criteria</b></p> <p>See entry of MacDonald 1985 for details</p>	<p><b>Interventions</b></p> <p>Intermittent auscultation (n = 6552 babies)</p> <p>Electronic fetal monitoring (EFM) (n = 6527 babies)</p>	<p><b>Details</b></p> <p>All 30 children from the original trial who survived following neonatal seizures and 125 (91%) of a further 138 children whose neurological status was judged to be abnormal, were considered. They underwent a general physical and detailed neurological examination by an experienced paediatrician who was blind to both the monitoring method and the nature of the neonatal neurological abnormality.</p> <p>In order to identify other cases, not originally identified as having abnormal neurological signs, data were sought from specialist remedial clinics in Ireland. Once a child was identified, information about the pregnancy, labour, delivery and neonatal period was extracted from the hospital case-record or trial data sheet. Then the children were divided based on allocation</p>	<p><b>Results</b></p> <p><b>Cerebral palsy (n/total)</b> Auscultation: 10/6552 (0.15) EFM: 12/6527 (0.18)</p> <p><b>Details of the cases</b> Note: - Auscultation group 3 were from the 21 babies with seizures that survived during the neonatal period 7 were identified via clinic notification</p> <p>- EFM group 4 were from the 9 babies with seizures that survived during the neonatal period 8 were identified via clinic notification</p> <p><u>a. Children with abnormal neurological signs during neonatal period</u> 30 of the 39 babies with neonatal seizures survived to be discharged from hospital; 3 from each group were then judged to have cerebral palsy at 4 years old.</p> <p>4 children (2 in each arm) had 'spastic quadriplegia with severe mental retardation'. There had been signs</p>	<p><b>Limitations</b></p> <p>Appropriate randomisation: Yes Allocation concealment: Yes Groups comparable at baseline: Yes Groups received same care (apart from intervention): Yes Blinding of participants: No Blinding of staff providing care: No Blinding of outcome assessors: Yes Missing data/loss to follow-up: Possible because apart from those babies with seizures/other symptoms after birth, other children were identified through specialist clinics in Ireland. This would not have covered any children who had moved away or possibly died Precise definition of outcomes: Yes Valid and reliable method of outcome assessment: Yes Intention-to-treat analysis performed: Yes</p> <p>Indirectness: in the original trial 22.5% of women were considered 'high risk'</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>To estimate the proportion of all cases of cerebral palsy that might possibly be associated with intrapartum asphyxia</p> <p><b>Study dates</b></p> <p>Recruitment into the original trial began on March 31st 1981 and ended on April 10th 1983</p> <p>Follow-up was at age 4 years</p> <p><b>Source of funding</b></p> <p>See entry on MacDonald 1985 for details of the trial</p>				<p>suggestive of asphyxia in 3 which were apparent both during labour and after the birth. The fourth child was born at 34 weeks' gestation with a 5-minute Apgar score of 8, then had severe respiratory distress syndrome following intraventricular haemorrhage and then post haemorrhage hydrocephalus.</p> <p>The other 2 children had mild spastic hemiplegias, and had a sequence of signs suggestive of asphyxia during labour and after birth.</p> <p>A seventh child with mild spastic hemiplegia was identified from among the 125 children who were formally reassessed because of neonatal neurologic abnormalities other than seizures. There had been transient abnormalities of tone, reflexes and behaviour, but they had resolved within 48 hours of birth.</p> <p><u>b. Identified from clinics</u> In 12 of the 15 cases (of which one was a twin), labour delivery and the neonatal period seemed normal. Of the 3 others, 1 (allocated EFM) had respiratory distress syndrome and pneumonia following spontaneous rupture of the</p>	<p><b>Other information</b></p> <p>This is a follow-up to MacDonald 1985</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				membranes and birth at 30 weeks. One (allocated auscultation) had an emergency caesarean section (CS) because of failed induction at 43 weeks and suspected intrauterine infection. The third (allocated auscultation) was discharged apparently well but later had severe gastroenteritis that had been complicated by cerebral oedema with seizures and later meningitis.	
<p><b>Full citation</b></p> <p>Kelso,I.M., Parsons,R.J., Lawrence,G.F., Arora,S.S., Edmonds,D.K., Cooke,I.D., An assessment of continuous fetal heart rate monitoring in labor. A randomized trial, American Journal of Obstetrics and Gynecology, 131, 526-532, 1978</p> <p><b>Ref Id</b></p> <p>164097</p> <p><b>Country/ies where the study was carried out</b></p> <p>England</p>	<p><b>Sample size</b></p> <p>N = 504</p> <p><b>Characteristics</b></p> <p><b><u>Maternal age/years (mean ± SD)</u></b></p> <p>Auscultation: 25.6 ± 5.0 EFM: 26.0 ± 4.9 (NS)</p> <p><b><u>Gestation/weeks (mean ± SD)</u></b></p> <p>Auscultation: 39.75 ± 1.18 EFM: 39.67 ± 1.32 (NS)</p> <p><b><u>Nulliparous (n/total)</u></b></p> <p>Auscultation: 134/251 EFM: 116/253</p> <p><b><u>Cervical assessment using Bishop score (n/total)</u></b></p>	<p><b>Interventions</b></p> <p>Auscultation (n = 251)</p> <p>EFM (n = 253)</p>	<p><b>Details</b></p> <p>All women under the care of the University Department at the Jessop Hospital for Women, Sheffield, admitted to the labour ward during the study period had their labours analysed. Women were admitted in spontaneous labour or to have labour induced. The study authors wanted to evaluate a non high-risk population; therefore, the exclusion criteria aimed to exclude high-risk women. All other women were allotted a sealed envelope when they were admitted, containing treatment allocation.</p> <p>Women allocated to continuous monitoring had a fetal scalp electrode attached, with or without an intrauterine pressure</p>	<p><b>Results</b></p> <p><b><u>Mode of birth (n/total)</u></b></p> <p><b><u>a. Spontaneous vaginal birth</u></b> Auscultation: 162/251 EFM: 158/253</p> <p><b><u>b. Forceps or ventouse birth</u></b> Auscultation: 78/251 EFM: 71/253</p> <p><b><u>c. Caesarean section</u></b> Auscultation: 11/251 (3 for fetal distress) EFM: 24/253 (4 for fetal distress)</p> <p><b><u>Perinatal death (n/total)</u></b> Auscultation: 1/251 EFM: 0/253 (Note: the woman was multiparous and admitted at 41 weeks' gestation in spontaneous labour. The labour</p>	<p><b>Limitations</b></p> <p>Appropriate randomisation: Unclear - method of randomisation is not reported Allocation concealment: Yes Groups comparable at baseline: Yes; however, there was a significantly shorter first and second stage of labour in the EFM arm Groups received same care (apart from intervention): Monitoring was internal; therefore, in order to fit the scalp electrode, women in the EFM arm were likely to have received an amniotomy to fit the electrode in cases where the membranes had not ruptured; this would not be necessary in the other arm of the trial. Blinding of participants: Not</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study type</b></p> <p>Randomised controlled trial</p> <p><b>Aim of the study</b></p> <p>To compare the usefulness of continuous fetal heart rate monitoring in labour using the dip area as a measure of fetal distress with or without intrauterine pressure recordings</p> <p><b>Study dates</b></p> <p>July 1976 to June 1977</p> <p><b>Source of funding</b></p> <p>The first author received a British Commonwealth Medical Fellowship. Financial assistance was also gained from Pye Dynamics, Ltd and Devices, Ltd</p>	<p>1 - 4 Auscultation: 43/251 EFM: 38/253</p> <p>5 - 8 Auscultation: 154/251 EFM: 151/253</p> <p>9 - 12 Auscultation: 54/251 EFM: 64/253 (NS)</p> <p><b><u>Type of labour (n/total)</u></b></p> <p>- Spontaneous Auscultation: 120/251 EFM: 132/253</p> <p>- Accelerated Auscultation: 69/251 EFM: 51/253</p> <p>- Induced Auscultation: 62/251 EFM: 70/253 (NS)</p> <p><b><u>Intra or postpartum pyrexia (n/total)</u></b></p> <p>Auscultation: 7/251 EFM: 8/253 (NS)</p> <p><b><u>Birth weight / grams (mean <math>\pm</math> SD)</u></b></p> <p>Auscultation: 3349 <math>\pm</math> 430 EFM: 3335 <math>\pm</math> 459</p>		<p>catheter, at the earliest convenient time. Oxytocin was given to all women when indicated.</p> <p>In women allocated to intermittent auscultation, the fetal heart rate (FHR) was counted every 15 minutes (or more frequently if indicated) during or immediately after a contraction. A Pinard fetal stethoscope was used, and the rate was counted for 1 full minute. If there was any difficulty hearing the sounds, an ultrasonic Doppler was used intermittently.</p> <p>A double-clamped section of the cord was collected at birth before the baby's first breath. Arterial and venous blood gas measurements were taken.</p> <p>Augmentation, using amniotomy alone or amniotomy with oxytocin infusion, was performed if the progress of the labour fell to the right of the nomogram. Decisions to perform caesarean section or instrumental birth were the responsibility of duty staff.</p> <p><b><u>Outcomes reported:</u></b></p> <p>1. Mode of birth: rate of spontaneous birth, forceps or ventouse, and caesarean section were reported</p>	<p>was slow despite an oxytocin infusion, and there were at least two separate episodes of fetal tachycardia [170 - 190 bpm]. After 12 hours and 45 minutes, meconium stained liquor was noted. The FHR was 190 bpm and the cervix was dilated. Forceps were applied to rotate the vertex. After birth, the baby was transferred to SCBU and intubated. The baby died of meconium aspiration at 4 hours)</p> <p><b><u>Abnormal neurologic signs (n/total)</u></b></p> <p>Auscultation: 3/251 EFM: 2/253 (Note: All 5 babies had depressed Apgar scores and were admitted to SCBU. In the EFM group: both babies were hypertonic at birth, but there were no symptoms at day 9 or week 6. in the auscultation group: the first baby was jittery and irritable for 3 days, but there were no abnormal neurological findings on day 6 or week 6. The second baby had a cyanotic attack and a left-sided convulsion at 6 hours after the birth. The baby was treated with phenobarbitone for 3 days, and there were no further convulsions, and no issues at day 12 or week 6. The third baby was 'stiff and irritable' at 11 hours and received phenobarbitone for 3 days, after</p>	<p>reported</p> <p>Blinding of staff providing care: Not reported</p> <p>Blinding of outcome assessors: Not reported</p> <p>Missing data/loss to follow-up: No</p> <p>Precise definition of outcomes: Yes</p> <p>Valid and reliable method of outcome assessment: Yes</p> <p>Intention-to-treat analysis performed: Yes</p> <p>Indirectness: 26% of women had induction of labour</p> <p><b>Other information</b></p> <p>CTG: internal</p> <p>2 other perinatal deaths were detailed in the article, but they were born to women excluded from the trial due to breech presentation.</p> <p><b><u>Length of labour (mean <math>\pm</math> SD)</u></b></p> <p><b><u>a. First stage / hours</u></b> Auscultation: 6.63 <math>\pm</math> 3.88 EFM: 5.94 <math>\pm</math> 3.36 (p &lt; 0.05)</p> <p><b><u>b. Second stage / minutes</u></b> Auscultation: 32.35 <math>\pm</math> 25.23 EFM: 28.01 <math>\pm</math> 21.00 (p &lt; 0.05)</p> <p><b><u>c. Third stage / minutes</u></b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><b>Inclusion criteria</b></p> <p>Admitted to the labour ward during the study period</p> <p><b>Exclusion criteria</b></p> <p>Breech presentation</p> <p>Multiple pregnancy</p> <p>Maternal age of 40 years or greater</p> <p>Previously mentally disabled or spastic child resulting from birth</p> <p>Previous perinatal death - cause unknown</p> <p>Previous severe fetal distress - Apgar score of 3 or less</p> <p>Hypertension with diastolic pressure 100 mmHg or 100 mmHg with proteinuria</p> <p>Two consecutive estrogen estimations outside 2 SD from the normal</p> <p>Anaemia of 8 g/dl or less</p> <p>Type 1 diabetes</p> <p>Admitted fully dilated and ready for birth</p>		<p>2. Perinatal death</p> <p>3. Admission to special care baby unit (SCBU)</p> <p>4. Abnormal neurological signs</p>	<p>which time there were no abnormal neurologic findings)</p> <p><b>Admission to SCBU (n/total)</b>                      Auscultation: 43/251                      EFM: 45/253</p> <p>Note: the indications for admission were as follows (n):                      -- infant depressed at birth                      Auscultation: 12                      EFM: 9                      -- birthweight less than 2500 g or considered preterm by attending paediatrician                      Auscultation: 7                      EFM: 6                      -- jaundiced - admitted for phototherapy                      Auscultation: 10                      EFM: 16                      -- treated maternal thyrotoxicosis euthyroid at time of labour                      Auscultation: 4                      EFM: 0                      -- maternal thrombocytopenia                      Auscultation: 1                      EFM: 0                      -- maternal pyrexia &gt; 38 degrees                      Auscultation: 1                      EFM: 0                      -- meconium aspiration                      Auscultation: 3                      EFM: 2                      -- congenital anomalies                      Auscultation: 1                      EFM: 2                      -- hypothermia</p>	<p>Auscultation: 6.66 ± 10.32                      EFM: 6.19 ± 8.13                      (NS)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	Missed			<p>Auscultation: 1 EFM: 4 -- other Auscultation: 3 EFM: 6</p> <p><b><u>Cord blood gas values</u></b> The authors reported that cord arterial and venous blood gas analysis was performed for 37 babies in each arm. There were no statistically significant differences in the proportion of babies with pH of 7.25 or less, or base deficit of 10 mmol/l or more. No further details were reported; therefore, this is not reported in the GRADE table.</p>	
<p><b>Full citation</b></p> <p>Leveno,K.J., Cunningham,F.G., Nelson,S., Roark,M., Williams,M.L., Guzick,D., Dowling,S., Rosenfeld,C.R., Buckley,A., A prospective comparison of selective and universal electronic fetal monitoring in 34,995 pregnancies, New England Journal of Medicine,N Engl J Med, 315, 615-619, 1986</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>N = 34,995  (However, the population of interest for this review is 14,618)</p> <p><b>Characteristics</b></p> <p>The following represent characteristics of the entire study population. Details of the low-risk subgroup are not reported separately.</p> <p><b>Parity (%)</b> - Nulliparous Selective: 39</p>	<p><b>Interventions</b></p> <p>Selective monitoring: intermittent auscultation for low-risk women and EFM for high-risk women (n = 7330)</p> <p>Universal monitoring: all women monitored with EFM (n = 7288)</p>	<p><b>Details</b></p> <p>This was a trial comparing the policy of all women being monitored using EFM (universal monitoring) with a policy of only monitoring high-risk women with EFM (selective monitoring). The trial employed these different policies during alternating months, and compared the results.</p> <p>The standard policy in the unit (Parkland Memorial Hospital) was a policy of only using EFM in high risk pregnancies (see details listed in inclusion criteria above). Women who had</p>	<p><b>Results</b></p> <p><b><u>Caesarean section for fetal distress (n/total (%))</u></b> Selective/auscultation: 28/7330 (0.4) Universal/EFM: 64/7288 (0.9) (p &lt; 0.01)</p> <p><b><u>Mortality (n/total (%))</u></b> <b>a. Intrapartum fetal death</b> Selective/auscultation: 0/7330 (0) Universal/EFM: 0/7288 (0) (NS)</p> <p><b>b. Neonatal death</b> Selective/auscultation: 5/7330 (0.1)</p>	<p><b>Limitations</b></p> <p>Appropriate randomisation: No - low risk women received auscultation or EFM on alternating months Allocation concealment: No Groups comparable at baseline: Unclear - there were no significant differences in the selective versus universal groups, but this detail was not reported for low-risk women Groups received same care (apart from intervention): Yes Blinding of participants: Unclear, but unlikely considering the intervention Blinding of staff providing care:</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>164091</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Quasi-randomised trial</p> <p><b>Aim of the study</b></p> <p>To compare the differences in perinatal outcome between universal and selective electronic fetal monitoring (EFM) in 34,995 births</p> <p><b>Study dates</b></p> <p>October 1st 1982 onwards, for a 36-month period</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p>Universal: 40</p> <p>- Multiparous Selective: 61 Universal: 60</p> <p><b><u>Prenatal care (%)</u></b> Selective: 81 Universal: 82</p> <p><b><u>Birth weight / g (%)</u></b></p> <p>- 500-999 Selective: 0.8 Universal: 0.8</p> <p>- 1000-1500 Selective: 1.2 Universal: 1.1</p> <p>- 1501-2000 Selective: 2.3 Universal: 2.5</p> <p>- 2001-2500 Selective: 7.2 Universal: 7.2</p> <p>- ≥ 2501 Selective: 88.5 Universal: 88.4</p> <p>There were no significant differences identified between the two groups</p> <p><b>Inclusion criteria</b></p> <p>Not reported for the study; however, the following</p>		<p>complications were transferred into a labour intensive unit with 5 beds (this continued throughout both parts of the trial). Most electronic monitoring was done in this unit. A maximum of seven portable electronic monitors were available during selective monitoring months.</p> <p>During universal monitoring months, 12 additional monitors were made available and installed in labour rooms. Therefore, a total of 19 monitors were available for a 20-bed unit. The policy during these months was to use EFM for every pregnancy in which the baby was viable.</p> <p>Other than the policy of selective or universal monitoring, there were no differences in care during the alternate months. Nursing personnel were in a ratio of 2 women to one nurse. Oxytocin was administered according to a strict protocol. Women admitted to single-bed labour rooms were visited every 30 minutes, and had the fetal heart rate measured using intermittent auscultation with a Doppler device or visual inspection of the trace.</p> <p>Nurses attending each birth</p>	<p>Universal/EFM: 4/7288 (0.1) (NS)</p> <p><b><u>Admission to intensive care nursery (n/total (%))</u></b> Selective/auscultation: 17/7330 (0.2) Universal/EFM: 25/7228 (0.3) (NS)</p> <p><b><u>Neonates with seizures (n/total (%))</u></b> Selective/auscultation: 3/7330 (0.4) Universal/EFM: 1/7288 (0.01) (NS)</p> <p>Note: non-significant p-values were not reported</p>	<p>No</p> <p>Blinding of outcome assessors: Unclear - no details were reported</p> <p>Missing data/loss to follow-up: Unclear</p> <p>Precise definition of outcomes: Yes</p> <p>Valid and reliable method of outcome assessment: Unclear at what point seizures were assessed and the reasons for admission to NICU</p> <p>Intention-to-treat analysis performed: Unclear</p> <p>Overall, this study is not well reported for the guideline comparison and population of interest. The data for low-risk women were reported for the comparison of selective versus universal monitoring, and therefore, the technical team made the assumption that this represents auscultation versus EFM, because according to the trial protocol, in 'selective' months low-risk women should all have received auscultation and in 'universal' months they should have received EFM. This assumption is corroborated by the assumption of a Cochrane review (Alfirevic 2013) who reported this trial for the same comparison</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>definitions are used to describe the different parts of the study population:</p> <p>High risk:</p> <ul style="list-style-type: none"> <li>- induction or augmentation of labour</li> <li>- dysfunctional labour (not defined)</li> <li>- abnormal fetal heart rate</li> <li>- presence of meconium in the amniotic fluid</li> <li>- other complications of pregnancy, including hypertension, vaginal bleeding, prolonged pregnancy, diabetes, twins, breech presentation and preterm labour</li> </ul> <p>Low risk:</p> <ul style="list-style-type: none"> <li>- single baby</li> <li>- cephalic presentation</li> <li>- spontaneous, uncomplicated labour</li> <li>- birth weight exceeding 2500 g</li> </ul> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>completed a perinatal data sheet, and research nurses assessed the data for consistency and completeness before it was stored electronically. Statistical analysis was done using chi-squared test or Fisher's exact test. Two sided p-values of 0.05 were considered significant</p>		<p><b>Other information</b></p> <p>Cardiotocograph (CTG): not reported whether monitoring was internal or external. Abnormal fetal heart rates were identified in 2.7% of selective/auscultation women and 7.6% of universal/EFM women (low risk). The difference was statistically significantly (<math>p &lt; 0.01</math>)</p>
<p><b>Full citation</b></p> <p>MacDonald,D., Grant,A., Sheridan-Pereira,M., Boylan,P., Chalmers,I., The Dublin randomized</p>	<p><b>Sample size</b></p> <p>N = 12,964</p>	<p><b>Interventions</b></p> <p>Intermittent auscultation (n = 6490)</p>	<p><b>Details</b></p> <p><u>Sample size calculation</u> A sample size calculation was based on adverse outcomes for babies, and the anticipated</p>	<p><b>Results</b></p> <p><u>Mode of birth and primary indication (n (%))</u> a. Caesarean section Auscultation: 144 (2.2)</p>	<p><b>Limitations</b></p> <p>Appropriate randomisation: Yes Allocation concealment: Yes Groups comparable at</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>controlled trial of intrapartum fetal heart rate monitoring, American Journal of Obstetrics and Gynecology, 152, 524-539, 1985</p> <p><b>Ref Id</b> 164093</p> <p><b>Country/ies where the study was carried out</b> Ireland</p> <p><b>Study type</b> Randomised controlled trial</p> <p><b>Aim of the study</b> To compare continuous electronic intrapartum fetal heart monitoring with a policy of intermittent auscultation</p> <p><b>Study dates</b> March 31st 1981 to April 10th 1983</p> <p><b>Source of funding</b></p>	<p><b>Characteristics</b></p> <p><b><u>Nulliparous n (%)</u></b> Auscultation: 1964 (39.3) Electronic fetal monitoring (EFM): 2015 (40.4)</p> <p><b><u>Receiving induction of labour (n (%))</u></b> Auscultation: 475 (9.5) EFM: 434 (8.7)</p> <p><b><u>Giving birth earlier than 37 weeks' gestation (n (%))</u></b> Auscultation: 133 (2.7) EFM: 156 (3.1)</p> <p><b><u>Considered high risk at the start of labour (n (%))</u></b> Auscultation: 1137 (22.7) EFM: 1106 (22.2)</p> <p>(Note: this was defined as maternal age of 40 years or more, diabetes, pre-eclampsia, chronic hypertension, renal disease, cardiac disease, previous stillbirth or neonatal death, previous child with neurological abnormality, previous low birthweight baby, bleeding in pregnancy requiring admission to hospital after the first trimester, induction of labour for pregnancy of more than 42 completed weeks' gestation, multiple pregnancy, breech</p>	EFM (n = 6474)	<p>population of 10,000 had 80% power to detect a statistically significant difference if the rate was reduced by half through more intensive monitoring. An interim analysis, after 4,000 cases, determined that recruitment should be extended to 13,000 to assess the difference on the most unambiguous set of outcomes (deaths and seizures). This would have 75% power to detect a 50% reduction. For practical reasons, data on umbilical venous acid-base status were limited to 1000 consecutive babies. The trial protocol pre-specified stratification by risk status and by time interval between entry to trial and birth (&lt; 1 hour, &gt; 1 hour).</p> <p><b><u>Study population</u></b> During the study period, 17381 women gave birth. 4356 were ineligible due to having an elective caesarean section (CS), suffering a fetal death before labour, delivering so rapidly after arrival (&lt; 1 hour from admission) that presence of meconium stained liquor and hence eligibility could not be assessed, less than 28 weeks, gross fetal abnormality, or meconium staining or no fluid. Out of the remaining 13,025 women eligible for inclusion,</p>	<p>- Failure to progress in labour: 88 (1.3) - Fetal distress: 10 (0.2) - Other: 46 (0.7)</p> <p>EFM: 158 (2.4) - Failure to progress in labour: 84 (1.3) - Fetal distress: 25 (0.4) - Other: 49 (0.7)</p> <p><b><u>b. Forceps birth</u></b> Auscultation: 407 (6.3) - Failure to advance: 313 (4.8) - Fetal distress: 75 (1.2) - Other: 19 (0.3)</p> <p>EFM: 528 (8.2) - Failure to advance: 323 (5.0) - Fetal distress: 190 (2.9) - Other: 15 (0.2)</p> <p><b><u>Admission to SCN (n/total (%))</u></b> Auscultation: 543/6554 (8.3) EFM: 547/6530 (8.4)</p> <p>(Note: in an analysis based only on the first 10,000 women recruited, it was reported that 2.7% of babies were admitted for reasons that might have been affected by intrapartum care.)</p> <p><b><u>Umbilical cord venous pH (n/total (%))</u></b> &lt; 7.05 Auscultation: 2/535 (0.4) EFM: 2/540 (0.4)</p>	<p>baseline: Yes Groups received same care (apart from intervention): Yes (because clear liquor had to be demonstrated to enter the trial; therefore, extra amniotomy was not required for EFM arm) Blinding of participants: No Blinding of staff providing care: No Blinding of outcome assessors: Yes for neonatal outcomes Missing data/loss to follow-up: For cord blood gas values, there were limited data; for other outcomes, more detail was collected in the first part of the trial than in the second (i.e the last 3,000 women) i.e. for 'other neurological abnormality' data were only collected for 10,094/13,084 (77%) of study babies. Precise definition of outcomes: Yes Valid and reliable method of outcome assessment: Yes Intention-to-treat analysis performed: Yes</p> <p>Indirectness: 22.5% of women were considered 'high risk'</p> <p><b>Other information</b></p> <p>CTG: monitoring was internal</p> <p>Rates of successful fetal blood sampling were 3.5% in the</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Medical Research Council of Ireland</p> <p>National Maternity Hospital Research Fund</p> <p>Wellcome Trust</p> <p>Department of Health and Social Security (supported the National Perinatal Epidemiology Unit [NPEU])</p>	<p>presentation in labour, and gestational age less than 34 completed weeks.)</p> <p><b>Inclusion criteria</b></p> <p>Live fetus of at least 28 weeks' gestation with no evidence of gross abnormality</p> <p>Diagnosis of labour made</p> <p>Amniotic fluid without significant meconium staining had been positively demonstrated, either at spontaneous rupture of membranes or early amniotomy</p> <p><b>Exclusion criteria</b></p> <p>Elective caesarean section</p> <p>Fetal death prior to the onset of labour</p>		<p>12,964 were entered into the trial and gave birth to 13,084 babies.</p> <p><b>Randomisation</b></p> <p>Randomisation was performed after eligibility had been confirmed through assessment of liquor. Allocation was done by opening the next in a series of serially numbered, sealed opaque envelopes.</p> <p><b>Monitoring in EFM arm</b></p> <p>Following randomisation, an electrode was applied to the fetal scalp and an external tocodynamometer was attached. If it was not possible to get a signal from the electrode, an external transducer was used. If the midwife was concerned about the trace, they first checked it using auscultation and then informed the nurse-midwife in charge of the labour ward. If the latter considered the trace to be abnormal, an obstetrician was called.</p> <p>The following fetal heart rate (FHR) patterns were considered to be suspicious:</p> <ul style="list-style-type: none"> <li>- marked tachycardia or bradycardia</li> <li>- moderate tachycardia or bradycardia with reduced variability</li> <li>- minimal variability (absent</li> </ul>	<p>7.05-7.09 Auscultation: 9/535 (1.7) EFM: 3/540 (0.6)</p> <p>7.10-7.20 Auscultation: 40/535 (7.5) EFM: 41/540 (7.6)</p> <p>&gt; 7.20 Auscultation: 484/535 (90.4) EFM: 494/540 (91.4)</p> <p><b>Neonatal morbidity (n/total (%))</b></p> <p><b>a. Need for intubation</b> Auscultation: 54/5058 (1.1) EFM: 58/5035 (1.2)</p> <p><b>b. Neonatal seizures (all women)</b> Auscultation: 27/6554 (0.4) EFM: 12/6530 (0.2)</p> <p>(Note: in 10/12 cases in the EFM arm and 24/27 in the auscultation arm, seizures were first noted within 48 hours of birth. In 4 out of the 5 later cases, the cause was unlikely to be due to birth event [meningitis at 28 weeks, 2 cases of complications of hyaline membrane disease, and 1 case of hypoglycemia] and in the fifth, the seizures were first noted at 56 hours of age)</p> <p><b>c. Neonatal seizures (women without pregnancy risk factors)*</b></p>	<p>auscultation group and 4.4% in the EFM group.</p> <p>97.7% of those allocated to auscultation received it throughout labour. In the EFM group, 80.7% received EFM throughout; birth was too rapid in 10.5%, 6.6% refused monitoring, and there were technical problems in 1.1% of cases.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>beat-to-beat variation, flat tracing)                      - late deceleration pattern                      - moderate and severe variable deceleration patterns                      - other confusing patterns with varying baselines which could not be clearly interpreted</p> <p>If any of these patterns had been present for at least 10 minutes and did not respond to measures such as changing position or adjusting transducers, then clinical action was taken. In the first stage of labour this was the taking of fetal scalp blood pH; in the second stage of labour the action was immediate birth.</p> <p>If the fetal scalp blood pH was less than 7.20 birth was actioned as soon as possible. If the pH was 7.20 - 7.25 and the FHR pattern remained suspicious, birth was also completed as soon as possible. If the FHR reverted to a normal pattern, the situation was managed expectantly. If the pH was over 7.25 and the trace stayed suspicious, scalp blood pH was measured 30 minutes to 1 hour later.</p> <p>Throughout the trial, tracings were reviewed by a single experienced observer, who was blinded to the outcome of the</p>	<p>Auscultation: 19/5015 (0.4)                      EFM: 7/5038 (0.1)</p> <p><u>d. Other neurological abnormality</u>                      Auscultation: 25/5058 (0.5)                      EFM: 16/5035 (0.3)</p> <p>(Note: This is abnormalities other than seizures and was only reported in survivors. In the auscultation group, 5 babies had 'simultaneous abnormalities of tone and reflex' and 20 babies had 'other abnormal neurological signs persisting for at least a week'. In the EFM arm, the numbers were 4 and 12 respectively.)</p> <p><u>e. Neonatal trauma</u>                      Auscultation: 66/5058 (1.3)                      EFM: 71/5035 (1.4)</p> <p>(Note: In decreasing order of prevalence: scalp laceration, abrasion or bruising; facial bruising, suffusion, forceps marks and conjunctival haemorrhage; cephalhematoma; other bruising; motor deficit in right arm; fractured clavicle; subdural haemorrhage and death; facial nerve injury)</p> <p>* Data from low risk women are reported in the GRADE table</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>baby following birth. The trace was classified according to whether the observer felt that it should or should not have prompted clinical action.</p> <p><b><u>Monitoring in auscultation arm</u></b>                      Women randomised to receive auscultation were managed according to the hospital's standard policy. The FHR was auscultated with a Pinard stethoscope for 60 seconds following a contraction. This was done at least every 15 minutes in the first stage and during every interval between contractions in the second stage. If there was an issue detecting the FHR with auscultation, intermittent Doppler ultrasound was used.</p> <p>If the FHR was &lt; 100 or &gt; 160 bpm during three contractions, and the abnormality did not respond to measures such as a change in posture or treatment of pyrexia, then clinical action was taken as above; i.e in the first stage of labour scalp pH was taken and a scalp clip attached, and in the second stage of labour, birth was expedited.</p> <p><b><u>Outcomes reported</u></b>                      1. Mode of birth</p>	<p><b><u>Perinatal death (n/total (%))</u></b>  <b><u>a. Total</u></b>                      Auscultation: 14/6554                      EFM: 14/6530</p> <p><b><u>b. Intrapartum stillbirth</u></b>                      Auscultation: 2/6554                      EFM: 3/6530</p> <p><b><u>c. Neonatal deaths</u></b>                      Auscultation: 12/6554                      EFM: 11/6530</p> <p>The following details are given about the primary causes of the deaths (n):                      -- Asphyxial conditions developing in labour                      Auscultation: 7                      EFM: 7                      -- Conditions associated with immaturity                      Auscultation: 4†                      EFM: 1                      -- Birth trauma                      Auscultation: 1                      EFM: 3*                      -- Other                      Auscultation: 2                      EFM: 3</p> <p>† in one of the babies in each of these groups, asphyxial conditions developing during labour may have been contributing factors but were not primary cause of death</p> <p><b><u>Stratified analyses</u></b>  <b><u>a. By risk status</u></b></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>2. Mortality: intrapartum deaths and deaths within 28 days (neonatal deaths) were examined by a pathologist blinded to allocation. Each case was classified by primary cause of death, and in cases where the primary cause was not 'asphyxial conditions developing during labour' they were reviewed to see if the conditions may have contributed</p> <p>3. Neurological abnormalities: Neurological assessments were made by a blinded neonatologist. The babies were considered to have had seizures if the neonatologist felt there was evidence of seizures of the following types: generalised tonic, multifocal clonic, focal clonic, or myoclonic. This did not include babies with 'subtle seizure activity' or 'jitteriness'.</p> <p>- During recruitment of the first 10,000 women, serial standardised assessments were made on all babies admitted to the special care nursery (SCN) and any babies on the ward who staff were concerned about. Any babies identified in these ways were examined within 48 hours of birth, then at 72 hours, at 7 days, and at discharge. Assessment of tone, movement, reflexes and behaviour was performed to classify babies into one of the following categories:</p>	<p>22.5% of women met the criteria for being high risk. Compared to the other participants of the trial, these women were 2.7 times more likely to have a caesarean section, and their babies were more than three times more likely to have an Apgar &lt; 4 at one minute, to be admitted to SCN or to die. Within the risk groups, there was little evidence of a differential effect of the two policies on outcome. In the case of neonatal seizures, the effect of EFM in preventing neonatal seizures was stronger in women without risk factors when compared to women with risk factors. However, the effect of monitoring on neonatal seizures that resulted in survival was not different in the two risk groups.</p> <p><u>Neonatal seizures (rate per 1000)</u></p> <p>- Pregnancy risk factors present Auscultation: 5.2 EFM: 3.4 Risk difference (RD): -1.8 per 1000</p> <p>- Pregnancy risk factors not present Auscultation: 3.8 EFM: 1.4 RD: - 2.4 per 1000</p> <p><u>b. By duration of labour</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>simultaneous abnormalities of both tone and reflexes, other neurological abnormalities persisting 1 week after birth, and other transient abnormalities resolved by 7 days</p> <p>- During recruitment of the last 3,000 women, the identification protocol was simplified and neonatologists only identified babies who had seizures in the neonatal period.</p> <p>4. Admission to special care nursery</p> <p>5. Umbilical cord blood gas values: Collection of blood samples only occurred during a 2-month period of the trial. A 15 cm section of cord was double clamped at birth and 3 ml of venous blood was aspirated anaerobically into a heparinised syringe.</p> <p><b><u>Follow-up and statistical analyses</u></b></p> <p>Babies who survived neonatal seizures or other abnormalities of tone and reflexes were followed up for at least 1 year, and seen by senior paediatricians who were not involved in the trial and were blinded to allocation.</p> <p>Chi-squared tests or t-tests of</p>	<p>The longer labours demonstrated a protective effect of EFM, whereas in the shorter labours, the risk of seizures was similar in the two monitoring arms.</p> <p><u>Neonatal seizures (rate per 1000)</u></p> <p>- Labour &lt; 5 hours Auscultation: 1.8 EFM: 1.6 RD: - 0.2 per 1000</p> <p>- Labour &gt; 5 hours Auscultation: 8.5 EFM: 2.4 RD: - 6.1 per 1000</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			statistical significance were used to compare groups.		
<p><b>Full citation</b></p> <p>Vintzileos,A.M., Antsaklis,A., Varvarigos,I., Papas,C., Sofatzis,I., Montgomery,J.T., A randomized trial of intrapartum electronic fetal heart rate monitoring versus intermittent auscultation, Obstetrics and Gynecology, 81, 899-907, 1993</p> <p><b>Ref Id</b></p> <p>164083</p> <p><b>Country/ies where the study was carried out</b></p> <p>Greece</p> <p><b>Study type</b></p> <p>Randomised controlled trial</p> <p><b>Aim of the study</b></p> <p>To determine whether the use of continuous electronic fetal</p>	<p><b>Sample size</b></p> <p>N = 1428</p> <p><b>Characteristics</b></p> <p><b><u>Maternal age/years (mean <math>\pm</math> SD)</u></b></p> <p>Auscultation: 26.6 <math>\pm</math> 5.1 EFM: 26.2 <math>\pm</math> 5.1 (NS)</p> <p><b><u>Nulliparous (n (%))</u></b></p> <p>Auscultation: 340 (50%) EFM: 408 (54.7%) (NS)</p> <p><b><u>Gestational age distribution/weeks (n (%))</u></b></p> <p>26-37 Auscultation: 57 (8.3) EFM: 48 (6.4) (NS)</p> <p>37-42 Auscultation: 608 (89.1) EFM: 686 (91.9) (NS)</p> <p>&gt; 42 Auscultation: 17 (2.4) EFM: 12 (1.6) (NS)</p> <p><b><u>Antepartum risk factors (n</u></b></p>	<p><b>Interventions</b></p> <p>Electronic fetal monitoring (n = 746)</p> <p>Intermittent auscultation (n = 682)</p>	<p><b>Details</b></p> <p>The study was performed in two university hospitals (total of 3000 births per year across the sites). Prior to the study, standard practice was intermittent auscultation, with only approximately 20% of women receiving continuous EFM. Intensive training sessions were given to all clinical personnel, although most were familiar with the use of EFM before the trial.</p> <p>The sample size calculation was based on showing a 2/3 decrease in perinatal mortality. This was based on background mortality rates and reported prevalence of perinatal asphyxia in the year prior to the study. It was calculated that 2210 patients in total were needed (based on alpha of 0.05 and 80% power).</p> <p>Eligible patients were randomised using a coin toss. Women in both arms had IV access secured after admission and labour in lateral or semi-Fowler position. There was one nurse for each woman in both</p>	<p><b>Results</b></p> <p><b><u>Mode of birth (n (%))</u></b></p> <p><b>a. Spontaneous vaginal</b> Auscultation: 561 (82.2) EFM: 571 (76.5)</p> <p><b>b. Vacuum extraction</b> Auscultation: 58 (8.5) EFM: 101 (13.5)</p> <p><b>c. Low forceps</b> Auscultation: 2 (0.3) EFM: 3 (0.4)</p> <p><b>d. Mid forceps</b> Auscultation: 2 (0.3) EFM: 0 (0)</p> <p><b>e. Caesarean section</b> Auscultation: 59 (8.6) - for fetal distress: 16 - reasons other than suspected fetal distress: 43 EFM: 71 (9.5) - for fetal distress: 40 - reasons other than suspected fetal distress: 31</p> <p><b><u>Admission to NICU (n (%))</u></b></p> <p><b>a. Total</b> Auscultation: 102 (14.9) EFM: 104 (13.9)</p> <p><b>b. Unrelated to prematurity</b></p>	<p><b>Limitations</b></p> <p>The trial was stopped after the third periodic review due to increasing mortality rates.</p> <p>Appropriate randomisation: Yes Allocation concealment: Yes Groups comparable at baseline: Yes. There were significant differences between the two groups in the proportion of women having spontaneous labour (higher in auscultation arm), augmented labour (higher in EFM arm) and induction of labour (higher in EFM arm). The duration of labour was also significantly longer in the EFM arm. However, the authors reported that this should have put the EFM arm at a disadvantage. Groups received same care (apart from intervention): Yes Blinding of participants: No Blinding of staff providing care: No Blinding of outcome assessors: No No for maternal outcomes, yes for neonatal outcomes, unclear for cord blood gas values (but unlikely to cause bias for this outcome, because it is</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>monitoring (EFM) alone during labour is associated with decreased perinatal mortality and morbidity when compared to intermittent auscultation, in a population with a relatively high perinatal mortality rate</p> <p><b>Study dates</b></p> <p>October 1st 1990 to June 30th 1991</p> <p><b>Source of funding</b></p> <p>Advanced Medical Systems provided financial support for the study</p>	<p><b>(%)</b></p> <p>Auscultation: 94 (13.7) EFM: 89 (11.9) (NS) (Note: antepartum risk factors were: hypertension, diabetes, premature rupture of membranes, suspected fetal growth restriction, oligohydramnios and vaginal bleeding)</p> <p><b>Meconium stained liquor (n (%))</b></p> <p>Auscultation: 84 (12.3) EFM: 112 (15) (NS)</p> <p><b>Presentation (n (%))</b></p> <p>- Vertex Auscultation: 670 (98.3) EFM: 733 (98.2) (NS)</p> <p>- Breech Auscultation: 11 (1.6) EFM: 12 (1.6) (NS)</p> <p>- Other Auscultation: 1 (0.1) EFM: 1 (0.1) (NS)</p> <p><b>Labour</b></p> <p>- Spontaneous Auscultation: 374 (54.8) EFM: 238 (31.9) (p = 0.0001)</p>		<p>groups.</p> <p>External fetal monitoring was performed using a tocodynamometer for recording uterine contractions and a Doppler ultrasound to monitor fetal heart rate. External monitoring was continued for as long as satisfactory tracings were obtained. Direct monitoring, by the insertion of a fetal scalp electrode, was indicated if the quality of the trace was not satisfactory. If the EFM trace was satisfactory, the decision to use internal monitoring was left to the managing clinician. The initial FHR trace was assessed at least every 15 minutes during the first stage of labour and every 5 minutes during the second stage.</p> <p>Women assigned to auscultation were monitored using a Doppler ultrasound device. The baseline heart rate was counted between contractions and then auscultated every 15 minutes during the first stage and every 5 minutes during the second stage. The FHR was measured during and immediately after the contraction, for at least 30 seconds afterwards. The auscultation lasted 1 minute. Uterine contraction was</p>	<p>Auscultation: 69/625 (11) EFM: 72/698 (10.3)</p> <p><b><u>Cord arterial pH &lt; 7.10 (n/total (%))</u></b></p> <p>Auscultation: 18/680 (2.6) EFM: 31/739 (4.1)</p> <p><b><u>Neonatal complications (n (%))</u></b></p> <p>a. None Auscultation: 594 (87.1) EFM: 639 (85.6)</p> <p>b. Hypoxic ischaemic encephalopathy Auscultation: 2 (0.3) EFM: 1 (0.1)</p> <p>c. Intraventricular haemorrhage Auscultation: 1 (0.1) EFM: 0 (0)</p> <p>d. Seizures Auscultation: 2 (0.3) EFM: 0 (0)</p> <p><b><u>e. Respiratory distress</u></b></p> <p>Auscultation: 40 (5.8) EFM: 55 (7.3)</p> <p>f. Hypotonia* Auscultation: 3 (0.4) EFM: 3 (0.4)</p> <p>g. Necrotizing enterocolitis* Auscultation: 0 (0) EFM: 2 (0.2)</p> <p>h. Sepsis*</p>	<p>biochemical)</p> <p>Missing data/loss to follow-up: Generally not; 0.6% of women had missing data for cord arterial pH</p> <p>Precise definition of outcomes: Yes</p> <p>Valid and reliable method of outcome assessment: Yes</p> <p>Intention-to-treat analysis performed: Yes</p> <p>Indirectness: This was not a completely low-risk population: 12.8% of women had antepartum risk factors, 7.4% labours were preterm and 12% were induced. (As these conditions are not mutually exclusive, the total proportion was considered low enough not to exclude the study)</p> <p><b>Other information</b></p> <p>CTG: monitoring was external for as long as traces were satisfactory</p> <p><b><u>Duration of labour (mean ± SD)</u></b></p> <p>a. First stage / hours</p> <p>Auscultation: 5.5 ± 3.7 EFM: 6.1 ± 4.3 (p = 0.006)</p> <p>b. Second stage / minutes Auscultation: 26.9 ± 16.9</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>- Augmented* Auscultation: 260 (38.1) EFM: 391 (58.4) (p = 0.0001)</p> <p>- Induced Auscultation: 48 (7) EFM: 117 (15.6)</p> <p>* The higher use of oxytocin for augmentation in the EFM group was related to the longer labours in the EFM arm</p> <p><b>Inclusion criteria</b></p> <p>Singleton living fetus</p> <p>Gestational age of 26 weeks or more</p> <p>Admitted in spontaneous labour or for induction of labour</p> <p><b>Exclusion criteria</b></p> <p>Known fetal congenital or chromosomal abnormalities</p>		<p>evaluated using palpation.</p> <p>In the EFM group, non-reassuring heart rate patterns were defined as:</p> <ul style="list-style-type: none"> <li>- late decelerations unrelated to supine hypotension or regional anaesthesia, which failed to respond to conservative measures</li> <li>- persistent prolonged decelerations of less than 80 beats per minute (bpm) lasting more than 2 minutes</li> <li>- severe variable decelerations (70 bpm or fewer lasting 60 seconds or more)</li> <li>- variable decelerations with a rising baseline and loss of variability</li> <li>- persistent fetal tachycardia (more than 160 bpm) associated with decreased variability (less than 5 bpm)</li> <li>- persistent decreased variability</li> <li>- sinusoidal FHR pattern (three to five cycles per minute, amplitude 5 to 15 bpm)</li> </ul> <p>In the auscultation group, non-reassuring heart rate patterns were defined if one or more of the following was present:</p> <ul style="list-style-type: none"> <li>- FHR during and immediately after a contraction repeatedly below 100 bpm, even if there was recovery to 120-160 before the next contraction (moderate decelerations when FHR was</li> </ul>	<p>Auscultation: 2 (0.3) EFM: 3 (0.4)</p> <p><u>i. Hyperbilirubinemia*</u> Auscultation: 26 (3.8) EFM: 31 (4.1)</p> <p><u>j. Hypoglycemia*</u> Auscultation: 4 (0.6) EFM: 5 (0.6)</p> <p><u>k. Other (including congenital abnormalities)*</u> Auscultation: 2 (0.3) (Note: Congenital heart disease; gastroschisis) EFM: 7 (0.9) (Note: Congenital heart disease (n = 2); cleft lip/palate (n = 1; duodenal atresia (n = 1); no further details given)</p> <p>* reported here as morbidities, as reported in the paper, but not reported in the GRADE table as they are unlikely to be affected by method of intrapartum monitoring</p> <p><b><u>Need for neonatal resuscitation (n (%))</u></b> Auscultation: 65 (9.5) EFM: 63 (8.4)</p> <p><b><u>Death of baby (n (%))</u></b></p> <p><u>a. Intrapartum fetal death</u> Auscultation: 2 (0.3) EFM: 0 (0)</p> <p><u>b. Neonatal death</u></p>	<p>EFM: 29.4 ± 18.6 (p = 0.01)</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>80-99 and severe when it was less than 80)                      - persistent baseline rate (between contractions) of less than 100 bpm                      - persistent baseline rate of more than 160 bpm                      In the presence of non-reassuring patterns, groups were managed similarly. Management was initially conservative, for example, stopping oxytocin, administering maternal oxygen, changing position, or increasing IV fluids. Fetal scalp pH, or crossing patients over from one group to another were not used. If the non-reassuring pattern persisted after 20 minutes of trying conservative methods, a surgical intervention (forceps, vacuum extraction or caesarean section) was performed.</p> <p>A data sheet was completed by the attending physicians which recorded maternal characteristics, and outcomes for the woman and baby. Most neonatal outcomes were collected by neonatologists blinded to allocation. Obstetric records and FHR data from both arms of the trial were reviewed throughout by two authors blinded to monitoring method. This was aimed at determining whether interpretation and management of FHR had been</p>	<p>Auscultation: 7 (1)                      EFM: 2 (0.26)</p> <p><u>c. Total perinatal death†</u>                      Auscultation: 9 (1.3)                      EFM: 2 (0.26)</p> <p>† of these, 6 in the auscultation group and 0 in the EFM group were reported as being due to fetal hypoxia. Note: the 2 deaths in the EFM group could not have been prevented by monitoring: one baby died of complex congenital heart disease and the other of haemorrhage and DIC due to trauma at the base of the tongue during intubation attempt for meconium suctioning; among the 9 deaths in the auscultation group, there was compliance with trial protocol and vaginal delivery in all 9. Details of deaths are reported below</p> <p>Clinical characteristics of the nine perinatal deaths in the auscultation group:                      Intrapartum (n = 2)                      - Both women were at term (39 weeks; 41 weeks)                      - Neither woman had risk factors and both were vertex presentation                      - One had meconium staining                      Neonatal (n = 7)                      - 2 out of 7 were preterm (26.3 weeks; 30 weeks)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>appropriate. If there was delayed or absent intervention after persistent non-reassuring patterns, or surgical intervention in the presence of reassuring patterns, this was recorded as 'failure to comply with protocol'.</p> <p>Data were reviewed every 3 months to detect trends in mortality. The continuing trend of increasing death in the auscultation group was compared with the year before the study, which did not show any peaks, and the study was stopped after the third review.</p> <p>Statistical analysis was done using chi-squared, Fisher's exact test, Student's t tests, ANOVA, and Mann-Whitney tests, where appropriate; <math>p &lt; 0.05</math> was considered significant.</p> <p><b>Outcomes reported</b></p> <ol style="list-style-type: none"> <li>1. Mode of birth: recorded on a data sheet by attending physician</li> <li>2. Admission to NICU: data collected by neonatologists blinded to allocation</li> <li>3. Neonatal morbidity: data collected by neonatologists blinded to allocation on development of complications such as neonatal death, ischaemic encephalopathy,</li> </ol>	<ul style="list-style-type: none"> <li>- Risk factors were present in 6 out of 7 (prematurity [2], PROM [3], gastroschisis [1]) and the presentation of the remaining baby was breech.</li> <li>- 3 had meconium staining</li> <li>- The two premature babies and the case of gastroschisis were considered to be deaths that were not related to hypoxia</li> </ul>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>neurologic abnormalities, seizures, intraventricular haemorrhage, sepsis, necrotising enterocolitis, respiratory distress syndrome (need for supplemental oxygen for over 24 hours), hyperbilirubinemia, hyperglycemia, and metabolic or other problems</p> <p>4. Cord blood gas values: following the birth, the cord was clamped and blood gases were measured from the artery and vein within 10 minutes of birth. Who collected these data was not clearly reported</p>		
<p><b>Full citation</b></p> <p>Wood,C., Renou,P., Oats,J., Farrell,E., Beischer,N., Anderson,I., A controlled trial of fetal heart rate monitoring in a low-risk obstetric population, American Journal of Obstetrics and Gynecology, 141, 527-534, 1981</p> <p><b>Ref Id</b></p> <p>164094</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 989</p> <p><b>Characteristics</b></p> <p>There were no significant differences in maternal age, parity, injections of opiate, use of other drugs, or ketones between the two groups</p> <p><b>Inclusion criteria</b></p> <p>None of the exclusion criteria</p>	<p><b>Interventions</b></p> <p>Standard care (n = 482)</p> <p>Electronic fetal monitoring (n = 507)</p>	<p><b>Details</b></p> <p>Randomisation was by randomised cards. In one of the study sites this did not work effectively because a significantly higher proportion of low parity patients were in the EFM group compared to the auscultation group. Cards were not in sealed envelopes. Parity was corrected by random elimination, leaving 927 of the original 989 patients in the trial. Results were analysed for both 927 and 989 patients, and the results were the same, so the former were reported by the study authors.</p>	<p><b>Results</b></p> <p><b><u>Mode of birth (n/total (%))</u></b></p> <p><b>a. Normal</b> Standard: 371/482 (77.0) EFM: 307/445 (69.0)</p> <p><b>b. Forceps</b> Standard: 101/482 (21.0) EFM: 120/445 (27.0)</p> <p><b>c. Caesarean section</b> Standard: 10/482 (2.1) EFM: 18/445 (4.0)</p> <p><b><u>Neonatal death</u></b> Standard: 0/482 EFM: 1/445</p>	<p><b>Limitations</b></p> <p>Appropriate randomisation: Allocation was by randomised cards Allocation concealment: No, cards were not in sealed envelopes Groups comparable at baseline: This was reported for the denominator of most of the outcomes, but for neurological symptoms/signs, due to issues with randomisation, there may be a difference in the proportion of primigravidas Groups received same care (apart from intervention): Yes (according to study authors)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Australia</p> <p><b>Study type</b></p> <p>Randomised controlled trial</p> <p><b>Aim of the study</b></p> <p>To determine the effects of fetal heart rate monitoring in low-risk women</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p><b>Exclusion criteria</b></p> <p>Past history of stillbirth or neonatal death</p> <p>Antepartum haemorrhage in more than one pregnancy</p> <p>Eclampsia</p> <p>Previous birth before 37 weeks' gestation</p> <p>Clinical signs of fetal distress of meconium stained liquor and fetal heart rate above 160 or below 12 between contractions</p> <p>Medical and obstetric complications of hypertension (145/90 mmHg)</p> <p>Proteinuria (on boiling)</p> <p>Proven renal disease, cyanotic heart disease, rhesus isoimmunisation, diabetes, jaundice of hepatosis, anaemia (Hb 9g/100 ml) at any stage of pregnancy</p> <p>Antepartum haemorrhage</p> <p>Low estriol excretion</p> <p>Polyhydramnios</p> <p>Multiple pregnancy</p>		<p>Control women were managed by staff in the standard way. Women randomised to EFM were managed in a similar way, with the addition of fetal monitoring. Management of labour and birth was the responsibility of the attending medical staff. If complications in labour indicated the need for monitoring among those randomised to standard care, this was performed, but the women remained in the standard care group for the analysis.</p> <p>Following randomisation, external CTG was performed until the time at which either an amniotomy was performed for obstetric reasons or vaginal examination was performed after the membranes had ruptured. At that point, a scalp electrocardiographic electrode was applied.</p> <p>FHR tracings were examined by a skilled, unbiased observer who reported on their type and significance to the medical staff, who then made the final decision concerning management of the labour. All staff were trained in the recognition and significance of FHR abnormalities, but there</p>	<p>(Note: the authors reported the following details: normal labour (9 hours), type 1 dips present in contractions for a couple of hours before delivery with the FHR slowing to 100 bpm. The baby was delivered by forceps, with the head being rotated when the cord prolapsed. The baby was born in poor condition, with Apgar scores of 1 and 3, and died after 2 days in the intensive care. Cause of death was shown to be hypoxic brain damage)</p> <p><b><u>Neurological symptoms and/or signs (n/total (%))</u></b></p> <p>Standard: 3/495 (0.6) EFM: 1/479 (0.2) (Note: the data reported for this outcome appear not to exclude the women that the authors reported that they would, because N = 974)</p> <p><b><u>Care of the baby (n/total (%))</u></b></p> <p><b>a. Need for isolette*</b> Standard: 29/480 (6.0) EFM: 40/443 (9.0)</p> <p><b>b. Need for nursery*</b> Standard: 48/474 (10.1) EFM: 59/443 (13.3)</p> <p><b>c. Need for phototherapy</b> Standard: 4/480 (0.8) EFM: 16/443 (3.6)</p> <p>* The article reported the</p>	<p>Blinding of participants: Not reported Blinding of staff providing care: Not reported Blinding of outcome assessors: Not reported Missing data/loss to follow-up: There are small amounts of missing data (&lt; 2%) for need for isolette, need for nursery, and neurological signs and symptoms Precise definition of outcomes: Type of neurological symptoms or signs were not reported (and the denominator does not match what the authors stated that they would analyse/report in the methods section) Valid and reliable method of outcome assessment: Unclear for neurological symptoms and signs as no details were reported Intention to treat analysis performed: Yes</p> <p>No details of what standard care involved were reported. However, judging by the discussion section of the article, this has been assumed to be by intermittent auscultation. This is supported by assumptions made by Cochrane reviewers, who included this study in a review of intermittent auscultation compared with EFM</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Breech presentation</p> <p>Premature labour (37 weeks)</p> <p>Prolonged pregnancy (42 weeks)</p> <p>Prolonged labour (24 hours)</p> <p>Known fetal malformation</p>		<p>were very few incidences of abnormal traces.</p>	<p>proportion of babies spending 0, 1, 2 and <math>\geq 3</math> days in isolette/nursery; therefore, the proportion of babies not spending 0 days is reported above</p>	<p><b>Other information</b></p> <p>CTG was external until membranes ruptured, and then internal.</p> <p>49 women in the standard care group received EFM due to meconium in the amniotic fluid or FHR abnormality detected by auscultation. No caesarean sections were prompted by the results of the traces. Babies with early, mid or late dips were delivered by forceps</p>

**G.3 Intermittent auscultation compared with cardiotocography in the presence of meconium stained liquor**

Study Details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Alfirevic,Zarko, Devane,Declan, Gyte,Gillian ML, Continuous cardiotocography (CTG) as a form of electronic fetal monitoring (EFM) for fetal assessment during labour, Cochrane Database of Systematic Reviews, -, 2013</p> <p><b>Ref Id</b></p> <p>200781</p> <p><b>Country(ies) where the study was done</b></p> <p>Various</p> <p><b>Study type</b></p> <p>Systematic review</p> <p><b>Aim of the study</b></p> <p>To evaluate the effectiveness of</p>	<p><b>Sample size</b></p> <p>n = 500 from two studies (Pakistan 1989, Melbourne 1976)</p> <p><b>Characteristics</b></p> <p>Twelve studies included in the systematic review but only two studies consisted of right population for this review: <u>Pakistan 1989</u></p> <p>Randomisation: women selecting sealed unnumbered envelopes</p> <p>Participants: high-risk women all with meconium stained liquor</p> <p>Intervention: cardiotocography (CTG) versus intermittent auscultation</p> <p>Outcomes: neonatal mortality, mode of birth, Apgar score</p>	<p><b>Intervention</b></p> <p>Intermittent auscultation: intermittent monitoring undertaken either by listening to the baby's heart rate using a fetal stethoscope (Pinard) or a hand-held Doppler device</p> <p>Continuous fetal monitoring: electronic fetal heart rate monitoring by means of cardiotocograph</p>	<p><b>Details</b></p> <p><u>Electronic searches</u></p> <p>The Cochrane Pregnancy and Childbirth Group's Trials Register was searched by contacting the Trials Search Co-ordinator. CENTRAL, MEDLINE were searched, and hand searching of 30 journals and conference proceedings was done. No language restrictions were applied.</p> <p><u>Selection of studies</u></p> <p>Two review authors independently assessed the full text of all potential studies for inclusion and methodological quality.</p> <p><u>Data extraction and management</u></p> <p>Two authors extracted the data</p>	<p><b>Results</b></p> <p><u>Caesarean section</u></p> <p>Continuous fetal monitoring: n = 74/275 (26.9%)</p> <p>Intermittent auscultation: n = 36/275 (13.1%)</p> <p>RR 2.11 (1.19 to 3.74)</p> <p><u>Caesarean section for abnormal FHR pattern and/or acidosis</u></p> <p>Continuous fetal monitoring: n = 47/275 (17.1%)</p> <p>Intermittent auscultation: n = 21/275 (7.6%)</p> <p>RR 2.24 (1.38 to 3.64)</p> <p><u>Instrumental vaginal birth</u></p> <p>Continuous fetal monitoring: n = 108/275 (39.3%)</p>	<p><b>Limitations</b></p> <p>Pakistan 1989: - data extracted from unpublished trial lodged with Cochrane centre - no allocation concealment</p> <p><b>Other information</b></p> <p>The systematic review is available online at: <a href="http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006066.pub2/full">http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006066.pub2/full</a></p>

Study Details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>continuous cardiotocography during labour</p> <p><b>Study dates</b></p> <p>Assessed as up-to-date: January 2013</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>Study period: 1988 - 1989</p> <p><u>Melbourne 1976</u></p> <p>Randomisation: cards in sealed numbered envelopes</p> <p>Participants: high-risk women (40% with meconium stained liquor)</p> <p>Intervention: continuous CTG versus intermittent auscultation</p> <p>Outcomes: mode of birth, oxytocin use, analgesia use, maternal infection, neonatal mortality and morbidity, umbilical cord blood gas</p> <p>Study period: April 1974 - April 1975</p> <p><b>Inclusion criteria</b></p> <p>Randomised and quasi-randomised controlled trials</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>separately and double checked it for discrepancies. Statistical analysis was done using RevMan. Where information was unclear, the reviewers attempted to contact the original authors.</p> <p><u>Assessment of risk of bias</u></p> <p>Two review authors independently assessed risk of bias using criteria from the Cochrane Handbook for Systematic Reviews of Interventions:</p> <ul style="list-style-type: none"> <li>- Selection bias</li> <li>- Allocation concealment</li> <li>- Blinding</li> <li>- Incomplete outcome data</li> <li>- Sequence generation</li> <li>- Other sources of bias</li> </ul> <p><u>Measures of effect</u></p> <p>Dichotomous outcomes were presented risk</p>	<p>Intermittent auscultation: n = 94/275 (34.2%) RR 1.16 (0.88 to 1.54)</p> <p><u>Spontaneous vaginal birth not achieved</u></p> <p>Continuous fetal monitoring: n = 182/275 (66.2%)</p> <p>Intermittent auscultation: n = 130/275 (47.3%) RR 1.4 (1.2 to 1.63)</p> <p><u>Perinatal death</u></p> <p>Continuous fetal monitoring: n = 5/275 (1.8%)*</p> <p>Intermittent auscultation: n = 6/275 (2.2%)* RR 0.83 (0.26 to 2.67)</p> <p><u>NICU admission</u></p> <p>Continuous fetal monitoring: n = 11/175 (6.3%)</p> <p>Intermittent auscultation: n = 30/175 (17.1%) RR 0.37 (0.19 to 71)</p>	

Study Details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>ratios with 95% confidence intervals. For continuous data, weighted mean differences were used. Fixed-effect analysis was performed in the absence of significant heterogeneity. In the presence of heterogeneity sensitivity analysis followed by random effects analysis was performed.</p> <p><u>Dealing with missing data</u> The authors investigated the effect of including trials with high levels of attrition using sensitivity analysis. Outcomes were assessed on an intention-to-treat basis, with the denominator being set as the number randomised minus any participants whose outcomes were known to be missing.</p>	<p><u>Infection/damage from scalp electrode</u> Continuous fetal monitoring: n = 1/100 (1%) Intermittent auscultation: n = 0/100 (0%) RR 3.00 (0.12 to 72.77)</p> <p><u>Neonatal seizure</u> Continuous fetal monitoring: n = 0/175 (0%) Intermittent auscultation: n = 4/175 (2.3%) RR 0.11 (0.01 to 2.05)</p>	



Study Details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p><u>Analysis</u>                      If high levels of heterogeneity (&gt; 50%) were identified, prespecified sensitivity analysis was performed according to the quality of the trials. Planned subgroup analyses:</p> <ol style="list-style-type: none"> <li>1. low risk (absence of identified risk factors)</li> <li>2. high risk of perinatal mortality and morbidity</li> <li>3. spontaneous onset of labour</li> <li>4. induction of labour</li> <li>5. preterm</li> <li>6. term</li> <li>7. singleton/twin pregnancy</li> <li>8. with and without fetal blood sampling (FBS)</li> <li>9. parity</li> </ol>		

### G.4 Interpretation of cardiocograph traces

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Cibils,L.A., Clinical significance of fetal heart rate patterns during labor. II. Late decelerations, American Journal of Obstetrics and Gynecology, 123, 473-494, 1975</p> <p><b>Ref Id</b></p> <p>195117</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To evaluate fetal heart rate (FHR) changes and patterns in two groups (with decelerations, no decelerations) in</p>	<p><b>Sample size</b></p> <p>n = 1304 records reviewed: n = 598 had no accelerations, n = 147 had late decelerations</p> <p><b>Characteristics</b></p> <p>Women in the no decelerations group were younger than women in the late decelerations group (22.8 years versus 25.1 years). Gestational age and duration of FHR recording were similar in the two groups</p> <p><b>Inclusion criteria</b></p> <p>Singleton pregnancy</p> <p>Cephalic presentation</p> <p>Direct or internal monitoring</p> <p>Minimum of 60 minutes recording prior to 2nd stage/decision to</p>	<p><b>Interventions</b></p> <p>60 minutes of FHR trace analysis (available prior to second stage of labour)</p>	<p><b>Details</b></p> <p>During the study period n = 1,304 records were reviewed manually and coded (details provided in a previously published paper). n = 598 (46%) had no decelerations of FHR which could be correlated in time with uterine contractions. n = 147 (11%) had FHR late decelerations</p>	<p><b>Results</b></p> <p>There is low likelihood of neonatal problems when there is no deceleration of FHR:</p> <p><u>Neonatal morbidity and/or death*</u></p> <p>Late decelerations group: 7%</p> <p>No decelerations group: 0.5%</p> <p>p &lt; 0.0001</p> <p>* no further details on neonatal mortality reported</p> <p>High numbers of mortality and morbidity present in neonates with low birthweight with late decelerations:</p> <p><u>Neonatal morbidity and/or death in low birthweight babies &lt; 2500g</u></p> <p>Late decelerations group: 15%</p> <p>No decelerations group: 5%</p> <p>p = ns</p> <p>A high percentage of babies with FHR late decelerations (50%) were distressed during labour and 33% born depressed (clinical distress defined as presence of meconium stained liquor, tachycardia, markedly irregular heart beat, no definition for "depressed" babies given)</p>	<p><b>Limitations</b></p> <p>Limited outcome data</p> <p>No exclusion criteria specified hence high risk of selection bias</p> <p>Women's demographic characteristics not reported</p> <p>Unclear how and by whom data were analysed</p> <p>No statistical analysis of data reported</p> <p><b>Other information</b></p> <p>Normal baseline FHR defined as 120 to 150 beats per minute (bpm)</p> <p>Tachycardia: &gt; 150 beats per minute</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>order to predict fetal condition at birth</p> <p><b>Study dates</b> June 1970 to 1974</p> <p><b>Source of funding</b> Not reported</p>	<p>perform a caesarean section</p> <p><b>Exclusion criteria</b> Not reported</p>				
<p><b>Full citation</b> Cibils, L.A., Clinical significance of fetal heart rate patterns during labor. V. Variable decelerations, American Journal of Obstetrics and Gynecology, 132, 791-805, 1978</p> <p><b>Ref Id</b> 195119</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b> n = 1304 records reviewed. n = 598 had no decelerations, n = 312 had variable decelerations</p> <p><b>Characteristics</b> Women in the no decelerations group were significantly younger than women in the late decelerations group (22.8 yr vs. 24.4 yr), had higher gestational age (39.4 wk vs. 38.6 wk) and longer duration of FHR recording (252 minutes vs. 223 minutes). Fetal</p>	<p><b>Interventions</b> FHR: variable decelerations  variable decelerations with late component ('variable with hypoxic component')</p>	<p><b>Details</b> From n = 1,304 records that were reviewed manually and coded (details provided in a previously published paper): n = 598 (46%) had no decelerations of FHR which could be correlated in time with uterine contractions; n = 312 had FHR variable decelerations (n = 18 women</p>	<p><b>Results</b> Cases with variable decelerations n = 312 Cases with no deceleration n = 598</p> <p>Association between variable deceleration and baseline alterations (tachycardia, saltatory or fixed FHR baselines):</p> <p><u>Saltatory fixed</u> No deceleration: 39% Variable decelerations: 25% p = ns</p> <p><u>Tachycardia</u> No decelerations: 5% Variable decelerations: 21% p &lt; 0.0005</p> <p><u>Sustained</u></p>	<p><b>Limitations</b> Limited outcome data  No exclusion criteria specified hence high risk of selection bias  Women's demographic characteristics not reported  Unclear how and by whom data were analysed</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>USA</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To evaluate fetal heart rate (FHR) changes and patterns in two groups (with decelerations, variable decelerations) in order to predict fetal condition at birth</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>weight was significantly higher in the no decelerations group compared with the variable decelerations group (3236 g vs. 2988 g). There were fewer normal and hypertensive women in the variable decelerations group, but there was a higher rate of women with other pathological conditions such as premature rupture of membranes.</p> <p><b>Inclusion criteria</b></p> <p>Singleton labours</p> <p>60 minutes of FHR trace available prior to second stage</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>had variable decelerations with a component of late deceleration in the recovery period, all of these cases had umbilical cord problems). The maternal condition and neonatal outcomes were compared in order to ascertain the clinical value of observed changes in FHR pattern.</p>	<p>No decelerations: 8% Variable decelerations: 21% p &lt; 0.0005</p> <p><u>Fetal distress</u> No decelerations: 4% Variable decelerations: 23% p &lt; 0.0005</p> <p><u>Neonatal death</u> No decelerations: 0.2% Variable decelerations: 2.2% p &lt; 0.0005</p> <p>Significant association between variable decelerations (with a hypoxic [late] component) and baseline alterations (tachycardia, saltatory or fixed FHR baselines):</p> <p><u>Saltatory fixed</u> Variable decelerations with late component: 39% Variable decelerations: 25% p &lt; 0.0005</p> <p><u>Tachycardia</u> Variable decelerations with late component: 61% Variable decelerations: 21% p &lt; 0.0005</p> <p><u>Sustained</u> Variable decelerations with late component: 67% Variable decelerations: 21% p &lt; 0.0005</p> <p><u>Fetal distress</u> Variable decelerations with late component:</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				78% Variable decelerations: 23% p < 0.0005  <u>Neonatal death</u> Variable decelerations with late component: 11% Variable decelerations: 2.2% p = ns	
<p><b>Full citation</b></p> <p>Cibils,L.A., Clinical significance of fetal heart rate patterns during labor. VI. Early decelerations, American Journal of Obstetrics and Gynecology, 136, 392-398, 1980</p> <p><b>Ref Id</b></p> <p>195120</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort</p>	<p><b>Sample size</b></p> <p>n = 1304 records reviewed. n= 598 had no accelerations, n = 247 had early decelerations</p> <p><b>Characteristics</b></p> <p>Women in the no decelerations group were younger than women in the early decelerations group (22.8 yr vs. 23.6 yr), had similar gestational ages (39.4 wk vs. 38.2 wk) and longer durations of FHR recording (252 minutes vs. 231 minutes). Fetal weight was significantly higher in the no decelerations group compared with the early decelerations</p>	<p><b>Interventions</b></p> <p>FHR: No decelerations  Early decelerations</p>	<p><b>Details</b></p> <p>From n = 1,304 records that were reviewed manually and coded (referred to a previous published paper): n = 598 (46%) had no decelerations of FHR which could be correlated in time with uterine contractions; n = 247 had FHR early decelerations prior to 2nd stage of labour. The maternal condition and neonatal outcomes were compared in</p>	<p><b>Results</b></p> <p><u>Transient tachycardia</u> Early decelerations group: 10% No decelerations groups: 5%</p> <p><u>Fetal distress (no definition provided)</u> Early decelerations group: 5% No decelerations groups: 4%</p> <p><u>Neonatal death</u> Early decelerations group: n = 1 (congenital heart disease) No decelerations groups: n = 1 (congenital malformation)</p>	<p><b>Limitations</b></p> <p>Limited outcome data  No exclusion criteria specified hence high risk of selection bias  Women's demographic characteristics not reported  Unclear how and by whom data were analysed</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To evaluate fetal heart rate (FHR) changes and patterns in two groups (no decelerations, early decelerations) in order to predict fetal condition at birth</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>group (3236 g vs. 3129 g).</p> <p><b>Inclusion criteria</b></p> <p>Singleton labours</p> <p>60 minutes of FHR trace available prior to second stage</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>order to ascertain the clinical value of observed changes in FHR pattern.</p>		
<p><b>Full citation</b></p> <p>Cibils,L.A., Votta,R., Clinical significance of fetal heart rate patterns during labor. IX: Prolonged pregnancy, Journal of Perinatal</p>	<p><b>Sample size</b></p> <p>707 post-term pregnancies (&gt; 14 days post estimated date of delivery [EDD])</p> <p><b>Characteristics</b></p> <p>No characteristics specified. It is specified that the relevant clinical informations has been</p>	<p><b>Interventions</b></p> <p>Fetal heart rate records</p>	<p><b>Details</b></p> <p>n = 707 pregnancies that passed the estimated date of delivery by 14 days were included in the study. This was assessed in women with good menstrual histories, who</p>	<p><b>Results</b></p> <p>No significant correlation between abnormal FHR patterns and pH:  n = 598 no decelerations  n = 147 traces with late decelerations</p> <p><u>Deceleration pattern</u>  Variable decelerations: 55%  No or early decelerations: 23%  Late deceleration: 17%</p> <p><u>Baseline frequency</u>  Normal: 71%</p>	<p><b>Limitations</b></p> <p>No exclusion criteria specified hence high risk of selection bias</p> <p>Women's demographic characteristics not reported</p> <p>Unclear how and by whom data were analysed</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Medicine, 21, 107-116, 1993</p> <p><b>Ref Id</b></p> <p>195122</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Case series</p> <p><b>Aim of the study</b></p> <p>To evaluate fetal heart rate (FHR) changes and patterns in women with prolonged labour in order to diagnose early fetal compromise</p> <p><b>Study dates</b></p> <p>July 1980 to December 1984</p>	<p>reported in a previously published paper.</p> <p><b>Inclusion criteria</b></p> <p>Post-term pregnancies (&gt; 14 days post EDD)</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>had dating examinations or confirmed by an ultrasound in the first trimester of pregnancy. All women had either internal or external continuous fetal monitoring. Data for this study were gathered prospectively. The observation was based on the interpretation of fetal heart rate and uterine contraction and their value as a tool to diagnose early fetal compromise or to prevent fetal deterioration by early intervention. Statistical analysis was performed</p>	<p>Tachycardia: 26% Bradycardia: 4%</p> <p><u>Baseline pattern</u> Normal: 75% Fixed: 8% Saltatory: 17%</p> <p>Acidemia (pH ≤ 7.20) could not be predicted from deceleration patterns in FHR trace: <u>FHR and umbilical cord pH</u> pH ≤ 7.20 Total n = 46 pH ≥ 7.21 Total n = 108</p> <p><u>No or early decelerations</u> pH ≤ 7.20 n = 11 (23%) pH ≥ 7.21 n = 25 (23%)</p> <p><u>Variable decelerations</u> pH ≤ 7.20 n = 17 (36%) pH ≥ 7.21 n = 48 (44%)</p> <p><u>Late decelerations</u> pH ≤ 7.20 n = 18 (39%) pH ≥ 7.21 n = 35 (32%)</p>	<p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p> <p>Not specified</p>			<p>using <math>\chi^2</math> method.</p>		
<p><b>Full citation</b></p> <p>Low, J.A., Cox, M.J., Karchmar, E.J., McGrath, M.J., Pancham, S.R., Piercy, W.N., The prediction of intrapartum fetal metabolic acidosis by fetal heart rate monitoring, American Journal of Obstetrics and Gynecology, 139, 299-305, 1981</p> <p><b>Ref Id</b></p> <p>195666</p> <p><b>Country/ies where the study was carried out</b></p> <p>Canada</p> <p><b>Study type</b></p> <p>Case series</p>	<p><b>Sample size</b></p> <p>n = 200 term infants with significant metabolic acidosis (base buffer &lt; 36.1 mEq/l)</p> <p>n = 200 term infants without metabolic acidosis (base buffer &gt; 36.1 mEq/l)</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <p>Women admitted and monitored in the intrapartum intensive-care unit.</p>	<p>All FHR variables</p>	<p><b>Details</b></p> <p>FHR characteristics during the 8 hours prior to delivery were studied in 200 women in whom the baby had evidence of a metabolic acidosis at birth (base buffer &lt; 36.1 mEq/l), and compared to those in 200 women in whom the baby had a normal acid-base at birth (base buffer &gt; 36.1 mEq/l). Fetal heart rate records were scored for each 20 minute period for a maximum of 24 twenty-minute cycles (8</p>	<p><b>Results</b></p> <p>There was no statistically significant difference between the two groups in regard to decrease frequency or absence of FHR accelerations in the 12 FHR trace cycles (4 hours before birth) indicating that fetal heart rate accelerations (as an independent variable) were not predictive of fetal acidosis (no synthesis of the statistical data provided).</p> <p>Total decelerations and variable decelerations in last hour prior to birth were significantly associated with acidosis. Late decelerations in the last hour prior to birth were significantly associated with neonatal acidosis. Variable decelerations only in last 20 minutes prior to birth were significantly associated with acidosis:</p> <p><u>Cycle 1 (20 min FHR trace 20 min before birth)</u></p> <p>Total decelerations: Index: n = 51/200 Control: n = 33/200 <math>p = 0.001</math></p> <p><u>Cycle 1 (20 min FHR trace 20 min before</u></p>	<p><b>Limitations</b></p> <p>No analysis on combining factors for prediction.</p> <p><b>Other information</b></p> <p>Baseline heart rate classified as normal: 120 to 160 beats per minute (bpm) Bradycardia: &lt; 120 bpm Tachycardia: &gt; 160 bpm</p> <p>Baseline variability: amplitude of oscillation as normal (6 to 25 bpm), decreased (3 to 5 bpm) and absent (&lt; 3 bpm)</p> <p>Accelerations: at least 15 bpm above the baseline. Normal (<math>\geq 2</math> acceleration in 20 min), decreased (1 acceleration in 20 min), absent (no accelerations in 20 min)</p> <p>Decelerations: fall in FHR in excess of 15 bpm. Total deceleration patterns were classified on the basis of frequency of contraction in 20 minute period. None (0% or 4% contractions associated with a deceleration), moderate (5% to 30% contractions associated with a deceleration), marked (&gt; 30% contractions associated with a deceleration)</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To evaluate the fetal heart rate (FHR) characteristics in predicting the presence of a metabolic acidosis</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>hours) prior to birth. All records were assessed by one of the two authors. The assessment was performed without knowledge of the clinical or laboratory data. In each 20 minute cycle the following characteristics were scored: baseline fetal heart rate, baseline FHR long term variability, FHR accelerations, FHR variable decelerations and FHR late decelerations.</p>	<p><u>birth)</u> Variable decelerations: Index: n = 38/200 Control: n = 30/200 p = 0.01</p> <p><u>Cycle 1 (20 min FHR trace 20 min before birth)</u> Late decelerations: Index: n = 78/200 Control: n = 23/200 p = 0.001</p> <p><u>Cycle 2 (20 min FHR trace 40 min before birth)</u> Total decelerations: Index: n = 42/200 Control: n = 30/200 p = 0.001</p> <p><u>Cycle 2 (20 min FHR trace 40 min before birth)</u> Variable decelerations: Index: n = 30/200 Control: n = 26/200 p = 0.2</p> <p><u>Cycle 2 (20 min FHR 40 min trace before birth)</u> Late decelerations: Index: n = 59/200 Control: n = 21/200 p = 0.001</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><u>Cycle 3 (20 min FHR trace 60 min before birth)</u>                      Total decelerations:                      Index: n = 35/200                      Control: n = 26/200                      p = 0.006</p> <p><u>Cycle 3 (20 min FHR trace 60 min before birth)</u>                      Variable decelerations:                      Index: n = 26/200                      Control: n = 24/200                      p = 0.3</p> <p><u>Cycle 3 (20 min FHR 60 min trace before birth)</u>                      Late decelerations:                      Index: n = 42/200                      Control: n = 21/200                      p = 0.01</p>	
<p><b>Full citation</b></p> <p>Low, J.A., Pancham, S.R., Piercy, W.N., Intrapartum fetal asphyxia: Clinical characteristics, diagnosis, and significance in relation to pattern of development, American Journal of Obstetrics and</p>	<p><b>Sample size</b></p> <p>Total n = 587</p> <p>n = 122 with significant metabolic acidosis (base buffer &lt; 36.1 mEq/l)</p>	<p><b>Interventions</b></p> <p>All FHR variables</p>	<p><b>Details</b></p> <p>Fetal heart rate records (obtained via a scalp electrode) were reviewed for each two hour period prior to birth in n = 587 women. Based on the serial acid base observations (maternal</p>	<p><b>Results</b></p> <p>There were no statistically significant differences between the two groups (asphyxia and normal group) at mid-labour (&gt; 2 hours prior to birth) in regard to pH, buffer base, and oxygen or carbon dioxide tension. However, the maternal pH, buffer base, and oxygen tension in the asphyxia group were all significantly lower compared to the normal group at two hours, one hour and 5 minutes prior to birth. The umbilical artery and vein buffer base was also</p>	<p><b>Limitations</b></p> <p>Unclear how and by who the records were assessed.</p> <p><b>Other information</b></p> <p>Baseline heart rate classified as normal: 120 to 160 beats per minute (bpm) bradycardia: &lt; 120 bpm, tachycardia: &gt; 160 bpm</p> <p>Baseline variability: amplitude of oscillation as normal (6 to 25 bpm), decreased (3 to 5 bpm) and absent (&lt; 3 bpm)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Gynecology, 129, 857-872, 1977</p> <p><b>Ref Id</b></p> <p>196822</p> <p><b>Country/ies where the study was carried out</b></p> <p>Canada</p> <p><b>Study type</b></p> <p>Case series</p> <p><b>Aim of the study</b></p> <p>To examine clinical circumstances related to development of intrapartum fetal asphyxia</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Supported by Ministry of Health grant</p>	<p>n = 465 without metabolic acidosis (base buffer &gt; 36.1 mEq/l)</p> <p><b>Characteristics</b></p> <p><u>Parity 0</u> Normal group: 61% Asphyxia terminal: 67% Asphyxia/one hour: 55% Asphyxia/two hours: 72%</p> <p><u>Parity ≥ 1</u> Normal group: 39% Asphyxia terminal: 33% Asphyxia one/hour: 45% Asphyxia two/hours: 28%</p> <p><u>Preterm neonates</u> Normal group: 11% Asphyxia terminal: 0% Asphyxia one/hour: 15% Asphyxia two/hours: 3%</p> <p><u>Preterm neonates</u> Normal group: 10% Asphyxia terminal: 0% Asphyxia one/hour: 15%</p>		<p>venous blood acid base, lactate, and pyruvate characteristics during the labour and birth, fetal acid base characteristics during the last half of labour and fetal acid base, lactate and pyruvate characteristics during the labour and birth), women were divided into the normal group or the asphyxia group. FHR observations were made on the total decelerations, and late decelerations in relation to the contractions in each two hour period. The baseline FHR was observed at six 20-minute intervals in a</p>	<p>significantly lower in the asphyxia group when compared with the normal group.</p> <p>Normal group n = 465 Asphyxia group n = 122 (terminal n = 46, one hour n = 40, two hours n = 36)</p> <p><u>Perinatal death</u> Normal group: n = 29/465 (16%) Asphyxia terminal: n = 1/46 (2%) Asphyxia one/hour: n = 0/40 (0%) Asphyxia two/hours: n = 1/36 (3%)</p> <p><u>Mode of birth</u> <u>Spontaneous low forceps</u> Normal group: n = 270/465 (58%) Asphyxia terminal: n = 14/46 (30%) Asphyxia/one hour: n = 14/40 (35%) Asphyxia/two hours: n = 11/36 (30%)</p> <p><u>Mid-forceps</u> Normal group: n = 133/465 (29%) Asphyxia terminal: n = 28/46 (61%) Asphyxia/one hour: n = 14/40 (35%) Asphyxia/two hours: n = 8/36 (22%)</p>	<p>Accelerations: at least 15 bpm above the baseline. Normal (≥ 2 accelerations in 20 min), decreased (1 acceleration in 20 min), absent (no accelerations in 20 min) Decelerations: fall in FHR in excess of 15 bpm. Total deceleration patterns were classified on the basis of frequency of contractions in 20 minute period. None (0% or 4% contractions associated with a deceleration), moderate (5% to 30% contractions associated with a deceleration), marked (&gt; 30% contractions associated with a deceleration)</p> <p>Total decelerations defined as percentage of contractions associated with a deceleration in each two-hour period. It was classified as moderate (5% to 29% of contractions were associated with a deceleration) and marked (&gt; 30% of contractions were associated with a deceleration)</p> <p>Late decelerations defined as percentage of contractions associated with a late deceleration in each two-hour period. It was classified as moderate (&lt; 10% of contractions were associated with a late deceleration) and marked (≥ 10% of contractions were associated with a late deceleration)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Asphyxia two/hours: 3%</p> <p><u>Post term gestation</u> Normal group: 10% Asphyxia terminal: 13% Asphyxia one/hour: 20% Asphyxia two/hours: 14%</p> <p><u>Medical complication (hypertension, diabetes, other)</u> Normal group: 15% Asphyxia terminal: 12% Asphyxia one/hour: 9% Asphyxia two/hours: 33%</p> <p><u>Meconium stained liquor</u> Normal group: 33% Asphyxia terminal: 35% Asphyxia one/hour: 45% Asphyxia two/hours: 50%</p> <p><u>Regional or local anaesthesia</u> Normal group: 90% Asphyxia terminal: 85% Asphyxia one/hour: 75% Asphyxia two/hours: 80%</p>		<p>two hour period. The normal acid base group as determined by a serial acid base study during birth included n = 465 women with a fetus with capillary blood buffer base of &gt; 1 SD below the normal mean, i.e. <math>\geq 40</math> mEq/l, and umbilical artery buffer base at delivery of &gt; 1 SD below the normal mean, i.e. <math>\geq 38.6</math> mEq/l.</p> <p>The fetal asphyxia group included n = 122 women in whom the baby at delivery had an umbilical artery buffer base of &lt; 2 SD below the normal mean, i.e. &lt; 36.1 mEq/L.</p>	<p><u>Caesarean section</u> Normal group: n = 55/465 (12%) Asphyxia terminal: n = 3/46 (6%) Asphyxia/one hour: n = 9/40 (22%) Asphyxia/two hours: n = 16/36 (44%)</p> <p><u>Marked patterns of total decelerations (8 hours prior to birth)</u> Normal group: 9% Asphyxia terminal: 29% Asphyxia/one hour: not reported Asphyxia/two hours: 20%</p> <p><u>Marked patterns of total decelerations (6 hours prior to birth)</u> Normal group: 13% Asphyxia terminal: 21% Asphyxia/one hour: 14% Asphyxia/two hours: 20%</p> <p><u>Marked patterns of total decelerations (4 hours prior to birth)</u> Normal group: 19% Asphyxia terminal: 30% Asphyxia/one hour: 37% Asphyxia/two hours: 39%</p> <p><u>Marked patterns of total decelerations (2 hours prior to birth)</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><b>Inclusion criteria</b></p> <p>Women admitted and monitored in the intrapartum intensive-care unit. The criteria for admission were maternal, fetal, or labour risk factors that could have been predictive of fetal asphyxia.</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>Duration of metabolic acidosis during labour were determined by the available serial fetal acid base observation in the second half of labour for each case. The criteria of developing metabolic acidosis during labour were a capillary blood buffer base of &lt; 1 SD below the normal mean in the last hour of labour, i.e. &lt; 40 mEq/l.</p> <p>The asphyxia group were divided into three groups based on the acid base characteristics during labour and delivery: terminal asphyxia (just before birth); asphyxia/one hour (one hour</p>	<p>Normal group: 34% Asphyxia terminal: 54% Asphyxia/one hour: 52% Asphyxia/two hours: 61%</p> <p><u>Moderate or marked patterns of late decelerations (8 hours prior to birth)</u> Normal group: 15% Asphyxia terminal: 9% Asphyxia/one hour: not reported Asphyxia/two hours: not reported</p> <p><u>Moderate or marked patterns of late decelerations (6 hours prior to birth)</u> Normal group: 18% Asphyxia terminal: 31% Asphyxia/one hour: 8% Asphyxia/two hours: 16%</p> <p><u>Moderate or marked patterns of late decelerations (4 hours prior to birth)</u> Normal group: 21% Asphyxia terminal: 26% Asphyxia/one hour: 26% Asphyxia/two hours: 27%</p> <p><u>Moderate or marked patterns of late decelerations (2 hours prior to birth)</u> Normal group: 31%</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			before birth); asphyxia/two hours (two hours before birth).	Asphyxia terminal: 59% Asphyxia/one hour: 59% Asphyxia/two hours: 68%	
<p><b>Full citation</b></p> <p>Maso,G., Businelli,C., Piccoli,M., Montico,M., De,Seta F., Sartore,A., Alberico,S., The clinical interpretation and significance of electronic fetal heart rate patterns 2 h before delivery: an institutional observational study, Archives of Gynecology and Obstetrics, 286, 1153-1159, 2012</p> <p><b>Ref Id</b></p> <p>275105</p> <p><b>Country/ies where the study was carried out</b></p> <p>Italy</p>	<p><b>Sample size</b></p> <p>n = 198</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <ul style="list-style-type: none"> <li>- Singleton</li> <li>- Term</li> <li>- Spontaneous and operative vaginal birth</li> <li>- External continuous FHR monitoring during the last 2 hours of labour was available</li> <li>- Short term neonatal outcomes were available</li> <li>- Low risk pregnancy (defined as cases without risk factors for the development of acidosis, cerebral palsy, perinatal death, and neonatal encephalopathy)</li> </ul>	<p><b>Interventions</b></p> <p>Intrapartum electronic fetal monitoring</p>	<p><b>Details</b></p> <p>Data collected (retrospective for 6 months) from a labour database of Maternal and Child Institute Burlo Garofolo in Italy. Based on the inclusion criteria, all cases with the last 2 hours continuous electronic fetal monitoring (EFM) before birth were included in the study. An obstetrician, blinded to neonatal outcomes, retrospectively reviewed the included cases. The tracings were</p>	<p><b>Results</b></p> <p>Umbilical artery pH value of 7.20 chosen as the cut off to define neonatal acidemia.</p> <p><b>Three EFM groups: normal, suspicious, pathological</b></p> <p><u>Normal</u> If all four FHR variables (baseline, variability, decelerations, accelerations) falls into reassuring category (see 'Other information')</p> <p><u>Suspicious</u> If one of the variables presented non reassuring characteristics and the reminder variables were reassuring (see 'Other information')</p> <p><u>Pathological</u> If more than two non-reassuring or more than one abnormal variable was respectively (see 'Other information')</p> <p><b>Mean pH values in the three EFM groups:</b></p> <p><u>Normal</u> pH 7.30 (95% CI 7.28 to 7.32)</p> <p><u>Suspicious</u> pH 7.25 (95% CI 7.23 to 7.27)</p> <p><u>Pathological</u> pH 7.20 (95% CI 7.17 to 7.13) p &lt; 0.001 (for all pairwise comparisons)</p>	<p><b>Limitations</b></p> <ul style="list-style-type: none"> <li>- Women characteristics not reported</li> <li>- Selective data reported</li> </ul> <p><b>Other information</b></p> <p><b>Categorisation of FHR:</b></p> <p><u>Reassuring</u> Baseline: 100-180 Variability: ≥ 5 Decelerations: none Accelerations: present</p> <p><u>Non-reassuring</u> Baseline: 110 -160 Variability: &lt; 5 for ≥ 40 but &lt; 90 min Decelerations: - repetitive (≥ 3) typical variable decelerations with over 50% of contractions - single prolonged &lt; 3 min Accelerations: the absence of accelerations with an otherwise normal FHR tracing is of uncertain significance</p> <p><u>Abnormal</u> Baseline: - 161 - 180 - &lt; 100 - &gt;180 - sinusoidal pattern</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study type</b></p> <p>Case series</p> <p><b>Aim of the study</b></p> <p>To evaluate the clinical significance of intrapartum fetal heart rate (FHR) monitoring in low-risk pregnancies</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p><b>Exclusion criteria</b></p> <p>Cases with risk factors for the development of acidosis, cerebral palsy, perinatal death, and neonatal encephalopathy</p>		<p>interpreted as normal, suspicious or pathological, according to specific guidelines of EFM and by grouping the different FHR patterns considering baseline, variability, presence of decelerations and bradycardia (see 'Other information' section).</p> <p>Analysis: Comparisons between groups were performed with Kruskal-Wallis test. Differences among categorical variables were evaluated using Fisher's exact test.</p>	<p><b>Mean BD mmol/L values in the three EFM groups:</b></p> <p><u>Normal</u> -3.35 (95% CI -4.19 to -2.50)</p> <p><u>Suspicious</u> -5.62 (95% CI -6.43 to -4.81)</p> <p><u>Pathological</u> -7.50 (95% CI -8.50 to -6.50) p &lt; 0.001 (for all pairwise comparisons)</p> <p><b>Composite dverse outcomes*:</b></p> <p><u>Normal</u> n = 0/51 (0%)</p> <p><u>Suspicious</u> n = 5/88 (5.7%)</p> <p><u>Pathological</u> n = 6/59 (10.1%) p = 0.005 (normal vs. pathological)</p> <p><b>Normal variability:</b></p> <p><u>pH &lt; 7.20</u> n = 3/51 (5.9%)</p> <p><u>pH &lt; 7.10</u> n = 0/51 (0%)</p> <p><u>PH &lt; 7.00</u> n = 0/51 (0%)</p> <p><u>BD mmol/l</u> 0/51 (0%)</p> <p><b>Normal variability and typical variable decelerations:</b></p> <p><u>pH &lt; 7.20</u> n = 18/63 (28.6%)</p> <p><u>pH &lt; 7.10</u> n = 6/63 (9.5%)</p> <p><u>PH &lt; 7.00</u> n = 1/63 (1.6%)</p> <p><u>BD mmol/l</u> 5/63 (7.9%)</p>	<p>- ≥ 10 min</p> <p>Variability: &lt; 5 for ≥ 40 to ≥ 90 min</p> <p>Decelerations:</p> <ul style="list-style-type: none"> <li>- either repetitive (≥ 3) atypical variable decelerations or late decelerations, with over 50% of contractions</li> <li>- single prolonged deceleration &gt; 3 min</li> </ul> <p>Accelerations: the absence of accelerations with an otherwise normal FHR tracing is of uncertain significance</p> <p><b><u>Normal, suspicious, pathological</u></b></p> <p><u>Normal</u> If all four FHR variables (baseline, variability, decelerations, accelerations) falls into reassuring category</p> <p><u>Suspicious</u> If one of the variables presented non reassuring characteristics and the reminder variables were reassuring</p> <p><u>Pathological</u> If more than two non-reassuring or more than one abnormal variable was respectively</p> <p><b>FHR features definitions:</b></p> <p><u>Atypical variable</u> Defined in the presence of at least one of the following conditions: loss of primary or secondary rise in the baseline rate; slow return to baseline FHR after the contraction; prolong secondary rise in the baseline rate; biphasic deceleration; loss of variability during deceleration; continuation of baseline rate at lower level</p> <p><u>Bradycardia</u> Defined as moderate or severe if persistent fall of baseline between 100 and 109 bpm was respectively observed over a time period of 5 to 10 min.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><b>Normal variability and atypical variable decelerations:</b>  <u>pH &lt; 7.20</u>                      n = 13/27 (48.2%)  <u>pH &lt; 7.10</u>                      n = 2/27 (7.4%)  <u>PH &lt; 7.00</u>                      n = 0/27 (0%)  <u>BD mmol/l</u>                      0/27 (0%)</p> <p><b>Moderate bradycardia</b>  <u>pH &lt; 7.20</u> n = 6/17 (35.3%)  <u>pH &lt; 7.10</u> n = 0/17 (0%)  <u>PH &lt; 7.00</u> n = 0/17 (0%)  <u>BD mmol/l</u> 0/17 (0%)</p> <p><b>Severe bradycardia</b>  <u>pH &lt; 7.20</u> n = 7/15 (46.7%)  <u>pH &lt; 7.10</u> n = 4/15 (26.7%)  <u>PH &lt; 7.00</u> n = 1/15 (6.7%)  <u>BD mmol/l</u> 2/15 (13.3%)</p> <p>*Composite neonatal outcomes: umbilical artery pH &lt; 7 and/or APGAR score &lt; 7 at 5 min and/or neonatal resuscitation in delivery room and admission to neonatal intensive care unit for distress at birth.</p>	
<p><b>Full citation</b></p> <p>Cahill,A.G.,                      Caughey,A.B.,                      Roehl,K.A.,                      Odibo,A.O.,                      Macones,G.A.,                      Terminal fetal</p>	<p><b>Sample size</b></p> <p>Terminal deceleration:                      n = 951                      No terminal deceleration n = 4,437</p>	<p><b>Interventions</b></p> <p>Electronic fetal monitoring</p>	<p><b>Details</b></p> <p>Data collected from all consecutive births at Washington University in</p>	<p><b>Results</b></p> <p><b><u>Terminal deceleration and neonatal outcomes</u></b>  <u>Arterial umbilical cord pH level of 7.10 or less</u>                      Terminal deceleration                      n = 12/951 (1.3%)</p>	<p><b>Limitations</b></p> <ul style="list-style-type: none"> <li>- Uneven number of participants in two groups</li> <li>- 30 min EFM traces just before birth were analysed</li> <li>- if trace was lost or discontinuous after the initiation of the terminal deceleration, it was</li> </ul>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>heart decelerations and neonatal outcomes, Obstetrics and Gynecology, 122, 1070-1076, 2013</p> <p><b>Ref Id</b> 298858</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Study type</b> Retrospective cohort study</p> <p><b>Aim of the study</b> To examine the incidence and characteristics of terminal fetal heart rate decelerations and to estimate their association with acidemia</p> <p><b>Study dates</b></p>	<p><b>Characteristics</b></p> <p>Groups were similar with respect to:</p> <ul style="list-style-type: none"> <li>- maternal age and race</li> <li>- body mass index</li> <li>- gestational age at delivery</li> <li>- use of regional anesthesia</li> <li>- induction in labour</li> </ul> <p>Women with a terminal deceleration were more likely to be nulliparous and, they were less likely to have a spontaneous vaginal birth. The mean BMI in both groups was &gt; 31.</p> <p><b>Inclusion criteria</b></p> <ul style="list-style-type: none"> <li>- singleton</li> <li>- vertex gestation at term (at or after 37 0/7 weeks),</li> <li>- labored, and reached complete dilation.</li> </ul> <p><b>Exclusion criteria</b></p> <ul style="list-style-type: none"> <li>- Multiple gestation</li> <li>- Fetus with a known congenital anomaly</li> <li>- Did not have sufficient electronic</li> </ul>		<p>St. Louis Medical Center during the study period. The institutional policy is one of universal EFM during labor and arterial umbilical cord gas pH level birth.</p> <p>Women's EFM trace from 30 minutes before birth was interpreted by two formally trained obstetric research nurses certified in EFM interpretation and blinded to clinical data and outcomes.. Electronic fetal monitoring was interpreted using the <i>Eunice Kennedy Shriver</i> National Institute of Child Health and Human Development and the American</p>	<p>Not terminal deceleration n = 45/4437 (1.0%) Adjusted* OR 1.2 (95% CI 0.6 to 2.3) P = 0.49</p> <p><u>Arterial umbilical cord pH level of 7.05 or less</u></p> <p>Terminal deceleration n = 4/951 (0.4%)</p> <p>Not terminal deceleration n = 13/4437 (0.3%) Adjusted* OR 1.4 (95% CI 0.5 to 4.4) P = 0.52</p> <p><u>Arterial umbilical cord pH level of 7.10 or less and base excess &lt; -8.0</u></p> <p>Terminal deceleration n = 11/951 (1.2%)</p> <p>Not terminal deceleration n = 39/4437 (0.9%) Adjusted* OR 1.3 (95% CI 0.7 to 2.6) P = 0.45</p> <p><u>Apgar score less than 7 at 5 minutes</u></p> <p>Terminal deceleration n = 4/951 (0.4%)</p> <p>Not terminal deceleration n = 51/4437 (1.2%) Adjusted* OR 0.4 (95% CI 0.1 to 1.1) P = 0.05</p> <p><u>Special care or NICU admission</u></p> <p>Terminal deceleration n = 42/951 (4.4%)</p> <p>Not terminal deceleration n = 228/4437 (5.2%) Adjusted* OR 0.8 (95% CI 0.6 to 1.2) P = 0.35</p> <p><u>Abruption composite</u></p> <p>Terminal deceleration n = 10/951 (1.1%)</p> <p>Not terminal deceleration n = 18/4437 (0.4%)</p>	<p>assumed that duration of terminal deceleration was until birth</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Between 2004 and 2008</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>fetal monitoring (EFM) recording during the 30 minutes before birth (less than 10 minutes of EFM during the 30 minutes before birth).</p>		<p>College of Obstetricians and Gynecologists three-tiered category system. Terminal deceleration, defined as a prolonged deceleration (15 bpm or more below baseline for 120 seconds (2 min) or more and fewer than 10 minutes) or bradycardia (&lt; 110 bpm for 10 minutes or more). The comparison made between women who had a terminal deceleration and those who did not.</p> <p>Interval interobserver reliability was performed. For presence of terminal decelerations, kappa</p>	<p>Adjusted* OR 2.6 (95% CI 1.2 to 5.6) P = 0.2</p> <p><b><u>Terminal deceleration characteristics by acidemia:</u></b>  <u>Number of babies born with acidemia.</u> n = 12/951 (1.3%)  <u>Number of babies born with no acidemia.</u> n = 939/951 (1.3%)</p> <p><u>Median time to birth (min SD)</u>  Acidemia 6.7 (SD 3.7 to 12.7)  No academia 3.2 (SD 2.5 to 4.6)  P&lt;.01</p> <p>For every additional 120 seconds of duration of the terminal deceleration beyond the first 120 seconds, there was a corresponding decrease in arterial umbilical cord pH level by 0.042 (95% CI 0.040 to 0.048; P&lt;.01). However, terminal deceleration characteristics, such as median or greatest depth and variability within the nadir, were not associated with risk of acidemia</p> <p><b><u>Bradycardia and terminal deceleration</u></b>  <u>Risk associated with Bradycardia among women with terminal deceleration:</u>  <u>Bradycardia duration of 10 minutes or more</u> n = 31/951  <u>Bradycardia duration of &lt; 10 minutes</u> n = 930/951  <u>Risk of acidemia (pH level of 7.10 or less):</u>  Bradycardia duration of 10 minutes or more n = 4/31 (12.9%)  Bradycardia duration of &lt; 10 minutes n = 8/920 (0.9%)  Adjusted OR 18.6 (5.0 to 68.9)  P &lt; 0.01  <u>Risk of acidemia (pH level of 7.05 or less):</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>coefficient was consistently more than 0.9. Detailed maternal and pregnancy data including obstetric history, pregnancy course and complications, medication exposure and acute events (including placental abruption, umbilical cord prolapse, and uterine rupture), physical examination, anesthesia type, delivery, and neonatal outcomes were also extracted. Use of internal monitors for fetal heart rate monitoring and contractions and umbilical cord gas arterial pH level, as well as CO<sub>2</sub> and</p>	<p>Bradycardia duration of 10 minutes or more n = 2/31 (6.5%) Bradycardia duration of &lt; 10 minutes n = 2/920 (0.2%) Adjusted* OR 46.0 (5.7 to 373.0) P &lt; 0.01</p> <p><u>Apgar score &lt; 7 at 5 min:</u> Bradycardia duration of 10 minutes or more n = 2/31 (6.5%) Bradycardia duration of &lt; 10 minutes n = 2/920 (0.2%) Adjusted* OR 67.0 (8.4 to 536.6) P &lt; 0.01</p> <p><u>Special care and NICU admission:</u> Bradycardia duration of 10 minutes or more n = 3/31 (10%) Bradycardia duration of &lt; 10 minutes n = 8/920 (0.9%) Adjusted* OR 11.4 (3.2 to 40.7) P &lt; 0.01</p> <p>* Adjusted for nulliparity Presence of bradycardia (10 minutes or more) was poorly predictive of acidemia, with a sensitivity of 33.3%, a specificity of 97.0%, and a positive predictive value of only 12.9%.</p> <p><u>Duration of terminal deceleration</u></p> <p><u>Predictive value of duration of terminal deceleration beyond 2 minutes for acidemia (pH level of 7.10 or less)</u> AUC (area under the curve) 0.78 (95% CI 0.60–0.94)</p> <p><u>Predictive value of duration of terminal deceleration cut-off of 4 minutes or more for acidemia (pH level of 7.10 or less)</u></p> <p>Sensitivity: 75.0% (95% CI 74.2 to 76.3%) Specificity: 64.0% (95% CI 62.8–65.1%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>base excess, also were recorded. The primary outcome was acidemia, defined as arterial umbilical cord gas pH level of 7.10 or less. Secondary outcomes included arterial umbilical cord gas pH level 7.05 or less, base excess more than -8, metabolic acidemia (pH level 7.10 or less and base excess more than -8), admission to the neonatal intensive care unit (level IV) or admission to the special care unit (level II), and Apgar score less than 7 at 5 minutes.</p> <p><u>Analysis:</u> For continuous variables Student <i>t</i> tests</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>and Mann-Whitney <i>U</i> tests were used and <math>\chi^2</math> and for dichotomous variables Fisher exact tests were used as appropriate. Stratified analyses were performed to identify potentially confounding factors, which were considered in multivariable analyses.</p> <p>To refine estimates of association between terminal decelerations and acidemia by eliminating nonsignificant factors, multivariable logistic regression was performed.</p> <p>To explore the risk of acidemia and other adverse</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>outcomes among women with terminal bradycardia a secondary analysis was performed. Linear regression was then used to estimate the incremental association between increasing terminal deceleration duration beyond 2 minutes and decreasing arterial umbilical cord pH level. To estimate the predictive ability of terminal deceleration duration and risk of acidemia, Receiver-operator characteristic curve analysis was used. STATA 10 special edition</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments															
			was used for the all analysis.																	
<p><b>Full citation</b></p> <p>Graham,E.M., Adami,R.R., McKenney,S.L., Jennings,J.M., Burd,I., Witter,F.R., Diagnostic accuracy of fetal heart rate monitoring in the identification of neonatal encephalopathy, Obstetrics and Gynecology, 124, 507-513, 2014</p> <p><b>Ref Id</b></p> <p>346212</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Case-control study</p>	<p><b>Sample size</b></p> <p>N=39 cases (neonates treated with whole-body hypothermia for suspected hypoxic-ischaemic encephalopathy) N=78 controls (matched to each neonate in the case group in a two-to-one fashion using the two subsequent births in the same hospital matched by gestational age within 1 weeks and mode of birth)</p> <p><b>Characteristics</b></p> <p>There was no difference in the following characteristics in the case and control groups: maternal age, parity, race, receiving oxytocin, pre-eclampsia, intrauterine growth restriction, oligohydramnios, abruption, histologic chorioamnionitis, histologic funisities,</p>	<p><b>Interventions</b></p> <p>Non-computer-assisted interpretation of the last hour of EFM tracing before birth</p>	<p><b>Details</b></p> <p>The last 1 hour of EFM tracing was reviewed independently by three obstetricians blinded to outcome using the National Institute of Child Health and Human Development and the American College of Obstetricians and Gynecologists three-tiered category system and definitions. Each reviewer assessed the last hour of tracing and assigned a category based on the most non-reassuring portion of the tracing and the</p>	<p><b>Results</b></p> <p><b>Odds ratio* (OR) with 95% CI of the following EFM features in the case group. (Last 1 hour tracing before birth.)</b>                      Reactive: OR 0.50 (0.22-1.12)                      Late decelerations: OR 1.10 (1.00-1.21)                      Early decelerations: OR 0.58 (0.35-0.94)                      Debt 30: 1.00 (1.00-1.00)                      Debt 60: 1.00 (1.00-1.00)                      *Adjusted for chorioamnionitis</p> <p><b>Diagnostic accuracy (95% CI) of the following EFM features or classifications to detect cases with whole-body hypothermia treatment for suspected hypoxic-ischaemic encephalopathy</b></p> <table border="1"> <thead> <tr> <th></th> <th>Suspected hypoxic-ischaemic encephalopathy</th> <th>No suspected encephalopathy</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Early decelerations</td> <td>NR</td> <td>NR</td> <td>NR</td> </tr> </tbody> </table>		Suspected hypoxic-ischaemic encephalopathy	No suspected encephalopathy	Totals	Early decelerations	NR	NR	NR	<p><b>Limitations</b></p> <p><b>Assessed with QUADAS-2 (for measures of diagnostic accuracy):</b>  <b>Patient selection:</b> High risk (case-control design)  <b>Index test(s)</b> (The index test in the study is the interpretation of the last hour of EFM tracing prior to birth): Low risk (3 reviewers assessed the trace and they were blinded to the results of the reference standard)  <b>Reference standard</b> (The reference standard in the study is the assessment of suspected hypoxic-ischaemic encephalopathy within 6 hours of birth): Low risk (the reference standard is likely to correctly classify the target condition)  <b>Flow and timing:</b> High risk (index test was performed before birth, reference tests were performed after birth, this means that differences in outcomes may be due to events in-between the two tests)  <b>Overall risk of bias: very serious risk of bias</b>  <b>Assessed with NICE 2012 guideline manual checklist for prognostic studies (for ORs):</b></p> <table border="1"> <tr> <td rowspan="2"><b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit</b></td> <td><u>Yes</u></td> <td>No</td> <td>Unclear</td> </tr> <tr> <td></td> <td></td> <td></td> </tr> </table>	<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit</b>	<u>Yes</u>	No	Unclear			
	Suspected hypoxic-ischaemic encephalopathy	No suspected encephalopathy	Totals																	
Early decelerations	NR	NR	NR																	
<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit</b>	<u>Yes</u>	No	Unclear																	

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments								
<p><b>Aim of the study</b></p> <p>To estimate the diagnostic accuracy of electronic heart rate abnormalities in the identification of neonates with encephalopathy treated with whole-body hypothermia</p> <p><b>Study dates</b></p> <p>Between January 1, 2007 and July 1, 2013</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p>histologic placental infarcts, birthweight, gender.</p> <p>The case group more often had clinical chorioamnionitis, nonreassuring fetal heart rate, and meconium, 1-minute Apgar score of less than 7, 5-minute Apgar score of less than 7, cord pH &lt;7.0 or base deficit &gt;12mM, respiratory distress, positive blood cultures, seizures and longer stay length of stay at hospital</p> <p><b>Inclusion criteria</b></p> <p>All neonates born with suspected hypoxic-ischaemic encephalopathy at two hospitals and treated with whole-body hypothermia within 6 hours of birth during the 6.5-year period from January 1, 2007 to July 1, 2013. Neonates were eligible for treatment with whole-body hypothermia if moderate to severe encephalopathy was</p>		<p>final category was assigned based on consensus among the reviewers.</p> <p>Each reviewer recorded the fetal heart rate (FHR), time with FHR greater than 160 bpm (tachycardia), or less than 110 bpm (bradycardia), number of accelerations, reactivity, total number of decelerations, and number of late, variable, or early decelerations. Reactivity was defined as the presence of at least two FHR accelerations that peaked (but did not necessarily remain) at least 15 bpm above the baseline and lasted 15 seconds during a 20-minute</p>	No early decelerations	NR	NR	NR	Totals	NR	NR	NR	<p><b>potential bias to the results</b></p> <p>All neonates born at two hospitals with suspected hypoxic-ischaemic encephalopathy treated with whole-body hypothermia within 6 hours of birth during the 6.5-year period from January 1, 2007 to July 1, 2013 were included. Neonates in the control group were matched to each neonate in the case group in a two-to-one fashion using the subsequent two deliveries in the same hospital matched by</p>				
				<p>Sensitivity 23.1% (11.7-39.7%)                  Specificity 94.9% (86.7-98.3%)                  Positive likelihood ratio** 4.53                  Negative likelihood ratio** 0.81</p> <p><u>Category III (versus category I)</u></p>					Suspected encephalopathy	No suspected encephalopathy	Totals					
				Category III	5	1	6									
				Category I (normal)	4	7	11									
				Totals	9	8	17									
				<p>Sensitivity** 55.6% (22.7-84.7%)                  Specificity** 87.5% (46.7-99.3%)                  Positive likelihood ratio** 4.44 (0.65-30.44)</p>												



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																																											
	<p>present at birth (manifested by lethargy, stupor, coma, decreased or no activity, distal flexion, complete extension, decerebrate posture, hypotonia or flaccidity, abnormal primitive reflexes, bradycardia, periodic breathing, apnoea, or seizures) and had a cord gas or early neonatal gas at less than 1 hour with pH 7.0 or less or base deficit greater than 16 mM. They were also eligible if the cord or early neonatal gas at less than 1 hour showed pH 7.01-7.15 and base deficit 10-15.9 mM if moderate to severe encephalopathy was present with evidence of an acute sentinel event, 10-minute Apgar score less than 5, or there was need for assisted ventilation initiated at birth with continuation for at least 10 minutes.</p> <p>Neonates in the control group were matched to each neonate in the case group in a two-to-one fashion using the</p>		<p>period that occurred any time during the last hour before birth. Variability was classified as absent (undetectable), minimal (amplitude range 5 bpm or less), moderate (amplitude range from 6-25 bpm) or marked (amplitude range greater than 25 bpm). Absent or minimal were considered as decreased variability. The number of prolonged decelerations lasting 2-10 minutes was recorded as well as the nadir and length of the most severe prolonged deceleration. Severe variable decelerations</p>	<p>Negative likelihood ratio** 0.51 (0.24-1.09)</p> <p><u>Category II (versus category I)</u></p> <table border="1" data-bbox="1021 360 1491 967"> <tr> <td data-bbox="1021 360 1111 560">a</td> <td data-bbox="1111 360 1267 560">Suspected encephalopathy</td> <td data-bbox="1267 360 1424 560">No suspected encephalopathy</td> <td data-bbox="1424 360 1491 560">Totals</td> </tr> <tr> <td data-bbox="1021 560 1111 679">Category II</td> <td data-bbox="1111 560 1267 679">30</td> <td data-bbox="1267 560 1424 679">70</td> <td data-bbox="1424 560 1491 679">100</td> </tr> <tr> <td data-bbox="1021 679 1111 879">Category I (normal)</td> <td data-bbox="1111 679 1267 879">4</td> <td data-bbox="1267 679 1424 879">7</td> <td data-bbox="1424 679 1491 879">11</td> </tr> <tr> <td data-bbox="1021 879 1111 967">Totals</td> <td data-bbox="1111 879 1267 967">34</td> <td data-bbox="1267 879 1424 967">77</td> <td data-bbox="1424 879 1491 967">111</td> </tr> </table> <p>Sensitivity** 88.2% (71.6-96.2%)                      Specificity** 9.1% (4.0-18.4%)                      Positive likelihood ratio** 0.97 (0.84-1.12)                      Negative likelihood ratio** 1.29 (0.40-4.19)</p> <p><u>Decreased variability</u></p> <table border="1" data-bbox="1021 1142 1491 1342"> <tr> <td data-bbox="1021 1142 1111 1342"></td> <td data-bbox="1111 1142 1267 1342">Suspected encephalopathy</td> <td data-bbox="1267 1142 1424 1342">No suspected encephalopathy</td> <td data-bbox="1424 1142 1491 1342">Totals</td> </tr> </table>	a	Suspected encephalopathy	No suspected encephalopathy	Totals	Category II	30	70	100	Category I (normal)	4	7	11	Totals	34	77	111		Suspected encephalopathy	No suspected encephalopathy	Totals	<table border="1" data-bbox="1514 272 2047 1377"> <tr> <td data-bbox="1514 272 1760 392">gestational age within 1 weeks and mode of birth</td> <td data-bbox="1760 272 1850 392"></td> <td data-bbox="1850 272 1962 392"></td> <td data-bbox="1962 272 2047 392"></td> </tr> <tr> <td data-bbox="1514 392 1760 847"><b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately represent the sample), sufficient to limit potential bias</b></td> <td data-bbox="1760 392 1850 847">Yes</td> <td data-bbox="1850 392 1962 847">No</td> <td data-bbox="1962 392 2047 847">Unclear</td> </tr> <tr> <td data-bbox="1514 847 1760 863">N/A</td> <td data-bbox="1760 847 1850 863"></td> <td data-bbox="1850 847 1962 863"></td> <td data-bbox="1962 847 2047 863"></td> </tr> <tr> <td data-bbox="1514 863 1760 1377"><b>The prognostic factor of interest is adequately measured in study participants, sufficient to limit potential bias</b></td> <td data-bbox="1760 863 1850 1377">Yes</td> <td data-bbox="1850 863 1962 1377">No</td> <td data-bbox="1962 863 2047 1377">Unclear</td> </tr> <tr> <td data-bbox="1514 1377 1760 1385">3 reviewers assessed the trace and they were blinded to the results of the reference standard</td> <td data-bbox="1760 1377 1850 1385"></td> <td data-bbox="1850 1377 1962 1385"></td> <td data-bbox="1962 1377 2047 1385"></td> </tr> </table>				gestational age within 1 weeks and mode of birth				<b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately represent the sample), sufficient to limit potential bias</b>	Yes	No	Unclear	N/A				<b>The prognostic factor of interest is adequately measured in study participants, sufficient to limit potential bias</b>	Yes	No	Unclear	3 reviewers assessed the trace and they were blinded to the results of the reference standard			
a	Suspected encephalopathy	No suspected encephalopathy	Totals																																													
Category II	30	70	100																																													
Category I (normal)	4	7	11																																													
Totals	34	77	111																																													
	Suspected encephalopathy	No suspected encephalopathy	Totals																																													
gestational age within 1 weeks and mode of birth																																																
<b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately represent the sample), sufficient to limit potential bias</b>	Yes	No	Unclear																																													
N/A																																																
<b>The prognostic factor of interest is adequately measured in study participants, sufficient to limit potential bias</b>	Yes	No	Unclear																																													
3 reviewers assessed the trace and they were blinded to the results of the reference standard																																																

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments											
	<p>two subsequent births in the same hospital matched by gestational age to within 1 week and mode of birth).</p> <p><b>Exclusion criteria</b></p> <p>Exclusion criteria for whole-body hypothermia treatment included greater than 6 hours of life, gestational age less than 35 weeks, severe growth restriction (birthweight less than 1800 g), major congenital anomaly, severe persistent pulmonary hypertension with anticipated need for extracorporeal membrane oxygenation, coagulopathy with active bleeding, and suspected sepsis with severe hemodynamic compromise requiring large doses of pressors</p>		<p>were those with a drop to less than 70 bpm or lasting greater than 60 seconds. The number of contractions in the last hour before birth were counted, and the ratio of late decelerations per contractions and variable decelerations per contractions were expressed as a percentage. Total deceleration area was calculated as the sum of the area within all decelerations in the final 30 minutes (debt30) and final 60 minutes (debt60) of the tracing as a measure of both quantity and severity.</p>	Decreased variability	13	15	18	No decreased variability	26	63	89	Totals	39	78	117	<p>using the NICHD classification. The reviewers were an obstetric resident (RRA), and two maternal-fetal medicine attendings, all of whom had passed the required EFM course. Categorical EFM tracing parameters were determined by consensus among the three reviewers, and continuous parameters were averaged.</p>			
				<p>Sensitivity** 33.3% (19.6-50.3%)                  Specificity** 80.8% (70.0-88.5%)                  Positive likelihood ratio** 1.73 (0.92-3.27)                  Negative likelihood ratio** 0.83 (0.66-1.04)</p>															
				<p><u>Reactivity</u></p> <table border="1" data-bbox="1021 975 1491 1174"> <tr> <td></td> <td>Suspected encephalopathy</td> <td>No suspected encephalopathy</td> <td>Totals</td> </tr> <tr> <td>Reactivity</td> <td>16</td> <td>48</td> <td>64</td> </tr> </table>					Suspected encephalopathy	No suspected encephalopathy	Totals	Reactivity	16	48	64				
	Suspected encephalopathy	No suspected encephalopathy	Totals																
Reactivity	16	48	64																
				<p><b>The outcome of interest is adequately measured in study participants, sufficient to limit potential bias</b></p>				Yes	No	Unclear									

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments			
			<p>The area within each deceleration was approximated as one-half (width in seconds x depth in bpm). Multiple variable logistic regression models were used to determine the diagnostic accuracy of EFM parameters in the identification of neonates with encephalopathy treated with whole-body hypothermia. Variables significant at a p-value of &lt;0.10 in bivariate analyses were used in the multiple variable regression</p>	No reactivity	23	30	53	<p>Clear and comprehensive criteria to assess moderate to severe encephalopathy at birth and eligibility for treatment with whole-body hypothermia due to suspected encephalopathy</p>			
Totals	39	78	117	<p>Sensitivity** 41.0% (26.0-57.8%)                      Specificity** 38.5% (27.9-50.2%)                      Positive likelihood ratio** 0.67 (0.44-1.01)                      Negative likelihood ratio** 1.53 (1.13-2.07)</p> <p>**Calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a></p>					<p><b>Important potential confounders are appropriately accounted for, limiting potential bias with respect to the prognostic factor of interest</b>                      ORs were adjusted for the presence of clinical chorioamnionitis, however not for other factors such</p>	Yes	<b>No</b>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments			
					as demographic characteristics or presence of meconium			
					<p><b>The statistical analysis is appropriate for the design of the study, limiting potential for the presentation of invalid results</b></p> <p>Multivariable logistic regression was appropriately conducted.</p>	Yes	No	Unclear
					<p><b>Risk of bias:</b></p>	No serious risk of bias	<p><b>Serious risk of bias</b></p>	Very serious risk of bias
					<p><b>Other information</b></p>			
Full citation	Sample size	Interventions	Details	Results	Limitations			

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
<p>Holzmann, M., Wretler, S., Cnattingius, S., Nordstrom, L., <i>Cardiotocography patterns and risk of intrapartum fetal acidemia</i>, Journal of Perinatal Medicine, 43, 473-479, 2015</p> <p><b>Ref Id</b> 446285</p> <p><b>Country/ies where the study was carried out</b> Sweden</p> <p><b>Study type</b> Prospective observational cohort study</p> <p><b>Aim of the study</b> To identify cardiotocography patterns associated with increased risk of intrapartum fetal acidemia</p>	<p>N= 1070 women in labour, 2134 fetal blood samples (FBSs)</p> <p><b>Characteristics</b></p> <p>Women who underwent FBS due to a CTG trace that was assessed as 'non-reassuring' by the attending physician during labour at Karolinska University Hospital, Stockholm. Median maternal age: 31 (range: 15 to 47) Median gestational age (weeks+days): 40+3 (range:34+1 to 42+4) Thick meconium: 75 (7.0%) Mode of birth: Spontaneous: 421 (39.4%); Ventouse: 349 (32.6%); Caesarean section: 300 (28.0%)</p> <p><b>Inclusion criteria</b></p> <p>Singleton pregnancy, &gt;=34 weeks of gestation, cephalic presentation, and indication for FBS according to the attending doctor</p>	<p><b>Intervention 1</b> Interpretation of cardiotocography tracing for the last 60 minutes prior to first FBS</p> <p><b>Intervention 2</b> Interpretation of cardiotocography tracing for the last 60 minutes prior to last FBS</p>	<p>All women had an admission CTG; with a normal test result and the woman being considered to be at low risk, intermittent CTG monitoring every 2 hours was recommended. Women considered to be at high risk, having epidural analgesia or oxytocin augmentation had continuous CTG monitoring. CTG interpretation followed the guidelines of the Swedish Society of Obstetrics and Gynecology (SFOG), based on the international classification system of the International Federation of Gynecology</p>	<p><b>Diagnostic accuracy (95% CI) of the following EFM features to detect fetal lactacidaemia (lactate&gt;4.8 mmol/l) at first FBS (negative test result is 'normal baseline and variability')</b></p> <p><u>Reduced variability</u></p> <table border="1"> <thead> <tr> <th></th> <th>Lactate &gt;4.8 mmol/l</th> <th>Lactate ≤4.8 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Reduced variability</td> <td>4</td> <td>150</td> <td>154</td> </tr> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> <tr> <td>Totals</td> <td>10</td> <td>386</td> <td>396</td> </tr> </tbody> </table> <p>Sensitivity 40.00% (13.69% to 72.63%) Specificity 61.14% (56.06% to 66.00%) Positive likelihood ratio 1.03 (0.48 to 2.22) Negative likelihood ratio 0.98 (0.59 to 1.63)</p> <p><u>Absent variability</u></p> <table border="1"> <thead> <tr> <th></th> <th>Lactate &gt;4.8 mmol/l</th> <th>Lactate ≤4.8 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Reduced variability	4	150	154	Normal baseline and variability	6	236	242	Totals	10	386	396		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals					<p><b>Assessed with QUADAS-2:</b></p> <p><b>Patient selection:</b> High risk (All women included in the study had received FBS, and FBS was only recommended by the attending physician if the CTG was non-reassuring. Therefore, even if some of the CTGs were later classified as normal when re-evaluated for the study by a senior obstetrician, they may have not been representative of the 'average' normal CTG)</p> <p><b>Index test:</b> High risk (Even if the senior obstetrician interpreting the CTGs was blinded to the outcome, it is known that FHR trace interpretation is difficult and can be subjective and therefore introduce bias; other studies rely on consensus across multiple reviewers for trace interpretation)</p> <p><b>Reference standard:</b> Low risk (For the last FBS, an exclusion criterion was active pushing prior to sampling because active pushing is known to increase the lactate concentration)</p> <p><b>Flow and timing:</b> Low risk (CTG trace interpretation was applied to the last 60 minutes prior to each FBS; both the index test and the reference standard were applied before birth)</p> <p><b>Overall risk of bias:</b> Very serious</p> <p><b>Other information</b></p>
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Reduced variability	4	150	154																										
Normal baseline and variability	6	236	242																										
Totals	10	386	396																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments			
<p><b>Study dates</b> February 2009-February 2011</p> <p><b>Source of funding</b> Not reported</p>	<p><b>Exclusion criteria</b> For the last sample in a particular woman, an exclusion criterion was active pushing prior to sampling</p>		<p>and Obstetrics (FIGO) from 1987. The attending physician decided upon FBS if the CTG trace was visually interpreted as non=reassuring . FBS was performed according to clinical routine; 5 µl of fetal scalp blood was collected after wiping dry from amniotic fluid and applying silicone gel. Analysis was done at the bedside using Lactate Pro™ (KDK Corp., Kyoto, Japan), calibrated every 50<sup>th</sup> analysis. Half of the women had more than one FBS. The study authors, therefore, included results for both the first</p>	Absent variability	4	28	32				
				Normal baseline and variability	6	236	242				
				Totals	10	264	274				
				<p>Sensitivity 40.00% (13.69% to 72.63%) Specificity 89.39% (84.88% to 92.72%) Positive likelihood ratio 3.77 (1.63 to 8.70) Negative likelihood ratio 0.67 (0.40 to 1.11)</p>							
				<p><u>Increased variability</u></p>							
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals				
				Increased variability	2	8	10				
				Normal baseline and variability	6	236	242				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
			<p>sample, including the total population that met the inclusion criteria, and included results from the last sample unless this failed to meet the inclusion criteria. A senior obstetrician (LN), blinded to the lactate concentration at sampling, interpreted all CTG tracings with focus on the last 60 minutes prior to each FBS. The study authors documented baseline FHR, variability, accelerations, type of decelerations, and duration of CTG pattern prior to FBS. Definitions published by FIGO were used, i.e. FHR (normal) 110–</p>	<table border="1" data-bbox="1021 276 1487 349"> <tr> <td>Totals</td> <td>8</td> <td>244</td> <td>252</td> </tr> </table> <p>Sensitivity 25.00 % (4.45% to 64.42%)                      Specificity 96.72% (93.40% to 98.47%)                      Positive likelihood ratio 7.63 (1.92 to 30.31)                      Negative likelihood ratio 0.78 (0.52 to 1.16)</p> <p><u>Bradycardic episode</u></p> <table border="1" data-bbox="1021 528 1487 724"> <tr> <td></td> <td>Lactate &gt;4.8 mmol/l</td> <td>Lactate ≤4.8 mmol/l</td> <td>Totals</td> </tr> <tr> <td>Bradycardic episode</td> <td>10</td> <td>36</td> <td>46</td> </tr> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> <tr> <td>Totals</td> <td>16</td> <td>272</td> <td>288</td> </tr> </table> <p>Sensitivity 62.50 % (35.87% to 83.72%)                      Specificity 86.76% (82.02% to 90.44%)                      Positive likelihood ratio 4.72 (2.90 to 7.68)                      Negative likelihood ratio 0.43 (0.23 to 0.81)</p> <p><u>Tachycardia</u></p>	Totals	8	244	252		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Bradycardic episode	10	36	46	Normal baseline and variability	6	236	242	Totals	16	272	288	
Totals	8	244	252																						
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																						
Bradycardic episode	10	36	46																						
Normal baseline and variability	6	236	242																						
Totals	16	272	288																						

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
			<p>150 beats per minute (bpm), bradycardia &lt;110 bpm, and tachycardia &gt;150 bpm. Variability: normal 5–25 bpm, reduced: 2–4 bpm, absent: &lt;2 bpm, and increased: &gt;25 bpm, accelerations: transient increase in FHR of ≥15 bpm for ≥15 seconds, and decelerations: transient episodes of slowing of FHR below baseline level of ≥15 bpm lasting ≥15 seconds. Severe variable decelerations were defined as having a variable shape, an abrupt fall from baseline FHR to nadir of deceleration, and a duration &gt;60 seconds.</p>	<table border="1" data-bbox="1021 276 1485 882"> <thead> <tr> <th data-bbox="1021 276 1167 475"></th> <th data-bbox="1173 276 1261 475">Lactate &gt;4.8 mmol/l</th> <th data-bbox="1267 276 1413 475">Lactate ≤4.8 mmol/l</th> <th data-bbox="1420 276 1485 475">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 480 1167 600">Tachycardia</td> <td data-bbox="1173 480 1261 600">10</td> <td data-bbox="1267 480 1413 600">114</td> <td data-bbox="1420 480 1485 600">124</td> </tr> <tr> <td data-bbox="1021 604 1167 804">Normal baseline and variability</td> <td data-bbox="1173 604 1261 804">6</td> <td data-bbox="1267 604 1413 804">236</td> <td data-bbox="1420 604 1485 804">242</td> </tr> <tr> <td data-bbox="1021 809 1167 882">Totals</td> <td data-bbox="1173 809 1261 882">16</td> <td data-bbox="1267 809 1413 882">350</td> <td data-bbox="1420 809 1485 882">366</td> </tr> </tbody> </table> <p data-bbox="1021 887 1485 1002">Sensitivity 62.50% (35.87% to 83.72%) Specificity 67.43% (62.21% to 72.26%) Positive likelihood ratio 1.92 (1.28 to 2.89) Negative likelihood ratio 0.56 (0.29 to 1.05)</p> <p data-bbox="1021 1023 1485 1054"><u>Tachycardia+reduced variability</u></p> <table border="1" data-bbox="1021 1059 1485 1299"> <thead> <tr> <th data-bbox="1021 1059 1223 1299"></th> <th data-bbox="1229 1059 1301 1299">Lactate &gt;4.8 mmol/l</th> <th data-bbox="1308 1059 1424 1299">Lactate ≤4.8 mmol/l</th> <th data-bbox="1431 1059 1485 1299">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 1059 1223 1299"></td> <td data-bbox="1229 1059 1301 1299"></td> <td data-bbox="1308 1059 1424 1299"></td> <td data-bbox="1431 1059 1485 1299"></td> </tr> </tbody> </table>		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Tachycardia	10	114	124	Normal baseline and variability	6	236	242	Totals	16	350	366		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals					
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Tachycardia	10	114	124																										
Normal baseline and variability	6	236	242																										
Totals	16	350	366																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
			<p>Late decelerations were defined as start of deceleration after a peak of contraction, uniform shape, and gradual fall to nadir of deceleration. Bradycardic episodes were defined as baseline FHR &lt;110 bpm for &gt;3 minutes occurring within 30 minutes before sampling, including prolonged decelerations lasting &lt;10 minutes and bradycardia for &gt;10 minutes. Simple variable decelerations (duration &lt;60 seconds) and early decelerations (starting before peak of contraction) were referred to the normal group</p>	<table border="1" data-bbox="1021 272 1491 635"> <tr> <td>Tachycardia+reduced variability</td> <td>9</td> <td>140</td> <td>149</td> </tr> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> <tr> <td>Totals</td> <td>15</td> <td>376</td> <td>391</td> </tr> </table> <p>Sensitivity 60.00% (32.89% to 82.54%)                      Specificity 62.76% (57.64% to 67.63%)                      Positive likelihood ratio 1.61 (1.04 to 2.49)                      Negative likelihood ratio 0.64 (0.34 to 1.19)</p> <p><u>Severe variable decelerations</u></p> <table border="1" data-bbox="1021 810 1491 1310"> <tr> <td></td> <td>Lactate &gt;4.8 mmol/l</td> <td>Lactate ≤4.8 mmol/l</td> <td>Totals</td> </tr> <tr> <td>Severe variable decelerations</td> <td>18</td> <td>109</td> <td>127</td> </tr> <tr> <td>Normal baseline</td> <td>6</td> <td>236</td> <td>242</td> </tr> </table>	Tachycardia+reduced variability	9	140	149	Normal baseline and variability	6	236	242	Totals	15	376	391		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Severe variable decelerations	18	109	127	Normal baseline	6	236	242	
Tachycardia+reduced variability	9	140	149																										
Normal baseline and variability	6	236	242																										
Totals	15	376	391																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Severe variable decelerations	18	109	127																										
Normal baseline	6	236	242																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 272 1487 389"> <tr> <td>and variability</td> <td></td> <td></td> <td></td> </tr> </table> <table border="1" data-bbox="1021 394 1487 469"> <tr> <td>Totals</td> <td>24</td> <td>345</td> <td>369</td> </tr> </table> <p data-bbox="1012 475 1496 587">Sensitivity 75.00 % (52.95% to 89.40%) Specificity 68.41 % (63.17% to 73.22%) Positive likelihood ratio 2.37 (1.80 to 3.14) Negative likelihood ratio 0.37 (0.18 to 0.73)</p> <p data-bbox="1012 616 1211 639"><u>Late decelerations</u></p> <table border="1" data-bbox="1021 644 1487 842"> <tr> <td></td> <td>Lactate &gt;4.8 mmol/l</td> <td>Lactate ≤4.8 mmol/l</td> <td>Totals</td> </tr> </table> <table border="1" data-bbox="1021 847 1487 1007"> <tr> <td>Late decelerations</td> <td>8</td> <td>50</td> <td>58</td> </tr> </table> <table border="1" data-bbox="1021 1011 1487 1209"> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> </table> <table border="1" data-bbox="1021 1214 1487 1289"> <tr> <td>Totals</td> <td>14</td> <td>286</td> <td>300</td> </tr> </table> <p data-bbox="1012 1295 1496 1378">Sensitivity 57.14% (29.65% to 81.19%) Specificity 82.52% (77.50% to 86.64%) Positive likelihood ratio 3.27 (1.95 to 5.49)</p>	and variability				Totals	24	345	369		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Late decelerations	8	50	58	Normal baseline and variability	6	236	242	Totals	14	286	300	
and variability																													
Totals	24	345	369																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Late decelerations	8	50	58																										
Normal baseline and variability	6	236	242																										
Totals	14	286	300																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<p>Negative likelihood ratio 0.52 (0.28 to 0.95)</p> <p><u>Severe variable decelerations+reduced variability</u></p> <table border="1" data-bbox="1021 389 1491 995"> <thead> <tr> <th data-bbox="1021 389 1234 632"></th> <th data-bbox="1234 389 1308 632">Lactate &gt;4.8 mmol/l</th> <th data-bbox="1308 389 1429 632">Lactate ≤4.8 mmol/l</th> <th data-bbox="1429 389 1491 632">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 632 1234 794">Severe variable decelerations+reduced variability</td> <td data-bbox="1234 632 1308 794">4</td> <td data-bbox="1308 632 1429 794">24</td> <td data-bbox="1429 632 1491 794">28</td> </tr> <tr> <td data-bbox="1021 794 1234 916">Normal baseline and variability</td> <td data-bbox="1234 794 1308 916">6</td> <td data-bbox="1308 794 1429 916">236</td> <td data-bbox="1429 794 1491 916">242</td> </tr> <tr> <td data-bbox="1021 916 1234 995">Totals</td> <td data-bbox="1234 916 1308 995">10</td> <td data-bbox="1308 916 1429 995">260</td> <td data-bbox="1429 916 1491 995">270</td> </tr> </tbody> </table> <p>Sensitivity 40.00% (13.69% to 72.63%)                      Specificity 90.77% (86.41% to 93.88%)                      Positive likelihood ratio 4.33 (1.85 to 10.13)                      Negative likelihood ratio 0.66 (0.40 to 1.10)</p> <p><u>Late decelerations+reduced variability</u></p> <table border="1" data-bbox="1021 1171 1491 1326"> <thead> <tr> <th data-bbox="1021 1171 1234 1326"></th> <th data-bbox="1234 1171 1308 1326">Lactate &gt;4.8 mmol/l</th> <th data-bbox="1308 1171 1429 1326">Lactate ≤4.8 mmol/l</th> <th data-bbox="1429 1171 1491 1326">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 1171 1234 1326"></td> <td data-bbox="1234 1171 1308 1326"></td> <td data-bbox="1308 1171 1429 1326"></td> <td data-bbox="1429 1171 1491 1326"></td> </tr> </tbody> </table>		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Severe variable decelerations+reduced variability	4	24	28	Normal baseline and variability	6	236	242	Totals	10	260	270		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals					
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Severe variable decelerations+reduced variability	4	24	28																										
Normal baseline and variability	6	236	242																										
Totals	10	260	270																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
					mmo l/l			
				Late decelerations+reduced variability	3	22	25	
				Normal baseline and variability	6	236	242	
				Totals	9	258	267	
				Sensitivity 33.33% (9.04% to 69.08%) Specificity 91.47% (87.20% to 94.46%) Positive likelihood ratio 3.91 (1.43 to 10.70) Negative likelihood ratio 0.73 (0.46 to 1.16)				
				<u>Severe variable decelerations+tachycardia</u>				
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	
				Severe variable decelerations+tachycardia	8	24	32	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 272 1487 389"> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> </table> <table border="1" data-bbox="1021 394 1487 469"> <tr> <td>Totals</td> <td>14</td> <td>260</td> <td>274</td> </tr> </table> <p data-bbox="1012 475 1496 587">Sensitivity 57.14% (29.65% to 81.19%) Specificity 90.77% (86.41% to 93.88%) Positive likelihood ratio 6.19 (3.42 to 11.20) Negative likelihood ratio 0.47 (0.26 to 0.87)</p> <p data-bbox="1012 616 1350 639"><u>Late decelerations+tachycardia</u></p> <table border="1" data-bbox="1021 644 1487 887"> <tr> <td></td> <td>Lactate &gt;4.8 mmol/l</td> <td>Lactate ≤4.8 mmol/l</td> <td>Totals</td> </tr> <tr> <td>Late decelerations+tachycardia</td> <td>6</td> <td>24</td> <td>30</td> </tr> <tr> <td>Normal baseline and variability</td> <td>6</td> <td>236</td> <td>242</td> </tr> <tr> <td>Totals</td> <td>12</td> <td>260</td> <td>272</td> </tr> </table> <p data-bbox="1012 1257 1496 1369">Sensitivity 50.00% (22.29% to 77.71%) Specificity 90.77% (86.41% to 93.88%) Positive likelihood ratio 5.42 (2.74 to 10.72) Negative likelihood ratio 0.55 (0.31 to 0.97)</p>	Normal baseline and variability	6	236	242	Totals	14	260	274		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Late decelerations+tachycardia	6	24	30	Normal baseline and variability	6	236	242	Totals	12	260	272	
Normal baseline and variability	6	236	242																										
Totals	14	260	274																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Late decelerations+tachycardia	6	24	30																										
Normal baseline and variability	6	236	242																										
Totals	12	260	272																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<p><b>Diagnostic accuracy (95% CI) of the following EFM features to detect fetal lactacidaemia (lactate&gt;4.8 mmol/l) at last FBS (negative test result is 'normal baseline and variability')</b></p> <p><u>Reduced variability</u></p> <table border="1" data-bbox="1021 472 1487 1035"> <thead> <tr> <th></th> <th>Lactate &gt;4.8 mmol/l</th> <th>Lactate ≤4.8 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Reduced variability</td> <td>5</td> <td>108</td> <td>113</td> </tr> <tr> <td>Normal baseline and variability</td> <td>9</td> <td>178</td> <td>187</td> </tr> <tr> <td>Totals</td> <td>14</td> <td>286</td> <td>300</td> </tr> </tbody> </table> <p>Sensitivity 35.7% (14.1-63.9%)                      Specificity 62.2% (61.2-63.6%)                      Positive likelihood ratio 0.95 (0.36-1.76)                      Negative likelihood ratio 1.03 (0.57-1.40)</p> <p><u>Absent variability</u></p> <table border="1" data-bbox="1021 1211 1487 1370"> <thead> <tr> <th></th> <th>Lactate &gt;4.8 mmol/l</th> <th>Lactate ≤4.8 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Reduced variability	5	108	113	Normal baseline and variability	9	178	187	Totals	14	286	300		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals					
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Reduced variability	5	108	113																										
Normal baseline and variability	9	178	187																										
Totals	14	286	300																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
				Absent variability	7	25	32	
				Normal baseline and variability	9	178	187	
				Totals	16	203	219	
				Sensitivity 43.8% (20.8-69.4%) Specificity 87.7% (82.2-91.7%) Positive likelihood ratio 3.55 (1.83-6.91) Negative likelihood ratio 0.64 (0.42-0.99)				
				<u>Increased variability</u>				
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	
				Increased variability	2	5	7	
				Normal baseline and variability	9	178	187	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
				<table border="1" data-bbox="1021 276 1491 347"> <tr> <td>Totals</td> <td>11</td> <td>183</td> <td>194</td> </tr> </table> <p data-bbox="1021 355 1491 467">                     Sensitivity 18.2% (3.2-52.2%)                      Specificity 97.3% (93.4-99.0%)                      Positive likelihood ratio 6.65 (1.45-30.51)                      Negative likelihood ratio 0.84 (0.64-1.11)                 </p> <p data-bbox="1021 496 1491 520"><u>Bradycardic episode</u></p> <table border="1" data-bbox="1021 528 1491 722"> <tr> <td></td> <td>Lactate &gt;4.8 mmol/l</td> <td>Lactate ≤4.8 mmol/l</td> <td>Totals</td> </tr> <tr> <td>Bradycardic episode</td> <td>12</td> <td>24</td> <td>36</td> </tr> <tr> <td>Normal baseline and variability</td> <td>9</td> <td>178</td> <td>187</td> </tr> <tr> <td>Totals</td> <td>21</td> <td>202</td> <td>223</td> </tr> </table> <p data-bbox="1021 1137 1491 1249">                     Sensitivity 57.1% (34.4-77.4%)                      Specificity 88.1% (82.6-92.1%)                      Positive likelihood ratio 4.81 (2.84-8.15)                      Negative likelihood ratio 0.49 (0.30-0.80)                 </p> <p data-bbox="1021 1273 1491 1297"><u>Tachycardia</u></p>	Totals	11	183	194		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Bradycardic episode	12	24	36	Normal baseline and variability	9	178	187	Totals	21	202	223	
Totals	11	183	194																						
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																						
Bradycardic episode	12	24	36																						
Normal baseline and variability	9	178	187																						
Totals	21	202	223																						



Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	
				Tachycardia	16	90	106	
				Normal baseline and variability	9	178	187	
				Totals	25	268	293	
Sensitivity 64.0% (42.6-81.3%) Specificity 66.4% (60.4-72.0%) Positive likelihood ratio 1.91 (1.36-2.67) Negative likelihood ratio 0.54 (0.32-0.92)								
<u>Tachycardia + reduced variability</u>								
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	
				Tachycardia +	7	121	128	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1019 272 1487 671"> <tr> <td>reduced variability</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Normal baseline and variability</td> <td>9</td> <td>178</td> <td>187</td> </tr> <tr> <td>Totals</td> <td>16</td> <td>299</td> <td>315</td> </tr> </table> <p data-bbox="1019 683 1487 794">Sensitivity 43.8% (20.8-69.4%) Specificity 59.3% (53.7-65.1%) Positive likelihood ratio 1.08 (0.61-1.92) Negative likelihood ratio 0.94 (0.61-1.46)</p> <p data-bbox="1019 821 1487 845"><u>Severe variable decelerations</u></p> <table border="1" data-bbox="1019 853 1487 1348"> <thead> <tr> <th></th> <th>Lactate &gt;4.8 mmol/l</th> <th>Lactate ≤4.8 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Severe variable decelerations</td> <td>21</td> <td>76</td> <td>97</td> </tr> <tr> <td>Normal baseline</td> <td>9</td> <td>178</td> <td>187</td> </tr> </tbody> </table>	reduced variability				Normal baseline and variability	9	178	187	Totals	16	299	315		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Severe variable decelerations	21	76	97	Normal baseline	9	178	187	
reduced variability																													
Normal baseline and variability	9	178	187																										
Totals	16	299	315																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Severe variable decelerations	21	76	97																										
Normal baseline	9	178	187																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1019 272 1494 472"> <tr> <td data-bbox="1019 272 1176 392">and variability</td> <td data-bbox="1176 272 1270 392"></td> <td data-bbox="1270 272 1415 392"></td> <td data-bbox="1415 272 1494 392"></td> </tr> <tr> <td data-bbox="1019 392 1176 472">Totals</td> <td data-bbox="1176 392 1270 472">30</td> <td data-bbox="1270 392 1415 472">254</td> <td data-bbox="1415 392 1494 472">284</td> </tr> </table> <p data-bbox="1019 472 1494 592">Sensitivity 70.0% (50.4-84.6%) Specificity 70.1% (64.0-75.6%) Positive likelihood ratio 2.34 (1.73-3.16) Negative likelihood ratio 0.43 (0.25-0.74)</p> <p data-bbox="1019 616 1211 639"><u>Late decelerations</u></p> <table border="1" data-bbox="1019 639 1494 847"> <tr> <td data-bbox="1019 639 1176 847"></td> <td data-bbox="1176 639 1270 847">Lactate &gt;4.8 mmol/l</td> <td data-bbox="1270 639 1415 847">Lactate ≤4.8 mmol/l</td> <td data-bbox="1415 639 1494 847">Totals</td> </tr> <tr> <td data-bbox="1019 847 1176 1007">Late decelerations</td> <td data-bbox="1176 847 1270 1007">11</td> <td data-bbox="1270 847 1415 1007">38</td> <td data-bbox="1415 847 1494 1007">49</td> </tr> <tr> <td data-bbox="1019 1007 1176 1214">Normal baseline and variability</td> <td data-bbox="1176 1007 1270 1214">9</td> <td data-bbox="1270 1007 1415 1214">178</td> <td data-bbox="1415 1007 1494 1214">187</td> </tr> <tr> <td data-bbox="1019 1214 1176 1294">Totals</td> <td data-bbox="1176 1214 1270 1294">0</td> <td data-bbox="1270 1214 1415 1294">216</td> <td data-bbox="1415 1214 1494 1294">236</td> </tr> </table> <p data-bbox="1019 1294 1494 1385">Sensitivity 55.0% (32.0-76.2%) Specificity 82.4% (76.5-87.1%) Positive likelihood ratio 3.13 (1.91-5.10)</p>	and variability				Totals	30	254	284		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Late decelerations	11	38	49	Normal baseline and variability	9	178	187	Totals	0	216	236	
and variability																													
Totals	30	254	284																										
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																										
Late decelerations	11	38	49																										
Normal baseline and variability	9	178	187																										
Totals	0	216	236																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																
				<p>Negative likelihood ratio 0.55 (0.34-0.89)</p> <p><u>Severe variable decelerations and + reduced variability</u></p> <table border="1" data-bbox="1021 389 1491 1161"> <thead> <tr> <th data-bbox="1021 389 1178 587"></th> <th data-bbox="1178 389 1272 587">Lactate &gt;4.8 mmol/l</th> <th data-bbox="1272 389 1417 587">Lactate ≤4.8 mmol/l</th> <th data-bbox="1417 389 1491 587">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 587 1178 874">Severe variable decelerations and + reduced variability</td> <td data-bbox="1178 587 1272 874">8</td> <td data-bbox="1272 587 1417 874">20</td> <td data-bbox="1417 587 1491 874">28</td> </tr> <tr> <td data-bbox="1021 874 1178 1082">Normal baseline and variability</td> <td data-bbox="1178 874 1272 1082">9</td> <td data-bbox="1272 874 1417 1082">178</td> <td data-bbox="1417 874 1491 1082">187</td> </tr> <tr> <td data-bbox="1021 1082 1178 1161">Totals</td> <td data-bbox="1178 1082 1272 1161">17</td> <td data-bbox="1272 1082 1417 1161">198</td> <td data-bbox="1417 1082 1491 1161">215</td> </tr> </tbody> </table> <p>Sensitivity 47.1% (23.9-71.5%)                      Specificity 89.9% (84.6-93.6%)                      Positive likelihood ratio 4.66 (2.42-8.95)                      Negative likelihood ratio 0.59 (0.38-0.92)</p> <p><u>Late decelerations + reduced variability</u></p>		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Severe variable decelerations and + reduced variability	8	20	28	Normal baseline and variability	9	178	187	Totals	17	198	215	
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																		
Severe variable decelerations and + reduced variability	8	20	28																		
Normal baseline and variability	9	178	187																		
Totals	17	198	215																		

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	
				Late decelerations + reduced variability	10	24	34	
				Normal baseline and variability	9	178	187	
				Totals	19	202	221	
				Sensitivity 52.6% (29.5-74.8%) Specificity 88.1% (82.6-92.1%) Positive likelihood ratio 4.43 (2.51-7.82) Negative likelihood ratio 0.54 (0.33-0.86)				
				<u>Severe variable decelerations + tachycardia</u>				
					Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
				<table border="1" data-bbox="1021 272 1491 799"> <tr> <td data-bbox="1021 272 1178 520">Severe variable decelerations + tachycardia</td> <td data-bbox="1178 272 1267 520">16</td> <td data-bbox="1267 272 1413 520">17</td> <td data-bbox="1413 272 1491 520">33</td> </tr> <tr> <td data-bbox="1021 520 1178 719">Normal baseline and variability</td> <td data-bbox="1178 520 1267 719">9</td> <td data-bbox="1267 520 1413 719">178</td> <td data-bbox="1413 520 1491 719">187</td> </tr> <tr> <td data-bbox="1021 719 1178 799">Totals</td> <td data-bbox="1178 719 1267 799">25</td> <td data-bbox="1267 719 1413 799">195</td> <td data-bbox="1413 719 1491 799">220</td> </tr> </table> <p data-bbox="1021 807 1491 919">Sensitivity 64.0% (42.6-81.3%) Specificity 91.3% (86.2-94.7%) Positive likelihood ratio 7.34 (4.27-12.61) Negative likelihood ratio 0.39 (0.23-0.67)</p> <p data-bbox="1021 943 1491 967"><u>Late decelerations + tachycardia</u></p> <table border="1" data-bbox="1021 975 1491 1377"> <tr> <td data-bbox="1021 975 1178 1174"></td> <td data-bbox="1178 975 1267 1174">Lactate &gt;4.8 mmol/l</td> <td data-bbox="1267 975 1413 1174">Lactate ≤4.8 mmol/l</td> <td data-bbox="1413 975 1491 1174">Totals</td> </tr> <tr> <td data-bbox="1021 1182 1178 1377">Late decelerations + tachycardia</td> <td data-bbox="1178 1182 1267 1377">10</td> <td data-bbox="1267 1182 1413 1377">20</td> <td data-bbox="1413 1182 1491 1377">30</td> </tr> </table>	Severe variable decelerations + tachycardia	16	17	33	Normal baseline and variability	9	178	187	Totals	25	195	220		Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals	Late decelerations + tachycardia	10	20	30	
Severe variable decelerations + tachycardia	16	17	33																						
Normal baseline and variability	9	178	187																						
Totals	25	195	220																						
	Lactate >4.8 mmol/l	Lactate ≤4.8 mmol/l	Totals																						
Late decelerations + tachycardia	10	20	30																						

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments								
				<table border="1" data-bbox="1019 272 1491 555"> <tr> <td data-bbox="1019 272 1178 475">Normal baseline and variability</td> <td data-bbox="1178 272 1270 475">9</td> <td data-bbox="1270 272 1417 475">178</td> <td data-bbox="1417 272 1491 475">187</td> </tr> <tr> <td data-bbox="1019 480 1178 555">Totals</td> <td data-bbox="1178 480 1270 555">19</td> <td data-bbox="1270 480 1417 555">198</td> <td data-bbox="1417 480 1491 555">217</td> </tr> </table> <p data-bbox="1019 560 1491 671">Sensitivity 52.6% (29.5-74.8%) Specificity 89.9% (84.6-93.6%) Positive likelihood ratio 5.21 (2.87-9.45) Negative likelihood ratio 0.53 (0.33-0.85)</p> <p data-bbox="1019 699 1491 783">Sensitivity, specificity and likelihood ratios calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a></p>	Normal baseline and variability	9	178	187	Totals	19	198	217	
Normal baseline and variability	9	178	187										
Totals	19	198	217										
<p data-bbox="190 932 392 963"><b>Full citation</b></p> <p data-bbox="190 986 392 1393">Liu, L., Tuuli, M. G., Roehl, K. A., Odibo, A. O., Macones, G. A., Cahill, A. G., Electronic fetal monitoring patterns associated with respiratory morbidity in term neonates, American Journal of Obstetrics &amp;</p>	<p data-bbox="392 932 656 963"><b>Sample size</b></p> <p data-bbox="392 986 656 1018">N=4736</p> <p data-bbox="392 1070 656 1102"><b>Characteristics</b></p> <p data-bbox="392 1125 656 1393">Compared to the group who had no respiratory morbidity (n=4561), the group that had respiratory morbidity (n=175) more often had pre-eclampsia, pregestational diabetes, were</p>	<p data-bbox="656 932 831 963"><b>Interventions</b></p> <p data-bbox="656 986 831 1102">EFM patterns in the last 30 minutes before birth</p>	<p data-bbox="831 932 1008 963"><b>Details</b></p> <p data-bbox="831 986 1008 1393">EFM was performed with the use of internal or external monitoring as clinically indicated. The primary outcome was neonatal respiratory morbidity, which was</p>	<p data-bbox="1008 932 1503 963"><b>Results</b></p> <p data-bbox="1008 986 1503 1161"><b>Adjusted* odds ratio (aOR) with 95% confidence interval (CI) of neonatal respiratory morbidity** in the presence of the following EFM characteristics in the last 30 minutes before birth in the whole sample (n=4736)</b></p> <p data-bbox="1008 1184 1503 1326">Ever baseline bradycardia &lt;110bpm: aOR 0.5 (0.1-3.4) Ever baseline &lt;120 bpm: aOR 0.7 (0.4-1.3) Ever baseline tachycardia &gt;160 bpm: aOR 2.9 (1.9-4.4)</p>	<p data-bbox="1503 932 2042 963"><b>Limitations</b></p> <p data-bbox="1503 986 2042 1050">According to NICE 2012 guidelines manual checklist for prognostic studies</p> <table border="1" data-bbox="1503 1050 2042 1393"> <tr> <td data-bbox="1503 1050 1749 1393"><b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential</b></td> <td data-bbox="1749 1050 1852 1393">Yes</td> <td data-bbox="1852 1050 1955 1393">No</td> <td data-bbox="1955 1050 2042 1393">Unclear</td> </tr> </table>	<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential</b>	Yes	No	Unclear				
<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential</b>	Yes	No	Unclear										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments								
<p>Gynecology, 213, 681.e1-6, 2015</p> <p><b>Ref Id</b> 446299</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Study type</b> Prospective cohort study</p> <p><b>Aim of the study</b> To identify electronic fetal monitoring patterns that are associated with neonatal respiratory morbidity</p> <p><b>Study dates</b> The study was conducted after approval from the Washington University School of medicine Human Research</p>	<p>nulliparous, had had previous caesarean section, had received prostaglandin, had not had vaginal birth, had caesarean birth, and had had maternal fever. No difference between the groups was observed in maternal age, gestational age at birth, labour type (spontaneous, augmented or induced), birthweight, percentage of maternal black race, percentage of gestational diabetes, and use of regional anaesthesia, Foley bulb, and oxytocin</p> <p><b>Inclusion criteria</b> Term, vertex, non-anomalous singleton pregnancies during labour at Washington University in St. Louis Missouri, USA</p> <p><b>Exclusion criteria</b> Neonates with &lt;10 minutes of EFM in the 30 minutes before birth.</p>		<p>defined as either any oxygen requirement at or after 6 hours of life or any mechanical ventilation in the first 24 hours. Because caesarean birth and maternal fever are both risk factors for increased neonatal respiratory morbidity, secondary analyses were performed that excluded those women who underwent caesarean birth and those with fever. Because mechanical ventilation is the most severe acute respiratory morbidity for a term infant, analyses were repeated to estimate which EFM patterns</p>	<p>Ever absent or minimal variability: aOR 1.3 (0.9-1.8) Mostly absent or minimal variability: aOR 1.1 (0.8-1.6) Always absent or minimal variability: aOR 1.2 (0.8-1.7) Mostly moderate variability: aOR 0.7 (0.5-1.0) Always moderate variability: aOR 0.7 (0.5-0.9) Ever marked variability: aOR 2.7 (1.5-5.0)</p> <p>Accelerations present: aOR 0.6 (0.4-0.9) Decelerations present: aOR 0.8 (0.5-1.2) Early decelerations: aOR 0.4 (0.1-1.1) Variable decelerations: aOR 0.8 (0.5-1.1) Late decelerations: aOR 0.8 (0.6-1.1) Prolonged decelerations: aOR 1.7 (1.3-2.4)</p> <p><b>Adjusted* OR (95% CI) of neonatal respiratory morbidity** in the presence of the following EFM characteristics in the last 30 minutes before birth <u>excluding caesarean birth (n=3994)</u></b></p> <p>Ever baseline tachycardia &gt;160 bpm: aOR 3.0 (1.8-5.1) Always moderate variability: aOR 0.7 (0.5-1.1) Ever marked variability: aOR 2.7 (1.3-5.7)</p> <p>Accelerations present: aOR 0.8 (0.5-1.2) Variable decelerations: aOR 3.4 (1.2-9.5) Prolonged decelerations: aOR 1.8 (1.2-2.8)</p> <p><b>Adjusted* OR (95% CI) of neonatal respiratory morbidity** in the presence of</b></p>	<table border="1"> <tr> <td rowspan="2"> <p><b>bias to the results</b></p> <p>Consecutive singleton, vertex, non-anomalous pregnancies were included. Mean gestational weeks in the sample was 38.9 (±1.3) and 38.9 (±1.2) (depending on the outcome finding) so a small portion of the births might be preterm. Also, the population is of both low- and high-risk pregnancies</p> </td> <td></td> <td></td> <td></td> </tr> <tr> <td> <p><b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately</b></p> </td> <td> <p><b>Yes</b></p> </td> <td> <p>No</p> </td> <td> <p>Unclear</p> </td> </tr> </table>	<p><b>bias to the results</b></p> <p>Consecutive singleton, vertex, non-anomalous pregnancies were included. Mean gestational weeks in the sample was 38.9 (±1.3) and 38.9 (±1.2) (depending on the outcome finding) so a small portion of the births might be preterm. Also, the population is of both low- and high-risk pregnancies</p>				<p><b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately</b></p>	<p><b>Yes</b></p>	<p>No</p>	<p>Unclear</p>
<p><b>bias to the results</b></p> <p>Consecutive singleton, vertex, non-anomalous pregnancies were included. Mean gestational weeks in the sample was 38.9 (±1.3) and 38.9 (±1.2) (depending on the outcome finding) so a small portion of the births might be preterm. Also, the population is of both low- and high-risk pregnancies</p>													
	<p><b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately</b></p>	<p><b>Yes</b></p>	<p>No</p>	<p>Unclear</p>									



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments			
<p>Protection Office (approval in 11/2014)</p> <p><b>Source of funding</b></p> <p>Supported in part by the National Institute of Child Health and Human Development</p>	<p>Gestational age &lt;37 weeks.</p> <p>Postnatal anomaly diagnosis</p>		<p>were associated with mechanical ventilation compared with those without morbidity. Multivariable logistic regression was performed in a backward step-wise fashion to refine estimates of the association between EFM characteristics and neonatal respiratory morbidity by controlling for confounding factors. Model fit of the final model (adjusted for maternal fever, parity, pregestational diabetes, previous caesarean birth, and pre-eclampsia) was tested with the Hosmer-Lemeshow goodness-of-fit test</p>	<p><b>the following EFM characteristics in the last 30 minutes before birth <u>excluding women with maternal fever (n=4647)</u></b></p> <p>Ever baseline tachycardia &gt;160 bpm: aOR 2.9 (1.9-4.6)</p> <p>Always moderate variability: aOR 0.7 (0.5-1.0)</p> <p>Ever marked variability: aOR 3.1 (1.7-5.7)</p> <p>Accelerations present: aOR 0.6 (0.4-0.9)</p> <p>Prolonged decelerations: aOR 1.8 (1.3-2.5)</p> <p><b>Adjusted* OR (95% CI) of neonatal <u>mechanical ventilation</u> (versus no respiratory morbidity) in the presence of the following EFM characteristics in the last 30 minutes before birth(n=4605)</b></p> <p>Ever baseline tachycardia &gt;160 bpm: aOR 3.1 (1.4-6.7)</p> <p>Always moderate variability: aOR 0.8 (0.4-1.40)</p> <p>Ever marked variability: aOR 2.2 (0.7-7.2)</p> <p>Accelerations present: aOR 0.4 (0.2-0.9)</p> <p>Prolonged decelerations: aOR 2.6 (1.4-4.7)</p> <p>*Adjusted for maternal fever, parity, pregestational diabetes, previous caesarean birth, pre-eclampsia</p> <p>**Neonatal respiratory morbidity defined as either any oxygen requirement at or after 6 hours of life or any mechanical ventilation in the first 24 hours after birth</p>	<p><b>represent the sample), sufficient to limit potential bias</b></p> <p>N/A</p>			
<p><b>The prognostic factor of interest is adequately measured in study participants, sufficient to limit potential bias</b></p> <p>EFM interpretation is known to be difficult and can be subject to bias. It is not reported if more than reviewer interpreted each tracing. Only the last 30 minutes of the EFM before</p>					<p>Yes</p>	<p>No</p>	<p><b>Unclear</b></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments			
					birth was considered			
					<b>The outcome of interest is adequately measured in study participants, sufficient to limit potential bias</b>	<u>Yes</u>	No	Unclear
					<b>Important potential confounders are appropriately accounted for, limiting potential bias with respect to the prognostic factor of interest</b>	<u>Yes</u>	No	Unclear
					<b>The statistical analysis is appropriate for the design of the study, limiting potential for the</b>	<u>Yes</u>	No	Unclear

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments												
					<table border="1"> <tr> <td><b>presentation of invalid results</b> Multiple variable logistic regression was conducted appropriately. However, it is unclear why the crude outcome was reported as relative risk and the adjusted one as odds ratio</td> <td></td> <td></td> <td></td> </tr> <tr> <td><b>Risk of bias:</b></td> <td><b>No serious risk of bias</b></td> <td>Serious</td> <td>Very serious</td> </tr> <tr> <td colspan="4"><b>Other information</b></td> </tr> </table>	<b>presentation of invalid results</b> Multiple variable logistic regression was conducted appropriately. However, it is unclear why the crude outcome was reported as relative risk and the adjusted one as odds ratio				<b>Risk of bias:</b>	<b>No serious risk of bias</b>	Serious	Very serious	<b>Other information</b>			
<b>presentation of invalid results</b> Multiple variable logistic regression was conducted appropriately. However, it is unclear why the crude outcome was reported as relative risk and the adjusted one as odds ratio																	
<b>Risk of bias:</b>	<b>No serious risk of bias</b>	Serious	Very serious														
<b>Other information</b>																	
<b>Full citation</b> Sharbaf,F.R., Amjadi,N., Alavi,A., Akbari,S.,	<b>Sample size</b> N=818 total (including both low- and high-risk populations, 328 high risk and 497 low risk)	<b>Interventions</b> Fetal heart rate (FHR) tracings obtained with a	<b>Details</b> The FHR tracings were interpreted by two	<b>Results</b> <b>Relative risk (RR) of the following perinatal outcomes in low- and high-risk and overall populations with</b>	<b>Limitations</b> Assessed with QUADAS-2: -Not described whether all women fitting the inclusion/exclusion criteria during the study period were selected.												

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																	
<p>Forghani,F., Normal and indeterminate pattern of fetal cardiotocography in admission test and pregnancy outcome, Journal of Obstetrics and Gynaecology Research, 40, 694-699, 2014</p> <p><b>Ref Id</b> 324863</p> <p><b>Country/ies where the study was carried out</b>  Iran</p> <p><b>Study type</b>  Prospective comparative study</p> <p><b>Aim of the study</b>  To evaluate the prognostic value of normal and indeterminate patterns of cardiotocography in admission tests and</p>	<p>n=659 normal tracing n=159 intermediate tracing</p> <p>N=492 low-risk sample n=410 normal tracing in low-risk sample n=82 intermediate tracing in low-risk sample</p> <p>N=326 high-risk sample n=249 normal tracing in high-risk sample n=77 intermediate tracing in high-risk sample</p> <p><b>Characteristics</b>  The mean age of the women was 26.6 (+- 5.1) years. The median gestational age at birth was 39 (34-42) weeks. Admission tests were: 659 (80.4%) normal, 159 (19.4%) indeterminate and two (0.2%) abnormal. 60% of the women were categorised as low-risk and 40% were categorised as high-risk.</p>	<p>non-stress test machine in early labour during a 20-40 minute period</p>	<p>obstetricians according to NICHHD recommendations resulting in normal, indeterminate, or abnormal categories based on baseline fetal heart rate, variability, acceleration and types of deceleration. Obstetricians were blinded to clinical conditions in order to avoid biased findings. When there was a disagreement, consensus was obtained with a perinatologist .</p> <p>Unfavourable outcome related to the women was only caesarean section due to non-reassuring fetal heart rate pattern. Non-reassuring fetal heart rate</p>	<p><b>indeterminate FHR tracing (according to NICHHD classification)</b> <u>CS due to non-reassuring fetal heart rate pattern</u> Overall: RR 3.8 (2.5-5.6) Low-risk group: RR 3.7 (2.1-6.9) High-risk group: RR 3.4 (2.0-5.7)</p> <p><u>Umbilical artery pH &lt;=7.2</u> Overall: RR 1.5 (0.8-2.8) Low-risk group: RR 1.05 (0.4-3.0) High-risk group: RR 1.9 (0.8-4.5)</p> <p><u>NICU admission</u> Overall: RR 2.3 (1.2-4.2) Low-risk group: RR 1.0 (0.3-3.4) High-risk group: RR 3.2 (1.5-6.9)</p> <p><u>NICU admission after excluding preterm birth</u> Overall: RR 2.0 (1.0-4.1) Low-risk group: RR 0.7 (0.2-3.1) High-risk group: RR 3.6 (1.4-9.2)</p> <p><b>Diagnostic accuracy of indeterminate FHR tracing (NICHHD classification) on different perinatal outcomes (NR = not reported)</b> <u>CS</u> <i>Mixed population (including both low- and high-risk samples)</i></p> <table border="1"> <tr> <td>a</td> <td>CS</td> <td>No CS</td> <td>Totals</td> </tr> <tr> <td>Indeterminate FHR category</td> <td>NR</td> <td>NR</td> <td>NR</td> </tr> </table>	a	CS	No CS	Totals	Indeterminate FHR category	NR	NR	NR	<p>-The study included gestational ages 35-36 weeks (preterm), while the guideline review is looking at term only (37-42 weeks).</p> <p>-Even though there were two independent FHR tracing reviewers who were blinded to clinical conditions, it is known that FHR tracing interpretation is difficult and can be subjective and therefore introduce bias.</p> <p>-Unlikely that the 'diagnosis' of outcomes would have been blinded to the FHR tracing interpretation.</p> <p>-Index test (CTG tracing) performed before birth and reference test (usual ascertainment of outcome) performed during/after birth might mean that differences in the test results are due to events after the index test.</p> <p>Assessed with NICE 2012 guidelines manual checklist for prognostic studies:</p> <table border="1"> <tr> <td rowspan="2">1.1</td> <td><b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential bias to the results</b></td> <td><b>Yes</b></td> <td>No</td> <td>Unclear</td> </tr> <tr> <td>The study population included singleton pregnancies with more than 34 weeks of gestation, intact membranes with both low- and high-risk</td> <td></td> <td></td> <td></td> </tr> </table>	1.1	<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential bias to the results</b>	<b>Yes</b>	No	Unclear	The study population included singleton pregnancies with more than 34 weeks of gestation, intact membranes with both low- and high-risk			
a	CS	No CS	Totals																			
Indeterminate FHR category	NR	NR	NR																			
1.1	<b>The study sample represents the population of interest with regard to key characteristics, sufficient to limit potential bias to the results</b>	<b>Yes</b>	No	Unclear																		
	The study population included singleton pregnancies with more than 34 weeks of gestation, intact membranes with both low- and high-risk																					

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments						
pregnancy outcomes	Obstetric characteristics of the women (n=818):		pattern was defined as abnormal patterns according to the NICHD recommendation. Fetal complications (neonatal death, umbilical cord artery pH <=7.2, 5-minute Apgar <7, thick meconium staining in liquor, admission to the neonatal intensive care unit, neonatal mortality and low birthweight) were assessed and compared in both groups	Normal FHR category	NR	NR	NR	pregnancies. However, since the proportion of preterm births (<37 weeks of gestation) in the study population is small (8.1%), and only includes late preterm births (35-36 weeks of gestation), this was not considered a serious risk of bias/serious indirectness. The findings are presented in the whole population (mix of low- and high-risk) as well as for low- and high-risk populations separately			
Study dates	%			Totals	NR	NR	NR				
March 2010 to February 2011	Nulliparous	64.2		Sensitivity 30.9% Specificity 86.3% Positive likelihood ratio* 2.26 Negative likelihood ratio* 0.80							
Source of funding	Preterm <37 wks	8.1		<i>Low-risk population</i>							
None reported	Post-date >41 wks	0.9			CS	No CS	Totals				
	Pregnancy-induced hypertension	8.3		Indeterminate FHR category	NR	NR	NR				
	Pre-eclampsia	8.4		Normal FHR category	NR	NR	NR				
	Gestational diabetes	4.5		Totals	NR	NR	NR				
	Intrauterine growth restriction	3.9		Sensitivity 28.6% Specificity 87.7% Positive likelihood ratio* 2.33 Negative likelihood ratio* 0.81							
	Decreased fetal movement	15.2		<i>High-risk population</i>							
	Decreased amniotic fluid	11.7		CS	No CS	Totals	1.2	<b>Loss to follow-up is unrelated to key characteristics (that is, the study data adequately represent the sample), sufficient to limit potential bias</b> N/A	Yes	No	Unclear
							1.3	<b>The prognostic factor of interest is adequately measured in study participants, sufficient to limit potential bias</b>	Yes	No	Unclear

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																
	Thick meconium staining	14.1		Indeterminate FHR category	<p>Only 20-40 minutes of trace in 'early labour' were considered. The tracings were interpreted by two obstetricians who were blinded to the clinical conditions. In case of disagreement of interpretation between the obstetricians, a consensus view would be provided by a perinatologist.</p> <p>Interpretation of CTG tracing is known to be difficult and can be subject to bias, however, two (and potentially three) different persons reviewed each tracing</p>																
	Non-reassuring fetal heart rate pattern	11.4		Normal FHR category																	
	Caesarean section (CS)	33.3		Totals																	
	CS due to non-reassuring fetal heart rate pattern	10.3		<p>Sensitivity 33.1%</p> <p>Specificity 83.4%</p> <p>Positive likelihood ratio* 1.99</p> <p>Negative likelihood ratio* 0.80</p> <p><u>Umbilical artery pH <math>\leq 7.2</math></u>  <i>Mixed population (including both low- and high-risk samples)</i></p>																	
	<p><b>Inclusion criteria</b></p> <p>Women admitted to the labour ward at the Women's Hospital, Tehran University of Medical Sciences between March 2010 and February 2011 with singleton pregnancies with gestational age of more than 34 weeks and intact membranes. Pregnancies were considered 'high risk' when there was a post-dated pregnancy (&gt;41 weeks),</p>			<table border="1"> <thead> <tr> <th></th> <th>Umbilical artery pH <math>\leq 7.2</math></th> <th>Umbilical artery pH <math>&gt; 7.2</math></th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Indeterminate FHR category</td> <td>13</td> <td>55</td> <td>68</td> </tr> <tr> <td>Normal FHR category</td> <td>19</td> <td>127</td> <td>146</td> </tr> <tr> <td>Totals</td> <td>32</td> <td>182</td> <td>214</td> </tr> </tbody> </table> <p>Sensitivity 40.6% (24.2-59.2%)</p>		Umbilical artery pH $\leq 7.2$	Umbilical artery pH $> 7.2$	Totals	Indeterminate FHR category	13	55	68	Normal FHR category	19	127	146	Totals	32	182	214	
	Umbilical artery pH $\leq 7.2$	Umbilical artery pH $> 7.2$	Totals																		
Indeterminate FHR category	13	55	68																		
Normal FHR category	19	127	146																		
Totals	32	182	214																		
				1.4	<p><b>The outcome of interest is adequately measured in study participants, sufficient to limit potential bias</b></p> <p>Yes</p> <p>No</p> <p>Unclear</p>																
				1.5	<p><b>Important potential confounders are appropriately accounted for, limiting potential bias with respect to the</b></p> <p>Yes</p> <p>No</p> <p>Unclear</p>																

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																											
	<p>oligohydramnios (amniotic fluid index &lt;=5), pregnancy-induced hypertension, gestational diabetes, pre-eclampsia, intra-uterine growth restriction or decreased fetal movements</p> <p><b>Exclusion criteria</b></p> <p>Women with active phase of labour, &lt;34 weeks of gestation and those with twin pregnancies, hydramnios or previous caesarean section who were not candidates for vaginal birth</p>			<p>Specificity 69.8% (62.5-76.2%) Positive likelihood ratio* 1.34 (0.84-2.16) Negative likelihood ratio* 0.85 (0.64-1.14)</p> <p><i>Low-risk population</i></p> <table border="1" data-bbox="1021 416 1491 655"> <thead> <tr> <th></th> <th>Umbilical artery pH &lt;=7.2</th> <th>Umbilical artery pH &gt;7.2</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Indeterminate FHR category</td> <td>4</td> <td>78</td> <td>82</td> </tr> <tr> <td>Normal FHR category</td> <td>22</td> <td>388</td> <td>410</td> </tr> <tr> <td>Totals</td> <td>26</td> <td>466</td> <td>492</td> </tr> </tbody> </table> <p>Sensitivity 26.7% (8.9-55.2%) Specificity 83.7% (80.0-86.8%) Positive likelihood ratio* 1.63 (0.69-3.87) Negative likelihood ratio* 0.88 (0.65-1.19)</p> <p><i>High-risk population</i></p> <table border="1" data-bbox="1021 1158 1491 1286"> <thead> <tr> <th></th> <th>Umbilical artery</th> <th>Umbilical artery</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Umbilical artery pH <=7.2	Umbilical artery pH >7.2	Totals	Indeterminate FHR category	4	78	82	Normal FHR category	22	388	410	Totals	26	466	492		Umbilical artery	Umbilical artery	Totals					<p><b>prognostic factor of interest</b></p> <p>Statistical methods for deriving relative risks were not described at all and whether or not the model adjusted for potential confounders is not reported. Presumably, the estimates are crude and therefore might be subject to serious risk of bias</p>			
	Umbilical artery pH <=7.2	Umbilical artery pH >7.2	Totals																													
Indeterminate FHR category	4	78	82																													
Normal FHR category	22	388	410																													
Totals	26	466	492																													
	Umbilical artery	Umbilical artery	Totals																													
				<p>1.6</p>	<p><b><u>The statistical analysis is appropriate for the design of the study, limiting potential for the presentation of invalid result</u></b></p> <p>Statistical methods for deriving relative risks were not described at all and whether or not the model adjusted for potential confounders is not reported. Presumably, the estimates are crude and therefore might be subject to serious risk of bias</p>	Yes	<b>No</b>	Unclear																								

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments											
					pH ≤7.2	pH >7.2		<table border="1"> <tr> <td colspan="2" data-bbox="1514 272 1921 475"><b>Overall risk of bias (for RR data):</b></td> <td data-bbox="1921 272 2029 475">No serious risk of bias</td> <td data-bbox="2029 272 2130 475">Serious risk of bias</td> <td data-bbox="2130 272 2233 475"><b><u>Very serious risk of bias</u></b></td> </tr> </table>	<b>Overall risk of bias (for RR data):</b>		No serious risk of bias	Serious risk of bias	<b><u>Very serious risk of bias</u></b>	<p data-bbox="1514 560 1713 587"><b>Other information</b></p>					
<b>Overall risk of bias (for RR data):</b>		No serious risk of bias	Serious risk of bias	<b><u>Very serious risk of bias</u></b>															
			Indeterminate FHR category	9	68	77													
			Normal FHR category	8	241	249													
			Totals	17	309	326													
				<p data-bbox="1016 724 1496 836">Sensitivity 52.9% (28.5-76.1%)                      Specificity 80.0% (72.9-82.4%)                      Positive likelihood ratio**2.41 (1.47-3.95)                      Negative likelihood ratio* 0.60 (0.36-1.00)</p> <p data-bbox="1016 863 1496 943"><u>NICU admission</u>                      Mixed population (including both low- and high-risk samples)</p> <table border="1"> <thead> <tr> <th data-bbox="1016 951 1180 1107"></th> <th data-bbox="1180 951 1296 1107">NICU admission</th> <th data-bbox="1296 951 1417 1107">No NICU admission</th> <th data-bbox="1417 951 1496 1107">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1016 1118 1180 1270">Indeterminate FHR category</td> <td data-bbox="1180 1118 1296 1270">15</td> <td data-bbox="1296 1118 1417 1270">144</td> <td data-bbox="1417 1118 1496 1270">159</td> </tr> </tbody> </table>					NICU admission	No NICU admission	Totals	Indeterminate FHR category	15	144	159				
	NICU admission	No NICU admission	Totals																
Indeterminate FHR category	15	144	159																



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 276 1487 472"> <tr> <td data-bbox="1021 276 1178 387">Normal FHR category</td> <td data-bbox="1184 276 1294 387">27</td> <td data-bbox="1301 276 1411 387">632</td> <td data-bbox="1417 276 1487 387">659</td> </tr> <tr> <td data-bbox="1021 392 1178 472">Totals</td> <td data-bbox="1184 392 1294 472">42</td> <td data-bbox="1301 392 1411 472">776</td> <td data-bbox="1417 392 1487 472">818</td> </tr> </table> <p data-bbox="1021 480 1487 592">Sensitivity 35.7% (22.0-52.0%) Specificity 81.4% (78.5-84.1%) Positive likelihood ratio* 1.92 (1.25-2.96) Negative likelihood ratio* 0.79 (0.63-1.00)</p> <p data-bbox="1021 616 1223 639"><i>Low-risk population</i></p> <table border="1" data-bbox="1021 647 1487 1169"> <thead> <tr> <th data-bbox="1021 647 1178 807"></th> <th data-bbox="1184 647 1294 807">NICU admission</th> <th data-bbox="1301 647 1411 807">No NICU admission</th> <th data-bbox="1417 647 1487 807">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 812 1178 971">Indeterminate FHR category</td> <td data-bbox="1184 812 1294 971">3</td> <td data-bbox="1301 812 1411 971">79</td> <td data-bbox="1417 812 1487 971">82</td> </tr> <tr> <td data-bbox="1021 976 1178 1136">Normal FHR category</td> <td data-bbox="1184 976 1294 1136">15</td> <td data-bbox="1301 976 1411 1136">395</td> <td data-bbox="1417 976 1487 1136">410</td> </tr> <tr> <td data-bbox="1021 1141 1178 1169">Totals</td> <td data-bbox="1184 1141 1294 1169">18</td> <td data-bbox="1301 1141 1411 1169">474</td> <td data-bbox="1417 1141 1487 1169">492</td> </tr> </tbody> </table> <p data-bbox="1021 1177 1487 1289">Sensitivity 16.7% (4.4-42.2%) Specificity 83.3% (79.6-86.5%) Positive likelihood ratio* 1.00 (0.35-2.86) Negative likelihood ratio* 1.00 (0.81-1.23)</p> <p data-bbox="1021 1313 1223 1337"><i>High-risk population</i></p>	Normal FHR category	27	632	659	Totals	42	776	818		NICU admission	No NICU admission	Totals	Indeterminate FHR category	3	79	82	Normal FHR category	15	395	410	Totals	18	474	492	
Normal FHR category	27	632	659																										
Totals	42	776	818																										
	NICU admission	No NICU admission	Totals																										
Indeterminate FHR category	3	79	82																										
Normal FHR category	15	395	410																										
Totals	18	474	492																										

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments																								
				<table border="1" data-bbox="1021 277 1491 799"> <thead> <tr> <th data-bbox="1021 277 1207 480"></th> <th data-bbox="1207 277 1317 480">NICU admission</th> <th data-bbox="1317 277 1424 480">No NICU admission</th> <th data-bbox="1424 277 1491 480">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 480 1207 596">Indeterminate FHR category</td> <td data-bbox="1207 480 1317 596">12</td> <td data-bbox="1317 480 1424 596">65</td> <td data-bbox="1424 480 1491 596">77</td> </tr> <tr> <td data-bbox="1021 596 1207 719">Normal FHR category</td> <td data-bbox="1207 596 1317 719">12</td> <td data-bbox="1317 596 1424 719">237</td> <td data-bbox="1424 596 1491 719">249</td> </tr> <tr> <td data-bbox="1021 719 1207 799">Totals</td> <td data-bbox="1207 719 1317 799">24</td> <td data-bbox="1317 719 1424 799">302</td> <td data-bbox="1424 719 1491 799">326</td> </tr> </tbody> </table> <p data-bbox="1021 807 1491 919">Sensitivity 50.0% (29.6-70.4%)                      Specificity 78.5% (73.3-82.9%)                      Positive likelihood ratio* 2.32 (1.47-3.66)                      Negative likelihood ratio* 0.64 (0.43-0.95)</p> <p data-bbox="1021 943 1491 1031"><u>NICU admission excluding preterm birth</u>  <i>Mixed population (including both low- and high-risk samples)</i></p> <table border="1" data-bbox="1021 1031 1491 1353"> <thead> <tr> <th data-bbox="1021 1031 1207 1353"></th> <th data-bbox="1207 1031 1317 1353">NICU admission excluding preterm birth</th> <th data-bbox="1317 1031 1424 1353">No NICU admission excluding</th> <th data-bbox="1424 1031 1491 1353">Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>					NICU admission	No NICU admission	Totals	Indeterminate FHR category	12	65	77	Normal FHR category	12	237	249	Totals	24	302	326		NICU admission excluding preterm birth	No NICU admission excluding	Totals					
	NICU admission	No NICU admission	Totals																													
Indeterminate FHR category	12	65	77																													
Normal FHR category	12	237	249																													
Totals	24	302	326																													
	NICU admission excluding preterm birth	No NICU admission excluding	Totals																													

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
						preterm birth		
				Indeterminate FHR category	NR	NR	NR	
				Normal FHR category	NR	NR	NR	
				Totals	NR	NR	NR	
				Sensitivity 31.3% Specificity 81.9% Positive likelihood ratio* 1.73 Negative likelihood ratio* 0.84				
				<i>Low-risk population</i>				
					NICU admission excluding preterm birth	No NICU admission excluding preterm birth	Totals	
				Indeterminate FHR category	NR	NR	NR	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 272 1491 392"> <tr> <td data-bbox="1021 272 1178 392">Normal FHR category</td> <td data-bbox="1178 272 1301 392">NR</td> <td data-bbox="1301 272 1424 392">NR</td> <td data-bbox="1424 272 1491 392">NR</td> </tr> <tr> <td data-bbox="1021 392 1178 472">Totals</td> <td data-bbox="1178 392 1301 472">NR</td> <td data-bbox="1301 392 1424 472">NR</td> <td data-bbox="1424 392 1491 472">NR</td> </tr> </table> <p data-bbox="1021 472 1491 592">Sensitivity 12.5% Specificity 83.2% Positive likelihood ratio* 0.74 Negative likelihood ratio* 1.05</p> <p data-bbox="1021 592 1491 647"><i>High-risk population</i></p> <table border="1" data-bbox="1021 647 1491 1015"> <tr> <td data-bbox="1021 647 1211 1015"></td> <td data-bbox="1211 647 1317 1015">NICU admission excluding preterm birth</td> <td data-bbox="1317 647 1424 1015">No NICU admission excluding preterm birth</td> <td data-bbox="1424 647 1491 1015">Totals</td> </tr> <tr> <td data-bbox="1021 1015 1211 1134">Indeterminate FHR category</td> <td data-bbox="1211 1015 1317 1134">NR</td> <td data-bbox="1317 1015 1424 1134">NR</td> <td data-bbox="1424 1015 1491 1134">NR</td> </tr> <tr> <td data-bbox="1021 1134 1211 1254">Normal FHR category</td> <td data-bbox="1211 1134 1317 1254">NR</td> <td data-bbox="1317 1134 1424 1254">NR</td> <td data-bbox="1424 1134 1491 1254">NR</td> </tr> <tr> <td data-bbox="1021 1254 1211 1334">Totals</td> <td data-bbox="1211 1254 1317 1334">NR</td> <td data-bbox="1317 1254 1424 1334">NR</td> <td data-bbox="1424 1254 1491 1334">NR</td> </tr> </table> <p data-bbox="1021 1334 1491 1372">Sensitivity 50.0%</p>	Normal FHR category	NR	NR	NR	Totals	NR	NR	NR		NICU admission excluding preterm birth	No NICU admission excluding preterm birth	Totals	Indeterminate FHR category	NR	NR	NR	Normal FHR category	NR	NR	NR	Totals	NR	NR	NR	
Normal FHR category	NR	NR	NR																										
Totals	NR	NR	NR																										
	NICU admission excluding preterm birth	No NICU admission excluding preterm birth	Totals																										
Indeterminate FHR category	NR	NR	NR																										
Normal FHR category	NR	NR	NR																										
Totals	NR	NR	NR																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<p>Specificity 79.9%                      Positive likelihood ratio* 2.49                      Negative likelihood ratio* 0.63</p> <p><u>Neonatal death</u>  <i>Mixed population (including both low- and high-risk samples)</i></p> <table border="1" data-bbox="1021 472 1491 994"> <thead> <tr> <th></th> <th>Neonatal death</th> <th>No neonatal death</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Indeterminate FHR category</td> <td>2</td> <td>157</td> <td>159</td> </tr> <tr> <td>Normal FHR category</td> <td>0</td> <td>659</td> <td>659</td> </tr> <tr> <td>Totals</td> <td>2</td> <td>816</td> <td>818</td> </tr> </tbody> </table> <p>Sensitivity 100% (19.8-100%)                      Specificity 80.8% (77.8-83.4%)                      Positive likelihood ratio* 5.2 (4.52-5.98)                      Negative likelihood ratio* 0 (NA)</p> <p><i>Low-risk population</i></p> <table border="1" data-bbox="1021 1169 1491 1326"> <thead> <tr> <th></th> <th>Neonatal death</th> <th>No neonatal death</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Neonatal death	No neonatal death	Totals	Indeterminate FHR category	2	157	159	Normal FHR category	0	659	659	Totals	2	816	818		Neonatal death	No neonatal death	Totals					
	Neonatal death	No neonatal death	Totals																										
Indeterminate FHR category	2	157	159																										
Normal FHR category	0	659	659																										
Totals	2	816	818																										
	Neonatal death	No neonatal death	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																												
				<table border="1" data-bbox="1021 272 1487 592"> <tr> <td data-bbox="1021 272 1205 392">Indeterminate FHR category</td> <td data-bbox="1205 272 1301 392">0</td> <td data-bbox="1301 272 1426 392">82</td> <td data-bbox="1426 272 1487 392">82</td> </tr> <tr> <td data-bbox="1021 392 1205 512">Normal FHR category</td> <td data-bbox="1205 392 1301 512">0</td> <td data-bbox="1301 392 1426 512">410</td> <td data-bbox="1426 392 1487 512">410</td> </tr> <tr> <td data-bbox="1021 512 1205 592">Totals</td> <td data-bbox="1205 512 1301 592">0</td> <td data-bbox="1301 512 1426 592">492</td> <td data-bbox="1426 512 1487 592">492</td> </tr> </table> <p data-bbox="1012 600 1496 711">Sensitivity NA** Specificity 83.3% (79.7-86.5)** Positive likelihood ratio 0 (NA)** Negative likelihood ratio 1.20 (NA)**</p> <p data-bbox="1012 735 1232 762"><i>High-risk population</i></p> <table border="1" data-bbox="1021 767 1487 1246"> <thead> <tr> <th data-bbox="1021 767 1205 927"></th> <th data-bbox="1205 767 1301 927">Neonatal death</th> <th data-bbox="1301 767 1426 927">No neonatal death</th> <th data-bbox="1426 767 1487 927">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 927 1205 1046">Indeterminate FHR category</td> <td data-bbox="1205 927 1301 1046">2</td> <td data-bbox="1301 927 1426 1046">75</td> <td data-bbox="1426 927 1487 1046">77</td> </tr> <tr> <td data-bbox="1021 1046 1205 1166">Normal FHR category</td> <td data-bbox="1205 1046 1301 1166">0</td> <td data-bbox="1301 1046 1426 1166">249</td> <td data-bbox="1426 1046 1487 1166">249</td> </tr> <tr> <td data-bbox="1021 1166 1205 1246">Totals</td> <td data-bbox="1205 1166 1301 1246">2</td> <td data-bbox="1301 1166 1426 1246">324</td> <td data-bbox="1426 1166 1487 1246">326</td> </tr> </tbody> </table> <p data-bbox="1012 1254 1496 1366">Sensitivity 100% (19.8-100%) Specificity 76.9% (71.8-81.3%) Positive likelihood ratio* 4.32 (3.54-5.27) Negative likelihood ratio* 0 (NA)</p>	Indeterminate FHR category	0	82	82	Normal FHR category	0	410	410	Totals	0	492	492		Neonatal death	No neonatal death	Totals	Indeterminate FHR category	2	75	77	Normal FHR category	0	249	249	Totals	2	324	326	
Indeterminate FHR category	0	82	82																														
Normal FHR category	0	410	410																														
Totals	0	492	492																														
	Neonatal death	No neonatal death	Totals																														
Indeterminate FHR category	2	75	77																														
Normal FHR category	0	249	249																														
Totals	2	324	326																														

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																						
				<p>* Calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a></p> <p>** Calculated by the NGA technical team using <a href="https://www.medcalc.org/calc/diagnostic_test.php">https://www.medcalc.org/calc/diagnostic_test.php</a></p> <p>Confidence intervals (CIs) calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a></p>																							
<p><b>Full citation</b></p> <p>Soncini, E., Paganelli, S., Vezzani, C., Gargano, G., Giovanni Battista, L. S., Intrapartum fetal heart rate monitoring: evaluation of a standardized system of interpretation for prediction of metabolic acidosis at delivery and neonatal neurological morbidity, Journal of Maternal-Fetal &amp; Neonatal Medicine, 27, 1465-9, 2014</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>N=314</p> <p><b>Characteristics</b></p> <p>The characteristics of the sample:</p> <table border="1"> <tr> <td></td> <td>n=314</td> </tr> <tr> <td>Maternal age in years, mean (SD)</td> <td>30 (5.2)</td> </tr> <tr> <td>Parity 1, %</td> <td>75.5</td> </tr> <tr> <td>Gravidity 1, %</td> <td>53.5</td> </tr> <tr> <td>Gestational age (GA) in weeks, mean (SD)</td> <td>40 (1.2)</td> </tr> </table>		n=314	Maternal age in years, mean (SD)	30 (5.2)	Parity 1, %	75.5	Gravidity 1, %	53.5	Gestational age (GA) in weeks, mean (SD)	40 (1.2)	<p><b>Interventions</b></p> <p>Continuous cardiotocography at least 1 hour and up to 5 hours before birth</p>	<p><b>Details</b></p> <p>FHR tracings were obtained by external transducer ultrasound and recorded Philips Series 50A fetal monitor and Philips Avalon FM 20 fetal monitor; the paper sliding speed was 1 cm/minute. All tracings recorded prior to birth were reviewed by a single expert observer who was blinded to umbilical blood pH, gas values and neonatal</p>	<p><b>Results</b></p> <p>Diagnostic accuracy of FHR tracing classifications on different perinatal outcomes with 95% CI (calculated by the NGA technical team)</p> <p><b>Category III (abnormal) versus Category I (normal)</b></p> <p><u>NICU admission</u></p> <table border="1"> <thead> <tr> <th></th> <th>NICU admission</th> <th>No NICU admission</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>Category III (abnormal)</td> <td>12</td> <td>19</td> <td>27</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> </tbody> </table>		NICU admission	No NICU admission	Total	Category III (abnormal)	12	19	27	Category I (normal)	0	108	108	<p><b>Limitations</b></p> <p>The study was assessed using QUADAS-2 checklist.</p> <ul style="list-style-type: none"> <li>-The study sample was selected and analysed retrospectively with specific inclusion/exclusion criteria, no random sampling</li> <li>-In the study setting, continuous CTG was only performed for labouring women with antenatal or intrapartum risk factors. Also umbilical cord blood sampling was only performed in cases of continuous fetal monitoring and operative birth. Therefore, it is assumed that all the included women are high risk, however, the details of the reason for 'high risk' were not reported</li> <li>-The interpretation of CTG tracings is known to be difficult and subjective and since only one expert reviewed the tracings (two others reviewed 10% of the tracings with good/excellent inter-observer agreement, kappa=0.77) it could be a biased interpretation</li> <li>-The diagnosis of outcomes was likely not done blinded to the index test (CTG tracing), thus, might introduce bias</li> <li>-Index test (CTG tracing) was performed before birth and reference test (ascertainment of outcome) performed during/after birth which might</li> </ul>
	n=314																										
Maternal age in years, mean (SD)	30 (5.2)																										
Parity 1, %	75.5																										
Gravidity 1, %	53.5																										
Gestational age (GA) in weeks, mean (SD)	40 (1.2)																										
	NICU admission	No NICU admission	Total																								
Category III (abnormal)	12	19	27																								
Category I (normal)	0	108	108																								

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
<p>446330</p> <p><b>Country/ies where the study was carried out</b></p> <p>Italy</p> <p><b>Study type</b></p> <p>Retrospective comparative study</p> <p><b>Aim of the study</b></p> <p>To assess the ability of the intrapartum fetal heart rate interpretation system developed in 2008 by the National Institute of Health and Human Development (NICHD) to predict fetal metabolic acidosis at delivery and neonatal neurological morbidity</p>	<p>Spontaneous birth, %</p> <p>Vacuum extraction, %</p> <p>Caesarean section (CS), %</p> <p>Birthweight in g, mean (SD)</p> <p>Small for gestational age (SGA), %</p> <p>1-minute Apgar &lt;7, %</p> <p>5-minute Apgar &lt;7, %</p> <p>Meconium-stained amniotic fluid, %</p> <p>NICU admission, %</p>	<p>37.6</p> <p>25.8</p> <p>36.6</p> <p>3411 (483)</p> <p>12.7</p> <p>17.8</p> <p>2.5</p> <p>36.6</p> <p>7.6</p>	<p>outcome. The analysis included both the dilutant period and the expulsive period, if available. In accordance with NICHD recommendations, both qualitative and quantitative analysis of the FHR tracing was performed. Baseline heart rate, baseline variability, presence of accelerations and decelerations, and uterine contractions were assessed. Tracings were further classified using a three-tier system: Category I (normal), Category II (indeterminate), Category III (abnormal). Trends in FHR</p>	<table border="1"> <tr> <td>Totals</td> <td>12</td> <td>127</td> <td>135</td> </tr> </table> <p>Sensitivity 100% (69.9-100%)                      Specificity 85.0% (77.4-90.5%)                      Positive likelihood ratio 6.68 (4.42-10.12)                      Negative likelihood ratio 0 (NA)</p> <p><u>Neonatal encephalopathy</u></p> <table border="1"> <tr> <td></td> <td>Neonatal encephalopathy</td> <td>No neonatal encephalopathy</td> <td>Totals</td> </tr> <tr> <td>Category III (abnormal)</td> <td>8</td> <td>23</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>8</td> <td>131</td> <td>139</td> </tr> </table> <p>Sensitivity 100% (59.8-100%)                      Specificity 82.4% (74.6-88.3%)                      Positive likelihood ratio 5.70 (3.93-8.25)</p>	Totals	12	127	135		Neonatal encephalopathy	No neonatal encephalopathy	Totals	Category III (abnormal)	8	23	31	Category I (normal)	0	108	108	Totals	8	131	139	<p>mean that events after the index test influenced the outcome independently of the index test</p> <p><b>Other information</b></p>
Totals	12	127	135																						
	Neonatal encephalopathy	No neonatal encephalopathy	Totals																						
Category III (abnormal)	8	23	31																						
Category I (normal)	0	108	108																						
Totals	8	131	139																						



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																						
<p><b>Study dates</b> August 2007 to May 2011</p> <p><b>Source of funding</b> None reported</p>	<table border="1"> <tr> <td>Neonatal encephalopathy, %</td> <td>3.5</td> </tr> <tr> <td>Moderate-severe neonatal encephalopathy, %</td> <td>1.6</td> </tr> <tr> <td>Death before NICU discharge, %</td> <td>1.0</td> </tr> </table> <p>The proportion of FHR tracings was as follows: Category I 34.4%, Category IIA 37.6%, Category IIB 18.2%, and Category III 9.8%. No statistically significant differences were found between groups in terms of parity, gestational age, oligohydramnios, induction of labour or mode of birth. Rate of operative delivery for suspected fetal distress increased significantly with worsening FHR pattern</p> <p><b>Inclusion criteria</b></p>	Neonatal encephalopathy, %	3.5	Moderate-severe neonatal encephalopathy, %	1.6	Death before NICU discharge, %	1.0		<p>patterns over time were quantified in minutes. Abnormal FHR patterns lasting longer than 30 minutes fell into Category III. Indeterminate FHR patterns lasting longer than 30 minutes fell within Category II. Otherwise, tracings were classified as Category I. When both indeterminate and abnormal FHR patterns were present in the same tracing, with each FHR pattern lasting under 30 minutes but overall total more than 30 minutes, it was classified as Category II. Category II was further divided into two subcategories</p>	<p>Negative likelihood ratio 0 (NA)</p> <p><u>Moderate-severe neonatal encephalopathy</u></p> <table border="1"> <thead> <tr> <th></th> <th>Moderate-severe neonatal encephalopathy</th> <th>No moderate-severe neonatal encephalopathy</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category III (abnormal)</td> <td>4</td> <td>27</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>4</td> <td>135</td> <td>139</td> </tr> </tbody> </table> <p>Sensitivity 100% (39.6-100%) Specificity 80.0% (72.1-86.2%) Positive likelihood ratio 5.00 (3.57-7.01) Negative likelihood ratio 0 (NA)</p> <p><u>Instrumental birth</u></p>		Moderate-severe neonatal encephalopathy	No moderate-severe neonatal encephalopathy	Totals	Category III (abnormal)	4	27	31	Category I (normal)	0	108	108	Totals	4	135	139	
Neonatal encephalopathy, %	3.5																										
Moderate-severe neonatal encephalopathy, %	1.6																										
Death before NICU discharge, %	1.0																										
	Moderate-severe neonatal encephalopathy	No moderate-severe neonatal encephalopathy	Totals																								
Category III (abnormal)	4	27	31																								
Category I (normal)	0	108	108																								
Totals	4	135	139																								

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
	<p>All labouring women, monitored with continuous cardiotocography, carrying singleton fetuses with cephalic presentation at <math>\geq 37</math> weeks of gestation whose umbilical artery blood gas and acid-base analysis at birth was available</p> <p><b>Exclusion criteria</b></p> <p>Cases with fetal malformation, arrhythmia, elective caesarean section, or absence of significant uterine contractions (fewer than 3 contractions in 10 minutes). Cases with no fetal heart rate tracing available in the last hour prior to birth</p>		<p>according to the 2010 American College of Obstetricians and Gynecologists management guidelines. Within this study, the authors denoted the two subcategories Category IIA and IIB. Tracings with moderate FHR variability or FHR accelerations were classified as Category IIA and tracings with minimal/absent baseline FHR variability and no FHR accelerations were classified as Category IIB. To assess the reproducibility of heart rate readings, a second and a third</p>	<table border="1" data-bbox="1021 304 1491 911"> <thead> <tr> <th></th> <th>Instrumental birth</th> <th>No instrumental birth</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category III (abnormal)</td> <td>19</td> <td>12</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>74</td> <td>34</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>93</td> <td>46</td> <td>139</td> </tr> </tbody> </table> <p>Sensitivity 20.4% (13.0-30.3%) Specificity 73.9% (58.6-85.2%) Positive likelihood ratio 0.78 (0.42-1.47) Negative likelihood ratio 1.08 (0.96-1.21)</p> <p><u>Instrumental birth for suspected fetal distress</u></p> <table border="1" data-bbox="1021 1169 1491 1342"> <thead> <tr> <th></th> <th>Instrumental birth for suspected</th> <th>No instrumental birth for suspected</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Instrumental birth	No instrumental birth	Totals	Category III (abnormal)	19	12	31	Category I (normal)	74	34	108	Totals	93	46	139		Instrumental birth for suspected	No instrumental birth for suspected	Totals					
	Instrumental birth	No instrumental birth	Totals																										
Category III (abnormal)	19	12	31																										
Category I (normal)	74	34	108																										
Totals	93	46	139																										
	Instrumental birth for suspected	No instrumental birth for suspected	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
			investigator further reviewed tracings independently in 10% of cases		d fetal distress	fetal distress		
Category III (abnormal)	18	13	31	Category I (normal)	24	84	108	
Totals	42	97	139	Sensitivity 42.9% (28.1-58.9%) Specificity 86.6% (77.8-92.4%) Positive likelihood ratio 3.20 (1.73-5.91) Negative likelihood ratio 0.66 (0.51-0.86)				
<u>Death before NICU discharge</u>					Death before NICU discharge	No death before NICU discharge	Totals	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																												
				<table border="1"> <tr> <td>Category III (abnormal)</td> <td>3</td> <td>28</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>3</td> <td>136</td> <td>139</td> </tr> </table> <p>Sensitivity 100% (31.0-100%)                      Specificity 79.4% (71.4-85.7%)                      Positive likelihood ratio 4.86 (3.49-6.76)                      Negative likelihood ratio 0 (NA)</p> <p><u>pH &lt;7</u></p> <table border="1"> <tr> <td></td> <td>pH &lt;7</td> <td>pH ≥7</td> <td>Totals</td> </tr> <tr> <td>Category III (abnormal)</td> <td>17</td> <td>14</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>17</td> <td>122</td> <td>139</td> </tr> </table>	Category III (abnormal)	3	28	31	Category I (normal)	0	108	108	Totals	3	136	139		pH <7	pH ≥7	Totals	Category III (abnormal)	17	14	31	Category I (normal)	0	108	108	Totals	17	122	139	
Category III (abnormal)	3	28	31																														
Category I (normal)	0	108	108																														
Totals	3	136	139																														
	pH <7	pH ≥7	Totals																														
Category III (abnormal)	17	14	31																														
Category I (normal)	0	108	108																														
Totals	17	122	139																														

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
				<p>Sensitivity 100% (77.1-100%)                      Specificity 88.5% (81.2-93.3%)                      Positive likelihood ratio 8.71 (5.32-14.27)                      Negative likelihood ratio 0 (NA)</p> <p><u>Base excess (BE) &lt;=-12 mmol/l</u></p> <table border="1" data-bbox="1021 472 1442 995"> <tr> <td></td> <td>BE &lt;=- 12 mmol/l</td> <td>BE &gt;= 12 mmol/l</td> <td>Totals</td> </tr> <tr> <td>Category III (abnormal)</td> <td>19</td> <td>12</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>3</td> <td>105</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>22</td> <td>117</td> <td>139</td> </tr> </table> <p>Sensitivity 86.4% (64.0-96.4%)                      Specificity 89.7% (82.4-94.4%)                      Positive likelihood ratio 8.42 (4.80-14.76)                      Negative likelihood ratio 0.15 (0.05-0.44)</p> <p><u>pH &lt;7 and BE &lt;=-12 mmol/l</u></p> <table border="1" data-bbox="1021 1225 1442 1315"> <tr> <td></td> <td>pH &lt;7 and BE</td> <td>pH ≥7 and BE</td> <td>Totals</td> </tr> </table>		BE <=- 12 mmol/l	BE >= 12 mmol/l	Totals	Category III (abnormal)	19	12	31	Category I (normal)	3	105	108	Totals	22	117	139		pH <7 and BE	pH ≥7 and BE	Totals	
	BE <=- 12 mmol/l	BE >= 12 mmol/l	Totals																						
Category III (abnormal)	19	12	31																						
Category I (normal)	3	105	108																						
Totals	22	117	139																						
	pH <7 and BE	pH ≥7 and BE	Totals																						

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1"> <thead> <tr> <th></th> <th>&lt;=-12 mmol/l</th> <th>&gt;-12 mmol/l</th> <th></th> </tr> </thead> <tbody> <tr> <td>Category III (abnormal)</td> <td>14</td> <td>17</td> <td>31</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>14</td> <td>125</td> <td>139</td> </tr> </tbody> </table> <p>Sensitivity 100% (73.2-100%)                      Specificity 86.4% (78.8-91.6%)                      Positive likelihood ratio 7.35 (4.73-11.44)                      Negative likelihood ratio 0 (NA)</p> <p><b>Category IIB (indeterminate B) versus                      Category I (normal)                      NICU admission</b></p> <table border="1"> <thead> <tr> <th></th> <th>NICU admission</th> <th>No NICU admission</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>9</td> <td>48</td> <td>57</td> </tr> </tbody> </table>		<=-12 mmol/l	>-12 mmol/l		Category III (abnormal)	14	17	31	Category I (normal)	0	108	108	Totals	14	125	139		NICU admission	No NICU admission	Totals	Category IIB	9	48	57	
	<=-12 mmol/l	>-12 mmol/l																											
Category III (abnormal)	14	17	31																										
Category I (normal)	0	108	108																										
Totals	14	125	139																										
	NICU admission	No NICU admission	Totals																										
Category IIB	9	48	57																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1"> <tr> <td data-bbox="1021 272 1137 432">Category I (normal)</td> <td data-bbox="1137 272 1267 432">0</td> <td data-bbox="1267 272 1397 432">108</td> <td data-bbox="1397 272 1491 432">108</td> </tr> <tr> <td data-bbox="1021 432 1137 512">Totals</td> <td data-bbox="1137 432 1267 512">9</td> <td data-bbox="1267 432 1397 512">156</td> <td data-bbox="1397 432 1491 512">165</td> </tr> </table> <p data-bbox="1021 544 1491 655">Sensitivity 100% (62.9-100%) Specificity 69.2% (61.3-76.2%) Positive likelihood ratio 3.25 (2.57-4.11) Negative likelihood ratio 0 (NA)</p> <p data-bbox="1021 679 1290 711"><u>Neonatal encephalopathy</u></p> <table border="1"> <thead> <tr> <th data-bbox="1021 743 1111 943"></th> <th data-bbox="1111 743 1267 943">Neonatal encephalopathy</th> <th data-bbox="1267 743 1424 943">No Neonatal encephalopathy</th> <th data-bbox="1424 743 1491 943">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 943 1111 1062">Category IIB</td> <td data-bbox="1111 943 1267 1062">3</td> <td data-bbox="1267 943 1424 1062">54</td> <td data-bbox="1424 943 1491 1062">57</td> </tr> <tr> <td data-bbox="1021 1062 1111 1262">Category I (normal)</td> <td data-bbox="1111 1062 1267 1262">0</td> <td data-bbox="1267 1062 1424 1262">108</td> <td data-bbox="1424 1062 1491 1262">108</td> </tr> <tr> <td data-bbox="1021 1262 1111 1342">Totals</td> <td data-bbox="1111 1262 1267 1342">3</td> <td data-bbox="1267 1262 1424 1342">162</td> <td data-bbox="1424 1262 1491 1342">165</td> </tr> </tbody> </table>	Category I (normal)	0	108	108	Totals	9	156	165		Neonatal encephalopathy	No Neonatal encephalopathy	Totals	Category IIB	3	54	57	Category I (normal)	0	108	108	Totals	3	162	165	
Category I (normal)	0	108	108																										
Totals	9	156	165																										
	Neonatal encephalopathy	No Neonatal encephalopathy	Totals																										
Category IIB	3	54	57																										
Category I (normal)	0	108	108																										
Totals	3	162	165																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																
				<p>Sensitivity 100% (31.0-100%)                      Specificity 66.7% (58.8-73.8%)                      Positive likelihood ratio 3.00 (2.41-3.73)                      Negative likelihood ratio 0 (NA)</p> <p><u>Moderate-severe neonatal encephalopathy</u></p> <table border="1" data-bbox="1021 499 1491 1145"> <thead> <tr> <th></th> <th>Moderate-severe neonatal encephalopathy</th> <th>No moderate-severe neonatal encephalopathy</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>1</td> <td>56</td> <td>57</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>1</td> <td>164</td> <td>165</td> </tr> </tbody> </table> <p>Sensitivity 100% (5.5-100%)                      Specificity 65.9% (58.0-73.0%)                      Positive likelihood ratio 2.93 (2.37-3.62)                      Negative likelihood ratio 0 (NA)</p> <p><u>Instrumental birth</u></p>		Moderate-severe neonatal encephalopathy	No moderate-severe neonatal encephalopathy	Totals	Category IIB	1	56	57	Category I (normal)	0	108	108	Totals	1	164	165	
	Moderate-severe neonatal encephalopathy	No moderate-severe neonatal encephalopathy	Totals																		
Category IIB	1	56	57																		
Category I (normal)	0	108	108																		
Totals	1	164	165																		



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 304 1491 826"> <thead> <tr> <th></th> <th>Instrumental birth</th> <th>No instrumental birth</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>30</td> <td>27</td> <td>57</td> </tr> <tr> <td>Category I (normal)</td> <td>74</td> <td>34</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>104</td> <td>61</td> <td>165</td> </tr> </tbody> </table> <p data-bbox="1021 863 1491 975">Sensitivity 28.9% (20.6-38.7%) Specificity 55.7% (42.5-68.2%) Positive likelihood ratio 0.65 (0.43-0.98) Negative likelihood ratio 1.28 (1.10-1.48)</p> <p data-bbox="1021 1031 1491 1054"><u>Instrumental birth for suspected fetal distress</u></p> <table border="1" data-bbox="1021 1086 1491 1366"> <thead> <tr> <th></th> <th>Instrumental birth for suspected fetal distress</th> <th>No instrumental birth for suspected fetal distress</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		Instrumental birth	No instrumental birth	Totals	Category IIB	30	27	57	Category I (normal)	74	34	108	Totals	104	61	165		Instrumental birth for suspected fetal distress	No instrumental birth for suspected fetal distress	Totals					
	Instrumental birth	No instrumental birth	Totals																										
Category IIB	30	27	57																										
Category I (normal)	74	34	108																										
Totals	104	61	165																										
	Instrumental birth for suspected fetal distress	No instrumental birth for suspected fetal distress	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																				
				<table border="1"> <tr> <td data-bbox="1019 272 1115 395">Category IIB</td> <td data-bbox="1115 272 1254 395">29</td> <td data-bbox="1254 272 1422 395">28</td> <td data-bbox="1422 272 1494 395">57</td> </tr> <tr> <td data-bbox="1019 395 1115 598">Category I (normal)</td> <td data-bbox="1115 395 1254 598">24</td> <td data-bbox="1254 395 1422 598">84</td> <td data-bbox="1422 395 1494 598">108</td> </tr> <tr> <td data-bbox="1019 598 1115 678">Totals</td> <td data-bbox="1115 598 1254 678">53</td> <td data-bbox="1254 598 1422 678">112</td> <td data-bbox="1422 598 1494 678">165</td> </tr> </table> <p data-bbox="1019 710 1494 821">Sensitivity 54.7% (40.6-68.2%) Specificity 75.0% (65.8-82.5%) Positive likelihood ratio 2.19 (1.46-3.28) Negative likelihood ratio 0.60 (0.45-0.82)</p> <p data-bbox="1019 869 1494 901"><u>Death before NICU discharge</u></p> <table border="1"> <tr> <td data-bbox="1019 933 1137 1136"></td> <td data-bbox="1137 933 1261 1136">Death before NICU discharge</td> <td data-bbox="1261 933 1391 1136">No death before NICU discharge</td> <td data-bbox="1391 933 1471 1136">Totals</td> </tr> <tr> <td data-bbox="1019 1136 1137 1257">Category IIB</td> <td data-bbox="1137 1136 1261 1257">0</td> <td data-bbox="1261 1136 1391 1257">57</td> <td data-bbox="1391 1136 1471 1257">57</td> </tr> </table>	Category IIB	29	28	57	Category I (normal)	24	84	108	Totals	53	112	165		Death before NICU discharge	No death before NICU discharge	Totals	Category IIB	0	57	57	
Category IIB	29	28	57																						
Category I (normal)	24	84	108																						
Totals	53	112	165																						
	Death before NICU discharge	No death before NICU discharge	Totals																						
Category IIB	0	57	57																						

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 279 1469 512"> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>0</td> <td>165</td> <td>165</td> </tr> </table> <p data-bbox="1012 549 1391 655">                     Sensitivity NA*                      Specificity 65.5% (57.7-72.7%)*                      Positive likelihood ratio 0 (NA)*                      Negative likelihood ratio 1.53 (NA)*                 </p> <p data-bbox="1012 687 1084 711"><u>pH &lt;7</u></p> <table border="1" data-bbox="1021 746 1391 1142"> <thead> <tr> <th></th> <th>pH &lt;7</th> <th>pH ≥7</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>7</td> <td>50</td> <td>57</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>7</td> <td>158</td> <td>165</td> </tr> </tbody> </table> <p data-bbox="1012 1179 1435 1286">                     Sensitivity 100% (56.1-100%)                      Specificity 68.4% (60.4-75.4%)                      Positive likelihood ratio 3.16 (2.51-3.97)                      Negative likelihood ratio 0 (NA)                 </p> <p data-bbox="1012 1318 1196 1342"><u>BE ≤-12 mmol/l</u></p>	Category I (normal)	0	108	108	Totals	0	165	165		pH <7	pH ≥7	Totals	Category IIB	7	50	57	Category I (normal)	0	108	108	Totals	7	158	165	
Category I (normal)	0	108	108																										
Totals	0	165	165																										
	pH <7	pH ≥7	Totals																										
Category IIB	7	50	57																										
Category I (normal)	0	108	108																										
Totals	7	158	165																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1"> <thead> <tr> <th></th> <th>BE &lt;=-12 mmol/l</th> <th>BE &gt;-12 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>14</td> <td>43</td> <td>57</td> </tr> <tr> <td>Category I (normal)</td> <td>3</td> <td>105</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>17</td> <td>148</td> <td>165</td> </tr> </tbody> </table> <p>Sensitivity 82.4% (55.8-95.3%)                      Specificity 71.0% (62.8-78.0%)                      Positive likelihood ratio 2.83 (2.03-3.96)                      Negative likelihood ratio 0.25 (0.09-0.70)</p> <p><u>pH &lt;7 and BE &lt;=-12 mmol/l</u></p> <table border="1"> <thead> <tr> <th></th> <th>pH &lt;7 and BE &lt;=-12 mmol/l</th> <th>pH ≥7 and BE &gt;-12 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIB</td> <td>4</td> <td>53</td> <td>57</td> </tr> </tbody> </table>		BE <=-12 mmol/l	BE >-12 mmol/l	Totals	Category IIB	14	43	57	Category I (normal)	3	105	108	Totals	17	148	165		pH <7 and BE <=-12 mmol/l	pH ≥7 and BE >-12 mmol/l	Totals	Category IIB	4	53	57	
	BE <=-12 mmol/l	BE >-12 mmol/l	Totals																										
Category IIB	14	43	57																										
Category I (normal)	3	105	108																										
Totals	17	148	165																										
	pH <7 and BE <=-12 mmol/l	pH ≥7 and BE >-12 mmol/l	Totals																										
Category IIB	4	53	57																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 272 1417 515"> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>4</td> <td>161</td> <td>165</td> </tr> </table> <p data-bbox="1021 547 1440 659">Sensitivity 100% (39.6-100%) Specificity 67.1% (59.2-74.2%) Positive likelihood ratio 3.04 (2.44-3.79) Negative likelihood ratio 0 (NA)</p> <p data-bbox="1021 683 1447 767"><b>Category IIA (indeterminate A) versus Category I (normal)</b> <u>NICU admission</u></p> <table border="1" data-bbox="1021 799 1485 1281"> <thead> <tr> <th></th> <th>NICU admission</th> <th>No NICU admission</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIA</td> <td>3</td> <td>115</td> <td>118</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>3</td> <td>223</td> <td>226</td> </tr> </tbody> </table> <p data-bbox="1021 1313 1346 1369">Sensitivity 100% (31.0-100%) Specificity 48.4% (41.7-55.2%)</p>	Category I (normal)	0	108	108	Totals	4	161	165		NICU admission	No NICU admission	Totals	Category IIA	3	115	118	Category I (normal)	0	108	108	Totals	3	223	226	
Category I (normal)	0	108	108																										
Totals	4	161	165																										
	NICU admission	No NICU admission	Totals																										
Category IIA	3	115	118																										
Category I (normal)	0	108	108																										
Totals	3	223	226																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																
				<p>Positive likelihood ratio 1.94 (1.71-2.20)                      Negative likelihood ratio 0 (NA)</p> <p><u>Neonatal encephalopathy</u></p> <table border="1" data-bbox="1021 416 1489 1023"> <thead> <tr> <th></th> <th>Neonatal encephalopathy</th> <th>No neonatal encephalopathy</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIA</td> <td>0</td> <td>118</td> <td>118</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>0</td> <td>226</td> <td>226</td> </tr> </tbody> </table> <p>Sensitivity NA*                      Specificity 47.8% (41.1-54.5%)*                      Positive likelihood ratio 0 (NA)*                      Negative likelihood ratio 2.09 (NA)*</p> <p><u>Moderate-severe neonatal encephalopathy</u></p>		Neonatal encephalopathy	No neonatal encephalopathy	Totals	Category IIA	0	118	118	Category I (normal)	0	108	108	Totals	0	226	226	
	Neonatal encephalopathy	No neonatal encephalopathy	Totals																		
Category IIA	0	118	118																		
Category I (normal)	0	108	108																		
Totals	0	226	226																		

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments
					Moderate-severe neonatal encephalopathy	No moderate-severe neonatal encephalopathy	Totals	
				Category IIA	0	118	118	
				Category I (normal)	0	108	108	
				Totals	0	226	226	
				Sensitivity NA* Specificity 47.8% (41.1-54.5%)* Positive likelihood ratio 0 (NA)* Negative likelihood ratio 2.09 (NA)*				
				<u>Instrumental birth</u>				
					Instrumental birth	No instrumental birth	Totals	

Study details	Participants	Interventions	Methods	Outcomes and Results				Comments																				
				<table border="1" data-bbox="1021 272 1491 678"> <tr> <td data-bbox="1021 272 1115 395">Category IIA</td> <td data-bbox="1115 272 1254 395">73</td> <td data-bbox="1254 272 1422 395">45</td> <td data-bbox="1422 272 1491 395">118</td> </tr> <tr> <td data-bbox="1021 395 1115 595">Category I (normal)</td> <td data-bbox="1115 395 1254 595">74</td> <td data-bbox="1254 395 1422 595">34</td> <td data-bbox="1422 395 1491 595">108</td> </tr> <tr> <td data-bbox="1021 595 1115 678">Totals</td> <td data-bbox="1115 595 1254 678">147</td> <td data-bbox="1254 595 1422 678">79</td> <td data-bbox="1422 595 1491 678">226</td> </tr> </table> <p data-bbox="1021 710 1491 821">Sensitivity 49.7% (41.4-58.0%) Specificity 43.0% (32.1-54.6%) Positive likelihood ratio 0.87 (0.68-1.12) Negative likelihood ratio 1.17 (0.96-1.42)</p> <p data-bbox="1021 874 1491 901"><u>Instrumental birth for suspected fetal distress</u></p> <table border="1" data-bbox="1021 933 1491 1337"> <tr> <td data-bbox="1021 933 1115 1217"></td> <td data-bbox="1115 933 1254 1217">Instrumental birth for suspected fetal distress</td> <td data-bbox="1254 933 1422 1217">No instrumental birth for suspected fetal distress</td> <td data-bbox="1422 933 1491 1217">Totals</td> </tr> <tr> <td data-bbox="1021 1217 1115 1337">Category IIA</td> <td data-bbox="1115 1217 1254 1337">50</td> <td data-bbox="1254 1217 1422 1337">68</td> <td data-bbox="1422 1217 1491 1337">118</td> </tr> </table>				Category IIA	73	45	118	Category I (normal)	74	34	108	Totals	147	79	226		Instrumental birth for suspected fetal distress	No instrumental birth for suspected fetal distress	Totals	Category IIA	50	68	118	
Category IIA	73	45	118																									
Category I (normal)	74	34	108																									
Totals	147	79	226																									
	Instrumental birth for suspected fetal distress	No instrumental birth for suspected fetal distress	Totals																									
Category IIA	50	68	118																									



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1021 272 1491 555"> <tr> <td data-bbox="1021 272 1115 475">Category I (normal)</td> <td data-bbox="1115 272 1256 475">24</td> <td data-bbox="1256 272 1420 475">84</td> <td data-bbox="1420 272 1491 475">108</td> </tr> <tr> <td data-bbox="1021 475 1115 555">Totals</td> <td data-bbox="1115 475 1256 555">74</td> <td data-bbox="1256 475 1420 555">152</td> <td data-bbox="1420 475 1491 555">226</td> </tr> </table> <p data-bbox="1021 587 1491 699">Sensitivity 67.6% (55.6-77.7%) Specificity 55.3% (47.0-63.3%) Positive likelihood ratio 1.51 (1.19-1.91) Negative likelihood ratio 0.59 (0.42-0.82)</p> <p data-bbox="1021 754 1491 783"><u>Death before NICU discharge</u></p> <table border="1" data-bbox="1021 810 1491 1377"> <thead> <tr> <th data-bbox="1021 810 1133 1013"></th> <th data-bbox="1133 810 1256 1013">Death before NICU discharge</th> <th data-bbox="1256 810 1391 1013">No death before NICU discharge</th> <th data-bbox="1391 810 1491 1013">Totals</th> </tr> </thead> <tbody> <tr> <td data-bbox="1021 1013 1133 1134">Category IIA</td> <td data-bbox="1133 1013 1256 1134">0</td> <td data-bbox="1256 1013 1391 1134">118</td> <td data-bbox="1391 1013 1491 1134">118</td> </tr> <tr> <td data-bbox="1021 1134 1133 1294">Category I (normal)</td> <td data-bbox="1133 1134 1256 1294">0</td> <td data-bbox="1256 1134 1391 1294">108</td> <td data-bbox="1391 1134 1491 1294">108</td> </tr> <tr> <td data-bbox="1021 1294 1133 1377">Totals</td> <td data-bbox="1133 1294 1256 1377">0</td> <td data-bbox="1256 1294 1391 1377">226</td> <td data-bbox="1391 1294 1491 1377">226</td> </tr> </tbody> </table>	Category I (normal)	24	84	108	Totals	74	152	226		Death before NICU discharge	No death before NICU discharge	Totals	Category IIA	0	118	118	Category I (normal)	0	108	108	Totals	0	226	226	
Category I (normal)	24	84	108																										
Totals	74	152	226																										
	Death before NICU discharge	No death before NICU discharge	Totals																										
Category IIA	0	118	118																										
Category I (normal)	0	108	108																										
Totals	0	226	226																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<p>Sensitivity NA*</p> <p>Specificity 47.8% (41.1-54.5%)*</p> <p>Positive likelihood ratio 0 (NA)*</p> <p>Negative likelihood ratio 2.09 (NA)*</p> <p><u>pH &lt;7</u></p> <table border="1" data-bbox="1021 499 1391 898"> <thead> <tr> <th></th> <th>pH &lt;7</th> <th>pH ≥7</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIA</td> <td>0</td> <td>118</td> <td>118</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>0</td> <td>226</td> <td>226</td> </tr> </tbody> </table> <p>Sensitivity NA*</p> <p>Specificity 47.8% (41.1-54.5%)*</p> <p>Positive likelihood ratio 0 (NA)*</p> <p>Negative likelihood ratio 2.09 (NA)*</p> <p><u>BE &lt;=-12 mmol/l</u></p> <table border="1" data-bbox="1021 1129 1417 1289"> <thead> <tr> <th></th> <th>BE &lt;=-12 mmol/l</th> <th>BE &gt;-12 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>		pH <7	pH ≥7	Totals	Category IIA	0	118	118	Category I (normal)	0	108	108	Totals	0	226	226		BE <=-12 mmol/l	BE >-12 mmol/l	Totals					
	pH <7	pH ≥7	Totals																										
Category IIA	0	118	118																										
Category I (normal)	0	108	108																										
Totals	0	226	226																										
	BE <=-12 mmol/l	BE >-12 mmol/l	Totals																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																								
				<table border="1" data-bbox="1019 272 1417 635"> <tr> <td>Category IIA</td> <td>2</td> <td>116</td> <td>118</td> </tr> <tr> <td>Category I (normal)</td> <td>3</td> <td>105</td> <td>108</td> </tr> <tr> <td>Totals</td> <td>5</td> <td>221</td> <td>226</td> </tr> </table> <p data-bbox="1019 667 1451 778">                     Sensitivity 40.0% (7.3-83.0%)                      Specificity 47.5% (40.8-54.3%)                      Positive likelihood ratio 0.76 (0.26-2.25)                      Negative likelihood ratio 1.26 (0.61-2.61)                 </p> <p data-bbox="1019 805 1317 831"> <u>pH &lt;7 and BE &lt;=-12 mmol/l</u> </p> <table border="1" data-bbox="1019 866 1417 1350"> <thead> <tr> <th></th> <th>pH &lt;7 and BE &lt;=-12 mmol/l</th> <th>pH ≥7 and BE &gt;-12 mmol/l</th> <th>Totals</th> </tr> </thead> <tbody> <tr> <td>Category IIA</td> <td>0</td> <td>118</td> <td>118</td> </tr> <tr> <td>Category I (normal)</td> <td>0</td> <td>108</td> <td>108</td> </tr> </tbody> </table>	Category IIA	2	116	118	Category I (normal)	3	105	108	Totals	5	221	226		pH <7 and BE <=-12 mmol/l	pH ≥7 and BE >-12 mmol/l	Totals	Category IIA	0	118	118	Category I (normal)	0	108	108	
Category IIA	2	116	118																										
Category I (normal)	3	105	108																										
Totals	5	221	226																										
	pH <7 and BE <=-12 mmol/l	pH ≥7 and BE >-12 mmol/l	Totals																										
Category IIA	0	118	118																										
Category I (normal)	0	108	108																										

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments				
				<table border="1" data-bbox="1019 276 1415 347"> <tr> <td>Totals</td> <td>0</td> <td>226</td> <td>226</td> </tr> </table> <p>Sensitivity NA*                      Specificity 47.8% (41.1-54.5%)*                      Positive likelihood ratio 0 (NA)*                      Negative likelihood ratio 2.09 (NA)*</p> <p>Sensitivity, specificity and likelihood ratios calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a> unless marked with *</p> <p>*Sensitivity, specificity and likelihood ratios calculated by the NGA technical team using <a href="https://www.medcalc.org/calc/diagnostic_test.php">https://www.medcalc.org/calc/diagnostic_test.php</a></p>	Totals	0	226	226	
Totals	0	226	226						
<p><b>Full citation</b></p> <p>Berkus,M.D., Langer,O., Samueloff,A., Xenakis,E.M., Field,N.T., Electronic fetal monitoring: what's reassuring?, Acta Obstetrica et Gynecologica Scandinavica, 78, 15-21, 1999</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>n = 2200 consecutive singleton term pregnancies</p> <p>n = 484/2200 (26%) with normal FHR trace during the last 30 minutes prior to delivery</p> <p><b>Characteristics</b></p> <p>There were no significant differences observed between the</p>	<p><b>Interventions</b></p> <p><b>Normal</b>                      Baseline 120–160 bpm                      Variability &gt; 5 bpm                      Presence of accelerations                      No variable or late decelerations</p> <p><b>Abnormal</b>                      Baseline 90–120 bpm or &gt; 160 bpm</p>	<p><b>Details</b></p> <p>A cohort of n = 2200 consecutive birth was examined and the fetal heart rate tracings analysed. Arterial blood gas was collected from 97.5% of the study population. Blood sample</p>	<p><b>Results</b></p> <p><u>Association between abnormal FHR tracing patterns and immediate adverse outcome (1st stage n = 224)</u></p> <p>Mild or moderate variable deceleration: not significant (ns)                      Decreased variability: ns                      Mild bradycardia: ns                      Tachycardia: ns                      Prolonged bradycardia: OR 1.9 (95% CI 1.3 to 3.7)                      Severe variable deceleration: ns                      late deceleration: ns</p> <p><u>Association between abnormal FHR tracing patterns and cord pH &lt; 7.15 &amp; 5 min apgar</u></p>	<p><b>Limitations</b></p> <p>No separate data for Apgar and pH</p> <p><b>Other information</b></p> <p><u>Reassuring (normal) trace defined as:</u>                      Any tracing with acceleration                      Had mild variables                      Had decreased variability                      Had mild bradycardia                      Had any above combination</p> <p><u>Non-reassuring (abnormal) trace defined as:</u>                      No acceleration                      Severe or late deceleration                      Prolonged bradycardia</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>196611</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To determine which combinations of fetal heart rate (FHR) pattern abnormalities are associated with normal outcome in term pregnancies</p> <p><b>Study dates</b></p> <p>From March to August 1991</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>reassuring and non-reassuring group in fetal gestational age, sex, birth weight, and fetal complications.</p> <p>Women with non-reassuring tracing were significantly older, more often primigravida, had more maternal illness (cardiovascular, thyroid, kidney disease or diabetes) and more caesarean section and instrumental birth.</p> <p>However, there was no statistically significant differences in pregnancy complications (hypertension, infection, post-date, substance abuse, meconium stained liquor).</p> <p><b>Inclusion criteria</b></p> <p>Term pregnancy (&gt; 36 weeks or birth weight &gt; 2500g)</p> <p>Live birth</p> <p>Singleton pregnancy</p>	<p>Variability &lt; 5 bpm</p> <p>No accelerations</p> <p>Any decelerations</p> <p>Prolonged bradycardia or any combination</p>	<p>was drawn immediately after birth and analysed within 30 minutes of birth. Every women entering the delivery room had FHR trace performed. The last 30 minutes of trace segment prior to delivery was analysed. All traces were obtained by scalp electrocardiography, and observers that analysed the data were blinded to birth outcomes.</p>	<p><u>score &lt; 7 (first stage n = 224)</u></p> <p>Mild or moderate variable deceleration: ns</p> <p>Decreased variability: ns</p> <p>Mild bradycardia: ns</p> <p>Tachycardia: ns</p> <p>Prolonged bradycardia: ns</p> <p>Severe variable deceleration: ns</p> <p>Late deceleration: ns</p> <p><u>Association between abnormal FHR tracing patterns and immediate adverse outcome (second stage n = 1635)</u></p> <p>Mild or moderate variable deceleration: ns</p> <p>Decreased variability: ns</p> <p>Mild bradycardia: ns</p> <p>Tachycardia: OR 1.9 (95% CI 1.2 to 2.8)</p> <p>Prolong bradycardia: ns</p> <p>Severe variable deceleration: ns</p> <p>Late deceleration: ns</p> <p><u>Association between abnormal FHR tracing patterns and cord pH &lt; 7.15 &amp; 5 min apgar score &lt; 7 (second stage n = 1635)</u></p> <p>Mild or moderate variable deceleration: ns</p> <p>Decreased variability: ns</p> <p>Mild bradycardia: ns</p> <p>Tachycardia: ns</p> <p>Prolonged bradycardia: OR 3.6 (95% CI 1.2 to 11)</p> <p>Severe variable deceleration: OR 2.4 (95% CI 1.2 to 4)</p> <p>Late deceleration: OR 6.9 (95% CI 2.1 to 23)</p> <p>Decreased variability: ≤ 5 bpm</p> <p>Mild bradycardia: 90 &lt; FHR &lt; 120 bpm</p> <p>Tachycardia: 120 &lt; FHR &lt; 160 bpm</p> <p>Prolonged bradycardia: &lt; 90 bpm, &gt; 2.5 min</p>	<p>Tachycardia any above combination</p> <p><u>Neonates were assessed to have immediate adverse outcomes if they:</u></p> <p>were admitted to level III, neonatal intensive care unit for &gt; 24 hours and required oxygen support (intubation &gt; 6 hrs, or &gt; 24 hrs of &gt; 40% oxygen supplementation)</p> <p>had significant complications (intracranial haemorrhage, neonatal death)</p> <p>experienced neurological sequelae (seizure, persistent hypotonia at discharge)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><b>Exclusion criteria</b></p> <p>Choriamnionitis</p> <p>Major congenital abnormalities</p>				
<p><b>Full citation</b></p> <p>Cardoso,C.G., Graca,L.M., Clode,N., A study on second-stage cardiotocographic patterns and umbilical blood acid-base balance in cases with first-stage normal fetal heart rates, Journal of Maternal-Fetal Investigation, 5, 144-147, 1995</p> <p><b>Ref Id</b></p> <p>197264</p> <p><b>Country/ies where the study was carried out</b></p> <p>Portugal</p> <p><b>Study type</b></p> <p>Cohort</p>	<p><b>Sample size</b></p> <p>n = 293 singleton term pregnancies. Normal 1st stage traces, analysed on all of second stage. Classified on modified Melchior and Barnard classification. n = 103 type 0 used as controls.</p> <p><b>Characteristics</b></p> <p>Instrumental vaginal birth performed in 10 cases of 0 type (9.7%), n =11 of type 1a (11.8%), n = 6 of type 1b (31.5%), n = 6 of 2a (16.6%), n = 9 of type 2b (69%), n = 10 of type 3 (71%) and n = 2 of type 4 (13.4). No other characteristics specified.</p>	<p><b>Interventions</b></p> <p><u>Type 0</u> Stable FHR during entire second stage</p> <p><u>Type 1a</u> Mild variable decelerations</p> <p><u>Type 1b</u> Moderate to severe variable decelerations or late decelerations with each contraction, returning to baseline inbetween</p> <p><u>Type 2a</u> Baseline 90–120 bpm with decelerations</p> <p><u>Type 2b</u></p>	<p><b>Details</b></p> <p>n = 293 cases in which FHR monitoring was obtained during the last hour of the 1st stage and entire 2nd stage were evaluated. Arterial and venous umbilical blood was obtained in all cases. n = 103 cases were included in type 0 (absence of FHR abnormalities during the 2nd stage) were used as a control group. FHR tracing was recorded via a spiral electrode applied to the</p>	<p><b>Results</b></p> <p><b><u>Umbilical artery acid base pH (2nd stage CTG types)</u></b></p> <p><u>Type 0</u> 7.24 ± 0.06</p> <p><u>Type 1a</u> 7.15 ± 0.07 p = ns</p> <p><u>Type 1b</u> 7.19 ± 0.07 p = 0.0001</p> <p><u>Type 2a</u> 7.19 ± 0.06 p = 0.0001</p> <p><u>Type 2b</u> 7.06 ± 0.07 p = 0.0001</p> <p><u>Type 3</u> 7.09 ± 0.06 p = 0.0001</p> <p><u>Type 4</u> 7.19 ± 0.07 p = 0.01</p> <p><b><u>Umbilical vein acid base pH (2nd stage CTG types)</u></b></p> <p><u>Type 0</u> 7.30 ± 0.06</p> <p><u>Type 1a</u></p>	<p><b>Limitations</b></p> <p>Unusual scoring system.</p> <p>Analysis not based on specific FHR abnormalities.</p> <p>Small numbers in more severe categories (2b: n = 13, 3: n = 14).</p> <p><b>Other information</b></p> <p>Beginning of 2nd stage: Defined as the moment of the initiation of pushing effort and full cervical dilatation</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To examine the correlation between fetal heart rate (FHR) patterns during the 2nd stage of labour and umbilical blood acid based parameters</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p><b>Inclusion criteria</b></p> <p>Singleton pregnancy</p> <p>Term pregnancy (37-42 weeks gestation)</p> <p>No maternal and fetal pathology</p> <p>Vertex birth</p> <p>Spontaneous or instrumental vaginal birth</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>	<p>Basal FHR below 90 bpm, usually with reduced variability</p> <p><u>Type 3</u></p> <p>Basal FHR below 90 bpm, low variability, accelerations with contractions</p> <p><u>Type 4</u></p> <p>Basal FHR below 90bpm during final moments of 2nd stage only</p>	<p>fetal head and uterine contractions were measured by tocodynamometer. Paper speed of the monitor was 1cm/min.</p> <p><b>Analysis</b></p> <p>Analysis of the tracing was independently interpreted and classified by two investigators that were blinded to the information regarding umbilical cord pH and cases.</p> <p>Acidemia was diagnosed when pH levels were more than one standard deviation below the mean level obtained in the control group. The 2nd stage of labour never exceeded 45 min</p>	<p>7.29 ± 0.07 p = ns</p> <p><u>Type 1b</u></p> <p>7.22 ± 0.07 p = 0.0001</p> <p><u>Type 2a</u></p> <p>7.26 ± 0.06 p = 0.001</p> <p><u>Type 2b</u></p> <p>7.12 ± 0.07 p = 0.0001</p> <p><u>Type 3</u></p> <p>7.15 ± 0.06 p = 0.0001</p> <p><u>Type 4</u></p> <p>7.24 ± 0.06 p = 0.004</p> <p><b>Early neonatal morbidity was found in n = 3 neonates:</b></p> <p><u>Case 1</u></p> <p>CTG pattern 1b Arterial pH 7.07 Morbidity: resuscitation Days in NICU: 2</p> <p><u>Case 2</u></p> <p>CTG pattern 2b Arterial pH 7.00 Morbidity: grunting Days in NICU: 7</p> <p><u>Case 3</u></p> <p>CTG pattern 2b Arterial pH 7.09 Morbidity: resuscitation Days in NICU: 4</p> <p>Arterial and venous pH values significantly lower in types 1b and below compared with controls.</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				Mean pH only < 7.20 in types 2b and 3.	
<p><b>Full citation</b></p> <p>Dellinger, E.H., Boehm, F.H., Crane, M.M., Electronic fetal heart rate monitoring: early neonatal outcomes associated with normal rate, fetal stress, and fetal distress, American Journal of Obstetrics and Gynecology, 182, 214-220, 2000</p> <p><b>Ref Id</b></p> <p>170635</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p>	<p><b>Sample size</b></p> <p>n = 898</p> <p>Normal pattern n = 627</p> <p>Stress pattern n = 263</p> <p>Distress pattern n = 8</p> <p><b>Characteristics</b></p> <p>Comparative characteristics not reported</p> <p><b>Inclusion criteria</b></p> <p>Singleton pregnancy</p> <p>&gt; 32 weeks gestation</p> <p><b>Exclusion criteria</b></p> <p>Presence of anomalies or arrhythmias</p> <p>Multiple pregnancy</p> <p>Gestational age &lt; 32 weeks</p>	<p><b>Interventions</b></p> <p><u>Normal pattern</u> 110–160 bpm, minimal to moderate variability, with or without accelerations</p> <p><u>Stress pattern</u> &gt; 160 bpm for &gt; 5 minutes, minimal to moderate variability, moderate to severe variable decelerations, late decelerations or sinusoidal pattern</p> <p><u>Distress pattern</u> &lt; 110 bpm for &gt; 5 minutes, moderate to severe variable decelerations with absent variability, late</p>	<p><b>Details</b></p> <p>Fetal heart rate data from all labouring women monitored at 2 institutions were examined. Tracings in the final hour before delivery were defined as normal, fetal stress, or fetal distress. Based on the standard care of the hospital all labouring women received electronic fetal heart monitoring. All tracings were stored after birth and reviewed at the later date by an observer blinded to the birth outcomes. The FHR</p>	<p><b>Results</b></p> <p>Total normal n = 627 Total stress n = 236 Total distress n = 8</p> <p><u>Umbilical pH &lt; 7.00</u> Normal n = 0/627 Stress n = 2/263 (1.6%) Distress n = 2/8 (28.5%) p &gt; 0.001</p> <p><u>NICU admission</u> Normal n = 29 Distress/Stress n = 25</p> <p><u>LSCS rate</u> Normal n = 75 Distress/Stress n = 4</p> <p><u>Stress/distress vs. normal</u> Sensitivity 68% Specificity 71% PPV 5% NPV 99%. Umbilical cord pH &lt; 7.00</p> <p><u>Stress/distress vs. normal</u> Sensitivity 100% Specificity 66% PPV 3% NPV 100%</p> <p><u>Results also on distress vs. normal</u> NPV for all outcomes &gt; 98%</p>	<p><b>Limitations</b></p> <p>Underpowered cohort due to imbalance between groups.</p> <p>Analysis between distress and normal for pH and Apgar highly specific but interpret with caution in view of numbers in each group.</p> <p><b>Other information</b></p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>To examine the ability of well-defined classification system for electronic fetal heart rate (FHR) tracing to predict early neonatal outcome</p> <p><b>Study dates</b></p> <p>One hospital: July 1993 to February 1994 One hospital: February to June 1995</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>Caesarean section before onset of labour</p> <p>Inability to obtain an adequate FHR tracing</p> <p>Traces were excluded from the study if <math>\geq 15</math> min during the final hour went untraced</p>	<p>decelerations with absent variability, 110–160 bpm with absent variability and no accelerations</p>	<p>tracing was evaluated for the one hour period preceding the birth.</p>		
<p><b>Full citation</b></p> <p>Ellison,P.H., Foster,M., Sheridan-Pereira,M., MacDonald,D., Electronic fetal heart monitoring, auscultation, and neonatal outcome,</p>	<p><b>Sample size</b></p> <p>Original cohort from Dublin RCT. Two groups of FHR traces: electronic fetal monitoring (EFM) alone n = 2362 and EFM plus neurological examination n = 135</p>	<p><b>Interventions</b></p> <p>All FHR variables</p>	<p><b>Details</b></p> <p>Data in this study are from a randomised control trial conducted in Dublin (comparing the effectiveness of electronic fetal monitoring</p>	<p><b>Results</b></p> <p>Correlation of specific fetal heart patterns to neonatal convulsions (n = 135):</p> <p><u>1<sup>st</sup> stage of labour</u></p> <p>Late deceleration r = 0.38, p &lt; 0.001 Severe variable deceleration r = -0.04, p = ns Marked tachycardia r = -0.02 Moderate variable decelerations r = -0.02 Early decelerations r = 0.01 Normal baseline and variability r = -0.05</p>	<p><b>Limitations</b></p> <p>No specifics of scoring for neurological examination specified</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>American Journal of Obstetrics and Gynecology, 164, 1281-1289, 1991</p> <p><b>Ref Id</b> 164084</p> <p><b>Country/ies where the study was carried out</b> Ireland</p> <p><b>Study type</b> Retrospective cohort study</p> <p><b>Aim of the study</b> To examine the relationship between a number of maternal, labour and delivery variables (including fetal heart rate [FHR] patterns) to neonatal outcomes</p> <p><b>Study dates</b></p>	<p><b>Characteristics</b> Not specified</p> <p><b>Inclusion criteria</b> Not specified</p> <p><b>Exclusion criteria</b> Heavily stained meconium liquor Decreased amniotic fluid Abnormal heart rate on admission</p>		<p>and auscultation in improving the health of fetus during delivery and birth). For the purpose of this review only data on electronic fetal monitoring will be reported. Data for electronic fetal heart monitoring were available for both the 1st and 2nd stages of labour. The fetal heart rate monitoring was interpreted by an obstetrician who was blinded to the women's characteristics and neonatal birth outcomes. All newborns were examined physically and neurologically by a physician. FHR patterns were recorded separately.</p>	<p><u>2<sup>nd</sup> stage of labour</u> Late decelerations <math>r = 0.38</math>, <math>p &lt; 0.001</math> Early decelerations <math>r = 0.01</math></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>March 1981 to April 1983</p> <p><b>Source of funding</b></p> <p>Not specified</p>			<p><u>Analysis</u></p> <p>Frequencies were reviewed for all variables, as well as distributions and skews. Pearson correlation and biserial correlations for dichotomous variables were obtained and reviewed for each sample.</p>		
<p><b>Full citation</b></p> <p>Gaffney,G., Flavell,V., Johnson,A., Squier,M., Sellers,S., Cerebral palsy and neonatal encephalopathy, Archives of Disease in Childhood Fetal and Neonatal Edition, 70, F195-F200, 1994</p> <p><b>Ref Id</b></p> <p>196440</p>	<p><b>Sample size</b></p> <p>141 case of cerebral palsy; UK hospital</p> <p><b>Characteristics</b></p> <p>No significant differences observed between the two groups (with neonatal encephalopathy [NE] and without neonatal encephalopathy) marital status, maternal disease, recurrent abortion, poor obstetric history, previous preterm birth, maternal smoking habit, and</p>	<p><b>Interventions</b></p> <p>Ominous FHR pattern</p>	<p><b>Details</b></p> <p>Children with cerebral palsy born during the study period were identified from the Oxford health regional register of childhood impairment. The children with cerebral palsy were divided into those with signs of neonatal</p>	<p><b>Results</b></p> <p><b><u>Findings on cardiocotograph (CTG) in mothers of children with cerebral palsy with or without neonatal encephalopathy</u></b></p> <p><u>Ominous first stage CTG</u>                      Without NE: n = 4/48 (8%)                      With NE: n = 13/27 (48%)                      OR 10.2 (2.9 to 36.4)</p> <p><u>Ominous second stage CTG</u>                      Without NE: n = 19/45 (42%)                      With NE: n = 21/25 (84%)                      OR 7.2 (2.1 to 24.4)</p> <p><u>Median duration of first stage abnormality (min)</u>                      Without NE: 48.5 (38 to 287)                      With NE: 200.0 (15 to 480)</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>Neonatal encephalopathy defined as: Depression at birth, based on a one minute apgar score of less than or equal to 6. Followed by evidence of neonatal neurological abnormality such as lethargy, coma, impaired respiration, seizures, and/or tone change</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To test the hypothesis that children born at term with cerebral palsy with signs of neurological dysfunction preceded by depression at birth (termed neonatal encephalopathy) differ from those without such signs in the frequency of antenatal and perinatal factors, and in the severity and characteristics of their impairment and disability</p>	<p>maternal age. More women in the 'without NE' group were primigravida compared with the 'with NE' group. Half the mothers of infants with neonatal encephalopathy (51/100) and mothers of infants with neonatal encephalopathy (20/41), had one or more complicating factors (antenatal infection, premature rupture of membranes, pre-eclampsia, severe pre-eclampsia, antepartum haemorrhage, previous infertility, induced conception, raised maternal serum alpha fetoprotein, polyhydramnios, reduced fetal movement, or complicated antenatal course). More women in the neonatal encephalopathy group had post-date pregnancy (&gt; 41 weeks), induction of labour, 2nd stage of labour exceeding &gt; 2 hours, meconium stained liquor, caesarean section or instrumental birth.</p>		<p>encephalopathy (with NE) and those without (without NE). This was based on the information recorded in the neonatal case notes. The clinical characteristics of the children in the study were described in terms of distribution of tone changes, as walking and non walking, and with or without intellectual deficit, vision loss, seizures, involuntary movement, or bulbar signs such as difficulty in swallowing.</p>	<p><math>p = 0.3</math></p> <p><u>Median duration of second stage abnormality (min)</u></p> <p>Without NE: 38 (8 to 287)</p> <p>With NE: 100.0 (12 to 480)</p> <p><math>p = 0.003</math></p> <p>Follow-on data: significant association with major and minor impairment in encephalopathy group.</p> <p>Quadraplegia (OR 4.8; 95% CI 2.2 to 10.5)</p> <p>Hemiplegia (OR 0.3; 95% CI 0.1 to 0.8)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study dates</b></p> <p>1984 to 1987</p> <p><b>Source of funding</b></p> <p>Funded by Oxford Regional Health Authority</p>	<p>There was no significant difference in augmentation use between the two groups.</p> <p><b>Inclusion criteria</b></p> <p>Singleton pregnancy</p> <p>Term pregnancy</p> <p><b>Exclusion criteria</b></p> <p>Children with major congenital abnormality</p> <p>Children in whom there was a definite postnatal cause for cerebral palsy such as meningitis or trauma</p>				
<p><b>Full citation</b></p> <p>Giannubilo,S.R., Buscicchio,G., Gentilucci,L., Palla,G.P., Tranquilli,A.L., Deceleration area of fetal heart rate trace and fetal acidemia at delivery: A case-control study, Journal of</p>	<p><b>Sample size</b></p> <p>Total n = electronic fetal monitoring (EFM) traces of 236 pregnancy n = 56 pregnancies met the inclusion criteria (Acidemia n = 26, Control = 30)</p> <p><b>Characteristics</b></p>	<p><b>Interventions</b></p> <p>EFM traces</p>	<p><b>Details</b></p> <p>From n = 410 third trimester cardiotocograph (CTG) tracings performed at the department of obstetrics and gynaecology, Belcolle Hospital during</p>	<p><b>Results</b></p> <p><u>Number of decelerations (&gt; 15bpm/15s) during the second stage of labour</u></p> <p>Acidemia: 8.03 ± 3.77 Control: 4.64 ± 3.84)</p> <p><u>Total deceleration area/cm<sup>2</sup>/hour</u></p> <p>Acidemia: 35.56 ± 11.87 Control: 17.81 ± 9.38</p>	<p><b>Limitations</b></p> <p>Small study with a large drop out</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Maternal-Fetal and Neonatal Medicine, 20, 141-144, 2007</p> <p><b>Ref Id</b> 158821</p> <p><b>Country/ies where the study was carried out</b>  Italy</p> <p><b>Study type</b>  Retrospective cohort</p> <p><b>Aim of the study</b>  To assess the correlation between the total deceleration area of the fetal heart rate (FHR) pre-delivery trace and intrapartum fetal acid-base status in a low risk population.</p> <p><b>Study dates</b>  January to August 2004</p>	<p><u>Maternal</u> There were no significant differences observed between the two groups (normal and abnormal pH at birth) in maternal age, gestational age at delivery, primiparity, length of second stage of labour or operative delivery rate. The length of first stage of labour was statistically significantly longer in controls compared with academic group <math>p &lt; 0.001</math>.</p> <p><u>Neonatal</u> There were also no significant differences observed in birth weight, baby's sex, apgar score 1 min <math>&lt; 7</math> and apgar score 5 min <math>&lt; 7</math>, or cord arterial pH. Cord base deficit was significantly higher in the academic group compared with controls <math>p &lt; 0.001</math>.</p> <p><u>CTG parameter</u> (Academic <math>n = 26</math>, Control <math>n = 30</math>) <u>Baseline heart rate</u> Academic <math>131.25 \pm 9.19</math> Control <math>136.25 \pm 10.14</math></p>		<p>the study period, <math>n = 236</math> with performed cord gas analysis were selected for inclusion. <math>n = 56</math> pregnancies met the inclusion criteria (Acidemia <math>n = 26</math>, Control = 30). CTG was performed during second stage of labour at least one hour without interruption. Umbilical blood gas performed by collecting blood samples from cord artery and the <math>pH &lt; 7.2</math> was considered abnormal. A base deficit <math>\geq 12</math> mmol/l was considered the threshold of the fetal metabolic acidosis at delivery. Hospital records of each newborn were evaluated for</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p> <p>Not reported</p>	<p><u>Number of decelerations &gt; 15 bpm/15</u>                      Academic 8.03 ± 3.77                      Control 4.64 ± 3.84</p> <p><u>Fetal deceleration area cm<sup>2</sup>/h</u>                      Academic 17.81 ± 9.38                      Control 35.56 ± 11.87</p> <p><b>Inclusion criteria</b></p> <p>Normal FHR pattern (normal variability, presence of accelerations)</p> <p>Singleton pregnancy</p> <p>Caucasian race</p> <p>Vertex presentation</p> <p>Vaginal birth, no labour augmentation</p> <p>Term birth &gt; 37 wks</p> <p><b>Exclusion criteria</b></p> <p>Technically uninterpretable trace</p> <p>Required emergency caesarean section (CS) because of maternal or fetal conditions (such</p>		<p>Apgar, weight and neonatal complication.</p> <p><u>Analysis</u>                      The deceleration area was calculated, after digital analysis, with Autocad System 2004. Statistical analysis performed with SPSS version 0.8 statistical package. Chi-square or Fisher's exact tests were used for comparison of proportions. Student's t-test was applied for comparisons of means.</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>as sign of placental insufficiency, cephalo-pelvic distribution)</p> <p>Previous CS</p> <p>Pre-existing heart or lung disease</p> <p>Carrying a baby with growth restriction or malformation</p>				
<p><b>Full citation</b></p> <p>Gilstrap,L.C.,III, Hauth,J.C., Hankins,G.D., Beck,A.W., Second-stage fetal heart rate abnormalities and type of neonatal acidemia, Obstetrics and Gynecology, 70, 191-195, 1987</p> <p><b>Ref Id</b></p> <p>195342</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p>	<p><b>Sample size</b></p> <p>n = 277 cases with known arterial cord pH samples and satisfactory second stage traces</p> <p><b>Characteristics</b></p> <p>White race: 83%</p> <p>Maternal age 20-29 years old: 71%</p> <p>Primiparous: 51%</p> <p><b>Inclusion criteria</b></p> <p>Term birth</p>	<p><b>Interventions</b></p> <p>Uncomplicated bradycardia or tachycardia</p>	<p><b>Details</b></p> <p>Cord pH was determined within 5 minutes of birth and specimens were obtained from either the umbilical artery or vein. Acidosis defined as a arterial cord pH of less than 7.20. Fetal heart rate tracings were obtained during the second stage via a scalp electrode. The tracing during the 2<sup>nd</sup> stage</p>	<p><b>Results</b></p> <p><b><u>Correlation of normal and abnormal traces and cord pH (mean ± SD)</u></b></p> <p>Normal (n = 129) 7.29 ± 0.6                      Tachycardia (n = 32) 7.25 ± 0.5 p &lt; 0.05                      Mild bradycardia (n = 53) 7.23 ± 0.7 p &lt; 0.05                      Moderate or severe bradycardia (n = 63) 7.22 ± 0.7 p &lt; 0.05</p>	<p><b>Limitations</b></p> <p>Unclear for how long abnormalities were present for</p> <p>Not consecutive cases, hence subject to selection bias</p> <p><b>Other information</b></p> <p><u>Uncomplicated bradycardia:</u>                      Mild (90–119 bpm)</p> <p>Moderate (60–89 bpm)</p> <p>Severe (&lt; 60 bpm)</p> <p>Tachycardia (&gt; 160 bpm)</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study type</b></p> <p>Cohort study</p> <p><b>Aim of the study</b></p> <p>To examine the incidence and type of acidaemia, degree of buffer base deficit, and immediate neonatal outcome in relation to baseline second stage fetal heart rate (FHR) patterns before delivery</p> <p><b>Study dates</b></p> <p>June 1985 to April 1986</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>Vaginal birth</p> <p>Vertex presentation</p> <p><b>Exclusion criteria</b></p> <p><u>Women with complication such as:</u></p> <p>Diabetes</p> <p>Chronic hypertension</p> <p>Preeclampsia</p> <p>Acute chorioamnionitis</p> <p>Significant medical illness</p> <p>Women with abnormal FHR such as late decelerations, moderate or severe variable decelerations, bradycardia and tachycardia</p>		<p>(before expulsion of head) was evaluated for baseline FHR abnormality and variability. Only women with either a normal FHR pattern or obvious baseline changes, consisting of bradycardia or tachycardia, were included.</p> <p>Analysis</p> <p>The FHR trace was independently analysed by both authors without knowledge of blood gas results. Traces were only included if the interpretation was in agreement (there was disagreement in &lt; 2% of the traces)</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Gilstrap,L.C.,III, Hauth,J.C., Toussaint,S., Second stage fetal heart rate abnormalities and neonatal acidosis, Obstetrics and Gynecology, 63, 209-213, 1984</p> <p><b>Ref Id</b></p> <p>195341</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Prospective cohort study</p> <p><b>Aim of the study</b></p> <p>To examine the correlation of baseline fetal heart rate (FHR) abnormalities in the last 10</p>	<p><b>Sample size</b></p> <p>n = 833 cases with cord pH samples and interpretable traces in the last 10 minutes of labour</p> <p><b>Characteristics</b></p> <p><b>Demographic characteristics:</b></p> <p>White race: 75%</p> <p>Maternal age 20-29 years old: 65%</p> <p>Primiparous: 85%</p> <p>Term pregnancy: 98%</p> <p><b>Inclusion criteria</b></p> <p>If a cord pH was obtained</p> <p>If there was satisfactory fetal heart tracing during the last minutes of 2<sup>nd</sup> stage</p> <p><b>Exclusion criteria</b></p>	<p>Uncomplicated bradycardia</p> <p>Uncomplicated tachycardia</p>	<p><b>Details</b></p> <p>All infants during the study period, whose delivery was by forceps, were included in the study. Cord pH was determined within 5 minutes of birth and specimens were obtained from either the umbilical artery or vein. Acidosis was defined as a pH of less than 7.20. Fetal heart rate tracings were obtained during the second stage via a scalp electrode. The tracing during the last 10 mins of delivery (before expulsion of the head) was evaluated for</p>	<p><b>Results</b></p> <p><b>Correlation of n = 833 normal and abnormal traces and cord pH</b></p> <p>Acidosis: Normal n = 19/430 (4%) Abnormal n = 80/403 (20%) p &lt; 0.001</p> <p><b>Association of mild bradycardia and umbilical cord pH</b></p> <p>Acidosis: Normal n = 19/430 (4%) Abnormal (with mild bradycardia [present 1-3 min in 17% and &gt; 3 in 20%]) n = 30/165 (18%) p &lt; 0.001</p> <p>Association of moderate bradycardia and umbilical cord pH</p> <p>Acidosis: Normal n = 19/430 (4%) Abnormal (with mild bradycardia [present 1-3 min in 25% and &gt; 3 in 29%]) n = 33/121 (27%) p &lt; 0.001</p> <p>Association of tachycardia (mild and marked) and umbilical cord pH</p> <p>Acidosis: Normal n = 19/430 (4%) Abnormal (with mild or marked tachycardia) n = 17/117 (18%) p &lt; 0.001</p> <p><b>Umbilical artery pH &lt; 7.20</b></p> <p>Mild tachycardia:</p>	<p><b>Limitations</b></p> <p>Not consecutive cases, high risk of selection bias</p> <p>Unclear how and by whom data were analysed</p> <p>Blood for cord pH was taken from umbilical artery or vein.</p> <p><b>Other information</b></p> <p><b>Uncomplicated bradycardia:</b></p> <p>Mild (90–119 bpm) Moderate (60–89 bpm) Severe (&lt; 60 bpm)</p> <p><b>Uncomplicated tachycardia</b></p> <p>Mild (160–179 bpm) Marked (&gt; 180 bpm)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>minutes of the second stage of labour with neonatal acid-base status</p> <p><b>Study dates</b></p> <p>August 1979 to January 1983</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>Women with significant FHR abnormalities during the 1<sup>st</sup> stage of labour such as: Decelerations Persistent pattern of bradycardia Tachycardia</p> <p>Women with significant FHR abnormalities, such as late or moderate or severe variable decelerations were excluded from the analysis</p>		<p>FHR abnormalities. Only women with either a normal FHR pattern or obvious baseline changes, consisting of bradycardia or tachycardia, were included.</p>	<p>&lt; 3 minutes: 4/42 (10%) &gt; 3 minutes: 9/54 (17%)</p> <p>Marked tachycardia: &lt; 3 minutes: 2/5 (40%) &gt; 3 minutes: 2/16 (13%)</p> <p>Mild bradycardia: &lt; 3 minutes: 19/110 (17%) &gt; 3 minutes: 11/55 (20%)</p> <p>Moderate to severe bradycardia: &lt; 3 minutes: 19/72 (26%) &gt; 3 minutes: 14/49 (29%)</p>	
<p><b>Full citation</b></p> <p>Hadar,A., Sheiner,E., Hallak,M., Katz,M., Mazor,M., Shoham-Vardi,I., Abnormal fetal heart rate tracing patterns during the first stage of labor: Effect on perinatal outcome, American Journal of Obstetrics and</p>	<p><b>Sample size</b></p> <p>n = 601 FHR tracing (pregnancies); n = 301 abnormal pattern, n = 300 normal pattern</p> <p><b>Characteristics</b></p> <p>Women with abnormal tracing were more often nulliparous and delivered infants with significantly lower birth weight, compared with women with normal tracing. There were no</p>	<p><b>Interventions</b></p> <p>Fetal heart rate tracing (normal vs. abnormal)</p>	<p><b>Details</b></p> <p>The perinatal outcomes of 301 infants born at 37 to 42 weeks of gestation with pathologic fetal heart rate patterns during the first stage of labour were compared with 300 infants with normal fetal heart rate tracing</p>	<p><b>Results</b></p> <p><u>Arterial pH 7.2</u> Abnormal FHR n = 48/301 (16%) Normal FHR n = 14/300 (4.7%) p &lt; 0.001</p> <p><u>Arterial pH 7.1</u> Abnormal FHR n = 10/301 (3.3%) Normal FHR n = 2/300 (0.7%) p &lt; 0.02</p> <p><u>Base deficit ≥ 12</u> Abnormal FHR n = 25/301 (8.3%) Normal FHR n = 7/300 (2.3%) p = 0.001</p> <p><u>Admission to NICU</u></p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>Tracings were interpreted with the use of National Institute of Child Health Development Research Planing Workshop Guideline (NICHD)</p> <p>Abnormal pH was defined as: pH 7.2 in 2 separate analyses</p> <p>Base deficit of ≥ 12 mmol/l was considered to be diagnostic of fetal metabolic acidosis at birth</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Gynecology, 185, 863-868, 2001</p> <p><b>Ref Id</b> 169256</p> <p><b>Country/ies where the study was carried out</b> Israel</p> <p><b>Study type</b> Cohort</p> <p><b>Aim of the study</b> To evaluate perinatal outcomes of infants who had pathologic fetal heart rate (FHR) tracings during the first stage of labour, in comparison with pregnancies with normal tracings.</p> <p><b>Study dates</b> January to June 2000</p>	<p>significant differences observed in FHR patterns in maternal age, ethnic origin, gravidity, gestational age and sex of the baby. Women with abnormal tracing had a significantly higher rate of oligohydramnios and oxytocin augmentation in labour. Women with abnormal FHR patterns had a significantly longer duration of 1st stage labour, and a higher incidence of thick meconium stained amniotic fluid.</p> <p><b>Inclusion criteria</b></p> <p>Low risk women</p> <p>Fetus at vertex presentation</p> <p>Normal FHR pattern</p> <p>Women with interpretable external fetal monitoring tracing during the labour and birth</p> <p>Cases with values taken immediately after birth</p>		<p>patterns. Data were collected prospectively and demographic information was obtained on each woman's admission to the hospital. The labour room team evaluated each woman's FHR tracing hourly and documented the results. The same obstetrician collected the data after assessing the FHR tracing and the delivery chart. The data were collected prospectively. Tracings were interpreted with the use of the National Institute of Child Health and Human Development fetal heart rate monitor</p>	<p>Abnormal FHR n = 4/301 (1.3%) Normal FHR n = 4/300 (1.3%) p &lt; 0.343</p> <p><u>Vacuum birth</u> Abnormal FHR n = 33/301 (11.0%) Normal FHR n = 12/300 (4%)</p> <p><u>Caesarean birth</u> Abnormal FHR n = 46/301 (15%) Normal FHR n = 20/300 (6.3%)</p> <p><u>Spontaneous vaginal birth</u> Abnormal FHR n = 222/301 (73.8%) Normal FHR n = 268/300 (89.3%)</p> <p><u>Factors associated with pathologic fetal heart rate monitoring during the first stage of labour in a multivariable analysis</u> Hydramnios: odds ratio 7.68 (95% CI, 1.75% to 33.63%), Oligohydramnios: odds ratio 2.74 (95% CI, 1.01% to 7.39%), Presence of meconium-stained amniotic fluid: odds ratio 1.91 (95% CI, 1.03% to 3.3%)</p> <p><u>Pathological fetal heart patterns during the 1st stage of labour (compared with normal tracing n = 300 associated with fetal acidosis (pH &lt; 7.2 and base deficit ≥ 12)</u> Late deceleration (yes/no): odds ratio 17.5 (95% CI, 1.6 to 185.7) p = 0.01 Variable deceleration &lt; 70 bpm (yes/no): odds ratio 3.9 (95% CI, 1.3 to 11.7) p = 0.01 Pathologic FHR during the 1st stage of labour (yes/no): odds ratio 2.86 (95% CI, 0.3 to 24.4) p = 0.336</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p>	<p><b>Exclusion criteria</b></p> <p>Congenital abnormalities</p> <p>Preexisting maternal heart or lung disease</p> <p>Fetuses with intrauterine growth retardation</p> <p>Women in need of emergency caesarean section</p> <p>Previous Caesarean section</p>		<p>guidelines. Umbilical cord blood was collected immediately after birth and all blood gas analysis performed within 10 minutes of birth.</p> <p><u>Analysis</u> SPSS version 8.0 package was used for the analysis. Chi square test used for comparison between the two groups for the categorical variable and Student's t-test was used for continuous variables with normal distribution. Multiple logistic regression was used to investigate the independent contribution of obstetric factors to abnormal fetal</p>		
<p>Not specified</p>					

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			heart patterns and to investigate the contribution of those factors to the occurrence of fetal acidosis (pH 7.2 and base deficit $\geq$ 12)		
<p><b>Full citation</b></p> <p>Heinrich,J., Elective fetal monitoring and obstetrical operative frequency, European Journal of Obstetrics, Gynecology, and Reproductive Biology, 14, 143-152, 1982</p> <p><b>Ref Id</b></p> <p>196602</p> <p><b>Country/ies where the study was carried out</b></p> <p>Germany</p> <p><b>Study type</b></p> <p>Cohort</p>	<p><b>Sample size</b></p> <p>n = 2694 unselected deliveries</p> <p>n = 5000 elective monitored women (additional group)</p> <p><b>Characteristics</b></p> <p>Unclear gestation range/risk range</p> <p><b>Inclusion criteria</b></p> <p>Not specified</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>	<p><b>Interventions</b></p> <p>All FHR variables. Grouped into scoring system</p> <p><u>Normal</u> Baseline 120–160 bpm; constant mild bradycardia; variability 10–25 bpm; sporadic variable declarations; accelerations; mild variable deceleration</p> <p><u>Warning</u> Tachycardia; variability &lt; 10 bpm or &gt; 25 bpm; periodic accelerations;</p>	<p><b>Details</b></p> <p>Digital display fetal monitors were used recording several tocometric parameters such as amplitude, frequency, base tonus and Montevideo units of labor. If the measured values exceeded an upper limit, an automatic alarm signal was activated. Arterial umbilical pH was carried out for all liveborns. The</p>	<p><b>Results</b></p> <p><u>Umbilical artery pH</u> Significant difference at pH &lt; 7.20 between severe and hypoxic categories compared to warning and normal categories.</p> <p><b><u>FHF parameter in the 2nd stage of labour (30 min antepartum) and pH of umbilical arteria</u></b></p> <p><u>Normal classification (n = 1080)</u> Normal pH (pH &gt; 7.20): 1043/1080 (96.6%) Preacidosis (pH 7.25 - 7.20): 27/1080 (2.5%) Acidosis (pH &lt; 7.20): 10/1080 (0.9)</p> <p><u>Warning symptoms (n = 1133)</u> Normal pH (pH &gt; 7.20): 1095/1133 (96.7%) Preacidosis (pH 7.25 - 7.20): 27/1133 (2.4%) Acidosis (pH &lt; 7.20): 11/1133 (0.9)</p> <p><u>Severe functional hemodynamic (n = 431)</u> Normal pH (pH &gt; 7.20): 357/431 (93.0%) Preacidosis (pH 7.25 - 7.20): 48/431 (11%) Acidosis (pH &lt; 7.20): 26/451 (6.0%)</p>	<p><b>Limitations</b></p> <p>Small numbers in hypoxic category</p> <p>Not possible to determine gestation or risk categories</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To evaluate the influence of fetal monitoring on obstetric operation rates with emphasis on fetal heart frequency (FHF).</p> <p><b>Study dates</b></p> <p>1977 to 1978 and 1979 to 1981 (additional group)</p> <p><b>Source of funding</b></p> <p>Not specified</p>		<p>moderate variable decelerations; early decelerations</p> <p><u>Severe</u> Transient bradycardia; severe variable decelerations; prolonged decelerations</p> <p><u>Hypoxia</u> Final bradycardia; variability 0–5 bpm; typical late decelerations</p>	<p>collected data included identification of the patient, results of medical history as well as of clinical and laboratory examinations and a final review of the course of pregnancy, delivery and post-partum period. The validity of the FHF-classification was demonstrated in 2694 unselected deliveries (June 1977/1978) by comparison with postnatal measurement of acid-base balance and Apgar scoring. The relation of obstetric operation rate, values of acid-base balance in umbilical arteria and</p>	<p><u>Hypoxia (n = 50)</u> Normal pH (pH &gt; 7.20): 30/50 (60.0%) Preacidosis (pH 7.25 - 7.20): 11/50 (22%) Acidosis (pH &lt; 7.20): 9/50 (18%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>FHF-parameters were also studied in an additional group of 5000 elective monitored patients (November 1979-1981).</p> <p>Data analysis The automated data analysis was made by means of a digital computer system (ES 1040).</p>		
<p><b>Full citation</b></p> <p>Honjo,S., Yamaguchi,M., Umbilical artery blood acid-base analysis and fetal heart rate baseline in the second stage of labor, Journal of Obstetrics and Gynaecology Research, 27, 249-254, 2001</p>	<p><b>Sample size</b></p> <p>n = 365</p> <p><b>Characteristics</b></p> <p>All subjects in the study were Japanese, no further characteristics were specified</p> <p><b>Inclusion criteria</b></p> <p>Term pregnancy (37 - 42 weeks)</p>	<p><b>Interventions</b></p> <p>FHR tracing with either normal or baseline abnormality consisting of bradycardia or tachycardia during the 2nd stage of labour</p>	<p><b>Details</b></p> <p>Data were collected from n = 365 newborns, born during the study period in maternity ward of a hospital in Takasaki city. Based on the hospital policy, umbilical cord artery blood was taken from</p>	<p><b>Results</b></p> <p>Umbilical arterial acidemia occurred in 54.1% of the newborns with moderate to severe bradycardia, in 27.3% with mild bradycardia, and in 19.3% with tachycardia, compared with only 1.3% of those with a normal FHR (p &lt; 0.001).</p> <p><b><u>Umbilical cord pH and blood gas analysis in newborn with normal and abnormal FHR tracing</u></b></p> <p><u>pH</u> Normal (n = 236) 7.31 ± 0.05 Tachycardia (n = 57) 7.22 ± 0.11 (p &lt; 0.001 as compared with normal)</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>The FHR definition proposed by the National Institute of Child Health and Human Development Research Planning Workshop was used: Abnormal tracing: - Baseline 110 - 160 bpm - Variability &lt; 5 bpm - No periodic deceleration - The baseline FHR was taken as approx. mean FHR rounded to increments of 5 bpm during a 10 minute segment</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<b>Ref Id</b> 195455  <b>Country/ies where the study was carried out</b> Japan  <b>Study type</b> Cohort  <b>Aim of the study</b> To evaluate the correlation between umbilical arterial acidemia and second-stage baseline fetal heart rate (FHR) abnormalities in Japanese newborn infants.  <b>Study dates</b> 1998 to 1999  <b>Source of funding</b> Not specified	Vertex presentation  Vaginal birth  <b>Exclusion criteria</b>  Women with complication such as: Diabetes  Pre-eclampsia  Multiple gestation  Chronic hypertension  Chorioamnionitis  Significant medical illness  Other pregnancy complications  Newborns with fetal heart rate abnormality during the 1st stage of labour including:  Late deceleration  Moderate or severe variable deceleration  Any persistent nonperiodic patterns of bradycardia, or tachycardia		all newborns for blood gas determinations within 5 minutes of birth. FHR monitoring was performed in the second stage. Fetal heart rate tracings were obtained for as long as possible during the second stage of labour. Babies with marked periodic FHR abnormalities were excluded from the analysis. Therefore, in this study FHR tracings with either normal or baseline abnormality consisting of bradycardia or tachycardia were evaluated.  The cord was clamped immediately after birth, and	Mild bradycardia (n = 11) $7.25 \pm 0.06$ (p < 0.01 as compared with normal) Moderate to severe bradycardia (n = 61) $7.18 \pm 0.06$ (p < 0.001 as compared with normal)  <u>Base excess</u> Normal (n = 236) - $5.2 \pm 2.8$ Tachycardia (n = 57) - $9.5 \pm 4.5$ (p < 0.001 as compared with normal) Mild bradycardia (n = 11) - $8.7 \pm 4.4$ (p < 0.05 as compared with normal) Moderate to severe bradycardia (n = 61) - $10.2 \pm 3.5$ (p < 0.001 as compared with normal)  <u>Number of newborns with an umbilical arterial pH &lt; 7.2 in different FHR patterns</u> Normal FHR pattern n = 3/236 (1.3%) Tachycardia n = 11/57 (19.3%) Mild bradycardia n = 3/11 (27.3%) Moderate to severe bradycardia n = 33/61 (54.1%) p < 0.001 (all 3 groups compared with normal group)	The baseline tachycardia and bradycardia was defined as: - Mild bradycardia: baseline FHR between 90 - 109 bpm for $\geq 10$ minutes - Moderate to severe bradycardia: baseline FHR < 90 bpm for $\geq 10$ minutes - Tachycardia: baseline FHR of 160 bpm for $\geq 10$ minutes  The decrease from the baseline was taken as $\geq 15$ bpm, lasting $\geq 2$ minutes, but < 10 minutes.  Newborn acidemia was defined as umbilical cord pH < 7.2, a pCO <sub>2</sub> 65 mmHg or lower, and bicarbonate 17.3 mmol/l or lower Metabolic acidemia was defined as an umbilical pH < 7.2, a pCO <sub>2</sub> 49.2 mmHg or lower, and bicarbonate 17.3 mmol/l, or lower

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			the blood samples were taken as soon afterwards as possible.		
<p><b>Full citation</b></p> <p>Krebs,H.B., Petres,R.E., Dunn,L.J., Smith,P.J., Intrapartum fetal heart rate monitoring. VI. Prognostic significance of accelerations, American Journal of Obstetrics and Gynecology, 142, 297-305, 1982</p> <p><b>Ref Id</b></p> <p>159500</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort study</p>	<p><b>Sample size</b></p> <p>n = 1996 fetal heart rate (FHR) traces</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <p>Term, singleton pregnancies</p> <p>&gt; 34 weeks gestation</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>	<p><b>Interventions</b></p> <p>Periodic variable and uniform accelerations</p>	<p><b>Details</b></p> <p>Fetal tracings were obtained from women in labour during the study period. The time of monitoring exceeded 2 hours and included at least 30 minutes of the first stage of labour. The FHR tracings were reviewed by the senior author. The average monitoring time was 6.2 hours. Indications for monitoring were preeclampsia and eclampsia (10.2%), meconium stained liquor</p>	<p><b>Results</b></p> <p>Mode of birth: Caesarean section: 16.2% (n = 241 in the 1st stage of labour, n = 83 in the second stage of labour)</p> <p><b><u>Prognostic significance of sporadic accelerations in the first 30 minutes of monitored labour: ≥ 3 accelerations per 30 minutes</u></b> <u>Perinatal mortality</u></p> <p>Elective n = 2 (0.2%) Non elective (with high risk factors) n = 4 (0.4%)</p> <p>P &gt; 0.5</p> <p><b><u>Prognostic significance of sporadic accelerations in the first 30 minutes of monitored labour: &lt; 3 accelerations per 30 minutes</u></b> <u>Perinatal mortality</u></p> <p>Elective n = 3 (2.8%) Non elective (with high risk factors) n = 12 (9.8%)</p>	<p><b>Limitations</b></p> <p>Unbalanced cohort with only 86 (4%) adverse outcomes.</p> <p>Not clear if the outcome assessors were blinded to outcomes.</p> <p>Unclear data analysis.</p> <p><b>Other information</b></p> <p>FHR scoring for internal FHR monitoring; for each of the criteria 0 to 2 points may be given so that a score of 0 to 10 may be obtained</p> <p><b><u>Baseline FHR</u></b> &lt; 100, &gt; 180 = 0 score 100 - 119, 161 - 180 = 1 score 120 - 160 = 2 score</p> <p><b><u>Variability (oscillatory amplitude [bpm])</u></b> &lt; 3 = 0 score 3 - 5 &gt; 25 = 1 score 6 - 25 = 2 score</p> <p><b><u>Variability (frequency [bpm])</u></b> &lt; 3 = 0 score</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To assess the prognostic value of accelerations in early labour and just prior to delivery</p> <p><b>Study dates</b></p> <p>January 1975 to June 1977</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>(14.2%), premature rupture of membranes (16.8%), and other high risk factors such as post-datism, intrauterine growth retardation, diabetes (7.1%), and oxytocin for indicated induction or augmentation (23%).</p> <p>Monitoring was elective in 46% of the women. The first and last 30 minutes of FHR tracing obtained from women in labour were evaluated.</p>	<p>P &lt; 0.05</p>	<p>3 - 6 = 1 score &gt; 6 = 2 score</p> <p><u>Acceleration/30 min</u> 0 = 0 score period, 1 - 4 sporadic = 1 score ≥ 5 sporadic = 2 score</p> <p><u>Deceleration/30 min</u> Late, severe variable, atypical variable = 0 score Mild variable, moderate variable = 1 score None, early deceleration, dip 0 = 2 score</p> <p>Acceleration defined: Transient increase in the FHR bpm above the baseline FHR. Sporadic accelerations occur independently from uterine contractions. Uniform sporadic accelerations have a rounded configuration, whereas variable sporadic accelerations differ from one another and abruptly leave and return to the baseline FHR. Periodic accelerations occur during the uterine contractions and are called uniform periodic accelerations. Variable accelerations are varied in shape and often develop notching, which widen, deepen, and progress into variable decelerations.</p>
<p><b>Full citation</b></p> <p>Larma,J.D., Silva,A.M., Holcroft,C.J., Thompson,R.E., Donohue,P.K., Graham,E.M., Intrapartum</p>	<p><b>Sample size</b></p> <p>Cases n = 107 Control n = 107</p> <p><b>Characteristics</b></p>	<p><b>Interventions</b></p> <p>Electronic fetal monitoring</p>	<p><b>Details</b></p> <p>Infants who were born with metabolic acidosis born in a single university were identified. The</p>	<p><b>Results</b></p> <p>Cases had a significant increase in late and prolonged decelerations/hour and late decelerations/contractions. Those fetuses with HIE had significant increases in bradycardia, decreased variability, and non reactivity but no difference in late or variable decelerations/hour.</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>Fetal metabolic acidosis and HIE are associated with significant increases in electronic fetal monitoring abnormalities, but their predictive ability to identify these conditions is low.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>electronic fetal heart rate monitoring and the identification of metabolic acidosis and hypoxic-ischemic encephalopathy, American Journal of Obstetrics and Gynecology, 197, 301-308, 2007</p> <p><b>Ref Id</b> 121224</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Study type</b> Case controlled study</p> <p><b>Aim of the study</b> To determine whether electronic fetal monitoring (EFM) can identify fetuses with metabolic acidosis and</p>	<p>The gestational age distribution: Born ≥ 37 weeks: 64% Born 29 - 36 weeks: 30% Born 24 - 28 weeks: 6% Born by caesarean section: 71%</p> <p><b>Inclusion criteria</b> All infants born with metabolic acidosis</p> <p><b>Exclusion criteria</b> Not specified</p>		<p>cases were 107 non anomalous chromosomally normal fetuses with an umbilical arterial pH &lt; 7.0 and base excess &lt; or = 12 mmol/l. Controls were the subsequent delivery that was matched by gestational age and mode of delivery. The last hour of the electronic fetal monitoring before delivery was evaluated by 3 obstetricians who were blinded to the outcome using a guideline developed by National Institute of Child Health and Human Development (NICHD) research planning workshop. Within the case group, n = 13</p>	<p><b>Identification of HIE (FHR parameters during the last hour before delivery)</b> <u>Time baselines &lt; 110 beats/min</u> Area under receiver operating characteristic curve: 0.56 Sensitivity: 15.4% Specificity: 98.9% Positive predictive values (PPV): 66.7%, Negative predictive values (NPV): 89.4%</p> <p><u>Baseline variability &lt; 5 beats/min</u> Area under receiver operating characteristic curve: 0.69 Sensitivity: 53.8% Specificity: 79.8% PPV: 26.9% NPV: 92.6%</p> <p><u>Non-reactive</u> Area under receiver operating characteristic curve: 0.65 Sensitivity: 92.3% Specificity: 61.7% PPV: 2.7% NPV: 82.9%</p> <p><u>all 3 abnormalities</u> Area under receiver operating characteristic curve: 0.82 Sensitivity: 7.7% Specificity: 98.9% Positive predictive values: 50.0% Negative predictive values: 88.6%</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>hypoxic-ischemic encephalopathy</p> <p><b>Study dates</b></p> <p>April 1991 to February 2006</p> <p><b>Source of funding</b></p> <p>Not specified</p>			<p>neonates had neurological complications (including 8 with seizures, n = 1 with grade 3 intra ventricular haemorrhage, n= 4 died). All 13 infants had clinical features that were consistent with at least Sarnat stage 2 (moderate hypoxic ischemic encephalopathy [HIE]). The EFM tracings of these 13 infants were compared with those of the other 94 infants with metabolic acidosis who had no neurologic injury.</p>		
<p><b>Full citation</b></p> <p>Low,J.A., Pickersgill,H., Killen,H.,</p>	<p><b>Sample size</b></p> <p>n = 166 term pregnancies with</p>	<p><b>Interventions</b></p> <p>Fetal heart rate patterns</p>	<p><b>Details</b></p> <p>The outcomes of n = 166 term pregnancies</p>	<p><b>Results</b></p> <p>Fetal asphyxial exposures were as follows: mild, n = 140; moderate, n = 22; and severe, n = 4.</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Derrick,E.J., The prediction and prevention of intrapartum fetal asphyxia in term pregnancies, American Journal of Obstetrics and Gynecology, 184, 724-730, 2001</p> <p><b>Ref Id</b></p> <p>197178</p> <p><b>Country/ies where the study was carried out</b></p> <p>Canada</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To examine the roles of clinical risk scoring, electronic fetal heart rate monitoring, and fetal blood gas and acid-base assessment in the prediction and prevention of intrapartum fetal</p>	<p>confirmed fetal asphyxia</p> <p><b>Characteristics</b></p> <p><b>Inclusion criteria</b></p> <p>Term pregnancies</p> <p>base deficit &gt; 12mmol/l</p> <p><b>Exclusion criteria</b></p>		<p>with biochemically confirmed fetal asphyxia (umbilical artery base deficit at delivery, &gt; 12 mmol/l) were examined. The population included n = 83 women who delivered by caesarean section matched with 83 women delivered vaginally. Antepartum and intrapartum clinical risk factors and neonatal complications were documented. Fetal assessments included fetal heart rate patterns in the fetal heart rate record and fetal capillary blood gas and acid-base assessments.</p>	<p><u>Mode of birth in mild fetal asphyxia</u> Caesarean section n = 67 (n 24/67 had meconium stained amniotic fluid) vaginal birth n = 73 (n = 32/67 had meconium stained amniotic fluid)</p> <p><u>Mode of birth in moderate or severe fetal asphyxia</u> Caesarean section n = 16 (n = 4/16 had meconium stained amniotic fluid) vaginal birth n = 10 (n = 4/10 had meconium stained amniotic fluid)</p> <p><b><u>Predictive and non-predictive FHR patterns according to mild fetal asphyxia vs moderate or severe fetal asphyxia</u></b></p> <p><u>Mild asphyxia</u> predictive pattern n = 89 Nonpredictive FHR pattern n = 25 No record n = 26</p> <p><u>Moderate or severe asphyxia</u> predictive pattern n = 20 Nonpredictive FHR pattern n = 4 No record n = 2</p> <p><u>Classification of FHR patterns in 26 pregnancies with moderate or severe asphyxia</u> Predictive n = 13 Suspect n = 7 Nonpredictive n = 3 No FHR monitoring record n = 3</p>	<p>Fetal asphyxia was classified as mild, moderate, or severe on the basis of umbilical artery base deficit (cutoff &gt; 12 mmol/l) and neonatal encephalopathy and other organ system complications</p> <p><b><u>FHR criteria predictive of fetal asphyxia:</u></b> Absent or minimal baseline variability and late or prolonged decelerations</p> <p>The FHR patterns are based on the findings in six 10 minute cycle of FHR recording: - Absent baseline variability, usually with repetitive cycles (<math>\geq 2</math>) of the late or prolonged deceleration - Repetitive cycles (<math>\geq 2</math>) of both minimal baseline variability and late or prolonged decelerations - Repetitive cycles (<math>\geq 2</math>) of either minimal baseline variability or late or prolonged deceleration - One cycle of either minimal baseline variability or late or prolonged decelerations - no cycle of either minimal baseline variability or late or prolonged decelerations</p> <p><b><u>Criteria for classification of FHR as predictive, suspect, and nonpredictive of fetal asphyxia on the basis of a 10 minute cycle of FHR recordings</u></b></p> <p><u>Predictive</u> Absent (cycle) <math>\geq 1</math> and late or prolonged decelerations <math>\geq 2</math> or Minimal (cycle) <math>\geq 2</math> and late or prolonged decelerations <math>\geq 2</math></p> <p><u>Suspect</u> Minimal (cycle) <math>\geq 2</math> and late or prolonged decelerations <math>\geq 0/1</math> or</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>asphyxia in term pregnancies.</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not specified</p>			<p>Each caesarean birth was matched with a vaginal birth on the basis of gestational age (<math>\pm</math> 1 week), birth weight (<math>\pm</math> 100g) and umbilical artery acid base deficit <math>&gt;</math> 12 mmol/l in the same year. The assessment of electronic FHR record was the interpretation of clinician in charge (outlined by medical record).</p> <p><u>Analysis</u> Statistical analysis included Student's t test. No further details provided</p>		<p>Minimal (cycle) <math>\geq</math> 0/1 and late or prolong decelerations <math>\geq</math> 2</p> <p><u>Nonpredictive</u> Minimal (cycle) 1 and late or prolong decelerations 0 or Minimal (cycle) 0 and late or prolong decelerations 1 or Minimal (cycle) 0 and late or prolong decelerations 0</p> <p><b><u>Classification of intrapartum fetal asphyxia</u></b></p> <p><u>Mild asphyxia</u> Metabolic acidosis (base deficit <math>\geq</math> 12 ): present Encephalopathy: minor* present or not present Cardiovascular, respiratory and renal complications: minor† present or not present</p> <p><u>Moderate asphyxia</u> Metabolic acidosis (Base deficit <math>\geq</math> 12 ): present Encephalopathy: moderate** present Cardiovascular, respiratory and renal complications: moderate †† or severe††† present or not present</p> <p><u>Severe asphyxia</u> Metabolic acidosis (Base deficit <math>\geq</math> 12 ): present Encephalopathy: severe* present** Cardiovascular, respiratory and renal complications: moderate †† or severe†† present</p> <p>* Irritability or jitteriness ** Profound lethargy or abnormal tone *** Coma or abnormal tone with seizure † Cardiovascular: with bradycardia (<math>\leq</math> 100 beats/min) or tachycardia (<math>\geq</math> 100 beats/min), respiratory: supplementary oxygen was required, †† Cardiovascular: with hypertention or hypotention, respiratory: if positive pressure or</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					ventilation > 24 hours were required, renal: elevation of serum creatinine level (> 100 mmol/l) ††† With abnormal electrocardiographic or echocardiographic findings, respiratory: if mechanical ventilation >24 hours were required, renal: anuria or oliguria (< 1 ml/kg per hour)
<p><b>Full citation</b></p> <p>Low,J.A., Victory,R., Derrick,E.J., Predictive value of electronic fetal monitoring for intrapartum fetal asphyxia with metabolic acidosis, Obstetrics and Gynecology, 93, 285-291, 1999</p> <p><b>Ref Id</b></p> <p>196968</p> <p><b>Country/ies where the study was carried out</b></p> <p><b>Study type</b></p> <p>Case control study</p> <p><b>Aim of the study</b></p> <p><u>Mean birth weight</u> Asphyxia group 3,412 ±</p>	<p><b>Sample size</b></p> <p>n = 71 term infants with base deficits &gt; 16 mmol/l n = 71 term infants with base deficits &lt; 8 mmol/l Studied over 4 hours prior to delivery (divided into 10-minute cycles)</p> <p><b>Characteristics</b></p> <p>No significant differences between the asphyxia and control group observed in maternal age, parity, medical and obstetric history or birth characteristics. Higher rate of meconium stained liquor in the asphyxia group compared with the control group (23/71 vs. 12/71 p = 0.05).</p> <p><u>Mean birth weight</u> Asphyxia group 3,412 ±</p>	<p><b>Interventions</b></p> <p>All FHR variables</p>	<p><b>Details</b></p> <p>A matched case control study conducted during the study period. n = 142 term infants who had the blood gas and acid base assessment at delivery were selected. Each case in the asphyxia group (infants with umbilical artery &gt; 16 mmol/l) was matched with a control infant whose umbilical artery base deficit was &lt; 8 mmol/l. Matching was performed based on the</p>	<p><b>Results</b></p> <p>Predictive value of abnormal FHR variables for acidosis <u>Absent baseline variability (&gt; 10 minutes) with late and/or prolonged decelerations:</u> sensitivity - 17% specificity - 98% positive predictive value (PPV) - 18 negative predictive value (NPV) - 98.3</p> <p><u>Minimal baseline variability (&gt; 20 minutes) and late and/or prolonged decelerations (&gt; 20 minutes):</u> sensitivity - 46% specificity - 89% PPV - 8 NPV - 98.7</p> <p><u>Minimal baseline variability (&gt; 20 minutes) or late decelerations and/or prolonged decelerations (&gt; 20 minutes):</u> sensitivity - 75% specificity - 57% positive predictive value - 3.5 negative predictive value - 99.1</p> <p><u>Minimal baseline variability (10 minutes) and/or late and/or prolonged decelerations (10 minutes):</u> sensitivity - 93%</p>	<p><b>Limitations</b></p> <p>Good NPV for all features individually. Poor specificity in combination. Baseline tachycardia, variable and early decelerations not discriminative features</p> <p><b>Other information</b></p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>To examine the predictive value of each fetal heart rate (FHR) variable and of patterns of FHR variables for fetal asphyxia during labour</p> <p><b>Study dates</b> May 1984 to May 1996</p> <p><b>Source of funding</b> Not specified</p>	<p>472 Control group 3,426 ± 459</p> <p><u>Caesarean section rate</u> Asphyxia group 23/71 Control group 11/71 p = 0.01</p> <p><b>Inclusion criteria</b> For infants in the asphyxia group: - Umbilical artery base deficit &gt; 16 mmol/l</p> <p>Infants in control group: - Umbilical artery base deficit &lt; 8 mmol/l</p> <p><b>Exclusion criteria</b> Not specified</p>		<p>birth weights (± 150 g) and gestational age (± 1 week). The control infant was the next one after the asphyxia case that met the criteria. The severity of asphyxia was classified as mild (n = 41), moderate (n = 17) or severe (n = 13) on the basis of short term outcome or expressed by newborn encephalopathy and other newborn organ system complications.</p>	<p>specificity - 29% PPV - 2.6 NPV - 99.5</p>	
<p><b>Full citation</b> Menihan,C.A., Phipps,M., Weitzen,S., Fetal heart rate patterns and sudden infant death syndrome, JOGNN - Journal of Obstetric,</p>	<p><b>Sample size</b> Cases n = 29 Controls n = 98</p> <p><b>Characteristics</b> There were no significant differences observed between the</p>	<p><b>Interventions</b> Electronic fetal heart monitoring (EFM)</p>	<p><b>Details</b> Data were obtained from 127 infants born during the study period at Women and Infants Hospital in Rhode Island. Thirty</p>	<p><b>Results</b> <b>FHR measures among fetuses ≥ 32 weeks</b>  <b><u>Baseline variability in 1st hour of tracing</u></b> <u>Increased or moderate</u> Cases n = 15 (57%) Controls n = 56 (78%) Unadjusted OR: not reported (NR)</p>	<p><b>Limitations</b>  <b>Other information</b> Statistical differences were found in demographic characteristics between sudden infant death syndrome mother-infant couples and their controls. However, no differences were detected in the intrapartum electronic fetal monitoring</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Gynecologic, and Neonatal Nursing, 35, 116-122, 2006</p> <p><b>Ref Id</b> 117077</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Study type</b> Retrospective case control study</p> <p><b>Aim of the study</b> To determine differences in electronic fetal monitoring patterns between infants who died of sudden infant death syndrome (SIDS) and controls.</p> <p><b>Study dates</b> Between 1990 and 1998</p>	<p>two groups in previous live birth, any obstetric and medical conditions (mixed population), maternal surgeries, medication and vitamins taken during pregnancy and prior infant birth weight &lt; 2500g.</p> <p>Compared with controls (n = 98), the mothers whose infants subsequently died of SIDS (n = 29), were younger (22 vs. 28 years; p &lt; 0.01), were more likely to receive Medicaid health insurance (odds ratio 4.6; confidence interval 1.9 to 11.2), were more likely to be unmarried (odds ratio 5.2; confidence interval 2.1 to 12.8), had less intention to breastfeed (26% vs. 57%), and were more likely to smoke (odds ratio 4.6; confidence interval 9 to 11.2).</p> <p><b>Inclusion criteria</b> Infants born between 1990 and 1998 who subsequently died of</p>		<p>two infants (n = 32) who had been born at the hospital were chosen as potential cases and the control infants for each of 32 SIDS cases were selected by computer, matching the day of birth for each case (unclear if mode of birth was matched). A total of 96 infants were identified for the control group. The birth certificates of each of 32 SIDS babies were reviewed by one of the researchers for confirmation of autopsy result. 29/32 infants were confirmed as SIDS and included in the study. The reasons for death in three</p>	<p><u>Minimal or absent</u> Cases n = 5 (45%) Controls n = 16 (23%) Unadjusted OR 1.2 (95% CI: NR)</p> <p><b><u>Baseline variability in last hour of tracing</u></b> <u>Increased or moderate</u> Cases n = 9 (45%) Controls n = 35 (49%) Unadjusted OR: NR</p> <p><u>Minimal or absent</u> Cases n = 11 (55%) Controls n = 36 (51%) Unadjusted OR 1.2 (95% CI 0.4 to 3.2)</p> <p><b><u>Fetal sleep cycles during tracing</u></b> <u>Present throughout tracing</u> Cases n = 1 (5%) Controls n = 14 (20) Unadjusted OR: NR</p> <p><u>50% -75% of tracing</u> Cases n = 7 (35%) Controls n = 24 (34%) Unadjusted OR 4.1 (95% CI 0.5 to 52.3)</p> <p><u>25% - 49% of tracing</u> Cases n = 4 (20%) Controls n = 11 (16%) Unadjusted OR 5.1 (95% CI 0.5 to 43.4)</p> <p><u>&lt; 25% of tracing</u> Cases n = 6 (30%) Controls n = 18 (26%) Unadjusted OR 4.7 (95% CI 0.6 to 139.6)</p> <p><u>Not present during tracing</u> Cases n = 2 (10%)</p>	<p>records, specifically in variability and sleep/wake cycles.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p> <p>Association of Women's Health, Obstetrics, and neonatal Nurses Philips Grant</p>	<p>sudden infant death syndrome (SIDS) and controls.</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>other infants were unclear - SIDS was listed as a possible diagnosis in their death certificate.</p> <p><u>Sample size</u> For the sample size calculation it assumed 50% of SIDS victims would have minimal or absent variability in the EFM readings, and 20% of controls would have minimal or absent variability in their EFM readings. Therefore 3 control per case incorporated and an alpha error of 0.05 and beta error of 20 included. Based on these assumptions, a sample size of 112 (28 cases and 84</p>	<p>Controls n = 3 (5%) Unadjusted OR 9.3 (95% CI: NR)</p> <p><b><u>Fetal sleep cycles (dichotomised)</u></b></p> <p><u>50% - 100% of tracing</u> Cases n = 8 (40%) Controls n = 38 (54%) Unadjusted OR: NR</p> <p><u>0% - 49% of tracing</u> Cases n = 12 (60%) Controls n = 32 (46%) Unadjusted OR 1.8 (95% CI 0.6 to 4.0)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			controls) was needed for the study.  <u>Data analysis</u> Data were analysed using Student's t test for continuous variables and chi-square and Fisher's exact test for categorical variables.		
<p><b>Full citation</b></p> <p>Murphy,K.W., Russell,V., Collins,A., Johnson,P., The prevalence, aetiology and clinical significance of pseudo-sinusoidal fetal heart rate patterns in labour, British Journal of Obstetrics and Gynaecology, 98, 1093-1101, 1991</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>n = 1520 women who had fetal monitoring during labour for various reason were reviewed Intervention n = 230 Control n = 100</p> <p><b>Characteristics</b></p> <p>The reasons for monitoring were (high risk and low risk population): Oxytocin (31%) Hypertensive disorder and intrauterine growth retardation (22%)</p>	<p><b>Interventions</b></p> <p>Sinusoidal and pseudo-sinusoidal patterns</p>	<p><b>Details</b></p> <p>Study conducted in John Radcliffe Hospital, Oxford, over a 6 month period in which all women who had continuous FHR monitoring in labour had their intrapartum CTGs inspected for the presence of sinusoidal or pseudo-</p>	<p><b>Results</b></p> <p>Intervention n = 230 with pseudo-sinusoidal patterns (n = 219 were minor and n = 11 intermediate patterns) Control n = 100 with no sinusoidal pattern</p> <p><u>Minor pseudo-sinusoidal n = 65/219 (30%)</u> Control group n = 26/100 (26%)</p> <p><u>Frequency distribution of minor pseudo sinusoidal patterns in the study group</u></p> <p>Number of pseudo sinusoidal episodes per subject n = 1 Number of subjects n = 94 (42%) Number of pseudo sinusoidal episodes per subject n =2 Number of subjects n = 71 (32%) Number of pseudo sinusoidal episodes per subject n = 3 Number of subjects n = 38 (17%) Number of pseudo sinusoidal episodes per</p>	<p><b>Limitations</b></p> <p>Unclear how and by whom data were analysed and if the assessor was blinded to the outcomes</p> <p><b>Other information</b></p> <p>Pseudo-sinusoidal pattern classification: - Minor when the amplitude of the oscillations was 5-15 beats/min - Intermediate at 16-24 beats/min - Major when the amplitude was <math>\geq 25</math> cycle frequency was 2-5 cycles/min for minor and intermediate patterns and 1-2 cycles/min for major patterns</p> <p>CTG classified as normal or abnormal according to the criteria suggested by Steer et al. (1989)</p> <p>Uterine hyper-stimulation: - When more than 15 contractions were present during a 30 min period</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>122221</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Prospective Cohort</p> <p><b>Aim of the study</b></p> <p>To investigate the prevalence of sinusoidal and pseudo-sinusoidal fetal heart rate (FHR) patterns in labour and the relation between the characteristics of the FHR pattern and fetal outcome.</p> <p><b>Study dates</b></p> <p>September 1987 to February 1988</p> <p><b>Source of funding</b></p>	<p>Epidural analgesia (15%)</p> <p>Breech (4%)</p> <p>Irregular FHR on auscultation (3%)</p> <p>Others (16%)</p> <p><b>Inclusion criteria</b></p> <p>All women who had fetal monitoring in labour during the study time (49% of all labours were monitored).</p> <p>Only cardiocographs (CTG) with pseudo-sinusoidal pattern which persisted <math>\geq 10</math> min were included</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>sinusoidal FHR patterns.</p> <p>Control: Every tenth women who was monitored during the study period and who did not have a sinusoidal or pseudo-sinusoidal FHR pattern was selected as a control. Intrapartum ultrasonography was undertaken in a small pseudo-sinusoidal episode in order to look for fetal sucking or mouth movements.</p> <p>Analysis: Both internal (electrocardiographic) and external (ultrasonic) recordings of FHR were analysed. The intrapartum</p>	<p>subject <math>n &gt; 4</math></p> <p>Number of subjects <math>n = 18</math> (8%)</p> <p><u>Caesarean section rates</u></p> <p>Minor pseudo-sinusoidal <math>n = 22/219</math> (10%)</p> <p>Control group <math>n = 12/100</math> (12%)</p> <p><math>p = ns</math></p> <p><u>Instrumental vaginal birth</u></p> <p>Minor pseudo-sinusoidal <math>n = 65/219</math> (30%)</p> <p>Control group <math>n = 26/100</math> (26%)</p> <p><math>p = ns</math></p> <p><u>Fetal sleep pattern present</u></p> <p>Minor pseudo-sinusoidal <math>n = 125/219</math> (57%)</p> <p>Control group <math>n = 51/100</math> (51%)</p> <p><math>p = ns</math></p> <p><u>Umbilical artery pH <math>&lt; 7.12</math> (measured in 67% of intervention group and 57% of the control group)</u></p> <p>Minor pseudo-sinusoidal <math>n = 20/147</math> (14%)</p> <p>Control group <math>n = 5/57</math> (9%)</p> <p><math>p = ns</math></p> <p><u>Admission to special care</u></p> <p>Minor pseudo-sinusoidal <math>n = 19</math> (9%)</p> <p>Control group <math>n = 4</math> (4%)</p> <p><math>p = ns</math></p> <p>Significant association with epidural analgesia (RR 1.84; 95% CI 1.24 to 2.76) and pethidine administration (RR 1.84; 95% CI 1.31 to 2.59) from multivariate analysis.</p>	<p>Data on pseudo sinusoidal traces divided into minor, moderate and severe categories depending on amplitude of oscillations and frequency of cycles.</p> <p>CTGs were classified as normal or abnormal according to criteria suggested by Steer et al. (1989)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Not specified			<p>CTGs were reviewed immediately after recordings were made. To compare the results between the study group and the control group univariate analyses were performed. The reviewers examined the association between the presence of pseudo-sinusoidal patterns and some variables. Multivariate analyses (logistic regression analysis) were performed.</p>		
<p><b>Full citation</b></p> <p>Nelson,K.B., Dambrosia,J.M., Ting,T.Y., Grether,J.K.,</p>	<p><b>Sample size</b></p> <p>n = 95 infants with cerebral palsy (CP) at aged 3 years with n</p>	<p><b>Interventions</b></p> <p>Continuous electronic fetal monitoring (EFM) (except</p>	<p><b>Details</b></p> <p>Data were collected from singleton children born</p>	<p><b>Results</b></p> <p><u>Heart rate patterns according to presence (n = 78) or absence of cerebral palsy (n = 300)</u></p>	<p><b>Limitations</b></p> <p>The findings on fetal monitoring record were those noted in the birth records, as indicated by the physicians attending the deliveries. No monitoring strips were available for this study.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Uncertain value of electronic fetal monitoring in predicting cerebral palsy, New England Journal of Medicine, 334, 613-618, 1996</p> <p><b>Ref Id</b> 171881</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Study type</b> Case control study</p> <p><b>Aim of the study</b> To investigate the usefulness of fetal monitoring as interpreted by the obstetrician at the time of birth of infants who were diagnosed with cerebral palsy</p>	<p>= 378 matched controls</p> <p><b>Characteristics</b></p> <p><u>Maternal parity (nulliparous)</u> Children with CP: n = 42 (54%) Controls: n = 144 (48%)</p> <p><u>Maternal gestational age (means)</u> Children with CP: 40 weeks Controls: n = 40 weeks</p> <p><u>Maternal age (mean)</u> Children with CP: 28 yr Controls: 27 yr</p> <p><u>Induction of labour</u> Children with CP: n = 13 (17%) Controls: n = 48 (16%)</p> <p><u>Internal monitoring</u> Children with CP: n = 45 (58%) Controls: n = 170 (57%)</p> <p><b>Inclusion criteria</b> Singleton infants with birth weight of 2500 grams or more</p>	<p>in 9% of CP cases and 13% of controls)</p>	<p>during the three-year study period in four counties in the San Francisco area. All weighed 2500 g or more at birth, survived to the age of three years, and had moderate or severe cerebral palsy. The inclusion or exclusion of each identified child was determined by means of a standardised clinical examination or extensive review of the medical records. Controls were randomly selected from the singleton children who met all the criteria for the case children except the diagnosis of cerebral palsy.</p>	<p><u>Tachycardia &gt; 160 bpm</u> Children with CP: n = 22 (28%) Control: n = 85 (28.3%) Odds ratio 1.0 (0.6 to 1.7)</p> <p><u>Tachycardia &gt; 180 bpm</u> Children with CP: n = 5 (6.4%) Control: n = 16 (5.3%) Odds ratio 1.3 (0.4 to 3.4)</p> <p><u>Bradycardia &lt; 100 bpm</u> Children with CP: n = 27 (34.6%) Control: n = 75 (25%) Odds ratio 1.5 (0.9 to 2.5)</p> <p><u>Bradycardia &lt; 80 bpm</u> Children with CP: n = 13 (16.7%) Control: n = 35 (11.7%) Odds ratio 1.5 (0.8 to 3)</p> <p><u>Multiple late decelerations</u> Children with CP: n = 11 (14.1%) Control: n = 12 (4.0%) Odds ratio 3.9 (1.7 to 9.3)</p> <p><u>Decreased beat to beat variability</u> Children with CP: n = 13 (16.7%) Control: n = 21 (7%) Odds ratio 2.7 (1.1 to 5.8)</p> <p><u>MLD/DV</u> Children with CP: n = 21 (26.9%) Control: n = 28 (9.3%) Odds ratio 3.6 (1.9 to 6.7)</p> <p><u>Association between multiple late decelerations, decreased variability or both with cerebral palsy in high and low risk populations</u> <u>Low</u></p>	<p>No actual definition of reduced beat-to-beat variability or multiple late decelerations.</p> <p>Duration of monitoring or specific heart-rate patterns not specified in the analysis.</p> <p><b>Other information</b> Cerebral palsy defined as chronic disability originating from central nervous system, characterised by aberrant control of movement or posture, appearing in early life, and not resulting from progressive disease</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study dates</b></p> <p>From 1983 to 1985</p> <p><b>Source of funding</b></p> <p>Supported in part by a cooperative agreement with the Center for Environmental Health and Injury Control, Centers for Disease Control and Prevention, in part by funds from the Comprehensive Environmental Response, Compensation, and Liability Act Trust Fund through an interagency agreement with the Agency for Toxic Substances and Disease Registry, Public Health Service, and in part by a training grant from the Department of Health and</p>	<p><b>Exclusion criteria</b></p> <p>Children in whom cerebral palsy was acquired after the first 28 days of life or through non-accidental head trauma in the first month and children with mild involvement or isolated hypotonia were not included.</p>		<p>Demographic data were extracted by nurses working at the California Birth Defects Monitoring Program who did not know whether the records were those of case or control children and did not know that the study was about cerebral palsy. The findings on fetal monitoring record were those noted in the birth records, as indicated by the physicians attending the deliveries. No monitoring strips were available for this study. Data collected on the highest fetal heart rate above 160 or 180 beats per minute, the</p>	<p>Sensitivity: 13.8 Specificity: 91.3 PPV: 0.05</p> <p><u>High</u> Sensitivity: 13.8 Specificity: 89.1 PPV: 0.25</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Human Services, Maternal and Child Health Bureau.			<p>lowest fetal heart rate below 100 or 80 beats per minute, and the presence or absence of multiple late decelerations (commonly defined as bradycardia occurring well after the onset of uterine contractions, although in this study the term was recorded as used by the clinicians involved) and decreased beat-to-beat variability in heart rate. Multiple late decelerations and decreased beat-to-beat variability were then combined into a single variable indicating the occurrence of either or both during labor.</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Ozden,S., Demirci,F., Significance for fetal outcome of poor prognostic features in fetal heart rate traces with variable decelerations, Archives of Gynecology and Obstetrics, 262, 141-149, 1999</p> <p><b>Ref Id</b></p> <p>197028</p> <p><b>Country/ies where the study was carried out</b></p> <p>Turkey</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To determine the clinical significance of the existence of poor prognostic features in fetal heart rate (FHR)</p>	<p><b>Sample size</b></p> <p>167 'randomly' selected FHR traces Study group n = 76 with variable decelerations. Divided to two groups poor cases with poor prognostic features (PPFs) (n = 45) and poor cases without PPFs (n = 31) Control group n = 91 normal traces</p> <p><b>Characteristics</b></p> <p>No significant differences observed between the two group in maternal age, gravidity, parity, and cervical dilatation.</p> <p><b>Inclusion criteria</b></p> <p>Singleton</p> <p>Term pregnancy</p> <p><b>Exclusion criteria</b></p> <p>Poorly documented gestational age</p> <p>Premature birth</p>	<p><b>Interventions</b></p> <p>Variable deceleration classified into 7 subtypes according to PPFs</p> <ol style="list-style-type: none"> <li>1. Loss of primary acceleration</li> <li>2. Loss of secondary acceleration</li> <li>3. Loss of variability during deceleration</li> <li>4. Slow return to baseline</li> <li>5. Biphasic deceleration</li> <li>6. Prolonged secondary acceleration</li> <li>7. Prolonged deceleration</li> </ol>	<p><b>Details</b></p> <p>Data for the study were collected from n = 167 randomly selected women with a singleton pregnancy at term. n = 96 women who had an FHR trace without pathological features were selected as a control group. The remaining 76 women had variable decelerations and their FHR traces were analysed for the existence of poor prognostic features. All the traces were analysed by one study author. Umbilical cord pH were taken for included women and pH &lt; 7.20 were</p>	<p><b>Results</b></p> <p><b>Mode of birth</b></p> <p><u>Vaginal birth</u> Study group: poor ( PPFs) n = 25/45 (55.6%); poor (- PPFs) n = 18/31 (58%) Control group n = 65/91 (71.4%) P = ns</p> <p><u>Caesarean section</u> Study group: poor ( PPFs) n = 20/45 (44.4%); poor (- PPFs) n = 13/31 (41.9%) Control group n = 26/91 (28.6%) P = ns</p> <p><u>pH</u> Study group: poor ( PPFs) n 7.18 - 0.08 poor (- PPFs) 7.24 - 0.08 Control group 7.27 - 0.06 P = 0.00001</p> <p><b>Comparison of vriable deceleration subgroups to the number of poor prognostic features for the neonatal outcomes</b></p> <p><u>Vaginal birth</u> Study group: PPF0 n = 18/31 (58%); PPF1 n = 9/13 (69%); PPF2 n = 7/12 (58%); PPF3 n = 5/8 (62%); PPF 4 4/12 (33%) p = ns (comparison between the group without PPF n = 31 and with PPF n = 45)</p> <p><u>Caesarean section</u> Study group: PPF0 n = 13/31 (42%); PPF1 n = 4/13 (31%); PPF2 n = 5/12 (42%); PPF3 n = 3/8 (37%); PPF 4 8/12 (67%) Caesarean section</p> <p><u>PH</u> Study group: PPF0 7.24 - 0.08; PPF1 7.20 -</p>	<p><b>Limitations</b></p> <p>Complex analysis</p> <p>Small sample size</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>traces with variable decelerations.</p> <p><b>Study dates</b></p> <p>From January 1995 to January 1996</p> <p><b>Source of funding</b></p> <p>Not specified</p>	Multiple pregnancy		<p>defined as acidemia.</p> <p><u>Analysis</u> Statistical analysis performed using SPSS. Kruskal Wallis one way ANOVA was used to compare cord blood gas value among the three groups.</p>	0.06; PPF2 7.15 - 0.09; PPF3 7.18 - 0.08; PPF 4 7.18 - 0.01 p = 0.02	
<p><b>Full citation</b></p> <p>Powell,O.H., Melville,A., MacKenna,J., Fetal heart rate acceleration in labor: excellent prognostic indicator, American Journal of Obstetrics and Gynecology, 134, 36-38, 1979</p> <p><b>Ref Id</b></p> <p>196676</p>	<p><b>Sample size</b></p> <p>n = 1677 monitored labours</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <p>Not specified</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>	<p><b>Interventions</b></p> <p>Uniform accelerations (&gt; 3 in 15 minutes &gt; 15 beats for &gt; 15s)</p>	<p><b>Details</b></p> <p>Infants born during the study period in a teaching hospital of the Eastern Virginia Medical school, who met the inclusion criteria, were included in the study. All labouring women had electronic fetal monitoring (EFM) routinely. 65%</p>	<p><b>Results</b></p> <p>Mortality rate of the hospital during the study period: 18.6/1000</p> <p>Mortality rate of group of monitored women during the study period: 14.9/1000</p> <p>Acceleration present in 935 women who were monitored</p> <p><u>Perinatal mortality</u> Acceleration present: n = 4 per 1000 Acceleration not present: n = 20 per 1000</p> <p>The 4 deaths in the "acceleration" group were due to pneumonia in one case (a term infant), due to intracranial haemorrhage in one case (a 37 week infant delivered by midforceps), and due to respiratory distress syndromes in two babies.</p>	<p><b>Limitations</b></p> <p>No population data presented.</p> <p>Unclear how and by whom the data were analysed.</p> <p>No inclusion/exclusion criteria specified.</p> <p>Unclear what percentage of premature labour and high risk pregnancies were included.</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort study</p> <p><b>Aim of the study</b></p> <p>To examine correlation between fetal heart rate (FHR) acceleration and neonatal outcomes</p> <p><b>Study dates</b></p> <p>January 1976 to December 1976</p> <p><b>Source of funding</b></p> <p>Not specified</p>			<p>of the study population gave birth in the private section and 35% in the usual section of the clinic. Only traces with uniform FHR acceleration patterns were included. The accelerations occurring in association with decelerations were excluded.</p>	<p>In the 20 babies who died in the "no accelerations" group, the deaths were often associated with hypoxia (such as: diabetes, post maturity, sepsis, preeclampsia) that were demonstrable in 16 babies. Two (n = 2) died from respiratory distress syndrome and two died with congenital abnormality syndrome.</p> <p>There was no difference in the presence of accelerations in vertex and non vertex presentations. n = 91 women had breech presentation. n = 76 were monitored and only n = 2 failed to show acceleration in labour. There was one death among breech births which was due to severe hypoxia in a vaginal birth and there were no accelerations present during labour for this baby.</p>	
<p><b>Full citation</b></p> <p>Roy,K.K., Baruah,J., Kumar,S., Deorari,A.K.,</p>	<p><b>Sample size</b></p> <p>Total n = 217</p>	<p><b>Interventions</b></p> <p>Caesarean section for non reassuring fetal heart rate</p>	<p><b>Details</b></p> <p>During the study period, a total of 3,148 women</p>	<p><b>Results</b></p> <p><b>Various fetal heart abnormalities indicated by CTG and its relation to immediate adverse neonatal outcomes</b></p>	<p><b>Limitations</b></p> <p>No definition for bradycardia, deceleration and non reassuring CTG provided.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Sharma, J.B., Karmakar, D., Cesarean section for suspected fetal distress, continuous fetal heart monitoring and decision to delivery time, Indian Journal of Pediatrics, 75, 1249-1252, 2008</p> <p><b>Ref Id</b> 60814</p> <p><b>Country/ies where the study was carried out</b> India</p> <p><b>Study type</b> Prospective observational study</p> <p><b>Aim of the study</b> To find out the efficacy of continuous fetal heart monitoring by analysing the cases of cesarean section for non</p>	<p><b>Characteristics</b> Not specified</p> <p><b>Inclusion criteria</b> Gestational age <math>\geq</math> 36 No fetal anomalies Non reassuring CTG not responding to conservative management (including changing the maternal position, intravenous hydration, and oxygen administration)</p> <p><b>Exclusion criteria</b> Abnormal presentation Multiple pregnancy Intrauterine growth restriction (IUGR) Caesarean section for other primary indications</p>	(FHR) detected by cardiotocograph (CTG)	<p>delivered in a maternity unit of whom 217 (6.8%) women underwent cesarean section for non-reassuring fetal heart trace in labor. The percentage of caesarean sections for various indications was 16.2%. The maternal demographic profile, specific types of abnormal fetal heart rate tracing and the decision to delivery time interval were noted. The decision time to perform a caesarean section was defined as when the senior resident on duty took the decision to perform the caesarean and exact delivery time. The</p>	<p><u>Persistent bradycardia n = 106/217 (48.8%)</u> 5 minutes Apgar &lt; 7 n = 16/106 Umbilical cord pH &lt; 7.10 n = 4/106 NICU admission n = 16/106</p> <p><u>Recurrent late deceleration n = 56 (25.8%)</u> 5 minutes Apgar &lt; 7 n = 10/56 Umbilical cord pH &lt; 7.10 n = 5/56 NICU admission n = 10/56</p> <p><u>Variable deceleration n = 38/217 (17.5%)</u> 5 minutes Apgar &lt; 7 n = 7/38 Umbilical cord pH &lt; 7.10 n = 4/38 NICU admission n = 7/38</p> <p><u>Decreased variability n = 17/217 (7.8%)</u> 5 minutes Apgar &lt; 7 n = nil Umbilical cord pH &lt; 7.10 n = nil NICU admission n = nil</p> <p><b>Overall findings for non-reassuring CTG and its relation to the neonatal outcomes</b> Decision to delivery interval (DDI): DDI <math>\leq</math> 30 min n = 121/217 DDI &gt; 30 min n = 96/217</p> <p><u>5 minutes apgar &lt; 7</u> DDI <math>\leq</math> 30 min n = 18/121 (14.8%) DDI &gt; 30 min n = 15/96 (15.6%) p = ns</p> <p><u>Arterial cord pH &lt; 7.10</u> DDI <math>\leq</math> 30 min n = 8/121 (6.6%) DDI &gt; 30 min n = 5/96 (5.2%) p = ns</p> <p><u>NICU admission for suspected birth asphyxia</u> DDI <math>\leq</math> 30 min n = 26/121 (21.4%) DDI &gt; 30 min n = 7/96 (7.2%) p &lt; 0.05</p>	<p>Unclear if the outcome assessors were blinded to the study groups allocation.</p> <p>Women's demographic characteristics not reported.</p> <p><b>Other information</b> Non-reassuring fetal heart rate detected by CTG did not correlate well with adverse neonatal outcome.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>reassuring fetal heart in labour, detected by cardiotocography (CTG) and correlating these cases with perinatal outcome.</p> <p><b>Study dates</b></p> <p>March 2002 to March 2007</p> <p><b>Source of funding</b></p> <p>Not specified</p>			<p>adverse immediate neonatal outcomes in terms of Apgar score &lt; 7 at 5 minutes, umbilical cord pH &lt; 7.10, neonates requiring immediate ventilation and NICU admissions were recorded. The correlation between non-reassuring fetal heart, decision to delivery interval and neonatal outcome were analysed.</p> <p><u>Data analysis</u> Statistical analysis was done using Student's t-test and chi square test where appropriate.</p>	<p><u>Fresh stillbirth</u> DDI ≤ 30 min n = 1*/121 (0.8%) DDI &gt; 30 min n = nil p &lt; 0.05</p> <p>*Death was due to placental abruption</p> <p><u>Born healthy</u> n = 184 (84.7%)</p>	
<b>Full citation</b>	<b>Sample size</b>	<b>Interventions</b>	<b>Details</b>	<b>Results</b>	<b>Limitations</b>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Salim,R., Garmi,G., Nachum,Z., Shalev,E., The impact of non-significant variable decelerations appearing in the latent phase on delivery mode: a prospective cohort study, Reproductive Biology and Endocrinology, 8, 81-, 2010</p> <p><b>Ref Id</b> 109319</p> <p><b>Country/ies where the study was carried out</b> Israel</p> <p><b>Study type</b> Prospective cohort</p> <p><b>Aim of the study</b> To estimate the impact of non-significant variable</p>	<p>Category I n = 251</p> <p>Category II NSV n = 186</p> <p>Category II SV n = 76</p> <p><b>Characteristics</b> There were no significant differences observed between the three groups in maternal age, parity and polyhydramnios.</p> <p><b>Inclusion criteria</b> Term pregnancy (<math>\geq 37</math>)</p> <p>In the latent phase of labour (defined as interval between the start of regular contractions combined with any cervical dynamics [dilating &gt; 4 cm])</p> <p>Singleton pregnancy</p> <p><b>Exclusion criteria</b> Fetal heart tracing abnormalities during the latent phase</p>	<p>Electronic fetal monitoring (EFM)</p>	<p>Variable deceleration was defined according to 2008 National Institute of Child Health and Human Development workshop. Variable decelerations were categorised as significant (SV) if fetal heart rate (FHR) reached 70 beats/min for one minute or more but less than 2 minutes, otherwise they were categorised as non-significant (NSV)</p> <p>Women were divided into three groups. All had a fetal heart rate tracing with normal baseline and variability:</p> <p>Study group (Category II</p>	<p>Total n = 1005</p> <p>Category II-NSV tracings (study group) n = 186</p> <p>Category II-SV n = 76</p> <p>Category I tracings n = 251</p> <p><b>Mode of birth</b> There was a statistically significant differences observed between the three groups in method of birth (category II-SV versus category I and category II-NSV) (<math>p = 0.0001</math>)</p> <p><u>Spontaneous vaginal birth</u> Control group (Category I): n = 238 (94.8%) Study group (Category II NSV): n = 166 (89.2%) Second control group (Category II SV): n = 40 (52.6%)</p> <p><u>Vacuum</u> Control group (Category I): n = 6 (2.4%) Study group (Category II NSV): n = 8 (4.3%) Second control group (Category II SV): 11 (14.5%)</p> <p><u>Caesarean</u> Control group (Category I): n = 7 (2.8%) Study group (Category II NSV): n = 12 (6.5%) Second control group (Category II SV): n = 25 (32.9%)</p> <p><u>Reasons for vacuum or caesarean delivery</u> There was a statistically significant difference observed between the three groups in reasons for vacuum or ceasarean delivery (category II-SV versus category I and category II-NSV) (<math>p = 0.0001</math>)</p>	<p><b>Other information</b> Fetal Heart interpretation categorisation from National Institute of Child Health and Human Development workshop 2008 (Macones et al., 2008):</p> <p><b>Category I</b> <u>Category I fetal heart rate (FHR) tracings include all of the following:</u> Baseline rate: 110–160 beats per minute (bpm) Baseline FHR variability: moderate Late or variable decelerations: absent Early decelerations: present or absent Accelerations: present or absent</p> <p><b>Category II</b> Category II FHR tracings include all FHR tracings not categorized as Category I or Category III. Category II tracings may represent an appreciable fraction of those encountered in clinical care. Examples of Category II FHR tracings include any of the following: <u>Baseline rate</u> Bradycardia not accompanied by absent baseline variability Tachycardia</p> <p><u>Baseline FHR variability</u> Minimal baseline variability Absent baseline variability not accompanied by recurrent decelerations Marked baseline variability</p> <p><b>Accelerations</b> Absence of induced accelerations after fetal stimulation</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>decelerations (NSV) appearing during the latent phase of labour on delivery mode and neonatal outcome.</p> <p><b>Study dates</b></p> <p>January to April 2009</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>Caesarean section without a trial of labour</p> <p>Women gave birth to infants with major malformation</p>		<p>NSV): women who had Category II tracing based on Institute of Child Health and Human Development (NICHD) categorisation system; women with NSV, episodic or recurrent, and normal base line and moderate variability</p> <p>Control group (Category I): women who had category I tracing based on NICHD categorisation</p> <p>Second control group (Category II-SV): women who had category II-SV tracing based on NICHD categorisation; women with significant variables (SV)</p>	<p><u>Indication for CS (not reassuring FHR monitoring)</u> Control group (Category I): n = 3 (23.1%) Study group (Category II NSV): n = 5 (25%) Second control group (Category II SV): n = 20 (55.6%)</p> <p><u>Indication for CS (failure to progress in the active or second stage)</u> Control group (Category I): n = 10 (76.9%) Study group (Category II NSV): n = 15 (75.0%) Second control group (Category II SV): n = 16 (44.4%)</p> <p><b>Neonatal outcomes</b> <u>Neonatal weight (g)</u> Control group (Category I): mean 3329 ± 392 Study group (Category II NSV): mean 3397 ± 439 Second control group (Category II SV): mean 3130 ± 487 p = 0.002 (category II-SV versus category I and category II-NSV)</p> <p><u>Neonatal born &lt; 2500 g</u> Control group (Category I): n = 2 (0.8%) Study group (Category II NSV): n = 1 (0.5%) Second control group (Category II SV): n = 4 (5.3%) p = 0.0001 (category II-SV versus category II-NSV)</p> <p><u>Apgar score at 5 min (out of 10)</u> Control group (Category I): mean 9.96 ± 0.23 Study group (Category II NSV): mean 9.90 ± 0.31 Second control group (Category II SV): mean 9.86 ± 0.39 p = 0.01</p>	<p><b>Periodic or episodic decelerations</b> Recurrent variable decelerations accompanied by minimal or moderate baseline variability Prolonged deceleration ≥ 2 minutes but ≤ 10 minutes Recurrent late decelerations with moderate baseline variability Variable decelerations with other characteristics, such as slow return to baseline, "overshoots," or "shoulders"</p> <p><b>Category III</b> <u>Category III FHR tracings include either:</u> Absent baseline FHR variability and any of the following: Recurrent late decelerations Recurrent variable decelerations Bradycardia Sinusoidal pattern</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p><u>Sample size</u> In order to show a difference of 10% in the rate of operative birth between the category I and category II-NSV tracing with an alpha of 0.05 and a power of 80% a sample size of 160 per group was required</p> <p><u>Analysis</u> One-way analysis of variance was used to compare the continuous demographic and clinical variables of the three groups. Significant group differences were tested (post-hoc). Backwards stepwise logistic regression using significant</p>	<p><u>Mean cord PH</u> Control group (Category I): 7.31 ± 0.07 Study group (Category II NSV): 7.31 ± 0.07 Second control group (Category II SV): 7.30 ± 0.08 p = 0.5</p> <p><u>Cord pH between 7.0 to 7.1</u> Control group (Category I): n = 2 (0.8%) Study group (Category II NSV): n = 7 (3.8%) Second control group (Category II SV): n = 4 (5.3%)</p> <p><u>Meconium stained amniotic fluid</u> Control group (Category I): n = 22(8.8%) Study group (Category II NSV): n = 26 (14%) Second control group (Category II SV): n = 15 (19.7%)</p> <p><u>Nuchal cord or true knot</u> Control group (Category I): n = 23 (9.2%) Study group (Category II NSV): n = 19 (10.2%) Second control group (Category II SV): n = 12 (15.8%) p = 0.3</p> <p><u>Neonatal death</u> Control group (Category I): n = 0 Study group (Category II NSV): n = 0 Second control group(Category II SV): n = 0</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>invariables was performed to determine which predicted operative delivery. P &lt; 0.05 was considered significant.</p> <p><u>Assessment</u> All traces were assessed by two obstetricians at the same time, both were blinded to the groups allocation and neonatal outcomes.</p>		
<p><b>Full citation</b></p> <p>Sameshima,H., Ikenoue,T., Predictive value of late decelerations for fetal acidemia in unselective low-risk pregnancies, American Journal of Perinatology, 22, 19-23, 2005</p>	<p><b>Sample size</b></p> <p>Carditocograph (CTG) trace of n = 5522 women with low-risk pregnancies</p> <p><b>Characteristics</b></p> <p><u>Average maternal age</u> No decelerations 28.4 ± 4.8 Occasional LD 30.0 ± 4.9</p>	<p><b>Interventions</b></p> <p>FHR via cardiotocograph (CTG) trace</p>	<p><b>Details</b></p> <p>Clinical significance of late decelerations (LD) of intrapartum fetal heart rate (FHR) monitoring to detect low pH (&lt; 7.1) in low-risk</p>	<p><b>Results</b></p> <p>Occasional LD n = 301/5522 Recurrent LD n = 99/5522</p> <p><u>Recurrent LD n = 99</u> Moderate variability and acceleration n = 64/99 Moderate variability without acceleration n = 16/99 Acceleration with minimal variability n = 3/99 Minimal variability without accelerations n = 16/99</p>	<p><b>Limitations</b></p> <p>Poor reporting of results</p> <p>Unclear if the outcome assessor was blinded to the outcomes</p> <p><b>Other information</b></p> <p>In low-risk pregnancies, information on LD combined with acceleration and baseline variability enables us to predict the potential incidence of fetal acidemia.</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Ref Id</b> 157246</p> <p><b>Country/ies where the study was carried out</b> Japan</p> <p><b>Study type</b> Retrospective cohort study</p> <p><b>Aim of the study</b> To evaluate the clinical significance of late decelerations (LD) of intrapartum fetal heart rate (FHR) monitoring to detect low pH (&lt; 7.1) in low-risk pregnancies.</p> <p><b>Study dates</b> 1995 to 2000</p> <p><b>Source of funding</b></p>	<p>Recurrent LD 38.8 ± 2.0 p = ns</p> <p><u>Average gestational age</u> No decelerations 38.5 ± 1.8 Occasional LD 38.8 ± 2.0 Recurrent LD 38.1 ± 2.5</p> <p>Average parity of the three groups 0.6 ± 0.9</p> <p><b>Inclusion criteria</b> Low risk pregnancies Cases with recurrent and occasional late deceleration (LD)</p> <p><b>Exclusion criteria</b> Premature birth &lt; 32 wk</p> <p>Multiple pregnancy Hypertensive disorders Pre-eclampsia or eclampsia Chronic hypertension</p>		<p>pregnancies was evaluated. Data collected from two secondary and two tertiary-level institutions where 10,030 women delivered. Among them, 5522 were low-risk pregnancies. The last 2 hours of FHR patterns before delivery were interpreted according to the guidelines of the National Institute of Child Health and Human Development. The correlation between the incidence of LD (occasional, &lt; 50%; recurrent, ≥ 50%) and severity (reduced baseline FHR accelerations and variability) of LD, and low</p>	<p>Blood gases and pH values deteriorated as the incidence of LD increased and as baseline accelerations or variability decreased. Positive predictive value for low pH (&lt; 7.1) was exponentially elevated from 0% at no decelerations, 1% in occasional LD, and &gt; 50% in recurrent LD with no baseline FHR accelerations and reduced variability.</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
Supported in part by Grant-in-Aid for Scientific Research from Ministry of Education, Japan	Collagen diseases Diabetes mellitus Thyroid dysfunction Cardiac, respiratory, renal disease Epilepsy Placenta praevia Coagulation disorders Intrauterine infection and chorioamnionitis Intrauterine growth restriction Fetal abnormalities Anomalies Hydrops fetalis Metabolic disorders Known congenital syndromes		pH (< 7.1) wasevaluated.  <u>Statistical analyses</u> Included a contingency table with chi <sup>2</sup> and Fisher's exact test, and one-way analysis of variance with the Bonferroni/Dunn test.		
<b>Full citation</b> Samueloff,A., Langer,O., Berkus,M.,	<b>Sample size</b> n = 2220 consecutive deliveries	<b>Interventions</b> Scoring FHR variability	<b>Details</b> Data were collected from follow up of n =	<b>Results</b> pH ≥ 7.20, <7.20	<b>Limitations</b> Variability not single useful predictor of outcome.

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Field,N., Xenakis,E., Ridgway,L., Is fetal heart rate variability a good predictor of fetal outcome?, Acta Obstetricia et Gynecologica Scandinavica, 73, 39-44, 1994</p> <p><b>Ref Id</b></p> <p>196845</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To investigate whether fetal heart rate (FHR) variability serves as a reliable single predictor of fetal outcome</p> <p><b>Study dates</b></p>	<p><b>Characteristics</b></p> <p>Maternal age (mean <math>\pm</math> SD) 27.4 <math>\pm</math> 6.04</p> <p>Complication in pregnancy (hypertension, diabetes, abrupto placenta, placenta previa, chorioamnionitis, previous caesarean section): 27. 34% Epidural: 47.3%</p> <p><b>Inclusion criteria</b></p> <p>Not specified</p> <p><b>Exclusion criteria</b></p> <p>&lt; 37 weeks gestation</p> <p>Twins</p> <p>Fetal malformation</p> <p>Stillbirth</p>	<p>using 5 scoring systems:</p> <p>A. FHR amplitude variability <math>\geq</math> 3 bpm &lt; 3 bpm</p> <p>B. FHR amplitude <math>\geq</math> 5bpm &lt; 5 bpm</p> <p>C. FHR frequency of oscillations <math>\geq</math> 3 bpm &lt; 3/min</p> <p>D. FHR frequency of oscillations <math>\geq</math> 5 bpm &lt; 5/min</p> <p>E. Combination of (amplitude frequency)/2. Value &lt; 3 scored as low and <math>\geq</math> 3 as high</p>	<p>2200 consecutive births during 1991 from a teaching hospital. Based on the hospital policy, every women entering the labour ward was connected to a fetal heartt monitor. Fetal heart variability data were obtained from n = 1816 women (the missing 7.8% of variability data was due to either imminent birth in which obtaining a trace was not possible or lost tracing).</p> <p><b>Analysis</b></p> <p>Three sections of the trace were analysed:</p> <p>1. early in labour for a period of 30 minutes,</p>	<p>Scoring method A: sensitivity 10.99%, specificity 93.80%, positive predictive value (PPV) 25.20%, negative predictive value (NPV) 84.74%</p> <p>Scoring method B: sensitivity 26.24%, specificity 78.93%, PPV 19.12%, NPV 84.93%</p> <p>Scoring method C: sensitivity 6.78%, specificity 95.18%, PPV 23.17%, NPV 84.48%</p> <p>Scoring method D: sensitivity 25.35%, specificity 90.52%, PPV 19.72%, NPV 85.11%</p> <p>Scoring method E: sensitivity 7.44%, specificity 96.30%, PPV 27.63%, NPV 84.58%</p> <p>Both amplitude and frequency methods poorly sensitive at lower limits (&lt; 3). Sensitivity increased by increasing limit to 5 in both scores but consequent drop in specificity. Combination method has low sensitivity.</p>	<p>Division of cases into normal and abnormal not balanced as non-matched.</p> <p>Hence, performance of tests affected.</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>During 1991</p> <p><b>Source of funding</b></p> <p>not specified</p>			<p>2. 30 minutes of tracing in the active phase</p> <p>3. throughout the entire 2nd stage in segments of 30 minutes (a maximum of three segments). In all deliveries with 2nd stage longer than 90 minutes, the last tracing prior to the delivery was analysed. A total of 4361 tracing segments were analysed by five maternal-fetal faculty members blinded to the maternal and neonatal outcomes.</p>		
<p><b>Full citation</b></p> <p>Sheiner,E., Hadar,A., Hallak,M., Katz,M., Mazor,M.,</p>	<p><b>Sample size</b></p> <p>n = 601</p> <p><b>Characteristics</b></p>	<p><b>Interventions</b></p> <p>Abnormal fetal heart rate tracing</p>	<p><b>Details</b></p> <p>Women were examined at the delivery suite. Based on the hospital</p>	<p><b>Results</b></p> <p><b><u>Pathologic FHR patterns during 2nd stage of labour (compared with normal tracing) associated with pH &lt; 7.2 (n = 57) and base deficit of ≥ 12 (n = 28)</u></b></p>	<p><b>Limitations</b></p> <p>Unclear if the assessors were blinded to the outcomes</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Shoham-Vardi, I., Clinical significance of fetal heart rate tracings during the second stage of labor, Obstetrics and Gynecology, 97, 747-752, 2001</p> <p><b>Ref Id</b> 196075</p> <p><b>Country/ies where the study was carried out</b> Israel</p> <p><b>Study type</b> Cohort</p> <p><b>Aim of the study</b> To examine the importance of abnormal FHR patterns during the second stage of labor in terms of pregnancy outcome</p> <p><b>Study dates</b></p>	<p>Women with abnormal FHR patterns were of significantly lower birth order and more often carried male fetuses compared with women with normal FHR patterns. The women with abnormal FHR tracings during the second stage of labour had a significantly higher rate of oligohydramnios and a non-significantly higher rate of hydramnios. No other significant differences were seen between the groups for anesthesia use, first and second stage duration, presence of meconium in amniotic fluid, cord problems, and birth weight.</p> <p><b>Inclusion criteria</b></p> <p>Low risk pregnancy</p> <p>Singleton gestation</p> <p>Vertex presentation</p> <p>Term delivery (greater than 37 completed weeks gestation)</p>		<p>policy, all labouring women had continuous fetal monitoring and the monitor patterns were checked and the findings documented hourly. The same obstetrician collected the data after carefully evaluating both the monitor files and the flow charts. Tracings were interpreted using the guidelines of the National Institute of Child Health and Human Development Research Planning Workshop.</p> <p>The cumulative depth of decelerations or bradycardia was classified by a nadir of</p>	<p><u>Variable decelerations ≥ 70 bpm</u> pH &lt; 7.2 OR 5.1 (95% CI 1.4 to 21.4) p = 0.008</p> <p>Base deficit of ≥ 12 OR 3.5 (95% CI 0.8 to 15.8) p = 0.101</p> <p><u>Variable decelerations &lt; 70 bpm</u> pH &lt; 7.2 OR 16.3 (95% CI 3.8 to 80.5) p &lt; 0.001</p> <p>Base deficit of ≥ 12 OR 10.5 (95% CI 1.9 to 56.4) p = 0.006</p> <p><u>Late decelerations</u> pH &lt; 7.2 OR 15.2 (95% CI 2.8 to 91.4) p &lt; 0.001</p> <p>Base deficit of ≥ 12 OR 17.3 (95% CI 2.9 to 101.9) p = 0.002</p> <p><u>Bradycardia ≥ 70 bpm</u> pH &lt; 7.2 OR 2.3 (95% CI 0.3 to 17.1) p = 0.390</p> <p>Base deficit of ≥ 12 OR 3.8 (95% CI 0.3 to 44.2) p = 0.282</p> <p><u>Bradycardia &lt; 70 bpm</u> pH &lt; 7.2 OR 26.6 (95% CI 5.2 to 150.3) p &lt; 0.001</p> <p>Base deficit of ≥ 12 OR 5.2 (95% CI 0.8 to 31.9) p = 0.007</p> <p><u>Bradycardia &lt; 70 bpm</u> pH &lt; 7.2 OR 2.2 (95% CI 0.3 to 17.1) p = 0.728</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>January to June 2000</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p><b>Exclusion criteria</b></p> <p>Uninterpretable tracings</p> <p>Immediate caesarean because of maternal or fetal indications, such as clinical evidence of cephalopelvic disproportion or placental insufficiency</p> <p>Previous caesarean section</p> <p>Pre-existing heart or lung disease</p> <p>Fetuses with known growth restriction or malformations</p>		<p>less than 100 but at least 70 beats per minute, and decelerations with a nadir less than 70 beats per minute.</p> <p>Information was collected about labor duration, performance of an episiotomy, mode of delivery (spontaneous, vacuum, or caesarean), neonatal sex, birth weight, presence of cord problems (nuchal cord or true knot of the cord), Apgar scores, and acid-base status (in particular, metabolic acidosis). The umbilical cord was clamped immediately after delivery. Arterial blood was drawn into</p>	<p>Base deficit of <math>\geq 12</math> OR 5.1 (95% CI 0.6 to 46.1) p = 0.098</p> <p><b><u>Pathologic FHR patterns during 2nd stage of labour (compared with normal tracing) associated with fetal acidosis (pH &lt; 7.2 and base deficit of <math>\geq 12</math>) n = 28</u></b></p> <p><u>Late decelerations</u> OR 3.9 (95% CI 1.1 to 13.1) p = 0.029</p> <p><u>Abnormal tracing during the 1st stage</u> OR 3.4 (95% CI 1.3 to 8.7) p = 0.011</p> <p><u>Bradycardia &lt; 70 bpm</u> OR 3.0 (95% CI 1.02 to 8.6) p = 0.045</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>a 2-ml plastic syringe that was flushed with heparin, and then transferred to the pH machine located in the delivery ward. The pH was considered abnormal when it was lower than 7.2. Base deficit of 12 mmol/l or greater was considered the threshold of fetal metabolic acidosis at delivery. Newborn morbidity included admission to the intensive care unit or delayed discharge from the hospital because of fetal indications. The local ethics institutional review board</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>approved the study.</p> <p><u>Analysis</u>                      Comparison of group means was performed with the SPSS version 8.0 statistical package (SPSS Inc., Chicago, IL). Chi-square or Fisher's exact test was used for comparison of proportions. Student's t-test was applied for comparison of means. <math>P &lt; 0.05</math> was considered statistically significant. Multiple logistic regression models were used to investigate the independent contributions of obstetric factors to abnormal FHR patterns during the second stage of labor</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>and to investigate the contributions of those patterns to selected fetal outcomes. Odds ratios (ORs) and their 95% confidence intervals (CIs) were calculated from the regression coefficients.</p>		
<p><b>Full citation</b></p> <p>Spencer, J.A., Badawi, N., Burton, P., Keogh, J., Pemberton, P., Stanley, F., The intrapartum CTG prior to neonatal encephalopathy at term: a case-control study, British Journal of Obstetrics and Gynaecology, 104, 25-28, 1997</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>Cases n = 55</p> <p>Controls n = 39</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <p>One or more of the following features present during the first week of life:</p> <ul style="list-style-type: none"> <li>- Seizures</li> <li>- Absent or altered responsiveness</li> </ul>	<p><b>Interventions</b></p> <p>Fetal heart rate patterns</p>	<p><b>Details</b></p> <p>All cases of neonatal encephalopathy developing during the first seven days of life in term infants were identified from five hospitals (two teaching and three peripheral) in Perth, Western Australia.</p> <p>One control per case was</p>	<p><b>Results</b></p> <p>Comparison of first and last sections of CTG between cases of neonatal encephalopathy and controls. Individual parameters and Krebs' score derived from 30 min sections. FIGO classification derived from 60 min sections.</p> <p>First CTG section Cases n = 38 Controls n = 35</p> <p><b><u>Late decelerations</u></b></p> <p><b>Cases</b></p> <p>Yes n = 2</p> <p>No n = 36</p> <p><b>Controls</b></p> <p>Yes n = 0</p> <p>No n = 35</p>	<p><b>Limitations</b></p> <p>Low intra-observer agreement</p> <p>No exclusion criteria or women's characteristics reported</p> <p><b>Other information</b></p> <p><b>FIGO FHR pattern</b></p> <p><u>Abnormal (pathological)</u></p> <p>Baseline FHR: &lt; 100, &gt; 170</p> <p>Variability (amplitude bpm): &lt; 5 for 40 min</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>197160</p> <p><b>Country/ies where the study was carried out</b></p> <p>Australia</p> <p><b>Study type</b></p> <p>Case control</p> <p><b>Aim of the study</b></p> <p>To compare cardiotocograph (CTG) records during labour in cases of neonatal encephalopathy and matched controls.</p> <p><b>Study dates</b></p> <p>Eight months during 1992</p> <p><b>Source of funding</b></p> <p>British council and The Royal Society and The Royal College of Obstetrician and Gynaecologists</p>	<p>- Abnormal muscular tone, feeding difficulties of central origin</p> <p>- Difficulty with central control of respiration</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>subsequently selected by matching for hospital of delivery, time and day of the week, sex, and maternal insurance status. All cases and controls had a neurological examination within the first seven days of birth. Clinical data were obtained from the obstetric case notes and a maternal questionnaire. The selected CTG traces were interpreted with out knowledge of the outcome. A note was made of baseline rate, amplitude and frequency of the variability, presence of accelerations, and presence and type of decelerations.</p>	<p><b><u>FHR acceleration</u></b></p> <p><u>Cases</u> Yes n = 16 No n = 22</p> <p><u>Controls</u> Yes n = 8 No n = 27</p> <p><b><u>FHR variability</u></b></p> <p><u>Cases</u> ≤ 5bpm n = 4 &gt; 5 bpm n = 34</p> <p><u>Controls</u> ≤ 5bpm n = 2 &gt; 5 bpm n = 33</p> <p><b><u>Krebs' score</u></b></p> <p><u>Cases</u> 0-3 n = 2 4-10 n = 36</p> <p><u>Controls</u> 0-3 n = 1 4-10 n = 34</p> <p><b><u>FIGO Classification</u></b></p> <p><u>Cases</u> Abnormal n =19 Normal n = 19</p> <p><u>Control</u> Abnormal n = 9 Normal n = 26</p> <p><u>First CTG section Cases n = 38 Controls n = 35</u></p> <p><b><u>Late decelerations</u></b></p> <p><u>Cases</u></p>	<p>Deceleration: severe variable, severe repeated early, prolonged, late or sinusoidal</p> <p><u>Suspicious</u></p> <p>Baseline FHR: 100 – 110, 150 - 170</p> <p>Variability (amplitude bpm): 5 – 10 for 40 min &gt; 25</p> <p>Deceleration/30 min: variable</p> <p><u>Normal</u></p> <p>Baseline FHR: 120 - 150</p> <p>Variability (amplitude bpm): 6 - 25</p> <p>Deceleration/30 min: none</p> <p>FHR scoring for internal FHR monitoring; for each of the criteria 0 to 2 points may be given so that a score of 0 to 10 may be obtained</p> <p>Abnormal: score 0 – 3</p> <p>Suspicious: score 4 – 6</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
(Ethicon travel grant)			<p>Krebs' intrapartum CTG score 9 for the first and last 30 min of the trace was calculated, as defined. The total score for each section of CTG was considered abnormal (score 0-3), suspicious (score 4-6) or normal (score 7-10) and these classifications were reduced to two groupings for analyses. The FIGO classification 3 was also determined for the first and last hour of each CTG. Half of the traces were reviewed on a second occasion, at least 10 days later. Intra-observer reproducibility was evaluated</p>	<p>Yes n = 17 No n = 19</p> <p><u>Controls</u> Yes n = 8 No n = 23</p> <p><b><u>FHR acceleration</u></b> <u>Cases</u> Yes n = 26 No n = 10</p> <p><u>Controls</u> Yes n = 15 No n = 16</p> <p><b><u>FHR variability</u></b> <u>Cases</u> ≤ 5bpm n = 14 &gt; 5 bpm n = 22</p> <p><u>Controls</u> ≤ 5bpm n = 4 &gt; 5 bpm n = 27</p> <p><b><u>Krebs' score</u></b> <u>Cases</u> 0-3 n = 19 4-10 n = 17</p> <p><u>Controls</u> 0-3 n = 10 4-10 n = 21</p> <p><b><u>IGO Classification</u></b> <u>Cases</u> Abnormal n =32 Normal n = 4</p> <p><u>Control</u></p>	<p>Normal: score 7 – 10</p> <p><u>Score 0</u></p> <p>Baseline FHR: &lt; 100, &gt; 180</p> <p>Variability (amplitude bpm): &lt; 3</p> <p>Variability (frequency bpm): &lt; 3</p> <p>Acceleration/30 min: 0</p> <p>Deceleration/30 min: late, severe variable, atypical variable = 0 score</p> <p><u>Score 1</u></p> <p>Baseline FHR: 100 - 119, 161 -180</p> <p>Variability (amplitude bpm): 3 - 5 &gt; 25</p> <p>Variability (frequency bpm): 3 - 6</p> <p>Acceleration/30 min: 1 -4</p> <p>Deceleration/30 min: moderate variable</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>using Cohen's Kappa. <u>Analysis</u> Associations between case-control status and binary explanatory variables were assessed using the x2 test for association, or Fisher's exact test if the expected cell count was 5 or less.</p>	<p>Abnormal n = 16 Normal n = 15</p> <p>Intra-observer reproducibility using Cohen's Kappa for the 1st and last sections of CTG traces (Krebs' score) First section: 0.58 (95% CI 0.30 to 0.87) Last section 0.40 (95% CI 0.16 to 0.62)</p> <p>Intra-observer reproducibility using Cohen's Kappa for the 1st and last sections of CTG traces (FIGO classification) First section: 0.47 (95% CI 0.24 to 0.70) Last section 0.33 (95% CI 0.12 to 0.55)</p>	<p><u>Score 2</u></p> <p>Baseline FHR: 120 - 160</p> <p>Variability (amplitude bpm): 6 - 25</p> <p>Variability (frequency bpm): &gt; 6</p> <p>Acceleration/30 min: &gt; 4</p> <p>Deceleration/30 min: none, early</p>
<p><b>Full citation</b></p> <p>Spencer,J.A., Johnson,P., Fetal heart rate variability changes and fetal behavioural cycles during labour, British Journal of Obstetrics and Gynaecology, 93, 314-321, 1986</p> <p><b>Ref Id</b></p> <p>174553</p>	<p><b>Sample size</b></p> <p>n = 301 consecutive fetal heart rate (FHR) recording</p> <p><b>Characteristics</b></p> <p><u>Prostagladine/oxytocin</u> Cycle present n = 163 (93%) No cycle present n = 110 (88%)</p> <p><u>pethidine/epidural</u> Cycle present n = 159 (90%) No cycle present n = 117 (94%)</p>	<p><b>Interventions</b></p> <p>FHR variability</p>	<p><b>Details</b></p> <p>During the study period all 1st stage cardiotocograph (CTG) recordings with ≥ 6 hour duration were analysed for cycles of low and high FHR variability episodes. Each episode was visually identified by the change in long term</p>	<p><b>Results</b></p> <p>Mode of birth in presence and on presence of FHR variability cycles <u>Instrumental vaginal birth</u> Cycle present n = 159 (90%) No cycle present n = 117 (94%)</p> <p><u>Caesarean section</u> Cycle present n = 70 (40%) No cycle present n = 51 (41%)</p>	<p><b>Limitations</b></p> <p>No demographic data reported.</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Case control study</p> <p><b>Aim of the study</b></p> <p>To evaluate the cycle of low and high fetal heart rate (FHR) and fetal behavioural cycles</p> <p><b>Study dates</b></p> <p>March 1983 to July 1983</p> <p><b>Source of funding</b></p> <p>Grant from DHSS and the MRC</p>	<p><b>Inclusion criteria</b></p> <p>Term birth</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>variability of <math>\geq 5</math> beats per minute maintained for <math>\geq 5</math> minutes duration. A complete cycle required both low and high FHR variability episodes with changes before and after. The actual variability during the quiet episode (episodes of low FHR variability) of cycles was recorded as <math>&gt; 5</math> or <math>&lt; 5</math> beats/min, and the predominant variability of CTG without cycle was also recorded as <math>&gt; 5</math> or <math>&lt; 5</math> beats/min. A minimum of two cycles required before a CTG was regarded as showing evidence of fetal</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			behavioural state changes.  Analysis: The CTG analysis was performed independently by two observers without knowledge of details of labour outcomes. All information were coded and SPSS were used for data analysis. Statistical comparison made using Student's t-test and chi square.		
<b>Full citation</b>  Williams,K.P., Galerneau,F., Intrapartum fetal heart rate patterns in the prediction of neonatal acidemia, American Journal of Obstetrics and	<b>Sample size</b>  n = 488 fetuses  <b>Characteristics</b>  Not specified  <b>Inclusion criteria</b>	<b>Interventions</b>  Fetal heart rate patterns	<b>Details</b>  Study population consisted of n = 488 women who had continuous electronic fetal monitoring during labor for the last 2	<b>Results</b>  Women with normal variability and accelerations, even in the presence of late decelerations or variable decelerations, maintained an umbilical artery pH 7.0 or greater in more than 97% of cases. In the presence of minimal/absent variability (amplitude < 5) for at least an hour, the incidence of significant acidemia (pH < 7.0) ranged from (12%-31%):	<b>Limitations</b>  <b>Other information</b>  Fetal Heart rate traces were assessed based on the National Institute of Child Health and Human Development guidelines for FHR monitoring  Neonatal acidosis defined as a pH of less than 7.0 at birth



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Gynecology, 188, 820-823, 2003</p> <p><b>Ref Id</b> 174581</p> <p><b>Country/ies where the study was carried out</b> Canada</p> <p><b>Study type</b> Cohort</p> <p><b>Aim of the study</b> To correlate changes in the intrapartum electronic fetal heart rate patterns with the development of significant neonatal acidemia.</p> <p><b>Study dates</b> January 1997 to January 2000</p>	<p>Term pregnancy (&gt; 37 weeks)</p> <p>Birth of neonates within 30 minutes of the bradycardia</p> <p>Continuous electronic fetal monitoring for 2 hours before the delivery</p> <p>Umbilical cord artery and cord blood gases done at birth</p> <p><b>Exclusion criteria</b> Fetal anomaly Multiple gestation</p>		<p>hours.</p> <p>Umbilical artery cord gas analysis performed at birth. One investigator blinded to the cord gas outcome reviewed all 488 tracings using the National Institute of Child Health and Human Development guidelines for fetal heart rate monitoring. The women were placed in six groups, depending on the absence or presence of normal variability (amplitude &gt; 5 beats) during the last hour of monitoring combined with the absence of decelerations or the presence of variable or late decelerations.</p>	<p><b><u>Outcome variable corelated with different intrapartum electronic fetal monitoring parameters</u></b></p> <p><u>Group 1 (normal variability) n = 42</u> Umbilical artery pH (mean ± SD) 7.24 ± 0.07 Base deficit (mean ± SD) 3.62 ± 3.16 Incidence of pH &lt; 7.0: 0% (p &lt; 0.05 vs. group 1, 2, 3) Incidence of pH &lt; 7.1: 9.5% Incidence of base deficit &lt; 16: 0% Incidence of base deficit &lt; 12: 2.4%</p> <p><u>Group 2 (normal variability and late decelerations) n = 173</u> Umbilical artery pH (mean ± SD) 7.18 ± 0.07 Base deficit (mean ± SD) -6.17 ± 3.14 Incidence of pH &lt; 7.0: 1.7% Incidence of pH &lt; 7.1: 13.3% Incidence of base deficit &lt; 16: 0% Incidence of base deficit &lt; 12: 4.6%</p> <p><u>Group 3 (normal variability and and variable decelerations) n = 219</u> Umbilical artery pH (mean ± SD) 7.18 ± 0.08 Base deficit (mean ± SD) -6.24 ± 3.6 Incidence of pH &lt; 7.0: 23% Incidence of pH &lt; 7.1: 9.1% Incidence of base deficit &lt; 16: 0.91% Incidence of base deficit &lt; 12: 5.5%</p> <p><u>Group 4 (decreased variability) n = 13</u> Umbilical artery pH (mean ± SD) 7.07 ± 0.2 Base deficit (mean ± SD) -9.8 ± 7.7 (p &lt; 0.05 vs. group 4 and 5) Incidence of pH &lt; 7.0: 31% (p &lt; 0.05 vs. group 1, 2, 3 and 6) Incidence of pH &lt; 7.1: 38.5% (p &lt; 0.05 group 1, 2, 3 and 6) Incidence of base deficit &lt; 16: 23.1% (p &lt;</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p>			<p>The relationship between changes in variability and the outcome variables of pH and base deficit in the six groups was assessed with analysis of variance and Chi Square test. Significance was set at the P &lt; 0.05 level.</p>	<p>0.05 group 1, 2, 3 and 6)                      Incidence of base deficit &lt; 12: 38.5% (p &lt; 0.05 group 1, 2, 3 and 6)</p> <p><u>Group 5 (decreased variability and late deceleration) n = 25</u>                      Umbilical artery pH (mean ± SD) 7.01 ± 0.14                      Base deficit (mean ± SD) -9.58 ± 6.14 (p &lt; 0.05 vs. group 4 and 5)                      Incidence of pH &lt; 7.0: 24% (p &lt; 0.05 vs. group 1, 2, 3 and 6)                      Incidence of pH &lt; 7.1: 44% (p &lt; 0.05 group 1, 2, 3 and 6)                      Incidence of base deficit &lt; 16: 24% (p &lt; 0.05 group 1, 2, 3 and 6)                      Incidence of base deficit &lt; 12: 32% (p &lt; 0.05 group 1, 2, 3 and 6)</p> <p><u>Group 6 (decreased variability and variable decelerations) n = 16</u>                      Umbilical artery pH (mean ± SD) 7.19 ± 0.14 (p &lt; 0.05 vs. group 2, 3, 4 and 5)                      Base deficit (mean ± SD) 3.37 ± 5.07                      Incidence of pH &lt; 7.0: 12.5%                      Incidence of pH &lt; 7.1: 18.8%                      Incidence of base deficit &lt; 16: 12.5%                      Incidence of base deficit &lt; 12: 12.5%</p> <p><b><u>Umbilical artery blood gas value in the absence of accelerations</u></b>  <u>Group 4 n = 8</u>                      Umbilical artery pH (mean ± SD) 6.97 ± 0.17                      Base deficit (mean ± SD) -13.06 ± 7.07                      Incidence of pH &lt; 7.0: 62.5%                      Incidence of pH &lt; 7.1: 62.5%                      Incidence of base deficit &lt; 16: 37.5%                      Incidence of base deficit &lt; 12: 62.5%</p> <p><u>Group 5 n = 19</u>                      Umbilical artery pH (mean ± SD) 7.01 ± 0.13</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>Base deficit (mean ± SD) -13.15 ± 6.64                      Incidence of pH &lt; 7.0: 31.6%                      Incidence of pH &lt; 7.1: 52.6%                      Incidence of based deficit &lt; 16: 26.3%                      Incidence of based deficit &lt; 12: 42.1%</p> <p><u>Group 6 n = 8</u>                      Umbilical artery pH (mean ± SD) 7.08 ± 0.2                      Base deficit (mean ± SD) -9.95 ± 6.25                      Incidence of pH &lt; 7.0: 25%                      Incidence of pH &lt; 7.1: 37.5%                      Incidence of base deficit &lt; 16: 25%                      Incidence of base deficit &lt; 12: 25%</p>	
<p><b>Full citation</b></p> <p>Williams,K.P.,                      Galerneau,F.,                      Fetal heart rate                      parameters                      predictive of                      neonatal                      outcome in the                      presence of a                      prolonged                      deceleration,                      Obstetrics and                      Gynecology, 100,                      951-954, 2002</p> <p><b>Ref Id</b></p> <p>174549</p> <p><b>Country/ies                      where the study                      was carried out</b></p>	<p><b>Sample size</b></p> <p>n = 186 women</p> <p><b>Characteristics</b></p> <p>Not specified</p> <p><b>Inclusion criteria</b></p> <p>Term pregnancy (&gt; 37 weeks)</p> <p>An identified prolonged deceleration/bradycardia for &gt; 2 minutes with fall &lt; 100 bpm</p> <p>Birth of neonates within 30 minutes of the bradycardia</p> <p>Continous electronic</p>	<p><b>Interventions</b></p> <p>Fetal heart rate tracing</p>	<p><b>Details</b></p> <p>Study's population consisted of n = 186 women with term gestations who had continuous electronic fetal monitoring for at least 2 hours before delivery, with an identified bradycardia during that period. Each woman had umbilical artery cord analysis done and delivery within 30</p>	<p><b>Results</b></p> <p><u><b>Outcome variable correlated with different intrapartum electronic fetal monitoring parameters</b></u></p> <p><u>Group 1 (normal variability and recovery) n = 128</u>                      Umbilical artery pH (mean ± SD) 7.17 ± 0.09                      Base deficit (mean ± SD) -6.54 ± 3.9                      Incidence of pH &lt; 7.0: 2% (p &lt; 0.05 vs. group 2 and 3)                      Incidence of pH &lt; 7.1: 22%                      Incidence of pH &lt; 7.0: 1%                      Incidence of pH &lt; 7.0: 5%                      P &lt; 0.001</p> <p><u>Group 2 (normal variability and no recovery) n = 40</u>                      Umbilical artery pH (mean ± SD) 7.13 ± 0.15                      Base deficit (mean ± SD) -7.15 ± 5.1                      Incidence of pH &lt; 7.0: 18%                      Incidence of pH &lt; 7.1: 33%                      Incidence of pH &lt; 7.0: 8%                      Incidence of pH &lt; 7.0: 13%</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>Fetal heart rate traces were assessed based on the National Institute of Child Health and Human Development guidelines for FHR monitoring</p> <p>Neonatal acidosis defined as a pH of less than 7.0 at birth</p> <p>Prolonged deceleration/bradycardia: &gt; 2 minutes with a fall to &lt; 100 bpm</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Canada</p> <p><b>Study type</b></p> <p>Cohort</p> <p><b>Aim of the study</b></p> <p>To correlate the presence of baseline variability and the duration of a prolonged deceleration/bradycardia in intrapartum fetal heart rate (FHR) tracings with the development of neonatal acidemia</p> <p><b>Study dates</b></p> <p>January 1997 to January 2000</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>fetal monitoring (EFM) for 2 hours before the delivery</p> <p>Umbilical cord artery and cord blood gases done at birth</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>minutes of that bradycardia. The last hour of all electronic monitoring tracings was reviewed by one investigator blinded to the cord gas outcome reviewed using the National Institute of Child Health and Human Development guidelines for FHR monitoring. The presence or absence of variability before the bradycardia and recovery of the bradycardia were assessed and women were categorised into four groups. Group 1 (n = 128 women) with normal variability and</p>	<p>P &lt; 0.001</p> <p><u>Group 3 (decreased variability and recovery) n = 9</u></p> <p>Umbilical artery pH (mean ± SD) 7.11 ± 0.11</p> <p>Base deficit (mean ± SD) -10.32 ± 3.68</p> <p>Incidence of pH &lt; 7.0: 44%</p> <p>Incidence of pH &lt; 7.1: 56%</p> <p>Incidence of pH &lt; 7.0: 11.1%</p> <p>Incidence of pH &lt; 7.0: 22%</p> <p>P &lt; 0.001</p> <p><u>Group 4 (decreased variability and no recovery) n = 9</u></p> <p>Umbilical artery pH (mean ± SD) 6.83 ± 0.16 (p &lt; 0.05 vs. group 1,2,3)</p> <p>Base deficit (mean ± SD) -20.17. ± 6.0 (p &lt; 0.05 vs. group 1,2,3)</p> <p>Incidence of pH &lt; 7.0: 78% (p &lt; 0.05 vs. group 1 and 2)</p> <p>Incidence of pH &lt; 7.1: 89% (p &lt; 0.05 vs. group 1)</p> <p>Incidence of pH &lt; 7.0: 78% (p &lt; 0.05 vs. group 1 and 2)</p> <p>Incidence of pH &lt; 7.0: 89% (p &lt; 0.05 vs. group 1 and 2)</p> <p>P &lt; 0.001</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>recovery before 10 minutes , group 2 (n = 40 women) with normal variability and no recovery within 10 minutes, group 3 (n = 9 women) with decreased variability and recovery within 10 minutes, and group 4 (n = 9 women) with decreased variability and no recovery within 10 minutes. Two cutoffs were used to define abnormal pH; a pH &lt; 7.0 and a pH &lt; 7.1. Two cutoffs were also used for base deficit, a base deficit &gt; -16 and a base deficit &gt; -12.</p> <p>Analysis Analysis of variance and the chi<sup>2</sup> test were used</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>to assess the relationship between the various groups. A multiple logistic regression model was developed with the parameters of amplitude and recovery used to predict pH at birth.</p>		
<p><b>Full citation</b></p> <p>Williams, K.P., Galerneau, F., Comparison of intrapartum fetal heart rate tracings in patients with neonatal seizures vs. no seizures: what are the differences?, Journal of Perinatal Medicine, 32, 422-425, 2004</p> <p><b>Ref Id</b></p> <p>121348</p>	<p><b>Sample size</b></p> <p>Seizure n = 25</p> <p>No seizure (controls) n = 25</p> <p><b>Characteristics</b></p> <p>There were no significant differences observed between the seizure and no seizure group in maternal age (32 ± 5 vs 34 ± 3), gravidity (2 ± 1 vs 2 ± 2), gestational age (39 ± 2 vs 38 ± 3) and neonatal birth weight.</p> <p><b>Inclusion criteria</b></p>	<p><b>Interventions</b></p> <p>Fetal heart rate parameters</p>	<p><b>Details</b></p> <p>The neonatal and antenatal records of the women who fit the inclusion criteria were reviewed. The cases with confirmed diagnoses of HIE (based on the clinical criteria and neuro-imaging) and cord pH &lt; 0.7 were chosen for the study. The intrapartum fetal heart rate tracings of</p>	<p><b>Results</b></p> <p><b><u>Incidence of fetal heart rate parameters (seizure n = 25, no seizure n = 25)</u></b></p> <p><u>Bradycardia</u></p> <p>Seizure n = 14 (56%)</p> <p>No seizure n = 21 (84%)</p> <p>Odds ratio 0.24 (0.06 to 0.92)</p> <p>p = 0.062</p> <p><u>Variable deceleration</u></p> <p>Seizure n = 9 (36%)</p> <p>No seizure = 15 (50%)</p> <p>Odds ratio 0.38 (0.12 to 1.18)</p> <p>p = 0.156</p> <p><u>Late decelerations</u></p> <p>Seizure n = 8 (32%)</p> <p>No seizure n = 13 (52%)</p> <p>Odds ratio 0.43 (0.14 to 1.37)</p> <p>p = 0.256</p> <p><u>Minimal/absent variability</u></p>	<p><b>Limitations</b></p> <p>Exclusion criteria not specified</p> <p>No definitions for all FHR features and abnormal FHR given</p> <p><b>Other information</b></p> <p>The tracing was reviewed in two 1 hour segments according to NICHD classification</p> <p>Minimal baseline variability: amplitude variation of ≤ 5 bpm</p> <p>Absent baseline variability: no amplitude variation</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p>Case control</p> <p><b>Aim of the study</b></p> <p>To examine which intrapartum fetal heart rate parameters in the presence of severe neonatal acidosis (pH &lt; 7.0) appropriately predicts the development of neonatal seizures in the context of hypoxic ischemic encephalopathy (HIE).</p> <p><b>Study dates</b></p> <p>January 1997 to January 2000</p> <p><b>Source of funding</b></p> <p>Not specified</p>	<p>Singleton pregnancy</p> <p>Term ≥ 37 weeks</p> <p>Presence of neonatal convulsions with 24 - 48 hours of birth secondary to hypoxic ischemic encephalopathy</p> <p><b>Exclusion criteria</b></p> <p>Not specified</p>		<p>neonates who developed neonatal seizures secondary to HIE were compared with matched neonates with similar pH (pH &lt; 0.7) and gestational age (&gt; 37) who did not develop seizures. All women had at least 2 hours of intrapartum fetal heart rate patterns available for review. The fetal heart rate parameters (prolonged deceleration, variable and late decelerations, variability, accelerations, fetal heart rate baseline and duration of the fetal heart rate abnormality) were reviewed.</p> <p><u>Analysis</u></p> <p>Comparison</p>	<p>Seizure n = 16 (64%) No seizure n = 9 (36%) Odds ratio 3.16 (1 to 10.03) p = 0.080</p> <p><u>Accelerations</u></p> <p>Seizure n = 6 (24%) No seizure = 12 (36%) Odds ratio 0.34 (0.10 to 1.15) p = 0.140</p> <p><u>Duration of abnormal FHR(min)</u></p> <p>Seizure 72 ± 12 No seizure 36 ± 18 p &lt; 0.001</p> <p><u>Baseline FHR (beats/min)</u></p> <p>Seizure 143 ± 11 No seizure 146 ± 16 p = 0.444</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			between the groups was done using chi-square and Fisher's exact test for nominal data, and Student's t-test for continuous data.		



## G.5 Care in labour as a result of cardiotocography

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Clark, S. L., Meyers, J. A., Frye, D. K., Garthwaite, T., Lee, A. J., Perlin, J. B., Recognition and response to electronic fetal heart rate patterns: impact on newborn outcomes and primary cesarean delivery rate in women undergoing induction of labor, American Journal of Obstetrics &amp; Gynecology, 212, 494.e1-6, 2015</p> <p><b>Ref Id</b></p> <p>391386</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 14398 charts reviewed in total</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion criteria</b></p> <p>Singleton, term (<math>\geq 37</math> weeks) pregnancies, undergoing induction of labour with oxytocin</p> <p><b>Exclusion criteria</b></p> <p>None reported</p>	<p><b>Interventions</b></p> <p>The protocol for intervention advocated a reduction in the dose of oxytocin according to the fetal heart rate pattern, or according to features of the uterine contractions. <u>Safety checks for fetal heart rate pattern</u></p> <p>In any 30 minute segment of CTG there should be:</p> <ul style="list-style-type: none"> <li>-at least one acceleration of 15 bpm for 15 seconds, or adequate variability present for at least 10 minutes</li> <li>-no more than one late deceleration</li> <li>-no more than 2 variable decelerations exceeding 60 seconds in duration and decreasing for more than 60 bpm</li> </ul> <p><u>Safety checks for uterine contractions:</u></p> <p>In any 30 minute segment of CTG there should be:</p> <ul style="list-style-type: none"> <li>-no more than 5 contractions in 10 minutes, for any 20 minute interval</li> <li>-no two contractions exceeding 120 seconds in duration</li> </ul>	<p><b>Details</b></p> <p>Chart reviews were conducted for all pregnancies which met the inclusion criteria. Each chart was examined by a regional nurse who was certified as a fetal heart rate monitor instructor by the Association of Women's Health, Obstetric and Neonatal Nurses. Each 30 minute section of CTG recorded during the infusion of oxytocin was examined for specific features (as described above). For every segment in which these reassuring features were not present, the chart was reviewed to assess whether oxytocin had been reduced or not. The chart</p>	<p><b>Results</b></p> <p><u>In the traces with non-reassuring fetal heart rate features:</u></p> <p><b>NICU admission</b></p> <p>Group in whom oxytocin was decreased, n/N: 91/2354 (3.8%)</p> <p>Group in whom oxytocin was not decreased, n/N: 276/5272 (5.2%)</p> <p>RR 0.74 (95% CI 0.58-0.93)</p> <p><b>Primary caesarean section</b></p> <p>Group in whom oxytocin was decreased, n/N: 630/2364 (26.6%)</p> <p>Group in whom oxytocin was not decreased, n/N: 923/5272 (17.5%)</p> <p>RR 1.52 (95% CI 1.39-1.66)</p> <p>Risk ratios (RRs) calculated by the NGA technical team using Review Manager version 5.3.</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>NICE 2012 guidelines manual checklist for cohort studies</p> <p><b>A. Selection bias</b></p> <p>A1 The method of allocation to treatment groups was unrelated to potential confounding factors: Unclear</p> <p>- Although participants were not 'allocated' to treatment groups, they were assigned to the groups retrospectively based on the interpretation of CTGs and data extraction from case notes. It is not clear whether those responsible for allocating CTGs to the two groups were aware of the neonatal outcome at the time of</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>USA</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To examine the clinical impact of specific fetal monitoring related procedures during induced labour</p> <p><b>Study dates</b></p> <p>April to September 2013</p> <p><b>Source of funding</b></p> <p>Not reported</p>		<p>-the uterus should palpate as soft between contractions</p> <p>-if an intrauterine pressure catheter is in place, the Montevideo units must calculate less than 300 mmHg and the baseline resting tone must be &lt; 25mmHg</p>	<p>was regarded as compliant if the oxytocin infusion had been reduced. The chart was regarded as non-compliant if the dose of oxytocin was not reduced, despite the absence of reassuring features. Charts had to be compliant throughout the entire duration of oxytocin infusion. The proportion of babies with adverse outcome (NICU admission; 1 minute Apgar score of &lt; 7; 5 minute Apgar score of &lt; 7 or primary caesarean section) was compared in the groups in whom oxytocin was reduced appropriately, to those in whom the oxytocin had not been reduced despite non-</p>		<p>allocation. This could affect how cases were allocated as compliant or non-compliant</p> <p>A2 Attempts were made within the design or analysis to balance the comparison groups for potential confounders: No - RRs were calculated by the NGA technical team based on the n/N provided by the study, therefore, the RRs are unadjusted for potential confounding factors and can cause high risk of bias</p> <p>A3 The groups were comparable at baseline, including all major confounding and prognostic factors: Unclear - no baseline characteristics reported</p> <p>Based on your answers to the above, in your opinion was selection bias</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			reassuring CTG features		<p>present? If so, what is the likely direction of its effect?                      High risk of bias  <b>B. Performance bias</b>                      B1 The comparison groups received the same care apart from the intervention(s) studied: Unclear                      B2 Participants receiving care were kept 'blind' to treatment allocation: n/a                      B3 Individuals administering care were kept 'blind' to treatment allocation: n/a                      Based on your answers to the above, in your opinion was performance bias present? If so, what is the likely direction of its effect?                      Unclear or unknown risk  <b>C. Attrition bias</b>                      C1 All groups were followed up for an equal length of time (or analysis was adjusted to allow</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>for differences in length of follow-up)                      Yes                      C2 a. How many participants did not complete treatment in each group? n/a                      b. The groups were comparable for treatment completion: n/a                      C3 a. For how many participants in each group were no outcome data available?                      - No Apgar data for 12 participants in the compliant group, and 18 in the non-compliant group (with regard to fetal heart rate)                      - No Apgar data for 3 participants in the compliant group, and 9 in the non-compliant group (with regard to contractions).                      b. The groups were comparable with respect to the availability of outcome data: Yes                      Based on your answers to the above, in your opinion was</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>attrition bias present? If so, what is the likely direction of its effect?                      Low risk of bias  <b>D. Detection bias</b>                      D1 The study had an appropriate length of follow-up: Yes                      D2 The study used a precise definition of outcome: Yes                      D3 A valid and reliable method was used to determine the outcome: Yes                      D4 Investigators were kept 'blind' to participants' exposure to the intervention: No                      D5 Investigators were kept 'blind' to other important confounding and prognostic factors: No                      Based on your answers to the above, in your opinion was detection bias present? If so, what is the likely direction of its effect?                      Low risk of bias</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments																												
<p><b>Full citation</b></p> <p>Lowe, B., Beckmann, M., Involving the consultant before fetal blood sampling, Australian &amp; New Zealand Journal of Obstetrics &amp; Gynaecology, 14, 14, 2016</p> <p><b>Ref Id</b></p> <p>458053</p> <p><b>Country/ies where the study was carried out</b></p> <p>Australia</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To compare neonatal outcomes</p>	<p><b>Sample size</b></p> <p>N = 4712 n = 2225 births prior to the new protocol being implemented n = 2487 births after the protocol was implemented</p> <p><b>Characteristics</b></p> <table border="1"> <thead> <tr> <th>Characteristic</th> <th>Before protocol introduction</th> <th>After protocol introduction</th> <th>p value</th> </tr> </thead> <tbody> <tr> <td>Maternal age, mean (SD)</td> <td>29.4 (5.6)</td> <td>29.6 (5.4)</td> <td>0.18</td> </tr> <tr> <td>BMI, median (IQR)</td> <td>23.1 (20.3, 27.0)</td> <td>23.1 (20.4, 26.9)</td> <td>0.56</td> </tr> <tr> <td>Nulliparity, n (%)</td> <td>1287 (57.8)</td> <td>1440 (57.9)</td> <td>0.97</td> </tr> <tr> <td>Gestational age at birth, mean (SD)</td> <td>39.5 (1.2)</td> <td>39.4 (1.2)</td> <td>0.08</td> </tr> <tr> <td>Birthweight (g), mean (SD)</td> <td>3497 (489)</td> <td>3479 (494)</td> <td>0.22</td> </tr> <tr> <td>Induction of labour, n (%)</td> <td>964 (43.3)</td> <td>1100 (44.2)</td> <td>0.53</td> </tr> </tbody> </table>	Characteristic	Before protocol introduction	After protocol introduction	p value	Maternal age, mean (SD)	29.4 (5.6)	29.6 (5.4)	0.18	BMI, median (IQR)	23.1 (20.3, 27.0)	23.1 (20.4, 26.9)	0.56	Nulliparity, n (%)	1287 (57.8)	1440 (57.9)	0.97	Gestational age at birth, mean (SD)	39.5 (1.2)	39.4 (1.2)	0.08	Birthweight (g), mean (SD)	3497 (489)	3479 (494)	0.22	Induction of labour, n (%)	964 (43.3)	1100 (44.2)	0.53	<p><b>Interventions</b></p> <p>A new hospital protocol was instigated whereby CTGs had to be reviewed by a consultant prior to a fetal blood sample being collected</p>	<p><b>Details</b></p> <p>Prior to the new protocol, CTGs were not routinely reviewed by a consultant before a fetal blood sample was collected. After implementing the new protocol, all CTGs were reviewed remotely by a consultant prior to the decision to collect a fetal blood sample. The criterion for fetal blood sampling was a pathological CTG</p>	<p><b>Results</b></p> <p><b>Fetal blood samples performed</b> Before protocol, n/N (%): 79/2225 (3.6) After protocol implemented, n/N (%): 43/2487 (1.7) RR 0.49 (95% CI 0.34-0.70)</p> <p><b>Acidosis (pH &lt;7.1)</b> Before protocol, n/N (%): 49/2225 (2.2) After protocol implemented, n/N (%): 20/2487 (0.8) RR 0.37 (95% CI 0.22-0.61)</p> <p><b>Admission to NICU</b> Before protocol, n/N (%): 98/2225 (4.4) After protocol implemented, n/N (%): 106/2487 (4.3) RR 0.97 (95% CI 0.74-1.27)</p> <p><b>Emergency caesarean section</b> Before protocol, n/N (%): 537/2225 (24.1) After protocol implemented, n/N (%): 559/2487 (22.5)</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p>NICE 2012 guidelines manual checklist for cohort studies</p> <p><b>A. Selection bias</b></p> <p>A1 The method of allocation to treatment groups was unrelated to potential confounding factors: No - The two separate groups comprised women giving birth during different time periods, therefore there are potentially confounders as well as the change in protocol that the study aimed to assess</p> <p>A2 Attempts were made within the design or analysis to balance the comparison groups for potential confounders: No</p>
Characteristic	Before protocol introduction	After protocol introduction	p value																														
Maternal age, mean (SD)	29.4 (5.6)	29.6 (5.4)	0.18																														
BMI, median (IQR)	23.1 (20.3, 27.0)	23.1 (20.4, 26.9)	0.56																														
Nulliparity, n (%)	1287 (57.8)	1440 (57.9)	0.97																														
Gestational age at birth, mean (SD)	39.5 (1.2)	39.4 (1.2)	0.08																														
Birthweight (g), mean (SD)	3497 (489)	3479 (494)	0.22																														
Induction of labour, n (%)	964 (43.3)	1100 (44.2)	0.53																														

Study details	Participants				Interventions	Methods	Outcomes and Results	Comments
following a change in hospital policy to consultant review of all CTG traces prior to collection of a fetal blood sample (FBS)	Oxytocic augmentation, n (%)	550 (24.7)	531 (21.3)	0.01			RR 0.93 (95% CI 0.84-1.03)	- Multiple variable analysis was done on only one outcome, otherwise ORs/RRs not reported and were calculated by the NGA technical using n/N reported. Therefore, most results are presenting unadjusted RRs and can be subject to bias since no adjustments for possible confounding variables were made A3 The groups were comparable at baseline, including all major confounding and prognostic factors: No - The majority of characteristics were not significantly different between the two groups. However, there was a significant reduction in the use of oxytocin during the second time period, which could affect the possible
	Epidural, n (%)	1106 (49.7)	1262 (50.7)	0.48			<b>Instrumental birth</b> Before protocol, n/N (%): 445/2225 (20) After protocol implemented, n/N (%): 439/2487 (17.6) RR 0.88 (95% CI 0.78-0.99)	
	FBS performed, n (%)	79 (3.5)	43 (1.7)	<0.01			<b>Emergency caesarean section due to fetal distress</b> Before protocol, n/N (%): 181/2225 (8.1) After protocol implemented, n/N (%): 165/2487 (6.6) RR 0.82 (95% CI 0.67-1.00)	
	Cord gas completed, n (%)	1006 (45.2)	1112 (44.7)	0.73			<b>Emergency caesarean section due to failure to progress</b> Before protocol, n/N (%): 230/2225 (10.3) After protocol implemented, n/N (%): 253/2487 (10.2) RR 0.98 (95% CI 0.83-1.17)	
<b>Study dates</b> Period 1: 1st May 2011 to 30th April 2012 Period 2 (following implementation of the new protocol): 1st May 2012 to 30th April 2013	<b>Inclusion criteria</b> All publically funded women with continuous CTG during labour, who gave birth during the study period							
<b>Source of funding</b> None reported	<b>Exclusion criteria</b> Preterm birth (< 37 weeks), fetal death in utero, known congenital abnormalities and multiple pregnancy							
							<b>Emergency caesarean section due to other reasons</b> Before protocol, n/N (%): 126/2225 (5.7)	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>After protocol implemented, n/N (%): 141/2487 (5.7) RR 1.00 (95% CI 0.79-1.26)</p> <p><b>Normal vaginal birth</b> Before protocol, n/N (%): 1231/2225 (55.3) After protocol implemented, n/N (%): 1460/2487 (58.7) RR 1.06 (95% CI 1.01-1.12)</p> <p><b>Fetal scalp lactate &gt; 4.8 mmol/l</b> Before protocol, n/N (%): 56/2225 (2.5) After protocol implemented, n/N (%): 36/2487 (1.4) RR 0.58 (95% CI 0.38-0.87)</p> <p>Risk ratios calculated by the NGA technical team using Review Manager version 5.3</p>	<p>need for FBS, as well as potentially affecting neonatal outcome Based on your answers to the above, in your opinion was selection bias present? If so, what is the likely direction of its effect? High risk of bias - potential confounders should be accounted for in the analysis</p> <p><b>B. Performance bias</b> B1 The comparison groups received the same care apart from the intervention(s) studied: Unclear - As above, the different time periods mean that care may have changed in other ways for the later group B2 Participants receiving care were kept 'blind' to treatment allocation: n/a</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>B3 Individuals administering care were kept 'blind' to treatment allocation: No                      Based on your answers to the above, in your opinion was performance bias present? If so, what is the likely direction of its effect?                      Unclear or unknown risk</p> <p><b>C. Attrition bias</b>                      C1 All groups were followed up for an equal length of time (or analysis was adjusted to allow for differences in length of follow-up):                      Yes                      C2 a. How many participants did not complete treatment in each group? n/a                      b. The groups were comparable for treatment completion: n/a                      C3 a. For how many participants in each group were no outcome data available? None reported</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>b. The groups were comparable with respect to the availability of outcome data: Yes                      Based on your answers to the above, in your opinion was attrition bias present? If so, what is the likely direction of its effect?                      Low risk of bias</p> <p><b>D. Detection bias</b>                      D1 The study had an appropriate length of follow-up: Yes                      D2 The study used a precise definition of outcome: Yes                      D3 A valid and reliable method was used to determine the outcome: Yes                      D4 Investigators were kept 'blind' to participants' exposure to the intervention: No                      D5 Investigators were kept 'blind' to other important confounding and prognostic factors: No</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					Based on your answers to the above, in your opinion was detection bias present? If so, what is the likely direction of its effect? Low risk of bias
<p><b>Full citation</b></p> <p>Katsuragi, S., Parer, J. T., Noda, S., Onishi, J., Kikuchi, H., Ikeda, T., Mechanism of reduction of newborn metabolic acidemia following application of a rule-based 5-category color-coded fetal heart rate management framework, Journal of Maternal-Fetal and Neonatal Medicine, 28,</p>	<p><b>Sample size</b></p> <p>N = 3907 overall. Number of women included in each of the two groups is not clear</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion criteria</b></p> <p>All births in a single institution during the study period</p> <p><b>Exclusion criteria</b></p> <p>Delivery by planned caesarean section</p>	<p><b>Interventions</b></p> <p>A 6 month training period was undertaken, during which time members of staff were trained in a new CTG management system. This was based on the NICHD categorisation and rule management system. CTGs were categorised into five colour coded tiers (with increasing severity: green, blue, yellow, orange and red) using 134 different combinations of variability, heart rate and graded decelerations. Each colour coded level had corresponding suggested interventions (ranging from patient positioning to immediate birth). The colour framework provides decision support only,</p>	<p><b>Details</b></p> <p>CTGs showing variable decelerations during the 10 minutes before birth were chosen for further analysis. The acid-base status of these neonates was compared before and after the training programme</p>	<p><b>Results</b></p> <p><b>Acidosis (pH &lt;7.15)</b> Before training, n/N (%): 11/688 (1.6) After training, n/N (%): 2/744 (0.2) RR 0.17 (95% CI 0.04-0.76)</p> <p><b>Acidosis (BE &lt; -12 mmol/l)</b> Before training, n/N (%): 11/688 (1.6) After training, n/N (%): 2/744 (0.2) RR 0.17 (95% CI 0.04-0.76)</p> <p>Risk ratios (RRs) calculated by the NGA technical team using Review Manager version 5.3</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><u>NICE 2012 guidelines manual checklist for cohort studies:</u> <b>A. Selection bias</b> A1 The method of allocation to treatment groups was unrelated to potential confounding factors: No – different time periods were studied A2 Attempts were made within the design or analysis to balance the comparison groups for potential confounders: No</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>1608-1613, 2015</p> <p><b>Ref Id</b></p> <p>446292</p> <p><b>Country/ies where the study was carried out</b></p> <p>Japan</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To assess neonatal outcomes before and after training with a rule-based, 5 category management system for CTG interpretation</p> <p><b>Study dates</b></p> <p>Baseline data were from 2003 to 2004. Follow</p>		<p>without dictating the decision</p> <p>All healthcare staff were trained with the new system over a 6 month period. Pre- and post-intervention assessment was not undertaken</p>			<p>RRs calculated by the NGA technical team, therefore, the RRs are unadjusted and are subject to bias because there is no adjustment for potential confounders</p> <p>A3 The groups were comparable at baseline, including all major confounding and prognostic factors: Unclear</p> <p>Based on your answers to the above, in your opinion was selection bias present? If so, what is the likely direction of its effect?</p> <p>High risk of bias</p> <p><b>B. Performance bias</b></p> <p>B1 The comparison groups received the same care apart from the intervention(s) studied: Unclear</p> <p>B2 Participants receiving care were kept 'blind' to</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>up data were from 2006 to 2007 (following a 6 month training period in 2005)</p> <p><b>Source of funding</b></p> <p>Institutional funding only</p>					<p>treatment allocation: n/a                      B3 Individuals administering care were kept 'blind' to treatment allocation: n/a                      Based on your answers to the above, in your opinion was performance bias present? If so, what is the likely direction of its effect?                      Low risk of bias</p> <p><b>C. Attrition bias</b>                      C1 All groups were followed up for an equal length of time (or analysis was adjusted to allow for differences in length of follow-up):                      Yes                      C2 a. How many participants did not complete treatment in each group? n/a                      b. The groups were comparable for treatment completion: n/a                      C3 a. For how many participants in each group were no outcome data</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>available? Not reported</p> <p>b. The groups were comparable with respect to the availability of outcome data: Unclear</p> <p>Based on your answers to the above, in your opinion was attrition bias present? If so, what is the likely direction of its effect? Unclear risk of bias</p> <p><b>D. Detection bias</b></p> <p>D1 The study had an appropriate length of follow-up: Yes</p> <p>D2 The study used a precise definition of outcome: Yes</p> <p>D3 A valid and reliable method was used to determine the outcome: Yes</p> <p>D4 Investigators were kept 'blind' to participants' exposure to the intervention: Unclear</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>D5 Investigators were kept 'blind' to other important confounding and prognostic factors: Unclear Based on your answers to the above, in your opinion was detection bias present? If so, what is the likely direction of its effect? Low risk of bias</p>

## G.6 Fetal scalp stimulation

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
<p><b>Full citation</b></p> <p>Anyaegbunam,A.M., Ditchik,A., Stoessel,R., Mikhail,M.S., Vibroacoustic stimulation of the fetus entering the second stage of labor, Obstetrics and Gynecology, 83, 963-966, 1994</p> <p><b>Ref Id</b></p> <p>202123</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To evaluate the fetal heart rate response to vibroacoustic stimulation of fetuses entering the second stage of labour as a predictor of neonatal outcome</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>July 1991 - July 1992</p>	<p><b>Sample size</b></p> <p>N = 632 Vibroacoustic stimulation (VAS) = 316 Sham stimulation = 316</p> <p><b>Characteristics</b></p> <p><u>Maternal age (years) - mean ± SD</u> VAS = 26 ± 4 Sham = 24 ± 3</p> <p><u>Nulliparous</u> VAS = 40.5% Sham = 44.6%</p> <p><u>Gestational age at delivery (weeks) - mean ± SD</u> VAS = 39 ± 1 Sham = 38 ± 2</p> <p><u>Birthweight (g) - mean ± SD</u> VAS = 3430 ± 438 Sham = 3363 ± 381</p> <p><u>Low arterial pH (&lt;7.20)</u> VAS = 5.7% Sham = 4.7%</p> <p><b>Inclusion Criteria</b></p>	<p><b>Tests</b></p> <p>5 seconds of fetal vibroacoustic stimulation</p>	<p><b>Methods</b></p> <p>Consecutive volunteers who met the study criteria were included. Women were assigned to the study or control group based on a pre-generated list of random numbers - allocation was to VAS if the next number was odd, and to sham stimulation if the number was even.</p> <p>A 5c electronic larynx (AT&amp;T, Special Needs Center, Parsippany, NJ) was placed above the symphysis on the mother's abdomen. The larynx was activated for 5 seconds, 30 seconds after a uterine contraction, and the fetal heart rate (FHR) trace was marked and the response recorded. In the sham stimulation group the artificial larynx was not activated but the FHR trace was marked in a similar fashion.</p> <p>FHR traces were interpreted by an investigator blinded to group allocation. An acceleration was defined as an increase over baseline of at least 15 bpm for at least 15 seconds. Those receiving VAS were stratified into 3 groups:</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis (umbilical) pH &lt; 7.20</u></b> 18/316 (6%)</p> <p><u>a. For umbilical cord pH &lt;7.20</u> All values calculated by NCC from data in Table 3 Sensitivity: 22.2% (3.02 to 41.43) Specificity: 77.18% (72.42 to 81.95) PPV: 5.56% (0 to 10.85) NPV: 94.26% (91.34 to 97.18) LR+: 0.97 (0.40 to 2.37) LR-: 1.01 (0.78 to 1.30)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> All values calculated by NCC from data in Table 3 Sensitivity: 30% (1.60 to 58.40) Specificity: 77.45% (72.77 to 82.13) PPV: 4.17% (0 to 8.78) NPV: 97.13% (95.04 to 99.23) LR+: 1.33 (0.50 to 3.51) LR-: 0.90 (0.60 to 1.36)</p> <p><b>Cord pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <th>Predictive Test +ve</th> <td>4</td> <td>68</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	4	68	<p><b>Limitations</b></p> <p>Only outcome data reported for those receiving the active intervention (VAS) - case series Allocation concealment unclear Period of FHR observation for qualifying acceleration following stimulus not reported Indirectness: All participants had reassuring FHR traces; unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Definition of positive stimulation test: no acceleration (selected by NCC, authors do not define positive stimulation test and do not report predictive accuracy statistics) For 2x2 table acceleration and acceleration followed by deceleration were considered a negative stimulation test result</p>
	Reference Test +ve	Reference Test -ve									
Predictive Test +ve	4	68									



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments												
<p><b>Source of funding</b></p> <p>Not reported</p>	<p>Gestational age <math>\geq 37</math> weeks, singleton fetus, reassuring heart rate patterns, cephalic presentation, absence of heavy meconium and fully dilated cervix</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>acceleration, initial acceleration followed by immediate deceleration, and no response.</p> <p>Samples of umbilical artery and vein blood were obtained at birth and tested for pH, carbon dioxide pressure, oxygen pressure and base deficit</p>	<table border="1"> <tr> <td><b>Predictive Test -ve</b></td> <td>14</td> <td>230</td> </tr> </table> <p><b>Apgar score</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>3</td> <td>69</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>7</td> <td>237</td> </tr> </tbody> </table>	<b>Predictive Test -ve</b>	14	230		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	3	69	<b>Predictive Test -ve</b>	7	237	
<b>Predictive Test -ve</b>	14	230															
	Reference Test +ve	Reference Test -ve															
<b>Predictive Test +ve</b>	3	69															
<b>Predictive Test -ve</b>	7	237															
<p><b>Full citation</b></p> <p>Arulkumaran,S., Ingemarsson,I., Ratnam,S.S., Fetal heart rate response to scalp stimulation as a test of fetal well-being in labour, Asia-Oceania Journal of Obstetrics and Gynaecology, 13, 131-135, 1987</p> <p><b>Ref Id</b></p> <p>201763</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 50</p> <p><b>Characteristics</b></p> <p>Suspicious trace = 32/50 (64%) Ominous trace = 18/50 (36%)</p> <p><b>Inclusion Criteria</b></p> <p>Women in the first stage of labour with cephalic presentation</p>	<p><b>Tests</b></p> <p>Fetal scalp stimulation for 15 seconds carried out with Allis' tissue forceps (closed to first ratchet)</p>	<p><b>Methods</b></p> <p>Fetal heart rate was monitored with a scalp electrode and the trace interpreted by two senior members of staff.</p> <p>Suspicious trace defined as: no accelerations and reduced baseline variability (5-10 bpm) or abnormal baseline rate or flat baseline (&lt; 5 bpm) or variable decelerations without ominous features. Ominous trace defined as: flat baseline and abnormal baseline rate or repeated late decelerations or repeated variable decelerations with ominous features (duration &gt; 60 seconds, beat loss &gt; 60 beats,</p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis</b> 4% (2/50)</p> <p><b>Predictive accuracy of no acceleration following fetal scalp stimulation (Allis clamp)</b></p> <p>a. For FBS pH &lt; 7.20 All values calculated by NCC from data in Table 1 Sensitivity: 100% (100 to 100) Specificity: 83.33% (72.79 to 93.88) PPV: 20% (0 to 44.79) NPV: 100% (100 to 100) LR+: 6 (3.19 to 11.30) LR-: 0 (NC)</p> <p>b. For caesarean section All values calculated by NCC from data in Table 2</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included, length of study period not reported Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: period of fetal heart rate observation for qualifying acceleration following stimulus not reported Outcome of interest is sufficiently measured in</p>												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
<p>Singapore</p> <p><b>Aim of the study</b></p> <p>To evaluate the response of the fetus to painful pinch stimulation of the scalp and its relation to fetal acid base balance when a suspicious or ominous fetal heart rate was encountered</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>slow recovery, rebound tachycardia, late deceleration component). Fetal heart rate changes were so classified if it persisted after corrective measures of alteration of position of the mother, hydration, oxygen inhalation and omission of oxytocin infusion.</p> <p>Scalp stimulation was carried out for 15 seconds when the fetal heart rate recording was at the baseline rate. The presence or absence of immediate fetal heart rate acceleration was noted. Acceleration was defined as at least 15 beats above the baseline for at least 15 seconds duration.</p> <p>Within 20 min of the test stimulation fetal blood sampling was performed with the mother in in the left lateral position. Management was according to FBS results and continued CTG trace.</p>	<p>Sensitivity: 60% (29.64 to 90.36)                      Specificity: 90% (80.70 to 99.30)                      PPV: 60% (29.64 to 90.36)                      NPV: 90% (80.70 to 99.30)                      LR+: 6 (2.08 to 17.29)                      LR-: 0.44 (0.21 to 0.96)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 549 1731 847"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>8</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>40</td> </tr> </tbody> </table> <p><b>Caesarean section</b></p> <table border="1" data-bbox="1346 959 1731 1257"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>6</td> <td>4</td> </tr> <tr> <td>Predictive Test -ve</td> <td>4</td> <td>36</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	8	Predictive Test -ve	0	40		Reference Test +ve	Reference Test -ve	Predictive Test +ve	6	4	Predictive Test -ve	4	36	<p>participants: yes                      Important potential confounders are accounted for: time between stimulation, fetal blood sample and delivery not reported                      Statistical analysis is appropriate for study design: yes                      Indirectness: unclear whether women were considered high risk</p> <p><b>Other information</b></p> <p>Authors define an acceleration as a positive stimulation test but do not report any accuracy statistics calculated using this definition. NCC calculated predictive values using no acceleration as definition of positive stimulation test, in line with other included studies.                      Two babies who had negative tests and acidotic scalp pH values had cord arterial pH values below 7.20 at birth but none had low Apgar score (&lt; 7) at 5 minutes.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	2	8																					
Predictive Test -ve	0	40																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	6	4																					
Predictive Test -ve	4	36																					

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
<p><b>Full citation</b></p> <p>Bartelsmeyer, J.A., Sadovsky, Y., Fleming, B., Petrie, R.H., Utilization of fetal heart rate acceleration following vibroacoustic stimulation in labor to predict fetal acidemia and base deficit levels, Journal of Maternal-Fetal Medicine, 4, 120-125, 1995</p> <p><b>Ref Id</b></p> <p>202115</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To evaluate if vibroacoustic stimulation can predict fetal scalp blood base deficit levels in addition to pH levels.</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>24-month period (study dates not reported)</p>	<p><b>Sample size</b></p> <p>N = 104</p> <p><b>Characteristics</b></p> <p><u>Gestational age (weeks) - mean ± SD, N</u></p> <p>15bpm x 15 sec acceleration = 38.8 ± 1.7, 52</p> <p>10bpm x 10 sec acceleration = 39.2 ± 2.3, 23</p> <p>No acceleration = 37.7 ± 3.1, 29</p> <p><u>Birth weight (g) - mean ± SD</u></p> <p>15bpm x 15 sec acceleration = 3343 ± 482, 52</p> <p>10bpm x 10 sec acceleration = 3339 ± 507, 23</p> <p>No acceleration = 2855 ± 872, 29</p> <p><b>Inclusion Criteria</b></p> <p>Women having fetal scalp blood sampling (FBS)</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>5 seconds of continuous fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>Women having FBS were studied over a 24 month period. Immediately prior to FBS fetal VAS was performed using a model 5C electronic artificial larynx (AT&amp;T Consumer Products, USA) which produces a mixed frequency sound of 81 Hz and 81 db measured at 1 m in air. A single stimulus was applied continuously for 5 seconds to the maternal abdomen one-third of the distance from the symphysis pubis to the umbilicus.</p> <p>Accelerations of the fetal heart rate (FHR) occurring within 20 seconds of VAS were recorded as a positive response. The amplitude and duration of acceleratory response was recorded and FHR traces interpreted by either of two investigators. FHR responses were classified in to three groups: FHR response of at least 15 bpm for 15 seconds, FHR response of at least 10 bpm for 10 seconds but less than 15 bpm for 15 seconds and no response.</p> <p>FHR was recorded by an internal scalp electrode. FBS</p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis</b></p> <p>14/104 (13%)</p> <p><b>Predictive value of no acceleration following VAS</b></p> <p>a. For fetal blood sample pH &lt; 7.20</p> <p>All values calculated by NCC from data in Table 4 (corresponds to sensitivity reported in text of paper)</p> <p>Sensitivity: 79% (57.08 to 100)</p> <p>Specificity: 52.22% (41.9 to 62.54)</p> <p>PPV: 20.37% (9.63 to 31.11)</p> <p>NPV: 94% (87.42 to 100)</p> <p>LR+: 1.64 (1.12 to 2.33)</p> <p>LR-: 0.41 (0.15 to 1.14)</p> <p>b. For Apgar score &lt; 7 at 5 min</p> <p>All values calculated by NCC from data in Table 2</p> <p>Sensitivity: 83.33% (53.51 to 100)</p> <p>Specificity: 52.04% (42.15 to 61.93)</p> <p>PPV: 9.62% (1.6 to 17.63)</p> <p>NPV: 98.08% (94.34 to 100)</p> <p>LR+: 1.74 (1.15 to 2.62)</p> <p>LR-: 0.32 (0.05 to 1.93)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>11</td> <td>43</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	11	43	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included</p> <p>Loss to follow-up is unrelated to key characteristics: no loss to follow up</p> <p>Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome</p> <p>Outcome of interest is sufficiently measured in participants: yes</p> <p>Important potential cofounders are accounted for: time between VAS and delivery not reported</p> <p>Statistical analysis is appropriate for study: yes</p> <p>Indirectness of population: based on gestational age mean and SD for 'no acceleration' population not all fetuses were delivered at term; unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration</p>
	Reference Test +ve	Reference Test -ve									
<b>Predictive Test +ve</b>	11	43									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments															
<p><b>Source of funding</b></p> <p>Not reported</p>			was performed immediately following VAS.	<table border="1"> <tr> <td><b>Predictive Test -ve</b></td> <td>3</td> <td>47</td> </tr> <tr> <td colspan="3"><b>Appgar score</b></td> </tr> <tr> <td></td> <td><b>Reference Test +ve</b></td> <td><b>Reference Test -ve</b></td> </tr> <tr> <td><b>Predictive Test +ve</b></td> <td>5</td> <td>47</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>1</td> <td>51</td> </tr> </table>	<b>Predictive Test -ve</b>	3	47	<b>Appgar score</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	<b>Predictive Test +ve</b>	5	47	<b>Predictive Test -ve</b>	1	51	For 2x2 table no response and FHR response of at least 10 bpm for 10 seconds but less than 15 bpm for 15 seconds were considered a positive stimulation test result
<b>Predictive Test -ve</b>	3	47																		
<b>Appgar score</b>																				
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																		
<b>Predictive Test +ve</b>	5	47																		
<b>Predictive Test -ve</b>	1	51																		
<p><b>Full citation</b></p> <p>Chauhan,S.P., Hendrix,N.W., Devoe,L.D., Scardo,J.A., Fetal acoustic stimulation in early labor and pathological fetal acidemia: a preliminary report, Journal of Maternal-Fetal Medicine, 8, 208-212, 1999</p> <p><b>Ref Id</b></p> <p>201734</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p>	<p><b>Sample size</b></p> <p>N = 271</p> <p><b>Characteristics</b></p> <p><u>Maternal age (years) - mean ± SD</u> 24.4 ± 6.0</p> <p><u>Nulliparous</u> 104/271 (82%)</p> <p><u>Mean gestational age (weeks) - mean ± SD</u> 39.1 ± 1.5</p> <p><u>Mean birth weight (g) - mean ± SD</u> 3328 ± 486</p>	3-seconds of vibroacoustic stimulation (VAS)	3-second fetal VAS was performed by placing the stimulator unit (Corometrics model 146, Wallingford, CT) over the symphysis. If no acceleration of fetal heart rate (FHR) occurred within 1 min of stimulation, additional pulses were applied at 1-min intervals with a maximum of 3 pulses. If 10 min after the third stimuli there was no acceleration (acceleration defined as an increase of 15 bpm lasting for at least 15 seconds) of FHR then the response was considered non-reactive.	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis</u></b></p> <p><u>a. pH &lt; 7.10</u> 8/271 (3.3%)</p> <p><u>b. pH &lt; 7.00</u> 4/271 (1.6%)</p> <p><b><u>Predictive value of no acceleration following VAS</u></b></p> <p><u>a. For umbilical pH &lt; 7.10</u> Values as reported in Table 2; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 44% (11.98 to 76.91) Specificity: 91% (87.79 to 94.65) PPV: 15% (1.41 to 28.21) NPV: 97.95 (96.17 to 99.73) LR+: 5.06 (2.21 to 11.59) LR-: 0.61 (0.34 to 1.09)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: not consecutive (women only included when one of the study authors was available) Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: 10-minute window for reaction to 3rd stimulus, compared with 1-min window for reaction to 1st and 2nd stimuli Outcome of interest is sufficiently measured in participants: results</p>															

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p><b>Aim of the study</b></p> <p>To determine if a non-reactive response to fetal acoustic stimulation in early labour can predict a significantly higher risk of umbilical arterial pH &lt; 7.10 or &lt; 7.00</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>6-month period (dates not reported)</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><b>Inclusion Criteria</b></p> <p>1] Singleton gestation                  2] In early active labour (cervical dilation of 5 cm or less)                  3] no contraindication to continue labour                  4] vertex presentation                  5] no narcotics                  6] umbilical arterial blood gas analysis within 30 min of delivery                  7] ≥ 37 weeks' gestational age</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>Immediately after birth a segment of umbilical cord was doubly clamped and umbilical arterial and venous blood samples were collected. Blood gas analyses were performed within 30 min of delivery.</p> <p>Caesarean delivery for fetal distress was undertaken if fetal bradycardia, late decelerations, or moderate to severe variable decelerations occurred and were unresponsive to conservative management such as changes in maternal position, hydration, supplemental oxygenation, transcervical amnioinfusion and use of tocolytics for intrauterine resuscitation. Scalp stimulation was performed prior to proceeding with urgent caesarean delivery for abnormal FHR. Scalp pH was not obtained due to nonavailability of the machine.</p> <p>Results of VAS were not used in the management of the woman's labour.</p>	<p><b>b. For umbilical pH &lt; 7.00</b></p> <p>Values as reported in Table 2; NCC calculated LR+, LR- and all confidence intervals                  Sensitivity: 50% (1 to 99)                  Specificity: 91% (87.14 to 94.13)                  PPV: 7% (0 to 17.29)                  NPV: 99.18 (98.05 to 100)                  LR+: 5.34 (1.87 to 15.24)                  LR-: 0.55 (0.21 to 1.47)</p> <p><b>c. For cesarean section</b></p> <p>Values as reported in Table 2; NCC calculated LR+, LR- and all confidence intervals                  Sensitivity: 37% (3.95 to 71.05)                  Specificity: 92% (87.39 to 94.35)                  PPV: 11% (0 to 22.97)                  NPV: 97% (96.17 to 99.73)                  LR+: 4.11 (1.55 to 10.87)                  LR-: 0.69 (0.40 to 1.18)</p> <p><b>Umbilical cord pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>4</td> <td>23</td> </tr> <tr> <td>Predictive Test -ve</td> <td>5</td> <td>239</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	4	23	Predictive Test -ve	5	239	<p>reported for pH &lt; 7.10 and &lt; 7.00 (standard definition is &lt; 7.20)                  Important potential confounders are accounted for: yes                  Statistical analysis is appropriate for study design: yes                  Indirectness: unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration</p> <p><u>Number of stimulations applied</u>                  One stimulation = 214/271 (78.9%)                  Two stimulations = 19/271 (7%)                  Three stimulations = 38/271 (14%)                  Of the 38 fetuses who received three stimulations, only 11 had an acceleration with 10 min of last VAS application (definition of response)</p> <p><u>Interval between first VAS to delivery</u>                  Full study population = 7.9 ± 6.9 hours</p>
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	4	23												
Predictive Test -ve	5	239												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<p><b>Umbilical cord pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>25</td> </tr> <tr> <td>Predictive Test -ve</td> <td>2</td> <td>242</td> </tr> </tbody> </table> <p><b>Caesarean section</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>24</td> </tr> <tr> <td>Predictive Test -ve</td> <td>5</td> <td>239</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	25	Predictive Test -ve	2	242		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	24	Predictive Test -ve	5	239	<p>Caesarean section for distress = <math>7.3 \pm 4.3</math> hours vs. No caesarean section = <math>7.9 \pm 6.9</math> hours                      Umbilical arterial pH <math>&lt; 7.10 = 7.2 \pm 6.0</math> hours vs. umbilical arterial pH <math>\geq 7.10 = 7.9 \pm 6.6</math> hours                      Umbilical arterial pH <math>&lt; 7.00 = 9.5 \pm 8.0</math> hours vs. umbilical arterial pH <math>\geq 7.00 = 8.0 \pm 6.9</math> hours</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	2	25																					
Predictive Test -ve	2	242																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	3	24																					
Predictive Test -ve	5	239																					
<p><b>Full citation</b></p> <p>Clark,S.L., Gimovsky,M.L., Miller,F.C., Fetal heart rate response to scalp blood sampling, American Journal of Obstetrics and Gynecology, 144, 706-708, 1982</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>N = 200</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion Criteria</b></p>	<p><b>Tests</b></p> <p>Endoscope placement and fetal scalp blood sampling (scalp puncture served as fetal scalp stimulation)</p>	<p><b>Methods</b></p> <p>The labour records of women who delivered at Los Angeles County/University of Southern California Women's Hospital during a 2-year period were reviewed. Intrapartum fetal heart rate tracings of 200 women who had undergone fetal scalp blood sampling were chosen sequentially.</p>	<p><b>Results</b></p> <p><b>Prevalence of FBS pH <math>&lt; 7.21</math></b>                      19/200 (10%)</p> <p><b>Predictive value of no acceleration following fetal scalp puncture for FBS pH <math>&lt; 7.21</math></b>                      All values calculated by NCC using data in Table I                      Sensitivity: 100% (100 to 100)                      Specificity: 93.37% (89.75 to 96.99)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included                      Loss to follow-up is unrelated to key characteristics: no loss to follow up                      Prognostic factor is adequately measured in</p>																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>201761</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To ascertain the correlation between fetal acid-base status and the ability of the fetus to manifest a reassuring fetal heart rate pattern in response to tactile stimulation provided by fetal blood sampling</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>A 2-year period (dates not reported)</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>Fetal heart rate tracings were reviewed blindly, without knowledge of the pH values obtained at the time of sampling. They were judged to be either reactive (demonstrating fetal heart rate acceleration of 15 bpm lasting 15 seconds) or non-reactive in response to endscope placement and scalp puncture.</p>	<p>PPV: 61.29% (44.14 to 78.44)                      NPV: 100% (100 to 100)                      LR+: 15.08 (8.73 to 26.06)                      LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>19</td> <td>12</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>0</td> <td>169</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	19	12	<b>Predictive Test -ve</b>	0	169	<p>participants: period of fetal heart rate observation for qualifying acceleration following stimulus not reported</p> <p>Outcome of interest is sufficiently measured in participants: yes</p> <p>Important potential confounders are accounted for: time between stimulation, fetal blood sampling and delivery not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: gestational age not reported - at least one woman was in pre-term labour (32 to 33 weeks' gestation); unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Definition of positive stimulation test: no acceleration (selected by NCC, authors do not define positive stimulation test and do not report predictive accuracy statistics)</p> <p>All FBS was performed during the first stage of labour.</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	19	12												
<b>Predictive Test -ve</b>	0	169												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
					<p><u>Mean (range) scalp pH</u>                      Acceleration in response to stimulation = 7.32 (7.21 to 7.42)                      No acceleration in response to stimulation = 7.16 (6.95 to 7.31)</p>						
<p><b>Full citation</b>                      Clark,S.L., Gimovsky,M.L., Miller,F.C., The scalp stimulation test: a clinical alternative to fetal scalp blood sampling, American Journal of Obstetrics and Gynecology, 148, 274-277, 1984</p> <p><b>Ref Id</b>                      202086</p> <p><b>Country/ies where the study was carried out</b>                      USA</p> <p><b>Aim of the study</b>                      To compare the correlation between heart rate accelerations in response to non-invasive tactile stimulation of the fetal scalp and subsequent pH obtained at scalp blood sampling</p>	<p><b>Sample size</b>                      N = 100</p> <p><b>Characteristics</b>  <u>Gestational age</u>                      Preterm (33 to 35 weeks) = 4/100 (4%)                      Term (37 to 41 weeks) = 76/100 (76%)                      Post-term (≥ 42 weeks) = 20/100 (20%)</p> <p><b>Inclusion Criteria</b>                      Fetuses with heart rate tracings indicating possible acidosis mandating scalp blood sampling</p> <p><b>Exclusion Criteria</b>                      Not reported</p>	<p><b>Tests</b>                      15 seconds of gentle digital pressure on the scalp through the dilated cervix, followed by transvaginal application on fetal scalp of Allis clamp closed to first ratchet and left in place for 15 seconds</p>	<p><b>Methods</b>                      100 fetuses with heart tracings indicating possible acidosis were prospectively enrolled by the clinical resident on the labour and delivery floor after review of the woman's clinical course and fetal heart rate (FHR) pattern.</p> <p>FHR response to each stimulation (15 seconds of gentle digital pressure followed by 15 seconds application of Allis clamp) was observed, followed by scalp blood sampling in the usual manner.</p> <p>Each tracing was reviewed by one of the authors without knowledge of the fetal scalp pH and was judged to be reactive or non-reactive to each stimulus as well as to the stimulus of the scalp puncture itself.</p>	<p><b>Results</b>  <u>Prevalence of acidosis pH &lt; 7.20</u>                      19/64 (30%)</p> <p><u>Predictive accuracy of no acceleration following fetal scalp stimulation (FSS) (Allis clamp) for FBS pH &lt; 7.20 [only in those fetuses who had not responded to initial digital FSS]</u>                      All values calculated by NCC from data presented in Fig 2                      Sensitivity: 100% (100 to 100)                      Specificity: 33.33% (19.56 to 47.11)                      PPV: 38.78% (25.13 to 52.42)                      NPV: 100% (100 to 100)                      LR+: 1.5 (1.22 to 1.84)                      LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>19</td> <td>30</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	19	30	<p><b>Limitations</b>                      Study sample represents population: unclear whether consecutive women were included                      Loss to follow-up is unrelated to key characteristics: no loss to follow up                      Prognostic factor is adequately measured in participants: period of FHR observation for qualifying acceleration following stimulus not reported                      Outcome of interest is sufficiently measured in participants: results not adequately reported digital stimulation                      Important potential confounders are accounted for: time between stimulation, FBS and delivery not reported                      Statistical analysis is appropriate for study design: yes - although data not sufficiently reported for</p>
	Reference Test +ve	Reference Test -ve									
<b>Predictive Test +ve</b>	19	30									



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments			
<p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>Reactive response was defined as an acceleration of fetal heart rate of 15 bpm lasting at least 15 seconds</p>	<table border="1" data-bbox="1346 272 1729 371"> <tr> <td data-bbox="1346 272 1464 371"><b>Predictive Test -ve</b></td> <td data-bbox="1464 272 1599 371">0</td> <td data-bbox="1599 272 1729 371">15</td> </tr> </table>	<b>Predictive Test -ve</b>	0	15	<p>digital scalp stimulation</p> <p>Indirectness of population: 76% of fetuses were delivered at term; fetuses had failed to respond to digital stimulation; unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Definition of positive stimulation test: no acceleration (selected by NCC, authors do not define positive stimulation test and do not report predictive accuracy statistics).</p> <p>2x2 table could not be calculated for digital fetal scalp stimulation. 2x2 table could be calculated for predictive accuracy of response to Allis clamp stimulation for the 64 fetuses who did not respond with an acceleration to digital stimulation.</p> <p>Data not reported for response to stimulation of scalp puncture.</p> <p>Data reported in Fig 2 (used to calculate 2x2</p>
<b>Predictive Test -ve</b>	0	15						

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					<p>table) specify percentage of fetuses with pH &lt; 7.20 and percentage of fetuses with pH &gt; 7.20. Unclear in which group fetuses with a pH of 7.20 were included.</p> <p>All women were in the first stage of labour.</p>
<p><b>Full citation</b></p> <p>Edersheim,T.G., Hutson,J.M., Druzin,M.L., Kogut,E.A., Fetal heart rate response to vibratory acoustic stimulation predicts fetal pH in labor, American Journal of Obstetrics and Gynecology, 157, 1557-1560, 1987</p> <p><b>Ref Id</b></p> <p>201764</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To examine the relationship between vibratory acoustic stimulation, direct fetal</p>	<p><b>Sample size</b></p> <p>N = 188 responses N = 127 women</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion Criteria</b></p> <p>≥ 34 weeks' gestation, active labour with ruptured membranes, and evidence of abnormal fetal heart rate tracings</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>3 seconds of fetal vibroacoustic stimulation (VAS) followed by the incision of fetal scalp blood sampling (FBS) serving as fetal scalp stimulation.</p>	<p><b>Methods</b></p> <p>FBS was performed where fetal heart rate (FHR) tracings were suspicious or equivocal. FBS was also performed with meconium plus FHR abnormality such as decreased beat-to-beat variability or fetal tachycardia.</p> <p>FHR was monitored continuously by Corometrics 112 fetal heart rate monitor. 60 seconds before FBS a single 3-second VAS was applied over the fetal vertex with the Western Electric Model 5c electronic artificial larynx.</p> <p>FHR was observed for 60 seconds and FBS was performed by standard puncture technique and analysed on a Corometrics 220 pH system. FHR response to both VAS and</p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis pH &lt; 7.20</b> 6/188 (3%) [acidotic samples, not fetuses]</p> <p><b>1. Predictive accuracy of an acceleration</b></p> <p><b>a. Following vibroacoustic stimulation for FBS pH &gt; 7.20</b> As reported in Table II and text of paper; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 63.7% (56.75 to 70.72) Specificity: 100% (100 to 100) PPV: 100% (100 to 100) NPV: 8.33% (1.95 to 14.72) LR+: NC LR-: 0.36 (0.30 to 0.44)</p> <p><b>b. Following fetal scalp stimulation for FBS pH &gt; 7.20</b> As reported in Table II and text of paper; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 43.4% (36.21 to 50.61) Specificity: 100% (100 to 100) PPV: 100 % (100 to 100)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear how many women were in preterm labour, unclear whether consecutive women were included Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome; Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between FBS and delivery not reported Statistical analysis is appropriate for study design: yes</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
<p>scalp stimulation, and fetal scalp blood pH</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>March 1985 - March 1986</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>fetal scalp stimulation was recorded and correlated with pH value obtained. An acceleration was defined as an increase in FHR above the baseline of 15bpm sustained for 15 seconds occurring within 60 seconds after either stimulation.</p>	<p>NPV: 5.5% (1.22 to 9.79) LR+: NC LR-: 0.57 (0.50 to 0.64)</p> <p><b>2. Predictive accuracy of no acceleration</b></p> <p><u>a. Following vibroacoustic stimulation for FBS pH &lt; 7.20</u> All values calculated by NCC using data presented in Table II Sensitivity: 100% (100 to 100) Specificity: 63.74% (56.75 to 70.72) PPV: 8.33% (1.95 to 14.72) NPV: 100% (100 to 100) LR+: 2.76 (2.27 to 3.24) LR-: 0 (NC)</p> <p><u>b. Following fetal scalp stimulation for FBS pH &lt; 7.20</u> All values calculated by NCC using data presented in Table II Sensitivity: 100% (100 to 100) Specificity: 43.41% (36.21 to 50.61) PPV: 5.5% (1.22 to 9.79) NPV: 100% (100 to 100) LR+: 1.77 (1.56 to 2.01) LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 1129 1729 1329"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <th>Predictive Test +ve</th> <td>116</td> <td>0</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	116	0	<p>Indirectness: unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Responses to both VAS and fetal scalp stimulation were recorded in 188 instances in 127 women</p> <p>Authors' definition of positive stimulation test: acceleration Authors' definition of positive fetal scalp test: no acidosis pH &gt; 7.20</p> <p>First set of predictive accuracy results in evidence table are as reported in the study Second set of predictive accuracy results were calculated by NCC with a recalculated 2x2 table using a definition of positive stimulation test being no acceleration and definition of positive fetal scalp test of acidosis pH &lt; 7.20, in line with other studies included in this review.</p>
	Reference Test +ve	Reference Test -ve									
Predictive Test +ve	116	0									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																														
				<table border="1"> <tr> <td data-bbox="1346 272 1464 371">Predictive Test -ve</td> <td data-bbox="1464 272 1599 371">66</td> <td data-bbox="1599 272 1733 371">6</td> </tr> <tr> <td colspan="3" data-bbox="1346 427 1733 459"><b>FBS pH</b></td> </tr> <tr> <td data-bbox="1346 483 1464 582"></td> <td data-bbox="1464 483 1599 582"><b>Reference Test +ve</b></td> <td data-bbox="1599 483 1733 582"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1346 590 1464 681">Predictive Test +ve</td> <td data-bbox="1464 590 1599 681">79</td> <td data-bbox="1599 590 1733 681">0</td> </tr> <tr> <td data-bbox="1346 689 1464 780">Predictive Test -ve</td> <td data-bbox="1464 689 1599 780">103</td> <td data-bbox="1599 689 1733 780">6</td> </tr> <tr> <td colspan="3" data-bbox="1346 836 1733 868"><b>FBS pH</b></td> </tr> <tr> <td data-bbox="1346 892 1464 991"></td> <td data-bbox="1464 892 1599 991"><b>Reference Test +ve</b></td> <td data-bbox="1599 892 1733 991"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1346 999 1464 1090">Predictive Test +ve</td> <td data-bbox="1464 999 1599 1090">6</td> <td data-bbox="1599 999 1733 1090">66</td> </tr> <tr> <td data-bbox="1346 1098 1464 1189">Predictive Test -ve</td> <td data-bbox="1464 1098 1599 1189">0</td> <td data-bbox="1599 1098 1733 1189">116</td> </tr> <tr> <td colspan="3" data-bbox="1346 1244 1733 1276"><b>FBS pH</b></td> </tr> </table>	Predictive Test -ve	66	6	<b>FBS pH</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	79	0	Predictive Test -ve	103	6	<b>FBS pH</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	6	66	Predictive Test -ve	0	116	<b>FBS pH</b>			
Predictive Test -ve	66	6																																	
<b>FBS pH</b>																																			
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																																	
Predictive Test +ve	79	0																																	
Predictive Test -ve	103	6																																	
<b>FBS pH</b>																																			
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																																	
Predictive Test +ve	6	66																																	
Predictive Test -ve	0	116																																	
<b>FBS pH</b>																																			

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
				<table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>6</td> <td>103</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>79</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	6	103	Predictive Test -ve	0	79	
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	6	103												
Predictive Test -ve	0	79												
<p><b>Full citation</b></p> <p>Elimian,A., Figueroa,R., Tejani,N., Intrapartum assessment of fetal well-being: a comparison of scalp stimulation with scalp blood pH sampling, Obstetrics and Gynecology, 89, 373-376, 1997</p> <p><b>Ref Id</b></p> <p>201856</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To determine if and to what extent the need for scalp pH sampling is decreased by the scalp stimulation test and whether redefinition of reactivity and</p>	<p><b>Sample size</b></p> <p>N = 108</p> <p><b>Characteristics</b></p> <p><u>Mean gestational age</u> 39.2 ± 1.7 weeks</p> <p><u>Mean birthweight</u> 3240 ± 579 g</p> <p><u>Mean maternal age</u> 24.2 ± 5.9 years</p> <p><u>Nulliparous</u> 73/108 (68%)</p> <p><u>Indications for FBS*</u> Moderate to severe variable decelerations = 84/108 (78%) Late decelerations = 12/108 (11%) Baseline tachycardia = 5/108 (5%) Baseline bradycardia = 3/108 (3%)</p>	<p><b>Tests</b></p> <p>15 seconds of gentle digital fetal scalp stimulation</p>	<p><b>Methods</b></p> <p>108 consecutive women were entered prospectively in to the study. The decision to perform fetal scalp blood sampling (FBS) was made by the attending senior resident in the labour and delivery suite after review of the woman's clinical course and FHR trace.</p> <p>15 seconds of digital fetal scalp stimulation was performed through the dilated cervix, followed 1 to 2 minutes later by FBS in the usual manner. Each FHR trace was marked at the time of both stimulations and judged to be reactive or non-reactive in response to both digital stimulation and scalp puncture.</p> <p>Reactive response defined as an acceleration of 15 bpm lasting at least 15 seconds. FHR reaction was then</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis pH &lt; 7.20</u></b> 15/108 (14%)</p> <p><b><u>Predictive value of no acceleration following digital fetal scalp stimulation (FSS) (first FSS intervention) for fetal blood sample pH &lt; 7.20</u></b> All values calculated by NCC from data in Table 1 (corresponds to sensitivity, specificity, PPV reported in text of paper) Sensitivity: 100% (100 to 100) Specificity: 54.84% (44.72 to 64.95) PPV: 26.32% (14.88 to 37.75) NPV: 100% (100 to 100) LR+: 2.21 (1.77 to 2.77) LR-: 0 (NC)</p> <p><b><u>Predictive value of no acceleration following scalp puncture (second FSS intervention) for fetal blood sample pH &lt; 7.20</u></b> Calculated by NCC from data in Table 1 (corresponds to sensitivity, specificity, PPV reported in text of</p>	<p><b>Limitations</b></p> <p>Study sample represents population: yes Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome; period of FHR observation for qualifying acceleration following stimulus not reported Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between stimulation, FBS and delivery not reported Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: 5% of women</p>									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
<p>presence of fetal heart rate (FHR) variability preceding scalp stimulation further decreased the need for fetal scalp blood sampling</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>January - September 1995</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>Decreased variability = 4/108 (4%)</p> <p>*percentage calculated by NCC-WCH, do not add up to 100% due to rounding up</p> <p><b>Inclusion Criteria</b></p> <p>FHR patterns, recorded by fetal scalp electrode, suggestive of possible acidosis</p> <p><b>Exclusion Criteria</b></p> <p>1] HIV positive or positive for hepatitis B surface antigen                  2] Herpes virus lesions                  3] Women in whom scalp was inaccessible for sampling</p>		<p>correlated with scalp blood pH values (using 220 pH system, Corometrics Medical Systems, Wallingford, CT, USA). Fetal acidosis defined as scalp pH &lt; 7.20</p>	<p>paper)</p> <p>Sensitivity: 100% (100 to 100)                  Specificity: 53.76% (43.63 to 63.9)                  PPV: 25.86% (14.59 to 27.13)                  NPV: 100% (100 to 100)                  LR+: 2.16 (1.73 to 2.69)                  LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>15</td> <td>42</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>51</td> </tr> </tbody> </table> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>15</td> <td>43</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>50</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	15	42	Predictive Test -ve	0	51		Reference Test +ve	Reference Test -ve	Predictive Test +ve	15	43	Predictive Test -ve	0	50	<p>were in pre-term labour (34-36 weeks); unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration.</p> <p>5/108 (4.6%) had a gestational age of 34-36 weeks.</p> <p>Where there was more than one FBS only the last sample was used for analysis.</p> <p>Variability of FHR was performed before scalp stimulation and confirmed by two of the authors blinded to scalp pH results - it is unclear whether FHR response (reactive or non-reactive) to stimulation was also assessed blindly.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	15	42																					
Predictive Test -ve	0	51																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	15	43																					
Predictive Test -ve	0	50																					
<b>Full citation</b>	<b>Sample size</b>	<b>Tests</b>	<b>Methods</b>	<b>Results</b>	<b>Limitations</b>																		
	N = 33																						

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>Ingemarsson,I., Arulkumaran,S., Reactive fetal heart rate response to vibroacoustic stimulation in fetuses with low scalp blood pH, British Journal of Obstetrics and Gynaecology, 96, 562-565, 1989</p> <p><b>Ref Id</b> 202006</p> <p><b>Country/ies where the study was carried out</b> Unclear</p> <p><b>Aim of the study</b> To describe fetal heart rate responses to vibroacoustic stimulation of the fetus in labour</p> <p><b>Study type</b></p> <p><b>Study dates</b> Not reported</p> <p><b>Source of funding</b> Not reported</p>	<p><b>Characteristics</b> Not reported</p> <p><b>Inclusion Criteria</b> Women undergoing fetal blood sampling because of suspicious or ominous fetal heart rate (FHR) traces in the first stage of labour</p> <p><b>Exclusion Criteria</b> Not reported</p>	5 seconds of fetal vibroacoustic stimulation (VAS)	<p>Women between 35 and 42 gestational weeks received fetal blood sampling (FBS). Before FBS a model 5C electronic artificial larynx (Western Electric, Bell Telephone) was applied to the maternal abdomen in the region of the fetal head for 5 seconds. A response was defined as reactive if the FHR showed an acceleration of 15 bpm for 15 seconds immediately after the sound stimulation. FBS was taken by one of the authors within 20 minutes of sound stimulation with the woman in the left lateral position. Cord artery blood was taken at caesarean section in 15 women when FBS was not possible due to high head and inadequate dilatation of the cervix. Acidosis was defined as pH &lt; 7.20</p> <p>Suspicious or ominous FHR traces showed late decelerations (intermittently or repeatedly), pronounced variable decelerations (depth &gt; 60 bpm or lasting for &gt; 60 seconds or both), tachycardia with late or variable decelerations, or reduced variability (&lt; 5 bpm lasting for &gt; 60 min) indicative of possible fetal acidosis</p>	<p><b>Prevalence of acidosis pH &lt;7.20</b> 4/51 (8%)</p> <p><b>Predictive accuracy of no acceleration following VAS</b> a. For FBS pH &lt;7.20</p> <p>All values calculated by NCC using data presented in Table 1 and 2 Sensitivity: 50% (1 to 99) Specificity: 68.97% (52.13 to 85.80) PPV: 18.18% (0 to 40.97) NPV: 90.91% (78.90 to 100) LR+: 1.61 (0.53 to 4.94) LR-: 0.73 (0.26 to 1.99)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>2</td> <td>9</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>2</td> <td>20</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	2	9	<b>Predictive Test -ve</b>	2	20	<p>Study sample represents population: unclear, characteristics not reported; unclear whether consecutive women were included</p> <p>Loss to follow-up is unrelated to key characteristics: no loss to follow up</p> <p>Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome</p> <p>Outcome of interest is sufficiently measured in participants: yes</p> <p>Important potential confounders are accounted for: time between stimulation, FBS and delivery not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: unclear whether any women were considered high risk</p> <p><b>Other information</b> Definition of positive stimulation test: no acceleration (selected by NCC, authors do not define positive stimulation test and do not report predictive accuracy statistics).</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	2	9												
<b>Predictive Test -ve</b>	2	20												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					<p>51 women were recruited in to the study but data for both stimulation test plus FBS test only reported for 33 women.</p> <p>Individual data are reported for 11 fetuses with no FHR response to VAS and no FHR response to FBS (the scalp puncture acting as the stimulus). These data were used to calculate predictive accuracy statistics for VAS (FBS pH &lt; 7.20). Results were the same for FBS and so predictive accuracy statistics for FBS (FBS pH &lt; 7.20) were not calculated.</p>
<p><b>Full citation</b></p> <p>Irion,O., Stuckelberger,P., Moutquin,J.M., Morabia,A., Extermann,P., Beguin,F., Is intrapartum vibratory acoustic stimulation a valid alternative to fetal scalp pH determination?, British Journal of Obstetrics and Gynaecology, 103, 642-647, 1996</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>N = 421 samples N = 253 women</p> <p><b>Characteristics</b></p> <p><u>Maternal age (years) - mean ± SD</u> 28.3 ± 4.4</p> <p><u>Gestational age (weeks) - mean ± SD</u> 39.1 ± 1.6</p>	<p><b>Tests</b></p> <p>5 seconds of fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>All fetal scalp blood samplings (FBS) for abnormal intrapartum fetal heart rate (FHR) tracings at &gt; 30 pregnancy weeks were consecutively included in the study.</p> <p>FHR abnormalities were the presence of at least one of the following: late decelerations, decreased baseline variability (beat-to-beat variability &lt; 5</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis</u></b> 31/421 (7.4%)</p> <p><b><u>1. Predictive accuracy of an acceleration following VAS</u></b> <b><u>a. For FBS pH &gt; 7.20</u></b> As reported in Table 3 of paper Sensitivity: 52% (47 to 57) Specificity: 77% (63 to 92) PPV: 97% (94 to 99) NPV: 11% (7 to 16) LR+: 2.29 (1.19 to 4.43) LR-: 0.62 (0.50 to 0.77)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: yes Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential</p>



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>201885</p> <p><b>Country/ies where the study was carried out</b></p> <p>Switzerland</p> <p><b>Aim of the study</b></p> <p>To determine the validity of fetal heart rate accelerations, either spontaneous or induced by vibratory acoustic stimulation, as an indicator of fetal wellbeing according to subsequent scalp pH values</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Over a 15 month period (dates not reported)</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><u>Operative delivery for fetal distress</u></p> <p>106/253 (42%)</p> <p>Forceps or vacuum extractor = 75/253 (30%)</p> <p>Caesarean section = 30/253 (12%)</p> <p>[one operative delivery not accounted for in text of study]</p> <p><b>Inclusion Criteria</b></p> <p>Abnormal intrapartum fetal heart rate tracings at &gt; 30 weeks' pregnancy</p> <p><b>Exclusion Criteria</b></p> <p>No cases were excluded</p>		<p>bpm for 20 min), severe variable decelerations, moderate or severe bradycardia (&lt; 100 bpm for 3 min), tachycardia (baseline rate &gt; 160 bpm).</p> <p>Every time FBS was deemed necessary, VAS was performed by applying a model 5C electronic artificial larynx (Western Electric, New York) to the maternal abdominal wall above the fetal vertex for 5 sec. FHR tracing was observed for at least 60 sec after VAS. FBS was performed by scalp puncture for pH determination within 5 min.</p> <p>Reactivity was defined as FHR acceleration of at least 15 bpm above the baseline level, lasting for at least 15 sec. Tracings were blindly assessed by one author for the presence of VAS-induced reactivity prior to FBS.</p>	<p><u>b. For FBS pH &gt; 7.25</u></p> <p>As reported in Table 3 of paper</p> <p>Sensitivity: 56% (51 to 62)</p> <p>Specificity: 65% (57 to 74)</p> <p>PPV: 78% (73 to 84)</p> <p>NPV: 40% (33 to 47)</p> <p>LR+: 1.63 (1.26 to 2.11)</p> <p>LR-: 0.67 (0.56 to 0.80)</p> <p><b>2. Predictive accuracy of no acceleration following VAS</b></p> <p><u>a. For FBS pH &lt; 7.20</u></p> <p>All values calculated by NCC using data presented in Table 2</p> <p>Sensitivity: 77.42% (62.70 to 92.14)</p> <p>Specificity: 51.54% (46.58 to 56.50)</p> <p>PPV: 11.27% (7.02 to 15.51)</p> <p>NPV: 96.63% (94.18 to 99.09)</p> <p>LR+: 1.60 (1.29 to 1.98)</p> <p>LR-: 0.44 (0.23 to 0.85)</p> <p><u>b. For FBS pH &lt; 7.25</u></p> <p>All values calculated by NCC using data presented in Table 2</p> <p>Sensitivity: 65.38% (57.21 to 73.56)</p> <p>Specificity: 56.01% (50.31 to 61.72)</p> <p>PPV: 39.91% (33.33 to 46.48)</p> <p>NPV: 78.37% (72.77 to 83.96)</p> <p>LR+: 1.49 (1.24 to 1.78)</p> <p>LR-: 0.62 (0.48 to 0.80)</p> <p><b>FBS pH</b></p>	<p>confounders are accounted for: time between FBS and delivery not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: unclear how many women were in preterm labour, unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Responses to both VAS and fetal scalp stimulation were recorded in 421 instances in 253 consecutive women</p> <p>Authors' definition of positive stimulation test: acceleration</p> <p>Authors' definition of positive fetal scalp test: no acidosis pH &gt; 7.20</p> <p>First set of predictive accuracy results in evidence table are as reported in the study</p> <p>Second set of predictive accuracy results were calculated by NCC with a recalculated 2x2 table using a definition of positive stimulation test being no acceleration and definition of positive fetal</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1" data-bbox="1346 272 1731 572"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>201</td> <td>7</td> </tr> <tr> <td>Predictive Test -ve</td> <td>189</td> <td>24</td> </tr> </tbody> </table> <p data-bbox="1346 632 1435 655">FBS pH</p> <table border="1" data-bbox="1346 684 1731 984"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>163</td> <td>45</td> </tr> <tr> <td>Predictive Test -ve</td> <td>128</td> <td>85</td> </tr> </tbody> </table> <p data-bbox="1346 1043 1435 1067">FBS pH</p> <table border="1" data-bbox="1346 1096 1731 1295"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>24</td> <td>189</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	201	7	Predictive Test -ve	189	24		Reference Test +ve	Reference Test -ve	Predictive Test +ve	163	45	Predictive Test -ve	128	85		Reference Test +ve	Reference Test -ve	Predictive Test +ve	24	189	<p data-bbox="1749 272 2033 384">scalp test of acidosis pH &lt; 7.20, in line with other studies included in this review.</p>
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	201	7																											
Predictive Test -ve	189	24																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	163	45																											
Predictive Test -ve	128	85																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	24	189																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments												
				<table border="1"> <tr> <td>Predictive Test -ve</td> <td>7</td> <td>201</td> </tr> </table> <p>FBS pH</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>85</td> <td>128</td> </tr> <tr> <td>Predictive Test -ve</td> <td>45</td> <td>163</td> </tr> </tbody> </table>	Predictive Test -ve	7	201		Reference Test +ve	Reference Test -ve	Predictive Test +ve	85	128	Predictive Test -ve	45	163	
Predictive Test -ve	7	201															
	Reference Test +ve	Reference Test -ve															
Predictive Test +ve	85	128															
Predictive Test -ve	45	163															
<p><b>Full citation</b></p> <p>Lazebnik,N., Neuman,M.R., Lysikiewicz,A., Dierker,L.R., Mann,L.I., Response of fetal heart rate to scalp stimulation related to fetal acid-base status, American Journal of Perinatology, 9, 228-232, 1992</p> <p><b>Ref Id</b></p> <p>202013</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 104</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion Criteria</b></p> <p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>The incision of fetal scalp blood sampling (FBS) served as fetal scalp stimulation</p>	<p><b>Methods</b></p> <p>Term fetuses during labour were studied by scalp pH. All fetuses were monitored by an internal scalp electrode and intrauterine pressure catheter. The timing of stimulation was marked on fetal heart tracings.</p> <p>Recordings of fetal heart rate (FHR) were digitised by tracing the curves on a digitising tablet (Houston Instruments DT-114). Data were then run through a computer program that sampled it every 0.5 seconds. The FHR was recorded, digitised and sampled for 15 to 25 minutes before and after</p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis pH &lt;7.20</b> 15/104 (14%)</p> <p><b>Predictive value of mean change in heart rate &lt; 15bpm following fetal scalp stimulation for fetal blood sample pH &lt; 7.20</b></p> <p>As reported in Table 4 of paper; NCC calculated confidence intervals, LR+ and LR- Sensitivity: 73% (50.95 to 95.71) Specificity: 17% (9.08 to 24.63) PPV: 13% (5.81 to 20.08) NPV: 79% (60.62 to 97.28) LR+: 0.88 (0.64 to 1.21) LR-: 1.58 (0.61 to 4.12)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between FBS and delivery was recorded but not reported</p>												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>USA</p> <p><b>Aim of the study</b></p> <p>To determine whether fetal scalp stimulation during active labour results in a fetal heart response, and whether the magnitude and direction of any change is related to fetal acid-base status</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>FBS. The 5 minutes immediately preceding FBS were omitted from the analysis. FHR was averaged for 5 minutes before the beginning of preparations for the FBS procedure and over 1 minute immediately following FBS to obtain pre- and post-stimulation mean heart rates.</p> <p>The effect of fetal scalp stimulation was examined by setting the time of scalp incision at zero and determining the FHR at 0.5 second intervals before and after the scalp incision from the digitised heart rate recordings.</p> <p>Subjects were divided in to three groups according FBS pH and mean and standard error of the heart rate for each group was determined for each 0.5 second sample point. These values were then plotted as a function of time for each group.</p>	<p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>11</td> <td>74</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>4</td> <td>15</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	11	74	<b>Predictive Test -ve</b>	4	15	<p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness of outcome: standard definition of acceleration not used; net difference in heart rate of more than 15 bpm was applied; population and inclusion and exclusion criteria not sufficiently reported to assess indirectness of population</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: mean increase in FHR &lt;15 bpm.</p> <p>Some fetuses underwent more than one scalp blood sampling; only the first sampling was used to avoid the effect of habituation.</p> <p>All fetuses with FBS pH &lt; 7.20 were tested at delivery for acidosis by cord blood gas analysis.</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	11	74												
<b>Predictive Test -ve</b>	4	15												
<p><b>Full citation</b></p> <p>Lin,C.C., Vassallo,B., Mittendorf,R., Is</p>	<p><b>Sample size</b></p> <p>N = 113</p>	<p><b>Tests</b></p>	<p><b>Methods</b></p> <p>3-seconds of VAS using an artificial larynx (model 5E,</p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis</b></p> <p>31/113 (27%)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear</p>									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>intrapartum vibroacoustic stimulation an effective predictor of fetal acidosis?, Journal of Perinatal Medicine, 29, 506-512, 2001</p> <p><b>Ref Id</b> 201886</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Aim of the study</b> The hypothesis is that intrapartum vibroacoustic stimulation is an effective predictor of fetal acidosis during labour</p> <p><b>Study type</b></p> <p><b>Study dates</b> 1 July 1995 - 30 April 1997</p> <p><b>Source of funding</b> Not reported</p>	<p><b>Characteristics</b></p> <p><u>Stage of labour</u> First stage = 53 Second stage = 60</p> <p><u>Gestational age</u> Term (≥ 37 weeks) = 94 Pre-term (≥ 34, &lt; 37 weeks) = 13 Very pre-term (&lt; 34 weeks) = 6</p> <p><b>Inclusion Criteria</b> Singleton gestations in active phase of first or second stage of labour and exhibiting abnormal fetal heart rate (FHR) patterns (moderate to severe variable decelerations or late decelerations, with or without baseline tachycardia or significantly decreased baseline variability).</p> <p>Women with known medical or obstetric complications, such as diabetes, hypertension, preeclampsia or fetal growth restriction were included.</p>	<p>3 seconds of fetal vibroacoustic stimulation (VAS)</p>	<p>AT&amp;T, Van Nuys, CA, USA) was applied to the maternal abdomen directly over the fetal head. For women in the second stage of labour VAS was applied to the suprapubic area, or if the fetal head was at plus two station or lower, directly to the fetal head on parietal or occiput area with a sterile latex glove covered VAS applicator.</p> <p>FHR response was monitored; a positive response was defined as 15bpm acceleration above baseline for a duration ≥ 15 seconds. No response or a deceleration after VAS suggested an acidotic fetus. A biphasic response, defined as an acceleration followed by a deceleration was considered equivocal.</p> <p>Scalp blood was obtained immediately following VAS testing during the first stage of labour. During the second stage of labour, one or several VAS testings were performed, so that the time intervals between the last VAS testing and the delivery of the fetus were within 15 minutes.</p> <p>Umbilical blood sample was obtained at delivery for fetal blood pH and blood gas analysis in every case by a</p>	<p><b>Predictive value of no acceleration following VAS</b></p> <p><u>a. For fetal blood sample pH &lt; 7.20</u> Values as reported in Table II; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 39% (21.56 to 55.86) Specificity: 93% (87.05 to 98.32) PPV: 67% (44.89 to 88.44) NPV: 80% (71.96 to 88.04) LR+: 5.29 (2.18 to 12.86) LR-: 0.66 (0.50 to 0.88)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 100% (100 to 100) Specificity: 86% (79.95 to 92.78) PPV: 17% (0 to 33.88) NPV: 100% (100 to 100) LR+: 7.33 (4.58 to 11.74) LR-: 0 (NC)</p> <p><u>c. For NICU admission</u> Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 55% (33.20 to 76.80) Specificity: 92% (87.11 to 97.84) PPV: 61% (38.59 to 83.63) NPV: 91% (84.64 to 96.42) LR+: 7.31 (3.23 to 16.51) LR-: 0.49 (0.30 to 0.79)</p> <p><u>d. For neonatal morbidity</u> Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals</p>	<p>whether consecutive women were included Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome; period of FHR observation for qualifying acceleration following stimulus was not reported Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between FBS and delivery for women in first stage of labour unclear Statistical analysis is appropriate for study design: yes Indirectness of population: 17% of women were in pre-term labour; high risk women were included (numbers not reported)</p> <p><b>Other information</b> While authors state a positive stimulation test was FHR acceleration, statistics reported are for no acceleration predicting acidosis (&lt; 7.20).</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
	<p><b>Exclusion Criteria</b></p> <p>Multiple gestation, congenital fetal malformations, gestational age &lt; 28 weeks and administration of narcotic analgesia to the mother within the last 3 hours</p>		<p>Corometric 220 pH System (Wallingford, CT).</p> <p>The decision to perform fetal scalp blood sampling or caeserean section was made by the attending physician or senior resident assessing the FHR tracing and reviewing the clinical course.</p>	<p>Sensitivity: 71% (37.96 to 105)                      Specificity: 88% (81.49 to 93.98)                      PPV: 28% (7.09 to 48.47)                      NPV: 98% (95.01 to 101)                      LR+: 5.82 (2.91 to 11.63)                      LR-: 0.33 (0.10 to 1.05)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 549 1733 847"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>12</td> <td>6</td> </tr> <tr> <td>Predictive Test -ve</td> <td>19</td> <td>76</td> </tr> </tbody> </table> <p><b>Appgar score</b></p> <table border="1" data-bbox="1346 959 1733 1257"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>15</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>95</td> </tr> </tbody> </table> <p><b>NICU admission</b></p>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	12	6	Predictive Test -ve	19	76		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	15	Predictive Test -ve	0	95	<p>Authors' definition of positive stimulation test: no acceleration.</p> <p>When more than one fetal blood pH value was obtained, only the last one was used for analysis.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	12	6																					
Predictive Test -ve	19	76																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	3	15																					
Predictive Test -ve	0	95																					

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<table border="1" data-bbox="1346 272 1729 571"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>11</td> <td>7</td> </tr> <tr> <td>Predictive Test -ve</td> <td>9</td> <td>86</td> </tr> </tbody> </table> <p data-bbox="1346 632 1559 655"><b>Neonatal morbidity</b></p> <table border="1" data-bbox="1346 683 1729 981"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>5</td> <td>13</td> </tr> <tr> <td>Predictive Test -ve</td> <td>2</td> <td>93</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	11	7	Predictive Test -ve	9	86		Reference Test +ve	Reference Test -ve	Predictive Test +ve	5	13	Predictive Test -ve	2	93	
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	11	7																					
Predictive Test -ve	9	86																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	5	13																					
Predictive Test -ve	2	93																					
<p><b>Full citation</b></p> <p>Polzin,G.B., Blakemore,K.J., Petrie,R.H., Amon,E., Fetal vibro-acoustic stimulation: magnitude and duration of fetal heart rate accelerations as a marker of fetal health, Obstetrics and Gynecology, 72, 621-626, 1988</p>	<p><b>Sample size</b></p> <p>N = 100</p> <p><b>Characteristics</b></p> <p><u>Gestational age (weeks)</u> - mean ± SD, N 15 bpm x 15 sec acceleration = 39.4 ± 1.9, 57 10 bpm x 10 sec acceleration = 39.1 ±</p>	<p><b>Tests</b></p> <p>5 seconds of continuous fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>Over a period of 20 months, when one of the study authors was available, 100 women were studied using the standard indications for fetal scalp blood sampling (FBS; late, moderate or severe variable fetal heart rate (FHR) decelerations, fetal tachycardia or bradycardia, or poor FHR variability longer than 30 minutes).</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis &lt; 7.20</u></b> 10/100 (10%)</p> <p><b><u>Predictive value of no acceleration following VAS</u></b> a. For fetal blood sample pH &lt; 7.20 All values calculated by NCC from data presented in Table 4 (see Other information) Sensitivity: 90% (71.41 to 100) Specificity: 84.44% (76.96 to 91.93) PPV: 39.13% (19.88 to 59.08)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: not consecutive (women only included when one of the study authors was available) Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in</p>																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p><b>Ref Id</b> 201800</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Aim of the study</b> To evaluate whether there are significant differences in the intrapartum fetal acid-base status according to the magnitude and duration of fetal heart rate accelerations in response to fetal vibroacoustic stimulation. The predictive value of these responses in the detection of the acidotic versus non-acidotic fetus during labour was also examined.</p> <p><b>Study type</b></p> <p><b>Study dates</b> Over a 20 month period (dates not reported)</p> <p><b>Source of funding</b> Not reported</p>	<p>2.5, 20 No acceleration = 38.3 ± 3.1, 23</p> <p><u>Birth weight (g) - mean ±SD, N</u> 15 bpm x 15 sec acceleration = 3289 ± 527, 57 10 bpm x 10 sec acceleration = 3043 ± 588, 20 No acceleration = 2703 ± 909, 23</p> <p><b>Inclusion Criteria</b> Active phase of labour, singleton gestation, vertex presentation</p> <p><b>Exclusion Criteria</b> Not reported</p>		<p>Immediately before FBS, VAS was performed using a Model 5C electronic artificial larynx (AT&amp;T Consumer Products, USA), which produced a mixed-frequency sound of 81 Hz and 81 db at 1 m in air. A single stimulus was applied continuously for 5 seconds to the maternal abdomen one-third of the distance from the symphysis pubis to the umbilicus. FHR accelerations, if they occurred, began within 20 seconds of the stimulus.</p> <p>FHR responses were classified in to three groups: FHR acceleration of ≥ 15 bpm lasting ≥ 15 seconds, 10-15 bpm lasting 10-15 seconds, or no acceleration.</p> <p>FBS was performed immediately after VAS, usually in the left lateral position. Mean pH values were derived from logarithmic tables.</p>	<p>NPV: 98.70 % (96.17 to 100) LR+: 5.79 (3.43 to 9.77) LR-: 0.11 (0.02 to 0.76)</p> <p><u>b. For fetal blood sample pH &lt; 7.25</u> All values calculated by NCC from data presented in Table 4 (see Other information) Sensitivity: 45.45% (24.65 to 66.26) Specificity: 83.33% (75.06 to 91.60) PPV: 43.48% (23.22 to 63.74) NPV: 84.41% (76.31 to 92.52) LR+: 2.73 (1.39 to 5.36) LR-: 0.65 (0.44 to 0.97)</p> <p><u>c. For Apgar score &lt; 7 at 5 minutes</u> All values calculated by NCC from data presented in Table 2 Sensitivity: 50% (9.99 to 90.01) Specificity: 57.45% (47.45 to 67.44) PPV: 6.98% (1 to 14.59) NPV: 94.74% (88.94 to 100) LR+: 1.18 (0.51 to 2.71) LR-: 0.87 (0.38 to 1.97)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>9</td> <td>14</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>76</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	9	14	Predictive Test -ve	1	76	<p>participants: unclear whether assessor blinded to outcome Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between FBS and delivery not reported Statistical analysis is appropriate for study design: yes Indirectness: based on gestational age mean and SD for 'no acceleration' population not all fetuses were delivered at term; unclear whether any women were considered high risk</p> <p><b>Other information</b> Authors' definition of positive stimulation test: no acceleration. Predictive accuracy statistics presented in Table 3 of study report do not account for the full study population - data for 10bpm x 10sec population not included with the no acceleration population. Therefore, data extracted for full study population from Table 4</p>
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	9	14												
Predictive Test -ve	1	76												



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<p><b>FBS pH</b></p> <table border="1" data-bbox="1346 379 1733 683"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>10</td> <td>13</td> </tr> <tr> <td>Predictive Test -ve</td> <td>12</td> <td>65</td> </tr> </tbody> </table> <p><b>Apgar score</b></p> <table border="1" data-bbox="1346 794 1733 1094"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>40</td> </tr> <tr> <td>Predictive Test -ve</td> <td>3</td> <td>54</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	10	13	Predictive Test -ve	12	65		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	40	Predictive Test -ve	3	54	<p>and all statistics calculated by NCC.</p> <p>For the 2x2 table no acceleration and FHR acceleration <math>\geq 10</math> bpm and 10 sec but <math>&lt; 15</math> bpm and 15 sec were considered a positive stimulation test result.</p> <p>In nearly all cases FHR was recorded by internal scalp electrode.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	10	13																					
Predictive Test -ve	12	65																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	3	40																					
Predictive Test -ve	3	54																					
<p><b>Full citation</b></p> <p>Sarno,A.P., Ahn,M.O., Phelan,J.P., Paul,R.H., Fetal acoustic stimulation in the early intrapartum period as a predictor of subsequent fetal condition, American Journal of Obstetrics and</p>	<p><b>Sample size</b></p> <p>N = 201</p> <p><b>Characteristics</b></p> <p><u>Maternal age (years) - mean <math>\pm</math> SD</u> 25.9 <math>\pm</math> 5.5</p>	<p><b>Tests</b></p> <p>3 seconds of fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>Consecutive women who met inclusion criteria were included over the study period, during periods of availability of the first author.</p> <p>Following admission electronic fetal monitoring was instituted.</p>	<p><b>Results</b></p> <p><b><u>Predictive value of no acceleration following VAS</u></b></p> <p>a. For Apgar score <math>&lt; 7</math> at 1 minute Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 24.1% (8.56 to 39.71) Specificity: 95.9% (92.98 to 98.88)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: included women who were considered high risk Loss to follow-up is unrelated to key characteristics: no loss to follow up</p>																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
<p>Gynecology, 162, 762-767, 1990</p> <p><b>Ref Id</b></p> <p>201730</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Aim of the study</b></p> <p>To evaluate the usefulness of fetal acoustic stimulation in the early intrapartum period as a predictor of subsequent fetal condition</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>1 August 1987 - 1 November 1987</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><u>Nulliparous</u> 74/201 (37%)</p> <p><u>Gestational age (weeks) - mean ± SD</u> 40.1 ± 2.2</p> <p><u>Duration of ruptured membranes (hours) - mean ± SD</u> 14.2 ± 17.0</p> <p><u>Duration of labour (hours) - mean ± SD</u> 17.4 ± 8.5</p> <p><b>Inclusion Criteria</b></p> <p>Gestational age ≥ 37 weeks, singleton fetus, vertex presentation, latent phase of labour (cervical dilatation ≤ 4 cm)</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>A 40-min baseline fetal heart rate (FHR) monitor tracing was obtained, then VAS was performed using a fetal acoustic stimulator (Corometrics model 146, Wallingford, CT, USA), sound level 82 dB at 1 m in air. The acoustic stimulator was placed on the maternal abdomen over the fetal vertex and a 3-second pulse of stimulation applied. If no acceleration of FHR was noted within 1 min an additional pulse was administered to a maximum of three pulses, each 1 minute apart.</p> <p>A reactive response was defined as one or more accelerations of the FHR 15 bpm from baseline, persisting for 15 seconds. A non-reactive response was defined as failure to elicit a qualifying acceleration after any of three separate stimuli and for 10 minutes after the last stimulus.</p> <p>Care was taken not to perform acoustic stimulation during or immediately after a uterine contraction to avoid periods of transient fetal hypoxia and for standardisation of the technique.</p> <p>The result of stimulation was blinded from the physicians</p>	<p>PPV: 50% (23.81 to 76.19) NPV: 88.2% (83.62 to 92.85) LR+: 5.93 (2.25 to 15.66) LR-: 0.79 (0.64 to 0.97)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 33.3% (0 to 71.05) Specificity: 93.8% (90.47 to 97.22) PPV: 14.3% (0 to 32.62) NPV: 97.9% (95.79 to 99.93) LR+: 5.42 (1.54 to 19.05) LR-: 0.71 (0.40 to 1.25)</p> <p><u>c. For caesarean delivery for fetal distress</u> Values as reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 31.2% (8.54 to 53.96) Specificity: 95.1% (92.04 to 98.24) PPV: 35.7% (10.61 to 60.81) NPV: 94.1% (90.75 to 97.49) LR+: 6.42 (2.44 to 16.89) LR-: 0.72 (0.52 to 1.01)</p> <p><b>Apgar score</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>7</td> <td>7</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	7	7	<p>Prognostic factor is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential cofounders are accounted for: time between VAS and delivery not reported Statistical analysis is appropriate for study: yes Indirectness of population: 118/201 (59%) had one or more complications of pregnancy [complications not reported]</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration.</p>
	Reference Test +ve	Reference Test -ve									
<b>Predictive Test +ve</b>	7	7									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																											
			<p>who managed the woman's labour. All FHR tracings were read by a single examiner without knowledge of the prior fetal acoustic stimulation results.</p> <p>Outcome was assessed by incidences of meconium staining, fetal distress requiring caesarean delivery, Apgar scores &lt; 7 at 1 and 5 minutes, subsequent abnormal FHR patterns and perinatal mortality.</p>	<table border="1"> <tr> <td data-bbox="1348 272 1464 371">Predictive Test -ve</td> <td data-bbox="1464 272 1599 371">22</td> <td data-bbox="1599 272 1733 371">165</td> </tr> <tr> <td colspan="3" data-bbox="1348 427 1733 459"><b>Apgar score</b></td> </tr> <tr> <td></td> <td data-bbox="1464 483 1599 582"><b>Reference Test +ve</b></td> <td data-bbox="1599 483 1733 582"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1348 582 1464 681">Predictive Test +ve</td> <td data-bbox="1464 582 1599 681">2</td> <td data-bbox="1599 582 1733 681">12</td> </tr> <tr> <td data-bbox="1348 681 1464 780">Predictive Test -ve</td> <td data-bbox="1464 681 1599 780">4</td> <td data-bbox="1599 681 1733 780">183</td> </tr> <tr> <td colspan="3" data-bbox="1348 836 1733 868"><b>Caesarean section</b></td> </tr> <tr> <td></td> <td data-bbox="1464 892 1599 991"><b>Reference Test +ve</b></td> <td data-bbox="1599 892 1733 991"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1348 991 1464 1090">Predictive Test +ve</td> <td data-bbox="1464 991 1599 1090">5</td> <td data-bbox="1599 991 1733 1090">9</td> </tr> <tr> <td data-bbox="1348 1090 1464 1189">Predictive Test -ve</td> <td data-bbox="1464 1090 1599 1189">11</td> <td data-bbox="1599 1090 1733 1189">176</td> </tr> </table>	Predictive Test -ve	22	165	<b>Apgar score</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	2	12	Predictive Test -ve	4	183	<b>Caesarean section</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	5	9	Predictive Test -ve	11	176	
Predictive Test -ve	22	165																														
<b>Apgar score</b>																																
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																														
Predictive Test +ve	2	12																														
Predictive Test -ve	4	183																														
<b>Caesarean section</b>																																
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																														
Predictive Test +ve	5	9																														
Predictive Test -ve	11	176																														
<p><b>Full citation</b></p> <p>Smith,C.V., Nguyen,H.N., Phelan,J.P., Paul,R.H., Intrapartum assessment of fetal well-being: a</p>	<p><b>Sample size</b></p> <p>N = 64</p>	<p><b>Tests</b></p> <p>≤ 3 seconds of fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>Immediately before fetal blood sampling (FBS) with the woman in the dorsal lithotomy position, transabdominal</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis pH &lt; 7.25</u></b> 18/64 (28%)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included</p>																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>comparison of fetal acoustic stimulation with acid-base determinations, American Journal of Obstetrics and Gynecology, 155, 726-728, 1986</p> <p><b>Ref Id</b> 201855</p> <p><b>Country/ies where the study was carried out</b> USA</p> <p><b>Aim of the study</b> To compare acoustically evoked accelerations of the fetal heart rate (FHR) with fetal acid-base status</p> <p><b>Study type</b></p> <p><b>Study dates</b> Not reported</p> <p><b>Source of funding</b> Not reported</p>	<p><b>Characteristics</b></p> <p><u>FHR abnormality indicating need for fetal blood sampling</u></p> <p>Intermittent late decelerations = 20/64 (31%) Severe variable decelerations = 14/64 (22%) Absent variability = 12/64 (19%) Tachycardia = 11/64 (17%) Repetitive late decelerations = 7/64 (11%)</p> <p><b>Inclusion Criteria</b></p> <p>Women with FHR tracings sufficiently abnormal to merit either fetal blood sampling (FBS) or immediate caesarean delivery for fetal distress</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>acoustic stimulation of the fetus was accomplished by a Model 5C electronic artificial larynx (Western Electric). The artificial larynx produces a vibratory acoustic stimulus of approximately 80 Hz and 82 dB, measured at 1 m in air. The stimulus was applied overlying the fetal vertex for ≤ 3 seconds. The response was termed reactive if an immediate acceleration of 15 bpm and 15 seconds was evident. If a qualifying acceleration was not present, the stimulus was repeated at 1-minute intervals for a maximum of three times. Fetal scalp sampling was then accomplished by existing protocol.</p> <p>In 15 cases where scalp sampling was not possible immediate cesarean delivery was performed. In all cases the fetus was delivered within 15 minutes of the stimulus. The arithmetic mean of the umbilical arterial and venous pH determinations was calculated.</p>	<p><u>Predictive value of no acceleration following VAS for fetal blood sample pH &lt; 7.25</u></p> <p>All values calculated by NCC from data in Table II Sensitivity: 100% (100 to 100) Specificity: 65.22% (51.45 to 78.98) PPV: 52.94% (36.16 to 69.72) NPV: 100% (100 to 100) LR+: 2.88 (1.94 to 4.27) LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td><b>Predictive Test +ve</b></td> <td>18</td> <td>16</td> </tr> <tr> <td><b>Predictive Test -ve</b></td> <td>0</td> <td>30</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	18	16	<b>Predictive Test -ve</b>	0	30	<p>Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome Outcome of interest is sufficiently measured in participants: yes Important potential cofounders are accounted for: length of stimulation not standardised (≤ 3 seconds); time between VAS and deliveries that were not caesarean births not reported Statistical analysis is appropriate for study: yes Indirectness: unclear whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Definition of positive stimulation test: no acceleration (selected by NCC, authors do not define positive stimulation test and do not report predictive accuracy statistics).</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	18	16												
<b>Predictive Test -ve</b>	0	30												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					Five fetuses that failed to respond to VAS did respond to the stimulus of FBS scalp puncture (data for scalp puncture not sufficiently reported to construct 2x2 table).
<p><b>Full citation</b></p> <p>Spencer, J.A., Predictive value of a fetal heart rate acceleration at the time of fetal blood sampling in labour, <i>Journal of Perinatal Medicine</i>, 19, 207-215, 1991</p> <p><b>Ref Id</b></p> <p>196967</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Aim of the study</b></p> <p>To present the results of a 1-year audit of all cases requiring fetal scalp blood sampling during labour at a major teaching hospital, with particular emphasis on the relationship between the fetal heart rate reaction at the time of fetal scalp</p>	<p><b>Sample size</b></p> <p>N = 138</p> <p><b>Characteristics</b></p> <p><u>Gestation <math>\geq</math> 37 weeks</u> 133/138 (96%)</p> <p><u>Nulliparous</u> 110/138 (80%)</p> <p><u>Mode of delivery</u> Normal vaginal delivery = 38/138 (27%) Operative vaginal delivery = 60/138 (43%) Caesarean section = 40/138 (30%)</p> <p><b>Inclusion Criteria</b></p> <p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>The incision of fetal scalp blood sampling served as fetal scalp stimulation</p>	<p><b>Methods</b></p> <p>Data were collected from all cases that required intrapartum fetal scalp blood sampling (FBS) due to concerns regarding the CTG during 1 year at the John Radcliffe Maternity Hospital, Oxford.</p> <p>Fetal heart rate (FHR) records were derived from the fetal electrocardiogram using a spiral electrode and an HP 8040 fetal monitor (Hewlett Packard, Uxbridge, UK). FHR reaction to FBS was noted to be either an acceleration (transient rise above baseline of more than 15 bpm for longer than 15 seconds), no response or a deceleration (transient fall below baseline of more than 15 bpm for longer than 15 seconds). Fetal scalp blood was collected into heparinised capillary tubes for immediate blood gas analysis</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis <math>&lt;</math> 7.20</u></b> 6/138 (4%)</p> <p><b><u>1. Predictive value of an acceleration following fetal scalp stimulation</u></b></p> <p><b><u>a. For fetal blood sample pH <math>\geq</math> 7.20</u></b> As reported in Table V; NCC calculated LR+, LR- and all confidence intervals Sensitivity: 52.3% (43.75 to 60.79) Specificity: 100% (100 to 100) PPV: 100% (100 to 100) NPV: 8.7% (2.05 to 15.34) LR+: NC LR-: 0.48 (0.40 to 0.57)</p> <p><b><u>b. For fetal blood sample pH <math>\geq</math> 7.25</u></b> All values calculated by NCC from data in Table IV Sensitivity: 53.57% (44.33 to 62.81) Specificity: 65.38% (47.10 to 83.67) PPV: 86.96% (79.01 to 94.90) NPV: 24.64% (14.47 to 34.81) LR+: 1.55 ((0.89 to 2.70) LR-: 0.71 (0.50 to 1.00)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: yes Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: unclear whether assessor blinded to outcome; period of FHR observation for qualifying acceleration following stimulus was not reported Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between stimulation, FBS and delivery not reported Statistical analysis is appropriate for study design: yes Indirectness: 96% delivered at term; unclear</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>blood sampling and the fetal scalp pH</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>using an ABL 3 (Radiometer, Copenhagen).</p> <p>Fetal pH was related to the FHR before the FBS and to the FHR reaction at the time.</p>	<p><b><u>2. Predictive value of no acceleration following fetal scalp stimulation</u></b></p> <p><b><u>a. For fetal blood sample pH &lt; 7.20</u></b> All values calculated by NCC from data in Table IV Sensitivity: 100% (100 to 100) Specificity: 52.27% (43.75 to 60.79) PPV: 8.70% (2.05 to 15.34) NPV: 100% (100 to 100) LR+: 2.10 (1.75 to 2.50) LR-: 0 (NC)</p> <p><b><u>b. For fetal blood sample pH &lt; 7.25</u></b> All values calculated by NCC from data in Table IV Sensitivity: 65.38% (47.10 to 83.67) Specificity: 53.57% (44.33 to 62.81) PPV: 24.64% (14.47 to 34.81) NPV: 86.96% (79.01 to 94.90) LR+: 1.41 (1.00 to 1.98) LR-: 0.87 (0.79 to 0.95)</p> <p><b><u>c. For Apgar score &lt; 7 at 1 minute</u></b> All values calculated by NCC from data in Table III Sensitivity: 54.00% (40.19 to 67.81) Specificity: 52.27% (41.84 to 62.71) PPV: 39.13% (27.61 to 50.65) NPV: 66.67% (55.54 to 77.79) LR+: 1.13 (0.81 to 1.58) LR-: 0.88 (0.61 to 1.26)</p> <p><b><u>d. For Apgar score &lt; 7 at 5 minutes</u></b> Calculated by NCC from data in Table III Sensitivity: 100% (100 to 100) Specificity: 50.36% (41.99 to 58.74) PPV: 1.45% (0 to 4.27%) NPV: 100% (100 to 100)</p>	<p>whether any women were considered high risk</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: acceleration (≥ 15 bpm above baseline for ≥ 15 seconds) Authors' definition of positive FBS pH; a. no acidosis ≥ 7.20; b. no acidosis ≥ 7.25</p> <p>First set of predictive accuracy results in evidence table are as reported in the study. Second set of predictive accuracy results were calculated by NCC with a recalculated 2x2 table using a definition of positive stimulation test being no acceleration and definition of positive fetal scalp test of a. acidosis pH &lt; 7.20 and b. acidosis pH &lt; 7.25 in line with other studies included in this review. Data for Apgar score &lt; 7 at 1 and 5 minutes was calculated from Apgar ≥ 7 at 1 and 5 minutes reported in Table III, to be in line with other</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<p>LR+: 2.01 (1.70 to 2.38) LR-: 0 (NC)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 440 1733 738"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>69</td> <td>0</td> </tr> <tr> <td>Predictive Test -ve</td> <td>63</td> <td>6</td> </tr> </tbody> </table> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 850 1733 1149"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>60</td> <td>9</td> </tr> <tr> <td>Predictive Test -ve</td> <td>52</td> <td>17</td> </tr> </tbody> </table> <p><b>FBS pH</b></p>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	69	0	Predictive Test -ve	63	6		Reference Test +ve	Reference Test -ve	Predictive Test +ve	60	9	Predictive Test -ve	52	17	<p>studies included in this review.</p> <p>Approximately 50% of labours were monitored by CTG because of perceived risk factors or the use of epidural analgesia.</p> <p>Only the first FBS on any single patient was included in the analysis.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	69	0																					
Predictive Test -ve	63	6																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	60	9																					
Predictive Test -ve	52	17																					

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>6</td> <td>63</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>69</td> </tr> </tbody> </table> <p><b>FBS pH</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>17</td> <td>52</td> </tr> <tr> <td>Predictive Test -ve</td> <td>9</td> <td>60</td> </tr> </tbody> </table> <p><b>Apgar score</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>27</td> <td>42</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	6	63	Predictive Test -ve	0	69		Reference Test +ve	Reference Test -ve	Predictive Test +ve	17	52	Predictive Test -ve	9	60		Reference Test +ve	Reference Test -ve	Predictive Test +ve	27	42	
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	6	63																											
Predictive Test -ve	0	69																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	17	52																											
Predictive Test -ve	9	60																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	27	42																											



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments												
				<table border="1"> <tr> <td>Predictive Test -ve</td> <td>23</td> <td>46</td> </tr> </table> <p>Apgar score</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>1</td> <td>68</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>69</td> </tr> </tbody> </table>	Predictive Test -ve	23	46		Reference Test +ve	Reference Test -ve	Predictive Test +ve	1	68	Predictive Test -ve	0	69	
Predictive Test -ve	23	46															
	Reference Test +ve	Reference Test -ve															
Predictive Test +ve	1	68															
Predictive Test -ve	0	69															
<p><b>Full citation</b></p> <p>Tannirandorn,Y., Wacharaprechanont,T., Phaosavasdi,S., Fetal acoustic stimulation for rapid intrapartum assessment of fetal well-being, Journal of the Medical Association of Thailand, 76, 606-612, 1993</p> <p><b>Ref Id</b></p> <p>201731</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 140</p> <p><b>Characteristics</b></p> <p><u>Nulliparous</u> 88/140 (63%)</p> <p><u>Gestational age (weeks) - mean (range)</u> 39.5 (37 - 43)</p> <p><u>Antenatal risk factors</u> Post-term (≥ 42 weeks) = 14/140 (10%) Poor weight gain = 11/140 (7.8%) Pre-eclampsia = 9/140 (6.4%)</p>	<p><b>Tests</b></p> <p>3-seconds of fetal vibroacoustic stimulation (VAS)</p>	<p><b>Methods</b></p> <p>After admission to the delivery room, blood pressure was monitored at 10-min intervals and a tocodynamometer and Doppler FHR transducer (Sonic Aid FM 3, Oxford, UK) were applied to the abdomen and adjusted for best signal.</p> <p>Fetal heart rate (FHR) and uterine contractions were recorded for 15 to 20 min. Acoustic stimulation was then performed using a fetal acoustic stimulator (Corometrics 146, CT, USA; sound level 82 dB at 1 m in air) placed on the maternal abdomen over the fetal head</p>	<p><b>Results</b></p> <p><b><u>Predictive value of no acceleration following VAS for poor perinatal outcome</u></b> Values as reported in Table 4; NCC caclulated LR+, LR- and all confidence intervals Sensitivity: 71.4% (37.96 to 100) Specificity: 99.2% (97.78 to 100) PPV: 83.3% (53.51 to 100) NPV: 98.5% (96.45 to 100) LR+: 95 (12.75 to 707.63) LR-: 0.29 (0.09 to 0.93)</p> <p><b>Poor perinatal outcome</b></p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether consecutive women were included Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: 15-minute window for reaction to 3rd stimulus, compared with 30-sec</p>												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>Thailand</p> <p><b>Aim of the study</b></p> <p>To evaluate the usefulness of fetal acoustic stimulation in the early intrapartum period as a rapid screening test to predict subsequent fetal condition</p> <p><b>Study type</b></p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>No antenatal care = 5/140 (3.6%)                      Oligohydramnios = 1/140 (0.7%)                      Others (poor obstetric history, intrauterine growth restriction, diabetes etc.) = 5/140 (3.6%)</p> <p><b>Inclusion Criteria</b></p> <p>Gestational age <math>\geq</math> 37 weeks, cephalic presentation, latent phase of labour (cervical dilatation <math>\leq</math> 3 cm), intact membranes</p> <p><b>Exclusion Criteria</b></p> <p>Women with spurious labour who had not been delivered within 24 hours of admission and those with twin pregnancies or known fetal abnormalities were excluded from analysis</p>		<p>and a 3-sec pulse of sound stimulation was applied. If no acceleration of the FHR was noted within 30 sec an additional pulse was administered to a maximum of 3 pulses, 30 seconds apart. Care was taken not to perform acoustic stimulation during or immediately after uterine contractions to avoid periods of transient fetal hypoxia and for standardisation of the technique.</p> <p>A reactive response to VAS was defined as one or more accelerations of FHR <math>\geq</math> 15 bpm from the baseline persisting for 15 seconds. A non-reactive response was defined as a failure to elicit a qualifying acceleration after any of three separate stimuli and for 15 min after the last stimulus. All VAS results were interpreted by a single examiner without knowledge of the perinatal outcome. Obstetricians managing the woman's labour were not informed of the results of VAS.</p> <p>Perinatal outcome was considered poor when there was perinatal death, a 5-min Apgar score <math>&lt;</math> 7, fetal distress requiring caesarean section, thick meconium stained amniotic fluid or admission to</p>	<table border="1" data-bbox="1346 272 1733 571"> <thead> <tr> <th data-bbox="1346 272 1464 379"></th> <th data-bbox="1464 272 1599 379">Reference Test +ve</th> <th data-bbox="1599 272 1733 379">Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td data-bbox="1346 379 1464 475"><b>Predictive Test +ve</b></td> <td data-bbox="1464 379 1599 475">5</td> <td data-bbox="1599 379 1733 475">1</td> </tr> <tr> <td data-bbox="1346 475 1464 571"><b>Predictive Test -ve</b></td> <td data-bbox="1464 475 1599 571">2</td> <td data-bbox="1599 475 1733 571">132</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	5	1	<b>Predictive Test -ve</b>	2	132	<p>window for reaction to 1st and 2nd stimuli; time between VAS and delivery not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness of population: 32% of women had one or more antenatal complication (10% had a gestational age <math>\geq</math> 42 weeks)</p> <p>Indirectness of outcome: composite of poor perinatal outcome</p> <p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	5	1												
<b>Predictive Test -ve</b>	2	132												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
			the neonatal intensive care unit.		
<p><b>Full citation</b></p> <p>Trochez,R.D., Sibanda,T., Sharma,R., Draycott,T., Fetal monitoring in labor: are accelerations good enough?, Journal of Maternal-Fetal and Neonatal Medicine, 18, 349-352, 2005</p> <p><b>Ref Id</b></p> <p>201769</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Aim of the study</b></p> <p>To investigate whether accelerations evoked by fetal scalp stimulation from routine vaginal examination prior to fetal blood sampling (FBS) predicted the absence of fetal acidosis at the time of the FBS</p> <p><b>Study type</b></p>	<p><b>Sample size</b></p> <p>N = 54</p> <p><b>Characteristics</b></p> <p><u>Mode of delivery</u> Spontaneous vertex = 17/54 (31%) Instrumental = 22/54 (41%) Emergency caesarean section = 15/54 (28%)</p> <p><b>Inclusion Criteria</b></p> <p>Term (&gt; 37 weeks gestation) singleton fetuses where FBS was obtained in labour</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>Fetal scalp stimulation during vaginal examination (method and duration of stimulation not reported)</p>	<p><b>Methods</b></p> <p>69 fetuses were identified during the study period but information retrieval was only possible in 54 (78%), in whom 70 scalp blood sample procedures were performed.</p> <p>The CTG traces for all of these fetuses were reviewed by an investigator blind to the outcome. A portion of the trace starting from the point of the vaginal examination, as indicated by routine markings made on the CTG by the attending midwife, was reviewed for accelerations. Accelerations were defined as an increase in fetal heart rate above the baseline of at least 15bpm for at least 15 seconds.</p> <p>The position of the presenting part was determined and recorded in all cases ensuring scalp stimulation.</p>	<p><b>Results</b></p> <p><b><u>Prevalence of acidosis <math>\leq</math> 7.20</u></b> 5/70 (7%)</p> <p><b><u>Predictive value of no acceleration fetal scalp stimulation</u></b></p> <p><b><u>a. For fetal blood sample pH <math>\leq</math> 7.20</u></b> As reported in Table I and Table II of paper Sensitivity: 40% (7.26 to 82.96) Specificity: 69.23% (56.4 to 79.76) PPV: 9.09% (2.52 to 27.81) NPV: 93.75% (83.16 to 97.85) LR+: 1.3 (0.27 to 6.24) LR-: 0.87 (0.44 to 1.70)</p> <p><b><u>b. For cord pH <math>\leq</math> 7.20</u></b> Calculated by NCC from data in Table III Sensitivity: 40% (-2.94 to 82.94) Specificity: 75.86% (60.29 to 91.44) PPV: 22.22% (-4.94 to 49.38) NPV: 88% (75.26 to 100) LR+: 1.66 (0.47 to 5.80) LR-: 0.79 (0.38 to 1.67)</p> <p><b><u>c. For Apgar score <math>&lt;</math> 7 at 5 minutes</u></b> Calculated by NCC from data in Table III Sensitivity: 50% (1 to 99) Specificity: 69.57% (56.27 to 82.86) PPV: 12.5% (-3.71 to 28.71) NPV: 94.12% (86.21 to 102.03)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: yes Loss to follow-up is unrelated to key characteristics: no loss to follow up Prognostic factor is adequately measured in participants: period of FHR observation for qualifying acceleration following stimulus not reported Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: time between stimulation, FBS and delivery reported only for acidotic babies, not whole study population Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: unclear whether any women were considered high risk</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
<p><b>Study dates</b></p> <p>November 2002 - November 2003</p> <p><b>Source of funding</b></p> <p>Not reported</p>				<p>LR+: 1.64 (0.56 to 4.80) LR-: 0.72 (0.26 to 1.95)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 437 1731 738"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>20</td> </tr> <tr> <td>Predictive Test -ve</td> <td>3</td> <td>45</td> </tr> </tbody> </table> <p><b>Cord pH</b></p> <table border="1" data-bbox="1346 850 1731 1152"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>7</td> </tr> <tr> <td>Predictive Test -ve</td> <td>3</td> <td>22</td> </tr> </tbody> </table> <p><b>Apgar score</b></p>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	20	Predictive Test -ve	3	45		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	7	Predictive Test -ve	3	22	<p><b>Other information</b></p> <p>Authors' definition of positive stimulation test: no acceleration</p> <p>43/54 (80%) had one scalp sampling, 6/54 (11%) had two and 5/54 (9%) had 3, giving a total of 70 FBS procedures.</p> <p>48/54 (89%) of women were in the first stage of labour with dilatation ranging from 5 to 9cm; 6/54 (11%) were at full dilatation.</p> <p>The five acidotic fetuses were all delivered within 30 minutes of scalp blood sampling; 4 by caesarean section and one by instrumental delivery.</p> <p>Cord pH data were not available for 16 fetuses; 7/16 had a positive FSS test result (no CTG acceleration), 9/16 had a negative FSS results (CTG acceleration)</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	2	20																					
Predictive Test -ve	3	45																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	2	7																					
Predictive Test -ve	3	22																					

Bibliographic details	Participants	Tests	Methods	Outcomes and results		Comments	
				Reference Test +ve	Reference Test -ve		
				Predictive Test +ve	2	14	
				Predictive Test -ve	2	32	
<p><b>Full citation</b></p> <p>Umstad,M., Bailey,C., Permezel,M., Intrapartum fetal stimulation testing, Australian and New Zealand Journal of Obstetrics and Gynaecology, 32, 222-224, 1992</p> <p><b>Ref Id</b></p> <p>201865</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Aim of the study</b></p> <p>To evaluate the usefulness of intrapartum fetal stimulation tests in routine clinical practice</p>	<p><b>Sample size</b></p> <p>N = 60</p> <p><b>Characteristics</b></p> <p>All fetuses were at least 36 weeks' gestation</p> <p><b>Inclusion Criteria</b></p> <p>Fetal heart rate (FHR) tracing significantly abnormal such that fetal scalp blood sampling (FBS) was indicated</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>3 seconds of fetal vibroacoustic stimulation (VAS) followed by the incision of FBS serving as fetal scalp stimulation</p>	<p><b>Methods</b></p> <p>Several minutes prior to FBS, a 3-second VAS was applied over the fetal head via a Corometrics Fetal Acoustic Stimulator (model 146), which generates a sound level of 82 db at 1 m in air.</p> <p>FBS was performed in the usual manner in either lithotomy (with appropriate tilt) or left lateral positions. A Corometrics Model 220 pH Analyzer was used to assess pH of both fetal capillary and umbilical artery blood samples.</p> <p>FHR traces were reported by one of the study authors who was blinded to the results of FBS, Apgar scores, mode of delivery and umbilical artery cord pH values. A reactive FHR response was defined as an acceleration <math>\geq</math> 15bpm for <math>\geq</math></p>	<p><b>Results</b></p> <p><b>Prevalence of acidosis &lt; 7.20</b></p> <p>8/60 (13%)</p> <p><b>Prevalence of acidosis &lt; 7.25</b></p> <p>23/60 (38%)</p> <p><b>Predictive value of no FHR acceleration following VAS</b></p> <p><b>a. For fetal blood sample pH &lt; 7.20</b></p> <p>As reported in Table 4; NCC calculated LR+, LR- and all confidence intervals</p> <p>Sensitivity:100% (100 to 100)</p> <p>Specificity: 59.6% (46.28 to 72.95)</p> <p>PPV: 27.6% (11.32 to 43.85)</p> <p>NPV: 100% (100 to 100)</p> <p>LR+: 2.48 (1.78 to 3.45)</p> <p>LR-: 0 (NC)</p> <p><b>b. For fetal blood sample pH &lt; 7.25</b></p> <p>As reported in Table 3; NCC calculated LR+, LR- and all confidence intervals</p> <p>Sensitivity: 100% (100 to 100)</p> <p>Specificity: 83.8% (71.91 to 95.66)</p> <p>PPV: 79.3% (64.57 to 94.05)</p> <p>NPV: 100% (100 to 100)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: yes</p> <p>Loss to follow-up is unrelated to key characteristics: no loss to follow up</p> <p>Prognostic factor is adequately measured in participants: yes - assessor blinded to outcome</p> <p>Outcome of interest is sufficiently measured in participants: yes</p> <p>Important potential confounders are accounted for: time between FBS and delivery not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p>Indirectness: unclear whether any women were considered high risk</p> <p><b>Other information</b></p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments						
<p><b>Study type</b></p> <p><b>Study dates</b></p> <p>6-month period (dates not reported)</p> <p><b>Source of funding</b></p> <p>The Royal Women's Hospital/3AW Clinical Research Foundation</p>			<p>15 seconds occurring within 60 seconds of the stimulus.</p> <p>Fetal scalp stimulation responses were assessed by determining the reaction to fetal scalp puncture with the guarded scalpel blade during FBS.</p>	<p>LR+: 6.17 (2.96 to 12.83) LR-: 0 (NC)</p> <p><b><u>Predictive value of no FHR acceleration following fetal scalp stimulation (scalp puncture)</u></b></p> <p><b>c. For fetal blood sample pH &lt; 7.20</b> As reported in Table 6 NCC calculated LR+, LR- and all confidence intervals Sensitivity: 62.5% (28.95 to 96.05) Specificity: 67.3% (54.56 to 80.06) PPV: 22.7% (5.22 to 40.24) NPV: 92.1% (83.53 to 101) LR+: 1.91 (0.98 to 3.71) LR-: 0.56 (0.22 to 1.39)</p> <p><b>d. For fetal blood sample pH &lt; 7.25</b> As reported in Table 5 NCC calculated LR+, LR- and all confidence intervals Sensitivity: 82.6% (67.12 to 98.10) Specificity: 91.9% (83.10 to 100) PPV: 86.4% (72.02 to 100) NPV: 89.5% (79.72 to 99.23) LR+: 10.19 (3.39 to 30.63) LR-: 0.19 (0.08 to 0.46)</p> <p><b>FBS pH</b></p> <table border="1" data-bbox="1346 1131 1731 1331"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>8</td> <td>21</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	8	21	<p>Authors' definition of positive stimulation test: no acceleration.</p> <p>Results of fetal stimulation tests were not used in the obstetric management of the women.</p>
	Reference Test +ve	Reference Test -ve									
Predictive Test +ve	8	21									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																					
				<table border="1"> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>31</td> </tr> </table> <p>FBS pH</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>23</td> <td>6</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>31</td> </tr> </tbody> </table> <p>FBS pH</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>5</td> <td>17</td> </tr> <tr> <td>Predictive Test -ve</td> <td>3</td> <td>35</td> </tr> </tbody> </table> <p>FBS pH</p>	Predictive Test -ve	0	31		Reference Test +ve	Reference Test -ve	Predictive Test +ve	23	6	Predictive Test -ve	0	31		Reference Test +ve	Reference Test -ve	Predictive Test +ve	5	17	Predictive Test -ve	3	35	
Predictive Test -ve	0	31																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	23	6																								
Predictive Test -ve	0	31																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	5	17																								
Predictive Test -ve	3	35																								

Bibliographic details	Participants	Tests	Methods	Outcomes and results			Comments
					Reference Test +ve	Reference Test -ve	
				Predictive Test +ve	19	3	
				Predictive Test -ve	4	34	



## G.7 Fetal blood sampling as an adjunct to cardiotocography

Study details	Participants	Intervention s	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Alfirevic,Z., Devane,D., Gyte,G.M., Continuous cardiotocography (CTG) as a form of electronic fetal monitoring (EFM) for fetal assessment during labour. [55 refs]Updated, Cochrane Database of Systematic Reviews, 5, CD006066-, 2013</p> <p><b>Ref Id</b></p> <p>65685</p> <p><b>Country/ies where the study was carried out</b></p> <p>Various</p> <p><b>Study type</b></p> <p>Systematic review of RCTs</p> <p><b>Aim of the study</b></p>	<p><b>Sample size</b></p> <p>Total number of studies included n = 13</p> <p>Number of studies reporting outcomes for CTG plus fetal blood sampling (FBS) intervention n = 8</p> <p><b>Characteristics</b></p> <p><b>Athens 1993</b> RCT; randomisation by tossing a coin on admission; women and obstetricians not blinded; neonatologists collecting data on neonatal outcomes were blinded Population: n = 1428 Inclusion: mixed-risk, women with a singleton pregnancy at ≥ weeks' gestation admitted in spontaneous labour or for induction of labour Exclusion: women with known fetal congenital or</p>	<p><b>Intervention s</b></p> <p>Intervention: continuous CTG during labour Control: no fetal heart monitoring Intermittent auscultation of fetal heart rate with Pinard or Doppler Intermittent CTG</p>	<p><b>Details</b></p> <p><u>Electronic searches</u> The Cochrane Pregnancy and Childbirth Group's Trials Register was searched by contacting the Trials Search Coordinator. CENTRAL, MEDLINE, EMBASE were searched and hand searching of journals and conference proceedings was performed. Dissertation abstracts and National Research Register was searched for accessing grey literature. No language restrictions were applied.</p> <p><u>Selection of studies</u> Two review authors independently assessed all potential studies for inclusion. There</p>	<p><b>Results</b></p> <p><b>Thirteen studies were identified and included in the systematic review but only eight (8) studies had CTG plus FBS as an intervention. Therefore outcomes related to those studies are reported here.</b></p> <p><u>Continuous CTG and FBS versus IA</u> <u>Neonatal seizures</u> No. studies: 5 n = 15004 Continuous CTG and FBS n = 7542 IA n = 7462 RR 0.49 (95% CI 0.29 to 0.84)</p> <p><u>Cerebral palsy</u> No. studies: 2 n = 13252 Continuous CTG and FBS n = 6609 IA n = 6643 RR 1.74 (95% CI 0.97 to 3.11)</p>	<p><b>Limitations</b></p> <p><u>Attrition bias reported by the review authors for the included studies</u></p> <p><b>Athens 1993</b> Attrition bias: (A) less than 3% of participants excluded Allocation concealment: no</p> <p><b>Copenhagen 1985</b> Attrition bias: (B) 3% to 9.9% of participants excluded (1061 women agreed to participate; 92 excluded) Allocation concealment: unclear</p> <p><b>Dallas 1986</b> Attrition bias: information not available Allocation concealment: no</p> <p><b>Denver 1976</b> Attrition bias: (A) less than 3% of participants excluded Allocation concealment: unclear</p> <p><b>Denver 1979</b> Attrition bias: (A) less than 3% of participants excluded Allocation concealment: unclear</p> <p><b>Dublin 1985</b> Attrition bias: (A) less than 3% of participants excluded FBS was performed when the duration of labour exceeded 8 hours. This occurred in 77/6474 (1.2%) of women in the CTG arm and 139/6486 (2.1%) of women in the IA arm</p> <p><b>Lund 1994</b> Attrition bias: (A) less than 3% of participants excluded Allocation concealment: unclear</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>To evaluate the effectiveness and safety of continuous cardiotocography (CTG) when used as a method to monitor fetal wellbeing during labour</p> <p><b>Study dates</b></p> <p>Assessed as up-to-date May 2013</p> <p><u>Included studies:</u></p> <p><b>Athens 1993</b> Study period: October 1990 to June 1991</p> <p><b>Copenhagen 1985</b> Study period: January 1981 to January 1982 (date women expected to give birth)</p> <p><b>Dallas 1986</b> Study period: not reported</p> <p><b>Denver 1976</b> Study period: not reported</p> <p><b>Denver 1979</b></p>	<p>chromosomal abnormalities</p> <p>Intervention: continuous CTG without FBS n = 746</p> <p>Comparison: intermittent auscultation (IA) n = 682</p> <p>CTG: external unless trace poor when internal CTG used</p> <p><b>Copenhagen 1985</b> RCT; randomisation by random sampling; method of randomisation unclear</p> <p>Population n = 969 women, high- and low-risk women, only women with diabetes excluded; 3 twin pairs in CTG group and 6 twin pairs in IA group</p> <p>Intervention: continuous CTG in conjunction with FBS (CTG: external or internal) n = 482</p> <p>Comparison: IA n = 487</p> <p><b>Dallas 1986</b> Quasi RCT; randomisation by alternate months; selective monitoring</p>		<p>was no disagreement regarding the eligibility for inclusion that needed to be resolved through consultation with a third person.</p> <p><u>Data extraction and management</u></p> <p>A form was designed to extract data, and two authors extracted them. Data were analysed in RevMan. Where information was unclear, the reviewers attempted to contact the original authors.</p> <p><u>Assessment of risk of bias</u></p> <p>Two review authors independently assessed risk of bias using criteria from the Cochrane Handbook for Systematic Reviews of Interventions:</p> <ul style="list-style-type: none"> <li>- Selection bias (allocation concealment)</li> </ul>	<p><u>Caesarean section</u></p> <p>No. studies: 6 n = 15074</p> <p>Continuous CTG and FBS n = 7582</p> <p>IA n = 7492</p> <p>RR 1.50 (1.10 to 2.06)</p> <p><u>Instrumental vaginal birth</u></p> <p>No. studies: 5 n = 14828</p> <p>Continuous CTG and FBS n = 7460</p> <p>IA n = 7368</p> <p>RR 1.25 (1.13 to 1.38)</p> <p><u>Cord blood acidosis</u></p> <p>No. studies: 1 n = 1075</p> <p>Continuous CTG and FBS n = 540</p> <p>IA n = 535</p> <p>RR 0.45 (0.16 to 1.29)</p> <p><u>Any pharmacological analgesia</u></p> <p>No. studies: 2 n = 828</p> <p>Continuous CTG and FBS n = 482</p> <p>IA n = 367</p>	<p><b>Melbourne 1976</b> Attrition bias: information not available; one obstetrician withdrew his participants from the trial; it was not clear whether this was pre- or post-randomisation nor how many participants were withdrawn</p> <p>Allocation concealment: yes</p> <p><b>Melbourne 1981</b> Attrition bias: (B) 3% to 9.9% of participants excluded</p> <p>Allocation concealment: no</p> <p><b>New Delhi 2006</b> No good information on study methodology</p> <p><b>Pakistan 1989</b> Attrition bias: (A) less than 3% of participants excluded</p> <p>Allocation concealment: no</p> <p>Data extracted from unpublished trial lodged with Cochrane centre</p> <p><b>Seattle 1987</b> Attrition bias: (D) more than 20% of participants excluded</p> <p>Allocation concealment: unclear</p> <p><b>Sheffield 1978</b> Attrition bias: (A) less than 3% of participants excluded</p> <p>Allocation concealment: unclear</p> <p><b>Other information</b></p> <p>The systematic review is available online at: <a href="http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006066.pub2/full">http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD006066.pub2/full</a></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Study period: July 1975 to July 1977</p> <p><b>Dublin 1985</b> Study period: March 1981 to April 1983</p> <p><b>Lund 1994</b> Study period: October 1989 to May 1991</p> <p><b>Melbourne 1976</b> Study period: March 1974 to April 1975</p> <p><b>Melbourne 1981</b> Study period: not reported</p> <p><b>New Delhi 2006</b> No good information on study methodology</p> <p><b>Pakistan 1989</b> Study period: 1988 to 1989</p> <p><b>Seattle 1987</b> Study period: November 1981 to February 1985</p> <p><b>Sheffield 1978</b> Study period: July 1976 to June 1977</p>	<p>(policy of using monitoring only in high-risk pregnancies) versus universal monitoring (use of a monitor for every pregnancy in which the fetus was considered viable, i.e. irrespective of risk status) Population: n = 34,995 women; data were extracted for 14,618 women with low-risk pregnancies; 7288 in universal monitoring group where all women monitored by CTG, and 7330 in selective monitoring where low-risk women monitored by IA Intervention: Continuous CTG (CTG: no information on external or internal) n = 7288 Comparison: IA n = 7330</p> <p><b>Denver 1976</b> RCT; randomised by sealed envelope Population n = 483; high-risk women; those with meconium stained fluid, needing</p>		<p>- Attrition bias - Blinding: lack of blinding was not considered to undermine the validity of the study - Incomplete outcome data - Other sources of bias</p> <p><u>Measures of effect</u> Dichotomous outcomes were presented as a risk ratios (RRs) with 95% confidence intervals (CIs). For continuous data, weighted mean differences and their 95% CI were used.</p> <p><u>Dealing with missing data</u> The review authors investigated the effect of including trials with high levels of attrition using sensitivity analysis. Outcomes were assessed on an intention-to-treat basis, with the denominator being the number randomised minus</p>	RR 0.99 (0.90 to 1.07)	

Study details	Participants	Intervention s	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b> Not reported</p>	<p>oxytocin or abnormal fetal heart tones during labour were eligible to participate Intervention: continuous CTG without FBS versus (CTG: internal) n = 242 Comparison: IA n = 241</p> <p><b>Denver 1979</b> RCT; randomisation by random numbers in sealed envelopes Population: n = 690 high-risk women participating with 5 pairs of twins Intervention 1: continuous CTG with FBS (CTG: external until internal feasible) n = 229 Intervention 2: continuous CTG without FBS (CTG: external until internal feasible) n = 230 Comparison: IA n = 231</p> <p><b>Dublin 1985</b> RCT; randomisation by opaque, sealed envelopes Population: n = 12,964; mixed risked</p>		<p>any participants whose outcomes were known to be missing. For the purpose of the sensitivity analysis 'high quality' was defined as a trial having allocation concealment classified as 'adequate'.</p> <p><u>Analysis</u> If high levels of heterogeneity (&gt; 50%) were identified, prespecified sensitivity analysis was done according to the quality of the trials. A random effects model was used as an overall summary where appropriate.</p> <p>Fixed-effect meta-analysis was used in the absence of substantial heterogeneity between the trials. Random effects meta-analyses were used where heterogeneity was</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>women at &gt; 28 weeks' gestation, in labour; total of 12,964 women participated                      Intervention: continuous CTG in conjunction with FBS versus (CTG: internal) n = 6474                      Comparison: IA n = 6490                      Attrition bias: (A) less than 3% of participants excluded                      Study period: March 1981 to April 1983</p> <p><b>Lund 1994</b>                      RCT; randomisation by shuffled opaque envelopes                      Population: n = 4044 women with low to moderate risk factors during labour                      Intervention: continuous CTG with FBS versus (CTG: no information on external or internal) n = 2029                      Comparison: intermittent CTG with FBS (CTG: no information on external or internal) n = 2015</p>		<p>present or suspected</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><b>Melbourne 1976</b>                      RCT; randomised by cards in sealed numbered envelopes                      Population: n = 350 high-risk women                      Intervention: continuous CTG with FBS (CTG: external) n = 175                      Comparison: intermittent auscultation n = 175</p> <p><b>Melbourne 1981</b>                      RCT; randomisation by cards; envelopes unsealed; biased randomisation in one of the participating hospitals; 62 low-parity women excluded post hoc to correct for inequality in randomisation                      Population: n = 989 low-risk women                      Intervention: continuous CTG without FBS (CTG: external until membranes ruptured then internal) n = 445                      Comparison: intermittent auscultation n = 482</p> <p><b>New Delhi 2006</b>                      RCT; no details on</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>how this was undertaken                      Population: n = 100 women who had had one previous low-transverse caesarean section; for this pregnancy, singleton and cephalic                      Intervention: continuous CTG n = 50                      Comparison: IA n = 50</p> <p><b>Pakistan 1989</b>                      RCT; randomisation by woman selecting a sealed, unnumbered envelope                      Population: n = 200 high-risk women (all participants had meconium stained liquor)                      Intervention: continuous CTG with FBS (external) n = 100                      Comparison: IA n = 100                      Attrition bias: (A) less than 3% of participants excluded                      Study period: 1988 to 1989</p> <p><b>Seattle 1987</b>                      RCT; randomisation</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>by numbered, sealed envelopes                      Population: n = 386 high-risk women                      Preterm labour (28-32 weeks' gestation), estimated fetal weight 700-1750 g                      Intervention:                      continuous CTG with FBS (CTG: external until rupture of membranes then internal) n = 188                      Comparison: IA n = 188</p> <p><b>Sheffield 1978</b>                      RCT; randomisation by sealed envelopes; details not reported                      Population: n = 504 women with mixed risk                      Intervention:                      continuous CTG without FBS versus (CTG: internal) n = 253                      Comparison: IA n = 251</p> <p><b>Inclusion criteria</b></p> <p>Randomised and quasi randomised studies comparing continuous</p>				



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>cardiotocography (CTG) with or without fetal blood sampling (FBS) with a) no fetal monitoring b) intermittent auscultation of the fetal heart rate using a Pinard stethoscope or hand-held Doppler device or intermittent CTG. Studies using less robust methods of allocation (for example, alternation) were not included</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>				
<p><b>Full citation</b></p> <p>Noren,H., Luttkus,A.K., Stupin,J.H., Blad,S., Arulkumaran,S., Erkkola,R., Luzietti,R., Visser,G.H., Yli,B., Rosen,K.G., Fetal scalp pH and ST analysis of the fetal ECG as an adjunct to cardiotocography to</p>	<p><b>Sample size</b></p> <p>Cases n = 97 (marked acidosis n = 53, moderate acidaemia n = 44) Control n = 97</p> <p><b>Characteristics</b></p> <p>There were statistically significant differences observed</p>	<p><b>Interventions</b></p> <p>STAN analysis plus electronic fetal monitoring (EFM) plus FBS</p>	<p><b>Details</b></p> <p>From a European Union multicentre study on clinical implementation of STAN methodology, 911 cases were identified where a scalp pH had been obtained. A total of n = 6999 cases were recorded during the study</p>	<p><b>Results</b></p> <p><u>Time between onset of significant ST events (FHR plus ST indication to intervene) and birth</u>  <u>FHR+ST events recorded within 16 minutes of birth</u>  <u>(cord artery pH ≥</u></p>	<p><b>Limitations</b></p> <p>Data from a previously published study used. Not clear how the observers assessed the data. Results reported poorly and inconsistently</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>predict fetal acidosis in labor--a multi-center, case controlled study, Journal of Perinatal Medicine, 35, 408-414, 2007</p> <p><b>Ref Id</b></p> <p>121268</p> <p><b>Country/ies where the study was carried out</b></p> <p>Norway</p> <p><b>Study type</b></p> <p>Retrospective cohort</p> <p><b>Aim of the study</b></p> <p>To assess the relationship between scalp pH (fetal blood sampling [FBS]) and ST analysis in situations of acidosis with special emphasis on the timing of cardiotocography (CTG), FBS and ST changes during labour</p>	<p>in two groups (cases and controls) on antenatal factors, primigravidae and cord pH. Significantly more operative births were observed in marked acidosis and moderate acadaemia cases compared with controls. Admission to neonatal care unit was significantly higher in marked acidosis cases compared with the matched control</p> <p><b>Inclusion criteria</b></p> <p>Pregnancy &gt; 36 weeks, high-risk pregnancy, women with suspicious or abnormal external CTG, induced or oxytocin-augmented labour or meconium stained liquor</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>period in maternity units and 911 cases were identified where a FBS was performed. Each ward had a research midwife responsible for education and data collection. The decision for need of FBS was left to the clinician in charge and time and pH reading was recorded.</p> <p>In 53 cases, marked cord artery acidosis was found (cord artery pH &lt; 7.06) and 44 cases showed moderate acidaemia at birth (pH 7.06-7.09). Comparisons were made with 97 control cases (pH ≥ 7.20).</p> <p><u>Intervention:</u></p> <p>Clinical management was guided by CTG interpretation supported by computerised ST waveform assessment (ST log) and or FBS</p>	<p><u>7.20</u> n = 17/28(61%)</p> <p><u>STAN indications recorded &gt;16 minutes (cord artery pH ≥ 7.20)</u> n = 13/69 (19%) OR 6.66 (2.53 to 17.55) P &lt; 0.001</p> <p><b><u>Distribution of FBS and ST guideline indication to intervene (marked acidosis)</u></b></p> <p><u>Women with abnormal FBS</u> Marked acidosis n = 24/53 (45%) Control n = 4/53 (7.5%)</p> <p><u>Number of samples with scalp pH &gt; 7.19</u> Marked acidosis n = 43 Control n = 53</p> <p><u>Number of samples with scalp pH 7.15 - 7.19</u> Marked acidosis n</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study dates</b></p> <p>October 2000 to June 2002</p> <p><b>Source of funding</b></p> <p>Not reported</p>			<p>according to the study protocol. The ST log automatically notified the staff if any ST events occurred and intervention was required in case of combined CTG and ST changes. Intervention was also indicated by occurrence of preterminal CTG (complete loss of variability and reactivity). No intervention was recommended if CTG was normal, irrespective of the ST. During the first stage of labour identification and alleviation of the cause of hypoxia was the intervention. If that was not possible operative birth was recommended. In the second stage of labour, if the ST changes appeared, immediate birth was recommended. In the event of</p>	<p>= 6 Control n = 1</p> <p><u>Number of samples with scalp pH &lt; 7.15</u> Marked acidosis n s 21 Control n = 3</p> <p><u>Number of adequately monitored</u> Marked acidosis n = 46/53 (86.8%) Control n = 42/53 (79.2%)</p> <p><u>ST indication</u> Marked acidosis n = 41/53 (77.4%) Control n = 20/53 (37.7%)</p> <p><u>No ST indication (adequately monitored)</u> Marked acidosis n = 5/46 (11%) Control n = 22/42 (52.4%)</p> <p><b><u>Distribution of FBS and ST guideline indication to intervene (moderate acidaemia)</u></b></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>abnormal CTG and normal ST during the second stage of labour, a maximum of 90 minutes was recommended before birth. FBS was optional during the first and second stages of labour. In the cases with no indication to intervene, the recording continued until the birth.</p> <p><u>Analysis:</u> The results were evaluated with medical statistical software. Student's t test or Mann-Whitney test were used for testing continuous variables. Fisher's exact test was used for discrete variables</p>	<p><u>Women with abnormal FBS</u> Moderate acidaemia n = 24/53 (45%) Control n = 4/53 (7.5%)</p> <p><u>Number of samples with scalp pH &gt; 7.19</u> Moderate acidaemia n = 57 Control n = 61</p> <p><u>Number of samples with scalp pH 7.15 - 7.19</u> Moderate acidaemia n = 10 Control n = 0</p> <p><u>Number of samples with scalp pH &lt; 7.15</u> Moderate acidaemia n = 13 Control n = 0</p> <p><u>Number of adequately monitored</u> Moderate acidaemia n = 40/44 (91%) Control n = 32/44 (72.7%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><u>ST indication</u>            Moderate            acidaemia n = 24/44 (54.5%)            Control n = 10/44 (22.7%)</p> <p><u>No ST indication (adequately monitored)</u>            Moderate            acidaemia n = 16/40 (40%)            Control n = 22/32 (68.8%)</p> <p><b><u>Cases with abnormal CTG and their relation to FBS and ST</u></b>  <u>Abnormal CTG patterns</u>            Normal ST n = 60/121 (49.6%)            Abnormal ST n = 61/121 (50.4%)</p> <p><u>Cases with an abnormal CTG and cord artery pH &lt; 7.10</u>            n = 84/121 (69%):            Abnormal ST n = 70/84 (83%)</p> <p><u>Abnormal FBS (&lt; 7.20)</u>            Normal ST n = 7*/60 (11.7%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>Abnormal ST n = 29/61 (47.5%)</p> <p><u>Normal FBS</u> Normal ST n = 50/60 (83.3%) Abnormal ST n = 12†/61 (19.7%)</p> <p><u>No FBS in connection with abnormal CTG</u> Normal ST n = 3‡/60 (5%) Abnormal ST n = 20/61 (32.8%)</p> <p>*All had FBS taken in the second stage of labour; n = 6 had respiratory acidosis with normal neonatal period; n = 1 had cord pH &gt;= 7.20 †n = 5/12 developed acidosis subsequently and n = 7 had a normal cord acid base ‡All developed acidosis</p> <p><b><u>FBS and ST indication of abnormality in cases with CTG changes noted at</u></b></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><b><u>the start of ST recording</u></b></p> <p><u>Total ST findings with normal FBS</u>                      Normal ST n = 43/44 (97.7%)                      Abnormal ST n = 1/44 (2.3%)</p> <p><u>Total ST findings with abnormal FBS</u>                      Normal ST n = 3/17 (17.6%)                      Abnormal ST n = 14/17 (82.4%)</p> <p><u>ST findings with normal FBS (marked acidosis)</u>                      Normal ST n = 14*/14 (100%)                      Abnormal ST n = 0/14 (0%)</p> <p><u>Total ST findings with abnormal FBS (marked acidosis)</u>                      Normal ST n = 2/7 (28.6%)                      Abnormal ST n = 5/7 (71.4%)</p> <p><u>ST findings with normal FBS (marked acidaemia)</u>                      Normal ST n = 29†/30 (96.7%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>Abnormal ST n = 1/30 (3.3%)</p> <p><u>ST findings with abnormal FBS (marked acidaemia)</u></p> <p>Normal ST n =1/10 (10%) Abnormal ST n = 9/10 (90%)</p> <p><u>Special care baby unit was associated with low Apgar scores (&lt; 7 at 5 minutes)</u></p> <p>Marked acidosis: 15/26 (58%) Moderate acidosis: 4/14 (26%) The corresponding rate for control group was 1 of 12 (8%)</p> <p>* n =11/14 subsequently developed ST changes and those that did not, ST changes were inadequately recorded † n = 2 developed</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				subsequent ST changes	
<p><b>Full citation</b></p> <p>Stein,W., Hellmeyer,L., Misselwitz,B., Schmidt,S., Impact of fetal blood sampling on vaginal delivery and neonatal outcome in deliveries complicated by pathologic fetal heart rate: a population based cohort study, Journal of Perinatal Medicine, 34, 479-483, 2006</p> <p><b>Ref Id</b></p> <p>121315</p> <p><b>Country/ies where the study was carried out</b></p> <p>Germany</p> <p><b>Study type</b></p> <p>Population-based cohort study</p>	<p><b>Sample size</b></p> <p>n = 49,560 births, 26% underwent FBS</p> <p><b>Characteristics</b></p> <p>No significant differences observed between the two groups in neonatal sex, birthweight &lt; 2.5 kg, birthweight &gt; 4 kg and maternal risk in pregnancy. Gestational age &gt; 40 weeks, maternal age &gt; 35 years, and additional risk factors at birth were significantly associated with FBS</p> <p><b>Inclusion criteria</b></p> <p>Pathologic fetal heart rate</p> <p>Singleton pregnancy</p>	<p><b>Interventions</b></p> <p>EFM plus FBS</p>	<p><b>Details</b></p> <p><u>Data collection</u> Data about the woman, pregnancy and birth were collected from the perinatal birth register of Hense, using an evaluated 76 item questionnaire. From 1990 to 2000, the perineal birth register of Hense recorded data of 589,609 births &gt; 35 weeks. Of these, 49,450 births fulfilled the inclusion criteria.</p> <p><u>Analysis</u> Bivariate analyses between the usage of FBS and the characteristics of the newborn, woman and birth were performed only on those records with no missing values for</p>	<p><b>Results</b></p> <p><u>Spontaneous birth (no presence of additional risk factor)</u> EFM + FBS n = 2191 (82%) EFM alone n = 7678 (76.7%) OR 1.41 (95% CI 1.27 to 1.58)</p> <p><u>Spontaneous birth (in presence of additional risk factor)</u> EFM + FBS n = 5912 (57.8%) EFM alone n = 13974 (52.4%) OR 1.24 (95% CI 1.19 to 1.30)</p> <p><u>Vaginal assisted birth (no presence of additional risk factor)</u> EFM + FBS n = 472 (16.8%) EFM alone n = 2336 (23.3%) OR not reported</p>	<p><b>Limitations</b></p> <p>Choice of treatment unrelated to confounders (selection bias): unclear Groups comparable at baseline: unclear Groups received same/similar care (apart from intervention): unclear Blinding of those assessing outcomes: no Missing data/loss to follow-up: unclear Precise definition of outcomes: yes Valid and reliable method of outcome assessment: unclear Intention-to-treat analysis performed: no</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To compare the impact of electronic fetal monitoring (EFM) alone versus EFM with additional fetal blood sampling (FBS) in vaginal births complicated by pathologic fetal heart rate (FHR)</p> <p><b>Study dates</b></p> <p>All births in Hesse between 1990 and 2000</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>Vaginal birth</p> <p>Cephalic presentation</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>any maternal covariates. To assess the effect of FBS in the births with pathological FHR on mode of birth and neonatal outcomes, univariate regression analysis was performed and odds ratios (ORs) and their corresponding 95% confidence intervals (95% CIs) were calculated</p>	<p><u>Vaginal assisted birth (in presence of additional risk factor)</u>                      EFM + FBS n = 4318 (42.2%)                      EFM alone n = 12679 (47.6%)                      OR not reported</p> <p><b><u>Neonatal outcomes</u></b></p> <p><u>Severe fetal acidosis (umbilical artery pH &lt; 7.0)</u>                      EFM + FBS n = 64 (0.5%)                      EFM alone n = 307 (0.91%)                      OR 0.55 (95% CI 0.42 to 0.72)</p> <p><u>Apgar score &lt; 5 after 7 minutes</u>                      EFM + FBS n = 78 (0.61%)                      EFM alone n = 314 (0.86%)                      OR 0.71 (95% CI 0.55 to 0.90)</p> <p><u>Admission to neonatal unit</u>                      EFM + FBS n = 1025 (8.0%)                      EFM alone n =</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				3220 (8.8%) OR 0.90 (95% CI 0.83 to 0.96)  <u>Reanimation</u> EFM + FBS n = 652 (5.1%) EFM alone n = 3220 (8.8%) OR 0.80 (95% CI 0.73 to 0.88)	
<b>Full citation</b> Becker, J.H., Westerhuis, M.E., Sterrenburg, K., van den Akker, E.S., van Beek, E., Bolte, A.C., van Dessel, T.J., Drogdrop, A.P., van Geijn, H.P., Graziosi, G.C., van Lith, J.M., Mol, B.W., Moons, K.G., Nijhuis, J.G., Oei, S.G., Oosterbaan, H.P., Porath, M.M., Rijnders, R.J., Schuitemaker, N.W., Wijnberger, L.D., Willekes, C., Visser, G.H., Kwee, A., Fetal blood sampling in addition	<b>Sample size</b> At least one FBS performed for n = 301 women; n = 224 complete ST recordings were available for assessment  <b>Characteristics</b> Not reported  <b>Inclusion criteria</b> Women in labour with a high-risk singleton pregnancy in cephalic position at term	<b>Interventions</b> FBS in conjunction with electronic fetal monitoring (EFM) and ST wave analysis	<b>Details</b> Data were used from women monitored in the STAN arm of a previously published multicentre randomised controlled trial; participants had been randomly assigned to monitoring by cardiotocography (CTG) combined with ST-analysis of the fetal electrocardiogram (ECG; index group) or CTG without ST-analysis (control group).	<b>Results</b> <b><u>FBS in births monitored by ST-analysis of the fetal ECG related to the trial protocol</u></b> <b><u>Number of FBS</u></b> According to trial protocol n = 171 Not according to trial protocol n = 126  <b><u>pH &gt; 7.25</u></b> According to trial protocol n = 112/171 (65.5%) Not according to trial protocol n = 96/126 (76.2%)  <b><u>pH 7.20 - 7.25</u></b> According to trial	<b>Limitations</b> A large number of women in whom at least one FBS was performed were excluded from the analysis for various reasons that were not reported. Data from a previously published trial were used  <b>Other information</b>

Study details	Participants	Intervention s	Methods	Outcomes and Results	Comments
<p>to intrapartum ST-analysis of the fetal electrocardiogram: evaluation of the recommendations in the Dutch STAN[REGISTERED] trial, BJOG: An International Journal of Obstetrics and Gynaecology, 118, 1239-1246, 2011</p> <p><b>Ref Id</b> 156994</p> <p><b>Country/ies where the study was carried out</b> Netherlands</p> <p><b>Study type</b> Prospective cohort study</p> <p><b>Aim of the study</b> To evaluate recommendations for additional fetal blood sampling (FBS) when using ST-</p>	<p><b>Exclusion criteria</b> Not reported</p>		<p>This study was on the women randomised to the index group in whom FBS was undertaken. In women in the index group, a scalp electrode was applied to the fetal head and connected to a STAN S21 or S31 fetal heart monitor (Neovinta Medical, Gothenburg, Sweden). Clinical management was guided by the STAN clinical guidelines. In the study protocol FBS was recommended in three situations: (1) start of STAN registration with an intermediary or abnormal CTG trace (2) abnormal CTG trace for more than 60 minutes without ST-events (3) poor ECG signal quality in the presence of an intermediary or abnormal CTG trace.</p>	<p>protocol n = 33/171 (19.3%) Not according to trial protocol n = 15/126 (12%)</p> <p><u>pH &lt; 7.20</u> According to trial protocol n = 17/171 (10%) Not according to trial protocol n = 10/126 (7.9%)</p> <p><u>Missing pH</u> According to trial protocol n = 9/171 (5.3%) Not according to trial protocol n = 5/126 (4%)</p> <p><b><u>FBS in births monitored by ST-analysis of the fetal ECG related to reasons according to the trial protocol</u></b> <u>Number of FBS</u> Total n = 171 Abnormal CTG (cardiotocography) at start n = 18 Intermediary CTG at start n = 9 Abnormal CTG &gt; 60 min without ST events n = 111</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>analysis of the fetal electrocardiogram</p> <p><b>Study dates</b></p> <p>January 2006 to July 2008</p> <p><b>Source of funding</b></p> <p>Funded by a grant from ZonMW, the Dutch Organisation for Health Research and Development</p>			<p>Poor signal quality was defined as absence of ST-information for more than 4 minutes or less than one average ECG-complex per minute within a period of 10 minutes. If FBS showed a pH &lt; 7.20, an immediate birth was advised. If the pH was between 7.20 and 7.25 the advice was to repeat FBS after 30 minutes. If the pH was &gt; 7.25, the fetal condition was considered well enough to continue labour. Presence of STAN abnormalities (defined in the protocol) was also an indication for immediate birth.</p> <p><u>Data analysis</u></p> <p>All STAN recordings of women in the index group in which at least one FBS was performed were assessed by two observers</p>	<p>Poor ECG signal quality n = 33</p> <p><u>pH &gt; 7.25</u></p> <p>Total n = 112</p> <p>Abnormal CTG at start n = 9</p> <p>Intermediary CTG at start n = 9</p> <p>Abnormal CTG &gt; 60 min without ST events n = 69</p> <p>Poor ECG signal quality n = 25</p> <p><u>pH 7.20 - 7.25</u></p> <p>Total n = 33</p> <p>Abnormal CTG at start n = 5</p> <p>Intermediary CTG at start n = 0</p> <p>Abnormal CTG &gt; 60 min without ST events n = 24</p> <p>Poor ECG signal quality n = 4</p> <p><u>pH &lt; 7.20</u></p> <p>Total n = 17</p> <p>Abnormal CTG at start n = 2</p> <p>Intermediary CTG at start n = 0</p> <p>Abnormal CTG &gt; 60 min without ST events n = 12</p> <p>Poor ECG signal quality n = 3</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>who examined whether or not additional FBS was performed according to the trial protocol. When there was disagreement, the opinion of a third observer was decisive. The observers were only provided with information on the timing of FBS, without knowledge of its result, other clinical parameters obtained during labour, or the neonatal outcome. For each FBS the following items had to be scored:</p> <p>(1) classification of the CTG as normal, intermediary, abnormal or (pre)terminal within a 60-minute period before performance of FBS</p> <p>(2) duration of an intermediary, abnormal or (pre)terminal CTG in minutes</p> <p>(3) interpretation of any ST-events; and</p>	<p><u>Missing pH</u> Total n = 9 Abnormal CTG at start n = 2 Intermediary CTG at start n = 0 Abnormal CTG &gt; 60 min without ST events n = 6 Poor ECG signal quality n = 1</p> <p><b><u>Relation of presence or absence of significant ST-events and preterminal CTG with results of FBS not taken according to protocol</u></b></p> <p><u>Indication to intervene (at least on significant ST events) Total n = 34</u> pH &lt; 7.20 n = 8 (23.5%) pH 7.20 - 7.25 n = 5 (14.7%) pH &gt; 7.25 n = 19 (60%) Missing value n = 2 (5.9%)</p> <p><u>No indication to intervene (total n = 92)</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>(4) judgement of whether FBS was performed according to the randomised controlled trial protocol.</p> <p>Observers evaluated whether the FBS was performed according to the trial protocol, and assessed the relation between pH result measured by FBS and the reason to perform FBS was described.</p> <p>In the cases of protocol violation (FBS not performed according to the trial protocol) the relation between pH results of FBS and ST-waveform interpretation regarding fetal indications to intervene, was evaluated. Fetal acidosis was defined as an FBS pH &lt; 7.20. Women were classified as being treated 'not</p>	<p>pH &lt; 7.20 n = 2 (2.2%)                      pH 7.20 - 7.25 n = 10 (11%)                      pH &gt; 7.25 n = 77 (83.7%)                      Missing value n = 3 (3.2%)</p> <p><u>Preterminal CTG (total n = 1)</u>                      pH &lt; 7.20 n = 1 (100%)                      pH 7.20 - 7.25 n = 0                      pH &gt; 7.25 n = 0                      Missing value n = 0</p> <p><b><u>Neonatal outcomes</u></b>  <b>FBS was taken according to the trial protocol</b>  <u>Neonates with metabolic acidosis at birth</u>                      n = 3                      One out of the three women had abnormal CTG for 36 minutes plus poor ECG quality before FBS with pH 7.9. In the other women (n=2), FBS was performed because of abnormal CTG &gt; 60 minutes and</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>according to trial protocol' if at least one of the FBSs was not performed according to the trial protocol. Metabolic acidosis for neonates was defined as an umbilical cord artery pH &lt; 7.05 and base deficit &gt; 12 mmol/l</p>	<p>result of FBS was normal but CTG abnormalities persisted. For one woman the time between FBS and birth was only 20 minutes; in the other it was 9 hours with an abnormal CTG for the last 115 minutes (FBS pH 7.32, umbilical cord artery pH 6.93)</p> <p><b>FBS was performed not according to the trial protocol</b>  <u>Neonates with metabolic acidosis at birth</u>                      n = 3                      In all three women earlier intervention was recommended based on significant ST-events. In one of these women multiple FBSs were performed because of an abnormal CTG-pattern (pH 7.38, 7.33, 7.31, 7.28 and 7.28). The final two FBSs were both</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>preceded by a significant ST-event. Abnormalities on CTG persisted thereafter and ST-analysis showed one more significant ST-event 76 minutes after the last FBS, during the second stage of labour. The time between the last FBS and birth was 114 minutes; after a failed vacuum extraction, caesarean section was performed and the baby was born with cord pH 6.95 and died because of severe asphyxia and encephalopathy</p>	

## G.8 Fetal blood sampling – time to result

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Annappa,R., Campbell,D.J., Simpson,N.A., Fetal blood sampling in labour and the decision to delivery interval, European Journal of Obstetrics, Gynecology, and Reproductive Biology, 141, 10-12, 2008</p> <p><b>Ref Id</b></p> <p>92285</p> <p><b>Country/ies where the study was carried out</b></p> <p>England</p> <p><b>Study type</b></p> <p>Prospective case series of consecutive attempts at fetal blood sampling (FBS)</p> <p><b>Aim of the study</b></p> <p>To determine the time interval from the decision to the result for fetal blood sampling (FBS) and the time from an abnormal pH to the birth of the baby</p> <p><b>Study dates</b></p> <p>April 1st 2006 to August 1st 2006</p>	<p><b>Sample size</b></p> <p>N = 107</p> <p>(This was the number of attempts to do FBS, involving 72 women)</p> <p><b>Characteristics</b></p> <p><b><u>BMI (n/total (%))</u></b></p> <p>≤ 25: 44/72 (61.1)</p> <p>&gt; 25: 28/72 (38.9)</p> <p><b><u>Cervical dilatation in cm (n/total (%))</u></b></p> <p>≤ 5: 27/72 (37.5)</p> <p>&gt; 5: 45/72 (62.5)</p> <p><b><u>Operator grade (n/total (%))</u></b></p> <p>SHO/SSHO: 41/72 (56.9)</p> <p>SPR/Senior Registrar: 31/72 (43.1)</p> <p><b>Inclusion criteria</b></p> <p>Consecutive attempts at FBS</p>	<p><b>Interventions</b></p> <p>Fetal blood sampling</p>	<p><b>Details</b></p> <p>Consecutive attempts at FBS over the study period were reported. Operators performed the procedure with women in either lithotomy or left lateral position. Fetal capillary blood samples were collected in a heparinised glass tube and analysed using a Bayer Rapid Lab 840 blood gas analyser.</p> <p>All details were recorded in a document designed for this audit. If a sample was taken but judged to be inadequate, another sample was taken; 107 attempts yielded 177 samples due to the need for repeat samples. The time interval was taken from the decision to perform FBS to the result of a successfully attained sample.</p> <p>Non-parametric tests were used for the analysis. The time from the decision to the result was compared for each factor using Mann-Whitney tests. Regression analysis was undertaken to investigate the factor, while controlling for other factors</p>	<p><b>Results</b></p> <p><b><u>Time from decision to the result of the FBS</u></b></p> <p>a. Median/minutes (IQR): 17 (11 - 22)</p> <p>b. Time taken &gt; 30 minutes (n/total (%)): 5/107 (4.7)</p> <p>[Note: the median time for preparation was 8 minutes (IQR 7 - 15), and the median time to perform the procedure was 10 minutes (IQR 9 - 16)]</p> <p><b><u>Factors affecting the time interval between decision to result of FBS/minutes (median (IQR))</u></b></p> <p>a. <b>BMI</b></p> <p>≤ 25: 13 (11 - 17)</p> <p>&gt; 25: 17 (14 - 22)</p> <p>(p &lt; 0.001)</p> <p>b. <b>Cervical dilatation</b></p> <p>≤ 5: 22 (16 - 25)</p> <p>&gt; 5: 15 (10 - 17)</p> <p>(p &lt; 0.0001)</p> <p>c. <b>Operator grade</b></p> <p>SHO/SSHO: 17 (17 - 22)</p> <p>SPR/Senior Registrar: 13 (10 - 17)</p> <p>(p &lt; 0.001)</p>	<p><b>Limitations</b></p> <p>Inclusion or exclusion criteria and characteristics of the study population were not reported in detail; therefore, it is not possible to establish whether women had low-risk pregnancies</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p> <p>None reported</p>	<p><b>Exclusion criteria</b></p> <p>None reported</p>			<p>These were all independent predictors in the regression model, when including all factors. No valid comparisons for position or epidural could be performed because 95% of women had epidural and 95% of women had FBS taken in the left lateral position</p> <p><b>Number of samples needed (n)</b>                      One: 46                      Two: 52                      Three: 9                      Failed to obtain sample: 2</p> <p>(Note: 23/177 (13%) of samples were inadequate for analysis)</p>	
<p><b>Full citation</b></p> <p>Tuffnell,D., Haw,W.L., Wilkinson,K., How long does a fetal scalp blood sample take?, BJOG: An International Journal of Obstetrics and Gynaecology, 113, 332-334, 2006</p> <p><b>Ref Id</b></p> <p>158858</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p>	<p><b>Sample size</b></p> <p>N=74 women and 100 samples</p> <p><b>Characteristics</b></p> <p>No description of the study population</p> <p><b>Inclusion criteria</b></p> <p>A series of 100 consecutive FBSs on vertex-presenting fetuses</p>	<p><b>Interventions</b></p> <p>FBS</p>	<p><b>Details</b></p> <p>The cases, including the timing of each result, were collected daily from the record in the micro blood analyser database. The clinical staff were aware of the audit and recorded time of decision to perform the test, the time the procedure was started and the operator grade. The operator also recorded the number of attempts for each FBS in the case notes. Those women in whom an FBS was attempted but an inadequate sample obtained were also included in the analysis</p>	<p><b>Results</b></p> <p>100 fetal scalp pH results on 74 babies were reviewed; 89 were successful and 11 were inadequate for the analysis. The median time interval between decision to perform the test and the results was 18 min (IQR 12–25). In 35 (39.5%) of the successful FBS, the time taken was &gt; than 20 minutes, and in eight (9%), it took &gt; than 30 minutes</p>	<p><b>Limitations</b></p> <p>Inclusion or exclusion criteria and characteristics of the study population were not reported in detail; therefore, it is not possible to establish whether women had low-risk pregnancies</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Case series (consecutive, prospective)</p> <p><b>Aim of the study</b></p> <p>To identify the time from a decision to perform a fetal blood sample (FBS) to the result of the test being available</p> <p><b>Study dates</b></p> <p>May 2004 to September 2004</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><b>Exclusion criteria</b></p> <p>Not reported</p>				
<p><b>Full citation</b></p> <p>Rimmer, S., Roberts, S. A., Heazell, A. E., Cervical dilatation and grade of doctor affects the interval between decision and result of fetal scalp blood sampling in labour, <i>Journal of Maternal-Fetal &amp; Neonatal Medicine</i>, 29, 2671-4, 2016</p> <p><b>Ref Id</b></p> <p>451292</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N=119 (n=207 procedures); n=112 (199) included in the analysis</p> <p><b>Characteristics</b></p> <p>No description of the study population</p> <p><b>Inclusion criteria</b></p>	<p><b>Interventions</b></p> <p>FSBS</p>	<p><b>Details</b></p> <p>From women who were eligible, 119 were selected randomly using a computer-generated randomisation list until at least 20 participants had been sampled from each grade of clinician and a minimum of 150 procedures overall. The case notes were identified and relevant information collected from these and the K2 Guardian electronic labour record system (Version 2.050.056.001, K2 Medical Systems, Plymouth, UK) using a standardised proforma. Seven participants for whom complete case notes could</p>	<p><b>Results</b></p> <p>The median time interval from the decision for FSBS to obtaining the result was 10 minutes (range 2–39 minutes). Fifteen samples (7.5%) took <math>\geq 20</math> min to obtain the sample. In four of these cases, the delay resulted from a senior grade of doctor having to perform the procedure after a junior doctor had been unsuccessful</p>	<p><b>Limitations</b></p> <p>Inclusion or exclusion criteria and characteristics of the study population were not reported in detail; therefore, it is not possible to establish whether women had low-risk pregnancies</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>UK</p> <p><b>Study type</b></p> <p>Case series (consecutive, retrospective)</p> <p><b>Aim of the study</b></p> <p>To determine the average time interval between decision to perform a fetal scalp blood sample (FSBS) and obtaining the result in a sufficiently large sample so that other influences on the speed of sampling such as cervical dilatation or grade of operator could be assessed</p> <p><b>Study dates</b></p> <p>April 2013 to May 2014</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p>All women who had a FSBS between April 2013 and May 2014 were eligible</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>not be located were excluded from the study</p>		

## G.9 Predictive value of fetal blood sampling

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Full citation</b></p> <p>Bakr,A.F., Al-Abd,M., Karkour,T., Fetal pulse oximetry and neonatal outcome: a study in a developing country, Journal of Perinatology, 25, 759-762, 2005</p> <p><b>Ref Id</b></p> <p>121095</p> <p><b>Country/ies where the study was carried out</b></p> <p>Egypt</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>To compare the diagnostic value of fetal pulse oximetry with that of fetal scalp blood gas for an abnormal neonatal outcome in cases with abnormal fetal heart rate tracings</p> <p><b>Study dates</b></p> <p>June 2001 to May 2002</p> <p><b>Source of funding</b></p> <p>None, institutional resources</p>	<p><b>Sample size</b></p> <p>N = 150</p> <p><b>Characteristics</b></p> <p>None reported</p> <p><b>Inclusion Criteria</b></p> <p>Abnormal fetal heart rate tracing (criteria not reported)</p> <p>Complete screening panel (fetal pulse oximetry, fetal scalp blood gas and umbilical cord blood gas)</p> <p><b>Exclusion Criteria</b></p> <p>None reported</p>	<p><b>Tests</b></p> <p>Fetal scalp pH analysis</p>	<p><b>Methods</b></p> <p>Informed consent was given by all participants before enrolment. Routine care was given to all patients. Women were monitored with a fetal oxygen saturation monitor and an average value of 30 minutes reading was calculated. A fetal scalp blood gas was taken. An umbilical cord gas sample was obtained shortly following birth, prior for the baby being moved from the delivery area.</p> <p>Abnormal neonatal outcome was defined as having any of the following:</p> <ul style="list-style-type: none"> <li>- Apgar score <math>\leq 7</math> at 5 minutes</li> <li>- Secondary respiratory distress</li> <li>- Transfer to NICU</li> <li>- Neonatal arterial blood pH <math>\leq 7.15</math></li> <li>- Neonatal death</li> </ul> <p>The diagnostic value of fetal blood sampling (FBS) and fetal pulse oximetry were compared</p>	<p><b>Results</b></p> <p><b><u>Predictive value of pH <math>\leq 7.20</math> (95% CI)</u></b></p> <p><b>a. For umbilical artery pH <math>\leq 7.15</math></b></p> <p>Sensitivity: 72% (58 to 82) Specificity: 53% (42 to 63) PPV: 57% (48 to 65)* [NCC: 51% (40 to 61)] NPV: 43% (35 to 51)* [NCC: 74% (63 to 85)] LR+: 1.54 (1.17 to 2.02)† LR-: 0.53 (0.34 to 0.83)†</p> <p><b>b. For abnormal neonatal outcome</b></p> <p>Sensitivity: 82% (65 to 91) Specificity: 52% (42 to 61) PPV: 57% (48 to 64)* [NCC: 36% (26 to 47)] NPV: 43% (35 to 51)* [NCC: 89% (82 to 97)] LR+: 1.69 (1.33 to 2.16)† LR-: 0.36 (0.18 to 0.71)†</p> <p>* values reported here are as reported in the study; however, the PPV and NPV values do not match the 2x2 data reported in the study. NCC calculations are reported in square brackets following study data. † calculated by the NCC-WCH technical team, as likelihood ratios were not reported in the study</p> <p><b>pH <math>\leq 7.2</math> for UA pH <math>\leq 7.15</math></b></p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear - no characteristics of the study population are reported Loss to follow-up is unrelated to key characteristics: no loss to follow-up Prognostic factors is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: no details about mode of birth or when they intervened are reported Statistical analysis is appropriate for study design: yes</p> <p>For PPV and NPV, calculations reported in the study are not consistent with the 2x2 data that are reported.</p> <p>Indirectness of population: not reported whether women were low risk in pregnancy. Also, it</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
			for their ability to predict umbilical cord blood pH $\leq$ 7.15 and abnormal neonatal outcome. Sensitivity, specificity and predictive values were calculated. (Note: this review deals only with FBS; therefore, data for fetal pulse oximetry are not reported)	<table border="1" data-bbox="1317 272 1751 576"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>43</td> <td>42</td> </tr> <tr> <td>Predictive Test -ve</td> <td>17</td> <td>48</td> </tr> </tbody> </table> <p data-bbox="1317 632 1682 683"><b>pH <math>\leq</math> 7.2 for abnormal neonatal outcome</b></p> <table border="1" data-bbox="1317 715 1751 1010"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>31</td> <td>54</td> </tr> <tr> <td>Predictive Test -ve</td> <td>7</td> <td>58</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	43	42	Predictive Test -ve	17	48		Reference Test +ve	Reference Test -ve	Predictive Test +ve	31	54	Predictive Test -ve	7	58	<p data-bbox="1760 272 2040 523">is likely that some women had an interval of longer than 1 hour between FBS and birth; however, the mean and SD suggest that the vast majority will have been an under an hour which is why the study was included</p> <p data-bbox="1760 619 1973 643"><b>Other information</b></p> <p data-bbox="1760 675 2033 778">The mean time lag between the fetal blood gas analysis and birth was <math>36.7 \pm 15.3</math> minutes.</p>
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	43	42																					
Predictive Test -ve	17	48																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	31	54																					
Predictive Test -ve	7	58																					
<p data-bbox="197 1026 338 1050"><b>Full citation</b></p> <p data-bbox="197 1082 622 1278">East,Christine E., Leader,Leo R., Sheehan,Penelope, Henshall,Naomi E., Colditz,Paul B., Intrapartum fetal scalp lactate sampling for fetal assessment in the presence of a non-reassuring fetal heart rate trace, Cochrane Database of Systematic Reviews, -, 2011</p> <p data-bbox="197 1302 271 1326"><b>Ref Id</b></p>	<p data-bbox="638 1026 786 1050"><b>Sample size</b></p> <p data-bbox="638 1082 875 1193">N = 2 trials N = 3348 mother and baby pairs</p> <p data-bbox="638 1249 819 1273"><b>Characteristics</b></p> <p data-bbox="638 1305 819 1329"><u><b>Westgren 1998</b></u> N = 341 Inclusion criteria:</p>	<p data-bbox="925 1026 999 1050"><b>Tests</b></p> <p data-bbox="925 1082 1016 1225">pH analysis Lactate analysis</p>	<p data-bbox="1032 1026 1137 1050"><b>Methods</b></p> <p data-bbox="1032 1082 1308 1385"><u><b>Searching and identification of studies</b></u> The Trials Search Co-ordinator was contacted to search the Cochrane Pregnancy and Childbirth Group's Trials Register (November 2009). At least 2 review authors independently assessed all potential studies for</p>	<p data-bbox="1317 1026 1406 1050"><b>Results</b></p> <p data-bbox="1317 1082 1751 1361"><u><b>ALL SAMPLES Mode of birth (n/total)</b></u> <u>a. Spontaneous vaginal birth</u> Lactate: 709/1667 pH: 709/1652 RR 0.91 (95% CI 0.67 to 1.24) Heterogeneity: <math>I^2 = 64\%</math> [therefore, random effects model was used] Test for overall effect: <math>Z = 0.62</math> (<math>p = 0.54</math>)</p>	<p data-bbox="1760 1026 1895 1050"><b>Limitations</b></p> <p data-bbox="1760 1082 2040 1361">This systematic review does not have any limitations.  Indirectness: it is unclear whether these women had low risk pregnancies; for most outcomes, time interval between FBS and birth is not reported.</p>																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>151307</p> <p><b>Country/ies where the study was carried out</b></p> <p>Sweden</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>To evaluate the effectiveness and risks of fetal scalp lactate sampling in the assessment of fetal well-being during labour, compared with no testing or alternative testing</p> <p><b>Study dates</b></p> <p>Review content was assessed as up-to-date in February 2010</p> <p><b>Source of funding</b></p> <p>Department of Obstetrics and Gynaecology and Pregnancy Research Centre, Department of Perinatal Medicine, University of Melbourne, Royal Women's Hospital, Australia</p> <p>School of Women's and Children's Health, University of New South Wales, Royal Hospital for Women, Randwick, Australia</p> <p>Perinatal Research Centre, University of Queensland, Royal Brisbane &amp; Women's Hospital, Australia</p>	<p>abnormal fetal heart rate during labour and fetal blood sample (FBS) deemed necessary by the attending physician</p> <p>Interventions: - pH analysis was performed in the delivery ward (35 microlitres using ABL 510) - lactate analysis was performed at bedside (5 microlitres using Lactate card)</p> <p>Cut-off action values: pH &lt; 7.20; lactate 2.9 - 3.09 mmol/l was deemed suspicious, and &gt; 3.08 mmol/l was deemed abnormal. No standard advice was given regarding action, so that clinician would consider whole clinical picture, not just one value</p> <p><b>Wiberg-Itzel 2008</b> N = 3007 randomised; N = 2992 analysed Inclusion criteria: singleton pregnancy, cephalic presentation at 34 or more weeks, clinical indication for fetal scalp blood analysis during labour Post-randomisation</p>		<p>inclusion.</p> <p><b>Data extraction and management</b> A form was designed to extract data. Two review authors did data extraction and data was entered into RevMan and checked for accuracy. If any data was unclear, an attempt was made to contact the study authors to provide details.</p> <p>Two review authors assessed risk of bias using criteria outlined in the Cochrane Handbook:</p> <ul style="list-style-type: none"> <li>- The method used to generate the allocation sequence</li> <li>- Allocation concealment</li> <li>- Blinding</li> <li>- Incomplete outcome data, including attrition and exclusions</li> <li>- Selective reporting bias</li> <li>- Other sources of bias</li> </ul> <p><b>Data analysis</b> Fixed-inverse variance meta-analysis was used for combining data, where the authors judged the trials' populations and methods to be sufficiently similar. Where there was</p>	<p>[2 studies: Westgren 1998; Wiberg-Itzel 2008]</p> <p><b>b. Assisted vaginal birth</b> Lactate: 415/1667 pH: 455/1652</p> <p>RR 0.90 (95% CI 0.81 to 1.01) Heterogeneity: <math>I^2 = 0.0\%</math> Test for overall effect: <math>Z = 1.73</math> (<math>p = 0.084</math>)</p> <p>[2 studies: Westgren 1998; Wiberg-Itzel 2008]</p> <p><b>c. Caesarean section</b> Lactate: 472/1667 pH: 432/1652</p> <p>RR 1.09 (95% CI 0.97 to 1.22) Heterogeneity: <math>I^2 = 0.0\%</math> Test for overall effect: <math>Z = 1.50</math> (<math>p = 0.13</math>)</p> <p>[2 studies: Westgren 1998; Wiberg-Itzel 2008]</p> <p><b>d. Operative delivery for non-reassuring fetal status</b> Lactate: 580/1496 pH: 571/1496</p> <p>RR 1.02 (95% CI 0.93 to 1.11) Heterogeneity: NA Test for overall effect: <math>Z = 0.34</math> (<math>p = 0.74</math>)</p> <p>[1 study: Wiberg-Itzel 2008]</p> <p><b>Neonatal death*</b> Lactate: 0/1496 pH: 3/1496</p>	<p>The following represent the review authors assessment of the risk of bias of the included studies:</p> <p><b>Westgren 1998</b> Adequate sequence generation: unclear, method not reported</p> <p>Adequate allocation concealment: yes</p> <p>Blinding: No blinding of participants; blinding of clinicians not feasible; no blinding of outcome assessors reported</p> <p>Incomplete outcome data: excludes women with protocol violations (n = 1 from lactate group, n = 13 from pH group)</p> <p>Selective reporting: unclear</p> <p>Other bias: unclear</p> <p><b>Wiberg-Itzel 2008</b> Adequate sequence generation: yes</p> <p>Adequate allocation concealment: yes</p> <p>Blinding: No blinding of participants; blinding of</p>



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
	<p>exclusion: multiple pregnancy, gestational age &lt; 34 weeks</p> <p>Interventions: - pH analysis was done using different blood gas analysers - Lactate was measured with the Lactate Pro</p> <p>Cut-off action values: - pH: normal &gt; 7.25, pre-acidaemia 7.21 - 7.25, acidaemia &lt; 7.21 - Lactate: normal &lt; 4.2 mmol/l, pre-acidaemia 4.2 - 4.8 mmol/l, acidaemia &gt; 4.8 mmol/l Following pre-acidaemia, the recommendation was for further sampling 20 - 30 minutes later if no other indications for intervention. Following acidaemia, management decisions were made by the attending clinicians</p> <p><b>Inclusion Criteria</b></p> <p>Published and unpublished randomised and quasi-randomised trials comparing fetal scalp lactate testing with no testing or alternative additional tests (e.g. pH, fetal pulse oximetry) to</p>		<p>suspected clinical or methodological heterogeneity between studies, sufficient to suggest that treatment effects could differ, the authors planned to use random effects meta-analysis. Where substantial heterogeneity was identified in a fixed effects meta-analysis, the analysis was repeated using random effects.</p> <p>There were planned subgroup analyses by stage of labour, gestation, and concurrent use of alternative tests; however, there were not sufficient data to do this.</p>	<p>RR 0.14 (95% CI 0.01 to 2.76) Heterogeneity: NA Test for overall effect: Z = 1.29 (p = 0.20)</p> <p>[1 trial: Wiberg-Itzel 2008]</p> <p>* Based on data reported in the full text of the trial, the causes of death were lung hypoplasia due to diaphragmatic hernia (n = 2) and congenital cardiac fibrosis (n = 1).</p> <p><b><u>Neonatal encephalopathy (n/total)†</u></b> Lactate: 6/1496 pH: 6/1496</p> <p>RR 1.00 (95% CI 0.32 to 3.09) Heterogeneity: NA Test for overall effect: Z = 0.0 (p = 1.0)</p> <p>[1 trial: Wiberg-Itzel 2008]</p> <p>† Based on data reported in the full text of the trial, this was hypoxic ischaemic encephalopathy. In the lactate group, 5 cases were mild and one was moderate. In the pH group, 4 cases were mild and 2 were moderate.</p> <p><b><u>Admission to NICU (n/total)</u></b> Lactate: 167/1496 pH: 164/1496</p> <p>RR 1.02 (95% CI 0.83 to 1.25) Heterogeneity: NA Test for overall effect: Z = 0.17 (p = 0.86)</p> <p>[1 trial: Wiberg-Itzel 2008]</p>	<p>clinicians not feasible; no blinding of outcome assessors reported</p> <p>Incomplete outcome data: There were post-randomisation exclusions for 8 of lactate group (twins n = 7, &lt; 34 weeks n = 5) and 7 of the pH group (twins n = 3, &lt; 34 weeks n = 4). All other data reported by intention to treat, but FBS was not undertaken in all women due to: - sampling or analysis failure (lactate: 18, pH: 155) - rapid delivery, need for expedited delivery, reassuring CTG, withdrew consent, no reason given (lactate: 81, pH: 106) There was incomplete umbilical cord blood gas analysis for the following outcomes: - metabolic acidaemia: lactate group 9%, pH group 12% - pH: lactate group 8%, pH group 12%</p> <p>Selective reporting: unclear</p> <p>Other bias: unclear</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
	<p>evaluate fetal status in the presence of a non-reassuring cardiotocograph (CTG) during labour</p> <p><b>Exclusion Criteria</b></p> <p>None reported</p>			<p><b><u>Apgar score &lt; 7 at 5 minutes (n/total)</u></b>                      Lactate: 50/1667                      pH: 44/1652</p> <p>RR 1.13 (95% CI 0.76 to 1.68)                      Heterogeneity: I<sup>2</sup> = 0.0%                      Test for overall effect: Z = 0.59 (p = 0.56)</p> <p>[2 trials: Westgren 1998; Wiberg-Itzel 2008]</p> <p><b><u>Metabolic acidaemia (umbilical artery pH &lt; 7.05 + base deficit &gt; 12 mmol/l)</u></b>                      Lactate: 44/1360                      pH: 47/1315</p> <p>RR 0.91 (95% CI 0.60 to 1.36)                      Heterogeneity: NA                      Test for overall effect: Z = 0.48 (p = 0.63)</p> <p>[1 trial: Wiberg-Itzel 2008]</p> <p><b><u>Cord blood gas values at birth</u></b>  <b><u>a. Umbilical artery pH &lt; 6.98 (n/total)</u></b>                      Lactate: 4/171                      pH: 8/156</p> <p>RR 0.46 (95% CI 0.14 to 1.49)                      Heterogeneity: NA                      Test for overall effect: Z = 1.30 (p = 0.19)</p> <p>[1 trial: Westgren 1998]</p> <p><b><u>b. Umbilical artery pH &lt; 7.00 (n/total)</u></b>                      Lactate: 21/1376                      pH: 24/1322</p> <p>RR 0.84 (95% CI 0.47 to 1.50)                      Heterogeneity: NA                      Test for overall effect: Z = 0.59 (p = 0.56)</p>	<p><b>Other information</b></p> <p><b><u>Success rate of fetal blood sampling (n/total (%))</u></b></p> <p>Lactate: 1478/1496 (97.8%)                      pH: 1341/1496 (89.6%)</p> <p>[1 trial: Wiberg-Itzel 2008]</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>[1 trial: Wiberg-Itzel 2008]</p> <p><u>c. Umbilical artery pH &lt; 7.10 (n/total)</u>                      Lactate: 121/1376                      pH: 131/1322</p> <p>RR 0.89 (95% CI 0.70 to 1.12)                      Heterogeneity: NA                      Test for overall effect: Z = 0.99 (p = 0.32)</p> <p>[1 trial: Wiberg-Itzel 2008]</p> <p><u>d. Umbilical artery lactate &gt; 4.68 mmol/l (n/total)‡</u>                      Lactate: 20/171                      pH: 29/156</p> <p>RR 0.63 (95% CI 0.37 to 1.07)                      Heterogeneity: NA                      Test for overall effect: Z = 1.72 (p = 0.085)</p> <p>[1 study: Westgren 1998]</p> <p><u>e. Umbilical artery base deficit (mean ± SD)</u>                      Lactate: 8 ± 3.8 [n = 171]                      pH: 8.7 ± 4.6 [n = 156]</p> <p>MD - 0.70 (95% CI - 1.62 to 0.22)                      Heterogeneity: NA                      Test for overall effect: Z = 1.49 (p = 0.14)</p> <p>[1 study: Westgren 1998]</p> <p><u>f. Umbilical artery base deficit &gt; 19.2‡</u>                      Lactate: 1/171                      pH: 3/156</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>RR 0.30 (0.03 to 2.89)                      Heterogeneity: NA                      Test for overall effect: Z = 1.04 (p = 0.30)</p> <p>[1 study: Westgren 1998]</p> <p>‡ According to the original trial paper, the thresholds used by Westgren were chosen according to the 1st or 99th centile of normal values, which are reported in another study</p> <p><b><u>SUB-GROUP ANALYSIS OF FBS TAKEN WITHIN 60 MINUTES OF DELIVERY</u></b>  <b><u>Operative delivery for non-reassuring fetal status</u></b>                      Lactate: 380/684                      pH: 257/508</p> <p>RR 1.10 (95% CI 0.98 to 1.22 )                      Heterogeneity: NA                      Test for overall effect: Z = 1.68 (p = 0.092)</p> <p>[1 study: Wiberg-Itzel et al., 2008)</p> <p><b><u>Apgar score &lt; 7 at 5 minutes</u></b>                      Lactate: 28/684                      pH: 21/508</p> <p>RR 0.99 (95% CI 0.57 to 1.72)                      Heterogeneity: NA                      Test for overall effect: Z = 0.03 (p = 0.97)</p> <p>[1 study: Wiberg-Itzel et al., 2008)</p> <p><b><u>Metabolic acidaemia (umbilical artery</u></b></p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p><b><u>pH &lt; 7.05 + base deficit &gt; 12 mmol/l</u></b>  <b><u>(n/total)</u></b>                      Lactate: 25/684                      pH: 20/508</p> <p>RR 0.93 (95% CI 0.52 to 1.65)                      Heterogeneity: NA                      Test for overall effect: Z = 0.25 (p = 0.80)</p> <p>[1 study: Wiberg-Itzel et al., 2008)</p> <p><b><u>Umbilical artery pH &lt; 7.00 (n/total)</u></b>                      Lactate: 10/684                      pH: 11/508</p> <p>RR 0.68 (95% CI 0.29 to 1.58)                      Heterogeneity: NA                      Test for overall effect: Z = 0.59 (p = 0.56 )</p> <p>[1 study: Wiberg-Itzel et al., 2008)</p>	
<p><b>Full citation</b></p> <p>Hon,E.H., Khazin,A.F., Paul,R.H.,                      Biochemical studies of the fetus. II. Fetal                      pH and apgar scores, Obstetrics and                      Gynecology,Obstet.Gynecol., 33, 237-                      255, 1969</p> <p><b>Ref Id</b></p> <p>159922</p> <p><b>Country/ies where the study was                      carried out</b></p> <p>USA</p>	<p><b>Sample size</b></p> <p>N = 194 patients</p> <p><b>Characteristics</b></p> <p>No details given</p> <p><b>Inclusion Criteria</b></p> <p>None reported</p> <p><b>Exclusion Criteria</b></p>	<p><b>Tests</b></p> <p>pH                      analysis</p>	<p><b>Methods</b></p> <p>Patients were monitored                      using electrocardiogram                      (ECG), fetal heart rate                      (FHR) patterns,                      monitoring of uterine                      contractions and blood                      pressure monitoring.                      Biochemical measures                      included maternal, fetal                      and neonatal pH, pO<sub>2</sub>,                      pCO<sub>2</sub>, base deficit,                      lactate, pyruvates and                      haemoglobin. 1392 fetal                      scalp samples were</p>	<p><b>Results</b></p> <p><b><u>Correlation between 1 minute Apgar                      scores and fetal blood pH at different                      intervals before birth</u></b>  <u>All samples</u>                      Apgar 7-10                      - Time interval (mean ± SD): 80.35                      ± 114.50                      - Apgar (mean ± SD): 8.56 ± 0.64                      - pH (mean ± SD): 7.28 ± 0.058                      - r: 0.0812                      - number of samples: 851                      - p-value: &lt; 0.05</p> <p>Apgar 1-6</p>	<p><b>Limitations</b></p> <p>No 2x2 data are available                      for samples taken within                      an hour of birth.</p> <p>Study sample represents                      population: unclear, as                      very few details are given                      Loss to follow-up is                      unrelated to key                      characteristics: unclear                      Prognostic factors are                      adequately measured in                      participants: yes                      Outcome of interest is</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>Not reported</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Supported in part by grants from the National Institute of Child Health and Human Development</p>	None reported		<p>obtained in total, of which 1117 samples were included in the study (194 patients).</p> <p>At the start of the study, pH was determined twice, once in early labour and once during late labour. However, during the later parts of the study, more frequent sampling was done, and reached as high as 28 per person.</p> <p>Apgar score was assessed as follows:</p> <ul style="list-style-type: none"> <li>- 7 - 10 was considered high</li> <li>- 6 or less was considered low</li> </ul> <p>A pH of 7.20 was used as the pH threshold.</p>	<p>- Time interval (mean <math>\pm</math> SD): 144.65 <math>\pm</math> 171.49</p> <p>- Apgar (mean <math>\pm</math> SD): 3.63 <math>\pm</math> 2.03</p> <p>- pH (mean <math>\pm</math> SD): 7.26 <math>\pm</math> 0.082</p> <p>- r: 0.3395</p> <p>- number of samples: 257</p> <p>- p-value: &lt; 0.005</p> <p><u>Within 60 minutes</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean <math>\pm</math> SD): 14.70 <math>\pm</math> 13.64</li> <li>- Apgar (mean <math>\pm</math> SD): 8.56 <math>\pm</math> 0.64</li> <li>- pH (mean <math>\pm</math> SD): 7.27 <math>\pm</math> 0.059</li> <li>- r: -0.0004</li> <li>- number of samples: 530</li> <li>- p-value: &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean <math>\pm</math> SD): 19.22 <math>\pm</math> 15.23</li> <li>- Apgar (mean <math>\pm</math> SD): 3.13 <math>\pm</math> 2.04</li> <li>- pH (mean <math>\pm</math> SD): 7.23 <math>\pm</math> 0.093</li> <li>- r: 0.4402</li> <li>- number of samples: 106</li> <li>- p-value: &lt; 0.005</li> </ul> <p><u>Within 45 minutes</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean <math>\pm</math> SD): 12.49 <math>\pm</math> 10.49</li> <li>- Apgar (mean <math>\pm</math> SD): 8.54 <math>\pm</math> 0.65</li> <li>- pH (mean <math>\pm</math> SD): 7.27 <math>\pm</math> 0.060</li> <li>- r: 0.0037</li> <li>- number of samples: 500</li> <li>- p-value: &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean <math>\pm</math> SD): 15.51 <math>\pm</math> 10.31</li> </ul>	<p>sufficiently measured in participants: yes</p> <p>Important potential confounders are accounted for: mode of birth is not reported</p> <p>Statistical analysis is appropriate for study design: yes</p> <p><b>Other information</b></p> <p>This study population appears to be the same as Khazin et al., but different data are reported</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>- Apgar (mean ± SD): 3.20 ± 2.00                      - pH (mean ± SD): 7.23 ± 0.089                      - r: 0.4248                      - number of samples: 96                      - p-value: &lt; 0.005</p> <p><u>Within 30 minutes</u>                      Apgar 7-10                      - Time interval (mean ± SD): 10.05 ± 7.15                      - Apgar (mean ± SD): 8.57 ± 0.64                      - pH (mean ± SD): 7.27 ± 0.060                      - r: 0.0203                      - number of samples: 456                      - p-value: &gt; 0.05</p> <p>Apgar 1-6                      - Time interval (mean ± SD): 13.50 ± 8.50                      - Apgar (mean ± SD): 3.23 ± 2.06                      - pH (mean ± SD): 7.22 ± 0.089                      - r: 0.4608                      - number of samples: 87                      - p-value: &lt; 0.005</p> <p><u>Within 15 minutes</u>                      Apgar 7-10                      - Time interval (mean ± SD): 7.28 ± 4.18                      - Apgar (mean ± SD): 8.61 ± 0.64                      - pH (mean ± SD): 7.27 ± 0.064                      - r: 0.0111                      - number of samples: 371                      - p-value: &gt; 0.05</p> <p>Apgar 1-6                      - Time interval (mean ± SD): 7.64 ± 4.25                      - Apgar (mean ± SD): 3.53 ± 2.17                      - pH (mean ± SD): 7.21 ± 0.104                      - r: 0.5490                      - number of samples: 53</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>- p-value: &lt; 0.005</p> <p><u>Within 5 minutes</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 2.87 ± 1.35</li> <li>- Apgar (mean ± SD): 8.58 ± 0.68</li> <li>- pH (mean ± SD): 7.25 ± 0.073</li> <li>- r: 0.0154</li> <li>- number of samples: 142</li> <li>- p-value: &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 2.71 ± 1.32</li> <li>- Apgar (mean ± SD): 3.47 ± 2.07</li> <li>- pH (mean ± SD): 7.23 ± 0.083</li> <li>- r: 0.7376</li> <li>- number of samples: 17</li> <li>- p-value: &lt; 0.005</li> </ul> <p><b><u>Correlation between 5 minute Apgar scores and fetal blood pH at different intervals before birth</u></b></p> <p><u>All samples</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 89.85 ± 118.90</li> <li>- Apgar (mean ± SD): 8.99 ± 0.74</li> <li>- pH (mean ± SD): 7.28 ± 0.060</li> <li>- r: 0.04343</li> <li>- number of samples: 1029</li> <li>- p-value: p &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 164.83 ± 240.04</li> <li>- Apgar (mean ± SD): 4.20 ± 1.57</li> <li>- pH (mean ± SD): 7.23 ± 0.097</li> <li>- r: 0.3485</li> <li>- number of samples: 79</li> </ul>	



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>- p-value: &lt;0.005</p> <p><u>Within 60 minutes:</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 15.52 ± 14.31</li> <li>- Apgar (mean ± SD): 9.11 ± 0.69</li> <li>- pH (mean ± SD): 7.27 ± 0.061</li> <li>- r: 0.0607</li> <li>- number of samples: 595</li> <li>- p-value: p &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 14.48 ± 8.69</li> <li>- Apgar (mean ± SD): 4.00 ± 1.82</li> <li>- pH (mean ± SD): 7/18 ± 0.098</li> <li>- r: 0.3880</li> <li>- number of samples: 41</li> <li>- p-value: &lt;0.01</li> </ul> <p><u>Within 45 minutes:</u></p> <p>Apgar 7-10</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 12.87 ± 10.63</li> <li>- Apgar (mean ± SD): 9.12 ± 0.68</li> <li>- pH (mean ± SD): 7.27 ± 0.06</li> <li>- r: 0.0019</li> <li>- number of samples: 555</li> <li>- p-value: p &gt; 0.05</li> </ul> <p>Apgar 1-6</p> <ul style="list-style-type: none"> <li>- Time interval (mean ± SD): 14.48 ± 8.69</li> <li>- Apgar (mean ± SD): 4.00 ± 1.82</li> <li>- pH (mean ± SD): 7/18 ± 0.098</li> <li>- r: 0.3880</li> <li>- number of samples: 41</li> <li>- p-value: &lt;0.01</li> </ul>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p><u>Within 30 minutes:</u>                      Apgar 7-10                      - Time interval (mean ± SD): 10.33 ± 7.35                      - Apgar (mean ± SD): 9.15 ± 0.67                      - pH (mean ± SD): 7.27 ± 0.06                      - r: 0.0044                      - number of samples: 503                      - p-value: p &gt; 0.05</p> <p>Apgar 1-6                      - Time interval (mean ± SD): 14.06 ± 8.38                      - Apgar (mean ± SD): 3.95 ± 1.81                      - pH (mean ± SD): 7.18 ± 0.096                      - r: 0.3591                      - number of samples: 40                      - p-value: &lt; 0.05</p> <p><u>Within 15 minutes:</u>                      Apgar 7-10                      - Time interval (mean ± SD): 7.27 ± 4.17                      - Apgar (mean ± SD): 9.22 ± 0.63                      - pH (mean ± SD): 7.27 ± 0.063                      - r: -0.0120                      - number of samples: 400                      - p-value: p &gt; 0.05</p> <p>Apgar 1-6                      - Time interval (mean ± SD): 8.31 ± 4.44                      - Apgar (mean ± SD): 4.21 ± 1.84                      - pH (mean ± SD): 7.16 ± 0.114                      - r: 0.4261                      - number of samples: 24                      - p-value: &lt; 0.05</p> <p><u>Within 5 minutes:</u>                      Apgar 7-10                      - Time interval (mean ± SD): 2.83 ± 1.34                      - Apgar (mean ± SD): 9.18 ± 0.65</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>- pH (mean ± SD): 7/25 ± 0.071                      - r: -0.0534                      - number of samples: 151                      - p-value: p &gt; 0.05</p> <p>Apgar 1-6                      - Time interval (mean ± SD): 3.31 ± 1.44                      - Apgar (mean ± SD): 4.25 ± 1.58                      - pH (mean ± SD): 7.18 ± 0.080                      - r: 0.6171                      - number of samples: 8                      - p-value: &lt; 0.05</p>	
<p><b>Full citation</b>                      Kerenyi,T.D., Falk,S., Mettel,R.D., Walker,B., Acid-base balance and oxygen saturation of fetal scalp blood during normal and abnormal labors, Obstetrics and Gynecology, 36, 398-404, 1970</p> <p><b>Ref Id</b>                      169762</p> <p><b>Country/ies where the study was carried out</b>                      USA</p> <p><b>Study type</b></p> <p><b>Aim of the study</b>                      Not stated</p>	<p><b>Sample size</b>                      N = 33                      (However, only 23 were taken within 1 hour of delivery and hence constitute the population of interest)</p> <p><b>Characteristics</b>                      Of the study population who had a fetal blood sample (FBS) taken within an hour of birth:                      8 had normal labours and gave birth to babies with an Apgar score of 6 or better, following a blood sample taken within 1 hour of birth (range 10 minutes to 55 minutes). Dilatation</p>	<p><b>Tests</b>                      pH analysis within 60 minutes of birth</p>	<p><b>Methods</b>                      Fetal blood sampling was done with the patient in the lithotomy position, after the membranes had either been ruptured artificially or had spontaneously ruptured. An endoscope was put through the os and pressed against the head. The scalp was cleaned and at the time of a contraction was sprayed with ethyl chloride to produce hyperaemia. A silicone preparation was applied to enhance blood beading. A puncture was made with a 2mm blade and blood was collected in a heparinised tube after suction was applied</p>	<p><b>Results</b>                      The following predictive value measures were calculated by the technical team, based on data reported in tables 1 - 3 of the paper. The calculations only include fetal scalp samples that were taken within 1 hour of birth (n = 23). There is missing data for 2 arterial samples.</p> <p><b>Predictive value of pH &lt; 7.10 (95% CI)</b></p> <p><u>a. For Apgar score &lt; 7 at 1 minute</u>                      Sensitivity: 25.00% (0.50 to 49.50)                      Specificity: 100 (NC)                      PPV: 100 (NC)                      NPV: 55.00% (33.20 to 76.80)                      LR+: infinite                      LR-: 0.75 (0.54 to 1.04)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u>                      Sensitivity: 66.67% (13.32 to 100)                      Specificity: 95.00% (85.45 to 100)                      PPV: 66.67% (13.32 to 100)                      NPV: 95.00% (85.45 to 100)                      LR+: 13.33 (1.68 to 105.79)</p>	<p><b>Limitations</b>                      Study sample represents population: Many of the women were not low risk; inclusion and exclusion criteria are not reported                      Loss to follow-up is unrelated to key characteristics: No loss to follow-up                      Prognostic factors are adequately measured in participants: There are missing data for between 4 and 5 (17 - 22%) out of the 23 women for base deficit values.                      Outcome of interest is sufficiently measured in participants: There are missing data for 2/23 arterial pH measurements                      Important potential confounders are</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p>was rim in one woman, 6-9 in 5 women and full in 2 women.</p> <p>7 had complicated labours and gave birth to babies with an Apgar score of 6 or better after an FBS within an hour of birth (range 1 minute - 40 minutes):</p> <p>Case 5: abnormal fetal heart rate (FHR), pitocin drip, secondary uterine inertia, - Full dilatation</p> <p>Case 15: Toxemia - Full dilatation</p> <p>Case 22: Relative cephalopelvic disproportion, eclamptic - Full dilatation</p> <p>Case 23: premature (2300 g), fetal tachycardia - Full dilatation</p> <p>Case 27: meconium staining - Full dilatation</p> <p>Case Elm 4: toxemia, relative chronic pulmonary disease (CPD), premature rupture of membranes (RoM), tachycardia, rim and full dilatation - Full dilatation</p> <p>Case 26: Class D diabetes - Full dilatation</p>		<p>by mouth. The sample was immediately analysed.</p> <p>Samples were taken periodically during labour. If any value was abnormal, the analysis was immediately repeated and the result compared to the maternal blood. As the series went on, maternal acid-base status was found to be a useful tool in determining whether acidosis started in the mother or the baby.</p> <p>At delivery, blood samples from the cord were collected before clamping. The clinical status of the baby was evaluated at 1 minute and 5 minutes.</p> <p>All patients delivered under local or regional anaesthesia, where possible. Patients received varying amounts of meperidine and scopolamine for analgesia.</p>	<p>LR-: 0.35 (0.07 to 1.74)</p> <p><u>c. For umbilical artery pH &lt; 7.10</u> Sensitivity: 33.33% (0 to 86.68) Specificity: 94.44% (83.86 to 100) PPV: 50.00% (0 to 100) NPV: 89.47% (75.67 to 100) LR+: 6.00 (0.50 to 72.21) LR-: 0.71 (0.31 to 1.58)</p> <p><b><u>Predictive value of pH ≤ 7.20 (95% CI)</u></b></p> <p><u>a. For Apgar score &lt; 7 at 1 minute</u> Sensitivity: 58.33% (30.44 to 86.23) Specificity: 72.73% (46.41 to 99.05) PPV: 70.00% (41.60 to 98.40) NPV: 61.54% (35.09 to 87.99) LR+: 2.14 (0.73 to 6.28) LR-: 0.57 (0.27 to 1.23)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> Sensitivity: 66.67% (13.32 to 100) Specificity: 60.00% (38.53 to 81.47) PPV: 20.00% (0 to 44.79) NPV: 92.31% (77.82 to 100) LR+: 1.67 (0.64 to 4.37) LR-: 0.56 (0.11 to 2.86)</p> <p><u>c. For umbilical artery pH &lt; 7.1</u> Sensitivity: 100% (NC) Specificity: 66.67% (44.89 to 88.44) PPV: 33.33% (2.5 to 64.13) NPV: 100% (NC) LR+: 3.00 (1.56 to 5.77) LR-: 0.00 (NC)</p> <p><b><u>Predictive value of pH ≤ 7.25 (95% CI)</u></b></p> <p><u>a. For Apgar score &lt; 7 at 1 minute</u> Sensitivity: 75.00% (50.50 to 99.50) Specificity: 9.09% (0 to 26.08)</p>	<p>accounted for: Mode of birth is not reported Statistical analysis is appropriate for study design: Yes</p> <p><b>Other information</b></p> <p><b><u>Further information about cases of low Apgar score at 5 minutes</u></b></p> <p><b>Case 14:</b> - Meconium staining, fetal tachycardia - Tested at 19 minutes before birth - Apgar of 2 at 1 minute and 5 at 5 minutes</p> <p><b>Case 18:</b> - Fetal distress, irregular and slow FHR - Tested at 25 minutes before birth - Baby was stillborn</p> <p><b>Case 30:</b> - Cephalopelvic disproportion, irregular FHR, caesarean section - Tested at 40 minutes before birth - Apgar of 4 at 1 minute and 6 at 5 minutes</p> <p><b><u>Further information about cases of low</u></b></p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
	<p>8 had complicated labours and gave birth to depressed babies within an hour of FBS (range 16 minutes to 40 minutes):</p> <p>Case 3: relative CPD, pitocin drip - 7 cm dilatation</p> <p>Case 12: CPD - Full dilatation</p> <p>Case 14: meconium staining, fetal tachycardia - 5-6 cm dilatation</p> <p>Case 18: fetal distress, irregular and slow FHR [still born] - Full dilatation</p> <p>Case 19: CPD, fetal distress, FHR 60, cord around shoulder - Full dilatation</p> <p>Case 24: prolonged RoM, amniotitis, fetal sepsis - Full dilatation</p> <p>Case Elm 3: toxemia, type II dips, CPD - Full dilatation</p> <p>Case 30: CPD, irregular FHR, caesarean - 7 cm dilatation</p> <p><b>Inclusion Criteria</b></p> <p>None reported</p>			<p>PPV: 47.37% (24.92 to 69.82) NPV: 25.00% (0 to 67.44) LR+: 0.83 (0.57 to 1.20) LR-: 2.75 (0.33 to 22.69)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> Sensitivity: 66.67% (13.32 to 100) Specificity: 15.00% (0 to 30.65) PPV: 10.53% (0 to 24.33) NPV: 75.00% (32.56 to 100) LR+: 0.78 (0.35 to 1.78) LR-: 2.22 (0.33 to 15.01)</p> <p><u>c. For umbilical artery pH &lt; 7.1</u> Sensitivity: 100% (NC) Specificity: 22.22% (3.02 to 41.43) PPV: 17.65% (0 to 35.77) NPV: 100% (NC) LR+: 1.29 (1.00 to 1.65) LR-: 0 (NC)</p> <p><b><u>Predictive value of base deficit &gt; 10 mEq/l (95% CI)</u></b></p> <p><u>a. For Apgar score &lt; 7 at 1 minute</u> Sensitivity: 25.00% (0 to 55.01) Specificity: 90.91% (73.92 to 100) PPV: 66.67% (13.32 to 100) NPV: 62.50% (38.78 to 86.22) LR+: 2.75 (0.30 to 25.35) LR-: 0.83 (0.53 to 1.28)</p> <p><b><u>b. For Apgar score &lt; 7 at 5 minutes</u></b> Sensitivity: 0 (NC) Specificity: 83.33% (66.12 to 100) PPV: 0 (NC) NPV: 93.75% (81.89 to 100) LR+: 0 (NC) LR-: 1.20 (0.98 to 1.48)</p> <p><u>c. For umbilical artery pH &lt; 7.10</u></p>	<p><b><u>arterial pH (&lt; 7.10) at birth</u></b></p> <p><b>Case 12:</b> - Cephalopelvic disproportion - Tested at 16 minutes before birth and had pH of 7.12 - Artery pH of 7.06</p> <p><b>Case 18:</b> - Fetal distress, irregular and slow FHR - Tested at 25 minutes before birth and had pH of 6.64 - Baby was stillborn and had arterial pH of 6.81</p> <p><b>Case Elm 3:</b> - Toxemia, type II dips, cephalopelvic disproportion - Tested at 25 minutes before birth and had pH of 7.15 - Artery pH of 7.08</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
	<p><b>Exclusion Criteria</b></p> <p>None reported</p>			<p>Sensitivity: 0 (NC)                      Specificity: 81.25% (62.12 to 100)                      PPV: 0 (NC)                      NPV: 86.67% (69.46 to 100)                      LR+: 0 (NC)                      LR-: 1.23 (0.97 to 1.56)</p> <p><b><u>Predictive value of base deficit &gt; 12 mEq/l (95% CI)</u></b></p> <p><b><u>a. For Apgar score &lt; 7 at 1 minute</u></b>                      Sensitivity: 25.00% (0 to 55.01)                      Specificity: 100% (NC)                      PPV: 100 (NC)                      NPV: 64.71% (41.99 to 87.42)                      LR+: infinite                      LR-: 0.75 (0.51 to 1.12)</p> <p><b><u>b. For Apgar score &lt; 7 at 5 minutes</u></b>                      Sensitivity: 0 (NC)                      Specificity: 88.89% (74.37 to 100)                      PPV: 0 (NC)                      NPV: 94.12 (82.93 to 100)                      LR+: 0 (NC)                      LR-: 1.13 (0.96 to 1.32)</p> <p><b><u>c. For umbilical artery pH &lt; 7.10</u></b>                      Sensitivity: 0 (NC)                      Specificity: 87.50% (71.29 to 100)                      PPV: 0 (NC)                      NPV: 87.50% (71.29 to 100)                      LR+: 0 (NC)                      LR-: 1.14 (0.95 to 1.38)</p> <p><b><u>Predictive value of base deficit &gt; 12.5 mEq/l (95% CI)</u></b></p> <p><b><u>a. For Apgar score &lt; 7 at 1 minute</u></b>                      Sensitivity: 12.50% (0 to 35.42)                      Specificity: 100 (NC)                      PPV: 100 (NC)                      NPV: 61.11% (38.59 to 83.63)</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
				<p>LR+: infinite LR-: 0.88 (0.67 to 1.14)</p> <p><u>b. For Apgar score &lt; 7 at 5 minutes</u> Sensitivity: 0 (NC) Specificity: 94.44% (83.86 to 100) PPV: 0 (NC) NPV: 94.44% (83.86 to 100) LR+: 0 (NC) LR-: 1.06 (0.95 to 1.18)</p> <p><u>c. For umbilical artery pH &lt; 7.10</u> Sensitivity: 0 (NC) Specificity: 93.75% (81.89 to 100) PPV: 0 (NC) NPV: 88.24% (72.92 to 100) LR+: 0 (NC) LR-: 1.07 (0.94 to 1.21)</p> <p><b>FBS pH &lt; 7.1 for Apgar &lt; 7 at 1 minute</b></p> <table border="1" data-bbox="1319 882 1749 1182"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>0</td> </tr> <tr> <td>Predictive Test -ve</td> <td>9</td> <td>11</td> </tr> </tbody> </table> <p><b>FBS pH &lt; 7.1 for arterial pH &lt; 7.10</b></p>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	0	Predictive Test -ve	9	11	
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	3	0												
Predictive Test -ve	9	11												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1" data-bbox="1319 272 1749 572"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>1</td> <td>1</td> </tr> <tr> <td>Predictive Test -ve</td> <td>2</td> <td>17</td> </tr> </tbody> </table> <p data-bbox="1319 632 1715 655"><b>FBS pH &lt;= 7.20 for arterial pH &lt; 7.1</b></p> <table border="1" data-bbox="1319 684 1749 984"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>6</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>12</td> </tr> </tbody> </table> <p data-bbox="1319 1043 1693 1094"><b>FBS pH &lt;= 7.20 for Apgar &lt; 7 at 1 minute</b></p> <table border="1" data-bbox="1319 1123 1749 1326"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>7</td> <td>3</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	1	1	Predictive Test -ve	2	17		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	6	Predictive Test -ve	0	12		Reference Test +ve	Reference Test -ve	Predictive Test +ve	7	3	
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	1	1																											
Predictive Test -ve	2	17																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	3	6																											
Predictive Test -ve	0	12																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	7	3																											



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																					
				<table border="1" data-bbox="1319 272 1749 371"> <tr> <td>Predictive Test -ve</td> <td>5</td> <td>8</td> </tr> </table> <p data-bbox="1319 427 1697 480"><b>FBS pH ≤ 7.20 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1" data-bbox="1319 509 1749 810"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>8</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>12</td> </tr> </tbody> </table> <p data-bbox="1319 866 1715 892"><b>FBS pH ≤ 7.25 for arterial pH &lt; 7.1</b></p> <table border="1" data-bbox="1319 920 1749 1222"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>14</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>4</td> </tr> </tbody> </table> <p data-bbox="1319 1278 1693 1331"><b>FBS pH ≤ 7.25 for Apgar &lt; 7 at 1 minute</b></p>	Predictive Test -ve	5	8		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	8	Predictive Test -ve	1	12		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	14	Predictive Test -ve	0	4	
Predictive Test -ve	5	8																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	2	8																								
Predictive Test -ve	1	12																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	3	14																								
Predictive Test -ve	0	4																								

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
				<table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>9</td> <td>10</td> </tr> <tr> <td>Predictive Test -ve</td> <td>3</td> <td>1</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	9	10	Predictive Test -ve	3	1	
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	9	10												
Predictive Test -ve	3	1												
				<p><b>FBS &lt;= 7.25 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>17</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>3</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	17	Predictive Test -ve	1	3	
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	2	17												
Predictive Test -ve	1	3												
				<p><b>FBS base deficit &gt; 10 for Apgar &lt; 7 at 1 minute</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>1</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	1				
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	2	1												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																					
				<table border="1" data-bbox="1319 272 1749 371"> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>10</td> </tr> </table> <p data-bbox="1319 427 1749 483"><b>FBS base deficit &gt; 10 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1" data-bbox="1319 507 1749 810"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>0</td> <td>3</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>15</td> </tr> </tbody> </table> <p data-bbox="1319 866 1749 922"><b>FBS base deficit &gt; 10 for arterial pH &lt; 7.10</b></p> <table border="1" data-bbox="1319 946 1749 1249"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>0</td> <td>3</td> </tr> <tr> <td>Predictive Test -ve</td> <td>2</td> <td>13</td> </tr> </tbody> </table> <p data-bbox="1319 1305 1749 1361"><b>FBS base deficit &gt; 12 for Apgar &lt; 7 at 1 minute</b></p>	Predictive Test -ve	6	10		Reference Test +ve	Reference Test -ve	Predictive Test +ve	0	3	Predictive Test -ve	1	15		Reference Test +ve	Reference Test -ve	Predictive Test +ve	0	3	Predictive Test -ve	2	13	
Predictive Test -ve	6	10																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	0	3																								
Predictive Test -ve	1	15																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	0	3																								
Predictive Test -ve	2	13																								

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1" data-bbox="1319 272 1749 572"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>2</td> <td>0</td> </tr> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>11</td> </tr> </tbody> </table> <p data-bbox="1319 632 1749 683"><b>FBS base deficit &gt; 12 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1" data-bbox="1319 711 1749 1011"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>0</td> <td>2</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>16</td> </tr> </tbody> </table> <p data-bbox="1319 1070 1749 1121"><b>FBS base deficit &gt; 12 for arterial pH &lt; 7.10</b></p> <table border="1" data-bbox="1319 1150 1749 1353"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>0</td> <td>2</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	2	0	Predictive Test -ve	6	11		Reference Test +ve	Reference Test -ve	Predictive Test +ve	0	2	Predictive Test -ve	1	16		Reference Test +ve	Reference Test -ve	Predictive Test +ve	0	2	
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	2	0																											
Predictive Test -ve	6	11																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	0	2																											
Predictive Test -ve	1	16																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	0	2																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																					
				<table border="1" data-bbox="1319 272 1749 373"> <tr> <td>Predictive Test -ve</td> <td>2</td> <td>14</td> </tr> </table> <p data-bbox="1319 427 1736 480">FBS base deficit &gt; 12.5 for Apgar &lt; 7 at 1 minute</p> <table border="1" data-bbox="1319 509 1749 810"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>1</td> <td>0</td> </tr> <tr> <td>Predictive Test -ve</td> <td>7</td> <td>11</td> </tr> </tbody> </table> <p data-bbox="1319 865 1736 917">FBS base deficit &gt; 12.5 for Apgar &lt; 7 at 5 minutes</p> <table border="1" data-bbox="1319 946 1749 1248"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>0</td> <td>1</td> </tr> <tr> <td>Predictive Test -ve</td> <td>1</td> <td>17</td> </tr> </tbody> </table> <p data-bbox="1319 1302 1736 1355">FBS base deficit &gt; 12.5 for arterial pH &lt; 7.10</p>	Predictive Test -ve	2	14		Reference Test +ve	Reference Test -ve	Predictive Test +ve	1	0	Predictive Test -ve	7	11		Reference Test +ve	Reference Test -ve	Predictive Test +ve	0	1	Predictive Test -ve	1	17	
Predictive Test -ve	2	14																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	1	0																								
Predictive Test -ve	7	11																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	0	1																								
Predictive Test -ve	1	17																								

Bibliographic details	Participants	Tests	Methods	Outcomes and results		Comments	
				Reference Test +ve	Reference Test -ve		
				Predictive Test +ve	0	1	
				Predictive Test -ve	2	15	
<p><b>Full citation</b></p> <p>Khazin,A.F., Hon,E.H., Quilligan,E.J., Biochemical studies of the fetus. 3. Fetal base and Apgar scores, Obstetrics and Gynecology, 34, 592-609, 1969</p> <p><b>Ref Id</b></p> <p>170426</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>Not reported</p> <p><b>Study dates</b></p> <p>Not reported</p>	<p><b>Sample size</b></p> <p>N = 194</p> <p><b>Characteristics</b></p> <p>80 patients had complications of pregnancy such as toxemia, Rh sensitisation, diabetes, premature rupture of membranes, clinically diagnosed fetal distress or post-dates (proportions of each are not reported)</p> <p><b>Inclusion Criteria</b></p> <p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>pH analysis</p>	<p><b>Methods</b></p> <p>Fetal blood samples were collected according to Saling's technique, but glass capillary tubes were used instead of plastic. Patients were monitored using electrocardiogram (ECG), fetal heart rate (FHR) patterns, monitoring of uterine contractions and blood pressure monitoring. Biochemical measures included maternal, fetal and neonatal pH, pO<sub>2</sub>, pCO<sub>2</sub>, base deficit, lactate, pyruvates and haemoglobin.</p> <p>Umbilical artery and vein blood was obtained before the first breath of the infant, from a doubly clamped segment of the umbilical cord.</p>	<p><b>Results</b></p> <p>The following calculations were performed by the technical team, based on 2x2 data reported in the text for 130 <b>babies</b> who had samples taken within 30 minutes of birth:</p> <p><b><u>Predictive accuracy (95% CI) of a fetal base deficit of &gt; 12.5 mEq/l for:</u></b></p> <p><b>a. 1-minute Apgar score &lt; 7</b>  Sensitivity: 31.82% (12.35 to 51.28)  Specificity: 92.59% (87.65 to 97.53)  PPV: 46.67% (21.42 to 71.91)  NPV: 86.96% (80.80 to 93.11)  LR+: 4.30 (1.74 to 10.62)  LR-: 0.74 (0.55 to 0.98)</p> <p><b>b. 5-minute Apgar score &lt; 7</b>  Sensitivity: 42.86% (6.20 to 79.52)  Specificity: 90.24% (85.00 to 95.49)  PPV: 20.00% (0 to 40.24)  NPV: 96.52% (93.17 to 99.87)  LR+: 4.39 (1.60 to 12.06)  LR-: 0.63 (0.33 to 1.21)</p> <p><b><u>Correlation between 1 minute Apgar score and fetal base-deficit at different intervals before birth</u></b></p>	<p><b>Limitations</b></p> <p>Study sample represents population: 80/194 women had complications in labour; very few other details about the population are reported  Loss to follow-up is unrelated to key characteristics: no loss to follow-up  Prognostic factors are adequately measured in participants: very few details about what happened to the babies during labour  Outcome of interest is sufficiently measured in participants: yes  Important potential confounders are accounted for: mode of birth is not reported  Statistical analysis is appropriate for study design: yes</p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Source of funding</b></p> <p>Supported in part by research grants from the National Institute of Child Health and Human Development, USPHS, and a grant from the Health Sciences Computing Facility</p>			<p>A radiometer microelectrode was done to determine pH. Fetal scalp blood samples were obtained during different stages of labour, and between 1 and 35 samples were taken per patient. Fetal base determinations were done on 602 samples taken from 140 patients (1 - 17 per patient).</p> <p>Apgar score at 1 and 5 minutes were taken. 1 - 6 was considered low, and 7 - 10 was considered high. This was first done for all samples, and then restricted to samples taken within the last 30 minutes of labour.</p> <p>To determine the impact of time interval between fetal base determination and birth on predictive values, correlation coefficients were taken for all samples, then restricted to those in the last 60, 45, 30, 15 and 5 minutes preceding birth.</p>	<p><u>All samples</u></p> <p>- Apgar 7 - 10 Time interval (mean <math>\pm</math> SD): 86.06 <math>\pm</math> 111.55 Apgar (mean <math>\pm</math> SD): 8.53 <math>\pm</math> 0.63 Base deficit / mEq/l (mean <math>\pm</math> SD): 7.91 <math>\pm</math> 2.80 number of samples: 472 r: -0.1459 p-value: &lt; 0.05</p> <p>- Apgar 1 - 6 Time interval (mean <math>\pm</math> SD): 194.54 <math>\pm</math> 225.81 Apgar (mean <math>\pm</math> SD): 3.29 <math>\pm</math> 2.08 Base deficit / mEq/l (mean <math>\pm</math> SD): 8.26 <math>\pm</math> 3.39 number of samples: 130 r: +0.0387 p-value: &gt; 0.05</p> <p><u>60 minutes before birth</u></p> <p>- Apgar 7 - 10 Time interval (mean <math>\pm</math> SD): 15.75 <math>\pm</math> 15.05 Apgar (mean <math>\pm</math> SD): 8.48 <math>\pm</math> 0.67 Base deficit / mEq/l (mean <math>\pm</math> SD): 8.27 <math>\pm</math> 2.95 number of samples: 277 r: -0.2002 p-value: &lt;0.005</p> <p>- Apgar 1 - 6 Time interval (mean <math>\pm</math> SD): 19.70 <math>\pm</math> 12.05 Apgar (mean <math>\pm</math> SD): 3.16 <math>\pm</math> 2.03 Base deficit / mEq/l (mean <math>\pm</math> SD): 9.75 <math>\pm</math> 3.85 number of samples: 45 r: -0.2056</p>	<p><b>Other information</b></p> <p><b><u>Further information about the false negatives (i.e. base deficit <math>\leq</math> 12.5 mEq/l but with a low Apgar score at 1 minute, table 5 in paper)</u></b></p> <p>1. - 2 samples taken, at 20 minutes and 16 minutes prior to birth - BD 11.1 - 11.3 - Late decelerations (+++), hyperactivity (+++) - Apgar scores: 2, 5</p> <p>2. - 5 samples taken, at between 320 and 18 minutes prior to birth - BD 8.8 - 10.3 - Variable decelerations (++) , Caput (+++) - Forceps applied with traction for 7 minutes - Apgar scores: 4, 7</p> <p>3. - 3 samples taken, at between 12 and 9 minutes prior to birth - BD 9.4 - 12.4 - Variable decelerations (+) - Shoulder dystocia, midforceps</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>p-value: &gt; 0.05</p> <p><u>45 minutes before birth</u>                      - Apgar 7 - 10                      Time interval (mean ± SD): 12.80 ± 11.04                      Apgar (mean ± SD): 8.47 ± 0.67                      Base deficit / mEq/l (mean ± SD): 8.32 ± 2.99                      number of samples: 257                      r: -0.1817                      p-value: &lt;0.005</p> <p>- Apgar 1 - 6                      Time interval (mean ± SD): 18.38 ± 10.59                      Apgar (mean ± SD): 3.26 ± 2.03                      Base deficit / mEq/l (mean ± SD): 9.72 ± 3.68                      number of samples: 43                      r: -0.2167                      p-value: &gt; 0.05</p> <p><u>30 minutes before birth</u>                      - Apgar 7 - 10                      Time interval (mean ± SD): 9.94 ± 7.50                      Apgar (mean ± SD): 8.52 ± 0.66                      Base deficit / mEq/l (mean ± SD): 8.39 ± 2.98                      number of samples: 230                      r: -0.1825                      p-value: &lt; 0.05</p> <p>- Apgar 1 - 6                      Time interval (mean ± SD): 14.59 ± 7.43                      Apgar (mean ± SD): 3.31 ± 2.15                      Base deficit / mEq/l (mean ± SD): 10.43 ± 3.31                      number of samples: 35                      r: -0.2664</p>	<p>- Apgar scores: 6, 9</p> <p>4.                      - 2 samples taken at between 24 and 22 minutes prior to birth                      - BD 7.2                      - Variable decelerations (++)                      - Twin A, variable decelerations with delivery                      - Apgar scores: 5, 9</p> <p>[Note: there was one further case, but the sample was taken outside the time of interest; therefore details have not been reported here]</p>



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>p-value: &gt; 0.05</p> <p><u>15 minutes before birth</u></p> <p>- Apgar 7 - 10                      Time interval (mean ± SD): 6.84 ± 4.06                      Apgar (mean ± SD): 8.58 ± 0.66                      Base deficit / mEq/l (mean ± SD): 8.28 ± 2.98                      number of samples: 185                      r: -0.1812                      p-value: &gt; 0.05</p> <p>- Apgar 1 - 6                      Time interval (mean ± SD): 8.58 ± 4.36                      Apgar (mean ± SD): 3.44 ± 2.55                      Base deficit / mEq/l (mean ± SD): 10.57 ± 3.36                      number of samples: 18                      r: -0.3553                      p-value: &gt; 0.05</p> <p><u>5 minutes before birth</u></p> <p>- Apgar 7 - 10                      Time interval (mean ± SD): 3.01 ± 1.37                      Apgar (mean ± SD): 8.61 ± 0.68                      Base deficit / mEq/l (mean ± SD): 8.49 ± 2.46                      number of samples: 81                      r: -0.0590                      p-value: &gt; 0.05</p> <p>- Apgar 1 - 6                      Time interval (mean ± SD): 1.75 ± 0.50                      Apgar (mean ± SD): 2.50 ± 2.38                      Base deficit / mEq/l (mean ± SD): 10.68 ± 1.08                      number of samples: 4                      r: -0.9259                      p-value:</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p><b><u>Correlation between 5 minute Apgar score and fetal base-deficit at different intervals before birth</u></b></p> <p><u>All samples</u></p> <p>- Apgar 7 - 10 Time interval (mean ± SD): 94.26 ± 114.80 Apgar (mean ± SD): 9.01 ± 0.70 Base deficit / mEq/l (mean ± SD): 7.97 ± 2.92 number of samples: 559 r: -0.0918 p-value: &lt; 0.05</p> <p>- Apgar 1 - 6 Time interval (mean ± SD): 307.45 ± 326.20 Apgar (mean ± SD): 4.65 ± 1.25 Base deficit / mEq/l (mean ± SD): 8.11 ± 3.27 number of samples: 43 r: -0.3210 p-value: &lt; 0.05</p> <p><u>60 minutes before birth</u></p> <p>- Apgar 7 - 10 Time interval (mean ± SD): 16.31 ± 14.94 Apgar (mean ± SD): 9.08 ± 0.68 Base deficit / mEq/l (mean ± SD): 8.35 ± 3.06 number of samples: 309 r: -0.0960 p-value: &gt; 0.05</p> <p>- Apgar 1 - 6 Time interval (mean ± SD): 16.31 ± 7.99 Apgar (mean ± SD): 4.62 ± 1.76 Base deficit / mEq/l (mean ± SD): 11.47 ± 3.18</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>number of samples: 13  r: -0.8362  p-value: &lt; 0.005</p> <p><u>45 minutes before birth</u>  - Apgar 7 - 10  Time interval (mean ± SD): 13.48 ± 11.25  Apgar (mean ± SD): 9.08 ± 0.68  Base deficit / mEq/l (mean ± SD): 8.38 ± 3.06</p> <p>number of samples: 287  r: -0.0663  p-value: &gt; 0.05</p> <p>- Apgar 1 - 6  Time interval (mean ± SD): 16.31 ± 7.99  Apgar (mean ± SD): 4.62 ± 1.76  Base deficit / mEq/l (mean ± SD): 11.47 ± 3.18</p> <p>number of samples: 13  r: -0.8362  p-value: &lt; 0.005</p> <p><u>30 minutes before birth</u>  - Apgar 7 - 10  Time interval (mean ± SD): 10.34 ± 7.61  Apgar (mean ± SD): 9.11 ± 0.64  Base deficit / mEq/l (mean ± SD): 8.51 ± 3.03</p> <p>number of samples: 253  r: -0.1383  p-value: &lt; 0.05</p> <p>- Apgar 1 - 6  Time interval (mean ± SD): 15.13 ± 7.05  Apgar (mean ± SD): 4.50 ± 1.78  Base deficit / mEq/l (mean ± SD): 11.84 ± 3.02</p> <p>number of samples: 12</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p>r: -0.8359 p-value: &lt; 0.005</p> <p><u>15 minutes before birth</u> - Apgar 7 - 10 Time interval (mean ± SD): 6.91 ± 4.07 Apgar (mean ± SD): 9.21 ± 0.58 Base deficit / mEq/l (mean ± SD): 8.36 ± 2.98 number of samples: 197 r: -0.1454 p-value: &gt; 0.05</p> <p>- Apgar 1 - 6 Time interval (mean ± SD): 9.75 ± 4.45 Apgar (mean ± SD): 4.33 ± 2.58 Base deficit / mEq/l (mean ± SD): 12.42 ± 4.12 number of samples: 6 r: -0.9366 p-value: &lt; 0.005</p> <p><u>5 minutes before birth</u> - Apgar 7 - 10 Time interval (mean ± SD): 2.96 ± 1.37 Apgar (mean ± SD): 9.21 ± 0.62 Base deficit / mEq/l (mean ± SD): 8.55 ± 2.44 number of samples: 84 r: -0.1517 p-value: 0.05</p> <p>- Apgar 1 - 6 Time interval (mean ± SD): 2.00 (NA) Apgar (mean ± SD): 6 (NA) Base deficit / mEq/l (mean ± SD): 11.80 (NA) number of samples: 1 r: NA p-value: NA</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<p>FBS base deficit &gt; 12.5 for Apgar &lt; 7 at 1 minute</p> <table border="1" data-bbox="1317 411 1751 711"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>7</td> <td>8</td> </tr> <tr> <td>Predictive Test -ve</td> <td>15</td> <td>100</td> </tr> </tbody> </table> <p>FBS base deficit &gt; 12.5 for Apgar &lt; 7 at 5 minutes</p> <table border="1" data-bbox="1317 849 1751 1149"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>12</td> </tr> <tr> <td>Predictive Test -ve</td> <td>4</td> <td>111</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	7	8	Predictive Test -ve	15	100		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	12	Predictive Test -ve	4	111	
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	7	8																					
Predictive Test -ve	15	100																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	3	12																					
Predictive Test -ve	4	111																					
<p><b>Full citation</b></p> <p>Kubli,F.W., Influence of labor on fetal acid-base balance, Clinical Obstetrics and Gynecology, 11, 168-191, 1968</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>N = 77</p> <p><b>Characteristics</b></p> <p>none reported</p>	<p><b>Tests</b></p> <p>pH within 30 minutes of birth</p>	<p><b>Methods</b></p> <p>Very few details are reported, as this is a further analysis of another study by Hon (referenced as not published). 77 patients</p>	<p><b>Results</b></p> <p>The following measures were calculated based on 2x2 data reported in table 2a of the paper.</p> <p><b>Predictive value of pH &lt; 7.20 for an Apgar &lt; 7 (reported as ≤ 6) at 1 minute</b></p>	<p><b>Limitations</b></p> <p>Study sample represents population: Unclear, exclusion and inclusion criteria are not reported and there are no characteristics reported</p>																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
<p>169765</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>Not reported</p> <p><b>Study dates</b></p> <p>1966 - 1967</p> <p><b>Source of funding</b></p> <p>Supported in part by Public Health Service Research Grant from the National Heart Institute and a Grant from DFG (Deutsche Forschungsgemeinschaft)</p>	<p><b>Inclusion Criteria</b></p> <p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>		<p>were selected in whom the last sample was done 30 minutes before birth. However, the authors report including 5 further patients with an abnormal pH value with or without depression.</p> <p>For all patients, continuous fetal heart rate monitoring was done and amniotic fluid pressure was recorded.</p>	<p>Sensitivity: 57.14% (31.22 to 83.07)                      Specificity: 84.13% (75.10 to 93.15)                      PPV: 44.44% (21.49 to 67.40)                      NPV: 89.83% (82.12 to 97.54)                      LR+: 3.60 (1.74 to 7.45)                      LR-: 0.51 (0.28 to 0.94)</p> <p><b><u>Correlation of fetal scalp measurements with umbilical cord measurements (r value)*</u></b></p> <p>a. pH: 0.76                      b. Base excess: 0.90</p> <p>Note: this relates to 31 samples from uncomplicated, spontaneous births where the FBS was done within 5 minutes of birth</p> <p>* There is some discrepancy between data reported in the text and in the figures; data from the text have been reported here</p> <p><b>FBS pH &lt; 7.20 for Apgar &lt; 7 at 1 minute</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>8</td> <td>10</td> </tr> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>53</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	8	10	Predictive Test -ve	6	53	<p>Loss to follow-up is unrelated to key characteristics: Unclear                      Prognostic factors are adequately measured in participants: Yes                      Outcome of interest is sufficiently measured in participants: Yes                      Important potential confounders are accounted for: No, there are very few details and mode of birth is not reported                      Statistical analysis is appropriate for study design: Unclear</p> <p>They restricted sample to those within 30 minutes, but then added a further 5 patients as they didn't have sufficient data. In general, this study is very badly reported.</p> <p><b>Other information</b></p> <p><b><u>Additional details about babies with low scalp pH but born vigorous ('false positives')</u></b>                      Note: The detail provided about the 'false positives' does not use the same threshold for high Apgar as the rest of the data reported; therefore, not all</p>
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	8	10												
Predictive Test -ve	6	53												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					<p>of the false positives have extra data reported for them.</p> <p>Out of the 7 babies with abnormal pH but an Apgar of at least 8:</p> <ul style="list-style-type: none"> <li>- 2 had unknown causes</li> <li>- In one, there was transient uterine hypertonus due to oxytocin over-dosage, which was associated with marked and prolonged late decelerations.</li> <li>- In the remaining 4 cases, the presence of severe or moderate cord compression was suggested.</li> </ul>
<p><b>Full citation</b></p> <p>Wiberg-Itzel,E., Lipponer,C., Norman,M., Herbst,A., Prebensen,D., Hansson,A., Bryngelsson,A.L., Christoffersson,M., Sennstrom,M., Wennerholm,U.B., Nordstrom,L., Determination of pH or lactate in fetal scalp blood in management of intrapartum fetal distress: randomised controlled multicentre trial, BMJ, 336, 1284-1287, 2008</p> <p><b>Ref Id</b></p> <p>116763</p>	<p><b>Sample size</b></p> <p>N = 3007 randomised</p> <p><b>Characteristics</b></p> <p><b><u>Maternal age/years (mean (range))</u></b></p> <p>pH: 33.0 (19 - 49) Lactate: 32.5 (19 - 48)</p> <p><b><u>Parity (n (%))</u></b></p> <p>- Nulliparous pH: 1179 (78.8) Lactate: 1155 (77.2)</p>	<p><b>Tests</b></p> <p>pH analysis</p> <p>Lactate analysis</p> <p>[data are reported for within 60 minutes of birth]</p>	<p><b>Methods</b></p> <p>Antenatal clinics gave information about the study to women who were late in pregnancy, and requested consent either then or when the woman was admitted in labour. If consent was not obtained, or the woman was distressed, she was cared for according to the protocols of the department she was in.</p>	<p><b>Results</b></p> <p>The following data was reported in the trial, and this was used to calculate the diagnostic accuracy data below.</p> <p><b><u>Incidence of metabolic acidaemia (n/total (%))</u></b></p> <p><b>a. Split by pH status</b></p> <p>&gt; 7.25: 7/281 (2.5) 7.25 - 7.21: 3/92 (3.3) &lt; 7.21: 10/135 (7.4)</p> <p><b>b. Split by lactate status</b></p> <p>&lt; 4.2: 6/344 (1.7) 4.2 - 4.8: 0/73 (0)</p>	<p><b>Limitations</b></p> <p>Study sample represents population: unclear whether these women were definitely low risk during their pregnancy Loss to follow-up is unrelated to key characteristics: Not applicable because there was no loss to follow-up. However, there are some missing data: samples for cord pH measurement were missing in 174 in pH</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>Sweden</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>To examine the effectiveness of pH analysis of fetal scalp blood compared with lactate analysis in identifying hypoxia in labour to prevent acidaemia at birth</p> <p><b>Study dates</b></p> <p>December 2002 to December 2005</p> <p><b>Source of funding</b></p> <p>Signhild Engqvists Stiftelse, Almanna BB's Minnesfond, the regional city council research and development foundations, the health and medical committee of the region Vastra Gotaland, and Medexa, Lomma, Sweden</p>	<p>- Multiparous pH: 317 (21.2) Lactate: 341 (22.8)</p> <p><b><u>Gestational age/weeks+days (mean (range))</u></b> pH: 40+2 (34+0 - 44+2) Lactate: 40+3 (34+0 - 43+6)</p> <p><b><u>Fetal weight</u></b> <b><u>a. Mean/grams (range)</u></b> pH: 3575 (1590 - 5680) Lactate: 3566 (1860 - 6110)</p> <p><b><u>b. Proportion with fetal weight &lt; 2500 (n/total)</u></b> pH: 39/1496 Lactate: 36/1496</p> <p><b><u>Use of STAN monitor (n (%))</u></b> pH: 393 (26.2) Lactate: 392 (26.2)</p> <p><b><u>Inclusion Criteria</u></b></p> <p>Singleton pregnancy</p> <p>Cephalic presentation</p> <p>Gestational age ≥ 34 weeks</p> <p>Non-reassuring fetal heart rate trace that the clinician in charge</p>		<p>3007 women were randomised, and then 15 were excluded as per exclusion criteria.</p> <p>An internet based system was used for randomisation and data entry. Randomisation was stratified by department, and also by the use of electrocardiogram (ECG) as an adjunct to cardiotocography (CTG). At the point that the clinician decided to sample fetal scalp blood, the woman was randomised to either pH or lactate analysis. If sampling or analysis failed, management was based on other clinical information. Any crossover was regarded as a protocol violation.</p> <p>Scalp blood was sampled one to nine times for each fetus. In the pH group, successful sampling or analysis was performed in 1008 fetuses, with a total of 1628 analyses of pH. In the lactate group, successful sampling was done in 1355 fetuses, with a total of 2301</p>	<p>&gt; 4.8: 19/267 (7.1)</p> <p><b><u>Incidence of pH &lt; 7.00 at birth (n/total (%))</u></b></p> <p><b><u>a. Split by pH status</u></b> &gt; 7.25: 4/281 (1.4) 7.25 - 7.21: 2/92 (2.2) &lt; 7.21: 5/135 (3.7)</p> <p><b><u>b. Split by lactate status</u></b> &lt; 4.2: 0/344 (0) 4.2 - 4.8: 0/73 (0) &gt; 4.8: 10/267 (3.7)</p> <p><b><u>Incidence of Apgar &lt; 7 at 5 minutes (n/total (%))</u></b></p> <p><b><u>a. Split by pH status</u></b> &gt; 7.25: 9/281 (3.2) 7.25 - 7.21: 2/92 (2.2) &lt; 7.21: 10/135 (7.4)</p> <p><b><u>b. Split by lactate status</u></b> &lt; 4.2: 4/344 (1.2) 4.2 - 4.8: 1/73 (1.4) &gt; 4.8: 23/267 (8.6)</p> <p>The following diagnostic accuracy measures were calculated by the technical team, based on the above data. They refer to fetuses in whom fetal scalp blood was collected within 60 minutes of birth.</p> <p><b><u>Predictive accuracy of scalp pH &lt; 7.21</u></b></p> <p><b><u>a. For metabolic acidaemia</u></b> Sensitivity: 50.00% (28.09 to 71.91) Specificity: 74.39% (70.51 to 78.26) PPV: 7.41% (2.99 to 11.83) NPV: 97.32% (95.68 to 98.96) LR+: 1.95 (1.23 to 3.10)</p>	<p>arm and 120 in lactate arm; however, it is unclear whether these came from the subset of the study population with measurements done within 60 minutes of birth. Prognostic factors is adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: not really applicable - women were randomised to receive lactate or pH Statistical analysis is appropriate for study design: yes</p> <p><b>Other information</b></p> <p>This study is also included in the Cochrane review (East et al., 2010) which has been included in this review. However, further data are available from the full text of the trial. Data that have been reported in the Cochrane review will not be reported here.</p> <p>There were 155 protocol violations in the pH group</p>



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
	<p>considered to be an indication for FBS</p> <p><b>Exclusion Criteria</b></p> <p>Multiple pregnancy</p> <p>Gestational age &lt; 34 weeks</p>		<p>analyses.</p> <p>End points were metabolic acidaemia in cord blood (defined as a pH &lt; 7.05 and base deficit &gt; 12 mmol/l) and pH &lt; 7.00. Base deficit was calculated with the algorithm used by Radiometer blood gas analysers.</p> <p>Lactate was measured using a microvolume test strip device (Lactate Pro). Various pH analysers were used, but regular quality checks were performed. Guidelines for interpreting blood gas were:</p> <ul style="list-style-type: none"> <li>- pH &gt; 7.25 or lactate &lt; 4.2 mmol/l: normal</li> <li>- pH 7.21 - 7.25 or lactate 4.2 - 4.8 mmol/l: pre-acidaemia</li> <li>- pH &lt; 7.21 or lactate &gt; 4.8 mmol/l: acidaemia</li> </ul> <p>The guidelines for pre-acidaemia were to repeat the sample in 20-30 minutes if there was no other indication for intervention. For fetuses with acidaemia, the decision about delivery was left to the clinician.</p>	<p>LR-: 0.67 (0.43 to 1.05)</p> <p><b>b. For umbilical artery pH &lt; 7.00</b> Sensitivity: 45.45% (16.03 to 74.88) Specificity: 73.84% (69.98 to 77.71) PPV: 3.70% (0.52 to 6.89) NPV: 98.39% (97.11 to 99.67) LR+: 1.74 (0.89 to 3.38) LR-: 0.74 (0.43 to 1.27)</p> <p><b>c. For Apgar &lt; 7 at 5 minutes</b> Sensitivity: 47.62% (26.26 to 68.98) Specificity: 74.33% (70.45 to 78.21) PPV: 7.41% (2.99 to 11.83) NPV: 97.05% (95.33 to 98.77) LR+: 1.86 (1.16 to 2.98) LR-: 0.70 (0.47 to 1.06)</p> <p><b><u>Diagnostic accuracy of scalp pH ≤ 7.25</u></b></p> <p><b>a. For metabolic acidaemia</b> Sensitivity: 65.00% (44.10 to 85.90) Specificity: 56.15% (51.74 to 60.55) PPV: 5.73% (2.70 to 8.75) NPV: 97.51% (95.69 to 99.33) LR+: 1.48 (1.06 to 2.08) LR-: 0.62 (0.34 to 1.14)</p> <p><b>b. For umbilical artery pH &lt; 7.00</b> Sensitivity: 63.64% (35.21 to 92.06) Specificity: 55.73% (51.37 to 60.10) PPV: 3.08% (0.83 to 5.33) NPV: 98.58% (97.19 to 99.96) LR+: 1.44 (0.91 to 2.27) LR-: 0.65 (0.30 to 1.43)</p> <p><b>c. For Apgar &lt; 7 at 5 minutes</b> Sensitivity: 57.14% (35.98 to 78.31) Specificity: 55.85% (51.44 to 60.26) PPV: 5.29% (2.38 to 8.2)</p>	<p>(146 failed FBS and 9 failed analysis) and 18 in the lactate group (all failed sampling). However, data for these women would not be incorporated in this data, as they could not be classified by pH or lactate value.</p> <p>No fetal scalp blood was collected in 106 women in the pH arm and 81 in the lactate arm. In most cases a reason was not provided, however, some were as a result of rapid delivery, expedited delivery, reassuring CTG or the withdrawal of consent.</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
			<p>A sample size calculation calculated that a total of 2872 participants would be needed to detect a 100% increase in metabolic acidaemia with lactate, compared to a prevalence of 1.6% in the pH arm, with 80% power. To show a 50% reduction, 2907 cases in each arm would be needed. For the endpoint of pH &lt; 7.00, 1141 cases in each arm were needed to detect a 50% decrease or increase.</p> <p>Interim analyses were done after 1400 and 2400 randomised cases. Following the second analysis, the independent steering committee recommended stopping the trial after 3000 cases.</p> <p>Data was analysed on an intention-to-treat basis. Chi-squared and relative risks were used to compared pH and lactate groups. p &lt; 0.05 was considered significant.</p>	<p>NPV: 96.80% (94.74 to 98.86) LR+: 1.29 (0.88 to 1.90) LR-: 0.77 (0.47 to 1.27)</p> <p><b><u>Diagnostic accuracy of scalp lactate &gt; 4.8 mmol/l</u></b></p> <p><b>a. For metabolic acidaemia</b> Sensitivity: 76.00% (59.26 to 92.74) Specificity: 62.37% (58.67 to 66.07) PPV: 7.12% (4.03 to 10.2) NPV: 98.56% (97.42 to 99.70) LR+: 2.02 (1.59 to 2.57) LR-: 0.38 (0.19 to 0.78)</p> <p><b>b. For umbilical artery pH &lt; 7.00</b> Sensitivity: 100% (100 to 100) Specificity: 61.87% (58.20 to 65.54) PPV: 3.75% (1.47 to 6.02) NPV: 100% (100 to 100) LR+: 2.62 (2.38 to 2.89) LR-: 0.00 (not calculable [NC])</p> <p><b>c. For Apgar &lt; 7 at 5 minutes</b> Sensitivity: 82.14% (67.96 to 96.33) Specificity: 62.80% (59.11 to 66.50) PPV: 8.61% (5.25 to 11.98) NPV: 98.80% (97.76 to 99.85) LR+: 2.21 (1.81 to 2.70) LR-: 0.28 (0.13 to 0.63)</p> <p><b><u>Diagnostic accuracy of scalp lactate ≥ 4.2 mmol/l</u></b></p> <p><b>a. For metabolic acidaemia</b> Sensitivity: 76.00% (59.26 to 92.74) Specificity: 51.29% (47.47 to 55.11) PPV: 5.59% (3.15 to 8.03) NPV: 98.26% (96.87 to 99.64) LR+: 1.56 (1.24 to 1.97) LR-: 0.47 (0.23 to 0.94)</p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
				<p><u>b. For umbilical artery pH &lt; 7.00</u>                      Sensitivity: 100% (100 to 100)                      Specificity: 51.04% (47.26 to 54.81)                      PPV: 2.94% (1.15 to 4.74)                      NPV: 100% (100 to 100)                      LR+: 2.04 (1.89 to 2.21)                      LR-: 0.00 (NC)</p> <p><u>c. For Apgar &lt; 7 at 5 minutes</u>                      Sensitivity: 85.71% (72.75 to 98.68)                      Specificity: 51.83% (48.01 to 55.65)                      PPV: 7.06% (4.34 to 9.78)                      NPV: 98.84% (97.70 to 99.97)                      LR+: 1.78 (1.50 to 2.11)                      LR-: 0.28 (0.11 to 0.69)</p> <p><b><u>Operative delivery due to fetal distress in women in whom fetal scalp blood was taken within 60 minutes of delivery (n/total (%))</u></b></p> <p><u>a. In women randomised to pH analysis</u>                      pH &gt; 7.25: 81/281 (28.8)                      pH 7.21 - 7.25: 58/92 (63.0)                      pH &lt; 7.21: 118/135 (87.4)</p> <p><b><u>b. In women randomised to lactate analysis</u></b>                      Lactate &lt; 4.2: 79/334 (23.0)                      Lactate 4.2 - 4.8: 50/73 (68.5)                      Lactate &gt; 4.8: 251/267 (94.0)</p> <p><b>FBS &lt; 7.21 for metabolic acidaemia</b></p>	

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>10</td> <td>125</td> </tr> <tr> <td>Predictive Test -ve</td> <td>10</td> <td>363</td> </tr> </tbody> </table> <p><b>FBS &lt; 7.21 for UA pH &lt; 7.00</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>5</td> <td>130</td> </tr> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>367</td> </tr> </tbody> </table> <p><b>FBS &lt; 7.21 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>10</td> <td>125</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	10	125	Predictive Test -ve	10	363		Reference Test +ve	Reference Test -ve	Predictive Test +ve	5	130	Predictive Test -ve	6	367		Reference Test +ve	Reference Test -ve	Predictive Test +ve	10	125	
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	10	125																											
Predictive Test -ve	10	363																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	5	130																											
Predictive Test -ve	6	367																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	10	125																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																														
				<table border="1"> <tr> <td data-bbox="1319 272 1451 371">Predictive Test -ve</td> <td data-bbox="1451 272 1603 371">11</td> <td data-bbox="1603 272 1749 371">362</td> </tr> <tr> <td colspan="3" data-bbox="1319 427 1749 459"><b>FBS &lt;= 7.25 for metabolic acidaemia</b></td> </tr> <tr> <td data-bbox="1319 483 1451 582"></td> <td data-bbox="1451 483 1603 582"><b>Reference Test +ve</b></td> <td data-bbox="1603 483 1749 582"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1319 590 1451 683">Predictive Test +ve</td> <td data-bbox="1451 590 1603 683">13</td> <td data-bbox="1603 590 1749 683">214</td> </tr> <tr> <td data-bbox="1319 691 1451 783">Predictive Test -ve</td> <td data-bbox="1451 691 1603 783">7</td> <td data-bbox="1603 691 1749 783">274</td> </tr> <tr> <td colspan="3" data-bbox="1319 839 1749 871"><b>FBS &lt;= 7.25 for pH &lt; 7.00</b></td> </tr> <tr> <td data-bbox="1319 895 1451 994"></td> <td data-bbox="1451 895 1603 994"><b>Reference Test +ve</b></td> <td data-bbox="1603 895 1749 994"><b>Reference Test -ve</b></td> </tr> <tr> <td data-bbox="1319 1002 1451 1094">Predictive Test +ve</td> <td data-bbox="1451 1002 1603 1094">7</td> <td data-bbox="1603 1002 1749 1094">220</td> </tr> <tr> <td data-bbox="1319 1102 1451 1195">Predictive Test -ve</td> <td data-bbox="1451 1102 1603 1195">4</td> <td data-bbox="1603 1102 1749 1195">277</td> </tr> <tr> <td colspan="3" data-bbox="1319 1251 1749 1307"><b>FBS &lt;= 7.25 for Apgar &lt; 7 at 5 minutes</b></td> </tr> </table>	Predictive Test -ve	11	362	<b>FBS &lt;= 7.25 for metabolic acidaemia</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	13	214	Predictive Test -ve	7	274	<b>FBS &lt;= 7.25 for pH &lt; 7.00</b>				<b>Reference Test +ve</b>	<b>Reference Test -ve</b>	Predictive Test +ve	7	220	Predictive Test -ve	4	277	<b>FBS &lt;= 7.25 for Apgar &lt; 7 at 5 minutes</b>			
Predictive Test -ve	11	362																																	
<b>FBS &lt;= 7.25 for metabolic acidaemia</b>																																			
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																																	
Predictive Test +ve	13	214																																	
Predictive Test -ve	7	274																																	
<b>FBS &lt;= 7.25 for pH &lt; 7.00</b>																																			
	<b>Reference Test +ve</b>	<b>Reference Test -ve</b>																																	
Predictive Test +ve	7	220																																	
Predictive Test -ve	4	277																																	
<b>FBS &lt;= 7.25 for Apgar &lt; 7 at 5 minutes</b>																																			

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
				<table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>12</td> <td>215</td> </tr> <tr> <td>Predictive Test -ve</td> <td>9</td> <td>272</td> </tr> </tbody> </table> <p>Lactate &gt; 4.8 for metabolic acidaemia</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>19</td> <td>248</td> </tr> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>411</td> </tr> </tbody> </table> <p>Lactate &gt; 4.8 for UA pH &lt; 7.00</p> <table border="1"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>10</td> <td>257</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	12	215	Predictive Test -ve	9	272		Reference Test +ve	Reference Test -ve	Predictive Test +ve	19	248	Predictive Test -ve	6	411		Reference Test +ve	Reference Test -ve	Predictive Test +ve	10	257	
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	12	215																											
Predictive Test -ve	9	272																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	19	248																											
Predictive Test -ve	6	411																											
	Reference Test +ve	Reference Test -ve																											
Predictive Test +ve	10	257																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																					
				<table border="1" data-bbox="1319 272 1749 371"> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>417</td> </tr> </table> <p data-bbox="1319 427 1666 480"><b>Lactate &gt; 4.8 for Apgar &lt; 7 at 5 minutes</b></p> <table border="1" data-bbox="1319 509 1749 810"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>23</td> <td>244</td> </tr> <tr> <td>Predictive Test -ve</td> <td>5</td> <td>412</td> </tr> </tbody> </table> <p data-bbox="1319 866 1637 919"><b>Lactate &gt;= 4.2 for metabolic acidaemia</b></p> <table border="1" data-bbox="1319 948 1749 1249"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>19</td> <td>321</td> </tr> <tr> <td>Predictive Test -ve</td> <td>6</td> <td>338</td> </tr> </tbody> </table> <p data-bbox="1319 1305 1671 1332"><b>Lactate &gt;= 4.2 for UA pH &lt; 7.00</b></p>	Predictive Test -ve	0	417		Reference Test +ve	Reference Test -ve	Predictive Test +ve	23	244	Predictive Test -ve	5	412		Reference Test +ve	Reference Test -ve	Predictive Test +ve	19	321	Predictive Test -ve	6	338	
Predictive Test -ve	0	417																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	23	244																								
Predictive Test -ve	5	412																								
	Reference Test +ve	Reference Test -ve																								
Predictive Test +ve	19	321																								
Predictive Test -ve	6	338																								

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
				<table border="1" data-bbox="1319 272 1749 572"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>10</td> <td>330</td> </tr> <tr> <td>Predictive Test -ve</td> <td>0</td> <td>344</td> </tr> </tbody> </table> <p data-bbox="1319 632 1682 683"><b>Lactate <math>\geq</math> 4.2 for Apgar <math>&lt;</math> 7 at 5 minutes</b></p> <table border="1" data-bbox="1319 715 1749 1011"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>24</td> <td>316</td> </tr> <tr> <td>Predictive Test -ve</td> <td>4</td> <td>340</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	10	330	Predictive Test -ve	0	344		Reference Test +ve	Reference Test -ve	Predictive Test +ve	24	316	Predictive Test -ve	4	340	
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	10	330																					
Predictive Test -ve	0	344																					
	Reference Test +ve	Reference Test -ve																					
Predictive Test +ve	24	316																					
Predictive Test -ve	4	340																					
<p data-bbox="192 1027 338 1054"><b>Full citation</b></p> <p data-bbox="192 1083 633 1249">Young,D.C., Gray,J.H., Luther,E.R., Peddle,L.J., Fetal scalp blood pH sampling: its value in an active obstetric unit, American Journal of Obstetrics and Gynecology,Am.J.Obstet.Gynecol., 136, 276-281, 1980</p> <p data-bbox="192 1275 271 1302"><b>Ref Id</b></p> <p data-bbox="192 1331 282 1358">159915</p>	<p data-bbox="633 1027 786 1054"><b>Sample size</b></p> <p data-bbox="633 1083 920 1302">N = 232 women (Note: the last scalp sample was taken less than 1 hour before birth in 95 women, and they constitute the true population of interest)</p>	<p data-bbox="920 1027 1028 1137"><b>Tests</b></p> <p data-bbox="920 1083 1028 1137">Fetal scalp pH</p>	<p data-bbox="1028 1027 1137 1054"><b>Methods</b></p> <p data-bbox="1028 1083 1312 1385">232 women had a total of 335 pH determinations done (mean 1.5 per patient, range 1 to 5). 98% of sampling was due to changes in fetal heart rate. 95% of the samples in the study were done with the patients in a modified Sims' position. A</p>	<p data-bbox="1312 1027 1406 1054"><b>Results</b></p> <p data-bbox="1312 1083 1756 1331">The following diagnostic accuracy measures have been calculated by the technical team, based on 2x2 data that was reported in the study. The data only relate to babies born within 1 hour of the fetal pH measurement. 136 babies who had a pH <math>\geq</math> 7.25 and were born over an hour after the measurement were not included for these calculations:</p> <p data-bbox="1312 1358 1697 1385"><b><u>Diagnostic accuracy for neonatal</u></b></p>	<p data-bbox="1756 1027 1899 1054"><b>Limitations</b></p> <p data-bbox="1756 1083 2042 1369">Study sample represents population: there was a high proportion of women who would not be considered low risk Loss to follow-up is unrelated to key characteristics: there was no loss to follow up Prognostic factor is</p>																		



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>Canada</p> <p><b>Study type</b></p> <p><b>Aim of the study</b></p> <p>To determine:</p> <ul style="list-style-type: none"> <li>- indications for fetal blood pH sampling</li> <li>- the incidence of fetal acidosis with each indication</li> <li>- incidence of neonatal depression related to fetal acidosis</li> <li>- complications of fetal blood sampling (FBS)</li> <li>- number of caesarean sections avoided</li> <li>- number of asphyxiated infants born less than 1 hour after fetal blood sampling</li> </ul> <p><b>Study dates</b></p> <p>January 1st 1978 to September 30th 1978</p> <p><b>Source of funding</b></p> <p>Life Insurance Association of Canada</p>	<p><b>Characteristics</b></p> <p><b><u>Time between last FBS and birth (n (%))</u></b></p> <p>&lt; 1 hour: 95 (40.9) 1 - 2 hours: 67 (28.9) &gt; 2 hours: 70 (30.2)</p> <p><b><u>Obstetric characteristics (n (%))</u></b></p> <p>Pre-eclampsia toxemia: 37 (16) Premature rupture of membranes: 23 (10) intrauterine growth restriction (IUGR): 19 (8) Prematurity: 9 (4) Post-maturity: 32 (14) Meconium-stained fluid: 77 (33) Oxytocin induced labour: 103 (44) Oral prostaglandin: 16 (7) Nulliparous: 162 (70) Epidural: 175 (75) Parenteral narcotic &lt; 6 hours: 53 (23)</p> <p><b><u>Indication for fetal blood sampling (n (%))</u></b></p> <p>Baseline: - Tachycardia: 14 (6) - Bradycardia: 15 (6)</p> <p>Decreased variability: 24 (10)</p> <p>Variable decelerations: - Mild: 22 (10)</p>		<p>Monoject Sterile Disposable Fetal Blood Sampling Kit was used for sample collection, and results were available within 10 minutes of sampling.</p> <p>The fetal heart trace in the hour before FBS were analysed and classified using ACOG Technical Bulletin 32, and in addition as follows:</p> <ul style="list-style-type: none"> <li>- Mild decelerations: less than 30 bpm in depth</li> <li>- Moderate decelerations: 30 - 60 bpm in depth</li> <li>- Severe decelerations: greater than 60 bpm in depth</li> <li>- Persistent decelerations: longer than 30 minutes and with more than 50% of contractions</li> <li>- Variable decelerations that did not return to baseline were considered indicative of late recovery</li> </ul> <p>The FHR tracings were reviewed by members of the Perinatal Medicine Division without knowledge of pH values, to try and estimate whom</p>	<p><b><u>depression (95% CI)</u></b></p> <p><b>a. pH &lt; 7.20</b> Sensitivity: 37.50% (3.95 to 71.05) Specificity: 96.59% (92.80 to 100) PPV: 50.00% (9.99 to 90.01) NPV: 94.44% (89.71 to 99.18) LR+: 11.00 (2.64 to 45.84) LR-: 0.65 (0.38 to 1.11)</p> <p><b>b. pH &lt; 7.25</b> Sensitivity: 50.00% (15.35 to 84.65) Specificity: 81.82% (73.76 to 89.88) PPV: 20.00% (2.47 to 37.53) NPV: 94.74% (89.72 to 99.76) LR+: 2.75 (1.21 to 6.26) LR-: 0.61 (0.30 to 1.23)</p> <p>The following data relate to the entire study population:</p> <p><b><u>Proportion of women having caesarean section (n/total (%))</u></b></p> <p>pH &lt; 7.20: 6/6 (100) - all 6 born within 1 hour of pH measurement</p> <p>pH 7.20 - 7.24: 7/14 (50) - all 14 born within 1 hour of pH measurement</p>	<p>adequately measured in participants: yes Outcome of interest is sufficiently measured in participants: yes Important potential confounders are accounted for: there were differences in the proportion of babies born by CS, and this is not reported for the sub-group of babies with normal pH but who were born within an hour Statistical analysis is appropriate for study design: yes</p> <p>Indirectness of population: yes, a high proportion of women were not low risk</p> <p><b>Other information</b></p> <p><b><u>Further information regarding babies with severe fetal acidosis (pH &lt; 7.20) in labour True positives (depressed at birth) Baby 1</u></b></p> <ul style="list-style-type: none"> <li>- had severe pre-eclamptic toxemia</li> <li>- fetal pH of 7.12</li> <li>- 32 minutes before birth</li> <li>- Apgar of 1 at 1 minute</li> </ul>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
	<p>- Moderate: 84 (36) - Severe: 38 (16)</p> <p>Late decelerations: - Mild: 19 (8) - Moderate: 5 (2)</p> <p>Early decelerations: 7 (3)</p> <p>Other indications: 4 (2)</p> <p><b>Inclusion Criteria</b></p> <p>All patients having fetal scalp blood pH sampling (98% were due to fetal heart rate changes)</p> <p><b>Exclusion Criteria</b></p> <p>None reported</p>		<p>they would have performed a caesarean on without knowledge of pH values. For this, only patients with less than full dilatation of the cervix and who subsequently delivered vaginally were included.</p> <p>Fetal acidosis was classified as: - Mild: pH 7.20 - 7.24 - Severe: &lt; 7.20</p> <p>Neonatal depression was defined as one of: - 1 minute Apgar less than 7 and the need for positive pressure resuscitation - 5 minute Apgar less than 7</p>	<p>pH ≥ 7.25: 40/212 (19) - 76 born within 1 hour, 66 born within 1-2 hours, 70 born over 2 hours later</p> <p>Note: the overall CS rate was 23%, of which 25% were performed for fetal distress.</p> <p><b>Complications of fetal blood sampling (n (%))</b></p> <p>Bleeding: - Haematoma: 6 (2.6) - Abrasions: 3 (1.3) - Ecchymosis: 1 (0.4) - Anaemia of unknown etiology: 1 (0.4)</p> <p>Infection: - Abscess: 1 (0.4) - Cellulitis: 1 (0.4) - Erythema: 1 (0.4) - Herpes: 1 (0.4)</p> <p>Total: 15 (6.5)</p> <p><b>FBS pH &lt; 7.20 for neonatal depression</b></p> <table border="1" data-bbox="1317 1077 1749 1380"> <thead> <tr> <th></th> <th>Reference Test +ve</th> <th>Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td>Predictive Test +ve</td> <td>3</td> <td>3</td> </tr> <tr> <td>Predictive Test -ve</td> <td>5</td> <td>85</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	Predictive Test +ve	3	3	Predictive Test -ve	5	85	<p>and 3 at 5 minutes</p> <p>- FHR tracing decelerations: persistent, mild, late - cord pH 7.21/7.11</p> <p><u>Baby 2</u></p> <p>- had meconium and died at about 4 hours - fetal pH of 6.74 - 37 minutes before birth - Apgar of 0 at 1 minute and 1 at 5 minutes - FHR tracing decelerations: persistent, moderate, late - cord pH 6.79/6.60</p> <p><u>Baby 3</u></p> <p>- post-mature, hypertension, prior stillbirth - fetal pH of 6.94 - 41 minutes before birth - Apgar of 1 at 1 minute and 4 at 5 minutes - FHR tracing decelerations: occasional severe, variable, late recovery, decreasing variability - cord pH 7.14/7.09</p> <p><b><u>False positives (normal Apgar scores)</u></b></p> <p><u>Baby 4</u></p> <p>- chronic active hepatitis - fetal pH of 7.19 - 58 minutes before birth - Apgar of 9 at 1 minute</p>
	Reference Test +ve	Reference Test -ve												
Predictive Test +ve	3	3												
Predictive Test -ve	5	85												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments									
				<p><b>FBS pH &lt; 7.25 for neonatal depression</b></p> <table border="1" data-bbox="1319 411 1747 711"> <thead> <tr> <th data-bbox="1319 411 1451 517"></th> <th data-bbox="1451 411 1603 517">Reference Test +ve</th> <th data-bbox="1603 411 1747 517">Reference Test -ve</th> </tr> </thead> <tbody> <tr> <td data-bbox="1319 517 1451 612"><b>Predictive Test +ve</b></td> <td data-bbox="1451 517 1603 612">4</td> <td data-bbox="1603 517 1747 612">16</td> </tr> <tr> <td data-bbox="1319 612 1451 711"><b>Predictive Test -ve</b></td> <td data-bbox="1451 612 1603 711">4</td> <td data-bbox="1603 612 1747 711">72</td> </tr> </tbody> </table>		Reference Test +ve	Reference Test -ve	<b>Predictive Test +ve</b>	4	16	<b>Predictive Test -ve</b>	4	72	<p>and 10 at 5 minutes                      - FHR tracing                      decelerations: persistent, moderate, variable late recovery                      - cord pH</p> <p><u>Baby 5</u>                      - true knot in cord                      - fetal pH of 7.19                      - 45 minutes before birth                      - Apgar of 9 at 1 minute and 10 at 5 minutes                      - FHR tracing                      decelerations: persistent mild late                      - cord pH 7.26/7.20</p> <p><u>Baby 6</u>                      - 32 weeks, pre-eclamptic toxaemia, abruptio placentae                      - fetal pH of 7.16                      - 38 minutes before birth                      - Apgar of 7 at 1 minute and 8 at 5 minutes                      - FHR tracing                      decelerations: persistent mild late                      - cord pH 7.19/7.17</p> <p><b><u>Further information regarding babies whose pH was <math>\geq</math> 7.25 but were born depressed (false negatives)</u></b></p> <p><u>Baby 1</u>                      - meconium, analgesic at 3 hours</p>
	Reference Test +ve	Reference Test -ve												
<b>Predictive Test +ve</b>	4	16												
<b>Predictive Test -ve</b>	4	72												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					<ul style="list-style-type: none"> <li>- fetal pH of 7.36</li> <li>- 54 minutes before birth (vaginal birth)</li> <li>- Apgar of 4 at 1 minute and 6 at 5 minutes</li> <li>- FHR tracing</li> <li>decelerations: moderate variable late recovery</li> <li>- cord pH 7.27/7.11</li> </ul> <p><u>Baby 2</u></p> <ul style="list-style-type: none"> <li>- meconium aspiration</li> <li>- fetal pH of 7.34</li> <li>- 50 minutes before birth (vaginal birth)</li> <li>- Apgar of 4 at 1 minute and 8 at 5 minutes</li> <li>- FHR tracing</li> <li>decelerations: moderate variable</li> <li>- cord pH 7.14/7.10</li> </ul> <p><u>Baby 3</u></p> <ul style="list-style-type: none"> <li>- IUGR</li> <li>- fetal pH of 7.25</li> <li>- 38 minutes before birth (vaginal birth)</li> <li>- Apgar of 4 at 1 minute and 6 at 5 minutes</li> <li>- FHR tracing</li> <li>decelerations: moderate variable late recovery</li> <li>- cord pH 7.25/7.02</li> </ul> <p><u>Baby 4</u></p> <ul style="list-style-type: none"> <li>- meconium</li> <li>- fetal pH of 7.37</li> <li>- 45 minutes before birth (vaginal birth)</li> <li>- Apgar of 6 at 1 minute</li> </ul>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
					and 9 at 5 minutes - FHR tracing decelerations: mild early - cord pH 7.37/7.34

**G.10 Women's experience of fetal monitoring**

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Parisaei,M., Harrington,K.F., Erskine,K.J., Maternal satisfaction and acceptability of foetal electrocardiographic (STAN[REGISTERED]) monitoring system, Archives of Gynecology and Obstetrics, 283, 31-35, 2011</p> <p><b>Ref Id</b></p> <p>134248</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Prospective questionnaire-based study</p> <p><b>Aim of the study</b></p> <p>To assess the acceptability of the fetal electrocardiographic (STAN®) monitoring system by women at a London Hospital</p> <p><b>Study dates</b></p> <p>November 2003 to June 2005</p>	<p><b>Sample size</b></p> <p>Total n = 125</p> <p><b>Characteristics</b></p> <p>Population consisted of women with high-risk pregnancy (diabetes, pre-eclampsia, previous caesarean section) or intrapartum risk factors (meconium stained liquor, oxytocin augmentation); 78% were believed to be low risk at their antenatal booking appointment</p> <p>Mean age (years): 28.8 (SD 6.3)</p> <p>Nulliparous: 75%</p> <p>Spoke English fluently: 83%</p> <p><u>Ethnicity</u></p> <p>African: 40%</p> <p>White: 30%</p> <p>Asian: 10%</p> <p>Other: 20%</p> <p><u>Intrapartum characteristics in cohort of women being monitored by STAN</u></p> <p>Induction of labour: 37%</p> <p>Meconium stained liquor: 50%</p> <p>Epidural use: 80%</p> <p>Fetal blood sampling performed: 13%</p>	<p><b>Interventions</b></p> <p>Fetal electrocardiographic (STAN) monitoring</p>	<p><b>Details</b></p> <p>A questionnaire was designed to assess women's acceptability for STAN. The study was conducted in a university hospital in East London with 4000 births per year. Women who had STAN monitoring were provided with information sheets about the study. Women were asked to fill in the questionnaire after the birth (the majority of women filled in the questionnaire on the day of the birth). The information sheet and the questionnaire were reviewed by a clinical psychologist; n = 125 women were monitored with STAN during the study period. The questionnaire consisted of 7 yes/no questions and space was provided for further comments.</p>	<p><b>Results</b></p> <p>1) Did the midwife(s) looking after you in labour explain the reasons why your baby was monitored continuously in labour? Yes: 93% (CI 85% to 98%)</p> <p>2) Did the doctor(s) looking after you in labour explain the reasons why your baby was monitored continuously in labour? Yes: 99% (CI 83% to 99.9%)</p> <p>3) Did you understand how the STAN system monitors your baby's wellbeing in labour? Yes: 95% (CI 87% to 99%)</p> <p>4) Did you think the STAN system is an acceptable additional way of monitoring your baby in labour? Yes: 95% (CI 87% to 99%)</p> <p>5) Did you feel reassured by having the STAN system as well as the CTG monitor in labour? Yes: 96% (CI 89% to 99%)</p> <p>6) Would you have the STAN system again in future labours if we needed</p>	<p><b>Limitations</b></p> <p>Unclear whether the questionnaire was a validated tool or not</p> <p>Unclear how the questionnaire was developed and by whom</p> <p>Questionnaire response rate was 61% (77/125)</p> <p>Unclear how the data were analysed and by whom</p> <p>Unclear what explanation was given to women about the reasons why the baby was monitored continuously in labour</p> <p>13.3% of study population had a language problem</p> <p>Unclear whether women received unbiased information about STAN and how it assesses the baby's wellbeing</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Source of funding</b></p> <p>Not reported</p>	<p>Syntocinon infusion utilised: 67%</p> <p>Spontaneous vaginal birth: 29%</p> <p>Emergency caesarean section (CS): 54% (215 of these were for fetal distress according to STAN clinical protocol)</p> <p><b>Inclusion criteria</b></p> <p>Term pregnancy (&gt; 37 weeks' gestation)</p> <p>Singleton pregnancy</p> <p><b>Exclusion criteria</b></p> <p>Multiple pregnancy</p> <p>Women with viral infection (HIV or hepatitis B and C)</p>		<p>Analysis: Dichotomous and categorical data were summarised using percentages and hypothesis tests. Continuous data were summarised using mean for normally distributed data and median for non-normal data</p>	<p>further information about your baby's wellbeing in labour?</p> <p>Yes: 93% (CI 85% to 98%)</p> <p>7) Would you recommend the STAN system to your friends who are going to be mothers?</p> <p>Yes: 89% (CI 80% to 95%); the majority would only do so if they were high risk and there was a need for continuous fetal monitoring</p>	
<p><b>Full citation</b></p> <p>Hindley,C., Hinliff,S.W., Thomson,A.M., Pregnant women's views about choice of intrapartum monitoring of the fetal heart rate: a questionnaire survey, International Journal of Nursing Studies, 45, 224-231, 2008</p> <p><b>Ref Id</b></p> <p>136975</p>	<p><b>Sample size</b></p> <p>Total n = 63</p> <p><b>Characteristics</b></p> <p><b><u>Antepartum sample Total n = 63</u></b></p> <p><u>Gestational age when questionnaire completed</u></p> <p>34-36 weeks 6 days n = 45</p> <p>37-40 weeks n = 18</p> <p><u>Age (years)</u></p> <p>Under 20 n = 3</p>	<p><b>Interventions</b></p> <p>Intrapartum electronic fetal monitoring (EFM)</p>	<p><b>Details</b></p> <p>A total of 63 pregnant women at low obstetric risk were approached to complete antepartum and postpartum questionnaires. The sample was recruited from two maternity hospitals (centre 1 n = 30; centre 2 n = 33). After gaining informed consent, women were</p>	<p><b>Results</b></p> <p><b><u>Women's preference for electronic fetal monitoring (EFM)</u></b></p> <p><u>Antenatal survey (n = 63)</u></p> <p>Women did not prefer one specific option. The majority preferred a combination of intermittent and continuous EFM n = 35/63 (56%)</p> <p><u>Postnatal survey (n = 38)</u></p> <p>Number of women received EFM n = 23/38 (61%)</p>	<p><b>Limitations</b></p> <p>Participants recruited from two different hospitals, the influence of different setting should be considered when interpreting the data</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Qualitative exploratory/descriptive</p> <p><b>Aim of the study</b></p> <p>To investigate women's view on intrapartum fetal monitoring techniques and informed choice</p> <p><b>Study dates</b></p> <p>Not specified</p> <p><b>Source of funding</b></p> <p>NHS, Northern region Research and Development Directorate</p>	<p>20-24 n = 14 25-29 n = 20 30-34 n = 20 35-39 n = 6</p> <p><b>Ethnicity</b> White n = 49 Other n = 12 Missing n = 2</p> <p><b>Jarman deprivation score</b> Low deprivation (30 - 39.99) n = 14 Not deprived (below 30) n = 48 Missing n = 1</p> <p><b>Educational qualifications</b> No recorded qualification n = 2 Secondary education qualification n = 9 Further education qualification n = 38 Higher education n = 14</p> <p><b>Parity</b> Primigravida n = 31 Multigravida n = 32</p> <p><b>Postpartum sample n = 38</b> <b>Completion of questionnaire in weeks postpartum</b> 0-2 weeks n = 24 3-4 weeks n = 8 &gt; 5 weeks n = 5 Missing n = 1</p> <p><b>Type of birth</b> Normal Instrumental Emergency caesarean section</p> <p><b>Analgesia</b> Epidural n = 8</p>		<p>asked to complete the first questionnaire between 34 and 40 weeks of pregnancy. Sixty-three (n = 63) women completed the antepartum questionnaire; 38 of them also completed the postpartum questionnaire.</p> <p><b>Questionnaire</b> A validated tool (from an informed choice across maternity care) was modified and used for women's preferences of fetal monitoring. The developed questionnaire was piloted with a small sample and modified according to the results. Themes chosen for the questionnaire were identified from a background literature review. The antepartum questionnaire contained 28 items and aimed to elicit information on women's knowledge and preferences of intrapartum fetal monitoring. The</p>	<p><b><u>Women's preference for mobility during labour</u></b> <b>Antenatal survey</b> Stay mobile or off the bed n = 46/63 (73%) <b>Postnatal survey</b> Women reported staying in bed n = 16/38 (40%)</p> <p><b><u>Women's preference for decision making on fetal monitoring</u></b> <b>Antenatal survey</b> Women wanted the final decision after considering midwife's view: antepartum n = 35/63 (56%); intrapartum n = 28/63 (44%) <b>Postnatal survey</b> Women had conceded decision making to midwife in intrapartum period n = 14/38 (38%)</p> <p><b><u>Choice/control preference</u></b> <b>Antenatal survey</b> Felt choice of being in control is important n = 61/63 (97%) Felt midwives did not facilitate a choice in intrapartum fetal method antenatally n = 59/63 (94%) Not received enough information and discussion to make a choice regarding fetal monitoring method n = 25/63 (40%)</p> <p><b><u>Importance of information</u></b> <b>Antenatal survey</b> Women were aware of different types of monitoring</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Narcotic n = 12                      Entonox n = 11                      Other n = 3                      None n = 4  <u>Age (years)</u>                      Under 20 n = 1                      20-24 n = 5                      25-29 n = 10                      30-34 n = 17                      35-39 n = 5  <u>Ethnicity</u>                      White n = 30                      Others n = 7                      Missing n = 1  <u>Jarman deprivation score</u>                      Low deprivation (30 - 39.99) n = 7                      Not deprived (below 30) n = 30                      Missing n = 1  <u>Parity</u>                      Primigravida n = 16                      Multigravida n = 22</p> <p><b>Inclusion criteria</b></p> <p>Women with no underlying medical condition (low-risk pregnancy)                      Predicted a vaginal birth</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>postpartum questionnaire had 21 items and asked for information about monitoring preferences for labour and actual monitoring outcomes</p> <p><u>Data collection</u>                      Women were approached at 34 weeks of their pregnancy at the antenatal clinic. The midwife was the first point of contact, referring suitable women to the researcher to discuss the study in detail. An information pack plus the questionnaire and a stamped envelope were given to women. Women who did not return their questionnaire were approached in their next antenatal visit and reminded about the study (only one reminder was permitted based on ethics committee's approval). Following women's birth of a healthy infant, they were sent the postpartum</p>	<p>n = 59/63 (94%)                      Knew all types of monitoring except Pinard stethoscope n = 46/63 (73%)                      Felt it is very important to have information on intrapartum fetal monitoring n = 54/63 (86%)  <u>Postnatal survey</u>                      Felt it is very important to have information on intrapartum fetal monitoring n = 15/38 (39%)  <b>Sources of information</b>  <u>Antenatal survey</u>                      Felt midwife had not explicitly given any information on monitoring n = 41/63 (65%)                      Felt had the information from media n = 36/63 (57%)                      Women relied on past experience n = 29/63 (46%)                      Felt had informed choice or partially had informed choice n = 25/63  <u>Postnatal survey</u>                      Felt that they have been given informed choice n = 15/38 (39%)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>questionnaire and stamped addressed envelope, together with a letter of congratulations. Women were not followed up if they failed to respond.</p> <p><u>Data analysis</u> The data were analysed using SPSS 10.1. The analysis of data was descriptive. Frequency count and cross-tabulations were used.</p>		
<p><b>Full citation</b> Shields,D., Fetal and maternal monitoring: maternal reactions to fetal monitoring, American Journal of Nursing, 78, 2110-2112, 1978</p> <p><b>Ref Id</b> 170538</p> <p><b>Country/ies where the study was carried out</b> Canada</p> <p><b>Study type</b> Prospective observational study</p>	<p><b>Sample size</b> Total n = 30</p> <p><b>Characteristics</b> Age: ranged from 17 to 42 years Married: n = 19, single: n = 9, separated: n = 2 White: n = 16 Black: n = 14 Primiparous: n = 18 Multiparous: n = 12 <u>Reason women were monitored</u> Failure to progress and oxytocin stimulation: n = 7 Induced labour: n = 18 Poor obstetrical history: n = 1</p>	<p><b>Interventions</b> Internal electronic fetal monitoring</p>	<p><b>Details</b> The time that women were monitored ranged from 1 hour to 12 hours (no more details about the monitoring machine reported). To assess the general attitudes of women regarding fetal monitoring, the study author developed a 'mood and feeling inventory'. The scale consisted of a list of adjectives that women marked according to their feelings in a scale ranging from 1 (not at</p>	<p><b>Results</b> <u>Scores</u> Women in positive range: n = 22 Women in negative range: n = 8 Highly negative category: n = 2 Highly positive category: n = 3</p> <p>One woman had a high negative score (-3.46). She expressed a high degree of negativity throughout the interview. She expressed that she received 'too little information about the equipment', and did not like</p>	<p><b>Limitations</b> Data and results poorly reported. Very old study, advances in technology should be considered when interpreting the data. A self-developed scale used with unclear validity; 18/30 women were multiparous</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To examine women's experience and reaction to fetal monitoring</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>Research on normal labour: n = 4</p> <p><u>Mode of birth</u></p> <p>Spontaneous vaginal birth: n = 8</p> <p>Forceps delivery: n = 13</p> <p>Vacuum extraction: n = 2</p> <p>Caesarean section: n = 7</p> <p><u>Mean length of labour</u></p> <p>Multiparous: n = 6 hours and 26 min</p> <p>Nulliparous: n = 12 hours and 9 min</p> <p>Mean duration of monitoring: 5 hours and 16 min</p> <p><b>Inclusion criteria</b></p> <p>Women who had internal fetal monitoring during labour and gave birth at term</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>all) to 6 (very much). The negative scale consisted of eight words; apprehensive, uneasy, tense, frightened, worried, upset, nervous. The positive scale consisted of six words; relaxed, confident, peaceful, comfortable, optimistic, calm. Women were asked to mark the scale regarding their feelings during fetal monitoring retrospectively (as they remembered). Women were interviewed by the author within 48 hours of the birth. Their positive or negative attitudes toward the monitoring experience were assessed. Interviews were carried out using an open-ended questionnaire.</p> <p><u>Analysis</u></p> <p>A positive and a negative response for each woman was tabulated and a mean score was calculated. The negative score was subtracted from the positive score and the difference served</p>	<p>the idea of attaching it to the baby's head. She felt that, the monitoring was not a good indicator of what was happening; while she was in severe pain, she was told by the nurse that the equipment showed mild pain. She also expressed that 'the head is the most important part and I was worried about brain damage because of the clamp'.</p> <p>The woman with the highest negative score (-3.75) said she 'felt like a battery being charged with all those wires and connections'. From three women who had a high positive score, one woman with a score of 4.17, said she 'Knew exactly what was going on and therefore was not afraid'. A woman with a score of 4.45, was a 'little frightened' but thought it was an 'exciting idea' compared with other labours and felt that 'monitoring seemed to make it shorter and more interesting'. The woman with the highest positive score of 4.87 thought monitoring was 'a fantastic, good idea'. No differences were observed between these five women with the rest of the study's population.</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>as an indication of an overall positive or negative reaction. The maximum difference of 5 that could happen between the positive and negative scores of an individual woman were divided into high, medium, or low, positive or negative and women were placed by their scores in those categories</p>	<p>When a Chi- square computation was performed between the inventory scores and the age, race, parity, marital status length of labour and length of monitoring, no significant difference in the results were observed.</p> <p><u>Understanding the reason for monitoring</u> (determined by comparing women's response to the reason for monitoring, to the reason given in the women's charts):                      Good understanding: n = 27                      Partially understood: n = 3 (n = 2/3 were women with high negative score)</p> <p><u>Information received</u>                      Adequate: n = 27 (20 said they had full information and 7 said they received as much as they requested)                      No adequate information received: n = 3</p> <p><u>Nurse's presence</u>                      All women expressed their desire about wanting nurses to stay with them all the time; n = 17 wanted nurses for supportive care; n = 6 expressed a desire for the nurse's presence as a person that could intervene in some way if necessary.</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p><u>Worries about monitoring</u>                      No worries: n =7                      Some worries (not the same as those during pregnancy): n = 11 (4 expressed fears related to the electrodes)                      Some worries (the same as those during pregnancy): n = 12 (fearing that baby would be deformed in some way or die)</p> <p><u>Complain about monitoring</u>                      Getting comfortable: the most frequent complaint was with regard to difficulty in getting comfortable. Some women were annoyed about the fact that when the electrode fell off, an additional vaginal examination was needed to reapply the electrode.                      Complaints about vaginal examination mainly related to privacy and too many people being present in the room.                      Noise of fetal heart beat: was considered discomforting by 2 women because of fears that it would stop (one expressed that she 'worried the whole time that the baby's heart would stop if the machine stopped').</p> <p><u>Caregiveres</u>                      Four (n = 4) women</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				expressed that the clinicians were the cause of some discomfort for them. Two of these women considered the facial expression of the physician frightening. The other 2 women thought that some staff were unfamiliar with the machine and they found this disquieting. One woman thought that the clinician had more interest in the machine than they did with her, she said 'they all came with the machine and they all left with the machine'	
<p><b>Full citation</b></p> <p>Hansen,P.K., Smith,S.F., Nim,J., Neldam,S., Osler,M., Maternal attitudes to fetal monitoring, European Journal of Obstetrics, Gynecology, and Reproductive Biology, 20, 43-51, 1985</p> <p><b>Ref Id</b></p> <p>171177</p> <p><b>Country/ies where the study was carried out</b></p> <p>Denmark</p> <p><b>Study type</b></p> <p>Prospective observational study</p>	<p><b>Sample size</b></p> <p>Total n = 655</p> <p><b>Characteristics</b></p> <p>A: preferred auscultation (AUS-P), B: preferred electronic fetal monitoring (EFM-P), C: undecided (UD), p (A:B), p (a:b:c)</p> <p><u>Number</u></p> <p>AUS-P: n = 212 EFM-P: n = 259 UD: 184</p> <p><u>Age (mean ± SD)</u></p> <p>AUS-P: 27.8 ± 4.7 EFM-P: 28.1 ± 5.1 UD: 26.3 ± 5.6</p> <p>p (A:B) = ns p (A:B:C) &lt; 0.001</p>	<p><b>Interventions</b></p> <p>EFM versus auscultation</p>	<p><b>Details</b></p> <p>Parallel to a randomised clinical trial concerning alternative methods of intrapartum fetal surveillance (electronic fetal monitoring [EFM] and auscultation [AUS]) an investigatory interview was carried conducted to examine women's views on fetal monitoring. The first interview was conducted when women were at 36 weeks' gestation. In the first semi-structured interview</p>	<p><b>Results</b></p> <p><b>Women's preference</b> EFM (electronic fetal monitoring) n = 39.5% AUS (auscultation) n = 32.4% UD (undecided) n = 28%</p> <p><b>Sources of information</b></p> <p><u>Antenatal classes</u> Total number: n = 326 AUS-P: 40% EFM-P: 38% UD: 22%</p> <p><u>Books</u> Total number: n = 130 AUS-P: 47% EFM-P: 35% UD: 22%</p>	<p><b>Limitations</b></p> <p>Unclear if the outcome assessors were blinded to the study groups allocation</p> <p>41% of study population were not available for the second interview; the reason was not reported</p> <p>Inclusion and exclusion criteria not reported</p> <p>Significantly more women in EFM-P group had high-risk pregnancy</p> <p>No subgroup analysis performed based on parity</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Aim of the study</b></p> <p>To examine women's views on of intrapartum fetal surveillance methods</p> <p><b>Study dates</b></p> <p>January to August 1981</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p><u>Pathological obesity</u> AUS-P: n = 0 EFM-P: n = 9 UD: n = 8 p (A:B) &lt; 0.01 p (A:B:C) &lt; 0.05</p> <p><u>High-risk pregnancy</u> AUS-P: n = 46 EFM-P: n = 109 UD: n = 49 p (A:B) &lt; 0.001 p (A:B:C) &lt; 0.001</p> <p>There were no statistically significant differences observed between the three groups on pre-eclampsia, bleeding in pregnancy, twins, anaemia, pathological HPL, pathological estriol, diabetes, previous sterility</p> <p><b>Inclusion criteria</b></p> <p>Not reported</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>women were told about the study and consent was obtained. They were asked about their knowledge of fetal monitoring during labour and their source of information. They were also asked about their preference and asked to state the advantages and disadvantages of the two different methods. The interview lasted about 20 minutes. Out of 665 participants, 655 were interviewed initially (ten declined to participate) and 385 were interviewed again. Women were asked to state their preference for EFM or AUS and also state the advantages and disadvantages of the two methods. All women who had the pre-birth interview, were interviewed again on the 2nd or 3rd day after the birth. The person that performed the 2nd interview was blinded to the women's preference stated at the first interview regarding fetal</p>	<p><u>Newspaper</u> Total number: n = 100 AUS-P: 45% EFM-P: 40% UD: 15%</p> <p><u>Doctors</u> Total number: n = 90 AUS-P: 59% EFM-P: 32% UD: 9%</p> <p><u>Parents (a monthly magazine from a lay/support movement)</u> Total number: n = 59 AUS-P: 66% EFM-P: 24% UD: 11%</p> <p><u>Radio and TV</u> Total number: n = 56 AUS-P: 36% EFM-P: 46% UD: 19%</p> <p><u>All with information of EFM</u> Total number: n = 560 AUS-P: 35% EFM-P: 41% UD: 24%</p> <p><u>Not heard of EFM</u> Total number: n = 95 AUS-P: 18% EFM-P: 32% UD: 51%</p> <p><b>Distribution of preference related to place</b></p>	<p>(nuliparous and multiparous women)</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>monitoring. The women were asked how their labour was monitored, what the advantages or disadvantages were of the method used and how they would want the fetal heart monitored in future labours/births.</p> <p><u>Analysis</u> Analysis of variance was used for the statistical evaluation of age and parity. Elsewhere X<sup>2</sup> statistics were used</p>	<p><b>of antenatal classes</b> <u>The department</u> Total number: n = 321 AUS-P: 31% EFM-P: 42% UD: 27%</p> <p><u>Women's liberation</u> Total number: n = 64 AUS-P: 70% EFM-P: 20% UD: 9%</p> <p><u>Public schools</u> Total number: n = 35 AUS-P: 35% EFM-P: 37% UD: 27%</p> <p><u>Private institution</u> Total number: n = 31 AUS-P: 26% EFM-P: 48% UD: 26%</p> <p><u>No birth preparing courses</u> Total number: n = 213 AUS-P: 21% EFM-P: 42% UD: 36%</p> <p><b>Advantages and disadvantages of AUS mentioned postpartum by AUS-P (n = 85) and EFM-P (n = 94) groups who had their labour monitored by auscultation</b> <u>No pain to the baby</u> AUS-P: 11% EFM-P: 3%</p>	



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>p &lt;0.05</p> <p><u>No discomfort from sensors and belt</u>                      AUS-P: 58%                      EFM-P: 30%                      p &lt;0.05</p> <p><u>Increased contact with personnel</u>                      AUS-P: 25%                      EFM-P: 15%                      p &lt;0.05</p> <p><u>More natural childbirth</u>                      AUS-P: 72%                      EFM-P: 45%                      p &lt;0.05</p> <p><b>Advantages and disadvantages of EFM mentioned postpartum by AUS-P (n = 36) and EFM-P (n = 66) groups who had their labour monitored by EFM</b></p> <p><u>EFM promoting husband involvement</u>                      AUS-P: 25%                      EFM-P: 45%                      p &lt; 0.05</p> <p><u>More positively influenced by EFM signal/trace</u>                      AUS-P: 31%                      EFM-P: 67%                      p &lt; 0.01</p> <p><u>Possibility of quick intervention</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
				<p>AUS-P: 44% EFM-P: 62% p &lt; 0.05</p> <p><u>Continuous precise surveillance</u> AUS-P: 45% EFM-P: 70% p &lt; 0.05</p> <p><u>Enforced mobility</u> AUS-P: 22% EFM-P: 20% p &lt; 0.05</p> <p><u>Technical milieu</u> AUS-P: 25% EFM-P: 3% p &lt; 0.05</p> <p><u>Disturbance from EFM signals</u> AUS-P: 20% EFM-P: 3% p &lt; 0.05</p> <p><u>Fear of the trauma to the baby</u> AUS-P: 5% EFM-P: 2% p &lt; 0.05</p>	
<p><b>Full citation</b></p> <p>Mangesi,L., Hofmeyr,G.J., Woods,D.L., - Assessing the preference of women for different methods of monitoring the fetal</p>	<p><b>Sample size</b></p> <p>Total n = 100 women</p> <p><b>Characteristics</b></p>	<p><b>Interventions</b></p> <p>Fetal stethoscope, cardiotocography (CTG), Doppler ultrasound monitor</p>	<p><b>Details</b></p> <p>Convenience sampling was used; women who were in the active phase of the first stage</p>	<p><b>Results</b></p> <p><u>First maternal preference:</u> Fetal stethoscope: 13/97 FHRM: 72/97 CTG: 12/97</p>	<p><b>Limitations</b></p> <p>No details of the women's characteristics reported Women provided with the study's information</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>heart in labour, - South African Journal of Obstetrics and Gynaecology, 15, 2009-</p> <p><b>Ref Id</b> 187897</p> <p><b>Country/ies where the study was carried out</b> South Africa</p> <p><b>Study type</b> Prospective cross-sectional study</p> <p><b>Aim of the study</b> To assess which method of fetal monitoring was preferred by labouring women</p> <p><b>Study dates</b> Not reported</p> <p><b>Source of funding</b> Not reported</p>	<p>Not reported</p> <p><b>Inclusion criteria</b> Women in first stage of active labour</p> <p><b>Exclusion criteria</b> Women in second stage of labour Twin pregnancy Preterm labour Evidence of fetal distress</p>	<p>(fetal heart rate monitor [FHRM])</p>	<p>of labour were recruited from a hospital (in the Eastern Cape province, South Africa) after the study was explained and verbal consent obtained (no further details were reported). A researcher spent approximately 30 minutes with each woman; 10 minutes were spent explaining the study and obtaining consent, 10 minutes were spent monitoring the fetal heart with the stereoscope and a Doppler device (FHRM), and for the last 10 minutes the fetal heart was monitored with a cardiotocograph and if the tracing was unsatisfactory a doctor was notified. Participants were asked to indicate their first and second preferred method.</p> <p><u>Data analysis</u> Data were recorded in a collecting sheet and then entered into Epi_Info 2002</p>	<p><u>Second maternal preference:</u> Fetal stethoscope: 58/97 FHRM: 17/97 CTG: 22/97 n = 2 women were unable to decide n = 1 loss of data</p> <p>The fetal stereoscope was disliked because of causing discomfort during the examination and CTG was disliked because it often confined women to the bed and the securing belt of the cardiotocograph restricted the woman's movement</p>	<p>when they were in labour Consent obtained verbally Intervention applied over very short period of time Not clear when participants were asked about their preference Poor reporting with limited information provided</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			computer software (no further detail reported)		
<p><b>Full citation</b></p> <p>McCourt, C., Technologies of birth and models of midwifery care, Revista Da Escola de Enfermagem Da Usp, 48 Spec No, 168-77, 2014</p> <p><b>Ref Id</b></p> <p>446553</p> <p><b>Country/ies where the study was carried out</b></p> <p>UK</p> <p><b>Study type</b></p> <p>Qualitative (the study author reported that she relied on questionnaire responses too, but the findings included for this review were obtained using qualitative methodology)</p> <p><b>Aim of the study</b></p> <p>The article focuses on the theme of birth technology and discusses the impact on women's embodiment in birth and sources of information women use about the status of their bodies, their labour and the babies. The overarching study explored how the impact of birth on</p>	<p><b>Sample size</b></p> <p>N=1403 (survey); 44 women were interviewed (20 had responded to the survey, 24 had not)</p> <p><b>Characteristics</b></p> <p>Not reported for the group of women that replied to the questionnaire. For the group that did not respond to the questionnaire, the authors targeted women in minority ethnic groups and young mothers</p> <p><b>Inclusion criteria</b></p> <p>For the interviews the author wrote to all women returning questionnaires in a particular time period including all those who were contactable until 20 interviews had been arranged.* The second group were women who had not returned the questionnaires but had not declined consent to take part. Because the author was concerned about possible skews in response</p>	<p><b>Interventions</b></p> <p>Continuous electronic fetal monitoring</p>	<p><b>Details</b></p> <p>The article draws on the evaluation of a pilot scheme for caseload midwifery, which was implemented in response to UK government policy recommendations on woman-centred care in 1993. The evaluation was performed using both a survey and semi-structured interviews.* The survey of women's responses to care was based on a detailed structured postal questionnaire about how women experience their care and whether the pattern of care affects their wellbeing. The study authors* also interviewed two groups of women, chosen as subsamples from the survey, using semi-structured interviews. The first group were women</p>	<p><b>Results</b></p> <p>The following quotations were cited from two interviews. "I could tell he was OK by the monitor I think" (Standard care, 418). "I kept asking questions though... but otherwise it was just through my husband... he was in the delivery suite and in the operating theatre... he had had quite a good idea, he had been able to look at the graphs, baby's heartbeat and my contractions, and even though maybe not knowing exactly what to read into the graphs" (Standard care, 424). The comments above were chosen by the author of the article as examples of her impression that the baby and the labour were perceived to some extent as being in the woman's body. The author specified that she built her impression from listening to the women's narratives and from observation of medical staff, although the</p>	<p><b>Limitations</b></p> <p>Aims of the research: Low risk of bias (clearly explained, with comprehensive background and rationale) Qualitative methodology: Low risk of bias (qualitative research is an appropriate methodology for the research goal) Research design: Low risk of bias (in relation to the group of women who had already responded to a questionnaire, the study author reported that interviews were carried out not only to check the validity of closed questionnaire responses but also to give a greater depth of response than could be obtained through a structured questionnaire) Recruitment strategy: Unclear risk of bias (in relation to the group of women that were interviewed, the study authors reported that all women returning the first postal questionnaire during a particular time period were</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>women's experiences may be mediated by a relational model of support achieved through a caseload model of midwifery care</p> <p><b>Study dates</b></p> <p>The study was conducted over a 2-year period from 1994 to 1996* *This information was reported in the companion paper: McCourt, C., Page, L., Hewison J., Vail, A., Evaluation of One-to-One Midwifery: Women's Responses to Care, Birth, 25:2, 73-80, 1998</p> <p><b>Source of funding</b></p> <p>Not reported</p>	<p>patterns, she targeted women who were less likely to respond to a written questionnaire – women in minority ethnic groups and young mothers (under 21 years*). All such women, who had not declined consent but had not returned a questionnaire, were contacted by letter, and all those who responded by letter or could be contacted by telephone were included.* *This information was reported in the paper: McCourt, C., Page, L., Hewison J., Vail, A., Evaluation of One-to-One Midwifery: Women's Responses to Care, Birth, 25:2, 73-80, 1998</p> <p><b>Exclusion criteria</b></p> <p>Not reported</p>		<p>who had responded to the survey by completing questionnaires. The other interviews were conducted for the group who had not returned the questionnaires but had not declined consent to take part, including one interview involving assistance of an interpreter. The interviews used a narrative approach; women were asked to tell their stories from first contact with maternity services. They were asked to reflect what they found most helpful or would like to change about each stage of care. The article used analysis of women's narrative accounts of labour and birth. Transcripts of interview tapes were analysed with computer-assisted text analysis.* The article is based mainly on analysis of the interviews but is also informed by the analysis of women's questionnaire</p>	<p>impressions were rarely articulated by the women. The authors wrote that many women and partners, and medical staff, focused attention on the monitor screen to try to understand their labour. This tendency was increased for women who had an epidural (these women could not feel their contractions and watched the monitor to see when contractions were taking place) and for women in standard care (these women were less satisfied with the information and support they received than those who experienced the caseload model of midwifery care) In addition to the main outcomes, the study authors reported that responses to CTG monitoring were ambiguous. In questionnaire responses women were least likely to be critical of receiving CTG monitoring since they perceived this to be important for the safety of the baby; however, no quotations from the women who participated in the study were reported in support of this</p>	<p>contacted and asked to participate until 20 interviews had been arranged,* however the time period was not specified and the authors did not specify how they chose this time period) Data collection: Low risk of bias (semi-structured interviews*) Relationship between researcher and participants: Unclear risk of bias (it was not reported whether the relationship between the researcher and the participants had been considered) Ethical issues: Low risk of bias (the original study was approved by the ethics committee of the hospitals concerned) Data analysis: Low risk of bias (the study authors reported that transcripts of all interviews were analysed with computer-assisted text analysis and that key emergent themes were developed through open coding; responses were then sorted to log the number of women providing comments in each category and the nature of the responses*) Statement of findings: Low risk of bias (the findings are explicit and there is adequate discussion of the evidence)</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
			<p>responses, which provided less depth but covered a broader scope of women. The article focuses mainly on women's experiences of birth and differences in the ways in which women recounted these experiences according to whether they were attended by a caseload midwife and whether they received a high or low level of technological intervention. The overall findings had been published previously by a larger group of authors but this article focused on a different aspect: birth technology.</p> <p>*This information is reported in the companion paper: McCourt, C., Page, L., Hewison J., Vail, A., Evaluation of One-to-One Midwifery: Women's Responses to Care, Birth, 25:2, 73-80, 1998</p>		<p>Research value: Unclear risk of bias (the study author did not discuss whether or how the findings could be transferred to other populations, and they did not identify new areas where research would be necessary)</p> <p>Overall quality rating based on the aforementioned considerations: Moderate</p> <p>*This information was reported in the companion paper: McCourt, C., Page, L., Hewison J., Vail, A., Evaluation of One-to-One Midwifery: Women's Responses to Care, Birth, 25:2, 73-80, 1998</p> <p>Note: limitations were assessed using the Critical Appraisal Skills Programme (CASP) Qualitative Research Checklist as recommended in the 2012 NICE guidelines manual</p> <p><b>Other information</b></p> <p>The article includes only limited information relating to the methods employed in the</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
					<p>study, however the author of the article reported that she used interviews from which overall findings had been published previously. Therefore it was possible to obtain more information from the following companion paper as referred to above:                      McCourt, C., Page, L., Hewison J., Vail, A., Evaluation of One-to-One Midwifery: Women's Responses to Care, Birth, 25:2, 73-80, 1998</p>

**G.11 Cardiocography with electrocardiogram analysis compared with cardiocography alone**

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Full citation</b></p> <p>Belfort, M. A., Saade, G. R., Thom, E., Blackwell, S. C., Reddy, U. M., Thorp, J. M., Tita, A. T. N., Miller, R. S., Peaceman, A. M., McKenna, D. S., Chien, E. K. S., Rouse, D. J., Gibbs, R. S., El-Sayed, Y. Y., Sorokin, Y., Caritis, S. N., VanDorsten, J. P., A randomized trial of intrapartum fetal ECG ST-segment analysis, New England Journal of Medicine, 373, 632-641, 2015</p> <p><b>Ref Id</b></p> <p>446127</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p>	<p><b>Sample size</b></p> <p>See Neilson 2015</p> <p><b>Characteristics</b></p> <p>11,108 randomised women with a single fetus &gt;36 weeks of gestation who were attempting vaginal birth and had cervical dilation between 2 and 7 cm; trial conducted at 16 university-based clinical centres in Eunice Kennedy Shriver National Institute of Child Health &amp; Human Development (NICHD) Maternal-Fetal Medicine Units (MFMU) Network</p> <p>Intervention: CTG plus fetal ECG (ST-segment analysis) (n=5532) versus CTG only (n=5576). Monitoring device used was STAN S31 (Neoventa Medical)</p>	<p><b>Interventions</b></p> <p>Intervention: CTG plus fetal ECG-ST analysis, n=5532 Control: CTG only, n=5576</p>	<p><b>Details</b></p> <p>See Neilson 2015</p>	<p><b>Results</b></p> <p>See Neilson 2015 for other outcomes</p> <p>1. <u>Spontaneous vaginal birth</u> CTG plus fetal ECG-ST analysis: 4269/5532 CTG only: 4348/5576</p> <p>2. <u>Apgar score ≤ 3 at 5 minutes</u> CTG plus fetal ECG-ST analysis: 17/5532 CTG only: 6/5576</p>	<p><b>Limitations</b></p> <p>Risk of bias: no details of randomisation procedure reported Participant blinding: not possible Outcome assessment blinding: protocol subcommittee that was unaware of study group assignment conducted chart review of all cases that met primary outcome criteria Attrition bias: full clinical data and valid umbilical blood gas results obtained from 96.5% of neonates</p> <p><b>Other information</b></p> <p>See Neilson 2015</p>



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Multicentre randomised controlled trial (RCT)</p> <p><b>Aim of the study</b></p> <p>To assess whether intrapartum fetal ECG ST-segment analysis in addition to conventional CTG modifies intrapartum and neonatal outcomes</p> <p><b>Study dates</b></p> <p>Recruitment from November 2010 to March 2014</p> <p><b>Source of funding</b></p> <p>Grants from NICHD and funding from Neoventa Medical</p>	<p><b>Inclusion criteria</b></p> <p>Women with a singleton fetus &gt;36 weeks of gestation who were attempting vaginal birth and had cervical dilation of between 2 and 7 cm</p> <p><b>Exclusion criteria</b></p> <p>Noncephalic presentation, planned caesarean birth, need for immediate birth, absent fetal heart-rate variability (amplitude range undetectable) or a sinusoidal pattern, minimal fetal heart-rate variability in the 20 minutes before randomization, or other fetal or maternal conditions that would preclude trial of labour or placement of scalp electrode</p>				
<p><b>Full citation</b></p> <p>Neilson, J. P., Fetal electrocardiogram (ECG) for fetal monitoring during labour, Cochrane</p>	<p><b>Sample size</b></p> <p>Total n = 27403 Electrocardiogram (ECG) plus cardiotocograph (CTG) n = 13711</p>	<p><b>Interventions</b></p> <p>Intervention n: CTG plus ECG</p>	<p><b>Details</b></p> <p><u>Electronic searches</u> The Cochrane Pregnancy and Childbirth Group's Trials Register was</p>	<p><b>Results</b></p> <p><u>1 Caesarean section</u> No. of studies: 7 total n = 27403</p>	<p><b>Limitations</b></p> <p><b>Quality of review</b></p> <ol style="list-style-type: none"> <li>1. Was an 'a priori' design provided? Yes</li> <li>2. Was there duplicate study selection and data extraction? Yes</li> </ol>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>Database of Systematic Reviews, 12, CD000116, 2015</p> <p><b>Ref Id</b></p> <p>446197</p> <p><b>Country/ies where the study was carried out</b></p> <p><b>Study type</b></p> <p>Cochrane systematic review</p> <p><b>Aim of the study</b></p> <p>To compare the effects of analysis of fetal ECG waveform during labour with alternative methods of fetal monitoring</p> <p><b>Study dates</b></p> <p>Updated to 23 September 2015</p> <p><b>Source of funding</b></p> <p>Supported by NIHR via Cochrane Infrastructure</p>	<p>CTG alone n = 13692</p> <p><b>Characteristics</b></p> <p><u>Amer-Wahlin 2001</u> 4966 women in labour at &gt; 36 weeks with singleton pregnancies, cephalic presentation and perceived need for continuous fetal heart rate monitoring via a fetal scalp electrode; high-risk pregnancies, suspicious or abnormal cardiotocography, induced labour, oxytocin augmentation, meconium-stained amniotic fluid or epidural analgesia. The trial took place between 1998 and 2000 in 3 Swedish centres, Lund, Malmo, Gothenburg.</p> <p>Intervention: CTG plus ST analysis of fetal ECG (2519 women) versus CTG alone (2477). The monitoring device was the STAN S21 (Neoventa Medical, Gothenburg) which incorporates an 'expert system' to provide</p>	<p>(ST or PR analysis)</p> <p>Control: CTG only</p>	<p>searched by the Trials Search Coordinator (September 23, 2015). CENTRAL, MEDLINE, EMBASE were searched, and hand searching of journals and conference proceedings was conducted. No language restrictions were applied. Weekly current awareness alert for a further of 44 journals, plus monthly BidMed Central email alerts, were also considered.</p> <p><u>Selection of studies</u> The review author (JPN) assessed all potential identified studies for inclusion.</p> <p><u>Data extraction and management</u> A form was designed to extract data and JPN extracted the data using the agreed form. The data were analysed in RevMan. Where information was unclear, JPN contacted the</p>	<p><u>1.1 ST analysis:</u> No. of studies: 6 n = 26446 ECG plus CTG n = 1810/13229 CTG alone n = 1779/13217 RR 1.02 (95% CI 0.96 to 1.08)</p> <p><u>1.2 PR analysis:</u> No. of studies: 1 n = 957 ECG plus CTG n = 79/482 CTG alone n = 98/475 RR 0.79 (95% CI 0.61 to 1.04)</p> <p><u>2 Cord pH &lt; 7.05 + base deficit &gt;12 mmol/l</u> No. of studies: 6 n = 25682</p> <p><u>2.1 ST analysis:</u> No. of studies: 6 n=25682 ECG plus CTG n = 81/12850 CTG alone n =</p>	<ol style="list-style-type: none"> <li>3. Was a comprehensive literature search performed? Yes</li> <li>4. Was the status of publication (i.e. grey literature) used as an inclusion criteria? No</li> <li>5. Was a list of studies (included and excluded) provided? Yes</li> <li>6. Were the characteristics of the included studies provided? Yes</li> <li>7. Was the scientific quality of the included studies assessed and documented? Yes</li> <li>8. Was the scientific quality of the included studies used appropriately in formulating conclusions? Yes</li> <li>9. Were the methods used to combine the findings of studies appropriate? Yes</li> <li>10. Was the likelihood of publication bias assessed? No</li> <li>11. Was the conflict of interest included? Yes</li> </ol> <p><b>Details of individual studies</b></p> <p><b>Amer-Wahlin 2001</b> A modified intention to treat analysis performed excluding non cephalic and preterm babies from the analysis.</p> <p><b>Belfort 2015</b> Unclear random sequence generation. Blinding of participants and study personnel not possible. Protocol subcommittee unaware of group assignment conducted chart review of all cases that met primary outcome criteria.</p> <p><b>Ojala 2006</b> n = 5 in CTG group and n = 78 in the ECG group had technical difficulties in achieving satisfactory monitoring.</p> <p><b>Strachan 2000</b> For unclear reason the results are reported for 92.2% of study's population. Subgroup analysis of babies born with a low arterial pH showed no action for fetal distress had been taken in nearly 75% of cases, suggesting study protocol violation within the trial groups.</p> <p><b>Westerhuis 2010</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>funding to Cochrane Pregnancy and Childbirth</p>	<p>advice to clinical staff. In this, it constitutes a technically more advanced system than used in the Westgate 1993 trial.</p> <p><u>Belfort 2015</u></p> <p>11,108 randomised women with a single fetus &gt;36 weeks of gestation who were attempting vaginal birth and had cervical dilation between 2 and 7 cm. Trial conducted at 16 university-based clinical centres in Eunice Kennedy Shriver National Institute of Child Health &amp; Human Development (NICHD) Maternal-Fetal Medicine Units (MFMU) Network. Intervention: CTG plus fetal ECG (ST-segment analysis) (n=5532) versus CTG alone (n=5576). Monitoring device was STAN S31 (Neovanta Medical).</p>		<p>original authors for further details.</p> <p><u>Assessment of risk of bias</u></p> <p>JPN assessed risk of bias using criteria from the Cochrane Handbook for Systematic Reviews of Interventions: - Sequence generation - Allocation concealment - Blinding - Incomplete outcome data - Selective reporting bias - Other sources of bias</p> <p><u>Measures of effect</u></p> <p>Dichotomous outcomes were presented as risk ratios (RR) with 95% confidence intervals (CIs). No continuous data analysed.</p> <p><u>Dealing with missing data</u></p> <p>Levels of attrition noted for included studies. Impact of including studies with high levels of missing data will be explored in future updates. Outcomes were assessed on an intention-to-treat</p>	<p>121/12832 RR 0.72 (95% CI 0.43 to 1.2)</p> <p><u>2.2 PR analysis:</u> No. of studies: 0</p> <p><u>3 Neonatal encephalopathy</u> No. of studies: 6 n = 26410</p> <p><u>3.1 ST analysis:</u> n = 26410 ECG plus CTG n = 12/13210 CTG alone n = 20/13200 RR 0.61 (95% CI 0.3 to 1.22)</p> <p><u>3.2 PR analysis:</u> No. of studies: 0</p> <p><u>4 Fetal blood sampling</u></p>	<p>There was no blinding for women or clinicians, and a secondary analysis on 61 babies with adverse outcomes (metabolic acidosis in umbilical cord artery, pH &lt; 7.00, sign of severe hypoxic ischaemic encephalopathy [HIE] and perinatal death) showed the trial protocol was violated in 11 (42%) and 13 (19%) cases of study and control group respectively.</p> <p><b>Other information</b></p> <p>The systematic review is available online at: <a href="http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000116.pub5/full">http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD000116.pub5/full</a></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><u>Ojala 2006</u> 1483 women randomised; 11 exclusions; clinical data available but blood gas data missing for 36. In labour at <math>\geq 36</math> weeks with singleton fetus, cephalic presentation, decision to perform amniotomy, no contraindication to scalp electrode. Sample size based on 50% reduction of umbilical artery pH <math>&lt; 7.10</math> Intervention: CTG plus ECG waveform analysis (STAN) (733 women) versus CTG (739 women). Fetal scalp sampling for pH estimation an option in either group. Recruitment in tertiary referral hospital in Finland 2003-4</p> <p><u>Strachan 2000</u> 957 women in labour with perceived need for continuous fetal heart rate monitoring (age <math>&gt; 35</math> years, maternal disease, adverse</p>		<p>basis as far as possible. The denominator for each outcome in each trial was the number randomised minus any participants whose outcomes were known to be missing. <u>Analysis</u> Heterogeneity was regarded high if <math>I^2 &gt; 30\%</math> and either <math>Tau^2 &gt; 0</math> or there was a low P value (<math>&lt; 0.10</math>) in the <math>Chi^2</math> test. A fixed-effect model was used for combining data where studies were assumed estimating the same underlying treatment effect. If substantial clinical or statistical heterogeneity was detected, a random effects meta analysis was used. Fixed-effect meta-analysis was used where trials were comparing the same intervention and the populations and methods were judged to be similar enough. Random</p>	<p><u>No. of studies: 5 n = 10628</u></p> <p><u>4.1 ST analysis:</u> No. of studies: 4 n = 9671 ECG plus CTG n = 449/4870 CTG alone n = 503/4801 RR 0.61 (95% CI 0.41 to 0.9)</p> <p><u>4.2 PR analysis:</u> No. of studies: 1 n = 957 ECG plus CTG n = 81/482 CTG alone n = 88/475 RR 0.91 (95% CI 0.69 to 1.19)</p> <p><u>5 Instrumental vaginal birth</u></p> <p><u>5.1 ST analysis</u> No. of studies = 6 n = 26,446 ECG plus CTG n = 1810/4870 CTG alone n =</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>obstetric history, prematurity, suspected fetal growth restriction, antepartum haemorrhage, breech presentation, multiple pregnancy, epidural analgesia, induction or augmentation of labour, abnormal cardiotocography, meconium, previous caesarean section). Results were only available for 957 women (92%) for reasons that are unclear. The trial took place in 5 centres: Nottingham and Dundee (UK), Hong Kong, Amsterdam (Netherlands) and Singapore</p> <p>Intervention: CTG plus fetal ECG (n = 482) versus CTG alone (n = 475).</p> <p><u>Vayssiere 2007</u> 799 women in labour at 36 weeks or more, with a single fetus with cephalic presentation, and either abnormal cardiotocographic trace</p>		<p>effects meta-analyses were used where heterogeneity was present or suspected. If substantial heterogeneity was detected, it was investigated using subgroup and sensitivity analysis</p>	<p>1489/13217 RR 1.02 (95% CI 0.96 to 1.08)</p> <p><u>5.2 PR analysis</u> No. of studies = 1 n = 957 ECG plus CTG n = 116/482 CTG alone n = 122/475 RR 0.94 (95% CI 0.75 to 1.17)</p> <p><u>6 Neonatal intubation</u>  No. of studies: 3 n=13501</p> <p><u>6.1 ST analysis</u>  No. of studies = 2 n = 12544  ECG plus CTG n = 51/6246 CTG alone n = 36/6298 RR 1.37 (95% CI 0.89 to 2.11)</p> <p><u>6.2. PR analysis</u></p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>or thick meconium-stained amniotic fluid. Exclusions included maternal infections that contraindicated scalp electrode attachment (e.g. HIV), cardiac malformation, severely abnormal cardiotocography at the time of recruitment was an option in both groups</p> <p>Intervention: CTG + fetal ECG (n = 399) versus CTG alone (n = 400). Scalp sampling for pH estimation</p> <p><u>Westerhuis 2010</u></p> <p>5681 women in labour with a singleton fetus in vertex position, a gestational age 36 weeks or greater and a medical indication for electronic fetal monitoring defined by either a high-risk pregnancy (induction or augmentation of labour, epidural anaesthesia, meconium-stained amniotic fluid) or non-reassuring fetal heart</p>			<p>No. of studies = 1 n = 957</p> <p>ECG plus CTG n = 6/482 CTG alone n = 8/475 RR 0.74 (95% CI 0.26 to 2.11)</p> <p><u>7 Admission to neonatal care unit</u> No. of studies: 7 n = 27367</p> <p><u>7.1 ST analysis:</u></p> <p>No. of studies: 6 n=26410 ECG plus CTG n = 1113/13210 CTG alone n = 1155/13200 RR 0.96 (95% CI 0.89 to 1.04)</p> <p><u>7.2 PR analysis</u> No. of studies: 1 n = 957 ECG plus CTG n = 22/482 CTG alone n = 28/475</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>rate Intervention group:</p> <p>CTG and ST-analysis. Control group: CTG.</p> <p><u>Westgate 1993</u> 2434 pregnant women, 1215 in cardiotocography alone arm, 1219 ST waveform and CTG arm. (More than 34 weeks of gestation with no gross fetal abnormality.) Intervention: CTG plus ST analysis (n =1219) versus CTG alone (n = 1215).</p> <p><b>Inclusion criteria</b></p> <p>Trials comparing analysis of any component of the fetal electrocardiographic (ECG) during labour with alternative fetal monitoring methods. Studies using less robust methods of allocation (for example,</p>			<p>RR 0.77 (95% CI 0.45 to 1.33)</p> <p><u>8 Fetal, perinatal or neonatal death</u> No. of studies: 7 n = 26446</p> <p><u>8.1 ST analysis</u></p> <p><u>Fetal or neonatal death</u> No. of studies: 6 n = 15338 ECG plus CTG n = 11/13229 CTG alone n = 6/13217 RR 1.71 (95% CI 0.67 to 4.33)</p> <p><u>8.2 PR analysis</u></p> <p><u>Perinatal death</u> No. of studies: 1 n = 957 ECG plus CTG n = 1/482 CTG alone n = 0/475 RR 2.96 (95% CI 0.12 to 72.39)</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	alternation) were not included  <b>Exclusion criteria</b>			<u>9 Apgar score &lt;7 at 5 minutes</u> No. of studies: 6 n = 16259  <u>9.1 ST analysis</u>  No. of studies: 5 n = 15302 ECG plus CTG n = 103/7678 CTG alone n = 1078/7624 RR 0.95 (95% CI 0.73 to 1.24)  <u>9.2 PR analysis</u>  No. of studies: 1 n = 957 ECG plus CTG n = 3/482 CTG alone n = 7/475 RR 0.42 (95% CI 0.11 to 1.62)	
<b>Full citation</b>  Olofsson, P., Ayres-de-Campos, D., Kessler, J., Tendal, B., Yli, B. M., Devoe, L., A critical appraisal of the	<b>Sample size</b>  No. of studies: 5, n=15363 CTG plus fetal ECG-ST (n=7702) versus CTG only (n=7661)	<b>Interventions</b>  Intervention: CTG plus fetal ECG-ST analysis	<b>Details</b>  No details reported of how studies were selected. Includes revised data from Amer-Wahlin 2011	<b>Results</b>  <u>1. Spontaneous vaginal birth</u> No. of studies: 5, n=15363	<b>Limitations</b>  <u>Quality of review</u>  1. Was an 'a priori' design provided? No 2. Was there duplicate study selection and data extraction? No



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p>evidence for using cardiotocography plus ECG ST interval analysis for fetal surveillance in labor. Part I: the randomized controlled trials, Acta Obstetrica et Gynecologica Scandinavica, 93, 556-68; discussion 568-9, 2014</p> <p><b>Ref Id</b> 446200</p> <p><b>Country/ies where the study was carried out</b></p> <p><b>Study type</b> Critical review of CTG plus fetal ECG-ST analysis randomised controlled trials (RCTs)</p> <p><b>Aim of the study</b> To assess the quality of 5 RCTs evaluating CTG plus fetal ECG ST analysis</p>	<p><b>Characteristics</b></p> <p><u>Westgate 1993</u> 2434 pregnant women, 1215 cardiotocography alone arm, 1219 ST waveform and CTG arm. (More than 34 weeks of gestation with no gross fetal abnormality.) Intervention: CTG plus ST analysis (n =1219) versus CTG alone (n = 1215). CTG plus fetal ECG-ST (n=1219) versus CTG only (n=1215)</p> <p><u>Amer-Wahlin 2001/2011</u> 4966 women in labour at &gt; 36 weeks with singleton pregnancies, cephalic presentation and perceived need for continuous fetal heart rate monitoring via a fetal scalp electrode; high-risk pregnancies (suspicious or abnormal cardiotocography, induced labour, oxytocin augmentation, meconium-stained amniotic fluid or epidural analgesia). The</p>	Control: CTG only	and Westerhuis 2011. Review addressed: (1) Power calculations, (2) Prestudy training, inclusion criteria, randomisation and recruitment pace, (3) Intrapartum management protocols, (4) Intrapartum interventions, (5) Cord blood and early neonatal metabolic acidosis, (6) Neonatal outcomes	<p>CTG plus fetal ECG-ST (n=7702) versus CTG only (n=7661) <u>Westgate 1993</u> CTG plus fetal ECG-ST: 875/1219 CTG only: 832/1215 RR 1.05 (95%CI 0.995, 1.1)</p> <p><u>Amer-Wahlin 2001/2011</u> CTG plus fetal ECG-ST: 2065/2519 CTG only: 1947/2447 RR 1.03 (95%CI 1.003, 1.059)</p> <p><u>Ojala 2006</u> CTG plus fetal ECG-ST: 616/733 CTG only: 625/739 RR 0.99 (95%CI 0.95, 1.04)</p> <p><u>Vayssiere 2007</u> CTG plus fetal ECG-ST: 183/399 CTG only: 179/400 RR 1.02 (95%CI 0.88, 1.19)</p> <p><u>Westerhuis 2010/2011</u> CTG plus fetal ECG-ST: 2038/2827 CTG only: 2018/2840 RR 1.01 (95%CI 0.98, 1.05)</p>	<ol style="list-style-type: none"> <li>3. Was a comprehensive literature search performed? No</li> <li>4. Was the status of publication (i.e. grey literature) used as an inclusion criteria? No</li> <li>5. Was a list of studies (included and excluded) provided? Not applicable, not a systematic review</li> <li>6. Were the characteristics of the included studies provided? Yes</li> <li>7. Was the scientific quality of the included studies assessed and documented? Yes</li> <li>8. Was the scientific quality of the included studies used appropriately in formulating conclusions? Yes</li> <li>9. Were the methods used to combine the findings of studies appropriate? Not applicable, meta-analysis not conducted</li> <li>10. Was the likelihood of publication bias assessed? No</li> <li>11. Was the conflict of interest included? Yes</li> </ol> <p><b>Details of individual studies:</b></p> <p><b>Amer-Wahlin 2001</b> A modified intention to treat analysis performed excluding non-cephalic and preterm babies from the analysis.</p> <p><b>Ojala 2006</b> n = 5 in CTG group and n = 78 in the ECG group had technical difficulties in achieving satisfactory monitoring.</p> <p><b>Strachan 2000</b></p> <p>For unclear reason the results are reported for 92.2% of the study's population. Subgroup analysis of babies born with a low arterial pH showed no action for fetal distress had been taken in nearly 75% of cases, suggesting study protocol violation within the trial groups.</p> <p><b>Westerhuis 2010</b> There was no blinding for women or clinicians, and a secondary analysis on 61 babies with adverse outcomes (metabolic acidosis in umbilical cord artery, pH &lt; 7.00, sign of severe hypoxic ischaemic encephalopathy [HIE] and</p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Study dates</b> From 1993 to 2011</p> <p><b>Source of funding</b> None reported</p>	<p>trial took place between 1998 and 2000 in 3 Swedish centres, Lund, Malmo, Gothenburg Intervention: CTG plus ST analysis of fetal ECG (2519 women) versus CTG alone (2477). The monitoring device was the STAN S21 (Neoventa Medical, Gothenburg) which incorporates an 'expert system' to provide advice to clinical staff. In this, it constitutes a technically more advanced system than used in the Westgate 1993 trial. CTG plus fetal ECG-ST (n=2519) versus CTG only (n=2447)</p> <p><u>Ojala 2006</u> 1483 women randomised; 11 exclusions; clinical data available but blood gas data missing for 36. In labour at <math>\geq 36</math> weeks with singleton fetus, cephalic presentation, decision to perform amniotomy, no contraindication to scalp electrode. Sample size based on</p>			<p><b><u>Overall (not reported in review article; calculated by NGA technical team in RevMan)</u></b> CTG plus fetal ECG-ST: n=7702 CTG only: n=7661 RR 1.02 (95%CI 1.0, 1.04)</p>	<p>perinatal death) showed the trial protocol was violated in 11 (42%) and 13 (19%) cases of study and control group respectively.</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>50% reduction of umbilical artery pH &lt; 7.10                      Intervention: CTG plus ECG waveform analysis (STAN) (733 women) versus CTG (739 women). Fetal scalp sampling for pH estimation an option in either group.                      Recruitment in tertiary referral hospital in Finland 2003-4                      CTG + fetal ECG-ST (n=733) versus CTG only (n=739)</p> <p><u>Vayssiere 2007</u>                      799 women in labor at 36 weeks or more, with a single fetus with cephalic presentation, and either abnormal cardiotocographic trace or thick meconium-stained amniotic fluid. Exclusions included maternal infections that contraindicated scalp electrode attachment (e.g. HIV), cardiac malformation, severely abnormal cardiotocography at the time of recruitment was an option in both groups</p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>Intervention: CTG + fetal ECG (n = 399) versus CTG alone (n = 400). Scalp sampling for pH estimation CTG + fetal ECG-ST (n=399) versus CTG only (n=400)</p> <p><u>Westerhuis 2010/2011</u> 5681 women in labour with a singleton fetus in vertex position, a gestational age 36 weeks or greater and a medical indication for electronic fetal monitoring. A medical indication is defined by either a high-risk pregnancy, induction or augmentation of labour, epidural anaesthesia, meconium-stained amniotic fluid or non-reassuring fetal heart rate Intervention group: CTG and ST-analysis. Control group: CTG. CTG + fetal ECG-ST (n=2832) versus CTG only (n=2849)</p> <p><b>Inclusion criteria</b></p>				

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>RCT of CTG plus fetal ECG-ST analysis studies</p> <p><b>Exclusion criteria</b></p> <p>None reported</p>				
<p><b>Full citation</b></p> <p>van Wijngaarden,W.J., Sahota,D.S., James,D.K., Farrell,T., Mires,G.J., Wilcox,M., Chang,A., Improved intrapartum surveillance with PR interval analysis of the fetal electrocardiogram: a randomized trial showing a reduction in fetal blood sampling, American Journal of Obstetrics and Gynecology, 174, 1295-1299, 1996</p> <p><b>Ref Id</b></p> <p>196803</p>	<p><b>Sample size</b></p> <p>N=214. CTG plus fetal ECG-PR interval analysis, n=112 (Included in analyses, n=84; &gt;37 week=76, 27-37 week=8) CTG only, n=102 (Included in analyses, n=100; &gt;37 week=92, 27-37 week=8) Excluded: Inability to obtain analysable fetal ECG waveform signal, n=8; non-availability of umbilical artery gas measurements, n=4; discontinuation of trial at woman's request, n=1; erroneous fetal ECG analyser settings by labour suite staff resulting in inverted waveform that did not provide any fetal ECG data, n=17</p>	<p><b>Interventions</b></p> <p>No. of participants, N=214 Intervention: CTG plus fetal ECG-PR interval analysis, n=112 Control: CTG only, n=102</p>	<p><b>Details</b></p> <p>Randomisation using PC-random number generator. All participants monitored by fetal ECG analyser system. Fetal ECG signal obtained by Copeland's fetal scalp electrode (Surgicraft, Redditch, UK) or a spiral scalp electrode (Corometrics Medical Systems, Wallingford, CT, USA), processed, and analysed with Nottingham fetal ECG analyser. Time-interval parameters displayed on video display in CTG plus fetal ECG-PR interval analysis</p>	<p><b>Results</b></p> <p><u>1 Number undergoing fetal blood sampling</u> CTG plus fetal ECG-PR interval analysis: 5/84 CTG only: 21/100 Intention to treat: CTG plus fetal ECG-PR interval analysis: 5/112 CTG only: 21/103</p> <p><u>2 Acidotic infants</u> CTG plus fetal ECG-PR interval analysis: 8/84 CTG only: 14/100 Intention to treat: CTG plus fetal ECG-PR interval analysis: 8/112 CTG only: 14/102</p> <p><u>3 Assisted births</u> CTG plus fetal ECG-PR interval analysis: 36/84 CTG only: 42/100 Intention to treat:</p>	<p><b>Limitations</b></p> <p>Allocation concealment: no details reported Participant blinding: not possible Outcome assessment blinding: all labour records, CTG, fetal ECG data, and biochemical data were reviewed and scrutinised according to signal quality, protocol adherence, and sample quality by a research fellow and research engineer at Queen's Medical Centre before analysis of outcomes Attrition bias: full clinical data available for 86% of sample</p> <p><b>Other information</b></p>

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
<p><b>Country/ies where the study was carried out</b></p> <p>UK, Hong Kong</p> <p><b>Study type</b></p> <p>Randomised prospective trial</p> <p><b>Aim of the study</b></p> <p>To test potential reduction in unnecessary fetal blood sampling in sample of high-risk labours using CTG plus fetal ECG-PR interval analysis versus CTG only</p> <p><b>Study dates</b></p> <p>Not reported clearly</p> <p><b>Source of funding</b></p> <p>None reported</p>	<p><b>Characteristics</b></p> <p>Compared fetal blood sampling rate and results in 214 'high-risk' parturients (where the fetus was at risk of acidosis) monitored by CTG plus fetal ECG-PR interval analysis or by CTG only in 3 teaching hospitals over period of 10 months (Queens Medical Centre, Nottingham, UK), 8 months (Ninewells Hospital, Dundee, UK) and 3 months (Prince of Wales Hospital, Hong Kong). Randomisation using PC-random number generator. All participants monitored by fetal ECG analyser system. Fetal ECG signal obtained by Copeland's fetal scalp electrode (Surgicraft, Redditch, UK) or a spiral scalp electrode (Corometrics Medical Systems, Wallingford, CT, USA), processed, and analysed with Nottingham fetal ECG analyser. Time-interval</p>		<p>group, whilst only electronic fetal monitoring information displayed in CTG only group. Labour management and decision making sole responsibility of on-call labour ward staff. Intervention with fetal blood sampling or birth in CTG only group according to established International Federation of Gynecology and Obstetrics (FIGO) guidelines in use at labour suites of each unit. Management in CTG plus ECG-PR interval analysis group based on: (1) electronic fetal monitoring; (2) conduction index: positive index &gt;20 minutes defined as 'abnormal'; (3) ratio index &gt;4% defined as 'abnormal'. If the CTG became abnormal (e.g. prolonged profound bradycardia) then an 'opt-out clause' allowing</p>	<p>CTG plus fetal ECG-PR interval analysis: 36/112 CTG only: 42/102</p> <p><u>4 Assisted births for presumed fetal distress</u> CTG plus fetal ECG-PR interval analysis: 7/84 CTG only: 16/100</p> <p>Intention to treat CTG plus fetal ECG-PR interval analysis: 7/112 CTG only: 16/102</p>	

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>parameters displayed on video display in CTG plus fetal ECG-ST analysis group, whilst only electronic fetal monitoring information displayed in CTG only group.</p> <p>Labour management and decision making sole responsibility of on-call labour ward staff.</p> <p>Intervention with fetal blood sampling or birth in CTG only group according to established International Federation of Gynecology and Obstetrics (FIGO) guidelines in use at labour suites of each unit.</p> <p>Management in CTG plus ECG-ST analysis group based on: (1) electronic fetal monitoring; (2) conduction index: positive index &gt;20 minutes defined as 'abnormal'; (3) ratio index &gt;4% defined as 'abnormal'. If CTG became abnormal (e.g. prolonged profound bradycardia), then an 'opt-out clause' allowing management</p>		<p>management based only on CTG was allowed.</p> <p>Abnormal fetal blood sampling result: pH&lt;=7.25</p> <p>Normal fetal blood sampling result: pH&gt;7.25</p> <p>Acidosis at birth: arterial umbilical cord pH&lt;=7.15 (1 SD below mean of population studied)</p>		

Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p>based only on CTG was allowed</p> <p><b>Inclusion criteria</b></p> <p>High-risk parturients. Since there was only one ECG analyser at each centre, if there was more than one eligible participant then the one thought to have greatest risk of fetal compromise was approached for recruitment.</p> <p>Definition of 'high risk':</p> <p>(1) Maternal factors: age &lt;16 or &gt;35 years; weight &lt;45 kg or &gt;90 kg; any disease with potential adverse effect on fetus. (2) Obstetric factors: poor obstetric history; intrauterine growth restriction; prematurity; antepartum haemorrhage. (3) Intrapartum factors: breech presentation; epidural anaesthesia; induction or augmentation of labour with oxytocin; trial of scar with labour; cardiotocographic abnormalities; meconium.</p>				



Study details	Participants	Interventions	Methods	Outcomes and Results	Comments
	<p><b>Exclusion criteria</b></p> <p>Women giving birth by elective caesarean section; cases in which &lt;1 hour of interpretable data expected; woman did not consent to trial; fetal ECG analyser at site not available</p>				

## G.12 Automated interpretation of cardiotocograph traces

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Full citation</b></p> <p>Chen, C. Y., Yu, C., Chang, C. C., Lin, C. W., Comparison of a novel computerized analysis program and visual interpretation of cardiotocography, PLoS ONE [Electronic Resource], 9, e112296, 2014</p> <p><b>Ref Id</b></p> <p>446257</p> <p><b>Country/ies where the study was carried out</b></p> <p>Taiwan</p> <p><b>Study type</b></p>	<p><b>Sample size</b></p> <p>N = 62 CTG traces</p> <p><b>Characteristics</b></p> <p>Mean gestational age 38 weeks (range 37-40) No other characteristics reported</p> <p><b>Inclusion Criteria</b></p> <p>Singleton pregnancies of <math>\geq 37</math> weeks' gestation. No medical complications in the woman and no known congenital abnormalities in the fetus</p> <p><b>Exclusion Criteria</b></p> <p>None reported</p>	<p><b>Tests</b></p> <p>A computerised algorithm for interpretation of the CTG was developed using LabVIEW 2010 software. This enabled detection of the baseline fetal heart rate, variability, accelerations and decelerations (number and timing). The NICHD 3 tier system for the classification of CTGs was used to define the traces as normal (category I), indeterminate (category II) or abnormal (category III)</p>	<p><b>Methods</b></p> <p>62 admission CTGs were obtained from a database including women admitted in early labour to a tertiary care university hospital. The duration of each trace was between 20 and 30 minutes. They were independently examined by 8 obstetricians with between 3 and 6 years of experience. Observers were asked to record the baseline heart rate, variability, number of accelerations, number and type of decelerations, uterine</p>	<p><b>Results</b></p> <p><b>Agreement between the computer algorithm and the eight obstetricians</b></p> <p>Baseline fetal heart rate, ICC (95% CI): 0.91 (0.88 - 0.94) Baseline variability, <math>\kappa</math> statistic (95% CI): 0.68 (0.51 - 0.84) Accelerations, ICC (95% CI): 0.85 (0.80 - 0.90) Early decelerations, ICC (95% CI): 0.78 (0.71 - 0.84) Late decelerations, ICC (95% CI): 0.67 (0.59 - 0.76) Variable decelerations, ICC (95% CI): 0.60 (0.51 - 0.70) Prolonged deceleration, <math>\kappa</math> statistic (95% CI): 0.82 (0.58 - 1.00) Recurrent deceleration, <math>\kappa</math> statistic (95% CI): 0.82 (0.67 - 0.97) Contraction frequency, ICC (95% CI): 0.97 (0.96 - 0.98)</p> <p><u>CTG categories</u></p> <p>Category I, <math>\kappa</math> statistic (95% CI): 0.91 (0.81 - 1.00) Category II, <math>\kappa</math> statistic (95% CI): 0.78 (0.63 - 0.93) Category III, <math>\kappa</math> statistic (95% CI): 0.50 (0.17 - 0.83) Overall categorisation, <math>\kappa</math> statistic (95% CI): 0.80 (0.67 - 0.94)</p> <p><b>Agreement between the eight obstetricians only</b></p> <p>Baseline fetal heart rate, ICC (95% CI): 0.91 (0.88 - 0.94) Baseline variability, <math>\kappa</math> statistic (95% CI): 0.67 (0.51 - 0.83) Accelerations, ICC (95% CI): 0.84 (0.79 - 0.89) Early decelerations, ICC (95% CI): 0.78 (0.71 - 0.84) Late decelerations, ICC (95% CI): 0.65 (0.56 - 0.74) Variable decelerations, ICC (95% CI): 0.59 (0.50 - 0.69) Prolonged deceleration, <math>\kappa</math> statistic (95% CI): 0.82 (0.58 - 1.00) Recurrent deceleration, <math>\kappa</math> statistic (95% CI): 0.82 (0.66 - 0.97) Contraction frequency, ICC (95% CI): 0.97 (0.96 - 0.98)</p> <p><u>CTG categories</u></p> <p>Category I, <math>\kappa</math> statistic (95% CI): 0.90 (0.81 - 1.00)</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS criteria</b></p> <p>1. Patient selection – high risk; selection of CTGs was not reported to be random or consecutive; cases were apparently chosen to ensure different classes of CTG were included 2. Index tests – low risk 3. Reference standard – low risk</p>

<b>Bibliographic details</b>	<b>Participants</b>	<b>Tests</b>	<b>Methods</b>	<b>Outcomes and results</b>	<b>Comments</b>
<p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To compare new computerised CTG analysis software with visual interpretation of the CTG</p> <p><b>Study dates</b></p> <p>CTGs were recorded between March and September 2011</p> <p><b>Source of funding</b></p> <p>No funding or support reported</p>			<p>contractions and category of CTG (according to the NICHD criteria)</p>	<p>Category II, <math>\kappa</math> statistic (95% CI): 0.78 (0.62 - 0.93)                      Category III, <math>\kappa</math> statistic (95% CI): 0.48 (0.15 - 0.80)                      Overall categorisation, <math>\kappa</math> statistic (95% CI): 0.80 (0.66 - 0.93)</p>	<p>4. Flow and timing – low risk</p>
<b>Full citation</b>	<b>Sample size</b>	<b>Tests</b>	<b>Methods</b>	<b>Results</b>	<b>Limitations</b>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																																																			
<p>Chung,T.K., Mohajer,M.P., Yang,Z.J., Chang,A.M., Sahota,D.S., The prediction of fetal acidosis at birth by computerised analysis of intrapartum cardiotocography, British Journal of Obstetrics and Gynaecology, 102, 454-460, 1995</p> <p><b>Ref Id</b> 197179</p> <p><b>Country/ies where the study was carried out</b> UK</p> <p><b>Study type</b> Retrospective cohort study</p>	<p>n = 73 CTG traces</p> <p><b>Characteristics</b></p> <table border="1"> <thead> <tr> <th>Characteristic</th> <th>Number</th> <th>Mean (range)</th> </tr> </thead> <tbody> <tr> <td>Maternal</td> <td></td> <td></td> </tr> <tr> <td>Maternal age (years)</td> <td></td> <td>26.6 (15-40)</td> </tr> <tr> <td>Primiparous</td> <td>50</td> <td></td> </tr> <tr> <td>Multiparous</td> <td>23</td> <td></td> </tr> <tr> <td>Labour and birth</td> <td></td> <td></td> </tr> <tr> <td>Induction of labour</td> <td>36</td> <td></td> </tr> <tr> <td>Duration of labour (hours)</td> <td></td> <td>9.53 (3-17)</td> </tr> <tr> <td>Epidural anaesthesia</td> <td>57</td> <td></td> </tr> <tr> <td>Nitrous oxide only</td> <td>7</td> <td></td> </tr> </tbody> </table>	Characteristic	Number	Mean (range)	Maternal			Maternal age (years)		26.6 (15-40)	Primiparous	50		Multiparous	23		Labour and birth			Induction of labour	36		Duration of labour (hours)		9.53 (3-17)	Epidural anaesthesia	57		Nitrous oxide only	7		<p>A CTG interpretation algorithm was designed by the study authors, which classified traces as normal or abnormal. An abnormal trace was defined by one or more of the following criteria.</p> <ol style="list-style-type: none"> <li>Tachycardia (fetal heart rate &gt;160 bpm) for more than 30 minutes during labour</li> <li>Bradycardia (fetal heart rate &lt;110 bpm) for</li> </ol>	<p>The categorisation of CTG traces as normal or abnormal by the computer algorithm was compared to the outcome of fetal acidosis. Acidosis was defined by an umbilical artery pH of less than 7.15, or by a base excess (BE) of less than -8mmol/l at birth. Results were reported as overall accuracy of the algorithm, as well as sensitivity and specificity</p>	<p><b>Diagnostic accuracy of computer algorithm for fetal acidosis, as defined by umbilical arterial pH of &lt;7.15</b> Sensitivity, % (95%CI): 87.5 (46.7 - 99.3)* Specificity, % (95% CI): 75.4 (62.9 - 84.9)* Positive likelihood ratio (95% CI): 3.55 (2.16 - 5.86)* Negative likelihood ratio (95% CI): 0.17 (0.03 - 1.05)*</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th rowspan="2"></th> <th colspan="2">Computer diagnosis</th> <th rowspan="2">Total</th> </tr> <tr> <th>Abnormal CTG</th> <th>Normal CTG</th> </tr> </thead> <tbody> <tr> <td rowspan="2">Reference standard</td> <td>Acidosis (pH &lt; 7.15)</td> <td>7</td> <td>1</td> <td>8</td> </tr> <tr> <td>No acidosis (pH ≥ 7.15)</td> <td>16</td> <td>49</td> <td>65</td> </tr> <tr> <td colspan="2">Total</td> <td>23</td> <td>50</td> <td>73</td> </tr> </tbody> </table> <p><b>Diagnostic accuracy of computer algorithm for fetal acidosis, as defined by base excess of less than -8mmol/l</b> Sensitivity, % (95% CI): 76.5 (49.8 - 92.2)* Specificity, % (95% CI): 82.1 (69.2 - 90.7)* Positive likelihood ratio (95% CI): 4.28 (2.30 - 7.96)* Negative likelihood ratio (95% CI): 0.29 (0.12 - 0.68)*</p>			Computer diagnosis		Total	Abnormal CTG	Normal CTG	Reference standard	Acidosis (pH < 7.15)	7	1	8	No acidosis (pH ≥ 7.15)	16	49	65	Total		23	50	73	<p>Selection of cases for CTG interpretation not well reported, and it was unclear whether a consecutive or random sampling approach was taken. Thresholds for fetal acidosis used differed from those pre-defined by the guideline committee as clinically significant</p> <p><b>Other information</b> <b>QUADAS 2 criteria</b> 1. Patient selection: Unclear risk - it is not clear how</p>
Characteristic	Number	Mean (range)																																																						
Maternal																																																								
Maternal age (years)		26.6 (15-40)																																																						
Primiparous	50																																																							
Multiparous	23																																																							
Labour and birth																																																								
Induction of labour	36																																																							
Duration of labour (hours)		9.53 (3-17)																																																						
Epidural anaesthesia	57																																																							
Nitrous oxide only	7																																																							
		Computer diagnosis		Total																																																				
		Abnormal CTG	Normal CTG																																																					
Reference standard	Acidosis (pH < 7.15)	7	1	8																																																				
	No acidosis (pH ≥ 7.15)	16	49	65																																																				
Total		23	50	73																																																				

Bibliographic details	Participants			Tests	Methods	Outcomes and results				Comments	
<p><b>Aim of the study</b></p> <p>To assess the ability of a computer software interpretation program to predict fetal acidosis at birth</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>	Other analgesia	9		<p>3. Low variation (standard deviation of the fetal heart rate of <math>\leq 3</math> bpm) for more than 60 minutes during labour</p> <p>4. More than five late decelerations (minimum of the fetal heart</p>				Computer diagnosis	Total	<p>CTG traces were selected for assessment</p> <p>2. Index test(s): Low risk</p> <p>3. Reference standard: Unclear risk - thresholds differ from those suggested by the guideline committee</p> <p>4. Flow and timing: Low risk</p>	
	Normal birth	39				Abnormal CTG	Normal CTG				
	Forceps birth	25				Reference standard	Acidosis (BE < -8 mmol/l)	13	4		17
	Caesarean section	9					No acidosis (BE $\geq$ -8mmol/l)	10	46		56
	Infant					Total		23	50		73
	Birthweight (g)		3226.25 (1500-4580)			*Sensitivity, specificity and likelihood ratios calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a>					
	Male infants	40									
	Female infants	33									
	Indication for fetal monitoring										
	Intrauterine growth restriction	10									
Pregnancy induced hypertension	9										

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
	<table border="1"> <tr> <td>Prolonged rupture of membranes</td> <td>2</td> <td></td> </tr> <tr> <td>Polyhydramnios</td> <td>2</td> <td></td> </tr> <tr> <td>Maternal anaemia</td> <td>3</td> <td></td> </tr> <tr> <td>Post term</td> <td>14</td> <td></td> </tr> <tr> <td>Meconium stained amniotic fluid</td> <td>6</td> <td></td> </tr> <tr> <td>Suspicious antepartum CTG</td> <td>9</td> <td></td> </tr> <tr> <td>Decreased fetal movements</td> <td>6</td> <td></td> </tr> <tr> <td>Other</td> <td>12</td> <td></td> </tr> </table>	Prolonged rupture of membranes	2		Polyhydramnios	2		Maternal anaemia	3		Post term	14		Meconium stained amniotic fluid	6		Suspicious antepartum CTG	9		Decreased fetal movements	6		Other	12		<p>5. rate occurring 20-60 seconds after the maximum of the contraction) during labour</p> <p>More than 10 variable decelerations (minimum of the FHR occurring more than 20 seconds prior to, or 60 seconds after, the maximum of</p>			
Prolonged rupture of membranes	2																												
Polyhydramnios	2																												
Maternal anaemia	3																												
Post term	14																												
Meconium stained amniotic fluid	6																												
Suspicious antepartum CTG	9																												
Decreased fetal movements	6																												
Other	12																												
	<p><b>Inclusion Criteria</b></p> <p>CTG traces were selected from women recruited at the University Hospital of Nottingham. Eligible women had a recognised indication for continuous fetal monitoring (in accordance with local</p>																												

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments															
	<p>management guidelines) and a labour that lasted for more than 3 hours</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p>the contraction) during labour</p>																		
<p><b>Full citation</b></p> <p>Costa, A., Santos, C., Ayres-de-Campos, D., Costa, C., Bernardes, J., Access to computerised analysis of intrapartum cardiotocographs improves clinicians' prediction of newborn umbilical artery blood pH, BJOG : an international journal of obstetrics and gynaecology,</p>	<p><b>Sample size</b></p> <p>N = 204 CTG traces  n = 104 randomised to receive computer analysis  n = 100 randomised to receive no computer analysis</p> <p><b>Characteristics</b></p> <table border="1" data-bbox="353 815 1106 1350"> <thead> <tr> <th data-bbox="353 815 669 938">Characteristic</th> <th data-bbox="669 815 857 938">Visual assessment n = 100</th> <th data-bbox="857 815 1106 938">Computerised assessment n = 104</th> </tr> </thead> <tbody> <tr> <td data-bbox="353 938 669 1023">Gestational age, weeks, mean (SD)</td> <td data-bbox="669 938 857 1023">39 (1)</td> <td data-bbox="857 938 1106 1023">39 (1)</td> </tr> <tr> <td data-bbox="353 1023 669 1107">Birth weight, g, mean (SD)</td> <td data-bbox="669 1023 857 1107">3362 (446)</td> <td data-bbox="857 1023 1106 1107">3282 (427)</td> </tr> <tr> <td data-bbox="353 1107 669 1192">Male births, n (%)</td> <td data-bbox="669 1107 857 1192">50 (50)</td> <td data-bbox="857 1107 1106 1192">46 (44)</td> </tr> <tr> <td data-bbox="353 1192 669 1350">Duration of assessed trace, minutes, median (minimum - maximum)</td> <td data-bbox="669 1192 857 1350">227 (60-770)</td> <td data-bbox="857 1192 1106 1350">213 (64-780)</td> </tr> </tbody> </table>	Characteristic	Visual assessment n = 100	Computerised assessment n = 104	Gestational age, weeks, mean (SD)	39 (1)	39 (1)	Birth weight, g, mean (SD)	3362 (446)	3282 (427)	Male births, n (%)	50 (50)	46 (44)	Duration of assessed trace, minutes, median (minimum - maximum)	227 (60-770)	213 (64-780)	<p><b>Tests</b></p> <p>The Omniview-SisPorto 3.5 system was used for CTG analysis</p>	<p><b>Methods</b></p> <p>Using computer-generated random numbers, traces were assigned to receive computer analysis by the Omniview SisPorto 3.5 system, or to no analysis (control group). The tracing printout in the study group had the baseline drawn on the fetal heart rate graph. Accelerations, decelerations, contractions and periods with abnormal</p>	<p><b>Results</b></p> <p><b>Accuracy of observers' prediction of umbilical arterial pH (within a margin of 0.1)</b>  For traces without computerised CTG analysis (control):  correct prediction of pH in 46% of cases (95% CI: 35% - 56%)  intraclass correlation coefficient = 0.29 (0.08 - 0.47)  For traces with computerised CTG analysis (intervention):  correct prediction of pH in 70% of cases (95% CI: 61% - 79%)  intraclass correlation coefficient = 0.52 (0.34 - 0.66)</p> <p><b>Agreement between the three observers in prediction of umbilical arterial pH</b>  For traces without computerised CTG analysis (control):  intraclass correlation coefficient = 0.43 (0.21 - 0.60)  For traces with computerised CTG analysis (intervention):  intraclass correlation coefficient = 0.70 (0.61 - 0.77)  (Study authors reported that the difference between these results was statistically significant; a p value was not reported)</p> <p><b>Agreement between the three observers in prediction of 5 minute Apgar score</b>  For traces without computerised CTG analysis (control):  intraclass correlation coefficient = 0.42 (0.25 to 0.57)  For traces with computerised CTG analysis (intervention):</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <ol style="list-style-type: none"> <li>1. Patient selection: Low risk</li> <li>2. Index tests: Low risk</li> <li>3. Reference standard: Low risk</li> <li>4. Flow and timing: Low risk</li> </ol>
Characteristic	Visual assessment n = 100	Computerised assessment n = 104																		
Gestational age, weeks, mean (SD)	39 (1)	39 (1)																		
Birth weight, g, mean (SD)	3362 (446)	3282 (427)																		
Male births, n (%)	50 (50)	46 (44)																		
Duration of assessed trace, minutes, median (minimum - maximum)	227 (60-770)	213 (64-780)																		

Bibliographic details	Participants			Tests	Methods	Outcomes and results	Comments
<p>117, 1288-1293, 2010</p> <p><b>Ref Id</b></p> <p>446136</p> <p><b>Country/ies where the study was carried out</b></p> <p>Portugal</p> <p><b>Study type</b></p> <p>Randomised controlled study</p> <p><b>Aim of the study</b></p> <p>To assess whether access to computerised CTG analysis improves clinicians' prediction of neonatal outcomes (umbilical artery pH and 5 minute Apgar score)</p>	<p>Cord artery pH, mean (SD) [21 missing values]</p>	<p>7.25 (0.08)</p>		<p>7.22 (0.08)</p>	<p>short term and long term variability were highlighted.</p> <p>The last alert elicited by the system was also displayed underneath the tracing.</p> <p>Traces in the control group showed only the standard fetal heart rate and uterine contraction signals.</p> <p>All traces were presented independently to three obstetricians with more than 5 years of experience in CTG interpretation.</p> <p>With the information that tracings had been recorded in term pregnancies, and that timings to birth were those previously</p>	<p>intraclass correlation coefficient = 0.55 (0.37 to 0.68) (Study authors reported that the difference between these results was statistically significant; a p value was not reported)</p>	
	<p>5-minute Apgar scores, median (minimum - maximum)</p>	<p>10 (8 - 10)</p>		<p>10 (6 - 10)</p>			
	<p>Caesarean birth, n (%)</p>	<p>12 (12)</p>		<p>15 (14)</p>			
	<p><b>Inclusion Criteria</b></p> <p>Singleton pregnancies of more than 36 weeks' gestation, cephalic presentation, absence of known fetal malformations, active phase of labour, generally accepted indication for internal fetal heart rate monitoring (poor signal quality, heavy meconium staining, high-risk pregnancy etc), a minimum of 60 minutes of trace duration, signal loss in the last hour &lt; 20%, no complications with the potential to influence fetal oxygenation occurring between tracing end and delivery (difficult vaginal or abdominal fetal extractions, cord prolapse, maternal hypotension, shoulder dystocia etc), and no anaesthetic complications taking place at the time of surgery</p> <p><b>Exclusion Criteria</b></p> <p>Time interval between tracing end and vaginal delivery exceeded 5 minutes, or interval between tracing end and caesarean birth exceeded 20 minutes</p>						



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Study dates</b> Not reported</p> <p><b>Source of funding</b> Not financially supported</p>			<p>mentioned (5 minutes for vaginal birth, 20 minutes for caesarean birth), the obstetricians were asked to estimate the newborns' umbilical arterial pH (to 2 decimal places) and 5 minute Apgar scores. A predicted pH of within 0.1 of the actual result was considered to be accurate, as was an Apgar score of within 1</p>		
<p><b>Full citation</b> Costa, M. A., Ayres-de-Campos, D., Machado, A. P., Santos, C. C., Bernardes, J., Comparison</p>	<p><b>Sample size</b> n = 50 CTG traces</p> <p><b>Characteristics</b> Not reported</p>	<p><b>Tests</b> The Omniview SisPorto 3.5 system was used to analyse the CTG traces and determine baseline fetal heart rate, accelerations,</p>	<p><b>Methods</b> Three clinicians (all with &gt; 5 years' experience of CTG interpretation) initially assessed the traces</p>	<p><b>Results</b> <b>Agreement on baseline estimation</b> Agreement between observers, ICC (95% CI): 0.87 (0.84 - 0.90) Observer 1 and computer, ICC (95% CI): 0.79 (0.48 - 0.89) Observer 2 and computer, ICC (95% CI): 0.88 (0.74 - 0.93) Observer 3 and computer, ICC (95% CI): 0.78 (0.27 - 0.91)</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS criteria</b></p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>of a computer system evaluation of intrapartum cardiotocographic events and a consensus of clinicians, Journal of Perinatal Medicine, 38, 191-5, 2010</p> <p><b>Ref Id</b></p> <p>457633</p> <p><b>Country/ies where the study was carried out</b></p> <p>Portugal</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To compare computer analysis of</p>	<p><b>Inclusion Criteria</b></p> <p>Singleton pregnancies of more than 36 weeks' gestation. Traces were recorded as part of a previously conducted randomised controlled trial. Included CTGs were of more than 60 minutes' duration with less than 10% signal loss</p> <p><b>Exclusion Criteria</b></p> <p>None reported</p>	<p>decelerations and contractions</p>	<p>independently. A second round of assessment was promoted for discordant segments of CTG, but without informing the clinicians of the other observers' results. Finally, a consensus meeting was held between all three clinicians to review the second round discordant segments. CTG segments which remained discordant after the third round were discarded from further analysis. Determination of agreement in baseline rate was assessed using the intraclass</p>	<p>Consensus of observers and computer, ICC (95% CI): 0.85 (0.46 - 0.93)</p> <p><b>Agreement on accelerations</b></p> <p>Agreement between observers, proportion of agreement (95% CI): 60% (48 - 66)</p> <p>Observer 1 and computer, proportion of agreement (95% CI): 68% (52 - 75)</p> <p>Observer 2 and computer, proportion of agreement (95% CI): 69% (55 - 76)</p> <p>Observer 3 and computer, proportion of agreement (95% CI): 65% (50 - 71)</p> <p>Consensus of observers and computer, proportion of agreement (95% CI): 71% (69 - 73)</p> <p><b>Agreement on decelerations</b></p> <p>Agreement between observers, proportion of agreement (95% CI): 65% (57 - 69)</p> <p>Observer 1 and computer, proportion of agreement (95% CI): 63% (51 - 68)</p> <p>Observer 2 and computer, proportion of agreement (95% CI): 62% (49 - 65)</p> <p>Observer 3 and computer, proportion of agreement (95% CI): 61% (51 - 68)</p> <p>Consensus of observers and computer, proportion of agreement (95% CI): 68% (66 - 70)</p> <p><b>Agreement on contractions</b></p> <p>Agreement between observers, proportion of agreement (95% CI): 93% (90 - 95)</p> <p>Observer 1 and computer, proportion of agreement (95% CI): 86% (83 - 88)</p> <p>Observer 2 and computer, proportion of agreement (95% CI): 84% (83 - 87)</p> <p>Observer 3 and computer, proportion of agreement (95% CI): 85% (81 - 90)</p> <p>Consensus of observers and computer, proportion of agreement (95% CI): 87% (85 - 89)</p>	<p>1. Patient selection - low risk</p> <p>2. Index tests - low risk</p> <p>3. Reference standard - low risk</p> <p>4. Flow and timing - low risk</p>

<b>Bibliographic details</b>	<b>Participants</b>	<b>Tests</b>	<b>Methods</b>	<b>Outcomes and results</b>	<b>Comments</b>				
<p>intrapartum CTG features using the Omniview SisPorto 3.5 system with interpretation by clinicians</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>None reported</p>			<p>correlation coefficient, the proportions of specific agreement and the limits of agreement. Agreement in determining accelerations, decelerations and contractions was assessed using the proportions of specific agreement and 95% confidence interval (CI)</p>						
<p><b>Full citation</b></p> <p>Keith, R. D., Beckley, S., Garibaldi, J. M., Westgate, J. A., Ifeachor, E. C., Greene, K. R., A multicentre comparative study of 17 experts and</p>	<p><b>Sample size</b></p> <p>n = 50 CTG traces</p> <p><b>Characteristics</b></p> <table border="1" data-bbox="349 1145 658 1315"> <thead> <tr> <th data-bbox="349 1145 613 1230">Characteristic</th> <th data-bbox="613 1145 658 1230">n</th> </tr> </thead> <tbody> <tr> <td data-bbox="349 1230 613 1315">Mode of birth</td> <td data-bbox="613 1230 658 1315"></td> </tr> </tbody> </table>	Characteristic	n	Mode of birth		<p><b>Tests</b></p> <p>The computerised system used in this study was developed by the study authors to assist clinical staff in their interpretation of CTG and consequent labour</p>	<p><b>Methods</b></p> <p>A panel of 17 experts was asked to independently score the CTGs to provide a reference standard for comparison with the computerised system. The</p>	<p><b>Results</b></p> <p>Agreement in scoring between the computerised system and experts: <math>\kappa = 0.31</math>                      Consistency in scoring for the computerised system: <math>\kappa = 0.98</math>                      The computer system identified the need for intervention for 2/3 cases of birth asphyxia, 2/4 cases of metabolic acidosis and 2/5 cases of acidosis.                      The computer system recommended no unnecessary intervention in all of the 11 cases with a good perinatal outcome (normal vaginal birth with an arterial pH &gt;7.15, venous pH &gt;7.20 and 5 minute Apgar score <math>\geq 9</math> with no resuscitation)</p>	<p><b>Limitations</b></p> <p>The system used in this article incorporated both CTG data and clinical information</p>
Characteristic	n								
Mode of birth									

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments														
<p>an intelligent computer system for managing labour using the cardiotocogram, British Journal of Obstetrics &amp; Gynaecology, 102, 688-700, 1995</p> <p><b>Ref Id</b> 457998</p> <p><b>Country/ies where the study was carried out</b> UK</p> <p><b>Study type</b> Retrospective cohort study</p> <p><b>Aim of the study</b> To investigate whether computer software</p>	<table border="1" data-bbox="353 304 658 882"> <tr> <td>Vaginal birth</td> <td>21</td> </tr> <tr> <td>Forceps birth</td> <td>13</td> </tr> <tr> <td>Caesarean section</td> <td>16</td> </tr> <tr> <td><b>Outcome</b></td> <td></td> </tr> <tr> <td>Birth asphyxia<sup>1</sup></td> <td>3</td> </tr> <tr> <td>Metabolic acidosis<sup>2</sup></td> <td>4</td> </tr> <tr> <td>Acidosis<sup>3</sup></td> <td>5</td> </tr> </table> <p><sup>1</sup> Cord arterial pH &lt; 7.05, base deficit ≥ 12 and Apgar score at 5 minutes of ≤7 with neonatal morbidity  <sup>2</sup> Cord arterial pH &lt; 7.05, base deficit ≥ 12 and Apgar score at 5 minutes of &gt;7 with no neonatal morbidity  <sup>3</sup> Cord arterial pH &lt; 7.05 and base deficit &lt;12 with no neonatal morbidity</p> <p><b>Inclusion Criteria</b> CTGs were selected from a database of 2400 high risk labours in which cord blood gas analysis and Apgar scores had been recorded</p> <p><b>Exclusion Criteria</b></p>	Vaginal birth	21	Forceps birth	13	Caesarean section	16	<b>Outcome</b>		Birth asphyxia <sup>1</sup>	3	Metabolic acidosis <sup>2</sup>	4	Acidosis <sup>3</sup>	5	<p>management. The system extracts relevant data from the CTG using numerical algorithms (including signal quality, baseline heart rate, heart rate variability, accelerations, the magnitude and timing of decelerations). These features are classified using additional algorithms and a small neural net. Relevant clinical information (such as cervical dilatation, risk factors and analgesia) is then considered. The system interprets all of these features using a database of over 400 rules which are used to recommend action</p>	<p>experts were asked to score 15 minute segments of CTG trace according to the following five-point protocol.</p> <ol style="list-style-type: none"> <li>1. I am not concerned for this fetus</li> <li>2. I have concerns for this fetus, but they are not sufficient to request fetal blood sampling (FBS); I may</li> </ol>		<p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <p>1. Patient selection: Unclear risk - selection of cases is not fully reported</p> <p>2. Index tests: Low risk</p> <p>3. Reference standard: Low risk</p> <p>4. Flow and timing: High risk - unclear how the system performed with regard to women who had an intervention for birth but a normal perinatal outcome</p>
Vaginal birth	21																		
Forceps birth	13																		
Caesarean section	16																		
<b>Outcome</b>																			
Birth asphyxia <sup>1</sup>	3																		
Metabolic acidosis <sup>2</sup>	4																		
Acidosis <sup>3</sup>	5																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>which integrates CTG interpretation and clinical features has a performance comparable to experts in the management of labour</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>The Mason Medical Research Foundation, the Northcott Devon Medical Foundation, the Science and Engineering Research Council, the South Western Region</p>	<p>Cases which had been previously reviewed by the computerised system or used to build its knowledge.</p>		<p>take some remedial action</p> <p>3. I am sufficiently concerned to request FBS or, if possible, a simple vaginal birth</p> <p>4. The information I have leads me to be seriously concerned for this fetus; I am not</p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
Health Authority and the Polytechnic Central Funding Council			<p>going to recommend immediate birth although I am thinking of expediting birth and will do so if things deteriorate further</p> <p>5. I am so concerned for this fetus that I want immediate birth</p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
			<p>A method was derived to identify agreement between any two sets of scoring sequences. This gave a value of 0 if no similarity was seen, and 1 if perfect concordance was present. The method incorporated a weighted agreement matrix which rewarded similar scores given to a particular segment, but heavily penalised widely differing scores. The method also awarded a partial agreement when two experts took a major decision close to each other, but not within the</p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																		
			same segment of CTG. The agreement between the system and each of the 17 experts was calculated for each case and averaged																				
<p><b>Full citation</b></p> <p>Mongelli, M., Dawkins, R., Chung, T., Sahota, D., Spencer, J.A., Chang, A.M., Computerised estimation of the baseline fetal heart rate in labour: the low frequency line, British Journal of Obstetrics and Gynaecology, 104, 1128-1133, 1997</p>	<p><b>Sample size</b></p> <p>n = 60 CTG traces</p> <p><b>Characteristics</b></p> <table border="1" data-bbox="353 842 819 1337"> <thead> <tr> <th>Characteristic</th> <th>%</th> <th>Mean (SD)</th> </tr> </thead> <tbody> <tr> <td>Nulliparous</td> <td>57</td> <td></td> </tr> <tr> <td>Induction of labour</td> <td>25</td> <td></td> </tr> <tr> <td>Operative birth</td> <td>55</td> <td></td> </tr> <tr> <td>Gestational age, weeks</td> <td></td> <td>39.8 (1.8)</td> </tr> <tr> <td>Birthweight, g</td> <td></td> <td>3373 (447)</td> </tr> </tbody> </table>	Characteristic	%	Mean (SD)	Nulliparous	57		Induction of labour	25		Operative birth	55		Gestational age, weeks		39.8 (1.8)	Birthweight, g		3373 (447)	<p><b>Tests</b></p> <p>The fetal electrocardiogram signal was collected using a fetal scalp electrode. A computer algorithm was developed to estimate the baseline fetal heart rate, with an aim to produce a low frequency line that would be stable under noisy conditions yet responsive to both sudden and gradual changes. Values outside the range of 30</p>	<p><b>Methods</b></p> <p>Sixty 40-minute segments of intrapartum CTG records were selected from 60 different women. Traces were chosen on the grounds of complexity and potential difficulty in interpretation. The tracings were reproduced and sent to 12 clinical experts for their estimation of the baseline.</p>	<p><b>Results</b></p> <p>The intraclass correlation between the computer and the panel of experts was in excess of 0.9. The 95% confidence interval (CI) for the difference in baseline between computer and experts was -12 to 15 bpm. The 95% CI for the difference in baseline between experts was -10 to 10 bpm</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <ol style="list-style-type: none"> <li>1. Patient selection - low risk</li> <li>2. Index test - low risk</li> <li>3. Reference standard - low risk</li> <li>4. Flow and timing - low risk</li> </ol>
Characteristic	%	Mean (SD)																					
Nulliparous	57																						
Induction of labour	25																						
Operative birth	55																						
Gestational age, weeks		39.8 (1.8)																					
Birthweight, g		3373 (447)																					



Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p><b>Ref Id</b> 196506</p> <p><b>Country/ies where the study was carried out</b> UK</p> <p><b>Study type</b> Retrospective cohort study</p> <p><b>Aim of the study</b> To develop a computerised algorithm for the determination of fetal heart rate baseline during labour</p> <p><b>Study dates</b> Not reported</p>	<p><b>Inclusion Criteria</b> All women had electronic fetal monitoring because of perceived high fetal risks. Traces for this study were selected because of complexity and potential difficulty in interpretation</p> <p><b>Exclusion Criteria</b> Not reported</p>	<p>to 240 bpm were considered as noise and excluded from analysis</p>	<p>Of these, 8 were NHS consultants or senior academics, and 4 were of senior registrar/lecturer status</p>		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																								
<p><b>Source of funding</b></p> <p>Not reported</p>																													
<p><b>Full citation</b></p> <p>Nielsen, P. V., Stigsby, B., Nickelsen, C., Nim, J., Computer assessment of the intrapartum cardiotocogram. II. The value of compared with visual assessment, Acta Obstetrica et Gynecologica Scandinavica, 67, 461-4, 1988</p> <p><b>Ref Id</b></p> <p>454968</p> <p><b>Country/ies where the</b></p>	<p><b>Sample size</b></p> <p>Not reported; 50 CTG records</p> <p><b>Characteristics</b></p> <p>Pregnant women in the first stage of labour</p> <p><b>Inclusion Criteria</b></p> <p>Not reported</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>The computer Cardiotocographic Assessment System (CAS)</p>	<p><b>Methods</b></p> <p>The CTGs were assessed both by 4 obstetricians and the computer system as being normal or pathological. The 4 obstetricians, all experienced in EFM, had been working in the same department, using EFM routinely in all births. They were informed of the incidence of compromised infants (one-third). The newborn was declared compromised if the 1-minute</p>	<p><b>Results</b></p> <p>Compared with individual obstetricians the computer system achieved the highest accuracy, and compared with the obstetrician obtaining the best accuracy, the result of the computer was significantly better. Computer assessment of 50 CTGs compared with 4 obstetricians' visual assessment</p> <table border="1" data-bbox="1290 703 1897 1241"> <thead> <tr> <th></th> <th>Fetal outcome</th> <th></th> <th></th> <th></th> <th></th> </tr> <tr> <th></th> <th>Normal</th> <th>Compromised</th> <th>Total</th> <th>Fisher's test</th> <th>Accuracy</th> </tr> </thead> <tbody> <tr> <td>Computer</td> <td>N 32</td> <td>5</td> <td>37</td> <td>&lt;0.001</td> <td>86%</td> </tr> <tr> <td></td> <td>P 2</td> <td>11</td> <td>13</td> <td></td> <td></td> </tr> </tbody> </table>		Fetal outcome						Normal	Compromised	Total	Fisher's test	Accuracy	Computer	N 32	5	37	<0.001	86%		P 2	11	13			<p><b>Limitations</b></p> <p><b>QUADAS 2 criteria</b></p> <p>1. Patient selection - High risk; selection of CTGs was not reported to be random or consecutive</p> <p>2. Index tests - High risk; not clear if the index test was interpreted without knowledge of the results of the reference standard</p> <p>3. Reference</p>
	Fetal outcome																												
	Normal	Compromised	Total	Fisher's test	Accuracy																								
Computer	N 32	5	37	<0.001	86%																								
	P 2	11	13																										

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																																										
<p><b>study was carried out</b></p> <p>Denmark</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To compare the accuracy of a computer Cardiotocographic Assessment System (CAS) with that of four very skilled obstetricians' using the same set of CTGs</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p>			<p>Apgar score was below 7, or the umbilical arterial blood was acidotic (pH &lt; 7.15 or standard base excess below - 10 meq/l), or primary resuscitation was needed. The CAS operates as follows.</p> <p>1) The first program automatically detects the CTG patterns (decelerations, accelerations, uterine contractions, baseline and resting tone) and describes these patterns by 17 variables (duration, amplitude, and area of each acceleration, deceleration, and contraction; level of baseline and resting tone;</p>	<table border="1"> <thead> <tr> <th data-bbox="1292 304 1417 472">Obstetricians</th> <th data-bbox="1417 304 1534 472">1 N</th> <th data-bbox="1534 304 1671 472">24</th> <th data-bbox="1671 304 1807 472">9</th> <th data-bbox="1807 304 1906 472">33</th> <th data-bbox="1906 304 2004 472">0.2</th> <th data-bbox="2004 304 2045 472">62%</th> </tr> </thead> <tbody> <tr> <td data-bbox="1292 472 1417 639"></td> <td data-bbox="1417 472 1534 639">1 P</td> <td data-bbox="1534 472 1671 639">10</td> <td data-bbox="1671 472 1807 639">7</td> <td data-bbox="1807 472 1906 639">17</td> <td data-bbox="1906 472 2004 639"></td> <td data-bbox="2004 472 2045 639"></td> </tr> <tr> <td data-bbox="1292 639 1417 807"></td> <td data-bbox="1417 639 1534 807">2 N</td> <td data-bbox="1534 639 1671 807">28</td> <td data-bbox="1671 639 1807 807">11</td> <td data-bbox="1807 639 1906 807">39</td> <td data-bbox="1906 639 2004 807">0.2</td> <td data-bbox="2004 639 2045 807">66%</td> </tr> <tr> <td data-bbox="1292 807 1417 975"></td> <td data-bbox="1417 807 1534 975">2 P</td> <td data-bbox="1534 807 1671 975">6</td> <td data-bbox="1671 807 1807 975">5</td> <td data-bbox="1807 807 1906 975">11</td> <td data-bbox="1906 807 2004 975"></td> <td data-bbox="2004 807 2045 975"></td> </tr> <tr> <td data-bbox="1292 975 1417 1142"></td> <td data-bbox="1417 975 1534 1142">3 N</td> <td data-bbox="1534 975 1671 1142">18</td> <td data-bbox="1671 975 1807 1142">5</td> <td data-bbox="1807 975 1906 1142">23</td> <td data-bbox="1906 975 2004 1142">0.1</td> <td data-bbox="2004 975 2045 1142">58%</td> </tr> <tr> <td data-bbox="1292 1142 1417 1294"></td> <td data-bbox="1417 1142 1534 1294">3 P</td> <td data-bbox="1534 1142 1671 1294">16</td> <td data-bbox="1671 1142 1807 1294">11</td> <td data-bbox="1807 1142 1906 1294">27</td> <td data-bbox="1906 1142 2004 1294"></td> <td data-bbox="2004 1142 2045 1294"></td> </tr> </tbody> </table>	Obstetricians	1 N	24	9	33	0.2	62%		1 P	10	7	17				2 N	28	11	39	0.2	66%		2 P	6	5	11				3 N	18	5	23	0.1	58%		3 P	16	11	27			<p>standard - Low risk</p> <p>4. Flow and timing - Low risk</p> <p><b>Other information</b></p>
Obstetricians	1 N	24	9	33	0.2	62%																																									
	1 P	10	7	17																																											
	2 N	28	11	39	0.2	66%																																									
	2 P	6	5	11																																											
	3 N	18	5	23	0.1	58%																																									
	3 P	16	11	27																																											

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																																	
<p>The development of the computer system was supported by the Danish Medical Research Council, grant numbers 12-3832, 5.52.13.16 and 12-3202</p>			<p>baseline variability; slope of the descending part of the deceleration, recovery time, and residual area for the ascending part; lag time and latency time.                      2) The second program calculates 1) the number, 2) the mean value, 3) standard deviation, 4) and trend of each of the 17 variables for a chosen epoch of the CTG. This calculation results in 17x4=68 subvariables but 12 of these contain only duplicate information, leaving 56 subvariables to be considered in the</p>	<table border="1" data-bbox="1290 304 1895 715"> <tr> <td data-bbox="1290 304 1417 472"><b>N</b></td> <td data-bbox="1417 304 1532 472">4 20</td> <td data-bbox="1532 304 1671 472">11</td> <td data-bbox="1671 304 1727 472">31</td> <td data-bbox="1727 304 1807 472">8.8</td> <td data-bbox="1807 304 1895 472">50%</td> </tr> <tr> <td data-bbox="1290 472 1417 639"><b>P</b></td> <td data-bbox="1417 472 1532 639">4 14</td> <td data-bbox="1532 472 1671 639">5</td> <td data-bbox="1671 472 1727 639">19</td> <td data-bbox="1727 472 1807 639"></td> <td data-bbox="1807 472 1895 639"></td> </tr> <tr> <td data-bbox="1290 639 1417 715"><b>Total</b></td> <td data-bbox="1417 639 1532 715">34</td> <td data-bbox="1532 639 1671 715">16</td> <td data-bbox="1671 639 1727 715">50</td> <td data-bbox="1727 639 1807 715"></td> <td data-bbox="1807 639 1895 715"></td> </tr> </table> <p data-bbox="1290 719 1895 775">N=CTGs assessed as normal; P=CTGs assessed as pathological</p> <table border="1" data-bbox="1290 799 1895 1297"> <thead> <tr> <th data-bbox="1290 799 1417 967"></th> <th data-bbox="1417 799 1565 967">Sensitivity (95% CI)</th> <th data-bbox="1565 799 1713 967">Specificity (95%)</th> <th data-bbox="1713 799 1807 967">LR+ (95% CI)</th> <th data-bbox="1807 799 1895 967">LR- (95% CI)</th> </tr> </thead> <tbody> <tr> <td data-bbox="1290 967 1417 1129"><b>Computer</b></td> <td data-bbox="1417 967 1565 1129">68.8 (41.48-87.87)</td> <td data-bbox="1565 967 1713 1129">94.12 (78.94-98.97)</td> <td data-bbox="1713 967 1807 1129">11.7 (2.93-46.67)</td> <td data-bbox="1807 967 1895 1129">0.33 (0.16-0.69)</td> </tr> <tr> <td data-bbox="1290 1129 1417 1297"><b>Obs 1</b></td> <td data-bbox="1417 1129 1565 1297">43.75 (20.75-69.45)</td> <td data-bbox="1565 1129 1713 1297">70.59 (52.33-84.29)</td> <td data-bbox="1713 1129 1807 1297">1.49 (0.69-3.19)</td> <td data-bbox="1807 1129 1895 1297">0.8 (0.50-1.26)</td> </tr> </tbody> </table>	<b>N</b>	4 20	11	31	8.8	50%	<b>P</b>	4 14	5	19			<b>Total</b>	34	16	50				Sensitivity (95% CI)	Specificity (95%)	LR+ (95% CI)	LR- (95% CI)	<b>Computer</b>	68.8 (41.48-87.87)	94.12 (78.94-98.97)	11.7 (2.93-46.67)	0.33 (0.16-0.69)	<b>Obs 1</b>	43.75 (20.75-69.45)	70.59 (52.33-84.29)	1.49 (0.69-3.19)	0.8 (0.50-1.26)	
<b>N</b>	4 20	11	31	8.8	50%																																	
<b>P</b>	4 14	5	19																																			
<b>Total</b>	34	16	50																																			
	Sensitivity (95% CI)	Specificity (95%)	LR+ (95% CI)	LR- (95% CI)																																		
<b>Computer</b>	68.8 (41.48-87.87)	94.12 (78.94-98.97)	11.7 (2.93-46.67)	0.33 (0.16-0.69)																																		
<b>Obs 1</b>	43.75 (20.75-69.45)	70.59 (52.33-84.29)	1.49 (0.69-3.19)	0.8 (0.50-1.26)																																		

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments															
			<p>assessment of the CTG.                      3) The third program calculates the probability of the CTG belonging to a compromised infant. This probability is calculated by a discriminant function, and a CTG is considered pathological if the probability is above 0.5. The computer system's calculation of the probability of a compromised infant was for each CTG based on the experience from the other 49 CTG thus excluding the possibility of 'self-recognition'. The best combination of subvariables was found by</p>	<table border="1"> <tbody> <tr> <td data-bbox="1292 304 1422 472"><b>Obs 2</b></td> <td data-bbox="1422 304 1568 472">31.3 (12.113-58.52)</td> <td data-bbox="1568 304 1713 472">82.4 (64.83-92.61)</td> <td data-bbox="1713 304 1809 472">1.77 (0.63-4.95)</td> <td data-bbox="1809 304 1899 472">0.83 (0.59-1.18)</td> </tr> <tr> <td data-bbox="1292 472 1422 639"><b>Obs 3</b></td> <td data-bbox="1422 472 1568 639">68.8 (41.48-87.87)</td> <td data-bbox="1568 472 1713 639">52.9 (35.40-69.84)</td> <td data-bbox="1713 472 1809 639">1.5 (0.90-2.38)</td> <td data-bbox="1809 472 1899 639">0.6 (0.27-1.29)</td> </tr> <tr> <td data-bbox="1292 639 1422 807"><b>Obs 4</b></td> <td data-bbox="1422 639 1568 807">31.3 (12.13-58.52)</td> <td data-bbox="1568 639 1713 807">58.8 (40.83-74.87)</td> <td data-bbox="1713 639 1809 807">0.8 (0.33-1.74)</td> <td data-bbox="1809 639 1899 807">1.2 (0.81-1.69)</td> </tr> </tbody> </table> <p>"Sensitivity, specificity and likelihood ratios calculated by the NGA technical team using <a href="http://vassarstats.net/clin1.html">http://vassarstats.net/clin1.html</a>"</p>	<b>Obs 2</b>	31.3 (12.113-58.52)	82.4 (64.83-92.61)	1.77 (0.63-4.95)	0.83 (0.59-1.18)	<b>Obs 3</b>	68.8 (41.48-87.87)	52.9 (35.40-69.84)	1.5 (0.90-2.38)	0.6 (0.27-1.29)	<b>Obs 4</b>	31.3 (12.13-58.52)	58.8 (40.83-74.87)	0.8 (0.33-1.74)	1.2 (0.81-1.69)	
<b>Obs 2</b>	31.3 (12.113-58.52)	82.4 (64.83-92.61)	1.77 (0.63-4.95)	0.83 (0.59-1.18)																
<b>Obs 3</b>	68.8 (41.48-87.87)	52.9 (35.40-69.84)	1.5 (0.90-2.38)	0.6 (0.27-1.29)																
<b>Obs 4</b>	31.3 (12.13-58.52)	58.8 (40.83-74.87)	0.8 (0.33-1.74)	1.2 (0.81-1.69)																

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
			minimising the average probability of misclassification		
<p><b>Full citation</b></p> <p>Parer, J.T., Hamilton, E.F., Comparison of 5 experts and computer analysis in rule-based fetal heart rate interpretation, American Journal of Obstetrics and Gynecology, 203, 451-457, 2010</p> <p><b>Ref Id</b></p> <p>169819</p> <p><b>Country/ies where the study was carried out</b></p>	<p><b>Sample size</b></p> <p>N = 30 CTG traces</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion Criteria</b></p> <p>Singleton, term pregnancies with umbilical blood gas analysis present</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>PeriCALM computer software was used for CTG analysis. This software follows a strict rule-based system to classify the CTG based on fetal heart rate baseline, variability, and decelerations (depth, duration and timing). The scoring system results in a five-level classification system for CTGs. For a CTG to be coded as green (category 1) all features must be within normal limits. Progressively</p>	<p><b>Methods</b></p> <p>The CTGs were shown to five experts who were asked to follow the same strict, rule-based system to interpret the traces. The experts were given a copy of the rules and encouraged to follow them, even if they disagreed with them, as the purpose of the study was to assess concordance when using the rules. The percentage of exact agreement (where the</p>	<p><b>Results</b></p> <p><b>Exact agreement with all clinical decisions</b>            Computer software, % (95% CI): 44.9% (43.4 - 46.5)*            Experts (average agreement for all experts), % (95% CI): 45.5% (42.1 - 48.4)</p> <p><b>Exact agreement with majority clinical decisions</b>            Computer software, % (95% CI): 56.8% (52.6 - 61.0)*            Computer software, κ statistic: 0.52 (no CI reported)            Experts (average agreement for all experts), % (95% CI): 56.7% (49.4 - 63.9)            Experts, κ statistic (95% CI): 0.58 (0.48 - 0.68)</p> <p><b>Close agreement with majority clinical decisions</b>            Computer software, % (95% CI): 83.1% (79.7 - 86.1)*            Experts (average agreement for all experts), % (95% CI): 88.6% (80.8 - 96.4)</p> <p>* Confidence interval (CI) calculated by the NGA technical team using <a href="http://statpages.info/confint.html">http://statpages.info/confint.html</a></p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <p>1. Patient selection - high risk; selection of CTGs is not well reported, and it may not represent the population in whom this method would be used</p> <p>2. Index tests - low risk</p> <p>3. Reference</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>USA</p> <p><b>Study type</b></p> <p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To measure agreement between five expert clinicians and a computerised method with a strict rule-based method of CTG interpretation</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>No external funding reported</p>		<p>abnormal traces are coded as blue, yellow, orange and red.</p>	<p>computer assigned exactly the same colour category as the observers) was calculated, using the individual scores of each expert. The percentage of majority agreement was also calculated to assess how often the computer agreed with the score given by the majority of experts for any particular CTG segment. Finally, the percentage of 'close' agreement was calculated (when the computer assigned scores <math>\pm 1</math> category of the majority agreement).</p>		<p>standard - unclear risk; a specific 'rule-based' system was used by the experts to interpret the CTG for this study; this is likely to differ from how experts interpret the CTG in clinical practise</p> <p>4. Flow and timing - low risk</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments																				
			Agreement between experts was calculated in the same way																						
<p><b>Full citation</b></p> <p>Taylor,G.M., Mires,G.J., Abel,E.W., Tsantis,S., Farrell,T., Chien,P.F., Liu,Y., The development and validation of an algorithm for real-time computerised fetal heart rate monitoring in labour, BJOG: An International Journal of Obstetrics and Gynaecology , 107, 1130-1137, 2000</p> <p><b>Ref Id</b></p>	<p><b>Sample size</b></p> <p>n = 24 CTG traces taken from a total of 30 labours</p> <p><b>Characteristics</b></p> <table border="1" data-bbox="353 703 913 1281"> <thead> <tr> <th>Characteristic</th> <th>n/N</th> <th>median (range)</th> <th>mean (SD)</th> </tr> </thead> <tbody> <tr> <td>Induction of labour</td> <td>16/30</td> <td></td> <td></td> </tr> <tr> <td>Maternal age, years</td> <td></td> <td>27.5 (18-35)</td> <td></td> </tr> <tr> <td>Primiparous</td> <td>16/30</td> <td></td> <td></td> </tr> <tr> <td>Duration of labour, minutes</td> <td></td> <td>484 (143 - 1155)</td> <td></td> </tr> </tbody> </table>	Characteristic	n/N	median (range)	mean (SD)	Induction of labour	16/30			Maternal age, years		27.5 (18-35)		Primiparous	16/30			Duration of labour, minutes		484 (143 - 1155)		<p><b>Tests</b></p> <p>Cardiotocograms were recorded using a fetal scalp electrode. A computer algorithm was developed to identify key features of the CTG, including baseline fetal heart rate, fetal heart variability, accelerations and decelerations</p>	<p><b>Methods</b></p> <p>The cardiotocograms were analysed independently by 7 reviewers, all of whom were senior obstetric staff (consultants or senior specialist registrars) who were actively involved in the labour ward. Each reviewer assessed the baseline heart rate, the number of accelerations and the number and type of decelerations. The inter-rater reliability of the components of</p>	<p><b>Results</b></p> <p><b>Inter-rater reliability between expert reviewers</b>                      Baseline fetal heart rate: intraclass correlation coefficient 0.93                      Number of decelerations: intraclass correlation coefficient 0.93                      Number of late decelerations: intraclass correlation coefficient 0.79                      Number of accelerations: intraclass correlation coefficient 0.27                      Baseline variability: <math>\kappa</math> statistic 0.27</p> <p><b>Validity of computerised algorithm when compared to expert reviewers</b>                      Baseline fetal heart rate: intraclass correlation coefficient 0.91 to 0.98                      Number of decelerations: intraclass correlation coefficient 0.82 to 0.92                      Number of late decelerations: intraclass correlation coefficient 0.68 to 0.85                      Number of accelerations: intraclass correlation coefficient 0.06 to 0.80                      Baseline variability: <math>\kappa</math> statistic 0.00 to 0.34</p>	<p><b>Limitations</b></p> <p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <ol style="list-style-type: none"> <li>1. Patient selection - Unclear risk; methods of participant recruitment are not reported</li> <li>2. Index tests - Low risk</li> <li>3. Reference standard - Low risk</li> <li>4. Flow and timing - Unclear risk; it is not clear why 30 CTGs were recorded, but only 24</li> </ol>
Characteristic	n/N	median (range)	mean (SD)																						
Induction of labour	16/30																								
Maternal age, years		27.5 (18-35)																							
Primiparous	16/30																								
Duration of labour, minutes		484 (143 - 1155)																							



Bibliographic details	Participants				Tests	Methods	Outcomes and results	Comments
197103  <b>Country/ies where the study was carried out</b>  UK  <b>Study type</b>  Prospective cohort study  <b>Aim of the study</b>  To develop and validate a computerised algorithm for the interpretation of the characteristics of the intrapartum CTG  <b>Study dates</b>  Not reported  <b>Source of funding</b>	Operative vaginal birth	6/30				the CTG for the expert reviewers, and the validity of the computer algorithm were assessed with the intra-class correlation coefficient for continuous variables (baseline heart rate, number of accelerations, number of decelerations), and by the kappa statistic for dichotomous variables (baseline variability). 24 CTGs were randomly chosen for review		'randomly' selected for use in the study; methods for random selection are not reported
	Caesarean section	7/30						
	Birthweight, g			3538 (526)				
	Gestational age, weeks			40.1 (1.6)				
	Admission to SCBU	0/30						
	SCBU: special care baby unit							
	<b>Inclusion Criteria</b>							
	Women in active labour or undergoing induction of labour							
	<b>Exclusion Criteria</b>							
	Not reported							

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
Not reported					
<p><b>Full citation</b></p> <p>Todros,T., Preve,C.U., Plazzotta,C., Biolcati,M., Lombardo,P., Fetal heart rate tracings: observers versus computer assessment, European Journal of Obstetrics, Gynecology, and Reproductive Biology, 68, 83-86, 1996</p> <p><b>Ref Id</b></p> <p>196732</p> <p><b>Country/ies where the study was carried out</b></p> <p>Italy</p> <p><b>Study type</b></p>	<p><b>Sample size</b></p> <p>N = 63 CTG recordings</p> <p><b>Characteristics</b></p> <p>Not reported</p> <p><b>Inclusion Criteria</b></p> <p>High- and low-risk pregnancies between 30 and 41 weeks of gestation</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p><b>Tests</b></p> <p>A 25 minute strip of CTG from each of 63 tracings was randomly chosen. The 2CTG computerised system was used to analyse the traces. The computer output variables included in the analysis were: baseline heart rate, the amplitude bandwidth around the baselines (a measure of long-term variability), the number of accelerations, and the number and timing of decelerations</p>	<p><b>Methods</b></p> <p>Four observers independently assessed the CTG traces for the same variables. Two of the observers were consultants with experience of reading CTGs (experts) and 2 were residents with 1 year of experience (non-experts)</p>	<p><b>Results</b></p> <p><b>Reproducibility among observers</b>  Baseline fetal heart rate: <math>\kappa</math> statistic 0.65  Variability: <math>\kappa</math> statistic 0.38  Accelerations: <math>\kappa</math> statistic 0.58  Number of decelerations: <math>\kappa</math> statistic 0.67  Type of decelerations: <math>\kappa</math> statistic 0.05</p> <p><b>Concordance between expert observers and the computer system</b>  Baseline fetal heart rate: <math>\kappa</math> statistic 0.18 to 0.48  Variability: <math>\kappa</math> statistic 0.16 to 0.74  Accelerations: <math>\kappa</math> statistic 0.58 to 0.64  Number of decelerations: <math>\kappa</math> statistic 0.41 to 0.45</p> <p><b>Concordance between non-expert observers and the computer system</b>  Baseline fetal heart rate: <math>\kappa</math> statistic 0.24 to 0.36  Variability: <math>\kappa</math> statistic 0.65 to 0.69  Accelerations: <math>\kappa</math> statistic 0.37 to 0.48  Number of decelerations: <math>\kappa</math> statistic 0.54</p>	<p><b>Limitations</b></p> <p>CTG traces used were from women at 30 to 41 weeks of gestation. It is unclear whether the recordings were all made intrapartum, or whether some were taken antenatally</p> <p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b>  1. Patient selection - high risk; CTGs included those from premature gestations,</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To assess the reproducibility of CTG interpretation among observers and between observers and a computerised system</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>The Italian National Research Council</p>					<p>and it is unclear whether all women were in labour at the time of monitoring</p> <p>2. Index tests - low risk</p> <p>3. Reference standard - low risk</p> <p>4. Flow and timing - low risk</p>
<p><b>Full citation</b></p> <p>Wolfberg,A.J ..</p>	<p><b>Sample size</b></p> <p>n = 30 CTG traces</p>	<p><b>Tests</b></p> <p>Mean fetal heart rate was</p>	<p><b>Methods</b></p> <p>Four perinatologists</p>	<p><b>Results</b></p> <p><b>Correlation between the computer analysis and the (average) expert interpretation of variability</b></p>	<p><b>Limitations</b></p> <p>Correlation was</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>Derosier,D.J., Roberts,T., Syed,Z., Clifford,G.D., Acker,D., Plessis,A.D., A comparison of subjective and mathematical estimations of fetal heart rate variability, Journal of Maternal-Fetal and Neonatal Medicine, 21, 101-104, 2008</p> <p><b>Ref Id</b></p> <p>169793</p> <p><b>Country/ies where the study was carried out</b></p> <p>USA</p> <p><b>Study type</b></p>	<p><b>Characteristics</b></p> <p>Apgar scores for all infants were greater than 6 at both 1 and 5 minutes, and there were no neonatal complications for any of the newborns</p> <p><b>Inclusion Criteria</b></p> <p>Women in labour who had a fetal scalp electrode positioned for clinical indications. Singleton pregnancies, between 35 and 41 weeks' gestation</p> <p><b>Exclusion Criteria</b></p> <p>Not reported</p>	<p>calculated over a 10 minute period for each of the CTG recordings. The variance was then calculated for the same period. The standard deviation was used as the computed measure of fetal heart rate variability</p>	<p>with recognised expertise in CTG interpretation were asked to assess the variability for the same 10 minute segments of CTG. They were asked to give a value for the variability (to the closest integer, not a range) and to rate the variability according to NICHD criteria as absent, minimal, moderate, marked or sinusoidal</p>	<p>Intraclass correlation coefficient 0.62 (range 0.27 to 0.68)</p> <p><b>Correlation between the expert interpretation of variability</b></p> <p>Intraclass correlation coefficient 0.44 (range 0.33 to 0.72)</p>	<p>reported for determining the absolute variability for the fetal heart rate (i.e. a specific value). The computerised results were not further categorised into categories of variability according to the NICHD criteria. Therefore the correlation between the computer and experts for different categories of variability was not reported</p>

Bibliographic details	Participants	Tests	Methods	Outcomes and results	Comments
<p>Retrospective cohort study</p> <p><b>Aim of the study</b></p> <p>To develop a computer algorithm to determine baseline fetal heart rate variability, and compare it to clinicians' interpretation</p> <p><b>Study dates</b></p> <p>Not reported</p> <p><b>Source of funding</b></p> <p>Not reported</p>					<p><b>Other information</b></p> <p><b>QUADAS 2 criteria</b></p> <p>1. Patient selection – unclear risk; insufficient data were reported with regard to selection of participants</p> <p>2. Index tests – low risk</p> <p>3. Reference standard – low risk</p> <p>4. Flow and timing – low risk</p>