

Ectopic pregnancy and miscarriage: diagnosis and initial management

[C] Evidence review for progestogens for preventing miscarriage

NICE guideline NG126 (update)

Evidence review underpinning recommendation 1.5.2 and research recommendations in the NICE guideline

July 2021

Draft for consultation

This evidence review was developed by the National Guideline Alliance which is a part of the Royal College of Obstetricians and Gynaecologists

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to [Notice of rights](#).

ISBN:

Contents

Evidence review for progestogens for preventing miscarriage	6
Review question	6
Introduction	6
Summary of the protocol	6
Methods and process	7
Effectiveness evidence.....	7
Summary of included studies.....	7
Summary of the evidence.....	8
Economic evidence	10
Summary of included economic evidence.....	10
Economic model.....	15
The committee’s discussion and interpretation of the evidence	15
Recommendations supported by this evidence review	19
References.....	20
Appendices	22
Appendix A Review protocols	22
Review protocol for review question: What is the clinical and cost- effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	22
Appendix B Literature search strategies	23
Literature search strategies for review question: What is the clinical and cost- effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	23
Appendix C Effectiveness	24
Study selection for: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?.....	24
Appendix D Evidence tables	25
Evidence tables for review question: What is the clinical and cost- effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	25
Appendix E Forest plots	26
Forest plots for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage? ...	26
Appendix F GRADE tables	27
GRADE tables for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	27
Appendix G Economic evidence study selection	28
Study selection for: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?.....	28
Appendix H Economic evidence tables	29

	Economic evidence tables for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	29
Appendix I	Economic model	33
	Economic model for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	33
Appendix J	Excluded studies	34
	Excluded studies for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	34
Appendix K	Research recommendations – full details	35
	Research recommendations for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?	35
K.1.1	Research recommendation.....	35
K.1.2	Why this is important.....	35
K.1.3	Rationale for research recommendation	35
K.1.4	Modified PICO table	36
K.1.5	Research recommendation.....	36
K.1.6	Why this is important.....	36
K.1.7	Rationale for research recommendation	36
K.1.8	Modified PICO table	37
Appendix L	Methodological limitations	38

1 Evidence review for progestogens for 2 preventing miscarriage

3 Review question

4 What is the clinical and cost-effectiveness of progestogens in improving outcomes in women
5 at risk of miscarriage?

6 Introduction

7 Miscarriage is a common complication of early pregnancy, affecting about 20% of all
8 pregnancies. It can have serious consequences for women, including psychological harm
9 and mental health complications such as anxiety, depression and post-traumatic stress
10 disorder. Progesterone is an essential hormone secreted by the corpus luteum that provides
11 early pregnancy support until placental production takes over at 10 to 12 weeks' gestation.
12 Low levels of circulating progesterone have been linked to impending miscarriage and the
13 presence of associated early pregnancy bleeding. It has been postulated that a lack of
14 progesterone is a cause of miscarriage rather than a secondary signal of failing pregnancy,
15 and that augmenting levels of progesterone may prevent miscarriage of euploid (without any
16 genetic abnormalities) fetuses. The aim of this review is to assess the clinical and cost-
17 effectiveness of progestogens in improving outcomes in women at risk of miscarriage.

18 Summary of the protocol

19 See Table 1 for a summary of the Population, Intervention, Comparison and Outcome
20 (PICO) characteristics of this review.

21 **Table 1: Summary of the protocol (PICO table)**

Population	Women at risk of miscarriage, for example: <ul style="list-style-type: none">• Women with threatened miscarriage (defined as early vaginal bleeding during the first trimester of pregnancy)• Women with a history of recurrent miscarriage (defined by the investigators).
Intervention	Progestogens: <ul style="list-style-type: none">• Oral dydrogesterone• Oral micronised progesterone• Vaginal micronised progesterone• Intramuscular 17-OH progesterone
Comparison	<ul style="list-style-type: none">• Any of the above progestogens compared to another• No treatment• Placebo
Outcome	Critical <ul style="list-style-type: none">• Live birth Important <ul style="list-style-type: none">• Miscarriage (birth before 24 weeks' gestation)• Preterm birth (birth before 37 weeks' gestation)• Stillbirth• Ectopic pregnancy• Congenital abnormalities• Adverse drug effects

1 *OH: hydroxy*

2 For further details see the review protocol in appendix A.

3 **Methods and process**

4 During the development of this guideline, a registered Cochrane protocol was identified
5 which matched the committee's intended protocol. The Cochrane review team completed
6 their review (Devall 2021) and presented their results to the committee which used them to
7 make recommendations. Cochrane's methods are closely aligned to standard NICE methods
8 and minor deviations (the use of the original Cochrane risk of bias tool and defining primary
9 and secondary outcomes as opposed to critical and important) relevant to the topic area
10 were highlighted to the committee and taken into account in discussions of the evidence.

11 See differences between the final Cochrane review (Devall 2021) and the Cochrane review
12 protocol (Devall 2020) in

13 <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>.

14 Declarations of interest were recorded according to [NICE's conflicts of interest policy](#).

15 **Effectiveness evidence**

16 **Included studies**

17 One Cochrane review (Devall 2021) including 7 randomised controlled trials (RCTs) (Chan
18 2020, Coomarasamy 2015, Coomarasamy 2019, Gerhard 1987, MacDonald 1972,
19 Shearman 1963, Siew 2018) was considered in this report. This review was used for
20 recommendation making by the committee as it was considered sufficiently relevant, high
21 quality and up to date.

22 The Cochrane review is summarised in Table 2 and the results of the review are provided in
23 the summary of the evidence in this report, however full details of the Cochrane review
24 including methods are available here:

25 <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

26 See the Cochrane review for the literature search strategy, study selection flowchart, forest
27 plots and GRADE summary of findings tables.

28 **Excluded studies**

29 See the Cochrane review for the list of excluded studies with reasons for their exclusions

30 **Summary of included studies**

31 Summaries of the studies that were included in this review are presented in Table 2.

1 **Table 2: Summary of included studies**

Study	Population	Comparisons	Outcomes
Devall 2021	Number of studies= 7 RCTs	Women with threatened miscarriage	<ul style="list-style-type: none"> • Live birth • Miscarriage (birth before 24 weeks gestational age) • Preterm birth (birth before 37 weeks gestational age) • Stillbirth • Ectopic pregnancy • Congenital abnormalities • Adverse drug effects
Systematic review	Number of women = 5682	<u>Vaginal micronised progesterone versus placebo</u> 2 RCTs, N=4090, UK, Germany, Coomarasany 2019, Gerhard 1987	
		<u>Dydrogesterone versus placebo</u> 1 RCT, N= 406, Hong Kong, Chan 2020	
		<u>Oral micronised progesterone versus dydrogesterone</u> 1 RCT, N=118, Singapore, Siew 2018	
		Women with recurrent miscarriage	
		<u>Vaginal micronised progesterone versus placebo</u> 1 RCT, N=826, UK, Coomarasany 2015	
		<u>Dydrogesterone versus placebo</u> 1 RCT, N=40, UK, MacDonald 1972	
		<u>17- hydroxyprogesterone versus placebo</u> 1 RCT, N=50, Australia, Shearman 1963	

2 *RCT(s): randomised controlled trial(s)*

3 See the Cochrane review for full evidence tables.

4

5 **Summary of the evidence**

6

7 Women with threatened miscarriage

8 Across all the comparisons identified in this Cochrane review, the majority showed no
 9 important difference for any outcomes between the interventions compared (for example,
 10 vaginal micronised progesterone versus placebo, dydrogesterone versus placebo, oral
 11 micronised progesterone versus dydrogesterone) for women with threatened miscarriage.

12 Evidence from pre-specified subgroups showed that there was no important benefit for
 13 vaginal micronised progesterone versus placebo in women with no previous miscarriages
 14 and early pregnancy bleeding in terms of the outcome of live birth. However, for this same

1 comparison, vaginal micronised progesterone showed an important benefit in terms of the
2 outcome of live birth for the subgroup of women with one or more previous miscarriages and
3 early pregnancy bleeding.

4 Typically, the outcomes for which no difference between interventions was found included
5 few studies and had seriously imprecise findings, therefore they should not be taken as
6 definitive evidence of no difference between the interventions. For the comparison vaginal
7 micronised progesterone versus placebo and dydrogesterone versus placebo, the outcome
8 miscarriage was precise and of high to moderate quality evidence, therefore this is indicative
9 that the true effect is similar to the estimated effect.

10 Women with recurrent miscarriage

11 Overall, vaginal micronised progesterone, dydrogesterone and 17-hydroxyprogesterone
12 appeared to have no important benefit over placebo for the outcomes assessed in women
13 with recurrent miscarriage.

14 The outcomes reported for the comparison vaginal micronised progesterone versus placebo
15 were generally of high quality, indicating that the true effect may be similar to the estimated
16 effect.

17 The outcomes reported for the comparison dydrogesterone and 17-hydroxyprogesterone
18 versus placebo were generally of low to very low quality evidence, indicating that the true
19 effect might be markedly different from the estimated effect.

20 Quality of the Cochrane review

21 Based on the methodological limitations assessed with the ROBIS tool to assess risk of bias
22 in systematic reviews and the modified PRISMA-NMA checklist, the Cochrane review was
23 deemed to be high quality. It was conducted following methodologically robust methods, and
24 a prospectively registered protocol. There were not important differences between the
25 included studies; all were conducted in the same setting and included women with similar
26 baseline characteristics in terms of age, gestational age and findings on ultrasonography.
27 The authors conducted pre-planned subgroup analyses for the factors that could have
28 interacted with treatment effects, such as age, number of previous miscarriages, regimen,
29 dosage and route of drug administration. As women with threatened miscarriage and women
30 with recurrent miscarriage are two very distinct populations, analyses were conducted
31 separately from the outset. There were no particular concerns for risk of bias in any of the
32 trials included, and the Cochrane review authors conducted a scientific
33 integrity/trustworthiness check, following the criteria agreed by the Cochrane Pregnancy and
34 Childbirth criteria, which provides an additional layer of quality assessment for the trials
35 included in the Cochrane review. Publication bias could not be fully ruled out, although it was
36 not possible to assess it formally in the Cochrane review.

37 The Cochrane review identified a gap in the evidence based. The authors had intended to
38 conduct a NMA, however interventions did not form a connected network, therefore only
39 indirect comparisons and pairwise analyses were performed. The authors defined in advance
40 the methods to conduct a NMA, which were considered to be appropriate.

41 For further details see the methodological limitations of the review in appendix L.

1 **Economic evidence**

2 **Included studies**

3 Two economic studies (reporting on results from the same randomised controlled trial) were
4 identified which were relevant to this question (Ogwulu 2019, Coomarasamy 2020).

5 As this review was based on a published Cochrane systematic review, no formal economic
6 literature search was undertaken but the included studies reported on an economic
7 evaluation that was undertaken alongside the PRISM RCT which was by far the biggest
8 study contributing to the clinical evidence in the Cochrane review.

9 **Summary of included economic evidence**

10 See

Table 3: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
Coomarasamy 2020 Progesterone to prevent miscarriage in women with early pregnancy bleeding: the PRISM RCT	Minor limitations	Directly applicable ¹	<p>Type of economic analysis: Cost-effectiveness analysis – economic evaluation conducted alongside a RCT</p> <p>Time horizon: < 1-year Trial duration</p> <p>Primary measure of outcome: Live births ≥ 34 weeks gestation</p>	£79 ²	0.022	£3,612 per live birth ²	<p>Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000 paired estimates of costs and outcomes suggested that there was a greater than 90% probability that progesterone was cost-effective for a cost-effectiveness threshold of £30,000 per live birth</p> <p>Deterministic sensitivity analysis was undertaken to assess:</p> <ul style="list-style-type: none"> • Fixed progesterone costs until a gestational age of 16 weeks

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
							<ul style="list-style-type: none"> • Imputation of missing primary care costs • Varying costs of inpatient night admissions • Varying cost of miscarriage management • Removing birth costs <p>The ICERs for these sensitivity analyses ranged from £3,282 per live birth to £4,977 per live birth</p>

¹ QALYs are not used as an outcome measure but the results can still be benchmarked using a cost per QALY approach as a live birth is likely to be valued much more highly than a QALY

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

Table 4: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac and at least 1 previous miscarriage

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
Ogwulu 2019 The cost-effectiveness of progesterone in preventing	Minor limitations	Directly applicable ¹	Type of economic analysis: Cost-effectiveness analysis – economic evaluation	-£337 ²	0.055	Progesterone dominant	Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
miscarriages in women with early pregnancy bleeding: an economic evaluation based on the PRISM trial			<p>conducted alongside a RCT</p> <p>Time horizon: < 1-year Trial duration</p> <p>Primary measure of outcome: Live births ≥ 34 weeks gestation</p>				paired estimates of costs and outcomes suggested that there was a greater than 90% probability that progesterone was cost-effective for a cost-effectiveness threshold of £20,000 per live birth

¹ QALYs are not used as an outcome measure but as the results showed dominance they can easily be interpreted for the purposes of decision making

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

Table 5: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac and 3 or more previous miscarriages

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
<p>Ogwulu 2019</p> <p>The cost-effectiveness of progesterone in preventing miscarriages in women with early pregnancy bleeding: an economic</p>	Minor limitations	Directly applicable ¹	<p>Type of economic analysis: Cost-effectiveness analysis – economic evaluation conducted alongside a RCT</p> <p>Time horizon: < 1-year</p>	£1,834 ²	0.15	£12,137 per live birth ²	Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000 paired estimates of costs and outcomes suggested that there was a greater than 90%

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
evaluation based on the PRISM trial			Trial duration Primary measure of outcome: Live births \geq 34 weeks gestation				probability that progesterone was cost-effective for a cost-effectiveness threshold of £20,000 per live birth

¹ QALYs are not used as an outcome measure but the results can still be benchmarked using a cost per QALY approach as a live birth is likely to be valued much more highly than a QALY

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

1 , Table 4 and Table 5 for the economic evidence profile of the included studies.
2

Table 3: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
Coomarasamy 2020 Progesterone to prevent miscarriage in women with early pregnancy bleeding: the PRISM RCT	Minor limitations	Directly applicable ¹	<p>Type of economic analysis: Cost-effectiveness analysis – economic evaluation conducted alongside a RCT</p> <p>Time horizon: < 1-year Trial duration</p> <p>Primary measure of outcome: Live births ≥ 34 weeks gestation</p>	£79 ²	0.022	£3,612 per live birth ²	<p>Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000 paired estimates of costs and outcomes suggested that there was a greater than 90% probability that progesterone was cost-effective for a cost-effectiveness threshold of £30,000 per live birth</p> <p>Deterministic sensitivity analysis was undertaken to assess:</p> <ul style="list-style-type: none"> • Fixed progesterone costs until a gestational age of 16 weeks

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
							<ul style="list-style-type: none"> • Imputation of missing primary care costs • Varying costs of inpatient night admissions • Varying cost of miscarriage management • Removing birth costs <p>The ICERs for these sensitivity analyses ranged from £3,282 per live birth to £4,977 per live birth</p>

¹ QALYs are not used as an outcome measure but the results can still be benchmarked using a cost per QALY approach as a live birth is likely to be valued much more highly than a QALY

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

Table 4: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac and at least 1 previous miscarriage

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
Ogwulu 2019 The cost-effectiveness of progesterone in preventing	Minor limitations	Directly applicable ¹	Type of economic analysis: Cost-effectiveness analysis – economic evaluation	-£337 ²	0.055	Progesterone dominant	Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
miscarriages in women with early pregnancy bleeding: an economic evaluation based on the PRISM trial			conducted alongside a RCT Time horizon: < 1-year Trial duration Primary measure of outcome: Live births ≥ 34 weeks gestation				paired estimates of costs and outcomes suggested that there was a greater than 90% probability that progesterone was cost-effective for a cost-effectiveness threshold of £20,000 per live birth

¹ QALYs are not used as an outcome measure but as the results showed dominance they can easily be interpreted for the purposes of decision making

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

Table 5: Economic evidence profile of progesterone for preventing miscarriage for women with bleeding in early pregnancy and ultrasound evidence of an intrauterine sac and 3 or more previous miscarriages

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
Ogwulu 2019 The cost-effectiveness of progesterone in preventing miscarriages in women with early pregnancy bleeding: an economic	Minor limitations	Directly applicable ¹	Type of economic analysis: Cost-effectiveness analysis – economic evaluation conducted alongside a RCT Time horizon: < 1-year	£1,834 ²	0.15	£12,137 per live birth ²	Probabilistic sensitivity analysis using non-parametric bootstrapping to generate 5000 paired estimates of costs and outcomes suggested that there was a greater than 90%

Study	Limitations	Applicability	Other comments	Incremental			Uncertainty
				Costs	Effect	Cost effectiveness	
evaluation based on the PRISM trial			Trial duration Primary measure of outcome: Live births ≥ 34 weeks gestation				probability that progesterone was cost-effective for a cost-effectiveness threshold of £20,000 per live birth

¹ QALYs are not used as an outcome measure but the results can still be benchmarked using a cost per QALY approach as a live birth is likely to be valued much more highly than a QALY

² Costs from a 2017-18 price year were updated for inflation to 2019/20 using an inflator of 1.05 derived from the hospital & community health services (HCHS) index and NHS Cost Inflation Index (NHSCII).

1 **Economic model**

2 No economic modelling was undertaken for this review because there was a recently
3 published UK economic evaluation alongside an RCT, which was considered
4 sufficiently high quality to support the committee in making recommendations.

5 **Evidence statements**

6 **Economic evidence statements**

7 One cost effectiveness analysis undertaken in the UK found that progesterone to
8 prevent miscarriage in early pregnancy had an incremental cost-effectiveness ratio of
9 £3,612 per live birth (price year 2019-20) relative to placebo. There was a greater
10 than 90% probability that progesterone was cost-effective at a cost-effectiveness
11 threshold of £30,000 per live birth. The economic analysis is directly applicable to the
12 NICE decision-making context, and is characterised by minor limitations.

13 One cost effectiveness analysis undertaken in the UK found that progesterone to
14 prevent miscarriage in early pregnancy in women with previous miscarriage
15 dominated placebo. There was a greater than 90% probability that progesterone was
16 cost-effective at a cost-effectiveness threshold of £20,000 per live birth. The
17 economic analysis is directly applicable to the NICE decision-making context, and is
18 characterised by minor limitations.

19 One cost effectiveness analysis undertaken in the UK found that progesterone to
20 prevent miscarriage in early pregnancy in women with 3 or more previous
21 miscarriages had an incremental cost-effectiveness ratio of £12,137 per live birth
22 (price year 2019-20) relative to placebo. There was a greater than 90% probability
23 that progesterone was cost-effective at a cost-effectiveness threshold of £20,000 per
24 live birth. The economic analysis is directly applicable to the NICE decision-making
25 context, and is characterised by minor limitations.

26 **The committee's discussion and interpretation of the evidence**

27 **The outcomes that matter most**

28 As the aim of this review was to determine if progestogens can prevent miscarriage,
29 the primary outcome was live birth, and the secondary outcomes were miscarriage
30 (defined as delivery before 24 weeks of gestation) and preterm birth (defined as birth
31 before 37 weeks of gestation). Other secondary outcomes to determine if the use of
32 progestogens in early pregnancy led to any adverse effects were stillbirth, ectopic
33 pregnancy, congenital abnormalities and adverse drug events.

34 **The quality of the evidence**

35 The quality of the evidence was high to very low, and it was typically downgraded
36 due to imprecision (small sample size and the 95% confidence intervals spanned
37 possible benefit and possible harm) and due to limitations in the study designs. The
38 methodological limitations of the Cochrane review were assessed using the ROBIS
39 tool to assess risk of bias in systematic reviews and the modified PRISMA-NMA
40 checklist. The overall quality of the Cochrane review was considered to be high (see
41 appendix L for further details).

1 The Cochrane authors reported that there were 13 studies awaiting classification
2 following assessment with their scientific integrity and trustworthiness checks (where
3 the Cochrane authors contact the primary study authors and request further details
4 about the study). The authors of the Cochrane review concluded that when all the
5 studies awaiting classification were included, the benefit of vaginal micronised
6 progesterone in the sub-group of women with early pregnancy bleeding and a
7 previous miscarriage remained (data not shown).

8 **Benefits and harms**

9 There was high quality evidence that in women with one or more previous
10 miscarriages and early pregnancy bleeding, 400 mg twice a day of vaginal
11 micronised progesterone increased the live birth rate compared to placebo, therefore
12 the committee based their recommendation on the use of progesterone in women
13 with threatened miscarriage on this evidence. The committee noted that there was no
14 evidence of harms for women or babies following the use of vaginal micronised
15 progesterone in this way, with no increase in risk of stillbirth, ectopic pregnancy,
16 congenital abnormalities or adverse drug reactions. However, they also noted that
17 the trials were not powered to detect very rare events and so the possibility of these
18 cannot be excluded.

19 The committee noted that the effect size observed in the subgroup of women with
20 one or more previous miscarriages and early pregnancy bleeding was larger than
21 that observed in the subgroup of women with no previous miscarriages and early
22 pregnancy bleeding. Therefore, the committee agreed that vaginal micronised
23 progesterone in women with no previous miscarriages and early pregnancy bleeding
24 would be of little or no benefit, however, according to this evidence, progesterone
25 would be beneficial in women with one of more previous miscarriages and early
26 pregnancy bleeding. This conclusion was further reinforced by the biological rationale
27 by which euploid miscarriages (that is, loss of chromosomally normal pregnancies)
28 are more likely to occur with increasing number of previous miscarriages. Euploid
29 miscarriages are associated with luteal phase defect, which is caused by low
30 progesterone levels and associated with vaginal bleeding, therefore it is biologically
31 plausible that the benefit of vaginal micronised progesterone is greater as the
32 number of previous miscarriages increases.

33 The committee noted that the evidence for the effectiveness of vaginal micronised
34 progesterone was in women with an ultrasound-confirmed intra-uterine pregnancy
35 and a confirmed fetal heartbeat. This was also the case for the population of women
36 included in the trial providing the highest weight (98.8%) to the meta-analysis for
37 vaginal micronised progesterone versus placebo reporting on the live birth outcome,
38 the Progesterone In Spontaneous Miscarriage (PRISM) trial (Coomarasamy 2019), in
39 which women had an intrauterine gestational sac visible on ultrasonography.

40 The committee discussed the relative merits of starting vaginal micronised
41 progesterone as early as possible (for example, after a positive home pregnancy
42 test) in women with a previous miscarriage and early pregnancy bleeding, compared
43 to starting after an ultrasound scan. The committee noted that there was no evidence
44 about the effectiveness and safety of progesterone started prior to ultrasound
45 confirmation of an intra-uterine pregnancy as no controlled trials had been conducted
46 to date on women with only a positive pregnancy test. Furthermore, there were also
47 concerns that recommending progesterone be commenced based only a pregnancy
48 confirmed using a pregnancy test, may lead to potential over-prescription of vaginal
49 micronised progesterone. As women in this early stage of their pregnancy would
50 usually present to their GP first, there would also be pressure on GPs to prescribe it
51 immediately, rather than referring women for an early pregnancy scan.

1 The committee discussed the fact that an intrauterine pregnancy can be detected on
2 an ultrasound scan as early as 4 weeks + 4 days of gestation, whereas the fetal
3 heartbeat may not be detected until about 5 weeks + 5 days of gestation. Therefore,
4 to avoid any potential delays in the commencement of vaginal micronised
5 progesterone in women with early pregnancy bleeding, the committee agreed to
6 recommend that it should be started once there is confirmation of an intra-uterine
7 pregnancy, and then if the fetal heartbeat is seen, the progesterone can be continued
8 up to 16 weeks. The committee noted that if heartbeat is not seen on an initial scan,
9 women are always offered a repeat scan in a maximum timeframe of 2 weeks.

10 The committee discussed that the recommendation to use vaginal micronised
11 progesterone would not be generalizable to women with a pregnancy of unknown
12 location or ectopic pregnancy as the evidence had excluded these pregnancies, but
13 that the use of ultrasound confirmation of pregnancy would enable these women to
14 be identified before the progesterone was prescribed.

15 The committee noted that threatened miscarriage can be very stressful and upsetting
16 regardless of the prognosis and type of pregnancy the woman has, and emphasised
17 that healthcare professionals providing care for these women will need to provide
18 accurate information about what to expect, as well as providing details of support and
19 advice available from relevant organisations. However, they agreed it was not
20 necessary to make a separate recommendation about this as it is already covered in
21 the support and information giving section of the guideline.

22 The committee discussed that the risk of miscarriage increases with maternal age
23 and whether the age of the woman could have a role on the effectiveness of vaginal
24 micronised progesterone, however no age effect was observed when data was split
25 between those above and below 35 years on prespecified subgroup analysis.

26 The committee noted that there are 4 types of progesterone preparations available in
27 the UK: micronised oral progesterone, micronised vaginal progesterone,
28 progesterone vaginal tablets and pessaries, and a progesterone vaginal gel. These
29 are indicated in women for supplementation of the luteal phase, for luteal phase
30 support and pre-menstrual syndrome or postnatal depression but are not approved
31 for the prevention of miscarriage.

32 The authors of the PRISM trial emphasised that the conclusions obtained about the
33 effectiveness of 400 mg twice a day of vaginal micronised progesterone cannot be
34 extrapolated to other progesterone preparations, doses or routes and it cannot be
35 assumed that non-micronised preparations would behave in the same way as
36 micronised progesterone as this preparation has an identical molecular structure to
37 natural progesterone. The committee also noted that the 2 main trials contributing to
38 the evidence (PRISM and Gerhard 1987) both used vaginal micronised
39 progesterone. Therefore, although vaginal micronised progesterone is not licenced in
40 the UK for the prevention of miscarriage, the committee agreed that it was necessary
41 to be specific about this formulation in their recommendations, and they also adopted
42 the dose used in the PRISM study. The review found no evidence of harm for the
43 prespecified outcomes, namely stillbirth, ectopic pregnancy, congenital abnormalities
44 or adverse drug reactions, however this should not be taken as definitive evidence as
45 the trials were not powered to detect very rare events.

46 The committee noted that progesterone pessaries (not micronised) are commonly
47 used in clinical practice for the prevention of miscarriage and were aware that there
48 is no available evidence to support their effectiveness, or evidence for any of the
49 other available preparations. As there was so little evidence for other progesterone
50 preparations the committee made a research recommendation to determine the

1 comparative effectiveness of other progesterone preparations for the prevention of
2 miscarriage.

3 The committee also discussed the duration of therapy and noted that the PRISM
4 study had continued therapy until 16 completed weeks of pregnancy. This decision to
5 use this duration in the PRISM study had been based on a consensus of a national
6 body of UK clinicians, and reflected that the progesterone was being used to treat an
7 assumed deficit in the pregnant woman's own endogenous progesterone production.
8 The committee discussed that in a normal pregnancy, the predominant source of
9 progesterone in the first 12 weeks of pregnancy is the corpus luteum and after that
10 the placenta is the main source of progesterone. So, if progesterone therapy was
11 intended to correct only a corpus luteal defect, then treatment for the first 12 weeks
12 of gestation would have been sufficient, but as placental production of progesterone
13 may also be sub-optimal it was felt necessary to continue the treatment to 16 weeks.

14 The PRISM trial is a well-powered multi-center randomised placebo-controlled trial,
15 conducted with methodological rigour, following a prespecified protocol. However, it
16 is worth noting that the primary analysis in the trial did not find any important
17 difference between groups, and that the only clinically important finding was found in
18 a prespecified subgroup analyses, which was not adjusted for multiplicity. Therefore,
19 this may raise concerns about a possible false positive result. Given the biological
20 rationale presented above, the small but important effect size the PRISM trial
21 showed, and the critical evaluation following guidelines for the interpretation of
22 subgroup analysis with which the trial has been assessed (Coomarasamy 2020), the
23 committee agreed that the evidence was a robust enough base for their
24 recommendation.

25 The committee discussed the evidence for women with unexplained recurrent
26 miscarriage (but no early pregnancy bleeding). Three studies had been included in
27 the Cochrane analysis for this population of women using three different
28 progestogens. The PROgesterone in recurrent MIScarriagE (PROMISE) trial
29 (Coomarasamy 2015) used vaginal micronised progesterone and is the largest
30 published in the field. However, it did not result in a significantly higher rate of live
31 births. The other studies used dydrogesterone and 17OH-progesterone and provided
32 little or no evidence for the outcome of live birth. The committee agreed that further
33 research is needed to assess the role of progestogens in women with unexplained
34 recurrent miscarriage, therefore the committee made a research recommendation to
35 this effect.

36 **Cost effectiveness and resource use**

37 An economic evaluation of the PRISM trial was reported in 2 papers (Ogwulu 2019,
38 Coomarasamy 2020). The primary analysis was in a population of women with early
39 pregnancy bleeding and an ultrasound confirmed intrauterine sac. The evaluation
40 assessed the cost-effectiveness of progesterone compared with placebo in
41 preventing pregnancy loss.

42 As part of the trial, maternal and neonatal hospital resource use was recorded from
43 randomisation to hospital discharge for each pregnancy. Health services self-
44 completed questionnaires were used to measure other resource use provided in
45 primary care and the community during the same period. The adjusted mean
46 difference in costs of £76 was not statistically significant with the intervention cost of
47 £204 partly offset by a non-significant reduction in hospital costs in the intervention
48 group. The intervention resulted in 2.2 (95% CI -0.4% to 5.0%; p = 0.08) additional
49 live births at ≥ 34 weeks' gestational age per 100 pregnancies. The incremental cost-
50 effectiveness ratio (ICER) was £3,305 per live birth. If a live birth was assumed to be

1 equivalent to approximately 25 discounted Quality Adjusted Life Years (QALYs) then
2 the ICER would be around £138 per QALY, which the committee noted would easily
3 fall within a £20,000 per QALY cost-effectiveness threshold often used to inform
4 recommendations for NICE guidelines. Probabilistic sensitivity analysis based on
5 5,000 simulations using non-parametric bootstrapping found that there was a greater
6 than 90% probability that progesterone would be considered cost-effective at a cost-
7 effectiveness threshold of £30,000 per live birth.

8 However, the committee noted that the analysis was based on a population that
9 included some women with a history of previous miscarriage for whom the evidence
10 of treatment benefit was clearer and therefore they did not consider there was
11 sufficient evidence to recommend progesterone for women without a history of
12 previous miscarriage.

13 A sub-group analysis in the PRISM trial suggested that progesterone was dominant
14 in women with a history of previous miscarriage, leading to a cost saving of £322 and
15 5.5 more live births per 100 pregnancies. Probabilistic sensitivity analysis indicated
16 that there was more than a 90% probability that this would be cost-effective at a cost-
17 effectiveness threshold of £20,000 per live birth. Therefore, the committee agreed
18 there was strong evidence to support a recommendation for progesterone to prevent
19 miscarriage in women with early pregnancy bleeding who have a history of previous
20 miscarriage.

21 A further sub-group analysis considered progesterone in the smaller sub-group of
22 women who had a history of 3 or more previous miscarriages. In this analysis
23 progesterone led to higher costs although, as in the other analyses, the difference
24 was not statistically significant and, reflecting the smaller sample, the confidence
25 intervals were much wider than for the other analyses. However, the committee
26 noted that the authors tentatively suggested that the higher costs of the progesterone
27 intervention in this group could be a result of its success with greater neonatal
28 intensive care costs because of averted miscarriage driving the increase in costs.
29 Despite the higher costs probabilistic sensitivity analysis indicated that progesterone
30 was likely to be cost-effective. However, the committee did not consider that the
31 results of this sub-group analysis warranted any additional recommendations for
32 women with a history of 3 or more miscarriages.

33 The committee noted that recommending progesterone for women with a history of
34 previous miscarriage would represent a change in practice for the NHS. However,
35 they did not consider that this change would have a significant resource impact.
36 Although the effect is likely to be small, successfully averted miscarriages are likely
37 to result in a reduction in subsequent conceptions, which would offset treatment
38 costs to some extent. Furthermore, although subject to considerable uncertainty as to
39 the precise effect, the subgroup analysis in women with a history of previous
40 miscarriage had suggested that the intervention may generate cost savings.

41 **Recommendations supported by this evidence review**

42 This evidence review supports recommendation 1.5.2 and the research
43 recommendations on the use of progesterone in recurrent miscarriage and the
44 effectiveness of other progesterone preparations at preventing miscarriage.

1 **References**

2 **Effectiveness**

3 **Chan 2020**

4 Chan DM, Cheung KW, Ko JK, Yung SS, Lai SF, Lam MT, Ng DY, Lee VC, Li RH,
5 Ng EH. Use of oral progestogen in women with threatened miscarriage in the first
6 trimester: a randomized double-blind controlled trial. *Human Reproduction*. 2021
7 Mar;36(3):587-95.

8 **Coomarasamy 2015**

9 Coomarasamy A, Williams H, Truchanowicz E, Seed PT, Small R, Quenby S, Gupta
10 P, Dawood F, Koot YE, Bender Atik R, Bloemenkamp KW. A randomized trial of
11 progesterone in women with recurrent miscarriages. *New England Journal of*
12 *Medicine*. 2015 Nov 26;373(22):2141-8.

13 **Coomarasamy 2019**

14 Coomarasamy A, Devall AJ, Cheed V, Harb H, Middleton LJ, Gallos ID, Williams H,
15 Eapen AK, Roberts T, Ogwulu CC, Goranitis I. A randomized trial of progesterone in
16 women with bleeding in early pregnancy. *New England Journal of Medicine*. 2019
17 May 9;380(19):1815-24.

18 **Devall 2020**

19 Devall AJ, Papadopoulou A, Haas DM, Price MJ, Coomarasamy A, Gallos ID.
20 Progestogens for preventing miscarriage: a network meta-analysis (Protocol).
21 *Cochrane Database of Systematic Reviews* 2020 (11).

22 **Devall 2021**

23 Devall AJ, Papadopoulou A, Podeseck M, Haas DM, Price MJ, Coomarasamy A,
24 Gallos ID. Progestogens for preventing miscarriage: a network meta-analysis.
25 *Cochrane Database of Systematic Reviews*. 2021(4).

26 **Gerhard 1987**

27 Gerhard I, Gwinner B, Eggert-Kruse W, Runnebaum B. Double-blind controlled trial
28 of progesterone substitution in threatened abortion. *Biological research in pregnancy*
29 *and perinatology*. 1987 Jan 1;8(1 1ST Half):26-34.

30 **MacDonald 1972**

31 MacDonald RR, Goulden R, Oakey RE. Cervical mucus, vaginal cytology and steroid
32 excretion in recurrent abortion. *Obstetrics & Gynecology*. 1972 Sep 1;40(3):394-402.

33 **Shearman 1963**

34 Shearman RP, Garrett WJ. Double-blind study of effect of 17-hydroxyprogesterone
35 caproate on abortion rate. *British medical journal*. 1963 Feb 2;1(5326):292.

36 **Slew 2018**

37 Siew JY, Allen JC, Hui CY, Ku CW, Malhotra R, Østbye T, Tan TC. The randomised
38 controlled trial of micronised progesterone and dydrogesterone (TRoMaD) for
39 threatened miscarriage. *European Journal of Obstetrics & Gynecology and*
40 *Reproductive Biology*. 2018 Sep 1;228:319-24.

1 **Economic**

2 **Coomarasamy 2020**

3 Coomarasamy A, Harb HM, Devall AJ, Cheed V, Roberts TE, Goranitis I, et al.
4 Progesterone to prevent miscarriage in women with early pregnancy bleeding: the
5 PRISM RCT. *Health Technol Assess* 2020;24(33).

6 **Ogwulu 2019**

7 Okeke Ogwulu CB, Goranitis I, Devall AJ, Cheed V, Gallos ID, Middleton LJ, Harb
8 HM, Williams HM, Eapen A, Daniels JP, Ahmed A, Bender-Atik R, Bhatia K,
9 Bottomley C, Brewin J, Choudhary M, Deb S, Duncan WC, Ewer AK, Hinshaw K,
10 Holland T, Izzat F, Johns J, Lumsden M, Manda P, Norman JE, Nunes N, Overton
11 CE, Kriedt K, Quenby S, Rao S, Ross J, Shahid A, Underwood M, Vaithilingham N,
12 Watkins L, Wykes C, Horne AW, Jurkovic D, Coomarasamy A, Roberts TE. The cost-
13 effectiveness of progesterone in preventing miscarriages in women with
14 early pregnancy bleeding: an economic evaluation based on the PRISM trial. *BJOG*
15 2020;127:757-767

16 **Other**

17 **Coomarasamy 2020**

18 Coomarasamy A, Devall AJ, Brosens JJ, Quenby S, Stephenson MD, Sierra S,
19 Christiansen OB, Small R, Brewin J, Roberts TE, Dhillon-Smith R. Micronized vaginal
20 progesterone to prevent miscarriage: a critical evaluation of randomized evidence.
21 *American journal of obstetrics and gynecology*. 2020 Jan 31.

22

1 **Appendices**

2 **Appendix A Review protocols**

3 **Review protocol for review question: What is the clinical and cost-**
4 **effectiveness of progestogens in improving outcomes in women at risk**
5 **of miscarriage?**

6 See Cochrane review

7 <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792/epdf/full>

Appendix B Literature search strategies

Literature search strategies for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix C Effectiveness

Study selection for: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix D Evidence tables

Evidence tables for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix E Forest plots

Forest plots for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix F GRADE tables

GRADE tables for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix G Economic evidence study selection

Study selection for: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

See NIHR technology appraisal and associated article in the British Journal of Obstetrics and Gynaecology (BJOG)

<https://www.journalslibrary.nihr.ac.uk/hta/hta24330#/abstract>

<https://obgyn.onlinelibrary.wiley.com/doi/epdf/10.1111/1471-0528.16068>

Appendix H Economic evidence tables

Economic evidence tables for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

Table 6: Economic evidence tables

Study country and type	Intervention and comparator	Study population, design and data sources	Costs and outcomes (descriptions and values)	Results	Comments
Coomarasamy 2020 UK Ogwulu 2019 UK Study type: Cost-effectiveness analysis Source of funding: UK National Institute for Health Research (NIHR) Health Technology Assessment programme	Intervention: 400 mg of progesterone (Twice daily vaginal 200 mg pessaries for the duration of treatment – treatment began as soon as possible after confirmation of an intrauterine pregnancy sac and within 4 days of vaginal bleeding and continued until a gestational age of 16 weeks Comparator: Placebo pessaries :	Women with bleeding in early pregnancy bleeding, with an intrauterine gestation sac visible on ultrasonography, aged 16–39 years at randomisation Modelling approach: Economic evaluation alongside PRISM RCT Source of baseline data: Control group in RCT Source of effectiveness data: RCT Source of resource use data:	Primary analysis Mean cost per participant: Intervention: £7,655 Control: £7,572 Difference: £76 Primary measure of outcome: Live birth at ≥ 34 weeks gestation Mean outcome per participant Intervention: 0.0747	Primary analysis ICERs: £3,305 per live birth Probability of being cost effective: >80% probability of being cost-effective as a cost-effectiveness threshold of £15,000 per live birth >90% probability of being cost-effective as a cost-effectiveness threshold of £30,000 per live birth Sensitivity analysis:	Currency: GBP Cost year: 2017-18 Time horizon: Duration of trial Discounting: None (N/A) Applicability: Directly applicable Limitations: Minor limitations Other comments: The results were presented in 2 papers

Study country and type	Intervention and comparator	Study population, design and data sources	Costs and outcomes (descriptions and values)	Results	Comments
		<p>RCT – data on hospital resource use from randomisation to discharge was collected using researcher completed data forms. Data on services provided in the community was collected retrospectively using health services self-completed questionnaires.</p> <p>Source of unit cost data:</p> <ul style="list-style-type: none"> • NHS Reference Costs 2016/17 • PSSRU • BNF 	<p>Control: 0.0725</p> <p>Difference: 0.022</p> <p>Sub-group analysis (Women with at least 1 previous miscarriage)</p> <p>Mean cost per participant</p> <p>Difference: -£322</p>	<p>Fixed progesterone costs until 16 weeks: £4,977 per live birth</p> <p>Imputation of primary care costs: £4,321 per live birth</p> <p>Varying costs of inpatient night admissions: £3,356 per live birth</p> <p>Varying costs of miscarriage management: £3,282 per live birth</p> <p>Removing birth costs: £3,743 per live birth</p> <p>Sub-group analysis (Women with at least 1 previous miscarriage)</p> <p>Progesterone dominates</p> <p>>90% probability of being cost-effective as a cost-effectiveness</p>	<p>but were based on the same PRISM RCT.</p> <p>Costs difference are adjusted means based on bootstrapped difference</p>

Study country and type	Intervention and comparator	Study population, design and data sources	Costs and outcomes (descriptions and values)	Results	Comments
			<p>Mean outcome per participant</p> <p>Difference: 0.055</p> <p><u>Sub-group analysis (Women with at 3 or more previous miscarriages)</u></p> <p>Mean cost per participant</p> <p>Difference: £1,754</p> <p>Mean outcome per participant</p> <p>Difference: 0.15</p>	<p>threshold of £20,000 per live birth</p> <p><u>Sub-group analysis (Women with at 3 or more previous miscarriages)</u></p> <p>£11,606 per live birth</p> <p>>90% probability of being cost-effective as a cost-effectiveness threshold of £20,000 per live birth</p>	

Study country and type	Intervention and comparator	Study population, design and data sources	Costs and outcomes (descriptions and values)	Results	Comments

BNF = British National Formulary; ICER = incremental cost-effectiveness ratio; PSSRU = Personal Social Services Research Unit; RCT = randomised controlled trial

Appendix I Economic model

Economic model for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

No economic analysis was conducted for this review question.

Appendix J Excluded studies

Excluded studies for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

Excluded effectiveness studies

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Appendix K Research recommendations – full details

Research recommendations for review question: What is the clinical and cost-effectiveness of progestogens in improving outcomes in women at risk of miscarriage?

K.1.1 Research recommendation

What is the clinical and cost-effectiveness of progesterone in improving outcomes in women with unexplained recurrent miscarriage?

K.1.2 Why this is important

Women with previous pregnancy losses have an increased risk of miscarriage in subsequent pregnancies. Progesterone is essential for maintaining a healthy pregnancy, and there is evidence that it is safe for both women and fetuses.

A recent randomised controlled trial assessed the effectiveness of micronised vaginal progesterone supplementation in women with 3 or more first-trimester losses and did not show a benefit with progesterone therapy use during the first trimester, concluding that there is not enough evidence to support its use in women with unexplained recurrent miscarriage. However, this trial was designed to look for a 10% difference in live birth outcomes in those who received progesterone versus those who did not receive it. A larger randomised controlled trial is required to determine if there is a smaller difference (for example 2.5% to 5%) which would still lead to a meaningful increase live births and reduce the trauma of a further miscarriage for a number of women.

K.1.3 Rationale for research recommendation

Table 7: Research recommendation rationale

Importance to 'patients' or the population	Miscarriage can have significant long-lasting negative effects on women, their partners and families. Progestogen supplementation may increase the opportunity to maintain a healthy pregnancy in some women.
Relevance to NICE guidance	The committee were unable to make recommendations on women with unexplained recurrent miscarriage as current evidence base does not show a benefit with progesterone use. A high-quality clinical trial with a large number of participants is needed to study small effect size. This would allow recommendations to be made in future guideline updates.
Relevance to the NHS	The management of miscarriage and its complications poses a significant burden to the NHS. Ensuring women at high risk of miscarriage are able to maintain a healthy pregnancy will lead to better outcomes for women and support the NHS goal to improve the care of women with unexplained recurrent miscarriage.
National priorities	High
Current evidence base	A number of small trials which have found a benefit and the PROMISE study which did not find a benefit.
Equality considerations	None known

PROMISE: PROgesterone in recurrent MIScarriageE

K.1.4 Modified PICO table

Table 8: Research recommendation modified PICO table

Population	Women with 3 or more unexplained previous miscarriages
Intervention	Progestogens: <ul style="list-style-type: none"> • Dydrogesterone • Micronised oral progesterone • Micronised vaginal progesterone • Non-micronised vaginal progesterone • 17-OH progesterone
Comparator	<ul style="list-style-type: none"> • Any of the above progestogens compared to another • No treatment • Placebo
Outcome	<ul style="list-style-type: none"> • Live birth • Miscarriage (birth before 24 weeks' gestation) • Preterm birth (birth before 37 weeks' gestation) • Stillbirth • Ectopic pregnancy • Congenital abnormalities • Adverse drug effects
Study design	Randomised, double-blind placebo controlled trial
Timeframe	36 months
Additional information	None

OH: hydroxy

K.1.5 Research recommendation

What is the clinical and cost-effectiveness of vaginal micronised progesterone versus other progesterone preparations in improving outcomes in women at risk of miscarriage?

K.1.6 Why this is important

Evidence from a recent randomised controlled trial showed a small but important benefit for the outcome of live birth when vaginal micronised progesterone was given to women with early pregnancy bleeding and a history of one or more previous miscarriages. However, there was not enough evidence available to assess whether other formulations of progesterone would lead to other beneficial outcomes in this group of women. Research is needed to identify whether there is a difference in the effectiveness of micronised versus non-micronised progesterone therapy in women with early pregnancy bleeding and a history of one or more previous miscarriages.

K.1.7 Rationale for research recommendation

Table 9: Research recommendation rationale

Importance to 'patients' or the population	Miscarriage can have a significant emotional and physical impact on women, their partners and families. The specific type of progesterone used may have a role in the attainment and maintenance of a healthy pregnancy.
---	--

Relevance to NICE guidance	A high-quality clinical trial will allow recommendations for a range of progestogen preparations to be made in future guideline updates.
Relevance to the NHS	The management of miscarriage and its complications poses a significant burden to the NHS. Ensuring women at high risk of miscarriage are able to maintain a healthy pregnancy will lead to better outcomes for women and support the NHS goal to improve the care of women with threatened miscarriage.
National priorities	High
Current evidence base	Minimal
Equality considerations	None known

K.1.8 Modified PICO table

Table 10: Research recommendation modified PICO table

Population	Women with early pregnancy bleeding and a history of one or more previous miscarriages
Intervention	<ul style="list-style-type: none"> • Vaginal micronised progesterone
Comparator	<ul style="list-style-type: none"> • Vaginal non-micronised progesterone preparations • Oral/subcutaneous/rectal progesterone preparations
Outcome	<ul style="list-style-type: none"> • Live birth • Miscarriage (birth before 24 weeks' gestation) • Preterm birth (birth before 37 weeks' gestation) • Stillbirth • Ectopic pregnancy • Congenital abnormalities • Adverse drug effects
Study design	Randomised, double-blind placebo controlled trial
Timeframe	36 months
Additional information	None

Appendix L Methodological limitations

The methodological limitations of the Cochrane review (Devall 2021) have been assessed using the ROBIS tool to assess risk of bias in systematic reviews and the modified PRISMA-NMA checklist

L.1.1 ROBIS tool to assess risk of bias in systematic reviews

Domain 1: Study eligibility criteria

- 1.1 Did the review adhere to pre-defined objectives and eligibility criteria? Y
 - 1.2 Were the eligibility criteria appropriate for the review question? Y
 - 1.3 Were eligibility criteria unambiguous? Y
 - 1.4 Were any restrictions in eligibility criteria based on study characteristics appropriate (for example, date, sample size, study quality, outcomes measured)? N
 - 1.5 Were any restrictions in eligibility criteria based on sources of information appropriate (for example, publication status or format, language, availability of data)? Y
- Concerns regarding specification of study eligibility criteria: LOW

Domain 2: Identification and selection of studies

- 2.1 Did the search include an appropriate range of databased/electronic sources for published and unpublished reports? Y
 - 2.2 Were methods additional to database searching used to identify relevant reports? Y
 - 2.3 Were the terms and structure of the search strategy likely to retrieve as many eligible studies as possible? Y
 - 2.4 Were restrictions based on date, publication format, or language appropriate? Y
 - 2.5 Were efforts made to minimise error in selection of studies? Y
- Concerns regarding methods used to identify and/or select studies: LOW

Domain 3: Data collection and study appraisal

- 3.1 Were efforts made to minimise error in data collection? Y
 - 3.2 Were sufficient study characteristics available for both review authors and reader to be able to interpret the results? Y
 - 3.3 Were all relevant study results collected for use in the synthesis? Y
 - 3.4 Was risk of bias (or methodological quality) formally assessed using appropriate criteria? Y
 - 3.5 Were efforts made to minimise error in risk of bias assessment? Y
- Concerns regarding methods used to collect data or appraise studies: LOW

Domain 4: synthesis and findings

- 4.1 Did the synthesis include all studies that it should? Y
 - 4.2 Were all pre-defined analyses reported or departures explained? Y
 - 4.3 Was the synthesis appropriate given the nature and similarity in the research questions, study designs and outcomes across included studies? Y
 - 4.4 Was between-study variation (heterogeneity) minimal or addressed in the synthesis? Y
 - 4.5 Were the findings robust, for example, as demonstrated through funnel plot or sensitivity analyses? Y
 - 4.6 Were biases in primary studies minimal or addressed in the synthesis? Y
- Concerns regarding the synthesis and findings: LOW

Risk of bias in the review

- A. Did the interpretation of findings address all of the concerns identified in Domains 1 to 4?
 - B. Was the relevance of identified studies to the review's research question appropriately considered?
 - C. Did the reviewers avoid emphasising results on the basis of statistical significance? Y
- Risk of bias in the review: LOW

L.1.2 Modified PRISMA-NMA checklist

1. Describe the reasons for the review in the context of what is already known, including mention of why a network meta-analysis has been conducted

Not applicable

2. Specify study characteristics (for example, PICOS, length of follow-up) and report characteristics (for example, years considered, language, publication status) used as criteria for eligibility, giving rationale. Clearly describe eligible treatments included in the treatment network, and note whether any have been clustered or merged into the same node (with justification).

Not applicable

3. Describe methods used to explore the geometry of the treatment network and potential biases related to it. This should include how the evidence base has been graphically summarised for presentation, and what characteristics were compiled and used to describe the evidence base to readers

Authors included a network diagram for each outcome and a qualitative description of the treatment network, which represented the available comparisons, number of trials, number of participants included, and the grading of the direct evidence.

The size of the nodes and the limited number of edges indicated a gap in the evidence base and the graphical representation of all comparisons showed that a NMA was not feasible (because the interventions did not form a connected network).

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

Quantitative metrics assessing features of the network geometry were not mentioned.

4. State the principal summary measures (for example, risk ratio, difference in means). Also describe the use of additional summary measures assessed, such as treatment rankings and surface under the cumulative ranking curve (SUCRA) values, as well as modified approaches used to present summary findings from meta-analyses

The principal summary measure of relative treatment effect of dichotomous outcomes for direct comparisons were risk ratios (RRs). The authors planned to rank and obtain a hierarchy of the treatments according to their SUCRA values and to evaluate each outcome to determine the confidence of the NMA output. The treatment hierarchy was going to be presented along with summary effect measures and 95% CIs, however, it was not possible to do this due to the paucity of evidence. As a result, modified approaches to represent the summary findings, such as additional graphical or tabular approaches, were not included.

5. Describe the methods of handling data and combining results of studies for each network meta-analysis. This should include, but not be limited to: a) Handling of multi-

arm trials; b) Selection of variance structure; c) Selection of prior distributions in Bayesian analyses; d) Assessment of model fit.

Methods for handling data:

- a) Cluster randomised trials were planned to be included and to use appropriate methods for adjusting sample size, combining results with individually randomised trials and exploring heterogeneity. However, no cluster randomised trials were included in the review.
- b) Multi-arm trials were planned to be included and treated as multiple independent comparisons. For global analyses, authors planned to combine different dosage, regimen and route of drug administration and to explore these in separate subgroup analyses. However, no multi-arm trials were included in the review.
- c) Missing data: authors planned to note levels of attrition and to explore high levels of missing data in sensitivity analyses. Intention-to-treat (ITT) analysis were prioritised.
- d) Statistical heterogeneity: this was planned to be estimated from the multivariate meta-analysis model, based on the magnitude of the heterogeneity variance parameter (T^2) and compared to empirical distributions for dichotomous variables.
- e) Clinical homogeneity was explored by describing the study population characteristics across trials. The appropriateness of the transitivity assumption was met for both direct and indirect comparisons.
- f) Inconsistency: this was planned to be done following local (such as node-splitting) and global (such as 'design by treatment' interaction model) approaches, along with the evaluation of clinical homogeneity described above.

Methods for combining results:

- g) Indirect treatment comparisons: data was extracted and set up using the augmented format by comparing all treatments with a reference treatment, and those studies without a treatment have one created. Network diagrams were created to verify if a NMA was feasible. NMAs would have been performed under a frequentist framework using multivariate random effect meta-analyses estimated using maximum likelihood estimation. NMAs were planned to be carried using Stata® statistical software. Indirect comparisons were produced where possible and estimated using Microsoft Excel®.

6. Describe the statistical methods used to evaluate the agreement of direct and indirect evidence in the treatment network(s) studied. Describe efforts taken to address inconsistency when found.

As a NMA could not be conducted, the authors did not evaluate the agreement of direct and indirect evidence in the treatment networks, however they noted in their protocol that their final step for assessing the quality of the NMA would include the consideration of coherence between direct and indirect effect estimates.

7. Describe methods of additional analyses if done, indicating which were pre-specified. This may include, but not be limited to, the following: e) Sensitivity or subgroup analyses; f) Meta-regression analyses; g) Alternative formulations of the treatment network; h) Use of alternative prior distributions for Bayesian analyses (if applicable).

As a NMA could not be conducted, the authors did not describe additional methods of analyses. However, in the protocol they describe that sensitivity analyses were

planned by evaluating model fit for: overall quality of the studies, randomisation unit and use of placebo.

No additional methods of additional analyses were described.

8. Provide a network graph of the included studies to enable visualisation of the geometry of the treatment network.

See Cochrane review

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013792.pub2/epdf/full>

9. Provide a brief overview of characteristics of the treatment network. This may include commentary on the abundance of trials and randomised patients for the different interventions and pairwise comparisons in the network, gaps of evidence in the treatment network, and potential biases reflected by the network structure (for example, publication bias).

Due to the paucity of evidence, the network of interventions was not connected and it did not contain closed loops. For this same reason, it was not possible to assess bias formally.

10. Present results of each meta-analysis done, including confidence/credible intervals. In larger networks, authors may focus on comparisons versus a particular comparator (for example, placebo or standard care). League tables and forest plots may be considered to summarise pairwise comparisons. If additional summary measures were explored (such as treatment rankings), these should also be presented.

Not applicable as NMA was not feasible.

11. Describe results from investigations of inconsistency. This may include such information as measures of model fit to compare consistency and inconsistency models, P values from statistical tests, or summary of inconsistency estimates from different parts of the treatment network.

As a NMA could not be conducted, the authors did not investigate inconsistency. However, in the protocol, they described that sensitivity analyses were planned by evaluating model fit for overall quality of the studies, randomisation unit and use of placebo.

12. Give results of additional analyses, if done (for example, sensitivity or subgroup analyses, meta-regression analyses, alternative network geometries studied, alternative choice of prior distributions for Bayesian analyses, and so forth).

Apart from the sensitivity analyses outlined above, no additional methods of additional analyses were described.

13. Discuss limitations at study and outcome level (for example, risk of bias), and at review level (for example, incomplete retrieval of identified research, reporting bias). Comment on the validity of the assumptions, such as transitivity and consistency. Comment on any concerns regarding network geometry (for example, avoidance of certain comparisons).

All 7 studies were judged to be at low risk of bias. In addition to the risk of bias assessment per study, authors also conducted a scientific integrity/trustworthiness check, which ensures that only studies deemed to be sufficiently trustworthy are included in the analysis. Publication bias could not be assessed formally due to the paucity of evidence, however if this was investigated informally it was not reported.

The results of the analysis may have been impacted by publication bias, for example where results had not been published because results were unfavourable to progestogen treatment. The source of funding was not reported for 2 of the studies and the remaining studies were funded by public bodies. As non-publication is more common in trials that received industry funding, publication bias cannot be completely ruled out, although the Cochrane review authors did attempt to minimise it by identifying all relevant studies regardless of language and publication status. This approach also minimises the risk of incomplete retrieval of identified research. Some of the Cochrane review authors were also authors of the largest trial including women with threatened miscarriage, however they did not participate in any decisions regarding the assessment for inclusion/exclusion of trials, risk of bias assessment or data extraction, and there were not conflicts of interest reported, so it is not anticipated that this would have introduced bias in the reviewing process.

The assumption of transitivity was considered to be valid as the different progesterone preparations are administered in a similar way across trials. Trials were also similar in terms of design and characteristics. The treatment network showed that there were not enough trials available to conduct a NMA, indicating potential gaps of evidence. For example, most trials compared an active treatment with placebo, but there was only 1 study providing evidence for a type of progesterone compared to another. Indirect treatments were possible, but network interventions were not connected, limiting the ability to compare multiple treatments simultaneously in a single analysis. For this reason, the committee made a research recommendation to assess the clinical and cost-effectiveness of vaginal micronised progesterone versus other progesterone preparations in women at risk of miscarriage.

Overall, no significant limitations were identified, and it was considered that the statistical methods planned for the NMA were appropriate.