

Antenatal care

[W] Maternal sleep position during pregnancy

NICE guideline NG201

Evidence reviews underpinning recommendations 1.3.24 to 1.3.25

August 2021

Final

These evidence reviews were developed by the National Guideline Alliance, which is a part of the Royal College of Obstetricians and Gynaecologists

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2021. All rights reserved. Subject to [Notice of Rights](#).

ISBN: 978-1-4731-4227-5

Contents

Contents	4
Maternal sleep position during pregnancy	6
Review question	6
Introduction	6
Summary of the protocol	6
Methods and process	6
Clinical evidence	7
Summary of clinical studies included in the evidence review	8
Quality assessment of studies included in the evidence review	13
Economic evidence	14
Summary of studies included in the economic evidence review.....	14
Economic model.....	14
Clinical evidence statements	14
The committee’s discussion of the evidence.....	18
References.....	21
Appendices	22
Appendix A – Review protocols	22
Review protocol for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?	22
Appendix B – Literature search strategies	26
Literature search strategies for review question: Is there an association between sleep position on going to sleep and stillbirth or having a small for gestational age baby?	26
Appendix C – Clinical evidence study selection	29
Clinical evidence study selection for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	29
Appendix D – Clinical evidence tables	30
Clinical evidence tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?	30
Appendix E – Forest plots.....	52
Forest plots for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?	52
Appendix F – GRADE tables	53
GRADE tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby? – evidence from IPD meta-analysis	53

GRADE tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby? – evidence from primary studies	56
Appendix G – Economic evidence study selection.....	69
Economic evidence study selection for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?	69
Appendix H – Economic evidence tables.....	70
Economic evidence tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	70
Appendix I – Health economic evidence profiles.....	72
Economic evidence profiles for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	72
Appendix J – Health economic analysis.....	73
Economic analysis for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	73
Appendix K – Excluded studies	74
Excluded studies for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	74
Clinical studies	74
Economic studies	78
Appendix L – Research recommendations	79
Research recommendations for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?.....	79

Maternal sleep position during pregnancy

Review question

Is there an association between sleep position on going to sleep and stillbirth or having a small for gestational age baby?

Introduction

It is possible that sleep position may affect the likelihood of stillbirth or having a small for gestational age (SGA) baby. If there is an effect of sleep position, it is important for women to be made aware of this association so that they can try and modify their sleeping position accordingly. It is also important to clarify which sleeping positions are not associated with worse outcomes to reduce unnecessary sleep pattern restriction, worry and anxiety for women. The aim of this review is to determine what the associations are between maternal sleeping positions and stillbirth or SGA.

Summary of the protocol

Please see Table 1 for a summary of the Population, Intervention, Comparison, Outcomes, Timing and Setting (PICOTS) characteristics of this review.

Table 1: Summary of the protocol (PRO table)

Population	All pregnant women
Index (prognostic factors)	<ul style="list-style-type: none">• Maternal sleep position on going to sleep after 24⁺⁰ weeks gestation:<ul style="list-style-type: none">○ Other¹○ Prone (in other words on tummy)○ Side<ul style="list-style-type: none">▪ Left lateral▪ Right lateral▪ Variable○ Sitting/propped○ Supine (in other words on back)
Comparator	Not applicable
Outcomes	<u>Outcomes for the baby</u> <ul style="list-style-type: none">• Stillbirth (fetal death after 24⁺⁰ weeks but before delivery)• Small for gestational age (SGA)
Timing	Any time after pregnancy
Setting	Sleep position and outcomes to be assessed in any relevant location.

Notes: ¹, This factor is intended to capture any sleep position not described by the other listed sleep positions.
Abbreviations: SGA: small for gestational age.

For further details see the review protocol in appendix A.

Methods and process

This evidence review was developed using the methods and process described in [Developing NICE guidelines: the manual 2014](#). Methods specific to this review question are described in the review protocol in appendix A.

In the first iteration of this evidence review, only primary evidence was identified and fully extracted. Between the first iteration and the update of this evidence review, two individual patient data (IPD) meta-analyses were published and identified. This report includes both the IPD meta-analysed outcomes and the original primary study outcomes, which were not meta-analysed at a study level.

Declarations of interest were recorded according to [NICE's conflicts of interest policy](#).

Clinical evidence

Included studies

One systematic review and meta-analysis of individual patient data (IPD) (Cronin 2019) including data from 5 case control studies (Gordon 2015, Heazell 2018, McCowan 2017, O'Brien 2019, Stacey 2011) reporting on the association between sleep position and stillbirth was included in this report.

One of the included primary studies was a phase 1 investigation whose aim was to identify modifiable risk factors for late-pregnancy stillbirth (Gordon 2015), whilst the remaining 4 studies were phase 2 investigations that examined whether specific going-to-sleep positions were associated with stillbirth. One study defined late stillbirth as fetal death after and including 32 weeks gestation (Gordon 2015), whilst the remaining 4 studies defined it as fetal death after and including 28 weeks gestation. One study was retrospective (O'Brien 2019), whilst the remaining studies were prospective. Four of the studies were multicentre studies, two of which were conducted in New Zealand (McCowan 2017, Stacey 2011), one in Australia (Gordon 2015), and one in the UK (Heazell 2018), whilst one study was an international online-based study (O'Brien 2019). All the studies used left lateral going-to-sleep position as the control arm to compare against the odds of stillbirth with other sleeping positions. Data on going-to-sleep position was reported relative to the participants' recollection of their going-to-sleep position on the last month (Gordon 2015), last night (Heazell 2018, Stacey 2011), and last night and last week (McCowan 2017, O'Brien 2019). One of the studies (McCowan 2017) only reported data according to whether late stillbirth occurred pre-term (between 28 and 36 weeks gestation) or term (greater and including 37 weeks gestation), whilst the remaining four studies reported data for still birth regardless of term status.

Linked to the analysis reported in Cronin 2019, a secondary analysis (Anderson 2019) used IPD from 4 of the included studies (Heazell 2018, Stacey 2011, McCowan 2017, Gordon 2015) and explored the association between the position in which pregnant women went to sleep and infant birth weight.

The included studies are summarised in Table 2 and Table 3. See also the literature search strategy in appendix B and the study selection flow chart in appendix C.

Excluded studies

Studies not included in this review and reasons for their exclusions are provided in appendix K.

Summary of clinical studies included in the evidence review

A summary of the studies that were included in this review is presented in Table 2 and Table 3.

Table 2: Characteristics of included studies for the association between maternal sleep position and stillbirth

Study ID Study design Country Size of cohort	Cases Number of participants	Controls Number of participants	Sleep positions assessed	Confounding factors adjusted for
Cronin 2019 SR with MA IPD IPD from 4 case control studies: Gordon 2015; Heazell 2017; McCowan 2017; Stacey 2011 1 nested case control: O'Brien 2019	Women with singleton pregnancies who had stillbirth at ≥ 28 weeks gestation N=851	Women with singleton pregnancies at ≥ 28 weeks gestation N=2257	Going-to-sleep position (last available position, within 2 weeks of estimated fetal death in cases) and stillbirth: <ul style="list-style-type: none"> • Left lateral (reference) • Supine • Right • Prone • Variable • Propped up • Don't remember 	<ul style="list-style-type: none"> • Maternal age • Earliest pregnancy BMI • Ethnicity • Parity • Education level • Marital status • Pre-existing hypertension or diabetes • Smoking • Recreational drug use • Fetal movements • Infant birthweight by customised centiles <p>Three confounders going-to-sleep duration, frequency of overnight toilet use, and day-time napping) analysed in sensitivity models</p>
Gordon 2015	Women with singleton pregnancies who had	Women at 32 weeks of gestation with singleton pregnancies matched for	Going-to-sleep position in last month and stillbirth:	<ul style="list-style-type: none"> • Age • BMI

Study ID Study design Country Size of cohort	Cases Number of participants	Controls Number of participants	Sleep positions assessed	Confounding factors adjusted for
Prospective population-based case-control study Australia N=295	stillbirth at ≥32 weeks gestation N=103	booking hospital and gestation who were recruited at the same period as cases N=192	<ul style="list-style-type: none"> • Left lateral (reference) • Other¹ • Right lateral • Supine 	<ul style="list-style-type: none"> • Level of education • Not in paid work • Parity • Sleep apnoea symptoms Smoking during pregnancy
Heazell 2017 Prospective population-based case-control study UK N=1024	Women with singleton pregnancies who had a stillbirth ≥28 weeks gestation with no known congenital anomaly N=291	Women with an ongoing pregnancy at the time of interview with no congenital anomaly N=733	Going-to-sleep position on last night and stillbirth: <ul style="list-style-type: none"> • Left lateral (reference) • Prone • Propped • Right lateral • Supine • Variable-lateral 	<ul style="list-style-type: none"> • Age • BMI • Ethnicity • Level of education • Other sleep related variables that were significant in univariable analysis (including sleep duration on last night, number of times visited toilet on last night, and number of maternal daytime naps in last 4 weeks) • Parity • Small-for-gestational age Smoking during pregnancy
McCowan 2017 Prospective population-based case-control study	Women with singleton pregnancies who had a stillbirth at ≥28 weeks gestation N=164	Women with singleton pregnancies with ongoing pregnancies in the participating health region N=569	Going-to-sleep position on last night and pre-term stillbirth (28-36 weeks gestation): <ul style="list-style-type: none"> • Left lateral (reference) • Propped • Other - Restless² 	<ul style="list-style-type: none"> • Age • BMI • Ethnicity • Deprivation index

Study ID Study design Country Size of cohort	Cases Number of participants	Controls Number of participants	Sleep positions assessed	Confounding factors adjusted for
New Zealand N=733			<ul style="list-style-type: none"> • Right lateral • Supine <p>Going-to-sleep position on last week and pre-term stillbirth (28-36 weeks gestation):</p> <ul style="list-style-type: none"> • Left lateral (reference) • Prone • Propped • Right lateral • Supine • Variable-lateral <p>Going-to-sleep position on last night and term stillbirth (≥ 37 weeks gestation):</p> <ul style="list-style-type: none"> • Left lateral • Other - Restless² • Right lateral • Propped • Supine <p>Going-to-sleep position on last week and term stillbirths (≥ 37 weeks gestation):</p> <ul style="list-style-type: none"> • Left lateral (reference) • Propped • Right lateral 	<ul style="list-style-type: none"> • Other sleep related variables significant in univariable analysis (hours of night time sleep on the last night, getting up to toilet during the last night, sleep during the daytime in the last week) • Parity • Small-for-gestational age status • Smoking during pregnancy

Study ID Study design Country Size of cohort	Cases Number of participants	Controls Number of participants	Sleep positions assessed	Confounding factors adjusted for
			<ul style="list-style-type: none"> • Supine • Variable-lateral 	
<p>O'Brien 2019³</p> <p>Nested case-control study in uncontrolled cohort of known size</p> <p>Anonymous international online survey</p> <p>N=633</p>	<p>Women with a singleton stillbirth baby of ≥ 28 weeks gestation in month before completing the survey</p> <p>N=153</p>	<p>Women who were either pregnant with ≥ 28 weeks gestation or had a live birth in week before completing the survey</p> <p>N=480</p>	<p>Gong-to-sleep position on last night and stillbirth:</p> <ul style="list-style-type: none"> • Left lateral (reference) • Propped • Right lateral • Supine • Variable-lateral <p>Going-to-sleep position on last month and stillbirth:</p> <ul style="list-style-type: none"> • Left lateral (reference) • Propped • Right lateral • Supine • Variable-lateral 	<ul style="list-style-type: none"> • Age • BMI • Country of residence • Ethnicity • Level of education • Parity • Smoking during pregnancy
<p>Stacey 2011</p> <p>Prospective population-based case-control study</p> <p>New Zealand</p> <p>N=465</p>	<p>Women with singleton pregnancies who had a stillbirth at ≥ 28 weeks gestation</p> <p>N=155</p>	<p>Women selected from the pregnancy registration list of the district health board, matched for gestation to cases</p> <p>N=310</p>	<p>Going-to-sleep position on last night and stillbirth:</p> <ul style="list-style-type: none"> • Left lateral (Reference) • Other⁴ • Right lateral • Supine 	<ul style="list-style-type: none"> • Age • BMI • Ethnicity • Parity • Smoking during pregnancy • Socioeconomic status

Notes: ¹, The category of 'Other' going-to-sleep position reported in this study included any going-to-sleep position other than left lateral, right lateral or supine. This categorisation was not the same as intended in the protocol and this position was therefore not included in the narrative analysis; ², Restless going-to-sleep position refers to women with frequently changing positions when going to sleep and could not remember the position just before falling asleep; ³, Women were recruited using an online survey system, 'Study of Trends and Risk Factors for Stillbirth', developed by an international consortium of clinicians and academics together with the Star Legacy Foundation and other stillbirth and parental support groups; ⁴, The category of 'Other' going-to-sleep position reported in this study included prone, sitting, sleeping on both lateral sides, and unsure or don't remember going-to-sleep position. This categorisation was not the same as intended in the protocol and this position was therefore not included in the narrative analysis. Abbreviations: N: total number of participants in the study or case or control; BMI: body mass index; CI: confidence interval; QUIPS, Quality in Prognostic Studies checklist; SGA: small for gestational age.

Table 3: Characteristics of included studies for the association between maternal sleep position and small for gestational age

Study ID Study design Country Size of cohort	Cases Number of participants	Controls Number of participants	Sleep position assessed	Confounding factors adjusted for
Anderson 2019 Subgroup analysis of IPD MA IPD from 4 case control studies: Gordon 2015; Heazell 2017; McCowan 2017; Stacey 2011	Women with singleton pregnancies who went to sleep in a supine position N=57	Women with singleton pregnancies who went to sleep in a non-supine position N=1703	Going-to-sleep position over last week/2 weeks/month (whichever was longest and available from primary study) and SGA: <ul style="list-style-type: none"> • Left lateral (Reference) • Other (not specified) • Right lateral • Supine 	<ul style="list-style-type: none"> • Study site • Maternal age • Height • Weight • Parity • Ethnicity • Preexisting diabetes • Preexisting hypertension • Antepartum haemorrhage • Gestational hypertensive disorder • Gestational diabetes • Cigarette smoking • Recreational drug use

Abbreviations: IPD MA: individual patient data meta-analysis; N: total number of participants in the study or case or control; BMI: body mass index; CI: confidence interval; SGA: small for gestational age; vs: versus

See the full evidence tables in appendix D.

Quality assessment of studies included in the evidence review

See the evidence profiles in appendix F.

Economic evidence

Included studies

A systematic review of the economic literature was conducted but no economic studies were identified which were applicable to this review question.

A single economic search was undertaken for all topics included in the scope of this guideline. See supplementary material 2 for details.

Excluded studies

Economic studies not included in this review are listed, and reasons for their exclusion are provided in appendix K.

Summary of studies included in the economic evidence review

No economic studies were identified which were applicable to this review question.

Economic model

No economic modelling was undertaken for this review because the committee agreed that other topics were higher priorities for economic evaluation.

Clinical evidence statements

Individual patient data meta-analysis results

Association between going-to-sleep position (last available position, within last 2 weeks) and stillbirth in comparison to left lateral going-to-sleep position

- High quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found an important association between supine going-to-sleep position and stillbirth: aOR 2.63 (95% CI 1.72 to 4.04).
- Low quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found no important association between right sided going-to-sleep position and stillbirth: aOR 1.04 (95% CI 0.83 to 1.31).
- Very low quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found no important association between prone going-to-sleep position and stillbirth: aOR 0.63 (95% CI 0.12 to 3.25).
- Very low quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found no important association between variable side going-to-sleep position and stillbirth: aOR 0.97 (95% CI 0.70 to 1.35).
- Very low quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found no important association between propped up going-to-sleep position and stillbirth: aOR 1.30 (95% CI 0.68 to 2.49).
- Moderate quality evidence from an IPD meta-analysis of 5 case control studies (N=3108) found an important association between not remembering going-to-sleep position and stillbirth: aOR 2.26 (95% CI 1.48 to 3.46).

Association between going-to-sleep position (last available position, within last 2 weeks) and small for gestational age in comparison to left lateral going-to-sleep position

- High quality evidence from an IPD meta-analysis of 4 case control studies (N=1760) found an important association between supine going-to-sleep position and babies being born small for gestational age: aOR 3.23 (95% CI 1.37 to 7.59).
- Very low quality evidence from an IPD meta-analysis of 4 case control studies (N=1760) found no important association between right sided going-to-sleep position and babies being born small for gestational age: aOR 1.05 (95% CI 0.58 to 1.90).
- Very low quality evidence from an IPD meta-analysis of 4 case control studies (N=1760) found no important association between other going-to-sleep position and babies being born small for gestational age: aOR 1.14 (95% CI 0.62 to 2.09).

Primary study results**Association between going-to-sleep position on last night and stillbirth in comparison to left lateral going-to-sleep position on last night**

- Very low quality evidence from 1 phase 2 prospective population-based case-control study (N=1024) examined whether there was an association between prone going-to-sleep position on last night and stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 1.01 (95%CI 0.13 to 7.81)
- Very low quality evidence from 3 phase 2 case-control studies (N=2122) - 2 non-nested prospective population-based studies and 1 retrospective nested study in an online cohort of known size - examined whether there was an association between right-lateral going-to-sleep position on last night and stillbirth compared to left lateral going-to-sleep position on last night. However, none of the 3 studies found any such association: aOR 0.67 (95%CI 0.44 to 1.02); aOR 1.11 (95%CI 0.70 to 1.77); aOR 1.74 (95%CI 0.98 to 3.01)
- Very low quality evidence from 2 phase 2 case-control studies - 1 non-nested prospective population-based study and 1 retrospective nested study in an online cohort of known size (N=1657) - examined whether there was an association between sitting/propped going-to-sleep position on last night and stillbirth compared to left lateral going-to-sleep position on last night. However, none of the studies found any such association: aOR 0.44 (95%CI 0.13 to 1.49); aOR 0.71(95%CI 0.22 to 2.30).
- Very low quality evidence from 3 phase 2 case-control studies (N=2122) - 2 non-nested prospective population-based studies and 1 retrospective nested study in an online cohort of known size - examined whether there was an association between supine going-to-sleep position on last night and stillbirth compared to left lateral going-to-sleep position on last night. Two studies found that supine going-to-sleep position was associated with an increase in stillbirth compared to left lateral going-to-sleep position (aOR 2.31 [95%CI 1.04 to 5.11]; aOR 2.54 [95%CI 1.04 to 6.18]) whereas no association was found in another study: aOR 1.05 (95%CI 0.32 to 3.50). However this latter study, which was powered to detect an association between supine going-to-sleep position and stillbirth assuming a 20% exposure frequency, reported an exposure frequency of only 2.3%.
- Very low quality evidence from 2 phase 2 case-control studies - 1 non-nested prospective population-based study and 1 retrospective nested study in an online cohort of known size (N=1657) - examined whether there was an association between variable-lateral going-to-sleep position on last night and stillbirth compared to left lateral going-to-sleep

position on last night. However, none of the studies found any such association: aOR 0.93 (95%CI 0.51 to 1.69); aOR 0.75 (95%CI 0.34 to 1.64).

Association between going-to-sleep position on last month of pregnancy and late stillbirth in comparison to left lateral going-to-sleep position

- Very low quality evidence from 2 phase 2 case-control studies (N=928) - 1 non-nested prospective population-based study and 1 retrospective nested study in an online cohort of known size - examined whether there was an association between right-lateral going-to-sleep position on last month and stillbirth compared to left lateral going-to-sleep position on last month. However, none of the studies found any such association: aOR 1.1 (95%CI 0.43 to 2.6); aOR 1.14 (95%CI 0.70 to 1.85).
- Very low quality evidence from 1 phase 2 case control study – a retrospective nested study in an online cohort of known size (N=633) - examined whether there was an association between sitting/propped going-to-sleep position on last month and stillbirth compared to left lateral going-to-sleep position on last month. However, no association was found in this study: aOR 1.20 (95%CI 0.39 to 3.68).
- Very low quality evidence from 2 case-control studies in (N=928) - 1 phase 1 non-nested prospective population-based study and 1 phase 2 retrospective nested study in an online cohort of known size - examined whether there was an association between supine going-to-sleep position on last month and stillbirth compared to left lateral going-to-sleep position on last month. One study found that supine going-to-sleep position was associated with an increase in stillbirth compared to left lateral going-to-sleep position: aOR 6.26 (95%CI 1.2 to 34.00) whereas no association was found in another study: aOR 0.37 (95%CI 0.04 to 3.12).
- Very low quality evidence from 1 phase 2 retrospective nested case control study in an online cohort of known size (N=633) examined whether there was an association between variable-lateral going-to-sleep position on last month and stillbirth compared to left lateral going-to-sleep position on last month. However, no association was found in this study: aOR 0.87 (95%CI 0.48 to 1.55).

Association between going-to-sleep position on last night or last week of pregnancy and pre-term stillbirth (28 to 36 weeks gestation) in comparison to left lateral going-to-sleep position on last night or last week

Going-to-sleep position on last night

- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between restless going-to-sleep position on last night and preterm stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 3.50 (95%CI 0.61 to 19.97).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between right-lateral going-to-sleep position on last night and preterm stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 0.96 (95%CI 0.48 to 1.94).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between sitting/propped going-to-sleep position on last night and preterm stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 4.37 (95%CI 0.11 to 178.86).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between supine

going-to-sleep position on last night and preterm stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 2.25 (95%CI 0.65 to 7.84).

Going-to-sleep position on last week

- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between prone going-to-sleep position on last week and preterm stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 10.71 (95%CI 0.43 to 268.28).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between right-lateral going-to-sleep position on last week and preterm stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 0.73 (95%CI 0.34 to 1.54).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between sitting/propped going-to-sleep position on last week and preterm stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 4.01 (95%CI 0.08 to 210.43).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between supine going-to-sleep position on last week and preterm stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 2.25 (95%CI 0.65 to 7.84).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between variable-lateral going-to-sleep position on last week and preterm stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 0.63 (95%CI 0.18 to 2.19).

Association between going-to-sleep position on last night or last week of pregnancy and term stillbirth (≥ 37 weeks gestation) in comparison to left going-to-sleep position on last night or last week

Going-to-sleep position on last night

- Very low quality evidence from 1 prospective phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between restless going-to-sleep position on last night and term stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 2.0 (95%CI 0.64, 6.21).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between right-lateral going-to-sleep position on last night and term stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 0.98 (95%CI 0.48 to 1.99).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between sitting/propped going-to-sleep position on last night and term stillbirth compared to left lateral going-to-sleep position on last night. However, no association was found in this study: aOR 1.02 (95%CI 0.17 to 5.97).

- Low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between supine going-to-sleep position on last night and term stillbirth compared to left lateral going-to-sleep position on last night. The study found that supine going-to-sleep position was associated with an increase in stillbirth compared to left lateral going-to-sleep position: aOR 10.26 (95%CI 3.01 to 35.04).

Going-to-sleep position on last week

- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between right-lateral going-to-sleep position on last week and term stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 0.95 (95%CI 0.48 to 1.89).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between sitting/propped going-to-sleep position on last week and term stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 2.64 (95%CI 0.47 to 14.81).
- Low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between supine going-to-sleep position on last week and term stillbirth compared to left lateral going-to-sleep position on last week. The study found that supine going-to-sleep position was associated with an increase in stillbirth compared to left lateral going-to-sleep position: aOR 12.73 (95%CI 2.92 to 55.46).
- Very low quality evidence from 1 phase 2 non-nested prospective population-based case-control study (N=733) examined whether there was an association between variable-lateral going-to-sleep position on last week and term stillbirth compared to left lateral going-to-sleep position on last week. However, no association was found in this study: aOR 1.11 (95%CI 0.49 to 3.01).

The committee's discussion of the evidence

Interpreting the evidence

The quality of the evidence

The quality of each primary study was appraised using the QUIPS checklist. The main reasons for downgrading the quality of studies were recall bias due to the time delay between recruitment and interview, and selection bias due to limited participation from all the eligible women. The overall quality of evidence for each outcome was assessed using a modified GRADE framework for prognostic reviews and ranged from very low to low. Although a phase 3 prospective cohort study is the best study design to establish the causal relationship between a prognostic factor and an outcome, the frequency of stillbirth is low and such a study is therefore not practically feasible to conduct. Therefore, no restrictions were initially placed on the appropriate types of study considered for this review with phase 1 and 2 studies initially rated as providing a moderate and high quality of evidence, respectively. The main reasons for downgrading the overall evidence for all sleep positions was risk of bias associated with each of the contributing studies (recall bias, selection bias), inconsistency (variation in effect estimates across studies), indirectness (sample not clearly representative of target population), imprecision (effect estimate has wide 95% confidence intervals) and publication bias (due to the small number of early phase studies reporting outcomes).

The IPD meta-analysis quality was appraised using the ROBIS checklist and both IPD meta-analyses were at low risk of bias. The individual outcomes within the IPD meta-analyses ranged in quality rating from high (for example supine sleeping position and its association with stillbirth) to very low. The IPD meta-analyses outcomes were typically downgraded due to imprecision in their estimates and publication bias (due to the small number of early phase primary studies reporting outcomes).

Benefits and harms

Overall, the evidence suggests that there may be an increased risk of stillbirth after 28+0 weeks and babies being born small for gestational age (SGA) associated with going to sleep on one's back (in other words in the supine position) compared to going to sleep on one's left-hand side – in both the primary studies and the IPD meta-analysis. The best estimate, from the IPD meta-analysis was that sleeping in a supine position approximately doubled the odds of stillbirth and trebled the odds of babies being born SGA.

While the quality of the evidence from the primary studies ranged depending on the timing and precise outcomes considered, the quality of the evidence from the IPD, particularly for the evidence around supine sleeping position, was relatively high. The evidence still may not be considered to be definitive evidence of causality between going to sleep position and stillbirth or SGA as the IPD meta-analysis does not overcome issues with the study design of the primary studies. However the committee agreed it was of sufficient quality to advise women to try to avoid going to sleep on their back after 28 weeks and inform women of the likely link with stillbirth, alongside a caveat that the evidence is uncertain. The committee chose to specifically highlight stillbirth as this is a more concerning outcome than babies being born SGA and they agreed that including SGA in the recommendations made the advice less clear.

The evidence also suggests that the risk of going to sleep in any other position is the same as that of going to sleep on one's left side (that is, there is not an increased likelihood of stillbirth associated with any other going-to-sleep position). The committee agreed that, as there has been some concern that there is such an association, it was important to highlight that the evidence does not support the view that there is an increased likelihood of stillbirth associated with going to sleep on one's right side irrespective of the time of reporting compared to going to sleep on one's left side. However the evidence for other sleep positions was generally an absence of definitive association (with serious imprecision) as opposed to definitive evidence of no association, therefore the committee did not include this information in the recommendations.

The committee discussed that the studies looked at the association between going to sleep position and outcomes because due to the nature of the observational studies relying on women's recall, it would not be possible to study the association between the woman's sleeping position and outcomes. However, the going to sleep position should be considered a proxy for sleeping position. Sleeping position is perhaps best controlled by controlling the going to sleep position. In addition, pillows or other props could be used to aid that the position stays when sleeping. This review did not assess the effectiveness of any interventions to modify sleeping position but in the committee agreed, based on their knowledge and experience, to recommend advising women to consider using for example pillows so that they can maintain their position when sleeping.

The committee noted that there may be a psychological impact of informing pregnant women of the potential link between sleeping on one's back and stillbirth or SGA and did not want to cause undue anxiety. The committee also noted the relatively low incidence of stillbirth (1 in every 244 births in the UK according to 2018 [Office for National Statistics \[ONS\] data](#)).

However on balance they agreed that the evidence was strong enough that women should be advised about the risk.

Cost effectiveness and resource use

No economic studies were identified which were applicable to this review question.

Professional time advising women on healthy behaviours is already current practice. Therefore, the recommendations to advise women on sleeping position, and the possible link between sleep position and adverse outcomes is unlikely to require any increase in clinician time. In turn, advising women against sleeping on their backs after 28 weeks of pregnancy may reduce adverse outcomes such as stillbirth and small for gestational age reducing future healthcare costs.

References

Anderson 2019

Anderson, N.H., Gordon, A., Li, M., Cronin RS, Thompson, J.M.D., Raynes-Greenow, C.H., Heazell, A.E.P., Stacey, T., Culling, V.M., Wilson, J., Askie, L.M., Mitchell, E.A., McCowan, L.M.E.. Association of Supine Going-to-Sleep Position in Late Pregnancy With Reduced Birth Weight. A Secondary Analysis of an Individual Participant Data Meta-analysis. *JAMA Network Open* 2019; 2(10):e1912614

Cronin 2019

Cronin RS, Li, M., Thompson, J.M.D., Gordon, A., Raynes-Greenow, C.H., Heazell, A.E.P., Stacey, T., Culling, V.M., Bowring, V. Anderson, N.H., O'Brien, L.M., Mitchell, E.A., Askie, L.M., McCowan, L.M.E.. An Individual Participant Data Meta-analysis of Maternal Going-to-Sleep Position, Interactions with Fetal Vulnerability, and the Risk of Late Stillbirth. *Eclinical Medicine* 2019; 10:49-57.

Gordon 2015

Gordon, A., Raynes-Greenow, C., Bond, D., Morris, J., Rawlinson, W., Jeffery, H. Sleep position, fetal growth restriction, and late-pregnancy stillbirth: The sydney stillbirth study, *Obstetrics and Gynecology*, 125, 347-355, 2015.

Heazell 2018

Heazell, A. E. P., Li, M., Budd, J., Thompson, J. M. D., Stacey, T., Cronin, R. S., Martin, B., Roberts, D., Mitchell, E. A., McCowan, L. M. E. Association between maternal sleep practices and late stillbirth - findings from a stillbirth case-control study, *BJOG: An International Journal of Obstetrics and Gynaecology*, 125, 254-262, 2018.

McCowan 2017

McCowan, L. M. E., Thompson, J. M. D., Cronin, R. S., Li, M., Stacey, T., Stone, P. R., Lawton, B. A., Ekeroma, A. J., Mitchell, E. A. Going to sleep in the supine position is a modifiable risk factor for late pregnancy stillbirth; Findings from the New Zealand multicentre stillbirth case-control study, *Plos ONE*, 12 (6): e0179396, 2017.

O'Brien 2019

O'Brien, L. M., Warland, J, Stacey, T, Heazell, A. E. P., Mitchell, E. A. Maternal sleep practices and stillbirth: Findings from an international case-control study, *Issues in Perinatal care*, 1-11, 2019.

Stacey 2011

Stacey, T., Thompson, J. M. D., Mitchell, E. A., Ekeroma, A. J., Zuccollo, J. M., McCowan, L. M. E. Association between maternal sleep practices and risk of late stillbirth: A case-control study, *BMJ*, 342: d3403, 2011.

Appendices

Appendix A – Review protocols

Review protocol for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

Table 4: Review protocol

Field (based on PRISMA-P)	Content
Review question	Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?
Type of review question	Prognostic factors review
Objective of the review	The aim of this review is to evaluate whether specific sleep positions are associated with still birth or having a small for gestational age baby after 24 ⁺⁰ weeks gestation.
Eligibility criteria – population	All pregnant women
Eligibility criteria – Risk factors (s)	<ul style="list-style-type: none"> • Maternal sleep position on going to sleep after 24⁺⁰ weeks gestation: <ul style="list-style-type: none"> ○ Other ○ Prone (i.e. tummy) ○ Side <ul style="list-style-type: none"> ▪ Left lateral ▪ Right lateral ▪ Variable ○ Sitting/propped ○ Supine (i.e. on back) <p>Notes: The position 'other' is intended to capture any other sleep position that is not listed. SGA is defined as having a birth weight below the 10th centile. Some studies will report this as Low Birth Weight (LBW) adjusted for Gestational Age (GA) rather than as SGA. For participants in the case group, sleep position reported may be usual sleep position or sleep position on the night before stillbirth.</p>
Eligibility criteria – Confounding factors	Analysis used by studies must adjust for confounding factors using logistic regression to conduct multivariable analysis.
Outcomes and prioritisation	<p>Model performance</p> <ul style="list-style-type: none"> • Discrimination <ul style="list-style-type: none"> ○ Concordance (C) statistic <p>Note: the C statistic is also known as 'area under the receiver operating characteristics curve' (AUC).</p>

Field (based on PRISMA-P)	Content
	<p>Outcomes</p> <ul style="list-style-type: none"> • Still birth (i.e. fetal death after 24⁺⁰ weeks but before delivery) • Small for gestational age (SGA) <p>Note: SGA is defined as having a birth weight below the 10th centile. Some studies will report this as Low Birth Weight (LBW) adjusted for Gestational Age (GA) rather than as SGA.</p>
Eligibility criteria – study design	<p>INCLUDE:</p> <ul style="list-style-type: none"> • Systematic reviews of prognostic studies • Prognostic observational studies <ul style="list-style-type: none"> ○ Prospective cohort studies ○ Retrospective cohort studies ○ Nested case-control studies within a cohort of known size <p>The following types of study design will be considered only if no studies of the above types are identified:</p> <ul style="list-style-type: none"> • Non-nested case-control studies
Other inclusion exclusion criteria	<p>EXCLUDE:</p> <p>POPULATION:</p> <ul style="list-style-type: none"> • Multiple pregnancy • Pregnancy with congenital anomalies • Women who had stillborn or live-born babies with congenital abnormalities <p>STUDY DESIGN:</p> <ul style="list-style-type: none"> • Cross-over studies • Cross-sectional studies • Non-comparative studies • Randomised and quasi-randomised controlled trials <p>LANGUAGE:</p> <ul style="list-style-type: none"> • Non-English <p>PUBLICATION STATUS:</p> <ul style="list-style-type: none"> • Conference abstract <p>INCLUDE</p> <p>COUNTRY:</p> <ul style="list-style-type: none"> • No restriction
Proposed sensitivity/sub-group analysis, or meta-regression	<p>If the studies are sufficiently similar to merit meta-analysis, subgroup analysis according to World Bank status of country (High-income countries; Low and middle-income countries) in which they were conducted in will be performed (see https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups for classification of countries).</p> <p>Statistical heterogeneity will be assessed by visually examining the forest plots and by calculating the I² inconsistency statistic (with an I² value ≥50% indicating serious heterogeneity, and ≥80% indicating very serious heterogeneity).</p>

Field (based on PRISMA-P)	Content
Selection process – duplicate screening/selection/analysis	Review questions selected as high priorities for health economic analysis (and those selected as medium priorities and where health economic analysis could influence recommendations) will be subject to dual weeding and study selection; any discrepancies above 10% of the dual weeded resources will be resolved through discussion between the first and second reviewers or by reference to a third person. All data extraction will quality assured by a senior reviewer. Draft excluded studies and evidence tables will be circulated to the Topic Group for their comments. Resolution of disputes will be by discussion between the senior reviewer, Topic Advisor and Chair.
Data management (software)	NGA STAR software will be used to generate bibliographies/citations, and to conduct study sifting and data extraction. Pairwise meta-analyses, if possible, will be performed using Cochrane Review Manager (RevMan5). For details please see Supplement 1: methods.
Information sources – databases and dates	Sources to be searched: Medline, Medline In-Process, CCTR, CDSR, DARE, HTA, Embase Limits: <ul style="list-style-type: none"> • Date limit: none • Apply standard animal/non-English language exclusion • Limit to prognostic studies in first instance but download all results.
Identify if an update	This is a new area in the guideline.
Author contacts	Developer: National Guideline Alliance.
Highlight if amendment to previous protocol	For details please see section 4.5 of Developing NICE guidelines: the manual .
Search strategy – for one database	For details please see appendix B.
Data collection process – forms/duplicate	A standardised evidence table format will be used, and published as appendix D (clinical evidence tables) or H (economic evidence tables).
Data items – define all variables to be collected	For details please see evidence tables in appendix D (clinical evidence tables) or H (economic evidence tables).
Methods for assessing bias at outcome/study level	Quality assessment of individual studies will be performed using the following checklists: <ul style="list-style-type: none"> • ROBIS for systematic reviews of prognostic studies • QUIPS for prognostic studies For details please see section 6.2 of Developing NICE guidelines: the manual . The risk of bias for the evidence for each prognostic factor (i.e. sleep position) will be evaluated using an adaptation of the ‘Grading of Recommendations Assessment, Development and Evaluation (GRADE) toolbox’: http://www.gradeworkinggroup.org/ . For further details as to how GRADE will be adapted, see the following article: <ul style="list-style-type: none"> • Huguet, A., Hayden, J. A., Stinson, J., McGrath, P. J., Chambers, C. T., Tougas, M. E., & Wozney, L. (2013). Judging the quality of evidence in reviews of prognostic factor research: adapting the GRADE framework. <i>Systematic reviews</i>, 2(1), 71.
Criteria for quantitative synthesis (where suitable)	Meta-analyses will be conducted for this review only if the same confounders are accounted for in the analyses, the same analytical methods are adapted, and the populations assessed are suitably similar. In all other cases, the results will be reported separately.
Methods for analysis – combining studies and exploring (in)consistency	The adjusted Risk Ratio or Odds Ratio and 95% confidence intervals will be plotted in RevMan if appropriate, although the results for each relative measure will be presented separately. Good model performance regarding discrimination will be defined as a C statistic >0.75 as suggested in Debray 2017: <ul style="list-style-type: none"> • Debray, T. P., Damen, J. A., Snell, K. I., Ensor, J., Hooft, L., Reitsma, J. B., ... & Moons, K. G. (2017). A guide to systematic review and meta-analysis of prediction model performance. <i>Bmj</i>, 356, i6460.

Field (based on PRISMA-P)	Content
	If a meta-analysis is conducted, inconsistency will be assessed by visual examination of the forest plots and the I^2 statistic (with $I^2 \geq 50\%$ indicating serious heterogeneity and $I^2 \geq 80\%$ indicating very serious heterogeneity).
Meta-bias assessment – publication bias, selective reporting bias	For details please see Supplement 1: methods and section 6.2 of Developing NICE guidelines: the manual . If sufficient relevant RCT evidence is available, publication bias will be explored using RevMan software to examine funnel plots. Trial registries will be examined to identify missing evidence: Clinical trials.gov, NIHR Clinical Trials Gateway.
Assessment of confidence in cumulative evidence	For details please see sections 6.4 and 9.1 of Developing NICE guidelines: the manual .
Rationale/context – Current management	For details please see the introduction to the evidence review.
Describe contributions of authors and guarantor	A multidisciplinary committee developed the guideline. The committee was convened by the National Guideline Alliance and chaired by Kate Harding in line with section 3 of Developing NICE guidelines: the manual . Staff from the National Guideline Alliance undertook systematic literature searches, appraised the evidence, conducted meta-analysis and cost-effectiveness analysis where appropriate, and drafted the guideline in collaboration with the committee. For details please see Supplement 1: methods.
Sources of funding/support	The National Guideline Alliance is funded by NICE and hosted by the Royal College of Obstetricians and Gynaecologists.
Name of sponsor	The National Guideline Alliance is funded by NICE and hosted by the Royal College of Obstetricians and Gynaecologists.
Roles of sponsor	NICE funds the National Guideline Alliance to develop guidelines for those working in the NHS, public health, and social care in England.
PROSPERO registration number	This protocol is not registered with PROSPERO.

AUC: area under the curve, CDSR: Cochrane Database of Systematic Reviews; CENTRAL: Cochrane Central Register of Controlled Trials; DARE: Database of Abstracts of Reviews of Effects; GRADE: Grading of Recommendations Assessment, Development and Evaluation; HTA: Health Technology Assessment; LBW: Low Birth Weight; NGA: National Guideline Alliance; NHS: National health service; NICE: National Institute for Health and Care Excellence; RCT: randomised controlled trial; RoB: risk of bias; SGA: small for gestational age

Appendix B – Literature search strategies

Literature search strategies for review question: Is there an association between sleep position on going to sleep and stillbirth or having a small for gestational age baby?

Database(s): Medline & Embase (Multifile)

Last searched on **Embase Classic+Embase** 1947 to 2020 September 08, **Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily** 1946 to September 08, 2020

Date of last search: 9th September 2020

Multifile database codes: emczd = Embase Classic+Embase; ppez= MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily

#	Searches
1	Pregnancy/ use ppez
2	Pregnant Women/ use ppez
3	pregnancy/ use emczd
4	pregnant woman/ use emczd
5	pregnan\$.tw,kw.
6	1 or 2 or 3 or 4 or 5
7	Sleep/ use ppez
8	sleep/ use emczd
9	sleep pattern/ use emczd
10	Sleep Apnea Syndromes/ use ppez
11	sleep disordered breathing/ use emczd
12	(sleep\$ adj5 position\$).tw,kw.
13	(sleep\$ adj3 (pattern\$ or practice\$ or duration or quality or quantity or deprivation\$ or disruption\$ or disturb\$)).tw,kw.
14	(sleep\$ adj (apnea\$ or apnoea\$)).tw,kw.
15	(pillow\$ and sleep\$).tw,kw.
16	7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15
17	maternal sleep/ use emczd
18	(matern\$ adj sleep\$).tw,kw.
19	6 and 16
20	17 or 18 or 19
21	(controlled clinical trial or pragmatic clinical trial or randomized controlled trial).pt. or drug therapy.fs. or (groups or placebo or randomi#ed or randomly or trial).ab.
22	crossover procedure/ or double blind procedure/ or randomized controlled trial/ or single blind procedure/ or (assign* or allocat* or crossover* or cross over* or ((doubl* or singl*) adj blind*) or factorial* or placebo* or random* or volunteer*).ti,ab.
23	meta-analysis/
24	meta-analysis as topic/
25	systematic review/
26	meta-analysis/
27	((meta analy* or metanaly* or metaanaly*).ti,ab.
28	((systematic or evidence) adj2 (review* or overview*)).ti,ab.
29	((systematic* or evidence*) adj2 (review* or overview*)).ti,ab.
30	(reference list* or bibliograph* or hand search* or manual search* or relevant journals).ab.
31	(search strategy or search criteria or systematic search or study selection or data extraction).ab.
32	(search* adj4 literature).ab.
33	(medline or pubmed or cochrane or embase or psychlit or psyclit or psychinfo or psycinfo or cinahl or science citation index or bids or cancerlit).ab.
34	cochrane.jw.
35	((pool* or combined) adj2 (data or trials or studies or results)).ab.
36	letter/
37	editorial/
38	news/
39	exp historical article/
40	Anecdotes as Topic/
41	comment/
42	case report/
43	(letter or comment*).ti.

#	Searches
44	36 or 37 or 38 or 39 or 40 or 41 or 42 or 43
45	randomized controlled trial/ or random*.ti,ab.
46	44 not 45
47	animals/ not humans/
48	exp Animals, Laboratory/
49	exp Animal Experimentation/
50	exp Models, Animal/
51	exp Rodentia/
52	(rat or rats or mouse or mice).ti.
53	46 or 47 or 48 or 49 or 50 or 51 or 52
54	letter.pt. or letter/
55	note.pt.
56	editorial.pt.
57	case report/ or case study/
58	(letter or comment*).ti.
59	54 or 55 or 56 or 57 or 58
60	randomized controlled trial/ or random*.ti,ab.
61	59 not 60
62	animal/ not human/
63	nonhuman/
64	exp Animal Experiment/
65	exp Experimental Animal/
66	animal model/
67	exp Rodent/
68	(rat or rats or mouse or mice).ti.
69	61 or 62 or 63 or 64 or 65 or 66 or 67 or 68
70	53 use ppez
71	69 use emczd
72	70 or 71
73	21 use ppez
74	22 use emczd
75	73 or 74
76	(or/23-24,27,29-34) use ppez
77	(or/25-28,30-35) use emczd
78	76 or 77
79	20 and 72
80	20 not 79
81	limit 80 to english language
82	75 or 78
83	81 and 82
84	predict.ti.
85	(validat* or rule*).ti,ab.
86	(predict* and (outcome* or risk* or model*)),ti,ab.
87	((history or variable* or criteria or scor* or characteristic* or finding* or factor*) and (predict* or model* or decision* or identifi* or prognos*)),ti,ab.
88	Logistic models/ use ppez
89	Statistical model/ use emczd
90	decision*.ti,ab.
91	88 or 89
92	90 and 91
93	(decision* and (model* or clinical*)),ti,ab.
94	(prognostic and (history or variable* or criteria or scor* or characteristic* or finding* or factor* or model*)),ti,ab.
95	(stratification or discrimination or discriminate or c statistic or "area under the curve" or AUC or calibration or indices or algorithm or multivariable).ti,ab.
96	ROC curve/ use ppez
97	Receiver operating characteristic/ use emczd
98	84 or 85 or 86 or 87 or 92 or 93 or 94 or 95 or 96 or 97
99	81 and 98
100	83 or 99

Database(s): Cochrane Library

Last searched on **Cochrane Database of Systematic Reviews**, Issue 9 of 12, September 2020, **Cochrane Central Register of Controlled Trials**, Issue 9 of 12, September 2020
Date of last search: 9th September 2020

#	Searches
#1	MeSH descriptor: [Pregnancy] this term only
#2	MeSH descriptor: [Pregnant Women] this term only
#3	(pregnan*):ti,ab,kw
#4	#1 or #2 or #3
#5	MeSH descriptor: [Sleep] this term only
#6	MeSH descriptor: [Sleep Apnea Syndromes] this term only
#7	((sleep* NEAR/5 position*)):ti,ab,kw
#8	((sleep* NEAR/3 (pattern* or practice* or duration or quality or quantity or deprivation* or disruption* or disturb*)):ti,ab,kw
#9	((sleep* NEXT (apnea* or apnoea*)):ti,ab,kw
#10	((pillow* and sleep*)):ti,ab,kw
#11	#5 or #6 or #7 or #8 or #9 or #10
#12	#4 AND #11
#13	((matern* NEXT sleep*)):ti,ab,kw
#14	#12 OR #13

Database(s): CRD: Database of Abstracts of Reviews of Effects (DARE), HTA Database

Date of last search: 9th September 2020

#	Searches
1	MeSH DESCRIPTOR Pregnancy EXPLODE ALL TREES IN DARE,HTA
2	MeSH DESCRIPTOR Pregnant Women EXPLODE ALL TREES IN DARE,HTA
3	(pregnan*) IN DARE, HTA
4	#1 OR #2 OR #3
5	MeSH DESCRIPTOR Sleep EXPLODE ALL TREES IN DARE,HTA
6	MeSH DESCRIPTOR Sleep Apnea Syndromes EXPLODE ALL TREES IN DARE,HTA
7	((sleep* NEAR position*)) IN DARE, HTA
8	((sleep* NEAR (pattern* or practice* or duration or quality or quantity or deprivation* or disruption* or disturb*))) IN DARE, HTA
9	((sleep* NEAR (apnea* or apnoea*))) IN DARE, HTA
10	((pillow* and sleep*)) IN DARE, HTA
11	#5 OR #6 OR #7 OR #8 OR #9 OR #10
12	#4 AND #11
13	((matern* NEXT sleep*)) IN DARE, HTA
14	#12 OR #13

Database(s): Cinahl Plus


Date of last search: 9th September 2020

#	Searches
S16	S14 NOT S15 Limiters - English Language
S15	PT anecdote or PT audiovisual or PT bibliography or PT biography or PT book or PT book review or PT brief item or PT cartoon or PT commentary or PT computer program or PT editorial or PT games or PT glossary or PT historical material or PT interview or PT letter or PT listservs or PT masters thesis or PT obituary or PT pamphlet or PT pamphlet chapter or PT pictorial or PT poetry or PT proceedings or PT "questions and answers" or PT response or PT software or PT teaching materials or PT website
S14	S12 OR S13
S13	TI (matern* N1 sleep*) OR AB (matern* N1 sleep*)
S12	S4 AND S11
S11	S5 OR S6 OR S7 OR S8 OR S9 OR S10
S10	TI (pillow* and sleep*) OR AB (pillow* and sleep*)
S9	TI (sleep* N1 (apnea* or apnoea*)) OR AB (sleep* N1 (apnea* or apnoea*))
S8	TI (sleep* N3 (pattern* or practice* or duration or quality or quantity or deprivation* or disruption* or disturb*)) OR AB (sleep* N3 (pattern* or practice* or duration or quality or quantity or deprivation* or disruption* or disturb*))
S7	TI (sleep* N5 position*) OR AB (sleep* N5 position*)
S6	(MH "Sleep Apnea Syndromes")
S5	(MH "Sleep")
S4	S1 OR S2 OR S3
S3	TI pregnan* or AB pregnan*
S2	(MH "Expectant Mothers")
S1	(MH "Pregnancy")

Appendix C – Clinical evidence study selection

Clinical evidence study selection for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

Figure 1: PRISMA flow chart for review question:



Appendix D – Clinical evidence tables

Clinical evidence tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

Table 5: Clinical evidence tables for IPD meta-analyses

Study details	Participants	Factors	Results	Comments
<p>Full citation</p> <p>Anderson NH, Gordon A, Li M, Cronin RS, Thompson JMD, Raynes-Greenow CH, Heazel AEP, Stacey T, Culling VM, Wilson J, Askie LM, Mitchell EA, McCowan LME</p> <p>Ref Id</p> <p>1121708</p> <p>Country/ies where the study was carried out</p> <p>NA</p> <p>Study type</p> <p>Systematic Review</p> <p>Study dates</p> <p>Searches to January 2018</p> <p>Consecutive recruitment</p> <p>No</p>	<p>4 eligible case control studies. Eligible participants comprised 1,760 women</p> <p>Inclusion criteria</p> <p>Control participants from 4 case control studies with gestational age at birth collected in weeks and days, gestation at study interview of 28 days and 0 days or more, gestation at birth less than or equal to 42 weeks and 6 weeks and data for usual going-to-sleep position up to 4 weeks before the interview.</p> <p>Exclusion criteria</p> <p>A further case-control study that was included in the CRIBBS IPD was excluded from the current analysis as this online survey collected gestational age in completed weeks only. Individual participants were also excluded if they had missing variables required for calculation of birth weight centiles</p> <p>Statistical method</p>	<p>Adjusted for study site and maternal age, height, weight, parity, ethnicity, preexisting diabetes, preexisting hypertension, antepartum hemorrhage, gestational hypertensive disorder, gestational diabetes, cigarette smoking, and recreational drug use.</p>	<p>Adjusted odds ratio (95% CI), vs left, INTERGROWTH-21st, <10th centile</p> <p>Other – 1.14 (0.62, 2.09) Right – 1.05 (0.58, 1.90) Supine - 3.23 (1.37, 7.59)</p> <p>Results also available for customised growth charts where supine is no longer statistically significantly associated with <10th centile (adjusted odds ratio 1.55 (0.72 to 3.35))</p> <p>Sleep position defined as usual position over the previous week/2 weeks or month (whichever was longest and available from study).</p>	<p>Limitations</p> <p>ROBIS</p> <p>No concerns over relevance</p> <p>Study eligibility criteria: low concern</p> <p>Identification and selection of studies: low concern</p> <p>Data collection and study appraisal: low concern (although not all eligible studies were included this was due to a lack of data appropriate for IPD meta-analysis and researchers made reasonable efforts to contact authors)</p> <p>Synthesis and findings: low concern (no formal efforts to consider between study variation or incorporate biases in primary studies but risk of bias assessment was done and as this was an IPD meta-analysis, between study heterogeneity was less relevant).</p> <p>Overall: low risk of bias</p>

Study details	Participants	Factors	Results	Comments
<p>Funding</p> <p>Funding for this research was provided by a Trans-Tasman Research Funding Grant by Cure Kids and Red Nose Australia in 2016 (grant 6601).</p>	<p>Birth weight and birth weight centiles were compared by maternal going-to-sleep position and adjusted for infant gestational age at birth and at time of interview, infant sex, and maternal age, height, weight, parity, ethnicity, preexisting diabetes, preexisting hypertension, antepartum hemorrhage, gestational hypertensive disorder, gestational diabetes, cigarette smoking, and recreational drug use. To account for possible study differences, multivariable analyses were also adjusted for individual studies as a covariate. For continuous outcomes (birth weight and birth weight centiles), a generalized linear model was used with predicted adjusted means obtained using least-squares means. For binary outcomes (birth weight centile <10th, <50th, and >90th) logistic regression was used, stratified by study, and aORs and 95% confidence intervals were reported.</p> <p>Demographics</p> <p>Age years Supine: 29.6 (5.5) years; Nonsupine: 30.3 (5.5) years</p> <p>Ethnicity: White: Supine 31 (2.8%); Nonsupine 1,074 (97.2%)</p> <p>Black: Supine 35 (97.2%); Nonsupine 1 (2.8%)</p> <p>South Asian: Supine 9 (4.3%); Nonsupine 202 (95.7%)</p>			

Study details	Participants	Factors	Results	Comments
	<p>South East/East Asian: Supine 5 (4.6); Nonsupine 104 (95.4)</p> <p>Maori: Supine 2 (1.9); Nonsupine 104 (98.1)</p> <p>Pacific: Supine 9 (5.9); Nonsupine 143 (94.1)</p> <p>Others: Supine 0 (0%); Nonsupine 41 (100%)</p> <p>Parity: 0: 19 (2.5%) supine; 749 (97.5%) nonsupine 1: 32 (5.0%) supine; 604 (95.0%) nonsupine ≥2: 6 (1.7%); 350 (98.3%)</p> <p>Education: Primary and/or secondary school: 26 (4.5%) supine; 554 (95.5%) nonsupine Trade school: 6 (2.7%) 220 (97.3%) nonsupine Tertiary (university and postgraduate): 25 (2.6%); 929 (97.4%)</p> <p>Earliest pregnancy BMI (kg/m²): 24.0 (21.0, 28.7) supine; 24.6 (22.0, 29.0) nonsupine</p>			
<p>Full citation</p> <p>Cronin RS, Li M, Thompson JMD, Gordon A, Raynes-Greenow CH, Heazell AEP, Stacey T, Culling VM, Bowring V, Anderson NH, O'Brien LM,</p>	<p>6 eligible case control studies; 5 provided individual level data (note the study that did not provide data, Lakshmi 2017 – did not meet the NGA protocol criteria for inclusion). Eligible</p>	<p>Adjusted for:</p> <ul style="list-style-type: none"> • Maternal age • Earliest pregnancy BMI • Ethnicity 	<p>Going to sleep position (last recorded, within 2 weeks of estimated fetal death in cases) and the odds of late stillbirth</p>	<p>Limitations</p> <p>ROBIS</p> <p>No concerns over relevance</p>

Study details	Participants	Factors	Results	Comments
<p>Mitchell EA, Askie LM, McCowan LME. An Individual Participant Data Meta-analysis of Maternal Going-to-Sleep Position, Interactions with Fetal Vulnerability, and the Risk of Late Stillbirth. The Lancet 2019; 10:49-57.</p> <p>Ref Id</p> <p>1242569</p> <p>Country/ies where the study was carried out</p> <p>NA</p> <p>Study type</p> <p>Systematic Review</p> <p>Study dates</p> <p>Searches to January 2018</p> <p>Consecutive recruitment</p> <p>No</p> <p>Funding</p> <p>Funding reported as funding for the individual studies included in the review from NHMRC Career Development Fellowship #1087062; NHMRC Early Career Fellowship #1089898; Cure Kid; American Sleep</p>	<p>participants comprised in 851 cases and 2,257 controls</p> <p>Inclusion criteria</p> <p>Eligible participants (stillbirth cases and pregnant controls) were extracted from the identified studies that provided maternal going-to-sleep position and late stillbirth data.</p> <p>Exclusion criteria</p> <p>Participant level exclusion criteria (multiple pregnancy, major congenital abnormality, gestation <28 weeks' when going-to-sleep position data during pregnancy was collected, termination of pregnancy at ≥28 weeks', and receiving a study intervention that may have affected going-to-sleep position) were applied during the analysis.</p> <p>Statistical method</p> <p>Potential confounders were all prespecified (maternal age, earliest pregnancy body mass index (BMI kg/m²), ethnicity, parity, education level, marital status, pre-existing hypertension or diabetes, smoking, recreational drug use, going-to-sleep position, fetal movements, and infant birthweight by customised centiles).</p> <p>A one-stage approach to IPD meta-analysis was used, so that the data from all the eligible studies were included in a single model. Logistic</p>	<ul style="list-style-type: none"> • Parity • Education level • Marital status • Obesity • Pre-existing hypertension or diabetes • Smoking • Recreational drug use • Fetal movements • Infant birthweight by customised centiles • Small for gestational age infant (<10th infant birthweight customised centile) • Term (≥37 weeks) vs preterm (<37 weeks) infant <p>Three confounders (going-to-sleep duration, frequency of overnight toilet use, and day-time napping) analysed in sensitivity models</p>	<p>Reference group was left-side going-to-sleep position.</p> <p>Adjusted OR (95% CI)</p> <p>Supine: 2.63 (1.72, 4.04) Right side: 1.04 (0.83, 1.31) Prone: 0.63 (0.12, 3.25) Variable sides: 0.97 (0.70, 1.35) Propped up: 1.30 (0.68, 2.49) Don't remember: 2.26 (1.48, 3.46)</p>	<p>Study eligibility criteria: low concern</p> <p>Identification and selection of studies: low concern</p> <p>Data collection and study appraisal: low concern (although not all eligible studies were included this was due to a lack of data appropriate for IPD meta-analysis and researchers made reasonable efforts to contact authors)</p> <p>Synthesis and findings: low concern (no formal efforts to consider between study variation or incorporate biases in primary studies but risk of bias assessment was done and as this was an IPD meta-analysis, between study heterogeneity was less relevant).</p> <p>Overall: low risk of bias</p>

Study details	Participants	Factors	Results	Comments
Medicine Foundation and ResMed	<p>regression models were used for the binary outcome. A fixed study effect and study site effect were included in the model specification as strata. Univariable analysis was performed to evaluate the association between going-to-sleep position and the odds of late stillbirth. During data acquisition, one prespecified confounder, alcohol intake during pregnancy, was found to be inconsistently collected across the studies and unable to be merged, and consequently omitted from the analysis. A multivariable model was developed incorporating prespecified confounders available in all the studies.</p> <p>Three confounders (going-to-sleep duration, frequency of overnight toilet use, and day-time napping) were only available in some of the studies, and were therefore analysed in sensitivity models. A sensitivity analysis was also conducted after exclusion of controls who reported their pregnancy going-to-sleep position after they had given birth. The interaction between going-to-sleep position and prespecified factors indicating a vulnerable pregnancy were assessed in bi-variable regression models. Estimates of risk of late stillbirth were reported as odds ratio (OR) with 95% confidence intervals.</p> <p>For missing data in each individual study, no imputation was undertaken. The population attributable risk (PAR) was calculated using the unadjusted OR for the primary outcome for supine going-to-sleep position and for other</p>			

Study details	Participants	Factors	Results	Comments
	<p>modifiable risk factors that were significant in multivariable analysis. Statistical analyses were performed using SAS, version 9.4 (SAS Institute Inc., Cary NC USA).</p> <p>Demographics</p> <p>Age years <20 yrs: 38 (4.5) cases; 78 (3.5) controls 20–29 yrs: 343 (40.3) cases; 905 (40.1) controls 30–39 yrs: 425 (49.9); 1190 (52.7) controls ≥40 yrs: 45 (5.3) cases; 84 (3.7) controls</p> <p>Ethnicity: White: 522 (61.3) cases; 1545 (68.5) controls Black: 22 (2.6) cases; 42 (1.9) controls South Asian: 90 (10.6) cases; 219 (9.7) controls South East/East Asian: 40 (4.7) cases; 111 (4.9) controls Maori: 46 (5.4) cases; 107 (4.7) controls Pacific: 91 (10.7) cases; 154 (6.8) controls Others: 40 (4.7) cases; 79 (3.5) controls</p> <p>Parity: Nulliparous: 446 (52.4) cases; 930 (41.2) controls 1–2: 292 (34.3) cases; 1110 (49.2) controls</p>			

Study details	Participants	Factors	Results	Comments
	<p>3-4: 87 (10.2) cases; 176 (7.8) controls ≥5: 26 (3.1) cases; 41 (1.8) controls</p> <p>Education: Primary: 187 (22.0) cases; 348 (15.4) controls Secondary: 161 (18.9) cases; 343 (15.2) controls University: 328 (38.5) cases; 1069 (47.4) Postgraduate: 73 (8.6) cases; 240 (10.6) controls Non-University Trade: 93 (10.9) cases; 249 (11.0) controls</p> <p>Earliest pregnancy BMI (kg/m²): 26.0 (22.5, 31.4) cases; 24.8 (22.0, 29.3) controls</p>			

Table 6: Clinical evidence tables for primary studies

Study details	Participants	Factors	Results	Comments
<p>Full citation</p> <p>Heazell, A. E. P., Li, M., Budd, J., Thompson, J. M. D., Stacey, T., Cronin, R. S., Martin, B., Roberts, D., Mitchell, E. A., McCowan, L. M. E., Association between maternal sleep practices and late stillbirth - findings from a stillbirth case-control study, BJOG: An International Journal of Obstetrics and Gynaecology, 125, 254-262, 2018</p>	<p>Cases</p> <p>N=291</p> <p>Diagnostic criteria</p> <p>ReCoDe classification system.</p> <p>Controls</p> <p>N=733</p> <p>Inclusion criteria</p>	<p>Factors</p> <p>Sleeping practices: Maternal going-to-sleep position in the last 4 weeks and on the night prior to stillbirth for cases and night before interview for controls (left side, supine, right side, tummy, variable side, propped up, or unknown); Duration of sleep;</p>	<p>Adjusted odds ratio</p> <p>Late stillbirth and supine going-to-sleep position on last night</p> <p>Data adjusted for: Last night going-to-sleep position, maternal age group, ethnicity, parity, level of education, BMI, birthweight centile, gestation, sleep duration, duration of daytime nap, study site. Reference group was left-side going-to-sleep position.</p>	<p>Limitations</p> <p>QUIPS tool - modified version</p> <p>Study population: High risk of bias (multicentre (41 maternity units); >50% for cases and controls did not participate, although authors reported that women in participation and non-participant groups had similar profiles of maternal age and ethnicity; sufficient information provided on flow of participants)</p> <p>Study attrition: Low risk of bias (>80% seen at assessment; reasons for non-completion provided)</p>

<p>Ref Id 936669</p> <p>Country/ies where the study was carried out UK</p> <p>Study type Prospective population-based case-control study</p> <p>Study dates April 2014 to March 2016</p> <p>Consecutive recruitment No</p> <p>Funding Funded by Action Medical Research, Cure Kids, and Sands.</p>	<ul style="list-style-type: none"> Cases: Women who had a stillbirth after 28 weeks' gestation with no known congenital anomaly; Controls: Women with an ongoing pregnancy at the time of interview. <p>Exclusion criteria</p> <ul style="list-style-type: none"> Women with multiple pregnancies; Controls: Women who subsequently delivered an infant with congenital abnormality or who had a stillborn baby; Maternal age <16 years; Women unable to provide consent. <p>Statistical method</p> <p>Power analysis To achieve 80% power and assuming 30% non-participation, 415 cases and 830 controls were required.</p> <p>Statistical analyses Differences between cases and controls for categorical data were analysed using chi-squared test. Continuous variables were compared using Wilcoxon rank-sum tests as the data were not normally distributed. Univariable logistic regression was conducted to evaluate the association between sleep practices and risk of late stillbirth.</p>	<p>Number of times up to the toilet during the last night; Daytime napping in the last 4 weeks.</p> <p>Other factors: Small for gestational age (<10th centile); Smoking during pregnancy; Obesity; Overweight.</p> <p>Still birth - OR (95% CI) On last night before stillbirth for cases or night before interview for controls, women with late stillbirth more likely to report sleeping in supine position on last night before stillbirth: 2.17 (1.15 to 4.08) Women with late stillbirth had increased likelihood of not being able to recall going-to-sleep position: 3.73 (1.67 to 8.32) Women with late stillbirth more likely to report right-side going-to-sleep position: 0.91 (0.65 to 1.26) SGA - OR (95% CI) 7.01 (33.6% to 56.8%) Sleep duration previous night (hours) - number (%) <5.49 Cases: 129 (44.3)</p>	<p>Maternal going-to-sleep position previous night (before stillbirth/interview) - number (%)</p> <p>Left Cases: 140 (48.1) Controls: 383 (53.3)</p> <p>Right Cases: 73 (25.1) Controls: 220 (30.0) Adjusted odds ratio (aOR)=0.67 (95% CI 0.44 to 1.02)</p> <p>Back Cases: 19 (6.5) Controls: 24 (3.3) aOR=2.31 (95% CI 1.04 to 5.11)</p> <p>Tummy Cases: 3 (1.0) Controls: 4 (0.5) aOR=1.01 (95% CI 0.13 to 7.81)</p> <p>Propped Cases: 9 (3.1) Controls: 15 (2.0) aOR=0.44 (95% CI 0.13 to 1.49)</p> <p>Variable Cases: 32 (11.0) Controls: 76 (10.4) aOR=0.93 (95% CI 0.51 to 1.69)</p> <p>Unknown Cases: 15 (5.2) Controls: 11 (1.5) aOR=3.33 (95% CI 1.13 to 9.84)</p> <p>c-statistic for final multivariable model 0.827 *Reference</p>	<p>Prognostic factor measurement: Moderate risk of bias (prospective data collection; definition of risk factors provided; validated measurement tool used, but potential for recall bias; adequate proportion of study sample completed data for prognostic factors; no imputation performed for missing data)</p> <p>Outcome measurement: Low risk of bias (validated measurement tool administered by research midwives and same for all participants; questionnaires included maternal health so blinding not possible)</p> <p>Confounding measurement and account: Low risk of bias (cases and controls matched and adjustments made for confounding variables)</p> <p>Analysis and reporting: Low risk of bias (statistical model appropriate and results reported in final multivariable model with point estimates and measures of variance)</p> <p>Other information</p> <p>Notes: 50 cases and 119 controls reported receiving advice about sleep; obtained from the internet, health professionals, literature, and friends and family.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>A multivariable logistic model was developed to incorporate ethnicity and level of education, variables associated with increased risk of stillbirth based on previous literature (age, BMI, parity, smoking, small-for-gestational-age (SGA) status), other sleep related variables significant in univariable analysis, and variables used to select cases and controls (gestation and maternity unit). Unconditional logistic regression was used to adjust for potential confounders. No imputation for missing data was undertaken. The c statistic was used to assess the area under the curve.</p> <p>Demographics</p> <p><u>Maternal age (years) - median (interquartile range; IQR)</u> Cases: 30.2 Controls: 30.5</p> <p><u>Ethnicity - number (%)</u></p> <p><u>White</u> Cases: 234 (80.4) Controls: 594 (81.0)</p> <p><u>Black</u> Cases: 12 (4.1) Controls: 29 (4.0)</p> <p><u>South Asian</u> Cases: 39 (13.4) Controls: 95 (13.0)</p> <p><u>Others</u> Cases: 6 (2.1) Controls: 15 (2.0)</p> <p><u>Parity - number (%)</u></p> <p><u>0</u> Cases: 167 (57.4) Controls: 296 (40.4)</p> <p><u>1 to 2</u> Cases: 92 (31.6)</p>	<p>Controls: 227 (31.0) <u>5.5 to 8.49</u> Cases: 121 (41.6) Controls: 413 (56.3) <u>8.5 to 9.49</u> Cases: 20 (6.9) Controls: 55 (7.5) <u>9.5+</u> Cases: 19 (6.5) Controls: 36 (4.9) <u>Number of times up to the toilet previous night - number (%)</u> <u>One or less</u> Cases: 91 (31.3) Controls: 120 (16.4) <u>Two or more</u> Cases: 199 (68.4) Controls: 613 (83.6) <u>Maternal daytime naps in previous 4 weeks - number (%)</u> <u>Never</u> Cases: 58 (19.9) Controls: 157 (21.4) <u>Occasionally</u> Cases: 49 (16.8) Controls: 153 (20.9) <u>1 to 2 per week</u> Cases: 47 (16.1) Controls: 180 (24.6) <u>3 to 4 per week</u> Cases: 44 (15.1) Controls: 110 (15.0) <u>5 to 6 per week</u> Cases: 22 (7.6) Controls: 39 (5.3) <u>Everyday</u> Cases: 71 (24.4) Controls: 93 (12.7) <u>Unknown</u> Cases: 0 (0.0)</p>		
--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--

	<p>Controls: 386 (52.7) 3+ Cases: 32 (11.0) Controls: 51 (7.0) <u>Level of Education - number (%)</u> <u>Graduate Education</u> Cases: 99 (34.0) Controls: 326 (31.84) <u>Further Education</u> Cases: 112 (38.5) Controls: 278 (27.15) <u>Secondary education to 16 years</u> Cases: 56 (19.2) Controls: 100 (9.77) <u>No formal educational qualification</u> Cases: 23 (7.9) Controls: 29 (2.83) <u>Body mass index - mean (IQR)</u> Cases: 26.91 (15.44 to 47.87) Controls: 26.02 (15.41 to 48.59) <u>Gestational age (for cases, gestational age at diagnosis of stillbirth; for controls at time of interview) - median (IQR)</u> Cases: 37 weeks 4 days (33 weeks 4 days to 39 weeks 5 days) Controls: 36 weeks 3 days (32 weeks 6 days to 38 weeks 5 days)</p>	<p>Controls: 1 (0.1)</p>		
<p>Full citation</p> <p>Stacey, T., Thompson, J. M. D., Mitchell, E. A., Ekeroma, A. J., Zuccollo, J. M., McCowan, L. M. E., Association between maternal sleep practices and risk of late stillbirth: A case-control study, <i>Bmj</i>, 342 (7811) (no pagination), 2011</p> <p>Ref Id</p>	<p>Cases</p> <p>N=155</p> <p>Diagnostic criteria</p> <p>PSANZ classification system</p> <p>Controls</p> <p>N=310</p> <p>Inclusion criteria</p>	<p>Factors</p> <p>Sleeping practices: Maternal sleep position at the time of going to sleep and on waking (left side, right side, back, and other (front, sitting up, both sides, and unsure or don't remember)) in the last month, week, and night of pregnancy;</p>	<p>Adjusted odds ratio</p> <p><u>Maternal sleeping position in last night of pregnancy and risk of late stillbirth</u> Results adjusted for age, ethnicity, overweight or obesity, parity, social deprivation level, smoking, regular sleep in daytime in last month of pregnancy; hours of night time sleep in last month of pregnancy; number of times up</p>	<p>Limitations</p> <p>QUIPS tool - modified version Study population: Moderate risk of bias (multicentre (all maternity units in Auckland region); recruitment rate 72% for cases and controls and although no significant differences in age, parity, or ethnicity between those who did and did not consent, there was potential for selection bias; sufficient information provided on flow of participants) Study attrition: Low risk of bias (>80% seen at assessment; authors reported</p>

<p>936670</p> <p>Country/ies where the study was carried out</p> <p>New Zealand</p> <p>Study type</p> <p>Prospective population-based case-control study</p> <p>Study dates</p> <p>July 2006 to June 2009</p> <p>Consecutive recruitment</p> <p>No</p> <p>Funding</p> <p>Cure Kids, the Nurture Foundation, and the Auckland District Health Board Trust Fund.</p>	<ul style="list-style-type: none"> • Women who gave birth to stillborn baby at or after 28 weeks of gestation in the Auckland region; • Controls were selected from the pregnancy registration list of the district health board, matched for gestation to cases. <p>Exclusion criteria</p> <ul style="list-style-type: none"> • Women whose baby died from a congenital abnormality; • Multiple pregnancies; • Women who had not been booked to deliver their baby within the Auckland region. <p>Statistical method</p> <p>Power analysis</p> <p>The authors reported that the study was powered to detect an odds ratio of 2 with 80% power and significance level of 5%, with a prevalence of the risk factor of $\geq 20\%$ in the control population.</p> <p>Statistical analyses</p> <p>Continuous data were compared using Student's <i>t</i>-test, and the Pearson correlation coefficient was used to assess the correlation between variables.</p> <p>A multivariable regression model was used to include maternal variables reported to be associated with increased risk of stillbirth, based on previous literature (age, BMI, ethnicity, parity, smoking, and socioeconomic status).</p>	<p>Sleeping regularly during the daytime in the last month; Usual duration of sleep at night during the last month; Frequency of getting up to the toilet. <u>Changes in sleeping position on last night of pregnancy and risk of late stillbirth - number (%)</u> <u>Left on going to sleep; left on waking up</u> Cases: 29 (19) Controls: 95 (31) Univariable OR (95% CI): 1.00 <u>Left on going to sleep; other on waking up</u> Cases: 13 (8) Controls: 37 (12) Univariable OR (95% CI): 1.15 (0.54 to 2.45) <u>Other on going to sleep; left on waking up</u> Cases: 2 (1) Controls: 11 (4) Univariable OR (95% CI): 0.60 (0.13 to 2.84) <u>Other on going to sleep; other on waking up</u> Cases: 111 (72) Controls: 167 (54) Univariable OR (95% CI): 2.28 (1.35 to 3.52) <u>Regular sleep in daytime (last month of</u></p>	<p>to the toilet during last night of pregnancy. Left side sleeping position used as reference group, OR=1.00 <u>Right side</u> adjusted odds ratio (aOR)=1.74 (95% CI 0.98 to 3.01)*** <u>Back (supine)</u> aOR=2.54 (95% CI 1.04 to 6.18);p=0.005 <u>Other</u> aOR=2.32 (95% CI 1.28 to 4.19)</p>	<p>that there was no missing data for variables included in the paper) Prognostic factor measurement: Moderate risk of bias (prospective data collection; definition of risk factors provided; validated measurement tool used, but potential for recall bias due to length of time between stillbirth and interview, 25 days on average, compared with controls who were asked about sleep practices on the previous night; adequate proportion of study sample completed data for prognostic factors; no missing data) Outcome measurement: Low risk of bias (cases selected from clinicians in participating centres and hospital birth records, with confirmation through use of New Zealand national registry; controls selected from pregnancy registration list of district health board of participating centre). The authors stated that it was not always possible to be certain as to the exact timing of fetal death, and in some cases the 'last night' was not the final night before fetal death or during which the baby died. Confounding measurement and account: Low risk of bias (cases and controls matched and adjustments made for confounding variables) Analysis and reporting: Low risk of bias (statistical model appropriate and results reported in final multivariable model with point estimates and measures of variance)</p> <p>Other information</p> <p>Auckland Stillbirth Study</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>Chi-squared statistics were used to assess the significance of variables in the models, and individual level odds ratios were estimated for each category and compared to a reference category.</p> <p>Demographics</p> <p><u>Maternal age (years) - number (%)</u></p> <p><u>≤20</u> Cases: 10 (6) Controls: 24 (8)</p> <p><u>20 to 34</u> Cases: 113 (73) Controls: 216 (70)</p> <p><u>≥35</u> Cases: 32 (21) Controls: 70 (23)</p> <p><u>Ethnicity - number (%)</u></p> <p><u>Maori</u> Cases: 19 (12) Controls: 46 (15)</p> <p><u>Pacific</u> Cases: 48 (31) Controls: 67 (22)</p> <p><u>European</u> Cases: 55 (35) Controls: 139 (45)</p> <p><u>Other</u> Cases: 33 (21) Controls: 58 (19)</p> <p><u>Parity - number (%)</u></p> <p><u>0</u> Cases: 77 (50) Controls: 144 (46)</p> <p><u>1 to 3</u> Cases: 62 (40) Controls: 156 (51)</p> <p><u>≥4</u> Cases: 16 (10) Controls: 10 (3)</p> <p><u>Social deprivation level - number (%)</u></p>	<p><u>pregnancy - number (%)</u></p> <p><u>Yes</u> Cases: 78 (50) Controls: 116 (37)</p> <p><u>No</u> Cases: 77 (50) Controls: 194 (63)</p> <p><u>Hours of nighttime sleep (last month of pregnancy) - number (%)</u></p> <p><u><6</u> Cases: 30 (19) Controls: 46 (15)</p> <p><u>6 to 8</u> Cases: 82 (53) Controls: 205 (66)</p> <p><u>>8</u> Cases: 43 (28) Controls: 59 (19)</p> <p><u>Number of times getting up to toilet during night last night of pregnancy- number (%)</u></p> <p><u>>1</u> Cases: 86 (55) Controls: 207 (67)</p> <p><u>≤1</u> Cases: 69 (45) Controls: 103 (33)</p>		
--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--

	<p><u>1 to 4</u> Cases: 91 (59) Controls: 218 (70)</p> <p><u>5 (most deprived)</u> Cases: 64 (41) Controls: 92 (30)</p> <p><u>BMI at booking - number (%)</u></p> <p><u><25</u> Cases: 55 (35) Controls: 156 (50)</p> <p><u>25 to 29.9</u> Cases: 39 (25) Controls: 67 (22)</p> <p><u>≥30</u> Cases: 61 (39) Controls: 87 (28)</p> <p><u>Smoking during pregnancy - number (%)</u> Cases: 46 (30) Controls: 66 (21)</p>			
<p>Full citation</p> <p>McCowan, L. M. E., Thompson, J. M. D., Cronin, R. S., Li, M., Stacey, T., Stone, P. R., Lawton, B. A., Ekeroma, A. J., Mitchell, E. A., Going to sleep in the supine position is a modifiable risk factor for late pregnancy stillbirth; Findings from the New Zealand multicentre stillbirth case-control study, Plos one, 12 (6) (no pagination), 2017</p> <p>Ref Id</p> <p>929927</p> <p>Country/ies where the study was carried out</p>	<p>Cases</p> <p>N=164</p> <p>Diagnostic criteria</p> <p>PSANZ classification system</p> <p>Controls</p> <p>N=569</p> <p>Inclusion criteria</p> <ul style="list-style-type: none"> • Cases: consenting women with a stillbirth at ≥28 weeks of gestation; • Controls: women with ongoing pregnancies in each participating health region. 	<p>Factors</p> <p>Maternal sleeping practices: Self-reported going-to-sleep position on last night (left side, right side, restless, supine (lying on the back), on the front, or propped); Self-reported usual going-to-sleep position in last week (left side, right side, variable side, supine (lying on the back), on the front or propped); Getting up to go to the toilet during the night on the last night; How many hours slept on last night;</p>	<p>Adjusted odds ratio</p> <p>Results adjusted for gestation at interview in controls and at diagnosis of stillbirth for cases, district health board, maternal age, ethnicity, parity, social deprivation level, earliest pregnancy BMI, marital status, smoking in pregnancy, baby birthweight centile, hours of night time sleep on the last night, getting up to toilet during the last night, sleep during the daytime in the last week, going to sleep position on last night and in the last week). <u>Going-to-sleep position on last night and pre-term stillbirth (≥28 to 36 weeks' gestation)</u> Note: One pre-term case and one pre-term control had an unknown sleep position on the</p>	<p>Limitations</p> <p>QUIPS tool - modified version Study population: High risk of bias (multicentre (all maternity units across 7 New Zealand health regions); rate of recruitment 65.9% for cases and 62.2% for controls, further information on flow of participants not provided; women of high parity were underrepresented in both groups, while Indian women were over-represented and Maori women under-represented in cases compared with eligible controls) Study attrition: Low risk of bias (>80% seen at assessment; no imputation performed for missing data) Prognostic factor measurement: Moderate risk of bias (prospective data collection; definition of risk factors provided; validated measurement tool used, but potential for recall bias due to length of time between</p>

<p>New Zealand</p> <p>Study type</p> <p>Prospective population-based case-control study</p> <p>Study dates</p> <p>February 2012 to December 2015</p> <p>Consecutive recruitment</p> <p>No</p> <p>Funding</p> <p>Health Research Council of New Zealand, Cure Kids, Mercia Barnes Trust, Nurture Foundation, and the University of Auckland Faculty Research Development Fund.</p>	<p>Exclusion criteria</p> <ul style="list-style-type: none"> • Women with multiple pregnancies; • Babies with major congenital abnormalities at any stage of the study. <p>Statistical method</p> <p>Power analysis</p> <p>To achieve 80% power and based on 70% participation, 415 cases and 830 controls were required for recruitment.</p> <p>Statistical analyses</p> <p>Chi-squared tests were used to compare differences between categorical data. Continuous data were compared using Wilcoxon rank-sum tests. Univariable analysis was conducted to assess the association between sleep practices and risk of late stillbirth. A multivariable model was used to incorporate ethnicity and deprivation index, variables associated with increased risk of stillbirth based on previous literature (age, BMI, parity, smoking, SGA status), other sleep variables significant in univariable analysis, and variables used to select cases and controls (gestation and District Health Boards). Unconditional logistic regression was used to adjust for potential confounders. No imputation was performed for missing data (women who could not recall their going-to-sleep position on the last night were excluded from the multivariable model). Stratified analysis was carried out by term (≥ 37 weeks') and pre-term (≥ 28 to</p>	<p>Frequency of sleeping during the daytime in the last week.</p> <p><u>Going to sleep position on the last night - number (%)#</u></p> <p><u>Left side</u></p> <p>Cases: 78 (47.6)</p> <p>Controls: 328 (57.6)</p> <p>Adjusted OR (95% CI): 1.00</p> <p><u>Right side</u></p> <p>Cases: 44 (26.8)</p> <p>Controls: 187 (32.9)</p> <p>Adjusted OR (95% CI): 0.92 (0.58 to 1.44)</p> <p><u>Restless</u></p> <p>Cases: 14 (8.5)</p> <p>Controls: 22 (3.9)</p> <p>Adjusted OR (95% CI): 1.98 (0.87 to 4.50)</p> <p><u>Supine</u></p> <p>Cases: 19 (11.6)</p> <p>Controls: 22 (3.9)</p> <p>Adjusted OR (95% CI): 3.67 (1.74 to 7.78)</p> <p><u>Propped</u></p> <p>Cases: 4 (2.4)</p> <p>Controls: 9 (1.6)</p> <p>Adjusted OR (95% CI): 1.11 (0.30 to 4.06)</p> <p><u>Going to sleep position on the last week - number (%)##</u></p> <p><u>Left side</u></p> <p>Cases: 85 (51.8)</p> <p>Controls: 302 (53.1)</p> <p>Adjusted OR (95% CI): -</p> <p><u>Right side</u></p> <p>Cases: 42 (25.6)</p>	<p>last night and was excluded from the multivariable model.</p> <p><u>Left side (reference group)</u></p> <p>Cases, n (%): 34 (50.0)</p> <p>Controls, n (%): 147 (58.3)</p> <p>Odds ratio: 1.00</p> <p><u>Right side</u></p> <p>Cases: 22 (32.4)</p> <p>Controls: 86 (34.1)</p> <p>Adjusted odds ratio (aOR)=0.96 (95% CI 0.48 to 1.94)</p> <p><u>Restless</u></p> <p>Cases: 4 (5.9)</p> <p>Controls: 4 (1.6)</p> <p>aOR=3.50 (95% CI 0.61 to 19.97)</p> <p><u>Back (Supine)</u></p> <p>Cases: 6 (8.8)</p> <p>Controls: 13 (5.2)</p> <p>aOR=3.12 (95% CI 0.97 to 10.05)</p> <p><u>Propped</u></p> <p>Cases: 1 (1.5)</p> <p>Controls: 1 (0.4)</p> <p>aOR=4.37 (95% CI 0.11 to 178.86)</p> <p><u>Going-to-sleep position on last week and pre-term stillbirth (≥ 28 to 36 weeks' gestation)</u></p> <p>Note: One pre-term control did not recall their sleep position in last week and was excluded from the multivariable model.</p> <p><u>Left side (reference group)</u></p> <p>Cases, n (%): 38 (55.9)</p> <p>Controls, n (%): 137 (54.4)</p> <p>OR: 1.00</p> <p><u>Right side</u></p> <p>Cases: 19 (27.9)</p> <p>Controls: 75 (29.8)</p> <p>aOR=0.73 (95% CI 0.34 to 1.54)</p>	<p>stillbirth and interview, 24 days on average; adequate proportion of study sample completed data for prognostic factors)</p> <p>Outcome measurement: Low risk of bias (clear definition of outcome provided, validated measurement tool administered by research midwives, setting and measurement similar for all participants; authors state that possible in some cases 'last night' was not the night before fetal death, or the night during which the baby died.)</p> <p>Confounding measurement and account: Low risk of bias (cases and controls matched and adjustments made for confounding variables)</p> <p>Analysis and reporting: Low risk of bias (statistical model appropriate and results reported in final multivariable model with point estimates and measures of variance)</p> <p>Other information</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>36 weeks') gestation. The c statistic was calculated to assess model performance.</p> <p>Demographics</p> <p><u>Maternal age (years) - number (%)</u></p> <p><u><20</u> Cases: 9 (5.5) Controls: 17 (3.0)</p> <p><u>20 to 39</u> Cases: 141 (86.0) Controls: 532 (93.5)</p> <p><u>≥40</u> Cases: 14 (8.5) Controls: 20 (3.5)</p> <p><u>Ethnicity - number (%)</u></p> <p><u>Maori</u> Cases: 26 (16.0) Controls: 58 (10.0)</p> <p><u>Pacific</u> Cases: 38 (23.2) Controls: 86 (15.1)</p> <p><u>Indian</u> Cases: 17 (10.4) Controls: 77 (13.5)</p> <p><u>Other Asian</u> Cases: 13 (7.9) Controls: 72 (12.7)</p> <p><u>European</u> Cases: 65 (39.6) Controls: 263 (46.2)</p> <p><u>Other</u> Cases: 5 (3.1) Controls: 13 (2.3)</p> <p><u>Parity - number (%)</u></p> <p><u>0</u> Cases: 76 (46.3) Controls: 245 (43.1)</p> <p><u>1 to 3</u> Cases: 80 (48.8) Controls: 308 (54.1)</p> <p><u>≥4</u></p>	<p>Controls: 171 (30.1) Adjusted OR (95% CI): 0.82 (0.52 to 1.30)</p> <p><u>Variable side</u> Cases: 15 (9.2) Controls: 72 (12.7) Adjusted OR (95% CI): 0.85 (0.44 to 1.65)</p> <p><u>Supine</u> Cases: 15 (9.2) Controls: 72 (12.7) Adjusted OR (95% CI): 3.46 (1.49 to 8.03)</p> <p><u>Propped</u> Cases: 5 (3.1) Controls: 7 (1.2) Adjusted OR (95% CI): 2.10 (0.57 to 7.71)</p> <p><u>On front</u> Cases: 1 (0.6) Controls: 0</p>	<p><u>Variable side</u> Cases: 4 (5.9) Controls: 27 (10.7) aOR=0.63 (95% CI 0.18 to 2.19)</p> <p><u>Supine</u> Cases: 5 (7.4) Controls: 11 (4.4) aOR=2.25 (95% CI 0.65 to 7.84)</p> <p><u>Propped</u> Cases: 1 (1.5) Controls: 1 (0.4) aOR=4.01 (95% CI 0.08 to 210.43)</p> <p><u>On front</u> Cases: 1 (1.5) Controls: 0</p> <p><u>Going-to-sleep position on last night and term stillbirth (≥37 weeks' gestation)</u> Note: Four term cases had an unknown sleep position on the last night and was excluded from the multivariable model.</p> <p><u>Left side</u> Cases, n (%): 44 (45.8) Controls, n (%): 181 (57.1) OR=1.00</p> <p><u>Right side</u> Cases: 22 (22.9) Controls: 101 (31.9) aOR=0.98 (95% CI 0.48 to 1.99)</p> <p><u>Restless</u> Cases: 10 (10.4) Controls: 18 (5.7) aOR=2.00 (95% CI 0.64 to 6.21)</p> <p><u>Back (Supine)</u> Cases: 13 (13.5) Controls: 9 (2.8)</p>	
--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

	<p>Cases: 8 (4.9) Controls: 16 (2.8)</p>		<p>aOR=10.26 (95% CI 3.01 to 35.04) <u>Propped</u> Cases: 3 (3.1) Controls: 8 (2.5) aOR=1.02 (95% CI 0.17 to 5.97) <u>Going-to-sleep position on last week and term stillbirth (≥37 weeks' gestation)</u> <u>Note: One term case did not recall sleep position in the last week and was excluded from the multivariable model.</u> <u>Left side (reference group)</u> Cases, n (%): 47 (49.0) Controls, n (%): 165 (52.1) OR=1.00 <u>Right side</u> Cases: 23 (24.0) Controls: 96 (30.3) aOR=0.95 (95% CI 0.48 to 1.89) <u>Variable side</u> Cases: 11 (11.5) Controls: 45 (14.2) aOR=1.11 (95% CI 0.49 to 3.01) <u>Back (Supine)</u> Cases: 10 (10.4) Controls: 5 (1.6) aOR=12.73 (95% CI 2.92 to 55.46) <u>Propped</u> Cases: 4 (4.2) Controls: 6 (1.9) aOR=2.64 (95% CI 0.47 to 14.81) <u>c statistic for final multivariable model</u> 0.736</p>	
--	----------------------------------------------	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

			Non-left versus left positions in the multivariable model resulted in non-significant increase in late stillbirth risk compared with the combined non-left positions: 1.35 (0.92 to 1.99).	
Full citation	Cases	Factors	Adjusted odds ratio	Limitations
Gordon, A., Raynes-Greenow, C., Bond, D., Morris, J., Rawlinson, W., Jeffery, H., Sleep position, fetal growth restriction, and late-pregnancy stillbirth: The Sydney stillbirth study, <i>Obstetrics and Gynecology</i> , 125, 347-355, 2015	N=103	Maternal sleeping practices: Sleep position: left; right; back; other.	<u>Stillbirth and supine sleep position in the last month</u> Multivariate model adjusted for maternal age group; maternal BMI; primiparous; not in paid work; sleep apnoea symptoms; smoking; suspected fetal growth restriction; education to high school or less; sleep position (left, right, back, other). Left-side going-to-sleep position is reference group.	QUIPS tool - modified version Study population: Low risk of bias (multicentre (9 hospitals in the Sydney metropolitan area); rate of recruitment 86% for cases and 84.6% for controls, further information on flow of participants provided) Study attrition: Low risk of bias (>80% seen at assessment; reasons for non-participation reported) Prognostic factor measurement: Moderate risk of bias (prospective data collection; definition of risk factors provided; interviewer-administered questionnaires used, but potential for recall bias due to time delay between recruitment and interview; adequate proportion of study sample completed data for prognostic factors) Outcome measurement: Low risk of bias (clear definition of outcome; Cases recruited using clinicians/research staff at participating hospitals and confirmed through perinatal mortality review committees of these hospitals; controls identified using hospital databases) Confounding measurement and account: Low risk of bias (cases and controls matched and adjustments made for confounding variables) Analysis and reporting: Moderate risk of bias (statistical model appropriate and results reported in final multivariable
Ref Id	Diagnostic criteria	Suspected fetal growth restriction: <10th percentile; <3rd percentile.	<u>Left side</u> Cases: 32 (31) Controls: 48 (25) Odds ratio=1	
938535	PSANZ classification system	Other factors: Maternal age; Maternal BMI; Primiparous; Not in paid work; Sleep apnoea symptoms; Smoking; Education to high school or less.	<u>Right side</u> Cases: 14 (13.6) Controls: 25 (13) Adjusted odds ratio (aOR)=1.1 (95% CI 0.43 to 2.6)	
Country/ies where the study was carried out	Controls	<u>SGA and stillbirth - OR (95% CI)</u> Fetuses who were stillborn were significantly more likely to be SGA using the 10th percentile: 3.8 (1.8 to 8.2) and less than the third percentile: 3.6 (1.2 to 10.9).	<u>Back</u> Cases: 10 (9.7) Controls: 4 (2.1) aOR=6.26 (95% CI 1.2 to 34)	
Australia	N=192		<u>Other</u> Cases: 47 (45.6) Controls: 115 (60) aOR=0.69 (95% CI 0.36 to 1.3)	
Study type	Inclusion criteria			
Prospective population-based case-control study	<ul style="list-style-type: none"> Cases: women with singleton pregnancies who experienced stillbirth at ≥ 32 weeks of gestation Controls: pregnant women at 32 weeks of gestation with singleton pregnancies who were matched for booking hospital and gestation (by estimated date of delivery) and recruited during the same period of time as women in the case group. 	Univariate analysis <u>Stillbirth and suspected fetal growth</u>		
Study dates	Exclusion criteria			
January 2006 to December 2011	<ul style="list-style-type: none"> Women identified as Aboriginal or Torres Strait Islander; 			
Consecutive recruitment				
No				

<p>Funding</p> <p>Stillbirth Foundation, Australia.</p>	<ul style="list-style-type: none"> Fetuses that had known lethal or chromosomal anomalies; Terminations of pregnancy. <p>Statistical method</p> <p>Power analysis To achieve 80% power, based on prevalence of 10% small for gestational age (SGA) to detect an odds ratio (OR) of 25 between cases and controls, approximately 100 women were required in the case group and 200 women in the controls group.</p> <p>Statistical analyses Univariate analysis was conducted using chi-squared tests for categorical data and Student's t-test for continuous variables. Conditional logistic regression was used to calculate adjusted ORs for a priori-specified risk factors and to account for matching within stratification. Risk factors identified as significant on univariate analysis, or associated with stillbirth in previous literature (even if non-significant), were included in the multivariate models. However, if previously known risk factors were present in too few patients as to make no difference to the multivariate model, they were not included. Reference categories for the multivariable models were defined as the groups likely to have the lowest risk.</p> <p>Demographics</p> <p>Maternal age (years) - number (%) <u><35</u> Cases: 73 (70.9)</p>	<p><u>restriction - OR (95% CI)</u> 8.3 (2.3 to 30)</p> <p><u>Stillbirth and supine sleeping over the last month - OR (95% CI)</u> 5.0 (1.5 to 16.5)</p>		<p>model with point estimates and measures of variance; however, study underpowered to assess interactions between risk factors)</p> <p>Other information</p>
----------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	----------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>Controls: 121 (63) <u>35 to 39</u> Cases: 22 (21.4) Controls: 53 (27.6) <u>≥40</u> Cases: 8 (9.4) Controls: 18 (7.8) BMI (kg/m²) - number (%) <u><25</u> Cases: 62 (62.6) Controls: 129 (67.9) <u>25 to 29.9</u> Cases: 22 (22.2) Controls: 44 (23.2) <u>≥30</u> Cases: 15 (15.2) Controls: 17 (8.9) Primiparous - number (%) Cases: 53 (51.5) Controls: 104 (54.2) Not in paid work - number (%) Cases: 26 (25.2) Controls: 18 (9.4) Smoker - number (%) Cases: 14 (13.6) Controls: 25 (13) Education to high school or less - number (%) Cases: 43 (41.7) Controls: 49 (25.5)</p>			
Full citation	Cases	Factors	Adjusted odds ratio	Limitations
O'Brien, Louise M., Warland, Jane, Stacey, Tomasina, Heazell, Alexander E. P., Mitchell, Edwin A., Maternal sleep practices and stillbirth: Findings from an international case-control study, BirthBirth, 0	N=153	Sleeping practices: Maternal going-to-sleep position in the month and on the night prior to stillbirth for cases and last month and last night before their pregnancy for controls (left side, supine, right side,	Late stillbirth and supine going-to-sleep position on last night <u>Maternal going-to-sleep position previous night (before stillbirth/interview) - number (%)</u> Left-hand going-to-sleep position used as reference group. <u>Left</u>	QUIPS tool - modified version Study population: High risk of bias (international retrospective online survey so potential for self-selection bias) Study attrition: Low risk of bias (<20% missing data; prior sample size calculation which was fulfilled) Prognostic factor measurement: Moderate risk of bias (retrospective data collection; definition
Ref Id	Controls			
	N=480			

<p>967090</p> <p>Country/ies where the study was carried out</p> <p>Various (see other information)</p> <p>Study type</p> <p>Nested case-control study in uncontrolled cohort</p> <p>Study dates</p> <p>September 2012 to August 2014</p> <p>Consecutive recruitment</p> <p>No</p> <p>Funding</p> <p>Not reported</p>	<p>Inclusion criteria</p> <ul style="list-style-type: none"> Cases: Women who had a singleton stillborn baby \geq 28 weeks gestation within 1 month of completing the survey; Controls: Women with an ongoing pregnancy (\geq 28 weeks gestation) or had delivered a living baby within the month before survey completion <p>Exclusion criteria</p> <ul style="list-style-type: none"> Women with multiple pregnancies; Women whose fetus with known congenital abnormality ; Maternal age $<$18 years; Women unable to provide consent. <p>Statistical method</p> <p>Power analysis</p> <p>Sample size was calculated based on the anticipated exposure of supine sleep. To achieve 80% power and assuming an exposure frequency of 20%, 144 cases and controls were required to detect the odds of stillbirth 3.0 among cases compared to controls.</p> <p>Statistical analyses</p> <p>Data were cleaned by two authors. Data analyses were performed using SPSS using cross-tabulations, chi-squared tests and logistic regressions to find unadjusted and adjusted odds ratio with</p>	<p>tummy, variable side, propped up); Duration of sleep; Number of times up to the toilet during the last night; Daytime napping in the last 4 weeks.</p> <p>Other factors: Small for gestational age ($<$10th centile); Smoking during pregnancy; Obesity; Overweight. <u>Still birth - OR (95% CI)</u> On last night before stillbirth for cases or night before interview for controls, women with late stillbirth more likely to report sleeping in supine position on last night before stillbirth: 2.17 (1.15 to 4.08) Women with late stillbirth had increased likelihood of not being able to recall going-to-sleep position: 3.73 (1.67 to 8.32) Women with late stillbirth more likely to report right-side going-to-sleep position: 0.91 (0.65 to 1.26) <u>SGA - OR (95% CI)</u> 7.01 (33.6% to 56.8%)</p>	<p>Cases: 75 (49) Controls: 193 (40.2) Odds ratio=1</p> <p><u>Right</u> Cases: 45 (29.4) Controls: 111 (23.1) Adjusted odds ratio (aOR)=1.11 (95% CI 0.70 to 1.77)</p> <p><u>Back (Supine)</u> Cases: 4 (2.6) Controls: 11 (2.3) aOR=1.11 (95% CI 0.70 to 1.77)</p> <p><u>Tummy</u> Cases: 0 (0) Controls: 4 (0.8)</p> <p><u>Propped</u> Cases: 4 (2.6) Controls: 15 (3.1) aOR=0.71 (95% CI 0.22 to 2.30)</p> <p><u>Variable</u> Cases: 10 (6.5) Controls: 39 (8.1) aOR=0.75 (95% CI 0.34 to 1.64)</p>	<p>of risk factors provided; validated measurement tool used, but potential for recall bias; adequate proportion of study sample completed data for prognostic factors; no imputation performed for missing data) Outcome measurement: Moderate risk of bias (all outcomes self-reported via online form so potential for false reporting) Confounding measurement and account: High risk of bias (cases and controls not matched - women in case group were more likely to be non-Caucasians and nullip; but adjustments made for confounding variables) Analysis and reporting: Low risk of bias (statistical model appropriate and results reported in final multivariable model with point estimates and measures of variance)</p> <p>Other information</p> <p>Participants recruited using anonymous online survey 'Study of Trends and Risk Factors for Stillbirth' (STARS), which was developed during first Stillbirth Summit in Minneapolis in 2011 by international consortium of clinicians and academics, Star Legacy Foundation and other stillbirth/parental support groups. International online survey included respondents from following countries: Australia (n=21), Bahrain (n=1), Canada (n=44), Finland (n=1), Germany (n=2), Greece (n=2), India (n=2), Israel (n=1), Italy (n=1), New Zealand (n=2), Philippines (n=2), South Africa (n=2), Sweden (n=1), Switzerland (n=1), UK (n=95), USA (n=448).</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>95% confidence interval. Univariable logistic regression was conducted to evaluate the association between sleep practices and risk of late stillbirth. A multivariable logistic model was developed to incorporate maternal age, education level, smoking, body mass index, parity, country of respondent, ethnicity.</p> <p>Demographics</p> <p><u>Maternal age (years) - mean±standard deviation</u> Cases: 31 ± 5.4 Controls: 30±4.8</p> <p><u>Ethnicity - number (%)</u> <u>Caucasian</u> Cases: 122 (79.7)* Controls: 430 (90.2) <u>Non-Caucasian</u> Cases: 31 (20.3) Controls: 47 (9.8)</p> <p><u>Parity - median (interquartile range)</u> Cases: 1(0-6) Controls: 1(0-10)*</p> <p><u>Level of Education - number (%)</u> <u>Graduate Education</u> Cases: 35 (22.9) Controls: 129 (26.9) <u>College-level Education</u> Cases: 81 (52.9) Controls: 242 (50.4) <u>High school or lower</u> Cases: 37 (24.2) Controls: 106 (22.1)</p> <p><u>Body mass index - median (IQR)</u> Cases: 27 (23-32) Controls: 25 (23-31)</p>	<p><u>Sleep duration last month (hours) - number (%)</u> <u></=6</u> Cases: 15 (9.8) Controls: 47 (9.8) <u>6.5 to 8.5</u> Cases: 86 (56.2) Controls: 283 (59) <u>9.0+</u> Cases: 45(29.4) Controls: 79 (16.5)</p> <p><u>Number of times getting up previous month - number (%)</u> <u>One or less</u> Cases: 47 (30.7) Controls: 129 (26.9) <u>Two or more</u> Cases: 98(64.1) Controls: 270 (56.3)</p> <p><u>Maternal daytime naps in previous 4 weeks - number (%)</u> <u>Never</u> Cases: 38 (24.8) Controls: 125 (26) <u>Occasionally</u> Cases: 41 (26.8) Controls: 108 (22.5) <u>Often/almost always</u> Cases: 67 (43.8) Controls: 168 (35)</p> <p><u>Excessive daytime sleepiness last month</u> Cases: 42 (27.5) Controls: 107 (22.3)</p>		
--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--

	<p><u>Gestational age (for cases, gestational age at diagnosis of stillbirth; for controls at time of interview) - median (IQR)</u> Cases: 37 weeks (34 - 39) Controls: 37 (32 - 39)</p>			
--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--	--	--

Abbreviations: aOR: adjusted odds ratio; BMI: body mass index; CI: confidence interval; IPD MA: individual patient data meta-analysis; IQR: interquartile range; N: total number of participants in the study or case or control; OR: odds ratio; SGA: small for gestational age; vs: versus;

Appendix E – Forest plots

Forest plots for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

No meta-analyses were performed and therefore no forest plots are presented.

Appendix F – GRADE tables

GRADE tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby? – evidence from IPD meta-analysis

Table 7: GRADE table for independent association between going-to-sleep position and stillbirth in comparison to left lateral going-to-sleep position – evidence from IPD meta-analysis

Phase of investigation	<p>No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.</p> <p>One IPD meta-analysis, including data from 5 case control studies^a, was included and the quality of evidence was thus initially rated as high.</p>
Results	<p>Adjusted odds ratio (95% CI), vs left sleeping position for stillbirth:</p> <p>Supine: 2.63 (1.72, 4.04) Right side: 1.04 (0.83, 1.31) Prone: 0.63 (0.12, 3.25) Variable sides: 0.97 (0.70, 1.35) Propped up: 1.30 (0.68, 2.49) Don't remember: 2.26 (1.48, 3.46)</p>
Study limitations	<p>The IPD meta-analysis was rated at low risk of bias.</p>
Inconsistency	<p>There was no formal assessment of inconsistency within the IPD meta-analysis however the results of the IPD meta-analysis were broadly consistent with the findings of the primary studies reported in the tables below. Therefore the evidence was not downgraded for inconsistency.</p>
Indirectness	<p>No serious indirectness in the majority of the evidence contributing to the IPD meta-analysis.</p>
Imprecision	<p>Supine: no serious imprecision Right side: serious imprecision (confidence intervals crossed the line of no effect) Prone: very serious imprecision (confidence intervals crossed the line of no effect and subjectively were considered very wide)</p>

	<p>Variable sides: very serious imprecision (confidence intervals crossed the line of no effect and subjectively were considered very wide)</p> <p>Propped up: very serious imprecision (confidence intervals crossed the line of no effect and subjectively were considered very wide)</p> <p>Don't remember: no serious imprecision</p>
Publication bias	The evidence was downgraded for publication bias as it came from a relatively small number of primary studies, some of which were in an early phase of investigation.
Moderate/large effect size	The evidence was upgraded if moderate (aOR > 2.5 or aOR < 0.25) or large effect (aOR > 4.25 or aOR < 0.4) sizes were estimated. This applied to the supine position outcome.
Overall quality	<ul style="list-style-type: none"> • Supine: HIGH • Right side: LOW • Prone: VERY LOW • Variable sides: VERY LOW • Propped up: VERY LOW • Don't remember: MODERATE

Notes: ^aCronin 2019. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

Table 8: GRADE table for independent association between going-to-sleep position and SGA in comparison to left lateral going-to-sleep position on last night – evidence from IPD meta-analysis

Phase of investigation	<p>No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.</p> <p>One IPD meta-analysis, including data from 4 case control studies^a, was included and the quality of evidence was thus initially rated as high.</p>
Results	<p>Adjusted odds ratio (95% CI), vs left sleeping position for SGA as defined INTERGROWTH-21st 10th centile:</p> <p>Supine: 3.23 (1.37, 7.59)</p> <p>Right side: 1.05 (0.58, 1.90)</p> <p>Other: 1.14 (0.62, 2.09)</p>

	Results also available for customised growth charts where supine is no longer statistically significantly associated with <10th centile (adjusted odds ratio 1.55 (0.72 to 3.35)).
Study limitations	The IPD meta-analysis was rated at low risk of bias.
Inconsistency	There was no formal assessment of inconsistency within the IPD meta-analysis however the results of the IPD meta-analysis were broadly consistent with the findings of the primary studies reported in the tables below (assuming factors leading to stillbirth and SGA are likely to be similar). Therefore the evidence was not downgraded for inconsistency.
Indirectness	No serious indirectness in the majority of the evidence contributing to the IPD meta-analysis.
Imprecision	Supine: no serious imprecision Right side: very serious imprecision (confidence intervals crossed 0.8 and 1.25) Other: very serious imprecision (confidence intervals crossed 0.8 and 1.25)
Publication bias	The evidence was downgraded for publication bias as it came from a relatively small number of primary studies, some of which were in an early phase of investigation.
Moderate/large effect size	The evidence was upgraded if moderate (aOR > 2.5 or aOR <0.25) or large effect (aOR > 4.25 or aOR < 0.4) sizes were estimated. This applied to the supine position outcome.
Overall quality	<ul style="list-style-type: none"> • Supine: HIGH • Right side: VERY LOW • Other: VERY LOW

Notes: ^aAnderson 2019. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

GRADE tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby? – evidence from primary studies

Table 9: GRADE table for independent association between going-to-sleep position on last night and stillbirth in comparison to left lateral going-to-sleep position on last night

Phase of investigation	<p>No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.</p> <p>Prone going-to-sleep position One phase 2 prospective population-based case control study^a was included in the review of prone going-to-sleep position on last night and the quality of evidence was thus initially rated as high.</p> <p>Right lateral, supine going-to-sleep position One phase 2 retrospective nested case-control study in a cohort of known size^b and two phase 2 prospective population-based case control studies^{a,c} were included in the review of right lateral going-to-sleep position or supine going-to-sleep position on last night and the quality of evidence was thus initially rated as high.</p> <p>Sitting/propped, variable-lateral going-to-sleep position One phase 2 retrospective nested case-control study in a cohort of known size^b and one phase 2 prospective population-based case control study^a were included in the reviews of variable-lateral or sitting/propped going-to-sleep position on last night and the quality of evidence was thus initially rated as high.</p>
Study limitations	<p>The three included phase 2 studies were assessed as being at overall low^c, moderate^a and high^b risk of bias. The main sources of bias in these studies were: high^{a,b} or moderate^c risk of selection bias regarding the study population due to study attrition, and moderate risk of recall bias^{a,b,c} for measurement of going-to-sleep position on last night given that there was a delay in all studies between birth outcome (i.e. stillbirth or live birth) and the interview in which sleep data was collected.</p>
Inconsistency	<p>Prone going-to-sleep position There was only one study (N=1024)^a contributing to the evidence of association between prone going-to-sleep position and stillbirth. No serious inconsistency.</p> <p>Right lateral going-to-sleep position</p>

Although none of the studies found a statistically significant association between right lateral going-to-sleep position and stillbirth, there is some inconsistency in the results: two of the studies found positive association between pregnant women who reported right lateral going-to-sleep position on the last night and still birth compared to left lateral going-to-sleep position (aOR 1.11 [95% CI 0.70 to 1.77]^b; aOR 1.74 [95% CI 0.98 to 3.01]^c) whilst one of the studies found negative association between those reporting right lateral going-to-sleep position on the last night and stillbirth (aOR 0.67 [95% CI 0.44 to 1.02]^a); in addition, although each of the 95% confidence intervals associated with the reported adjusted odds ratios crossed the line of no effect, there was minimal overlap in two of the studies^{a,c}. The inconsistency in the results regarding the association between right-lateral going-to-sleep position on the last night and stillbirth was therefore considered to be serious.

Sitting/propped going-to-sleep position

None of the studies reported a statistically significant association between sitting/propped going-to-sleep position and stillbirth and the adjusted odds ratios were similar and there was significant overlap of 95% confidence interval between the two (aOR 0.44 [95%CI 0.13 to 1.49]^a; aOR 0.71 [95%CI 0.22 to 2.30]^b). Thus, the inconsistency in the results regarding the association between variable going-to-sleep position on the last night and stillbirth was considered to be minimal and the evidence was not downgraded.

Supine going-to-sleep position

Two studies^{a,c} reported a statistically significant association between supine going-to-sleep position and stillbirth compared to left lateral going-to-sleep position whereas another study^b did not find any significant association. There was inconsistency in the results: two of the studies^{a,c} reported a statistically significant association between supine going-to-sleep position and stillbirth compared to left lateral going-to-sleep position (aOR 2.31 [95% CI 1.04 to 5.11]^a; aOR 2.54 [95% CI 1.04 to 6.18]^c) whereas another study^b showed a positive but not a statistically significant association (aOR 1.05 [95% CI 0.32 to 3.50]^a). The inconsistency in the results regarding the association between supine going-to-sleep position on the last night and stillbirth was therefore considered to be serious.

Variable-lateral going-to-sleep position

None of the studies reported a statistically significant association between variable-lateral going-to-sleep position and stillbirth and the adjusted odds ratios were similar and there was significant overlap of 95% confidence interval between the two (aOR 0.93 [95%CI 0.51 to 1.69]^a; aOR 0.75 [95%CI 0.34 to 1.64]^b). Thus, the inconsistency in the results regarding the association between variable-lateral going-to-sleep position on the last night and stillbirth was considered to be minimal and the evidence was not downgraded.

Indirectness

Two of the included studies were multi-centre studies and included all consenting pregnant women who experienced stillbirth in high-income countries (New Zealand^a and the UK^c) and can therefore be

	<p>considered to be representative of the target population. One of the included studies^b was an anonymous international online survey that recruited participants through the use of web-based advertising, social media, and word of mouth. This study, unlike the other two studies^{a,c}, also restricted participants to those who were at least 18 years old. The evidence was therefore downgraded for serious indirectness in the population of interest. The evidence regarding indirectness in the studied prognostic factor (i.e. going-to-sleep position) and outcome of interest (i.e. stillbirth) was not considered to be serious.</p>
<p>Imprecision</p>	<p>Prone going-to-sleep position</p> <p>One study^a examined the relationship between prone going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. The confidence interval around the estimated effect size was wide (0.13 to 7.81), not indicating an association between prone going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. Therefore, the imprecision was considered to be very serious.</p> <p>Right lateral going-to-sleep position</p> <p>Three studies^{a,b,c} examined the association between right lateral going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. No significant effect of right lateral going-to-sleep position on the last night on stillbirth compared to left lateral going-to-sleep position was found and the 95% confidence intervals were relatively wide ([0.44 to 1.02]^a, [0.70 to 1.77]^b and [0.98 to 3.01]^c) with none of the studies indicating an association between right lateral going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. The evidence was therefore downgraded for serious imprecision due to substantial uncertainty in the effect estimate of each included study.</p> <p>Sitting/propped going-to-sleep position</p> <p>Two studies^{a,b} evaluated the association between sitting/propped going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. No significant association was found in either study and the 95% confidence intervals were wide ([0.13 to 1.49]^a and [0.22 to 2.30]^b), not indicating an association between sitting/propped going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. Therefore, the imprecision of the effect was considered to be very serious.</p> <p>Supine going-to-sleep position</p> <p>The 95% confidence intervals in all studies were wide^{a,b,c} ([1.04 to 5.11]^a, [0.32 to 3.50]^b and [1.04 to 6.18]^c), with 2 studies^{a,c} finding a statistically significant positive association between supine going-to-sleep position and stillbirth, and one study^b not finding a significant association, compared to left lateral going-to-sleep position. Thus, the results are inconclusive regarding the association and the imprecision was considered to be very serious.</p> <p>Variable-lateral going-to-sleep position</p>

	<p>No significant association was found in either study between variable-lateral going-to-sleep position and stillbirth and the 95% confidence intervals around the effect estimates were relatively wide [(0.51 to 1.69)^a and (0.34 to 1.64)^b], not indicating an association between variable-lateral going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. Therefore, the imprecision was considered to be very serious.</p>
Publication bias	<p>Although these studies were not ‘pure’ cohort studies, the use of a case control design is appropriate given the low frequency of stillbirth. All three studies used multivariate analysis to adjust for potential confounders such as age, BMI, ethnicity and smoking status during pregnancy. However, since the association between any going-to-sleep position on the last night stillbirth compared to left lateral going-to-sleep position was only explored in 3 studies, the evidence was downgraded for publication bias.</p>
Moderate/large effect size	<p>Prone going-to-sleep position The estimated adjusted effect size in the study was small (aOR=1.01)^a and the 95% confidence interval around the estimate crossed the line of no effect, therefore the evidence was not upgraded.</p> <p>Right-hand going-to-sleep position The estimated adjusted effect sizes in the three studies were all small with two studies favouring left lateral going-to-sleep position (aOR=1.11^b; aOR=1.74^c) compared to right-side position, and one study favouring right-side going-to-sleep position compared to left lateral position (aOR 0.67^a), therefore the evidence was not upgraded.</p> <p>Sitting/propped going-to-sleep position The odds of stillbirth for sitting/propped going-to-sleep position in two included studies were small compared to left lateral going-to-sleep position (aOR=0.44^a; aOR=0.71^b) and the 95% confidence intervals crossed the line of no effect, therefore the evidence was not upgraded.</p> <p>Supine going-to-sleep position The odds of stillbirth for supine going-to-sleep position in two studies were moderate to large (aOR=2.31^a; aOR=2.54^c) whereas the odds in another study was found to be small (aOR=1.05^b). In the first two studies^{a,c}, the 95% confidence intervals did not cross the line of no effect, indicating the positive association between supine going-to-sleep position and stillbirth, whilst the 95% confidence interval in another study^b crossed the line of no effect. Thus, because of variability in the results, the evidence for an independent effect of supine going-to-sleep position on stillbirth was not upgraded.</p> <p>Variable-lateral going-to-sleep position The estimated going-to-sleep position in two included studies were small (aOR=0.93^a; aOR=0.75^b) and the 95% confidence intervals crossed the line of no effect, therefore the evidence was not upgraded.</p>

Overall quality	<ul style="list-style-type: none"> • Prone: VERY LOW • Right-lateral: VERY LOW • Sitting/Propped: VERY LOW • Supine: VERY LOW • Variable-lateral: VERY LOW
------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Notes: ^aHeazell 2018; ^bO'Brien 2019; ^cStacey 2011. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

Table 10: GRADE table for independent association between going-to sleep position on last month of pregnancy and late stillbirth in comparison to left lateral going-to-sleep position on last month

Phase of investigation	<p>No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.</p> <p>Right-lateral going-to-sleep position, supine going-to-sleep position One phase 1 prospective case control study^a and one phase 2 nested case-control studies in a cohort of known size^b were included in the review of right lateral or supine going-to-sleep position on last month and the quality of evidence was thus initially rated as moderate.</p> <p>Variable-lateral going-to-sleep position, sitting/propped going-to-sleep position One phase 2 nested case-control study in a cohort of known size^b was included in the review of variable-lateral or sitting/propped going-to-sleep position on last month and the quality of evidence was thus initially rated as high.</p>
Study limitations	<p>The included phase 1 study and phase 2 study were all assessed as being at overall low^a and high^b risk of bias. The main sources of bias in these studies were: high risk of selection bias due to unclear information on study population^b, and incomparable baseline characteristics between cases and controls; moderate risk of recall bias^{a,b} for measurement of going-to-sleep position on last night given that there was a delay in all studies between birth outcome (i.e. stillbirth or live birth) and the interview in which sleep data was collected.</p>
Inconsistency	<p>Right-lateral going-to-sleep position None of the studies found a statistically significant association between right-lateral going-to-sleep position and stillbirth. There is minimal inconsistency in the results (aOR 1.1 [95% CI 0.43 to 2.6]^a; aOR 1.14 [95% CI 0.70 to 1.85]^b). The inconsistency in the results of these two studies was therefore not considered to be serious.</p>

	<p>Sitting/propped going-to-sleep position, variable-lateral going-to-sleep position</p> <p>There was only one study^b (N=633^b) contributing to the evidence of association between going-to-sleep position on last month and stillbirth. No serious inconsistency.</p> <p>Supine going-to-sleep position</p> <p>One study reported a statistically significant positive association between supine going-to-sleep position and stillbirth (aOR 6.26 [95%CI 1.15 to 34.00])^a whereas another study reported negative association although the result was not statistically significant (aOR 0.37 [95%CI 0.04 to 3.12])^b. Thus, the inconsistency of the results between these two studies was considered to be very serious.</p>
Indirectness	<p>One of the included studies was a multi-centre study and included all consenting pregnant women who experienced stillbirth in high-income countries (Australia^a) and can therefore be considered to be representative of the target population. Another study^b was an anonymous international online survey that recruited participants through the use of web-based advertising, social media, and word of mouth. This study, unlike the other study^a, also restricted participants to those who were at least 18 years old. The evidence was therefore downgraded for serious indirectness in the population of interest. The evidence in this study regarding indirectness in the studied prognostic factor (i.e. going-to-sleep position) and outcome of interest (i.e. stillbirth) was not considered to be serious.</p>
Imprecision	<p>Right-lateral going-to-sleep position</p> <p>Two studies^{a,b} examined the relationship between right-lateral going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. The 95% confidence intervals around the estimated effect size crossed the line of no effect and were considered to be wide^b (0.70 to 1.85)^b and very wide^a (0.43 to 2.6)^a, not indicating an association between right-lateral going-to-sleep position and stillbirth compared to left lateral going-to-sleep position. Overall the imprecision was considered to be serious.</p> <p>Sitting/propped going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate in this study crossed the line of no effect and was considered to be very wide (0.39 to 3.68)^b therefore, the imprecision was considered to be very serious.</p> <p>Supine going-to-sleep position</p> <p>One study (N=295)^a reported statistically significant positive association (aOR=6.26)^a although the 95% confidence interval was considered to be very wide (1.15 to 34.00)^a. However, another study (N=542)^b did not find an association and the 95% confidence interval ranged from negative association (0.04)^b to positive association (3.12)^b. Thus, the imprecision was considered to be very serious.</p> <p>Variable-lateral going-to-sleep position</p>

	The 95% confidence interval around the effect estimate in this study crossed the line of no effect and was considered to be wide (0.48 to 1.55) ^b therefore, the imprecision was considered to be serious.
Publication bias	Although these studies were not 'pure' cohort studies, the use of a case control design is appropriate given the low frequency of stillbirth. Both studies used multivariate analysis to adjust for potential confounders such as age, BMI, ethnicity and smoking status during pregnancy. However, since the association between any going-to-sleep position on the last month and late stillbirth compared to left lateral going-to-sleep position was only explored in 2 studies, the evidence was downgraded for publication bias.
Moderate/large effect size	<p>Right-lateral going-to-sleep position The estimated adjusted effect sizes in the two studies^{a,b} were small (aOR=1.1^a; aOR=1.14^b) and therefore not upgraded.</p> <p>Sitting/propped going-to-sleep position The estimated adjusted effect size in the study^b was small (aOR=1.20^a) and therefore not upgraded.</p> <p>Supine going-to-sleep position There were significant differences in the effect estimates reported by two included studies^{a,b} where the estimated adjusted effect size in one study was large (aOR=6.26)^a and showed positive association whereas the adjusted effect estimate in another study was moderate (aOR=0.37)^b and showed negative association. Given the inconsistency, the evidence was not upgraded.</p> <p>Variable-lateral going-to-sleep position The estimated adjusted effect size in the study was small (aOR=0.87)^b and therefore not upgraded.</p>
Overall quality	<p>Right-lateral going-to-sleep position: VERY LOW</p> <p>Sitting/propped going-to-sleep position: VERY LOW</p> <p>Supine going-to-sleep position: VERY LOW</p> <p>Variable-lateral going-to-sleep position: VERY LOW</p>

Notes: ^aGordon 2015; ^bO'Brien 2019. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

Table 11: GRADE table for independent association between going-to-sleep position on last night or last week of pregnancy and preterm stillbirth (28 to 36 weeks gestation) in comparison to left lateral going-to-sleep position on last night or last week

Phase of investigation	No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.
-------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	One phase 2 prospective population-based case-control study ^a was included in the review of the association between going-to-sleep position on last night or last week and preterm stillbirth (defined as fetal death between 28 to 36 weeks gestation). Thus, the quality of evidence was initially rated as high.
Study limitations	The included phase 2 study was assessed as being at overall moderate ^a risk of bias. The main sources of bias in this study were: moderate risk of selection bias regarding the study population due to study attrition ^a , and moderate risk of recall bias for measurement of going-to-sleep position on last night or last week ^a given that there was a delay in all studies between birth outcome (i.e. stillbirth or live birth) and the interview in which sleep data was collected.
Inconsistency	There was only one study (N=733) ^a contributing to the evidence of association between going-to-sleep position on last night or last week and preterm stillbirth (defined as fetal death between 28 and 36 weeks gestation). No serious inconsistency.
Indirectness	The included study was a multi-centre study and included all consenting pregnant women who experienced stillbirth in high-income country (New Zealand ^a) and can therefore be considered to be representative of the target population. In addition, the evidence regarding indirectness in the studied prognostic factor (i.e. going-to-sleep position) and outcome of interest (i.e. preterm stillbirth) was not considered to be serious.
Imprecision	<p><u>Going-to-sleep position on last night</u></p> <p>Restless going-to-sleep position^b The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.61 to 19.97)^a therefore the imprecision was considered to be very serious.</p> <p>Right-lateral going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.48 to 1.94)^a therefore the imprecision was considered to be serious.</p> <p>Sitting/propped going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.11 to 178.86)^a, therefore, the imprecision was considered to be very serious.</p> <p>Supine going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.97 to 10.05)^a, therefore, the imprecision was considered to be serious.</p> <p><u>Going-to-sleep position on last week</u></p>

	<p>Prone going-to-sleep position The 95% confidence interval around the effect estimate^a crossed the line of no effect and was considered to be very wide (0.45 to 278.58)^a, therefore the imprecision was considered to be very serious.</p> <p>Right-lateral going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.34 to 1.54)^a, therefore the imprecision was considered to be serious.</p> <p>Sitting/propped going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.08 to 210.43)^a, therefore the imprecision was considered to be very serious.</p> <p>Supine going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.65 to 7.84)^a, therefore the imprecision was considered to be very serious.</p> <p>Variable-lateral going-to-sleep position The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.18 to 2.19)^a, therefore the imprecision was considered to be very serious.</p>
Publication bias	Although the included study was not a 'pure' cohort study, the use of a case control design is appropriate given the low frequency of stillbirth. This study used multivariate analysis to adjust for potential confounders such as age, BMI, ethnicity and smoking status during pregnancy. However, since the association between any going-to-sleep position on the last night or last week and preterm stillbirth compared to left lateral going-to-sleep position is only explored in 1 study, the evidence was downgraded for publication bias.
Moderate/large effect size	<p><u>Going-to-sleep position on last night</u></p> <p>Restless going-to-sleep position^b The estimated adjusted effect size in this study was moderate (aOR=3.50)^a however this was not statistically significant and was therefore not upgraded.</p> <p>Right-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=0.96)^a and was therefore not upgraded.</p>

	<p>Sitting/propped going-to-sleep position The estimated adjusted effect size in this study was moderate (aOR=4.37)^a however this was not statistically significant and was therefore not upgraded.</p> <p>Supine going-to-sleep position The estimated adjusted effect size in this study was small (aOR=2.25)^a and was therefore not upgraded.</p> <p><u>Going-to-sleep position on last week</u></p> <p>Prone going-to-sleep position The estimated adjusted effect size in this study was large (aOR=10.71)^a however this was not statistically significant and was therefore not upgraded.</p> <p>Right-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=0.73)^a and was therefore not upgraded.</p> <p>Sitting/propped going-to-sleep position The estimated adjusted effect size in this study was moderate (aOR=4.01)^a however this was not statistically significant and was therefore not upgraded.</p> <p>Supine going-to-sleep position The estimated adjusted effect size in this study was small (aOR=2.25)^a and therefore not upgraded.</p> <p>Variable-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=0.63)^a and therefore not upgraded.</p>
Overall quality	<p><u>Going-to-sleep position on last night</u> Restless going-to-sleep position: VERY LOW Right-lateral going-to-sleep position: VERY LOW Sitting/propped going-to-sleep position: VERY LOW Supine going-to-sleep position: VERY LOW</p> <p><u>Going-to-sleep position on last week</u></p>

	Prone going-to-sleep position: VERY LOW Right-lateral going-to-sleep position: VERY LOW Sitting/propped going-to-sleep position: VERY LOW Supine going-to-sleep position: VERY LOW Variable-lateral going-to-sleep position: VERY LOW
--	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Notes: ^aMcCowan 2018; 'Restless' going-to-sleep position refers to women who frequently change positions when going to sleep and could not remember the position they had just before falling asleep. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

Table 12: GRADE table for independent association between going-to-sleep position on last night or last week of pregnancy and term stillbirth (≥37 weeks gestation) in comparison to left lateral going-to-sleep position

Phase of investigation	<p>No restrictions were placed on phase of investigation. Given the relatively low frequency of stillbirth, a phase 3 prospective cohort study that aims to examine the role of sleep position and its effect on still birth is not feasible. Therefore no restrictions were placed on the phase of investigation.</p> <p>One phase 2 prospective population-based case-control study^a was included in the review of the association between going-to-sleep position on last night or last week and term stillbirth (defined as fetal death after 36 weeks gestation). Thus, the quality of evidence was initially rated as high.</p>
Study limitations	<p>The included phase 2 study was assessed as being at overall moderate risk of bias.^a The main sources of bias in this study were: moderate risk of selection bias regarding the study population due to study attrition, and moderate risk of recall bias for measurement of going-to-sleep position on last night or last week given that there was a delay in all studies between birth outcome (i.e. stillbirth or live birth) and the interview in which sleep data was collected.</p>
Inconsistency	<p>There was only one study (N=733)^a contributing to the evidence of association between going-to-sleep position on last night or last week and term stillbirth (defined as fetal death after 36 weeks gestation). No serious inconsistency.</p>
Indirectness	<p>The included study was a multi-centre study and included all consenting pregnant women who experienced stillbirth in a high-income country (New Zealand^a) and can therefore be considered to be representative of the target population. In addition, the indirectness in the studied prognostic factor (i.e. going-to-sleep position) and outcome of interest (i.e. term stillbirth) was not considered to be serious.</p>
Imprecision	<p><u>Going-to-sleep position on last night</u></p> <p>Restless going-to-sleep position^b</p> <p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.64 to 6.21)^a therefore the imprecision was considered to be very serious.</p> <p>Right-lateral going-to-sleep position</p>

	<p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.48 to 1.99)^a, therefore the imprecision was considered to be serious.</p> <p>Sitting/propped going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.17 to 5.97)^a, therefore the imprecision was considered to be very serious.</p> <p>Supine going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate did not cross the line of no effect but was considered to be wide (3.01 to 35.04)^a therefore the imprecision was considered to be serious.</p> <p><u>Going-to-sleep position on last week</u></p> <p>Right-lateral going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.47 to 1.89)^a therefore the imprecision was considered to be serious.</p> <p>Sitting/propped going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be very wide (0.47 to 14.81)^a therefore the imprecision was considered to be very serious.</p> <p>Supine going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate did not cross the line of no effect but was considered to be very wide (2.92 to 55.46)^a therefore, the imprecision was considered to be serious.</p> <p>Variable-lateral going-to-sleep position</p> <p>The 95% confidence interval around the effect estimate crossed the line of no effect and was considered to be wide (0.49 to 3.01)^a therefore, the imprecision was considered to be serious.</p>
Publication bias	<p>The included study used multivariate analysis to adjust for potential confounders such as age, BMI, ethnicity and smoking status during pregnancy.^a Although this study was a non-nested case control study, this design is appropriate given the low frequency of stillbirth. However, this is the only study to have reported results relative to time of stillbirth. The association between any going-to-sleep position on the last night or last week and term stillbirth compared to left lateral going-to-sleep position was therefore considered to not be adequately explored by the identified study and the evidence was downgraded for publication bias.</p>

<p>Moderate/large effect size</p>	<p><u>Going-to-sleep position on last night</u></p> <p>Restless going-to-sleep position^b The estimated adjusted effect size in this study was small (aOR=2.0)^a and was therefore not upgraded.</p> <p>Right-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=0.98)^a and was therefore not upgraded.</p> <p>Sitting/propped going-to-sleep position The estimated adjusted effect size in this study was small (aOR=1.02)^a and was therefore not upgraded.</p> <p>Supine going-to-sleep position The estimated adjusted effect size in this study was large (aOR=10.26)^a. In addition, this adjusted effect estimate was statistically significant. The evidence was therefore upgraded.</p> <p><u>Going-to-sleep position on last week</u></p> <p>Right-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=0.95)^a and was therefore not upgraded.</p> <p>Sitting/propped going-to-sleep position The estimated adjusted effect size in this study was moderate (aOR=2.64)^a however this adjusted effect estimate was not statistically significant and was therefore not upgraded.</p> <p>Supine going-to-sleep position The estimated adjusted effect size in this study was large (aOR=12.73)^a. In addition, this adjusted effect estimate was statistically significant. The evidence was therefore upgraded.</p> <p>Variable-lateral going-to-sleep position The estimated adjusted effect size in this study was small (aOR=1.11)^a and was therefore not upgraded.</p>
<p>Overall quality</p>	<p><u>Going-to-sleep position on last night</u></p> <p>Restless going-to-sleep position: VERY LOW Right-lateral going-to-sleep position: VERY LOW Sitting/propped going-to-sleep position: VERY LOW Supine going-to-sleep position: LOW</p> <p><u>Going-to-sleep position on last week</u></p> <p>Right-lateral going-to-sleep position: VERY LOW Sitting/propped going-to-sleep position: VERY LOW Supine going-to-sleep position: LOW Variable-lateral going-to-sleep position: VERY LOW</p>

Notes: ^aMcCowan 2018; ^b, 'Restless' going-to-sleep position refers to women who frequently change positions when going to sleep and could not remember the position they had just before falling asleep. Abbreviations: aOR, adjusted odds ratio; CI, confidence interval.

Appendix G – Economic evidence study selection

Economic evidence study selection for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

A single economic search was undertaken for all topics included in the scope of this guideline. No economic studies were identified which were applicable to this review question. See supplementary material 2 for details.

Appendix H – Economic evidence tables

Economic evidence tables for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

No economic evidence was identified which was applicable to this review question.

Appendix I – Health economic evidence profiles

Economic evidence profiles for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

No evidence was identified which was applicable to this review question.

Appendix J – Health economic analysis

Economic analysis for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

No economic analysis was conducted for this review question.

Appendix K – Excluded studies

Excluded studies for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

Clinical studies

Table 13: Excluded studies

Study	Reason for exclusion
Bei, B., Neemia, D., Shen, L., Fulgoni, C., Blumfield, M. L., Drummond, S. P., Newman, L. K., Manber, R., A brief, automated cognitive behavioral program prevents sleep disturbance and insomnia in late pregnancy: A randomized controlled trial, <i>Sleep</i> , 41 (Supplement 1), A151, 2018	Conference abstract
Cronin, R. S., Chelimo, C., Mitchell, E. A., Okesene-Gafa, K., Thompson, J. M. D., Taylor, R. S., Hutchison, B. L., McCowan, L. M. E., Survey of maternal sleep practices in late pregnancy in a multi-ethnic sample in South Auckland, New Zealand, <i>BMC pregnancy and childbirth</i> , 17 (1) (no pagination), 2017	No relevant outcomes reported
Cronin, R. S., Li, M., Thompson, J. M. D., Gordon, A., Raynes-Greenow, C., Heazell, A. E. P., Stacey, T., Culling, V., Bowring, V., Askie, L., Mitchell, E. A., McCowan, L. M. E., An individual participant data meta-analysis of going-to-sleep position, interactions with fetal vulnerability and the risk of late stillbirth, <i>Journal of Paediatrics and Child Health</i> , 54 (Supplement 1), 3, 2018	Conference abstract
Gaudet, L., Simon, A., Pratt, M., Heslehurst, N., Hayes, L., Flynn, A., Velez, M. P., Smith, G., Skidmore, B., Hutton, B., Rybak, N., Walker, M., Predictors of adverse pregnancy outcomes in obese pregnant women a systematic review, <i>International Journal of Gynecology and Obstetrics</i> , 143 (Supplement 3), 600-601, 2018	Conference abstract
Heazell, A. E. P., Cronin, R. S., Li, M., Thompson, J. M. D., Gordon, A., Raynes-Greenow, C., Stacey, T., Culling, V., Bowring, V., Askie, L., Mitchell, E. A., McCowan, L. M. E., Going to sleep position and risk of late stillbirth: A systematic review and metaanalysis, <i>Journal of Paediatrics and Child Health</i> , 54 (Supplement 1), 24-25, 2018	Conference abstract
Heazell, A. E. P., Li, M., Thompson, J. M. D., Budd, J., Cronin, R., Mitchell, E., Stacey, T., Roberts, D., Martin, B., McCowan, L. M. E., Going to sleep supine and reduced sleep duration are risk factors for late stillbirth: Findings from the MiNESS Case-Control Study, <i>BMC Pregnancy and Childbirth</i> . Conference:	Conference abstract

Study	Reason for exclusion
International Stillbirth Alliance Conference, 17, 2017	
Hsu, Christine, Sleep Positions Can Predict Women's Risk of Stillbirth, Inside Childbirth Education, 9-9, 2013	Newsletter
Kempler, L., Sharpe, L., Bartlett, D., Sleep education during pregnancy for new mothers, BMC Pregnancy & Childbirth, 12, 155, 2012	Protocol
Kichler, A., Alzubaidi, M., Emery, J., Gabbard, S., Use of a positional therapy device significantly improves nocturnal gastroesophageal reflux disease symptoms in pregnant women, American Journal of Gastroenterology, 1), S703-S704, 2015	Conference abstract
Koken, G. N., Kanat-Pektas, M., Kose, S. K., Ario, D. T., Yilmazer, M., Maternal blood pressure and dominant sleeping position may affect placental localization, Journal of Maternal-Fetal and Neonatal Medicine, 27, 1564-1567, 2014	No relevant outcomes
Lakshmi, Sujatha Thankappan et al. , Risk factors for still birth: a hospital based case control study, International Journal of Reproduction, Contraception, Obstetrics and Gynecology, 6, 970-974, 2017	Incorrect comparison (non-left vs left)
Lee, K.A., Gay, C.L., Sleep in late pregnancy predicts length of labor and type of delivery, American Journal of Obstetrics and Gynecology, 191, 2041-2046, 2004	No relevant outcomes
Li, M., McCowan, L. M. E., Thompson, J. M. D., Cronin, R. S., Anderson, N., Stacey, T., Stone, P., Lawton, B. A., Ekeroma, A. J., Mitchell, E. A., Supine going-to-sleep position is a major risk factor for term stillbirth: Findings from the New Zealand multicentre stillbirth case-control study, BMC Pregnancy and Childbirth. Conference: International Stillbirth Alliance Conference, 17, 2017	Conference abstract
Li, M., Mitchell, E. A., Cronin, R. S., Thompson, J. M. D., Gordon, A., Raynes-Greenow, C., Heazell, A. E. P., Stacey, T., Culling, V., Bowring, V., Askie, L., McCowan, L. M. E., Environmental tobacco smoke exposure is associated with increased late stillbirth: Findings from the collaborative individual participant data (IPD) sleep and stillbirth (CRIBSS) meta-analysis, Journal of Paediatrics and Child Health, 54 (Supplement 1), 31, 2018	Conference abstract
Li, M., Thompson, J. M. D., Cronin, R. S., Gordon, A., Raynes-Greenow, C., Heazell, A. E. P., Stacey, T., Culling, V., Bowring, V., Mitchell, E. A., McCowan, L. M. E., Askie, L., The Collaborative IPD of Sleep and Stillbirth (Cribss): is maternal going-to-sleep position a risk factor for late stillbirth and does maternal sleep	Protocol

Study	Reason for exclusion
position interact with fetal vulnerability? An individual participant data meta-analysis study protocol, <i>BMJ open</i> , 8, e020323, 2018	
Li, R., Zhang, J., Zhou, R., Liu, J., Dai, Z., Liu, D., Wang, Y., Zhang, H., Li, Y., Zeng, G., Sleep disturbances during pregnancy are associated with cesarean delivery and preterm birth, <i>Journal of maternal-fetal & neonatal medicine</i> , 30, 733â–738, 2017	Sleep position not examined in study
Lillis, T. A., Hamilton, N. A., Pressman, S. D., Khou, C. S., The Association of Daytime Maternal Napping and Exercise With Nighttime Sleep in First-Time Mothers Between 3 and 6 Months Postpartum, <i>Behavioral sleep medicine</i> , 16, 527-541, 2018	Sleep position not examined in study
McCowan, L., Contribution of maternal going-to-sleep position and fetal movements to late stillbirth, <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 57 (Supplement 1), 19-20, 2017	Conference abstract
McCowan, Lesley M. E., Cronin, Robin S., Gordon, Adrienne, O'Brien, Louise, Heazell, Alexander E. P., Prospective Evaluation of Maternal Sleep Position Through 30 Weeks of Gestation and Adverse Pregnancy Outcomes, <i>Obstetrics & Gynecology</i> , 135, 218-218, 2020	Commentary
McIntyre, J. P. R., Ingham, C. M., Hutchinson, B. L., Thompson, J. M. D., McCowan, L. M., Stone, P. R., Veale, A. G., Cronin, R., Stewart, A. W., Ellyett, K. M., Mitchell, E. A., A description of sleep behaviour in healthy late pregnancy, and the accuracy of self-reports, <i>BMC Pregnancy and Childbirth</i> , 16 (1) (no pagination), 2016	Non-comparative study
McIntyre, J. P. R., Stone, P. R., Mitchell, E. M., Veale, A. G., How healthy women sleep in late pregnancy; A video and portable polysomnography study, <i>Sleep and Biological Rhythms</i> , 11, 44, 2013	Conference abstract
Morokuma, S., Shimokawa, M., Kato, K., Sanefuji, M., Shibata, E., Tsuji, M., Senju, A., Kawamoto, T., Kusuhara, K., Maternal sleep and small for gestational age infants in the Japan Environment and Children's Study: a cohort study, <i>BMC research notes</i> , 10, 394, 2017	Sleep position not examined in study
O'Brien, Louise M., Warland, Jane, Typical sleep positions in pregnant women, <i>Early Human Development</i> , 90, 315-317, 2014	Non-comparative study
Owusu, Jocelynn T., Anderson, Frank J., Coleman, Jerry, Oppong, Samuel, Seffah, Joseph D., Aikins, Alfred, O'Brien, Louise M., Association of maternal sleep practices with pre-eclampsia, low birth weight, and stillbirth among Ghanaian women, <i>International Journal of Gynecology & Obstetrics</i> , 121, 261-265, 2013	Cross-sectional study

Study	Reason for exclusion
Paine, S. J., Signal, T. L., Sweeney, B., Priston, M., Muller, D., Smith, A. A., Huthwaite, M., Lee, K., Gander, P. H., Ethnic differences in sleep across pregnancy: A cohort study, <i>Sleep and Biological Rhythms</i> , 2), 63, 2013	Conference abstract
Robertson, N., Okano, S., Kumar, S., Sleep in the supine position during pregnancy is associated with fetal cerebral redistribution, <i>Journal of Clinical Medicine</i> , 9, 1-12, 2020	Insufficient adjustment for confounders, insufficient detail on outcomes
Saarenmaa-Heikkila, O., Lehto, U., Kylliainen, A., Stenberg, T., Paunio, T., Paavonen, J., CHILD SLEEP-The Finnish birth cohort study: The effect of maternal sleep during pregnancy on a newborn wellbeing and a mother's labor experience, <i>Sleep Medicine</i> , 1), e42, 2013	Conference abstract
Silver, Robert M. M. D., Hunter, Shannon M. S., Reddy, Uma M. M. D. M. P. H., Facco, Francesca M. D., Gibbins, Karen J. M. D., Grobman, William A. M. D. M. B. A., Mercer, Brian M. M. D., Haas, David M. M. D. M. S., Simhan, Hyagriv N. M. D., Parry, Samuel M. D., Wapner, Ronald J. M. D., Louis, Judette M. D., Chung, Judith M. M. D., Pien, Grace M. D., Schubert, Frank P. M. D., Saade, George R. M. D., Zee, Phyllis M. D., Redline, Susan M. D., Parker, Corette B. DrPH, Silver, Robert M., Prospective Evaluation of Maternal Sleep Position Through 30 Weeks of Gestation and Adverse Pregnancy Outcomes, <i>Obstetrics & Gynecology</i> , 134, 667-676, 2019	Primary outcome was a composite adverse pregnancy outcome including stillbirth, hypertensive disorders of pregnancy (mild, severe, or superimposed preeclampsia; eclampsia; or antepartum gestational hypertension), and a small-for-gestational-age (SGA) newborns
Silver, Robert M., Reddy, Uma M., Gibbins, Karen J., Prospective Evaluation of Maternal Sleep Position Through 30 Weeks of Gestation and Adverse Pregnancy Outcomes, <i>Obstetrics & Gynecology</i> , 135, 218-219, 2020	Looked at sleep position in early pregnancy
Stacey, T., Mitchell, E. A., Sleep position and risk of late stillbirth, <i>BMC Pregnancy and Childbirth</i> . Conference: Stillbirth Summit, 12, 2011	Conference presentation of included study (Stacey 2011)
Stacey, T., Thomsson, J., Mitchell, E., McCowan, L., Maternal sleep practices: Possible risk factor for late stillbirth, <i>Archives of Disease in Childhood: Fetal and Neonatal Edition</i> , 1), Fa1, 2011	Conference abstract
Warland, J., Dorrian, J., Kember, A. J., Phillips, C., Borazjani, A., Morrison, J. L., O'Brien, L. M., Modifying maternal sleep position in late pregnancy through positional therapy: A feasibility study, <i>Journal of clinical sleep medicine</i> , 14, 1387-1397, 2018	No relevant outcomes
Warland, J., Dorrian, J., Morrison, J. L., O'Brien, L. M., Maternal sleep during pregnancy and poor fetal outcomes: A scoping review of the literature with meta-analysis, <i>Sleep Medicine Reviews</i> , 41, 197-219, 2018	Scoping review - references checked, no additional relevant studies (Lakshmi 2017 not included because it reported non-left vs left sleeping position)

Study	Reason for exclusion
Warland, J., Heazell, A. E. P., Collins, J. H., Huberty, J. L., Kliman, H. J., McGregor, J. A., Mitchell, E. A., O'Brien, L. M., Parast, M., Peesay, M., Stacey, T., Wimmer, L. J., An international internet survey of the experiences of 1,714 mothers with a late stillbirth: The STARS cohort study, BMC Pregnancy and Childbirth, 15 (1) (no pagination), 2015	Companion article to O'Brien 2017, no relevant data
Warland, J., Mitchell, E. A., A triple risk model for unexplained late stillbirth, BMC Pregnancy & Childbirth BMC Pregnancy Childbirth, 14, 142, 2014	Description of risk model for sudden infant death syndrome
Warland, J., Mitchell, E. A., O'Brien, L. M., Novel strategies to prevent stillbirth, Seminars In Fetal & Neonatal Medicine Semin Fetal Neonatal Med, 22, 146-152, 2017	Non-systematic review of novel strategies to prevent stillbirth
Woods, J. R., Heazell, A. E. P., Stillbirth: is it preventable?, Obstetrics, Gynaecology and Reproductive Medicine, 28, 148-154, 2018	Non-systematic review of stillbirth prevention strategies

Economic studies

One excluded list was created for all economic studies in this guideline. See supplementary material 2 for further information.

Appendix L – Research recommendations

Research recommendations for review question: Is there an association between sleep position on going to sleep and still birth or having a small for gestational age baby?

No research recommendations were made for this review question.