

Intrapartum care for healthy women and babies

Review questions

NICE guideline CG190 (update)

Review questions

March 2023

Early submission

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2023. All rights reserved. Subject to [Notice of Rights](#).

ISBN:

Contents

Review questions	5
-------------------------------	----------

Review questions

Evidence review	Review question
A	1.1 What are the benefits and risks of different places of birth for women at different BMI thresholds?
B	2.1 What is the optimum timeframe between a mother reporting possible PRoM and face-to-face clinical review?
C	3.1 What is the effectiveness of injected water papules for pain relief during labour?
D	3.2 What is the effectiveness of remifentanyl administered by intravenous patient-controlled analgesia (PCA) compared to other intramuscular opioids?
E	4.1 What is the effectiveness of Programmed Intermittent Epidural Bolus compared to other methods of maintaining epidural analgesia?
F	6.1 What is the effectiveness of altering the dose of intravenous oxytocin to reduce excessive frequency of uterine contractions? 6.2 What is the optimum dose at which oxytocin should be restarted if stopped due to an abnormality in the CTG?
G	7.1 What is the most effective position for birth in women with an epidural in situ? 7.2 What is the most effective position for birth in women without an epidural in situ?
H	7.3 What are the benefits and risks of the different pushing techniques (immediate, spontaneous, delayed, directed) in the second stage of labour in women with and without regional analgesia?
I	7.4 What is the effectiveness of perineal care in the second stage of labour (for example, massage, hands-on support and warm compresses) for reducing perineal trauma and tears?
J	7.5 What is the effectiveness of prophylactic antibiotics for preventing postnatal infections in assisted vaginal birth?
K	8.1 What are the benefits and risks associated with active management compared to physiological management in the third stage of labour?
L	8.2 Is intravenous administration of oxytocin more effective than intramuscular administration in the active management of the third stage of labour?
M	8.3 What is the effectiveness of uterotonics for the prevention of postpartum haemorrhage?
N	8.4 What is the optimum position for the baby during delayed cord clamping (including after instrumental and caesarean birth)?
O	8.5 What is the effectiveness of pharmacological treatments for the management of postpartum haemorrhage?