

Intrapartum care

Glossary and abbreviations

NICE guideline NG235

Supplement 2

September 2023

Final

*These supplements were developed by
NICE*

Disclaimer

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The recommendations in this guideline are not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

Local commissioners and/or providers have a responsibility to enable the guideline to be applied when individual health professionals and their patients or service users wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with compliance with those duties.

NICE guidelines cover health and care in England. Decisions on how they apply in other UK countries are made by ministers in the [Welsh Government](#), [Scottish Government](#), and [Northern Ireland Executive](#). All NICE guidance is subject to regular review and may be updated or withdrawn.

Copyright

© NICE 2023. All rights reserved. Subject to [Notice of Rights](#)

ISBN: 978-1-4731-5406-3

Contents

Glossary.....	5
Abbreviations	18

Glossary

Term	Definition
Abstract	Summary of a study, which may be published alone or as an introduction to a full scientific paper.
Alongside midwifery unit (AMU)	A place to give birth where the birth is overseen by midwives, but which is next to or in the same hospital as an obstetric unit.
Area under the curve (AUC)	Summary measure of the accuracy of a diagnostic test.
Arm (of a clinical study)	Subsection of individuals within a study who receive one particular intervention, for example placebo arm.
Assisted birth (or delivery)	The birth of a baby which requires assistance, involving either a ventouse suction cap or forceps.
Association	Statistical relationship between 2 or more events, characteristics or other variables. The relationship may or may not be causal.
Attrition bias	Systematic differences between comparison groups for withdrawal or exclusion of participants from a study.
Baseline	The initial set of measurements at the beginning of a study (after run-in period where applicable) with which subsequent results are compared.
Bias	Influences on a study that can make the results look better or worse than they really are. Bias can occur by chance, deliberately or as a result of systematic errors in the design and execution of a study. It can also occur at different stages in the research process, for example during the collection, analysis, interpretation, publication or review of research data. For examples see Confounding factor, Performance bias, Publication bias Selection bias.
Blinding	The practice of keeping the investigators or subjects of a study ignorant of the group to which a subject has been assigned. For example, a clinical trial in which the participating patients or their doctors are unaware of whether they (the patients) are taking the experimental drug or a placebo (dummy treatment). The purpose of 'blinding' or 'masking' is to protect against bias. See also double-blind study and single-blind study.
Caesarean birth or section, C-section, C-birth	A surgical procedure used to deliver a baby through incisions in the abdomen and uterus
Case series	Report of a number of cases of a given disease, usually covering the course of the disease and the response to treatment. There is no comparison (control) group of patients.
Case-control study	A study to find out the cause(s) of a disease or condition. This is done by comparing a group of patients who have the disease or condition (cases) with a group of people who do not have it (controls) but who are otherwise as similar as possible (in characteristics thought to be unrelated to the causes of the disease or condition). This means the researcher can look for aspects of their lives that differ to see if they may cause the condition. Such studies are retrospective because they look back in time from the outcome to the possible causes of a disease or condition.
Cephalic	The baby is positioned head-down, ready to be born head first through the vagina.
Clinical effectiveness	How well a specific test or treatment works when used in the 'real world' (for example when used by a doctor with a patient at home), rather than in a carefully controlled clinical trial. Trials that assess clinical

Term	Definition
	effectiveness are sometimes called management trials. Clinical effectiveness is not the same as efficacy.
Clinical efficacy	The extent to which an intervention is active when studied under controlled research conditions.
Clinician	A healthcare professional who provides patient care. For example a doctor, nurse or physiotherapist.
Cochrane Review	The Cochrane Library consists of a regularly updated collection of evidence-based medicine databases including the Cochrane Database of Systematic Reviews (reviews of RCTs prepared by the Cochrane Collaboration).
Cohort	A group of people sharing some common characteristic (e.g. patients with the same disease), followed up in a research study for a specified period of time.
Cohort study	A study with 2 or more groups of people – cohorts – with similar characteristics. One group receives a treatment, is exposed to a risk factor or has a particular symptom and the other group does not. The study follows their progress over time and records what happens.
Comorbidities	The presence of more than one disease or health condition in an individual at a given time
Comparative group	The group in the study who do not receive the treatment/procedure or who receive the norm treatment. This group is used to measure against the treatment/procedure being investigated.
Concealment of allocation	The process used to ensure that the person deciding to enter a participant into an RCT does not know the comparison group into which that individual will be allocated. This is distinct from blinding and is aimed at preventing selection bias. Some attempts at concealing allocation are more prone to manipulation than others and the method of allocation concealment is used as an assessment of the quality of a trial.
Confidence interval (CI)	<p>There is always some uncertainty in research. This is because a small group of patients is studied to predict the effects of a treatment on the wider population. The confidence interval is a way of expressing how certain we are about the findings from a study, using statistics. It gives a range of results that is likely to include the 'true' value for the population. The CI is usually stated as '95% CI', which means that the range of values has a 95 in 100 chance of including the 'true' value. For example, a study may state that "based on our sample findings, we are 95% certain that the 'true' population blood pressure is not higher than 150 and not lower than 110". In such a case the 95% CI would be 110 to 150.</p> <p>A wide confidence interval indicates a lack of certainty about the true effect of the test or treatment – often because a small group of patients has been studied. A narrow confidence interval indicates a more precise estimate (for example if a large number of patients have been studied).</p>
Confounding factor	Something that influences a study and can result in misleading findings if it is not understood or appropriately dealt with. For example, a study of heart disease may look at a group of people who exercise regularly and a group who do not exercise. If the ages of the people in the 2 groups are different, then any difference in heart disease rates between the 2 groups could be because of age rather than exercise. Therefore age is a confounding factor.
Continuous outcome	Data with a potentially infinite number of possible values within a given range. Height, weight and blood pressure are examples of continuous variables.

Term	Definition
Contraindicated	A situation in which a medication or treatment should not be administered
Control group	A group of people in a study who do not receive the treatment or test being studied. Instead, they may receive the standard treatment (sometimes called 'usual care') or a dummy treatment (placebo). The results for the control group are compared with those for a group receiving the treatment being tested. The aim is to check for any differences. Ideally, the people in the control group should be as similar as possible to those in the treatment group, to make it as easy as possible to detect any effects due to the treatment.
Controlled cord traction	Pulling on the cord while applying counter pressure on the uterus beneath the pubic bone, to help deliver the placenta.
Cost–benefit analysis (CBA)	Cost-benefit analysis is one of the tools used to carry out an economic evaluation. The costs and benefits are measured using the same monetary units (for example UK pounds) to see whether the benefits exceed the costs.
Cost–consequence analysis (CCA)	Cost-consequence analysis is one of the tools used to carry out an economic evaluation. This compares the costs (such as treatment and hospital care) with the consequences (such as health outcomes) of a test or treatment with a suitable alternative. Unlike cost–benefit analysis or cost-effectiveness analysis, it does not attempt to summarise outcomes in a single measure (such as the quality adjusted life year) or in financial terms. Instead, outcomes are shown in their natural units (some of which may be monetary) and it is left to decision-makers to determine whether, overall, the treatment is worth carrying out.
Cost-effectiveness analysis (CEA)	Cost-effectiveness analysis is one of the tools used to carry out an economic evaluation. The benefits are expressed in non-monetary terms related to health, such as symptom-free days, heart attacks avoided, deaths avoided or life years gained (that is, the number of years by which life is extended as a result of the intervention).
Cost-effectiveness model	An explicit mathematical framework which is used to represent clinical decision problems and incorporate evidence from a variety of sources in order to estimate the costs and health outcomes.
Cost-minimisation analysis (CMA)	Cost-minimisation analysis is a type of economic evaluation which can be used when the alternatives being compared have equivalent clinical effectiveness. The costs of alternatives are compared in order to determine which is the cheapest.
Cost–utility analysis (CUA)	Cost–utility analysis is one of the tools used to carry out an economic evaluation. The benefits are assessed in terms of both quality and duration of life, and expressed as quality adjusted life years (QALYs). See also Utility.
Credible interval (CrI)	The Bayesian equivalent of a confidence interval.
Cross-over study design	A study comparing two or more interventions in which the participants, upon completion of the course of one treatment, are switched to another. A problem with this study design is that the effects of the first treatment may carry over into the period when the second is given. Therefore a crossover study should include an adequate 'wash-out' period, which means allowing sufficient time between stopping one treatment and starting another so that the first treatment has time to wash out of the patient's system.
Cross-sectional study	The observation of a defined set of people at a single point in time or time period – a snapshot. (This type of study contrasts with a longitudinal study, which follows a set of people over a period of time.)

Term	Definition
Decision analysis	An explicit quantitative approach to decision-making under uncertainty, based on evidence from research. This evidence is translated into probabilities, and then into diagrams or decision trees which direct the clinician through a succession of possible scenarios, actions and outcomes.
Diagnostic study	A study to assess the effectiveness of a test or measurement in terms of its ability to accurately detect or exclude a specific disease.
Dichotomous outcomes	Outcome that can take one of 2 possible values, such as dead/alive, smoker/non-smoker, present/not present (also called binary data).
Discounting	Costs and perhaps benefits incurred today have a higher value than costs and benefits occurring in the future. Discounting health benefits reflects individual preference for benefits to be experienced in the present rather than the future. Discounting costs reflects individual preference for costs to be experienced in the future rather than the present.
Dominance	A health economics term. When comparing tests or treatments, an option that is both less effective and costs more is said to be 'dominated' by the alternative.
Double-blind study	A study in which neither the subject (patient) nor the observer investigator/ clinician) is aware of which treatment or intervention the subject is receiving. The purpose of blinding is to protect against bias.
Drop-out	A participant who withdraws from a trial before the end.
Economic evaluation	An economic evaluation is used to assess the cost effectiveness of healthcare interventions (that is, to compare the costs and benefits of a healthcare intervention to assess whether it is worth doing). The aim of an economic evaluation is to maximise the level of benefits – health effects – relative to the resources available. It should be used to inform and support the decision-making process; it is not supposed to replace the judgement of healthcare professionals. There are several types of economic evaluation: cost–benefit analysis, cost–consequence analysis, cost-effectiveness analysis, cost-minimisation analysis and cost–utility analysis. They use similar methods to define and evaluate costs, but differ in the way they estimate the benefits of a particular drug, programme or intervention.
Effect (as in effect measure, treatment effect, estimate of effect, effect size)	A measure that shows the magnitude of the outcome in 1 group compared with that in a control group. For example, if the absolute risk reduction is shown to be 5% and it is the outcome of interest, the effect size is 5%. The effect size is usually tested, using statistics, to find out how likely it is that the effect is a result of the treatment and has not just happened by chance.
Effectiveness	How beneficial a test or treatment is under usual or everyday conditions.
Effectiveness reviews	Evaluation of how beneficial a test or treatment is under everyday conditions.
Efficacy	How beneficial a test, treatment or public health intervention is under ideal conditions (for example in a laboratory).
Endometritis	Inflammation of the lining of the uterus, usually caused by infection.
Epidemiological study	The study of a disease within a population, defining its incidence and prevalence and examining the roles of external influences (for example infection, diet) and interventions.
EQ-5D (EuroQoL 5 dimensions) or EQ-VAS	A standardised instrument used to measure health-related quality of life. It provides a single index value for health status.
EQ-VAS (EuroQoL Visual analogue scale)	A standardised instrument used to measure health-related quality of life, using a visual analogue scale.

Term	Definition
Evidence	Information on which a decision or guidance is based. Evidence is obtained from a range of sources including RCTs, observational studies, expert opinion (of clinical professionals or patients).
Evidence based	The process of systematically finding, appraising and using research findings as the basis for clinical decisions.
Evidence table	A table summarising the results of a collection of studies which, taken together, represent the evidence supporting a particular recommendation or series of recommendations in a guideline.
Exclusion criteria (clinical study)	Criteria that define who is not eligible to participate in a clinical study.
Exclusion criteria (literature review)	Explicit standards used to decide which studies should be excluded from consideration as potential sources of evidence.
Expectant management	A management approach, also called 'wait and watch', when no medical or surgical treatment is given. The aim is to see if labour will start naturally.
Extended dominance	If Option A is both more clinically effective than Option B and has a lower cost per unit of effect when both are compared with a do-nothing alternative, then Option A is said to have extended dominance over Option B. Option A is therefore more cost effective and should be preferred, other things remaining equal.
Extrapolation	An assumption that the results of studies of a specific population will also hold true for another population with similar characteristics.
False negative	A diagnostic test result that incorrectly indicates that an individual does not have the disease of interest, when they do actually have it.
False positive	A diagnostic test result that incorrectly indicates that an individual has the disease of interest, when they actually do not have it.
Fixed-effect model	In meta-analysis, a model that calculates a pooled effect estimate using the assumption that all observed variation between studies is caused by random sample variability. Studies are assumed to be estimating the same overall effect.
Follow-up	Observation over a period of time of an individual, group or initially defined population whose appropriate characteristics have been assessed in order to observe changes in health status or health-related variables.
Forest plot	A graphical representation of the individual results of each study included in a meta-analysis together with the combined meta-analysis result. The plot also allows readers to see the heterogeneity among the results of the studies. The results of individual studies are shown as squares centred on each study's point estimate. A horizontal line runs through each square to show each study's confidence interval. The overall estimate from the meta-analysis and its confidence interval are shown at the bottom, represented as a diamond. The centre of the diamond represents the pooled point estimate, and its horizontal tips represent the confidence interval.
Freestanding midwifery unit (FMU)	A place to give birth where the birth is overseen by midwives, and which is separate from or at a different location to an obstetric unit.
Generalisability	The extent to which the results of a study hold true for groups that did not participate in the research.
Gestational age	The length of a pregnancy, measured in weeks from the first day of the last menstrual cycle.
Gold standard	A method, procedure or measurement that is widely accepted as being the best available to test for or treat a disease.

Term	Definition
GRADE, GRADE profile	A system developed by the GRADE Working Group to address the short-comings of present grading systems in healthcare. The GRADE system uses a common, sensible and transparent approach to grading the quality of evidence. The results of applying the GRADE system to clinical trial data are displayed in a table known as a GRADE profile.
Harms	Adverse effects of an intervention.
Health economics	Study or analysis of the cost of using and distributing healthcare resources.
Health-related quality of life (HRQoL)	A measure of the effects of an illness to see how it affects someone's day-to-day life.
Heterogeneity	The term is used in meta-analyses and systematic reviews to describe when the results of a test or treatment (or estimates of its effect) differ
Hyperstimulation	This is overactivity of the uterus. It is variously defined as uterine tachysystole (more than 5 contractions per 10 minutes for at least 20 minutes) and uterine hypersystole/hypertonicity (a contraction lasting at least 2 minutes). These may or may not be associated with changes in the fetal heart rate pattern (persistent decelerations, tachycardia or increased/decreased short term variability).
Imprecision	Results are imprecise when studies include relatively few patients and few events and thus have wide confidence intervals around the estimate of effect.
Incidence	The incidence of a disease is the rate at which new cases occur in a population during a specified period.
Inclusion criteria (clinical study)	Specific criteria that define who is eligible to participate in a clinical study.
Inclusion criteria (literature review)	Explicit criteria used to decide which studies should be considered as potential sources of evidence.
Incremental cost	The extra cost linked to using one test or treatment rather than another. Or the additional cost of doing a test or providing a treatment more frequently.
Incremental cost effectiveness ratio (ICER)	The difference in the mean costs in the population of interest divided by the differences in the mean outcomes in the population of interest for one treatment compared with another.
Incremental net benefit (INB)	The value (usually in monetary terms) of an intervention net of its cost compared with a comparator intervention. The INB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the INB is calculated as: (£20,000×QALYs gained) minus incremental cost.
Indirectness	The available evidence is different to the review question being addressed, in terms of population, intervention, comparison and outcome (PICO).
Instrumental birth (or delivery)	The birth of a baby which requires assistance, involving either a ventouse suction cap or forceps.
Intention-to-treat analysis (ITT)	An assessment of the people taking part in a clinical trial, based on the group they were initially (and randomly) allocated to. This is regardless of whether or not they dropped out, fully complied with the treatment or switched to an alternative treatment. Intention-to-treat analyses are often used to assess clinical effectiveness because they mirror actual practice: that is, not everyone complies with treatment and the treatment people receive may be changed according to how they respond to it.
Internal validity	How well an experiment is done and if it is clear that the variable being tested is what is causing the measured effect.

Term	Definition
Intervention	In medical terms this could be a drug treatment, surgical procedure, diagnostic or psychological therapy. Examples of public health interventions could include action to help someone to be physically active or to eat a more healthy diet.
Length of stay	The total number of days a patient stays in hospital.
Licence	See Product licence.
Likelihood ratio	The likelihood ratio combines information about the sensitivity and specificity. It tells you how much a positive or negative result changes the likelihood that a patient would have the disease. The likelihood ratio of a positive test result (LR+) is sensitivity divided by (1 minus specificity).
Lost to follow-up	Patients who have withdrawn from the clinical trial at the point of follow-up.
Markov model	A method for estimating long-term costs and effects for recurrent or chronic conditions, based on health states and the probability of transition between them within a given time period (cycle).
Mean	An average value, calculated by adding all the observations and dividing by the number of observations.
Mean difference	In meta-analysis, a method used to combine measures on continuous scales (such as weight), where the mean, standard deviation and sample size in each group are known. The weight given to the difference in means from each study (for example how much influence each study has on the overall results of the meta-analysis) is determined by the precision of its estimate of effect.
Median	The value of the observation that comes half-way when the observations are ranked in order.
Meta-analysis	A method often used in systematic reviews. Results from several studies of the same test or treatment are combined to estimate the overall effect of the treatment.
Methodology	Systematic, theoretical analysis of the methods applied to a field of study.
Minimal important difference (MID)	Threshold for clinical importance which represents the minimal important difference for benefit or for harm; for example the threshold at which drug A is less effective than drug B by an amount that is clinically important to patients.
Mixed parity	A study result which include both nulliparous and multiparous women
Morbidity	A diseased condition or state
Multidisciplinary team	A team with members from different healthcare professions (including for example, oncology, pathology, radiology, nursing)
Multiparous	A woman who has given birth to at least one child.
Multivariate model	A statistical model for analysis of the relationship between 2 or more predictors, (independent) variables and the outcome (dependent) variable.
Net monetary benefit (NMB)	The value (usually in monetary terms) of an intervention net of its cost. The NMB can be calculated for a given cost-effectiveness (willingness to pay) threshold. If the threshold is £20,000 per QALY gained then the NMB is calculated as: (£20,000×QALYs gained) minus cost.
Network meta-analysis (NMA)	Meta-analysis in which multiple treatments (that is, 3 or more) are being compared using both direct comparisons of interventions within RCTs and indirect comparisons across trials based on a common comparator.
Non-randomised	When subjects of a study are not allocated to a specific treatment/group at random.

Term	Definition
Number needed to treat (NNT)	The average number of patients who need to be treated to get a positive outcome. For example, if the NNT is 4, then 4 patients would have to be treated to ensure 1 of them gets better. The closer the NNT is to 1, the better the treatment. For example, if you give a stroke prevention drug to 20 people before 1 stroke is prevented, the number needed to treat is 20.
Nulliparous	A woman who has never given birth.
Observational study	Individuals or groups are observed or certain factors are measured. No attempt is made to affect the outcome. For example, an observational study of a disease or treatment would allow 'nature' or usual medical care to take its course. Changes or differences in one characteristic (for example whether or not people received a specific treatment or intervention) are studied without intervening. There is a greater risk of selection bias than in experimental studies.
Obstetric unit (OU)	A place to give birth which is in a hospital environment and where a medical team (obstetricians, anaesthetists and neonatologists/paediatricians) and medical equipment are available, as well as access to operating theatres.
Occult	Hidden, or difficult to observe directly
Odds ratio (OR)	<p>Odds are a way to represent how likely it is that something will happen (the probability). An odds ratio compares the probability of something in one group with the probability of the same thing in another.</p> <p>An odds ratio of 1 between 2 groups would show that the probability of the event (for example a person developing a disease, or a treatment working) is the same for both. An odds ratio greater than 1 means the event is more likely in the first group. An odds ratio less than 1 means that the event is less likely in the first group.</p> <p>Sometimes probability can be compared across more than 2 groups – in this case, one of the groups is chosen as the 'reference category' and the odds ratio is calculated for each group compared with the reference category. For example, to compare the risk of dying from lung cancer for non-smokers, occasional smokers and regular smokers, non-smokers could be used as the reference category. Odds ratios would be worked out for occasional smokers compared with non-smokers and for regular smokers compared with non-smokers.</p> <p>See also Confidence interval, Relative risk.</p>
Opportunity cost	The loss of other healthcare programmes displaced by investment in or introduction of another intervention. This may be best measured by the health benefits that could have been achieved had the money been spent on the next best alternative healthcare intervention.
Outcome	The impact that a test, treatment, policy, programme or other intervention has on a person, group or population. Outcomes from interventions to improve the public's health could include changes in knowledge and behaviour related to health, societal changes (for example a reduction in crime rates) and a change in people's health and wellbeing or health status. In clinical terms, outcomes could include the number of patients who fully recover from an illness or the number of hospital admissions, and an improvement or deterioration in someone's health, functional ability, symptoms or situation. Researchers should decide what outcomes to measure before a study begins.
p value	The p value is a statistical measure that indicates whether or not an effect is statistically significant. For example, if a study comparing 2 treatments found that one seems more effective than the other, the p value is the probability of obtaining these results by chance. By convention, if the p value is below 0.05 (that is, there is less than a 5%

Term	Definition
	probability that the results occurred by chance) it is considered that there probably is a real difference between treatments. If the p value is 0.001 or less (less than a 1% probability that the results occurred by chance), the result is seen as highly significant. If the p value shows that there is likely to be a difference between treatments, the confidence interval describes how big the difference in effect might be.
Pairwise analysis	A process of comparing entities in pairs to judge which of each entity is preferred, or has a greater amount of some quantitative property.
Parity	The number of times a woman has given birth
Performance bias	Systematic differences between intervention groups in care provided apart from the intervention being evaluated. Blinding of study participants (both the recipients and providers of care) is used to protect against performance bias.
Perineum	The area between the anus and genitals, extending from the vaginal opening to the anus.
Placebo	A fake (or dummy) treatment given to participants in the control group of a clinical trial. It is indistinguishable from the actual treatment (which is given to participants in the experimental group). The aim is to determine what effect the experimental treatment has had over and above any placebo effect caused because someone has received (or thinks they have received) care or attention.
Placebo effect	A beneficial (or adverse) effect produced by a placebo and not due to any property of the placebo itself.
Post-hoc analysis	Statistical analyses that are not specified in the trial protocol and are generally suggested by the data.
Postpartum haemorrhage	Heavy bleeding after birth. PPH can be primary or secondary: <ul style="list-style-type: none"> • Primary PPH is the loss of 500 mL or more of blood within the first 24 hours after the birth. Primary PPH can be minor (500–1000 mL loss) or major (more than 1000 mL loss). • Secondary PPH is abnormal or heavy vaginal bleeding between 24 hours and 12 weeks after the birth.
Power (statistical)	The ability to demonstrate an association when one exists. Power is related to sample size; the larger the sample size, the greater the power and the lower the risk that a possible association could be missed.
Premature rupture of membranes (PROM)	The rupture of the amniotic sac before labour begins. Also known as 'waters breaking'. If this occurs before 37 weeks it is defined as preterm prelabour rupture of membranes (PPROM)
Prevalence	The prevalence of a disease is the proportion of a population that are cases at a point in time.
Primary care	Healthcare provided in a community setting, outside hospitals (secondary care). It includes a range of services provided by GPs, nurses, health visitors, midwives and other healthcare professionals and allied health professionals such as dentists, pharmacists and opticians.
Primary outcome	The outcome of greatest importance, usually the one in a study that the power calculation is based on.
Primiparous	A woman who is giving birth for the first time
Product licence	An authorisation from the Medicines and Healthcare Products Regulatory Agency (MHRA) to market a medicinal product, and which is defined in the Summary of Product Characteristics.
Prognosis	A probable course or outcome of a disease. Prognostic factors are patient or disease characteristics that influence the course. Good prognosis is associated with low rate of undesirable outcomes; poor prognosis is associated with a high rate of undesirable outcomes.

Term	Definition
Prognostic factors	Disease characteristics that influence the course of the disease and which are used to predict the likely outcome
Prospective study	A research study in which the health or other characteristic of participants is monitored (or 'followed up') for a period of time, with events recorded as they happen. This contrasts with retrospective studies.
Protocol (review)	A document written prior to commencing a review that details exactly how evidence to answer a review question will be obtained and synthesised. It defines in detail the population of interest, the interventions, the comparators/controls and the outcomes of interest (PICO).
Publication bias	Publication bias occurs when researchers publish the results of studies showing that a treatment works well and don't publish those showing it did not have any effect. If this happens, analysis of the published results will not give an accurate idea of how well the treatment works. This type of bias can be assessed by a funnel plot.
Quality adjusted life year (QALY)	A measure of the state of health of a person or group in which the benefits, in terms of length of life, are adjusted to reflect the quality-of-life. One QALY is equal to 1 year of life in perfect health. QALYS are calculated by estimating the years of life remaining for a patient following a particular treatment or intervention and weighting each year with a quality-of-life score (on a scale of 0 to 1). It is often measured in terms of the person's ability to perform the activities of daily life, and freedom from pain and mental disturbance.
Quality of life	See Health-related quality of life.
Random effect model	In meta-analysis, a model that calculates a pooled effect estimate using the assumption that each study is estimating a different true treatment effect due to real differences between studies. Observed variation in effects are therefore caused by a combination of random sample variability (within-study variation) and heterogeneity between studies (between-study variation). The overall effects is an average of the estimated true study effects.
Randomisation	Assigning participants in a research study to different groups without taking any similarities or differences between them into account. For example, it could involve using a random numbers table or a computer-generated random sequence. It means that each individual (or each group in the case of cluster randomisation) has the same chance of receiving each intervention.
Randomised controlled trial (RCT)	A study in which a number of similar people are randomly assigned to 2 (or more) groups to test a specific drug or treatment. One group (the experimental group) receives the treatment being tested, the other (the comparison or control group) receives an alternative treatment, a dummy treatment (placebo) or no treatment at all. The groups are followed up to see how effective the experimental treatment was. Outcomes are measured at specific times and any difference in response between the groups is assessed statistically. This method is also used to reduce bias.
Recruitment bias	When proper randomisation is not achieved when recruiting individuals, meaning that the sample obtained may not be representative of the population intended to be analysed.
Reference standard	The test that is considered to be the best available method to establish the presence or absence of the outcome – this may not be the one that is routinely used in practice.
Regimen	A plan or regulated course of treatment

Term	Definition
Relative risk (RR)	The ratio of the risk of disease or death among those exposed to certain conditions compared with the risk for those who are not exposed to the same conditions (for example the risk of people who smoke getting lung cancer compared with the risk for people who do not smoke). If both groups face the same level of risk, the relative risk is 1. If the first group had a relative risk of 2, subjects in that group would be twice as likely to have the event happen. A relative risk of less than 1 means the outcome is less likely in the first group. Relative risk is sometimes referred to as risk ratio.
Reporting bias	See Publication bias.
Resource implication	The likely impact in terms of finance, workforce or other NHS resources.
Retrospective study	A research study that focuses on the past and present. The study examines past exposure to suspected risk factors for the disease or condition. Unlike prospective studies, it does not cover events that occur after the study group is selected.
Review question	The plan or set of steps to be followed in a study. A protocol for a systematic review describes the rationale for the review, the objectives and the methods that will be used to locate, select and critically appraise studies, and to collect and analyse data from the included studies.
Rhombus of Michaelis	The kite-shaped area on the lower back, marked by the two dimples of venus on either side, the lower end of the crease over the spine, and the top of the gluteal crease
Ruptured membranes	The rupture of the amniotic sac, which normally occurs spontaneously at the beginning or in the early stages of labour. Known as 'the waters breaking'.
Secondary care	Services provided by multidisciplinary team in the hospital, as opposed to the General Practitioner and the primary care team
Secondary outcome	An outcome used to evaluate additional effects of the intervention deemed a priori as being less important than the primary outcomes.
Selection bias	Selection bias occurs if: <ul style="list-style-type: none"> • the characteristics of the people selected for a study differ from the wider population from which they have been drawn; or • there are differences between groups of participants in a study in terms of how likely they are to get better.
Sensitivity	How well a test detects the thing it is testing for. If a diagnostic test for a disease has high sensitivity, it is likely to pick up all cases of the disease in people who have it (that is, give a 'true positive' result). But if a test is too sensitive it will sometimes also give a positive result in people who don't have the disease (that is, give a 'false positive'). For example, if a test were developed to detect if a woman is 6 months pregnant, a very sensitive test would detect everyone who was 6 months pregnant but would probably also include those who are 5 and 7 months pregnant. If the same test were more specific (sometimes referred to as having higher specificity), it would detect only those who are 6 months pregnant and someone who was 5 months pregnant would get a negative result (a 'true negative'). But it would probably also miss some people who were 6 months pregnant (that is, give a 'false negative'). Breast screening is a 'real-life' example. The number of women who are recalled for a second breast screening test is relatively high because the test is very sensitive. If it were made more specific, people who don't have the disease would be less likely to be called back for a second test but more women who have the disease would be missed.
Sensitivity analysis	A means of representing uncertainty in the results of an analysis. Uncertainty may arise from missing data, imprecise estimates or

Term	Definition
	<p>methodological controversy. Sensitivity analysis also allows for exploring the generalisability of results to other settings. The analysis is repeated using different assumptions to examine the effect on the results.</p> <ul style="list-style-type: none"> • One-way simple sensitivity analysis (univariate analysis) – each parameter is varied individually in order to isolate the consequences of each parameter on the results of the study. • Multi-way simple sensitivity analysis (scenario analysis) – 2 or more parameters are varied at the same time and the overall effect on the results is evaluated. • Threshold sensitivity analysis – the critical value of parameters above or below which the conclusions of the study will change are identified. • Probabilistic sensitivity analysis – probability distributions are assigned to the uncertain parameters and are incorporated into evaluation models based on decision analytical techniques (for example Monte Carlo simulation).
Shoulder dystocia	When either the anterior, or less commonly the posterior, fetal shoulder impacts on the maternal symphysis, or sacral promontory, respectively and delays the delivery of the baby's body.
Significance (statistical)	A result is deemed statistically significant if the probability of the result occurring by chance is less than 1 in 20 ($p < 0.05$).
Significant meconium	This is defined as dark green or black amniotic fluid that is thick or tenacious, or any meconium-stained amniotic fluid containing lumps of meconium.
Single blind study	A study in which either the subject (patient/participant) or the observer (clinician/investigator) is not aware of which treatment or intervention the subject is receiving.
Specificity	<p>The proportion of true negatives that are correctly identified as such. For example, in diagnostic testing the specificity is the proportion of non-cases correctly diagnosed as non-cases. In terms of literature searching a highly specific search is generally narrow and aimed at picking up the key papers in a field and avoiding a wide range of papers.</p> <p>See also Sensitivity.</p>
Stakeholder	<p>An organisation with an interest in a topic on which NICE is developing a clinical guideline or piece of public health guidance. Organisations that register as stakeholders can comment on the draft scope and the draft guidance. Stakeholders may be:</p> <ul style="list-style-type: none"> • manufacturers of drugs or equipment • national patient and carer organisations • NHS organisations • organisations representing healthcare professionals.
Standard deviation (SD)	A measure of the spread or dispersion of a set of observations, calculated as the average difference from the mean value in the sample.
Subgroup analysis	An analysis in which the intervention effect is evaluated in a defined subset of the participants in a trial, or in complementary subsets.
Systematic review	A review in which evidence from scientific studies has been identified, appraised and synthesised in a methodical way according to predetermined criteria. It may include a meta-analysis.
Systemic therapy/treatment	Medicine, usually given by mouth or injection, to treat the whole body rather than targeting one specific area
Time horizon	The time span over which costs and health outcomes are considered in a decision analysis or economic evaluation.
Treatment allocation	Assigning a participant to a particular arm of a trial.

Term	Definition
True negative	A diagnostic test result that correctly indicates that an individual does not have the disease of interest when they actually do not have it.
True positive	A diagnostic test result that correctly indicates that an individual has the disease of interest when they do actually have it.
Univariate	Analysis which separately explores each variable in a data set.
Uterotonics	Pharmacological agents used to induce contractions or greater tonicity of the uterus. Uterotonics are used both to induce labour and to reduce postpartum haemorrhage.
Utility	In health economics, a utility is the measure of the preference or value that an individual or society places upon a particular health state. It is generally a number between 0 (representing death) and 1 (perfect health). The most widely used measure of benefit in cost-utility analysis is the quality-adjusted life year, but other measures include disability-adjusted life years (DALYs) and healthy year equivalents (HYEs).

Abbreviations

Abbreviation	Definition
AE	Adverse event
AMSTAR	Assessing the Methodological Quality of Systematic Reviews
AMU	Alongside midwifery unit
aOR	Adjusted odds ratio
aRR	Adjusted risk ratio
AUC	Area under the curve
AUROC	Area under the receiver operating curve
BID	Twice a day
BMI	Body mass index
BNF	British National Formulary
CASP	Critical Appraisal Skills Programme
CB	Caesarean birth
CDSR	Cochrane Database of Systematic Reviews
CEAC	Cost-effectiveness acceptability curves
CEI	Continuous epidural infusion
CENTRAL	Cochrane Central Register of Controlled Trials
CG	Clinical guideline
CI	Confidence interval; continuous infusion
CrI	Credible interval
CS	Caesarean section
DARE	Database of Abstracts of Reviews of Effects
DIC	Deviance information criterion
eMIT	Electronic market information tool
EQ-VAS	EuroQoL Visual Analog Scale
FMU	Freestanding midwifery unit
FN	False negative
FP	False positive
g	Gramme
GA	Gestational age
GC	Guideline committee
GP	General Practitioner
GRADE	Grading of Recommendations Assessment, Development and Evaluation
Hb	Haemoglobin
h, hr	Hour
HIE	Hypoxic ischaemic encephalopathy
HR	Hazard ratio, heart rate
HRG	Healthcare Resource Group
HRQoL	Health-related quality of life
HTA	Health Technology Assessment
ICER	Incremental cost-effectiveness ratio
IHTA	International Health Technology Assessment
IM	Intramuscular

Abbreviation	Definition
IPD	Individual patient data
IQR	Interquartile range
ITT	Intention to treat
ITU	Intensive therapy unit
IU	International units
IV	Intravenous
k	Number of studies or publications
LOR	Logs odds ratio
LR-	Negative likelihood ratio
LR+	Positive likelihood ratio
M	Mean
MD	Mean difference
mg	Milligram
MID	Minimally important difference
mL	Millilitre
mmHg	Millimetres of mercury
mmol	Millimole
MR	Mean ratio
N, n	Number of participants
N/A	Not applicable
N/C	Not calculable
NGA	National Guideline Alliance
NHS	National Health Service
NICE	National Institute of Health and Care Excellence
NICU	Neonatal intensive care unit
NIHR	National Institute of Health Research
NMA	Network meta-analysis
NMB	Net monetary benefit
NMPA	National maternal and perinatal audit
NNH	Number needed to harm
NNT	Number needed to treat
NUU	Neonatal unit
NR, nr	Not reported
NRCT	Non-randomised controlled trial
ns	Not significant
OD	Once a day
O:E	Observed: expected
O-E	Observed minus expected
OECD	Organisation for economic co-operation and development
ONS	Office for National Statistics
OR	Odds ratio
OU	Obstetric unit
PCA	Patient-controlled analgesia
PCEA	Patient-controlled epidural analgesia

Abbreviation	Definition
PICO	Population, intervention, comparison, outcome
PICOTS	Population, intervention, comparator, outcome, timing and setting
PIEB	Programmed intermittent epidural bolus
PO	By mouth, orally
POR	Peto odds ratio
PPH	Postpartum haemorrhage
PRESS	Peer review of electronic search strategies
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
(P)PROM	(Preterm) prelabour rupture of membranes
PSA	Probabilistic sensitivity analysis
QALY	Quality-adjusted life year
QoL	Quality of life
RCOG	Royal College of Obstetricians and Gynaecologists
RCT	Randomised controlled trial
RD	Risk difference
ROBINS-I	Risk of bias in non-randomized studies of interventions
ROB(IS)	Risk of bias (in systematic reviews)
ROC	Receiver operating characteristics
RR	Relative risk/risk ratio
SCBU	Special care baby unit
SD	Standard deviation
SE	Standard error
Sens	Sensitivity
Spec	Specificity
SR	Systematic review
TA	Technology appraisal
TENS	Transcutaneous electrical nerve stimulation.
TID	Three times a day
TN	True negative
TP	True positive
TSU	Technical support unit
TXA	Tranexamic acid
UKMidSS	UK Midwifery study system
UME	Unrelated mean effects
VAS	Visual analogue scale
wk	Week