

DRAFT FOR CONSULTATION

Fractures (complex): assessment and management

Complex fractures: assessment and management of
complex fractures

Clinical Guideline <...>

Appendices I - P

August 2015

Draft for Consultation

*Commissioned by the National Institute for
Health and Care Excellence*

Disclaimer

Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and/or their guardian or carer.

Copyright

National Clinical Guideline Centre, 2015

Funding

National Institute for Health and Care Excellence

Contents

Appendices.....	5
Appendix I: Forest plots	5
Appendix J: Excluded clinical studies	24
Appendix K: Excluded economic studies.....	45
Appendix L: Cost analysis for open fractures	46
Appendix M: Research recommendations	68
Appendix N: NICE technical team.....	72
Appendix O: Additional cost data.....	73
Appendix P: Qualitative study checklist.....	81
References.....	83

1

Appendices

2

Appendix I: Forest plots

3

I.1 Open fractures


4

I.1.1 Limb salvage

5


Secondary upper limb amputation in adults

Figure 1: MESS in detecting the need for secondary upper limb amputation in adults



6


Figure 2: MESI in detecting need for secondary upper limb amputation in adults



7


Secondary lower limb amputation in adults

Figure 3: MESS in detecting the need for secondary lower limb amputation in adults




8

Figure 4: MESI in detecting the need for secondary lower limb amputation in adults




9

Figure 5: PSI in detecting need for secondary lower limb amputation in adults




1

Figure 6: LSI in detecting the need for secondary lower limb amputation in adults




2

Figure 7: NISSA in detecting the need for secondary lower limb amputation in adults




3

Figure 8: Ganga in detecting the need for secondary lower limb amputation in adults



4


Figure 9: HFS '97 in detecting the need for secondary lower limb amputation in adults



5


Primary/secondary upper limb amputation in children

Figure 10: MESS in detecting the need for primary/secondary upper limb amputation in children




6

Figure 11: MESI in detecting need for primary/secondary lower limb amputation in children




1

Figure 12: LSI in detecting need for primary/secondary lower limb amputation in children




2

Figure 13: PSI in detecting need for primary/secondary lower limb amputation in CHILDREN




3

Figure 14: HFS '98 in detecting need for primary/secondary lower limb amputation in children



4


Figure 15: NISSA in detecting need for primary/secondary lower limb amputation in children



5


Primary/secondary lower limb amputation in adults

Figure 16: MESS in detecting need for primary/secondary lower limb amputation in adults




6

Figure 17: Ganga in detecting need for primary/secondary lower limb amputation in adults




1

Figure 18: PSI in detecting need for primary/secondary lower limb amputation in adults




2

Figure 19: NISSA in detecting need for primary/secondary lower limb amputation in adults




3

Figure 20: LSI in detecting need for primary/secondary lower limb amputation in adults




4

Figure 21: HFS '98 in detecting need for primary/secondary lower limb amputation in adults



5


Figure 22: HFS/HFS '97 in detecting need for primary/secondary lower limb amputation in adults



1 **I.1.2 Arterial shunts**


2 **Shunt versus definitive vascular repair**

Figure 23: Mortality




3

Figure 24: Amputation




4

Figure 25: Compartment syndrome



5


Figure 26: Other vascular surgery



6 **I.1.3 MDT**


7 **Combined orthoplastic approach versus non-combined approach**

Figure 27: Amputation




1

Figure 28: flap failure




2

Figure 29: Deep infection



3


Figure 30: Enneking limb score (higher better)



1 **I.1.4 Optimal timing of debridement**

2 **Deep surgical site infection**

Figure 31: Early versus delayed/late debridement multivariate analysis results (odds ratio)




Test for subgroup differences: Chi² = 0.08, df = 1 (P = 0.78), I² = 0%

Noumi 2005 is adjusted for: age, sex, Gustilo type, fracture grade by AO type, fracture site, reamed versus unreamed nailing, existence of multiple trauma and existence of floating knee injury. Harley 2002 is adjusted for: male gender, age and Gustilo grade.

3

Figure 32: Early versus delayed/late debridement multivariate analysis results (relative risk)




Test for subgroup differences: Not applicable

Adjusted for: The entire data set was used. This was assumed to be the content of the baseline characteristics table; age, ISS, RTS, SBP, lactate and Gustilo grade.

4


Figure 33: Delayed versus earlier debridement multivariate analysis results (adjusted OR)



1

Amputation

Figure 34: Debridement on hospital day 0 versus other timings in open tibial fractures, multivariate analysis results




Adjusted for: age, sex, race, economic characteristics, injury severity scale score, comorbidities, associated injuries/procedure (arterial injury, tibial nerve injury, complicated open wound, fasciotomy, dislocation (knee or ankle)), admission type, location, bed size, hospital teaching status, hospital volume open tibial fractures per year, median household income and mechanism of injury.

2 I.1.5 Fixation


3 Definitive fixation and immediate cover versus definitive fixation and staged cover

Figure 35: Deep infection – RCT results




4

Figure 36: Deep infection - cohorts




1

Figure 37: Amputation - cohorts



2


Figure 38: Further unplanned surgery



3


Definitive fixation and immediate cover versus staged fixation and staged cover

Figure 39: Flap failure (total or partial) -cohort



4


Figure 40: Deep infection – cohort



1 **I.1.6 Cover**


2 **Immediate versus 3 days**

Figure 41: Deep infection




3 **Immediate versus 7 days**

Figure 42: Deep infection (RCT)



4 **Figure 43: Deep infection (cohorts)**



5 **Figure 44: Amputation (cohorts)**




1

2

Immediate versus more than 7 days


Figure 45: Infection (not specified if deep)



3

More than 14 days versus less than 3 days

Figure 46: Deep infection



4

Figure 47: Osteomyelitis






Figure 48: Flap take backs




1 **More than 5 days versus less than 5 days**

Figure 49: Deep infection



2 **Timing as a continuous variable**


Figure 50: Odds of deep infection per day delay in cover



1 **I.1.7 Definitive dressings after debridement**


2 **NPWT versus standard dressing**

Figure 51: Deep infection



3


Figure 52: Wound healed at 30 days



Appearance of 100% granulation tissue over the wound

4

Figure 53: Quality of life at 3 months (SF36 physical component)




1 **I.2 Pelvic fractures**

2 **I.2.1 Pelvic haemorrhage control**

3 **LAP versus TAE**

Figure 54: In-hospital mortality




4 **I.3 Pilon fractures**

5 **I.3.1 Pilon early fixation**

6 **MIXED OPEN/CLOSED**


7 **Definitive fixation within 24 hours versus temp fixation and definitive fixation at more than 7 days**

Figure 55: Number of surgeries




8

Figure 56: Function - AOFAS




9

Figure 57: Function - SMFA




1

Figure 58: People with unplanned surgery



2

Figure 59: Return to normal activities




3

Temp fixation and definitive fixation at more than 24 hours to 7 days versus temp fixation and definitive fixation at more than 7 days


4

Figure 60: Deep infection




5

Figure 61: Unplanned surgery




6

Figure 62: Function - foot function index



7

Figure 63: Function – musculoskeletal function assessment score




1 **CLOSED only**

2 **Temp fixation and definitive fixation at more than 24 hours to 7 days versus temp fixation and**
3 **definitive fixation at more than 7 days**


4

Figure 64: Deep infection




5

Figure 65: Function (poor/fair)n



6


Figure 66: Hospital stay (days)



1 **I.3.2 Pilon fixation**


2 **RCT data**

Figure 67: Surgical site infection




3

Figure 68: Osteomyelitis




4

Figure 69: Ankle Fusion




5

Figure 70: Unplanned further surgery (continuous)




6

Figure 71: Unplanned further surgery (dichotomous)




1

Figure 72: Wound breakdown



2


Figure 73: Amputation



3

Cohort Data

Figure 74: Health-related quality of life (SF-36 functional Score)



I.4 Other

I.4.1 Detecting compartment syndrome

Diagnostic RCT review

Continuous compartment pressure monitoring versus no compartment pressure monitoring

Figure 75: Sensory loss





Figure 76: Contracture



Appendix J: Excluded clinical studies

J.1 Open fractures

J.1.1 Limb salvage

Table 1: Studies excluded from the clinical review

Reference	Reason for exclusion
Adegbehingbe, 2006 ⁴	No usable accuracy data
Agel, 2014 ⁵	No accuracy data
Bevevino, 2014 ⁴²	Modelling study
Bosse, 2002 ⁵⁰	No accuracy data
Clement, 2014 ⁶⁹	No accuracy data
Dua, 2014A ⁹⁶	No accuracy data
Dua, 2014 ⁹⁵	Not assessing accuracy of prediction tools
Fochtman, 2014 ¹¹⁶	No accuracy data
Fodor, 2012 ¹¹⁷	Review – reference list examined
Gregory, 1985 ¹³⁴	Developmental study
Guraya, 2004 ¹⁴⁰	No accuracy data for salvage group
Hierner, 1995 ¹⁶³	No accuracy data
Higgins, 2010 ¹⁶⁴	Review –references examined
Hoogendoorn, 2002 ¹⁷²	Review –references examined
Howe, 1987 ¹⁷⁸	Developmental study
Krettek, 2001a ²²³	Erratum – related to author names
Ly, 2008 ²⁴²	Not predicting amputation
Mackenzie, 2006 ²⁴⁴	Not assessing accuracy of prediction tools
Osullivan, 1997 ²⁹³	No accuracy data
Poole, 1994 ³²²	Not assessing accuracy of prediction tools
Shanmuganathan, 2008 ³⁶⁵	Review –references examined
Sharma, 2003	Amputations appeared to be made on the basis of the MESS score
Swionkowski, 2002 ³⁸⁷	Developmental study
Zaraca, 2011 ⁴³³	Not an accuracy study

J.1.2 Antibiotics

Table 2: Studies excluded from the clinical review

Reference	Reason for exclusion
Alarabi 2007 ⁷	Inadequate adjustment for confounders
Alarabi 2008 ⁸	Corrigendum for ALARABI2007
Bremmer 2012 ⁵²	Abstract
Gonzalez 2014 ¹²⁸	Did not look at time of antibiotic administration
Grote 2012 ¹³⁶	Not in English and was not ordered
Hatfield 2012 ¹⁵³	Conference abstract
Hauser 2006 ¹⁵⁴	Review, not systematic

Reference	Reason for exclusion
Hughes 1993 ¹⁸⁰	Review, not systematic
Leonidou 2014 ²³⁴	No adjustment for key confounders; age and contamination data given but not for each group
Malik 2004 ²⁴⁸	Study does not report outcomes by timing of antibiotics
Mccaul 2013 ²⁵⁹	Conference abstract
Patzakis 1983 ³¹¹	Study does not report outcomes by timing of antibiotics
Rojczyk 1979 ³⁴⁶	Not in English
Thomas 2013 ³⁹²	Study does not report outcomes by timing of antibiotics
Zumsteg 2014 ⁴⁴⁰	Unclear MVA inputs. No baseline characteristics provided.

J.1.3 Dressings before debridement

Table 3: Studies excluded from the clinical review

Study	Reason for exclusion
Back 2013 ²³	Systematic review is not relevant to review question or unclear PICO
Blum 2012 ⁴⁹	Incorrect interventions. Post-debridement treatments
Calhoun 1993 ⁶⁰	Not review population. Chronic osteomyelitis
Contractor 2008 ⁷¹	Systematic review is not relevant to review question or unclear PICO
Halvorson 2011 ¹⁴⁴	Incorrect study design. Non-comparative study
Kazakos 2009 ²⁰⁶	Incorrect interventions. Plasma rich platelet gel
Keating 1996 ²⁰⁷	Incorrect interventions. Post-debridement treatment
Keen 2012 ²¹⁰	Incorrect study design
Liu 2012 ²³⁷	Incorrect interventions. After debridement
Moehring 2000 ²⁷⁶	Incorrect interventions. Took place after debridement
Moues 2004 ²⁸¹	Not guideline condition
Ogbemudia 2010 ²⁹⁷	Not review population
Rasool 2013 ³³²	Dressings applied after debridement
Rinker 2008 ³⁴²	Incorrect study design
Runkel 2011 ³⁵⁴	Systematic review is not relevant to review question or unclear PICO
Stannard 2009 ³⁸⁰	Incorrect interventions. After debridement
Stannard 2010 ³⁷⁹	Study design not relevant to review. Review
Tang 2010 ³⁸⁹	Incorrect interventions. Post-debridement treatment
Wright 2007 ⁴²³	Incorrect study design
Yuenyongviwat 2011 ⁴³⁰	Incorrect interventions. Only standard dressings

J.1.4 Arterial shunts

Table 4: Studies excluded from the clinical review

Study	Reason for exclusion
Al-salman 1997 ¹⁰	Incorrect interventions
Asensio 2006 ¹⁷	Incorrect interventions
Ball 2009 ²⁸	Incorrect interventions
Ball 2009 ²⁷	Incorrect interventions
Barros d'sa 2006 ³³	Failed to adjust for time to initial vascular repair

Study	Reason for exclusion
Cavadas 2009 ⁶⁴	Macroreplantations case series
Chambers 2006 ⁶⁵	Incorrect interventions
Fox 2008 ¹²⁰	Failed to adjust for time to initial vascular repair
Gifford 2009 ¹²³	failed to adjust for time to initial vascular repair
Granchi 2000 ¹³²	Incorrect interventions
Laohapensang 1994 ²²⁹	Incorrect interventions
Nichols 1986 ²⁸⁶	Failed to adjust for time to initial vascular repair
Reber 1999 ³³³	Incorrect interventions
Subramanian 2008 ³⁸⁴	Incorrect interventions. case series
Taller 2008 ³⁸⁸	Incorrect interventions

J.1.5 MDT

Table 5: Studies excluded from the clinical review

Reference	Reason for exclusion
British Orthopaedic Association, British Association of Plastic Surgeons. The Early Management of Severe Tibial Fractures: The Need for Combined Plastic and Orthopaedic Management : [a Report by the BOA/BAPS Working Party on Severe Tibial Injuries]. Associations; 1993. Available from: http://books.google.co.uk/books?id=swxDHQAACAAJ	Not available from any sources
Court-Brown, Cross AT, Hahn DM, Marsh DR, Willett K, Quaba AAWF et al. A report by the British Orthopaedic Association/British Association of Plastic Surgeons Working Party on the Management of Open Tibial Fractures September 1997. British Journal of Plastic Surgery. 1997; 50(8):570-583	Review article. References checked and one article ordered
Godina M. early microsurgical reconstruction of complex trauma of the extremities. Plast Reconst Surg 1986; 78:285-92	Mixed population containing people without open fractures
Green AR. The courage to co-operate: the team approach to open fractures of the lower limb. Annals of the Royal College of Surgeons of England. 1994; 76(6):365-366	Review. References checked and two articles ordered
Levin LS. The reconstructive ladder: An orthoplastic approach. Orthopedic Clinics of North America. 1993; 24(3):393-409	Review article. References checked and no articles ordered
Moda SK et al. The role of early flap coverage in the management of open fractures of both bones of the	Not relevant to the protocol

Reference	Reason for exclusion
leg. Injury 1994; 25: 83-5 (was in the search)	
Nayagam S, Graham K, Pearse M, Nanchahal J. Reconstructive surgery in limbs: the case for the orthoplastic approach. <i>Annals of Plastic Surgery</i> . 2011; 66(1):6-8	Review article. References checked and no articles ordered
Rahman S, Trickett R, Pallister I. From guidelines to standards of care: Increasing workload, but diminishing patient burden in open tibial fractures. <i>International Journal of Surgery</i> . 2013; 11(8):682	Abstract. Not relevant to protocol
Stammers J, Williams D, Hunter J, Vesely M, Nielsen D. The impact of trauma centre designation on open tibial fracture management. <i>Annals of the Royal College of Surgeons of England</i> . 2013; 95(3):184-187	Groups are not as on protocol. The 'tertiary' group contained 14/15 with a non-combined approach and the 'primary' group had 21/29 with a combined approach. There was sufficient departure from the protocol in these groups to exclude rather than downgrade for indirectness. For example, the 8 in the primary group that were not managed with a combined approach were very uncomplicated orthopaedic cases, and this may have contributed considerably to any advantage in the primary group. There was no sub-grouping of results within the study to allow us to extract only the results pertaining to the cases correlating with the protocol.

J.1.6 Optimal timing of debridement

Table 6: Studies excluded from the clinical review

Reference	Reason for exclusion
Ashford 2004 ¹⁸	Inadequate adjustment for confounders
Alarabi 2007 ⁷	Inadequate adjustment for confounders
Alhilli 2010 ⁹	Inadequate adjustment of confounders.
Arti 2012 ¹⁶	Systematic review does not meet protocol criteria.
Bednar 1993 ³⁹	Inadequate adjustment for confounders.
Dellinger 1988 ⁸⁸	Does not meet the protocol criteria.
Gougoulas 2009 ¹³¹	Systematic review does not look at debridement timing.
Ibrahim 2014 ¹⁸⁶	Systematic review does not meet protocol criteria.
Ikem 2001 ¹⁸⁷	No Surgical debridement timing comparisons
Ikem 2006 ¹⁸⁸	No Surgical debridement timing comparisons
Jacob 1992 ¹⁹⁰	No Surgical debridement timing comparisons
Khatod 2003 ²¹⁵	Inadequate adjustment of confounders.
Kindsfater 1995 ²¹⁷	Confounders are not compared between treatment groups
Kreder 1995 ²²²	Insufficient data reported by debridement time. Unclear reporting.
Kurylo 2011 ²²⁵	Inadequate adjustment of confounders.
Leonidou 2014 ²³⁴	No adjustment for key confounders; age and contamination data given but not for each group
Mclain 1991 ²⁶²	Inadequate adjustment of confounders.
Patzakis 1989 ³¹⁰	Inadequate adjustment of confounders.
Pollak 2010 ³²⁰	Does not meet the protocol criteria.

Reference	Reason for exclusion
Reuss 2007 ³³⁷	Inadequate adjustment of confounders.
Schenker 2012 ³⁶⁴	Systematic Review. References checked for eligibility
Schenker 2012 ³⁶⁴	Inadequate adjustment of confounders.
Skaggs 2000 ³⁷⁰	Inadequate adjustment for confounders.
Skaggs 2005 ³⁷¹	Inadequate adjustment for confounders.
Spencer 2004 ³⁷⁵	Inadequate adjustment of confounders.
Srouf 2015 ³⁷⁷	No multivariable analysis for Gustillo grade in the light of large differences in proportions of people with each grade across groups
Sungaran 2007 ³⁸⁵	Inadequate adjustment of confounders.
Tripuraneni 2008 ³⁹⁹	Inadequate adjustment of confounders.
Wei 2014 ⁴¹⁶	Inadequate adjustment of confounders across the debridement timing groups
Yusof 2013 ⁴³²	Inadequate adjustment of confounders.
Zumsteg 2014 ⁴⁴⁰	No baseline characteristics provided. Unclear what variables were in the logistic regression.

J.1.7 Fixation

Table 7: Studies excluded from the clinical review

Reference	Reason for exclusion
ALLISON2011 ¹¹	No comparison between immediate (early) versus staged (delayed) closure
BALDWIN2009 ²⁶	No comparison between immediate (early) versus staged (delayed) closure
BARTLETT1997 ³⁴	No comparison between immediate (early) versus staged (delayed) closure
BERRY2004 ⁴¹	No comparison between immediate (early) versus staged (delayed) closure
BHANDARI2001 ⁴³	No comparison between immediate (early) versus staged (delayed) closure
BLICK1989 ⁴⁸	No comparison between immediate (early) versus staged (delayed) closure
BREUGEM2006 ⁵³	SR does not meet our inclusion criteria
BUCKLEY1996 ⁵⁵	Inadequate adjustment for confounders
BURGESS1987 ⁵⁷	No comparison between immediate (early) versus staged (delayed) closure
BYRD1981 ⁵⁸	No comparison between immediate (early) versus staged (delayed) closure
BYRD1985 ⁵⁹	Inadequate adjustment for confounders
CAUDLE1987	Inadequate adjustment for confounders
CIERNY1983 ⁶⁷	Inadequate adjustment for confounders
COX1970 ⁷⁴	No comparison between immediate (early) versus staged (delayed) closure
CULLEN1996 ⁷⁸	No comparison between immediate (early) versus staged (delayed) closure
DALLEYRAND 2014 ⁸¹	Unknown when the fracture fixation was carried out in relation to the cover timing

Reference	Reason for exclusion
DAVISSEARS2012 ⁸³	No comparison between immediate (early) versus staged (delayed) closure
DELONG1999 ⁸⁹	Inadequate adjustment for confounders
DONG2011 ⁹³	No comparison between immediate (early) versus staged (delayed) closure
EDWARDS1983 ¹⁰¹	Not a systematic review
EDWARDS1988 ¹⁰²	No comparison between immediate (early) versus staged (delayed) closure
ERDMANN1997 ¹⁰⁸	No comparison between immediate (early) versus staged (delayed) closure
FERRERA1999 ¹¹³	No comparison between immediate (early) versus staged (delayed) closure
FISCHER1991 ¹¹⁴	No comparison between immediate (early) versus staged (delayed) closure
FRANCEL1992 ¹²¹	Inadequate adjustment for confounders
GLASS2009A ¹²⁴	No comparison between immediate (early) versus staged (delayed) closure
GODINA1986 ¹²⁶	Inadequate adjustment for confounders
GOPAL2000 ¹²⁹	Inadequate adjustment for confounders
GOUGOULIAS2009A ¹³⁰	No comparison between immediate (early) versus staged (delayed) closure
GREENBAUM2001 ¹³³	No comparison between immediate (early) versus staged (delayed) closure
GRIMARD1996 ¹³⁵	Inadequate adjustment for confounders
GUSTILO1976 ¹⁴¹	No comparison between immediate (early) versus staged (delayed) closure
HAASBEEK1995 ¹⁴³	No comparison between immediate (early) versus staged (delayed) closure
HAMMER1992 ¹⁴⁵	No comparison between immediate (early) versus staged (delayed) closure
HARLEY2002 ¹⁴⁷	Inadequate adjustment for confounders
HARRIS2006 ¹⁴⁹	No comparison between immediate (early) versus staged (delayed) closure
HARVEY2002 ¹⁵⁰	No comparison between immediate (early) versus staged (delayed) closure
HARWOOD2006 ¹⁵¹	No comparison between immediate (early) versus staged (delayed) closure
HAS1995 ¹⁵²	No comparison between immediate (early) versus staged (delayed) closure
HEE2001 ¹⁵⁵	No comparison between immediate (early) versus staged (delayed) closure
HEIER2003 ¹⁵⁶	No comparison between immediate (early) versus staged (delayed) closure
HELLAND1996 ¹⁵⁸	No comparison between immediate (early) versus staged (delayed) closure
HENLEY1998 ¹⁶⁰	No comparison between immediate (early) versus staged (delayed) closure
HERNIGOU2013 ¹⁶¹	No comparison between immediate (early) versus staged (delayed) closure

Reference	Reason for exclusion
	closure
HOFFMANN2013 ¹⁶⁷	No comparison between immediate (early) versus staged (delayed) closure
HOHMANN2007 ¹⁶⁸	Incorrect comparison
HONG1998 ¹⁷¹	No comparison between immediate (early) versus staged (delayed) closure
HOU2011 ¹⁷⁶	Inadequate adjustment for confounders
HULL2008 ¹⁸¹	Not a systematic review
HULSKER2011 ¹⁸³	Incorrect study design included
HUTCHINSON2012 ¹⁸⁴	No comparison between immediate (early) versus staged (delayed) closure
HUTSON2010 ¹⁸⁵	No comparison between immediate (early) versus staged (delayed) closure
JONES2003 ¹⁹⁵	No comparison between immediate (early) versus staged (delayed) closure
JOSHI2006 ¹⁹⁶	No comparison between immediate (early) versus staged (delayed) closure
KAI1998 ¹⁹⁸	No comparison between immediate (early) versus staged (delayed) closure
KAKAR2007 ²⁰⁰	No comparison between immediate (early) versus staged (delayed) closure
KAMATH2012 ²⁰²	Inadequate adjustment for confounders
KEELING2008 ²⁰⁹	No comparison between immediate (early) versus staged (delayed) closure
KESEMENLI2004 ²¹²	No comparison between immediate (early) versus staged (delayed) closure
KIM2012 ²¹⁶	Outcomes of the protocol not reported
KINZEL2006 ²¹⁸	Does not meet our protocol
KREDER1995 ²²²	Outcomes of the protocol not reported
KULSHRESTHA2008 ²²⁴	No comparison between immediate (early) versus staged (delayed) closure
LAUGHLIN1993 ²³⁰	No comparison between immediate (early) versus staged (delayed) closure
LENARZ2010 ²³²	No comparison between immediate (early) versus staged (delayed) closure
LEONG1988 ²³³	Inadequate adjustment for confounders
LERNER2006	No comparison between immediate (early) versus staged (delayed) closure
LOWENBERG1996 ²³⁹	No comparison between immediate (early) versus staged (delayed) closure
MACK2013 ²⁴³	No comparison between immediate (early) versus staged (delayed) closure
METSEMAKERS2015 ²⁶⁸	Did not cover the review question
MIN2011 ²⁶⁹	Inadequate adjustment for confounders
MODA1994 ²⁷²	No comparison between immediate (early) versus staged (delayed) closure
NAIQUE2006 ²⁸²	Inadequate adjustment for confounders

Reference	Reason for exclusion
PAPAKOSTIDIS2011 ³⁰⁷	No comparison between immediate (early) versus staged (delayed) closure
PARK2007 ³⁰⁸	No comparison between immediate (early) versus staged (delayed) closure
PARRETT2006 ³⁰⁹	No comparison between immediate (early) versus staged (delayed) closure
POLLAK2000 ³¹⁹	Intervention does not meet the protocol
POLLAK2010 ³²⁰	Outcomes did not meet the protocol
Radoicic 2014 ³²⁹	Comparison covered by an RCT. Cohorts with severe contamination were also used in that comparison to allow for fact that the RCT did not include high contamination (Grade III), but Radoicic2014 only involved open fractures with Grade I and II contamination, so was not included.
RAJASEKARAN2009 ³³⁰	No comparison between immediate (early) versus staged (delayed) closure
RAO2010 ³³¹	No comparison between immediate (early) versus staged (delayed) closure
RINKER2005 ³⁴³	Population does not match the protocol (includes patients who do not have open fractures)
RINKER2008 ³⁴²	Inadequate adjustment for confounders
ROMMENS1986 ³⁴⁸	No comparison between immediate (early) versus staged (delayed) closure
RUSSELL1990 ³⁵⁵	No comparison between immediate (early) versus staged (delayed) closure
SHEPHERD1998 ³⁶⁶	Inadequate adjustment for confounders
STALEKAR2003 ³⁷⁸	Inadequate adjustment for confounders
STANNARD2010 ³⁷⁹	Not a systematic review
STEIERT2009 ³⁸¹	Inadequate adjustment for confounders
SWANSON1991 ³⁸⁶	Inadequate adjustment for confounders
THO1994 ³⁹¹	No outcomes for comparison between immediate (early) versus staged (delayed) closure
TORCHIA1996 ³⁹⁵	Inadequate adjustment for confounders
TOWNLEY2010 ³⁹⁷	No comparison between immediate (early) versus staged (delayed) closure
WEBB2007 ⁴¹⁵	MVA variables unclear
WIDENFALK1979 ⁴¹⁸	No comparison between immediate (early) versus staged (delayed) closure
WOOD2012 ⁴²²	SR does not match our protocol.
YAREMCHUK1987	No comparison between immediate (early) versus staged (delayed) closure
YOKOYAMA1995 ⁴²⁸	Inadequate controlling for confounders
YOKOYAMA2006A ⁴²⁹	Inadequate controlling for confounders
YUSOF2013 ⁴³²	No comparison between immediate (early) versus staged (delayed) closure
ZATTI2000 ⁴³⁴	No comparison between immediate (early) versus staged (delayed) closure
ZIRAN2004 ⁴³⁸	No comparison between immediate (early) versus staged (delayed) closure

Reference	Reason for exclusion
	closure

J.1.8 Cover

Table 8: Studies excluded from the clinical review

Reference	Reason for exclusion
ALLISON2011 ¹¹	No comparison between immediate (early) versus staged (delayed) closure
BALDWIN2009 ²⁶	No comparison between immediate (early) versus staged (delayed) closure
BARTLETT1997 ³⁴	No comparison between immediate (early) versus staged (delayed) closure
BERRY2004 ⁴¹	No comparison between immediate (early) versus staged (delayed) closure
BHANDARI2001 ⁴³	No comparison between immediate (early) versus staged (delayed) closure
BLICK1989 ⁴⁸	No comparison between immediate (early) versus staged (delayed) closure
BREUGEM2006 ⁵³	SR does not meet our inclusion criteria
BUCKLEY1996 ⁵⁵	Inadequate adjustment for confounders
BURGESS1987 ⁵⁷	No comparison between immediate (early) versus staged (delayed) closure
BYRD1981 ⁵⁸	No comparison between immediate (early) versus staged (delayed) closure
BYRD1985 ⁵⁹	Inadequate adjustment for confounders
CAUDLE1987	Inadequate adjustment for confounders
CIERNY1983 ⁶⁷	Inadequate adjustment for confounders
COX1970 ⁷⁴	No comparison between immediate (early) versus staged (delayed) closure
CULLEN1996 ⁷⁸	No comparison between immediate (early) versus staged (delayed) closure
DAVISSEARS2012 ⁸³	No comparison between immediate (early) versus staged (delayed) closure
DELONG1999 ⁸⁹	Inadequate adjustment for confounders
DONG2011 ⁹³	No comparison between immediate (early) versus staged (delayed) closure
EDWARDS1983 ¹⁰¹	Not a systematic review
EDWARDS1988 ¹⁰²	No comparison between immediate (early) versus staged (delayed) closure
ERDMANN1997 ¹⁰⁸	No comparison between immediate (early) versus staged (delayed) closure
FERRERA1999 ¹¹³	No comparison between immediate (early) versus staged (delayed) closure
FISCHER1991 ¹¹⁴	No comparison between immediate (early) versus staged (delayed) closure
FRANCEL1992 ¹²¹	Inadequate adjustment for confounders
GLASS2009A ¹²⁴	No comparison between immediate (early) versus staged (delayed) closure

Reference	Reason for exclusion
GODINA1986 ¹²⁶	Inadequate adjustment for confounders
GOPAL2000 ¹²⁹	Inadequate adjustment for confounders
GOUGOULIAS2009A ¹³⁰	No comparison between immediate (early) versus staged (delayed) closure
GREENBAUM2001 ¹³³	No comparison between immediate (early) versus staged (delayed) closure
GRIMARD1996 ¹³⁵	Inadequate adjustment for confounders
GUSTILO1976 ¹⁴¹	No comparison between immediate (early) versus staged (delayed) closure
HAASBEEK1995 ¹⁴³	No comparison between immediate (early) versus staged (delayed) closure
HAMMER1992 ¹⁴⁵	No comparison between immediate (early) versus staged (delayed) closure
HARLEY2002 ¹⁴⁷	Inadequate adjustment for confounders
HARRIS2006 ¹⁴⁹	No comparison between immediate (early) versus staged (delayed) closure
HARVEY2002 ¹⁵⁰	No comparison between immediate (early) versus staged (delayed) closure
HARWOOD2006 ¹⁵¹	No comparison between immediate (early) versus staged (delayed) closure
HAS1995 ¹⁵²	No comparison between immediate (early) versus staged (delayed) closure
HEE2001 ¹⁵⁵	No comparison between immediate (early) versus staged (delayed) closure
HEIER2003 ¹⁵⁶	No comparison between immediate (early) versus staged (delayed) closure
HELLAND1996 ¹⁵⁸	No comparison between immediate (early) versus staged (delayed) closure
HENLEY1998 ¹⁶⁰	No comparison between immediate (early) versus staged (delayed) closure
HERNIGOU2013 ¹⁶¹	No comparison between immediate (early) versus staged (delayed) closure
HOFFMANN2013 ¹⁶⁷	No comparison between immediate (early) versus staged (delayed) closure
HONG1998 ¹⁷¹	No comparison between immediate (early) versus staged (delayed) closure
HOU2011 ¹⁷⁶	Inadequate adjustment for confounders
HULL2008 ¹⁸¹	Not a systematic review
HULSKER2011 ¹⁸³	Incorrect study design included
HUTCHINSON2012 ¹⁸⁴	No comparison between immediate (early) versus staged (delayed) closure
HUTSON2010 ¹⁸⁵	No comparison between immediate (early) versus staged (delayed) closure
JONES2003 ¹⁹⁵	No comparison between immediate (early) versus staged (delayed) closure
JOSHI2006 ¹⁹⁶	No comparison between immediate (early) versus staged (delayed) closure
KAI1998 ¹⁹⁸	No comparison between immediate (early) versus staged (delayed) closure

Reference	Reason for exclusion
	closure
KAKAR2007 ²⁰⁰	No comparison between immediate (early) versus staged (delayed) closure
KAMATH2012 ²⁰²	Inadequate adjustment for confounders
KEELING2008 ²⁰⁹	No comparison between immediate (early) versus staged (delayed) closure
KESEMENLI2004 ²¹²	No comparison between immediate (early) versus staged (delayed) closure
KIM2012 ²¹⁶	Outcomes of the protocol not reported
KINZEL2006 ²¹⁸	Does not meet our protocol
KREDER1995 ²²²	Outcomes of the protocol not reported
KULSHRESTHA2008 ²²⁴	No comparison between immediate (early) versus staged (delayed) closure
LAUGHLIN1993 ²³⁰	No comparison between immediate (early) versus staged (delayed) closure
LENARZ2010 ²³²	No comparison between immediate (early) versus staged (delayed) closure
LEONG1988 ²³³	Inadequate adjustment for confounders
LERNER2006	No comparison between immediate (early) versus staged (delayed) closure
LOWENBERG1996 ²³⁹	No comparison between immediate (early) versus staged (delayed) closure
MACK2013 ²⁴³	No comparison between immediate (early) versus staged (delayed) closure
MIN2011 ²⁶⁹	Inadequate adjustment for confounders
MODA1994 ²⁷²	No comparison between immediate (early) versus staged (delayed) closure
NAIQUE2006 ²⁸²	Inadequate adjustment for confounders
PAPAKOSTIDIS2011 ³⁰⁷	No comparison between immediate (early) versus staged (delayed) closure
PARK2007 ³⁰⁸	No comparison between immediate (early) versus staged (delayed) closure
PARRETT2006 ³⁰⁹	No comparison between immediate (early) versus staged (delayed) closure
POLLAK2000 ³¹⁹	Intervention does not meet the protocol
RAJASEKARAN2009 ³³⁰	No comparison between immediate (early) versus staged (delayed) closure
RAO2010 ³³¹	No comparison between immediate (early) versus staged (delayed) closure
RINKER2005 ³⁴³	Population does not match the protocol (includes patients who do not have open fractures)
RINKER2008 ³⁴²	Inadequate adjustment for confounders
ROMMENS1986 ³⁴⁸	No comparison between immediate (early) versus staged (delayed) closure
RUSSELL1990 ³⁵⁵	No comparison between immediate (early) versus staged (delayed) closure
SHEPHERD1998 ³⁶⁶	Inadequate adjustment for confounders
STANNARD2010 ³⁷⁹	Not a systematic review

Reference	Reason for exclusion
STEIERT2009 ³⁸¹	Inadequate adjustment for confounders
SWANSON1991 ³⁸⁶	Inadequate adjustment for confounders
THO1994 ³⁹¹	No outcomes for comparison between immediate (early) versus staged (delayed) closure
TORCHIA1996 ³⁹⁵	Inadequate adjustment for confounders
TOWNLEY2010 ³⁹⁷	No comparison between immediate (early) versus staged (delayed) closure
WEBB2007 ⁴¹⁵	MVA variables unclear
WIDENFALK1979 ⁴¹⁸	No comparison between immediate (early) versus staged (delayed) closure
WOOD2012 ⁴²²	SR does not match our protocol.
YAREMCHUK1987	No comparison between immediate (early) versus staged (delayed) closure
YOKOYAMA1995 ⁴²⁸	Inadequate controlling for confounders
YOKOYAMA2006A ⁴²⁹	Inadequate controlling for confounders
YUSOF2013 ⁴³²	No comparison between immediate (early) versus staged (delayed) closure
ZATTI2000 ⁴³⁴	No comparison between immediate (early) versus staged (delayed) closure
ZIRAN2004 ⁴³⁸	No comparison between immediate (early) versus staged (delayed) closure
JENKINSON2014 ¹⁹³	No timing data
EGOL2005 ¹⁰³	Non comparative study
SCHEMITSCH2012 ³⁶³	No timing data
LIU2012 ²³⁷	No immediate cover group

J.1.9 Definitive dressings after debridement

Table 9: Studies excluded from the clinical review

Study	Reason for exclusion
Back 2013 ²³	Systematic review is not relevant to review question or unclear PICO
Blum 2012 ⁴⁹	Study design not relevant to review. Cohort study
Calhoun 1993 ⁶⁰	Not review population. Chronic osteomyelitis
Contractor 2008 ⁷¹	Systematic review is not relevant to review question or unclear PICO
Halvorson 2011 ¹⁴⁴	Incorrect study design. Non-comparative study
Kazakos 2009 ²⁰⁶	Incorrect interventions. Plasma rich platelet gel
Keating 1996 ²⁰⁷	Incorrect intervention: bead group not given concomitant IV antibiotics
Keen 2012 ²¹⁰	Incorrect study design
Liu 2012 ²³⁷	Study design not relevant to review. Cohort study
Moehring 2000 ²⁷⁶	Incorrect interventions: antibiotic beads not given alongside IV antibiotics
Moues 2004 ²⁸¹	Not guideline condition
Ogbemudia 2010 ²⁹⁷	Not review population
Rinker 2008 ³⁴²	Incorrect study design
Runkel 2011 ³⁵⁴	Systematic review is not relevant to review question or unclear PICO

Study	Reason for exclusion
Stannard 2010 ³⁷⁹	Study design not relevant to review. Review
Tang 2010 ³⁸⁹	Incorrect study design. Case series
Wright 2007 ⁴²³	Incorrect study design
Yuenyongviwat 2011 ⁴³⁰	Incorrect interventions. Only standard dressings

J.2 Pelvic fractures

J.2.1 Transfer to MTC

Table 10: Studies excluded from the clinical review

Study	Reason for exclusion
Bouzat 2013 ⁵¹	Article in French
Demetriades 2005 ⁹⁰	Does not compare intervention of interest directly with each other

J.2.2 Decision for pelvic binders

Table 11: Studies excluded from the clinical review

Reference	Reason for exclusion
Baumann 2011 ³⁵	Evaluated eFAST - not a risk tool and eFAST not used pre-hospital
Reynolds 2014A ³³⁸	Abstract only

J.2.3 Timing of log roll

Table 12: Studies excluded from the clinical review

Reference	Reason for exclusion
Block2001	Review

J.2.4 Pelvic imaging

Table 13: Studies excluded from the clinical review

Reference	Reason for exclusion
Dormagen 2010 ⁹⁴	Inappropriate study design: Diagnosis of arterial injury
Duane 2008 ⁹⁷	Inappropriate study design: Initial CT as reference standard
Falchi 2004 ¹¹¹	Systematic review no meta-analysis: Used to source references only
Guillamondegui 2003 ¹³⁸	Inappropriate study design: Initial CT as reference standard
Harley 1982 ¹⁴⁸	Inappropriate study design: Initial CT+X-ray as reference standard
Henes 2012 ¹⁵⁹	Inappropriate study population: Low or moderate energy pelvic fractures in elderly population
Holmes 2012 ¹⁶⁹	Inappropriate study design: Initial CT as reference standard
Kirby 2010 ²¹⁹	Inappropriate study design: MRI as reference standard
Obaid 2006 ²⁹⁵	Inappropriate study design: Initial CT as reference standard
O'Shea 2006 ²⁹²	Inappropriate study design: Post-operative imaging
O'Toole 2001 ²⁹⁴	Inappropriate study design: Inter-rater reliability of imaging strategies
Magid 1986 ²⁴⁷	Inappropriate study design: Initial CT as reference standard
Nuchtern 2015 ²⁸⁹	Inappropriate study population: Low or moderate energy pelvic fractures

Reference	Reason for exclusion
	in elderly population
Paydar 2013 ³¹²	Case series: No comparative or diagnostic accuracy data
Potter 1994 ³²³	Inappropriate study design: Surgical findings as reference standard, but evidence of missed fractures with gold standard
Resnik 1992 ³³⁶	Inappropriate study design: Initial CT as reference standard
Robertson 1995 ³⁴⁵	Inappropriate study design: Initial CT as reference standard
Their 2005 ³⁹⁰	Inappropriate study design: Initial CT as reference standard
Vo2004 ⁴¹¹	Inappropriate study design: Initial CT as reference standard
Yugueros1995 ⁴³¹	Inappropriate study design: No relevant index test

J.2.5 Pelvic cystourethrogram

Table 14: Studies excluded from the clinical review

Reference	Reason for exclusion
Carroll 1983 ⁶³	Incorrect population: all patients had bladder rupture
Deck 2000 ⁸⁶	CT scanner utilised not MDCT
Deck 2001 ⁸⁷	CT scanner utilised not MDCT
Haas 1999 ¹⁴²	CT scanner utilised not MDCT
Horstman 1991 ¹⁷⁴	CT scanner utilised not MDCT
Kailidou 2005 ¹⁹⁹	No separate data for bladder injury
Kane 1989 ²⁰³	CT scanner utilised not MDCT
Luckhoff 2011 ²⁴⁰	Incorrect diagnostic test: urethral injury
Marks 2012 ²⁵²	Case report
Mokoena 1995 ²⁷⁷	Prognostic factor study
Morey 2001 ²⁷⁸	Not a diagnostic accuracy or effectiveness study
Morgan 2000 ²⁷⁹	Prognostic factor study
Pao 2000 ³⁰⁵	CT scanner utilised not MDCT
Peng 1999 ³¹³	CT scanner utilised not MDCT
Quagliano 2006 ³²⁸	One CT scanner utilised not MDCT
Rehm 1991 ³³⁵	No accuracy data presented
Spencer Netto 2008 ³⁷⁴	Incorrect population: all patients had bladder/urethral injuries
Stengel 2012 ³⁸²	No separate data for bladder injury
Udekwu 1996 ⁴⁰²	CT scanner utilised not MDCT
Ziran 2005 ⁴³⁹	Not a diagnostic accuracy or effectiveness study

J.2.6 Pelvic haemorrhage control

Table 15: Studies excluded from the clinical review

Study	Reason for exclusion
Abrassart 2013 ³	Groups in the study not adjusted for confounders
Akbar 2012 ⁶	Internal fixation is not used to treat pelvic haemorrhage
Anandakumar 2013 ¹²	Compared angiogram (with some that had EA) against no angiogram.
Baylis 2004 ³⁶	No intervention of interest

Study	Reason for exclusion
Beard 1988 ³⁷	No comparison of intervention in the study
Biffi 2001 ⁴⁵	study does not report outcomes separately for the interventions
Burgess 1990 ⁵⁶	No comparison of interventions in the study
Clamp 2011 ⁶⁸	Review
Cook 2002 ⁷²	No direct comparison of interventions in the study
Croce 2007 ⁷⁵	POD vs. Ex fixation. POD is not an invasive technique
Cullinane 2011 ⁷⁹	Review
Davis 2008 ⁸⁴	Review
Ertel 2001 ¹⁰⁹	No direct comparison of interventions in the study
Evers 1989 ¹¹⁰	Groups not adjusted for confounders
Flint 1990 ¹¹⁵	No outcomes reported for interventions and groups in the study not adjusted for confounders
Goins 1992 ¹²⁷	No comparison of interventions in the study
Grubor 2011 ¹³⁷	Groups were not adjusted for confounders and data was not reported for groups separately
Hu 2013 ¹⁷⁹	Review
Keel 2005 ²⁰⁸	review
Lai 2008 ²²⁶	case-series with no relevant data
Langford 2013 ²²⁸	Review
Lustenberger 2011 ²⁴¹	looks at intervention pelvic c-clamp followed by pelvic packing
Mauffrey 2014 ²⁵⁵	Review
Mlyncek 2005 ²⁷¹	Review
O'flanagan 1992 ²⁹¹	Internal fixation is a technique to stabilise fracture, not control pelvic haemorrhage
Osborn 2009 ³⁰²	half of the angiogram group did not undergo embolisation
Pizanis 2013 ³¹⁶	c-clamp compared against non-invasive techniques in study
Plaisier 2000 ³¹⁷	No comparison of interventions in the study
Ricci 2014 ³³⁹	Review
Richardson 1982 ³⁴¹	Case-series study
Ruchholtz 2004 ³⁵³	Groups in the study not adjusted for confounders
Sadri 2005 ³⁵⁶	Groups not adjusted for confounders
Sriussadaporn 2002 ³⁷⁶	Groups in the study not adjusted for confounders
Uchida 2011 ⁴⁰¹	Groups in the study not adjusted for confounders
Van veen 1995 ⁴⁰⁵	No direct comparisons between interventions in the study
Verbeek 2008 ⁴⁰⁸	Review
Vigdorchik 2012 ⁴¹⁰	Internal fixation device only
Waikukul 1999 ⁴¹²	external fixation compared with a non-invasive conventional method of treatment
Yang 2008 ⁴²⁶	No direct comparison of interventions in the study
Zhao 2011 ⁴³⁷	Review

J.3 Pilon fractures

J.3.1 Pilon early fixation

Table 16: Studies excluded from the clinical review

Study	Reason for exclusion
Anglen 1999 ¹³	Incorrect interventions
Bacon 2008 ²⁴	Inappropriate comparison
Bacon 2008 ²⁴	Both groups with same timing
Binda 2011 ⁴⁶	Abstract
Blauth 2001 ⁴⁷	No adjustment for open or closed
Calori 2010 ⁶¹	Incorrect interventions
Court-brown 1999 ⁷³	Incorrect interventions
Cronier 2012 ⁷⁶	Non-systematic review
Crutchfield 1995 ⁷⁷	Incorrect interventions
Gulabi 2012 ¹³⁹	Incorrect interventions
Horn 2011 ¹⁷³	Incorrect interventions
Kapukaya 2005 ²⁰⁴	Case series
Katsenis 2009 ²⁰⁵	Incorrect interventions
Ketz 2012 ²¹⁴	Incorrect interventions
Ketz 2012 ²¹³	Case series
Korkmaz 2013 ²²⁰	Incorrect interventions
Li 2012 ²³⁶	Incorrect interventions
Mandracchia 1999 ²⁵⁰	Non-systematic review
Marsh 1995 ²⁵⁴	Incorrect interventions
Mauffrey 2011 ²⁵⁶	Non-systematic review
Mcferran 1992 ²⁶⁰	Incorrect interventions
Okcu 2004 ²⁹⁹	Incorrect interventions
Papadokostakis 2008 ³⁰⁶	Systematic review is not relevant to review question or unclear PICO
Pollak 2003 ³²¹	The staging/timing categories in protocol were not evaluated in this cohort study
Pugh 1999 ³²⁵	Incorrect interventions
Purghel 2012 ³²⁷	Non-systematic review
Richard 2012 ³⁴⁰	Both groups had same category of delay to definitive treatment
Salton 2007 ³⁶⁰	No protocol outcomes
Sirkin 1999 ³⁶⁹	Incorrect interventions
Trumble 1993 ⁴⁰⁰	Incorrect interventions
Vasiliadis 2009 ⁴⁰⁷	Not relevant to protocol
Wang 2010 ⁴¹³	Both groups had same delay to definitive treatment
Watson 2000 ⁴¹⁴	Incorrect interventions
Wyrsh 1996 ⁴²⁵	No analysis for staging or timing
Zeng 2011 ⁴³⁵	No outcomes reported for group comparison

J.3.2 Pilon fixation

Table 17: Studies excluded from the clinical review

Reference	Reason for exclusion
Anglen1999 ¹³	Mixed groups (temporary external fixation)
Babis1997 ²²	Inadequate reporting of confounders.
Bacon 2008 ²⁴	Non-randomised study. Does not report health-related quality of life.
Baloch2009 ²⁹	Cancelled order- descriptive study and was unable to be found
Blauth2001 ⁴⁷	Inadequate reporting of confounders
Binda2011 ⁴⁶	Abstract
Calori2010 ⁶¹	Includes case series.
Crutchfield1995 ⁷⁷	Inadequate reporting of confounders
Davidovitch 2011 ⁸²	Non-randomised study. Does not report health-related quality of life.
Elkhechen2012 ¹⁰⁴	Duplication - not ordered
Endres2004 ¹⁰⁶	Not in English
Gulabi2012 ¹³⁹	Inadequate reporting of confounders
Harris2006 ¹⁴⁹	Inadequate adjustment of confounders. Unbalanced for age, and grade at baseline.
Helfet1994 ¹⁵⁷	Unclear who had initial ext. fixation. Inadequate reporting of confounders.
Joveniaux2010 ¹⁹⁷	Not pilon fracture specific
Kendig1997 ²¹¹	Not a systematic review
Korkmaz2013 ²²⁰	Inadequate reporting of confounders
Koulouvaris2007 ²²¹	Interventions were not those specified in protocol
Marsh1999 ²⁵³	Not a systematic review
Ovadia 1986 ³⁰³	Comparator was a mixed treatment. No adjustment for key confounders
Pierce1979 ³¹⁴	All internal fixation
Pollack 2003 ³²¹	Non-randomised study. Does not report health-related quality of life.
Pugh1999 ³²⁵	Inadequate reporting of confounders
Puha2014 ³²⁶	Doesn't meet protocol comparisons
Ristiniemi2011 ³⁴⁴	Not pilon fracture specific
Salmenkivi1999 ³⁵⁹	Not in English
Watson2000 ⁴¹⁴	Inadequate reporting of confounders
Willet2008 ⁴¹⁹	Cochrane protocol.
Williams1998 ⁴²⁰	Does not meet protocol comparisons
Williams2004 ⁴²¹	All patients had external fixation with limited internal fixation.
Wyrsh1996 ⁴²⁵	Not a proper RCT. Inadequate adjustment of confounders at baseline.
Zeng2011 ⁴³⁵	No baseline characteristics (age included)

J.4 Other

J.4.1 Identifying vascular compromise

Table 18: Studies excluded from the clinical review

Reference	Reason for exclusion
-----------	----------------------

Reference	Reason for exclusion
Eastman2006 ¹⁰⁰	Not extremity trauma.
Fry1993 ¹²²	No baseline characteristics.
Levy2005 ²³⁵	Not a systematic review
Lord1974 ²³⁸	Not a systematic review
Maclean1964 ²⁴⁵	Only includes case reports
Pieroni2009 ³¹⁵	Not a systematic review
Redmond2008 ³³⁴	Not a systematic review
Romano2012 ³⁴⁷	Not a systematic review
Rose1987 ³⁴⁹	27% had blunt injury – rest were penetrating. No mention of fractures.
Rose1988 ³⁵⁰	Not a diagnostic paper.

J.4.2 Detecting compartment syndrome

Table 19: Studies excluded from the clinical review

Reference	Reason for exclusion
BROOKER1979 ⁵⁴	Mixed population (extremity wounds)
HANSEN2013 ¹⁴⁶	Not RCT/diagnostic accuracy study
KALYANI2011 ²⁰¹	Systematic review (irrelevant inclusion criteria)
MCQUEEN1996B ²⁶³	Not RCT/No sensitivity or specificity data
MITTLMEIER1991 ²⁷⁰	Not RCT/No sensitivity or specificity data
OGUNLUSI2005A ²⁹⁸	Not RCT/No sensitivity or specificity data
OVRE1998 ³⁰⁴	Not RCT/No sensitivity or specificity data
ROYLE1992 ³⁵²	Not RCT/diagnostic accuracy study
SAKIA2008 ³⁵⁸	Not RCT/No sensitivity or specificity data
TRIFFITT1992 ³⁹⁸	Not RCT/No sensitivity or specificity data
UPPAL1992 ⁴⁰⁴	Not RCT/No sensitivity or specificity data
WHITNEY2014 ⁴¹⁷	Not RCT/No sensitivity or specificity data

J.4.3 Splinting of lower limb long bone fractures

Table 20: Studies excluded from the clinical review

Reference	Reason for exclusion
CHU2003 ⁶⁶	Incorrect interventions: not box splint
IRAJPOUR2012 ¹⁸⁹	Incorrect interventions: not box splint
LEMBO1975 ²³¹	Study not in English
PODESZWA2004 ³¹⁸	Incorrect interventions: not box splint
SHORT1984 ³⁶⁷	Incorrect interventions: not box splint
THOMAS1981 ³⁹³	Incorrect interventions: not box splint

J.4.4 Hip reduction

Table 21: Studies excluded from the clinical review

Reference	Reason for exclusion
Ashley, 1972 ¹⁹	Review; references screened

Reference	Reason for exclusion
Barquet, 1982 ³¹	Inadequate adjustments for key confounders
Barquet, 1982 ³²	Only some with open reduction and analysis not sub-grouped for these.
Bergman, 1994 ⁴⁰	Unrelated to review question
Bhandari, 2006 ⁴⁴	Dislocation reduction was closed; only fracture repair was open
de Palma, 2014 ⁸⁵	No analysis of timing
Dwyer, 2006 ⁹⁹	Inadequate adjustments for key confounders
Epstein, 1974 ¹⁰⁷	Inadequate adjustments for key confounders
Fordyce, 1971 ¹¹⁸	Case report
Herwig-Kempers, 1993 ¹⁶²	Unrelated to review question
Hillyard, 2003 ¹⁶⁵	Unclear if reductions were open
Hougaard, 1986 ¹⁷⁷	Only some with open reduction and analysis not sub-grouped for these.
Jacob, 1987 ¹⁹¹	Only some with open reduction and analysis not sub-grouped for these.
Marchetti, 1996 ²⁵¹	Timing was <24 hours versus >24 hours
McKee, 1998 ²⁶¹	Inadequate adjustments for key confounders
Mehlman, 2000 ²⁶⁵	Only some with open reduction and analysis not sub-grouped for these.
Mehta, 2008 ²⁶⁶	Inadequate adjustments for key confounders
Moed, 2000 ²⁷⁴	Dislocation reduction was closed; only fracture repair was open
Moed, 2002 ²⁷⁵	Dislocation reduction was closed; only fracture repair was open
Morsy Drch, 2001 ²⁸⁰	Dislocation reduction was closed; only fracture repair was open
Rosenthal, 1979 ³⁵¹	Only some with open reduction and analysis not sub-grouped for these.
Sahin, 2003 ³⁵⁷	Only some with open reduction and analysis not sub-grouped for these.
Sanders, 2010 ³⁶²	Review; references screened
Sturrock, 1899 ³⁸³	Unrelated to review question; archaic
Toni, 1985 ³⁹⁴	Inadequate adjustments for key confounders
Upadhyay, 1981 ⁴⁰³	Timing not considered
Viale, 2005 ⁴⁰⁹	Only some with open reduction and analysis not sub-grouped for these.
Yang, 1991 ⁴²⁷	Only some with open reduction and analysis not sub-grouped for these.
Zha, 2013 ⁴³⁶	Dislocation reduction was closed; only fracture repair was open

J.4.5 Full-body CT

Table 22: Studies excluded from the clinical review

Reason for exclusion	Reason for exclusion
Beck 2012 ³⁸	Review article but no RCT's included
Caputo 2014 ⁶²	Review article but no RCT's included
Ptak 2001 ³²⁴	Retrospective study and does not include outcomes of interest
Saltzherr 2009 ³⁶¹	Correspondence article
Van vugt 2012 ⁴⁰⁶	All studies included in the review were not RCT's

J.4.6 Documentation of open fracture wound photographs

Table 23: Studies excluded from the clinical review

Reference	Reason for exclusion
Solan MC, Calder JDF, Gibbons CER, Ricketts DM. ³⁷³ Photographic wound documentation after open fracture. <i>Injury, Int J care Injured</i> 2001; 32: 33-35	Not relevant to review question

J.4.7 Documentation of neurovascular compromise

Table 24: Studies excluded from the clinical review

Study	Reason for exclusion
Johnston-walker 2011 ¹⁹⁴	Not primary research
Mayne 2013 ²⁵⁸	No clinical outcomes linked to completeness of neurovascular documentation
Wright 2007 ⁴²⁴	No clinical outcomes linked to completeness of documentation recording neurovascular compromise

J.4.8 Information and support

Table 25: Studies excluded from the clinical review

Reference	Reason for exclusion
Aravind 2010 ¹⁴	No information themes
Archibald 2003 ¹⁵	No information themes
Atchison 2005 ²⁰	Not about patients' thoughts and feelings about information desired. This paper concerned whether patients recalled being given specific information
Azam 2011 ²¹	Non qualitative
Bagely 2011 ²⁵	Non qualitative
Congdon 1994 ⁷⁰	Hip fracture
Elliot 2014 ¹⁰⁵	Hip fracture
Glenny 2013 ¹²⁵	Hip fracture
Hommel 2012 ¹⁷⁰	Hip fracture
Hossieny 2012 ¹⁷⁵	Non qualitative
Jariwala 2004 ¹⁹²	Did not reveal participants' feelings about information desired.
Lam 2011 ²²⁷	Non qualitative
Macleod 2005 ²⁴⁶	Hip fracture
Malin Malmgren 2014 ²⁴⁹	Hip fracture
Mayich 2013 ²⁵⁷	Non qualitative
Meadows 2005 ²⁶⁴	Non-trauma population – fragility fractures
Meredith 1993 ²⁶⁷	Non qualitative
Modin 2009 ²⁷³	No information themes
Nielsen 2013 ²⁸⁷	Non-trauma population – fragility fractures
Olson 1990 ³⁰¹	Spinal fractures

Reference	Reason for exclusion
O'Toole 2001 ²⁹⁴	Non qualitative
Shyu 2010 ³⁶⁸	Non qualitative
Toscan 2012 ³⁹⁶	Hip fracture

Appendix K: Excluded economic studies

K.1 Pelvic fractures

K.1.1 Pelvic imaging

Table 26: Excluded studies

Reference	Reason for exclusion
Feeney 2011 ¹¹²	<p>This study was assessed as partially applicable with very serious limitations.</p> <p>It is a retrospective cost comparison looking at the savings involved if patients who are haemodynamically unstable and already having a CT are withheld a pelvic X-ray. The study is from a US perspective which is not particularly applicable to the UK setting. It does not include any health effects or downstream costs/consequences and is thus of limited usefulness.</p>
Barleben 2011 ³⁰	<p>This study was assessed as partially applicable with very serious limitations.</p> <p>It is a prospective cost comparison looking at the savings involved if an algorithm is used which outlines that patients who are undergoing a CT should only receive a pelvic X-ray if they fulfil certain haemodynamic/physiologic criteria. The study is from a US perspective which is not particularly applicable to the UK setting. It does not include quality of life and is thus of limited usefulness.</p>

Appendix L: Cost analysis for open fractures

L.1 Introduction

L.1.1 Background

A fracture that breaks through the skin is called an open fracture. There are different grades of open fracture defined by the Gustilo-Anderson fracture classification system, each of which depends on the level of tissue damage and whether there is a vascular injury that requires repair. Those with vascular compromise require emergency treatment to re-vascularise the limb and so are not relevant to the analysis for the timing of debridement. For those who do not have vascular compromise, the main concern is that the wound can become infected. If this infection is only superficial then it can be treated easily with a course of antibiotics. However, if this infection becomes deep then the treatment may require a series of additional procedures which can greatly increase costs. There is also a risk that the limb would require amputation, which would have a further cost and quality of life impact. In some cases deep infection could even result in death.

Treatment for open fractures has three main stages: debridement, fixation and soft tissue cover. Debridement, which is performed by an orthopaedic surgeon, involves cleaning the wound and removing any contaminated, unsalvageable or dead tissue. The timing of debridement is known to affect the risk of infection, and a clinical review (see chapter 6.7 of the complex fractures full guideline) was undertaken to identify to what extent early debridement can improve outcomes. In some hospitals across the UK, debridement is performed with the support of a plastic surgeon. The presence of a plastic surgeon in theatre allows the orthopaedic surgeon to utilise the expert knowledge of the plastic surgeon regarding the quantity of soft tissue that can be removed while still allowing for a successful cover procedure to be performed once fixation is complete. The rationale for this is that without this expert input, the orthopaedic surgeon may be too cautious and try to preserve as much tissue as possible in order to aid the later cover procedure. However, this can lead to a higher risk of infection as some contaminated tissue may still remain. The presence of a plastic surgeon can therefore help to reduce this risk but comes with the additional staffing cost for the duration of the procedure.

Fixation can involve one definitive procedure or it can be staged with an initial temporary fixator followed by later definitive fixation. These fixation procedures are performed by an orthopaedic surgeon and some form of fixation will occur immediately following debridement, whether it is definitive or temporary. Definitive fixation will only be delayed if a temporary fixator is applied immediately after debridement.

After debridement and definitive fixation have been completed, the open wound needs to be closed and covered by the surrounding soft tissue. This cover procedure is performed by a plastic surgeon and can be done immediately after definitive fixation or it can be delayed. Depending on the extent of the tissue damage, either a local flap procedure or a longer free flap procedure maybe required; the need for which can only be determined following the initial debridement. Until soft tissue cover has been successfully achieved, there is a risk of acquiring infection. Therefore the timing of the intervention is important in order to reduce this risk and the risk of other adverse events that require costly treatment and can have long term quality of life implications. The optimal timing of soft tissue cover may require a service delivery change by increasing the number of surgery lists, that is, the number of theatre days dedicated to orthoplastic procedures each week.

L.1.1.1 Exploration of TARN

Initially, a discrete event simulation model was planned, using the TARN database as a source of data to estimate the treatment effect of the interventions outlined above (see more detail on this in Appendix O:). The TARN database was explored primarily to find the effect of the timing of debridement, with and without the presence of a plastic surgeon, on the risk of deep infection and subsequent amputation as key outcomes. The analysis would also incorporate the number of theatre sessions the operations may take place in. In other words, this was attempting to inform the cost effectiveness of questions 1, 2 and 4 above. A brief overview of this model is provided below.

The population will include adults and children with open fractures. The strategies included in the model can be seen in Table 27.

Table 27: Proposed model strategies

Procedure combination	Theatre session 1	Theatre session 2	Theatre session 3
Option 1	A. Debridement B. Definitive fixation C. Definitive soft tissue cover	NA	NA
Option 2	A. Debridement B. Definitive fixation	C. Definitive cover	NA
Option 3	A. Debridement B. Temporary fixation	C. Definitive fixation D. Definitive soft tissue cover	NA
Option 4	A. Debridement B. Temporary fixation	C. Definitive fixation	D. Definitive soft tissue cover

Other components of the strategies include:

- The time of the initial debridement (<6 hours, 6-12 hours, 12-24 hours, >24 hours.)
- The presence of a plastic surgeon at the initial debridement

Outcomes included are:

- Time to death
- Time to deep infection
- Flap failure
- Amputation
- Length of hospital stay
- Number of unplanned operations (between debridement and cover)
- Time to soft tissue cover

Confounders that it felt should be adjusted for include:

- Age
- Grade of fracture (Gustilo Anderson)
- Upper/lower limb
- ISS
- The type and timing of prophylactic antibiotics that are given.
- Type of dressings used pre-debridement and post debridement.
- Type of definitive fixation (Temporary fixation should always be external).
- Method of soft tissue cover used (local flap or free flap).
- Polytrauma

- Major Trauma Centre or Trauma Unit

This model approach however was not developed further, after initial exploration of TARN for other guidelines on the trauma suite revealed that TARN data had limitations that deemed it not appropriate to use for our research purposes. For further detail on these limitations please see Major trauma economic model in appendix M of the major trauma guideline.

In addition to these limitations, for the complex fracture guideline, TARN was not felt to be appropriate because of difficulties in analysing certain codes in the database. A key limitation of the TARN database is that there is no direct link between each specific injury code and the related procedure code. This makes it difficult to identify the time of debridement for a specific open fracture in a patient with polytrauma. Although a clinician may be able to identify the sequence of procedures when looking at the data for each individual, it is not feasible to do this for 25,000 records and the computational coding cannot be adjusted to identify this accurately.

Another key limitation is that some of the procedure codes are not specific enough. This makes it difficult to identify the indication for an amputation for instance. We were interested in amputations resulting from deep infection as an outcome of debridement, however, patients who had an amputation due to an unsalvageable limb, compartment syndrome or a vascular injury would also have the same amputation code. This means that the analysis cannot accurately assess the effect on the risk of amputation that the timing of debridement has. Furthermore, the severity of the infection is not clearly defined, as it is not specified whether the infection is deep or superficial. This also makes it difficult to accurately assess our important outcomes.

The GDG thought a costing analysis would be helpful and alongside the limited clinical evidence identified would help them make a recommendation. The model was therefore downgraded to a costing analysis but was extended to look at the cost implications of the timing of definitive soft tissue cover also. The remainder of this appendix discusses the costing analyses that were undertaken.

L.1.2 Overview of analyses

The aim of this analysis is to inform the GDG of the cost implications for the open fracture questions that relate to the timing of the initial debridement, the provision of plastic surgery services for the initial debridement and the increase in the number of surgery lists made available for definitive soft tissue cover. Three separate analyses are presented in section L.2 to L.4 to help answer the questions outlined in section L.1.3 below.

This analysis was intended to focus on issues of additional plastic surgery services and therefore will not specifically look at the costs of fixation. However, a costing comparing different numbers of theatre sessions is included in section L.4 to demonstrate the cost of performing procedures in either one or more stages. This captures the staff cost implications when fixation and/or soft tissue cover is staged, but the cost of metal implants and fixation devices is not included in this analysis.

L.1.3 Questions and comparators relating to each analysis

The first analysis in section L.2 will address the two questions below:

1. What is the optimal timing of the initial debridement of open fractures?
 1. <6 hours
 2. 6 – 12 hours
 3. 12 – 24 hours

4. >24 hours
2. Is the presence of an orthopaedic surgeon and plastic surgeon at the initial surgical excision and stabilisation of an open fracture clinically and cost effective?
 1. Orthopaedic and plastic surgeon present in theatre
 2. Only orthopaedic surgeon present in theatre

The second analysis in section L.3 will address the question below:

3. What is the most clinically and cost effective time to achieve definitive soft tissue cover in open fractures?
 1. Immediate
 2. 1 day
 3. 3 days
 4. 7 days
 5. >7 days

The third analysis in section L.4 will address the trade-offs in staff time when fixation and/or cover is staged. This will help to inform the impact on cost for the question below and how that is affected by the presence of a plastic surgeon at debridement.

4. Is the use of initial definitive fixation and cover more clinically and cost effective in the management of open fractures compared to with staged fixation and cover?

L.1.4 Population

The population assessed are patients presenting with an open fracture that requires plastic surgery to cover the open wound after initial procedures have been performed.

L.2 Debridement cost analysis

L.2.1 Methods

The key cost impact for earlier debridement is that there will be an increased need for surgery during premium time, when theatre staff receive a higher rate of pay. For consultants, this is defined as 7pm until 7am on weekdays and all day on weekends and public holidays. Nurses and radiographers have different hours for premium time which are defined as 8pm until 6am on weekdays and all day Saturday. On Sundays and public holidays, a higher premium rate is paid for these staff. Registrars have a different arrangement as well and their premium time hours are from 7pm to 8am on weekdays and all day on weekends and public holidays. The salary enhancement for work performed in premium time will increase the cost of treatment for a proportion of patients when the time to debridement falls within this period. According to the clinical review, the outcomes of deep infection and amputation are reduced when debridement is performed earlier and so costs saved here could outweigh the cost of the increased salary for premium time work. Further detail on the times of premium time and the enhancement rates can be seen in Table 28 below. As the premium time bounds are slightly different for different staff, it was assumed for simplicity that no procedures would be performed during the hours where there is discrepancy between the premium and non-premium times.

Having a plastic surgeon at debridement adds another consultant and registrar salary to the theatre staffing with the out of hours enhancements as discussed above. The evidence suggests that there is

a reduction in the number of people acquiring a deep infection and subsequent amputation, so this costing will assess the net cost or cost saving when having a plastic surgeon present for debridement at each particular delay to debridement as outlined above.

This costing will include the costs of the core theatre staff with the addition of the relevant surgeons where appropriate for the intervention. Enhancements to salaries will be added for a proportion of patients who would be expected to have debridement out of hours. For the analyses including the presence of a plastic surgeon, additional time will be added for the plastic surgeon to travel in for a call out when procedures are performed in premium time. Only the salary of the plastic surgeon is included in this extra hour for travel. No travel expenses have been calculated.

The costs of the adverse events, based on the risks identified from the clinical review, will also be calculated and combined with the staffing costs to give an overall cost for each strategy.

L.2.2 Inputs

L.2.2.1 Resource use and unit costs of interventions

The costs incurred by the core non-surgical staff required to be available in theatre are presented in Table 29 below. These have been calculated from data published in PSSRU 2014.⁸⁰ The costs below include oncosts; qualifications; staff and non-staff overheads; and capital overheads. Oncosts were calculated using the HMRC national insurance rates for 2014-2015¹⁶⁶ and a superannuation rate provided by PSSRU⁸⁰. The total hourly cost of staff presented in PSSRU 2014⁸⁰ did not match the sum of the individual components presented and so we have used our own calculations in this analysis. However, the values we calculated were very similar to those presented in the publication.

The third and fourth columns of Table 29 show the hourly costs during premium time. As there is a further enhanced rate is paid to nurses on Sundays and public holidays and so this is separated into another column.

Table 28: Enhancement multiplier for premium time pay

Staff role	Premium time excluding Sundays and public holidays	Sundays and public holidays	Source	Comments
Consultant	1.33	1.33	Consultant contract. ²⁸⁴	
Registrar	1.50	1.50	Banding of junior doctors. ²⁸³	Assumed to be band 1A to account for the additional cost of unsocial hours.
Nurse and allied professionals (Agenda for Change bands 4-9)	1.30	1.60	Agenda for Change service handbook. ²⁸⁵	Applies to all core theatre staff outlined in Table 29 below.

Table 29: Core theatre staff costs

Staff role	Cost per hour (normal hours)	Cost per hour (premium time excluding Sundays and public holidays)	Cost per hour (Sundays and public holidays)	Source
				Error! Reference source not found.

Staff role	Cost per hour (normal hours)	Cost per hour (premium time excluding Sundays and public holidays)	Cost per hour (Sundays and public holidays)	SourceError! Reference ource not found.
Consultant anaesthetist	£139	£159	£159	Consultant medical, calculated using PSSRU 2014 data
Operating department practitioner	£49	£57	£65	Senior staff nurse, calculated using PSSRU 2014 data
Scrub nurse	£49	£57	£65	Senior staff nurse, calculated using PSSRU 2014 data
Running nurse	£41	£47	£54	Staff nurse, calculated using PSSRU 2014 data
Radiographer	£38	£44	£50	Hospital radiograph, calculated using PSSRU 2014 data
Recovery nurse	£41	£47	£54	Staff nurse, calculated using PSSRU 2014 data
TOTAL	£358	£412	£446	

The hourly cost of consultant surgeons during different hours is shown in Table 30 below. This cost applies to both orthopaedic and plastic surgeons and includes oncosts, qualifications and overheads.

Table 30: Surgeon staff costs

Cost per hour	Normal hours	Premium time including Sundays and public holidays
Consultant orthopaedic or plastic surgeon	£140	£161
Orthopaedic and plastic registrars	£58	£72

Source: PSSRU 2014

Costs Include oncosts, qualifications and overheads

The hourly cost of theatre staffing is shown in Table 31 below. This table shows the total cost for the core staff alone as well as with orthopaedic surgeons and with both orthopaedic surgeons and plastic surgeons (it was GDG opinion that there would be a consultant and registrar of each specialty), based on the costs reported in the two tables above. These are also shown for premium rate times as well as non-premium rate times.

Table 31: Theatre costs per hour

Input	Normal hours	Premium time excluding Sundays and public holidays	Sundays and public holidays
Core theatre staff	£358	£412	£446
Core staff plus orthopaedics ^a	£556	£644	£678
Core staff plus orthopaedics and plastics ^(a)	£754	£877	£911

Costs Include oncosts, qualifications and overheads

(a) Orthopaedics and plastics includes one consultant and one registrar for each speciality.

The duration of debridement (including call out time) and the proportion of people who are expected to be debrided during premium time are shown in Table 32 below. These values were estimated by the GDG. A sensitivity analysis will assess how robust the overall costs are to changes in these values.

Table 32: Duration of debridement and proportion requiring out of hours

	Timing of debridement from injury			
	6 hours	12 hours	24 hours	>24 hours
Duration of debridement (hours) ^(a)	3			
Additional hours for call out	1			
Proportion of injuries debrided in premium time (exc. Sunday)	0.2	0.1	0	0
Proportion of injuries debrided on a Sunday	0.1	0.05	0	0

(a) Includes an hour to perform debridement and two hours of theatre preparation and cleaning time

L.2.2.2 Resource use and unit costs of complications

Unit costs for the treatment for deep infection and amputation are shown in Table 33 below.

Table 33: Complications treatment costs

Outcome	Value	Source
Treatment of deep infection	£20,000	GDG assumption
Amputation procedure	£8,589	NHS Reference Costs 2013-2014 (HRG code = YQ22B)

The ranges for the cost of two common types of prosthesis are shown in Table 34 below. To estimate the expected cost of prosthesis, the midpoints for each range were calculated and the midpoint of the two midpoints was used as the base case value. Only leg prostheses were used due to an expected higher demand for leg prostheses and the importance of providing prostheses for ambulatory support.

Table 34: Prosthesis costs

Outcome	Mean Value	Source
Transtibial prosthesis	£2,350	GDG member contact – based on the midpoint between the lower and upper range limits (£700 and £4,000)
Transfemoral prosthesis	£4,750	GDG member contact – based on the midpoint between the lower and upper range limits (£1,500 and £8,000)
Total average	£3,550	

The number of prosthetics required, the lifetime of each prosthetic limb and the expected life years remaining for the patient is shown in Table 35 below. These values were used to calculate the expected number of prosthetic limbs required over a lifetime and the overall lifetime cost of these. These values are also presented in Table 35.

Table 35: Prosthetics resource use and lifetime cost

Outcome	Value	Source
Number of different prosthetics required at any time	2	GDG member contact
Life of prosthetics (years)	3	GDG member contact
Mean age at injury	45	GDG assumption
Mean age of death	83	Office for National Statistics ²⁹⁶
Life years remaining for patient	38	Calculated from above
Expected number of prosthetics over a lifetime	25	Calculated from above
Lifetime prosthetics cost	£92,300	Calculated from above
Discounted lifetime prosthetics cost ^(a)	£53,479	Calculated from above

(a) Discounted at an annual rate of 3.5%

L.2.2.3 Clinical effectiveness data

Table 36: Baseline data and odds ratios for deep infection

Inputs from clinical review	Data	Source
Baseline risk of deep infection (< 6 hours)	5.56%	Noumi 2005 ²⁸⁸
Odds ratio for deep infection per hour of delay	1.033	Hull 2014 ¹⁸²

The baseline risk of deep infection from Noumi et al.²⁸⁸ was converted into a baseline odds value (see section L.2.3 for more detail on computations). The odds ratio from Hull et al.¹⁸² (cross ref to review) was then applied to calculate the odds of infection for each debridement time. These odds values were then converted back into risks for each time point. These are shown in Table 37 below.

Table 37: Risks for deep infection

Duration of delay to debridement	Risk of deep infection
6 hours	5.56%
12 hours	6.75%
24 hours	9.97%
48 hours	21.7%

Risks were converted from odds calculated using the odds ratio from Hull 2014 and the baseline risk from Noumi 2005 in the table above.

The risk of amputation following deep infection is shown in Table 38 below. This value is multiplied by the risk of deep infection at the relevant time point to calculate the risk of amputation for the open fracture population.

Table 38: Risks for amputation

Input	Risk of deep infection	Source
-------	------------------------	--------

Input	Risk of deep infection	Source
Risk of amputation given deep infection has occurred	10%	GDG assumption

With the presence of a plastic surgeon, the risk of deep infection at each time points is reduced by 62%. (see Table 39). There is no difference in the risk of amputation if a plastic surgeon is present because this is only dependent on the risk of deep infection.

Table 39: Relative risks with a plastic surgeon present

Outcome	Data	Source
Deep infection	0.38	Naique 2006 ²⁸²
Amputation	N/A	GDG assumption

L.2.3 Computations

1. The total costs for each debridement strategy were calculated as illustrated by the equation below:

$$T_{ij} = (CORE_N r_{i,N} + CORE_P r_{i,P} + CORE_S r_{i,S} + ORTH_N r_{i,N} + ORTH_P r_{i,P} + ORTH_S r_{i,S}) \cdot t_{deb} + j \cdot (PLAST_N r_{i,N} + PLAST_P r_{i,P} + PLAST_S r_{i,S}) \cdot (t_{deb} + t_{call})$$

Where i represents the time of debridement strategy and j represents the plastic surgeon strategy ($j \in \{0 = \text{not present}, 1 = \text{present}\}$)

CORE, **ORTH** and **PLAST** denote the cost per hour of the core staff, orthopaedics and plastics respectively. The subscripts **N**, **P** and **S** specify whether these costs are in normal hours, premium time (excluding Sundays and public holidays) or Sundays and public holidays respectively.

The factors $r_{i,N}$, $r_{i,P}$ and $r_{i,S}$ represent the probabilities that debridement is performed in normal hours, premium time (excluding Sundays and public holidays), and Sundays (and public holidays) respectively.

The factors t_{deb} and t_{call} represent the duration that the costs are applied to i.e. the duration of debridement and the duration of additional travel time given for a plastic surgeon call out respectively.

2. Odds were converted into risks using the equation in the example below:

$a = \text{number of patients with event}$

$b = \text{number of patients without event}$

$$ODDS = \frac{a}{b}, \quad RISK = \frac{a}{a+b}$$

$$RISK = \frac{ODDS}{1 + ODDS}$$

L.2.4 Sensitivity analyses

A number of sensitivity analyses were undertaken to test the robustness of the results.

SA1: Relative risk of deep infection for the presence of a plastic surgeon threshold analysis

This analysis is a threshold analysis, meaning a value is altered until a certain condition is met. In this case, the relative risk of deep infection with a plastic surgeon present is altered until the total costs of the less than 6 hour debridement strategy, with and without a plastic surgeon present, are equal.

This was only calculated for debridement at less than 6 hours. The threshold value will increase for the later strategies due to the increasing risk of deep infection and reduced out of hours costs i.e. a smaller proportion of the larger number of infections is required to be reduced to cover the additional staffing cost of the plastic surgeons.

SA2: Increasing the proportion of patients whose debridement is performed in premium time by 50%

This analysis multiplies the proportion of patients expected to be debrided out of hours by 1.5 and assesses the effect on the overall costs.

SA3: Baseline risk of deep infection threshold analysis

This analysis is similar to SA1 but the value that is varied to find the threshold is the risk of deep infection. This threshold has only been calculated for the point where debridement at less than 6 hours becomes equally costly whether plastics are present or not. This risk will change the risks for each time of debridement when the odds ratio is applied.

SA4: Reducing the odds ratio of deep infection per hour of delay to debridement to 1.01

This analysis reduces the odds ratio of delay to debridement to 1.01, from the base case analysis value of 1.033, and assesses the effect of the overall costs.

SA5: Increasing prosthetic cost to £6,000

This analysis increases the cost of a single prosthesis from £3,550 to £6,000 and assesses the effect on the overall costs.

SA6: Cost of deep infection threshold analysis

This analysis is similar to SA1 but the value that is varied to find the threshold is the cost of deep infection. This threshold has only been calculated for the point where debridement at less than 6 hours becomes equally costly whether plastics are present or not.

L.2.5 Results

L.2.5.1 Base case analysis

The results of the cost analysis are reported in Table 40 for the analysis without the plastic surgeon at debridement and Table 41 including plastics at debridement.

Table 40: Cost of debridement without plastic surgeon present

	Timing of debridement			
	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Theatre staff cost	£1,758	£1,713	£1,668	£1,668

	Timing of debridement			
	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Complications cost	£1,379	£1,657	£2,375	£4,677
TOTAL	£3,137	£3,370	£4,043	£6,345

Table 41: Cost of debridement with plastic surgeon present

	Timing of debridement			
	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Theatre staff cost	£2,453	£2,358	£2,263	£2,263
Complications cost	£524	£630	£903	£1,777
TOTAL	£2,978	£2,988	£3,166	£4,041

As can be seen from above, the results show that at each time point, it is cheaper to have a plastic surgeon present at each time point because the increase in the plastic surgeon cost is more than covered by the savings from the reduced adverse events. However, both with and without a plastic surgeon, as the time to debridement increases, there is an increasing cost due to the increasing risks of adverse events.

L.2.5.2 Sensitivity analyses

SA1: Relative risk of deep infection for the presence of a plastic surgeon threshold analysis

The threshold for the relative risk of deep infection for the presence of a plastic surgeon at which the two strategies become cost neutral for early debridement (< 6 hours) is 0.50 compared to the base case value of 0.38 as shown in Table 39 above.

SA2: Increasing the proportion of patients whose debridement is performed in premium time by 50%

Table 42: Results of SA2

Overall cost	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Without plastics	£3,182	£3,393	£4,043	£6,345
With plastics	£3,073	£3,036	£3,166	£4,041

When a plastic surgeon is present, the cheapest strategy has now become 6 to 12 hours delay to debridement. However, the difference between this strategy and debridement within 6 hours is very small.

SA3: Baseline risk of deep infection threshold analysis

The presence of a plastic surgeon becomes cost neutral for early debridement (<6 hours) when the baseline risk of deep infection is decreased to 4.28% compared to the base case value of 5.56% as shown in Table 37 above. The resulting risks for the othertimes of debridement are shown in Table 43 below. These are based on the new baseline risk and the original relative risk for each hour of delay to debridement.

Table 43: Updated risks for deep infection

Duration of delay to debridement	Risk of deep infection
6 hours	4.28%
12 hours	5.15%
24 hours	7.43%
48 hours	14.88%

Risks were converted from odds calculated using the odds ratio from Hull 2014 and the baseline risk from Noumi 2005 in the table above.

SA4: Reducing the odds ratio of deep infection per hour of delay to debridement to 1.01

Table 44: Results of SA4

Overall cost	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Without plastics	£3,137	£3,173	£3,301	£3,708
With plastics	£2,978	£2,913	£2,884	£3,038

This has substantially reduced the costs of later debridement (>24 hours) without a plastic surgeon because the risks of deep infection were particularly high for this group. Overall for debridement later than 6 hours, the costs have decreased due to a lower risk of infection and therefore amputation. The costs without the presence of a plastic surgeon are now not as high compared to a plastic surgeon being present.

SA5: Increasing prosthetic cost to £6,000

Table 45: Results of SA5

Overall cost	<6 hours	6 to 12 hours	12 to 24 hours	>24 hours
Without plastics	£3,331	£3,604	£4,378	£7,004
With plastics	£3,051	£3,077	£3,293	£4,291

As the lifetime costs of prosthetics is a particularly costly downstream resource, an increase in the costs of this will have an impact on the results because it increases the adverse event costs. However, the savings from reducing the adverse events still outweigh the staff costs from having a plastic surgeon present.

SA6: Threshold analysis for the cost of deep infection treatment

The threshold for the cost of deep infection treatment, at which the presence of a plastic surgeon becomes equally costly as without, for debridement at less than 6 hours, is £15,107 compared to the base case value of £20,000. Therefore, even if the base case estimate of £20,000 is an overestimate, the presence of a plastic surgeon at early debridement is still likely to be cheaper than without a plastic surgeon.

L.2.6 Discussion

L.2.6.1 Summary of results

The results of this cost analysis show that the costs of earlier debridement due to increased out of hours surgery are small in comparison to the costs saved from the complications avoided. This is also the case with the presence of a plastic surgeon. The additional cost of adding a plastic surgeon along

with an additional registrar is far outweighed by the costs saved from complications avoided by having the expertise of the plastic surgeon available in theatre.

SA1 shows that the relative risk of deep infection would have to go up by 0.12 for the presence of a plastic surgeon to have an equal cost to early debridement without plastics. This is a fairly large increase from 0.38, however, 0.5 is still within the lower end of the confidence interval around this parameter, please see section L.2.6.2 for further discussion on this. -+

SA2 shows that increasing the proportion of patients who will have debridement performed out of hours by 50% makes the 6 to 12 hour strategy slightly cheaper than debridement at less than 6 hours, when a plastic surgeon is present. This minimal difference is likely to be outweighed by the health related quality of life benefits from reducing deep infections and amputations and so debridement in less than 6 hours with a plastic surgeon present is still likely to remain cost effective.

SA3 shows that the risk of deep infection for debridement in less than 6 hours has to reduce from 5.56% to 4.28% to make the presence of a plastic surgeon equally costly to without plastics for debridement at that time. This is only a small change in absolute risk but the base case value is fairly small to begin with and so proportionately it is larger than it may appear.

SA4 shows a similar result to SA2 with debridement at 6 to 12 hours becoming slightly cheaper than less than 6 hours when the plastic surgeon is present. Again, this difference in cost is not large and so taking health related quality of life into account is likely to still favour debridement at less than 6 hours with a plastic surgeon present.

SA5 showed that the overall results were robust to increasing the cost of a prosthetic, as debridement in less than 6 hours with a plastic surgeon present remained the cheapest option.

SA6 showed that the cost of deep infection would have to reduce by almost £5,000 for the presence of a plastic surgeon to be cost neutral for debridement at less than 6 hours. This shows the overall costs are fairly robust to this variable and that the overall conclusions remain.

L.2.6.2 Limitations and interpretation

This analysis only considers the key cost impacts for the debridement of open fractures. It does not explicitly evaluate the health related quality of life implications relating to deep infection and amputation and no mortality has been assumed post injury.

Although the key costs have been included, there are some costs that were difficult to accurately evaluate such as the ongoing cost of rehabilitation including physiotherapy and the support required for patients to become accustomed with a prosthetic limb. Also, further downstream resource use such as potential re-operations for amputees has not been included. However, had these costs been included, they are likely to favour the presence of a plastic surgeon where the risks of adverse events are reduced, and also favour earlier debridement where these risks are smaller.

The data included for the relative risk of deep infection when the plastic surgeon is present is from a very low quality study²⁸². This was a key parameter in the analysis; however it was subject to a threshold analysis to find the value at which the strategy becomes cost neutral. It is important to note that this parameter had a large confidence interval (0.047 – 3.037) and therefore the RR used in the analysis and in turn the impact on results is uncertain. However, it was felt by the clinical experts that there are costs and benefits that have not been taken account of in this analysis such as; the resource use associated with infection, and the detriment to quality of life. Therefore, this relative risk could increase further, resulting in a more costly approach if the plastic surgeon is present, yet still remain a *cost effective* strategy.

No probabilistic analysis has been undertaken as this is a simple cost analysis with a small number of parameters. However, a range of deterministic sensitivity analyses have been performed to assess any uncertainty in the inputs.

No data was found for length of hospital stay and so has not been included in the analysis. However, this is thought to increase for those with deep infection and so is likely to favour earlier debridement with a plastic surgeon present.

L.3 Soft tissue cover cost analysis

L.3.1 Methods

To provide definitive soft tissue cover within a specified time requires the availability of a plastic surgeon within the specified time. Soft tissue cover procedures are lengthy and so it is not appropriate for them to be performed by an on-call surgeon. Therefore the cost implications will be assessed by evaluating the cost of additional trauma surgery lists each week to allow for the surgery within a specific timeframe. It is assumed in this analysis that the surgery lists are only used for patients with open fractures who require plastic surgery. There may, of course, be other patients who can benefit from these additional resources and so the results presented are likely to overestimate the cost per patient and underestimate the overall clinical benefits. This is considered further in the discussion of the results and the conclusion.

The costing analysis for this question will include the core theatre staff, an orthopaedic surgeon and a plastic surgeon. Deep infection, amputation (including prosthetics) and length of hospital stay will also be calculated based on the risks from the clinical review, and these will be combined with the staff costs to give an overall cost for each intervention.

The average cost per patient will be presented assuming that all patients who require plastic surgery for definitive soft tissue cover will be transported to the nearest Major Trauma Centre where these skills are available.

L.3.2 Epidemiology

Table 46: Open fractures requiring plastic surgery

	Data	Source
Annual incidence (per 100,000 population)	5.16	BOA/BAPS report ⁹⁸
Population of England (millions)	53.0	2011 Census, Office for National Statistics ¹
Expected number in England per year	2,779	Calculated from above
Number of MTCs in England	26	NHS Major Trauma Centres Map ²
Expected number of fractures per MTC per year	105	Calculated from above

L.3.3 Inputs

L.3.3.1 Resource use and unit costs

The same resource use and unit costs presented in sections L.2.2.1 and (a) were used for this analysis. These values were used to calculate the cost of a theatre list as described in Table 47. The number of lists required each week to meet the each timing of cover strategy are given in Table 48 below shows the number of surgery lists required each week to facilitate soft tissue cover within a certain number of days as per the interventions listed in our review questions.

Table 48 below.

Table 47: Theatre list costs

Input	Data
Cost of theatre staff per hour ^(a)	£754
Number of hours per theatre list	8
Cost per theatre list	£6,035
Annual cost (for each theatre list per week)	£313,841

(a) Calculated in the debridement analysis in Table 31.

Table 48 below shows the number of surgery lists required each week to facilitate soft tissue cover within a certain number of days as per the interventions listed in our review questions.

Table 48: Number of lists required

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1

L.3.3.2 Clinical effectiveness data

Clinical data on risk of deep infection associated with the delay to soft tissue cover procedure were obtained from the systematic review conducted for this guideline (Section 6.9 in the full guideline). The main data used for this analysis are reported in the Table below.

Table 49: Risks of deep infection by delay to cover

Delay	Risk	Source
≤ 3 days	4.17%	Liu 2012 ²³⁷
4 to 7 days	7.69%	Liu 2012 ²³⁷
>7 days	21.4%	Liu 2012 ²³⁷

Based on the data reported in Liu 2012, to estimate the risk of deep infection per day, a line of best fit was fitted using the midpoints of the ranges used in the study (more detail on this can be found in section L.3.4). The obtained risk estimates are reported in the table below.

Table 50: Risks of deep infection by delay to cover (estimated from line of best fit)

Delay to cover	Risk	Source
1 day	3.33%	Estimate based on Liu 2012 ²³⁷ data
2 days	4.39%	Estimate based on Liu 2012 ²³⁷ data
3 days	5.45%	Estimate based on Liu 2012 ²³⁷ data
4 days	6.51%	Estimate based on Liu 2012 ²³⁷ data
7 days	9.68%	Estimate based on Liu 2012 ²³⁷ data

Similarly to the previous analysis, a proportion of patients experiencing deep infection would also have amputation. The same data used in the previous analysis was applied.

From the systematic review conducted for this guideline, we also obtained data on the length of stay associated with the delay to the soft tissue cover procedure {CROSS REFER TO SYST REVIEW}. The main data used for this analysis are reported in the tables below.

Table 51: Length of stay by delay to cover

Delay	Data	Source
≤ 3 days	20.0 days	Liu 2012 ²³⁷
4 to 7 days	24.8 days	Liu 2012 ²³⁷
>7 days	36.2 days	Liu 2012 ²³⁷

Based on the data reported in Liu 2012, to estimate the average length of stay per day of delay, a line of best fit was fitted using the midpoints of the ranges used in the study (more detail on this can be found in section L.3.4). The obtained length of stay estimates are reported in the table below.

Table 52: Length of stay by delay to cover (estimated from line of best fit)

Delay	Data	Source
1 day	19.9 days	Estimate based on Liu 2012 ²³⁷ data
2 days	20.9 days	Estimate based on Liu 2012 ²³⁷ data
3 days	21.9 days	Estimate based on Liu 2012 ²³⁷ data
4 days	22.8 days	Estimate based on Liu 2012 ²³⁷ data
7 days	25.7 days	Estimate based on Liu 2012 ²³⁷ data

L.3.4 Computations

The risk of complications and length of hospital stay for each day of delay was estimated by fitting a line of best fit through the midpoints of the ranges given in the studies. The best fit was achieved by using the Solver package in Microsoft Excel to minimise the square of the errors between the midpoints and the line while varying the gradient and constant term of the line.

The total cost per patient for each strategy was calculated as:

$$\text{Cost per patient strategy } x = N_{\text{list } x} * \text{Cost}_{(7\text{days})} + \text{CostComplications} * P \text{ Complication}_x + \text{CostLoS} * \text{LoS}_x$$

Where:

$N_{\text{list } x}$ is the number of lists required for strategy x

$\text{Cost}_{(7\text{days})}$ is the cost of lists per patient for the strategy '7 days'

CostComplications is the cost of complications

$P \text{ Complication}_x$ is the probability of complications for strategy x

CostLoS is the cost per day of hospital stay

and LoS_x is the LoS for strategy x.

L.3.5 Sensitivity analyses

A number of sensitivity analyses were undertaken to test the robustness of the results.

SA1: Risk of amputation for those with deep infection increased to 50%

This analysis increases the risk of amputation to 50% of those who have deep infection and assess the effect on the overall results.

SA2: Increasing cost of prosthetics to £6,000

This analysis increases the cost of prosthetics to £6,000 and assesses the effect on the overall results.

SA3: Reducing the cost of deep infection to £15,000

This analysis reduces the cost of deep infection to £15,000 and assesses the effect on the overall results.

L.3.6 Results

L.3.6.1 Base case analysis

Table 53: Theatre list costs on a population level

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per year	£2,196,889	£1,255,365	£941,524	£627,682	£313,841
Cost of outcomes per year	£663,471	£720,332	£777,194	£834,056	£1,004,642
Total cost	£2,860,359	£1,975,697	£1,718,718	£1,461,739	£1,318,483

Table 54: Theatre list costs per patient

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per person	£20,900	£11,943	£8,957	£5,971	£2,986
Cost of outcomes per person	£6,312	£6,853	£7,394	£7,935	£9,558
Total cost	£27,212	£18,796	£16,351	£13,906	£12,543

The results above show that the more theatre lists that are provided, the more this will cost overall. Although this also reduces the cost of adverse events by providing cover quicker, the increased cost of additional lists is higher than the cost savings from reduced adverse events.

L.3.6.2 Sensitivity analyses

SA1: Risk of amputation for those with deep infection increased to 50%

Table 55: Results of SA1 on a population level

Time to cover (days)	1	2	3	4	7
----------------------	---	---	---	---	---

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per year	£2,196,889	£1,255,365	£941,524	£627,682	£313,841
Cost of outcomes per year	£750,412	£834,904	£919,397	£1,003,889	£1,257,367
Total cost	£2,947,300	£2,090,269	£1,860,921	£1,631,572	£1,571,208

Table 56: Results of SA1 per patient

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per person	£20,900	£11,943	£8,957	£5,971	£2,986
Cost of outcomes per person	£7,139	£7,943	£8,747	£9,550	£11,962
Total cost	£28,039	£19,886	£17,704	£15,522	£14,948

The cost of outcomes has increased as the number of amputations has increased. The increase is larger for later time points.

SA2: Increasing cost of prosthetics to £6,000

Table 57: Results of SA2 on a population level

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per year	£2,196,889	£1,255,365	£941,524	£627,682	£313,841
Cost of outcomes per year	£676,395	£737,365	£798,334	£859,304	£1,042,212
Total cost	£2,873,284	£1,992,730	£1,739,858	£1,486,986	£1,356,053

Table 58: Results of SA2 per patient

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per person	£20,900	£11,943	£8,957	£5,971	£2,986
Cost of outcomes per person	£6,435	£7,015	£7,595	£8,175	£9,915
Total cost	£27,335	£18,958	£16,552	£14,146	£12,901

SA3: Reducing the cost of deep infection to £15,000

Table 59: Results of SA3 on a population level

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per year	£2,196,889	£1,255,365	£941,524	£627,682	£313,841
Cost of outcomes per year	£645,961	£697,258	£748,556	£799,853	£953,744
Total cost	£2,842,850	£1,952,623	£1,690,079	£1,427,535	£1,267,586

Table 60: Results of SA3 per patient

Time to cover (days)	1	2	3	4	7
Number of lists needed per week	7	4	3	2	1
Cost of list(s) per person	£20,900	£11,943	£8,957	£5,971	£2,986
Cost of outcomes per person	£6,145	£6,633	£7,121	£7,609	£9,073
Total cost	£27,045	£18,576	£16,078	£13,581	£12,059

L.3.7 Discussion

L.3.7.1 Summary of results

The results showed that the cost of providing an 8 hour orthoplastic theatre list is estimated to be £6,035. The GDG believed that current practice was to have two dedicated theatre lists per week, which would cost an estimated £627,682 per year. This can only guarantee soft tissue cover within four days and the estimated complication cost for this is £834,056 per year. Increasing the number of lists to three per week at an annual cost of £941,524 is estimated to reduce the cost of complications to £777,194. Hence, the overall annual costs for two and three lists per week are £1,461,739 and £1,718,718 respectively. There is, therefore, an increase in costs overall for performing soft tissue cover within 72 hours but this needs to be considered along with the health related quality of life benefits that come with the reduction in complications.

The analysis estimated the mean number of patients who present to a major trauma centre (either directly or indirectly) with an open fracture requiring plastic surgery as 105. On the assumption that these surgery lists will only be used for these patients, the estimated cost per patient for two lists and three lists per week respectively would be £13,906 and £16,351; an increase of £2,445 per patient for the additional list. This would require a mean increase in QALYs per patient of 0.12 in order to be cost effective. Over the estimated life years remaining of 38 years that was used in the model, this equates to a mean increase in utility of 0.003 for the duration of that period. Also taking into account the fact that this is based on assuming that staff are only working when a patient arrives, this is a conservative estimate. In reality, the staff can perform other elective surgical work that can be cancelled at short notice to accommodate any emergency arrivals. Therefore additional lists can be of benefit to other patient groups, which have not been considered here, thus cost effectiveness of additional lists may be underestimated. Increasing the number of lists per week to

four would add an additional £2,445 to the overall cost and therefore require an additional 0.12 QALYs in order to be cost effective. Providing a list every day would cost an additional £8,469 per person compared to four lists per week and this would require an additional 0.42 QALYs in order to be cost effective in comparison to four lists a week.

The results remained robust to changes in all sensitivity analyses undertaken and the conclusions did not change.

L.3.7.2 Limitations and interpretation

The evidence of risks for timing of cover is based on free flaps for people with Gustilo-Anderson grade 3 fractures and so may overestimate the risks for the whole population.

This analysis assumes that all patients have the delay to cover specified in our review protocol. In reality, some patients will arrive at a time that allows for earlier treatment and so this will overestimate the risks. This applies to all strategies; however there is less variability possible for the earlier strategies i.e. with a list every day the patient can be delayed between 0 and 24 hours depending on their arrival time and for a list every 3 days a patient could be delayed between 0 and 72 hours depending on their arrival time. Assuming that they are delayed for the maximum time may slightly favour the earlier strategy.

L.4 Multiple theatre sessions cost analysis

L.4.1 Methods

The three main types of surgical procedure performed on patients with an open fracture (debridement, fixation and cover) can be performed in one theatre session or across two or three. The first session is always debridement and an initial fixation (temporary or definitive) but a later second session can be used to perform definitive fixation and cover if only temporary fixation was used initially and a third session can be used to perform definitive cover at a later time. The use of multiple theatre sessions increases the time needed for preparation and so increases the costs. However, if the soft tissue cover procedure is performed in the same session following fixation, then there is an inefficient use of the plastic surgeon that is made available for the entire theatre session but is only needed for part of it. This is because the plastic surgeon has to do the final procedure for soft tissue cover as soon as the orthopaedic surgeon has completed the definitive fixation. The plastic surgeon cannot therefore perform any other work in the time before this procedure in case the work takes longer than anticipated.

This cost analysis will evaluate these trade-offs together to assess whether the inefficient use of the plastic surgeons time could actually be cost saving. It is based on the assumed durations of the procedures and preparation of the GDG and uses the staff costs as presented previously. No clinical outcomes are included in this analysis.

L.4.2 Inputs

L.4.2.1 Resource use and unit costs

The same resource use and unit costs presented in section L.2.2.1 were used for this analysis to cost surgery time during normal hours. The duration of each procedure performed for an open fracture is shown in Table 61 below.

Table 61: Duration of procedures

Procedure	Duration (hours)
Debridement	1
Temporary fixation	2
Definitive fixation	3
Local flap	2
Theatre session preparation	2

L.4.3 Results

L.4.3.1 Base case analysis

Table 62: Cost of different pathways with and without plastics present at debridement

Pathway	Cost without plastics present	Cost with plastics present
Strategy 1: Debridement, definitive fixation and definitive cover in one theatre session	£6,035.41	£6,035.41
Strategy 2: Debridement and definitive fixation in one theatre session followed by definitive cover in a later session	£5,561.09	£6,156.04
Strategy 3: Debridement and temporary fixation in one theatre session followed by definitive fixation and definitive cover in a later session.	£7,664.89	£8,259.85
Strategy 4: Debridement and temporary fixation in one theatre session followed by definitive fixation in a later session and definitive cover in a third session.	£7,785.53	£8,380.48

As can be seen from the table above, all strategies are more expensive with a plastic surgeon present except for the first strategy with all procedures in one session, which is equally costly. This is because the plastic surgeon has to be available to perform the soft tissue cover regardless of whether they are present for debridement or not.

Strategy 2 is cheaper if the plastic surgeon is not present at debridement as it removes the plastic surgeon salary cost during debridement and fixation, while adding a smaller cost of the additional preparation time for the second session. When the plastic surgeon is present for debridement strategy 2 is slightly more expensive than 1 as the only difference in cost is from the additional preparation time.

Strategy 3 has a large increase in cost compared to strategy 2 as there is the additional cost of the temporary fixation procedure. The cost of the plastic surgeon for the duration of the definitive fixation procedure still applies.

Strategy 4 has a small increase in cost compared to strategy 3 as there is the additional cost for preparing a third theatre sessions but there is no cost of having the plastic surgeon available during any fixation procedure.

L.4.4 Discussion

L.4.4.1 Summary of results

This cost analysis shows that if a plastic surgeon is not present for debridement then it is cheaper to have definitive cover in a separate session so that the plastic surgeon's time is used more efficiently. If the plastic surgeon is to be present for debridement however, it is then cheaper to have all procedures performed in the same session. This is because the additional preparation time for a second theatre session for definitive cover outweighs the inefficient use of the plastic surgeon while definitive fixation is performed between debridement and definitive cover.

If definitive fixation is delayed to the second session along with definitive cover then the inefficient use of the plastic surgeon still applies as well as the additional preparation time. This is therefore more expensive. Having definitive fixation in a second session and definitive cover in a third is even more expensive due to having another preparation time added for the third theatre session.

L.4.4.2 Limitations and interpretation

This analysis is based on assumptions of the durations of procedures and preparation time provided by the GDG. This analysis does not take into account the different costs of surgical implants required across each strategy.

L.5 Conclusion

From analysis 1: Debridement performed within 6 hours of injury with a plastic surgeon present in theatre is the most likely cost effective strategy and may even be cost saving.

From analysis 2: One theatre list per week is not enough to meet the demand based on the incidence of open fractures and so two lists a week is generally regarded as current practice in the UK. The increase in costs per person for three surgery lists compared to two only requires a small improvement in health related quality of life, which is potential feasible. A further list would add the same overall cost but there may not be as much benefit to be gained and so the cost effectiveness of this is less certain. A list every day would add a much greater cost and so the uncertainty in cost effectiveness of this based on the clinical evidence available is much more uncertain. The incidence of open fractures is low and is an important consideration with regards to the cost effectiveness. However there may be other population that would benefit from additional theatre lists.

From analysis 3: If a plastic surgeon is to be present at debridement as the first analysis suggest is cost saving, performing all procedures in one session can save further costs. However, this may not always be possible due to the restrictions of the conclusions to the other analyses.

The analyses are inter-related as they reflect different parts of the same pathway. It may be possible that a costly change in strategy in one part of the pathway could be offset by savings made via a change in strategy in another part of the pathway. For the overall conclusions of these analyses and the discussion given by the GDG, please see the link to evidence section for treatment of open fractures {please see section 6.9.6 of the Complex fracturesfull guideline}.

Appendix M: Research recommendations

M.1 Cystourethrogram

Research question: How accurate is the first CT scan with contrast (trauma scan) for detecting bladder injuries in people with suspected bladder injuries after a traumatic incident?

Why this is important: Bladder injuries usually occur in people with high-energy pelvic fractures after a traumatic incident. Currently people with suspected bladder injuries have a CT scan with intravenous contrast (a trauma scan) to diagnose non-bladder injuries. People who do not have injuries needing urgent treatment may then either be given another CT scan or a fluoroscopic cystogram to check for bladder injury. People with injuries needing urgent treatment (for example, bleeding or a neurological injury) are taken to the resuscitation room after the initial CT scan (trauma scan). Once the person's condition is stabilised they are taken to either the CT or fluoroscopy suite for a retrograde cystogram to check for bladder injury. The Guideline Committee agreed that these strategies are accurate for the diagnosis of bladder injuries, but felt that there were advantages to a strategy that did not involve a second set of images. The Guideline Committee was interested in whether the first CT scan with intravenous contrast (trauma scan) could accurately diagnose bladder injuries.

Criteria for selecting high priority research recommendations:

PICO question	<p>Population</p> <ul style="list-style-type: none"> People with suspected bladder injury after a traumatic incident. This would include multiply injured patients. <p>Index test</p> <ul style="list-style-type: none"> Trauma CT with IV contrast (no additional scanning). The contrast should be administered as early as is safely possible. <p>Reference standards</p> <ul style="list-style-type: none"> Later imaging: cystogram (CT / conventional fluoroscopy) Later clinical and surgical findings <p>Outcomes</p> <ul style="list-style-type: none"> Diagnostic accuracy <p>Stratify/subgroup</p> <ul style="list-style-type: none"> Pelvic fracture type (or no fracture)
Importance to patients or the population	If the initial trauma CT with contrast is found to have the requisite diagnostic accuracy for diagnosing bladder injuries then it would be a much faster strategy than the two scan approach currently in place. The GDG agreed that earlier definitive diagnosis of bladder injuries would lead to better outcomes for patients. The better outcomes would be realised through faster diagnosis of bladder injury, no dedicated further imaging for bladder injury that could impede or delay treatment of the patient and increase their radiation burden.
Relevance to NICE guidance	It would inform the Complex Fractures guideline question around the most effective method for diagnosis of bladder injuries.
Relevance to the NHS	Accurately identifying the injury using only one scan (the initial trauma CT scan) would mean management decisions would be made faster because the need for additional investigations is negated. This would lead to less downstream resource use in terms of imaging and staff time – which have an opportunity cost, and also potentially improve outcomes because; the injury can be

	diagnosed quicker, and also because in the context of a multiply injured patient – there is less time spent exploring the bladder injury which could be at the detriment of the other injuries the patient has.
National priorities	There are no specific national priorities pertaining to the diagnostic imaging of people with suspected bladder injuries.
Current evidence base	The studies included in the diagnostic accuracy review did not encompass the strategy proposed in this research recommendation. They investigated the accuracy of a dedicated cystogram for diagnosing bladder injuries. In addition they were relatively old studies, published in 2006 or earlier, but using two, four or occasionally 16 slice multi-detector CT machines. Ten years ahead and modern CT scanners can be dual source and reach 128 slices. The possibilities for diagnosis that may not have been considered using the previous generations of scanners may now be a reality.
Equality	This research recommendation would potentially benefit all children, young people and adults who are involved in a traumatic incident and are suspect of bladder injury.
Study design	A diagnostic accuracy study would be the most appropriate form of research methodology for this question.
Feasibility	The research would be very feasible, with low cost and no serious technical issues. It would require very little change in practice as the index test and reference standards are part of current clinical practice.
Other comments	Those interpreting the cystogram (reference standard) should be blinded to the bladder injury results of the trauma CT with IV contrast (index test). The timing of administration of contrast is important because very early administration could allow some of the contrast to reach the bladder, increasing the accuracy of the scan.
Importance	This research recommendation is of high importance: the research is essential to inform future updates of key recommendations in the guideline

M.2 Pilon fractures

Research question: In adults with closed pilon fractures what method of fixation provides the best clinical and cost effective outcomes as assessed by function and incidence of major complications at 2 years? (stratified for timing of definitive surgery early [<36hrs] vs later [>36hrs])

Why this is important: Pilon fractures involve a significant proportion of the weight-bearing surface of the distal tibia. The damaged joint surface is vulnerable to degeneration. Therefore, the injury can lead to long-term disability, most commonly arthritis with pain and stiffness. Surgery can improve outcomes, allowing reduction and fixation of the fracture and early movement of the ankle joint. However, it has a high incidence of serious complications, particularly related to the vulnerability of the soft tissues around the ankle. The potential for life-changing adverse consequences of both the injury and its treatment is known, but the best management strategy to minimise these consequences is unclear.

Criteria for selecting high priority research recommendations:

PICO question	<ul style="list-style-type: none"> • Population: adults with closed pilon fractures • Intervention: fine wire frame fixation vs. internal fixation with plates and screws vs. spanning external fixation (each augmented by joint reconstruction as required)
---------------	---

	<ul style="list-style-type: none"> • Comparator: see above • Outcomes: function, health-related quality of life, major complications
Importance to patients or the population	The best management of patients with pilon fractures is unknown. Although relatively rare injuries, pilon fractures are associated with a high rate of early complications and have inevitable long-term effects on patients' function and health-related quality of life. Therefore, research which identifies the optimal management strategy is of vital importance for patients.
Relevance to NICE guidance	Research which identifies the best method and timing of surgical fixation of pilon fractures addresses a key area identified in the scope of the NICE guidelines for complex fractures.
Relevance to the NHS	The identification of the most clinical and cost-effective method of surgical fixation and timing/staging of that fixation, would improve the outcome for patients and reduce the long-term costs associated with ankle arthritis and the need for further surgery.
National priorities	Pilon fractures affect the main weight-bearing portion of the ankle joint leading to early arthritis. Arthritis of the ankle has life-long effects in terms of mobility, pain and the patients' ability to perform their work and recreational activities. Improving the diagnosis and treatment of patients with this injury has been identified as a research priority by the Orthopaedic Trauma Society and Arthritis Research UK.
Current evidence base	<p>Current evidence for the type of fixation is very limited. The two existing RCTs and the single cohort study are all at very high risk of bias. Furthermore, one of the RCTs described a method of fixation which is no longer used in the NHS as it has been associated with a higher incidence of complication such as wound breakdown. In addition, the cohort study did not specify the type of external fixation system used. Evidence for all outcomes included in the review were either imprecise or reported with very low event rates. The GDG felt the quality of the evidence underlined the need for further research in this area.</p> <p>Current evidence concerning the timing of fixation is also very limited and imprecise. The three non-randomised studies included a mixture of closed and open pilon fractures, so the timing of fixation was confounded by the extent of the soft-tissue injury. For the non-randomised study looking at closed fractures alone, there was insufficient statistical power to detect a difference between groups in the key outcome of deep infection. Again, the GDG felt the quality of the evidence underlined the need for further research.</p>
Equality	This research recommendation would potentially benefit all groups of patients.
Study design	A randomised controlled trial with stratification for timing would be the most appropriate form of research methodology for this question.
Feasibility	The research would be feasible. Although this is a relatively rare injury, the current UK model for the management of complex fractures means that pilon fractures are increasingly concentrated in a smaller number of specialist centres where there is expertise in each of the different methods of fixation.
Other comments	Potential funders of this study may include the National Institute for Health Research and Arthritis Research UK.
Importance	This research recommendation is of high importance. Pilon fractures have a high risk of early complications and cause long-term disability. The current evidence base does not allow NICE to make a clear recommendation regarding the most clinically effective and cost effective method of fixation, nor the timing of fixation. The research is essential to inform future updates of key recommendations in the guideline.

Appendix N: NICE technical team

Name	Role
Sharon Summers-Ma	Guideline Lead
Phil Alderson	Clinical Advisor
Steven Barnes	Clinical Lead
Ross Maconachie	Health Economist
Ben Doak	Guideline Commissioning Manager
Thomas Feist	Guideline Coordinator
Anne-Louise Clayton	Editor

Appendix O: Additional cost data

O.1 Assessment of cost effectiveness for diagnostic interventions and prognostic tools

The cost effectiveness of a diagnostic modality stems from how accurately it can identify people with the injury and rule out people without the injury, as well as the true prevalence of the condition within the population being imaged. For the major trauma population, who are subject to polytrauma, systemic injury and fast deterioration, cost effectiveness of a diagnostic intervention is also impacted by the trade-off between time efficiency and accuracy of the intervention, as well as the potential for incidental findings. In the absence of economic evidence for a diagnostic review, the GDG were routinely asked to consider the below when assessing cost effectiveness of a diagnostic modality for a particular indication. The same considerations were applied to prognostic reviews on risk tools. Aspects of note are detailed in the respective Evidence and Link to Recommendation section of each review.

Impact of sensitivity, specificity and prevalence on the cost effectiveness of a diagnostic intervention

A modality or risk tool with a low sensitivity will lead to more false negatives (i.e. people missed or incorrectly predicted to have low risk and therefore do not need onward management). This will impact, resource use as well as health outcomes because these people who have been missed could therefore deteriorate, which in turn leads to longer hospital stay or higher mortality. All else being equal and assuming onward management is cost effective, a diagnostic intervention with a higher sensitivity than alternatives will be cost effective.

A modality or risk tool with a low specificity will lead to more false positives (i.e. people incorrectly labelled as having a condition or at high risk needing onward management). This will impact resource use as this leads to unnecessary treatment (which may carry potential harm). All else being equal and assuming onward management is costly and may carry harm, a diagnostic intervention with a higher specificity than alternatives will be cost effective.

Prevalence is important in the consideration of cost effectiveness. If the traumatic injury or condition being investigated is not common within the population suspected of the injury, prevalence of the injury is low. This indicates that a high proportion of people will be investigated, incurring cost, without any benefit. The lower the prevalence of the condition within the population tested the less cost effective the diagnostic intervention will be, regardless of its accuracy.

Incidental findings and cost effectiveness

When employing a diagnostic modality for a particular population group, there is normally “indirect benefit” afforded to other population groups through incidental findings. The incidental findings are of particular relevance for the trauma population for two reasons. Firstly due to the potential for poly-trauma (i.e. chest trauma and major haemorrhage are not mutually exclusive conditions). Secondly, and importantly, one injury may have systemic symptoms, signs and complications (i.e. blood may collect elsewhere to the injury site). Without consideration of potential for incidental findings, the overall benefit from undertaking the diagnostic intervention and therefore cost effectiveness may be underestimated. The sensitivity of the diagnostic intervention to find ANY injury increases as you increase the number and type of injuries that you are trying to identify with one diagnostic test. Furthermore, predictive power of finding ANY injury increases as the proportion of patients with injury in the pool that you are testing increases. Where appropriate onward

management of the type of injury you are assessing is similar, i.e. in systemic injury, cost effectiveness of the diagnostic modality is increased.

On the other hand, if incidental findings are taken into consideration of cost effectiveness, it also needs to be acknowledged that the potential of definitively ruling out ANY injury decreases (that is to say specificity and negative predictive power decreases). If onward management is costly and risky (for example surgery or interventional radiology) then this can decrease the cost effectiveness of the diagnostic intervention.

Radiation risk and cost effectiveness

Please refer to the chapter in Spinal injuries.

A concern raised around imaging is the risk of radiation. This was incorporated in a sensitivity analysis in the Spinal Injuries guideline model. The cost per patient on average is low, and particularly when time preference is taken into account (i.e. discounting of future costs and benefits), the costs and health risks are minimal. None the less, all else being equal, the diagnostic test with least radiation risk will be the most cost effective.

The trade-off between time efficiency and accuracy

Some modalities such as CT may take more time (from time of presentation) to undertake than others, particularly when issues such as scheduling and reporting are taken into account. Clinicians may need time to decide whether they should undertake these modalities only following a primary assessment (whether this is clinical or prior imaging such as x-ray). Thus there is potentially a trade-off between the quicker (and sometimes more readily available modalities) yet less accurate modalities, versus taking a bit more time for a more precise diagnosis. It is assumed that as net benefit increases (due to lack of deterioration), net cost will decrease (i.e. due to reduced length of stay, less complicated and costly treatment).

The service delivery costs of enabling timely diagnostic intervention (such as providing 24/7 CT) were considered outside the remit of this guideline and further considered in Guidance for Trauma Services (CG XXX). Where appropriate this guideline cross references these considerations. The trade-off between time efficiency and accuracy is therefore reflected in determining net clinical benefit, rather than in determination of net cost.

Consideration of overall resource use and costs of a diagnostic strategy

In the absence of economic evidence, the intervention cost of the diagnostic modality, as well the cost associated with each diagnostic outcome (in terms of the indicated onward management), was considered. The total cost of a diagnostic strategy was considered as the sum of the intervention cost and the product of each diagnostic outcome and the respective costs of indicated onward management. Costs of each diagnostic strategy are offset by the net clinical benefit that the strategy brings (i.e. through incidental findings or through time efficient management).

O.1.1 Full body CT

Table 63: Imaging costs ⁹¹

Resource	Description	National average unit cost	Lower Quartile Unit Cost	Upper Quartile Unit Cost	Notes
X-ray	Direct Access Plain Film (DAPF)	£28	£22	£33	The number of data submissions for this code was 153, with 5,254,817 units of activity (examinations)
CT	Computerised Tomography Scan, one area, no contrast, 19 years and over (RA08A)	£60	£62	£62	The number of data submissions for this code was 4, with 70 units of activity (examinations)
	Computerised Tomography Scan, one area, with post contrast only, 19 years and over (RA09A)	£71	£71	£71	The number of data submissions for this code was 1, with 10 units of activity (examinations)
	Computerised Tomography Scan, one area, pre and post contrast (RA10Z)	£301	£301	£301	The number of data submissions for this code was 1, with 1 unit of activity (examinations)
	Computerised Tomography Scan, two areas without contrast (RA11Z)	£58	£58	£58	The number of data submissions for this code was 1, with 12 units of activity (examinations)
	Computerised Tomography Scan, two areas with contrast (RA12Z)	£76	£72	£72	The number of data submissions for this code was 2, with 22 units of activity (examinations)
	Computerised Tomography Scan, more than three areas (RA14Z)	£146	£102	£190	The number of data submissions for this code was 2, with 2 units of activity (examinations)

(a) For CT, the costs are from the 'trauma and orthopaedics' service description.

(b) Note for CT, there is no category under the trauma and orthopaedics service description for below 19 years of age.

(c) The number of data submissions for the activity level recorded for CT indicate that the unit cost was likely to be reflective of the costs only incurred by a few providers. This may explain why the ultrasound of more than 20 minutes costs less than the ultrasound of less than 20 minutes.

(d) Note that for some of the modalities the lower and upper quartile costs are the same, however it is reported here as it is reported in NHS reference costs 2012-13.

(e) Where the number of submissions and activity levels is low, this may imply that the cost is not likely to be representative of the national average.

O.1.2 Pelvic imaging

Table 64: Imaging costs ⁹¹

Resource	Description	National average unit cost	Lower Quartile Unit Cost	Upper Quartile Unit Cost	NOTES
X-ray	Direct Access Plain Film (DAPF)	£28	£22	£33	The number of data submissions for this code was 153, with 5,254,817 units of activity (examinations)
CT	Computerised Tomography Scan, one area, no contrast, 19 years and over (RA08A)	£60	£62	£62	The number of data submissions for this code was 4, with 70 units of activity (examinations)
	Computerised Tomography Scan, one area, with post contrast only, 19 years and over (RA09A)	£71	£71	£71	The number of data submissions for this code was 1, with 10 units of activity (examinations)

(a) For CT, the costs are from the 'trauma and orthopaedics' service description.

(b) Note for CT, there is no category under the trauma and orthopaedics service description for below 19 years of age.

(c) The number of data submissions for the activity level recorded for CT indicate that the unit cost was likely to be reflective of the costs only incurred by a few providers.

(d) Note that for some of the modalities the lower and upper quartile nit costs are the same, however it is reported here as it is reported in NHS reference costs 2012-13.

O.1.3 Cystourethrogram

Table 65: Imaging costs⁹²

Resource	Description	National average unit cost	Lower Quartile Unit Cost	Upper Quartile Unit Cost	NOTES
Fluoroscopy	Contrast Fluoroscopy Procedures, less than 20 minutes	£69	£40	£86	The number of data submissions for this code was 119, with 48,617 units of activity (examinations)
CT	Computerised Tomography Scan, one area, no contrast, 19 years and over	£80	£62	£97	The number of data submissions for this code was 124, with 90,108 units of activity (examinations)
	Computerised Tomography Scan, one area, with post contrast only, 19 years and over	£91	£70	£105	The number of data submissions for this code was 116, with 18,505 units of activity (examinations)

(a) The costs here differ from those in the tables above because the costs for this question were gathered when the latest version on NHS reference costs had been published (NHS reference costs 2013/14)

(b) The number of data submissions for the activity level recorded for CT indicate that the unit cost was likely to be reflective of the the national average.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16

O.2 Limb Salvage

Table 66: Amputation cost⁹²

Resource	Description	National average unit cost	Lower quartile unit cost	Upper quartile unit cost	Notes
Amputation	Amputation of Single Limb with CC Score 0-9 (HRG: YQ22B)	£8,589	£6,439	£10,358	Data submissions for this code was 112, with 1,378 units of activity

(a) The number of data submissions for the activity level recorded indicate that the unit cost was likely to be reflective of the the national average.

This is the acute care cost associated with an amputation. Further lifetime resource use would include further re-operations and the prodthetics.

O.3 Arterial shunts

See the previous section for amputation costs.

1 **O.4 Pelvic haemorrhage control**

2 **Table 67: Interventional radiology costs⁹²**

Intervention/ Diagnosis	Reference cost HRG	National average unit cost	Lower Quartile Unit Cost	Upper Quartile Unit Cost	Average cost of excess bed day	Lower Quartile Unit Cost	Upper Quartile Unit Cost	Weighted national average	Weighted average length of stay	NOTES
Percutaneous Transluminal Embolisation of Blood Vessel	Percutaneous Transluminal Embolisation of Blood Vessel with CC Score 3+ (YR21A); as recorded for Non- Elective Inpatients long stay	£5,465	£2,779	£6,958	£259	£203	£284	£5,987	9.92	The number of data submissions for this code was 92, with 492 units of activity.
Percutaneous Transluminal Embolisation of Blood Vessel	Percutaneous Transluminal Embolisation of Blood Vessel with CC Score 0-2 (YR21B); as recorded for Non- Elective Inpatients long stay	£3,691	£2,370	£4,335	£329	£225	£391	£4,232	4.41	The number of data submissions for this code was 57, with 130 units of activity.
Percutaneous Transluminal Embolisation of Blood Vessel	Weighted for complications and co morbidities for HRG codes: YR21A, YR21B and ; as recorded for Non-Elective Inpatients long stay							£5,620	8.77	

3 (a) The number of data submissions for the activity level recorded indicate that the unit cost was likely to be reflective of the the national average.

O.5 Detecting compartment syndrome

See section O.2 for amputation costs.

Table 68: Fasciotomy costs ⁹²

Resource	Description	National average unit cost	Lower quartile unit cost	Upper quartile unit cost	Notes
Fasciotomy	Minor Knee Procedures for Trauma, Category 2, without CC (HA25C)	£3,477	£2,333	£4,297	Data submissions for this code was 112, with 265 units of activity

(a) The number of data submissions for the activity level recorded indicate that the unit cost was likely to be reflective of the the national average.

1
2
3
4
5
6
7
8
9
10
11
12

Appendix P: Qualitative study checklist

P.1 Information and support

Table 69: <Insert Table Title here>

Link to GRADE criteria	Question	Forsberg 2014 ¹¹⁹	Sleney 2014 ³⁷²	Okonta 2011 ³⁰⁰	O'Brien 2010 ²⁹⁰
Limitations of evidence	Is a qualitative study/survey an appropriate approach?	✓	✓	✓	✓
Limitations of evidence	Is the study clear in what it seeks to do?	✓	✓	✓	✓
Limitations of evidence	How defensible/rigorous is the research design/methodology?	?	✓	✓	✓
Limitations of evidence	How well was the data collection carried out?	✓	✓	✓	✓
Limitations of evidence	Is the role of the researcher clearly described?	✗	✓	✗	✓
Limitations of evidence	Is the context clearly described?	✓	✓	✗	✓
Limitations of evidence	Were the methods reliable?	✓	✓	✓	✓
Limitations of evidence	Is the data analysis sufficiently rigorous?	?	✓	?	✓
Limitations of evidence	Are the data rich (for qualitative study and open ended survey questions)?	✓	✓	✓	✓
Limitations of evidence	Is the analysis reliable?	?	✓	?	✓
Limitations of evidence/ Applicability of evidence/ Sufficiency of evidence	Are the findings convincing?	✓	✓	✓	✓
Applicability of evidence	Are the findings relevant to the aims of the study?	✓	✓	✓	✓
Limitations of evidence/ Applicability of evidence/ Sufficiency of evidence	Are the conclusions adequate?	✓	✓	✓	✓

References

- 1 2011 Census: Population Estimates for the United Kingdom, 27 March 2011. 2012. Available from: http://www.ons.gov.uk/ons/dcp171778_292378.pdf
- 2 Major Trauma Centres. 2012. Available from: <http://www.nhs.uk/NHSEngland/AboutNHSservices/Emergencyandurgentcareservices/Documents/2012/map-of-major-trauma-centres-2012.pdf>
- 3 Abrassart S, Stern R, Peter R. Unstable pelvic ring injury with hemodynamic instability: what seems the best procedure choice and sequence in the initial management? *Orthopaedics and Traumatology, Surgery and Research*. 2013; 99(2):175-182
- 4 Adegbehingbe OO, Akinyoola AL, Oginni LM. Predictive factors for primary amputation in trauma patients in a Nigerian university teaching hospital. *East African Medical Journal*. 2006; 83(10):539-544
- 5 Agel J, Rockwood T, Barber R, Marsh JL. Potential predictive ability of the orthopaedic trauma association open fracture classification. *Journal of Orthopaedic Trauma*. 2014; 28(5):300-306
- 6 Akbar M, Arshad RM, Hanif M, Rana RE. Treatment of open book pelvic fractures: Comparison between internal and external fixation. *Pakistan Journal of Medical and Health Sciences*. 2012; 6(3):662-665
- 7 Al-Arabi YB, Nader M, Hamidian-Jahromi AR, Woods DA. The effect of the timing of antibiotics and surgical treatment on infection rates in open long-bone fractures: a 9-year prospective study from a district general hospital. *Injury*. 2007; 38(8):900-905
- 8 Al-Arabi YB, Nader M, Hamidian-Jahromi AR, Woods DA. Corrigendum to "The effect of the timing of antibiotics and surgical treatment on infection rates in open long-bone fractures: A 9-year prospective study from a district general hospital" [*Injury* 38 (8) (2007) 900-905] (DOI:10.1016/j.injury.2007.02.043). *Injury*. 2008; 39(3):381
- 9 Al-Hilli AB, Salih DS. Early or delayed surgical treatment in compound limb fractures due to high velocity missile injuries: a 5-year retrospective study from Medical City in Baghdad. *Iowa Orthopaedic Journal*. 2010; 30:94-98
- 10 al-Salman MM, al-Khawashki H, Sindigki A, Rabee H, al-Saif A, al-Salman Fachartz F. Vascular injuries associated with limb fractures. *Injury*. 1997; 28(2):103-107
- 11 Allison P, Dahan-Oliel N, Jando VT, Yang SS, Hamdy RC. Open fractures of the femur in children: analysis of various treatment methods. *Journal of Children's Orthopaedics*. 2011; 5(2):101-108
- 12 Anandakumar V, Hussein FK, Varuun B, Zhu R. Predictive parameters for angiography and embolization in the bleeding pelvic fracture. *Journal of Clinical Orthopaedics and Trauma*. 2013; 4(2):70-74
- 13 Anglen JO. Early outcome of hybrid external fixation for fracture of the distal tibia. *Journal of Orthopaedic Trauma*. 1999; 13(2):92-97
- 14 Aravind M, Shauver MJ, Chung KC. A qualitative analysis of the decision-making process for patients with severe lower leg trauma. *Plastic and Reconstructive Surgery*. 2010; 126(6):2019-2029

- 1 15 Archibald G. Patients' experiences of hip fracture. *Journal of Advanced Nursing*. 2003; 44(4):385-
2 392
- 3 16 Arti HR. Comparison of early versus delayed debridement in open fractures. *Pakistan Journal of*
4 *Medical Sciences*. 2012; 28(5):856-859
- 5 17 Asensio JA, Kuncir EJ, Garcia-Nunez LM, Petrone P. Femoral vessel injuries: analysis of factors
6 predictive of outcomes. *Journal of the American College of Surgeons*. 2006; 203(4):512-520
- 7 18 Ashford RU, Mehta JA, Cripps R. Delayed presentation is no barrier to satisfactory outcome in the
8 management of open tibial fractures. *Injury*. 2004; 35(4):411-416
- 9 19 Ashley RK, Larsen LJ, James PM. Reduction of dislocation of the hip in older children: a
10 preliminary report. *Journal of Bone and Joint Surgery American Volume*. 1972; 54(3):545-550
- 11 20 Atchison KA, Black EE, Leathers R, Belin TR, Abrego M, Gironde MW et al. A qualitative report of
12 patient problems and postoperative instructions. *Journal of Oral and Maxillofacial Surgery*. 2005;
13 63(4):449-456
- 14 21 Azam N, Harrison M. Patients' perspectives on injuries. *Emergency Medicine Journal*. 2011;
15 28(7):601-603
- 16 22 Babis GC, Vayanos ED, Papaioannou N, Pantazopoulos T. Results of surgical treatment of tibial
17 plafond fractures. *Clinical Orthopaedics and Related Research*. 1997;(341):99-105
- 18 23 Back DA, Scheuermann-Poley C, Willy C. Recommendations on negative pressure wound therapy
19 with instillation and antimicrobial solutions'- 'when, where and how to use: "What does the
20 evidence show? *International Wound Journal*. 2013; 10(S1):32-42
- 21 24 Bacon S, Smith WR, Morgan SJ, Hasenboehler E, Philips G, Williams A et al. A retrospective
22 analysis of comminuted intra-articular fractures of the tibial plafond: Open reduction and
23 internal fixation versus external Ilizarov fixation. *Injury*. 2008; 39(2):196-202
- 24 25 Bagley CHM, Hunter AR, Bacarese-Hamilton IA. Patients' misunderstanding of common
25 orthopaedic terminology: the need for clarity. *Annals of the Royal College of Surgeons of*
26 *England*. 2011; 93(5):401-404
- 27 26 Baldwin KD, Babatunde OM, Russell Huffman G, Hosalkar HS. Open fractures of the tibia in the
28 pediatric population: a systematic review. *Journal of Children's Orthopaedics*. 2009; 3(3):199-208
- 29 27 Ball CG, Kirkpatrick AW, Rajani RR, Wyrzykowski AD, Dente CJ, Vercruysse GA et al. Temporary
30 intravascular shunts: when are we really using them according to the NTDB? *American Surgeon*.
31 2009; 75(7):605-607
- 32 28 Ball CG, Rozycki GS, Feliciano DV. Upper extremity amputations after motor vehicle rollovers.
33 *Journal of Trauma*. 2009; 67(2):410-412
- 34 29 Baloch MK, Bakhsh K, Masood AQ, Khan A, Ahmed J. Early outcome of operative fixation of tibial
35 pilon fractures. *Medical Forum Monthly*. 2009; 20(3):13-17
- 36 30 Barleben A, Jafari F, Rose JJ, Dolich M, Malinoski D, Lekawa M et al. Implementation of a cost-
37 saving algorithm for pelvic radiographs in blunt trauma patients. *Journal of Trauma*. 2011;
38 71(3):582-584

- 1 31 Barquet A. A vascular necrosis following traumatic hip dislocation in childhood: factors of
2 influence. *Acta Orthopaedica Scandinavica*. 1982; 53(5):809-813
- 3 32 Barquet A. Traumatic anterior dislocation of the hip in childhood. *Injury*. 1982; 13(5):435-440
- 4 33 Barros D'Sa AAB, Harkin DW, Blair PHB, Hood JM, McIlrath E. The Belfast approach to managing
5 complex lower limb vascular injuries. *European Journal of Vascular and Endovascular Surgery*.
6 2006; 32(3):246-256
- 7 34 Bartlett CS, Weiner LS, Yang EC. Treatment of type II and type III open tibia fractures in children.
8 *Journal of Orthopaedic Trauma*. 1997; 11(5):357-362
- 9 35 Bauman M, Marinaro J, Tawil I, Crandall C, Rosenbaum L, Paul I. Ultrasonographic determination
10 of pubic symphyseal widening in trauma: the FAST-PS study. *Journal of Emergency Medicine*.
11 2011; 40(5):528-533
- 12 36 Baylis TB, Norris BL. Pelvic fractures and the general surgeon. *Current Surgery*. 2004; 61(1):30-35
- 13 37 Beard JD, Davidson CM, Scott DJ, Turner AG. Pelvic injuries associated with traumatic abduction
14 of the leg. *Injury*. 1988; 19(5):353-356
- 15 38 Beck JJW. Trauma imaging in and out of conflict: A review of the evidence. *Radiography*. 2012;
16 18(4):292-295
- 17 39 Bednar DA, Parikh J. Effect of time delay from injury to primary management on the incidence of
18 deep infection after open fractures of the lower extremities caused by blunt trauma in adults.
19 *Journal of Orthopaedic Trauma*. 1993; 7(6):532-535
- 20 40 Bergman NJ. Reduction of posterior dislocation of the hip. *Tropical Doctor*. 1994; 24(3):134-135
- 21 41 Berry GK, Stevens DG, Kreder HJ, McKee M, Schemitsch E, Stephen DJG. Open fractures of the
22 calcaneus: a review of treatment and outcome. *Journal of Orthopaedic Trauma*. 2004; 18(4):202-
23 206
- 24 42 Bevevino AJ, Dickens JF, Potter BK, Dworak T, Gordon W, Forsberg JA. A model to predict limb
25 salvage in severe combat-related open calcaneus fractures. *Clinical Orthopaedics and Related
26 Research*. 2014; 472(10):3002-3009
- 27 43 Bhandari M, Guyatt GH, Swiontkowski MF, Schemitsch EH. Treatment of open fractures of the
28 shaft of the tibia. *Journal of Bone and Joint Surgery - British Volume*. 2001; 83(1):62-68
- 29 44 Bhandari M, Matta J, Ferguson T, Matthys G. Predictors of clinical and radiological outcome in
30 patients with fractures of the acetabulum and concomitant posterior dislocation of the hip.
31 *Journal of Bone and Joint Surgery British Volume*. 2006; 88(12):1618-1624
- 32 45 Biffl WL, Smith WR, Moore EE, Gonzalez RJ, Morgan SJ, Hennessey T et al. Evolution of a
33 multidisciplinary clinical pathway for the management of unstable patients with pelvic fractures.
34 *Annals of Surgery*. 2001; 233(6):843-850
- 35 46 Binda T, Faluomi M, D'Angelo F, Cherubino P. Tibial pilon fractures: Results in the medium- to
36 long-term. *Journal of Orthopaedics and Traumatology*. 2011; 12:S91
- 37 47 Blauth M, Bastian L, Krettek C, Knop C, Evans S. Surgical options for the treatment of severe tibial
38 pilon fractures: a study of three techniques. *Journal of Orthopaedic Trauma*. 2001; 15(3):153-160

- 1 48 Blick SS, Brumback RJ, Lakatos R, Poka A, Burgess AR. Early prophylactic bone grafting of high-
2 energy tibial fractures. *Clinical Orthopaedics and Related Research*. 1989;(240):21-41
- 3 49 Blum ML, Esser M, Richardson M, Paul E, Rosenfeldt FL. Negative pressure wound therapy
4 reduces deep infection rate in open tibial fractures. *Journal of Orthopaedic Trauma*. 2012;
5 26(9):499-505
- 6 50 Bosse MJ, MacKenzie EJ, Kellam JF, Burgess AR, Webb LX, Swiontkowski MF et al. An analysis of
7 outcomes of reconstruction or amputation after leg-threatening injuries. *New England Journal of*
8 *Medicine*. 2002; 347(24):1924-1931
- 9 51 Bouzat P, Broux C, Ageron F-X, Gros I, Levrat A, Thouret J-M et al. Impact of a trauma network on
10 mortality in patients with severe pelvic trauma. *Annales Francaises D'Anesthesie Et De*
11 *Reanimation*. 2013; 32(12):827-832
- 12 52 Bremmer DN, Miller AD, Bookstaver PB, Cairns M, Lindley KT, Koon J. Retrospective review of
13 antibiotic prophylaxis in open lower extremity fractures. *Pharmacotherapy*. 2012; 32(10):e292-
14 e293
- 15 53 Breugem CC, Strackee SD. Is there evidence-based guidance for timing of soft tissue coverage of
16 grade III B tibia fractures? *International Journal of Lower Extremity Wounds*. 2006; 5(4):261-270
- 17 54 Brooker AFJ, Pezeshki C. Tissue pressure to evaluate compartmental syndrome. *Journal of*
18 *Trauma*. 1979; 19(9):689-691
- 19 55 Buckley SL, Smith GR, Sponseller PD, Thompson JD, Robertson WWJ, Griffin PP. Severe (type III)
20 open fractures of the tibia in children. *Journal of Pediatric Orthopaedics*. 1996; 16(5):627-634
- 21 56 Burgess AR, Eastridge BJ, Young JW, Ellison TS, Ellison PSJ, Poka A et al. Pelvic ring disruptions:
22 effective classification system and treatment protocols. *Journal of Trauma*. 1990; 30(7):848-856
- 23 57 Burgess AR, Poka A, Brumback RJ, Flagle CL, Loeb PE, Ebraheim NA. Pedestrian tibial injuries.
24 *Journal of Trauma*. 1987; 27(6):596-601
- 25 58 Byrd HS, Cierny G, Tebbetts JB. The management of open tibial fractures with associated soft-
26 tissue loss: external pin fixation with early flap coverage. *Plastic and Reconstructive Surgery*.
27 1981; 68(1):73-82
- 28 59 Byrd HS, Spicer TE, Cierny G, III. Management of open tibial fractures. *Plastic and Reconstructive*
29 *Surgery*. 1985; 76(5):719-730
- 30 60 Calhoun JH, Anger DM, Ledbetter BR, Cobos JA, Mader JT. The Ilizarov fixator and
31 polymethylmethacrylate-antibiotic beads for the treatment of infected deformities. *Clinical*
32 *Orthopaedics and Related Research*. 1993;(295):13-22
- 33 61 Calori GM, Tagliabue L, Mazza E, de Bellis U, Pierannunzii L, Marelli BM et al. Tibial pilon
34 fractures: which method of treatment? *Injury*. 2010; 41(11):1183-1190
- 35 62 Caputo ND, Stahmer C, Lim G, Shah K. Whole-body computed tomographic scanning leads to
36 better survival as opposed to selective scanning in trauma patients: A systematic review and
37 meta-analysis. *Journal of Trauma and Acute Care Surgery*. 2014; 77(4):534-539
- 38 63 Carroll PR, McAninch JW. Major bladder trauma: The accuracy of cystography. *Journal of*
39 *Urology*. 1983; 130(5):887-888

- 1 64 Cavadas PC, Landin L, Ibanez J. Temporary catheter perfusion and artery-last sequence of repair
2 in macroreplantations. *Journal of Plastic, Reconstructive and Aesthetic Surgery*. 2009;
3 62(10):1321-1325
- 4 65 Chambers LW, Green DJ, Sample K, Gillingham BL, Rhee P, Brown C et al. Tactical surgical
5 intervention with temporary shunting of peripheral vascular trauma sustained during Operation
6 Iraqi Freedom: one unit's experience. *Journal of Trauma*. 2006; 61(4):824-830
- 7 66 Chu RS, Browne GJ, Lam LT. Traction splinting of femoral shaft fractures in a paediatric
8 emergency department: time is of the essence? *Emergency Medicine*. 2003; 15(5-6):447-452
- 9 67 Cierny G, III, Byrd HS, Jones RE. Primary versus delayed soft tissue coverage for severe open tibial
10 fractures. A comparison of results. *Clinical Orthopaedics and Related Research*. 1983;(178):54-63
- 11 68 Clamp JA, Moran CG. Haemorrhage control in pelvic trauma. *Trauma*. 2011; 13(4):300-316
- 12 69 Clement ND, MacDonald D, Ahmed I, Patton JT, Howie CR. Total femoral replacement for salvage
13 of periprosthetic fractures. *Orthopedics*. 2014; 37(9):e789-e795
- 14 70 Congdon JG. Managing the incongruities: the hospital discharge experience for elderly patients,
15 their families, and nurses. *Applied Nursing Research*. 1994; 7(3):125-131
- 16 71 Contractor D, Amling J, Brandoli C, Tosi LL. Negative pressure wound therapy with reticulated
17 open cell foam in children: an overview. *Journal of Orthopaedic Trauma*. 2008; 22(10
18 Suppl):S167-S176
- 19 72 Cook RE, Keating JF, Gillespie I. The role of angiography in the management of haemorrhage from
20 major fractures of the pelvis. *Journal of Bone and Joint Surgery - Series B*. 2002; 84(2):178-182
- 21 73 Court-Brown, Walker C, Garg A, McQueen MM. Half-ring external fixation in the management of
22 tibial plafond fractures. *Journal of Orthopaedic Trauma*. 1999; 13(3):200-206
- 23 74 Cox GW, Evans EB. Compound fracture of the tibia. *Southern Medical Journal*. 1970; 63(12):1409-
24 1414
- 25 75 Croce MA, Magnotti LJ, Savage SA, Wood GW, Fabian TC. Emergent pelvic fixation in patients
26 with exsanguinating pelvic fractures. *Journal of the American College of Surgeons*. 2007;
27 204(5):935-2
- 28 76 Cronier P, Steiger V, Rammelt S. Early open reduction and internal fixation of Pilon fractures. *Fuss
29 Und Sprunggelenk*. 2012; 10(1):12-26
- 30 77 Crutchfield EH, Seligson D, Henry SL, Warnholtz A. Tibial pilon fractures: a comparative clinical
31 study of management techniques and results. *Orthopedics*. 1995; 18(7):613-617
- 32 78 Cullen MC, Roy DR, Crawford AH, Assenmacher J, Levy MS, Wen D. Open fracture of the tibia in
33 children. *Journal of Bone and Joint Surgery - American Volume*. 1996; 78(7):1039-1047
- 34 79 Cullinane DC, Schiller HJ, Zielinski MD, Bilaniuk JW, Collier BR, Como J et al. Eastern Association
35 for the Surgery of Trauma practice management guidelines for hemorrhage in pelvic fracture--
36 update and systematic review. *Journal of Trauma*. 2011; 71(6):1850-1868

- 1 80 Curtis L. Unit costs of health and social care 2014. Canterbury: Personal Social Services Research
2 Unit, University of Kent; 2014. Available from: <http://www.pssru.ac.uk/project-pages/unit->
3 [costs/2014/index.php](http://www.pssru.ac.uk/project-pages/unit-costs/2014/index.php)
- 4 81 D'Alleyrand JC, Manson TT, Dancy L, Castillo RC, Bertumen JB, Meskey T et al. Is time to flap
5 coverage of open tibial fractures an independent predictor of flap-related complications? *Journal*
6 *of Orthopaedic Trauma*. 2014; 28(5):288-293
- 7 82 Davidovitch RI, Elkhechen RJ, Romo S, Walsh M, Egol KA. Open reduction with internal fixation
8 versus limited internal fixation and external fixation for high grade pilon fractures (OTA type
9 43C). *Foot and Ankle International*. 2011; 32(10):955-961
- 10 83 Davis Sears E, Davis MM, Chung KC. Relationship between timing of emergency procedures and
11 limb amputation in patients with open tibia fracture in the United States, 2003 to 2009. *Plastic*
12 *and Reconstructive Surgery*. 2012; 130(2):369-378
- 13 84 Davis JW, Moore FA, McIntyre J, Cocanour CS, Moore EE, West MA. Western trauma association
14 critical decisions in trauma: Management of pelvic fracture with hemodynamic instability. *Journal*
15 *of Trauma - Injury, Infection and Critical Care*. 2008; 65(5):1012-1015
- 16 85 de Palma L, Santucci A, Verdenelli A, Bugatti MG, Meco L, Marinelli M. Outcome of unstable
17 isolated fractures of the posterior acetabular wall associated with hip dislocation. *European*
18 *Journal of Orthopaedic Surgery and Traumatology*. 2014; 24(3):341-346
- 19 86 Deck AJ, Shaves S, Talner L, Porter JR. Computerized tomography cystography for the diagnosis of
20 traumatic bladder rupture. *Journal of Urology*. 2000; 164(1):43-46
- 21 87 Deck AJ, Shaves S, Talner L, Porter JR. Current experience with computed tomographic
22 cystography and blunt trauma. *World Journal of Surgery*. 2001; 25(12):1592-1596
- 23 88 Dellinger EP, Miller SD, Wertz MJ, Grypma M, Droppert B, Anderson PA. Risk of infection after
24 open fracture of the arm or leg. *Archives of Surgery*. 1988; 123(11):1320-1327
- 25 89 DeLong WG, Jr., Born CT, Wei SY, Petrik ME, Ponzio R, Schwab CW. Aggressive treatment of 119
26 open fracture wounds. *Journal of Trauma*. 1999; 46(6):1049-1054
- 27 90 Demetriades D, Martin M, Salim A, Rhee P, Brown C, Chan L. The effect of trauma center
28 designation and trauma volume on outcome in specific severe injuries. *Annals of Surgery*. 2005;
29 242(4):512-519
- 30 91 Department of Health. NHS reference costs 2012-13. 2013. Available from:
31 <https://www.gov.uk/government/publications/nhs-reference-costs-2012-to-2013> [Last accessed:
32 12 March 2015]
- 33 92 Department of Health. NHS reference costs 2013-14. 2014. Available from:
34 <http://www.gov.uk/government/publications/nhs-reference-costs-2013-to-2014> [Last accessed:
35 12 March 2015]
- 36 93 Dong JI, Zhou Ds. Management and outcome of open pelvic fractures: a retrospective study of 41
37 cases. *Injury*. 2011; 42(10):1003-1007
- 38 94 Dormagen JB, Totterman A, Roise O, Sandvik L, Klow NE. Efficacy of plain radiography and
39 computer tomography in localizing the site of pelvic arterial bleeding in trauma patients. *Acta*
40 *Radiologica*. 2010; 51(1):107-116

- 1 95 Dua A, Desai SS, Shah JO, Lasky RE, Charlton-Ouw KM, Azizzadeh A et al. Outcome predictors of
2 limb salvage in traumatic popliteal artery injury. *Annals of Vascular Surgery*. 2014; 28(1):108-114
- 3 96 Dua A, Patel B, Desai SS, Holcomb JB, Wade CE, Coogan S et al. Comparison of military and
4 civilian popliteal artery trauma outcomes. *Journal of Vascular Surgery*. 2014; 59(6):1628-1632
- 5 97 Duane TM, Dechert T, Wolfe LG, Brown H, Aboutanos MB, Malhotra AK et al. Clinical examination
6 is superior to plain films to diagnose pelvic fractures compared to CT. *American Surgeon*. 2008;
7 74(6):476-480
- 8 98 Duckworth T, Morgan BP. A Report by the BOA/BAPS Working Party on The Management of Open
9 Tibial Fractures. *British Orthopaedic Association and British Association of Plastic Surgeons*.
10 1997;1-45
- 11 99 Dwyer AJ, John B, Singh SA, Mam MK. Complications after posterior dislocation of the hip.
12 *International Orthopaedics*. 2006; 30(4):224-227
- 13 100 Eastman AL, Chason DP, Perez CL, McAnulty AL, Minei JP. Computed tomographic angiography
14 for the diagnosis of blunt cervical vascular injury: is it ready for primetime? *Journal of Trauma*.
15 2006; 60(5):925-929
- 16 101 Edwards CC. Staged reconstruction of complex open tibial fractures using Hoffmann external
17 fixation. *Clinical decisions and dilemmas. Clinical Orthopaedics and Related Research*.
18 1983;(178):130-161
- 19 102 Edwards CC, Simmons SC, Browner BD, Weigel MC. Severe open tibial fractures. Results treating
20 202 injuries with external fixation. *Clinical Orthopaedics and Related Research*. 1988;(230):98-
21 115
- 22 103 Egol KA, Tejwani NC, Capla EL, Wolinsky PL, Koval KJ. Staged management of high-energy
23 proximal tibia fractures (OTA types 41): the results of a prospective, standardized protocol.
24 *Journal of Orthopaedic Trauma*. 2005; 19(7):448-456
- 25 104 Elkhechen RJ. Open reduction with internal fixation versus limited internal fixation and external
26 fixation for high grade pilon fractures (OTA Type 43C). *Foot and Ankle International*. 2012;
27 33(9):vi
- 28 105 Elliott J, Forbes D, Chesworth BM, Ceci C, Stolee P. Information sharing with rural family
29 caregivers during care transitions of hip fracture patients. *International Journal of Integrated*
30 *Care*. 2014; 14:e018
- 31 106 Endres T, Grass R, Biewener A, Barthel S, Zwipp H. [Advantages of minimally-invasive reposition,
32 retention, and Ilizarov-(hybrid)fixation for pilon-tibial-fractures fractures with particular
33 emphasis on C2/C3 fractures]. *Der Unfallchirurg*. 2004; 107(4):273-284
- 34 107 Epstein HC. Posterior fracture-dislocations of the hip; long-term follow-up. *Journal of Bone and*
35 *Joint Surgery American Volume*. 1974; 56(6):1103-1127
- 36 108 Erdmann MW, Court-Brown, Quaba AA. A five year review of islanded distally based
37 fasciocutaneous flaps on the lower limb. *British Journal of Plastic Surgery*. 1997; 50(6):421-427
- 38 109 Ertel W, Keel M, Eid K, Platz A, Trentz O. Control of severe hemorrhage using C-clamp and pelvic
39 packing in multiply injured patients with pelvic ring disruption. *Journal of Orthopaedic Trauma*.
40 2001; 15(7):468-474

- 1 110 Evers BM, Cryer HM, Miller FB. Pelvic fracture hemorrhage. Priorities in management. Archives of
2 Surgery. 1989; 124(4):422-424
- 3 111 Falchi M, Rollandi GA. CT of pelvic fractures. European Journal of Radiology. 2004; 50(1):96-105
- 4 112 Feeney J, Jayaraman V, Luk S, Shapiro D, Virk M, Twohig M et al. Retrospective review of the
5 costs of routine pelvic X-rays in a trauma setting. American Surgeon. 2011; 77(3):337-341
- 6 113 Ferrera PC, Hill DA. Good outcomes of open pelvic fractures. Injury. 1999; 30(3):187-190
- 7 114 Fischer MD, Gustilo RB, Varecka TF. The timing of flap coverage, bone-grafting, and
8 intramedullary nailing in patients who have a fracture of the tibial shaft with extensive soft-tissue
9 injury. Journal of Bone and Joint Surgery - American Volume. 1991; 73(9):1316-1322
- 10 115 Flint L, Babikian G, Anders M, Rodriguez J, Steinberg S. Definitive control of mortality from severe
11 pelvic fracture. Annals of Surgery. 1990; 211(6):703-707
- 12 116 Fochtman A, Mittlbock M, Binder H, Kottstorfer J, Hajdu S. Potential prognostic factors
13 predicting secondary amputation in third-degree open lower limb fractures. Journal of Trauma
14 and Acute Care Surgery. 2014; 76(4):1076-1081
- 15 117 Fodor L, Sobec R, Sita-Alb L, Fodor M, Ciuce C. Mangled lower extremity: can we trust the
16 amputation scores? International Journal of Burns and Trauma. 2012; 2(1):51-58
- 17 118 Fordyce AJ. Open reduction of traumatic dislocation of the hip in a child. Case report and a
18 review of the literature. British Journal of Surgery. 1971; 58(9):705-707
- 19 119 Forsberg A, Soderberg S, Engstrom A. People's experiences of suffering a lower limb fracture and
20 undergoing surgery. Journal of Clinical Nursing. 2014; 23(1-2):191-200
- 21 120 Fox CJ, Gillespie DL, Cox ED, Kragh JFJ, Mehta SG, Salinas J et al. Damage control resuscitation for
22 vascular surgery in a combat support hospital. Journal of Trauma. 2008; 65(1):1-9
- 23 121 Francel TJ, Vander Kolk CA, Hoopes JE, Manson PN, Yaremchuk MJ. Microvascular soft-tissue
24 transplantation for reconstruction of acute open tibial fractures: timing of coverage and long-
25 term functional results. Plastic and Reconstructive Surgery. 1992; 89(3):478-487
- 26 122 Fry WR, Smith RS, Sayers DV, Henderson VJ, Morabito DJ, Tsoi EK et al. The success of duplex
27 ultrasonographic scanning in diagnosis of extremity vascular proximity trauma. Archives of
28 Surgery. 1993; 128(12):1368-1372
- 29 123 Gifford SM, Aidinian G, Clouse WD, Fox CJ, Porras CA, Jones WT et al. Effect of temporary
30 shunting on extremity vascular injury: an outcome analysis from the Global War on Terror
31 vascular injury initiative. Journal of Vascular Surgery. 2009; 50(3):549-6
- 32 124 Glass GE, Pearse M, Nanchahal J. The ortho-plastic management of Gustilo grade IIIB fractures of
33 the tibia in children: a systematic review of the literature. Injury. 2009; 40(8):876-879
- 34 125 Glenny C, Stolee P, Sheiban L, Jaglal S. Communicating during care transitions for older hip
35 fracture patients: family caregiver and health care provider's perspectives. International Journal
36 of Integrated Care. 2013; 13:e044
- 37 126 Godina M. Early microsurgical reconstruction of complex trauma of the extremities. Plastic and
38 Reconstructive Surgery. 1986; 78(3):285-292

- 1 127 Goins WA, Rodriguez A, Lewis J, Brathwaite CE, James E. Retroperitoneal hematoma after blunt
2 trauma. *Surgery, Gynecology and Obstetrics*. 1992; 174(4):281-290
- 3 128 Gonzalez A, Suva D, Dunkel N, Nicodeme JD, Lomessy A, Lauper N et al. Are there clinical
4 variables determining antibiotic prophylaxis-susceptible versus resistant infection in open
5 fractures? *International Orthopaedics*. 2014; 38(11):2323-2327
- 6 129 Gopal S, Majumder S, Batchelor AG, Knight SL, De Boer P, Smith RM. Fix and flap: the radical
7 orthopaedic and plastic treatment of severe open fractures of the tibia. *Journal of Bone and Joint
8 Surgery - British Volume*. 2000; 82(7):959-966
- 9 130 Gougoulas N, Khanna A, Maffulli N. Open tibial fractures in the paediatric population: A
10 systematic review of the literature. *British Medical Bulletin*. 2009; 91(1):75-85
- 11 131 Gougoulas N, Khanna A, McBride DJ, Maffulli N. Management of calcaneal fractures: systematic
12 review of randomized trials. *British Medical Bulletin*. 2009; 92(1):153-167
- 13 132 Granchi T, Schmittling Z, Vasquez J, Schreiber M, Wall M. Prolonged use of intraluminal arterial
14 shunts without systemic anticoagulation. *American Journal of Surgery*. 2000; 180(6):493-497
- 15 133 Greenbaum B, Zions LE, Ebramzadeh E. Open fractures of the forearm in children. *Journal of
16 Orthopaedic Trauma*. 2001; 15(2):111-118
- 17 134 Gregory RT, Gould RJ, Pecllet M. The Mangled Extremity Syndrome (M.E.S.): A severity grading
18 system for multisystem injury of the extremity. *Journal of Trauma*. 1985; 25(12):1147-1150
- 19 135 Grimard G, Naudie D, Laberge LC, Hamdy RC. Open fractures of the tibia in children. *Clinical
20 Orthopaedics and Related Research*. 1996;(332):62-70
- 21 136 Grote S, Polzer H, Prall WC, Gill S, Shafizadeh S, Banerjee M et al. [Prevention of infection in the
22 current treatment of open fractures: an evidence-based systematic analysis]. *Der Orthopade*.
23 2012; 41(1):32-42
- 24 137 Grubor P, Milicevic S, Biscevic M, Tanjga R. Selection of treatment method for pelvic ring
25 fractures. *Medicinski Arhiv*. 2011; 65(5):278-282
- 26 138 Guillamondegui OD, Mahboubi S, Stafford PW, Nance ML. The utility of the pelvic radiograph in
27 the assessment of pediatric pelvic fractures. *Journal of Trauma*. 2003; 55(2):236-240
- 28 139 Gulabi D, Toprak O, Sen C, Avci CC, Bilen E, Saglam F. The mid-term results of treatment for tibial
29 pilon fractures. *Ulusal Travma Ve Acil Cerrahi Dergisi*. 2012; 18(5):429-435
- 30 140 Guraya SY. Extremity vascular trauma in Pakistan. *Saudi Medical Journal*. 2004; 25(4):498-501
- 31 141 Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-
32 five open fractures of long bones: retrospective and prospective analyses. *Journal of Bone and
33 Joint Surgery - American Volume*. 1976; 58(4):453-458
- 34 142 Haas CA, Brown SL, Spirnak JP. Limitations of routine spiral computerized tomography in the
35 evaluation of bladder trauma. *Journal of Urology*. 1999; 162(1):51-52
- 36 143 Haasbeek JF, Cole WG. Open fractures of the arm in children. *Journal of Bone and Joint Surgery -
37 British Volume*. 1995; 77(4):576-581

- 1 144 Halvorson J, Jinnah R, Kulp B, Frino J. Use of vacuum-assisted closure in pediatric open fractures
2 with a focus on the rate of infection. *Orthopedics*. 2011; 34(7):e256-e260
- 3 145 Hammer R, Lidman D, Nettelblad H, Ostrup L. Team approach to tibial fracture. 37 consecutive
4 type III cases reviewed after 2-10 years. *Acta Orthopaedica Scandinavica*. 1992; 63(5):471-476
- 5 146 Hansen EN, Manzano G, Kandemir U, Mok JM. Comparison of tissue oxygenation and
6 compartment pressure following tibia fracture. *Injury*. 2013; 44(8):1076-1080
- 7 147 Harley BJ, Beaupre LA, Jones CA, Dulai SK, Weber DW. The effect of time to definitive treatment
8 on the rate of nonunion and infection in open fractures. *Journal of Orthopaedic Trauma*. 2002;
9 16(7):484-490
- 10 148 Harley JD, Mack LA, Winquist RA. CT of acetabular fractures: Comparison with conventional
11 radiography. *American Journal of Roentgenology*. 1982; 138(3):413-417
- 12 149 Harris AM, Patterson BM, Sontich JK, Vallier HA. Results and outcomes after operative treatment
13 of high-energy tibial plafond fractures. *Foot and Ankle International*. 2006; 27(4):256-265
- 14 150 Harvey EJ, Agel J, Selznick HS, Chapman JR, Henley MB. Deleterious effect of smoking on healing
15 of open tibia-shaft fractures. *American Journal of Orthopedics*. 2002; 31(9):518-521
- 16 151 Harwood PJ, Giannoudis PV, Probst C, Krettek C, Pape HC. The risk of local infective complications
17 after damage control procedures for femoral shaft fracture. *Journal of Orthopaedic Trauma*.
18 2006; 20(3):181-189
- 19 152 Has B, Jovanovic S, Wertheimer B, Mikolasevic I, Grdic P. External fixation as a primary and
20 definitive treatment of open limb fractures. *Injury*. 1995; 26(4):245-248
- 21 153 Hatfield J, Arthur A, Phillips M, Howard Z, Shear M. Time savings by rapid EMS antibiotic therapy
22 for fractures. *Air Medical Journal*. 2012; 31(5):225
- 23 154 Hauser CJ, Adams CAJ, Eachempati SR, Council of the Surgical Infection Society. Surgical Infection
24 Society guideline: prophylactic antibiotic use in open fractures: an evidence-based guideline.
25 *Surgical Infections*. 2006; 7(4):379-405
- 26 155 Hee HT, Wong HP, Low YP, Myers L. Predictors of outcome of floating knee injuries in adults: 89
27 patients followed for 2-12 years. *Acta Orthopaedica Scandinavica*. 2001; 72(4):385-394
- 28 156 Heier KA, Infante AF, Walling AK, Sanders RW. Open fractures of the calcaneus: soft-tissue injury
29 determines outcome. *Journal of Bone and Joint Surgery - American Volume*. 2003; 85-
30 A(12):2276-2282
- 31 157 Helfet DL, Koval K, Pappas J, Sanders RW, DiPasquale T. Intraarticular "pilon" fracture of the tibia.
32 *Clinical Orthopaedics and Related Research*. 1994;(298):221-228
- 33 158 Helland P, Boe A, Molster AO, Solheim E, Hordvik M. Open tibial fractures treated with the Ex-fi-
34 re external fixation system. *Clinical Orthopaedics and Related Research*. 1996;(326):209-220
- 35 159 Henes FO, Nuchtern JV, Groth M, Habermann CR, Regier M, Rueger JM et al. Comparison of
36 diagnostic accuracy of Magnetic Resonance Imaging and Multidetector Computed Tomography
37 in the detection of pelvic fractures. *European Journal of Radiology*. 2012; 81(9):2337-2342

- 1 160 Henley MB, Chapman JR, Agel J, Harvey EJ, Whorton AM, Swiontkowski MF. Treatment of type II,
2 IIIA, and IIIB open fractures of the tibial shaft: a prospective comparison of unreamed
3 interlocking intramedullary nails and half-pin external fixators. *Journal of Orthopaedic Trauma*.
4 1998; 12(1):1-7
- 5 161 Hernigou J, Schuind F. Smoking as a predictor of negative outcome in diaphyseal fracture healing.
6 *International Orthopaedics*. 2013; 37(5):883-887
- 7 162 Herwig-Kempers A, Veraart BE. Reduction of posterior dislocation of the hip in the prone
8 position. *Journal of Bone and Joint Surgery British Volume*. 1993; 75(2):328
- 9 163 Hierner R, Betz AM, Comtet JJ, Berger AC. Decision making and results in subtotal and total lower
10 leg amputations: reconstruction versus amputation. *Microsurgery*. 1995; 16(12):830-839
- 11 164 Higgins TF, Klatt JB, Beals TC. Lower Extremity Assessment Project (LEAP)--the best available
12 evidence on limb-threatening lower extremity trauma. *Orthopedic Clinics of North America*.
13 2010; 41(2):233-239
- 14 165 Hillyard RF, Fox J. Sciatic nerve injuries associated with traumatic posterior hip dislocations.
15 *American Journal of Emergency Medicine*. 2003; 21(7):545-548
- 16 166 HM Revenue and Customs. Rates and thresholds for employers. 2014. Available from:
17 <https://www.gov.uk/rates-and-thresholds-for-employers-2014-to-2015> [Last accessed: 12 March
18 2015]
- 19 167 Hoffmann MF, Jones CB, Sietsema DL, Tornetta P, Koenig SJ. Clinical outcomes of locked plating
20 of distal femoral fractures in a retrospective cohort. *Journal of Orthopaedic Surgery and*
21 *Research*. 2013; 8:43
- 22 168 Hohmann E, Tetsworth K, Radziejowski MJ, Wiesniewski TF. Comparison of delayed and primary
23 wound closure in the treatment of open tibial fractures. *Archives of Orthopaedic and Trauma*
24 *Surgery*. 2007; 127(2):131-136
- 25 169 Holmes JF, Wisner DH. Indications and performance of pelvic radiography in patients with blunt
26 trauma. *American Journal of Emergency Medicine*. 2012; 30(7):1129-1133
- 27 170 Hommel A, Kock ML, Persson J, Werntoft E. The Patient's View of Nursing Care after Hip Fracture.
28 *ISRN Nursing*. 2012; 2012:863291
- 29 171 Hong SW, Seah CS, Kuek LB, Tan KC. Soft tissue cover in compound and complicated tibial
30 fractures using microvascular flaps. *Annals of the Academy of Medicine, Singapore*. 1998;
31 27(2):182-187
- 32 172 Hoogendoorn JM, van der Werken C. The mangled leg. Decision-making based on scoring
33 systems and outcome. *European Journal of Trauma*. 2002; 28(1):1-10
- 34 173 Horn C, Dobeles S, Vester H, Schaffler A, Lucke M, Stockle U. Combination of interfragmentary
35 screws and locking plates in distal meta-diaphyseal fractures of the tibia: a retrospective, single-
36 centre pilot study. *Injury*. 2011; 42(10):1031-1037
- 37 174 Horstman WG, McClennan BL, Heiken JP. Comparison of computed tomography and
38 conventional cystography for detection of traumatic bladder rupture. *Urologic Radiology*. 1991;
39 12(4):188-193

- 1 175 Hossieny P, Carey Smith R, Yates P, Carroll G. Efficacy of patient information concerning casts
2 applied post-fracture. ANZ Journal of Surgery. 2012; 82(3):151-155
- 3 176 Hou Z, Irgit K, Strohecker KA, Matzko ME, Wingert NC, DeSantis JG et al. Delayed flap
4 reconstruction with vacuum-assisted closure management of the open IIIB tibial fracture. Journal
5 of Trauma. 2011; 71(6):1705-1708
- 6 177 Hougaard K, Thomsen PB. Traumatic posterior dislocation of the hip--prognostic factors
7 influencing the incidence of avascular necrosis of the femoral head. Archives of Orthopaedic and
8 Traumatic Surgery Archiv Fur Orthopadische Und Unfall-Chirurgie. 1986; 106(1):32-35
- 9 178 Howe HRJ, Poole GVJ, Hansen KJ, Clark T, Plonk GW, Koman LA et al. Salvage of lower extremities
10 following combined orthopedic and vascular trauma. A predictive salvage index. American
11 Surgeon. 1987; 53(4):205-208
- 12 179 Hu P, Zhang YZ. Surgical hemostatic options for damage control of pelvic fractures. Chinese
13 Medical Journal. 2013; 126(12):2384-2389
- 14 180 Hughes S. Prevention of infection in orthopaedic surgery. Prescribers' Journal. 1993; 33(5):191-
15 195
- 16 181 Hull P. The management of open tibial fractures. European Journal of Orthopaedic Surgery and
17 Traumatology. 2008; 18(6):441-447
- 18 182 Hull PD, Johnson SC FAU, Stephen DJ FAU, Kreder HJ FAU, Jenkinson RJ. Delayed debridement of
19 severe open fractures is associated with a higher rate of deep infection. Bone and Joint Journal.
20 2014; 96B(3):379-384
- 21 183 Hulsker CC, Kleinveld S, Zonnenberg CB, Hogervorst M, Bekerom MP. Evidence-based treatment
22 of open ankle fractures. Archives of Orthopaedic and Trauma Surgery. 2011;(3):1545-1553
- 23 184 Hutchinson AJP, Frampton AE, Bhattacharya R. Operative fixation for complex tibial fractures.
24 Annals of the Royal College of Surgeons of England. 2012; 94(1):34-38
- 25 185 Hutson JJJ, Dayicioglu D, Oeltjen JC, Panthaki ZJ, Armstrong MB. The treatment of gustilo grade
26 IIIB tibia fractures with application of antibiotic spacer, flap, and sequential distraction
27 osteogenesis. Annals of Plastic Surgery. 2010; 64(5):541-552
- 28 186 Ibrahim T, Riaz M, Hegazy A, Erwin PJ, Tleyjeh IM. Delayed surgical debridement in pediatric open
29 fractures: a systematic review and meta-analysis. Journal of Children's Orthopaedics. 2014;
30 8(2):135-141
- 31 187 Ikem IC, Oginni LM, Bamgboye EA. Open fractures of the lower limb in Nigeria. International
32 Orthopaedics. 2001; 25(6):386-388
- 33 188 Ikem IC, Oginni LM, Ogunlusi JD. Determinants of management outcome in open tibia fractures
34 in Ile-Ife. Nigerian Journal of Surgical Research. 2006; 8(1-2):81-85
- 35 189 Irajpour A, Kaji NS, Nazari F, Azizkhani R, Zadeh AH. A comparison between the effects of simple
36 and traction splints on pain intensity in patients with femur fractures. Iranian Journal of Nursing
37 and Midwifery Research. 2012; 17(7):530-533
- 38 190 Jacob E, Erpelding JM, Murphy KP. A retrospective analysis of open fractures sustained by U.S.
39 military personnel during Operation Just Cause. Military Medicine. 1992; 157(10):552-556

- 1 191 Jacob JR, Rao JP, Ciccarelli C. Traumatic dislocation and fracture dislocation of the hip. A long-
2 term follow-up study. *Clinical Orthopaedics and Related Research*. 1987;(214):249-263
- 3 192 Jariwala AC, Kandasamy, Abboud RJ, Wigderowitz CA. Patients and the Internet: a demographic
4 study of a cohort of orthopaedic out-patients. *Surgeon*. 2004; 2(2):103-106
- 5 193 Jenkinson RJ, Kiss A, Johnson S, Stephen DJG, Kreder HJ. Delayed wound closure increases deep-
6 infection rate associated with lower-grade open fractures: a propensity-matched cohort study.
7 *Journal of Bone and Joint Surgery - American Volume*. 2014; 96(5):380-386
- 8 194 Johnston-Walker E, Hardcastle J. Neurovascular assessment in the critically ill patient. *Nursing in
9 Critical Care*. 2011; 16(4):170-177
- 10 195 Jones BG, Duncan RDD. Open tibial fractures in children under 13 years of age--10 years
11 experience. *Injury*. 2003; 34(10):776-780
- 12 196 Joshi D, Singh D, Ansari J, Lal Y. Immediate open reduction and internal fixation in open ankle
13 fractures. *Journal of the American Podiatric Medical Association*. 2006; 96(2):120-124
- 14 197 Joveniaux P, Ohl X, Harisboure A, Berrichi A, Labatut L, Simon P et al. Distal tibia fractures:
15 management and complications of 101 cases. *International Orthopaedics*. 2010; 34(4):583-588
- 16 198 Kai H, Yokoyama K, Shindo M, Itoman M. Problems of various fixation methods for open tibia
17 fractures: experience in a Japanese level I trauma center. *American Journal of Orthopedics*. 1998;
18 27(9):631-636
- 19 199 Kailidou E, Pikoulis E, Katsiva V, Karavokyros IG, Athanassopoulou A, Papakostantinou I et al.
20 Contrast-enhanced spiral CT evaluation of blunt abdominal trauma. *JBR-BTR*. 2005; 88(2):61-65
- 21 200 Kakar S, Tornetta P. Segmental tibia fractures: a prospective evaluation. *Clinical Orthopaedics
22 and Related Research*. 2007; 460:196-201
- 23 201 Kalyani BS, Fisher BE, Roberts CS, Giannoudis PV. Compartment syndrome of the forearm: a
24 systematic review. *Journal of Hand Surgery - American Volume*. 2011; 36(3):535-543
- 25 202 Kamath JB, Shetty MS, Joshua TV, Kumar A, Harshvardhan, Naik DM. Soft tissue coverage in open
26 fractures of tibia. *Indian Journal of Orthopaedics*. 2012; 46(4):462-469
- 27 203 Kane NM, Francis IR, Ellis JH. The value of CT in the detection of bladder and posterior urethral
28 injuries. *AJR American Journal of Roentgenology*. 1989; 153(6):1243-1246
- 29 204 Kapukaya A, Subasi M, Arslan H. Management of comminuted closed tibial plafond fractures
30 using circular external fixators. *Acta Orthopaedica Belgica*. 2005; 71(5):582-589
- 31 205 Katsenis D, Dendrinou G, Kouris A, Savas N, Schoinochoritis N, Pogiatis K. Combination of fine
32 wire fixation and limited internal fixation for high-energy tibial plateau fractures: functional
33 results at minimum 5-year follow-up. *Journal of Orthopaedic Trauma*. 2009; 23(7):493-501
- 34 206 Kazakos K, Lyras DN, Verettas D, Tilkeridis K, Tryfonidis M. The use of autologous PRP gel as an
35 aid in the management of acute trauma wounds. *Injury*. 2009; 40(8):801-805
- 36 207 Keating JF, Blachut PA, O'Brien PJ, Meek RN, Broekhuysen H. Reamed nailing of open tibial
37 fractures: does the antibiotic bead pouch reduce the deep infection rate? *Journal of Orthopaedic
38 Trauma*. 1996; 10(5):298-303

- 1 208 Keel M, Trentz O. (ii) Acute management of pelvic ring fractures. *Current Orthopaedics*. 2005;
2 19(5):334-344
- 3 209 Keeling JJ, Gwinn DE, Tintle SM, Andersen RC, McGuigan FX. Short-term outcomes of severe open
4 wartime tibial fractures treated with ring external fixation. *Journal of Bone and Joint Surgery -*
5 *American Volume*. 2008; 90(12):2643-2651
- 6 210 Keen JS, Desai PP, Smith CS, Suk M. Efficacy of hydrosurgical debridement and nanocrystalline
7 silver dressings for infection prevention in type II and III open injuries. *International Wound*
8 *Journal*. 2012; 9(1):7-13
- 9 211 Kendig RJ. Operative treatment of fractures of the tibial plafond. A randomized, prospective
10 study. *Journal of Bone and Joint Surgery - American Volume*. 1997; 79(12):1893-1894
- 11 212 Kesemenli CC, Kapukaya A, Subasi M, Arslan H, Necmioglu S, Kayikci C. Early prophylactic
12 autogenous bone grafting in type III open tibial fractures. *Acta Orthopaedica Belgica*. 2004;
13 70(4):327-331
- 14 213 Ketz J, Sanders R. Results of staged posterior fixation in the treatment of high-energy tibial pilon
15 fractures. *Fuss Und Sprunggelenk*. 2012; 10(1):27-36
- 16 214 Ketz J, Sanders R. Staged posterior tibial plating for the treatment of Orthopaedic Trauma
17 Association 43C2 and 43C3 tibial pilon fractures. *Journal of Orthopaedic Trauma*. 2012;
18 26(6):341-347
- 19 215 Khatod M, Botte MJ, Hoyt DB, Meyer RS, Smith JM, Akeson WH. Outcomes in open tibia
20 fractures: relationship between delay in treatment and infection. *Journal of Trauma*. 2003;
21 55(5):949-954
- 22 216 Kim JW, Oh CW, Jung WJ, Kim JS. Minimally invasive plate osteosynthesis for open fractures of
23 the proximal tibia. *Clinics in Orthopedic Surgery*. 2012; 4(4):313-320
- 24 217 Kindsfater K, Jonassen EA. Osteomyelitis in grade II and III open tibia fractures with late
25 debridement. *Journal of Orthopaedic Trauma*. 1995; 9(2):121-127
- 26 218 Kinzel V, Skirving AP, Wren MN, Zellweger R. Sideswipe injuries to the elbow in Western
27 Australia. *Medical Journal of Australia*. 2006; 184(9):447-450
- 28 219 Kirby MW, Spritzer C. Radiographic detection of hip and pelvic fractures in the emergency
29 department. *AJR American Journal of Roentgenology*. 2010; 194(4):1054-1060
- 30 220 Korkmaz A, Ciftdemir M, Ozcan M, Copuroglu C, Saridogan K. The analysis of the variables,
31 affecting outcome in surgically treated tibia pilon fractured patients. *Injury*. 2013; 44(10):1270-
32 1274
- 33 221 Koulouvaris P, Stafylas K, Mitsionis G, Vekris M, Mavrodontidis A, Xenakis T. Long-term results of
34 various therapy concepts in severe pilon fractures. *Archives of Orthopaedic and Trauma Surgery*.
35 2007; 127(5):313-320
- 36 222 Kreder HJ, Armstrong P. A review of open tibia fractures in children. *Journal of Pediatric*
37 *Orthopaedics*. 1995; 15(4):482-488

- 1 223 Krettek C, Kontopp H, Tscherne H, Seekamp A. Erratum: Hannover Fracture Scale '98 - Re-
2 evaluation and new perspectives of an established extremity salvage score (Injury 32; 4 (317-328)
3 (2001) PII: S0020138300002011). Injury. 2001; 32(7):611
- 4 224 Kulshrestha V. Incidence of infection after early intramedullary nailing of open tibial shaft
5 fractures stabilized with pinless external fixators. Indian Journal of Orthopaedics. 2008;
6 42(4):401-409
- 7 225 Kurylo JC, Axelrad TW, Tornetta P, Jawa A. Open fractures of the distal radius: the effects of
8 delayed debridement and immediate internal fixation on infection rates and the need for
9 secondary procedures. Journal of Hand Surgery - American Volume. 2011; 36(7):1131-1134
- 10 226 Lai CH, Kam CW. Bleeding pelvic fractures: Updates and controversies in acute phase
11 management. Hong Kong Journal of Emergency Medicine. 2008; 15(1):36-42
- 12 227 Lam K, Chan SWC, Lam SC. Level of psychological distress and social support among patients with
13 limb fractures in Hong Kong. Journal of Clinical Nursing. 2011; 20(5-6):784-793
- 14 228 Langford JR, Burgess AR, Liporace FA, Haidukewych GJ. Pelvic fractures: part 2. Contemporary
15 indications and techniques for definitive surgical management. Journal of the American Academy
16 of Orthopaedic Surgeons. 2013; 21(8):458-468
- 17 229 Laohapensang K, Prathnadi P, Lumley JSP. Intraluminal shunting for limb-threatening vascular
18 injuries. Asian Journal of Surgery. 1994; 17(4):371-376
- 19 230 Laughlin RT, Smith KL, Russell RC, Hayes JM. Late functional outcome in patients with tibia
20 fractures covered with free muscle flaps. Journal of Orthopaedic Trauma. 1993; 7(2):123-129
- 21 231 Lembo DE, Gorosito I, Nieto J, Venturelli E. Use of external splints in the treatment of exposed
22 fractures of the leg. Boletines y Trabajos De La Sociedad Argentina De Ortopedia y
23 Traumatologia. 1975; 40(3):305-306
- 24 232 Lenarz CJ, Watson JT, Moed BR, Israel H, Mullen JD, Macdonald JB. Timing of wound closure in
25 open fractures based on cultures obtained after debridement. Journal of Bone and Joint Surgery -
26 American Volume. 2010; 92(10):1921-1926
- 27 233 Leong CM, Leong AP, Low BY. Management of open tibia fractures. Singapore Medical Journal.
28 1988; 29(1):42-44
- 29 234 Leonidou A, Kiraly Z, Gality H, Apperley S, Vanstone S, Woods DA. The effect of the timing of
30 antibiotics and surgical treatment on infection rates in open long-bone fractures: a 6-year
31 prospective study after a change in policy. Strategies in Trauma and Limb Reconstruction. 2014;
32 9(3):167-171
- 33 235 Levy BA, Zlowodzki MP, Graves M, Cole PA. Screening for extremity arterial injury with the
34 arterial pressure index. American Journal of Emergency Medicine. 2005; 23(5):689-695
- 35 236 Li Y, Liu L, Tang X, Pei F, Wang G, Fang Y et al. Comparison of low, multidirectional locked nailing
36 and plating in the treatment of distal tibial metadiaphyseal fractures. International Orthopaedics.
37 2012; 36(7):1457-1462
- 38 237 Liu DSH, Sofiadellis F, Ashton M, MacGill K, Webb A. Early soft tissue coverage and negative
39 pressure wound therapy optimises patient outcomes in lower limb trauma. Injury. 2012;
40 43(6):772-778

- 1 238 Lord RS, Irani CN. Assessment of arterial injury in limb trauma. *Journal of Trauma*. 1974;
2 14(12):1042-1053
- 3 239 Lowenberg DW, Feibel RJ, Louie KW, Eshima I. Combined muscle flap and Ilizarov reconstruction
4 for bone and soft tissue defects. *Clinical Orthopaedics and Related Research*. 1996;(332):37-51
- 5 240 Luckhoff C, Mitra B, Cameron PA, Fitzgerald M, Royce P. The diagnosis of acute urethral trauma.
6 *Injury*. 2011; 42(9):913-916
- 7 241 Lustenberger T, Meier C, Benninger E, Lenzlinger PM, Keel MJB. C-clamp and pelvic packing for
8 control of hemorrhage in patients with pelvic ring disruption. *Journal of Emergencies, Trauma,*
9 *and Shock*. 2011; 4(4):477-482
- 10 242 Ly TV, Trivison TG, Castillo RC, Bosse MJ, MacKenzie EJ, LEAP Study Group. Ability of lower-
11 extremity injury severity scores to predict functional outcome after limb salvage. *Journal of Bone*
12 *and Joint Surgery American Volume*. 2008; 90(8):1738-1743
- 13 243 Mack AW, Freedman BA, Groth AT, Kirk KL, Keeling JJ, Andersen RC. Treatment of open proximal
14 femoral fractures sustained in combat. *Journal of Bone and Joint Surgery - American Volume*.
15 2013; 95(3):e13
- 16 244 MacKenzie EJ, Bosse MJ. Factors influencing outcome following limb-threatening lower limb
17 trauma: lessons learned from the Lower Extremity Assessment Project (LEAP). *Journal of the*
18 *American Academy of Orthopaedic Surgeons*. 2006; 14(10 Spec No.):S205-S210
- 19 245 MacLean LD. The diagnosis and treatment of arterial injuries of the upper extremity. *Surgical*
20 *Clinics of North America*. 1964; 44:1037-1047
- 21 246 Macleod M, Chesson RA, Blackledge P, Hutchison JD, Ruta N. To what extent are carers involved
22 in the care and rehabilitation of patients with hip fracture? *Disability and Rehabilitation*. 2005;
23 27(18-19):1117-1122
- 24 247 Magid D, Fishman EK, Brooker AFJ, Mandelbaum BR, Siegelman SS. Multiplanar computed
25 tomography of acetabular fractures. *Journal of Computer Assisted Tomography*. 1986; 10(5):778-
26 783
- 27 248 Malik MHA, Harwood P, Diggle P, Khan SA. Factors affecting rates of infection and nonunion in
28 intramedullary nailing. *Journal of Bone and Joint Surgery - British Volume*. 2004; 86(4):556-560
- 29 249 Malin Malmgren RN, Eva TÃfÃ¶rnvall RN, Inger Jansson RN. Patients with hip fracture:
30 Experiences of participation in care. *International Journal of Orthopaedic and Trauma Nursing*.
31 2014; 18(3):143-150
- 32 250 Mandracchia VJ, Evans RD, Nelson SC, Smith KM. Pilon fractures of the distal tibia. *Clinics in*
33 *Podiatric Medicine and Surgery*. 1999; 16(4):743-767
- 34 251 Marchetti ME, Steinberg GG, Coumas JM. Intermediate-term experience of Pipkin fracture-
35 dislocations of the hip. *Journal of Orthopaedic Trauma*. 1996; 10(7):455-461
- 36 252 Marks W, Dawid S, Lasek J, Witkowski Z, Golabek-Dropiewska K, Stasiak M. Posterior urethra
37 rupture: contrast-enhanced computed tomography scan and urethrocytography
38 demonstrations. *Case Reports in Urology*. 2012; 2012:109589

- 1 253 Marsh JL. External fixation is the treatment of choice for fractures of the tibial plafond. *Journal of*
2 *Orthopaedic Trauma*. 1999; 13(8):583-585
- 3 254 Marsh JL, Bonar S, Nepola JV, DeCoster TA, Hurwitz SR. Use of an articulated external fixator for
4 fractures of the tibial plafond. *Journal of Bone and Joint Surgery American Volume*. 1995;
5 77(10):1498-1509
- 6 255 Mauffrey C, Cuellar Iii DO, Pieracci F, Hak DJ, Hammerberg EM, Stahel PF et al. Strategies for the
7 management of haemorrhage following pelvic fractures and associated trauma-induced
8 coagulopathy. *Bone and Joint Journal*. 2014; 96-B(9):1143-1154
- 9 256 Mauffrey C, Vasario G, Battiston B, Lewis C, Beazley J, Seligson D. Tibial pilon fractures: a review
10 of incidence, diagnosis, treatment, and complications. *Acta Orthopaedica Belgica*. 2011;
11 77(4):432-440
- 12 257 Mayich DJ, Tieszer C, Lawendy A, McCormick W, Sanders D. Role of patient information handouts
13 following operative treatment of ankle fractures: a prospective randomized study. *Foot and*
14 *Ankle International*. 2013; 34(1):2-7
- 15 258 Mayne AI, Perry DC, Stables G, Dhotare S, Bruce CE. Documentation of neurovascular status in
16 supracondylar fractures and the development of an assessment proforma. *Emergency Medicine*
17 *Journal*. 2013; 30(6):480-482
- 18 259 McCaul JK, McCaul MG. Pre-hospital antibiotics for open fractures: Is there time? A descriptive
19 study. *African Journal of Emergency Medicine*. 2013; 3(4 Suppl.1):S20
- 20 260 McFerran MA, Smith SW, Boulas HJ, Schwartz HS. Complications encountered in the treatment of
21 pilon fractures. *Journal of Orthopaedic Trauma*. 1992; 6(2):195-200
- 22 261 McKee MD, Garay ME, Schemitsch EH, Kreder HJ, Stephen DJ. Irreducible fracture-dislocation of
23 the hip: a severe injury with a poor prognosis. *Journal of Orthopaedic Trauma*. 1998; 12(4):223-
24 229
- 25 262 McLain RF, Steyers C, Stoddard M. Infections in open fractures of the hand. *Journal of Hand*
26 *Surgery - American Volume*. 1991; 16(1):108-112
- 27 263 McQueen MM, Court-Brown. Compartment monitoring in tibial fractures. The pressure threshold
28 for decompression. *Journal of Bone and Joint Surgery - British Volume*. 1996; 78(1):99-104
- 29 264 Meadows LM, Mrkonjic L, Lagendyk L. Women's perceptions of future risk after low-energy
30 fractures at midlife. *Annals of Family Medicine*. 2005; 3(1):64-69
- 31 265 Mehlman CT, Hubbard GW, Crawford AH, Roy DR, Wall EJ. Traumatic hip dislocation in children.
32 Long-term followup of 42 patients. *Clinical Orthopaedics and Related Research*. 2000;(376):68-79
- 33 266 Mehta S, Routt MLC. Irreducible fracture-dislocations of the femoral head without posterior wall
34 acetabular fractures. *Journal of Orthopaedic Trauma*. 2008; 22(10):686-692
- 35 267 Meredith P, Gillham NR. NHS provision for the treatment of ankle fractures: a patient satisfaction
36 study. *Journal of the Royal Society of Medicine*. 1993; 86(6):332-335
- 37 268 Metsemakers WJ, Handojo K, Reynders P, Sermon A, Vanderschot P, Nijs S. Individual risk factors
38 for deep infection and compromised fracture healing after intramedullary nailing of tibial shaft
39 fractures: A single centre experience of 480 patients. *Injury*. 2015; 46(4):740-745

- 1 269 Min W, Ding BC, Tejwani NC. Staged versus acute definitive management of open distal humerus
2 fractures. *Journal of Trauma*. 2011; 71(4):944-947
- 3 270 Mittlmeier T, Machler G, Lob G, Mutschler W, Bauer G, Vogl T. Compartment syndrome of the
4 foot after intraarticular calcaneal fracture. *Clinical Orthopaedics and Related Research*.
5 1991;(269):241-248
- 6 271 Mlyncek M, Uharcsek P, Obert A. The management of a life-threatening pelvic hemorrhage in
7 obstetrics and gynecology. *Journal of Gynecologic Surgery*. 2005; 21(2):43-53
- 8 272 Moda SK, Kalra GS, Gupta RS, Maggu NK, Gupta RK, Kalra MK. The role of early flap coverage in
9 the management of open fractures of both bones of the leg. *Injury*. 1994; 25(2):83-85
- 10 273 Modin M, Ramos T, Stomberg MW. Postoperative impact of daily life after primary treatment of
11 proximal/distal tibiafracture with Ilizarov external fixation. *Journal of Clinical Nursing*. 2009;
12 18(24):3498-3506
- 13 274 Moed BR, Carr SE, Watson JT. Open reduction and internal fixation of posterior wall fractures of
14 the acetabulum. *Clinical Orthopaedics and Related Research*. 2000;(377):57-67
- 15 275 Moed BR, WillsonCarr SE, Watson JT. Results of operative treatment of fractures of the posterior
16 wall of the acetabulum. *Journal of Bone and Joint Surgery American Volume*. 2002; 84-A(5):752-
17 758
- 18 276 Moehring HD, Gravel C, Chapman MW, Olson SA. Comparison of antibiotic beads and
19 intravenous antibiotics in open fractures. *Clinical Orthopaedics and Related Research*.
20 2000;(372):254-261
- 21 277 Mokoena T, Naidu AG. Diagnostic difficulties in patients with a ruptured bladder. *British Journal*
22 *of Surgery*. 1995; 82(1):69-70
- 23 278 Morey AF, Iverson AJ, Swan A, Harmon WJ, Spore SS, Bhayani S et al. Bladder rupture after blunt
24 trauma: guidelines for diagnostic imaging. *Journal of Trauma*. 2001; 51(4):683-686
- 25 279 Morgan DE, Nallamala LK, Kenney PJ, Mayo MS, Rue LW. CT cystography: radiographic and
26 clinical predictors of bladder rupture. *AJR American Journal of Roentgenology*. 2000; 174(1):89-
27 95
- 28 280 Morsy Drch HA. Preliminary results and complications following limited open reduction and
29 percutaneous screw fixation of displaced fractures of the actabulum. *Injury*. 2001; 32(SUPPL.
30 1):45-50
- 31 281 Moues CM, Vos MC, van den Bemd GJ, Stijnen T, Hovius SE. Bacterial load in relation to vacuum-
32 assisted closure wound therapy: a prospective randomized trial. *Wound Repair and*
33 *Regeneration*. 2004; 12(1):11-17
- 34 282 Naique SB, Pearse M, Nanchahal J. Management of severe open tibial fractures: the need for
35 combined orthopaedic and plastic surgical treatment in specialist centres. *Journal of Bone and*
36 *Joint Surgery - British Volume*. 2006; 88(3):351-357
- 37 283 NHS EMPLOYERS. Terms and Conditions of Service NHS Medical and Dental Staff (England) 2002,
38 2002. Available from:
39 [http://www.nhsemployers.org/~media/Employers/Documents/Pay%20and%20reward/Terms_a](http://www.nhsemployers.org/~media/Employers/Documents/Pay%20and%20reward/Terms_and_Conditions_of_Service_NHS_Medical_and_Dental_Staff_300813_bt.pdf)
40 [nd_Conditions_of_Service_NHS_Medical_and_Dental_Staff_300813_bt.pdf](http://www.nhsemployers.org/~media/Employers/Documents/Pay%20and%20reward/Terms_and_Conditions_of_Service_NHS_Medical_and_Dental_Staff_300813_bt.pdf)

- 1 284 NHS EMPLOYERS. Terms and Conditions – Consultants (England) 2003. The terms and conditions
2 set out in this document shall incorporate, and be read, subject to any amendments which are
3 from time to time the subject of negotiation by the appropriate negotiation bodies and are
4 approved by the Secretary of State after considering the results of such negotiations. Any
5 amendments should be published., 2003. Available from:
6 http://www.nhsemployers.org/~media/Employers/Documents/Pay%20and%20reward/Consultant_Contract_V9_Revised_Terms_and_Conditions_300813_bt.pdf
7
- 8 285 NHS Staff Council. NHS terms and conditions of service handbook, 2015. Available from:
9 http://www.nhsemployers.org/~media/Employers/Documents/Pay%20and%20reward/AfC_tc_of_service_handbook_fb.pdf
10
- 11 286 Nichols JG, Svoboda JA, Parks SN. Use of temporary intraluminal shunts in selected peripheral
12 arterial injuries. *Journal of Trauma*. 1986; 26(12):1094-1096
- 13 287 Nielsen D, Huniche L, Brixen K, Sahota O, Masud T. Handling knowledge on osteoporosis--a
14 qualitative study. *Scandinavian Journal of Caring Sciences*. 2013; 27(3):516-524
- 15 288 Noumi T, Yokoyama K, Ohtsuka H, Nakamura K, Itoman M. Intramedullary nailing for open
16 fractures of the femoral shaft: evaluation of contributing factors on deep infection and nonunion
17 using multivariate analysis. *Injury*. 2005; 36(9):1085-1093
- 18 289 Nuchtern JV, Hartel MJ, Henes FO, Groth M, Jauch SY, Haegele J et al. Significance of clinical
19 examination, CT and MRI scan in the diagnosis of posterior pelvic ring fractures. *Injury*. 2015;
20 46(2):315-319
- 21 290 O'Brien L, Presnell S. Patient experience of distraction splinting for complex finger fracture
22 dislocations. *Journal of Hand Therapy*. 2010; 23(3):249-260
- 23 291 O'Flanagan SJ, Fulton G, O'Beirne J, McElwain JP. Operative fixation of unstable pelvic ring
24 injuries in polytrauma patients. *Irish Journal of Medical Science*. 1992; 161(2):39-41
- 25 292 O'Shea K, Quinlan JF, Waheed K, Brady OH. The usefulness of computed tomography following
26 open reduction and internal fixation of acetabular fractures. *Journal of Orthopaedic Surgery*.
27 2006; 14(2):127-132
- 28 293 O'Sullivan ST, O'Sullivan M, Pasha N, O'Shaughnessy M, O'Connor TP. Is it possible to predict limb
29 viability in complex Gustilo IIIB and IIIC tibial fractures? A comparison of two predictive indices.
30 *Injury*. 1997; 28(9-10):639-642
- 31 294 O'Toole GC, O'Grady P, Beddy P, McElwain JP. A critical appraisal of the out-patient fracture
32 clinic: is communication really the key? *Irish Medical Journal*. 2001; 94(2):41-42
- 33 295 Obaid AK, Barleben A, Porral D, Lush S, Cinat M. Utility of plain film pelvic radiographs in blunt
34 trauma patients in the emergency department. *American Surgeon*. 2006; 72(10):951-954
- 35 296 Office for National Statistics. Interim life tables 2010-2012. 2013. Available from:
36 <http://www.ons.gov.uk/ons/rel/lifetables/interim-life-tables/2010-2012/index.html> [Last
37 accessed: 24 October 2013]
- 38 297 Ogbemudia AO, Bafor A, Edomwonyi E, Enemudo R. Prevalence of pin tract infection: the role of
39 combined silver sulphadiazine and chlorhexidine dressing. *Nigerian Journal of Clinical Practice*.
40 2010; 13(3):268-271

- 1 298 Ogunlusi JD, Oginni LM, Ikem IC. Compartmental pressure in adults with tibial fracture.
2 International Orthopaedics. 2005; 29(2):130-133
- 3 299 Okcu G, Aktuglu K. Intra-articular fractures of the tibial plafond. A comparison of the results using
4 articulated and ring external fixators. Journal of Bone and Joint Surgery British Volume. 2004;
5 86(6):868-875
- 6 300 Okonta HI, Malemo KL, Ogunbanjo GA. The experience and psychosocial needs of patients with
7 traumatic fractures treated for more than six months at doctors on call for service hospital,
8 Goma, Democratic republic of Congo. South African Family Practice. 2011; 53(2):189-192
- 9 301 Olson B, Ustanko L. Self-care needs of patients in the halo brace. Orthopaedic Nursing. 1990;
10 9(1):27-52
- 11 302 Osborn PM, Smith WR, Moore EE, Cothren CC, Morgan SJ, Williams AE et al. Direct
12 retroperitoneal pelvic packing versus pelvic angiography: A comparison of two management
13 protocols for haemodynamically unstable pelvic fractures. Injury. 2009; 40(1):54-60
- 14 303 Ovadia DN, Beals RK. Fractures of the tibial plafond. Journal of Bone and Joint Surgery American
15 Volume. 1986; 68(4):543-551
- 16 304 Ovre S, Hvaal K, Holm I, Stromsoe K, Nordsletten L, Skjeldal S. Compartment pressure in nailed
17 tibial fractures. A threshold of 30 mmHg for decompression gives 29% fasciotomies. Archives of
18 Orthopaedic and Trauma Surgery. 1998; 118(1-2):29-31
- 19 305 Pao DM, Ellis JH, Cohan RH, Korobkin M. Utility of routine trauma CT in the detection of bladder
20 rupture. Academic Radiology. 2000; 7(5):317-324
- 21 306 Papadokostakis G, Kontakis G, Giannoudis P, Hadjipavlou A. External fixation devices in the
22 treatment of fractures of the tibial plafond: a systematic review of the literature. Journal of Bone
23 and Joint Surgery British Volume. 2008; 90(1):1-6
- 24 307 Papakostidis C, Kanakaris NK, Pretel J, Faour O, Morell DJ, Giannoudis PV. Prevalence of
25 complications of open tibial shaft fractures stratified as per the Gustilo-Anderson classification.
26 Injury. 2011; 42(12):1408-1415
- 27 308 Park HJ, Uchino M, Nakamura M, Ueno M, Kojima Y, Itoman M et al. Immediate interlocking
28 nailing versus external fixation followed by delayed interlocking nailing for Gustilo type IIIB open
29 tibial fractures. Journal of Orthopaedic Surgery. 2007; 15(2):131-136
- 30 309 Parrett BM, Matros E, Pribaz JJ, Orgill DP. Lower extremity trauma: trends in the management of
31 soft-tissue reconstruction of open tibia-fibula fractures. Plastic and Reconstructive Surgery. 2006;
32 117(4):1315-4
- 33 310 Patzakis MJ, Wilkins J. Factors influencing infection rate in open fracture wounds. Clinical
34 Orthopaedics and Related Research. 1989;(243):36-40
- 35 311 Patzakis MJ, Wilkins J, Moore TM. Use of antibiotics in open tibial fractures. Clinical Orthopaedics
36 and Related Research. 1983;(178):31-35
- 37 312 Paydar S, Ghaffarpasand F, Foroughi M, Saberi A, Dehghankhalili M, Abbasi H et al. Role of
38 routine pelvic radiography in initial evaluation of stable, high-energy, blunt trauma patients.
39 Emergency Medicine Journal. 2013; 30(9):724-727

- 1 313 Peng MY, Parisky YR, Cornwell EE, Radin R, Bragin S. CT cystography versus conventional
2 cystography in evaluation of bladder injury. *AJR American Journal of Roentgenology*. 1999;
3 173(5):1269-1272
- 4 314 Pierce ROJ, Heinrich JH. Comminuted intra-articular fractures of the distal tibia. *Journal of*
5 *Trauma*. 1979; 19(11):828-832
- 6 315 Pieroni S, Foster BR, Anderson SW, Kertesz JL, Rhea JT, Soto JA. Use of 64-row multidetector CT
7 angiography in blunt and penetrating trauma of the upper and lower extremities. *Radiographics*.
8 2009; 29(3):863-876
- 9 316 Pizanis A, Pohlemann T, Burkhardt M, Aghayev E, Holstein JH. Emergency stabilization of the
10 pelvic ring: Clinical comparison between three different techniques. *Injury*. 2013; 44(12):1760-
11 1764
- 12 317 Plaisier BR, Meldon SW, Super DM, Malangoni MA. Improved outcome after early fixation of
13 acetabular fractures. *Injury*. 2000; 31(2):81-84
- 14 318 Podeszwa DA, Mooney III JF, Cramer KE, Mendelow MJ. Comparison of Pavlik harness application
15 and immediate spica casting for femur fractures in infants. *Journal of Pediatric Orthopaedics*.
16 2004; 24(5):460-462
- 17 319 Pollak AN, McCarthy ML, Burgess AR. Short-term wound complications after application of flaps
18 for coverage of traumatic soft-tissue defects about the tibia. The Lower Extremity Assessment
19 Project (LEAP) Study Group. *Journal of Bone and Joint Surgery - American Volume*. 2000; 82-
20 A(12):1681-1691
- 21 320 Pollak AN, Jones AL, Castillo RC, Bosse MJ, MacKenzie EJ, LEAP Study Group. The relationship
22 between time to surgical debridement and incidence of infection after open high-energy lower
23 extremity trauma. *Journal of Bone and Joint Surgery - American Volume*. 2010; 92(1):7-15
- 24 321 Pollak AN, McCarthy ML, Bess RS, Agel J, Swiontkowski MF. Outcomes after treatment of high-
25 energy tibial plafond fractures. *Journal of Bone and Joint Surgery - American Volume*. 2003; 85-
26 A(10):1893-1900
- 27 322 Poole GV, Agnew SG, Griswold JA, Rhodes RS. The mangled lower extremity: can salvage be
28 predicted? *American Surgeon*. 1994; 60(1):50-55
- 29 323 Potter HG, Montgomery KD, Heise CW, Helfet DL. MR imaging of acetabular fractures: value in
30 detecting femoral head injury, intraarticular fragments, and sciatic nerve injury. *AJR American*
31 *Journal of Roentgenology*. 1994; 163(4):881-886
- 32 324 Ptak T, Rhea JT, Novelline RA. Experience with a continuous, single-pass whole-body
33 multidetector CT protocol for trauma: The three-minute multiple trauma CT scan. *Emergency*
34 *Radiology*. 2001; 8(5):250-256
- 35 325 Pugh KJ, Wolinsky PR, McAndrew MP, Johnson KD. Tibial pilon fractures: a comparison of
36 treatment methods. *Journal of Trauma*. 1999; 47(5):937-941
- 37 326 Puha B, Petreus T, Berea G, Sirbu PD, Puha G, Alexa O. Surgical approach in difficult tibial pilon
38 fractures. *Chirurgia*. 2014; 109(1):104-110
- 39 327 Purghele F, Badea MR, Ciuvica R, Jemna C. Surgical treatment in tibial plafond fractures-our
40 experience. *European Orthopaedics and Traumatology*. 2012; 3(4):239-242

- 1 328 Quagliano PV, Delair SM, Malhotra AK. Diagnosis of blunt bladder injury: A prospective
2 comparative study of computed tomography cystography and conventional retrograde
3 cystography. *Journal of Trauma*. 2006; 61(2):410-412
- 4 329 Radoicic D, Micic I, Dasic Z, Kosutic M. Does timing of surgery affect the outcome of open
5 articular distal humerus fractures. *European Journal of Orthopaedic Surgery and Traumatology*.
6 2014; 24(5):777-782
- 7 330 Rajasekaran S, Dheenadhayalan J, Babu JN, Sundararajan SR, Venkatramani H, Sabapathy SR.
8 Immediate primary skin closure in type-III A and B open fractures: results after a minimum of five
9 years. *Journal of Bone and Joint Surgery - British Volume*. 2009; 91(2):217-224
- 10 331 Rao P, Schaverien MV, Stewart KJ. Soft tissue management of children's open tibial fractures--a
11 review of seventy children over twenty years. *Annals of the Royal College of Surgeons of England*.
12 2010; 92(4):320-325
- 13 332 Rasool G, Ahmed MU, Iqbal M, Khwaja Z. Vacuum assisted wound closure and normal saline
14 dressing in treatment of Gustilo type II, type IIIa and IIIb open fracture of tibia. *Rawal Medical*
15 *Journal*. 2013; 38(4):382-384
- 16 333 Reber PU, Patel AG, Sapio NL, Ris HB, Beck M, Kniemeyer HW. Selective use of temporary
17 intravascular shunts in coincident vascular and orthopedic upper and lower limb trauma. *Journal*
18 *of Trauma*. 1999; 47(1):72-76
- 19 334 Redmond JM, Levy BA, Dajani KA, Cass JR, Cole PA. Detecting vascular injury in lower-extremity
20 orthopedic trauma: the role of CT angiography. *Orthopedics*. 2008; 31(8):761-767
- 21 335 Rehm CG, Mure AJ, O'Malley KF, Ross SE. Blunt traumatic bladder rupture: the role of retrograde
22 cystogram. *Annals of Emergency Medicine*. 1991; 20(8):845-847
- 23 336 Resnik CS, Stackhouse DJ, Shanmuganathan K, Young JW. Diagnosis of pelvic fractures in patients
24 with acute pelvic trauma: efficacy of plain radiographs. *AJR American Journal of Roentgenology*.
25 1992; 158(1):109-112
- 26 337 Reuss BL, Cole JD. Effect of delayed treatment on open tibial shaft fractures. *American Journal of*
27 *Orthopedics*. 2007; 36(4):215-220
- 28 338 Reynolds SL, Hernandez J, Hogg M, Runyon M. Do physicians identify clinically significant
29 fractures of the hip and pelvis in stable, alert patients after blunt trauma? *Academic Emergency*
30 *Medicine*. 2014; 21(5 Suppl.1):S241-S242
- 31 339 Ricci WM, Linn M, Gardner M, McAndrew C. What's new in orthopaedic trauma. *Journal of Bone*
32 *and Joint Surgery - American Volume*. 2014; 96(14):1222-1230
- 33 340 Richards JE, Magill M, Tressler MA, Shuler FD, Kregor PJ, Obrebsky WT et al. External fixation
34 versus ORIF for distal intra-articular tibia fractures. *Orthopedics*. 2012; 35(6):e862-e867
- 35 341 Richardson JD, Harty J, Amin M, Flint LM. Open pelvic fractures. *Journal of Trauma*. 1982;
36 22(7):533-538
- 37 342 Rinker B, Amspacher JC, Wilson PC, Vasconez HC. Subatmospheric pressure dressing as a bridge
38 to free tissue transfer in the treatment of open tibia fractures. *Plastic and Reconstructive*
39 *Surgery*. 2008; 121(5):1664-1673

- 1 343 Rinker B, Valerio IL, Stewart DH, Pu LLQ, Vasconez HC. Microvascular free flap reconstruction in
2 pediatric lower extremity trauma: a 10-year review. *Plastic and Reconstructive Surgery*. 2005;
3 115(6):1618-1624
- 4 344 Ristiniemi J, Luukinen P, Ohtonen P. Surgical treatment of extra-articular or simple intra-articular
5 distal tibial fractures: external fixation versus intramedullary nailing. *Journal of Orthopaedic
6 Trauma*. 2011; 25(2):101-105
- 7 345 Robertson DD, Sutherland CJ, Chan BW, Hodge JC, Scott WW, Fishman EK. Depiction of pelvic
8 fractures using 3D volumetric holography: comparison of plain X-ray and CT. *Journal of Computer
9 Assisted Tomography*. 1995; 19(6):967-974
- 10 346 Rojczyk M, Malottke R. [The effect of antibiotic prophylaxis in the treatment of open fractures].
11 *Hefte Zur Unfallheilkunde*. 1979; 138:355-357
- 12 347 Romano L, Pinto A, Niola R, Stavolo C, Cinque T, Daniele S et al. Bleeding due to pelvic fractures
13 in female patients: pictorial review of multidetector computed tomography imaging. *Current
14 Problems in Diagnostic Radiology*. 2012; 41(3):83-92
- 15 348 Rommens P, Broos P, Gruwez JA. Operative results in 124 open fractures of the tibial shaft. *Der
16 Unfallchirurg*. 1986; 89(3):127-131
- 17 349 Rose SC, Moore EE. Angiography in patients with arterial trauma: correlation between
18 angiographic abnormalities, operative findings, and clinical outcome. *American Journal of
19 Roentgenology*. 1987; 149(3):613-619
- 20 350 Rose SC, Moore EE. Trauma angiography of the extremity: the impact of injury mechanism on
21 triage decisions. *Cardiovascular and Interventional Radiology*. 1988; 11(3):136-139
- 22 351 Rosenthal RE, Coker WL. Posterior fracture-dislocation of the hip: an epidemiologic review.
23 *Journal of Trauma*. 1979; 19(8):572-581
- 24 352 Royle SG. The role of tissue pressure recording in forearm fractures in children. *Injury*. 1992;
25 23(8):549-552
- 26 353 Ruchholtz S, Waydhas C, Lewan U, Pehle B, Taeger G, Kuhne C et al. Free abdominal fluid on
27 ultrasound in unstable pelvic ring fracture: Is laparotomy always necessary? *Journal of Trauma -
28 Injury, Infection and Critical Care*. 2004; 57(2):278-286
- 29 354 Runkel N, Krug E, Berg L, Lee C, Hudson D, Birke-Sorensen H et al. Evidence-based
30 recommendations for the use of negative pressure wound therapy in traumatic wounds and
31 reconstructive surgery: Steps towards an international consensus. *Injury*. 2011; 42(Suppl.1):S1-
32 S12
- 33 355 Russell GG, Henderson R, Arnett G. Primary or delayed closure for open tibial fractures. *Journal
34 of Bone and Joint Surgery - British Volume*. 1990; 72(1):125-128
- 35 356 Sadri H, Nguyen-Tang T, Stern R, Hoffmeyer P, Peter R. Control of severe hemorrhage using C-
36 clamp and arterial embolization in hemodynamically unstable patients with pelvic ring
37 disruption. *Archives of Orthopaedic and Trauma Surgery*. 2005; 125(7):443-447
- 38 357 Sahin V, Karakas ES, Aksu S, Atlihan D, Turk CY, Halici M. Traumatic dislocation and fracture-
39 dislocation of the hip: a long-term follow-up study. *Journal of Trauma*. 2003; 54(3):520-529

- 1 358 Saikia KC, Bhattacharya TD, Agarwala V. Anterior compartment pressure measurement in closed
2 fractures of leg. *Indian Journal of Orthopaedics*. 2008; 42(2):217-221
- 3 359 Salmenkivi J, Lindahl J, Hirvensalo E, Takala A. Traditional internal fixation compared to external
4 hybrid fixation in operative treatment of pilon tibial fractures. *Suomen Ortopedia Ja*
5 *Traumatologia*. 1999; 22(3):219-224
- 6 360 Salton HL, Rush S, Schuberth J. Tibial plafond fractures: limited incision reduction with
7 percutaneous fixation. *Journal of Foot and Ankle Surgery*. 2007; 46(4):261-269
- 8 361 Saltzherr TP, Goslings JC. Effect on survival of whole-body CT during trauma resuscitation. *Lancet*.
9 2009; 374(9685):198-199
- 10 362 Sanders S, Tejwani N, Egol KA. Traumatic hip dislocation--a review. *Bulletin of the NYU Hospital*
11 *for Joint Diseases*. 2010; 68(2):91-96
- 12 363 Schemitsch EH, Bhandari M, Guyatt G, Sanders DW, Swiontkowski M, Tornetta P et al. Prognostic
13 factors for predicting outcomes after intramedullary nailing of the tibia. *Journal of Bone and Joint*
14 *Surgery - American Volume*. 2012; 94(19):1786-1793
- 15 364 Schenker ML, Yannascoli S, Baldwin KD, Ahn J, Mehta S. Does timing to operative debridement
16 affect infectious complications in open long-bone fractures? A systematic review. *Journal of Bone*
17 *and Joint Surgery - American Volume*. 2012; 94(12):1057-1064
- 18 365 Shanmuganathan R. The utility of scores in the decision to salvage or amputation in severely
19 injured limbs. *Indian Journal of Orthopaedics*. 2008; 42(4):368-376
- 20 366 Shepherd LE, Costigan WM, Gardocki RJ, Ghiassi AD, Patzakis MJ, Stevanovic MV. Local or free
21 muscle flaps and unreamed interlocked nails for open tibial fractures. *Clinical Orthopaedics and*
22 *Related Research*. 1998;(350):90-96
- 23 367 Short J, Upadhyay SS. Does simple traction and functional bracing affect the outcome of a
24 fractured femur as compared with the Thomas' splint method? *Physiotherapy*. 1984; 70(9):350-
25 354
- 26 368 Shyu YL, Chen M, Wu C, Cheng H. Family caregivers' needs predict functional recovery of older
27 care recipients after hip fracture. *Journal of Advanced Nursing*. 2010; 66(11):2450-2459
- 28 369 Sirkin M, Sanders R, DiPasquale T, Herscovici DJ. A staged protocol for soft tissue management in
29 the treatment of complex pilon fractures. *Journal of Orthopaedic Trauma*. 1999; 13(2):78-84
- 30 370 Skaggs DL, Kautz SM, Kay RM, Tolo VT. Effect of delay of surgical treatment on rate of infection in
31 open fractures in children. *Journal of Pediatric Orthopaedics*. 2000; 20(1):19-22
- 32 371 Skaggs DL, Friend L, Alman B, Chambers HG, Schmitz M, Leake B et al. The effect of surgical delay
33 on acute infection following 554 open fractures in children. *Journal of Bone and Joint Surgery -*
34 *American Volume*. 2005; 87(1):8-12
- 35 372 Slaney J, Christie N, Earthy S, Lyons RA, Kendrick D, Towner E. Improving recovery - Learning from
36 patients' experiences after injury: A qualitative study. *Injury*. 2014; 45(1):312-319
- 37 373 Solan MC, Calder JD, Gibbons CE, Ricketts DM. Photographic wound documentation after open
38 fracture. *Injury*. 2001; 32(1):33-35

- 1 374 Spencer Netto FAC, Hamilton P, Kodama R, Scarpelini S, Ortega SJ, Chu P et al. Retrograde
2 urethrocytography impairs computed tomography diagnosis of pelvic arterial hemorrhage in the
3 presence of a lower urologic tract injury. *Journal of the American College of Surgeons*. 2008;
4 206(2):322-327
- 5 375 Spencer J, Smith A, Woods D. The effect of time delay on infection in open long-bone fractures: a
6 5-year prospective audit from a district general hospital. *Annals of the Royal College of Surgeons*
7 of England. 2004; 86(2):108-112
- 8 376 Sriussadaporn S, Sirichindakul B, Pak-Art R, Tharavej C. Pelvic fractures: experience in
9 management of 170 cases at a university hospital in Thailand. *Journal of the Medical Association*
10 of Thailand. 2002; 85(2):200-206
- 11 377 Srour M, Inaba K, Okoye O, Chan C, Skiada D, Schnuriger B et al. Prospective evaluation of
12 treatment of open fractures: effect of time to irrigation and debridement. *JAMA Surgery*. 2015;
13 150(4):332-336
- 14 378 Stalekar H, Fuckar Z, Ekl D, Sustic A, Loncarek K, Ledic D. Primary vs secondary wound
15 reconstruction in Gustilo type III open tibial shaft fractures: follow-up study of 35 cases. *Croatian*
16 *Medical Journal*. 2003; 44(6):746-755
- 17 379 Stannard JP, Singanamala N, Volgas DA. Fix and flap in the era of vacuum suction devices: What
18 do we know in terms of evidence based medicine? *Injury*. 2010; 41(8):780-786
- 19 380 Stannard JP, Volgas DA, Stewart R, McGwin J, Alonso JE. Negative pressure wound therapy after
20 severe open fractures: A prospective randomized study. *Journal of Orthopaedic Trauma*. 2009;
21 23(8):552-557
- 22 381 Steiert AE, Gohritz A, Schreiber TC, Krettek C, Vogt PM. Delayed flap coverage of open extremity
23 fractures after previous vacuum-assisted closure (VAC) therapy - worse or worth? *Journal of*
24 *Plastic, Reconstructive and Aesthetic Surgery*. 2009; 62(5):675-683
- 25 382 Stengel D, Ottersbach C, Matthes G, Weigeldt M, Grundei S, Rademacher G et al. Accuracy of
26 single-pass whole-body computed tomography for detection of injuries in patients with major
27 blunt trauma. *CMAJ*. 2012; 184(8):869-876
- 28 383 Sturrock CA. A Method for the Reduction of Dislocations of the Hip. *BMJ*. 1899; 1(1997):845
- 29 384 Subramanian A, Vercruyse G, Dente C, Wyrzykowski A, King E, Feliciano DV. A decade's
30 experience with temporary intravascular shunts at a civilian level I trauma center. *Journal of*
31 *Trauma*. 2008; 65(2):316
- 32 385 Sungaran J, Harris I, Mourad M. The effect of time to theatre on infection rate for open tibia
33 fractures. *ANZ Journal of Surgery*. 2007; 77(10):886-888
- 34 386 Swanson TV, Szabo RM, Anderson DD. Open hand fractures: prognosis and classification. *Journal*
35 *of Hand Surgery - American Volume*. 1991; 16(1):101-107
- 36 387 Swiontkowski MF, MacKenzie EJ, Bosse MJ, Kellam JF, Burgess AR, Webb LX et al. Factors
37 influencing the decision to amputate or reconstruct after high-energy lower extremity trauma.
38 *Journal of Trauma*. 2002; 52(4):641-649
- 39 388 Taller J, Kamdar JP, Greene JA, Morgan RA, Blankenship CL, Dabrowski P et al. Temporary
40 vascular shunts as initial treatment of proximal extremity vascular injuries during combat

- 1 operations: the new standard of care at Echelon II facilities? *Journal of Trauma*. 2008; 65(3):595-
2 603
- 3 389 Tang J, Guo Wc, Yu L, Zhao Sh. Clinical efficacy of artificial skin combined with vacuum sealing
4 drainage in treating large-area skin defects. *Chinese Journal of Traumatology*. 2010; 13(5):289-
5 292
- 6 390 Their MEA, Bensch FV, Koskinen SK, Handolin L, Kiuru MJ. Diagnostic value of pelvic radiography
7 in the initial trauma series in blunt trauma. *European Radiology*. 2005; 15(8):1533-1537
- 8 391 Tho KS, Chiu PL, Krishnamoorthy S. Grade III open ankle fractures--a review of the outcome of
9 treatment. *Singapore Medical Journal*. 1994; 35(1):57-58
- 10 392 Thomas SH, Arthur AO, Howard Z, Shear ML, Kadzielski JL, Vrahas MS. Helicopter emergency
11 medical services crew administration of antibiotics for open fractures. *Air Medical Journal*. 2013;
12 32(2):74-79
- 13 393 Thomas TL, Meggitt BF. A comparative study of methods for treating fractures of the distal half of
14 the femur. *Journal of Bone and Joint Surgery - British Volume*. 1981; 63-B(1):3-6
- 15 394 Toni A, Gulino G, Baldini N, Gulino F. Clinical and radiographic long term results of acetabular
16 fractures associated with dislocations of the hip. *Italian Journal of Orthopaedics and*
17 *Traumatology*. 1985; 11(4):443-454
- 18 395 Torchia ME, Lewallen DG. Open fractures of the patella. *Journal of Orthopaedic Trauma*. 1996;
19 10(6):403-409
- 20 396 Toscan J, Mairs K, Hinton S, Stolee P, InfoRehab Research Team. Integrated transitional care:
21 patient, informal caregiver and health care provider perspectives on care transitions for older
22 persons with hip fracture. *International Journal of Integrated Care*. 2012; 12:e13
- 23 397 Townley WA, Nguyen DQA, Rooker JC, Dickson JK, Goroszeniuk DZ, Khan MS et al. Management
24 of open tibial fractures - a regional experience. *Annals of the Royal College of Surgeons of*
25 *England*. 2010; 92(8):693-696
- 26 398 Triffitt PD, Konig D, Harper WM, Barnes MR, Allen MJ, Gregg PJ. Compartment pressures after
27 closed tibial shaft fracture. Their relation to functional outcome. *Journal of Bone and Joint*
28 *Surgery - British Volume*. 1992; 74(2):195-198
- 29 399 Tripuraneni K, Ganga S, Quinn R, Gehlert R. The effect of time delay to surgical debridement of
30 open tibia shaft fractures on infection rate. *Orthopedics*. 2008; 31(12)
- 31 400 Trumble TE, Schmitt SR, Vedder NB. Internal fixation of pilon fractures of the distal radius. *Yale*
32 *Journal of Biology and Medicine*. 1993; 66(3):179-191
- 33 401 Uchida K, Kokubo Y, Yayama T, Nakajima H, Miyazaki T, Negoro K et al. Fracture of the pelvic ring:
34 A retrospective review of 224 patients treated at a single institution. *European Journal of*
35 *Orthopaedic Surgery and Traumatology*. 2011; 21(4):251-257
- 36 402 Udekwu PO, Gurkin B, Oller DW. The use of computed tomography in blunt abdominal injuries.
37 *American Surgeon*. 1996; 62(1):56-59
- 38 403 Upadhyay SS, Moulton A. The long-term results of traumatic posterior dislocation of the hip.
39 *Journal of Bone and Joint Surgery British Volume*. 1981; 63B(4):548-551

- 1 404 Uppal GS, Smith RC, Sherk HH, Mooar P. Accurate compartment pressure measurement using the
2 Intervenous Alarm Control (IVAC) Pump. Report of a technique. *Journal of Orthopaedic Trauma*.
3 1992; 6(1):87-89
- 4 405 Van Veen IHPA, Van Leeuwen AAM, Van PT, Van Luyt PA, Bode PJ, Van Vugt AB. Unstable pelvic
5 fractures: A retrospective analysis. *Injury*. 1995; 26(2):81-85
- 6 406 Van Vugt R, Kool DR, Deunk J, Edwards MJR. Effects on mortality, treatment, and time
7 management as a result of routine use of total body computed tomography in blunt high-energy
8 trauma patients. *Journal of Trauma and Acute Care Surgery*. 2012; 72(3):553-559
- 9 407 Vasiliadis ES, Grivas TB, Psarakis SA, Papavasileiou E, Kaspiris A, Triantafyllopoulos G. Advantages
10 of the Ilizarov external fixation in the management of intra-articular fractures of the distal tibia.
11 *Journal of Orthopaedic Surgery and Research*. 2009; 4:35
- 12 408 Verbeek D, Sugrue M, Balogh Z, Cass D, Civil I, Harris I et al. Acute management of
13 hemodynamically unstable pelvic trauma patients: time for a change? Multicenter review of
14 recent practice. *World Journal of Surgery*. 2008; 32(8):1874-1882
- 15 409 Vialle R, Odent T, Pannier S, Pauthier F, Laumonier F, Glorion C. Traumatic hip dislocation in
16 childhood. *Journal of Pediatric Orthopedics*. 2005; 25(2):138-144
- 17 410 Vigdorichik JM, Esquivel AO, Jin X, Yang KH, Onwudiwe NA, Vaidya R. Biomechanical stability of a
18 supra-acetabular pedicle screw internal fixation device (INFIX) vs external fixation and plates for
19 vertically unstable pelvic fractures. *Journal of Orthopaedic Surgery and Research*. 2012; 7:31
- 20 411 Vo NJ, Gash J, Browning J, Hutson RK. Pelvic imaging in the stable trauma patient: is the AP pelvic
21 radiograph necessary when abdominopelvic CT shows no acute injury? *Emergency Radiology*.
22 2004; 10(5):246-249
- 23 412 Waikakul S, Harnroongroj T, Vanadurongwan V. Immediate stabilization of unstable pelvic
24 fractures versus delayed stabilization. *Journal of the Medical Association of Thailand*. 1999;
25 82(7):637-642
- 26 413 Wang C, Li Y, Huang L, Wang M. Comparison of two-staged ORIF and limited internal fixation with
27 external fixator for closed tibial plafond fractures. *Archives of Orthopaedic and Trauma Surgery*.
28 2010; 130(10):1289-1297
- 29 414 Watson JT, Moed BR, Karges DE, Cramer KE. Pilon fractures. Treatment protocol based on
30 severity of soft tissue injury. *Clinical Orthopaedics and Related Research*. 2000;(375):78-90
- 31 415 Webb LX, Bosse MJ, Castillo RC, MacKenzie EJ, Kellam JE, Trivison TG et al. Analysis of surgeon-
32 controlled variables in the treatment of limb-threatening type-III open tibial diaphyseal fracture.
33 *Journal of Bone and Joint Surgery - American Volume*. 2007; 89(5):923-928
- 34 416 Wei Sj, Cai Xh, Wang Hs, Qi Bw, Yu Ax. A comparison of primary and delayed wound closure in
35 severe open tibial fractures initially treated with internal fixation and vacuum-assisted wound
36 coverage: a case-controlled study. *International Journal of Surgery*. 2014; 12(7):688-694
- 37 417 Whitney A, O'Toole RV, Hui E, Sciadini MF, Pollak AN, Manson TT et al. Do one-time
38 intracompartmental pressure measurements have a high false-positive rate in diagnosing
39 compartment syndrome? *Journal of Trauma and Acute Care Surgery*. 2014; 76(2):479-483

- 1 418 Widenfalk B, Ponten B, Karlstrom G. Open fractures of the shaft of the tibia: analysis of wound
2 and fracture treatment. *Injury*. 1979; 11(2):136-143
- 3 419 Willett KM, Pandit H, Upadhyay A. Internal versus external fixation for treating distal tibial pilon
4 fractures in adults. *Cochrane Database of Systematic Reviews*. 2008;(1)
- 5 420 Williams TM, Marsh JL, Nepola JV, DeCoster TA, Hurwitz SR, Bonar SB. External fixation of tibial
6 plafond fractures: is routine plating of the fibula necessary? *Journal of Orthopaedic Trauma*.
7 1998; 12(1):16-20
- 8 421 Williams TM, Nepola JV, Decoster TA, Hurwitz SR, Dirschl DR, Marsh JL. Factors affecting outcome
9 in tibial plafond fractures. *Clinical Orthopaedics and Related Research*. 2004;(423):93-98
- 10 422 Wood T, Sameem M, Avram R, Bhandari M, Petrisor B. A systematic review of early versus
11 delayed wound closure in patients with open fractures requiring flap coverage. *Journal of*
12 *Trauma and Acute Care Surgery*. 2012; 72(4):1078-1085
- 13 423 Wright BA, Roberts CS, Seligson D, Malkani AL, McCabe SJ. Cost of antibiotic beads is justified: a
14 study of open fracture wounds and chronic osteomyelitis. *Journal of Long-Term Effects of*
15 *Medical Implants*. 2007; 17(3):181-185
- 16 424 Wright E. Evaluating a paediatric neurovascular assessment tool. *Journal of Orthopaedic Nursing*.
17 2007; 11(1):20-29
- 18 425 Wyrsh B, McFerran MA, McAndrew M, Limbird TJ, Harper MC, Johnson KD et al. Operative
19 treatment of fractures of the tibial plafond. A randomized, prospective study. *Journal of Bone*
20 *and Joint Surgery - American Volume*. 1996; 78(11):1646-1657
- 21 426 Yang J, Gao Jm, Hu P, Li Ch, Zhao Sh, Lin X. Application of damage control orthopedics in 41
22 patients with severe multiple injuries. *Chinese Journal of Traumatology*. 2008; 11(3):157-160
- 23 427 Yang RS, Tsuang YH, Hang YS, Liu TK. Traumatic dislocation of the hip. *Clinical Orthopaedics and*
24 *Related Research*. 1991;(265):218-227
- 25 428 Yokoyama K, Itoman M, Shindo M, Kai H. Contributing factors influencing type III open tibial
26 fractures. *Journal of Trauma*. 1995; 38(5):788-793
- 27 429 Yokoyama K, Uchino M, Nakamura K, Ohtsuka H, Suzuki T, Boku T et al. Risk factors for deep
28 infection in secondary intramedullary nailing after external fixation for open tibial fractures.
29 *Injury*. 2006; 37(6):554-560
- 30 430 Yuenyongviwat V, Tangtrakulwanich B. Prevalence of pin-site infection: the comparison between
31 silver sulfadiazine and dry dressing among open tibial fracture patients. *Journal of the Medical*
32 *Association of Thailand*. 2011; 94(5):566-569
- 33 431 Yugueros P, Sarmiento JM, Garcia AF, Ferrada R. Unnecessary use of pelvic x-ray in blunt trauma.
34 *Journal of Trauma*. 1995; 39(4):722-725
- 35 432 Yusof NM, Khalid KA, Zulkifly AH, Zakaria Z, Amin MAM, Awang MS et al. Factors associated with
36 the outcome of open tibial fractures. *Malaysian Journal of Medical Sciences*. 2013; 20(5):47-53
- 37 433 Zaraca F, Ponzoni A, Stringari C, Ebner JA, Giovannetti R, Ebner H. Lower extremity traumatic
38 vascular injury at a level II trauma center: an analysis of limb loss risk factors and outcomes.
39 *Minerva Chirurgica*. 2011; 66(5):397-407

- 1 434 Zatti G, Bini A, Surace MF, Cherubino P. The surgical treatment of fractures of the proximal end
2 of the tibia: a review of cases as related to prognostic factors. *La Chirurgia Degli Organi Di*
3 *Movimento*. 2000; 85(4):371-380
- 4 435 Zeng Xt, Pang Gg, Ma Bt, Mei Xi, Sun X, Wang J et al. Surgical treatment of open pilon fractures.
5 *Orthopaedic Surgery*. 2011; 3(1):45-51
- 6 436 Zha GC, Sun JY, Dong SJ. Predictors of clinical outcomes after surgical treatment of displaced
7 acetabular fractures in the elderly. *Journal of Orthopaedic Research*. 2013; 31(4):588-595
- 8 437 Zhao XG. Emergency management of hemodynamically unstable pelvic fractures. *Chinese Journal*
9 *of Traumatology*. 2011; 14(6):363-366
- 10 438 Ziran BH, Darowish M, Klatt BA, Agudelo JF, Smith WR. Intramedullary nailing in open tibia
11 fractures: a comparison of two techniques. *International Orthopaedics*. 2004; 28(4):235-238
- 12 439 Ziran BH, Chamberlin E, Shuler FD, Shah M. Delays and difficulties in the diagnosis of lower
13 urologic injuries in the context of pelvic fractures. *Journal of Trauma*. 2005; 58(3):533-537
- 14 440 Zumsteg JW, Molina CS, Lee DH, Pappas ND. Factors influencing infection rates after open
15 fractures of the radius and/or ulna. *Journal of Hand Surgery - American Volume*. 2014; 39(5):956-
16 961
- 17
- 18
- 19
- 20