

# Chapter 25 Admission through elderly care assessment units

Emergency and acute medical care in over 16s: service delivery and organisation

*NICE guideline <number>*

*July 2017*

*Draft for consultation*

*Developed by the National Guideline Centre,  
hosted by the Royal College of Physicians*



**Disclaimer**

Healthcare professionals are expected to take NICE clinical guidelines fully into account when exercising their clinical judgement. However, the guidance does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of each patient, in consultation with the patient and, where appropriate, their guardian or carer.

**Copyright**

© National Institute for Health and Care Excellence, 2017. All rights reserved.

# Contents

<b>25 Admission through ECAU .....</b>	<b>5</b>
25.1 Introduction .....	5
25.2 Review question: Does admission or assessment through an elderly care assessment unit (ECAU) improve patient outcomes and hospital resource usage? .....	5
25.3 Clinical evidence.....	5
25.4 Economic evidence .....	10
25.5 Evidence statements.....	12
25.6 Recommendations and link to evidence.....	13
<b>Appendices.....</b>	<b>23</b>
Appendix A: Review protocol .....	23
Appendix B: Clinical article selection .....	24
Appendix C: Forest plots .....	25
Appendix D: Clinical Evidence tables.....	27
Appendix E: Economic evidence tables .....	33
Appendix F: GRADE tables .....	34
Appendix G: Excluded clinical studies .....	37
Appendix H: Excluded economic studies.....	40

# 1 25 Admission through Elderly Care Assessment Units

## 2 25.1 Introduction

3 Older patients are more likely to be admitted as an AME, and to stay longer in hospital. This is due to  
 4 a higher proportion of multi-morbidity, frailty, and polypharmacy than in younger patients. Hospital  
 5 services have adapted to the growing pressure from older patients, by introducing liaison services,  
 6 such as Frail Older Persons' Assessment and Liaison (FOPAL) services. These are now widespread,  
 7 and share characteristics such as medication review and the use of Comprehensive Geriatric  
 8 Assessment.

9 However, it is not clear whether there are additional benefits from admitting patients to a specialised  
 10 elderly care assessment unit (ECAU). Theoretical advantages could include better planning of  
 11 investigation and diagnosis, multiprofessional working, and dedicated discharge teams. The question  
 12 is important because of the potential for large reductions in length of stay, and quality of care.

## 13 25.2 Review question: Does admission or assessment through an elderly 14 care assessment unit (ECAU) improve patient outcomes and 15 hospital resource usage?

16 For full details see review protocol in Appendix A.

17 **Table 1: PICO characteristics of review question**

<b>Population</b>	Elderly people (65 years and over) with a suspected or confirmed AME.
<b>Intervention</b>	Assessment and management during admission (by GP referral, or via ED or community): <ul style="list-style-type: none"> <li>• through an elderly care/frailty Assessment Unit.</li> <li>• through an elderly care Assessment Area (defined area within the AMU).</li> <li>• by a visiting elderly care team (geriatrician team) in AMU.</li> </ul>
<b>Comparison</b>	Direct admission to generalist ward care from ED, community, or by GP referral (inpatient care only); direct admission to AMU without geriatric team involvement.
<b>Outcomes</b>	<ul style="list-style-type: none"> <li>• Quality of life (CRITICAL)</li> <li>• Length of stay (CRITICAL)</li> <li>• Mortality (CRITICAL)</li> <li>• Readmissions up to 30 days (IMPORTANT)</li> <li>• Avoidable adverse events (CRITICAL)</li> <li>• Delayed transfers of care (IMPORTANT)</li> <li>• A&amp;E 4 hour waiting target (IMPORTANT)</li> <li>• Patient and/or carer satisfaction (CRITICAL)</li> </ul>
<b>Study design</b>	Systematic reviews (SRs) of RCTs, RCTs, observational studies only to be included if no relevant SRs or RCTs are identified.

## 18 25.3 Clinical evidence

19 Four before-after studies were identified<sup>12,18,31,99</sup>, where assessment and management during  
 20 admission through an elderly care assessment unit, frailty unit, or by a geriatric team were compared  
 21 with either direct admission to a generalist ward or management through an AMU without geriatric  
 22 team involvement. Evidence from these studies is summarised in the clinical evidence summary  
 23 below (Table 3, Table 4 and Table 5). See also the study selection flow chart in Appendix B, study

1 evidence tables in Appendix D, forest plots in Appendix C, GRADE tables in Appendix F and excluded  
 2 studies list in Appendix G.

3 **Table 2: Summary of studies included in the review**

Study	Intervention and comparison	Population	Outcomes
Cardwell 2016 <sup>12</sup>  Before and after study  UK setting: single centre ED	'Front door' assessment of all over 65s with frailty – multidisciplinary team at the front desk in the ED with access to 8 care-of-the-elderly inpatient beds and 2 23-hour beds in the clinical decisions unit adjacent to the ED; team used a frailty index to screen between 9am-5pm Monday to Friday, those identified as frail entered the frail elderly pathway developed in the hospital.  Versus  Usual care - no screening for frailty, ED processed the admissions in the same way as for all adult age groups – directed to the Acute Medical Receiving Unit as clinically appropriate.	n=16,061 patients >65 presenting to ED.  Exclusion criteria: stroke, high level of care needed, on renal dialysis, obvious requirement for specialist care such as recent chemotherapy or a myocardial infarction.	Readmission (7-day and 28-day).
Conroy 2014 <sup>18</sup>  Before and after study  UK setting: teaching hospital	Emergency frailty unit - embedded comprehensive geriatric assessment service within the ED.  Versus  Usual care – emergency decisions unit, no routine input from specialists trained in geriatric medicine.	n=4647 patients ≥65 years attending the ED.	Re-admission.
Ellis 2012 <sup>31</sup>  Before and after study	Acute Care for Elders (ACE) Unit situated adjacent to the emergency department	n=422 patients attending the ED.  Inclusion criteria: >65 years	Length of stay.  Re-admission.

Study	Intervention and comparison	Population	Outcomes
<p>UK setting: district general hospital</p>	<p>and medical receiving unit, designed to deliver rapid assessment for patients deemed by non-specialists to require admission as a form of clinical decision unit.</p> <p>Versus</p> <p>Medical receiving unit – use of standardised screening and assessment tools, multidimensional assessment by a multidisciplinary team and proactive discharge planning.</p>	<p>with 1 or more of the following:</p> <ul style="list-style-type: none"> <li>- functional impairment (acute or chronic),</li> <li>- cognitive impairment (acute or chronic),</li> <li>- falls or other geriatric syndromes,</li> <li>- care home patients.</li> </ul> <p>Exclusion criteria: functionally independent patients or those with only single organ pathology requiring specialist input.</p>	<p>Mortality.</p>
<p>Taylor 2016<sup>99</sup> Before and after study UK Setting: Urban teaching hospital</p>	<p>Comprehensive older persons evaluation (COPE) zone within the emergency assessment unit, twice daily multidisciplinary team meetings, patients identified as potentially fit for discharge kept on COPE zone, otherwise transferred to geriatric medicine ward.</p> <p>Versus</p> <p>Admission to the emergency assessment unit after being referred from the ED or a GP, patients requiring geriatrician input seen by a daily in-reaching service.</p>	<p>n=811 medical patients &gt;75 years admitted to the emergency assessment unit.</p>	<p>Mortality (in-patient and 30-day).</p> <p>Re-admission.</p>

1

2

**Table 3: Clinical evidence summary: admission through ECAU versus direct admission**

Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with direct admission	Risk difference with ECAU (95% CI)
Readmission no. of patients readmitted	5096 (2 studies) 30 days	⊕⊖⊖⊖ VERY LOW <sup>a,b</sup> due to risk of bias, imprecision	RR 0.78 (0.67 to 0.92)	Moderate	
				143 per 1000	31 fewer per 1000 (from 11 fewer to 47 fewer)
Mortality no. of patients dying	422 (1 study) 12 months	⊕⊖⊖⊖ VERY LOW <sup>b,a</sup> due to imprecision	RR 0.86 (0.68 to 1.1)	Moderate	
				420 per 1000	59 fewer per 1000 (from 134 fewer to 42 more)
Length of stay mean length of stay	422 (1 study)	⊕⊖⊖⊖ VERY LOW <sup>a</sup> due to risk of bias			The mean length of stay in the intervention groups was 0.5 higher (3.29 lower to 4.29 higher)

(a) All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

(b) Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

**Table 4: Clinical evidence summary: admission through ECA area within AMU versus direct admission**

Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with direct admission	Risk difference with ECA area within AMU (95% CI)
In-patient mortality no. of patients dying in hospital	811 (1 study)	⊕⊖⊖⊖ VERY LOW <sup>b</sup> due to imprecision	RR 1.11 (0.71 to 1.75)	Moderate	
				80 per 1000	9 more per 1000 (from 23 fewer to 60 more)
30 day mortality no. of patients dying within 30 days of discharge	811 (1 study) 30 days	⊕⊖⊖⊖ VERY LOW <sup>b</sup> due to imprecision	RR 0.83 (0.46 to 1.51)	Moderate	
				55 per 1000	9 fewer per 1000 (from 30 fewer to 28 more)



Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with direct admission	Risk difference with ECA area within AMU (95% CI)
Readmission no. of patients readmitted	742 (1 study) 30 days	⊕⊖⊖⊖ VERY LOW <sup>a,b</sup> due to risk of bias, imprecision	RR 0.96 (0.71 to 1.3)	Moderate	
				189 per 1000	8 fewer per 1000 (from 55 fewer to 57 more)

(a) All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

(b) Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

**Table 5: Clinical evidence summary: admission by a visiting elderly care team versus direct admission**

Outcomes	No of Participants (studies) Follow up	Quality of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with direct admission	Risk difference with ECA area within AMU (95% CI)
Readmission no. of patients readmitted to hospital	9293 (1 study) 28 days	⊕⊖⊖⊖ VERY LOW <sup>a</sup> due to risk of bias	RR 0.67 (0.61 to 0.74)	Moderate	
				195 per 1000	64 fewer per 1000 (from 51 fewer to 76 fewer)
Readmission no. of patients readmitted to hospital	9293 (1 study) 7 days	⊕⊖⊖⊖ VERY LOW <sup>a</sup> due to risk of bias	RR 0.33 (0.27 to 0.40)	Moderate	
				88 per 1000	59 fewer per 1000 (from 53 fewer to 64 fewer)

(a) All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

## 1 **25.4 Economic evidence**

### 2 **Published literature**

3 One health economic study was identified with the relevant comparison and has been included in  
4 this review<sup>13</sup>. This is described in the health economic evidence profile below (Table 6) and the health  
5 economic evidence table in Appendix F.

6 The economic article selection protocol and flow chart for the whole guideline can found in the  
7 guideline's Appendix 41A and Appendix 41B.

**Table 6: Health economic evidence profile: Elderly care assessment unit versus usual care**

Study	Applicability	Limitations	Other comments	Incremental cost	Incremental effects	Cost-effectiveness	Uncertainty
Cardwell 2016 <sup>13</sup> (Scotland)	Partially applicable <sup>(a)</sup>	Potentially serious limitations <sup>(b)</sup>	Retrospective cohort study <b>Intervention 1:</b> Frail older people's pathway (FOPP) - Frailty MDT team 9am-5pm. Those assessed to be frail in the ED were put on the frail person's pathway. <b>Intervention 2:</b> No FOPP.	-£287	NA	NA	NR

Abbreviations: NA not applicable; NR not reported.

(a) Only cost comparison – only indicators of health were process outcomes like re-attendance and re-admission. Usual care was not described.

(b) The study was observational study, with no control for case-mix or time trend. No statistical or sensitivity analysis was undertaken. Only hospital costs included.

## 1 25.5 Evidence statements

### 2 Clinical

3 Four studies comprising 21,941 people evaluated the role of admission or assessment through an  
4 ECAU, frailty unit or by a geriatric team compared with either direct admission to a generalist ward  
5 or management through an AMU without geriatric team involvement for improving outcomes in  
6 secondary care in elderly people (65 years and over) with AMEs.

7 The evidence suggested that admission through ECAUs provides a benefit in reduction of  
8 readmissions (2 studies, very low quality) and mortality (1 study, very low quality). However, the  
9 evidence suggested there was no effect on length of stay (1 study, very low quality).

10 One study comprising 811 people evaluated the role of admission through an ECA area within the  
11 AMU compared to direct admission. The evidence suggested there was no difference in readmission,  
12 in-patient mortality or 30 day mortality (very low quality).

13 One study comprising 9293 people evaluated for assessment and management during admission by  
14 an elderly care team compared to direct admission. The evidence suggested a benefit in reduction of  
15 the number of readmissions at 7 days and 28 days (1 study, very low quality).

### 16 Economic

17 One cost comparison showed that an elderly care assessment unit was cost saving compared with  
18 usual care (cost difference: £287 per patient). This study was assessed to be partially applicable with  
19 potentially serious limitations.  
20

## 1 25.6 Recommendations and link to evidence

<b>Recommendations</b>	-
<b>Research recommendations</b>	<b>RR13. What is the most clinically and cost-effective way to configure services to assess frail older people who present to hospital with a medical emergency?</b>
Relative values of different outcomes	<p>The guideline committee considered 5 outcomes were critical for inclusion in this review: mortality, patient and/or carer satisfaction, quality of life, avoidable adverse events and length of hospital stay.</p> <p>Number of readmissions within 30 days, delayed transfers of care and compliance with the A&amp;E 4 hour waiting target were all considered to be important outcomes.</p>
Trade-off between benefits and harms	<p>Four studies comprising 21,941 people evaluated the role of admission or assessment through an elderly care or frailty assessment unit (ECAU), an elderly care assessment area within the AMU or by an elderly care team, compared with either direct admission to a general medical ward or management through an AMU without elderly care team involvement, for improving outcomes in secondary care in older people (65 years and over) with an acute medical emergency.</p> <p>The evidence suggested that admission through ECAUs provides a benefit in reduction of readmissions and mortality. However, the evidence suggested there was no effect on length of stay. No evidence was identified for the outcomes of patient and/or carer satisfaction, quality of life, avoidable adverse events and delayed transfers of care or compliance with the A&amp;E 4 hour waiting target.</p> <p>One study evaluated the role of admission through an ECA area within the AMU compared to direct admission. The evidence suggested there was no difference in readmission, in-patient mortality or 30 day mortality.</p> <p>The evidence suggested there was no effect on readmission. No evidence was identified for the outcomes patient and/or carer satisfaction, quality of life, length of stay, avoidable adverse events, delayed transfers of care or compliance with the ED 4-hour emergency access target.</p> <p>For assessment and management during admission by a multidisciplinary frail elderly team, evidence suggested a benefit in reduction of the number of readmissions at 7 and 28 days. No evidence was identified for mortality, patient and/or carer satisfaction, quality of life, avoidable adverse events, length of stay, delayed transfers of care or compliance with the ED 4-hour access target.</p> <p>It was agreed that the evidence was not strong enough to make a recommendation and the committee therefore opted to make a research recommendation.</p> <p>The committee noted a research recommendation would be particularly beneficial given that nationally, the development of older person care units/acute frailty units are being encouraged alongside acute medical assessment units<sup>75,93,95</sup>.</p> <p>Further research should consider whether the provision of these units in parallel to an acute medical unit (AMU) is beneficial, whether both services can be combined into 1 unit or whether the presence of a multidisciplinary frail elderly team reviewing identified patients on the AMU is sufficient.</p>
Trade-off between net effects and costs	<p>One of the before and after studies referred to above, which evaluated assessment and management during admission by a multidisciplinary frail elderly team, had estimated the cost impact. The cost of the staff per year (£300,000) was more than offset by cost savings from reduced length of stay, avoided admissions and reduced readmissions (£4.9 million). The net savings amounted to £287 per patient assessed. As there was only a single study, the comparator was not clearly described and the design was subject to a high risk of bias, the committee decided that a research recommendation was needed to provide more evidence on ECAUs before a practice</p>

	recommendation could be made.
Quality of evidence	<p>The evidence was graded very low quality for all outcomes due to risk of bias and imprecision.</p> <p>Nationally, patients who are admitted to hospitals with an ECAU often come directly from the community or from the community via the emergency department rather than via AMU. In many of the papers identified, the patients were admitted from another hospital ward in order to undergo discharge planning and therefore these papers were excluded as this was not considered relevant to the review question.</p> <p>The committee noted that these studies were heterogeneous models of care and their study design meant that case mix was not taken into consideration. The committee also noted the limitation of before and after study designs in this context, as the NHS evolves rapidly and outcomes were likely to be affected by a whole-system change rather than just the interventions themselves. One study was limited by a small population of included patients (less than 500 cases).</p> <p>The economic evaluation was only partially applicable because it did not evaluate health outcomes. It had potentially serious limitations because it was based on an observational before and after study, with no control for case-mix or time trend. Furthermore, no statistical or sensitivity analysis was undertaken and only hospital costs included.</p>
Other considerations	<p>ECAUs are diverse in structure, process and staff composition, and are often focused on discharge planning and rehabilitation with a prime aim of maintaining patients in their own environment. The committee noted that ECAU services are being developed and implemented, but they have not been well-evaluated. Research should concentrate on providing evidence for the optimal structure of care within the boundaries of funding available within the NHS. Research should also focus on the cost to the whole health economy.</p> <p>The key question is what is the optimal configuration for care for the frail elderly? The focus must be on the delivery of care required and important patient outcomes in the contexts of the financial constraints to the NHS. It may be more than 1 type of configuration is required and that depends on the local demographics and current infrastructure. As the number of frail elderly patients is only going to increase, identifying this is crucial hence the reason for the research recommendation.</p> <p>The committee noted that there are 2 NICE guidelines which have recommendations on Comprehensive Geriatric Assessment (CGA): a multi-disciplinary process which can be conducted during admission but which focuses on discharge planning and long-term follow-up. The recommendations are as follows:</p> <ul style="list-style-type: none"> <li>• The guideline for transition between inpatient hospital settings and community or care home settings for adults with social care needs (NG27)<sup>69</sup> recommends ‘start a comprehensive assessment of older people with complex needs at the point of admission and preferably in a specialist unit for older people’.</li> <li>• The guideline on Multimorbidity: clinical assessment and management (NG56)<sup>70</sup> refers to the recommendation above from NICE guideline SC712.</li> </ul>

1  
2

## References

- 1 Allen S, Bartlett T, Ventham J, McCubbin C, Williams A. Benefits of an older persons' assessment and liaison team in acute admissions areas of a general hospital. *Pragmatic and Observational Research*. 2010; 1:1-6
- 2 Applegate W, Deyo R, Kramer A, Meehan S. Geriatric evaluation and management: current status and future research directions. *Journal of the American Geriatrics Society*. 1991; 39(9 Pt 2):2S-7S
- 3 Applegate WB, Miller ST, Graney MJ, Elam JT, Burns R, Akins DE. A randomized, controlled trial of a geriatric assessment unit in a community rehabilitation hospital. *New England Journal of Medicine*. 1990; 322(22):1572-1578
- 4 Basic D, Conforti DA. A prospective, randomised controlled trial of an aged care nurse intervention within the Emergency Department. *Australian Health Review*. 2005; 29(1):51-59
- 5 Becker PM, McVey LJ, Saltz CC, Feussner JR, Cohen HJ. Hospital-acquired complications in a randomized controlled clinical trial of a geriatric consultation team. *JAMA - Journal of the American Medical Association*. 1987; 257(17):2313-2317
- 6 Bloch F, Kiffel C, Guilmineau F, Bellamy V, Brunetti N, Patry C et al. Impact of the intervention of a Mobile Geriatric Assessment Team on the diagnosis of significant comorbidities in elderly patients hospitalised after a hip fracture. *European Geriatric Medicine*. 2013; 4(5):310-313
- 7 Borenstein JE, Aronow HU, Bolton LB, Dimalanta MI, Chan E, Palmer K et al. Identification and team-based interprofessional management of hospitalized vulnerable older adults. *Nursing Outlook*. 2016; 64(2):137-145
- 8 Braude P, Goodman A, Elias T, Babic-Illman G, Challacombe B, Harari D et al. Evaluation and establishment of a ward-based geriatric liaison service for older urological surgical patients: Proactive care of Older People undergoing Surgery (POPS)-Urology. *BJU International*. 2016;
- 9 Burke K, Ringer P, St.John PD, McKenzie J-A. Introduction of a geriatric program assessment team in an inner city community and tertiary care teaching hospital. *Geriatrics Today: Journal of the Canadian Geriatrics Society*. 2001; 4(2):71-74
- 10 Campbell LJ, Cole KD. Geriatric assessment teams. *Clinics in Geriatric Medicine*. 1987; 3(1):99-110
- 11 Cape RD, Gibson SJ. The influence of clinical problems, age and social support on outcomes for elderly persons referred to regional aged care assessment teams. *Australian and New Zealand Journal of Medicine*. 1994; 24(4):378-385
- 12 Cardwell S, Clifton E, and Bond P. Older people in acute care: Data review and cost analysis: A pathway for frail older people in the emergency department. Healthcare Improvement Scotland, 2016. Available from: [http://www.healthcareimprovementscotland.org/our\\_work/person-centred\\_care/opac\\_improvement\\_programme/frailty\\_pathway\\_data\\_review.aspx](http://www.healthcareimprovementscotland.org/our_work/person-centred_care/opac_improvement_programme/frailty_pathway_data_review.aspx)
- 13 Cardwell S, Clifton E, and Bond P. Older people in acute care: data review and cost analysis: a pathway for frail older people in the emergency department. Healthcare Improvement Scotland, 2016. Available from: [http://www.healthcareimprovementscotland.org/our\\_work/person-centred\\_care/opac\\_improvement\\_programme/frailty\\_pathway\\_data\\_review.aspx](http://www.healthcareimprovementscotland.org/our_work/person-centred_care/opac_improvement_programme/frailty_pathway_data_review.aspx)

- 1 14 Cavalieri TA, Chopra A, Gray-Miceli D, Shreve S, Waxman H, Forman LJ. Geriatric assessment  
2 teams in nursing homes: do they work? *Journal of the American Osteopathic Association*. 1993;  
3 93(12):1269-1272
- 4 15 Cefalu CA, Colbourne G, Duffy M, Johnson E, Lestter M, Wright J. A university-affiliated  
5 community hospital inpatient geriatrics program functioning in an administrative and  
6 educational capacity. *Journal of the American Geriatrics Society*. 1997; 45(3):355-360
- 7 16 Cohen HJ, Feussner JR, Weinberger M, Carnes M, Hamdy RC, Hsieh F et al. A controlled trial of  
8 inpatient and outpatient geriatric evaluation and management. *New England Journal of*  
9 *Medicine*. 2002; 346(12):905-912
- 10 17 Collard AF, Bachman SS, Beatrice DF. Acute care delivery for the geriatric patient: an innovative  
11 approach. *QRB Quality Review Bulletin*. 1985; 11(6):180-185
- 12 18 Conroy SP, Ansari K, Williams M, Laithwaite E, Teasdale B, Dawson J et al. A controlled evaluation  
13 of comprehensive geriatric assessment in the emergency department: the 'Emergency Frailty  
14 Unit'. *Age and Ageing*. 2014; 43(1):109-114
- 15 19 Conroy SP, Stevens T, Parker SG, Gladman JRF. A systematic review of comprehensive geriatric  
16 assessment to improve outcomes for frail older people being rapidly discharged from acute  
17 hospital: 'interface geriatrics'. *Age and Ageing*. 2011; 40(4):436-443
- 18 20 Covinsky KE, Palmer RM, Kresevic DM, Kahana E, Counsell SR, Fortinsky RH et al. Improving  
19 functional outcomes in older patients: lessons from an acute care for elders unit. *Joint*  
20 *Commission Journal on Quality Improvement*. 1998; 24(2):63-76
- 21 21 DasGupta PK. Developing an active geriatric service in Scunthorpe. *Public Health*. 1980;  
22 94(3):155-160
- 23 22 Del Giudice E, Ferretti E, Omiciuolo C, Sceusa R, Zanata C, Manganaro D et al. The hospital-based,  
24 post-acute geriatric evaluation and management unit: the experience of the acute geriatric unit  
25 in Trieste. *Archives of Gerontology and Geriatrics*. 2009; 49 Suppl 1:49-60
- 26 23 Denewet N, De Breucker S, Luce S, Kennes B, Higuete S, Pepersack T. Comprehensive geriatric  
27 assessment and comorbidities predict survival in geriatric oncology. *Acta Clinica Belgica*. 2016;  
28 71(4):206-213
- 29 24 Edmans J, Bradshaw L, Gladman JRF, Franklin M, Berdunov V, Elliott R et al. The Identification of  
30 Seniors at Risk (ISAR) score to predict clinical outcomes and health service costs in older people  
31 discharged from UK acute medical units. *Age and Ageing*. 2013; 42(6):747-753
- 32 25 Edmans J, Bradshaw L, Franklin M, Gladman J, Conroy S. Specialist geriatric medical assessment  
33 for patients discharged from hospital acute assessment units: randomised controlled trial. *BMJ*.  
34 2013; 347:f5874
- 35 26 Edmans J, Conroy S, Harwood R, Lewis S, Elliott RA, Logan P et al. Acute medical unit  
36 comprehensive geriatric assessment intervention study (AMIGOS). *Trials*. 2011; 12:200
- 37 27 Ekdahl AW, Sjostrand F, Ehrenberg A, Oredsson S, Stavenow L, Wisten A et al. Frailty and  
38 comprehensive geriatric assessment organized as CGA-ward or CGA-consult for older adult  
39 patients in the acute care setting: A systematic review and meta-analysis. *European Geriatric*  
40 *Medicine*. 2015; 6(6):523-540



- 1 28 Ekdahl AW, Alwin J, Eckerblad J, Husberg M, Jaarsma T, Mazya AL et al. Long-Term Evaluation of  
2 the Ambulatory Geriatric Assessment: A Frailty Intervention Trial (AGe-FIT): Clinical Outcomes  
3 and Total Costs After 36 Months. *Journal of the American Medical Directors Association*. 2016;  
4 17(3):263-268
- 5 29 Ekdahl AW, Wirehn AB, Alwin J, Jaarsma T, Unosson M, Husberg M et al. Costs and Effects of an  
6 Ambulatory Geriatric Unit (the AGe-FIT Study): A Randomized Controlled Trial. *Journal of the  
7 American Medical Directors Association*. 2015; 16(6):497-503
- 8 30 Elliott RA, Martinac G, Campbell S, Thorn J, Woodward MC. Pharmacist-led medication review to  
9 identify medication-related problems in older people referred to an Aged Care Assessment  
10 Team: a randomized comparative study. *Drugs and Aging*. 2012; 29(7):593-605
- 11 31 Ellis G, Jamieson C-A, Alcorn M, Devlin V. An Acute Care for Elders (ACE) unit in the emergency  
12 department. *European Geriatric Medicine*. 2012; 3(4):261-263
- 13 32 Ellis G, Whitehead M, Robinson D, O'Neill D, Langhorne P. Comprehensive geriatric assessment  
14 for older adults admitted to hospital: A systematic review. *Cochrane Database of Systematic  
15 Reviews*. 2006;(4):no
- 16 33 Ellis G, Langhorne P. Comprehensive geriatric assessment for older hospital patients. *British  
17 Medical Bulletin*. 2004; 71:45-59
- 18 34 Ellis G, Marshall T, Ritchie C. Comprehensive geriatric assessment in the emergency department.  
19 *Clinical Interventions in Aging*. 2014; 9:2033-2043
- 20 35 Ellis G, Whitehead MA, O'Neill D, Langhorne P, Robinson D. Comprehensive geriatric assessment  
21 for older adults admitted to hospital. *Cochrane Database of Systematic Reviews*. 2011; Issue  
22 7:CD006211. DOI:10.1002/14651858.CD006211.pub2
- 23 36 Ellis G, Whitehead MA, Robinson D, O'Neill D, Langhorne P. Comprehensive geriatric assessment  
24 for older adults admitted to hospital: meta-analysis of randomised controlled trials. *BMJ*. 2011;  
25 343:d6553
- 26 37 Epstein AM, Hall JA, Fretwell M, Feldstein M, DeCiantis ML, Tognetti J et al. Consultative geriatric  
27 assessment for ambulatory patients. A randomized trial in a health maintenance organization.  
28 *JAMA - Journal of the American Medical Association*. 1990; 263(4):538-544
- 29 38 Faul AC, Yankeelov PA, Rowan NL, Gillette P, Nicholas LD, Borders KW et al. Impact of geriatric  
30 assessment and self-management support on community-dwelling older adults with chronic  
31 illnesses. *Journal of Gerontological Social Work*. 2009; 52(3):230-249
- 32 39 Fletcher AE, Jones DA, Bulpitt CJ, Tulloch AJ. The MRC trial of assessment and management of  
33 older people in the community: objectives, design and interventions [ISRCTN23494848]. *BMC  
34 Health Services Research*. 2002; 2(1):21
- 35 40 Fox J, Pattison T, Wallace J, Pradhan S, Gaillemine O, Feilding E et al. Geriatricians at the front  
36 door: The value of early comprehensive geriatric assessment in the emergency department.  
37 *European Geriatric Medicine*. 2016; 7(4):383-385
- 38 41 Fretwell MD, Cutler C, Epstein AM. Outpatient geriatric assessment in a Health Maintenance  
39 Organization. Its structure, practice, and clinical implications. *Clinics in Geriatric Medicine*. 1987;  
40 3(1):185-191

- 1 42 Fretwell MD, Raymond PM, McGarvey ST, Owens N, Traines M, Silliman RA et al. The Senior Care  
2 Study. A controlled trial of a consultative/unit-based geriatric assessment program in acute care.  
3 Journal of the American Geriatrics Society. 1990; 38(10):1073-1081
- 4 43 Germain M, Knoeffel F, Wieland D, Rubenstein LZ. A geriatric assessment and intervention team  
5 for hospital inpatients awaiting transfer to a geriatric unit: a randomized trial. Aging. 1995;  
6 7(1):55-60
- 7 44 Gerritsen JC, Van der Ende PC, Wolffensperger EW, Boom C. Evaluation of a geriatric assessment  
8 unit. International Journal of Geriatric Psychiatry. 1995; 10(3):207-217
- 9 45 Gladman J, Edmans J. Acute Medical Unit Comprehensive Geriatric Assessment Intervention  
10 Study (AMIGOS). UK Clinical Research Network. 2012;
- 11 46 Graf CE, Zekry D, Giannelli S, Michel JP, Chevalley T. Efficiency and applicability of comprehensive  
12 geriatric assessment in the emergency department: a systematic review. Aging Clinical and  
13 Experimental Research. 2011; 23(4):244-254
- 14 47 Gregersen M, Pedersen ABL, Damsgaard EM. Comprehensive geriatric assessment increases 30-  
15 day survival in the aged acute medical inpatients. Danish Medical Journal. 2012; 59(6):A4442
- 16 48 Harari D, Martin FC, Buttery A, O'Neill S, Hopper A. The older persons' assessment and liaison  
17 team 'OPAL': evaluation of comprehensive geriatric assessment in acute medical inpatients. Age  
18 and Ageing. 2007; 36(6):670-675
- 19 49 Harari D, Hopper A, Dhese J, Babic-Illman G, Lockwood L, Martin F. Proactive care of older people  
20 undergoing surgery ('POPS'): designing, embedding, evaluating and funding a comprehensive  
21 geriatric assessment service for older elective surgical patients. Age and Ageing. 2007; 36(2):190-  
22 196
- 23 50 Harris RD, Henschke PJ, Popplewell PY, Radford AJ, Bond MJ, Turnbull RJ et al. A randomised  
24 study of outcomes in a defined group of acutely ill elderly patients managed in a geriatric  
25 assessment unit or a general medical unit. Australian and New Zealand Journal of Medicine.  
26 1991; 21(2):230-234
- 27 51 Heath JM, Kobylarz FA, Brown M, Castano S. Interventions from home-based geriatric  
28 assessments of adult protective service clients suffering elder mistreatment. Journal of the  
29 American Geriatrics Society. 2005; 53(9):1538-1542
- 30 52 Hernandez-Vian O, Moreno-Ramos C, Sanchez-Garcia A, Lopez-Gomez MJ, Ortiz-Alvarez E,  
31 Balboa-Blanco E. [Evaluation of the care of the elderly program in frail elderly individuals with  
32 COPD in primary care centers in Sabadell (Spain)]. Enfermeria Clinica. 2007; 17(3):109-116
- 33 53 Hogan DB. Impact of geriatric consultation services for elderly patients admitted to acute care  
34 hospitals. Canadian Journal on Aging. 1990; 9(1):35-44
- 35 54 Hogan DB, Cape RD. A geriatric assessment unit in a long-term care facility. Canadian Journal of  
36 Public Health. 1984; 75(4):301-303
- 37 55 Horgan AM, Leighl NB, Coate L, Liu G, Palepu P, Knox JJ et al. Impact and feasibility of a  
38 comprehensive geriatric assessment in the oncology setting: a pilot study. American Journal of  
39 Clinical Oncology. 2012; 35(4):322-328

- 1 56 Humphries S. Diagnosis and disability in Geriatric Assessment Team clients: a pilot study.  
2 Australian Health Review. 1992; 15(1):22-34
- 3 57 Jones DM, Song X, Rockwood K. Operationalizing a frailty index from a standardized  
4 comprehensive geriatric assessment. Journal of the American Geriatrics Society. 2004;  
5 52(11):1929-1933
- 6 58 Kamel SJ, Jarrett P, MacDonald E. Effectiveness of geriatric evaluation and management units in  
7 caring for older adults. Geriatrics Today: Journal of the Canadian Geriatrics Society. 2005;  
8 8(3):104-109
- 9 59 Karppi P. Effects of a geriatric inpatient unit on elderly home care patients: a controlled trial.  
10 Aging. 1995; 7(3):207-211
- 11 60 Karppi P, Tilvis R. Effectiveness of a Finnish geriatric inpatient assessment. Two-year follow up of  
12 a randomized clinical trial on community-dwelling patients. Scandinavian Journal of Primary  
13 Health Care. 1995; 13(2):93-98
- 14 61 Kay G, MacTavish M, Moffatt C, Lau G. Development and evaluation of a geriatric assessment  
15 unit in a community hospital. Perspectives. 1992; 16(3):2-9
- 16 62 Kergoat MJ, Latour J, Lebel P, Leclerc BS, Leduc N, Beland F et al. Quality-of-care processes in  
17 geriatric assessment units: principles, practice, and outcomes. Journal of the American Medical  
18 Directors Association. 2012; 13(5):459-463
- 19 63 Kircher TTJ, Wormstall H, Muller PH, Schwarzler F, Buchkremer G, Wild K et al. A randomised trial  
20 of a geriatric evaluation and management consultation services in frail hospitalised patients. Age  
21 and Ageing. 2007; 36(1):36-42
- 22 64 Landefeld CS, Palmer RM, Kresevic DM, Fortinsky RH, Kowal J. A randomized trial of care in a  
23 hospital medical unit especially designed to improve the functional outcomes of acutely ill older  
24 patients. New England Journal of Medicine. 1995; 332(20):1338-1344
- 25 65 Landi F, Onder G, Tua E, Carrara B, Zuccalá G, Gambassi G et al. Impact of a new assessment  
26 system, the MDS-HC, on function and hospitalization of homebound older people: a controlled  
27 clinical trial. Journal of the American Geriatrics Society. 2001; 49(10):1288-1293
- 28 66 Lightbody E, Baldwin R. Inpatient geriatric evaluation and management did not reduce mortality  
29 but reduced functional decline. Evidence-Based Mental Health. 2002; 5(4):109
- 30 67 McDowell BJ, Silverman M, Martin D, Musa D, Keane C. Identification and intervention for  
31 urinary incontinence by community physicians and geriatric assessment teams. Journal of the  
32 American Geriatrics Society. 1994; 42(5):501-505
- 33 68 McVey LJ, Becker PM, Saltz CC, Feussner JR, Cohen HJ. Effect of a geriatric consultation team on  
34 functional status of elderly hospitalized patients. A randomized, controlled clinical trial. Annals of  
35 Internal Medicine. 1989; 110(1):79-84
- 36 69 National Institute for Health and Care Excellence. Transition between inpatient hospital settings  
37 and community or care home settings for adults with social care needs, 2015. Available from:  
38 [https://www.nice.org.uk/guidance/ng27/resources/transition-between-inpatient-hospital-  
settings-and-community-or-care-home-settings-for-adults-with-social-care-needs-  
1837336935877](https://www.nice.org.uk/guidance/ng27/resources/transition-between-inpatient-hospital-<br/>39 settings-and-community-or-care-home-settings-for-adults-with-social-care-needs-<br/>40 1837336935877)

- 1 70 National Institute for Health and Care Excellence. Multimorbidity: clinical assessment and  
2 management. NICE guideline 56. London. National Institute for Health and Care Excellence, 2016.  
3 Available from: <https://www.nice.org.uk/guidance/ng56>
- 4 71 National Institute for Health and Clinical Excellence. The guidelines manual. London: National  
5 Institute for Health and Clinical Excellence; 2012. Available from:  
6 <http://publications.nice.org.uk/the-guidelines-manual-pmg6/>
- 7 72 Naughton BJ, Moran MB, Feinglass J, Falconer J, Williams ME. Reducing hospital costs for the  
8 geriatric patient admitted from the emergency department: a randomized trial. *Journal of the*  
9 *American Geriatrics Society*. 1994; 42(10):1045-1049
- 10 73 Nikolaus T, Specht-Leible N, Bach M, Oster P, Schlierf G. A randomized trial of comprehensive  
11 geriatric assessment and home intervention in the care of hospitalized patients. *Age and Ageing*.  
12 1999; 28(6):543-550
- 13 74 Nipp R, Sloane R, Rao AV, Schmader KE, Cohen HJ. Role of pain medications, consultants, and  
14 other services in improved pain control of elderly adults with cancer in geriatric evaluation and  
15 management units. *Journal of the American Geriatrics Society*. 2012; 60(10):1912-1917
- 16 75 Oliver D, Foot C, and Humphries R. Making our health and care systems fit for an ageing  
17 population. Kings Fund, 2016. Available from:  
18 [https://www.kingsfund.org.uk/sites/files/kf/field/field\\_publication\\_file/making-health-care-](https://www.kingsfund.org.uk/sites/files/kf/field/field_publication_file/making-health-care-systems-fit-ageing-population-oliver-foot-humphries-mar14.pdf)  
19 [systems-fit-ageing-population-oliver-foot-humphries-mar14.pdf](https://www.kingsfund.org.uk/sites/files/kf/field/field_publication_file/making-health-care-systems-fit-ageing-population-oliver-foot-humphries-mar14.pdf)
- 20 76 Owen C, Tiwari D. Impact of early comprehensive geriatric assessment in the acute medical unit.  
21 *Clinical Medicine*. 2015; 15(Suppl 3):s13
- 22 77 Parker G, Bhakta P, Katbamna S, Lovett C, Paisley S, Parker S et al. Best place of care for older  
23 people after acute and during subacute illness: a systematic review. *Journal of Health Services*  
24 *Research and Policy*. 2000; 5(3):176-189
- 25 78 Phibbs CS, Holty JE, Goldstein MK, Garber AM, Wang Y, Feussner JR et al. The effect of geriatrics  
26 evaluation and management on nursing home use and health care costs: results from a  
27 randomized trial. *Medical Care*. 2006; 44(1):91-95
- 28 79 Pitner J. Specialty geriatric evaluation and management teams reduce adverse drug reactions.  
29 *Consultant Pharmacist*. 2004; 19(11):1042-1049
- 30 80 Popplewell PY, Henschke PJ. What is the value of a Geriatric Assessment Unit in a teaching  
31 hospital? A comparative study of the management of elderly inpatients. *Australian Health*  
32 *Review*. 1983; 6(2):23-25
- 33 81 Reuben DB, Borok GM, Wolde-Tsadiq G, Ershoff DH, Fishman LK, Ambrosini VL et al. A  
34 randomized trial of comprehensive geriatric assessment in the care of hospitalized patients. *New*  
35 *England Journal of Medicine*. 1995; 332(20):1345-1350
- 36 82 Riley CG. A geriatric assessment unit: the first twelve months. *New Zealand Medical Journal*.  
37 1974; 80(528):435-442
- 38 83 Rockwood K, Howlett S, Stadnyk K, Carver D, Powell C, Stolee P. Responsiveness of goal  
39 attainment scaling in a randomized controlled trial of comprehensive geriatric assessment.  
40 *Journal of Clinical Epidemiology*. 2003; 56(8):736-743

- 1 84 Rosenberg T. Acute hospital use, nursing home placement, and mortality in a frail community-  
2 dwelling cohort managed with Primary Integrated Interdisciplinary Elder Care at Home. *Journal*  
3 *of the American Geriatrics Society*. 2012; 60(7):1340-1346
- 4 85 Rubenstein LZ, Josephson K, Wieland GD, Pietruszka F, Tretton C, Strome S et al. Geriatric  
5 assessment on a subacute hospital ward. *Clinics in Geriatric Medicine*. 1987; 3(1):131-143
- 6 86 Rubenstein LZ, Josephson KR, Harker JO, Miller DK, Wieland D. The Sepulveda GEU Study  
7 revisited: long-term outcomes, use of services, and costs. *Aging*. 1995; 7(3):212-217
- 8 87 Rubenstein LZ, Josephson KR, Wieland GD, English PA, Sayre JA, Kane RL. Effectiveness of a  
9 geriatric evaluation unit. A randomized clinical trial. *New England Journal of Medicine*. 1984;  
10 311(26):1664-1670
- 11 88 Saltvedt I, Mo ES, Fayers P, Kaasa S, Sletvold O. Reduced mortality in treating acutely sick, frail  
12 older patients in a geriatric evaluation and management unit. A prospective randomized trial.  
13 *Journal of the American Geriatrics Society*. 2002; 50(5):792-798
- 14 89 Saltvedt I, Saltnes T, Mo ES, Fayers P, Kaasa S, Sletvold O. Acute geriatric intervention increases  
15 the number of patients able to live at home. A prospective randomized study. *Aging Clinical and*  
16 *Experimental Research*. 2004; 16(4):300-306
- 17 90 Saltvedt I, Jordhoy M, Opdahl Mo ES, Fayers P, Kaasa S, Sletvold O. Randomised trial of in-  
18 hospital geriatric intervention: impact on function and morale. *Gerontology*. 2006; 52(4):223-230
- 19 91 Saltvedt I, Spigset O, Ruths S, Fayers P, Kaasa S, Sletvold O. Patterns of drug prescription in a  
20 geriatric evaluation and management unit as compared with the general medical wards: a  
21 randomised study. *European Journal of Clinical Pharmacology*. 2005; 61(12):921-928
- 22 92 Saltz CC, McVey LJ, Becker PM, Feussner JR, Cohen HJ. Impact of a geriatric consultation team on  
23 discharge placement and repeat hospitalization. *Gerontologist*. 1988; 28(3):344-350
- 24 93 Sheffield Teaching Hospital NHS Trust. Improving the flow of older people. The Health  
25 Foundation: Inspiring Improvement, 2013. Available from: [https://www.england.nhs.uk/wp-](https://www.england.nhs.uk/wp-content/uploads/2013/08/sheff-study.pdf)  
26 [content/uploads/2013/08/sheff-study.pdf](https://www.england.nhs.uk/wp-content/uploads/2013/08/sheff-study.pdf)
- 27 94 Silverman M, Musa D, Martin DC, Lave JR, Adams J, Ricci EM. Evaluation of outpatient geriatric  
28 assessment: a randomized multi-site trial. *Journal of the American Geriatrics Society*. 1995;  
29 43(7):733-740
- 30 95 Silvester KM, Mohammed MA, Harriman P, Girolami A, Downes TW. Timely care for frail older  
31 people referred to hospital improves efficiency and reduces mortality without the need for extra  
32 resources. *Age and Ageing*. 2014; 43(4):472-477
- 33 96 Soejono CH. The impact of 'comprehensive geriatric assessment (CGA)' implementation on the  
34 effectiveness and cost (CEA) of healthcare in an acute geriatric ward. *Acta Medica Indonesiana*.  
35 2008; 40(1):3-10
- 36 97 Stewart M, Suchak N, Scheve A, Popat-Thakkar V, David E, Laquinte J et al. The impact of a  
37 geriatrics evaluation and management unit compared to standard care in a community teaching  
38 hospital. *Maryland Medical Journal*. 1999; 48(2):62-67

- 1 98 Stuck AE, Aronow HU, Steiner A, Alessi CA, BüLa CJ, Gold MN et al. A trial of annual in-home  
2 comprehensive geriatric assessments for elderly people living in the community. *New England*  
3 *Journal of Medicine*. 1995; 333(18):1184-1189
- 4 99 Taylor JK, Gaillemín OS, Pearl AJ, Murphy S, Fox J. Embedding comprehensive geriatric  
5 assessment in the emergency assessment unit: the impact of the COPE zone. *Clinical Medicine*.  
6 2016; 16(1):19-24
- 7 100 Teasdale TA, Shuman L, Snow E, Luchi RJ. A comparison of placement outcomes of geriatric  
8 cohorts receiving care in a geriatric assessment unit and on general medicine floors. *Journal of*  
9 *the American Geriatrics Society*. 1983; 31(9):529-534
- 10 101 Toseland RW, O'Donnell JC, Engelhardt JB, Hendler SA, Richie JT, Jue D. Outpatient geriatric  
11 evaluation and management. Results of a randomized trial. *Medical Care*. 1996; 34(6):624-640
- 12 102 Trentini M, Semeraro S, Motta M, Italian Study Group for Geriatric Assessment and  
13 Management. Effectiveness of geriatric evaluation and care. One-year results of a multicenter  
14 randomized clinical trial. *Aging*. 2001; 13(5):395-405
- 15 103 Van Craen K, Braes T, Wellens N, Denhaerynck K, Flamaing J, Moons P et al. The effectiveness of  
16 inpatient geriatric evaluation and management units: a systematic review and meta-analysis.  
17 *Journal of the American Geriatrics Society*. 2010; 58(1):83-92
- 18 104 White SJ, Powers JS, Knight JR, Harrell D, Varnell L, Vaughn C et al. Effectiveness of an inpatient  
19 geriatric service in a university hospital. *Journal of the Tennessee Medical Association*. 1994;  
20 87(10):425-428
- 21 105 Williams ME, Williams TF, Zimmer JG, Hall WJ, Podgorski CA. How does the team approach to  
22 outpatient geriatric evaluation compare with traditional care: a report of a randomized  
23 controlled trial. *Journal of the American Geriatrics Society*. 1987; 35(12):1071-1078
- 24 106 Winograd CH, Gerety MB, Lai NA. A negative trial of inpatient geriatric consultation. Lessons  
25 learned and recommendations for future research. *Archives of Internal Medicine*. 1993;  
26 153(17):2017-2023
- 27 107 Wong BJ, Vogenberg FR, Gilbert HD, Dupee RM. Effectiveness of a pharmacist on a geriatric  
28 assessment team. *P and T*. 1996; 21(3):135-144
- 29 108 Wooldridge DB, McInnis JB, Nelson R, Piller J, Scott S, Whiting P. A geriatric evaluation and  
30 management (GEM) program: evaluation of patient outcomes. *Aging*. 1995; 7(3):251-254
- 31  
32

# 1 Appendices

## 2 Appendix A: Review protocol


3 **Table 7: Review protocol: Assessment through ECAU**

Review question	Admission through ECAU
Guideline condition and its definition	Acute medical emergencies. Definition: people with suspected or confirmed acute medical emergencies or at risk of an acute medical emergency.
Review population	Elderly people (65 years and over) with a suspected or confirmed AME.
Interventions and comparators: generic/class; specific/drug	<ul style="list-style-type: none"> <li>• Assessment and management during admission through an elderly care/frailty assessment unit.</li> <li>• Assessment and management during admission through an elderly care assessment area.</li> <li>• Assessment and management during admission by a geriatric team.</li> </ul>
Comparison	<ul style="list-style-type: none"> <li>• No assessment and management through the ECAU: <ul style="list-style-type: none"> <li>○ Direct admission to a general medical ward from ED or by community or GP referral (inpatient care only).</li> <li>○ Admission through the AMU without geriatric team involvement.</li> </ul> </li> </ul>
Outcomes	<ul style="list-style-type: none"> <li>- Mortality during the study period (Dichotomous) CRITICAL</li> <li>- Patient and/or carer satisfaction during the study period (Dichotomous) CRITICAL</li> <li>- Length of stay during the study period (Continuous) CRITICAL</li> <li>- Adverse event rates during the study period (Dichotomous) CRITICAL</li> <li>- Quality of life during the study period (Continuous) CRITICAL</li> <li>- Readmission (up to 30 days) during the study period (Dichotomous) IMPORTANT</li> <li>- A&amp;E 4 hour waiting target met during the study period (Dichotomous) IMPORTANT</li> <li>- Delayed transfers of care during the study period (Dichotomous) IMPORTANT</li> </ul>
Study design	Systematic reviews (SRs) of RCTs, RCTs, observational studies only to be included if no relevant SRs or RCTs are identified.
Unit of randomisation	Patient. Hospital. Ward.
Crossover study	Not permitted.
Minimum duration of study	Not defined.
Population stratification	None.
Reasons for stratification	Not applicable.
Subgroup analyses if there is heterogeneity	- Older than 85 years (85 years and younger; older than 85 years); effects may be different in this subgroup.
Search criteria	Databases: Medline, Embase, the Cochrane Library. Date limits for search: None. Language: English language only.

1

## Appendix B: Clinical article selection

Figure 1: Flow chart of clinical article selection for the review of assessment through ECAU



2

3




1

# Appendix C: Forest plots


## C.1 Admission through ECAU versus direct admission

**Figure 2: Readmission (30 days)**




3

**Figure 3: Mortality**



4


**Figure 4: Length of stay**



5

## C.2 Admission through ECA area within AMU versus direct admission

**Figure 5: Mortality (in-patient)**



7

**Figure 6: Mortality (30-day)**



8


**Figure 7: Readmission (30 days)**



1


## 2 C.3 Admission by an elderly care team versus direct admission

**Figure 8: Readmission (28-day)**



3

**Figure 9: Readmission (7-day)**



4

5

## Appendix D: Clinical Evidence tables

Study	Cardwell 2016 <sup>12</sup>
Study type	Before and after study
Number of studies (number of participants)	1 (n=16,061)
Countries and setting	Conducted in United Kingdom; setting: single centre ED
Line of therapy	Not applicable
Duration of study	Other: 6 months before the intervention and the same 6 months after
Method of assessment of guideline condition	Adequate method of assessment/diagnosis
Stratum	Admission through the AMU with care from a visiting elderly care team (geriatrician team): NA
Subgroup analysis within study	Not applicable
Inclusion criteria	Over 65 attending the ED between 9am and 5pm Monday - Friday
Exclusion criteria	Stroke, high level of care needed, on renal dialysis
Recruitment/selection of patients	Consecutive patients meeting the inclusion criteria during the study period
Age, gender and ethnicity	Age - Other: over 65s. Gender (M:F): not reported. Ethnicity: not reported
Further population details	1. Older than 85 years: Not applicable/Not stated/Unclear
Indirectness of population	No indirectness: n/a
Interventions	<p>(n=8084) Intervention 1: Assessment and management through the ECAU at any part in the clinical pathway i.e. direct admission to EAU from GP, ED, or community referral. 'Front door' assessment of all over 65s with frailty - multidisciplinary team (consultant geriatrician, consultant in emergency medicine, emergency department nursing staff, specialist nurses from IC&amp;ES, elderly mental health liaison nurse, local GP, pharmacist, physiotherapist, advanced nurse practitioner and admin staff) at the front desk in the ED with access to 8 care-of-the-elderly inpatient beds and 2 23-hour beds in the clinical decisions unit adjacent to the ED; team used a frailty index to screen between 9am-5pm Monday to Friday, those identified as frail entered the frail elderly pathway developed in the hospital. Duration: 6 months. Concurrent medication/care: n/a</p> <p>(n=7977) Intervention 2: No assessment and management through the ECAU at any part in the clinical pathway - Direct admission to a general medical ward from ED or by community or GP referral (inpatient care only). Usual care - no screening for frailty, ED processed the admissions in the same way as for all adult age groups – directed to the Acute Medical Receiving Unit as clinically appropriate. Duration: 6 months. Concurrent medication/care: NA</p>

Study	Cardwell 2016 <sup>12</sup>
Funding	-- (QuEST)
RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ASSESSMENT AND MANAGEMENT THROUGH THE ECAU AT ANY PART IN THE CLINICAL PATHWAY I.E. DIRECT ADMISSION TO EAU FROM GP, ED, OR COMMUNITY REFERRAL versus DIRECT ADMISSION TO A GENERAL MEDICAL WARD FROM ED OR BY COMMUNITY OR GP REFERRAL (INPATIENT CARE ONLY)	
Protocol outcome 1: Readmission (up to 30 days) - Actual outcome for Admission through the AMU with care from a visiting elderly care team (geriatrician team): 28-day readmission at 28 days; Group 1: 620/4746, Group 2: 885/4547; Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness - Actual outcome for Admission through the AMU with care from a visiting elderly care team (geriatrician team): 7-day readmission at 7 days; Group 1: 138/4746, Group 2: 402/4547; Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness	
Protocol outcomes not reported by the study	Mortality; Patient and/or carer satisfaction; Length of stay; Adverse event rates; Quality of life; A&E 4 hour waiting target met; Delayed transfers of care

Study	Conroy 2014 <sup>18</sup>
Study type	Before and after study
Number of studies (number of participants)	1 (n=4647)
Countries and setting	Conducted in United Kingdom; setting: ED East Midlands, UK
Line of therapy	Not applicable
Duration of study	Other: 2010-2012
Method of assessment of guideline condition	Adequate method of assessment/diagnosis
Stratum	Admission through an Elderly care/frailty Assessment Unit: n/a
Subgroup analysis within study	Not applicable: n/a
Inclusion criteria	All patients presenting to the ED
Exclusion criteria	Not reported
Recruitment/selection of patients	Consecutive patients presenting to the ED during the study period
Age, gender and ethnicity	Age - Other: 65+. Gender (M:F): Define. Ethnicity: not reported

Study	Conroy 2014 <sup>18</sup>
Further population details	1. Older than 85 years: Not applicable/Not stated/Unclear (638 in the control group and 753 in the intervention group were over 85 years).
Indirectness of population	No indirectness: n/a
Interventions	(n=2490) Intervention 1: Assessment and management through the ECAU at any part in the clinical pathway i.e. direct admission to EAU from GP, ED, or community referral. Emergency frailty unit - embedded comprehensive geriatric assessment service within the ED. Duration: July 2011 - June 2012. Concurrent medication/care: not reported  (n=2184) Intervention 2: No assessment and management through the ECAU at any part in the clinical pathway - Admission through the AMU. Emergency decisions unit - no routine input from specialists trained in geriatric medicine. Duration: 12 months (2010). Concurrent medication/care: not reported
Funding	Funding not stated
<p>RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ASSESSMENT AND MANAGEMENT THROUGH THE ECAU AT ANY PART IN THE CLINICAL PATHWAY I.E. DIRECT ADMISSION TO EAU FROM GP, ED, OR COMMUNITY REFERRAL versus ADMISSION THROUGH THE AMU</p> <p>Protocol outcome 1: Readmission (up to 30 days) - Actual outcome for Admission through an Elderly care/frailty Assessment Unit: 30 day readmission rate at 30 days; Group 1: 221/2490, Group 2: 254/2184; Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness</p>	
Protocol outcomes not reported by the study	Mortality; Patient and/or carer satisfaction; Length of stay; Adverse event rates; Quality of life; A&E 4 hour waiting target met; Delayed transfers of care

Study	Ellis 2012 <sup>31</sup>
Study type	Before and after study
Number of studies (number of participants)	1 (n=422)
Countries and setting	Conducted in United Kingdom; setting: district general hospital, Scotland
Line of therapy	Not applicable
Duration of study	--: Oct 2009 - February 2010
Method of assessment of guideline condition	Adequate method of assessment/diagnosis
Stratum	Admission through an Elderly care/frailty Assessment Unit: n/a

Study	Ellis 2012 <sup>31</sup>
Subgroup analysis within study	Not applicable: n/a
Inclusion criteria	Over 65 with 1 or more of the following: functional impairment (acute or chronic), cognitive impairment (acute or chronic), falls or other geriatric syndromes, care home patients
Exclusion criteria	Functionally independent patients or those with only single organ pathology requiring specialist input
Recruitment/selection of patients	Consecutive patients meeting the inclusion criteria during the study period
Age, gender and ethnicity	Age - Other: mean age 80.5 before ACE, mean age 81.1 after ACE. Gender (M:F): before ACE 59.4% female, after ACE 63.2% female. Ethnicity: not reported
Further population details	1. Older than 85 years: Not applicable/Not stated/Unclear (some patients were over 85 but unclear what proportion).
Indirectness of population	No indirectness: n/a
Interventions	(n=210) Intervention 1: Assessment and management through the ECAU at any part in the clinical pathway i.e. direct admission to EAU from GP, ED, or community referral. Acute care for elders unit - situated adjacent to the ED and medical receiving unit, designed to deliver rapid and thorough CGA for patients deemed by non-specialists to require admission as a form of clinical decision unit. Duration: December 2009 to February 2010. Concurrent medication/care: not reported  (n=212) Intervention 2: No assessment and management through the ECAU at any part in the clinical pathway - Admission through the AMU. Medical receiving unit - use of standardised screening and assessment tools, multidimensional assessment by a multidisciplinary team and proactive discharge planning. Duration: October to December 2009. Concurrent medication/care: not reported
Funding	No funding
RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ASSESSMENT AND MANAGEMENT THROUGH THE ECAU AT ANY PART IN THE CLINICAL PATHWAY I.E. DIRECT ADMISSION TO EAU FROM GP, ED, OR COMMUNITY REFERRAL versus ADMISSION THROUGH THE AMU	
<p>Protocol outcome 1: Mortality</p> <p>- Actual outcome for Admission through an Elderly care/frailty Assessment Unit: mortality at 12 months; Group 1: 76/210, Group 2: 89/212; Risk of bias: All domain - Low, Selection - High, Blinding - Low, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness</p> <p>Protocol outcome 2: Length of stay</p> <p>- Actual outcome for Admission through an Elderly care/frailty Assessment Unit: mean total length of stay at hospital stay; Group 1: mean 12.7 days (SD 21.01); n=210, Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness</p>	

Study	Ellis 2012 <sup>31</sup>
Protocol outcome 3: Readmission (up to 30 days) - Actual outcome for Admission through an Elderly care/frailty Assessment Unit: 30 day readmissions at 30 days; Group 1: 33/210, Group 2: 36/212; Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness	
Protocol outcomes not reported by the study	Patient and/or carer satisfaction; Adverse event rates; Quality of life; A&E 4 hour waiting target met; Delayed transfers of care

Study	Taylor 2016 <sup>99</sup>
Study type	Before and after study
Number of studies (number of participants)	1 (n=811)
Countries and setting	Conducted in United Kingdom; setting: large urban teaching hospital, UK
Line of therapy	Unclear
Duration of study	Intervention + follow up
Method of assessment of guideline condition	Adequate method of assessment/diagnosis
Stratum	Admission through an Elderly care Assessment Area (defined area) within the AMU: n/a
Subgroup analysis within study	Not applicable: n/a
Inclusion criteria	Patients over 75 years admitted to the emergency assessment unit
Exclusion criteria	Not reported
Recruitment/selection of patients	Consecutive patients meeting the inclusion criteria during the study period
Age, gender and ethnicity	Age - Median (range): pre-intervention 85(75-101), post-intervention 84 (75-101). Gender (M:F): M:F 293:518. Ethnicity: not reported
Further population details	1. Older than 85 years: Not applicable/Not stated/Unclear
Indirectness of population	No indirectness: n/a
Interventions	(n=413) Intervention 1: Assessment and management through the ECAU at any part in the clinical pathway i.e. direct admission to EAU from GP, ED, or community referral. Comprehensive older person's evaluation (COPE) zone - within the emergency assessment unit, twice daily MDT meeting, and patients identified as potentially fit for discharge kept on COPE zone, otherwise transferred to geriatric medicine ward. Duration: 1 month (September 2014). Concurrent medication/care: not reported

Study	Taylor 2016 <sup>99</sup>
	(n=398) Intervention 2: No assessment and management through the ECAU at any part in the clinical pathway - Admission through the AMU. Medical patients admitted to the emergency assessment unit after being referred from the ED or a GP, patients requiring geriatrician input were seen by a daily in-reaching service. Duration: 1 month (September 2013). Concurrent medication/care: not reported
Funding	Funding not stated
<p>RESULTS (NUMBERS ANALYSED) AND RISK OF BIAS FOR COMPARISON: ASSESSMENT AND MANAGEMENT THROUGH THE ECAU AT ANY PART IN THE CLINICAL PATHWAY I.E. DIRECT ADMISSION TO EAU FROM GP, ED, OR COMMUNITY REFERRAL versus ADMISSION THROUGH THE AMU</p> <p>Protocol outcome 1: Mortality            - Actual outcome for Admission through an Elderly care Assessment Area (defined area) within the AMU: in-patient deaths at admission; Group 1: 37/413, Group 2: 32/398; Risk of bias: Low; Indirectness of outcome: No indirectness            - Actual outcome for Admission through an Elderly care Assessment Area (defined area) within the AMU: mortality at 30 days; Group 1: 19/413, Group 2: 22/398; Risk of bias: All domain - Low, Selection - High, Blinding - Low, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness, Comments: NA; Baseline details: greater proportion of males in intervention group</p> <p>Protocol outcome 2: Readmission (up to 30 days)            - Actual outcome for Admission through an Elderly care Assessment Area (defined area) within the AMU: readmission at 30 days; Group 1: 68/376, Group 2: 69/366; Risk of bias: All domain - High, Selection - High, Blinding - High, Incomplete outcome data - Low, Outcome reporting - Low, Measurement - Low, Crossover - Low; Indirectness of outcome: No indirectness, Comments: NA; Baseline details: greater proportion of males in intervention group</p>	
Protocol outcomes not reported by the study	Patient and/or carer satisfaction; Length of stay; Adverse event rates; Quality of life; A&E 4 hour waiting target met; Delayed transfers of care



## Appendix E: Economic evidence tables

Study	Cardwell 2016 <sup>13</sup>			
Study details	Population & interventions	Costs	Health outcomes	Cost-effectiveness
<p><b>Economic analysis:</b> CC</p> <p><b>Study design:</b> Cohort study</p> <p><b>Approach to analysis:</b> 6 months prospective cohort in 2014 compared with the same 6 months in previous year</p> <p><b>Perspective:</b> NHS hospital</p> <p><b>Time horizon/Follow-up:</b> 28 days</p> <p><b>Discounting:</b> No discounting.</p>	<p><b>Population:</b> Patients age&gt;65 attending the ED (excluding those with an obvious specialist pathway (stroke, renal dialysis, ITU). A large district general hospital located just outside Kilmarnock. Mean age: NR % male: NR</p> <p><b>Intervention 1:</b> Frail older people's pathway (FOPP) - Frailty MDT team 9am-5pm. Those assessed to be frail in the ED were put on the frail person's pathway. (n=8,084)</p> <p><b>Intervention 2:</b> No FOPP. (n=7,977)</p>	<p><b>Incremental Costs (2-1) (mean per patient):</b> Intervention: +£19 LOS: -£67 Admission: -£63 Re-attendance -£11 Re-admission: -163 Total: -£287 (95% CI: NR; p=NR)</p> <p><b>Currency &amp; cost year:</b> 2014? UK pounds</p> <p><b>Cost components incorporated:</b> Bed days, admissions, re-attendances, re-admissions</p>	NA	<p>NA</p> <p><b>Analysis of uncertainty:</b> NR</p>
<b>Data sources</b>				
<b>Health outcomes:</b> NA. <b>Quality-of-life weights:</b> NA <b>Cost sources:</b> Agenda for change pay scales and 'NHS bed-day cost for each ward'.				
<b>Comments</b>				
<b>Source of funding:</b> QuEST, NHS Scotland <b>Applicability and limitations:</b> Only cost comparison – only indicators of health were process outcomes like re-attendance and re-admission. Usual care was not described. The study was observational study, with no control for case-mix or time trend. No statistical or sensitivity analysis undertaken. Only hospital costs included. <b>Other:</b>				
<b>Overall applicability:</b> <sup>(a)</sup> Partially applicable <b>Overall quality:</b> <sup>(b)</sup> Potentially serious limitations				

Abbreviations: CC: Comparative costs; 95% CI: 95% confidence interval; NA: not applicable; NR: not reported.

(a) Directly applicable/Partially applicable/Not applicable.

(b) Minor limitations/Potentially serious limitations/Very serious limitations.

# Appendix F: GRADE tables

**Table 8: Clinical evidence profile: admission through ECAU versus direct admission**

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ECAU	direct admission	Relative (95% CI)	Absolute		
<b>Readmission (30-day) (follow-up 30 days; assessed with: no. of patients readmitted)</b>												
2	observational studies	serious <sup>1</sup>	no serious inconsistency	no serious indirectness	serious <sup>2</sup>	none	254/270 0 (9.4%)	14.3%	RR 0.78 (0.67 to 0.92)	31 fewer per 1000 (from 11 fewer to 47 fewer)	⊕○○○ VERY LOW	IMPORTANT
<b>Mortality (12 months) (follow-up 12 months; assessed with: no. of patients dying)</b>												
1	observational studies	no serious risk of bias <sup>1</sup>	no serious inconsistency	no serious indirectness	serious <sup>2</sup>	none	76/210 (36.2%)	42%	RR 0.86 (0.68 to 1.1)	59 fewer per 1000 (from 134 fewer to 42 more)	⊕○○○ VERY LOW	CRITICAL
<b>Length of stay (measured with: mean length of stay; Better indicated by lower values)</b>												
1	observational studies	serious <sup>1</sup>	no serious inconsistency	no serious indirectness	no serious imprecision	none	210	212	-	MD 0.5 higher (3.29 lower to 4.29 higher)	⊕○○○ VERY LOW	CRITICAL

<sup>1</sup> All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

<sup>2</sup> Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

**Table 9: Clinical evidence profile: admission through ECA area within AMU versus direct admission**

Quality assessment							No of patients		Effect		Quality	Importance
No of	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other	ECA area	direct	Relative	Absolute		

studies						considerations	within AMU	admission	(95% CI)			
<b>In-patient mortality (assessed with: no. of patients dying in hospital)</b>												
1	observational studies	no serious risk of bias <sup>1</sup>	no serious inconsistency	no serious indirectness	very serious <sup>2</sup>	none	37/413 (9%)	8%	RR 1.11 (0.71 to 1.75)	9 more per 1000 (from 23 fewer to 60 more)	⊕○○○ VERY LOW	CRITICAL
<b>30 day mortality (follow-up 30 days; assessed with: no. of patients dying within 30 days of discharge)</b>												
1	observational studies	no serious risk of bias <sup>1</sup>	no serious inconsistency	no serious indirectness	very serious	none	19/413 (4.6%)	5.5%	RR 0.83 (0.46 to 1.51)	9 fewer per 1000 (from 30 fewer to 28 more)	⊕○○○ VERY LOW	CRITICAL
<b>Readmission (30-day) (follow-up 30 days; assessed with: no. of patients readmitted )</b>												
1	observational studies	Serious <sup>1</sup>	no serious inconsistency	no serious indirectness	very serious	none	68/376 (18.1%)	18.9%	RR 0.96 (0.71 to 1.3)	8 fewer per 1000 (from 55 fewer to 57 more)	⊕○○○ VERY LOW	IMPORTANT

<sup>1</sup> All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

<sup>2</sup> Downgraded by 1 increment if the confidence interval crossed 1 MID or by 2 increments if the confidence interval crossed both MIDs.

**Table 10: Clinical evidence profile: admission by an elderly care team versus direct admission**

Quality assessment							No of patients		Effect		Quality	Importance
No of studies	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	ECA area within AMU	direct admission	Relative (95% CI)	Absolute		
<b>Readmission (assessed with: no. of patients readmitted within 28 days)</b>												
1	observational studies	serious <sup>1</sup>	no serious inconsistency	no serious indirectness	no serious imprecision	none	620/4746 (13.1%)	19.5%	RR 0.67 (0.61 to 0.74)	64 fewer per 1000 (from 51 fewer to 76 fewer)	⊕○○○ VERY LOW	IMPORTANT
<b>Readmission (assessed with: no. of patients readmitted within 7 days)</b>												
1	observational studies	serious <sup>1</sup>	no serious inconsistency	no serious indirectness	no serious imprecision	none	138/4746 (2.9%)	8.8%	RR 0.33 (0.27 to 0.40)	59 fewer per 1000 (from 53 fewer to 64 fewer)	⊕○○○ VERY LOW	IMPORTANT

									0.40)	fewer)	LOW	
--	--	--	--	--	--	--	--	--	-------	--------	-----	--

<sup>1</sup> All non-randomised studies automatically downgraded due to selection bias. Studies may be further downgraded by 1 increment if other factors suggest additional high risk of bias, or 2 increments if other factors suggest additional very high risk of bias.

1

## Appendix G: Excluded clinical studies

2

**Table 11: Studies excluded from the clinical review**

Study	Exclusion reason
Allen 2010 <sup>1</sup>	n<250
Applegate 1990 <sup>3</sup>	Not guideline condition (inclusion – medically stable). CGA not focused on admission
Applegate 1991 <sup>2</sup>	Literature review
Basic 2005 <sup>4</sup>	Not guideline condition (inclusion – medically stable)
Becker 1987 <sup>5</sup>	Inappropriate comparison - multidimensional evaluation conducted by geriatric consultation team (GCT) for both intervention and control and treatment provided by GCT only for intervention group
Bloch 2013 <sup>6</sup>	Incorrect interventions
Borenstein 2016 <sup>7</sup>	Intervention not focused on admission (similar length of stay within the unit to a general medical ward)
Braude 2016 <sup>8</sup>	Incorrect population –Surgical patients. Study assessed ward based geriatric liaison service for older urological surgical patients
Burke 2001 <sup>9</sup>	No comparator
Campbell 1987 <sup>10</sup>	Study design (literature review)
Cape 1994 <sup>11</sup>	No comparator
Cavaliere 1993 <sup>14</sup>	Incorrect interventions (nursing home)
Cefalu 1997 <sup>15</sup>	No comparator
Cohen 2002 <sup>16</sup>	Inappropriate intervention. Not focussed on admission
Collard 1985 <sup>17</sup>	Incorrect interventions
Conroy 2011 <sup>19</sup>	Systematic review: study designs inappropriate
Covinsky 1998 <sup>20</sup>	Article on patients perspective on an acute care for elders unit
Dasgupta 1980 <sup>21</sup>	Outcome reporting (data cannot be extracted)
Del giudice 2009 <sup>22</sup>	Incorrect intervention -post-acute geriatric evaluation and management unit. Not focused on admission
Denewet 2016 <sup>23</sup>	Incorrect population –oncology patients. Study evaluated CGA for predicting survival in geriatric oncology
Edmans 2011 <sup>26</sup>	CGA not focused on admission (discharge)
Edmans 2013 <sup>24</sup>	Study design (prognostic)
Edmans 2013 <sup>25</sup>	CGA not focused on admission (discharge)
Ekdahl 2015 <sup>29</sup>	Incorrect intervention (comprehensive geriatric assessment provided by an ambulatory geriatric care unit in outpatient setting)
Ekdahl 2015 <sup>29</sup>	Outpatient setting- Comprehensive geriatric assessment (CGA) provided by an ambulatory geriatric care unit (AGU)
Ekdahl 2015 <sup>27</sup>	Systematic review is not relevant to review question or unclear PICO
Ekdahl 2016 <sup>28</sup>	Incorrect intervention and setting- CGA in a geriatric ambulatory unit in a municipality
Elliott 2012 <sup>30</sup>	Incorrect interventions (home care)
Ellis 2004 <sup>33</sup>	Systematic review. Checked for relevant references
Ellis 2006 <sup>32</sup>	Protocol for Cochrane review
Ellis 2011 <sup>35</sup>	Systematic review: study designs inappropriate

Study	Exclusion reason
Ellis 2011 <sup>36</sup>	Systematic review. Ordered relevant references
Ellis 2014 <sup>34</sup>	Descriptive literature review
Epstein 1990 <sup>37</sup>	Incorrect interventions (ambulatory care)
Faul 2009 <sup>38</sup>	Incorrect interventions (community-based)
Fletcher 2002 <sup>39</sup>	Incorrect interventions (community-based)
Fox 2016A <sup>40</sup>	No comparator
Fretwell 1987 <sup>41</sup>	Incorrect interventions (community-based)
Fretwell 1990 <sup>42</sup>	Incorrect population (patients included when transferred out of intensive and coronary-care units). Included out-patient follow-up
Germain 1995 <sup>43</sup>	Inappropriate comparison (ECAU compared to ECAU + team)
Gerritsen 1995 <sup>44</sup>	No comparator
Gladman 2012 <sup>45</sup>	CGA not focused on admission (discharge)
Graf 2011 <sup>46</sup>	Systematic review: study designs inappropriate
Gregersen 2012 <sup>47</sup>	Incorrect comparison (geriatric department compared with general medical department)
Harari 2007 <sup>49</sup>	Not guideline condition (elective surgical admissions)
Harari 2007 <sup>48</sup>	Observational study n<250
Harris 1991 <sup>50</sup>	Geriatric assessment unit not focused on admission (similar length of stay within the unit to a general medical ward)
Heath 2005 <sup>51</sup>	Incorrect interventions (home care)
Hernandez-vian 2007 <sup>52</sup>	Non-English
Hogan 1984 <sup>54</sup>	No comparator
Hogan 1990 <sup>53</sup>	Literature review
Horgan 2012 <sup>55</sup>	No comparator
Humphries 1992 <sup>56</sup>	Incorrect interventions. No comparison
Jones 2004 <sup>57</sup>	Incorrect interventions (community-based)
Kamel 2005 <sup>58</sup>	Systematic review is not relevant to review question or unclear PICO
Karppi 1995 <sup>59</sup>	Inappropriate comparison (home-care). CGA not focused on admission (discharge)
Karppi 1995 <sup>60</sup>	Inappropriate comparison (home-care)
Kay 1992 <sup>61</sup>	Not guideline condition (inclusion – medically stable)
Kergoat 2012 <sup>62</sup>	No comparator
Kircher 2007 <sup>63</sup>	Incorrect intervention (not focussed on admission)
Landefeld 1995 <sup>64</sup>	Inappropriate intervention. Not focussed on admission
Landi 2001 <sup>65</sup>	Incorrect interventions (home-care)
Lightbody 2002 <sup>66</sup>	Commentary on Cohen 2002
Mcdowell 1994 <sup>67</sup>	Incorrect interventions (out-patient)
McVey 1989 <sup>68</sup>	Not review population
Naughton 1994 <sup>72</sup>	Incorrect interventions (not focussed on admission)
Nikolaus 1999 <sup>73</sup>	Incorrect intervention (not focussed on admission); not review population (patients had to be stable)
Nipp 2012 <sup>74</sup>	Not review population
Owen 2015 <sup>76</sup>	No comparator
Parker 2000 <sup>77</sup>	Systematic review: study designs inappropriate

Study	Exclusion reason
Phibbs 2006 <sup>78</sup>	CGA not focused on admission (in-patient, discharge, and out-patient)
Pitner 2004 <sup>79</sup>	Incorrect intervention (not focussed on admission)
Popplewell 1983 <sup>80</sup>	Geriatric assessment unit not focused on admission (similar length of stay within the unit to a general medical ward)
Reuben 1995 <sup>81</sup>	CGA not focused on admission (patients included 24-72 after admission)
Riley 1974 <sup>82</sup>	Descriptive
Rockwood 2003 <sup>83</sup>	Incorrect interventions (community-based)
Rosenberg 2012 <sup>84</sup>	Incorrect interventions (home-care)
Rubenstein 1984 <sup>87</sup>	Not AME patients- patients still in the hospital 1 week after admission for acute care included in the study i.e. after stabilisation of their acute problems
Rubenstein 1987 <sup>85</sup>	Not AME patients (sub-acute)
Rubenstein 1995 <sup>86</sup>	Not AME patients- Only medically stable patients included
Saltvedt 2002 <sup>88</sup>	Incorrect intervention (not focussed on admission)
Saltvedt 2004 <sup>89</sup>	Incorrect interventions(not focussed on admission)
Saltvedt 2005 <sup>91</sup>	Incorrect intervention (not focussed on admission)
Saltvedt 2006 <sup>90</sup>	Incorrect intervention (not focussed on admission)
Saltz 1988 <sup>92</sup>	Not review population
Silverman 1995 <sup>94</sup>	Incorrect interventions (out-patient)
Soejono 2008 <sup>96</sup>	Incorrect intervention (not focussed on admission)
Stewart 1999 <sup>97</sup>	Incorrect intervention (not focussed on admission)
Stuck 1995 <sup>98</sup>	Incorrect intervention (home-care)
Teasdale 1983 <sup>100</sup>	Geriatric assessment unit not focused on admission (rehabilitation)
Toseland 1996 <sup>101</sup>	Incorrect interventions (out-patient)
Trentini 2001 <sup>102</sup>	Incorrect setting -outpatient. The study assessed effectiveness of outpatient elderly care based on CGA
Van Craen 2010 <sup>103</sup>	Systematic review. Ordered relevant references
White 1994 <sup>104</sup>	Inappropriate population- medically stable elderly patients at risk for function decline or with rehab potential
Williams 1987 <sup>105</sup>	Incorrect interventions (out-patient)
Winograd 1993 <sup>106</sup>	Incorrect intervention (not focussed on admission)
Wong 1996 <sup>107</sup>	Inappropriate comparison (team compared to team + pharmacist)
Wooldridge 1995 <sup>108</sup>	Incorrect intervention (not focussed on admission)

1  
2

1  
2  
3

## **Appendix H: Excluded economic studies**

No studies were excluded.