

# NATIONAL INSTITUTE FOR HEALTH AND CLINICAL EXCELLENCE

## Final appraisal determination

### Alendronate, etidronate, risedronate, raloxifene, strontium ranelate and teriparatide for the secondary prevention of osteoporotic fragility fractures in postmenopausal women

NOTE: The amendments made in this final appraisal determination (published originally as technology appraisal guidance 161 in October 2008) are the consequence of reconsideration by the Appraisal Committee following Her Majesty's High Court of Justice order of March 2009.

## 1 Guidance

This guidance relates only to treatments for the secondary prevention of fragility fractures in postmenopausal women who have osteoporosis and have sustained a clinically apparent osteoporotic fragility fracture. Osteoporosis is defined by a T-score<sup>1</sup> of  $-2.5$  standard deviations (SD) or lower on dual-energy X-ray absorptiometry (DXA) scanning. However, the diagnosis may be assumed in women aged 75 years or older if the responsible clinician considers a DXA scan to be clinically inappropriate or unfeasible.

This guidance assumes that women who receive treatment have an adequate calcium intake and are vitamin D replete. Unless clinicians are confident that women who receive treatment meet these criteria, calcium and/or vitamin D supplementation should be considered.

---

<sup>1</sup> T-score relates to the measurement of bone mineral density (BMD) using central (hip and/or spine) DXA scanning and is expressed as the number of standard deviations (SD) from peak BMD.

NICE is developing a clinical guideline on 'Osteoporosis: assessment of fracture risk and the prevention of osteoporotic fractures in individuals at high risk' (see [www.nice.org.uk](http://www.nice.org.uk)). This technology appraisal guidance should be read in the context of the clinical guideline when it is available.

This guidance does **not** cover the following:

- The use of alendronate, etidronate, risedronate, raloxifene, strontium ranelate or teriparatide for the secondary prevention of osteoporotic fragility fractures in women with normal bone mineral density (BMD) or osteopenia (that is, women with a T-score between  $-1$  and  $-2.5$  SD below peak BMD).
- The use of these drugs for the secondary prevention of osteoporotic fragility fractures in women who are on long-term systemic corticosteroid treatment.

These groups will be covered within future guidance produced by the Institute.

- 1.1 Alendronate is recommended as a treatment option for the secondary prevention of osteoporotic fragility fractures in postmenopausal women who are confirmed to have osteoporosis (that is, a T-score of  $-2.5$  SD or below). In women aged 75 years or older, a DXA scan may not be required if the responsible clinician considers it to be clinically inappropriate or unfeasible.

When the decision has been made to initiate treatment with alendronate, the preparation prescribed should be chosen on the basis of the lowest acquisition cost available.

1.2 Risedronate and etidronate are recommended as alternative treatment options for the secondary prevention of osteoporotic fragility fractures in postmenopausal women:

- who are unable to comply with the special instructions for the administration of alendronate, or have a contraindication to or are intolerant of alendronate (as defined in section 1.6) **and**
- who also have a combination of T-score, age and number of independent clinical risk factors for fracture (see section 1.5) as indicated in the following table.

**T-scores (SD) at (or below) which risedronate or etidronate is recommended when alendronate cannot be taken**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
50–54	– <sup>a</sup>	–3.0	–2.5
55–59	–3.0	–3.0	–2.5
60–64	–3.0	–3.0	–2.5
65–69	–3.0	–2.5	–2.5
70 or older	–2.5	–2.5	–2.5

<sup>a</sup> Treatment with risedronate or etidronate is not recommended

If a women aged 75 years or older has not previously had her BMD measured, a DXA scan may not be required if the responsible clinician considers it to be clinically inappropriate or unfeasible.

In deciding between risedronate and etidronate, clinicians and patients need to balance the overall proven effectiveness profile of the drugs against their tolerability and adverse effects in individual patients.

1.3 Strontium ranelate and raloxifene are recommended as alternative treatment options for the secondary prevention of osteoporotic fragility fractures in postmenopausal women:

- who are unable to comply with the special instructions for the administration of alendronate and either risedronate or etidronate, or have a contraindication to or are intolerant of alendronate and either risedronate or etidronate (as defined in section 1.6) **and**
- who also have a combination of T-score, age and number of independent clinical risk factors for fracture (see section 1.5) as indicated in the following table.

**T-scores (SD) at (or below) which strontium ranelate or raloxifene is recommended when alendronate and either risedronate or etidronate cannot be taken**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
50–54	– <sup>a</sup>	–3.5	–3.5
55–59	–4.0	–3.5	–3.5
60–64	–4.0	–3.5	–3.5
65–69	–4.0	–3.5	–3.0
70–74	–3.0	–3.0	–2.5
75 or older	–3.0	–2.5	–2.5

<sup>a</sup> Treatment with raloxifene or strontium ranelate is not recommended

If a woman aged 75 years or older who has one or more independent clinical risk factors for fracture or indicators of low BMD has not previously had her BMD measured, a DXA scan may not be required if the responsible clinician considers it to be clinically inappropriate or unfeasible.

For the purposes of this guidance, indicators of low BMD are low body mass index (defined as less than 22 kg/m<sup>2</sup>), medical conditions such as ankylosing spondylitis, Crohn’s disease, conditions that result in prolonged immobility, and untreated premature menopause<sup>2</sup>.

<sup>2</sup> Rheumatoid arthritis is also a medical condition indicative of low BMD.

In deciding between strontium ranelate and raloxifene, clinicians and patients need to balance the overall proven effectiveness profile of these drugs against their tolerability and other effects in individual patients.

1.4 Teriparatide is recommended as an alternative treatment option for the secondary prevention of osteoporotic fragility fractures in postmenopausal women:

- who are unable to take alendronate and either risedronate or etidronate, or have a contraindication to or are intolerant of alendronate and either risedronate or etidronate (as defined in section 1.6), **or** who have a contraindication to, or are intolerant of strontium ranelate (as defined in section 1.7), **or** who have had an unsatisfactory response (as defined in section 1.8) to treatment with alendronate, risedronate or etidronate **and**
- who are 65 years or older and have a T-score of  $-4.0$  SD or below, or a T-score of  $-3.5$  SD or below plus more than two fractures, **or** who are aged 55–64 years and have a T-score of  $-4$  SD or below plus more than two fractures.

1.5 For the purposes of this guidance, independent clinical risk factors for fracture are parental history of hip fracture, alcohol intake of 4 or more units per day, and rheumatoid arthritis.

1.6 For the purposes of this guidance, intolerance of alendronate, risedronate or etidronate is defined as persistent upper gastrointestinal disturbance that is sufficiently severe to warrant discontinuation of treatment, and that occurs even though the instructions for administration have been followed correctly.

1.7 For the purposes of this guidance, intolerance of strontium ranelate is defined as persistent nausea or diarrhoea, either of which warrants discontinuation of treatment.

- 1.8 For the purposes of this guidance, an unsatisfactory response is defined as occurring when a woman has another fragility fracture despite adhering fully to treatment for 1 year and there is evidence of a decline in BMD below her pre-treatment baseline.
- 1.9 Women who are currently receiving treatment with one of the drugs covered by this guidance, but for whom treatment would not have been recommended according to sections 1.1 to 1.4, should have the option to continue treatment until they and their clinicians consider it appropriate to stop.

## **2 Clinical need and practice**

- 2.1 Osteoporosis is a progressive, systemic skeletal disorder characterised by low bone mass and micro-architectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture.
- 2.2 Bone formation exceeds bone resorption in youth, but by the third decade of life there is a gradual loss of bone mass. Osteoporosis is therefore usually an age-related disease. It can affect both sexes, but women are at greater risk because the decrease in oestrogen production after the menopause accelerates bone loss to a variable degree.
- 2.3 The World Health Organization (WHO) has established diagnostic criteria for osteoporosis based on the measurement of BMD, expressed as the T-score, which is the number of SD below the mean BMD of young adults at their peak bone mass:
- normal BMD: T-score of  $-1$  SD or above
  - osteopenia: T-score of between  $-1$  and  $-2.5$  SD
  - osteoporosis: T-score of  $-2.5$  SD or below
  - established (severe) osteoporosis: T-score of  $-2.5$  SD or below with one or more associated fractures.

- 2.4 T-score measurements vary depending on the site and method of investigation. Measurement of BMD using central (hip and/or spine) DXA scanning can estimate fracture risk.
- 2.5 It is estimated that more than 2 million women have osteoporosis (that is, have a T-score of  $-2.5$  SD or below) in England and Wales. Osteoporosis is most common in older white women. After the menopause, the prevalence of osteoporosis increases markedly with age, from approximately 2% at 50 years rising to more than 25% at 80 years.
- 2.6 Fragility fracture is the clinically apparent and relevant outcome in osteoporosis (referred to as 'osteoporotic fragility fracture' in the following text). It is often referred to as a low-trauma fracture; that is, a fracture sustained as the result of a force equivalent to the force of a fall from a height equal to, or less than, that of an ordinary chair. In the absence of fracture, osteoporosis is asymptomatic and often remains undiagnosed. Osteoporotic fragility fractures occur most commonly in the vertebrae, hip and wrist, and are associated with substantial disability, pain and reduced quality of life.
- 2.7 In women aged over 50 years, the lifetime risk of a vertebral fracture is estimated to be one in three, and that of hip fracture one in five. Postmenopausal women with an initial fracture are at substantially greater risk of subsequent fractures. For instance, a woman with a vertebral fracture has an increased relative risk (RR) of 4.4 for a further vertebral fracture, 2.3 for a hip fracture, and 1.4 for a wrist fracture.
- 2.8 It is estimated that annually there are 180,000 osteoporosis-related symptomatic fractures in England and Wales. Of these, 70,000 are hip fractures, 25,000 are clinical vertebral fractures, and 41,000 are wrist fractures.

- 2.9 After a hip fracture, a high proportion of women are permanently unable to walk independently or to perform other activities of daily living and, consequently, many are unable to live independently. Hip fractures are also associated with increased mortality; estimates of the relative mortality risk vary from 2 to greater than 10 in the 12 months following hip fracture. However, it is unclear to what extent this can be attributed to fracture alone as opposed to pre-existing comorbidity.
- 2.10 Vertebral fractures can be associated with curvature of the spine and loss of height and can result in pain, breathing difficulties, gastrointestinal problems and difficulties in performing activities of daily living. It is thought that the majority of vertebral fractures (50–70%) do not come to clinical attention. Vertebral fractures are also associated with increased mortality; UK-specific data indicate a 4.4–fold increase in mortality related to vertebral fracture. However, as with hip fractures, it is unclear to what extent this may be due to comorbidities.
- 2.11 In addition to increasing age and low BMD, other clinical factors have been associated with increased fracture risk. Some of these clinical risk factors are at least partly independent of BMD, and include parental history of hip fracture, alcohol intake of 4 or more units per day, long-term systemic use of corticosteroids (which is not covered in this guidance), and rheumatoid arthritis.
- 2.12 Factors that are known to be indicators of low BMD include low body mass index (BMI) (defined as less than 22 kg/m<sup>2</sup>), and medical conditions such as ankylosing spondylitis, Crohn's disease, conditions that result in prolonged immobility, and untreated premature menopause.
- 2.13 A full review of the risk factors associated with osteoporosis is being carried out for the development of the NICE clinical guideline



'Osteoporosis: assessment of fracture risk and the prevention of osteoporotic fractures in individuals at high risk' (see [www.nice.org.uk](http://www.nice.org.uk)).

### 3 The technologies

#### **Bisphosphonates: alendronate, etidronate, risedronate**

- 3.1 The bisphosphonates alendronate, etidronate and risedronate are inhibitors of bone resorption and increase BMD by altering osteoclast activation and function.
- 3.2 Alendronate is an oral bisphosphonate that has a UK marketing authorisation as a once-weekly preparation (70 mg) for the treatment of postmenopausal osteoporosis. It also has a marketing authorisation at a daily dose of 10 mg for the treatment of osteoporosis in postmenopausal women to prevent fractures. Non-proprietary alendronate (Teva UK) costs £4.12 for four 70 mg tablets and £8.30 for twenty-eight 10 mg tablets (excluding VAT; NHS Drug Tariff, 24 February 2008). At these prices the drug costs for 1 year are £53.56 for once-weekly (70 mg) tablets and £108.20 for daily (10 mg) tablets. Proprietary alendronate (Fosamax; Merck Sharp & Dohme) is priced at £22.80 for four 70 mg tablets) and £23.12 for twenty-eight 10 mg tablets (excluding VAT; 'British national formulary' [BNF] edition 54). At these prices, the drug costs for 1 year are £296.40 for once-weekly (70 mg) tablets and £301.39 for daily (10 mg) tablets. Costs may vary in different settings because of negotiated procurement discounts.
- 3.3 Etidronate (Didronel; Procter & Gamble UK) is an oral bisphosphonate that has a UK marketing authorisation for the treatment of osteoporosis. The drug is administered in 90-day cycles, with each cycle consisting of etidronate (400 mg/day) for 14 days followed by calcium carbonate (1.25 g/day) for the remaining 76 days. The price per 90-day pack is £21.12 (excluding

VAT; BNF 54), which equates to a yearly cost of £85.65. Costs may vary in different settings because of negotiated procurement discounts.

- 3.4 Risedronate (Actonel; Procter & Gamble UK) is an oral bisphosphonate that has a UK marketing authorisation at a dosage of 5 mg/day or 35 mg/week for the treatment of postmenopausal osteoporosis, to reduce the risk of vertebral fractures, and for the treatment of established postmenopausal osteoporosis, to reduce the risk of hip fractures. Prices are £19.10 for twenty-eight 5 mg tablets and £20.30 for four 35 mg tablets (excluding VAT; BNF 54), which equates to yearly costs of £248.98 for the daily treatment or £264.63 for the once-weekly treatment. Costs may vary in different settings because of negotiated procurement discounts.
- 3.5 Gastrointestinal side effects are common with oral bisphosphonates. In people with oesophageal abnormalities and other factors that delay oesophageal transit or emptying, risedronate should be used cautiously and alendronate is contraindicated. For full details of side effects and contraindications, see the summaries of product characteristics.
- 3.6 Bisphosphonates have relatively complex instructions for administration. Alendronate and risedronate must be taken with 200 ml and 120 ml of water, respectively. Before and immediately after administration patients should not eat or drink, and must remain upright for stipulated time periods. Etidronate should be taken with water at the midpoint of a 4-hour fast (that is, 2 hours after and 2 hours before food, vitamins with mineral supplements such as iron, calcium supplements, laxatives containing magnesium, or antacids containing calcium or aluminium).

### **Selective oestrogen receptor modulator: raloxifene**

- 3.7 Selective oestrogen receptor modulators (SERMs) are drugs with selective activity in various organ systems, acting as weak oestrogen-receptor agonists in some systems and as oestrogen antagonists in others. The aim of treatment with SERMs is to maximise the beneficial effects of oestrogen on bone and to minimise the adverse effects on the breast and endometrium.
- 3.8 Raloxifene (Evista; Eli Lilly) has a marketing authorisation for the treatment of osteoporosis in postmenopausal women. The recommended dosage is 60 mg/day. The prices of 28- and 84-tablet packs are £17.06 and £59.59, respectively (excluding VAT; BNF 54), which equate to yearly costs of £222.39 and £258.93, respectively. Costs may vary in different settings because of negotiated procurement discounts.
- 3.9 Raloxifene is contraindicated in people with a history of venous thromboembolism (VTE), hepatic impairment, cholestasis, severe renal impairment, unexplained uterine bleeding or endometrial cancer. Raloxifene should not be co-administered with systemic oestrogens and, in patients with breast cancer, it should not be used for osteoporosis treatment or prevention until treatment of the breast cancer, including adjuvant treatment, has been completed. Raloxifene is associated with an increased risk of venous thromboembolic events, particularly during the first 4 months of treatment, which is similar to the reported risk associated with hormone replacement therapy. For full details of side effects and contraindications, see the summary of product characteristics.

### **Strontium ranelate**

- 3.10 Strontium ranelate (Protelos; Servier Laboratories) is a divalent strontium salt of ranelic acid (strontium is an element with properties similar to calcium). It is thought to have a dual effect on bone metabolism, increasing bone formation and decreasing bone

resorption. It has a UK marketing authorisation for the treatment of postmenopausal osteoporosis to reduce the risk of vertebral and hip fractures. The recommended dose is one 2 g sachet taken daily as a suspension in water. The price of a 28-sachet pack is £25.60 (excluding VAT; BNF 54), which equates to a yearly cost of £333.71. Costs may vary in different settings because of negotiated procurement discounts.

- 3.11 The absorption of strontium ranelate is reduced by food, milk and products derived from milk. It should therefore be administered between meals, ideally at bedtime and preferably at least 2 hours after eating.
- 3.12 The summary of product characteristics states that strontium ranelate is not recommended in patients with severe renal impairment and that it should be used with caution in patients at increased risk of VTE. Treatment with strontium ranelate should be discontinued during treatment with oral tetracycline or quinolone antibiotics. For full details of side effects, drug interactions and contraindications, see the summary of product characteristics.

### **Parathyroid hormone: teriparatide**

- 3.13 Teriparatide (Forsteo; Eli Lilly & Company) is a recombinant fragment of human parathyroid hormone and, as an anabolic agent, it stimulates new formation of bone and increases resistance to fracture.
- 3.14 Teriparatide has a marketing authorisation in the UK for the treatment of established osteoporosis in postmenopausal women. The recommended dose is 20 micrograms administered once daily by subcutaneous injection in the thigh or abdomen. Patients taking teriparatide must receive training in the injection technique. At the time of appraisal, the maximum total duration of treatment was restricted, by the marketing authorisation, to 18 months (see the

summary of product characteristics for current information). The price of a 28-day pre-filled pen is £271.88 (excluding VAT; BNF 54), which equates to a yearly cost of £3544.15.

- 3.15 Particular contraindications include pre-existing hypercalcaemia, severe renal impairment, metabolic bone diseases other than primary osteoporosis (including hyperparathyroidism and Paget's disease of bone), unexplained elevations of alkaline phosphatase, and previous radiation treatment to the skeleton. For full details of side effects and contraindications, see the summary of product characteristics.

## **4 Evidence and interpretation**

The Appraisal Committee (appendix A) considered evidence from a number of sources (appendix B).

### **4.1 Clinical effectiveness**

#### **Efficacy**

- 4.1.1 The Assessment Group for this appraisal (School of Health and Related Research, University of Sheffield [ScHARR]) reviewed data from published randomised controlled trials (RCTs) in postmenopausal women in which fracture or health-related quality of life was an endpoint and where one of the six drugs of interest was compared with a relevant comparator, such as no treatment, placebo, or one of the other included interventions. The majority of studies used placebo or no treatment as a control. Most studies ensured that women in all trial arms had normal calcium levels (that is, normal serum concentrations) or adequate supplementation, and some studies used additional dietary supplementation with vitamin D.

- 4.1.2 For this appraisal, reductions in RR associated with treatment were pooled regardless of the baseline BMD and fracture status of the

participants in the studies. It was also assumed that these reductions in RR remained constant at all ages, although little evidence was available for the effectiveness of the drugs in women aged 80 years or older.

- 4.1.3 For vertebral fractures, some studies used clinical (that is, symptomatic) fractures as their endpoint whereas others used fractures that were identified radiographically. Vertebral fractures identified radiographically, which are termed 'radiographic fractures' or 'morphometric fractures', include both symptomatic and asymptomatic fractures. There are different definitions of a vertebral radiographic fracture, but those definitions that require a 20% reduction in vertebral height are generally recognised as producing more reliable results than those that require a 15% reduction.
- 4.1.4 For non-vertebral fracture types, individual data on hip, leg, pelvis, wrist, hand, foot, rib and humerus fractures were sometimes provided, whereas some studies only presented data for all non-vertebral fractures grouped together.

#### *Alendronate*

- 4.1.5 Sixteen RCTs of alendronate in postmenopausal women were included in the assessment report: two studies in women with low or normal BMD; one in women with osteopenia; eight in women with osteopenia or osteoporosis; four in women with osteoporosis; and one in women with established osteoporosis. Overall, 15 studies compared alendronate with placebo or with no treatment. All the studies were conducted in women who had adequate levels of calcium, from either dietary intake or calcium supplementation.
- 4.1.6 Two studies, one comparing alendronate with oestrogen alone or with oestrogen and alendronate combined, and the other comparing alendronate with teriparatide, found no statistically

significant differences between the groups in numbers of clinically apparent fractures of any type in women with osteoporosis.

However, back pain was reported less frequently by women in the teriparatide group compared with women in the alendronate group (6% versus 19%,  $p = 0.012$ ).

- 4.1.7 In addition to the 16 RCTs, a 2-year study demonstrated the equivalence of weekly and daily doses of alendronate, in terms of clinical fracture incidence and gastrointestinal adverse events. However, this study was not included in the analysis because it did not include the specified comparators.
- 4.1.8 The meta-analysis for alendronate relative to placebo, carried out by the Assessment Group, resulted in an RR of vertebral fracture of 0.56 (95% confidence interval [CI] 0.46 to 0.68, four RCTs,  $n = 7039$ ), an RR of hip fracture of 0.62 (95% CI 0.40 to 0.98, three RCTs,  $n = 7455$ ), an RR of wrist fracture of 0.67 (95% CI 0.34 to 1.31, four RCTs,  $n = 7931$ ) and an RR for other non-vertebral fractures of 0.81 (95% CI 0.68 to 0.97, six RCTs,  $n = 9973$ ).
- 4.1.9 A post-hoc analysis of data from the largest study on alendronate, the 'Fracture intervention trial' (FIT) RCT (non-vertebral fracture population), suggested that alendronate may be less effective at reducing fractures in women with T-scores above (that is, better than)  $-2.5$  SD than in women with osteoporosis. These results were not statistically significant.
- 4.1.10 Gastrointestinal adverse events, including nausea, dyspepsia, mild oesophagitis/gastritis and abdominal pain, were reported in at least one third of the participants in studies of alendronate. However, only one study found the increased frequency of these symptoms to be statistically significant relative to placebo. This is consistent with post-marketing studies that indicate that approximately one third of alendronate users experience gastrointestinal adverse

events. To avoid oesophagitis, the summary of product characteristics now recommends that alendronate should be taken on rising for the day, with a full glass of water. It is possible that these instructions were not followed in all of the studies, particularly the earlier ones.

- 4.1.11 Prescription-event monitoring studies in patients for whom alendronate was prescribed (n = 11,916) by GPs in England demonstrated a high incidence of dyspepsia, particularly in the first month of treatment. Consultations for dyspepsia ranged from 32.2 per 1000 patient-months in the first month of treatment to 10.9 per 1000 patient-months in months 2 to 6. Because these studies lacked a comparator, it is not possible to assess the extent to which these rates of upper gastrointestinal events may be above baseline levels in those not taking bisphosphonates.
- 4.1.12 One study reported health-related quality of life outcomes. At 12 months there were statistically significant improvements in the alendronate group compared with the control group in scores for pain, social isolation, energy level and physical ability.

#### *Etidronate*

- 4.1.13 Twelve RCTs of etidronate in postmenopausal women were reviewed: three studies in women with low-to-normal BMD; two in women with osteopenia or osteoporosis; one in women with osteoporosis; one in women with osteoporosis or established osteoporosis; and five in women with established osteoporosis. Four studies included active comparators, and eight compared etidronate with placebo or with no treatment (although in six of these, study participants in all arms received calcium, either alone or with vitamin D). Some studies did not use the exact treatment regimen that currently has a UK marketing authorisation (that is, 90-day cycles of etidronate 400 mg/day for 14 days, followed by



calcium carbonate 1.25 g/day for the remaining 76 days). None of the studies reported health-related quality of life outcomes.

- 4.1.14 The meta-analysis of RCTs for etidronate relative to placebo carried out by the Assessment Group resulted in an RR of vertebral fracture of 0.40 (95% CI 0.20 to 0.83, three RCTs, n = 341), an RR of hip fracture of 0.50 (95% CI 0.05 to 5.34, two RCTs, n = 180), and an RR for other non-vertebral fractures of 1.04 (95% CI 0.64 to 1.69, four RCTs, n = 410). There were no data for wrist fracture.
- 4.1.15 An observational study in a general practice setting in the UK reported on fracture rates in people with a diagnosis of osteoporosis who were receiving etidronate compared with those who were not taking a bisphosphonate. People taking etidronate had an RR of non-vertebral fracture of 0.80 (95% CI 0.70 to 0.92). The RR of hip fracture was 0.66 (95% CI 0.51 to 0.85) and that of wrist fracture was 0.81 (95% CI 0.58 to 1.14).
- 4.1.16 Higher rates of gastrointestinal adverse effects were found in the etidronate groups of four RCTs, although the differences were not always statistically significant. However, non-RCT evidence and testimonies from clinical specialists and patient experts suggested that etidronate may be associated with fewer gastrointestinal adverse effects than other bisphosphonates.
- 4.1.17 The systematic review carried out by SchARR in 2006 identified a cohort study conducted in the UK that indicated that etidronate may be associated with a much lower rate of upper gastrointestinal adverse effects than alendronate or risedronate.

#### *Risedronate*

- 4.1.18 Seven RCTs of risedronate in postmenopausal women were reviewed: one study in women with normal BMD; one in women with osteopenia; one in women with osteopenia or osteoporosis;

one in women with osteoporosis or specific risk factors for hip fracture, such as a recent fall; and three in women with established osteoporosis. All compared risedronate with placebo (although, with the exception of those in the normal BMD study, all women also received calcium) and none reported on health-related quality of life outcomes.

- 4.1.19 The meta-analysis for risedronate relative to placebo, carried out by the Assessment Group, resulted in an RR of vertebral fracture of 0.61 (95% CI 0.50 to 0.75, three RCTs, n = 2301), an RR of hip fracture of 0.74 (95% CI 0.59 to 0.93, three RCTs, n = 11,770), an RR of wrist fracture of 0.68 (95% CI 0.43 to 1.08, two RCTs, n = 2439) and an RR for other non-vertebral fractures of 0.76 (95% CI 0.64 to 0.91, five RCTs, n = 12,399).
- 4.1.20 In all the studies, rates of gastrointestinal adverse events were similar in the risedronate and placebo groups.
- 4.1.21 Prescription-event monitoring studies in patients for whom risedronate was prescribed (n = 13,643) by GPs in England suggested a high incidence of dyspepsia, particularly in the first month of treatment. Consultations for dyspepsia ranged from 26.9 per 1000 patient-months in the first month of treatment to 8.1 per 1000 patient-months in months 2 to 6.

*Alendronate and risedronate: meta-analysis*

- 4.1.22 A meta-analysis of pooled data from the alendronate and risedronate studies, carried out by SchARR in 2006, resulted in an RR of vertebral fracture of 0.58 (95% CI 0.51 to 0.67, seven RCTs, n = 9340), an RR of hip fracture of 0.71 (95% CI 0.58 to 0.87, six RCTs, n = 19,233), an RR of wrist fracture of 0.69 (95% CI 0.45 to 1.05, six RCTs, n = 1037), and an RR for other non-vertebral fractures of 0.78 (95% CI 0.69 to 0.88, 11 RCTs, n = 22,372).

*Raloxifene*

- 4.1.23 Three RCTs of raloxifene in postmenopausal women were identified, but only two were included in the Assessment Group's meta-analysis: the largest study (the 'Multiple outcomes of raloxifene evaluation' [MORE] study) was carried out in women with osteoporosis, of whom 37% had a vertebral fracture at entry, and a smaller study was conducted in women with established osteoporosis. Both compared raloxifene with placebo (in both studies, women in both arms received calcium and vitamin D). Both studies examined raloxifene at dosages of 60 mg/day (the dosage specified in the UK marketing authorisation for treatment of postmenopausal osteoporosis) and 120 mg/day. Neither reported on health-related quality of life outcomes. The mean age of women in the studies was 67–68 years. The MORE study was extended further to assess fracture, breast cancer, and cardiovascular and uterine safety outcomes. A third study examined the additive effect of raloxifene compared with placebo in women with a femoral neck T-score of  $-2$  SD or below, with or without prior fracture, who were also receiving fluoride, calcium and vitamin D. Because of the use of fluoride as a co-intervention, these results were not included in the Assessment Group's meta-analysis.
- 4.1.24 The meta-analysis for raloxifene relative to placebo, carried out by the Assessment Group, resulted in an RR of vertebral fracture of 0.65 (95% CI 0.53 to 0.79, one RCT, n = 4551), an RR of hip fracture of 1.13 (95% CI 0.66 to 1.96, two RCTs, n = 6971), an RR of wrist fracture of 0.89 (95% CI 0.68 to 1.15, one RCT, n = 6828), and an RR for other non-vertebral fractures of 0.92 (95% CI 0.79 to 1.07, one RCT, n = 6828).
- 4.1.25 The most serious adverse effect associated with raloxifene was the approximately three-fold increased risk of VTE. Statistically significantly higher incidences of hot flushes, arthralgia, dizziness,

leg cramps, influenza-like symptoms, endometrial cavity fluid, peripheral oedema and worsening diabetes were also found with raloxifene compared with placebo. The impact of raloxifene on cardiovascular disease is unclear, but there is evidence that it lowers serum concentrations of fibrinogen as well as both total and low-density lipoprotein (LDL) cholesterol, levels (that is, serum concentrations) without increasing high-density lipoprotein (HDL) cholesterol.

- 4.1.26 The MORE study shows that raloxifene protects against breast cancer, with the RR at 4 years for all types of breast cancer reported as 0.38 (95% CI 0.24 to 0.58), and that for invasive breast cancer as 0.28 (95% CI 0.17 to 0.46).

#### *Strontium ranelate*

- 4.1.27 Three RCTs of strontium ranelate in postmenopausal women were identified: one study in women with osteoporosis and two in women with osteoporosis or established osteoporosis. All three studies compared strontium ranelate with placebo, and provided calcium and vitamin D supplementation to ensure an adequate intake.
- 4.1.28 The Assessment Group reported the results of a published meta-analysis that gave an RR for vertebral fracture of 0.60 (95% CI 0.53 to 0.69, two RCTs, n = 6551) and an RR for all non-vertebral fractures (including wrist fracture) of 0.84 (95% CI 0.73 to 0.97, two RCTs, n = 6551). Efficacy in reducing the rate of hip fracture was established in one study; the RR for hip fracture in the whole study population was 0.85 (95% CI 0.61 to 1.19, one RCT, n = 4932). A post-hoc subgroup analysis in women over 74 years of age with a T-score of  $-2.4$  SD resulted in an RR for hip fracture of 0.64 (95% CI 0.41 to 0.98, one RCT, n = 1977).
- 4.1.29 In general, strontium ranelate was not associated with an increased risk of adverse effects and for the most part adverse effects were

mild and transient; nausea, diarrhoea and creatine kinase elevations were the most commonly reported. A serious adverse event associated with strontium ranelate treatment was an increased incidence (RR = 1.42) of VTE and pulmonary embolism. This finding has been investigated further with the extension of ongoing studies and by post-marketing surveillance.

- 4.1.30 One study published results on health-related quality of life outcomes. It reported that strontium ranelate had quality of life benefits compared with placebo, as assessed by the QUALIOST osteoporosis-specific questionnaire and by the general health perception score of the short form (SF)-36 general scale.

#### *Teriparatide*

- 4.1.31 Three RCTs of teriparatide in postmenopausal women were considered: one small study compared teriparatide with alendronate in women with osteoporosis (but was not targeted at women with fractures), and two were placebo-controlled (although study participants also received vitamin D either with calcium or with nutritional advice to ensure adequate calcium intake). The largest trial was conducted in women with established osteoporosis, and the other in women who either had established osteoporosis or had osteoporosis and had been receiving hormone replacement therapy for at least 2 years.
- 4.1.32 For vertebral fractures (using a 20% reduction in vertebral height as the fracture definition) and grouped non-vertebral fractures in women with established osteoporosis, the largest placebo-controlled RCT found RRs of 0.35 (95% CI 0.22 to 0.55) and 0.65 (95% CI 0.43 to 0.98), respectively, in favour of teriparatide. When considered separately, the study did not demonstrate that teriparatide prevents hip and wrist fractures in women with established osteoporosis (RR for hip fractures 0.5; 95% CI 0.09 to 2.73; RR for wrist fractures 0.54; 95% CI 0.22 to 1.35). In this

placebo-controlled trial, teriparatide reduced the incidence of new or worsened back pain reported as an adverse event.

- 4.1.33 Data from a follow-up observational study cited in the manufacturer's submission (published in abstract form or available as an unpublished manuscript only) suggest that 18 months after the end of treatment with teriparatide there was a 41% reduction in vertebral fracture risk compared with placebo ( $p = 0.004$ ). Further data from the same study 31 months after the end of treatment with teriparatide suggest that proportionally fewer women who had received teriparatide reported non-vertebral fractures compared with those who had received placebo (13.3% in the placebo group; 8.5% in the 20 micrograms/day teriparatide group; 7.3% in the 40 micrograms/day teriparatide group;  $p = 0.03$  for both treatment groups versus placebo). No information was given on vertebral fractures for the 31-month follow-up.
- 4.1.34 The study comparing 40 micrograms/day teriparatide (twice the dose specified in the marketing authorisation) with 10 mg/day alendronate found an RR of non-vertebral fracture in women with osteoporosis of 0.30 (95% CI 0.09 to 1.05). The study did not provide data on vertebral fractures. Back pain was reported less frequently in the teriparatide group (6% versus 19%,  $p = 0.012$ ).
- 4.1.35 Nausea and headaches occurred more frequently with 40 micrograms/day teriparatide in the main placebo-controlled trial. In the smaller placebo-controlled trial, a proportion of women taking teriparatide were reported to suffer mild discomfort at the injection site. A systematic review of parathyroid hormone reported that treatment in a small proportion of women was associated with hypercalcaemia.

## Persistence and compliance

### *Bisphosphonates*

- 4.1.36 Data from 14 RCTs indicated that between 81% and 100% of patients persisted with bisphosphonates in the first year of treatment, with lower rates of persistence of between 51% and 89% in the third year of treatment (eight RCTs).
- 4.1.37 A prescription-event monitoring study of patients for whom alendronate was prescribed (n = 11,916) by GPs in England indicated that 24% discontinued treatment within 1 year. In a similar study of patients for whom risedronate was prescribed (n = 11,742) in primary care in England, 30% appeared to have discontinued treatment within 6 months. In another 12 studies reviewed, persistence at 1 year ranged from 16% to 90%.

### *Raloxifene*

- 4.1.38 Paid claims data from the USA suggested that only 18% of women starting raloxifene treatment continued to take their medication uninterrupted, and an investigation of a pharmacy prescription database indicated that only 44% were continuing treatment at the end of year 2.

### *Strontium ranelate*

- 4.1.39 Compliance data were reported for two RCTs of strontium ranelate and were similar in the strontium ranelate and placebo arms (ranging from 83% to 93%) at up to 3 years.

### *Teriparatide*

- 4.1.40 The main placebo-controlled RCT reported that adherence with injections varied from 79% to 83% and that there were no statistically significant differences between the teriparatide and placebo groups. The smaller placebo-controlled trial found that, after 3 years, 78% of women receiving teriparatide completed treatment, compared with 100% on placebo.

**Acid-suppressive medication and fracture risk**

- 4.1.41 Two cohort and two case–control studies reported on a potential relationship between acid-suppressive medication (proton pump inhibitors or histamine H2 receptor antagonists) and fracture risk. One of the case–control studies, which used the UK General Practice Research Database (GPRD), found that 1 year or more of acid-suppressive medication was associated with an increase in fracture risk. The other case–control study reported a reduction of fracture risk associated with use of histamine H2 receptor antagonists, and that use of other acid-suppressive medication might increase fracture risk. Both studies, however, were unable to demonstrate convincingly that fracture risk was independent of underlying disease that might determine differences in fracture risk.
- 4.1.42 A prospective cohort study excluded women taking medication for fracture prevention and reported an increase in non-vertebral fracture in those taking acid-suppressive medication compared with those who were not. Findings appeared similar for users of proton pump inhibitors or histamine H2 receptor antagonists, but differences in fracture risk were not statistically significant for those using proton pump inhibitors compared with those not using acid-suppressive medication. One large retrospective cohort study using the UK GPRD compared women taking acid-suppressive medication plus bisphosphonates with those taking bisphosphonates alone. This GPRD study reported an increase in fracture risk for some fracture sites with concomitant use of acid-suppressive medication and bisphosphonates, but a reduction in risk for other fracture sites. The information on patients included in this GPRD study was incomplete and details of adjustments for confounders were not reported. The two cohort studies were not fully published, and their analysis may have been prone to confounding.



## 4.2 *Cost effectiveness*

### Manufacturers' models

- 4.2.1 For proprietary alendronate, compared with no treatment, the manufacturer's model provided an incremental cost-effectiveness ratio (ICER) of £3135 per quality-adjusted life year (QALY) gained for 70-year-old women with a T-score below  $-1.6$  SD. The manufacturer's results were more favourable than the results of Assessment Group's 2003 model. This could be because the manufacturer's model was not adjusted for baseline fracture prevalence, or because it used different utilities for vertebral fractures, different efficacy data, different risk groups, and a longer time horizon.
- 4.2.2 For etidronate, compared with no treatment, the manufacturer's model provided an ICER of £18,634 per QALY gained for 70-year-old women with a T-score below  $-2.5$  SD. The manufacturer's model included morphometric vertebral fractures and corticosteroid use as risk factors for further fractures. It is unclear whether the manufacturer's ICER was for women with or without a prior osteoporotic fragility fracture.
- 4.2.3 For risedronate, compared with no treatment, the manufacturer provided data from two models. The ICER derived from the manufacturer's own model was £577 per QALY gained for women aged 74 years. In the second model provided by the manufacturer, which was commissioned from an external body, the ICER was higher, varying from £35,800 per QALY gained in women aged 60 years to £4800 per QALY gained in women aged 80 years, for women with a prior vertebral osteoporotic fragility fracture and a T-score of  $-2.5$  SD. For women at slightly higher risk of fracture, the ICERs were £18,600 per QALY gained or less for all age groups. The ICER calculated using the manufacturer's own model was difficult to verify from the information given. The ICERs generated

by the second model were more consistent with the figures provided by the Assessment Group's 2003 model, although they did differ somewhat. This may be because of different cost and RR inputs.

- 4.2.4 For raloxifene, compared with no treatment, the manufacturer provided data for different age groups and different risk levels. All of the analyses included the breast cancer benefits. It was not clear how the different risk levels were defined. The ICERs ranged from £12,000 to £22,000 per QALY gained and were slightly more favourable than the Assessment Group's 2003 analysis, even when the Assessment Group included the breast cancer benefits. In the Assessment Group's 2003 model, the RR for the breast cancer effect was higher (0.38) than the RR for invasive breast cancer used in the manufacturer's model (0.28), and the breast cancer risk was adjusted for the association between low BMD and decreased risk of breast cancer. Additionally, the manufacturer's model was not adjusted for baseline fracture prevalence, and included different utilities for vertebral fractures, different efficacy data, different risk groups, and a longer time horizon than the Assessment Group's model.
- 4.2.5 For strontium ranelate, compared with no treatment, the manufacturer provided two models: one developed in-house and the other commissioned from an external body. The first model showed that, for women aged over 75 years with previous fractures and a T-score of  $-2.5$  SD, strontium ranelate was cost-effective at a maximum acceptable incremental cost-effectiveness ratio of £30,000 per QALY gained. The results of this model were comparable with those generated by the Assessment Group's 2005 model. The second model resulted in an ICER of £6341 per QALY gained for 70-year-old women with a previous vertebral fracture and a T-score of  $-2.5$  SD, decreasing to £5002 per QALY gained in

women aged 80 years. The manufacturer's results were more favourable than the Assessment Group's 2005 results because different modelling assumptions were used. For example, fewer health-state transition possibilities were incorporated. Compared with the Assessment Group's model, the manufacturer's model used more favourable efficacy data for hip fracture from a subgroup of women aged over 74 years, and slightly more favourable efficacy data for wrist and proximal humerus fracture. Higher hip-fracture costs were used in the manufacturer's model.

- 4.2.6 For teriparatide, compared with no treatment, the manufacturer provided ICERs for women aged 69 years. For women with fractures that had occurred more than 6 months previously (historical fracture), the ICER was £35,400 per QALY gained and for women with a more recent fracture the ICER was £28,863 per QALY gained. The manufacturer supplied additional economic analyses with ICERs of £18,845 and £12,106 per QALY gained for historical and recent fracture, respectively, based on changes to the assumptions of sustained efficacy for non-vertebral fractures and of the RR for specific risk groups. The manufacturer's model and the Assessment Group's 2003 model differed in a number of assumptions. The manufacturer's model was not adjusted for baseline fracture prevalence and used different utilities. The Assessment Group's 2003 model used more favourable assumptions on the duration of sustained efficacy after the end of treatment.

### **The Assessment Group's model**

- 4.2.7 The Assessment Group provided a cost–utility model with two components (described in detail in the 2005 Strontium Ranelate Assessment Report). As a first step, the model calculated absolute fracture risk from the epidemiological literature on a number of independent clinical risk factors. These data were prepared under

the auspices of the WHO and were provided for this appraisal under an academic-in-confidence agreement. As a second step, the model applied RR reductions for fracture taken from the meta-analysis described in section 4.1.22. A single estimate of efficacy was used for alendronate and risedronate based on pooled data for these two drugs. Following advice from the Osteoporosis Guideline Development Group (see [www.nice.org.uk](http://www.nice.org.uk)), it was assumed that RRs remained constant across all ages, T-scores and fracture status. The most recent analyses carried out by SchARR were based on the price of non-proprietary alendronate in February 2008 (£53.56 per year for once-weekly 70 mg tablets; £108.20 per year for daily 10 mg tablets).

- 4.2.8 All osteoporotic fragility fractures in women aged 50 years or older were included in the modelling. The RR for hip fracture was assumed to apply also to pelvis and other femoral fractures. The RR for non-vertebral fracture was assumed to apply also to proximal humerus, rib, sternum, scapula, tibia, fibula and wrist fractures. Where confidence intervals for RRs spanned unity, it was assumed that there was no effect of treatment, except in the case of strontium ranelate in a subgroup of older women. In this case, an RR of 0.85 for hip fracture was used to acknowledge an effect reported in a subgroup of the study. The model used UK-specific epidemiological data on femoral neck BMD.
- 4.2.9 The model assumed an initial utility in the year of fracture and a higher utility in subsequent years. The time horizon for predicting morbidity was 10 years, consisting of 5 years of treatment with sustained efficacy plus 5 years of linear decline to no effect. However, treatment-related decreases in mortality rate extended beyond the 10-year time horizon. For this, the life expectancy for a woman at the threshold T-score for osteoporosis was calculated from standard life tables, and any increase in mortality rate due to

fracture would continue until death or an age of 110 years. In the base case, vertebral-fracture utility was assumed to be lower than hip-fracture utility, and a sensitivity analysis was carried out in which the utility for vertebral fracture was assumed to be the same as that for hip fracture. The percentage of women assumed to move from community living to a nursing home following a hip fracture increased with increasing age. An age-dependent gradient of hip-fracture risk was used, and an association between vertebral or proximal humerus fracture and increased mortality in women with osteoporosis was included. No follow-up BMD scans were included in the model; this reflects current clinical practice in the UK.

- 4.2.10 The model included an assumption about the costs and disutility associated with treatment-related side effects for all drugs, based on the findings of prescription-event monitoring studies in patients treated with alendronate. For the base case, the model assumed 50% persistence with treatment. In addition to the base case, the Assessment Group undertook a number of sensitivity analyses using alternative assumptions including: persistence with treatment (25% or 75% at 5 years); reduction in the efficacy of the drugs at reducing the risk of fracture associated with risk factors other than age, prior fracture and low BMD to 0% or 50% (with a consequent upward adjustment of the RR for the risk factors of age, prior fracture and low BMD); disutility of vertebral fracture; updated fracture costs; and the disutility and costs of treatment-related side effects. It was assumed that women who experience bisphosphonate-related side effects had 91% of the utility of women who do not have such side effects. In the base case analysis for all the drugs under consideration this was applied to 2.35% of women in the first treatment month and 0.35% of women thereafter and, in sensitivity analyses for bisphosphonates, to 24% of women in the first treatment month and 3.5% of women

thereafter. In the case of strontium ranelate, the effect on VTE was not included in the model. Discount rates of 6% per year for costs and 1.5% per year for health benefits were applied, in accordance with NICE methods relevant to this appraisal.

- 4.2.11 For raloxifene, 4-year follow-up data from the MORE study were used, and it was assumed that women with low BMD have a lower breast cancer risk than women with normal BMD. The cost effectiveness was modelled excluding the breast cancer benefit, the risk of VTE and the effect on cardiovascular events.
- 4.2.12 The independent clinical risk factors for fracture used in the model were based on the data prepared under the auspices of the WHO (see section 4.2.7) and included BMI, prior fracture, previous or current use of corticosteroids, parental history of fracture, current smoking, alcohol intake of more than 2 units per day, and rheumatoid arthritis. The study provided prevalence data for the different risk factors, and risk ratios for hip fracture and osteoporotic fracture for each risk factor, including T-score and age. Using these risk ratios, absolute risk of fracture was calculated.
- 4.2.13 The estimates of cost effectiveness were generated for different levels of absolute risk derived from a large number of combinations of T-score (in bands 0.5 SD wide), age and number of independent clinical risk factors for fracture. For practical reasons relating to the number of potential combinations, single-point RRs of fracture, calculated from the log-normal efficacy distributions, were used in the model. Results were presented for population groups categorised according to age, T-score and number of independent clinical risk factors.
- 4.2.14 Women with a fracture who present to clinicians require a DXA scan for osteoporosis to be established. Therefore, the Assessment Group also estimated the impact of DXA scanning on the cost

effectiveness of the drugs. This required both a calculation of the ICER for treatment and a calculation of the distribution of risk assessment cost over the population who would benefit from treatment. A net-benefit approach was used to do this. The net-benefit approach is analogous to the more traditional cost per QALY gained approach, but also requires a value of willingness to pay (WTP) for an additional QALY gained. For the calculation of the net benefit of an intervention, the WTP is first multiplied by the incremental QALY gained associated with the intervention, then the incremental cost associated with the intervention is subtracted. For this appraisal, the total net benefit for each age group and DXA scanning approach was calculated by subtracting the cost of DXA scanning from the net benefit of treating all women who can be treated cost effectively.

4.2.15 A stepped net-benefit approach was used to estimate, in reverse order, the cost effectiveness of risk assessment, DXA scanning and treatment of women with a prior fracture. Two WTP values, £20,000 or £30,000 per QALY gained, were applied in the modelling.

- Step 1. ICERs for treatment versus no treatment were calculated for each intervention for various combinations of age, T-score and number of independent clinical risk factors for fracture (see section 4.2.12). The net benefit of treatment per woman was calculated using the following formula:

Net benefit = £30,000 (or £20,000) × incremental QALYs – incremental costs.

For women for whom the ICER for treatment was more than £30,000 (or £20,000) per QALY gained, the net benefit was set to zero.

- Step 2. The net benefit per woman was multiplied by the number of women in the population estimated to fall within each

combination of age, T score and number of independent clinical risk factors for fracture (based on the data used to develop the algorithm prepared for the WHO). The net benefits for each group were then added together to give a total net benefit of treatment for women with no, one, two or three independent clinical risk factors within each age group.

- Step 3. The cost of DXA scanning all of the women in each age/independent clinical risk factor group was subtracted from the net benefit of treatment for that group (calculated as described in step 2). This provides the net benefit of treatment and DXA scanning for the group, assuming that the number of independent clinical risk factors is known. A positive net benefit indicates that DXA scanning of women in that age/independent clinical risk factor group and treating those groups of women in whom the ICER for treatment is £30,000 (or £20,000) or less provides an ICER for the entire strategy of less than £30,000 (or £20,000) per QALY gained.
- Step 4. When the resulting values of net benefit of treatment and scanning were negative they were set to zero. For each age group, the total net benefit of scanning and treatment was calculated by adding together the net benefits for each age/independent clinical risk factor group. The cost of opportunistic assessment for all women in this age group was then subtracted to give the net benefit of risk assessment, scanning and treatment. A positive net benefit indicates an ICER of less than £30,000 (or £20,000) per QALY gained for risk assessment, DXA scanning and treating women (at a specific T-score related to the ICER for treatment only) of that particular group. Cost per QALY gained data were presented for each strategy.



*The Assessment Group's model: results for alendronate*

- 4.2.16 First, the Assessment Group calculated ICERs (cost per QALY gained for alendronate compared with no treatment) without identification costs for all combinations of age, T-score and number of independent clinical risk factors for fracture. The cost per QALY gained, compared with no treatment, became more favourable with increasing age and number of independent clinical risk factors, and decreasing T-score (that is, with increasing annual absolute risk of fracture).
- 4.2.17 Then, the Assessment Group presented the results of the economic analyses in the form of identification and treatment strategies (based on age, T-score and number of independent clinical risk factors for fracture) that resulted in an ICER of £30,000 or less (cost per QALY gained compared with no treatment). The analyses shown below included the following assumptions: persistence at 5 years set to 50%; the efficacy of bisphosphonates on fracture risks associated with factors other than age, BMD and prior fracture status set to 50% of that observed for the total population in the trials (with a consequent upward adjustment of the RR associated with age, BMD and prior fracture); costs set to health resource group values including home-help costs; utility multiplier associated with vertebral fracture set to 0.792 in the first year of fracture and 0.909 in subsequent years (as for hip fracture); costs of bisphosphonate-related gastrointestinal symptoms incurred over 5 years; utility multiplier associated with bisphosphonate-related gastrointestinal symptoms set to 0.91 (included utility losses for non-compliant patients); and alendronate at a cost of £53.56 or £108.20 per year.
- 4.2.18 For alendronate priced at £53.56 per year (once-weekly treatment), and when assuming that 24% of women in the first treatment month

and 3.5% of women thereafter experienced bisphosphonate-related side effects, the model produced the following results:

- A strategy of risk assessment, DXA scanning and treatment with alendronate resulted in an ICER of less than £30,000 per QALY gained for all women aged 55 years or older with confirmed osteoporosis (that is, a T-score of  $-2.5$  SD), and for postmenopausal women aged 50–54 years with confirmed osteoporosis and two independent clinical risk factors for fracture.

4.2.19 In a sensitivity analysis for alendronate priced at £53.56 per year (with other assumptions as in section 4.2.17 and 4.2.18), acid-suppressive medication was assumed to affect fracture risk. The data inputs for this were taken from one GPRD study (see section 4.1.41) and represent the midpoint values pooled for patients using acid-suppressive medication. This sensitivity analysis produced the following results:

- A strategy of risk assessment, DXA scanning and treatment with alendronate in women younger than 55 years resulted in an ICER of more than £30,000 per QALY gained.
- A strategy of risk assessment, DXA scanning and treatment with alendronate resulted in an ICER of less than £30,000 per QALY gained for all women aged 65 years or older with confirmed osteoporosis (that is, a T-score of  $-2.5$  SD or below), for postmenopausal women aged 60–64 years with confirmed osteoporosis and one independent clinical risk factor for fracture, and postmenopausal women aged 55–59 years with confirmed osteoporosis and two independent clinical risk factors for fracture.

The ICER for treatment with alendronate (but excluding identification costs) for a woman aged 60–64 years with a T-score

of  $-2.5$  SD (using the assumptions described in sections 4.2.17 and 4.2.18) was £9005 per QALY gained without acid-suppressive medication and £21,656 per QALY gained with acid-suppressive medication. If this woman had an independent clinical risk factor for fracture, the ICERs would be £3969 per QALY gained without and £12,250 per QALY gained with acid-suppressive medication.

4.2.20 For alendronate priced at £108.20 per year (daily treatment), and when assuming that 24% of women were experiencing bisphosphonate-related side effects in the first treatment month and 3.5% of women thereafter, the model produced the following results:

- A strategy of risk assessment, DXA scanning and treatment with alendronate in women younger than 55 years resulted in an ICER of more than £30,000 per QALY gained.
- A strategy of risk assessment, DXA scanning and treatment with alendronate resulted in an ICER of less than £30,000 per QALY gained for women aged 65 years or older with confirmed osteoporosis (that is a T-score of  $-2.5$  SD or below), for postmenopausal women aged 60–64 years with confirmed osteoporosis and one independent clinical risk factor for fracture, and for postmenopausal women aged 55–59 years with confirmed osteoporosis and two independent clinical risk factors.

*The Assessment Group's model: results for other drugs*

4.2.21 Risedronate, raloxifene and strontium ranelate were dominated by alendronate (based on the price of £53.56 per year for alendronate); that is, these three drugs have a higher acquisition cost than alendronate, but are not more efficacious. Analyses were conducted as for alendronate (see section 4.2.17). For risedronate, base case assumptions for bisphosphonate-related side effects were modelled; that is 2.35% of women in the first treatment month and 0.35% thereafter experienced side effects (see section 4.2.10).

In addition a sensitivity analysis was performed, using the assumption that 24% of women in the first treatment month and 3.5% of women thereafter experienced bisphosphonate-related side effects. For raloxifene and strontium ranelate, base-case assumptions for side effects were used. In previous economic modelling and before the most recent price reduction for non-proprietary alendronate, etidronate's cost effectiveness was comparable to non-proprietary alendronate, but the calculations were based on a weaker clinical evidence base than for alendronate. Therefore the modelling for etidronate was not updated after the most recent price reduction for alendronate.

- 4.2.22 For risedronate, raloxifene, strontium ranelate and teriparatide, additional analyses were conducted to explore identification and treatment strategies that could be cost effective for these interventions when compared with no intervention. All results showed less favourable cost effectiveness than non-proprietary alendronate. For example, for women aged 55–59 years with an independent clinical risk factor for fracture, the ICERs (without considering costs related to risk assessment and DXA scanning) for risedronate and strontium ranelate (each compared with no treatment) were more than £40,000 and £55,000 per QALY gained, respectively. For these two groups of women, treatment with weekly non-proprietary alendronate, including risk assessment and DXA scanning costs, resulted in an ICER of less than £30,000 per QALY gained.

*The Assessment Group's model: results for other drugs in second-line use*

- 4.2.23 Further analyses were carried out assuming second-line use, that is, costs for risk assessment or DXA scanning were excluded because BMD was assumed to be known from the first-line management.

4.2.24 In the economic modelling carried out for this appraisal in 2006, lower ages and higher T-scores resulted in ICERs of less than £30,000 per QALY gained for etidronate compared with risedronate; that is, etidronate was more cost effective than risedronate. Because of the concerns expressed about the weaker clinical evidence base for etidronate, the modelling for this bisphosphonate was not updated.

4.2.25 For risedronate in second-line use, when assuming that 2.35% of women in the first treatment month and 0.35% of women thereafter experienced bisphosphonate-related side effects, the model produced the following results:

- Treatment with risedronate in women who have the combinations of T-score, age and number of independent clinical risk factors for fracture indicated in the table below resulted in an ICER of less than £30,000 per QALY gained. Including women aged 50–54 years with no independent clinical risk factors for fracture increased the ICER to more than £30,000 per QALY gained.

**T-scores (SD) at (or below) which risedronate in second-line use resulted in an ICER of less than £30,000 per QALY gained**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
50–54	– <sup>a</sup>	–3.0	–2.5
55–59	–3.0	–3.0	–2.5
60–64	–3.0	–3.0	–2.5
65–69	–3.0	–2.5	–2.0 <sup>b</sup>
70–74	–2.0 <sup>b</sup>	–2.0 <sup>b</sup>	–1.0 <sup>b</sup>
75 or older	–2.0 <sup>b</sup>	–1.5 <sup>b</sup>	–0.5 <sup>b</sup>

<sup>a</sup> ICER more than £30,000 per QALY gained

<sup>b</sup> Women with osteopenia are not included in the guidance (see sections 1 and 4.3.8)

4.2.26 For strontium ranelate in second-line use, the model produced the following results:

- Treatment with strontium ranelate in women who have the combinations of T-score, age and number of independent clinical risk factors for fracture indicated in the table below resulted in an ICER of less than £30,000 per QALY gained. Including women aged 50–54 years with no independent clinical risk factors for fracture increased the ICER to more than £30,000 per QALY gained.

**T-scores (SD) at (or below) which strontium ranelate in second-line use resulted in an ICER of less than £30,000 per QALY gained**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
50–54	– <sup>a</sup>	–3.5	–3.5
55–59	–4.0	–3.5	–3.5
60–64	–4.0	–3.5	–3.5
65–69	–4.0	–3.5	–3.0
70–74	–3.0	–3.0	–2.0 <sup>b</sup>
75 or older	–3.0	–2.5	–2.0 <sup>b</sup>

<sup>a</sup> ICER more than £30,000 per QALY gained

<sup>b</sup> Women with osteopenia are not included in the guidance (see section 1 and 4.3.8)

4.2.27 For raloxifene in second-line use, using base-case assumptions on side effects, the model produced the following results:

- Treatment with raloxifene in women younger than 70 years resulted in an ICER of more than £30,000 per QALY gained.
- Treatment with raloxifene in women who have the combinations of T-score, age and number of independent clinical risk factors for fracture indicated in the table below resulted in an ICER of less than £30,000 per QALY gained.

**T-scores (SD) at (or below) which raloxifene in second-line use resulted in an ICER of less than £30,000 per QALY gained**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
70–74	-5.0	-4.5	-4.0
75 or older	-4.5	-4.0	-3.5

4.2.28 For teriparatide, the model produced the following results.

- Treatment with teriparatide in women who have the combinations of T-score, age and number of independent clinical risk factors for fracture indicated in the table below resulted in an ICER of less than £30,000 per QALY gained. Including women aged 50–54 years with no independent clinical risk factors for fracture would increase the ICER to more than £30,000 per QALY gained.

**T-scores (SD) at (or below) which teriparatide in second-line use resulted in an ICER of less than £30,000 per QALY gained**

Age (years)	Number of independent clinical risk factors for fracture (section 1.5)		
	0	1	2
50–54	– <sup>a</sup>	-4.0	-4.0
55–59	-4.5	-4.5	-4.0
60–64	-4.5	-4.5	-4.0
65–69	-5.0	-4.5	-4.5
70–74	-4.5	-4.5	-3.5
75 or older	-4.5	-4.0	-3.5

<sup>a</sup> ICER more than £30,000 per QALY gained

4.2.29 If it was assumed that acid-suppressive medication affects fracture risk, the ICER for treatment with risedronate (compared with no treatment, but excluding identification costs) for a woman aged 70 years with a T-score of –3 SD increased from £12,273 to £17,848 per QALY gained (using base-case assumptions about side effects). The corresponding ICER for strontium ranelate was

£28,026 per QALY gained compared with no treatment (using base-case assumptions about side effects). For a women aged 70 years with a T-score of  $-3.5$  SD and one independent clinical risk factor for fracture, the ICER for risedronate increased from £3028 to £7688 per QALY gained when acid-suppressive medication was assumed to affect fracture risk (using base-case assumptions about side effects). The corresponding ICER for strontium ranelate was £14,986 per QALY gained compared with no treatment (using base-case assumptions about side effects).

### **Consultee comments on the Assessment Group's economic model**

4.2.30 Following the outcome of the judicial review and the court ruling of March 2009, NICE was able to offer the Assessment Group's executable economic model for consultation. Consultees and commentators who requested the model and returned the necessary confidentiality undertakings received a CD-ROM containing the executable version of the economic model, a document with instructions for running the model and a pro-forma for commenting on the model. Comments on the Assessment Group's model were received from Servier Laboratories (the manufacturer of strontium ranelate), the Bone Research Society (BRS), the National Osteoporosis Society (NOS) and the Society for Endocrinology. Comments received from each of these consultees are summarised in the sections below.

4.2.31 These four consultees expressed the view that the documentation provided with the Assessment Group's model was insufficient, that the model supplied to them was incomplete and that some inputs could not be altered. They also stated that the application of the fracture risk algorithm developed under the auspices of the WHO could not be assessed. They felt that the model could not be validated and that its validity had not been demonstrated in documents made available during development of the guidance.



- 4.2.32 Servier commented that the fracture risks entered in the Assessment Group's model differed from estimates that Servier calculated using the FRAX fracture risk calculation tool (see section 4.3.37 for further information about the FRAX tool). Servier commented that mortality risk associated with clinical risk factors had been omitted from the model. In Servier's opinion, these differences called into question whether the WHO fracture risk algorithm had been applied correctly in the Assessment Group's model.
- 4.2.33 Other comments questioned the use of a fixed value for BMI in the model. Consultees commented that no clear explanation was provided of the rationale for the choice of BMI value, that a range of BMI values should have been used, and that the use of a fixed BMI value resulted in underestimation of the cost effectiveness of treatment for some women at risk of fracture.
- 4.2.34 Servier commented on the selection and weighting of the independent clinical risk factors for fracture used in the Assessment Group's model. Servier, BRS and NOS suggested that the risk associated with alcohol intake was incorrect in the model and that this would have adversely affected estimation of the cost effectiveness of treatment for women at risk of fracture. They suggested that a threshold alcohol intake of 3 or more units per day, as used in the FRAX fracture risk calculation tool, should have been applied. They also stated that the Assessment Group's model and the guidance were inconsistent with each other, and that these differences resulted in the risk of fracture being underestimated in the model. Servier also noted that the Assessment Group's model gave an equal weighting to each of the independent clinical risk factors for fracture. Servier suggested that this was a less precise approach than that used in the FRAX tool, which used different weightings (some higher and some lower than those in the

Assessment Group's model) for each fracture risk for specific risk factors. Servier stated that the FRAX tool assesses fracture risk and cost effectiveness more accurately and 'deals more fairly' with variation between women at risk of fracture. Servier also noted that one of the risk multipliers for fracture risk included in the Assessment Group's model was not consistent with that given in the assessment report.

- 4.2.35 Servier and NOS noted that the Assessment Group's model had a time horizon limited to 10 years and criticised how mortality beyond 10 years had been taken into account in the economic evaluation. Servier expressed the view that, as a consequence of this, the model was inaccurate and underestimated the cost effectiveness of treatment for women at risk of fracture. Servier also identified two values ('wristbonusat2.5' and 'phbonusat2.5', related to QALY calculations) that were included in the model, but not described in assessment reports.
- 4.2.36 Servier commented that using the same disutility for side effects associated with strontium ranelate and bisphosphonates was not correct, as the side effects of strontium ranelate are different from those of the bisphosphonates.
- 4.2.37 BRS and NOS thought that the proportion of women with low BMD in England and Wales was substantially underestimated in the Assessment Group's model. These consultees were also concerned that although both smoking and previous or current glucocorticoid (corticosteroid) use had been included as additional independent clinical risk factors for fracture in the Assessment Group's model, they were not defined as risk factors in the recommendations (see section 1.5). In addition, both consultees thought that interactions between several clinical risk factors were not incorporated in the model, thereby reducing the cost

effectiveness of treatment for women at risk of fracture, especially younger women.

- 4.2.38 All four consultees commented on elements of the Assessment Group's economic evaluation that had been considered and agreed by the Appraisal Committee before it directed the Assessment Group to develop the economic model using specific assumptions. These Committee-directed assumptions included the compliance rate, costs associated with fracture, utility values used for vertebral fracture, and the strategy for identifying women at high risk. Servier also commented on the discount rates used in the model.
- 4.2.39 Servier reported that it had prepared a 'comparative' model which was run using assumptions similar to those in the Assessment Group's model. This model was referred to in a report to support the mathematical foundation of revised analyses discussed as part of Servier's comments on the Assessment Group's model. This report was made available to the Appraisal Committee to inform its consideration of comments by Servier on the DSU report (see below and section 4.3).

**Decision Support Unit (DSU) report on consultee comments on the Assessment Group's economic model**

- 4.2.40 The DSU was commissioned to review the comments from consultees on the Assessment Group's executable economic model and report to the Appraisal Committee. The DSU considered issues that were relevant to the economic model. Key issues were grouped under the common themes of model transparency and ability to assess its validity, methodology (approach) and model inputs.
- 4.2.41 The DSU assessed comments on the transparency and validity of the Assessment Group's model. With regard to the consultees' observation that some model inputs were fixed and that in their

view the model provided for consultation was incomplete and not fully executable, the DSU confirmed that certain inputs were intentionally fixed and the ability to alter these inputs was not a feature of the model or necessary for some for some parameters with minimal uncertainty that are commonly fixed in other economic models. In response to comments on the consultees' inability to assess the application of the WHO algorithm, the DSU explained that the WHO algorithm itself was not embedded within the model. The DSU confirmed that absolute fracture risks were correctly calculated using the WHO algorithm before being entered into the model. The DSU noted that documentation had been provided to consultees in the form of publicly available reports and peer-reviewed manuscripts produced by the Assessment Group, and that instructions on the operation of the Assessment Group's model were also offered to consultees and commentators.

4.2.42 With regard to comments on the modelling approach adopted in the Assessment Group's model, the DSU responded by confirming that alcohol consumption of more than 2 units per day was included in the model, and that the coefficients used in the model were consistent with the WHO algorithm (as supplied to the Assessment Group at the time the model was developed). The DSU also explored how the model considered corticosteroid-related fracture risk, and confirmed that corticosteroid use was included in the model and that the coefficient used for this risk factor was consistent with that calculated using the WHO algorithm. The DSU noted that the fracture risk of women using corticosteroids would have contributed to the overall fracture risk of the whole modelled population and thereby reduced the ICER associated with the treatment of all women at risk of fracture.

4.2.43 The DSU confirmed that each clinical risk factor for fracture was given equal weighting in the model. In response to consultee

comments expressing the view that this was a less precise approach than that used in the FRAX tool, the DSU noted two points. Firstly, no individual risk calculation tool was publicly available when the model was developed. Secondly, the DSU referred to the 2005 Strontium Ranelate Assessment Report, which compared suggested treatment thresholds (combinations of age, T-score and number of independent clinical risk factors for fracture) from the Assessment Group's model with treatment thresholds indicated by absolute fracture risk. The DSU suggested that the use of absolute fracture risk alone did not accurately predict cost effectiveness, and therefore would not provide a robust basis for the Committee's decision-making.

4.2.44 Consultee comments on the modelling approach also addressed the time horizon and population data used and the grouping of age in 5-year bands. The DSU confirmed that the consequences of fracture were considered beyond 10 years and provided further explanation of the modelling approach. The DSU additionally undertook exploratory analyses of the impact of mortality after the 10-year time horizon and of incorporating mortality associated with vertebral fracture and proximal humerus fracture. They reported that the change in the results produced by the model was minimal when mortality risk beyond 10 years was doubled. The DSU also confirmed that UK epidemiological data from a study by Holt et al. were used in the Assessment Group's model, and undertook an exploratory analysis around the assumptions of the distribution of T-scores used in the model. For some age bands modelled, the T-scores did not follow a statistically normal distribution, but the DSU noted that the assumption of a normal distribution made it more likely that treatments for women at risk of fracture would be judged to be cost effective. The DSU considered a comment on the calculation of cost-effectiveness estimates averaged for the 5-year age bands implemented in the model. It disagreed with the

alternative suggested by the consultee and noted that the Committee had considered and agreed that initial identification by age band was a workable strategy for selecting women at risk of fracture in clinical practice. It also noted that alternative strategies (which did not use age bands) may in fact be more resource-consuming and less likely to be judged as cost effective.

4.2.45 The DSU reviewed consultee comments on inputs used in the Assessment Group's model. It confirmed that the WHO algorithm (as supplied) had been correctly implemented in the model to produce estimates of fracture risk for each T-score band. The DSU suggested that the differences in the estimates of fracture risk obtained using the FRAX fracture risk calculation tool and the Assessment Group's model did not necessarily suggest that the WHO algorithm had been incorrectly applied (see section 4.3.37), and that these differences could occur for a number of reasons. For example, the use of a midpoint age to represent an age band of 5 years could lead to differences in estimates of fracture risk. The DSU confirmed that no increase in mortality associated with clinical risk factors was used in the model. The DSU suggested that inclusion of such mortality effects would be likely to increase the ICERs for women with those clinical risk factors. The DSU explained that this is because fewer QALY benefits would accrue in the model for women who die of causes related to risk factors. In response to a further comment from Servier, the DSU agreed that, for women without clinical risk factors, the inclusion of these mortality effects in the model may have the opposite effect (that is, a decrease in ICERs). Therefore the overall effect of including the increased mortality associated with clinical risk factors would be small.

4.2.46 The DSU also confirmed that a fixed value for BMI of 26 kg/m<sup>2</sup> was used in the Assessment Group's model. This was the mean BMI in

the UK epidemiological dataset from the Holt study used in the model. In an exploratory analysis using the WHO algorithm, the DSU showed that a using BMI of 26 kg/m<sup>2</sup> resulted in higher estimated fracture risk than a BMI of 20 or 32 kg/m<sup>2</sup> when BMD is known, and this was confirmed by the estimates supplied by one consultee. The DSU suggested that the BMI value used in the model may favour treatment of women at risk of fracture compared with alternative BMI values. The DSU also pointed out that BMI is a weak predictor of fracture when BMD is known (as specified in the identification strategy in the guidance).

4.2.47 The DSU investigated the risk multipliers used for fracture risk in the Assessment Group's model and the consultee comment that interactions between clinical risk factors had been omitted. It confirmed that the risk multipliers used for fracture risk had been correctly calculated from the WHO algorithm and that all interactions between risk factors had been included. The DSU also noted that the inconsistency between one of the risk multipliers for fracture risk included in the Assessment Group's model compared with the assessment report was the result of a typographical error. Accordingly there was no impact on the results of the model.

4.2.48 The DSU did not respond in detail to comments on assumptions in the model that had already been documented and agreed by the Appraisal Committee and which were available to consultees and commentators earlier in the development of the appraisal guidance. The DSU did, however, list these issues in its report and cited where they had been considered by the Committee or had been available for comment during development of the guidance. Features of the economic evaluation previously discussed and agreed by the Committee included the following (the sections of this document where these points are covered are given in parentheses):

- discount rates used in model (4.2.10)
- treatment compliance (4.2.10)
- costs associated with fracture (4.2.17)
- strategy for identifying women at high risk of fracture (4.2.17)
- utility values used for vertebral fracture (4.2.17 and 4.3.13)
- equal disutility for the side effects of strontium ranelate and bisphosphonates (4.2.10)
- sensitivity analyses on disutility (4.3.15).

4.2.49 The DSU concluded that, in its view, adequate documentation on the Assessment Group's model had been provided for consultees. It highlighted that the WHO algorithm used to generate estimates of fracture risk was not integrated within the Assessment Group's model; rather, the fracture risks derived from the algorithm were entered into the model. Comparisons with fracture risks derived using the FRAX fracture risk calculation tool were made by several consultees on the basis that the WHO algorithm supplied to the Assessment Group and the FRAX tool are assumed to be identical. The DSU could not verify these analyses without access to the FRAX algorithm. The DSU agreed that some parameters in the Assessment Group's model were fixed. These included those with minimal uncertainty, as well as those that are commonly fixed in other economic models. Sensitivity analyses conducted by the DSU suggested that none of the consultees' suggestions relating to the modelling approach would lead to significant improvements in the cost effectiveness of treatment for women at risk of fracture. The DSU concluded that, in its view, no issues raised by consultees would either affect the validity of the Assessment Group's model or raise significant doubts about the appropriateness of using the model to inform deliberations of the Committee.



### **4.3 Consideration of the evidence**

- 4.3.1 The Appraisal Committee reviewed the data available on the clinical and cost effectiveness of alendronate, etidronate, risedronate, strontium ranelate, raloxifene and teriparatide, having considered evidence on the nature of osteoporosis and the value placed on the benefits of these drugs by women with the condition, those who represent them, and clinical specialists. It also considered the consultation comments received in response to the previous appraisal consultation documents, the extra analysis undertaken by SchARR in November 2006 and February 2008, and comments received from consultees and commentators after an appeal against an earlier final appraisal determination was upheld in December 2007. Following the outcome of a judicial review and court ruling in March 2009, the Committee considered the comments received from consultees after release of the Assessment Group's executable economic model, a report by the DSU reviewing these comments, and responses from the consultees to the DSU report. It also took into account the effective use of NHS resources.
- 4.3.2 The Committee considered the extent to which NICE technology appraisal 87 should be updated in the light of the introduction of a new drug (strontium ranelate), new pricing for alendronate and etidronate, and new cost-effectiveness modelling developed as part of the technology appraisal on primary prevention.
- 4.3.3 The Committee considered the clinical effectiveness data for the bisphosphonates alendronate, etidronate and risedronate, strontium ranelate, raloxifene and teriparatide. It noted that all these drugs have proven efficacy in reducing the incidence of vertebral fragility fractures in women with osteoporosis, but that there were differences between the drugs as to the degree of certainty that treatment results in a reduction in hip fracture

(considered a crucial goal in osteoporosis management). In the case of alendronate and risedronate, the Committee accepted that there was sufficiently robust evidence to suggest a reduction in hip-fracture risk. The Committee noted that the available RCTs for etidronate were of insufficient size to show statistically significant reductions in hip-fracture risk, but that observational data lent support to a reduction in hip-fracture risk.

- 4.3.4 The Committee noted that strontium ranelate was effective in preventing vertebral and non-vertebral fractures, and the drug resulted in a non-significant 15% reduction in hip-fracture risk. The Committee was also aware of the result of a post-hoc subgroup analysis showing a statistically significant reduction in the incidence of hip fractures in women over the age of 74 years who had a T-score of  $-2.4$  SD or below.
- 4.3.5 The Committee noted that the evidence for raloxifene showed an effect on risk of vertebral fractures, but did not show any effect on risk of hip fractures. In addition, there was evidence for a beneficial side effect of raloxifene on the incidence of breast cancer.
- 4.3.6 The Committee noted that teriparatide was effective in preventing vertebral and grouped non-vertebral fractures in women with osteoporosis who have had a fracture, compared with placebo. The Committee also considered the favourable findings for teriparatide from one head-to-head RCT of teriparatide and alendronate, and that it conferred relatively favourable back-pain relief. However, the Committee was concerned about the small size of the head-to-head study, the fact that the study was not targeted at women with fractures, and the high dose of teriparatide used. Therefore it considered that the evaluation of the overall advantages of teriparatide over bisphosphonates requires more research to establish relative clinical effectiveness.

- 4.3.7 The Committee did not consider it appropriate to include recommendations for women on long-term corticosteroid treatment because this group is at greatly increased risk of fracture, and therefore requires special consideration, particularly if they have had a prior fracture. The Appraisal Committee therefore felt that it would be disadvantageous for this group to be included in the current guidance. Recommendations for this group of women will be made within future guidance produced by NICE.
- 4.3.8 Recommendations for the treatment of women with osteopenia (T-score of between  $-1$  and  $-2.5$  SD below peak BMD) were not made for two reasons. Firstly, it was agreed after the scope was issued in 2002 that the outcome in this appraisal should be 'the prevention of osteoporotic fractures' and this has been understood by the Committee to be a fragility fracture experienced by a women with osteoporosis, not osteopenia. Secondly, not all of the drugs under appraisal have a UK marketing authorisation for treatment of women with osteopenia. Recommendations for these groups of women will be made within future guidance produced by NICE.
- 4.3.9 The Committee noted that fracture risk is clearly related to age, low BMD and previous fracture. The Committee accepted that most other risk factors (see sections 2.11 and 2.12) were likely to be associated with an increased fracture risk. The Committee was concerned that there was not sufficient evidence for a proven treatment effect on fracture risk related to risk factors other than low BMD, age and prior fracture. The Committee therefore concluded that preventative drug treatment should be targeted at women whose absolute risk of fracture is driven by low BMD and age, and that the recommendations should be made on the basis of age and BMD in the form of T-scores below which treatment is recommended.

### Cost-effectiveness modelling

- 4.3.10 The Committee acknowledged the efforts of the Assessment Group to build on the model used previously, particularly in using epidemiological data and a fracture risk algorithm developed under the auspices of the WHO to calculate transition probabilities and to model the identification approaches. The Committee concluded that the Assessment Group's model was likely to give the best estimates of cost effectiveness because it used data from a wide age range (age 50–75 years or older), and was updated to use all osteoporotic fracture sites, more recent utility values, prevalence and risk-factor data, and an adjusted prevalence of fractures in the average population. Although the Assessment Group's model considered a shorter time period (10 years for predicting morbidity, see section 4.2.9) than the manufacturers' models, the Committee thought that this was appropriate considering the age groups involved and the uncertainties around health effects over a longer period.
- 4.3.11 The Committee discussed the assumptions underpinning the economic modelling undertaken by the Assessment Group. It noted that the most recent modelling explored some of the uncertainties identified by the Committee surrounding the results of the previous modelling; these related to the costs and disutility associated with treatment-related side effects and to non-compliance or non-persistence with treatment in a proportion of patients. The Committee also noted the effect of the recent price reductions for non-proprietary alendronate (70 mg weekly and 10 mg daily doses) on the cost effectiveness of the drug.
- 4.3.12 The Committee considered the base-case assumptions and those used in the additional analyses. The Committee noted that the costs associated with fractures used in the base-case analysis were those used in the original assessment report developed in

2003 and considered that these were likely to be outdated. The Committee agreed that costs based on health resource groups, including home-help costs, were likely to provide the most accurate reflection of the cost of fractures to the NHS and personal social services, and it decided to incorporate these costs into the base-case analysis.

4.3.13 The Committee considered the utility multiplier used in the base-case analysis for the first year after a vertebral fracture and noted that it was based on a hospitalised patient group and not on a typical group of patients with vertebral fractures. Consequently it was considerably lower than the utility value modelled for a hip fracture. Although the Committee acknowledged that vertebral fracture can lead to greatly reduced quality of life, it considered that its true value would not greatly outweigh the utility decrement associated with a hip fracture. Therefore, the Committee considered it reasonable to assume that the disutility in the first year after a vertebral fracture was equivalent to the disutility in the first year after a hip fracture and decided to include this assumption in the base-case analysis.

4.3.14 The Committee was not persuaded that the drugs under consideration had been unequivocally shown to reduce fracture risk that was attributable to independent clinical risk factors not mediated through low BMD and age. The Committee concluded that the uncertainty surrounding the efficacy of the drugs on independent clinical risk factors not mediated through low BMD and age should be factored into its decision-making by using an analysis that assumed 50% efficacy of the drugs on fractures associated with independent clinical risk factors other than age and low BMD. Although the Committee recognised that 50% was necessarily an arbitrary figure, the use of either 0% or 100% was considered both extreme and less plausible. In the analysis

accepted by the Committee, the assumption of 50% efficacy of the drugs on fractures associated with other independent clinical risk factors was adjusted by using a correspondingly greater efficacy of the drugs on fractures associated with the key independent clinical risk factors (age, BMD and prior fracture).

- 4.3.15 The Committee considered the assumptions used in the modelling for the side effects of bisphosphonates, in which women who experience bisphosphonate-related side effects had 91% of the utility of women who did not have such side effects. In the base case, this was applied to 2.35% of patients in the first treatment month and 0.35% of patients thereafter. Taking into account the persistence data (sections 4.1.36 and 4.1.37) and the comments received from consultees and commentators that about 25–30% of women experience gastrointestinal side effects when first taking a bisphosphonate, the Committee agreed that it was important to consider the results of a sensitivity analysis assuming that 24% of women were experiencing bisphosphonate-related side effects in the first treatment month and 3.5% of women thereafter.
- 4.3.16 The Committee acknowledged that the modelling made assumptions necessary about the value of a QALY gained that could be considered an acceptable use of NHS resources. The Committee considered that women who have already sustained an osteoporotic fracture live with the pain and distress caused by the fracture. The Committee considered that women with an osteoporotic fracture constitute a different population from the primary prevention population and that there were some factors that justified considering a higher ICER range in line with the 'Guide to the methods of technology appraisal' (see [www.nice.org.uk](http://www.nice.org.uk)).
- 4.3.17 The Committee discussed a number of concerns surrounding other issues that were not represented in the model but which may have had an impact on the cost-effectiveness estimates. These included:

possible long-term adverse effects of bisphosphonates on the formation of new bone; the likelihood that DXA scanning outside a clinical trial environment would not be as effective as in the clinical trials; and the possibility that the proportion of women who experience side effects may exceed the model's base-case assumptions. Finally, the Committee noted that current discount rates used by the Treasury, the Department of Health and NICE result in a cost-effectiveness calculation less favourable to the drugs than the discount rates used in the analysis considered by the Committee. Although a quantitative analysis of the uncertainties surrounding all these issues was not available, the Committee agreed that, for first-line treatment with a bisphosphonate, these uncertainties could be collectively approximated through the sensitivity analysis for side effects (see section 4.3.15). The Committee was persuaded, however, that the results of the sensitivity analysis need only apply to the first-line treatment with a bisphosphonate, because many of the factors that led to the adoption of the sensitivity analysis did not apply for second-line treatment.

**Alendronate**

4.3.18 The Committee considered the results of the economic model following the price reduction for non-proprietary alendronate, the newly included assumptions and the sensitivity analyses (see sections 4.3.9 to 4.3.15). The Committee agreed that, when considering the use of alendronate as a first-line treatment, the sensitivity analysis that captured the uncertainties in the economic model was the most appropriate (see section 4.3.15). This led the Committee to conclude that alendronate (based on the price of £53.56 per year for once-weekly treatment) would be an appropriate use of NHS resources for secondary preventative treatment in postmenopausal women with fragility fractures and confirmed osteoporosis (that is, a T-score of  $-2.5$  SD or below).

The Committee was advised by the clinical specialists from the Guideline Development Group for the NICE clinical guideline on osteoporosis that, in women aged 75 years or older with a prior fracture, a DXA scan may not be required if the responsible clinician considers it to be clinically inappropriate or unfeasible. This is because a very high proportion of these women would be likely to have a T-score of  $-2.5$  SD or below.

- 4.3.19 The Committee noted that the prices of different brands of alendronate vary greatly and concluded that alendronate should be prescribed on the basis of the lowest acquisition cost available.

**Considerations for the other drugs under appraisal**

- 4.3.20 The Committee noted that risedronate, etidronate, raloxifene and strontium ranelate were dominated by alendronate (based on the price of £53.56 per year for alendronate); that is, these drugs have a higher acquisition cost than alendronate, but are not more efficacious. The Committee was also aware that, for women for whom weekly non-proprietary alendronate could be recommended based on cost effectiveness, the ICERs for risedronate and strontium ranelate were very high, even without inclusion of identification costs (see examples in section 4.2.22).

- 4.3.21 The Committee considered an approach where the higher costs of risedronate, strontium ranelate and teriparatide were incorporated into the analysis by combining costs based on the estimated use of alendronate, risedronate and strontium ranelate and teriparatide. However, the overall cost effectiveness of such a combined approach for fracture prevention would be less favourable than that of alendronate. As a consequence, some women who would be eligible for treatment with alendronate under the recommendations in section 1.1 would not be offered treatment using such a combined approach. For this reason, the Committee did not consider the combined approach to be appropriate.



- 4.3.22 The Committee considered treatment options available for a woman who is intolerant to alendronate or unable to comply with instructions for administration despite reasonable measures to support continuation of alendronate treatment. The Committee noted that all other treatment options have higher acquisition costs and/or different effectiveness profiles, which would reduce the cost effectiveness of preventive treatment if these drugs were used. The Committee observed that the identification costs associated with finding women who could be cost-effectively treated with one of the other drugs would be negligible, because they would have already undergone an assessment and had a DXA scan in order to be assessed for first-line treatment with alendronate. Therefore, it agreed that the recommendations for this situation should be based on the modelling that excluded identification costs. The Committee also agreed that, when considering second-line or subsequent treatment, the base-case assumptions for side effects could be applied; that is, a 0.91 utility multiplier should be applied to 2.35% of patients in the first treatment month and 0.35% of patients thereafter.
- 4.3.23 The Committee considered women who cannot take alendronate because of a contraindication or a disability that prevents them from complying with the instructions for administration. Because such a contraindication or disability would be known before the risk assessment, this would comprise a first-line treatment situation, where identification costs are included. Alternative drugs become cost effective at a higher age and lower BMD in a first-line treatment situation, compared with a second-line treatment situation where identification costs are not included. However, such an approach was considered inappropriate by the Committee because it would unfairly disadvantage women who cannot take alendronate because of a contraindication or a disability. Therefore the Committee concluded that women who cannot take alendronate

for these reasons should have access to alternative drugs in the same way as women who cannot tolerate alendronate (that is second-line treatment, where the analysis excluded identification and assessment costs).

#### *Risedronate*

4.3.24 The Committee concluded that risedronate could be recommended for women who are unable to comply with the special instructions for the administration of alendronate, or have a contraindication to or are intolerant of alendronate, and who have a T-score of  $-2.5$  SD or below plus a combination of age and number of independent clinical risk factors for fracture where treatment with risedronate resulted in an ICER of less than £30,000 per QALY gained without the consideration of identification costs, as outlined in section 4.2.25. The Committee agreed that in women aged 75 years or older, where the T-score needed to make treatment cost-effective was  $-2.5$  SD or below, a DXA scan may not be required if the clinician considers it to be clinically inappropriate or unfeasible (see section 4.3.18).

4.3.25 Having reviewed the evidence on independent clinical risk factors for fractures and the views of the clinical specialists, the Committee agreed that the appropriate independent clinical risk factors indicating an increased risk of fracture were: parental history of hip fracture, alcohol intake of 4 or more units per day, and rheumatoid arthritis. The Committee noted that long-term systemic corticosteroid use is also a relevant clinical risk factor; women who are on long-term systemic corticosteroid treatment will be considered within future guidance produced by NICE.

#### *Etidronate*

4.3.26 The Committee considered the cost effectiveness of etidronate and noted that in previous modelling etidronate had a better cost-effectiveness profile than risedronate; since then there has been no

change in the evidence base that would affect the relative position of these two drugs. In view of its concerns surrounding the clinical evidence base for etidronate, and taking into account the views of clinical specialists and consultees, the Committee decided that etidronate should not be recommended in preference to risedronate. However, the Committee agreed that guidance on etidronate should be included in the recommendations, and concluded that it can be recommended as an alternative treatment option for women who cannot take alendronate, as outlined for risedronate in section 4.3.24. In deciding between risedronate and etidronate, clinicians and patients need to balance the overall effectiveness profile of the drugs against their tolerability and adverse effects in individual patients.

#### *Strontium ranelate*

- 4.3.27 The Committee did not accept the estimate of efficacy for strontium ranelate in preventing hip fracture from the post-hoc subgroup analysis, but it accepted the statistically non-significant RR of 0.85 for hip fracture to acknowledge an effect on this important type of fracture.
- 4.3.28 The Committee concluded that strontium ranelate could be recommended for women who are unable to comply with the special instructions for the administration of alendronate and either risedronate or etidronate, or have a contraindication to or are intolerant of alendronate and either risedronate or etidronate, and who have a T-score of  $-2.5$  SD or below plus a combination of age and number of independent clinical risk factors for fracture where treatment with strontium ranelate resulted in an ICER of less than £30,000 per QALY gained, without the consideration of identification costs, as outlined in section 4.2.26. The Committee agreed that in women aged 75 years or older, where the T-score needed to make treatment cost-effective was  $-2.5$  SD or below, a

DXA scan may not be required if the clinician considers it to be clinically inappropriate or unfeasible (see 4.3.18).

- 4.3.29 The Committee agreed a definition of alendronate, risedronate or etidronate intolerance as: persistent upper gastrointestinal disturbance that is sufficiently severe to warrant discontinuation of treatment and that occurs even though the instructions for administration have been followed correctly.

#### *Raloxifene*

- 4.3.30 The Committee discussed the reported benefits of raloxifene on breast cancer risk, and heard from the clinical specialists that the possibility of preventing vertebral fractures and breast cancer simultaneously could be attractive particularly to younger postmenopausal women. The Committee also heard from the specialists that evidence on the effect of raloxifene in reducing cardiovascular risk is not considered to be robust and that there is some concern over the increased risk of VTE (see section 4.1.25).
- 4.3.31 The Committee noted that a higher proportion of the overall benefit associated with raloxifene was attributable to its effect on the prevention of breast cancer than to its effect on the prevention of osteoporotic fragility fractures. The Committee agreed that, in principle, the side effects of using a drug should be considered; however, there were a number of reasons why the Committee considered that the breast cancer benefit should not be the sole factor in deciding whether raloxifene is a cost-effective option for treatment for the secondary prevention of osteoporotic fragility fractures, as follows:
- From the evidence presented, raloxifene was not as effective as the bisphosphonates for treating osteoporosis..
  - Full assessment of raloxifene's effect on the prevention of breast cancer and its cost effectiveness in this indication would require

consideration of how it compares with other drugs that could be used for breast cancer prevention.

4.3.32 The Committee noted that second-line treatment with raloxifene did not result in ICERs lower than £30,000 per QALY gained for women younger than 70 years, and for older women the T-scores at which ICERs were lower than £30,000 per QALY gained were very low. However, the Committee concluded that, the possible benefits in addition to fracture prevention meant that, in cases where women are unable to comply with the special instructions for the administration of alendronate and either risedronate or etidronate, or have contraindications to or are intolerant of alendronate and either risedronate or etidronate, raloxifene could be recommended for the same groups of women for whom treatment with strontium ranelate resulted in an ICER of less than £30,000 per QALY gained without the consideration of identification costs, as outlined in section 4.3.28. The Committee considered that in the younger women in these groups, raloxifene was a plausible choice. When deciding between strontium ranelate and raloxifene, clinicians and patients need to balance the overall proven effectiveness profile of these two drugs against their tolerability and other effects in individual patients.

#### *Teriparatide*

4.3.33 The Committee noted the very high ICER for teriparatide when compared with pooled results for alendronate and risedronate in an analysis carried out by SchARR before the latest price reduction for alendronate, and that there has been no change in the cost effectiveness evidence for teriparatide since. Noting the most recent modelling results for teriparatide, the Committee concluded that a change from the recommendations for teriparatide in NICE technology appraisal 87 for women aged 65 years and older is not warranted. Furthermore, the Committee considered that the

updated modelling indicated that women aged 55–64 years who have a T score of –4 SD or below and more than two fractures could be cost-effectively treated with teriparatide.

### **Women who cannot take alendronate**

4.3.34 The Committee carefully considered the position of women who cannot take alendronate because of a condition which either makes alendronate contraindicated or which prevents individuals from complying with the instructions for administration for alendronate. In doing so the Committee noted that at least some women in this patient group were likely to be 'disabled' as defined by the Disability Discrimination Act 1995. The Committee was aware of its duties under that Act to avoid unlawful discrimination, to have due regard to the need to promote equality of opportunity for disabled people, and the need to take steps to take account of disabled people's disabilities, as well as its broader legal duties to ensure that its guidance is fair and reasonable.

4.3.35 The Committee noted that the drugs other than alendronate are cost effective only for patients at higher risk of fracture than the risk levels at which alendronate is cost effective. If these other drugs are recommended for use by patients who cannot take alendronate, only when those patients meet the criteria at which these alternative drugs become cost effective, these patients will not receive preventative treatment unless they are at higher risk of fracture than the risk levels at which alendronate is recommended. The Committee therefore considered whether, for women who cannot receive alendronate, the other drugs should be recommended at the same risk levels as alendronate (that is using the criteria established as being cost effective for alendronate) in order to provide access to preventative treatment for all patients with the same level of risk. The Committee reviewed the ICERs for risedronate and strontium ranelate within the criteria established to

be cost effective for alendronate. The Committee noted that the prices for risedronate and strontium ranelate are approximately five to six times higher than the price for non-proprietary weekly alendronate, and that the ICERs for these drugs compared with no treatment were very high. For example, the ICER for strontium ranelate for women aged 55–59 years with an independent clinical risk factor for fracture was approximately £55,000 per QALY gained (see section 4.2.22). The Committee noted that strontium ranelate would be the most likely choice to be considered for women who are unable to comply with the instructions for administration of alendronate, because the instructions for administration of alendronate and risedronate are similar. The Committee took the view that recommending drugs other than alendronate using the same criteria as alendronate for women who cannot take alendronate would not be justified in this case because of the very high ICERs for the alternative drugs. In reaching this decision the Committee had regard to the fact that the impact of refusing the more favourable recommendation is that there is no generally recommended preventative treatment for a particular group of patients who are at the lower end of fracture risk for which treatment was considered, but that the alternative drugs are recommended when these patients are at higher risk of fracture.

- 4.3.36 The Committee considered that it is important to maximise the number of patients who are able to take alendronate. Some women will be unable to take alendronate in any circumstances because of contraindication, intolerance or inability to comply with the instructions for administration. However some women who have a disability that makes it difficult for them to comply with the instructions for administration of alendronate would be able to receive the drug if they received assistance in taking it. The Committee concluded that all reasonable steps should be taken to provide women who have a disability that makes it difficult for them

to comply with the instructions for administration of alendronate, with such practical support and assistance with administration (for example through district nurse visits or other home support services), as will enable them to take the drug.

#### **FRAX fracture risk calculation tool**

4.3.37 The Committee was aware of the availability of the FRAX internet-based tool, which can be used to calculate a 10-year absolute risk of fracture, developed under the auspices of the WHO. This assessment tool was based on the same epidemiological data that were used in the Assessment Group's model. However, the Committee was not persuaded that recommendations about treatment should be based on absolute risk as calculated using FRAX. Firstly, the Committee did not agree that all clinical risk factors included in the WHO algorithm were appropriate (see sections 4.2.12 and 4.3.9). Secondly, the Committee was aware that absolute fracture risk is not directly related to cost-effectiveness as outlined in the strontium ranelate assessment report issued in 2005. This is because absolute fracture risk is the total for all fracture sites, but different fracture sites have different impacts on quality of life, costs and mortality. Therefore, cost-effectiveness is dependent on the contribution from each fracture site to the total fracture risk. Thirdly, the Committee had agreed that treatment benefit had not been proven for fracture risk associated with all independent clinical risk factors (section 4.3.9). Therefore, the Committee concluded that using a combination of T-score, age and number of independent clinical risk factors for fracture is more appropriate for defining treatment recommendations in this appraisal

#### **Evidence on use of acid suppressive medication and fracture risk**

4.3.38 The Committee was made aware of data indicating that acid-suppressive medication leads to a small increase in fracture risk



and that co-administration of acid-suppressive medication and bisphosphonates may lead to an increased fracture risk compared with bisphosphonate administration alone. The Committee was not persuaded by this evidence; it noted that the data are observational and have not been reported in full, and are different for different fracture sites and for different acid suppressors. Furthermore, the Committee was informed, during consultation, of analyses showing that acid-suppressive medication given in addition to risedronate did not increase fracture risk. The Committee concluded that caution should be exercised when considering the evidence about co-prescription of acid-suppressive medication and bisphosphonates.

- 4.3.39 The Committee noted sensitivity analyses that included the assumption of an increase in fracture risk for women for whom acid-suppressive medications are co-prescribed (see section 4.2.19). The analysis for treatment strategies did not decrease the T-scores at which the ICERs for alendronate fell below £30,000 to the T-scores established for strategies including strontium ranelate or raloxifene. The Committee also noted that the ICERs for treatment compared with no treatment for an individual woman with a relevant combination of age and T-score were not more favourable for strontium ranelate than for risedronate even if an effect of acid-suppressive medication was assumed. The Committee considered that the evidence for this effect was not sufficiently robust. However, it concluded that the relative positions of alendronate, risedronate and strontium ranelate would remain unchanged even if an effect of acid-suppressive medication was assumed (see section 4.3.35). The Committee therefore concluded that it was not necessary to change its recommendations (section 1) to take account of acid-suppressive medication.

### **Calcium and vitamin D prerequisites for treatment**

4.3.40 The Committee discussed the effect of calcium and vitamin D on the clinical effectiveness of the drugs considered. In the studies that formed the basis of this guidance, all participants were said to have adequate calcium and vitamin D levels. The Committee appreciated that the general population, particularly the elderly population, cannot be assumed to have an adequate dietary intake of calcium and vitamin D. It was also considered important to note that adequate levels (normal serum concentrations) of calcium and vitamin D are needed to ensure optimum effects of the treatments for osteoporosis. The Committee concluded that calcium and/or vitamin D supplementation should be provided to women who receive osteoporosis treatment unless clinicians are confident that the women have an adequate calcium intake and are vitamin D replete. The Committee suggested that the forthcoming NICE clinical guideline ([www.nice.org.uk](http://www.nice.org.uk)) could specify how such assessments should be made and what supplementation should be prescribed.

### **Consultation on the Assessment Group's economic model**

4.3.41 Following the outcome of the judicial review and the court ruling of March 2009, the Appraisal Committee considered the comments received from consultees on the Assessment Group's executable economic model, a report by the DSU reviewing these comments, and responses from the consultees to the DSU report.

4.3.42 The Committee considered the comments from consultees that the Assessment Group's model was not sufficiently transparent, lacked adequate documentation and could not be validated. It noted the number of consultations that took place during the appraisal guidance development, that the consultation documents had included descriptions of the model, and that assumptions and parameter values used had also been provided to consultees. The

Committee was aware that instructions on how to run the model were released with the model and that consultees were able to run the model with changed input parameters. The Committee was satisfied with the exploration by the DSU of the functionality and validity of the model. The Committee noted that Servier stated that it had constructed its own economic model in order to validate the Assessment Group's model and to demonstrate the mathematical rationale to support its comments. The Committee noted that the results from Servier's model were very similar to those from the Assessment Group's model when similar assumptions and parameter inputs were used. The Committee was not persuaded by the consultees' doubt about the validity of the model, particularly since differences between the results obtained using Servier's model and the Assessment Group's model were largely because of differences in the assumptions used.

- 4.3.43 The Committee considered the comments from consultees that some inputs in the Assessment Group's model could not be changed and that it was unclear how fracture risk was calculated. The Committee noted that some of the fixed input parameters were inputs that do not need changing (such as the discount rate and standard mortality rates). Other fixed input values, such as the BMI and issues around the time horizon, were discussed separately (see sections 4.3.46 and 4.3.49 respectively). The Committee concluded that it was reasonable for some inputs in the model to be fixed. The Committee noted that fracture risks were calculated by the Assessment Group using the WHO algorithm in a separate spreadsheet and then entered into the model. It understood that the WHO algorithm itself was provided to the Assessment Group in 2005 as academic in confidence and that at that time NICE did not have permission from the owner of the algorithm to release it to consultees. The Committee understood that although the WHO fracture risk algorithm itself was not embedded in the economic

model, the model could not be released because the algorithm could have been back-calculated from the fracture risks entered in the model and because the numbers of women with risk factors from the algorithm were included in the model.

4.3.44 The Committee considered the comments from consultees that the fracture risks entered into the model, calculated using the WHO algorithm, were different from fracture risks estimated using the FRAX tool. The Committee was aware that some differences could be because of the Assessment Group's use of midpoint ages in each 5-year age grouping. It also heard that the Assessment Group had verified the application of the WHO algorithm as provided in 2005, including all interactions between clinical risk factors, and was satisfied that the DSU had adequately assessed its application as being correct in the model. Because neither the DSU nor NICE has access to the algorithm used for the construction of the FRAX tool, the Committee was not in a position to comment further on differences between the two ways of estimating fracture risk. It concluded that differences between fracture risk estimates produced using the FRAX tool and those used in the Assessment Group's model were not in themselves a reason to doubt the correct use of the WHO algorithm within the Assessment Group's model.

4.3.45 The Committee considered the comments from consultees that mortality associated with clinical risk factors had been omitted from the Assessment Group's model, and noted the confirmation from the DSU that this was the case. It was persuaded that the inclusion of such additional mortality effects would increase the complexity of the model, and may increase the ICERs for the treatment of women with such clinical risk factors, but decrease the ICERs for the treatment of women without such risk factors. The Committee agreed that the overall effect of including mortality associated with

clinical risk factors in the model was unlikely to lead to a marked change the overall results.

- 4.3.46 The Committee reviewed the consultee comments relating to the fixed BMI value of 26 kg/m<sup>2</sup> used in the Assessment Group's model. It noted the rationale for selecting this value (see section 4.2.46). It also noted that in the DSU's exploratory analysis using the WHO algorithm, no increase in fracture risk was identified for women with a higher or lower BMI when BMD was known. The Committee was aware of its recommendation to assess BMD in all women under the age of 75 years for whom treatment is being considered, and noted that BMI is a weak predictor of fracture when BMD is known. Therefore the Committee concluded that the use of a fixed BMI value of 26 kg/m<sup>2</sup> did not lead to an unfavourable assessment of the cost effectiveness of the interventions.
- 4.3.47 The Committee considered comments from consultees that the fracture risk associated with alcohol consumption used in the model was incorrect. It noted that the DSU had determined that the WHO algorithm had been correctly implemented, and understood that alcohol consumption of more than 2 units per day was included as a risk factor in the model. The Committee also noted that in its recommendations it had chosen to use a higher level of alcohol consumption in the risk identification strategy, because only alcohol consumption of 4 or more units per day was identified as a statistically significant risk factor for fracture for women – the population considered in the guidance. The Committee also considered a consultee comment that stated that it was unclear whether smoking and corticosteroid use had been included in the model as risk factors. It noted that the DSU had determined that the WHO algorithm had been correctly implemented with regard to both smoking and corticosteroid use in the model. The Committee noted

that assessment and treatment of corticosteroid-induced osteoporosis will be covered in the NICE clinical guideline on osteoporosis, and that the effect of smoking in women was not statistically significant when assessing risk of osteoporotic fractures taken as a whole. The Committee was therefore satisfied that risks associated with corticosteroids, smoking and alcohol consumption had been faithfully applied in the Assessment Group's model, and agreed that the levels of alcohol consumption and smoking that should be used in the risk identification strategy were a matter for the Committee to consider and determine. The Committee took the view that it is not appropriate to identify women at high risk of fracture on the basis of risks that were not statistically significant (such as smoking and consumption of fewer than 4 units of alcohol per day) and that, in addition, the impact of these risk factors could arguably be approached by a strategy of smoking cessation and reducing alcohol consumption. The Committee noted comments from the consultees that the Assessment Group's model should have been amended to reflect the Committee's agreed inclusion of risk factors. However, the Committee took the pragmatic view that such amendments would have added unnecessarily to the mathematical complexity of an already complex clinical situation. It noted that women who had taken corticosteroids were included in the model and therefore contributed to the underlying fracture risk, with the effect of reducing the ICERs for the treatment of the population of women considered in the recommendations.

- 4.3.48 The Committee considered consultee comments that giving equal weighting to different clinical risk factors for fracture in the Assessment Group's model was inaccurate. The Committee considered the complex results presented originally in the 2005 Strontium Ranelate Assessment Report related to the inclusion of different risk factors and combinations of risk factors. The Committee noted that it had previously agreed that a clinically

workable risk identification and treatment strategy should include the grouping of risk factors as the only practical way forward. At the time of the model's development, no individual risk calculation tool was available. Even if such a tool had been used in the development of the guidance, the prediction of cost effectiveness from overall absolute fracture risk alone, as suggested by consultees, would not be appropriate, for two reasons. Firstly, risk factors have different effects on different fracture types, and the cost effectiveness of treatment depends on the relative contributions of each risk factor to fracture risk. Secondly, the effectiveness of the drugs in reducing fracture risk was limited to only some of the clinical risk factors (age, T-score of  $-2.5$  SD or below and prior fracture). The Committee heard from the DSU that there was considerable uncertainty about the cost effectiveness of treating women based on absolute risk alone (see section 4.3.37). Therefore, the Committee concluded that, when developing the guidance, simplification of the model was justified in order to produce workable recommendations.

- 4.3.49 The Committee reviewed a comment from a consultee that the methods used to model effects beyond 10 years were not adequately described. It noted that the DSU confirmed that consequences beyond 10 years were considered in the Assessment Group's model, and an expanded description of the methods used was provided in an annex to the DSU report. The Committee also noted that the DSU carried out a sensitivity analysis in order to establish the impact of any possible underestimation of the mortality risk after the 10-year time horizon. It noted that doubling of the mortality risk led to only very small changes in the results. The Committee therefore concluded that mortality effects beyond the 10-year time horizon had been reasonably accounted for in the model and that sufficient

description of these methods had been made available to consultees.

- 4.3.50 The Committee considered comments from consultees that the population data on the distribution of BMD (T-score) were not appropriate. It noted the DSU response confirming that the UK epidemiological dataset from the Holt study had been correctly implemented in the Assessment Group's model, and that the assumptions about the normality of the distributions used were likely to favour treatment for women at risk of fracture. The Committee also noted that the particular UK epidemiological dataset used in the model had been originally suggested by consultees for this appraisal. The Committee concluded that the population data had been used appropriately in the model.
- 4.3.51 The Committee considered a comment from a consultee that using a single estimate of cost effectiveness for 5-year age groupings of women at risk of fracture could exclude women from being offered treatment. It noted that this identification method was a Committee decision, and that identification strategies based on other factors could make treatments less cost effective.
- 4.3.52 The Committee reviewed comments from a consultee that the application of the same disutility for the side effects associated with strontium ranelate and bisphosphonates was not correct, as the side effects of strontium ranelate are different from those of the bisphosphonates. The Committee was aware that the side effects observed for strontium ranelate in the clinical trials did not include gastrointestinal effects, but did include an increased risk of VTE. Because the increased risk of VTE was not included in the Assessment Group's model, the Committee had agreed that it was appropriate to include a disutility equivalent to the bisphosphonate base-case side-effect disutility to take account of this adverse effect.



- 4.3.53 The Committee reviewed comments from consultees about model assumptions or inputs that the Committee had directed the Assessment Group to use. It noted that issues such as treatment compliance, discount rates, costs of fracture, utility values for vertebral fracture and side-effect profiles used in the model had been considered and agreed by the Committee and reported in the guidance. The Committee also agreed that it had considered identification strategies for women at risk of fracture and, noting the advice of clinical specialists, it had recommended that women should have their BMD assessed by DXA scanning, except in certain circumstances as defined in the guidance. The Committee concluded that views expressed by consultees on the choice of modelling assumptions, input parameters and risk identification strategy were not about the operation of the Assessment Group's model, but were about Committee decisions that had already been discussed during development of the guidance.
- 4.3.54 The Committee also considered the consultees' view that the FRAX tool provides a 'mechanism to compute cost-effectiveness' according to clinical risk factors and that each of the current recommendations covers a wide range of absolute risk values, depending on the individual risk factors involved. The Committee understood that the FRAX tool is not an economic model, but a tool to estimate fracture risk. The Committee acknowledged that the current set of recommendations involved necessary simplifications from the more complex algorithm used to develop the Assessment Group's model. It was also aware that a direct prediction of cost effectiveness from absolute fracture risk alone would be inappropriate (see sections 4.3.37 and 4.3.48).
- 4.3.55 The Committee concluded that the Assessment Group had provided an executable economic model and had implemented the WHO algorithm (as supplied) correctly. The Committee agreed with

the DSU's comments that alterations to the modelling approach, as suggested by consultees, would not lead to significant improvements in the cost effectiveness of treatment for women at risk of fracture. The Committee confirmed that the model provided a suitable framework to allow it to make recommendations on the cost-effective use of treatment for women at risk of fracture. The Committee noted that assumptions used in the Assessment Group's model had been considered and agreed by the Committee in developing the guidance. It agreed that it would not be useful to request further analysis from the Assessment Group at this stage. The Committee further agreed that any exploration of how absolute fracture risk could be used in making treatment decisions would require a new assessment and appraisal. Therefore the Committee concluded that the recommendations based on the Assessment Group's model were appropriate, and that the recommendations should remain unchanged.

- 4.3.56 The Committee noted the comments from some consultees that the guidance should be reviewed soon because the price of some of the appraised drugs had changed. The Committee noted that any possible price reductions could be offset by the use of the currently applicable discount rate, and that any review should also take into consideration how NICE might assess diagnostic tools such as absolute fracture risk prediction tools.

## **5 Implementation**

- 5.1 The Secretary of State and the Welsh Assembly Minister for Health and Social Services have issued directions to the NHS on implementing NICE technology appraisal guidance. When a NICE technology appraisal recommends use of a drug or treatment, or other technology, the NHS must provide funding and resources for it within 3 months of the guidance being published. If the

Department of Health issues a variation to the 3-month funding

direction, details will be available on the NICE website. The NHS is not required to fund treatments that are not recommended by NICE.

5.2 NICE has developed tools to help organisations put this guidance into practice (listed below). These are available on our website ([www.nice.org.uk/TA161](http://www.nice.org.uk/TA161)).

- Slides highlighting key messages for local discussion.
- Costing report and costing template to estimate the savings and costs associated with implementation.
- Audit support for monitoring local practice.

## **6 Recommendations for further research**

6.1 Given the evidence that the benefits of one of the bisphosphonates (alendronate) may continue for several years after the end of treatment, the Committee recommends that research should be carried out to define the optimal duration of treatment with individual bisphosphonates.

6.2 The Committee recommends research into the long-term effects of bisphosphonates on bone quality, given the inhibitory effects on bone resorption of these drugs.

## **7 Related NICE guidance**

### **Published**

- Alendronate, etidronate, risedronate, raloxifene and strontium ranelate for the primary prevention of osteoporotic fragility fractures in postmenopausal women. NICE technology appraisal guidance 160 (2008). Available from: [www.nice.org.uk/TA160](http://www.nice.org.uk/TA160) (Note that an amended final appraisal determination for this guidance is being issued in November 2009.)

## **Under development**

NICE is developing the following guidance (details available from [www.nice.org.uk](http://www.nice.org.uk)):

- Denosumab for the prevention of osteoporotic fractures in postmenopausal women. NICE technology appraisal guidance (publication date to be confirmed).
- Osteoporosis: assessment of fracture risk and the prevention of osteoporotic fractures in individuals at high risk. NICE clinical guideline (publication date to be confirmed).

## **8 Review of guidance**

- 8.1 The guidance on these technologies will be considered for review by the Guidance Executive in July 2010. The Guidance Executive will decide whether the technology should be reviewed based on information gathered by NICE, and in consultation with consultees and commentators.

Andrew Stevens  
Chair, Appraisal Committee  
November 2009

## **Appendix A: Appraisal Committee members, guideline representatives and NICE project team**

### **A *Appraisal Committee members***

The Appraisal Committee is one of NICE's standing advisory committees. Its members are appointed for a 3-year term. A list of the Committee members who took part in the discussions for this appraisal appears below. The Appraisal Committee meets three times a month except in December, when there are no meetings. There are four Appraisal Committees, each with a chair and vice chair. Each Committee considers its own list of technologies, and ongoing topics are not moved between the branches.

Committee members are asked to declare any interests in the technology to be appraised. If it is considered there is a conflict of interest, the member is excluded from participating further in that appraisal.

The minutes of each Appraisal Committee meeting, which include the names of the members who attended and their declarations of interests, are posted on the NICE website.

#### **Dr Amanda Adler (2009)**

Consultant Physician, Addenbrooke's Hospital

#### **Professor Keith Abrams (2006–2009)**

Professor of Medical Statistics, University of Leicester

#### **Ms Julie Acred (2004–2005)**

Chief Executive, Derby Hospitals NHS Foundation Trust

#### **Dr Ray Armstrong (2008)**

Consultant Rheumatologist, Southampton General Hospital

#### **Dr Jeff Aronson (2006–2009)**

Reader in Clinical Pharmacology, University Department of Primary Health Care, University of Oxford

**Dr Darren Ashcroft (2004–2008)**

Senior Clinical Lecturer, School of Pharmacy and Pharmaceutical Sciences,  
University of Manchester

**Professor David Barnett (2004–2008)**

Professor of Clinical Pharmacology, University of Leicester

**Dr Peter Barry (2004–2009)**

Consultant in Paediatric Intensive Care, Leicester Royal Infirmary

**Dr Michael Boscoe (2009)**

Consultant Cardiothoracic Anaesthetist, Royal Brompton and Harefield NHS  
Foundation Trust

**Professor Stirling Bryan (2006–2008)**

Head, Department of Health Economics, University of Birmingham

**Mr Brian Buckley (2004–2006)**

Vice Chairman, InContact

**Professor John Cairns (2006–2009)**

Public Health and Policy, London School of Hygiene and Tropical Medicine

**Professor David Chadwick (2005–2006)**

Professor of Neurology, Walton Centre for Neurology and Neurosurgery

**Dr Peter I Clark (2004–2006)**

Honorary Chairman, Association of Cancer Physicians

**Ms Donna Covey (2004–2005)**

Chief Executive, Asthma UK

**Dr Mike Davies (2004–2008)**

Consultant Physician, University Department of Medicine and Metabolism,  
Manchester Royal Infirmary

**Mr Richard Devereaux-Phillips (2004–2006)**

Public Affairs Manager, Medtronic Ltd

**Professor Jack Dowie (2004–2009)**

Health Economist, London School of Hygiene and Tropical Medicine

**Lynn Field (2006–2008)**

Nurse Director, Pan Birmingham Cancer Network

National Institute for Health and Clinical Excellence

**Professor Gary A Ford (2004–2005)**

Professor of Pharmacology of Old Age/Consultant Physician, Royal Victoria Infirmary, Newcastle upon Tyne

**Professor Christopher Fowler (2006–2008)**

Professor of Surgical Education, Barts and The London School of Medicine and Dentistry, Queen Mary, University of London

**Dr Fergus Gleeson (2004–2009)**

Consultant Radiologist, Churchill Hospital, Oxford

**Ms Sally Gooch (2004–2009)**

Independent Nursing and Healthcare Consultant

**Professor Trisha Greenhalgh (2004–2005)**

Professor of Primary Health Care, University College London

**Mrs Barbara Greggains (2006–2008)**

Lay member

**Mrs Eleanor Grey (2009)**

Lay member

**Mr Sanjay Gupta (2005–2008)**

Former Service Manager in Stroke, Gastroenterology, Diabetes and Endocrinology, Basildon and Thurrock University Hospitals Foundation NHS Trust

**Ms Linda Hands (2004–2005)**

Consultant Vascular Surgeon, John Radcliffe Hospital, Oxford

**Professor Philip Home (2005–2006)**

Professor of Diabetes Medicine, University of Newcastle upon Tyne

**Dr Neil Iosson (2009)**

General Practitioner

**Dr Peter Jackson (2005–2006)**

Clinical Pharmacologist, University of Sheffield

**Professor Peter Jones (2004–2006)**

Professor of Statistics and Dean, Faculty of Natural Science, Keele University

**Professor Robert Kerwin (2004–2005)**

Professor of Psychiatry and Clinical Pharmacology, Institute of Psychiatry, London

**Dr Mike Laker (2005–2007)**

Medical Director, Newcastle Hospitals NHS Trust

**Ms Joy Leavesley (2004)**

Senior Clinical Governance Manager, Whittington Hospital

**Dr Ruth Lesirge (2004)**

Lay member

**Ms Rachel Lewis (2004–2006)**

Nurse Adviser to the Department of Health

**Mr Terence Lewis (2006–2009)**

Lay member

**Dr George Levvy (2005–2006)**

Lay member

**Professor Gary McVeigh (2006–2008)**

Professor of Cardiovascular Medicine, Queens University, Belfast

**Professor Jonathan Michaels (2004–2006)**

Professor of Vascular Surgery, University of Sheffield

**Dr Ruairidh Milne (2004–2009)**

Senior Lecturer in Health Technology Assessment, National Coordinating Centre for Health Technology, University of Southampton

**Dr Neil Milner (2004–2009)**

General Practitioner, Tramways Medical Centre, Sheffield

**Dr Rubin Minhas (2004–2009)**

General Practitioner, CHD Clinical Lead, Medway PCT

**Mr Stephen Palmer (2009)**

Senior Research Fellow, Centre for Health Economics, University of York

**Dr John Pounsford (2006–2009)**

Consultant Physician, Frenchay Hospital, Bristol



**Mr Philip Pugh (2009)**

Strategic Development Lead for HCAI & AMR, Health Protection Agency

**Dr Rosalind Ramsay (2006–2008)**

Consultant Psychiatrist, Adult Mental Health Services, Maudsley Hospital, London

**Dr Christa Roberts (2006–2008)**

UK Country Manager, Abbott Vascular

**Dr John Rodriguez (2009)**

Assistant Director of Public Health, NHS Eastern and Coastal Kent

**Dr Florian Alexander Ruths (2009)**

Consultant Psychiatrist and Cognitive Therapist, Maudsley Hospital, London

**Dr Stephen Saltissi (2006–2009)**

Consultant Cardiologist, Royal Liverpool University Hospital

**Mr Miles Scott (2004–2006)**

Chief Executive, Bradford Teaching Hospitals NHS Foundation Trust

**Mr Navin Sewak (2009)**

Primary Care Pharmacist, NHS Hammersmith and Fulham

**Dr Lindsay Smith (2005–2009)**

General Practitioner, East Somerset Research Consortium

**Mr Roderick Smith (2006–2009)**

Finance Director, West Kent PCT

**Mr Cliff Snelling (2006–2009)**

Lay member

**Mr Malcolm Stamp (2004)**

Chief Executive, Addenbrooke's NHS Trust

**Professor Ken Stein (2004–2009)**

Professor of Public Health, Peninsula College of Medicine and Dentistry, University of Exeter

**Professor Andrew Stevens (Chair) (2004–2009)**

Professor of Public Health, University of Birmingham

**Dr Rod Taylor (2006–2009)**

Associate Professor in Health Services Research, Peninsula Medical School, Universities of Exeter & Plymouth

***B Guideline representatives***

The following individuals, representing the Guideline Development Group responsible for developing NICE's clinical guideline related to this topic, were invited to attend Appraisal Committee meetings to observe and to contribute as advisers to the Committee.

- Professor Cameron G Swift (2008–2009), King's College London School of Medicine Clinical Age Research Unit King's College Hospital, London
- Dr Maggie Westby (2008–2009), Senior Research and Development Fellow, National Clinical Guideline Centre for Acute and Chronic Conditions
- Professor Juliet Compston (2005-2007), Professor of Bone Medicine, University of Cambridge School of Clinical Medicine and Addenbrooke's NHS Trust
- Dr Peter Selby (2005–2007), Consultant Physician, Central Manchester and Manchester Children's University Hospitals NHS Trust
- Professor David Barlow (2005–2007), Executive Dean of Medicine, University of Glasgow

***C NICE project team***

Each technology appraisal is assigned to a team consisting of one or more health technology analysts (who act as technical leads for the appraisal), a technical adviser and a project manager.

**Dr Ruairaidh Hill**

Technical Lead

**Fiona Rinaldi**

Technical Adviser

**Jeremy Powell**

Project Manager

## Appendix B: Sources of evidence considered by the Committee

A Extra analysis reports were prepared by the Decision Support Unit, the School of Health and Related Research, University of Sheffield (ScHARR).

- Stevenson M, Wailoo A. A review of comments submitted by consultees on the economic model, August 2009.
- Stevenson M. Analyses of cost-effective BMD scanning and treatment strategies for generic alendronate, and the cost-effectiveness of risedronate and strontium ranelate in those people who would be treated with generic alendronate, February 2008.
- Lloyd Jones M. Critique of evidence put forward by Servier suggesting an association between acid-suppressive medication and fracture risk, February 2008.

B The assessment reports for this appraisal were prepared by the School of Health and Related Research, University of Sheffield, (ScHARR).

- Stevenson M. Analyses of cost-effective BMD scanning and treatment strategies for generic alendronate, and the cost-effectiveness of risedronate and strontium ranelate in those people who would be treated with generic alendronate, February 2008.
- Lloyd Jones M. Critique of evidence put forward by Servier suggesting an association between acid-suppressive medication and fracture risk, February 2008.
- Stevenson M. Analyses of cost-effective BMD scanning and treatment strategies for generic alendronate, risedronate, strontium ranelate, raloxifene and teriparatide following corrections to the methodology associated with lower efficacy in some risk factors, November 2006.
- Stevenson M, Davis S. Addendum to the assessment report: analyses of the cost-effectiveness of pooled alendronate and risedronate, compared with strontium ranelate, raloxifene, etidronate and teriparatide, September 2006.
- Stevenson M, Lloyd Jones M, Davis S et al. Analyses of the cost-effectiveness of pooled alendronate and risedronate, compared with strontium ranelate, raloxifene, etidronate and teriparatide, July 2006.

- Lloyd Jones M, Wilkinson A. Adverse effects and persistence with therapy in patients taking oral alendronate, etidronate or risedronate: systematic reviews, July 2006.
- Stevenson M, Davis S, Lloyd Jones M et al. The clinical effectiveness and cost effectiveness of strontium ranelate for the prevention of osteoporotic fragility fractures in postmenopausal women, July 2005.
- Stevenson M, Davis S. Addendum to the assessment report: the clinical effectiveness and cost effectiveness of technologies for the primary prevention of osteoporotic fragility fractures in postmenopausal women, July 2005.
- Stevenson M, Lloyd Jones M, de Nigris E et al. The clinical effectiveness and cost effectiveness of prevention and treatment of osteoporosis, December 2003.

C The following organisations accepted the invitation to participate in this appraisal as consultees and commentators. They were invited to comment on the draft scope, assessment report and the appraisal consultation document (ACD). They were also invited to comment on the Assessment Group's economic model and on the 2009 DSU report. Organisations listed in I and II were also invited to make written submissions and have the opportunity to appeal against the final appraisal determination.

I Manufacturer/sponsors:

- Alliance for Better Bone Health
- Eli Lilly & Company
- Merck Sharp & Dohme
- Proctor & Gamble UK
- Servier Laboratories
- Teva UK

II Professional/specialist and patient/carer groups:

- Arthritis and Musculoskeletal Alliance
- Bone Research Society (formerly Bone and Tooth Society)
- British Geriatrics Society
- British Menopause Society
- British Orthopaedic Association
- British Society for Rheumatology
- Department of Health
- Institute for Ageing and Health
- National Osteoporosis Society

- National Rheumatoid Arthritis Society
- Primary Care Rheumatology Society
- RADAR (The Royal Association for Disability and Rehabilitation)
- Royal College of General Practitioners
- Royal College of Nursing
- Royal College of Pathologists
- Royal College of Physicians
- Society for Endocrinology
- Southwark Primary Care Trust
- The Society and The College of Radiographers
- Women's Health
- Women's Health Concern
- Women's Nutritional Advisory Service

III Commentator organisations (without the right of appeal)

- British National Formulary
- National Collaborating Centre for Nursing and Supportive Care
- NHS Quality Improvement Scotland
- Novartis Pharmaceuticals UK
- Research Institute for the Care of the Elderly
- Strakan Group
- Roche Products
- Nycomed UK
- Welsh Assembly Government

D The following individuals were selected from clinical specialist and patient expert nominations from the non-manufacturer/sponsor consultees and commentators. They participated in the Appraisal Committee discussions and provided evidence to inform the Appraisal Committee's deliberations. They gave their expert personal view on alendronate, etidronate, risedronate, raloxifene, strontium ranelate and teriparatide for the secondary prevention of osteoporotic fragility fractures in postmenopausal women by attending the initial Committee discussion and/or providing written evidence to the Committee.

- Mrs Jackie Parrington, Deputy Chief Executive, National Osteoporosis Society, nominated by the National Osteoporosis Society – patient expert
- Mrs Anthea Franks, nominated by the National Osteoporosis Society – patient expert

- Professor Juliet Compston, Professor of Bone Medicine, University of Cambridge School of Clinical Medicine and Addenbrooke's NHS Trust, nominated by the Royal College of Physicians – clinical specialist
- Dr RM Francis, Reader in Medicine (Geriatrics) and Honorary Consultant Physician, British Geriatrics Society, nominated by the British Geriatrics Society and the National Osteoporosis Society – clinical specialist
- Dr Caje Moniz, Consultant and Clinical Director, King's Healthcare NHS Trust, nominated by the National Osteoporosis Society – clinical specialist
- Dr Peter Selby, Consultant Physician, Central Manchester and Manchester Children's University Hospitals NHS Trust, nominated by the Society of Endocrinology and the National Osteoporosis Society – clinical specialist